

CATALAN MATROIDS ARE EHRHART POSITIVE

YIMING CHEN, YAO LI, AND MING YAO

ABSTRACT. We show that the Ehrhart polynomials of (a, b) -Catalan matroids are positive combinations of the Ehrhart polynomials of uniform matroids. Since Ferroni has shown that uniform matroids are Ehrhart positive, we obtain the Ehrhart positivity of (a, b) -Catalan matroids.

1 Introduction

Catalan matroids are a subfamily of Schubert matroids, first studied by Ardila [1], and independently by Bonin, de Mier, and Noy [2]. Bonin, de Mier, and Noy [2] also introduced (a, b) -Catalan matroids, which reduce to Catalan matroids when $a = b = 1$. In [5], Fan and Li provided a formula for the Ehrhart polynomials of Schubert matroids. As an application, they obtained a recursive formula for the Ehrhart polynomials of (a, b) -Catalan matroids, and conjectured that (a, b) -Catalan matroids are Ehrhart positive.

Although Ferroni [10] showed that not all matroids are Ehrhart positive, disproving conjectures of De Loera, Haws and Köppe [15] and Castillo and Liu [3], various specific subclasses of matroids turn out to be Ehrhart positive. For instance, uniform matroids, minimal matroids, sparse paving Schubert matroids, panhandle matroids, and all matroids of rank two are Ehrhart positive, see [4, 5, 8, 9, 11]. In this note, we express the Ehrhart polynomials of (a, b) -Catalan matroids as positive combinations of Ehrhart polynomials of uniform matroids, and thus obtain the Ehrhart positivity of (a, b) -Catalan matroids.

Suppose that $M = ([n], \mathcal{B})$ is a matroid with ground set $[n] := \{1, 2, \dots, n\}$ and base set \mathcal{B} . The matroid polytope $\mathcal{P}(M)$ of M is the convex hull

$$\mathcal{P}(M) = \text{conv}\{e_B : B \in \mathcal{B}\}, \quad (1.1)$$

where $e_B = e_{b_1} + \dots + e_{b_k}$ for $B = \{b_1, \dots, b_k\} \subseteq [n]$, and e_i is the i -th standard basis of \mathbb{R}^n . Given a polytope \mathcal{P} and a positive integer t , the t -dilation of \mathcal{P} is defined as $t\mathcal{P} = \{t\alpha : \alpha \in \mathcal{P}\}$. Let $i(\mathcal{P}, t) = |t\mathcal{P} \cap \mathbb{Z}^n|$ denote the number of lattice points in $t\mathcal{P}$. For integral polytopes, $i(\mathcal{P}, t)$ is a polynomial in t , called the Ehrhart polynomial of \mathcal{P} . For simplicity, we write $i(M, t)$ for $i(\mathcal{P}(M), t)$.

For $a, b \geq 1$, let $U_n^{a,b} := U_{nb, n(a+b)}$ denote the uniform matroid on $[n(a+b)]$ with bases $\{B : B \subseteq [n(a+b)], |B| = nb\}$ and $C_n^{a,b}$ be the (a, b) -Catalan matroid on $[n(a+b)]$. When $a = b = 1$, write $U_n := U_{n, 2n}^{1,1}$ and $C_n := C_n^{1,1}$ for short.

Recall that a sequence of positive integers $\lambda = (\lambda_1, \dots, \lambda_k)$ is called a partition of n , denoted as $\lambda \vdash n$, if $\lambda_1 \geq \dots \geq \lambda_k \geq 1$ and $\lambda_1 + \dots + \lambda_k = n$. Let

$$z_\lambda = \prod_{i=1}^n i^{\alpha_i} \alpha_i!,$$

where α_i is the number of appearances of i in λ . A sequence of positive integers $\gamma = (\gamma_1, \dots, \gamma_k)$ is called a composition of n , if $\gamma_1 + \dots + \gamma_k = n$. Define

$$i(U_\gamma^{a,b}, t) := \prod_{i=1}^k i(U_{\gamma_i}^{a,b}, t).$$

Theorem 1.1. *We have*

$$i(C_n, t) = \sum_{\lambda \vdash n} \frac{1}{z_\lambda} i(U_\lambda, t). \quad (1.2)$$

Moreover, we can also obtain the Ehrhart polynomials of (a, b) -Catalan matroids.

Corollary 1.2. *Let a, b be positive integers, we have*

$$i(C_n^{a,b}, t) = \sum_{\lambda \vdash n} \frac{1}{z_\lambda} i(U_\lambda^{a,b}, t). \quad (1.3)$$

As will be seen, the proofs of Theorem 1.1 and Corollary 1.2 are almost the same, for notational simplicity, we just give a proof of Theorem 1.1.

Remark 1. *Since our study of the Ehrhart polynomial of Catalan matroids is based on the subdivision of matroid polytopes, it follows that for any valuative invariant f (e.g., the Tutte polynomial, the Kazhdan–Lusztig polynomial, or Whitney polynomial and so on, see [12]), we have*

$$f(C_n^{a,b}, t) = \sum_{\lambda \vdash n} \frac{1}{z_\lambda} f(U_\lambda^{a,b}, t).$$

The following corollary follows easily from Theorem 1.1.

Corollary 1.3. *The volume of $\mathcal{P}(C_n)$ is given by*

$$\text{vol}(\mathcal{P}(C_n)) = \frac{1}{n(2n-1)!} A(2n-1, n-1), \quad (1.4)$$

where $A(n, k)$ denotes the Eulerian number, which counts the number of permutations on $[n]$ with exactly $k-1$ descents. Similarly, the volume of $\mathcal{P}(C_n^{a,b})$ is given by

$$\text{vol}(\mathcal{P}(C_n^{a,b})) = \frac{1}{n(2n-1)!} A(n(a+b)-1, nb). \quad (1.5)$$

2 Preliminaries

A *matroid* is a pair $M = (E, \mathcal{I})$ consisting of a finite set E , and a collection \mathcal{I} of subsets of E , satisfies: (i) $\emptyset \in \mathcal{I}$; (ii) If $J \in \mathcal{I}$ and $I \subseteq J$, then $I \in \mathcal{I}$; and (iii) If $I, J \in \mathcal{I}$ and $|I| < |J|$, then there exists $j \in J \setminus I$ such that $I \cup \{j\} \in \mathcal{I}$. By (ii), a matroid M is determined by the collection \mathcal{B} of maximal independent sets, called the bases of M . By (iii), all the bases have the same cardinality, called the rank of M , denoted as $\text{rk}(M)$. So we can write $M = (E, \mathcal{B})$. The *dual* of M is a matroid $M^* = (E, \mathcal{B}^*)$ where $\mathcal{B}^* = \{E \setminus B : B \in \mathcal{B}(M)\}$. The *rank function* $\text{rk}_M : 2^E \rightarrow \mathbb{Z}$ of M is defined as

$$\text{rk}_M(T) = \max\{|T \cap B| : B \in \mathcal{B}\}, \quad \text{for } T \subseteq E.$$

The *closure* of a set $A \subseteq E$ is the biggest set that contains A and has the same rank. A set is a *flat* if its closure is same as itself. A set $A \subseteq E$ is called *separable* if there are disjoint subsets R and T such that $A = R \cup T$ and $\text{rk}_M(A) = \text{rk}_M(R) + \text{rk}_M(T)$. Otherwise, A is called *inseparable*.

Let $[n] := \{1, 2, \dots, n\}$ and S be a subset of $[n]$. The *Schubert matroid* $\text{SM}_n(S)$ is the matroid with ground set $[n]$ and bases

$$\{T \subseteq [n] : T \leq S\},$$

where $T \leq S$ means that: $|T| = |S|$ and the i -th smallest element of T is less than or equal to that of S for $1 \leq i \leq |T|$. The *indicator vector* of S is the 0-1 vector $\mathbb{I}(S) = (i_1, \dots, i_n)$, where $i_j = 1$ if $j \in S$, and 0 otherwise. Without loss of generality, we can assume that $1 \notin S$ and the largest element of S is n . Thus $i_1 = 0$ and $i_n = 1$. For simplicity, write $\mathbb{I}(S) = (0^{r_1}, 1^{r_2}, \dots, 0^{r_{2m-1}}, 1^{r_{2m}})$, where 0^{r_1} represents r_1 copies of 0's, 1^{r_2} represents r_2 copies of 1's, etc. Then S can be written as a sequence of positive integers $r(S) = (r_1, r_2, \dots, r_{2m})$ of length $2m$. It is easy to see that given such an integer

sequence r , there is a unique set S whose indicator vector $\mathbb{I}(S)$ can be written in this way. For example, let $S = \{2, 5, 7, 8, 9\} \subseteq [9]$, then $\mathbb{I}(S) = (0, 1, 0^2, 1, 0, 1^3)$ and $r(S) = (1, 1, 2, 1, 1, 3)$.

Let $r(S) = (n - k, k), (1, k - 1, n - k - 1, 1), (k - 1, 1, 1, n - k - 1)$, we obtain the uniform matroids $U_{k,n}$ [8], the minimal matroids $T_{k,n}$ [9], and the sparse paving Schubert matroids $\mathbf{Sp}_{k,n}$ [5], respectively. For any positive integers a, b, c, d , $r(S) = (a, b, c, d)$ corresponds to the notched rectangle matroids [5], which are conjectured to be Ehrhart positive. In particular, $r(S) = (a, b, c, 1)$ corresponds to the

panhandle matroids $\mathbf{Pan}_{b+1, a+b, n}$ [4, 14], where $n = a + b + c + 1$. Moreover, let $r(S) = \overbrace{(a, b, a, b, \dots, a, b)}^{2n}$, we obtain the (a, b) -Catalan matroids $C_n^{a,b}$ introduced by Bonin, de Mier and Noy [2, Definition 3.7]. Notice that $C_n^{a,b}$ is a matroid on $[n(a + b)]$. When $a = b = 1$, $C_n^{1,1}$ is the n -th Catalan matroid C_n as first studied by Ardila [1].

It is well known that the matroid polytope $\mathcal{P}(M)$ defined in (1.1) is a generalized permutohedron parametrized by the rank function of M , see, for example, Fink, Mészáros and St. Dizier [13]. More precisely,

$$\mathcal{P}(M) = \left\{ x \in \mathbb{R}_{\geq 0}^n : \sum_{i \in [n]} x_i = \mathbf{rk}_M([n]) \text{ and } \sum_{i \in T} x_i \leq \mathbf{rk}_M(T) \text{ for } T \subsetneq [n] \right\}. \quad (2.1)$$

Feichtner and Sturmfels [7] provided a minimal system of inequalities for all matroid polytopes (2.1). In order to give a minimal system of inequalities for Schubert matroid polytopes, we recall the combinatorial algorithm of computing the rank function $\mathbf{rk}_S(T)$ of a Schubert matroid $\mathbf{SM}_n(S)$ provided by Fan and Guo [6, Theorem 3.3]. Given a subset $S = \{s_1, \dots, s_k\}$ on $[n]$, view S as $n \times 1$ grids such that there is a box in row j for $j \in S$, and the other rows are empty. The left diagram in Figure 2.1 is an illustration for $S = \{2, 4, 5, 6, 8\} \subseteq [8]$.

For a permutation $w = w_1 w_2 \dots w_t$ of elements of $T \subseteq [n]$, define $\mathcal{F}_S(w)$ to be the filling of S with w as follows. For j from 1 to t , put w_j into the first (from top to bottom) empty box whose row index is larger than or equal to w_j . If there are no such empty boxes, then w_j does not appear in the filling and skip to w_{j+1} . Let $|\mathcal{F}_S(w)|$ denote the number of non-empty boxes of S after filling the elements of w . The right diagram in Figure 2.1 is the filling $\mathcal{F}_S(w)$ of $S = \{2, 4, 5, 6, 8\}$ with $w = 67185$, where $|\mathcal{F}_S(w)| = 4$.




Figure 2.1: The filling $\mathcal{F}_S(w)$ for $w = 67185$.

Theorem 2.1 ([6]). *Let $w = w_1 w_2 \dots w_t$ be any permutation of elements of $T \subseteq [n]$. Then*

$$\mathbf{rk}_S(T) = |\mathcal{F}_S(w)|. \quad (2.2)$$

We have the following minimal system of inequalities for Schubert matroid polytopes.

Lemma 2.2. *For any $S = \{s_1, s_2, \dots, s_k\} \subseteq [n]$, we have*

$$\mathcal{P}(\mathbf{SM}_n(S)) = \left\{ x \in \mathbb{R}_{\geq 0}^n : \sum_{i \in [n]} x_i = k, \sum_{i=1}^{s_j} x_i \geq j, \text{ for } j = 1, 2, \dots, k \right\}. \quad (2.3)$$

Proof. By Feichtner and Sturmfels [7, Proposition 2.6], the facets of any matroid polytope $\mathcal{P}(M)$ is

$$x_F := \sum_{e \in F} x_e \leq \text{rk}_M(F),$$

where $x_e \geq 0$ and F is a *facet* of M , i.e., F is an inseparable flat of M and $F^c = [n] \setminus F$ is inseparable in M^* .

For $i = 0, 1, \dots, k-1$, let

$$D_i = \{s_i + 1, s_i + 2, \dots, s_k\},$$

where $s_0 = 0$. It is easy to check that D_i is a facet of $\text{SM}_n(S)$, and $\text{rk}_S(D_i) = k - i$. Moreover, since $x_1 + \dots + x_n = k$, the inequality $\sum_{e \in D_i} x_e \leq \text{rk}_S(D_i) = k - i$ is equivalent to $x_1 + \dots + x_{s_i} \geq i$.

Let F be any facet of $\text{SM}_n(S)$. We aim to show that $F \in \{D_0, D_1, \dots, D_{k-1}\}$. For $i = 0, 1, \dots, k-1$, let

$$D'_i = \{s_i + 1, s_i + 2, \dots, s_{i+1}\},$$

where $s_0 = 0$. Firstly, we show that $D'_{k-1} \subseteq F$. Suppose that $D'_{k-1} \cap F = \emptyset$. Then $D'_{k-1} \subset F^c$, so we can separate F^c into two disjoint subsets: D'_{k-1} and $F^c \setminus D'_{k-1}$. It's easy to check that

$$\text{rk}_{M^*}(F^c) = \text{rk}_{M^*}(D'_{k-1}) + \text{rk}_{M^*}(F^c \setminus D'_{k-1}),$$

which implies that F^c is separable in M^* , contradiction. Consequently, $D'_{k-1} \cap F \neq \emptyset$. If there exists $b \in D'_{k-1}$ but $b \notin F$, then we claim that

$$\text{rk}_S(F) = \text{rk}_S(F \cup \{b\}),$$

which contradicts F being a flat. In fact, choose any element $a \in D'_{k-1} \cap F$, arrange a and b as the last two elements of a permutation of $F \cup \{b\}$. By Theorem 2.1, since a and b can only occupy the box in row s_k of S (i.e., the last row of S) and a appears before b , we see that b will not appear in the filling. Thus the claim follows. Then we have $D'_{k-1} \subseteq F$.

If $D'_i \cap F \neq \emptyset$ for any $i = 0, 1, \dots, k-1$, then using similar arguments as above, one can show that $F = [n]$ recursively, namely, $F = D_0$. Let j be the largest index such that $D'_j \cap F = \emptyset$. That is, $D'_i \cap F \neq \emptyset$ for $i \geq j+1$. Similarly, we can obtain $D'_i \subseteq F$ for $i \geq j+1$. Since $D_{j+1} = D'_{j+1} \cup \dots \cup D'_k$, we obtain $D_{j+1} \subseteq F$. We claim that $F = D_{j+1}$.

Suppose to the contrary $F \setminus D_{j+1} \neq \emptyset$. We aim to show that F is separable, that is,

$$\text{rk}_S(F) = \text{rk}_S(F_1) + \text{rk}_S(F_2),$$

where $F_1 = F \setminus D_{j+1}$ and $F_2 = D_{j+1}$.

Arrange the elements of F decreasingly, and then apply Theorem 2.1. Since the elements of D_{j+1} occupy all the boxes from row s_{j+2} to row s_k , and can not be filled into the first s_{j+1} rows, we see that $\text{rk}_S(F)$ is determined by how many elements of F_1 can be filled into the first s_{j+1} rows of S . For convenience, let $S' = \{s_1, \dots, s_{j+1}\}$, then we have $\text{rk}_S(F) = \text{rk}_{S'}(F_1) + \text{rk}_S(F_2)$.

To prove F is separable, it suffices to show that $\text{rk}_{S'}(F_1) = \text{rk}_S(F_1)$. Clearly, $\text{rk}_{S'}(F_1) \leq \text{rk}_S(F_1)$. Assume that $\text{rk}_{S'}(F_1) < \text{rk}_S(F_1)$. Then the box in row s_{j+1} must be occupied in any filling of S' with elements of F_1 . This would imply F is not a flat. Actually, for any $e \in D'_j$, let us consider $\text{rk}_S(F \cup \{e\})$. Arrange the elements of F decreasingly and append e at the end, and then apply Theorem 2.1. Since the last $k - j - 1$ rows are all occupied by elements in D_{j+1} , and the box in row s_{j+1} is occupied by an element of F_1 , the element e will not appear in the filling. That is, $\text{rk}_S(F \cup \{e\}) = \text{rk}_S(F)$, which is a contradiction. \blacksquare

3 Proof of Theorem 1.1

Recall that the Catalan matroid C_n corresponds to $r(S) = \overbrace{(1, 1, \dots, 1, 1)}^{2n}$. By Lemma 2.2, we have

$$\mathcal{P}(C_n) = \left\{ x \in \mathbb{R}_{\geq 0}^{2n} : \sum_{i \in [2n]} x_i = n, \sum_{i \in [2j]} x_i \geq j, \text{ for } j = 1, 2, \dots, n-1 \right\}. \quad (3.1)$$

For $m = 0, 1, \dots, n-1$, define

$$\mathcal{P}(C_{n,m}) := \left\{ x \in \mathbb{R}_{\geq 0}^{2n} : \sum_{i \in [2n]} x_i = n, \sum_{i \in [2j]+2m} x_i \geq j, \text{ for } j = 1, 2, \dots, n-1 \right\}, \quad (3.2)$$

where, for simplicity, x_i is understood as x_{i-2n} if $i > 2n$. Clearly, $\mathcal{P}(C_n) = \mathcal{P}(C_{n,0})$. Moreover, it is well known that

$$\mathcal{P}(U_n) = \left\{ x \in \mathbb{R}_{\geq 0}^{2n} : \sum_{i \in [2n]} x_i = n \right\}.$$

Lemma 3.1. *We have*

$$\mathcal{P}(U_n) = \bigcup_{j=1}^n \mathcal{P}(C_{n,j-1}). \quad (3.3)$$

Proof. It is obvious that $\mathcal{P}(C_{n,j-1}) \subseteq \mathcal{P}(U_n)$ for any $j = 1, \dots, n$. Let $(x_1, \dots, x_{2n}) \in \mathcal{P}(U_n)$, i.e., $x_1 + \dots + x_{2n} = n$, we aim to show that $(x_1, \dots, x_{2n}) \in \bigcup_{j=1}^n \mathcal{P}(C_{n,j-1})$.

For $i = 1, 2, \dots, n$, let $c_i = x_{2i-1} + x_{2i} - 1$. If $c_1 + \dots + c_k \geq 0$ for all $k = 1, 2, \dots, n$, then it is clear that $(x_1, \dots, x_{2n}) \in \mathcal{P}(C_n) = \mathcal{P}(C_{n,0})$. Suppose that there exists some index k such that $c_1 + \dots + c_k < 0$. Let $m = \min\{c_1, c_1 + c_2, c_1 + c_2 + c_3, \dots, c_1 + \dots + c_n\}$. Then $m < 0$. Choose the smallest index l such that $c_1 + \dots + c_l = m$. We claim that $(x_1, \dots, x_{2n}) \in \mathcal{P}(C_{n,l})$.

Actually, the sequence (c_1, c_2, \dots, c_n) can be viewed as a path from $(0, 0)$ to $(n, 0)$ such that each c_i is a step from $(i-1, c_{i-1})$ to (i, c_i) , where $c_0 = 0$. By cutting off the first l steps of (c_1, \dots, c_n) and appending them to the end, we obtain $(c_{l+1}, c_{l+2}, \dots, c_n, c_1, \dots, c_l)$. It is easy to see that $(c_{l+1}, c_{l+2}, \dots, c_n, c_1, \dots, c_l)$ is a path whose steps never go below the x -axis. That is, $(c_{l+1}, c_{l+2}, \dots, c_n, c_1, \dots, c_l) \in \mathcal{P}(C_{n,0})$, which implies $(x_1, \dots, x_{2n}) \in \mathcal{P}(C_{n,l})$. \blacksquare

For any subset $A \subseteq [n]$, denote

$$Q(A) = \bigcap_{j \in A} \mathcal{P}(C_{n,j-1}).$$

By (3.1) and the inclusion-exclusion principle, we have

$$i(U_n, t) = \sum_{A \subseteq [n]} (-1)^{|A|-1} \cdot i(Q(A), t). \quad (3.4)$$

Assume that $A = \{a_1, \dots, a_r\} \subseteq [n]$ arranged increasingly, let $\lambda(A)$ be the partition obtained by arranging the integers

$$a_1, a_2 - a_1, \dots, a_r - a_{r-1}, n - a_r + a_1$$

decreasingly. Actually, if we put $1, 2, \dots, n$ on a circle, i.e., 1 and n are viewed as adjacent to each other, then the elements of $\lambda(A)$ are gaps between the a_i 's. For example, let $A = \{1, 2, 4, 5\} \subseteq [7]$, then $\lambda(A) = (3, 2, 1, 1)$.

Lemma 3.2. *Let $\lambda \vdash n$ be a partition of n and $A \subseteq [n]$ such that $\lambda(A) = \lambda$. Then*

$$i(Q(A), t) = i(C_\lambda, t).$$

Proof. Suppose that $A = \{a_1, \dots, a_r\}_\leq$ and by convention let $a_0 = 0$. By (3.2), we have

$$\begin{aligned} Q(A) &= \bigcap_{i=1}^r \left\{ x \in \mathbb{R}_{\geq 0}^{2n} : \sum_{k \in [2n]} x_k = n, \quad \sum_{k \in [2j] + 2a_i - 2} x_k \geq j, \text{ for } 1 \leq j \leq n-1 \right\} \\ &= \left\{ x \in \mathbb{R}_{\geq 0}^{2n} : \sum_{k \in [2n]} x_k = n, \quad \sum_{k \in [2j] + 2a_i - 2} x_k \geq j, \text{ for } 1 \leq i \leq r \text{ and } 1 \leq j \leq n-1 \right\}. \end{aligned} \quad (3.5)$$

Notice that there are $r(n-1)$ inequalities in (3.5). For any $1 \leq i \leq r-1$, since

$$\sum_{k \in [2j] + 2a_i - 2} x_k \geq j, \text{ for } 1 \leq j \leq n-1,$$

let $j = a_{i+1} - a_i$, we have

$$x_{2a_i-1} + x_{2a_i} + \dots + x_{2a_{i+1}-2} \geq a_{i+1} - a_i. \quad (3.6)$$

By letting $j = n - a_{i+1} + a_i$ in

$$\sum_{k \in [2j] + 2a_{i+1} - 2} x_k \geq j, \text{ for } 1 \leq j \leq n-1,$$

we get

$$x_{2a_{i+1}-1} + \dots + x_{2n+2a_i-2} = x_{2a_{i+1}-1} + \dots + x_{2n} + x_1 + \dots + x_{2a_i-2} \geq n - a_{i+1} + a_i,$$

which is equivalent to

$$x_{2a_i-1} + \dots + x_{2a_{i+1}-2} \leq a_{i+1} - a_i. \quad (3.7)$$

Combining (3.6) and (3.7), we obtain that

$$x_{2a_i-1} + \dots + x_{2a_{i+1}-2} = a_{i+1} - a_i. \quad (3.8)$$

Moreover, by putting $j = n - a_r + a_1$ in

$$\sum_{k \in [2j] + 2a_r - 2} x_k \geq j, \text{ for } 1 \leq j \leq n-1,$$

we have

$$x_{2a_r-1} + \dots + x_{2n+2a_1-2} = x_{2a_r-1} + \dots + x_{2n} + x_1 + \dots + x_{2a_1-2} \geq n - a_r + a_1. \quad (3.9)$$

By plugging $j = a_r - a_1$ into

$$\sum_{k \in [2j] + 2a_1 - 2} x_k \geq j, \text{ for } 1 \leq j \leq n-1,$$

we find

$$x_{2a_1-1} + \dots + x_{2a_r-2} \geq a_r - a_1,$$

which is equivalent to

$$x_1 + \dots + x_{2a_1-2} + x_{2a_r-1} + \dots + x_{2n} \leq n - a_r + a_1. \quad (3.10)$$

Combining (3.9) and (3.10), we obtain that

$$x_1 + \cdots + x_{2a_1-2} + x_{2a_r-1} + \cdots + x_{2n} = n - a_r + a_1. \quad (3.11)$$

Therefore, we have

$$\begin{cases} x_{2a_i-1} + \cdots + x_{2a_{i+1}-2} = a_{i+1} - a_i, & \text{for } 1 \leq i \leq r-1, \\ x_1 + \cdots + x_{2a_1-2} + x_{2a_r-1} + \cdots + x_{2n} = n - a_r + a_1, \\ x_{2a_i-1} + \cdots + x_{2a_i-2+2j} \geq j, & \text{for } 1 \leq j \leq n-1 \text{ and } 1 \leq i \leq r. \end{cases} \quad (3.12)$$

On the other hand, it's easy to see that (3.12) implies (3.5), so we conclude that (3.5) is equivalent to (3.12), namely, $Q(A)$ is the set of nonnegative solutions of equation (3.12).

We proceed to show that the set of nonnegative solutions of equation (3.12) is isomorphic to the direct sum of some Catalan matroid polytopes. For $1 \leq i \leq r-1$, by (3.1),

$$\mathcal{P}(C_{a_{i+1}-a_i}) = \left\{ x \in \mathbb{R}_{\geq 0}^{2n} : \sum_{k \in [2(a_{i+1}-a_i)]} x_k = a_{i+1} - a_i, \sum_{k \in [2j]} x_k \geq j, \text{ for } 1 \leq j \leq a_{i+1} - a_i - 1 \right\},$$

which is isomorphic to

$$\left\{ x \in \mathbb{R}_{\geq 0}^{2n} : \sum_{k=2a_i-1}^{2a_{i+1}-2} x_k = a_{i+1} - a_i, \sum_{k \in [2j]+2a_i-2} x_k \geq j, \text{ for } 1 \leq j \leq a_{i+1} - a_i - 1 \right\}. \quad (3.13)$$

Notice that in (3.13), if $a_{i+1} - a_i \leq j \leq n-1$, since $x_{2a_{i+1}-1} + \cdots + x_{2a_{i+1}-2+2k} \geq k$ for any $1 \leq k \leq n-1$, we see that the inequality

$$\sum_{k \in [2j]+2a_i-2} x_k = a_{i+1} - a_i + \sum_{k=2a_{i+1}-1}^{2j+2a_i-2} x_k \geq a_{i+1} - a_i + (j - a_{i+1} + s_i) = j$$

automatically holds. That is to say, $\mathcal{P}(C_{a_{i+1}-a_i})$ is isomorphic to

$$\left\{ x \in \mathbb{R}_{\geq 0}^{2n} : \sum_{k=2a_i-1}^{2a_{i+1}-2} x_k = a_{i+1} - a_i, \sum_{k \in [2j]+2a_i-2} x_k \geq j, \text{ for } 1 \leq j \leq n-1 \right\}.$$

Similarly,

$$\mathcal{P}(C_{n-a_r+a_1}) = \left\{ x \in \mathbb{R}_{\geq 0}^{2n} : \sum_{k \in [2(n-a_r+a_1)]} x_k = n - a_r + a_1, \sum_{k \in [2j]} x_k \geq j, \text{ for } 1 \leq j \leq n - a_r + a_1 - 1 \right\}.$$

After shifting all the indices by $2a_r - 2$, we see that $\mathcal{P}(C_{n-a_r+a_1})$ is isomorphic to

$$\left\{ x \in \mathbb{R}_{\geq 0}^{2n} : \sum_{2a_r-1}^{2n+2a_1-2} x_k = n - a_r + a_1, \sum_{k \in [2j]+2a_r-2} x_k \geq j, \text{ for } 1 \leq j \leq n - a_r + a_1 - 1 \right\}. \quad (3.14)$$

Notice that in (3.14), if $n - a_r + a_1 \leq j \leq n-1$, since $x_{2a_1-1} + \cdots + x_{2a_1-2+2k} \geq k$ for any $1 \leq k \leq n-1$, we see that the inequality

$$\sum_{k \in [2j]+2a_r-2} x_k = n - a_r + a_1 + \sum_{k=2a_1-1}^{2j-2n+2a_r-2} x_k \geq n - a_r + a_1 + (j - n + a_r - a_1) = j$$

always holds. Thus $\mathcal{P}(C_{n-a_r+a_1})$ is isomorphic to

$$\left\{ x \in \mathbb{R}_{\geq 0}^{2n} : \sum_{k=1}^{2a_1-2} x_k + \sum_{k=1}^{2n} x_k = n - a_r + a_1, \sum_{k \in [2j]+2a_r-2} x_k \geq j, \text{ for } 1 \leq j \leq n-1 \right\}.$$

Therefore, the set of nonnegative solutions of equation (3.12) is isomorphic to the Cartesian product of the Catalan matroid polytopes $\mathcal{P}(C_{n-a_r+a_1})$ and $\mathcal{P}(C_{a_{i+1}-a_i})$ for $1 \leq i \leq r-1$. That is,

$$Q(A) = \mathcal{P}(C_{a_2-a_1}) \times \mathcal{P}(C_{a_3-a_2}) \times \cdots \times \mathcal{P}(C_{a_r-a_{r-1}}) \times \mathcal{P}(C_{n-a_r+a_1}). \quad (3.15)$$

Therefore,

$$i(Q(A), t) = i(C_{a_2-a_1}, t) \cdots i(C_{a_r-a_{r-1}}, t) \cdot i(C_{n-a_r+a_1}, t) = i(C_\lambda, t),$$

as required. ■

For a partition $\lambda = (\lambda_1, \dots, \lambda_k)$ of n , let $\ell(\lambda)$ denote the number of parts of λ , i.e., $\ell(\lambda) = k$. We can also write $\lambda = (1^{\alpha_1}, \dots, n^{\alpha_n})$, where α_i is the number of appearances of i in λ . Denote $p(\lambda)$ by the number of permutations of the parts of λ , that is,

$$p(\lambda) = \frac{\ell(\lambda)!}{\alpha_1! \alpha_2! \cdots \alpha_n!}.$$

For a composition $\gamma = (\gamma_1, \dots, \gamma_k)$ of n , if there exists an integer $m > 0$ such that $\gamma_i = \gamma_{i+m}$ for all $1 \leq i \leq k-m$, then we call m a period of γ . Let $T(\gamma)$ denote the least period of γ . If there does not exist such a positive integer m , set $T(\gamma) = k$. For instance, if $\gamma = (2, 2, 1, 2, 2, 1, 2, 2, 1)$, then $T(\gamma) = 3$. If $\gamma = (2, 1, 3, 2)$, then $T(\gamma) = 4$.

Lemma 3.3. *Let $\lambda \vdash n$ be a partition. Then we have*

$$\#\{A \subseteq [n] : \lambda(A) = \lambda\} = \frac{n}{\ell(\lambda)} p(\lambda).$$

Proof. Let $\lambda = (\lambda_1, \dots, \lambda_k)$. Given a permutation $\gamma = (\lambda_{i_1}, \lambda_{i_2}, \dots, \lambda_{i_k})$ of λ , we can construct n subsets $A \subseteq [n]$ such that $\lambda(A) = \lambda$ as below

$$\{j, j + \lambda_{i_1}, j + \lambda_{i_1} + \lambda_{i_2}, \dots, j + \lambda_{i_1} + \cdots + \lambda_{i_{k-1}}\}, \quad (3.16)$$

where $1 \leq j \leq n$ and any λ_m is understood as λ_{m-n} if $m > n$. It is easy to see that all subsets $A \subseteq [n]$ such that $\lambda(A) = \lambda$ can be generated in this way. However, there are repetitions in (3.16). To be exact, it is easy to see that the set

$$\{j', j' + \lambda_{i_1}, j' + \lambda_{i_1} + \lambda_{i_2}, \dots, j' + \lambda_{i_2} + \cdots + \lambda_{i_{k-1}}\},$$

where $j' = j + \lambda_{i_1} + \cdots + \lambda_{i_{T(\gamma)}}$ is the same as the set in (3.16). Therefore, for a given a permutation γ of λ , each subset A of $[n]$ such that $\lambda(A) = \lambda$ is generated $\frac{\ell(\lambda)}{T(\gamma)}$ times.

On the other hand, different permutations of λ may generate the same subset A . More precisely, for a shifting $\gamma' = (\lambda_{i_2}, \lambda_{i_3}, \dots, \lambda_{i_k}, \lambda_{i_1})$ of γ , if $\gamma' \neq \gamma$, then we can construct the following subset

$$\{j'', j'' + \lambda_{i_2}, j'' + \lambda_{i_2} + \lambda_{i_3}, \dots, j'' + \lambda_{i_2} + \cdots + \lambda_{i_k}\}, \quad (3.17)$$

where $j'' = j + \lambda_{i_1}$. Since $j'' + \lambda_{i_2} + \cdots + \lambda_{i_k} = j + n$, the two sets in (3.16) and (3.17) are the same. Each permutation γ of λ can shift $T(\gamma)$ different times, all generating the same subset. Thus the total number of repetitions of each A is $\ell(\lambda)$.

Since there are $p(\lambda)$ permutations of λ , and for each permutation of λ , we constructed n subsets A as in (3.16), but there are $\ell(\lambda)$ copies of A constructed in this way, we conclude that the total number of subsets A with $\lambda(A) = \lambda$ is $\frac{n}{\ell(\lambda)} p(\lambda)$. ■

Suppose that $\tau = (\tau_1, \dots, \tau_k)$ and $\sigma = (\sigma_1, \dots, \sigma_s)$ are two compositions of n . We say that τ and σ are *equivalent*, denoted as $\tau \sim \sigma$, if they have the same multiset of parts. Obviously, if $\tau \sim \sigma$, then $i(U_\tau, t) = i(U_\sigma, t)$. We say that τ is a *refinement* of σ , denoted as $\tau \prec \sigma$, if each σ_i can be obtained by adding a sequence of consecutive factors of τ . More precisely, there exists i_1, i_2, \dots, i_{s-1} such that

$$\sigma_1 = \tau_1 + \dots + \tau_{i_1}, \sigma_2 = \tau_{i_1+1} + \dots + \tau_{i_2}, \dots, \sigma_s = \tau_{i_{s-1}+1} + \dots + \tau_k.$$

Let $F(\lambda)$ be the set of refinements of λ . For instance, let $\lambda = (3, 2)$, then $F(\lambda) = \{(3, 2), (2, 1, 2), (1, 1, 1, 2), (3, 1, 1), (2, 1, 1, 1), (1, 1, 1, 1, 1)\}$.

For a partition $\lambda \vdash n$, let K_λ denote the set of permutations on $[n]$ with *cycle type* λ . Recall that a permutation $w \in S_n$ has cycle type $\lambda = (1^{\alpha_1}, \dots, n^{\alpha_n})$ means that when converted into cycles, w has α_i cycles of length i , for $i = 1, 2, \dots, n$. It is well known that

$$|K_\lambda| = \frac{n!}{z_\lambda},$$

see, for example, Sagan [17].

Proof of Theorem 1.1. By (3.4), Lemma 3.2 and Lemma 3.3, we have

$$i(U_n, t) = \sum_{\lambda \vdash n} (-1)^{\ell(\lambda)-1} \frac{np(\lambda)}{\ell(\lambda)} i(C_\lambda, t). \quad (3.18)$$

Then

$$i(C_n, t) = \frac{1}{n} i(U_n, t) + \sum_{\lambda \vdash n, \lambda \neq (n)} (-1)^{\ell(\lambda)} \frac{p(\lambda)}{\ell(\lambda)} i(C_\lambda, t). \quad (3.19)$$

Since $z_{(n)} = \frac{1}{n}$ and $i(U_{(n)}, t) = i(U_n, t)$, by (3.19), it suffices to show that

$$\sum_{\lambda \vdash n, \lambda \neq (n)} (-1)^{\ell(\lambda)} \frac{p(\lambda)}{\ell(\lambda)} i(C_\lambda, t) = \sum_{\lambda \vdash n, \lambda \neq (n)} \frac{1}{z_\lambda} i(U_\lambda, t). \quad (3.20)$$

We make induction on n . By induction,

$$\begin{aligned} & \sum_{\lambda \vdash n, \lambda \neq (n)} (-1)^{\ell(\lambda)} \frac{p(\lambda)}{\ell(\lambda)} i(C_\lambda, t) \\ &= \sum_{\lambda \vdash n, \lambda \neq (n)} (-1)^{\ell(\lambda)} \frac{p(\lambda)}{\ell(\lambda)} \prod_{j=1}^{\ell(\lambda)} \left(\sum_{\sigma^j \vdash \lambda_j} \frac{1}{z_{\sigma^j}} i(U_{\sigma^j}, t) \right) \\ &= \sum_{\lambda \vdash n, \lambda \neq (n)} (-1)^{\ell(\lambda)} \frac{p(\lambda)}{\ell(\lambda)} \sum_{\sigma \in F(\lambda)} \frac{1}{z_{\sigma^1} z_{\sigma^2} \cdots z_{\sigma^{\ell(\lambda)}}} i(U_\sigma, t), \end{aligned} \quad (3.21)$$

where, given a refinement $\sigma \in F(\lambda)$ of λ , there is a unique way to cut σ into $\ell(\lambda)$ factors $\sigma^1, \sigma^2, \dots, \sigma^{\ell(\lambda)}$ such that each $\sigma^i \vdash \lambda_i$ is a partition of λ_i .

In order to prove (3.20), we need to show that for any partition μ of n with $\mu \neq (n)$, the coefficient of $i(U_\mu, t)$ in (3.21) is equal to $1/z_\mu$. Since $i(U_\sigma, t) = i(U_\mu, t)$ if $\sigma \sim \mu$, the coefficient of $i(U_\mu, t)$ in (3.21) is

$$\sum_{\lambda \vdash n, \lambda \neq (n)} (-1)^{\ell(\lambda)} \frac{p(\lambda)}{\ell(\lambda)} \sum_{\sigma \in F(\lambda), \sigma \sim \mu} \frac{1}{z_{\sigma^1} z_{\sigma^2} \cdots z_{\sigma^{\ell(\lambda)}}}.$$

That is, we need to show that

$$\sum_{\lambda \vdash n, \lambda \neq (n)} (-1)^{\ell(\lambda)} \frac{p(\lambda)}{\ell(\lambda)} \sum_{\sigma \in F(\lambda), \sigma \sim \mu} \frac{1}{z_{\sigma^1} z_{\sigma^2} \cdots z_{\sigma^{\ell(\lambda)}}} = \frac{1}{z_\mu},$$

which is equivalent to

$$\sum_{\lambda \vdash n, \lambda \neq (n)} (-1)^{\ell(\lambda)} \frac{p(\lambda)}{\ell(\lambda)} \sum_{\sigma \in F(\lambda), \sigma \sim \mu} \frac{\lambda_1!}{z_{\sigma^1}} \frac{\lambda_2!}{z_{\sigma^2}} \cdots \frac{\lambda_{\ell(\lambda)}!}{z_{\sigma^{\ell(\lambda)}}} \cdot \binom{n}{\lambda_1, \dots, \lambda_{\ell(\lambda)}} = \frac{n!}{z_\mu}.$$

Therefore, it is enough to show that for any partition $\mu = (\mu_1, \dots, \mu_r) \vdash n, \mu \neq (n)$, there holds

$$\sum_{\lambda \vdash n, \lambda \neq (n)} (-1)^{\ell(\lambda)} \frac{p(\lambda)}{\ell(\lambda)} \sum_{\sigma \in F(\lambda), \sigma \sim \mu} |K_{\sigma^1}| |K_{\sigma^2}| \cdots |K_{\sigma^{\ell(\lambda)}}| \cdot \binom{n}{\lambda_1, \dots, \lambda_{\ell(\lambda)}} = |K_\mu|. \quad (3.22)$$

Suppose that $w \in K_\mu$, that is, when converted in cycles, w has cycle type μ . In particular, w has r cycles. We need to show that the total number of appearances of w in the left-hand side of (3.22) is equal to 1. To this end, we discuss according to the number of parts of λ in the left-hand side of (3.22). We claim that for any fixed $k = \ell(\lambda) \geq 2$, w is counted in

$$\sum_{\lambda \vdash n, \ell(\lambda) = k} (-1)^k \frac{p(\lambda)}{k} \sum_{\sigma \in F(\lambda), \sigma \sim \mu} |K_{\sigma^1}| |K_{\sigma^2}| \cdots |K_{\sigma^k}| \cdot \binom{n}{\lambda_1, \dots, \lambda_k} \quad (3.23)$$

by $(-1)^k (k-1)! S(r, k)$ times, where $S(r, k)$ is Stirling number of the second kind.

Let $\lambda = (\lambda_1, \dots, \lambda_k) = (1^{\alpha_1}, \dots, n^{\alpha_n}) \vdash n$. Notice that if $\sigma \in F(\lambda)$, then σ admits a unique decomposition into factors $\sigma^1, \dots, \sigma^k$ such that $\sigma^i \vdash \lambda_i$. Choose an ordered partition of $[n]$, i.e.,

$$T_1 = \{j_1, \dots, j_{\lambda_1}\}, T_2 = \{j_{\lambda_1+1}, \dots, j_{\lambda_1+\lambda_2}\}, \dots, T_k = \{j_s, \dots, j_n\}, \quad (3.24)$$

where $s = n - (\lambda_1 + \dots + \lambda_{k-1}) + 1$, such that $|T_i| = \lambda_i$, $T_i \cap T_j = \emptyset$ for $i \neq j$ and $T_1 \cup \dots \cup T_k = [n]$. There are $\binom{n}{\lambda_1, \dots, \lambda_k}$ such ordered partitions of $[n]$. For $1 \leq i \leq k$, let T_i form a permutation of cycle type σ^i , there are $|K_{\sigma^i}|$ ways. Suppose that w is counted in (3.23) at least once. Then each T_i in (3.24) must be a union of some cycles of w . Moreover, although each T_i can form $|K_{\sigma^i}|$ different permutations of cycle type σ^i , there is exactly one way to obtain the cycles of w . Thus w is counted exactly once by a given ordered partition of $[n]$ in (3.24).

Assume that $|T_i| = |T_j|$ for some $i < j$ and T_i forms a permutation of cycle type σ^i , T_j forms a permutation of cycle type σ^j . Then we can exchange the positions of T_i and T_j in (3.24) to obtain k ordered disjoint subsets of $[n]$:

$$T_1, \dots, T_j, \dots, T_i, \dots, T_k,$$

and let them form k permutations of cycle types $\sigma^1, \dots, \sigma^j, \dots, \sigma^i, \dots, \sigma^k$, respectively. Clearly, w can also be obtained once in this way. In other words, for a given $\lambda \vdash n$ of length k , w is counted in

$$\sum_{\sigma \in F(\lambda), \sigma \sim \mu} |K_{\sigma^1}| |K_{\sigma^2}| \cdots |K_{\sigma^k}| \cdot \binom{n}{\lambda_1, \dots, \lambda_k}$$

by $\alpha_1! \alpha_2! \cdots \alpha_n! = k! / p(\lambda)$ times.

To prove the claim, we need to sum over all partitions $\lambda \vdash n$ with $\ell(\lambda) = k$. If w is counted once by (3.23), then each T_i in (3.24) is a union of some cycles of w . In other words, for a given $w \in K_\mu$, we can merge the r cycles of w to obtain k sets T_i in (3.24). If we run over all possible ways of ‘‘cycles merging’’ of w , then we actually obtain all partitions of n with exactly k parts. Obviously, there are $S(r, k)$ such ways. Therefore, w is counted in (3.23) by

$$(-1)^k \frac{p(\lambda)}{k} \frac{k!}{p(\lambda)} S(r, k) = (-1)^k (k-1)! S(r, k)$$

times.

By summing for all $k \geq 2$ and noticing that $S(r, k) = kS(r-1, k) + S(r-1, k-1)$, we conclude that the total number of $w \in K_\mu$ counted in the left-hand side of (3.22) is

$$\begin{aligned}
& \sum_{k=2}^r (-1)^k (k-1)! S(r, k) \\
&= \sum_{k=2}^r (-1)^k (k! S(r-1, k) + (k-1)! S(r, k-1)) \\
&= S(r-1, 1) + (-1)^r n! S(r-1, n) + \sum_{k=2}^{r-1} (k! S(r-1, k) - k! S(r-1, k)) \\
&= S(r-1, 1) \\
&= 1,
\end{aligned}$$

as desired. ■

For example, let $\mu = (2, 2, 1, 1)$ and $w = (1, 4)(2, 5)(3)(6) \in K_\mu$. We enumerate how many times w is counted in (3.23). The set of partitions $\lambda \vdash 6, \lambda \neq (6)$ such that there exists a refinement $\sigma \in F(\lambda)$ with $\mu \sim \sigma$ is

$$\{(2, 2, 1, 1), (3, 2, 1), (2, 2, 2), (4, 1, 1), (5, 1), (4, 2), (3, 3)\}.$$

For $\lambda = (2, 2, 1, 1)$, among the $\binom{6}{2, 2, 1, 1} = 180$ possible ordered partitions of $\{1, \dots, 6\}$, there are four that can be used to obtain w , i.e.,

$$\begin{aligned}
T_1 &= \{1, 4\}, T_2 = \{2, 5\}, T_3 = \{6\}, T_4 = \{3\}, \\
T_1 &= \{1, 4\}, T_2 = \{2, 5\}, T_3 = \{3\}, T_4 = \{6\}, \\
T_1 &= \{2, 5\}, T_2 = \{1, 4\}, T_3 = \{6\}, T_4 = \{3\}, \\
T_1 &= \{2, 5\}, T_2 = \{1, 4\}, T_3 = \{3\}, T_4 = \{6\}.
\end{aligned}$$

For each of ordered partition above, we let T_1, T_2 form permutations with cycle type $\sigma^1 = \sigma^2 = (2)$ and let T_3, T_4 form permutations with cycle type $\sigma^3 = \sigma^4 = (1)$. Thus w is counted $(-1)^{\ell(\lambda)} \frac{p(\lambda)}{\ell(\lambda)} \cdot 4 = 6$ times in (3.23) for $\lambda = (2, 2, 1, 1)$.

For $\lambda = (3, 2, 1)$, there are four ordered partitions of $\{1, \dots, 6\}$ that can be used to obtain w , i.e.,

$$\begin{aligned}
T_1 &= \{1, 4, 3\}, T_2 = \{2, 5\}, T_3 = \{6\}, \\
T_1 &= \{1, 4, 6\}, T_2 = \{2, 5\}, T_3 = \{3\}, \\
T_1 &= \{2, 5, 3\}, T_2 = \{1, 4\}, T_3 = \{6\}, \\
T_1 &= \{2, 5, 6\}, T_2 = \{1, 4\}, T_3 = \{3\}.
\end{aligned}$$

For each of them, let T_1 form a permutation of cycle type $\sigma^1 = (2, 1)$, T_2, T_3 form permutations of cycle types $\sigma^2 = (2), \sigma^3 = (1)$, respectively. Thus w is counted $(-1)^{\ell(\lambda)} \frac{p(\lambda)}{\ell(\lambda)} \cdot 4 = -8$ times in (3.23) for $\lambda = (3, 2, 1)$.

Similarly, w is counted -2 times for $\lambda = (4, 1, 1)$, -2 times for $\lambda = (2, 2, 2)$, 3 times for $\lambda = (4, 2)$, 2 times for $\lambda = (5, 1)$, and 2 times for $\lambda = (3, 3)$, respectively. In total, w is counted $6-8-2-2+3+2+2 = 1$ time.

As an extension of the preceding discussion, we are now at a position to consider the volume of $\mathcal{P}(C_n)$ and $\mathcal{P}(C_n^{a,b})$.

Proof of Corollary 1.3. It is clear that

$$\dim(\mathcal{P}(C_{n,j-1})) = \dim(\mathcal{P}(C_n)) = \dim(\mathcal{P}(U_n)) = n - 1.$$

Since the intersection $Q(A) = \bigcap_{j \in A} \mathcal{P}(C_{n,j-1})$ is a face of $\mathcal{P}(C_{n,j-1})$ for any $j \in A$, the dimension of $Q(A)$ is smaller than $n - 1$. Consequently, for any subset $B \subseteq [n]$, we have

$$\sum_{j \in B} \text{vol}(\mathcal{P}(C_{n,j-1})) = \text{vol}(\bigcup_{j \in B} \mathcal{P}(C_{n,j-1})).$$

By Lemma 3.1, we deduce that

$$\text{vol}(\mathcal{P}(U_n)) = \sum_{j=1}^n \text{vol}(\mathcal{P}(C_{n,j-1})) = n \text{vol}(\mathcal{P}(C_n)).$$

Let $A(n, k)$ be the Eulerian number, which is the number of permutations on $[n]$ with $k - 1$ descents. It is well known that

$$\text{vol}(\mathcal{P}(U_{k,n})) = \frac{1}{(n-1)!} A(n-1, k-1),$$

see, for example, [18, Exercise 4.59(b)]. Therefore,

$$\text{vol}(\mathcal{P}(U_n)) = \frac{1}{(2n-1)!} A(2n-1, n-1).$$

Thus (1.4) holds. Similarly, (1.5) follows. ■

4 Concluding remarks

Recall that notched rectangle matroids are Schubert matroids $\text{SM}_n(S)$ with $r(S) = (a, b, c, d)$. In particular, $r(S) = (a, b, c, 1)$ corresponds to the panhandle matroid $\text{Pan}_{b+1, a+b, a+b+c+1}$, which was conjectured to be Ehrhart positive in [14], and proved in [4]. When $c = 1$, $r(S) = (a, b, 1, 1)$ corresponds to the panhandle matroid $\text{Pan}_{b+1, a+b, a+b+2}$.

For simplicity, write $i(r(S), t)$ for $i(\text{SM}_n(S), t)$. By [5, Theorem 1.5], there holds

$$\begin{aligned} i((a, b, a, b), t) &= \frac{1}{2} i(U_{2b, 2a+2b}, t) + \frac{1}{2} i(U_{b, a+b}, t)^2, \\ i((a, a, b, b), t) &= \frac{1}{2} i(U_{a+b, 2a+2b}, t) + \frac{1}{2} i(U_{a, 2a}, t) i(U_{b, 2b}, t), \\ i((1, 1, a, a+1), t) &= \frac{1}{2} (t+2) i(U_{a+1, 2a+2}, t). \end{aligned}$$

In this paper, we express $i(C_n^{a,b}, t)$ in terms of the Ehrhart polynomials of uniform matroids. Therefore, it seems feasible to express other families of Ehrhart polynomials in terms of the Ehrhart polynomials of uniform matroids. McGinnis [16] showed that $\text{Pan}_{b+1, a+b, a+b+2}$ is Ehrhart positive by using intricate combinatorial arguments. We propose the following Conjecture, which manifestly shows the Ehrhart positivity of $\text{Pan}_{b+1, a+b, a+b+2}$.

Conjecture 4.1. *Let a, b be positive integers, we have*

$$i(\text{Pan}_{b+1, a+b, a+b+2}, t) = i((1, 1, a, b), t) = \left(\frac{a+1}{a+b+1} t + 1 \right) \cdot i(U_{b, a+b+1}, t).$$

We remark that the Schubert matroid $\text{SM}_n(S)$ is paving if and only if $r(S) = (a, 1, b, c)$. In particular, $\text{SM}_n(S)$ is a paving panhandle matroid if and only if $r(S) = (1, 1, a, b)$.

References

- [1] F. Ardila, The Catalan matroid, *J. Combin. Theory Ser. A* 104 (2003), 49–62.
- [2] J.E. Bonin, A. de Mier and M. Noy, Lattice path matroids: enumerative aspects and Tutte polynomials, *J. Combin. Theory Ser. A* 104 (2003), 63–94.
- [3] F. Castillo and F. Liu, Berline-Vergne valuation and generalized permutohedra. *Discrete Comput. Geom.*, 60(4) (2018):885–908.
- [4] D. Deligeorgaki, D. McGinnis, A. R. Vindas-Meléndez, Ehrhart bounds for panhandle and paving matroids through enumeration of chain forests, arXiv:2311.01640.
- [5] N.J.Y. Fan and Y. Li, On the Ehrhart polynomial of Schubert matroids, *Discrete Comput. Geom.* 71 (2024), no. 2, 587–626.
- [6] N.J.Y. Fan and P.L. Guo, Vertices of Schubitopes, *J. Combin. Theory, Series A*, 177(2021), 105311.
- [7] E. Feichtner and B. Sturmfels, Matroid polytopes, nested sets and bergman fans, *Portugaliae Mathematica. Nova Srie*, 62(2005), 437–468.
- [8] L. Ferroni, Hypersimplices are Ehrhart positive, *J. Combin. Theory, series A*, 178(2021), 105365.
- [9] L. Ferroni, On the Ehrhart polynomial of minimal matroids, *Discrete Comput. Geom.*, 68(2022), 255–273.
- [10] L. Ferroni, Matroids are not Ehrhart positive, *Adv. Math.*, 402(2022), 108337.
- [11] L. Ferroni, K. Jochemko and B. Schröter, Ehrhart polynomials of rank two matroids, *Adv. Appl. Math.*, 141(2022), 102410.
- [12] L. Ferroni and B. Schröter, Valuative invariants for large classes of matroids, *J. Lond. Math. Soc.* (2) 110 (2024), no. 3, Paper No. e12984, 86.
- [13] A. Fink, K. Mészáros and A. St. Dizier, Schubert polynomials as integer point transforms of generalized permutahedra, *Adv. Math.* 332 (2018), 465–475.
- [14] D. Hanely, J. L. Martin, D. McGinnis, D. Miyata, G. D. Nasr, A. R. Vindas-Meléndez, and M. Yin, Ehrhart theory of paving and panhandle matroids, *Adv. Geom.* 23 (2023), no.4, 501–526.
- [15] J. A. De Loera, D. C. Haws and M. Köppe. Ehrhart polynomials of matroid polytopes and poly-matroids, *Discrete Comput. Geom.* 42 (2009), no. 4, 670–702.
- [16] D. McGinnis, A combinatorial formula for the Ehrhart coefficients of a certain class of weighted multi-hypersimplices, arXiv:2303.04113.
- [17] B. E. Sagan, *The Symmetric Group: Representations, Combinatorial Algorithms, and Symmetric Functions*, Springer–Verlag, New York, 1991.
- [18] R. Stanley, *Enumerative Combinatorics. Volume 1. Cambridge Studies in Advanced Mathematics*, vol. 49. Cambridge University Press, Cambridge, 1997.

Y.M. CHEN, DEPARTMENT OF MATHEMATICS, SICHUAN UNIVERSITY, CHENGDU 610064, P.R. CHINA.
Email address: ym.chen@stu.scu.edu.cn

Y. LI, DEPARTMENT OF MATHEMATICS, SICHUAN UNIVERSITY, CHENGDU 610064, P.R. CHINA.
Email address: liyao@stu.pku.edu.cn

M. YAO, DEPARTMENT OF MATHEMATICS, SICHUAN UNIVERSITY, CHENGDU 610064, P.R. CHINA.
Email address: yaom@stu.scu.edu.cn