Splitting Answer Set Programs with respect to Intensionality Statements (Extended Version)

Jorge Fandinno* and Yuliya Lierler*

University of Nebraska at Omaha {jfandinno,ylierler}@unomaha.edu

Abstract

Splitting a logic program allows us to reduce the task of computing its stable models to similar tasks for its subprograms. This can be used to increase solving performance and to prove the correctness of programs. We generalize the conditions under which this technique is applicable, by considering not only dependencies between predicates but also their arguments and context. This allows splitting programs commonly used in practice to which previous results were not applicable.

Introduction

Answer set programming (ASP; Lifschitz 2008) is a declarative logic programming paradigm well-suited for solving knowledge-intensive search problems. Its success relies on the combination of a rich knowledge representation language with efficient solvers for finding solutions to problems expressed in this language (Lifschitz 2019). Solutions for logic programs in ASP are called "stable models". Lifschitz and Turner (1994) introduced a fundamental splitting method (or, simply, splitting) in the theory of ASP. Often, splitting is used to reduce the task of computing the stable models of a logic program to the task of computing these of its subprograms. Also, the splitting is used to understand the meaning of a program in terms of its smaller components and it has become a key instrument in constructing proofs of correctness for logic programs (Cabalar, Fandinno, and Lierler 2020). The original condition required to split a logic program was generalized by Oikarinen and Janhunen (2008) in the context of propositional programs. Also, when stable models were defined for arbitrary first-order formulas (Pearce and Valverde 2005; Ferraris, Lee, and Lifschitz 2007) and infinitary propositional formulas (Truszczyński 2012), the splitting method was extended to these new settings (Ferraris et al. 2009; Harrison and Lifschitz 2016). However, when dealing with first-order formulas (logic programs with variables are often identified with first-order formulas of a restricted syntactic form), the splitting condition is not sufficiently general for some applications.

For instance, consider the following simplified fragment of the blocks world encoding (Lifschitz 2002):

$on(B,L,T+1) \leftarrow$	on(B,L,T),	$not \ non(B, L, T$	+1) (1)
--------------------------	------------	---------------------	---------

 $on(B, L, T+1) \leftarrow move(B, L, T)$ (2)

 $non(B, L', T) \leftarrow on(B, L, T), \ location(L'), \ L \neq L'$ (3)

We are interested in splitting this fragment so that one subprogram contains instances of these rules for all time points before a (positive) threshold t and another contains rules corresponding to the time points at or after t. This form of splitting will help, for example, understanding iterative solving of planning problems (Gebser et al. 2019) and arguing correctness of action descriptions (that frequently result in rules of the form presented in our blocks world example). However, the proposed splitting is impossible under the currently available Splitting Theorem for first-order formulas (Ferraris et al. 2009). The splitting relies on identifying "non-circularly" dependent rules of a program. Ferraris et al. (2009) define dependencies in terms of predicate symbols. In our example, on/3 depends on itself in rule (1).

Here, we formulate a more general condition for the applicability of the splitting method that considers a refined version of the dependency graph that takes into account not only the predicate dependencies but also their arguments and context. Among others, this makes it applicable to the discussed example. The notion of *intensionality statement* introduced in this work is of key importance. This is a refinement of the idea of intensional and extensional predicates (Ferraris, Lee, and Lifschitz 2011) originally stemming from databases. It provides us with means for specifying arguments of the predicates for which these are seen as intensional or extensional. Thus, the same predicate symbol may be both intensional and extensional depending on its context. This refinement is the basis for a new Splitting result.

The rest of the paper is organized as follows. After reviewing some preliminaries, we introduce the notion of intensionality statement. Then, we present the key ideas for the new Splitting Theorem in the context of logic programs. Finally, we generalize this result to first-order theories.

^{*}These authors contributed equally.

Copyright © 2023, Association for the Advancement of Artificial Intelligence (www.aaai.org). All rights reserved.

Preliminaries

We start by reviewing a many-sorted first-order language. The use of a many-sorted language is motivated by its ability to formalize commonly used features of logic programs such as arithmetic operations (Fandinno et al. 2020; Lifschitz 2021) and aggregate expressions (Fandinno, Hansen, and Lierler 2022). We follow the presentation by Fandinno and Lifschitz (2022, Appendix A). After the presentation of many-sorted language, the review the logic of here-and-there follows. Syntax and semantics of disjunctive logic programs concludes this section.

Many-sorted first-order theories. A (many-sorted) signature consists of symbols of three kinds-sorts, function constants, and predicate constants. A reflexive and transitive subsort relation is defined on the set of sorts. A tuple s_1, \ldots, s_n $(n \ge 0)$ of argument sorts is assigned to every function constant and to every predicate constant; in addition, a value sort is assigned to every function constant. Function constants with n = 0 are called *object con*stants. We assume that for every sort, an infinite sequence of object variables of that sort is chosen. Terms over a (many-sorted) signature σ are defined recursively as usual. Atomic formulas over σ are either (i) expressions of the form $p(t_1, \ldots, t_n)$, where p is a predicate constant and t_1, \ldots, t_n are terms such that their sorts are subsorts of the argument sorts s_1, \ldots, s_n of p, or (ii) expressions of the form $t_1 = t_2$, where t_1 and t_2 are terms such that their sorts have a common supersort. (First-order) formulas over signature σ are formed from atomic formulas and the 0-place connective \perp (falsity) using the binary connectives \land, \lor, \rightarrow and the quantifiers \forall , \exists . The other connectives are treated as abbreviations: $\neg F$ stands for $F \rightarrow \bot$ and $F \leftrightarrow G$ stands for $(F \to G) \land (G \to F).$

An *interpretation* I of a signature σ assigns

- a non-empty *domain* $|I|^s$ to every sort s of I, so that $|I|^{s_1} \subseteq |I|^{s_2}$ whenever s_1 is a subsort of s_2 ,
- a function f^I from $|I|^{s_1} \times \cdots \times |I|^{s_n}$ to $|I|^s$ to every function constant f with argument sorts s_1, \ldots, s_n and value sort s, and
- a Boolean-valued function p^I on |I|^{s₁} ×···× |I|^{s_n} to every predicate constant p with argument sorts s₁,..., s_n.

If I is an interpretation over a signature σ then by σ^{I} we denote the signature obtained from σ by adding, for every element d of domain $|I|^{s}$, its name d^{*} as an object constant of sort s. The interpretation I is extended to σ^{I} by defining $(d^{*})^{I} = d$. The value t^{I} assigned by an interpretation I of σ to a ground term t over σ^{I} and the satisfaction relation (denoted, \models) between an interpretation of σ and a sentence over σ^{I} are defined recursively, in the usual way (Lifschitz, Morgenstern, and Plaisted 2008, Section 1.2.2). An interpretation is called a *model* of a *theory* – a (possibly infinite) set of sentences – when it satisfies every sentence in this theory. Some of the examples considered contain integers, function constants such as + and comparison predicate constants such as \leq or > (used utilizing infix notation common in arithmetic). We call these function and

predicate constants *arithmetic*. In these examples, we assume that (i) the underlying signature contains a sort integer and (ii) interpretations of special kind are considered so that they interpret arithmetic function and predicate constants as customary in arithmetic (see, for example, treatment of the integer sort by Fandinno et al. (2020) and Lifschitz (2021)). If d is a tuple d_1, \ldots, d_n of domains elements of I then d^* stands for the tuple d_1^*, \ldots, d_n^* of their names. If t is a tuple t_1, \ldots, t_n of ground terms then t^I is the tuple t_1^I, \ldots, t_n^I of values assigned to them by I.

Here-and-there. The first-order logic of here-and-there, Pearce and Valverde (2004, 2005), introduced by forms a monotonic base for stable model semantic (Gelfond and Lifschitz 1988, 1991). The many-sorted case was recently studied by Fandinno and Lifschitz (2022). By At(I) we denote the set of ground atoms of the form $p(\mathbf{d}^*)$ such that $I \models p(\mathbf{d}^*)$, where p is a predicate symbol and d is a tuple of elements of domains of I. An *HT-interpretation* of σ is a pair $\langle \mathcal{H}, I \rangle$, where I is an interpretation of σ , and \mathcal{H} is a subset of At(I). (In terms of Kripke models with two worlds, \mathcal{H} describes the predicates in the here-world and I captures the there-world). The satisfaction relation \models_{ht} between HT-interpretation $\langle \mathcal{H}, I \rangle$ of σ and a sentence F over σ^I is defined recursively:

- $\langle \mathcal{H}, I \rangle \models_{\text{tr}} p(\mathbf{t}), \text{ if } p((\mathbf{t}^I)^*) \in \mathcal{H}$
- $\langle \mathcal{H}, I \rangle \models_{\text{tt}} t_1 = t_2 \text{ if } t_1^I = t_2^I;$
- $\langle \mathcal{H}, I \rangle \models_{\operatorname{ht}} F \wedge G \text{ if } \langle \mathcal{H}, I \rangle \models_{\operatorname{ht}} F \text{ and } \langle \mathcal{H}, I \rangle \models_{\operatorname{ht}} G;$
- $\bullet \ \langle \mathcal{H}, I \rangle \models_{\overline{\mathrm{ht}}} F \lor G \text{ if } \langle \mathcal{H}, I \rangle \models_{\overline{\mathrm{ht}}} F \text{ or } \langle \mathcal{H}, I \rangle \models_{\overline{\mathrm{ht}}} G;$
- $\langle \mathcal{H}, I \rangle \models_{\overline{ht}} F \to G \text{ if (i) } \langle \mathcal{H}, I \rangle \not\models_{\overline{ht}} F \text{ or } \langle \mathcal{H}, I \rangle \models_{\overline{ht}} G, \text{ and}$ (ii) $I \models F \to G;$
- $\langle \mathcal{H}, I \rangle \models_{\overline{ht}} \forall X F(X) \text{ if } \langle \mathcal{H}, I \rangle \models_{\overline{ht}} F(d^*) \text{ for each } d \in |I|^s$, where s is the sort of X;
- $\langle \mathcal{H}, I \rangle \models_{\overline{ht}} \exists X F(X) \text{ if } \langle \mathcal{H}, I \rangle \models_{\overline{ht}} F(d^*) \text{ for some } d \in |I|^s$, where s is the sort of X.

If $\langle \mathcal{H}, I \rangle \models_{\overline{ht}} F$ holds, we say that $\langle \mathcal{H}, I \rangle$ satisfies F and that $\langle \mathcal{H}, I \rangle$ is an *HT-model* of F. If two formulas have the same HT-models then we say that they are *HT-equivalent*.

It is easy to see that (At(I), I) is an HT-model of a sentence F whenever I is a model of F. About a model Iof a theory Γ , we say it is *stable* if, for every proper subset \mathcal{H} of At(I), HT-interpretation $\langle \mathcal{H}, I \rangle$ does not satisfy Γ . In application to theories of a single sort, this definition is equivalent to the original definition by Pearce and Valverde (2004, 2005). In addition, if the theory is finite, then this definition of a stable model is also equivalent to the definition of such model in terms of the operator SM (Ferraris, Lee, and Lifschitz 2007, 2011), when all predicate are considered to be intensional. We generalize the distinction between intensional and extensional predicates in the next section.

Disjunctive logic programs. A disjunctive rule is a formula of the form $Head \leftarrow Body$ where Head is a list of atomic formulas and Body is a list of *literals*, that is, atomic formulas possibly preceded by one or two occurrences of negation *not*. A disjunctive program is a (possibly infinite) set of disjunctive rules. We identify each rule $Head \leftarrow Body$

with the universal closure of the formula $B \to H$, where B is the conjunction of all literals in *Body* after replacing *not* by \neg ; and H is the disjunction of all atomic formulas in *Head*. We often write rules as formulas $B \to H$ omitting the reference to the universal closure. For instance, rule (1) is understood as the universal closure of formula

$$on(B, L, T) \land \neg non(B, L, T) \to on(B, L, T+1).$$
(4)

We may also write rule (1) in this form. We assume that variable T in rules (1-3) is of sort integer. In examples, we assume that any variable that is used as argument of an arithmetic function is of this sort. Under these assumptions, the definition of stable models of sets of sentences stated earlier also applies to (possible infinite) disjunctive logic programs with arithmetic. Note that these assumptions are enough to showcase all examples in this paper. For a more general characterization of how sorts are assigned to variables, we refer to Fandinno et al. (2020) and Lifschitz (2021).

Stable Models with Intensionality Statements

The distinction between *extensional* and *intensional* predicates (Ferraris, Lee, and Lifschitz 2011) describes the inherent meaning of a group of rules in a precise way; and it is similar to the distinction between input and non-input atoms by Oikarinen and Janhunen (2008). It is sometimes convenient to treat words extensional and intensional as synonymous to words *input* and *defined*, respectively. Intuitively, the meaning of input predicate symbols is not fixed within the considered group of rules; these rules may constrain the interpretations of such predicate symbols but they do not "define" them. To the contrary, intensional predicate symbols can be viewed as defined by the group. For instance, consider rules (1-3) that we denote as Π_{block} . Let us elaborate on the meaning of Π_{block} . Intuitively, Π_{block} captures the behavior of the property on/3 that changes when relevant movement occurs and otherwise it obeys the commonsense law of inertia (i.e., that things stay as they are unless forced to change). To obtain such a reading of Π_{block} predicates move/3 and location/3 should be declared as extensional. On the other hand, predicates on/3 and non/3 should be declared as intensional as their behavior is defined by rules in Π_{block} . It looks as such separation of predicate symbols allows us to identify Π_{block} with its intuitive meaning and yet the devil is in the details. The reading that we have stated fails to mention the special case of the initial situation. At the initial state, i.e., when T = 0, we do not assume that rules in Π_{block} define on/3, rather that these values are specified elsewhere. Thus, we would like to declare that on/3 is extensional in the initial situation (when T = 0) and intensional in all other situations (when $T \neq 0$). However, the granularity desired for this example is not currently possible because the distinction between extensional and intensional by Ferraris, Lee, and Lifschitz (2011) is made at the predicate level, disregarding the context provided by their arguments. We address this issue here.

We generalize the idea of distinguish between extensional and intensional predicate symbols by allowing the possibility to declare circumstances under which a predicate symbol is considered to be intensional (if these circumstances are not the case the predicate symbol is considered extensional). This is achieved by associating a first-order formula with each predicate constant. For instance, in Π_{block} associating predicate symbol on/3 with the formula $T \neq 0$ will result in proper treatment of its rules respecting not only their core meaning but also the corner case of the initial situation.

Formally, we identify each predicate symbol p/n with an atom of the form $p(X_1, \ldots, X_n)$, where X_1, \ldots, X_n are pairwise distinct variables of appropriate sorts. An *intensionality statement* λ over a signature σ is a function mapping each predicate symbol p/n in σ to a formula $F(X_1, \ldots, X_n)$ such that

• X_1, \ldots, X_n are the only free variables in $F(X_1, \ldots, X_n)$,

• every predicate symbol q/m in F satisfies $\lambda(q/m) \equiv \bot$. We abbreviate $\lambda(p/n)(X_1, ..., X_n)$ as $\lambda^p(X_1, ..., X_n)$ when arity n is clear from the context. We say that a predicate symbol is (purely) *intensional/extensional* when it is associated with a valid/unsatisfiable formula, respectively.

Consider, for instance, predicate symbol on/3 and let $\lambda^{on}(B, L, T)$ be formula $T \neq 0$. Intuitively, this intensionality statement states that all ground atoms formed by predicate constant p with the third argument different from 0 are intensional; if the third argument is 0, then they are extensional. The complete intensionality statement, which we call β , for our motivating example follows:

$$\beta^{on}(B, L, T) \text{ is } T \neq 0$$

$$\beta^{non}(B, L, T) \text{ is } \top$$

$$\beta^{move}(B, L, T) \text{ is } \perp$$

$$\beta^{location}(L) \text{ is } \perp$$

$$\beta^{\prec}(X_1, X_2) \text{ is } \perp \text{ for any comparison } \prec$$
(5)

It states that predicate *move* and all comparison symbols are purely extensional, *non* is purely intensional and *on* is intensional under all circumstances except the initial situation.

Let σ be a signature and λ be an intensionality statement over σ . By $EM(\lambda)$, we denote the set consisting of a sentence of the form

$$\forall \mathbf{X} \left(\neg \lambda^p(\mathbf{X}) \to p(\mathbf{X}) \lor \neg p(\mathbf{X}) \right)$$

for every predicate symbol p/n in σ , where **X** is a tuple of variables of the appropriate length and sort (in the sequel we adopt this convention and use **X** to denote tuples of variables). For a theory Γ over σ , we say that an interpretation I is λ -stable if it is a stable model of $\Gamma \cup EM(\lambda)$. Note that, if every predicate symbol is intensional in λ , then the λ -stable models coincide with the stable models.

For example, let F be the rule

$$p(X,1) \to p(X,2) \tag{6}$$

and let $\lambda^p(X_1, X_2)$ be formula $X_2 = 2$ (we assume both arguments of p/2 be of sort integer). The intensionality statement formula $EM(\lambda)$ is the universal closure of

$$\neg (X_2 = 2) \rightarrow p(X_1, X_2) \lor \neg p(X_1, X_2).$$

The following sets of ground atoms correspond to the four λ -stable models of F with domain $\{1, 2\}$

 $\emptyset \quad \{p(1,1), p(1,2)\} \quad \{p(2,1), p(2,2)\} \\ \{p(1,1), p(2,1), p(1,2), p(2,2)\},$

where we list the atoms that are true in these models (as customary in logic programming). As a more elaborated example, we can see that an interpretation I that satisfies

$$\begin{aligned} location^{I} &= \{l_{1}, l_{2}\} & move^{I} &= \{(b, l_{2}, 0)\} \\ on^{I} &= \{(b, l_{1}, 0)\} \cup \{(b, l_{2}, t') \mid t' > 0\} \\ non^{I} &= \{(b, l_{2}, 0)\} \cup \{(b, l_{1}, t') \mid t' > 0\} \end{aligned}$$

is a β -stable model of Π_{block} .

Strong Equivalence with Intensionality Statements

Informally, two programs are strongly equivalent if one can replace the other in any context without changing the stable models of the program (Lifschitz, Pearce, and Valverde 2001, 2007). Let Γ_1 and Γ_2 be theories over the same signature σ . We say that sets Γ_1 and Γ_2 are strongly equivalent with respect to an intensionality statement λ (or, simply λ -strongly equivalent) if $\Gamma_1 \cup \Delta$ and $\Gamma_2 \cup \Delta$ have the same λ -stable models for any theory Δ over σ .

By definition of λ -stable model, we immediately obtain the following sufficient condition for λ -strong equivalence.

Proposition 1 If $\Gamma_1 \cup EM(\lambda)$ and $\Gamma_2 \cup EM(\lambda)$ have the same HT-models, then Γ_1 and Γ_2 are λ -strongly equivalent.

Proposition 1 allows us to conclude that we can replace, without changing the β -stable models, rule (1) in Π_{block} by rules

$$on(B, L, T+1) \leftarrow on(B, L, T),$$
(7)
$$not \ non(B, L, T+1), \ T < t$$

$$on(B, L, T+1) \leftarrow on(B, L, T),$$

$$not \ non(B, L, T+1), \ T \ge t$$
(8)

where t is any positive integer. Note that (7) and (8) are obtained from rule (1) by respectively adding T < t and $T \ge t$ to its body. In fact (1) is λ -strong equivalent to to $\{(7), (8)\}$ for every λ where comparison symbols are extensional. Analogous rewritings can be performed on the remaining rules of Π_{block} . By $\Pi_{block}^{<}$ we denote the program obtained from Π_{block} by adding T < t to the body of rules (1) and (2), and adding $T \le t$ to the body of rule (3). By $\Pi_{block}^{>}$ we denote the program obtained from Π_{block} by adding T < t to the body of rule (3). By $\Pi_{block}^{>}$ we denote the program obtained from Π_{block} by adding $T \ge t$ to the body of rule (3). Programs Π_{block} and $\Pi_{block} \cup \Pi_{block}^{>}$ are β -strongly equivalent. We use program $\Pi_{block}^{<} \cup \Pi_{block}^{>}$ to showcase our Splitting method in the following section. In particular, the proposed method allows us to split this program into its two subcomponents $\Pi_{block}^{<}$ and $\Pi_{block}^{>}$.

Splitting Theorem with Intensionality Statements

To make the key results of this work comprehensible, we describe our Splitting method using two settings. First, we introduce the core ideas of the method in a restricted context, where considered theories are composed of disjunctive rules. Then, we generalize the method to arbitrary theories.

Splitting Disjunctive Programs

The statement of the proposed Splitting Theorem refers to the concept of the predicate dependency graph defined below. This definition, requires the notion of a *partition* of an intensionality statement. Formally, given two intensionality statement, λ_1 and λ_2 , over a signature σ by $\lambda_1 \sqcup \lambda_2$ we denote the intensionality statement defined as

$$(\lambda_1 \sqcup \lambda_2)^p (X_1, \dots, X_n) = \lambda_1^p (X_1, \dots, X_n) \lor \lambda_2^p (X_1, \dots, X_n).$$

for every predicate symbol p in the signature. By $\lambda_1 \sqcap \lambda_2$, we denote the intensionality statements defined as

$$(\lambda_1 \sqcap \lambda_2)^p (X_1, \dots, X_n) = \lambda_1^p (X_1, \dots, X_n) \land \lambda_2^p (X_1, \dots, X_n).$$

for every predicate symbol p. Intensionality statements λ_1 and λ_2 are *equivalent* if the universal closure of formula

$$\lambda_1^p(X_1,\ldots,X_n) \leftrightarrow \lambda_2^p(X_1,\ldots,X_n)$$

is logically valid for every predicate symbol p. If λ_1 and λ_2 are equivalent, then we write $\lambda_1 \equiv \lambda_2$. By λ_{\top} and λ_{\perp} we denote the intensionality statements where all predicate symbols are intensional and extensional, respectively. Intensionality statements λ_1 and λ_2 are called *disjoint* if $\lambda_1 \sqcap \lambda_2 \equiv \lambda_{\perp}$. A set of intensionality statements $\{\lambda_1, \ldots, \lambda_k\}$ is called a *partition* of some intensionality statement λ if $\lambda_1 \sqcup \ldots \sqcup \lambda_k \equiv \lambda$ and $\lambda_i \sqcap \lambda_j \equiv \lambda_{\perp}$ for all $i \neq j$. For instance, we can see that $\{\beta_1, \beta_2\}$ is a partition of intensionality statement β , where β_1 and β_2 are obtained from β by modifying the formulas associated with predicate constants *on* and *non* as follows:

$$\beta_1^{on}(B, L, T) \text{ is } T \neq 0 \land T \leq t$$

$$\beta_1^{non}(B, L, T) \text{ is } T \leq t$$

$$\beta_2^{on}(B, L, T) \text{ is } T > t$$

$$\beta_2^{non}(B, L, T) \text{ is } T > t.$$
(9)

An occurrence of a predicate constant, or any other subexpression, in a formula is called *negated* if it occurs in the scope of negation (i.e in the antecedent of any implication of the form $F \to \bot$); *nonnegated* otherwise. As an example, predicate constant *on* occurs nonnegated in the antecedent of (4), while *non* does not. Given a program II (understood as a set of sentences) and a partition Λ of some intensionality statement λ , the (*directed*) graph of dependencies with *respect to* Λ , denoted $\mathfrak{G}_{\Lambda}(\Pi)$, is defined as follows:

- Its vertices are pairs (p, λ_i) such that p is a predicate symbol occurring in Π, λ_i ∈ Λ is an intensionality statement and ∃X λ_i^p(X) is satisfiable.
- It has an edge from (p, λ_i) to (q, λ_j) when for some rule $B \to H$ of Π ,
 - $p(\mathbf{t})$ occurs in H, and
 - there is a nonnegated occurrence of $q(\mathbf{r})$ in B and
 - the following sentence is satisfiable

$$\exists \mathbf{X} \left(B \wedge p(\mathbf{t}) \wedge \lambda_i^p(\mathbf{t}) \wedge \lambda_j^q(\mathbf{r}) \right), \tag{10}$$

where **X** are the free variables in $B \rightarrow H$.

We say that a partition $\Lambda = \{\lambda_1, \ldots, \lambda_k\}$ is *separable* (on $\mathfrak{G}_{\Lambda}(\Pi)$) when

every infinite walk v_1, v_2, \ldots of $\mathfrak{G}_{\Lambda}(\Pi)$ visits at most one λ_i infinitely many times, that is, there is some $i \in \{1, \ldots, k\}$ s.t $\{l \mid v_l = (p, \lambda_j)\}$ is finite for all $j \neq i$.

Note that, if (the signature of) Π is finite, every possible infinite walk must contain a cycle and, thus, this condition is equivalent to saying that every cycle is confined to one component of the partition. In general, checking cycles is not enough for infinite programs, even when these programs contain no variables (Harrison and Lifschitz 2016).

Continuing with our running example, let $\Pi'_{block} = \Pi^{<}_{block} \cup \Pi^{>}_{block}$ and $\Lambda_{block} = \{\beta_1, \beta_2\}$ be partition of β . Then, graph $\mathfrak{G}_{\Lambda_{block}}(\Pi'_{block})$ consists of vertices

 $(on, \beta_1), (on, \beta_2), (non, \beta_1), (non, \beta_2)$

and contains five edges: one leading from (on, β_1) to itself, another one leading from (on, β_2) to itself, one leading from (on, β_2) to (on, β_1) —induced by rule (8)— and two leading from (non, β_i) to (on, β_i) with $1 \le i \le 2$. Note that there is no edge from (on, β_1) to (on, β_2) . For instance, we can see that rules (7) and (8) do not induce such edge because the sentence of the form (10) corresponding to these rules contain conjuncts $T + 1 \le t$ and T > t, which make them unsatisfiable. Conjunct $T + 1 \le t$ is obtained from $\beta_1^{on}(B, L, T+1)$ corresponding to the head occurrence of on/3. The other conjunct is obtained from $\beta_2^{on}(B, L, T)$ corresponding to its body occurrence. Therefore, partition Λ_{block} is separable.

A program Π is *negative* on some intensionality statement λ if, for every rule $B \to H$ of Π and every atom $p(\mathbf{t})$ occurring in H, the following sentence is unsatisfiable

$$\exists \mathbf{X} (B \wedge p(\mathbf{t}) \wedge \lambda^p(\mathbf{t})),$$

where **X** are the free variables in $B \to H$. For instance, programs $\Pi_{block}^{<}$ and $\Pi_{block}^{>}$ are negative on β_2 and β_1 , respectively. We provide a part of an argument for the claim that $\Pi_{block}^{<}$ is negative on β_2 . In particular, we argue that rule (7) is negative on β_2 . This rule has on(B, L, T + 1) in its head and T < t in its body. It is sufficient to show that a formula of the form

 $\exists BLT \, (F \wedge T < t \wedge \beta_2^{on}(B, L, T+1)),$

is unsatisfiable. Recall that $\beta_2^{on}(B, L, T+1)$ is T+1 > t.

Theorem 1 (Splitting disjunctive programs) Let $\Pi = \Pi_1 \cup \dots \cup \Pi_n$ be a disjunctive program and $\Lambda = \{\lambda_1, \dots, \lambda_n\}$ be a partition of λ such that

- Λ is separable on $\mathfrak{G}_{\Lambda}(\Pi)$; and
- each Π_i is negative on λ_j for all $j \neq i$.

Then, for any interpretation I, the following two statements are equivalent

- *I* is a λ -stable model of Π , and
- *I* is a λ_i -stable model of \prod_i for all $1 \le i \le n$.

Recall that partition Λ_{block} is separable on $\mathfrak{G}_{\Lambda_{block}}(\Pi'_{block})$ and that programs Π^{\leq}_{block} and Π^{\geq}_{block} are negative on β_2 and β_1 , respectively. Therefore, the theorem on Splitting disjunctive programs allows us to conclude that the β -stable models of Π'_{block} are those interpretations that are at the same time β_1 -stable models of Π_{block}^{\leq} and β_2 -stable models of Π_{block}^{\geq} . As a final note, recall that Π_{block} and Π'_{block} are β -strongly equivalent and, thus, they have the same β -stable models. Hence, the statement that we can split program Π_{block} into programs Π_{block}^{\leq} and Π_{block}^{\geq} is a consequence of these two facts stemming from presented Theorem 1 and Proposition 1, respectively.

Splitting Arbitrary Theories

We now generalize Theorem 1 on Splitting disjunctive programs to the case of arbitrary sets of sentences. We start by motivating this generalization and then proceed to its formalization. Complex formulas not fitting into the syntax of disjunctive rules naturally appear as a result of translating some common constructs of logic programs into first-order formulas. These constructs include basic arithmetic operations such as division, which are part of the ASP-Core-2 (Calimeri et al. 2012), and more advanced features such as intervals, or conditional literals. Intervals and conditional literals are part of the language of the solver clingo (Gebser et al. 2015) and are commonly used in practice. For instance, the following rule contains a conditional literal holds(B) : body(R, B) in its body:

 $holds(H) \leftarrow head(R, H), holds(B) : body(R, B).$ (11) Hansen and Lierler (2022) showed that this rule can be understood as an abbreviation for the following first-order sentence containing a nested universal quantifier and a nested implication

$$\forall RH \big(head(R,H) \land$$

 $\forall B(body(R, B) \rightarrow holds(B)) \rightarrow holds(H))$. (12) Rules of with condition literals are common in metaprogramming (Kaminski et al. 2021), where reification of constructs are utilized to build ASP-based reasoning engines that *may* go beyond ASP paradigm itself. Kaminski et al. (2021) presented multiple examples of meta-programming. Here, we use rule (11) to showcase a simple use of this technique. Indeed, it can be used to express the meaning of a *definite* rule – a disjunctive rule whose *Head* consists of single atomic formula and *Body* contains no occurrences of negation *not* – encoded as a set of facts. For example, definite program

$$\begin{array}{l}
a \leftarrow b \\
b \leftarrow c
\end{array} \tag{13}$$

can be encoded by the following facts complemented with rule (11)

$$head(r1, a) \quad body(r1, b)$$
 (14)

$$head(r2, b) \quad body(r2, c).$$
 (15)

Clearly, we can split the program listed in (13) into two subprograms, each consisting of a single rule. We may expect the possibility of splitting its respective meta encoding as well. Consider sentences

$$\forall X (head(r1, X) \land \\ \forall W (body(r1, W) \to holds(W)) \to holds(X))$$

$$\forall X (head(r2, X) \land \\ \forall W (body(r2, W) \to holds(W)) \to holds(X))$$

$$(17)$$

corresponding to a "partially instantiated" portion of the meta-encoding of program (13). Ideally, the program corresponding to sentences (14-17) would be "identified" with subprograms (14,16) and (15,17) by splitting. Yet, Theorem 1 cannot support such a claim as (16) and (17) are not disjunctive rules. This claim is also not supported by other Splitting Theorems in the literature (Ferraris et al. 2009; Harrison and Lifschitz 2016) due to the "positive nonnegated" dependency induced by these rules in the predicate hold/1 (we define the concept of positive nonnegated dependency below).

Formalization. The first thing to note is that one direction of the Splitting Theorem always holds without the need to inspect the dependency graph.

Proposition 2 Let $\Gamma = \Gamma_1 \cup \cdots \cup \Gamma_n$ be some theory and let $\Lambda = \{\lambda_1, \ldots, \lambda_n\}$ be a partition of intensionality statement λ . If I is a λ -stable model of Γ , then I is a λ_i -stable model of Γ_i for all $1 \leq i \leq n$.

For the other direction, we have to generalize the notions of separability and of being negative to the case of arbitrary sentences. Similar to the Splitting Theorem by Ferraris et al. (2009), this generalization relies on the notions of strictly positive, positive nonnegated, and negative nonnegated occurrence of an expression. An occurrence of an expression is called *positive* if the number of implications containing that occurrence in the antecedent is even; and strictly positive if that number is 0. It is called *negative* if that number is odd. As in the case of logic programs, an occurrence of an expressions is called *negated* if it belongs to a subformula of the form $\neg F$ (that is, $F \rightarrow \bot$); and *nonnegated* otherwise. A rule is a strictly positive occurrence of a formula of the form $B \rightarrow H$. Therefore, the rules of a set of disjunctive rules are exactly all its disjunctive rules, but this definition also covers arbitrary nested rules. For instance, sentence $(c \rightarrow (a \rightarrow b)) \lor d$ contains two occurrences of rules, namely, $a \rightarrow b$ and $c \rightarrow (a \rightarrow b)$.

To extend Theorem 1 to arbitrary formulas, we use these notions to make the construction of a counterpart of sentence (10) recursive over the formula. In addition, we ought to incorporate in this construction a *context* that carries information about the rest of the program. To observe the need for this context, note that whether rules (16) and (17) can be separated into differ subprograms depends on the extension of head/2 and body/2. For instance, adding the fact body(r2, a) to our program creates a dependency that cannot be broken. In fact, this new program has the same meaning as a non-splittable program consisting of rules $a \leftarrow b$ and $b \leftarrow c$, a.

Given a formula F with free variables \mathbf{X} and a theory Ψ , by F^{Ψ} we denote formula F itself if $\Psi \cup \{\exists \mathbf{X} F\}$ is satisfiable; and \bot otherwise. Let now $p(\mathbf{t})$ be a strictly positive (resp. positive nonnegated or negative nonnegated) "distinguished" occurrence of predicate p in F and \mathbf{Y} a list of variables not occurring in F of the same length and sorts as \mathbf{t} . We recursively build formula $Sp_{\Psi}(F)$ (resp. $Pnn_{\Psi}(F)$) and $Nnn_{\Psi}(F)$) for this occurrence as described next; the construction of these three formulas only differs in the case of implication, so we use the metavariable $Fun_{\Psi}(F)$ for the common cases:

- If $\Psi \cup \{\exists \mathbf{X} F\}$ is unsatisfiable, then $Fun_{\Psi}(F) = \bot$.
- If Ψ ∪ {∃X F} is satisfiable and the distinguished occurrence p(t) does not occur in F, then Fun_Ψ(F) = F.

Otherwise,

- $Fun_{\Psi}(p(\mathbf{t})) = p(\mathbf{t}) \wedge \mathbf{Y} = \mathbf{t};$
- $Fun_{\Psi}(F_1 \wedge F_2) = Fun_{\Psi}(F_1) \wedge Fun_{\Psi}(F_2);$
- $Fun_{\Psi}(F_1 \vee F_2) = Fun_{\Psi}(F_i)$ with F_i containing the occurrence (*i* is 1 or 2);
- $Fun_{\Psi}(\forall XF) = Fun_{\Psi}(\exists XF) = \exists X(Fun_{\Psi}(F));$
- $Sp_{\Psi}(F_1 \to F_2) = F_1^{\Psi} \wedge Sp_{\Psi}(F_2);$
- $Pnn_{\Psi}(F_1 \to F_2) =$ $\begin{cases} Nnn_{\Psi}(F_1) & \text{if } F_1 \text{ contains the occurrence} \\ F_1^{\Psi} \wedge Pnn_{\Psi}(F_2) & \text{otherwise} \end{cases}$
- $Nnn_{\Psi}(F_1 \rightarrow F_2) =$ $\begin{cases}
 Pnn_{\Psi}(F_1) & \text{if } F_1 \text{ contains the occurrence} \\
 F_1^{\Psi} \wedge Nnn_{\Psi}(F_2) & \text{otherwise}
 \end{cases}$

Note that $Pnn_{\Psi}(F)$ and $Nnn_{\Psi}(F)$ are mutually recursive due to the case of implication. Examples illustrating the construction of these formulas follow the next definition.

Given theories Γ and Ψ and a partition Λ of some intensionality statement λ , the *(directed) graph of positive dependencies with respect to* Λ *and under context* Ψ , denoted $\mathfrak{G}_{\Lambda,\Psi}(\Gamma)$, is defined as follows:

- Its vertices are pairs (p, λ_i) such that p is a predicate symbol, λ_i ∈ Λ is an intensionality statement and theory Ψ ∪ {∃X λ_i^p(X)} is satisfiable.
- It has an edge from (p, λ_i) to (q, λ_j) when for some rule B → H of Γ, the following conditions hold
 - there is a strictly positive occurrence p(t) in H,
 - there is a positive nonnegated occurrence of $q(\mathbf{r})$ in B,
 - the theory below is satisfiable,
- $\Psi \cup \left\{ \exists \mathbf{XYZ} \left(Pnn_{\Psi}(B) \land Sp_{\Psi}(H) \land \lambda_{i}^{q}(\mathbf{Y}) \land \lambda_{i}^{p}(\mathbf{Z}) \right) \right\},$ (18)

where **X** are the free variables in $B \to H$, and **Y** and **Z** respectively are the free variables in formulas $Pnn_{\Psi}(B)$ and $Sp_{\Psi}(H)$ that are not in **X**; in the construction of $Sp_{\Psi}(H)$ and $Pnn_{\Psi}(B)$ we consider occurrences $p(\mathbf{t})$ in H and $q(\mathbf{r})$ in B, respectively.

Take sentence $F_1 = q \lor (r \land p)$ and rule $F_1 \to p$. Note that p occurs positively nonnegated in F_1 . We consider this occurrence p when constructing formulas $Pnn_{(\cdot)}(F_1)$ below. Assume that p is intensional in λ (i.e., $\lambda^p \equiv \top$) and that Λ is a partition of λ . On the one hand, if we consider the empty context, then $Pnn_{\emptyset}(F_1)$ is $r \land p$ and the dependency graph of $F_1 \to p$ with respect to Λ and the empty context contains a reflexive edge on vertex (p, λ) . On the other hand, if we take context $\Psi_1 = \{\neg r\}$, then $Pnn_{\Psi_1}(F_1)$ is \bot and the dependency graph with respect to Λ and Ψ_1 is empty. This example shows that, even on propositional formulas, the use of context leads to less dependencies than previous approaches such as (Ferraris et al. 2009; Harrison and Lifschitz 2016). This leads to the fact that the Splitting Theorem introduced in the sequel is applicable to more theories as it relies on this new notion of dependencies. This difference may appear even using the empty context. Take a theory consisting of formula $(q \lor (\bot \land p)) \to p$. It turns out that p never contributes any edge in our approach under any context. For the same theory, p always forms a dependency in the earlier papers. Consider now a small variation. Let p be a unary predicate and let $F_2 = q \lor \exists X (r \land p(X))$. Note that $Pnn_{\emptyset}(F_2) =$ $\exists X(r \land p(X) \land Y = X)$ and $Pnn_{\Psi_1}(F_2) = \bot$, where we construct these formulas for occurrence p(X). Under the empty context, analogously to the previous example, the dependency graph of $F_2 \rightarrow p(1)$ contains a reflexive edge on vertex (p, λ) . Under context Ψ_1 , the dependency graph is empty.

Construction of $Sp_{\Psi}(F)$, $Pnn_{\Psi}(F)$ and $Nnn_{\Psi}(F)$ reference a "distinguished" occurrence, as there may be multiple occurrences of the same atomic formula. For instance, there are two occurrences of formula p(X) in sentence F_3 defined as $\exists X(X = a \land p(X)) \lor (X = b \land p(X));$ for this sentence $Pnn_{\emptyset}(F_3)$ is $\exists X(X = a \land p(X) \land Y = X)$, when the first occurrence of p(X) is considered; and $\exists X(X=b \land p(X) \land Y=X)$, when the second one is considered. Take Ψ_2 to be the theory containing sentence $\neg p(b)$ only, then $Pnn_{\Psi_2}(F_3)$ remains unmodified for the first occurrence of p, but becomes \perp for the second one. As a consequence, in a context where p(b) is false, the first occurrence may generate an edge in the dependency graph, while the second occurrence does not generate any. Note that in all considered examples so far, $Sp_{(\cdot)}(\cdot)$ is the same as $Pnn_{(\cdot)}(\cdot)$ due to the lack of implications.

We now explore the meta encoding example introduced earlier. Let B denote the antecedent of rule (16) and H denote its consequent holds(X). Under the empty context, $Pnn_{\emptyset}(B)$ for the only occurrence of holds/1 in B is

$$head(r1, X) \land \exists W (body(r1, W) \land holds(W) \land Y = W);$$

and $Sp_{\emptyset}(H)$ for the only occurrence of holds/1 in H is $holds(X) \wedge X = Z$. Let γ_1 and γ_2 be intensionality statements where *head* and *body* are extensional and

$$\gamma_1^{holds}(X_1)$$
 is $X_1 = a$ $\gamma_2^{holds}(X_1)$ is $X_1 = b$.

Then, the singleton theory with the existential closure of

$$Pnn_{\emptyset}(B) \wedge Sp_{\emptyset}(H) \wedge \gamma_{j}^{holds}(Y) \wedge \gamma_{i}^{holds}(Z)$$
(19)

is satisfiable for any $i, j \in \{1, 2\}$. For a theory containing rule (16), the graph of positive dependencies with respect to $\{\gamma_1, \gamma_2\}$ and under the empty context contains an edge from vertex (*holds*, γ_1) to vertex (*holds*, γ_2), and vice-versa; together with reflexive edges in both vertices. Let Ψ_3 be the theory consisting of the universal closures of formulas

$$\begin{aligned} head(r1,X) \leftrightarrow X &= a \\ body(r1,X) \leftrightarrow X &= b \\ body(r2,X) \leftrightarrow X &= c. \end{aligned}$$

Then, $Sp_{\Psi_3}(H) = Sp_{\emptyset}(H)$ and $Pnn_{\Psi_3}(B) = Pnn_{\emptyset}(B)$. The union of Ψ_3 and the singleton theory consisting of the existential closure of formula (19) is satisfiable only with i = 1 and j = 2. Therefore, under this context, the graph of positive dependencies from the previous example contains an edge from $(holds, \gamma_1)$ to $(holds, \gamma_2)$ but not vice-versa. There are no reflexive edges either. In fact, under this context the positive dependency graph of (16-17) only contains this edge.

Let us now generalize the notion of being a negative occurrence to the case of arbitrary theories. We say that a theory Γ is Ψ -negative on some intensionality statement λ if, for every rule $B \to H$ of Γ and every strictly positive occurrence $p(\mathbf{t})$ in H, the following theory is unsatisfiable $\Psi \cup \{\exists \mathbf{XY} (B \land Sp_{\Psi}(H) \land \lambda^{p}(\mathbf{Y}))\}$, where \mathbf{X} consists of the free variables in $B \to H$; \mathbf{Y} consists of the free variables in $Sp_{\Psi}(H)$ that do not occur in \mathbf{X} ; and we consider occurrence $p(\mathbf{t})$ in H in the construction of $Sp_{\Psi}(H)$. We also say that theory Ψ is an λ -approximator of Γ if all the λ -stable models of Γ are models of Ψ .

Theorem 2 (Splitting Theorem) Let $\Gamma = \Gamma_1 \cup ... \cup \Gamma_n$, λ be an intensionality statement, and Ψ be an λ -approximator of Γ . Let $\Lambda = {\lambda_1, ..., \lambda_n}$ be a partition of λ such that

- Λ is separable on $\mathfrak{G}_{\Lambda,\Psi}(\Gamma)$; and
- each Γ_i is Ψ -negative on λ_j for all $j \neq i$.

Then, the following two statements are equivalent

- *I* is a λ -stable model of Π , and
- *I* is a model of Ψ and a λ_i -stable model of Π_i for all $i \in \{1, \ldots, n\}$.

Continuing with our meta encoding example, in addition to γ_1 and γ_2 , we define γ_3 to be the intensionality statement where head/2, body/2 are intensional and holds/1 is extensional. Let γ be $\gamma_1 \sqcup \gamma_2 \sqcup \gamma_3$. It is easy to see that $\{\gamma_1, \gamma_2, \gamma_3\}$ is a partition of γ . Let Γ_1 and Γ_2 be the singleton theories consisting of (16) and (17), respectively; let Γ_3 consist of facts in (14-15), and let Γ denote $\Gamma_1 \cup \Gamma_2 \cup \Gamma_3$. Partition $\Lambda = \{\gamma_1, \gamma_2, \gamma_3\}$ is separable on $\mathfrak{G}_{\Lambda, \Psi_3}(\Gamma)$. We also can see that Γ_i is negative on γ_j under context Ψ_3 for all $j \neq i$. Therefore, by Theorem 2, the γ -stable models of Γ are the models of Ψ_3 that are γ_i -stable models of Γ_i for all $i \in \{1, 2, 3\}$. The intent of context Ψ_3 is to carry information from one part of the theory into another. In our example all models of Ψ_3 are γ_3 -stable models of Γ_3 . Hence, we can simply say that the γ -stable models of Γ are the interpretations that are γ_i -stable models of Γ_i for all $i \in \{1, 2, 3\}$.

It is worth noting that the empty theory approximates any theory. When such theory is considered in the presented theorem, it more closely resembles the Splitting Theorem by Ferraris et al. (2009). In general, we can use the theory itself, its completion (Ferraris, Lee, and Lifschitz 2011; Fandinno and Lifschitz 2022) or the completion of a part of it as a "more precise" approximator.

Conclusions

The concept of intensionality statements introduced here provides us with a new granularity on considering semantics of logic programs and its subcomponents. It also paves a way to the refinement of earlier versions of the Spltting method. We generalized the conditions under which this method can be applied to first-order theories and show how the resulting approach covers more programs commonly used in practice. This generalization comes at a price. The conditions of the Splitting Theorem by Ferraris et al. (2009) are syntactic, while our conditions rely on verification of semantic properties. In fact, deciding whether there is an edge in our dependency graph is, in general, undecidable. For instance, a program containing rules $p \leftarrow q, t = 0$ and $q \leftarrow p$, where t is a polynomial, can be split into subprograms each containing of one of these rules only if the Diophantine equation in the body has no solutions. However, we illustrated that for many practical problems the Splitting result of this paper is applicable. In the future, we will investigate the possibility to utilize first-order theorem provers for checking the required semantic conditions.

Acknowledgements

We are thankful to Vladimir Lifschitz, Torsten Schaub and the anonymous reviewers for their comments on earlier versions of this draft. The first author is partially supported by the Nebraska EPSCoR 95-3101-0060-402 grant.

References

Cabalar, P.; Fandinno, J.; and Lierler, Y. 2020. Modular Answer Set Programming as a Formal Specification Language. *Theory and Practice of Logic Programming*, 20(5): 767–782.

Calimeri, F.; Faber, W.; Gebser, M.; Ianni, G.; Kaminski, R.; Krennwallner, T.; Leone, N.; Ricca, F.; and Schaub, T. 2012. ASP-Core-2: Input language format.

Fandinno, J.; Hansen, Z.; and Lierler, Y. 2022. Axiomatization of Aggregates in Answer Set Programming. In *Proceedings of the Thirty-sixth National Conference on Artificial Intelligence (AAAI'22)*, 5634–5641. AAAI Press.

Fandinno, J.; and Lifschitz, V. 2022. Verification of Locally Tight Programs. Under consideration for publication in Theory and Practice of Logic Programming.

Fandinno, J.; Lifschitz, V.; Lühne, P.; and Schaub, T. 2020. Verifying Tight Logic Programs with anthem and vampire. *Theory and Practice of Logic Programming*, 20(5): 735–750.

Ferraris, P.; Lee, J.; and Lifschitz, V. 2007. A New Perspective on Stable Models. In Veloso, M., ed., *Proceedings of the Twentieth International Joint Conference on Artificial Intelligence (IJCAI'07)*, 372–379. AAAI/MIT Press.

Ferraris, P.; Lee, J.; and Lifschitz, V. 2011. Stable models and circumscription. *Artificial Intelligence*, 175(1): 236–263.

Ferraris, P.; Lee, J.; Lifschitz, V.; and Palla, R. 2009. Symmetric Splitting in the General Theory of Stable Models. In Boutilier, C., ed., *Proceedings of the Twenty-first International Joint Conference on Artificial Intelligence (IJCAI'09)*, 797–803. AAAI/MIT Press.

Gebser, M.; Harrison, A.; Kaminski, R.; Lifschitz, V.; and Schaub, T. 2015. Abstract Gringo. *Theory and Practice of Logic Programming*, 15(4-5): 449–463.

Gebser, M.; Kaminski, R.; Kaufmann, B.; and Schaub, T. 2019. Multi-shot ASP solving with clingo. *Theory and Practice of Logic Programming*, 19(1): 27–82.

Gelfond, M.; and Lifschitz, V. 1988. The Stable Model Semantics for Logic Programming. In Kowalski, R.; and Bowen, K., eds., *Proceedings of the Fifth International Conference and Symposium of Logic Programming (ICLP'88)*, 1070–1080. MIT Press.

Gelfond, M.; and Lifschitz, V. 1991. Classical Negation in Logic Programs and Disjunctive Databases. *New Generation Computing*, 9: 365–385.

Hansen, Z.; and Lierler, Y. 2022. Semantics for Conditional Literals via the SM Operator. In *LPNMR*, volume 13416 of *Lecture Notes in Computer Science*, 259–272. Springer.

Harrison, A.; and Lifschitz, V. 2016. Stable models for infinitary formulas with extensional atoms. *Theory and Practice of Logic Programming*, 16(5-6): 771–786.

Kaminski, R.; Romero, J.; Schaub, T.; and Wanko, P. 2021. How to Build Your Own ASP-based System?! *Theory and Practice of Logic Programming*, 1–63.

Lifschitz, V. 2002. Answer set programming and plan generation. *Artificial Intelligence*, 138(1-2): 39–54.

Lifschitz, V. 2008. What Is Answer Set Programming? In Fox, D.; and Gomes, C., eds., *Proceedings of the Twenty-third National Conference on Artificial Intelligence* (AAAI'08), 1594–1597. AAAI Press.

Lifschitz, V. 2019. Answer Set Programming. Springer-Verlag.

Lifschitz, V. 2021. Transforming Gringo Rules into Formulas in a Natural Way. 421–434.

Lifschitz, V.; Morgenstern, L.; and Plaisted, D. 2008. Knowledge Representation and Classical Logic. In van Harmelen, F.; Lifschitz, V.; and Porter, B., eds., *Handbook of Knowledge Representation*, 3–88. Elsevier.

Lifschitz, V.; Pearce, D.; and Valverde, A. 2001. Strongly equivalent logic programs. *ACM Transactions on Computational Logic*, 2(4): 526–541.

Lifschitz, V.; Pearce, D.; and Valverde, A. 2007. A Characterization of Strong Equivalence for Logic Programs with Variables. In Baral, C.; Brewka, G.; and Schlipf, J., eds., *Proceedings of the Ninth International Conference on Logic Programming and Nonmonotonic Reasoning (LPNMR'07)*, volume 4483 of *Lecture Notes in Artificial Intelligence*, 188– 200. Springer-Verlag.

Lifschitz, V.; and Turner, H. 1994. Splitting a logic program. In *Proceedings of the Eleventh International Conference on Logic Programming*, 23–37. MIT Press.

Oikarinen, E.; and Janhunen, T. 2008. Achieving Compositionality of the Stable Model Semantics for smodels Programs. *Theory and Practice of Logic Programming*, 8(5-6): 717–761.

Pearce, D.; and Valverde, A. 2004. Towards a First Order Equilibrium Logic for Nonmonotonic Reasoning. In *Proceedings of European Conference on Logics in Artificial Intelligence (JELIA)*, 147–160.

Pearce, D.; and Valverde, A. 2005. A First Order Nonmonotonic Extension of Constructive Logic. *Studia Logica*, 30(2-3): 321–346.

Truszczyński, M. 2012. Connecting First-Order ASP and the Logic FO(ID) through Reducts. In Erdem, E.; Lee, J.; Lierler, Y.; and Pearce, D., eds., *Correct Reasoning: Essays* on Logic-Based AI in Honour of Vladimir Lifschitz, volume 7265 of Lecture Notes in Computer Science, 543–559. Springer-Verlag.

Locally Splittable Theories

This section introduces auxiliary definitions and results that will allow us to prove the Splitting Theorem stated in the main part of the paper.

Given a subset \mathcal{A} of At(I), we say that I is a \mathcal{A} -stable model of Γ if it is a stable model of $\Gamma \cup EM(I, \mathcal{A})$ where $EM(I, \mathcal{A})$ is the set containing disjunction $p(\mathbf{d}^*) \vee \neg p(\mathbf{d}^*)$ for every ground atom $p(\mathbf{d}^*)$ in $At(I) \setminus \mathcal{A}$. We can relate \mathcal{A} -stable models with λ -stable models as follows. By $At(I, \lambda)$ we denote the set of ground atoms of the form $p(\mathbf{d}^*)$ such that

$$I \models p(\mathbf{d}^*) \land \lambda^p(\mathbf{d}^*)$$

where p is a predicate symbol and d is a tuple of elements of domains of I.

Proposition 3 *I* is a λ -stable model of Γ iff *I* is a $At(I, \lambda)$ -stable model of Γ .

Proof. It is enough to show that $\langle \mathcal{H}, I \rangle \models_{\overline{ht}} EM(\lambda)$ iff $\langle \mathcal{H}, I \rangle \models_{\overline{ht}} EM(At(I, \lambda))$. Furthermore, for this is enough to show that, for every predicate symbol p,

$$\langle \mathcal{H}, I \rangle \models_{\overline{\mathrm{ht}}} \forall \mathbf{X} \left(\neg \lambda^p(\mathbf{X}) \to p(\mathbf{X}) \lor \neg p(\mathbf{X}) \right)$$
 (20)

iff

$$\langle \mathcal{H}, I \rangle \models_{ht} p(\mathbf{d}^*) \lor \neg p(\mathbf{d}^*) \text{ for every } p(\mathbf{d}^*) \text{ in } At(I) \setminus At(I, \lambda)$$
(21)

On the one hand, condition (20) holds iff

$$\langle \mathcal{H}, I \rangle \models_{\overline{\mathrm{ht}}} \neg \lambda^p(\mathbf{d}^*) \to p(\mathbf{d}^*) \lor \neg p(\mathbf{d}^*)$$

for every tuple of domain elements d. This holds iff one of the following conditions hold:

$$I \models_{\overline{ht}} \lambda^p(\mathbf{d}^*) \tag{22}$$

$$\langle \mathcal{H}, I \rangle \models_{\mathrm{ht}} p(\mathbf{d}^*) \lor \neg p(\mathbf{d}^*)$$
 (23)

for every tuple of domain elements d. Pick some $p(\mathbf{d}^*)$ in At(I). We proceed by cases. *Case 1*. Assume that $I \not\models p(\mathbf{d}^*)$. Then, $\langle \mathcal{H}, I \rangle \models_{\overline{ht}} \neg p(\mathbf{d}^*)$ and this implies (23). Hence, the result holds. *Case 2*. Assume that $I \not\models \lambda^p(\mathbf{d}^*)$. Then, condition (20) holds iff (23) and the result holds. *Case 3*. Assume that $I \models p(\mathbf{d}^*) \land \lambda^p(\mathbf{d}^*)$. This implies that $p(\mathbf{d}^*)$ belongs to $At(I, \lambda)$ and, thus, condition (21) vacuously follow. This also implies (22) and, thus, condition (20) follows.

For a sentence F over σ , an interpretation I of σ and a subset \mathcal{A} of At(I), we define the set $Sp_I(F)$ of (*strictly*) *positive atoms of* F with respect to I. Elements of this set are atoms of the signature σ^I that have the form $p(\mathbf{d}^*)$.¹ This set is defined recursively, as follows. If F is not satisfied by I then $Sp_I(F) = \emptyset$ (note how \bot is such a formula). Otherwise,

- $Sp_I(p(\mathbf{t})) = \{p((\mathbf{t}^I)^*)\};$
- $Sp_I(F_1 \wedge F_2) = Sp_I(F_1 \vee F_2) = Sp_I(F_1) \cup Sp_I(F_2);$

 $^{^1 \}text{For the definitions of the signature } \sigma^I$ and names $d^*,$ see Section .

- $Sp_I(F_1 \to F_2) = \begin{cases} Sp_I(F_2) & \text{if } I \models F_1 \\ \emptyset & \text{otherwise} \end{cases}$;
- $Sp_I(\forall XF(X)) = Sp_I(\exists XF(X))$ $\bigcup_{d \in |I|^s} Sp_I(F(d^*))$ if X is a variable of sort s.

We also define the sets $Pnn_I(F)$ and $Nnn_I(F)$ of positive nonnegated and negative nonnegated atoms of F with respect to I. These sets are defined recursively, as follows. If F is not satisfied by I then $Pnn_I(F) = Nnn_I(F) = \emptyset$. Otherwise,

- $Pnn_I(p(\mathbf{t})) = \{p((\mathbf{t}^I)^*)\};$
- $Pnn_I(F_1 \wedge F_2) = Pnn_I(F_1 \vee F_2) =$ $Pnn_I(F_1) \cup Pnn_I(F_2);$
- $Pnn_I(F_1 \to F_2) =$ $\begin{cases}
 Nnn_I(F_1) \cup Pnn_I(F_2) & \text{if } I \models F_1 \\
 \emptyset & \text{otherwise};
 \end{cases}$
- $Pnn_I(\forall XF(X)) = Pnn_I(\exists XF(X)) = \bigcup_{d \in |I|^s} Pnn_I(F(d^*))$ if X is a variable of sort s.
- $Nnn_I(p(\mathbf{t})) = \emptyset;$
- $Nnn_I(F_1 \wedge F_2) = Nnn_I(F_1 \vee F_2) =$ $Nnn_I(F_1) \cup Nnn_I(F_2);$
- $Nnn_I(F_1 \to F_2) =$ $\begin{cases}
 Pnn_I(F_1) \cup Nnn_I(F_2) & \text{if } I \models F_1 \\
 \emptyset & \text{otherwise};
 \end{cases}$
- $Nnn_I(\forall XF(X)) = Nnn_I(\exists XF(X)) = \bigcup_{d \in |I|^s} Nnn_I(F(d^*))$ if X is a variable of sort s.

Recall that a strictly positive occurrence of an implication $F_1 \to F_2$ in a formula F is called a *rule* of F. The *rules* of a theory Γ are all the rules of its sentences. An *instance* of a rule $F_1 \to F_2$ for an interpretation I of σ is a sentence over σ^I that can be obtained from $F_1 \to F_2$ by substituting names d^* for its free variables.

Given a set Γ of sentences, the graph of positive dependencies with respect to an interpretation I and set A of atoms, $\mathfrak{G}_{I,A}(\Gamma)$ is the directed graph defined as follows:

- Its vertices are elements of A.
- It has an edge from $p(\mathbf{d}^*)$ to $q(\mathbf{e}^*)$ iff, for some instance $F_1 \to F_2$ of a rule of Γ , $p(\mathbf{d}^*) \in Sp_I(F_2)$ and $q(\mathbf{e}^*) \in Pnn_I(F_1)$.

If Γ is a singleton $\{F\}$, then we write $\mathfrak{G}_{I,\mathcal{A}}(F)$ instead of $\mathfrak{G}_{I,\mathcal{A}}(\{F\})$. We say that a partition $\{\mathcal{A}_1, \mathcal{A}_2\}$ of a set of atoms $\mathcal{A} \subseteq At(I)$ is *separable* on $\mathfrak{G}_{I,\mathcal{A}}(\Gamma)$ if every infinite walk v_1, v_2, \ldots of $\mathfrak{G}_{I,\mathcal{A}}(\Gamma)$ visits almost one \mathcal{A}_i infinitely many times, that is, there is $i \in \{1, 2\}$ such that $\{k \mid v_k \in \mathcal{A}_i\}$ is finite.

A formula F is said to be *negative* on an interpretation Iand a subset A of At(I) if $Sp_I(F)$ and A are disjoint. A set of sentences Γ is said to be *negative* on an interpretation Iand a subset A if it is negative on all its sentences.

Proof of the Splitting Lemma

Lemma 1 Let Γ be a first-order theory, I be an interpretation and A be a subset of At(I). Let $\{A_1, A_2\}$ be a partition of A. If I is a A-stable model of Γ , then I is a A_i -stable model of Γ for all $1 \le i \le 2$. *Proof.* Assume that *I* is a \mathcal{A} -stable model of Γ and pick any strict subset \mathcal{H} of \mathcal{A}_i . Since $\mathcal{A}_i \subseteq \mathcal{A}$, it follows that \mathcal{H} is also a strict subset of \mathcal{A} . Furthermore, since *I* is a \mathcal{A} -stable model of Γ , it follows that $\langle \mathcal{H}, I \rangle \not\models_{ht} \Gamma \cup EM(I, \mathcal{A})$. Furthermore, $\mathcal{A}_i \subseteq \mathcal{A}$ implies $EM(I, \mathcal{A}_i) \supseteq EM(I, \mathcal{A})$ and, thus, this implies $\langle \mathcal{H}, I \rangle \not\models_{ht} \Gamma \cup EM(I, \mathcal{A}_i)$. Consequently, *I* is a \mathcal{A}_i -stable model of Γ . \Box

The following result by Fandinno and Lifschitz (2022) will be useful in the next Lemma.

Proposition 4

=

- (a) If $\langle \mathcal{H}, I \rangle \models_{\overline{ht}} F$ then $I \models F$.
- (b) For any sentence F that does not contain intensional symbols, $\langle \mathcal{H}, I \rangle \models_{ht} F$ iff $I \models F$.
- (c) For any subset S of \mathcal{H} such that the predicate symbols of its members do not occur in F, $\langle \mathcal{H} \setminus S, I \rangle \models_{ht} F$ iff $\langle \mathcal{H}, I \rangle \models_{ht} F$.

Lemma 2 For any HT-interpretation $\langle \mathcal{H}, I \rangle$ and any sentence F over σ^I , if $I \models F$ and $Sp_I(F) \subseteq \mathcal{H}$ then $\langle \mathcal{H}, I \rangle \models_{ht} F$.

Proof. By induction on the size of F. *Case 1:* F is $p(\mathbf{t})$. Then the assumption $Sp_I(F) \subseteq \mathcal{H}$ and the claim $\langle \mathcal{H}, I \rangle \models_{\mathrm{ht}} F$ turn into the condition $p((\mathbf{t}^I)^*) \in \mathcal{H}$. *Case 2:* F is $F_1 \wedge F_2$. Then from the assumption $I \models F$ we conclude that $I \models F_i$ for i = 1, 2. On the other hand,

$$Sp_I(F_i) \subseteq Sp_I(F) \subseteq \mathcal{H}.$$

By the induction hypothesis, it follows that $\langle \mathcal{H}, I \rangle \models_{\overline{ht}} F_i$, and consequently $\langle \mathcal{H}, I \rangle \models_{\overline{ht}} F$. *Case 3*: F is $F_1 \lor F_2$. Similar to Case 2. *Case 4*: F is $F_1 \to F_2$. Since $I \models F_1 \to F_2$, we only need to check that $\langle \mathcal{H}, I \rangle \models_{\overline{ht}} F_1$ or $\langle \mathcal{H}, I \rangle \models_{\overline{ht}} F_2$. *Case 4.1*: $I \models F_1$. Since $I \models F_1 \to F_2$, it follows that $I \models F_2$. On the other hand,

$$Sp_I(F_2) = Sp_I(F) \subseteq \mathcal{H}.$$

By the induction hypothesis, it follows that $\langle \mathcal{H}, I \rangle \models_{\overline{ht}} F_2$. *Case 4.2:* $I \not\models F_1$. By Proposition 4(a), it follows that $\langle \mathcal{H}, I \rangle \models_{\overline{ht}} F_1$. *Case 5:* F is $\forall X G(X)$, where X is a variable of sort s. Then, for every element d of $|I|^s$, $I \models G(d^*)$ and

$$Sp_I(G(d^*)) \subseteq Sp_I(F) \subseteq \mathcal{H}.$$

By the induction hypothesis, it follows that $\langle \mathcal{H}, I \rangle \models_{\overline{ht}} G(d^*)$. Consequently $\langle \mathcal{H}, I \rangle \models_{\overline{ht}} \forall X G(X)$. *Case 6:* F is $\exists X G(X)$. Similar to Case 5.

Lemma 3 If \mathcal{A} is disjoint from $Sp_I(F)$ and $I \models F$, then $\langle At(I) \setminus \mathcal{A}, I \rangle \models_{\overline{ht}} F$.

Proof. Since \mathcal{A} is disjoint from $Sp_I(F)$ and $Sp_I(F) \subseteq At(I)$, it follows that $Sp_I(F) \subseteq At(I) \setminus \mathcal{A}$. Since $I \models F$, the result follows from Lemma 2.

Lemma 4 Let I be an interpretation, \mathcal{B}_1 and \mathcal{B}_2 be subsets of At(I) and F be a sentence.

- 1. If \mathcal{B}_2 is disjoint from $Pnn_I(F)$ and $\langle \mathcal{H}_1, I \rangle \models_{\overline{ht}} F$, then $\langle \mathcal{H}_2, I \rangle \models_{\overline{ht}} F$,
- 2. If \mathcal{B}_2 is disjoint from $Nnn_I(F)$ and $\langle \mathcal{H}_2, I \rangle \models_{\overline{ht}} F$, then $\langle \mathcal{H}_1, I \rangle \models_{\overline{ht}} F$,

where $\mathcal{H}_1 = At(I) \setminus \mathcal{B}_1$ and $\mathcal{H}_2 = At(I) \setminus (\mathcal{B}_1 \cup \mathcal{B}_2)$.

Proof. The proof is by structural induction on *F*.

In case that F is an atom $p(\mathbf{t})$. Since $\langle \mathcal{H}_i, I \rangle \models_{\text{It}} F$, it follows that $p(\mathbf{t}^I)$ belongs to \mathcal{H}_i .

- 1. Then, $p(\mathbf{t}^{I})$ does not belong to \mathcal{B}_{1} . Since \mathcal{B}_{2} is disjoint from $Pnn_{I}(F)$, this means that $\mathcal{B}_{1} \cup \mathcal{B}_{2}$ is disjoint from $Pnn_{I}(F) = Sp_{I}(F)$, and, thus, condition 1 follows directly from Lemma 2.
- 2. Since $p(\mathbf{t}^I)$ belongs to $\mathcal{H}_2 \subseteq \mathcal{H}_1$, it follows that $\langle \mathcal{H}_1, I \rangle \models_{\overline{ht}} p(\mathbf{t})$. Therefore, condition 2 is also satisfied.

In case that F is an implication of form $F_1 \rightarrow F_2$.

- 1. Assume that \mathcal{B}_2 is disjoint from $Pnn_I(F)$ and $\langle \mathcal{H}_1, I \rangle \models_{\overline{ht}} F$. The latter implies $I \models F_1$ and, thus, $Pnn_I(F) = Nnn_I(F_1) \cup Pnn_I(F_2)$. Then,
 - \mathcal{B}_2 is disjoint from $Nnn_I(F_1)$, and
 - \mathcal{B}_2 is disjoint from $Pnn_I(F_2)$.

Furthermore, $\langle \mathcal{H}_1, I \rangle \models_{\overline{ht}} F$ implies that $\langle \mathcal{H}_1, I \rangle \not\models_{\overline{ht}} F_1$ or $\langle \mathcal{H}_1, I \rangle \models_{\overline{ht}} F_2$. By induction hypothesis, this implies that $\langle \mathcal{H}_2, I \rangle \not\models_{\overline{ht}} F_1$ or $\langle \mathcal{H}_2, I \rangle \models_{\overline{ht}} F_2$. In addition, $\langle \mathcal{H}_1, I \rangle \models_{\overline{ht}} F$ also implies $I \models F_1 \to F_2$. Therefore, $\langle \mathcal{H}_2, I \rangle \models_{\overline{ht}} F$.

- 2. Assume that \mathcal{B}_2 is disjoint from $Nnn_I(F)$ and $\langle \mathcal{H}_1, I \rangle \models_{\overline{ht}} F$. The latter implies $I \models F_1$ and, thus, $Pnn_I(F) = Pnn_I(F_1) \cup Nnn_I(F_2)$. Then,
 - \mathcal{B}_2 is disjoint from $Pnn_I(F_1)$, and
 - \mathcal{B}_2 is disjoint from $Nnn_I(F_2)$.

Furthermore, $\langle \mathcal{H}_2, I \rangle \models_{\overline{ht}} F$ implies $\langle \mathcal{H}_2, I \rangle \models_{\overline{ht}} F_1$ or $\langle \mathcal{H}_2, I \rangle \models_{\overline{ht}} F_2$. By induction hypothesis, this implies that $\langle \mathcal{H}_1, I \rangle \models_{\overline{ht}} F_1$ or $\langle \mathcal{H}_1, I \rangle \models_{\overline{ht}} F_2$. In addition, $\langle \mathcal{H}_2, I \rangle \models_{\overline{ht}} F$ also implies $I \models F_1 \to F_2$. Therefore, $\langle \mathcal{H}_1, I \rangle \models_{\overline{ht}} F$.

Finally, the cases in which F is of forms $F_1 \wedge F_2$, $F_1 \vee F_2$, $\forall X F_1(X)$ and $\exists X F_1(X)$ follow directly by induction hypothesis.

Lemma 5 Let I be an interpretation, $\{\mathcal{B}_1, \mathcal{B}_2\}$ be partition of some set of ground atoms \mathcal{X} and F be a sentence such that there are no edges from \mathcal{B}_1 to \mathcal{B}_2 in $\mathfrak{G}_{I,\mathcal{X}}(F)$. Let $\mathcal{H}_1 =$ $At(I) \setminus \mathcal{B}_1$ and $\mathcal{H}_2 = At(I) \setminus (\mathcal{B}_1 \cup \mathcal{B}_2)$. If $\langle \mathcal{H}_2, I \rangle \models_{\text{Int}} F$, then $\langle \mathcal{H}_1, I \rangle \models_{\text{Int}} F$.

Proof. The proof is by structural induction on F. In case that F is an atom $p(\mathbf{t})$, the result follows because $\mathcal{H}_2 \subseteq \mathcal{H}_1$.

In case that F is of form $F_1 \to F_2$. Then, from $\langle \mathcal{H}_2, I \rangle \models_{\overline{ht}} F_1 \to F_2$ we can conclude that $I \models F_1 \to F_2$. Assume that $\langle \mathcal{H}_1, I \rangle \models_{\overline{ht}} F_1$. Then, $I \models F_1$ and $I \models F_2$ follows. We just need to show that $\langle \mathcal{H}_1, I \rangle \models_{\overline{ht}} F_2$. We proceed by cases.

- If B₁ is disjoint from Sp_I(F₂) then, by Lemma 3 and the fact I ⊨ F₂, we get ⟨H₁, I⟩ ⊨ F₂.
- Otherwise, since \mathcal{B}_1 is not disjoint from $Sp_I(F_2)$, it must be that \mathcal{B}_2 is disjoint from $Pnn_I(F_1)$, because there are no edges from \mathcal{B}_1 to \mathcal{B}_2 in $\mathfrak{G}_{I,\mathcal{X}}(F)$. Then, by Lemma 4 and $\langle \mathcal{H}_1, I \rangle \models_{\overline{ht}} F_1$, we get $\langle \mathcal{H}_2, I \rangle \models_{\overline{ht}} F_1$. Since $\langle \mathcal{H}_2, I \rangle \models_{\overline{ht}} F_1 \to F_2$, this implies that $\langle \mathcal{H}_2, I \rangle \models_{\overline{ht}}$

 F_2 . Then, by induction hypothesis, we get $\langle \mathcal{H}_1, I \rangle \models_{ht} F_2$. Note that $\mathfrak{G}_{I,\mathcal{X}}(F_2)$ is a subgraph of $\mathfrak{G}_{I,\mathcal{X}}(F)$.

Finally, the cases in which F is of forms $F_1 \wedge F_2$, $F_1 \vee F_2$, $\forall X F(X)$ and $\exists X F(X)$ follow directly by induction hypothesis.

Lemma 6 Let Γ be a first-order theory, I be an interpretation and A be a subset of At(I). Let $\{A_1, A_2\}$ be a partition of A satisfying the following condition:

for every non-empty subset \mathcal{X} of \mathcal{A} , there is some $1 \leq i \leq 2$ and non-empty subset \mathcal{B} of $\mathcal{X} \cap \mathcal{A}_i$ such that there are no edges from \mathcal{B} to $\mathcal{X} \setminus \mathcal{B}$ in $\mathfrak{G}_{I,\mathcal{X}}(\Gamma)$.

If I is a A_i -stable model of Γ for all $1 \leq i \leq 2$, then I is a A-stable model of Γ .

Proof. Assume that I is a \mathcal{A}_i -stable model of Γ for all $1 \leq i \leq 2$ and pick any strict subset \mathcal{H} of At(I). It is enough to show that $\langle \mathcal{H}, I \rangle \not\models_{\mathrm{ht}} \Gamma \cup EM(I, \mathcal{A})$. Let $\mathcal{X} = At(I) \setminus \mathcal{H}$. If \mathcal{X} is not a subset of \mathcal{A} , then $\langle \mathcal{H}, I \rangle \not\models_{\mathrm{ht}} EM(I, \mathcal{A})$. Hence, we assume without loss of generality that $\mathcal{X} \subseteq \mathcal{A}$ and we show that $\langle \mathcal{H}, I \rangle \not\models_{\mathrm{ht}} \Gamma$.

By hypothesis, there is some $1 \leq i \leq 2$ and non-empty subset \mathcal{B} of $\mathcal{X} \cap \mathcal{A}_i$ such that there are no edges from \mathcal{B} to $\mathcal{X} \setminus \mathcal{B}$ in $\mathfrak{G}_{I,\mathcal{X}}(\Gamma)$. Let $\mathcal{H}_i = At(I) \setminus \mathcal{B}$ and $\mathcal{C} = \mathcal{X} \setminus \mathcal{B}$. Since $\mathcal{B} \subseteq \mathcal{X}$, we get that $\mathcal{X} = \mathcal{B} \cup \mathcal{C}$. We also get that

there are no edges from
$$\mathcal{B}$$
 to \mathcal{C} in $\mathfrak{G}_{I,\mathcal{X}}(\Gamma)$. (24)

Furthermore, since \mathcal{B} is non-empty, it follows that $\mathcal{H}_i \subset At(I)$ and, since I is a \mathcal{A}_i -stable model of Γ , we immediately get that $\langle \mathcal{H}_i, I \rangle \not\models_{\operatorname{ht}} \Gamma \cup EM(I, \mathcal{A}_i)$. Furthermore, since $\mathcal{B} \subseteq \mathcal{X} \cap \mathcal{A}_i \subseteq \mathcal{A}_i$, it follows that $\langle \mathcal{H}_i, I \rangle \models_{\operatorname{ht}} EM(I, \mathcal{A}_i)$. Therefore,

$$\langle \mathcal{H}_i, I \rangle \not\models_{\mathrm{ft}} \Gamma.$$
 (25)

By Lemma 5, facts (24) and (25) imply that

$$\langle At(I) \setminus (\mathcal{B} \cup \mathcal{C}), I \rangle \not\models_{ht} \Gamma.$$
 (26)

Since $\mathcal{H} \subseteq At(I)$ and $\mathcal{X} = At(I) \setminus \mathcal{H}$, we get that $\mathcal{H} = At(I) \setminus \mathcal{X} = At(I) \setminus (\mathcal{B} \cup \mathcal{C})$. Therefore, we can rewrite (26) as $\langle \mathcal{H}, I \rangle \not\models_{ht} \Gamma$.

Lemma 7 Let $\{X_1, X_2\}$ be a separable partition of some non-empty \mathcal{X} . Then, there is a vertex $v \in \mathcal{X}$ such that the set of vertices reachable $\mathfrak{G}_{I,\mathcal{X}}(\Gamma)$ from v is a subset of \mathcal{X}_i for some $1 \leq i \leq 2$.

Proof. Suppose, for the sake of contradiction, that for every vertex $v \in \mathcal{X}$, the set of vertices reachable from vertex v in $\mathfrak{G}_{I,\mathcal{X}}(\Gamma)$ is neither a subset of \mathcal{X}_1 nor \mathcal{X}_2 . That is, for every $v_{i,1} \in \mathcal{X}_1$, there is $v_{i,k_i} \in \mathcal{X}_2$ such that $v_{i,1}, \ldots, v_{i,k_i}$ is a walk in $\mathfrak{G}_{I,\mathcal{X}}(\Gamma)$; and for every $w_{j,1} \in \mathcal{X}_1$, there is $w_{j,l_j} \in \mathcal{X}_2$ such that $w_{j,1}, w_{j,2}, \ldots, w_{j,l_j}$ is a walk in $\mathfrak{G}_{I,\mathcal{X}}(\Gamma)$. By taking vertices that satisfy $v_{i,k_i} = w_{i,1}$ and $w_{j,l_j} = v_{i+1,1}$, we can build the infinite walk

$$v_{1,1}, v_{1,2}, \ldots, v_{1,k_1}, w_{1,2}, \ldots, w_{1,l_1}, v_{2,2}, \ldots$$

that visits both \mathcal{X}_1 and \mathcal{X}_2 infinitely many times. This is a contradiction with the fact that $\{\mathcal{X}_1, \mathcal{X}_2\}$ is separable. Hence, there is a vertex $v \in \mathcal{X}$ such that the set of vertices reachable $\mathfrak{G}_{I,\mathcal{X}}(\Gamma)$ from v is a subset of \mathcal{X}_i for some $1 \leq i \leq 2$. **Lemma 8** Let $\{A_1, A_2\}$ be a separable partition of A. Then,

for every non-empty subset \mathcal{X} of \mathcal{A} , there is some $1 \leq i \leq 2$ and non-empty subset \mathcal{B} of $\mathcal{X} \cap \mathcal{A}_i$ such that there are no edges from \mathcal{B} to $\mathcal{X} \setminus \mathcal{B}$ in $\mathfrak{G}_{I,\mathcal{X}}(\Gamma)$.

Proof. Let $\mathcal{X}_i = \mathcal{X} \cap \mathcal{A}_i$. Then, $\{\mathcal{X}_1, \mathcal{X}_2\}$ is a separable partition of \mathcal{X} and, by Lemma 7, there is a vertex $v \in \mathcal{X}$ such that the set of vertices \mathcal{B} reachable $\mathfrak{G}_{I,\mathcal{X}}(\Gamma)$ from v is a subset of \mathcal{X}_i for some i. Clearly, there are no edges from \mathcal{B} to $\mathcal{X} \setminus \mathcal{B}$ in $\mathfrak{G}_{I,\mathcal{X}}(\Gamma)$.

Lemma 9 (Splitting lemma with infinitary graph) Let Γ be a first-order theory, I be an interpretation and A be a subset of At(I). If $\{A_1, A_2\}$ is a partition of A that is separable on $\mathfrak{G}_{I,\mathcal{A}}(F)$, then the following two statements are equivalent

1. I is a A-stable model of Γ , and

2. *I* is a A_i -stable model of Γ for all $1 \le i \le 2$.

Proof. Condition 1 implies condition 2 by Lemma 1. Furthermore, by Lemma 8, if follows that

for every non-empty subset \mathcal{X} of \mathcal{A} , there is some $1 \leq i \leq 2$ and non-empty subset \mathcal{B} of $\mathcal{X} \cap \mathcal{A}_i$ such that there are no edges from \mathcal{B} to $\mathcal{X} \setminus \mathcal{B}$ in $\mathfrak{G}_{I,\mathcal{X}}(\Gamma)$.

Then, by Lemma 6, it follows that Condition 2 implies condition 1. $\hfill \Box$

Lemma 10 Let Γ_1, Γ_2 be two first-order theories. If \mathcal{A} is a set of ground atoms that is disjoint from $Sp_I(\Gamma_2)$, then I is an \mathcal{A} -stable model of $\Gamma_1 \cup \Gamma_2$ iff I is an \mathcal{A} -stable model of Γ_1 and $I \models \Gamma_2$.

Proof. For the if direction. Assume that I is an A-stable model of Γ_1 and $I \models \Gamma_2$. We get that $I \models \Gamma_1 \cup \Gamma_2$. Furthermore since I is an A-stable model of Γ_1 , for any $\mathcal{H} \subset At(I)$, we get that $\langle \mathcal{H}, I \rangle \not\models \Gamma_1$ and, thus, $\langle \mathcal{H}, I \rangle \not\models \Gamma_1 \cup \Gamma_2$. Therefore, A-stable model of $\Gamma_1 \cup \Gamma_2$.

For the only if direction. Assume that I is a A-stable model of $\Gamma_1 \cup \Gamma_2$. Then $I \models \Gamma_1$ and $I \models \Gamma_2$. Furthermore, since A is disjoint from $Sp_I(\Gamma_2)$, by Lemma 2 it follows that $\langle \mathcal{H}, I \rangle \not\models \Gamma_1$ for any $\mathcal{H} \subset At(I)$. Hence, I is an A-stable model of Γ_1 .

Theorem 3 (Splitting Theorem with infinitary graph)

Let $\Gamma = \Gamma_1 \cup \Gamma_2$ be a set of sentences, I be an interpretation, and A be a subset of At(I), and $\{A_1, A_2\}$ be a partition of A such that

- $\{A_1, A_2\}$ is separable on $\mathfrak{G}_{I,\mathcal{A}}(\Gamma)$;
- Γ_1 is negative on A_2 ; and
- Γ_2 is negative on \mathcal{A}_1 .

Then the following two statements are equivalent

- *1. I* is a A-stable model of Γ , and
- 2. *I* is a A_i -stable model of Γ_i for all $i \in \{1, 2\}$.

Proof. By Lemma 9 and the fact that $\Gamma = \Gamma_1 \cup \Gamma_2$, it follows that condition 1 below is equivalent to the conjunction of conditions a and b:

1. *I* is a A-stable model of Γ ;

a. *I* is a A₁-stable model of Γ₁ ∪ Γ₂;
b. *I* is a A₂-stable model of Γ₁ ∪ Γ₂.

Furthermore, since Γ_1 is negative on \mathcal{A}_2 , we get that $Sp_{\Gamma_1}(I)$ and \mathcal{A}_2 are disjoint. Similarly, we get that $Sp_{\Gamma_2}(I)$ and \mathcal{A}_1 are also disjoint. By Lemma 10, this implies condition a and b are respectively equivalent to:

a'. *I* is a A_1 -stable model of Γ_1 and $I \models \Gamma_2$, and b'. *I* is a A_2 -stable model of Γ_2 and $I \models \Gamma_1$.

It remains to see that if I is a stable model of Γ , then it satisfies Γ and, thus, the conjunction of conditions a' and b' is equivalent to

2. *I* is a \mathcal{A}_i -stable model of Γ_i for all $i \in \{1, 2\}$.

Proof of the Splitting Theorem with Intensionality Statements

Lemma 11 If $\Lambda = \{\lambda_1, \lambda_2\}$ is a partition of λ , then $\{At(I, \lambda_1), At(I, \lambda_2)\}$ is a partition of $At(I, \lambda)$.

Proof. Since $\Lambda = \{\lambda_1, \lambda_2\}$ is a partition of λ , we have

$$\lambda^{p}(\mathbf{d}^{*}) \leftrightarrow \lambda_{1}^{p}(\mathbf{d}^{*}) \lor \lambda_{2}^{p}(\mathbf{d}^{*})$$
(27)

$$\neg(\lambda_1^p(\mathbf{d}^*) \land \lambda_2^p(\mathbf{d}^*)) \tag{28}$$

are satisfied by every interpretation I, every predicate symbol p and list **d** of domain elements of the appropriate length and sorts. Then,

$$\begin{aligned} At(I,\lambda) &= \{ p(\mathbf{d}^*) \mid I \models p(\mathbf{d}^*) \land \lambda^p(\mathbf{d}^*) \} \\ &= \{ p(\mathbf{d}^*) \mid I \models p(\mathbf{d}^*) \land (\lambda_1^p(\mathbf{d}^*) \lor \lambda_2^p(\mathbf{d}^*)) \} \\ &= \{ p(\mathbf{d}^*) \mid I \models p(\mathbf{d}^*) \land \lambda_1^p(\mathbf{d}^*) \} \\ &\cup \{ p(\mathbf{d}^*) \mid I \models p(\mathbf{d}^*) \land \lambda_2^p(\mathbf{d}^*)) \} \\ &= At(I,\lambda_1) \cup At(I,\lambda_2) \end{aligned}$$

Finally, $At(I, \lambda_1) \cap At(I, \lambda_2) = \emptyset$ follows from (28).

Lemma 12 Let Γ be a set of basic rules. If $\Lambda = \{\lambda_1, \lambda_2\}$ is a partition of λ and $p(\mathbf{d}^*)$ is a vertex in $\mathfrak{G}_{I,At(I,\lambda)}(\Gamma)$ such that $p(\mathbf{d}^*)$ belongs to $At(I,\lambda_i)$ with $i \in \{1,2\}$, then (p,λ_i) is a vertex in $\mathfrak{G}_{\Lambda}(\Gamma)$ and $I \models \lambda_i^p(\mathbf{d}^*) \land \neg \lambda_i^p(\mathbf{d}^*)$ with $j \neq i$.

Proof. Since Λ is a partition of λ , by Lemma 11, $At(I, \lambda_1)$ and $At(I, \lambda_2)$ form a partition of $At(I, \lambda)$. Furthermore, since $p(\mathbf{d}^*)$ is a vertex in $\mathfrak{G}_{I,At(I,\lambda)}(\Gamma)$, by definition, we get that $p(\mathbf{d}^*)$ belongs to $At(I, \lambda)$. Hence, $p(\mathbf{d}^*)$ belongs to $At(I, \lambda_i)$ for exactly one $i \in \{1, 2\}$ and, thus, $I \models p(\mathbf{d}^*) \wedge \lambda_i(\mathbf{d}^*)$. We also get that $I \not\models p(\mathbf{d}^*) \wedge \lambda_j(\mathbf{d}^*)$ with $j = \{1, 2\}$ such that $j \neq i$. Therefore, $I \models \lambda_i(\mathbf{d}^*) \wedge \neg \lambda_j(\mathbf{d}^*)$. Finally, since $I \models p(\mathbf{d}^*) \wedge \lambda_i(\mathbf{d}^*)$, we get $\exists \mathbf{X} \lambda_i^p \not\equiv \bot$. This implies that (p, λ_i) is a vertex of $\mathfrak{G}_{\Lambda}(\Gamma)$.

Given a vertex $p(\mathbf{d}^*)$ in $\mathfrak{G}_{I,At(I,\lambda)}(\Gamma)$, by $[p(\mathbf{d}^*)]$ we denote the vertex (p, λ_i) in $\mathfrak{G}_{\Lambda}(\Gamma)$ such that $I \models \lambda_i^p(\mathbf{d}^*)$. Note that, by Lemma 12, such vertex exists and it is unique.

Lemma 13 Let Γ be a set of rules. If $\Lambda = \{\lambda_1, \lambda_2\}$ is a partition of λ and (v, w) is an edge in $\mathfrak{G}_{I,At(I,\lambda)}(\Gamma)$, then ([v], [w]) is an edge in $\mathfrak{G}_{\Lambda}(\Gamma)$. *Proof.* Since Γ is a set of rules and (v, w) is an edge in $\mathfrak{G}_{I,At(I,\lambda)}(\Gamma)$, there is an instance $F_1 \to F_2$ of a rule such that $v \in Sp_I(F_2)$ and $w \in Sp_I(F_1)$. Let $v = p(\mathbf{d}^*)$ and $w = q(\mathbf{e}^*)$. Then, $v \in Sp_I(F_2)$ and $w \in Sp_I(F_1)$ imply that

$$I \models F_1 \wedge F_2 \wedge p(\mathbf{d}^*) \wedge q(\mathbf{e}^*) \tag{29}$$

Let $[w] = (p, \lambda_v)$ and $[v] = (q, \lambda_w)$. From Lemma 12, we get $I \models \lambda_v^p(\mathbf{d}^*) \land \lambda_w^q(\mathbf{e}^*)$. This plus (29) imply

$$I \models F_1 \land p(\mathbf{d}^*) \land \lambda_v^p(\mathbf{d}^*) \land \lambda_w^q(\mathbf{e}^*)$$
(30)

Let $B \to H$ be the rule of which $F_1 \to F_2$ is an instance, **X** be the free variables in $B \to H$ and **x** be names of domain elements such that $(B \to H)_{\mathbf{x}}^{\mathbf{X}} = F_1 \to F_2$. Then, there is a positive occurrence of p in H within a literal $p(\mathbf{t})$ and a positive occurrence of q in B within a literal $q(\mathbf{r})$ such that $\mathbf{d} = (\mathbf{t}_{\mathbf{x}}^{\mathbf{X}})^I$ and $\mathbf{e} = (\mathbf{r}_{\mathbf{x}}^{\mathbf{X}})^I$. Therefore (30) implies

$$I \models B_{\mathbf{x}}^{\mathbf{X}} \land p(\mathbf{t}_{\mathbf{x}}^{\mathbf{X}}) \land \lambda_{v}^{p}(\mathbf{t}_{\mathbf{x}}^{\mathbf{X}}) \land \lambda_{w}^{q}(\mathbf{r}_{\mathbf{x}}^{\mathbf{X}})$$

and, this implies that

$$\exists \mathbf{X} (B \wedge p(\mathbf{t}) \wedge \lambda_v^p(\mathbf{t}) \wedge \lambda_w^q(\mathbf{r}))$$

is satisfiable. Consequently, ([v], [w]) is an edge in $\mathfrak{G}_{\Lambda}(\Gamma)$.

Lemma 14 Let F be a sentence. Let Ψ be a theory and I be an model of Ψ . Then, the following conditions hold:

- 1. If $p(\mathbf{d}^*)$ belongs to $Sp_I(F)$, then there is some a strictly positive occurrence $p(\mathbf{t})$ of predicate symbol p in F such that I satisfies $Sp_{\Psi}(F)_{\mathbf{d}^*}^{\mathbf{Y}}$ where \mathbf{Y} are the free variables in $Sp_{\Psi}(F)$.
- 2. If $p(\mathbf{d}^*)$ belongs to $Pnn_I(F)$, then there is some a positive nonnegated occurrence $p(\mathbf{t})$ of predicate symbol pin F such that I satisfies $Pnn_{\Psi}(F)_{\mathbf{d}^*}^{\mathbf{Y}}$ where \mathbf{Y} are the free variables in $Pnn_{\Psi}(F)$.
- If p(d*) belongs to Nnn_I(F), then there is some a negative nonnegated occurrence p(t) of predicate symbol p in F such that I satisfies Nnn_Ψ(F)^Y_{d*} where Y are the free variables in Nnn_Ψ(F).

Proof. The proof follows by induction on the size of the formula. Note that the fact that $p(\mathbf{d}^*)$ belongs to $Sp_I(F)$, $Pnn_I(F)$ or $Nnn_I(F)$ implies that $I \models F$. Since I is a model of Ψ , this implies that $\Psi \cup \{\exists \mathbf{X} F\}$ is satisfiable and, thus, we are not in the case where $Sp_{\Psi}(F)$ (resp. $Pnn_I(F)$ or $Nnn_I(F)$) is \bot by definition.

Case 1. F is an atom, then F is of the form $p(\mathbf{t})$ with $I \models p(\mathbf{t})$ and $\mathbf{d} = \mathbf{t}^I$. Hence, there is a strictly positive occurrence $p(\mathbf{t})$ of predicate p in F. Any strictly positive occurrence is also a positive nonnegated occurrence. Furthermore $Sp_{\Psi}(F)_{\mathbf{d}^*}^{\mathbf{Y}} = Pnn_{\Psi}(F)_{\mathbf{d}^*}^{\mathbf{Y}}$ is $p(\mathbf{t}) \wedge \mathbf{d}^* = \mathbf{t}$, which is satisfied by I. Note that if F is an atom $Nnn_I(F)$ is the empty set by definition. Therefore, the third condition holds vacuous.

Case 2. F is of the form $F_1 \wedge F_2$. Then, $p(\mathbf{d}^*)$ belongs to $Sp_I(F_i)$ for some $1 \leq i \leq 2$. By induction hypothesis, there is some a positive occurrence $p(\mathbf{t})$ of predicate symbol p in F_i such that I satisfies $Sp_{\Psi}(F_i)_{\mathbf{d}^*}^{\mathbf{Y}}$. Remains observing that $Sp_{\Psi}(F) \equiv Sp_{\Psi}(F_i) \wedge F_j$ with $1 \leq j \leq 2$ and $i \neq j$ and that $I \models F_j$ because $I \models F$. The same reasoning applies to $Pnn_I(F)$ and $Nnn_I(F)$.

Case 3. F is of the form $F_1 \vee F_2$. Then, $p(\mathbf{d}^*)$ belongs to $Sp_I(F_i)$ for some $1 \leq i \leq 2$. By induction hypothesis, this implies that there is some a positive occurrence $p(\mathbf{t})$ of predicate symbol p in F_i such that I satisfies $Sp_{\Psi}(F_i)_{\mathbf{d}^*}^{\mathbf{Y}}$. Remains observing that $Sp_{\Psi}(F) = Sp_{\Psi}(F_i)$. The same reasoning applies to $Pnn_I(F)$ and $Nnn_I(F)$.

Case 4. F is of the form $F_1 \rightarrow F_2$. Note that $p(\mathbf{d}^*)$ belonging to $Sp_I(F)$ (resp. $Pnn_I(F)$ or $Nnn_I(F)$) implies that (i) $I \models F_1$ and that (ii)

- 1. $p(\mathbf{d}^*)$ belongs to $Sp_I(F_2)$,
- 2. $p(\mathbf{d}^*)$ belongs to $Nnn_I(F_1) \cup Pnn_I(F_2)$, and
- 3. $p(\mathbf{d}^*)$ belongs to $Pnn_I(F_1) \cup Nnn_I(F_2)$.

Since $I \models F$, (i) implies that $I \models F_1 \land F_2$ and, since I is a model of Ψ , this implies $F_1^{\Psi} = F_1$.

1. By induction hypothesis, (ii) implies that there is some a positive occurrence $p(\mathbf{t})$ of predicate symbol p in F_2 such that I satisfies $Sp_{\Psi}(F_2)_{\mathbf{d}^*}^{\mathbf{Y}}$. Hence, I satisfies

$$Sp_{\Psi}(F)_{\mathbf{d}^*}^{\mathbf{Y}} = F_1^{\Psi} \wedge Sp_{\Psi}(F_2)_{\mathbf{d}^*}^{\mathbf{Y}} = F_1 \wedge Sp_{\Psi}(F_2)_{\mathbf{d}^*}^{\mathbf{Y}}$$

2. By induction hypothesis, (ii) implies that there is some negative nonnegated occurrence of $p(\mathbf{t})$ of predicate symbol p in F_1 such that I satisfies $Nnn_{\Psi}(F_1)_{\mathbf{d}^*}^{\mathbf{Y}}$ or some positive nonnegated occurrence of $p(\mathbf{t})$ of predicate symbol p in F_2 such that I satisfies $Pnn_{\Psi}(F_2)_{\mathbf{d}^*}^{\mathbf{Y}}$. In both cases there is a positive nonnegated occurrence of $p(\mathbf{t})$ of predicate symbol p in F. Furthermore, if such occurrence happens in F_1 , then

$$Pnn_{\Psi}(F)_{\mathbf{d}^*}^{\mathbf{Y}} = Nnn_{\Psi}(F_1)_{\mathbf{d}^*}^{\mathbf{Y}}$$

and, thus, condition 2 holds. Otherwise,

$$Pnn_{\Psi}(F)_{\mathbf{d}^{*}}^{\mathbf{Y}} = F_{1}^{\Psi} \wedge Pnn_{\Psi}(F_{2})_{\mathbf{d}^{*}}^{\mathbf{Y}} = F_{1} \wedge Pnn_{\Psi}(F_{2})_{\mathbf{d}^{*}}^{\mathbf{Y}}$$

and, thus, condition 2 also holds.

3. This case is analogous to the previous case.

Case 5. F is of the form $\forall XG(X)$. Then, $p(\mathbf{d}^*)$ belongs to $Sp_I(G(d^*))$ for some domain element of the appropriate sort. By induction hypothesis, there is some positive occurrence $p(\mathbf{r})$ of predicate symbol p in $G_{d^*}^X$ such that I satisfies $Sp_{\Psi}(G_{d^*}^X)_{\mathbf{d}^*}^{\mathbf{Y}}$ for every domain element d of the appropriate sort. Then, there is an occurrence $p(\mathbf{t})$ the occurrence on G corresponding to $p(\mathbf{r})$ in $G_{d^*}^X$ such that $Sp_{\Psi}(G)_{d^*\mathbf{d}^*}^{X\mathbf{Y}}$ for every domain element d of the appropriate sort and, thus, I satisfies $\forall X Sp_{\Psi}(G)_{\mathbf{d}^*}^{\mathbf{Y}}$. Case 6. F is of the form $\exists XG(X)$. Then, $p(\mathbf{d}^*)$ belongs to $Sp_I(G(d^*))$ for some domain element of the appropriate sort. By induction hypothesis, there is some positive occurrence $p(\mathbf{r})$ of predicate symbol p in $G_{d^*}^X$ such that I satisfies $Sp_{\Psi}(G_{d^*}^X)_{\mathbf{d}^*}^{\mathbf{Y}}$ for some domain element d of the appropriate sort. Then, there is an occurrence $p(\mathbf{t})$ the occurrence on G corresponding to $p(\mathbf{r})$ in $G_{d^*}^X$ such that $Sp_{\Psi}(G)_{d^*\mathbf{d}^*}^{X\mathbf{Y}}$ for some domain element d of the appropriate sort and, thus, I satisfies $\exists X Sp_{\Psi}(G)_{\mathbf{d}^*}^{\mathbf{Y}}$. The same reasoning applies to $Pnn_I(F)$ and $Nnn_I(F)$. \square

Lemma 15 Let Γ and Ψ be two set theories and I be a model of Ψ . If $\Lambda = \{\lambda_1, \lambda_2\}$ is a partition of λ and $p(\mathbf{d}^*)$ is a vertex in $\mathfrak{G}_{I,At(I,\lambda)}(\Gamma)$ such that $p(\mathbf{d}^*)$ belongs to $At(I,\lambda_i)$ with $i \in \{1,2\}$, then (p,λ_i) is a vertex in $\mathfrak{G}_{\Lambda,\Psi}(\Gamma)$ and

$$I \models \lambda_i^p(\mathbf{d}^*) \land \neg \lambda_i^p(\mathbf{d}^*)$$

with $j \neq i$.

Proof. Since Λ is a partition of λ , by Lemma 11, $At(I, \lambda_1)$ and $At(I, \lambda_2)$ form a partition of $At(I, \lambda)$. Furthermore, since $p(\mathbf{d}^*)$ is a vertex in $\mathfrak{G}_{I,At(I,\lambda)}(\Gamma)$, by definition, we get that $p(\mathbf{d}^*)$ belongs to $At(I, \lambda)$. Hence, $p(\mathbf{d}^*)$ belongs to $At(I, \lambda_i)$ for exactly one $i \in \{1, 2\}$ and, thus, $I \models p(\mathbf{d}^*) \wedge \lambda_i(\mathbf{d}^*)$. We also get that $I \not\models p(\mathbf{d}^*) \wedge \lambda_j(\mathbf{d}^*)$ with $j = \{1, 2\}$ such that $j \neq i$. Therefore, $I \models \lambda_i(\mathbf{d}^*) \wedge \neg \lambda_j(\mathbf{d}^*)$. Finally, since $I \models p(\mathbf{d}^*) \wedge \lambda_i(\mathbf{d}^*)$ and I is a model of Ψ , we get $\Psi \cup \{\exists \mathbf{X}\lambda_i^p\}$ is satisfiable. Therefore (p, λ_i) is a vertex of $\mathfrak{G}_{\Lambda,\Psi}(\Gamma)$.

Given a vertex $p(\mathbf{d}^*)$ in $\mathfrak{G}_{I,At(I,\lambda)}(\Gamma)$, by $[p(\mathbf{d}^*)]$ we denote the vertex (p, λ_i) in $\mathfrak{G}_{\Lambda}(\Gamma)$ such that $I \models \lambda_i^p(\mathbf{d}^*)$. Note that, by Lemma 15, such vertex exists and it is unique.

Lemma 16 Let Γ and Ψ be two theories and I be a model of Ψ . If $\Lambda = {\lambda_1, \lambda_2}$ is a partition of λ and (v, w) is an edge in $\mathfrak{G}_{I,At(I,\lambda)}(\Gamma)$, then ([v], [w]) is an edge in $\mathfrak{G}_{\Lambda,\Psi}(\Gamma)$.

Proof. Since (v, w) is an edge in $\mathfrak{G}_{I,At(I,\lambda)}(\Gamma)$, there is a rule $B \to H$ of Γ with free variables \mathbf{X} and a tuple of domain elements \mathbf{x} such that $v \in Sp_I(H_{\mathbf{x}}^{\mathbf{X}})$ and $w \in Pnn_I(B_{\mathbf{x}}^{\mathbf{X}})$. Let $v = p(\mathbf{d}^*)$ and $w = q(\mathbf{e}^*)$. By Lemma 14, this implies that

- there is some a strictly positive occurrence $p(\mathbf{t})$ of predicate symbol p in H such that I satisfies $Sp_{\Psi}(H_{\mathbf{x}}^{\mathbf{X}})_{\mathbf{d}^*}^{\mathbf{Y}}$ where \mathbf{Y} are the free variables in $Sp_{\Psi}(H_{\mathbf{x}}^{\mathbf{X}})$.
- there is some a positive nonnegated occurrence $q(\mathbf{s})$ of predicate symbol q in B such that I satisfies $Pnn_{\Psi}(B_{\mathbf{x}}^{\mathbf{X}})_{\mathbf{e}^*}^{\mathbf{Z}}$ where \mathbf{Z} are the free variables in $Pnn_{\Psi}(B_{\mathbf{x}}^{\mathbf{X}})$.

Let $[w] = (p, \lambda_v)$ and $[v] = (q, \lambda_w)$. From Lemma 15, we get $I \models \lambda_v^p(\mathbf{d}^*) \wedge \lambda_w^q(\mathbf{e}^*)$ and, thus,

$$I \models Pnn_{\Psi}(B_{\mathbf{x}}^{\mathbf{X}})_{\mathbf{d}^{*}}^{\mathbf{Y}} \wedge Sp_{\Psi}(H_{\mathbf{x}}^{\mathbf{X}})_{\mathbf{e}^{*}}^{\mathbf{Z}} \wedge \lambda_{v}^{p}(\mathbf{d}^{*}) \wedge \lambda_{w}^{q}(\mathbf{e}^{*})$$

which implies that

$$I \models \exists \mathbf{XYZ} \big(Pnn_{\Psi}(B) \land Sp_{\Psi}(H) \land \lambda_{v}^{p}(\mathbf{Y}) \land \lambda_{w}^{q}(\mathbf{Z}) \big)$$

Since *I* is a model of Ψ , this implies that (18) is satisfiable. Consequently, ([v], [w]) is an edge in $\mathfrak{G}_{\Lambda}(\Gamma)$.

Lemma 17 Let Γ and Ψ be two theories and I be a model of Ψ . If $\Lambda = \{\lambda_1, \lambda_2\}$ is a partition of λ that is separable on $\mathfrak{G}_{\Lambda,\Psi}(\Gamma)$, then $\{At(I, \lambda_1), At(I, \lambda_2)\}$ is a partition of $At(I, \lambda)$ and it is separable on $\mathfrak{G}_{I,At(I,\lambda)}(\Gamma)$ for any interpretation I.

Proof. Assume that Λ is a partition of λ that is separable on $\mathfrak{G}_{\Lambda,\Psi}(\Gamma)$. Then, by Lemma 11, we get that $\{At(I,\lambda_1), At(I,\lambda_2)\}$ is a partition of $At(I,\lambda)$. Let us show that this partition is separable on $\mathfrak{G}_{I,At(I,\lambda)}(\Gamma)$.

Pick any infinite walk v_1, v_2, v_3, \ldots of $\mathfrak{G}_{I,At(I,\lambda)}(\Gamma)$. By Lemma 16, we get that $[v_1], [v_2], [v_3], \ldots$ is an infinite walk of $\mathfrak{G}_{\Lambda,\Psi}(\Gamma)$. In addition, by Lemma 15, each $v_k \in At(I, \lambda_i)$ satisfies $[v_k] = (p, \lambda_i)$ for some predicate symbol p. Hence,

$$\{k \mid v_k \in At(I, \lambda_i)\} = \{k \mid [v_k] = (p, \lambda_i)\}\$$

for $1 \le i \le 2$. Since Λ is separable on $\mathfrak{G}_{\Lambda,\Psi}(\Gamma)$, at least one of this sets if finite and, therefore, $\{At(I,\lambda_1), At(I,\lambda_2)\}$ is separable on $\mathfrak{G}_{I,At(I,\lambda)}(\Gamma)$.

Lemma 18 Let Γ and Ψ be two theories and I be a model of Ψ . If Γ is Ψ -negative on λ , then Γ is negative on I and $At(I, \lambda)$.

Proof. Assume that Γ is Ψ -negative in λ and suppose, for the sake of contradiction, that Γ is not negative on I and $At(I, \lambda)$, that is, that there is some ground atom $p(\mathbf{d}^*)$ that belongs to $Sp_I(\Gamma) \cap At(I, \lambda)$. Since $p(\mathbf{d}^*)$ that belongs to $At(I, \lambda)$, it follows that

$$I \models p(\mathbf{d}^*) \land \lambda^p(\mathbf{d}^*)$$

Furthermore, since $p(\mathbf{d}^*)$ belongs to $Sp_I(\Gamma)$, there is a rule $B \to H$ of Γ with free variables \mathbf{X} and a tuple \mathbf{x} of names of domain elements of the appropriate length and sorts such that $I \models B_{\mathbf{x}}^{\mathbf{X}}$ and $p(\mathbf{d}^*)$ belongs to $Sp_I(H_{\mathbf{x}}^{\mathbf{X}})$. By Lemma 14, there is a strictly positive occurrence $p(\mathbf{t})$ of p in H such that I satisfies $Sp_{\Psi}(H_{\mathbf{x}}^{\mathbf{X}})_{\mathbf{d}^*}^{\mathbf{Y}}$ where \mathbf{Y} are the free variables in $Sp_{\Psi}(H_{\mathbf{x}}^{\mathbf{X}})$. Hence,

$$I \models B_{\mathbf{x}}^{\mathbf{X}} \wedge Sp_{\Psi}(H_{\mathbf{x}}^{\mathbf{X}})_{\mathbf{d}^*}^{\mathbf{Y}} \wedge \lambda^p(\mathbf{d}^*)$$

which implies that

$$I \models \Psi \cup \{ \exists \mathbf{X} \mathbf{Y} (B \land Sp_{\Psi}(H) \land \lambda^{p}(\mathbf{Y})) \}$$

This implies that Γ is Ψ -negative, which is a contradiction with the assumption. Consequently, Γ is negative on I and $At(I, \lambda)$.

Lemma 19 Let $\Gamma = \Gamma_1 \cup ... \cup \Gamma_n$ and Ψ be two theories, and let and $\Lambda = \{\lambda_1, ..., \lambda_n\}$ be a partition of λ such that

- Λ is separable on $\mathfrak{G}_{\Lambda,\Psi}(\Gamma)$; and
- each Γ_i is Ψ -negative on λ_j for all $j \neq i$.

Then, for any interpretation I that satisfies Ψ , the following two statements are equivalent

- *I* is a λ -stable model of Π , and
- *I* is a λ_i -stable model of \prod_i for all $1 \le i \le n$.

Proof. Assume that n = 2. The proof for n > 3 follows by induction on n. Pick any model I of Ψ . Since Λ is a partition of λ that is separable on $\mathfrak{G}_{\Lambda,\Psi}(\Gamma)$, by Lemma 17, it follows that $\{At(I, \lambda_1), At(I, \lambda_2)\}$ is a partition of $At(I, \lambda)$ and it is separable on $\mathfrak{G}_{I,At(I,\lambda)}(\Gamma)$ for any interpretation I. Similarly, since Γ_1 is negative on λ_2 and Γ_2 is negative on λ_1 , by Lemma 18, it follows that Γ_1 is negative on I and $At(I, \lambda_2)$ and Γ_2 is negative on I and $At(I, \lambda_1)$. Therefore, the conditions of Theorem 3 are met and, thus, the following two statements are equivalent

- 1. *I* is a A-stable model of Γ , and
- 2. *I* is a A_i -stable model of Γ_i for all $i \in \{1, 2\}$.

Finally, the result follows by Proposition 3.

Proof of Theorem 2. From Proposition 2, if *I* is a λ -stable model of Γ , then *I* is a λ_i -stable model of Γ_i for all $1 \leq i \leq n$. Since Ψ is an λ -approximator of Γ , by definition, it also follows that *I* is a model of Ψ . *The other direction*. Pick any model *I* of Ψ such that *I* a λ_i -stable model of Γ_i for all $1 \leq i \leq n$. From Lemma 19, this implies that *I* is a λ -stable model of Γ

Splitting Theorem for Disjuntive Logic Programs

Lemma 20 Let *F* be a sentence and $p(\mathbf{t})$ be a strictly positive occurrence in *F*. Then, $Sp_{\emptyset}(F) \models F \land p(\mathbf{t})$.

Proof. By induction in the size of the formula. *Case 1.* If F is an atom $p(\mathbf{t})$, then $Sp_{\emptyset}(F) = p(\mathbf{t}) \wedge \mathbf{Y} = \mathbf{t}$ and the result holds. *Case 2.* If $F = F_1 \wedge F_2$, then $Sp_{\emptyset}(F) = Sp_{\emptyset}(F_1) \wedge Sp_{\emptyset}(F_2) = Sp_{\emptyset}(F_i) \wedge F_j$ with F_i be subformula containing the occurrence $p(\mathbf{t})$ and j = 3 - i. By induction hypothesis, we get that $Sp_{\emptyset}(F_i) \models F_i \wedge p(\mathbf{t})$. Therefore, the result holds. *Case 3.* If $F = F_1 \vee F_2$, then $Sp_{\emptyset}(F) = Sp_{\emptyset}(F_i)$ with F_i containing the occurrence. By induction hypothesis, we get that $Sp_{\emptyset}(F_i) \models F_i \wedge p(\mathbf{t})$. Therefore, the result holds. *Cases 4 and 5.* If $F = \forall XG$ or $F = \exists XG$, then $Sp_{\emptyset}(F) = \exists XSp_{\emptyset}(G)$. By induction hypothesis, we get that $Sp_{\emptyset}(G_x^X) \models G_x^X \wedge p(\mathbf{t})$. Therefore, the result holds. *Case 6.* If $F = F_1 \rightarrow F_2$, then

$$Sp_{\emptyset}(F) = F_1^{\emptyset} \wedge Sp_{\emptyset}(F_2) \tag{31}$$

If F_1 is unsatisfiable, then (31) F_1^{\emptyset} is \perp and, thus, $Sp_{\emptyset}(F)$ is equivalent to \perp . Therefore, the result holds. Otherwise, F_1^{\emptyset} is F_1 and, thus, (31) is $F_1 \wedge Sp_{\emptyset}(F_2)$. By induction hypothesis, we get that $Sp_{\emptyset}(F_2) \models F_2 \wedge p(\mathbf{t})$. As a result, we can see that $Sp_{\emptyset}(F) \models F_1 \wedge F_2 \wedge p(\mathbf{t}) \models F \wedge p(\mathbf{t})$ and the result holds.

Lemma 21 Let F be a implication-free sentence and $p(\mathbf{t})$ be a positive nonnegated occurrence in F. Then, $Pnn_{\emptyset}(F) \models F \land p(\mathbf{t}).$

Proof. If F is implication-free, then $Pnn_{\emptyset}(F) = Pnn_{\emptyset}(F)$ and the result follows by Lemma 20.

Lemma 22 Let Π be a logic program and Λ be a partition of some intensionality statement. Then, $\mathfrak{G}_{\Lambda,\emptyset}(\Pi)$ is a subgraph of $\mathfrak{G}_{\Lambda}(\Pi)$.

Proof. Pick any vertex (p, λ) be a of $\mathfrak{G}_{\Lambda,\emptyset}(\Pi)$. Then, $\exists \mathbf{X} \lambda^p(\mathbf{X})$ is satisfiable and, thus, (p, λ) is a vertex of $\mathfrak{G}_{\Lambda}(\Pi)$. Pick an edge from (p, λ_i) to (q, λ_j) in $\mathfrak{G}_{\Lambda,\emptyset}(\Pi)$. Then, there is some rule $B \to H$ in Π , a strictly positive occurrence of p in H of the form $p(\mathbf{t})$, a positive nonnegated occurrence of q in B of the form $q(\mathbf{r})$ and

$$\Psi \cup \left\{ \exists \mathbf{XYZ} \left(Pnn_{\Psi}(B) \land Sp_{\Psi}(H) \land \lambda_{j}^{q}(\mathbf{Y}) \land \lambda_{i}^{p}(\mathbf{Z}) \right) \right\}$$
(32)

is satisfiable, where **X** are the free variables in $B \to H$, and **Y** and **Z** respectively are the free variables in formulas $Pnn_{\Psi}(B)$ and $Sp_{\Psi}(H)$ that are not in **X**. Note that, since Π is a program, it follows that H is a disjunction of the form $H_1 \vee \ldots \vee H_n$ with $p(\mathbf{t}) = H_i$ for some $i \in \{1, \ldots, n\}$. By construction, this implies that $Sp_{\Psi}(H)$ is $p(\mathbf{t}) \wedge \mathbf{t} = \mathbf{Y}$. Similarly, we have that B is a conjunction of the form $B_1 \wedge \ldots \wedge B_m$ with $q(\mathbf{r}) = B_j$ for some $j \in \{1, \ldots, m\}$. Therefore, there is a interpretation I that satisfies

$$\exists \mathbf{X} \left(B \wedge p(\mathbf{t}) \wedge \lambda_i^q(\mathbf{t}) \wedge \lambda_i^p(\mathbf{r}) \right)$$

Hence, there is also an edge from (p, λ_i) to (q, λ_j) in $\mathfrak{G}_{\Lambda}(\Pi)$.

Lemma 23 Let Π be a logic program and λ be a intensionality statement. If Π is negative on λ , then it is \emptyset -negative on λ .

Proof. Pick any rule $B \to H$ in Π and any strictly positive occurrence $p(\mathbf{t})$ in H. Suppose, for the sake of contradiction, that $\exists \mathbf{XY} (B \land SP(H) \land \lambda^p(\mathbf{Y}))$ is satisfiable. Since Π is a program, it follows that H is a disjunction of the form $H_1 \lor \ldots \lor H_n$ with $p(\mathbf{t}) = H_i$ for some $i \in \{1, \ldots, n\}$. By construction, this implies that $Sp_{\Psi}(H)$ is $p(\mathbf{t}) \land \mathbf{t} = \mathbf{Y}$. Hence, there is a interpretation I that satisfies $\exists \mathbf{X} (B \land p(\mathbf{t}) \land \lambda^p(\mathbf{t}))$. However, since Π is negative on λ , it follows that $\exists \mathbf{X} (B \land p(\mathbf{t}) \land \lambda^p(\mathbf{t}))$ is unsatisfiable. Therefore, $\exists \mathbf{XY} (B \land SP(H) \land \lambda^p(\mathbf{Y}))$ is unsatisfiable and Π is \emptyset -negative on λ .

Proof of Theorem 1. Since Π is a program, by Lemmas 22 and 23, we get that

• Λ is separable on $\mathfrak{G}_{\Lambda}(\Pi)$; and

• each Π_i is negative on λ_j under context Ψ for all $j \neq i$.

The result follows directly from Theorem 2 taking the empty context. $\hfill \Box$