

Accurate, provable, and fast nonlinear tomographic reconstruction: A variational inequality approach

Mengqi Lou[§], Kabir Aladin Verchand^{§,||}, Sara Fridovich-Keil^{‡,†}, Ashwin Pananjady^{§,†}

Georgia Institute of Technology, School of Industrial and Systems Engineering[§] and
School of Electrical and Computer Engineering[†]
University of Cambridge, Statistical Laboratory^{||}
Stanford University, Department of Electrical Engineering[‡]

March 27, 2025

Abstract

We consider the problem of signal reconstruction for computed tomography (CT) under a nonlinear forward model that accounts for exponential signal attenuation, a polychromatic X-ray source, general measurement noise (e.g. Poisson shot noise), and observations acquired over multiple wavelength windows. We develop a simple iterative algorithm for single-material reconstruction, which we call EXACT (EXtragradiant Algorithm for Computed Tomography), based on formulating our estimate as the fixed point of a monotone variational inequality. We prove guarantees on the statistical and computational performance of EXACT under practical assumptions on the measurement process. We also consider a recently introduced variant of this model with Gaussian measurements, and present sample and iteration complexity bounds for EXACT that improve upon those of existing algorithms. We apply our EXACT algorithm to a CT phantom image recovery task and show that it often requires fewer X-ray projection exposures, lower source intensity, and less computation time to achieve similar reconstruction quality to existing methods.

1 Introduction and methodology

X-ray computed tomography (CT) is a core imaging modality in modern medicine, and is also widely used in diverse nonmedical applications ranging from security inspection to nondestructive testing. CT typically works by rotating an X-ray source and detector along a prespecified trajectory around the subject, and recording the total energy or number of photons that are transmitted to the detector at each angle along the tomographic trajectory. One then solves the inverse problem of reconstructing the underlying 2D or 3D image of the subject from these measurements. Many CT reconstruction techniques have been proposed across multiple communities (see, e.g., [Elbakri and Fessler, 2002](#); [Geyer et al., 2015](#); [Schirra et al., 2013](#); [Sawatzky et al., 2014](#); [Schmidt et al., 2017](#); [Alvarez and Macovski, 1976](#); [Jadue et al., 2024](#)), and several algorithms are now built into commercial CT scanners (see, e.g., [Fu et al., 2016](#)). In this paper, we focus on photon-counting CT detectors ([Taguchi and Iwanczyk, 2013](#); [Persson et al., 2020](#); [Danielsson et al., 2021](#); [Nakamura et al., 2023](#)), though aspects of our methodology should be immediately transferrable to so-called “energy-integrating” detectors ([Sukovic and Clinthorne, 1999](#); [Wang and Pelc, 2012](#)).

The starting point of any reconstruction algorithm is a forward model for CT measurements; see [Tang \(2023\)](#); [Fu et al. \(2016\)](#); [Gjesteby et al. \(2016\)](#) for an overview and discussion of the importance of modeling and artifacts induced by preprocessing. The forward model for photon-

counting CT (see Mory et al. (2018, Eq. (2)) or Schmidt et al. (2023, Eq.(3))) is given by

$$y_{\omega,i}(x) = I_{\omega,i} \sum_{j=1}^W s_{\omega,i,j} \exp \left(- \sum_{m=1}^M \sum_{k=1}^d \mu_{m,j} A_{i,k} x_{k,m} \right). \quad (1)$$

Note that our notation differs slightly from the aforementioned papers, and we use the following convention:

- ω indexes over wavelength windows in the X-ray detector;
- i indexes over the n angles used to collect X-ray projection measurements, where each angle corresponds to a ray through the imaging target;
- $I_{\omega,i}$ is the (random) number of incident photons (produced by the X-ray source) in wavelength window ω at angle i ; we denote its mean by the *intensity* $\bar{I}_{\omega,i} := \mathbb{E}[I_{\omega,i}]$;
- $y_{\omega,i}$ is the (random) number of transmitted photons that arrive at the X-ray detector in wavelength window ω at angle i ;
- j indexes over W individual X-ray wavelengths that fall inside wavelength window ω ;
- $s_{\omega,i,j}$ is the normalized spectral sensitivity of the detector; $\sum_j s_{\omega,i,j} = 1$;
- m indexes over the M basis materials that are expected to be present in the imaging target;
- k indexes over the d total pixels or voxels in the imaging target;
- $\mu_{m,j}$ is the mass attenuation coefficient for basis material m at X-ray wavelength j ;
- $A_{i,k}$ is the entry of the projection matrix (derived from the Radon transform) denoting the fractional contribution of pixel/voxel k to the line integral at angle i — each such scalar is nonnegative;
- $\{x_{k,m}\}$ represents the unknown target image with k indexing spatial location (pixel or voxel index) and m indexing basis material; each scalar $x_{k,m}$ is nonnegative.

In this forward model, the strictly positive scalars $s_{\omega,i,j}$ and $\mu_{m,j}$ can be estimated in a calibration step, as can the average intensities $\bar{I}_{\omega,i}$. Accordingly, we assume that these quantities are known. For notational convenience, we assume that the detector sensitivity $s_{\omega,i,j}$ and average incident photon count $\bar{I}_{\omega,i}$ depend only on wavelength (governed by ω and j) and not on angle i , to write the forward model (in expectation) as

$$\mathbb{E}[y_{\omega,i}(x)] = \bar{I}_{\omega} \sum_{j=1}^W s_{\omega,j} \exp \left(- \sum_{m=1}^M \sum_{k=1}^d \mu_{m,j} A_{i,k} x_{k,m} \right).$$

We focus on a special case of this forward model that corresponds to a single basis material ($M = 1$), allowing us to remove the indexing over m and thereby write the model as

$$\mathbb{E}[y_{\omega,i}(x)] = \bar{I}_{\omega} \sum_{j=1}^W s_{\omega,j} \exp \left(- \sum_{k=1}^d \mu_j A_{i,k} x_k \right).$$

The restriction to a single basis material is fundamental to our proposed optimization strategy – it is a reasonable assumption in CT applications to material science (Chandel et al., 2014; Sietins et al., 2021), and in medical CT it approximates perfusion imaging of a single contrast agent against a known background (Pourmorteza et al., 2016) as well as imaging of the human body without any contrast agents.

To arrive at the model we study in this paper, we restrict to a single detector window for notational convenience,¹ thereby removing our indexing over ω to write

$$\mathbb{E}[y_i(x)] = \bar{I} \sum_{j=1}^W s_j \exp(-\mu_j \langle a_i, x \rangle), \quad (2)$$

where $a_i \in \mathbb{R}^d$ is the row of A corresponding to projection angle i and $x \in \mathbb{R}^d$ is the target image or volume. In our analysis, we consider both a general model in which the projection vectors a_i are entrywise nonnegative (as they are in practice), as well as a model in which we can prove more precise results and compare directly with recent work, which assumes Gaussian measurements a_i . We unify these models using the convention of Fridovich-Keil et al. (2024); Charisopoulos and Willett (2024) and work with the forward model

$$\mathbb{E}[y_i(x)] = \bar{I} \sum_{j=1}^W s_j \exp(-\mu_j \langle a_i, x \rangle_+). \quad (3)$$

Here, we use the shorthand $t_+ = \max\{t, 0\}$ for any scalar $t \in \mathbb{R}$; the thresholding operation ensures that the ray projection values are always nonnegative even if the entries of a_i may be negative, as is possible in the Gaussian model.

In practice, we may have prior information about the signal, which we encode by assuming that $x \in X \subseteq \mathbb{R}^d$. For example, the set with small total variation (TV) norm (Strong and Chan, 2003; Osher et al., 2005) is one such structured set which is particularly suited to imaging applications; we use this TV constraint along with a nonnegativity constraint in our simulations. Prior work on signal recovery under models (1) and (3) is discussed in detail in Section 1.3.

1.1 Formal model and approach

As described above, we are interested in recovering an unknown signal from a nonlinear measurement model of the form (3). We assume throughout that our structured set $X \subseteq \mathbb{R}^d$ is a closed, convex set. Henceforth, we denote the ground truth signal by x_* , and our task is to recover $x_* \in X$ from measurement vectors $\{a_i\}_{i=1}^n$ and the associated responses $\{y_i\}_{i=1}^n$. Formally, we rewrite the measurement model as

$$\mathbb{E}[y_i|a_i] = \bar{I} \sum_{j=1}^W s_j \exp(-\mu_j \langle a_i, x_* \rangle_+) \quad \text{for } i = 1, \dots, n. \quad (4)$$

The parameters \bar{I} and $\{s_j, \mu_j\}_{j=1}^W$ are known and satisfy

$$\bar{I} > 0, \quad s_j \in (0, 1], \quad \sum_{j=1}^W s_j = 1, \quad \text{and} \quad \mu_j > 0 \quad \text{for all } j = 1, \dots, W.$$

¹This restriction is made purely for convenience and is not fundamental to our approach. Indeed, in our CT simulations we do include multiple (distinct but overlapping) detector windows, as visualized in Figure 1.

Besides the conditional expectation relation in Eq. (4), we do not place any assumptions on the conditional distribution $y_i|a_i$. This flexibility allows us to cover many practical noise models. A salient such example is the Poisson shot-noise model (Schmidt et al., 2023), in which $I_{\omega,i}$ in Eq. (1) is given by a Poisson random variable with intensity $\bar{I}_{\omega,i}$. In our notation, we have

$$y_i \stackrel{\text{i.i.d.}}{\sim} \text{Poi}\left(\bar{I} \sum_{j=1}^W s_j \exp(-\mu_j \langle a_i, x_\star \rangle_+)\right). \quad (5)$$

In the CT literature, there are other noise assumptions for this problem, including those that model pulse-pileup or additional noise induced due to instrumentation errors (Danielsson et al., 2021; Persson et al., 2020).

For notational convenience in the rest of the paper, we define the function $h : \mathbb{R} \rightarrow \mathbb{R}$ via

$$h(t) := \sum_{j=1}^W s_j \exp(-\mu_j t_+), \quad \text{for all } t \in \mathbb{R}, \quad (6)$$

where we recall that $t_+ = \max(0, t)$ denotes the ReLU operation. Using this shorthand, our model is given by

$$\mathbb{E}[y_i|a_i] = \bar{I} \cdot h(\langle a_i, x_\star \rangle).$$

Our approach is based on defining a vector-valued map $F : X \rightarrow \mathbb{R}^d$, given by

$$F(x) = \frac{1}{n} \sum_{i=1}^n [y_i - \bar{I} \cdot h(\langle a_i, x \rangle)] \cdot a_i, \quad \text{for all } x \in X. \quad (7)$$

By construction, the map F is continuous. Two crucial observations that will motivate our methodological development are:

- (a) The *expectation* of F — call this operator \bar{F} — has a zero at x_\star , in that $\bar{F}(x_\star) = 0$; and
- (b) The operator F is also *monotone*, in that for all $x, x' \in \mathbb{R}^d$, we have

$$\langle F(x) - F(x'), x - x' \rangle \geq 0.$$

Indeed, a short calculation to verify this fact is provided below; for any $x, x' \in \mathbb{R}^d$, we have

$$\langle F(x) - F(x'), x - x' \rangle = \frac{1}{n} \sum_{i=1}^n \bar{I} \cdot [h(\langle a_i, x' \rangle) - h(\langle a_i, x \rangle)] \cdot [\langle a_i, x \rangle - \langle a_i, x' \rangle] \geq 0, \quad (8)$$

where the inequality follows since the function $h(\cdot)$ is nonincreasing.

Given these observations, we now frame the problem as one of finding a fixed point of the variational inequality defined by the operator F ; see the classical references (Kinderlehrer and Stampacchia, 2000; Rockafellar and Wets, 2009) for background on variational inequalities. We are interested in a point $\hat{x} \in X$ that satisfies

$$\langle F(\hat{x}), x - \hat{x} \rangle \geq 0 \text{ for all } x \in X. \quad (9)$$

With the goal of converging to \hat{x} — which one should expect to be close to the zero x_\star of \bar{F} — we study the projected extragradient method (Korpelevich, 1976) for this problem, which proceeds as follows: With an initialization $x_1 \in X$, execute the steps

$$x_{t+0.5} = \mathcal{P}_X(x_t - \gamma_t F(x_t)), \quad (10a)$$

$$x_{t+1} = \mathcal{P}_X(x_t - \gamma_t F(x_{t+0.5})), \text{ for } t = 1, 2, \dots \quad (10b)$$

where γ_t is the step size at iterate t , and $\mathcal{P}_X : \mathbb{R}^d \rightarrow X$ is the ℓ_2 -projection operator onto X . Ignoring the cost of these projection operations, the rest of iteration (10) can be implemented in linear time. Furthermore, owing to the convexity of X , the projection step can typically be implemented via an efficient algorithm. For example, the total variation projection that we employ in our experiments is very fast in practice.

Our methodology is inspired by algorithmic ideas that were proposed for other monotone single-index models (Gallant, 1990; Kalai and Sastry, 2009; Kakade et al., 2011) as well as the recent formalism of these methods using variational inequalities due to Juditsky and Nemirovski (2019, 2020). We refer to this extragradient algorithm (10) applied to the specific CT forward model (4) as EXACT: EXtragradient Algorithm for Computed Tomography.

1.2 Additional notation

For a positive integer k , we let $[k]$ denote the set of natural numbers less than or equal to k . For $a, b \in \mathbb{R}$, we let $a \vee b = \max\{a, b\}$ and $a \wedge b = \min\{a, b\}$. For notational convenience, we define $A \in \mathbb{R}^{n \times d}$ to be the data matrix whose i -th row is the measurement vector a_i for each $i \in [n]$, and let $\Sigma = A^\top A \in \mathbb{R}^{d \times d}$ denote the associated correlation matrix. We let I_d denote the identity matrix of dimension $d \times d$. For a symmetric and positive semidefinite matrix $M \in \mathbb{R}^{d \times d}$ and a closed set $S \subseteq \mathbb{R}^d$, we define the restricted minimum and maximum eigenvalues as

$$\lambda_{\min}(M, S) = \min_{v \in S} \frac{\|Mv\|_2}{\|v\|_2} \quad \text{and} \quad \lambda_{\max}(M, S) = \max_{v \in S} \frac{\|Mv\|_2}{\|v\|_2}. \quad (11)$$

For two sets $S_1, S_2 \subseteq \mathbb{R}^d$, we define their Minkowski sum $S_1 + S_2 = \{s_1 + s_2 : s_1 \in S_1, s_2 \in S_2\}$, and denote their Minkowski difference by $S_1 - S_2$. For $r > 0$, we let $\mathbb{B}(r) := \{x \in \mathbb{R}^d : \|x\|_2 \leq r\}$ denote the ℓ_2 ball of radius r . For $v \in \mathbb{R}^d$ and $r > 0$, we let $\mathbb{B}(r; v) := \{x \in \mathbb{R}^d : \|x - v\|_2 \leq r\}$ denote the ℓ_2 ball of radius r centered at v . For $v \in \mathbb{R}^d \setminus \{0\}$, we let $P_v = vv^\top / \|v\|_2^2$ and $P_v^\perp = I_d - vv^\top / \|v\|_2^2$ denote $d \times d$ projection matrices that when applied to a vector, project it onto the subspace that is parallel and orthogonal to v , respectively. We let $\mathbb{1}\{\cdot\}$ denote the indicator function, which takes value 1 when its argument is true and 0 otherwise. We denote by $\mathcal{N}(\mu, M)$ a multivariate normal distribution with mean μ and covariance matrix M . For two sequences of non-negative reals $\{f_n\}_{n \geq 1}$ and $\{g_n\}_{n \geq 1}$, we use $f_n \lesssim g_n$ to indicate that there is a universal positive constant C such that $f_n \leq Cg_n$ for all $n \geq 1$. The relation $f_n \gtrsim g_n$ implies that $g_n \lesssim f_n$, and we say that $f_n \asymp g_n$ if both $f_n \lesssim g_n$ and $f_n \gtrsim g_n$ hold simultaneously. We also use standard order notation $f_n = \mathcal{O}(g_n)$ to indicate that $f_n \lesssim g_n$ and $f_n = \tilde{\mathcal{O}}(g_n)$ to indicate that $f_n \lesssim g_n \log^c n$, for a universal constant $c > 0$. We say that $f_n = \Omega(g_n)$ (resp. $f_n = \tilde{\Omega}(g_n)$) if $g_n = \mathcal{O}(f_n)$ (resp. $g_n = \tilde{\mathcal{O}}(f_n)$). The notation $f_n = o(g_n)$ is used when $\lim_{n \rightarrow \infty} f_n/g_n = 0$, and $f_n = \omega(g_n)$ when $g_n = o(f_n)$. Throughout, we use c, C to denote universal positive constants, and their values may change from line to line. All logarithms are with respect to the natural base unless otherwise stated.

1.3 Contributions and comparison to related work

With this setup in hand, we are now in a position to describe our contributions and to compare them with prior work.

1. **Convergence guarantees for general CT model.** We provide a rigorous convergence analysis for EXACT (Equation (10)) under the assumption that the measurement vectors and true signal have nonnegative entries, which is indeed the case in CT applications. Our results accommodate an arbitrary noise model, and we show that the eventual recovery error

depends explicitly on a functional of this noise process. We emphasize that our results provide (to our knowledge) the first rigorous guarantees for signal reconstruction in the practical CT measurement model (2) including realistic nonnegative measurements, a polychromatic X-ray source, multiple detector windows, and a general noise model. In more detail, we show that the iterates $\{x_t\}$ generated by EXACT converge linearly to within a small neighborhood of x_* when the restricted smallest eigenvalue of the matrix Σ is positive, specifically, when $\lambda_{\min}(\Sigma, X - X) > 0$. Note that such a condition is necessary even for the identifiability of the linear variant of the model, i.e., if we observed linear functions of the signal x_* using the classical Radon transform. For our detailed guarantees and examples see Theorem 2 and the accompanying discussion.

For comparison, we note that several other algorithms have been proposed for signal recovery in the model (2), based on logarithmic transformations as preprocessing (Maaß et al., 2009; Niu et al., 2014), conjugate gradient methods (Cai et al., 2013), separable quadratic surrogates (Long and Fessler, 2014; Weidinger et al., 2016; Mechlem et al., 2017), and the ADMM algorithm (Barber et al., 2016; Schmidt et al., 2017). See the paper by Mory et al. (2018) for a detailed empirical comparison of these methods on some synthetic and clinical reconstruction tasks. Preprocessing-based methods are known to perform poorly with noise since they are based on heuristic inversion of nonlinearities in the model (Mory et al., 2018), and all the iterative methods, save for one, do not come equipped with theoretical guarantees. The exception is the coSSCIR method by Schmidt et al. (2017), which has since been applied to a variety of CT reconstruction tasks (Sidky et al., 2018; Schmidt et al., 2022; Rizzo et al., 2022). Rigorous guarantees on its performance have been proved under assumptions such as restricted strong convexity of the associated Poisson negative log-likelihood objective function (Barber and Sidky, 2024). However, to our knowledge, this assumption has not been rigorously verified when the observations are generated² by either model (1) or model (2).

Besides our theoretical contribution, we also implement our algorithm in practice. In Section 3, we present a suite of experiments for recovering a phantom image from measurements of the form (3), comparing EXACT with algorithms that come with provable guarantees in at least some setting — in particular, coSSCIR (Schmidt et al., 2017) as well as two first-order methods (Fridovich-Keil et al., 2024; Charisopoulos and Willett, 2024) that we discuss shortly. Overall, our algorithm achieves the lowest recovery error among all these methods while also having the smallest running time. In particular, our approach often yields comparable reconstruction quality, both visually and quantitatively, using an order of magnitude lower X-ray dose compared to these baselines.

2. **Convergence guarantees for Gaussian model.** To further explore the statistical and algorithmic properties of our algorithm, we study convergence guarantees for EXACT with Gaussian measurements. Indeed, Fridovich-Keil et al. (2024) introduced a monochromatic version of the stylized model (3) under i.i.d. Gaussian measurements a_i in order to provide rigorous sample and iteration complexity guarantees on signal recovery. They studied the gradient descent algorithm on the mean squared error objective in the case with a single X-ray wavelength and noiseless measurements. A follow-up paper by Charisopoulos and Willett (2024) considered the same setting and proposed a different first-order method based on optimizing the nonsmooth ℓ_1 loss, and showed improved sample and iteration complexity

²The only exception to this that we are aware of is the monochromatic case, where the Poisson negative log-likelihood is a convex function (Barber and Sidky, 2024). Note that this function is nonconvex in the polychromatic models (2) and (3).

guarantees, again in the monochromatic, noiseless setting with Gaussian measurements.

Our guarantees in this setting are easiest to describe and compare with prior work when there is no structure and no noise. In the proportional regime, where $n \asymp d$, we show that it takes EXACT (Equation (10))

$$t = \mathcal{O}\left(\exp(24\|x_\star\|_2)\|x_\star\|_2^4 \log\left(\frac{5\|x_\star\|_2}{\epsilon}\right)\right) \text{ iterations to achieve } \|x_t - x_\star\| \leq \epsilon.$$

On the other hand, in the regime $n \asymp \|x_\star\|_2^4 d$, it takes EXACT

$$t = \mathcal{O}\left(\|x_\star\|_2^4 \log\left(\frac{5\|x_\star\|_2}{\epsilon}\right)\right) \text{ iterations to achieve } \|x_t - x_\star\| \leq \epsilon.$$

This guarantee (see Theorem 3 for details) reveals an interesting trade-off between statistical performance and computational efficiency. It indicates that practitioners can use fewer samples—specifically, $\mathcal{O}(d)$ —but this will result in increased computational time. Conversely, using more samples—approximately $\mathcal{O}(\|x_\star\|_2^4 d)$ —reduces computation time significantly.

Comparing with prior work in the monochromatic, noiseless setting with Gaussian measurements, we find that EXACT requires fewer samples and converges faster than the gradient descent method developed in Fridovich-Keil et al. (2024) as well as the non-smooth optimization method introduced by Charisopoulos and Willett (2024). Concretely, EXACT needs $\mathcal{O}(d)$ samples for recovery (without multiplicative factors in $\|x_\star\|_2$), whereas Fridovich-Keil et al. (2024) and Charisopoulos and Willett (2024) require $\mathcal{O}(\exp(\|x_\star\|_2)d)$ and $\mathcal{O}(\|x_\star\|_2^4 d)$ samples, respectively. There are also significant gains in iteration complexity. In the regime $n \gtrsim \|x_\star\|_2^4 d$, achieving ϵ -accurate signal recovery using EXACT requires $\mathcal{O}(\|x_\star\|_2^4 \log(5\|x_\star\|_2/\epsilon))$ iterations. This represents an improvement over the iteration complexity presented by Fridovich-Keil et al. (2024), which has a multiplicative factor scaling as $\exp(\|x_\star\|_2)$, and by Charisopoulos and Willett (2024), in which the iteration complexity is scaled by a polynomial factor of $\|x_\star\|_2$. For a detailed comparison, see Table 1.

2 Theoretical results

We begin by giving a general-purpose recovery guarantee for the EXACT algorithm in the presence of noisy measurements. Recall the continuous and monotone map F defined in Eq. (7). To state our general result in Theorem 1 to follow, we assume that F is L -Lipschitz continuous, in that it satisfies

$$\|F(x) - F(x')\|_2 \leq L\|x - x'\|_2 \quad \text{for all } x, x' \in X. \quad (12a)$$

We also assume that F is ν -strongly pseudo-monotone around the point x_\star , in that

$$\langle F(x) - F(x_\star), x - x_\star \rangle \geq \nu\|x - x_\star\|_2^2 \quad \text{for all } x \in X. \quad (12b)$$

Note that we always have $L \geq \nu \geq 0$. Conditions (12a) and (12b) will be verified concretely in the results to follow under variants of the following assumption³ that X is a bounded convex set, i.e.,

Assumption 1. *For some finite scalar $R > 0$, we have*

$$X \subseteq \{x \in \mathbb{R}^d : \|x - x_\star\|_2 \leq R\}.$$

³Assumption 1 can likely be relaxed by assuming that the algorithm is initialized at some fixed point, say 0.

Such an assumption is weak, and only requires some rough knowledge of the signal strength $\|x_\star\|_2$ to set a suitable value of the maximum radius R and thereby define the constraint set X . Note that X can obey further structural assumptions in addition to satisfying Assumption 1.

A crucial object that enters our guarantees is the vector

$$\zeta = F(x_\star) = \frac{1}{n} \sum_{i=1}^n (y_i - \bar{I} \cdot h(\langle a_i, x_\star \rangle)) \cdot a_i, \quad (13)$$

where the function h was defined in Eq. (6). Note that $\mathbb{E}[\zeta] = 0$ according to model (4) and in the noiseless case we have $\zeta = 0$. We further define the following quantity that will be a convenient measure of statistical error in the problem:

$$\text{Err}(x_\star, X) := \sup_{x, x' \in X} \left\langle \zeta, \frac{x - x'}{\|x - x'\|_2} \right\rangle. \quad (14)$$

This error quantity is a random variable that provides a measure how the noise interacts with the geometry of the constraint set X . For concreteness, let us explicitly evaluate $\text{Err}(x_\star, X)$ in some special cases.

Example 1. *In the noiseless case, we have $\zeta = 0$, and therefore, the error quantity $\text{Err}(x_\star, X) = 0$ pointwise for any x_\star and X .* ♣

Example 2. *Consider the Poisson shot-noise model (5), and suppose we are in the unstructured setting with $X = \mathbb{B}(R)$. Then there exists a universal positive constant C such that with probability at least $1 - \delta$, we have*

$$\text{Err}(x_\star, X) \leq \frac{\left(\bar{I} \cdot \sum_{i=1}^n \|a_i\|_2^2 \cdot h(\langle a_i, x_\star \rangle) \right)^{1/2}}{n} + C \cdot \frac{\bar{I} (\|AA^\top\|_F)^{1/2}}{n} \cdot \log(2/\delta).$$

We provide a proof of this claim in Section A.1. ♣

We are now ready to state our first result, which is a convergence result for monotone VIs.

Theorem 1. *Suppose that the data $\{(a_i, y_i)\}_{i=1}^n$ satisfy the single-index relationship (4), Assumption 1 holds, and that the operator F in Eq. (7) is both L -Lipschitz (12a) as well as ν -strongly pseudo-monotone (12b). Then the iterates $\{x_t\}$ generated according to EXACT (10) with step size $\gamma_t = 1/(4L)$ and any $x_1 \in X$ satisfy, for all $t > 0$,*

$$\|x_{t+1} - x_\star\|_2 \leq \underbrace{\left(1 - \frac{\nu}{8L}\right)^{t/2} \|x_1 - x_\star\|_2}_{\text{Optimization error}} + \underbrace{\frac{4\text{Err}(x_\star, X)}{\nu}}_{\text{Statistical error}}.$$

While the quantity $\text{Err}(x_\star, X)$ is random, the conclusion of Theorem 1 is deterministic. An immediate consequence is that for each $\epsilon \in (0, 1)$, if algorithm (10) is run for

$$t \geq \frac{16L}{\nu} \log \left(\frac{\|x_1 - x_\star\|_2}{\epsilon} \right) \quad \text{iterations, then} \quad \|x_{t+1} - x_\star\|_2 \leq \epsilon + \frac{4\text{Err}(x_\star, X)}{\nu}. \quad (15)$$

Since we can set ϵ to be arbitrarily small, the eventual estimation error is controlled by the strong pseudo-monotonicity parameter ν and the quantity $\text{Err}(x_\star, X)$ that depends on the geometry of the set X . The convergence is geometric, and the rate is governed by the condition number L/ν . Note that our convergence guarantee decomposes into a sum of two terms, the first capturing the

optimization error and the second capturing the eventual statistical error. We provide the proof of Theorem 1 in Section 4.1. Since we impose strong pseudo-monotonicity only around x_* and not the sample fixed point \hat{x} , the proof is non-standard and requires careful decompositions and control of the various error terms. Finally, let us remark that while Theorem 1 is stated for the CT model (4), it should really be thought of as a result for the extragradient algorithm if applied to an operator F of the form (7), if this operator happens to be Lipschitz and strongly pseudo-monotone. Accordingly, this result may find broader applications in the study of monotone single-index models.

2.1 Positive measurements and positive signal

In this section, we assume that the measurement vectors and the signal x_* all have non-negative entries. Note that in computed tomography applications, the measurement vector a_i is a set of known coefficients derived from the Radon transform (Radon, 1986), whose entries are non-negative.

Recall that we use $X - X := \{x - x' : x, x' \in X\}$ to denote the Minkowski difference between the set X and itself. Recall the definition of restricted eigenvalues from Eq. (11).

Assumption 2. *For each $i \in [n]$, the measurement vector a_i has nonnegative entries. Furthermore, the signal x_* has nonnegative entries. The structured set satisfies $X \subseteq \{x \in \mathbb{R}^d : x \geq 0\}$. The correlation matrix $\Sigma = A^\top A$ satisfies $\lambda_{\min}(\Sigma, X - X) > 0$.*

Under Assumption 2, we verify in Section 4.2 that the strong pseudo-monotonicity parameter ν and the Lipschitz parameter L satisfy

$$\nu \geq \bar{I} \cdot \frac{\lambda_{\min}(\Sigma, X - X)}{n} \sum_{j=1}^W s_j \mu_j \exp\left(-\mu_j \max_{x \in X, i \in [n]} \langle a_i, x \rangle\right) \quad \text{and} \quad L \leq \bar{I} \cdot \frac{\lambda_{\max}(\Sigma, \mathbb{R}^d)}{n} \sum_{j=1}^W s_j \mu_j,$$

respectively. To reduce the notational burden, we define the quantity

$$\kappa = \exp\left(\max_{j \in [W]} \mu_j \cdot \max_{x \in X, i \in [n]} \langle a_i, x \rangle\right) \cdot \frac{\lambda_{\max}(\Sigma, \mathbb{R}^d)}{\lambda_{\min}(\Sigma, X - X)}, \quad (16)$$

which is an upper bound on the condition number L/ν . Under Assumptions 1 and 2, the scalar κ is guaranteed to be nonnegative and finite. As a consequence, applying Theorem 1 yields the following result.

Theorem 2. *Suppose that the data $\{(a_i, y_i)\}_{i=1}^n$ satisfy the single-index relationship (4), and recall the model parameters $\{s_j, \mu_j\}_{j=1}^W$ and \bar{I} in Eq. (4) and the error quantity $\text{Err}(x_*, X)$ in Eq. (14). Suppose Assumption 1 holds for some $R > 0$, Assumption 2 holds, and the iterates $\{x_t\}$ are generated according to EXACT (10) with step size*

$$\gamma_t = \frac{n}{4\bar{I} \cdot \lambda_{\max}(\Sigma, \mathbb{R}^d) \cdot \sum_{j=1}^W s_j \mu_j} \quad \text{and} \quad x_1 \in X.$$

Then, for each $\epsilon \in (0, 1)$, as long as $t \geq 16\kappa \cdot \log\left(\frac{\|x_1 - x_\|_2}{\epsilon}\right)$, we have*

$$\|x_t - x_*\|_2 \leq \epsilon + \frac{n}{\lambda_{\min}(\Sigma, X - X)} \cdot \frac{4\text{Err}(x_*, X)}{\bar{I}} \cdot \frac{1}{\sum_{j=1}^W s_j \mu_j \exp\left(-\mu_j \max_{x \in X, i \in [n]} \langle a_i, x \rangle\right)}.$$

We provide the proof of Theorem 2 in Section 4.2. The bulk of the proof involves computing bounds on the Lipschitz parameter L and the strong pseudo-monotonicity parameter ν . Equipped with these estimates, our convergence guarantee is an immediate consequence of our general-purpose result in Theorem 1. Two additional remarks are in order.

First, we emphasize that Theorem 2 is applicable to real CT observation models in which the measurements are derived from the Radon transform and consist of non-negative entries. To our knowledge, this constitutes the first provable recovery guarantee in the measurement model (4) with realistic modeling assumptions. By contrast, previous work focusing on rigorous guarantees studies non-convex formulations of the problem and provides guarantees under either an unrealistic Gaussian measurement assumption (Fridovich-Keil et al., 2024; Charisopoulos and Willett, 2024) or a restricted strong convexity assumption that has not been verified in applications to CT (Barber and Sidky, 2024).

Second, let us elaborate on our sample size condition, considering the noiseless case — where $\zeta = 0$ and $\text{Err}(x_\star, X) = 0$ — for simplicity. From Theorem 2, we obtain that the iterates $\{x_t\}$ converge linearly to the ground truth x_\star as long as the minimum restricted eigenvalue is positive, i.e., $\lambda_{\min}(\Sigma, X - X) > 0$. Note that this condition would have been necessary even *without* nonlinearities in the model. In the unstructured setting where $X = \mathbb{B}(R; x_\star)$, the condition $\lambda_{\min}(\Sigma, X - X) > 0$ typically holds once $n \geq d$, i.e., we have more observations than unknowns. When X captures meaningful structure in the signal, the condition $\lambda_{\min}(\Sigma, X - X) > 0$ can hold with $n \ll d$, allowing for compressive sensing.

2.2 Gaussian measurements

In this section, we consider the model in which measurements are assumed to be Gaussian random vectors. The main attractions of this assumption are its analytical tractability and its facilitation of comparison to previous theoretical results (Fridovich-Keil et al., 2024; Charisopoulos and Willett, 2024). Let us state this assumption concretely:

Assumption 3. *The measurements are generated at random, according to $\{a_i\}_{i=1}^n \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, I_d)$.*

Recalling our notation for projection matrices, we first define a few quantities that are useful in the statement of our results. Define the geometric quantity $\bar{\omega}(x_\star, X)$, which depends on the local behavior of the set X around x_\star as

$$\bar{\omega}(x_\star, X) := \mathbb{E} \left\{ \sup_{v \in X} \frac{\langle P_{x_\star}^\perp v, g \rangle}{\|P_{x_\star}^\perp v\|_2} \right\}, \quad \text{where } g \sim \mathcal{N}(0, I_d). \quad (17)$$

Note that $\bar{\omega}(x_\star, X)$ has a natural interpretation as the Gaussian width of a certain set intersected with the unit shell. Next, we define $\gamma_\star \in (0, +\infty)$ as the solution of the following fixed point equation:

$$\mathbb{E} \left\{ \frac{|h'(6\|x_\star\|_2 G)|^2 \cdot \mathbb{1}\{G > 0\}}{[|h'(6\|x_\star\|_2 G)| + \gamma_\star]^2} \cdot \int_{-G/4}^{G/4} \frac{t^2 e^{-t^2/2}}{\sqrt{2\pi}} dt \right\} = \frac{16\bar{\omega}(x_\star, X)^2}{n}. \quad (18)$$

We prove in Appendix A.2 that there exists a universal positive constant C such that if $n \geq C\bar{\omega}(x_\star, X)^2$, then γ_\star is unique and strictly positive. We further define the scalar quantity

$$\rho(\|x_\star\|_2) := \left(\sum_{j=1}^W \frac{s_j \mu_j}{(\mu_j \|x_\star\|_2)^4 \vee 1} \right)^{-1}, \quad (19)$$

where we recall the notation $a \vee b = \max(a, b)$. Note that $\rho(\|x_\star\|_2)$ grows at most polynomially in $\|x_\star\|_2$; in particular, we have

$$\rho(\|x_\star\|_2) = \mathcal{O}(\|x_\star\|_2^4 \vee 1), \quad (20)$$

where in the notation $\mathcal{O}(\cdot)$ we hide dependence on the set of constants $\{s_j, \mu_j\}_{j=1}^W$. We are now ready to state our main result for Gaussian measurements, which is the first such result that incorporates measurement noise.

Theorem 3. *Suppose that the data $\{(a_i, y_i)\}_{i=1}^n$ satisfy the single-index relationship (4). Recall the model parameters $\{s_j, \mu_j\}_{j=1}^W$ and \bar{I} in Eq. (4) and the error quantity $\text{Err}(x_\star, X)$ in Eq. (14). Further, suppose Assumption 1 holds with $R = 4\|x_\star\|_2$, Assumption 3 holds, and the iterates $\{x_t\}$ are generated according to EXACT (10) with step size*

$$\gamma_t = \frac{(n+d)/n}{40\bar{I} \sum_{j=1}^W s_j \mu_j} \quad \text{and} \quad x_1 \in \mathbb{B}(4\|x_\star\|_2; x_\star).$$

Recall the scalars $\bar{\omega}(x_\star, X)$, $\rho(\|x_\star\|_2)$ and γ_\star defined in Eqs. (17), (19) and (18), respectively. Then there exists a tuple of universal, positive constants (c, C, C_1, C_2) such that if

$$n \geq C \cdot \bar{\omega}(x_\star, X)^2 \quad \text{and} \quad \sqrt{\frac{\log(n)}{n}} \leq c \frac{\rho(\|x_\star\|_2)^{-1}}{\sum_{j=1}^W s_j \mu_j}, \quad (21)$$

then the following holds with probability at least $1 - 16 \exp(-\bar{\omega}(x_\star, X)^2/64) - 12n^{-10} - 2 \exp(-d)$. For all $\epsilon \in (0, 1)$, as long as

$$t \geq C_1 \frac{n+d}{n} \left(\sum_{j=1}^W s_j \mu_j \right) \cdot \frac{(\gamma_\star + \sum_{j=1}^W s_j \mu_j)^2}{\gamma_\star^2} \cdot \log\left(\frac{\|x_1 - x_\star\|_2}{\epsilon}\right) \cdot \rho(\|x_\star\|_2), \quad (22)$$

we have

$$\|x_t - x_\star\|_2 \leq \epsilon + C_2 \frac{\text{Err}(x_\star, X)}{\bar{I}} \cdot \frac{(\gamma_\star + \sum_{j=1}^W s_j \mu_j)^2}{\gamma_\star^2} \cdot \rho(\|x_\star\|_2).$$

We provide the proof of Theorem 3 in Section 4.3. We now carefully unpack Theorem 3. Note that condition (21) necessitates that

$$n = \Omega(\bar{\omega}(x_\star, X)^2) \quad \text{and} \quad n = \Omega(\|x_\star\|_2^8 \log(\|x_\star\|_2 \vee 1)),$$

where in the notation $\Omega(\cdot)$ we hide dependence on the set of constants $\{s_j, \mu_j\}_{j=1}^W$. Thus, perfect recovery is possible as soon as the sample size n scales linearly in $\bar{\omega}(x_\star, X)^2$, provided it is also larger than $\|x_\star\|_2^8 \log(\|x_\star\|_2 \vee 1)$.

We next consider two regimes to elucidate the result, noting that the theorem applies more generally. In the first regime, suppose $n \asymp \bar{\omega}(x_\star, X)^2$. As we show in Appendix A.3, the scalar γ_\star , which is the solution of Eq. (18), satisfies

$$\gamma_\star \geq \sum_{j=1}^W s_j \mu_j \exp(-12\mu_j \|x_\star\|_2) \quad \text{if} \quad n \gtrsim \bar{\omega}(x_\star, X)^2. \quad (23)$$

As a consequence, Theorem 3 guarantees that for all $\epsilon \in (0, 1)$, after

$$t = \mathcal{O}\left(\frac{d+n}{n} \cdot \exp\left(24 \max_{j \in [W]} \mu_j \|x_\star\|_2\right) \log\left(\frac{\|x_1 - x_\star\|_2}{\epsilon}\right) \cdot (\|x_\star\|_2 \vee 1)^4\right)$$

iterations, we obtain

$$\|x_t - x_\star\|_2 \leq \epsilon + \frac{\text{Err}(x_\star, X)}{\bar{I}} \cdot \mathcal{O}\left(\exp\left(24 \max_{j \in [W]} \mu_j \|x_\star\|_2\right) \cdot (\|x_\star\|_2 \vee 1)^4\right).$$

While the sample complexity in this regime scales linearly in $\bar{\omega}(x_\star, X)^2$, this comes at a cost: The iteration complexity and the eventual estimation error increase exponentially in $\|x_\star\|_2$.

Next, we consider a regime in which the sample size increases polynomially in $\|x_\star\|_2$. In particular, we show in Appendix A.4 that

$$\gamma_\star \geq \sum_{j=1}^W s_j \mu_j \quad \text{if} \quad n \gtrsim \frac{\left(\sum_{j=1}^W s_j \mu_j\right)^2}{\sum_{j=1}^W s_j^2 \mu_j^2} \left(\max_{j \in [W]} (\mu_j \|x_\star\|_2)^4 \vee 1\right) \cdot \bar{\omega}(x_\star, X)^2. \quad (24)$$

In this regime, Theorem 3 guarantees that for each $\epsilon \in (0, 1)$, after

$$t = \mathcal{O}\left(\frac{d+n}{n} \cdot \log\left(\frac{\|x_1 - x_\star\|_2}{\epsilon}\right) (\|x_\star\|_2 \vee 1)^4\right)$$

iterations, we obtain

$$\|x_t - x_\star\|_2 \leq \epsilon + \frac{\text{Err}(x_\star, X)}{\bar{I}} \cdot \mathcal{O}\left((\|x_\star\|_2 \vee 1)^4\right).$$

Notably, this iteration complexity and the eventual estimation error only grows polynomially (and not exponentially) in $\|x_\star\|_2$. Thus, a slight increase in the sample size can lead to significant benefits in computational complexity.

Let us conclude with a comparison of Theorem 3 for EXACT with the gradient descent (GD) method developed in Fridovich-Keil et al. (2024) and the nonsmooth optimization methods (PolyakSGM and AdPolyakSGM) developed in Charisopoulos and Willett (2024) — we collect this comparison of both the iteration and sample complexity of all algorithms in Table 1. We note that a comparison can only be made in the noiseless, unstructured setting where $X = \mathbb{B}(4\|x_\star\|_2; x_\star)$ — in this case, we have $\bar{\omega}(x_\star, X)^2 \asymp d$ — and the monochromatic setting where $W = 1$ and $\mu_1 = 1$, since the two related works (Fridovich-Keil et al., 2024; Charisopoulos and Willett, 2024) only study this setting. From this table, it is clear that our method has the best sample and iteration complexity. In particular, our method improves both the iteration and sample complexity by a polynomial factor of $\|x_\star\|_2$.

3 Experimental results

We present simulations under both the practical CT model with a phantom image, and the Gaussian measurement model. Quantifying reconstruction quality is itself a nuanced task (Cheng et al., 2019); for simplicity we quantify image quality using Euclidean distance or root mean squared error (RMSE) relative to a known ground truth phantom, augmented with visual comparisons for qualitative evaluation.

Method	Sample complexity	Iteration complexity
EXACT	$\mathcal{O}(\ x_\star\ _2^4 \cdot d \vee \ x_\star\ _2^8 \log(\ x_\star\ _2 \vee 1))$	$\mathcal{O}\left(\ x_\star\ _2^4 \log\left(\frac{5\ x_\star\ _2}{\epsilon}\right)\right)$
	$\mathcal{O}(d \vee \ x_\star\ _2^8 \log(\ x_\star\ _2 \vee 1))$	$\mathcal{O}\left(\exp(24\ x_\star\ _2)\ x_\star\ _2^4 \log\left(\frac{5\ x_\star\ _2}{\epsilon}\right)\right)$
PolyakSGM AdPolyakSGM	$\mathcal{O}(\ x_\star\ _2^4 \cdot d)$	$\mathcal{O}\left(\ x_\star\ _2^6 \log\left(\frac{\ x_\star\ _2}{\epsilon}\right)\right)$
GD	$\mathcal{O}\left(\frac{\exp(c_1\ x_\star\ _2)}{\ x_\star\ _2} \cdot d\right)$	$\mathcal{O}\left(\exp(c_2\ x_\star\ _2) \log\left(\frac{\ x_\star\ _2}{\epsilon}\right)\right)$

Table 1. Iteration and sample complexity of iterative reconstruction methods with Gaussian measurements and unstructured signal, where we consider the noiseless, monochromatic ($W = 1$) case in order to make a comparison. The corresponding results for PolyakSGM and AdPolyakSGM were obtained by [Charisopoulos and Willett \(2024\)](#) and for GD by [Fridovich-Keil et al. \(2024\)](#). EXACT can achieve $\mathcal{O}(d)$ sample complexity without additional multiplicative factors in $\|x_\star\|_2$. It also has state of the art iteration complexity for a slightly larger sample complexity $\mathcal{O}(\|x_\star\|_2^4 d)$. All in all, the proposed method (10) improves the best-known results in the Gaussian monochromatic model along both axes of sample complexity and iteration complexity.

3.1 Photon-counting CT simulation

Our photon-counting CT simulation experiments are modeled on those from [Barber and Sidky \(2024\)](#). Our imaging target is a simple simulated PMMA phantom consisting of a disc with 4 circular regions of interest (ROIs), each with different density of PMMA. The target image has $d = 25^2 = 625$ pixels, the simulated X-ray photon-counting detector has 3 energy windows, and the simulated X-ray source outputs $W = 50$ different X-ray wavelengths. The default simulation setting is based on the setting of [Barber and Sidky \(2024\)](#), and uses an average source intensity of $\bar{I} = 10^6$ photons per detector cell and $n = (50 \text{ projection exposures}) \times (50 \text{ detector cells}) = 2500$ total measurements. The source spectrum, detector sensitivity, and target image are shown in [Figure 1](#).




Figure 1. Spectral density of the simulated X-ray source (left), detector sensitivity for 3 simulated windows (middle), and target phantom made of simulated PMMA with 4 regions of interest (right, colorbar in Hounsfield Units).

In each experiment, we compare our proposed extragradient algorithm EXACT (“Extragradient”) defined by Equation (10) with the square loss gradient descent approach (“MSE GD”) of Fridovich-Keil et al. (2024), the ADMM approach of Barber and Sidky (2024) and the Polyak subgradient descent (“PolyakSGM”) of Charisopoulos and Willett (2024), where we endow the latter with oracle knowledge of the loss incurred by the target image. Our simulated measurements are corrupted by Poisson noise, though only the ADMM baseline leverages specific knowledge of this noise model (by minimizing the Poisson negative log likelihood objective). For all algorithms, we constrain reconstructions to lie in the intersection of the positive orthant and the total variation (TV) ball; for clean comparison the TV constraint for all methods is derived from oracle knowledge of the target image TV. For all methods, at each iteration t we consider the average \bar{x}_t of the previous $t/2$ iterates⁴. We run all algorithms until convergence, defined as the first iteration at which $\|\bar{x}_t - \bar{x}_{t-1}\|_2 \leq 10^{-5}$. We quantify reconstruction quality using root mean square error (RMSE) of \bar{x}_T , where T is the iteration at which the algorithm converged. We tune the step size for EXACT, scaling it inversely with the source intensity according to Theorem 2. For MSE GD we tune the step size at the default setting, 50 projection exposures with source intensity 10^6 photons, and use the same step size for all experiments. For ADMM and PolyakSGM we use hyperparameters suggested in their respective papers ($\sigma = 100$ for ADMM and $\eta = 1$ for PolyakSGM), after verifying that these yield strong performance in the default setting.




Figure 2. RMSE (left) and runtime (right), varying the number of projection exposures while the X-ray source outputs on average 10^6 photons per detector cell. Until the number of projection exposures drops below 10, at which point all algorithms fail, our proposed Extragradient method EXACT (defined in Equation (10)) converges fastest and to lowest error.

We begin with a sufficiently high source intensity, on average 10^6 photons per detector cell, to mimic the setup of Barber and Sidky (2024). We then gradually reduce the number of simulated projection exposures to reduce X-ray dose. Results are reported in Figure 2, including variation over 10 random seeds controlling the Poisson noise. EXACT (“Extragradient,” Equation (10)) yields the lowest RMSE and fastest convergence time even down to 10 projection exposures, while all methods fail with 5 projection exposures. Visual comparisons using 10 projection exposures are shown in Figure 3.

Since EXACT produces a visually flawless reconstruction (up to image compression artifacts in our visualization) with 10^6 photons and 10 projection exposures (Figure 3), we next hold the

⁴In their analysis of ADMM, Barber and Sidky (2024) likewise consider the average iterate. Due to the convexity of the ℓ_2 -norm, note that our guarantees for the $t/2$ -th iterate of EXACT also apply to the average iterate at time t .




Figure 3. Full reconstructions (in Hounsfield Units, HU) and zoom-ins (bottom row) after convergence of each algorithm using 10 projection exposures and an average source intensity of 10^6 photons per detector cell, a total of $5\times$ reduction in X-ray dose compared to the 50-projection exposure default setting in which all algorithms perform well. The EXACT (“Extragradient”) and MSE GD reconstructions are visually faithful to the target in Figure 1, the ADMM reconstruction is perceptually accurate but exhibits less uniformity in the constant regions, and the PolyakSGM reconstruction exhibits more severe artifacts that obscure part of the lowest-contrast ROI.

number of projection exposures fixed at 10 and gradually decrease the intensity of the X-ray source. RMSE and runtime results are summarized in Figure 4, and reconstructions with 10^5 photons are shown in Figure 5. This setting corresponds to a $50\times$ reduction in X-ray dose compared to the default setting of 50 projection exposures and 10^6 photons in which all algorithms succeed.




Figure 4. RMSE (left) and runtime (middle, right), varying the source intensity while the number of projection exposures is fixed at 10, the smallest value for which recovery is possible in Figure 2. All algorithms suffer higher reconstruction error as source intensity and thus SNR is decreased. The rightmost plot is a zoom-in of the center plot to illustrate the runtimes of the three fastest algorithms.

3.2 Gaussian model

In this section, we evaluate the performance of EXACT (Algorithm (10)) under Gaussian measurements (see Assumption 3) and in the monochromatic setting where $W = 1$ and $\mu_1 = 1$. We consider noiseless observations, i.e., $y_i = \exp\{-\langle a_i, x_\star \rangle\}$, set $d = 100$, use step size $\gamma_t = 0.25$, and vary the number of samples $n \in \{5d, 10d, 20d, 40d, 80d\}$. We set $\|x_\star\|_2 = 3$.




Figure 5. Full reconstructions (in Hounsfield Units, HU) and zoom-ins (bottom row) after convergence of each algorithm using 10 projection exposures and an average source intensity of 10^5 photons per detector cell, a total of $50\times$ reduction in X-ray dose compared to the 50-projection exposure, 10^6 photons default setting in which all algorithms perform well. The EXACT (“Extragradient”) reconstruction exhibits minor artifacts at the edges of several ROIs, the MSE GD and ADMM reconstructions exhibit artifacts that partially obscure the lowest-contrast ROI, and the PolyakSGM reconstruction exhibits more severe artifacts that obscure several ROIs.




Figure 6. Estimation error over iterations for different sample size n . The experiment is conducted in the setting of Gaussian measurements, noiseless observations, and monochromatic model where $W = 1$, $\bar{I} = 1$ and $\mu_1 = 1$. In this simulation, we have $d = 100$ and $\|x_*\|_2 = 3$, and we set the step size $\gamma_t = 0.25$ in Eq. (10). As predicted theoretically, we observe that increasing the number of measurements can dramatically improve runtime, up to a threshold of diminishing returns.

From Figure 6, we observe that as the sample size increases, it takes fewer iterations to achieve machine precision. This finding aligns with the theoretical results presented in Theorem 3 and its associated comments, which state that a larger sample size can significantly reduce iteration complexity. Moreover, we also note that once the sample size surpasses a certain threshold, the improvement in iteration complexity from further increasing the sample size becomes less pronounced.

4 Proofs

In this section, we provide proofs of all our main results. Auxiliary technical lemmas can be found in the appendix.

4.1 Proof of Theorem 1

Recall that F is ν -strongly pseudo-monotone around the ground truth x_* , i.e.,

$$\langle F(x) - F(x_*), x - x_* \rangle \geq \nu \cdot \|x - x_*\|_2^2 \quad \text{for all } x \in X,$$

and also Lipschitz on the set X , i.e.,

$$\|F(x) - F(x')\|_2 \leq L \cdot \|x - x'\|_2 \quad \text{for all } x, x' \in X.$$

By the definition of substep (10b), we have

$$\begin{aligned} \|x_{t+1} - x_*\|_2^2 &= \|\mathcal{P}_X(x_t - \gamma_t F(x_{t+0.5})) - \mathcal{P}_X(x_*)\|_2^2 \\ &\stackrel{(i)}{\leq} \langle x_{t+1} - x_*, x_t - \gamma_t F(x_{t+0.5}) - x_* \rangle \\ &= \langle x_{t+1} - x_*, x_t - x_* \rangle - \gamma_t \langle F(x_{t+0.5}), x_{t+1} - x_* \rangle \\ &= \frac{1}{2} \|x_{t+1} - x_*\|_2^2 + \frac{1}{2} \|x_t - x_*\|_2^2 - \frac{1}{2} \|x_{t+1} - x_t\|_2^2 \\ &\quad - \gamma_t \langle F(x_{t+0.5}), x_{t+1} - x_{t+0.5} \rangle - \gamma_t \langle F(x_{t+0.5}), x_{t+0.5} - x_* \rangle \end{aligned} \quad (25)$$

where in step (i) we use 1-co-coercivity of the projection operator onto a convex set, whereby

$$\|\mathcal{P}_X(x) - \mathcal{P}_X(x')\|_2^2 \leq \langle \mathcal{P}_X(x) - \mathcal{P}_X(x'), x - x' \rangle \quad \text{for all } x, x' \in X.$$

Next, we bound the latter two inner product terms on the RHS. By strong monotonicity of F around x_* , we have $\langle F(x_{t+0.5}) - F(x_*), x_{t+0.5} - x_* \rangle \geq \nu \cdot \|x_{t+0.5} - x_*\|_2^2$, and rearranging terms yields

$$\begin{aligned} -\gamma_t \langle F(x_{t+0.5}), x_{t+0.5} - x_* \rangle &\leq -\gamma_t \langle F(x_*), x_{t+0.5} - x_* \rangle - \nu \gamma_t \|x_{t+0.5} - x_*\|_2^2 \\ &\leq \gamma_t \|x_{t+0.5} - x_*\|_2 \cdot \mathbf{Err}(x_*, X) - \frac{1}{2} \nu \gamma_t \|x_t - x_*\|_2^2 + \nu \gamma_t \|x_{t+0.5} - x_t\|_2^2, \end{aligned} \quad (26)$$

where in the last step we use $\zeta = F(x_*)$ and recall the definition of $\mathbf{Err}(x_*, X)$ (14). In addition, we also use the elementary inequality $\|x_t - x_*\|_2^2 \leq 2\|x_t - x_{t+0.5}\|_2^2 + 2\|x_{t+0.5} - x_*\|_2^2$.

We next turn to bound the inner product term $-\gamma_t \langle F(x_{t+0.5}), x_{t+1} - x_{t+0.5} \rangle$. By using the first-order optimality conditions for the projection step used in $x_{t+0.5} = \mathcal{P}_X(x_t - \gamma_t F(x_t))$, we obtain

$$\langle x_{t+0.5} - x_t + \gamma_t F(x_t), x - x_{t+0.5} \rangle \geq 0 \quad \text{for all } x \in X.$$

Substituting $x = x_{t+1}$ above and rearranging the terms yields

$$\begin{aligned} -\gamma_t \langle F(x_t), x_{t+1} - x_{t+0.5} \rangle &\leq \langle x_{t+0.5} - x_t, x_{t+1} - x_{t+0.5} \rangle \\ &= \frac{1}{2} \|x_t - x_{t+1}\|_2^2 - \frac{1}{2} \|x_{t+0.5} - x_t\|_2^2 - \frac{1}{2} \|x_{t+1} - x_{t+0.5}\|_2^2. \end{aligned} \quad (27)$$

Proceeding now to the term of interest and performing some algebraic manipulation, we have

$$\begin{aligned}
-\gamma_t \langle F(x_{t+0.5}), x_{t+1} - x_{t+0.5} \rangle &= -\gamma_t \langle F(x_{t+0.5}) - F(x_t), x_{t+1} - x_{t+0.5} \rangle - \gamma_t \langle F(x_t), x_{t+1} - x_{t+0.5} \rangle \\
&\stackrel{(i)}{\leq} \gamma_t \|F(x_{t+0.5}) - F(x_t)\|_2 \cdot \|x_{t+1} - x_{t+0.5}\|_2 - \gamma_t \langle F(x_t), x_{t+1} - x_{t+0.5} \rangle \\
&\stackrel{(ii)}{\leq} \gamma_t L \|x_{t+0.5} - x_t\|_2 \cdot \|x_{t+1} - x_{t+0.5}\|_2 - \gamma_t \langle F(x_t), x_{t+1} - x_{t+0.5} \rangle \\
&\stackrel{(iii)}{\leq} \frac{\gamma_t^2 L^2}{2} \|x_{t+0.5} - x_t\|_2^2 + \frac{1}{2} \|x_{t+1} - x_{t+0.5}\|_2^2 - \gamma_t \langle F(x_t), x_{t+1} - x_{t+0.5} \rangle \\
&\stackrel{(iv)}{\leq} \frac{\gamma_t^2 L^2}{2} \|x_{t+0.5} - x_t\|_2^2 + \frac{1}{2} \|x_t - x_{t+1}\|_2^2 - \frac{1}{2} \|x_{t+0.5} - x_t\|_2^2, \tag{28}
\end{aligned}$$

where step (i) follows from the Cauchy–Schwarz inequality, step (ii) from the fact that F is L -Lipschitz, step (iii) from the AM-GM inequality, and step (iv) from Eq. (27).

Substituting Ineqs. (28) and (26) into inequality (25), we obtain

$$\begin{aligned}
\|x_{t+1} - x_\star\|_2^2 &\leq (1 - \nu\gamma_t) \cdot \|x_t - x_\star\|_2^2 + (\gamma_t^2 L^2 + 2\nu\gamma_t - 1) \cdot \|x_{t+0.5} - x_t\|_2^2 \\
&\quad + 2\gamma_t \|x_{t+0.5} - x_\star\|_2 \cdot \text{Err}(x_\star, X).
\end{aligned}$$

To bound the last term on the RHS, we apply the inequality $2ab \leq a^2 + b^2$ with $a = \gamma_t \|x_{t+0.5} - x_\star\|_2 \cdot \sqrt{\frac{\nu L}{2}}$ and $b = \text{Err}(x_\star, X) \cdot \sqrt{\frac{2}{\nu L}}$ to obtain

$$\begin{aligned}
2\gamma_t \|x_{t+0.5} - x_\star\|_2 \cdot \text{Err}(x_\star, X) &\leq \frac{\gamma_t^2 \nu L}{2} \|x_{t+0.5} - x_\star\|_2^2 + \frac{2}{\nu L} \cdot \text{Err}(x_\star, X)^2 \\
&\leq \gamma_t^2 \nu L \left(\|x_{t+0.5} - x_t\|_2^2 + \|x_t - x_\star\|_2^2 \right) + \frac{2}{\nu L} \cdot \text{Err}(x_\star, X)^2.
\end{aligned}$$

Combining with the other two terms, we have

$$\begin{aligned}
\|x_{t+1} - x_\star\|_2^2 &\leq (1 - \nu\gamma_t + \nu L \gamma_t^2) \cdot \|x_t - x_\star\|_2^2 \\
&\quad + (\gamma_t^2 L^2 + 2\nu\gamma_t - 1 + \nu L \gamma_t^2) \cdot \|x_{t+0.5} - x_t\|_2^2 + \frac{2\text{Err}(x_\star, X)^2}{\nu L}.
\end{aligned}$$

By setting $\gamma_t = \frac{1}{4L}$, we obtain

$$1 - \nu\gamma_t + \nu L \gamma_t^2 = 1 - \frac{\nu}{4L} + \frac{\nu}{16L} \leq 1 - \frac{\nu}{8L},$$

and

$$\gamma_t^2 L^2 + 2\nu\gamma_t - 1 + \nu L \gamma_t^2 = \frac{1}{16} + \frac{\nu}{2L} - 1 + \frac{\nu}{16L} < 0,$$

where we use the fact that $\nu \leq L$. Putting the pieces together yields

$$\|x_{t+1} - x_\star\|_2^2 \leq \left(1 - \frac{\nu}{8L}\right) \cdot \|x_t - x_\star\|_2^2 + \frac{2\text{Err}(x_\star, X)^2}{\nu L}.$$

Unrolling the recursion over time thus yields

$$\|x_{t+1} - x_\star\|_2^2 \leq \left(1 - \frac{\nu}{8L}\right)^t \cdot \|x_1 - x_\star\|_2^2 + \frac{16\text{Err}(x_\star, X)^2}{\nu^2},$$

and taking square-roots of both sides and using the inequality $\sqrt{a+b} \leq \sqrt{a} + \sqrt{b}$ completes the proof. \square

4.2 Proof of Theorem 2

Our proof proceeds by first bounding the strong pseudo-monotonicity parameter ν and Lipschitzness parameter L , and then we apply Theorem 1.

Strong pseudo-monotonicity. By definition, the strong pseudo-monotonicity constant around x_* is bounded below by the strong monotonicity constant. For any pair $x, x' \in X$, we have

$$\langle F(x) - F(x'), x - x' \rangle = \bar{I} \cdot \frac{1}{n} \sum_{i=1}^n [h(\langle a_i, x' \rangle) - h(\langle a_i, x \rangle)] \cdot [\langle a_i, x \rangle - \langle a_i, x' \rangle].$$

By the mean value theorem, for any $x, x' \in X$, there exists some $\min\{\langle a_i, x \rangle, \langle a_i, x' \rangle\} \leq t_i \leq \max\{\langle a_i, x \rangle, \langle a_i, x' \rangle\}$ such that

$$h(\langle a_i, x' \rangle) - h(\langle a_i, x \rangle) = h'(t_i) \cdot [\langle a_i, x' \rangle - \langle a_i, x \rangle] = |h'(t_i)| \cdot [\langle a_i, x \rangle - \langle a_i, x' \rangle],$$

where $h'(t_i) \leq 0$ is the derivative of h at t_i (recall that h is nonincreasing). Consequently, we obtain

$$\begin{aligned} \langle F(x) - F(x'), x - x' \rangle &\geq \bar{I} \cdot \frac{1}{n} \sum_{i=1}^n |h'(t_i)| \cdot [\langle a_i, x' \rangle - \langle a_i, x \rangle]^2 \\ &\geq \frac{\bar{I}}{n} \min_{1 \leq i \leq n} |h'(t_i)| \cdot \sum_{i=1}^n [\langle a_i, x' \rangle - \langle a_i, x \rangle]^2 \\ &\geq \frac{\bar{I}}{n} \cdot \min_{1 \leq i \leq n} |h'(t_i)| \cdot \lambda_{\min}(\Sigma, X - X) \cdot \|x - x'\|_2^2, \end{aligned}$$

By noting that

$$|h'(t_i)| = \sum_{j=1}^W s_j \mu_j \exp(-\mu_j \cdot t_i) \quad \text{and} \quad 0 \leq t_i \leq \max_{\tilde{x} \in X} \langle a_i, \tilde{x} \rangle,$$

we obtain the bound

$$\min_{1 \leq i \leq n} |h'(t_i)| \geq \sum_{j=1}^W s_j \mu_j \cdot \exp\left(-\mu_j \max_{\tilde{x} \in X, 1 \leq i \leq n} \langle a_i, \tilde{x} \rangle\right).$$

Putting the pieces together, we obtain

$$\langle F(x) - F(x'), x - x' \rangle \geq \bar{I} \cdot \frac{\lambda_{\min}(\Sigma, X - X)}{n} \cdot \sum_{j=1}^W s_j \mu_j \cdot \exp\left(-\mu_j \max_{\tilde{x} \in X, 1 \leq i \leq n} \langle a_i, \tilde{x} \rangle\right) \cdot \|x - x'\|_2^2.$$

Since the above bound holds for any $x, x' \in X$, we conclude that

$$\nu \geq \bar{I} \cdot \frac{\lambda_{\min}(\Sigma, X - X)}{n} \cdot \sum_{j=1}^W s_j \mu_j \cdot \exp\left(-\mu_j \max_{\tilde{x} \in X, 1 \leq i \leq n} \langle a_i, \tilde{x} \rangle\right). \quad (29)$$

Lipschitz constant. Let us now show that F obeys the Lipschitz property, noting that Assumption 2 is not required for this claim. Define a vector $w \in \mathbb{R}^n$ with $w_i = h(\langle a_i, x \rangle) - h(\langle a_i, x' \rangle)$. We have

$$\begin{aligned} \|F(x) - F(x')\|_2^2 &= \frac{\bar{I}^2}{n^2} \left\| \sum_{i=1}^n (h(\langle a_i, x \rangle) - h(\langle a_i, x' \rangle)) a_i \right\|_2^2 \\ &= \frac{\bar{I}^2}{n^2} \|\mathbf{A}^\top w\|_2^2 \leq \frac{\bar{I}^2 \lambda_{\max}(\Sigma, \mathbb{R}^d)}{n^2} \cdot \|w\|_2^2 \end{aligned}$$

We may now write the squared norm of w as

$$\begin{aligned} \|w\|_2^2 &= \sum_{i=1}^n [h(\langle a_i, x \rangle) - h(\langle a_i, x' \rangle)]^2 \\ &= \sum_{i=1}^n h'(t_i)^2 \cdot [\langle a_i, x \rangle - \langle a_i, x' \rangle]^2 \\ &\stackrel{(i)}{\leq} \left(\sum_{j=1}^W s_j \mu_j \right)^2 \cdot \sum_{i=1}^n [\langle a_i, x \rangle - \langle a_i, x' \rangle]^2 \\ &\leq \left(\sum_{j=1}^W s_j \mu_j \right)^2 \cdot \lambda_{\max}(\Sigma, \mathbb{R}^d) \cdot \|x - x'\|_2^2 \end{aligned}$$

where in step (i) we use for all $t \in \mathbb{R}$,

$$|h'(t)| = \sum_{j=1}^W s_j \mu_j \exp\{-\mu_j(t)_+\} \cdot \mathbb{1}\{t > 0\} \leq \sum_{j=1}^W s_j \mu_j.$$

Putting the two pieces together yields for all $x, x' \in \mathbb{R}^d$,

$$\|F(x) - F(x')\|_2 \leq \bar{I} \cdot \frac{\lambda_{\max}(\Sigma, \mathbb{R}^d)}{n} \cdot \left(\sum_{j=1}^W s_j \mu_j \right) \cdot \|x - x'\|_2. \quad (30)$$

In particular, we have shown that F is Lipschitz with parameter $L = \bar{I} \cdot \frac{\lambda_{\max}(\Sigma, \mathbb{R}^d)}{n} \cdot \left(\sum_{j=1}^W s_j \mu_j \right)$.

Putting together the pieces. To conclude, we apply Theorem 1, using the lower bound on ν given by Eq. (29) and the the upper bound on L given by Eq. (30). \square

4.3 Proof of Theorem 3

As in the previous proof, the bulk of the argument relies on bounding the parameters (ν, L) but now in the Gaussian measurement model.

Proposition 1. *Let γ_\star be the solution of the fixed point equation (18) and let $\bar{\omega}(x_\star, X)$ be defined in Eq. (17). Suppose Assumption 1 holds with $R \leq 4\|x_\star\|_2$ and Assumption 3 holds. Then for all sample size n satisfying condition (21), there exists a universal positive constant c such that with probability at least $1 - 16 \exp(-\bar{\omega}(x_\star, X)^2/64) - 12n^{-10} - 2 \exp(-d)$, the following statements hold:*

(a) The Lipschitzness parameter L (12a) satisfies

$$L \leq 10\bar{I} \cdot \frac{d+n}{n} \cdot \sum_{j=1}^W s_j \mu_j.$$

(b) The strong pseudo-monotonicity parameter ν (12b) satisfies

$$\nu \geq c\bar{I} \cdot \left(\frac{\gamma_\star}{\gamma_\star + \sum_{j=1}^W s_j \mu_j} \right)^2 \cdot \left(\sum_{j=1}^W \frac{s_j \mu_j}{(\mu_j \|x_\star\|_2)^4 \vee 1} \right).$$

We provide the proof of Proposition 1 in Section 4.3.1. Applying Theorem 1 in conjunction with the lower bound on ν and the upper bound on L given by Proposition 1 completes the proof of Theorem 3. \square

4.3.1 Proof of Proposition 1(a)

Under Assumption 3, applying Vershynin (2018, Theorem 4.4.5) yields

$$\lambda_{\max}(\Sigma) = \|A\|_{\text{op}}^2 \leq 10(d+n), \quad \text{with probability } \geq 1 - 2e^{-d}.$$

Combining this inequality with inequality (30) yields the desired result.

4.3.2 Proof of Proposition 1(b)

Define the shorthand

$$\mathcal{P}(A) = \inf_{x \in X, x \neq x_\star} \frac{\frac{1}{n} \sum_{i=1}^n [h(a_i^\top x_\star) - h(a_i^\top x)] \cdot [a_i^\top x - a_i^\top x_\star]}{\|x - x_\star\|_2^2}, \quad (31)$$

and note that

$$\inf_{x \in X, x \neq x_\star} \frac{\langle F(x) - F(x_\star), x - x_\star \rangle}{\|x - x_\star\|_2^2} = \bar{I} \cdot \mathcal{P}(A).$$

To prove the claimed bound, it suffices to show that the following holds with probability at least $1 - 16 \exp(-\bar{\omega}(x_\star, X)^2/64) - 12n^{-10}$:

$$\mathcal{P}(A) \geq \nu.$$

We first reduce the high-dimensional infimum in Eq. (31) to a low-dimensional infimum by employing the Gaussian minmax theorem; we will use a variant of the result in Gordon (1985, 1988); Thrampoulidis et al. (2018). Let $g, \tilde{g} \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, I_d)$ be Gaussian random vectors chosen independently of the data $\{a_i\}_{i=1}^n$. We define the random quantity

$$\omega(x_\star, X) = \sup_{v \in X} \frac{\langle P_{x_\star}^\perp v, g \rangle}{\|P_{x_\star}^\perp v\|_2}, \quad (32)$$

whose expectation is equal to the previously defined scalar $\bar{\omega}(x_\star, X)$ (Eq. (17)). We also define the random variable

$$\mathcal{D}(Ax_\star, g, \tilde{g}) := \inf_{\substack{\alpha \in \mathbb{R}, \beta \geq 0, \\ 0 < (\alpha - \|x_\star\|_2)^2 + \beta^2 \leq (4\|x_\star\|_2)^2}} \frac{\sup_{\gamma \geq 0} \left[\frac{1}{n} \sum_{i=1}^n \mathcal{T}_i(\alpha, \beta, \gamma) - \frac{\gamma \beta^2}{7n} \cdot \omega(x_\star, X)^2 \right]}{(\alpha - \|x_\star\|_2)^2 + \beta^2}, \quad (33)$$

where for each $i \in [n]$, the random variable $\mathcal{T}_i(\alpha, \beta, \gamma)$ is defined as

$$\begin{aligned} \mathcal{T}_i(\alpha, \beta, \gamma) := & \frac{1}{7} \frac{\gamma \cdot |h'(6a_i^\top x_\star)|}{\gamma + |h'(6a_i^\top x_\star)|} \cdot \left(\left(\frac{\alpha}{\|x_\star\|_2} - 1 \right) a_i^\top x_\star + \beta \tilde{g}_i \right)^2 \\ & \cdot \mathbb{1} \left\{ -6a_i^\top x_\star \leq \beta \tilde{g}_i + \frac{\alpha}{\|x_\star\|_2} a_i^\top x_\star \leq 6a_i^\top x_\star, a_i^\top x_\star > 0 \right\}. \end{aligned} \quad (34)$$

The notation $\mathcal{D}(Ax_\star, g, \tilde{g})$ reflects that \mathcal{D} depends on the matrix A only through the one-dimensional projection Ax_\star . We next state an auxiliary lemma that relates quantity $\mathcal{P}(A)$ (31) and quantity $\mathcal{D}(Ax_\star, g, \tilde{g})$ (33).

Lemma 1. *Let $\mathcal{P}(A)$ and $\mathcal{D}(Ax_\star, g, \tilde{g})$ be defined as in Eq. (31) and Eq. (33), respectively. Then*

$$\Pr \{ \mathcal{P}(A) \leq \tau \} \leq 2 \Pr \{ \mathcal{D}(Ax_\star, g, \tilde{g}) \leq \tau \} \quad \text{for all } \tau \in \mathbb{R}.$$

We provide the proof of Lemma 1 in Appendix B.1. Finally, we claim that there exists a universal and positive constant c such that with probability at least $1 - 8 \exp(-\bar{\omega}(x_\star, X)^2/64) - 6n^{-10}$,

$$\mathcal{D}(Ax_\star, g, \tilde{g}) \geq c \left(\frac{\gamma_\star}{\gamma_\star + \sum_{j=1}^W s_j \mu_j} \right)^2 \cdot \left(\sum_{j=1}^W \frac{s_j \mu_j}{(\mu_j \|x_\star\|_2)^4 \vee 1} \right). \quad (35)$$

Combining this inequality with Lemma 1, we obtain that with probability at least $1 - 16 \exp(-\bar{\omega}(x_\star, X)^2/64) - 12n^{-10}$,

$$\mathcal{P}(A) \geq c \left(\frac{\gamma_\star}{\gamma_\star + \sum_{j=1}^W s_j \mu_j} \right)^2 \cdot \left(\sum_{j=1}^W \frac{s_j \mu_j}{(\mu_j \|x_\star\|_2)^4 \vee 1} \right).$$

This concludes the proof of strong pseudo-monotonicity. It remains to verify claim (35).

Proof of Claim (35). To reduce the notational burden, we let

$$\Gamma(\alpha, \beta) := \sup_{\gamma \geq 0} \left[\frac{1}{n} \sum_{i=1}^n \mathcal{T}_i(\alpha, \beta, \gamma) - \frac{\gamma \cdot \beta^2}{7n} \cdot \omega(x_\star, X)^2 \right]. \quad (36)$$

We claim that there exists a universal and positive constant c such that uniformly for all (α, β) satisfying $\beta \geq 0$ and $0 < (\alpha - \|x_\star\|_2)^2 + \beta^2 \leq (4\|x_\star\|_2)^2$, the following holds with probability at least $1 - 16 \exp(-\bar{\omega}(x_\star, X)^2/64) - 12n^{-10}$.

$$\text{If } \beta > 0, \text{ then } \frac{\Gamma(\alpha, \beta)}{(\alpha - \|x_\star\|_2)^2 + \beta^2} \geq c \left(\frac{\gamma_\star}{\gamma_\star + \sum_{j=1}^W s_j \mu_j} \right)^2 \left(\sum_{j=1}^W \frac{s_j \mu_j}{(\mu_j \|x_\star\|_2)^4 \vee 1} \right); \quad (37a)$$

$$\text{If } \beta = 0, \text{ then } \frac{\Gamma(\alpha, 0)}{(\alpha - \|x_\star\|_2)^2} \geq c \left(\sum_{j=1}^W \frac{s_j \mu_j}{(\mu_j \|x_\star\|_2)^4 \vee 1} \right). \quad (37b)$$

Note that Claim (35) immediately follows by the definition of $\mathcal{D}(Ax_\star, g, \tilde{g})$ in Eq. (33). We now turn to verify inequality (37a) and inequality (37b).

Verifying inequality (37a). We first focus on computing the inner supremum over $\gamma > 0$ in Eq. (33). For each $\alpha \in \mathbb{R}$ and $\beta > 0$, let $\bar{\gamma}(\alpha, \beta)$ be the solution of the inner $\sup_{\gamma \geq 0}$ in Eq. (33), i.e.,

$$\bar{\gamma}(\alpha, \beta) := \arg \sup_{\gamma \geq 0} \left[\frac{1}{n} \sum_{i=1}^n \mathcal{T}_i(\alpha, \beta, \gamma) - \frac{\gamma \cdot \beta^2}{7n} \cdot \omega(x_\star, X)^2 \right], \quad (38)$$

where by convention we choose the smallest element of the arg sup if the set is not a singleton. The following auxiliary lemma states some useful properties of $\bar{\gamma}(\alpha, \beta)$.

Lemma 2. *Let $\bar{\gamma}(\alpha, \beta)$ be defined in Eq. (38) and let γ_* be defined in Eq. (18). Let $\bar{\omega}(x_*, X)$ and $\omega(x_*, X)$ be defined in Eq. (17) and Eq. (32), respectively. There exists a universal and positive constant C such that if $n \geq C \cdot \bar{\omega}(x_*, X)$, then uniformly for all*

$$\alpha \in \mathbb{R}, \quad \beta > 0, \quad \text{and} \quad 0 < (\alpha - \|x_*\|_2)^2 + \beta^2 \leq (4\|x_*\|_2)^2, \quad (39)$$

we have with probability at least $1 - 8 \exp(-\bar{\omega}(x_, X)^2/64)$, the scalar $\bar{\gamma}(\alpha, \beta)$ is finite, and furthermore*

$$\bar{\gamma}(\alpha, \beta) \geq \gamma_* > 0. \quad (40)$$

We provide the proof of Lemma 2 in Appendix B.2.

Note that $\mathcal{T}_i(\alpha, \beta, \gamma)$ defined in Eq. (34) is a concave function of γ for each $i \in [n]$. Then we have either $\bar{\gamma}(\alpha, \beta) = 0$ or $\bar{\gamma}(\alpha, \beta)$ must satisfy the first-order optimality condition, i.e.,

$$\frac{1}{n} \sum_{i=1}^n \frac{\partial \mathcal{T}_i(\alpha, \beta, \gamma)}{\partial \gamma} - \frac{\beta^2}{7n} \cdot \omega(x_*, X)^2 = 0.$$

But on the event guaranteed by Lemma 2, we have $\bar{\gamma}(\alpha, \beta) \geq \gamma_* > 0$ with probability at least $1 - 8 \exp(-\bar{\omega}(x_*, X)^2/64)$. Thus, the first-order optimality condition must hold, and evaluating the partial derivative $\partial \mathcal{T}_i(\alpha, \beta, \gamma)/\partial \gamma$, this is equivalent to the condition

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n \frac{|h'(6a_i^\top x_*)|^2 \cdot ((\alpha/\|x_*\|_2 - 1)a_i^\top x_* + \beta \tilde{g}_i)^2}{(\bar{\gamma}(\alpha, \beta) + |h'(6a_i^\top x_*)|)^2} \cdot \mathbb{1} \left\{ -6a_i^\top x_* \leq \beta \tilde{g}_i + \frac{\alpha}{\|x_*\|_2} a_i^\top x_* \leq 6a_i^\top x_*, a_i^\top x_* > 0 \right\} \\ = \frac{\beta^2 \omega(x_*, X)^2}{n}. \end{aligned} \quad (41)$$

Thus, recalling the definition of $\Gamma(\alpha, \beta)$ in Eq. (36), we have

$$\begin{aligned} \Gamma(\alpha, \beta) &= \frac{1}{n} \sum_{i=1}^n \mathcal{T}_i(\alpha, \beta, \bar{\gamma}(\alpha, \beta)) - \frac{\bar{\gamma}(\alpha, \beta) \cdot \beta^2}{7n} \cdot \omega(x_*, X)^2 \\ &\stackrel{(i)}{=} \frac{1}{7n} \sum_{i=1}^n \left(\frac{\bar{\gamma}(\alpha, \beta)}{\bar{\gamma}(\alpha, \beta) + |h'(6a_i^\top x_*)|} \right)^2 \cdot |h'(6a_i^\top x_*)| \cdot [(\alpha/\|x_*\|_2 - 1)a_i^\top x_* + \beta \tilde{g}_i]^2 \\ &\quad \cdot \mathbb{1} \left\{ -6a_i^\top x_* \leq \beta \tilde{g}_i + \frac{\alpha}{\|x_*\|_2} a_i^\top x_* \leq 6a_i^\top x_*, a_i^\top x_* > 0 \right\}. \end{aligned} \quad (42)$$

where in step (i), we have used the definition (34) of \mathcal{T}_i and the first-order optimality condition (41).

Next, note that since $\bar{\gamma}(\alpha, \beta) \geq \gamma_*$ and $|h'(t)| \leq \sum_{j=1}^W s_j \mu_j$ for all $t > 0$, we have

$$\frac{\bar{\gamma}(\alpha, \beta)}{\bar{\gamma}(\alpha, \beta) + |h'(6a_i^\top x_*)|} \geq \frac{\gamma_*}{\gamma_* + \sum_{j=1}^W s_j \mu_j} \quad \text{for} \quad a_i^\top x_* > 0. \quad (43)$$

Also note that for (α, β) satisfying condition (39), we have

$$\begin{aligned} \mathbb{1} \left\{ -6a_i^\top x_* \leq \beta \tilde{g}_i + \frac{\alpha}{\|x_*\|_2} a_i^\top x_* \leq 6a_i^\top x_*, a_i^\top x_* > 0 \right\} &\geq \mathbb{1} \left\{ -a_i^\top x_* \leq \beta \tilde{g}_i \leq a_i^\top x_*, a_i^\top x_* > 0 \right\} \\ &\geq \mathbb{1} \left\{ -\frac{a_i^\top x_*}{4\|x_*\|_2} \leq \tilde{g}_i \leq \frac{a_i^\top x_*}{4\|x_*\|_2}, a_i^\top x_* > 0 \right\}. \end{aligned} \quad (44)$$

Substituting the inequality in the display above and inequality (43) into inequality (42), we obtain that with probability at least $1 - 8 \exp(-\bar{\omega}(x_*, X)^2/64)$ the following holds for all (α, β) satisfying condition (39):

$$\begin{aligned} \Gamma(\alpha, \beta) &\geq \left(\frac{\gamma_\star}{\gamma_\star + \sum_{j=1}^W s_j \mu_j} \right)^2 \times \\ &\quad \frac{1}{7n} \sum_{i=1}^n |h'(6a_i^\top x_\star)| \cdot [(\alpha/\|x_\star\|_2 - 1)a_i^\top x_\star + \beta \tilde{g}_i]^2 \cdot \mathbb{1} \left\{ -\frac{a_i^\top x_\star}{4\|x_\star\|_2} \leq \tilde{g}_i \leq \frac{a_i^\top x_\star}{4\|x_\star\|_2}, a_i^\top x_\star > 0 \right\} \\ &=: \left(\frac{\gamma_\star}{\gamma_\star + \sum_{j=1}^W s_j \mu_j} \right)^2 \cdot \left((\alpha - \|x_\star\|_2)^2 \cdot \frac{1}{7n} \sum_{i=1}^n \theta_i + \beta^2 \cdot \frac{1}{7n} \sum_{i=1}^n \vartheta_i + 2(\alpha - \|x_\star\|_2)\beta \cdot \frac{1}{7n} \sum_{i=1}^n \xi_i \right), \end{aligned} \quad (45)$$

where in the last step we have expanded the square and used the following shorthand:

$$\theta_i = |h'(6a_i^\top x_\star)| \left(\frac{a_i^\top x_\star}{\|x_\star\|_2} \right)^2 \cdot \mathbb{1} \left\{ -\frac{a_i^\top x_\star}{4\|x_\star\|_2} \leq \tilde{g}_i \leq \frac{a_i^\top x_\star}{4\|x_\star\|_2}, a_i^\top x_\star > 0 \right\}, \quad (46a)$$

$$\vartheta_i = |h'(6a_i^\top x_\star)| \tilde{g}_i^2 \cdot \mathbb{1} \left\{ -\frac{a_i^\top x_\star}{4\|x_\star\|_2} \leq \tilde{g}_i \leq \frac{a_i^\top x_\star}{4\|x_\star\|_2}, a_i^\top x_\star > 0 \right\}, \quad (46b)$$

$$\xi_i = |h'(6a_i^\top x_\star)| \frac{a_i^\top x_\star}{\|x_\star\|_2} \tilde{g}_i \cdot \mathbb{1} \left\{ -\frac{a_i^\top x_\star}{4\|x_\star\|_2} \leq \tilde{g}_i \leq \frac{a_i^\top x_\star}{4\|x_\star\|_2}, a_i^\top x_\star > 0 \right\}. \quad (46c)$$

Note that all random variables defined in Eq. (46) are independent of the tuple (α, β, γ) . The following auxiliary lemma controls these random variables with high probability.

Lemma 3. *Let $\{\theta_i, \vartheta_i, \xi_i\}_{i=1}^n$ be defined in Eq. (46). Suppose the sample size n satisfies condition (21). Then there exists a pair of universal and positive constants (c_1, C_1) such that with probability at least $1 - 6n^{-10}$, we have*

$$\frac{1}{n} \sum_{i=1}^n \theta_i \wedge \frac{1}{n} \sum_{i=1}^n \vartheta_i \geq c_1 \sum_{j=1}^W \frac{s_j \mu_j}{(\mu_j \|x_\star\|_2)^4 \vee 1} \quad \text{and} \quad \left| \frac{1}{n} \sum_{i=1}^n \xi_i \right| \leq C_1 \sum_{j=1}^W s_j \mu_j \sqrt{\frac{\log(n)}{n}}.$$

We provide the proof of Lemma 3 in Appendix B.3. Substituting the bound of Lemma 3 into inequality (45) yields that with probability at least $1 - 8 \exp(-\bar{\omega}(x_\star, X)^2/64) - 6n^{-10}$,

$$\begin{aligned} \Gamma(\alpha, \beta) &\geq \frac{1}{7} \left(\frac{\gamma_\star}{\gamma_\star + \sum_{j=1}^W s_j \mu_j} \right)^2 \times \\ &\quad \left([(\alpha - \|x_\star\|_2)^2 + \beta^2] \cdot c_1 \sum_{j=1}^W \frac{s_j \mu_j}{(\mu_j \|x_\star\|_2)^4 \vee 1} - 2C_1 |\alpha - \|x_\star\|_2| \beta \sum_{j=1}^W s_j \mu_j \sqrt{\frac{\log(n)}{n}} \right) \\ &\stackrel{(i)}{\geq} \frac{1}{7} \left(\frac{\gamma_\star}{\gamma_\star + \sum_{j=1}^W s_j \mu_j} \right)^2 \left(\frac{c_1}{2} \sum_{j=1}^W \frac{s_j \mu_j}{(\mu_j \|x_\star\|_2)^4 \vee 1} \right) \left((\alpha - \|x_\star\|_2)^2 + \beta^2 \right), \end{aligned} \quad (47)$$

where in step (i) we use the AM-GM inequality and condition (21) so that

$$2C_1 |\alpha - \|x_\star\|_2| \beta \sum_{j=1}^W s_j \mu_j \sqrt{\frac{\log(n)}{n}} \leq \frac{c_1}{2} [(\alpha - \|x_\star\|_2)^2 + \beta^2] \cdot \sum_{j=1}^W \frac{s_j \mu_j}{(\mu_j \|x_\star\|_2)^4 \vee 1},$$

This concludes the proof of inequality (37a)

Verifying inequality (37b). By the definitions of $\Gamma(\alpha, \beta)$ (36) and $\mathcal{T}_i(\alpha, \beta, \gamma)$ (34), we obtain

$$\begin{aligned} \Gamma(\alpha, 0) &= \sup_{\gamma \geq 0} \left[\frac{1}{7n} \sum_{i=1}^n \frac{\gamma |h'(6a_i^\top x_\star)|}{\gamma + |h'(6a_i^\top x_\star)|} \frac{(a_i^\top x_\star)^2}{\|x_\star\|_2^2} \cdot (\alpha - \|x_\star\|_2)^2 \cdot \mathbb{1} \left\{ -6\|x_\star\|_2 \leq \alpha \leq 6\|x_\star\|_2, a_i^\top x_\star > 0 \right\} \right] \\ &\stackrel{(i)}{=} \frac{1}{7n} \sum_{i=1}^n |h'(6a_i^\top x_\star)| \frac{(a_i^\top x_\star)^2}{\|x_\star\|_2^2} \cdot (\alpha - \|x_\star\|_2)^2 \cdot \mathbb{1} \left\{ a_i^\top x_\star > 0 \right\} \\ &\stackrel{(ii)}{\geq} \frac{1}{7n} \sum_{i=1}^n \theta_i \cdot (\alpha - \|x_\star\|_2)^2, \end{aligned}$$

where in step (i), the supremum is achieved when $\gamma \uparrow \infty$, since the condition $-6\|x_\star\|_2 \leq \alpha \leq 6\|x_\star\|_2$ always holds due to the assumption $(\alpha - \|x_\star\|_2)^2 \leq (4\|x_\star\|_2)^2$. Step (ii) follows from definition of θ_i in Eq. (46). To conclude, we use the high-probability lower bound on $\frac{1}{n} \sum_{i=1}^n \theta_i$ guaranteed by Lemma 3 to obtain the desired inequality (37b). \square

5 Discussion

We presented the first provable and efficient algorithm for single-material reconstruction from spectral CT measurements, which was based on carefully constructing a variational inequality whose fixed point approximated the desired signal. We presented explicit bounds on the sample and iteration complexity of our EXACT algorithm both for general measurements and Gaussian measurements, handling noise in the measurements and structure in the signal. Several downstream questions are interesting from both the theoretical and methodological points of view — we point out a few such questions below.

Let us start with some interesting theoretical questions. First, understanding the fundamental (i.e. information-theoretic) limits of signal recovery in this problem — say under the Poisson noise model (5) — is an important direction, since it would provide guidance on whether the performance of our estimator can be improved. A second family of theoretical questions for the algorithm when run on Gaussian data is to characterize sharp rates of convergence and sample complexity (i.e. with exact constant factors). Note that our algorithm itself is a *general first-order method* in the nomenclature of Celentano et al. (2020), so we may be able to prove a so-called “state evolution” that characterizes its iterate-by-iterate performance as well as its eventual error. On a complementary note, information-theoretic lower bounds for the problem should be derivable via the techniques in Barbier et al. (2019). When does the eventual error of our algorithm exactly match the information-theoretic lower bound in the Gaussian model? To our knowledge, in spite of considerable research on monotone single-index models with Gaussian data (e.g., Balabdaoui et al., 2019; Pananjady and Foster, 2021) and although there has been some recent interest in proving state evolutions for algorithms that go beyond first-order methods (see, e.g., Chandrasekher et al., 2023, 2024; Lou et al., 2024; Kaushik et al., 2024), such questions have not received attention for algorithms that find fixed-points of (strongly) monotone VIs.

From a methodological viewpoint also, there are several interesting open questions. The first and most important open question is how to design a variant of our methodology for the multi-material recovery problem in Eq. (1). Note that in that case, the natural analog of the operator F that we use is no longer monotone, so we would need to take a different approach to designing a provably convergent algorithm. Our analysis also opens up interesting measurement design questions, such as how to optimally allocate a limited photon budget by trading off source intensity \bar{I} and number

of measurements n to yield the highest reconstruction accuracy. Future analysis may also consider detector non-idealities such as *pulse pileup* and *charge sharing*, in which some incident photons are not recorded or are recorded as multiple photons with incorrect energies. For example, reducing the source intensity \bar{I} increases shot noise but also mitigates pulse pileup by reducing the likelihood that a photon arrives during the sensor’s dead time. Incorporating these effects into algorithm design and analysis may inform refinements to the reconstruction process and measurement design setup.

Acknowledgments

ML, KAV and AP were supported in part by the National Science Foundation through grants CCF-2107455 and DMS-2210734, and by research awards from Adobe, Amazon, and Mathworks. SFK was supported in part by the NSF Mathematical Sciences Postdoctoral Research Fellowship under award number 2303178. We thank Amir Pourmorteza for helpful discussions.

References

- R. E. Alvarez and A. Macovski. Energy-selective reconstructions in X-ray computerised tomography. *Physics in Medicine & Biology*, 21(5):733, 1976.
- F. Balabdaoui, C. Durot, and H. Jankowski. Least squares estimation in the monotone single index model. *Bernoulli*, 25(4B):3276–3310, 2019.
- R. F. Barber and E. Y. Sidky. Convergence for nonconvex ADMM, with applications to CT imaging. *Journal of Machine Learning Research*, 25(38):1–46, 2024.
- R. F. Barber, E. Y. Sidky, T. G. Schmidt, and X. Pan. An algorithm for constrained one-step inversion of spectral CT data. *Physics in Medicine & Biology*, 61(10):3784, 2016.
- J. Barbier, F. Krzakala, N. Macris, L. Miolane, and L. Zdeborová. Optimal errors and phase transitions in high-dimensional generalized linear models. *Proceedings of the National Academy of Sciences*, 116(12):5451–5460, 2019.
- C. Cai, T. Rodet, S. Legoupil, and A. Mohammad-Djafari. A full-spectral Bayesian reconstruction approach based on the material decomposition model applied in dual-energy computed tomography. *Medical physics*, 40(11):111916, 2013.
- M. Celentano, A. Montanari, and Y. Wu. The estimation error of general first order methods. In *Conference on Learning Theory*, pages 1078–1141. PMLR, 2020.
- C. Chandel, P. K. Srivastava, and P. Mahajan. Micromechanical analysis of deformation of snow using X-ray tomography. *Cold Regions Science and Technology*, 101:14–23, 2014.
- K. A. Chandrasekher, A. Pananjady, and C. Thrampoulidis. Sharp global convergence guarantees for iterative nonconvex optimization with random data. *The Annals of Statistics*, 51(1):179–210, 2023.
- K. A. Chandrasekher, M. Lou, and A. Pananjady. Alternating minimization for generalized rank-1 matrix sensing: Sharp predictions from a random initialization. *Information and Inference: A Journal of the IMA*, 13(3):iaae025, 2024.

- V. Charisopoulos and R. Willett. Nonlinear tomographic reconstruction via nonsmooth optimization. *arXiv preprint arXiv:2407.12984*, 2024.
- Y. Cheng, E. Abadi, T. B. Smith, F. Ria, M. Meyer, D. Marin, and E. Samei. Validation of algorithmic CT image quality metrics with preferences of radiologists. *Medical Physics*, 46(11):4837–4846, 2019.
- M. Danielsson, M. Persson, and M. Sjölin. Photon-counting X-ray detectors for CT. *Physics in Medicine & Biology*, 66(3):03TR01, 2021.
- I. A. Elbakri and J. A. Fessler. Statistical image reconstruction for polyenergetic X-ray computed tomography. *IEEE Transactions on Medical Imaging*, 21(2):89–99, 2002.
- S. Fridovich-Keil, F. Valdivia, G. Wetzstein, B. Recht, and M. Soltanolkotabi. Gradient descent provably solves nonlinear tomographic reconstruction. *arXiv preprint arXiv:2310.03956*, 2024.
- L. Fu, T.-C. Lee, S. M. Kim, A. M. Alessio, P. E. Kinahan, Z. Chang, K. Sauer, M. K. Kalra, and B. De Man. Comparison between pre-log and post-log statistical models in ultra-low-dose CT reconstruction. *IEEE Transactions on Medical Imaging*, 36(3):707–720, 2016.
- S. I. Gallant. Perceptron-based learning algorithms. *IEEE Transactions on Neural Networks*, 1(2):179–191, 1990.
- L. L. Geyer, U. J. Schoepf, F. G. Meinel, J. W. Nance Jr, G. Bastarrika, J. A. Leipsic, N. S. Paul, M. Rengo, A. Laghi, and C. N. De Cecco. State of the art: Iterative CT reconstruction techniques. *Radiology*, 276(2):339–357, 2015.
- L. Gjestebj, B. De Man, Y. Jin, H. Paganetti, J. Verburg, D. Giantsoudi, and G. Wang. Metal artifact reduction in CT: Where are we after four decades? *IEEE Access*, 4:5826–5849, 2016.
- Y. Gordon. Some inequalities for Gaussian processes and applications. *Israel Journal of Mathematics*, 50:265–289, 1985.
- Y. Gordon. On Milman’s inequality and random subspaces which escape through a mesh in \mathbb{R}^n . In *Geometric Aspects of Functional Analysis: Israel Seminar (GAFA) 1986–87*, pages 84–106. Springer, 1988.
- F. Götze, H. Sambale, and A. Sinulis. Concentration inequalities for polynomials in α -sub-exponential random variables. *Electronic Journal of Probability*, 26:1–22, 2021.
- N. M. Jadue, M. Nagare, J. S. Maltz, G. T. Buzzard, and C. A. Bouman. MACE CT reconstruction for modular material decomposition from energy resolving photon-counting data. In *Medical Imaging 2024: Physics of Medical Imaging*, volume 12925, pages 11–22. SPIE, 2024.
- A. Juditsky and A. Nemirovski. *Statistical inference via convex optimization*. Princeton University Press, 2020.
- A. B. Juditsky and A. S. Nemirovski. Signal recovery by stochastic optimization. *Automation and Remote Control*, 80(10):1878–1893, 2019.
- S. M. Kakade, V. Kanade, O. Shamir, and A. Kalai. Efficient learning of generalized linear and single index models with isotonic regression. *Advances in Neural Information Processing Systems*, 24, 2011.

- A. T. Kalai and R. Sastry. The Isotron algorithm: High-dimensional isotonic regression. In *COLT*, volume 1, page 9, 2009.
- C. Kaushik, J. Romberg, and V. Muthukumar. Precise asymptotics of reweighted least-squares algorithms for linear diagonal networks. *arXiv preprint arXiv:2406.02769*, 2024.
- D. Kinderlehrer and G. Stampacchia. *An introduction to variational inequalities and their applications*. SIAM, 2000.
- G. M. Korpelevich. The extragradient method for finding saddle points and other problems. *Math. USSR - Izv.*, 12:747–756, 1976.
- Y. Long and J. A. Fessler. Multi-material decomposition using statistical image reconstruction for spectral CT. *IEEE transactions on medical imaging*, 33(8):1614–1626, 2014.
- M. Lou, K. A. Verchand, and A. Pananjady. Hyperparameter tuning via trajectory predictions: Stochastic prox-linear methods in matrix sensing. *arXiv preprint arXiv:2402.01599*, 2024.
- C. Maaß, M. Baer, and M. Kachelrieß. Image-based dual energy CT using optimized precorrection functions: A practical new approach of material decomposition in image domain. *Medical physics*, 36(8):3818–3829, 2009.
- K. Mechlem, S. Ehn, T. SELLERER, E. Braig, D. Münzel, F. Pfeiffer, and P. B. Noël. Joint statistical iterative material image reconstruction for spectral computed tomography using a semi-empirical forward model. *IEEE transactions on medical imaging*, 37(1):68–80, 2017.
- C. Mory, B. Sixou, S. Si-Mohamed, L. Boussel, and S. Rit. Comparison of five one-step reconstruction algorithms for spectral CT. *Physics in Medicine & Biology*, 63(23):235001, 2018.
- Y. Nakamura, T. Higaki, S. Kondo, I. Kawashita, I. Takahashi, and K. Awai. An introduction to photon-counting detector CT (PCD CT) for radiologists. *Japanese Journal of Radiology*, 41(3):266–282, 2023.
- T. Niu, X. Dong, M. Petrongolo, and L. Zhu. Iterative image-domain decomposition for dual-energy CT. *Medical physics*, 41(4):041901, 2014.
- S. Osher, M. Burger, D. Goldfarb, J. Xu, and W. Yin. An iterative regularization method for total variation-based image restoration. *Multiscale Modeling & Simulation*, 4(2):460–489, 2005.
- A. Pananjady and D. P. Foster. Single-index models in the high signal regime. *IEEE Trans. Inform. Theory*, 67(6):4092–4124, 2021.
- M. Persson, A. Wang, and N. J. Pelc. Detective quantum efficiency of photon-counting CdTe and Si detectors for computed tomography: A simulation study. *Journal of Medical Imaging*, 7(4):043501–043501, 2020.
- A. Pourmorteza, H. Dang, J. Siewerdsen, and J. Stayman. Reconstruction of difference in sequential CT studies using penalized likelihood estimation. *Physics in Medicine & Biology*, 61(5):1986, 2016.
- J. Radon. On the determination of functions from their integral values along certain manifolds. *IEEE Transactions on Medical Imaging*, 5(4):170–176, 1986. doi: 10.1109/TMI.1986.4307775.

- B. M. Rizzo, E. Y. Sidky, and T. G. Schmidt. Material decomposition from unregistered dual kV data using the cOSSCIR algorithm. In *7th International Conference on Image Formation in X-Ray Computed Tomography*, volume 12304, pages 539–544. SPIE, 2022.
- R. T. Rockafellar and R. J.-B. Wets. *Variational analysis*, volume 317. Springer Science & Business Media, 2009.
- A. Sawatzky, Q. Xu, C. O. Schirra, and M. A. Anastasio. Proximal ADMM for multi-channel image reconstruction in spectral X-ray CT. *IEEE Transactions on Medical Imaging*, 33(8):1657–1668, 2014.
- C. O. Schirra, E. Roessl, T. Koehler, B. Brendel, A. Thran, D. Pan, M. A. Anastasio, and R. Proksa. Statistical reconstruction of material decomposed data in spectral CT. *IEEE Transactions on Medical Imaging*, 32(7):1249–1257, 2013.
- T. G. Schmidt, R. F. Barber, and E. Y. Sidky. A spectral CT method to directly estimate basis material maps from experimental photon-counting data. *IEEE Transactions on Medical Imaging*, 36(9):1808–1819, 2017.
- T. G. Schmidt, B. A. Sammut, R. F. Barber, X. Pan, and E. Y. Sidky. Addressing CT metal artifacts using photon-counting detectors and one-step spectral CT image reconstruction. *Medical Physics*, 49(5):3021–3040, 2022.
- T. G. Schmidt, E. Y. Sidky, X. Pan, R. F. Barber, F. Grönberg, M. Sjölin, and M. Danielsson. Constrained one-step material decomposition reconstruction of head CT data from a silicon photon-counting prototype. *Medical Physics*, 50(10):6008–6021, 2023.
- E. Y. Sidky, R. F. Barber, T. Gilat-Schmidt, and X. Pan. Three material decomposition for spectral computed tomography enabled by block-diagonal step-preconditioning. *arXiv preprint arXiv:1801.06263*, 2018.
- J. M. Sietins, J. C. Sun, and D. B. K. Jr. Fiber orientation quantification utilizing X-ray micro-computed tomography. *Journal of Composite Materials*, 55(8):1109–1118, 2021.
- D. Strong and T. Chan. Edge-preserving and scale-dependent properties of total variation regularization. *Inverse Problems*, 19(6):S165, 2003.
- P. Sukovic and N. H. Clinthorne. Design of an experimental system for dual energy X-ray CT. In *1999 IEEE Nuclear Science Symposium. Conference Record. 1999 Nuclear Science Symposium and Medical Imaging Conference (Cat. No. 99CH37019)*, volume 2, pages 1021–1022. IEEE, 1999.
- K. Taguchi and J. S. Iwanczyk. Vision 20/20: Single photon counting X-ray detectors in medical imaging. *Medical Physics*, 40(10):100901, 2013.
- X. Tang. *Spectral Multi-detector Computed Tomography (sMDCT): Data Acquisition, Image Formation, Quality Assessment and Contrast Enhancement*. CRC Press, 2023.
- C. Thrampoulidis, E. Abbasi, and B. Hassibi. Precise error analysis of regularized M -estimators in high dimensions. *IEEE Trans. Inform. Theory*, 64(8):5592–5628, 2018.
- R. Vershynin. *High-dimensional Probability: An Introduction with Applications in Data Science*, volume 47. Cambridge University Press, 2018.

M. J. Wainwright. *High-Dimensional Statistics: A Non-Asymptotic Viewpoint*. Cambridge University Press, 2019.

A. S. Wang and N. J. Pelc. A comparison of dual kV energy integrating and energy discriminating photon counting detectors for dual energy X-ray imaging. In *Medical Imaging 2012: Physics of Medical Imaging*, volume 8313, pages 279–291. SPIE, 2012.

T. Weidinger, T. M. Buzug, T. Flohr, S. Kappler, and K. Stierstorfer. Polychromatic iterative statistical material image reconstruction for photon-counting computed tomography. *International journal of biomedical imaging*, 2016(1):5871604, 2016.

Appendix

A Proof of statements in the main text

We first present calculations for several statements that were made in the main text.

A.1 Proof of claim in Example 2

Since $X = \mathbb{R}^d$, we obtain

$$\sup_{x, x' \in X} \left\langle \zeta, \frac{x - x'}{\|x - x'\|_2} \right\rangle = \|\zeta\|_2.$$

For convenience, define the noise variables

$$R_i = y_i - \bar{I} \cdot h(\langle a_i, x_\star \rangle) \quad \text{for } i = 1, 2, \dots, n,$$

which are all independent and zero-mean, and drawn according to model (5).

To proceed, note that

$$\|\zeta\|_2^2 = \frac{1}{n^2} \left\| A^\top R \right\|_2^2 = \frac{1}{n^2} R^\top A A^\top R,$$

and that

$$\mathbb{E} \left[R^\top A A^\top R \right] = \sum_{i=1}^n \|a_i\|_2^2 \mathbb{E}[R_i^2] = \bar{I} \sum_{i=1}^n \|a_i\|_2^2 \cdot h(\langle a_i, x_\star \rangle).$$

Next, note that $\{R_i\}_{i=1}^n$ are i.i.d. zero-mean random variables according to model (5). Moreover, $\{R_i\}_{i=1}^n$ are sub-exponential random variables, with bounded Orlicz ψ_1 -norm (Vershynin, 2018). Indeed, we have for all $0 < t \leq 1$,

$$\begin{aligned} \mathbb{E}[\exp(t|R_i|)] &\leq \mathbb{E}[\exp(t|y_i|)] \cdot \exp(t\bar{I}h(\langle a_i, x_\star \rangle)) \\ &\leq \mathbb{E}[\exp(t|y_i|)] \cdot \exp(t\bar{I}h(\langle a_i, x_\star \rangle)) \\ &= \exp(\bar{I}h(\langle a_i, x_\star \rangle)(e^t - 1)) \cdot \exp(t\bar{I}h(\langle a_i, x_\star \rangle)) \\ &\stackrel{(i)}{\leq} \exp(\bar{I}(e^t - 1 + t)) \\ &\leq \exp(\bar{I}(e + 1)t), \end{aligned}$$

where in step (i) we use $h(t) \leq 1$ for all $t \in \mathbb{R}$ and in the last step we use $e^t \leq 1 + e \cdot t$ for $t \in (0, 1)$. Substituting $t = (10(e + 1)\bar{I})^{-1}$, we have $\mathbb{E} \left[\exp \left(\frac{|R_i|}{10(e+1)\bar{I}} \right) \right] \leq e^{1/10} \leq 2$, thereby proving that $\|R_i\|_{\psi_1} \lesssim \bar{I}$.

We now apply the Hanson-Wright inequality for 1-sub-exponential random variables (Götze et al., 2021, Proposition 1.1), which yields that for all $t > 0$, we have

$$\Pr \left\{ \left| R^\top AA^\top R - \mathbb{E} \left[R^\top AA^\top R \right] \right| \geq t \right\} \leq 2 \exp \left(-c \min \left(\frac{t^2}{(C\bar{I})^4 \|AA^\top\|_{\mathbb{F}}^2}, \left(\frac{t}{(C\bar{I})^2 \|AA^\top\|_{\text{op}}} \right)^{1/2} \right) \right),$$

where c and C are two universal and positive constants. Some algebra yields that

$$R^\top AA^\top R - \mathbb{E} \left[R^\top AA^\top R \right] \lesssim (\bar{I})^2 \cdot \left(\|AA^\top\|_{\mathbb{F}} \sqrt{\log(2/\delta)} + \|AA^\top\|_{\text{op}} \log^2(2/\delta) \right)$$

with probability at least $1 - \delta$. Substituting for the expectation and performing some algebraic simplifications proves the claim. \square

A.2 Proof of uniqueness of fixed-point Eq. (18)

We define a function $\psi : [0, +\infty) \mapsto \mathbb{R}$ as

$$\psi(\gamma) = \mathbb{E} \left\{ \frac{|h'(6\|x_\star\|_2 G)|^2 \cdot \mathbb{1}\{G > 0\}}{[|h'(6\|x_\star\|_2 G)| + \gamma]^2} \cdot \int_{-G/4}^{G/4} \frac{t^2 e^{-t^2/2}}{\sqrt{2\pi}} dt \right\}.$$

Note that $\psi(\gamma)$ is a strictly decreasing function of γ ,

$$\lim_{\gamma \rightarrow +\infty} \psi(\gamma) = 0 \quad \text{and} \quad \psi(0) = \int_0^{+\infty} \int_{-\tau/4}^{\tau/4} \frac{t^2 e^{-t^2/2}}{\sqrt{2\pi}} dt \cdot \frac{e^{-\tau^2/2}}{\sqrt{2\pi}} d\tau > 0,$$

and $\psi(0)$ is a universal and positive constant. Consequently, for each $n \geq (16/\psi(0)) \cdot \bar{\omega}(x_\star, X)^2$, the equation $\psi(\gamma) = 16\bar{\omega}(x_\star, X)^2/n$ has a unique and strictly positive solution. \square

A.3 Proof of inequality (23)

We obtain

$$\begin{aligned} \mathbb{E} \left\{ \frac{|h'(6\|x_\star\|_2 G)|^2 \mathbb{1}\{G > 0\}}{[|h'(6\|x_\star\|_2 G)| + \gamma_\star]^2} \int_{-G/4}^{G/4} \frac{t^2 e^{-t^2/2}}{\sqrt{2\pi}} dt \right\} &\geq \int_1^2 \frac{|h'(6\|x_\star\|_2 \tau)|^2}{[|h'(6\|x_\star\|_2 \tau)| + \gamma_\star]^2} \cdot \int_{-\tau/4}^{\tau/4} \frac{t^2 e^{-t^2/2}}{\sqrt{2\pi}} dt \cdot \frac{e^{-\tau^2/2}}{\sqrt{2\pi}} d\tau \\ &\geq \frac{|h'(12\|x_\star\|_2)|^2}{[|h'(12\|x_\star\|_2)| + \gamma_\star]^2} \cdot \int_{-1/4}^{1/4} \frac{t^2 e^{-t^2/2}}{\sqrt{2\pi}} dt \cdot \frac{e^{-2}}{\sqrt{2\pi}}. \end{aligned}$$

Let $c_0 = \int_{-1/4}^{1/4} \frac{t^2 e^{-t^2/2}}{\sqrt{2\pi}} dt \cdot \frac{e^{-2}}{\sqrt{2\pi}}$, and note that it is a universal constant. Combining the inequality in the display above with definition (18), we obtain

$$\frac{16\bar{\omega}(x_\star, X)^2}{n} \geq c_0 \cdot \frac{|h'(12\|x_\star\|_2)|^2}{[|h'(12\|x_\star\|_2)| + \gamma_\star]^2}.$$

Thus, as long as $n \geq (4 \times 16/c_0) \cdot \bar{\omega}(x_\star, X)^2$, we obtain

$$\gamma_\star \geq |h'(12\|x_\star\|_2)| = \sum_{j=1}^W s_j \mu_j \exp(-12\mu_j \|x_\star\|_2).$$

\square

A.4 Proof of inequality (24)

Note that $|h'(t)| \leq \sum_{j=1}^W s_j \mu_j$ for all $t > 0$. Thus,

$$\begin{aligned} \frac{16\bar{\omega}(x_*, X)^2}{n} &= \mathbb{E} \left\{ \frac{|h'(6\|x_*\|_2 G)|^2 \mathbb{1}\{G > 0\}}{[|h'(6\|x_*\|_2 G)| + \gamma_*]^2} \int_{-G/4}^{G/4} \frac{t^2 e^{-t^2/2}}{\sqrt{2\pi}} dt \right\} \\ &\stackrel{(i)}{\gtrsim} \frac{\mathbb{E} \left\{ |h'(6\|x_*\|_2 G)|^2 \mathbb{1}\{G > 0\} e^{-G^2/16} G^3 \right\}}{[\sum_{j=1}^W s_j \mu_j + \gamma_*]^2}. \end{aligned}$$

where step (i) follows because $\int_{-G/4}^{G/4} \frac{t^2 e^{-t^2/2}}{\sqrt{2\pi}} dt \geq \frac{e^{-G^2/16}}{\sqrt{2\pi}} \frac{(G/4)^3}{3}$ for all $G > 0$. Continuing, note that for $G > 0$

$$|h'(6\|x_*\|_2 G)|^2 \geq \sum_{j=1}^W s_j^2 \mu_j^2 \exp(-12\mu_j \|x_*\|_2 G).$$

We thus obtain

$$\begin{aligned} \mathbb{E} \left\{ |h'(6\|x_*\|_2 G)|^2 \mathbb{1}\{G > 0\} e^{-G^2/16} G^3 \right\} &\geq \frac{1}{\sqrt{2\pi}} \sum_{j=1}^W s_j^2 \mu_j^2 \int_0^{+\infty} \exp(-12\mu_j \|x_*\|_2 t - t^2/16 - t^2/2) t^3 dt \\ &\gtrsim \sum_{j=1}^W s_j^2 \mu_j^2 \int_0^{\frac{1}{6\mu_j \|x_*\|_2} \wedge \frac{1}{6}} \exp(-12\mu_j \|x_*\|_2 t - t^2) t^3 dt \\ &\stackrel{(i)}{\gtrsim} \sum_{j=1}^W s_j^2 \mu_j^2 \int_0^{\frac{1}{6\mu_j \|x_*\|_2} \wedge \frac{1}{6}} t^3 dt \\ &\gtrsim \sum_{j=1}^W s_j^2 \mu_j^2 \frac{1}{(\mu_j \|x_*\|_2)^4 \vee 1}, \end{aligned}$$

where in step (i) we use for $0 < t \leq \frac{1}{6\mu_j \|x_*\|_2} \wedge \frac{1}{6}$,

$$\exp(-12\mu_j \|x_*\|_2 t - t^2) \geq e^{-2-1/36} \gtrsim 1.$$

Putting all pieces together yields that there exists a universal and positive constant C such that

$$C \frac{\bar{\omega}(x_*, X)^2}{n} \geq \frac{\sum_{j=1}^W \frac{s_j^2 \mu_j^2}{(\mu_j \|x_*\|_2)^4 \vee 1}}{[\sum_{j=1}^W s_j \mu_j + \gamma_*]^2}.$$

Consequently, if n satisfies

$$C \frac{\bar{\omega}(x_*, X)^2}{n} \leq \sum_{j=1}^W \frac{s_j^2 \mu_j^2}{(\mu_j \|x_*\|_2)^4 \vee 1} \cdot \frac{1}{(2 \sum_{j=1}^W s_j \mu_j)^2},$$

then $\sum_{j=1}^W s_j \mu_j + \gamma_* \geq 2 \sum_{j=1}^W s_j \mu_j$ and thus $\gamma_* \geq \sum_{j=1}^W s_j \mu_j$. \square

B Proof of auxiliary lemmas

In this section, we provide proofs of all the auxiliary lemmas that were stated in Section 4.

B.1 Proof of Lemma 1

Introducing an auxiliary variable $u \in \mathbb{R}^n$, we may write

$$\begin{aligned} \mathcal{P}(A) &\stackrel{(i)}{=} \inf_{\substack{x \in X, u \in \mathbb{R}^n, \\ x \neq x_\star}} \sup_{\lambda \in \mathbb{R}^n} \frac{\frac{1}{n} \sum_{i=1}^n [h(a_i^\top x_\star) - h(u_i)] \cdot [u_i - a_i^\top x_\star]}{\|x - x_\star\|_2^2} + \frac{1}{n} \langle \lambda, Ax - u \rangle \\ &\stackrel{(ii)}{=} \inf_{\substack{x \in X, u \in \mathbb{R}^n, \\ x \neq x_\star}} \sup_{\lambda \in \mathbb{R}^n} \frac{\frac{1}{n} \sum_{i=1}^n [h(a_i^\top x_\star) - h(u_i)] \cdot [u_i - a_i^\top x_\star]}{\|x - x_\star\|_2^2} + \frac{1}{n} \langle \lambda, AP_{x_\star} x - u \rangle + \frac{1}{n} \langle \lambda, AP_{x_\star}^\perp x \rangle \end{aligned}$$

where in step (i), the scalar u_i is the i -th entry of $u \in \mathbb{R}^n$, and the inner sup is imposing the constraints $a_i^\top x = u_i$ for all $i \in [n]$. In step (ii), we use the decomposition $x = P_{x_\star} x + P_{x_\star}^\perp x$, where we recall that P_{x_\star} is the projection matrix onto the span of x_\star , and $P_{x_\star}^\perp$ is the projection matrix onto the orthogonal complement of this subspace. Note that $AP_{x_\star}^\perp$ and AP_{x_\star} are independent of each other since entries of A are i.i.d. standard Gaussian random variables.

The decomposition above allows us to invoke the CGMT (see, e.g., (Thramopoulos et al., 2018) although we will use the version in Proposition 1 of Chandrasekher et al. (2023)) to derive a surrogate for $\mathcal{P}(A)$. Specifically, let $g, \tilde{g} \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, I_d)$ and be independent of $\{a_i\}_{i=1}^n$ and define

$$\begin{aligned} \mathcal{Q}(Ax_\star, g, \tilde{g}) &= \inf_{\substack{x \in X, u \in \mathbb{R}^n, \\ x \neq x_\star}} \sup_{\lambda \in \mathbb{R}^n} \frac{\frac{1}{n} \sum_{i=1}^n [h(a_i^\top x_\star) - h(u_i)] \cdot [u_i - a_i^\top x_\star]}{\|x - x_\star\|_2^2} + \frac{1}{n} \langle \lambda, AP_{x_\star} x - u \rangle \\ &\quad + \frac{1}{n} \|\lambda\|_2 \cdot \langle P_{x_\star}^\perp x, g \rangle + \frac{1}{n} \|P_{x_\star}^\perp x\|_2 \cdot \langle \lambda, \tilde{g} \rangle. \end{aligned} \quad (48)$$

By employing part (a) of Proposition 1 in Chandrasekher et al. (2023), we obtain

$$\Pr \{ \mathcal{P}(A) \leq \tau \} \leq 2 \Pr \{ \mathcal{Q}(Ax_\star, g, \tilde{g}) \leq \tau \} \quad \text{for all } \tau \in \mathbb{R}.$$

In view of the above, it suffices to prove the following lower bound on \mathcal{Q} :

$$\mathcal{Q}(Ax_\star, g, \tilde{g}) \geq \mathcal{D}(Ax_\star, g, \tilde{g}).$$

To prove this lower bound, let us carefully manipulate the minmax problem (48) defining \mathcal{Q} . First, note that we may take the inner supremum in (48) by separately optimizing over the norm and direction of λ . Taking the supremum over the direction of $\lambda \in \mathbb{R}^n$, we obtain

$$\begin{aligned} \mathcal{Q}(Ax_\star, g, \tilde{g}) &= \inf_{\substack{x \in X, u \in \mathbb{R}^n, \\ x \neq x_\star}} \sup_{\lambda \in \mathbb{R}^n} \frac{\frac{1}{n} \sum_{i=1}^n [h(a_i^\top x_\star) - h(u_i)] \cdot [u_i - a_i^\top x_\star]}{\|x - x_\star\|_2^2} \\ &\quad + \frac{\|\lambda\|_2}{n} \|AP_{x_\star} x - u + \|P_{x_\star}^\perp x\|_2 \cdot \tilde{g}\|_2 + \frac{\|\lambda\|_2}{n} \langle P_{x_\star}^\perp x, g \rangle. \end{aligned}$$

Continuing, note that by definition of $\omega(x_\star, X)$ (32), we obtain

$$\langle P_{x_\star}^\perp x, g \rangle \geq -\|P_{x_\star}^\perp x\|_2 \cdot \omega(x_\star, X).$$

Putting the pieces together yields and letting $\gamma = \|\lambda\|_2$, we obtain

$$\begin{aligned} \mathcal{Q}(Ax_\star, g, \tilde{g}) &\geq \inf_{\substack{x \in X, u \in \mathbb{R}^n, \\ x \neq x_\star}} \sup_{\gamma \geq 0} \frac{\frac{1}{n} \sum_{i=1}^n [h(a_i^\top x_\star) - h(u_i)] \cdot [u_i - a_i^\top x_\star]}{\|x - x_\star\|_2^2} \\ &\quad + \frac{\gamma}{n} \left(\|AP_{x_\star} x - u + \|P_{x_\star}^\perp x\|_2 \cdot \tilde{g}\|_2 - \|P_{x_\star}^\perp x\|_2 \cdot \omega(x_\star, X) \right). \end{aligned}$$

In the above program, the decision variable x only enters through two low dimensional functionals $\langle x, x_\star \rangle$ and $\|P_{x_\star}^\perp x\|_2$. This suggests that we should define the scalar decision variables

$$\alpha = \langle x, x_\star \rangle / \|x_\star\|_2 \quad \text{and} \quad \beta = \|P_{x_\star}^\perp x\|_2.$$

Note that by Assumption 1, we have

$$\begin{aligned} & \{(\alpha, \beta) : \alpha = \langle x, x_\star \rangle / \|x_\star\|_2, \beta = \|P_{x_\star}^\perp x\|_2, x \in X, x \neq x_\star\} \\ & \subseteq \{(\alpha, \beta) : \alpha \in \mathbb{R}, \beta \geq 0, 0 < (\alpha - \|x_\star\|_2)^2 + \beta^2 \leq (4\|x_\star\|_2)^2\}. \end{aligned}$$

The above infimum over x can therefore be lower bounded using a low-dimensional infimum, thereby obtaining the bound

$$\begin{aligned} \mathcal{Q}(Ax_\star, g, \tilde{g}) \geq & \inf_{\substack{\alpha \in \mathbb{R}, \beta \geq 0, u \in \mathbb{R}^n, \\ 0 < (\alpha - \|x_\star\|_2)^2 + \beta^2 \leq (4\|x_\star\|_2)^2}} \sup_{\gamma \geq 0} \frac{\frac{1}{n} \sum_{i=1}^n [h(a_i^\top x_\star) - h(u_i)] \cdot [u_i - a_i^\top x_\star]}{(\alpha - \|x_\star\|_2)^2 + \beta^2} \\ & + \frac{\gamma}{n} \left(\|Ax_\star \alpha / \|x_\star\|_2 - u + \beta \tilde{g}\|_2 - \beta \omega(x_\star, X) \right). \quad (49) \end{aligned}$$

The rest of our proof is dedicated to lower-bounding the objective function in Eq. (49) in a particular form. To do so, it is convenient to define a function $f : \mathbb{R} \times \mathbb{R} \mapsto \mathbb{R}$ as

$$f(a_i^\top x_\star, u_i) := \frac{1}{7} |h'(6a_i^\top x_\star)| \cdot [u_i - a_i^\top x_\star]^2 \cdot \mathbb{1}\{a_i^\top x_\star > 0, -6a_i^\top x_\star \leq u_i \leq 6a_i^\top x_\star\}. \quad (50)$$

We will make use of the following relation:

$$[h(a_i^\top x_\star) - h(u_i)] \cdot [u_i - a_i^\top x_\star] \geq f(a_i^\top x_\star, u_i) \quad \text{for all } u_i \in \mathbb{R} \text{ and } a_i^\top x_\star \in \mathbb{R}. \quad (51)$$

We defer the proof of Claim (51) to the end of the proof, and use it to manipulate our lower bound for \mathcal{Q} . Applying Claim (51) to Eq. (49), we obtain

$$\begin{aligned} & \mathcal{Q}(Ax_\star, g, \tilde{g}) \\ & \geq \inf_{\substack{\alpha \in \mathbb{R}, \beta \geq 0, u \in \mathbb{R}^n, \\ 0 < (\alpha - \|x_\star\|_2)^2 + \beta^2 \leq (4\|x_\star\|_2)^2}} \sup_{\gamma \geq 0} \frac{\frac{1}{n} \sum_{i=1}^n f(a_i^\top x_\star, u_i)}{(\alpha - \|x_\star\|_2)^2 + \beta^2} + \frac{\gamma}{n} \left(\|Ax_\star \alpha / \|x_\star\|_2 - u + \beta \tilde{g}\|_2 - \beta \omega(x_\star, X) \right) \\ & \stackrel{(i)}{=} \inf_{\substack{\alpha \in \mathbb{R}, \beta \geq 0, u \in \mathbb{R}^n, \\ 0 < (\alpha - \|x_\star\|_2)^2 + \beta^2 \leq (4\|x_\star\|_2)^2}} \sup_{\gamma \geq 0} \frac{\frac{1}{n} \sum_{i=1}^n f(a_i^\top x_\star, u_i)}{(\alpha - \|x_\star\|_2)^2 + \beta^2} + \frac{\gamma}{7n} \left(\frac{\|Ax_\star \alpha / \|x_\star\|_2 - u + \beta \tilde{g}\|_2^2 - \beta^2 \omega(x_\star, X)^2}{(\alpha - \|x_\star\|_2)^2 + \beta^2} \right). \end{aligned}$$

To obtain step (i), we use the following reasoning: By virtue of taking the supremum over $\gamma \geq 0$, we are imposing the constraint $\|Ax_\star \alpha / \|x_\star\|_2 - u + \beta \tilde{g}\|_2 = \beta \omega(x_\star, X)$ in the first inequality. On the other hand, and by the same logic, the inner supremum in the second inequality is imposing the constraint

$$\frac{\|Ax_\star \alpha / \|x_\star\|_2 - u + \beta \tilde{g}\|_2^2}{(\alpha - \|x_\star\|_2)^2 + \beta^2} = \frac{\beta^2 \omega(x_\star, X)^2}{(\alpha - \|x_\star\|_2)^2 + \beta^2},$$

where we note that the denominator in the expression above is always nonzero by assumption.

Continuing, by the max-min inequality, we can switch $\inf_{u \in \mathbb{R}^n}$ and $\sup_{\gamma \geq 0}$, and obtain

$$\mathcal{Q}(Ax_\star, g, \tilde{g}) \geq \inf_{\substack{\alpha \in \mathbb{R}, \beta \geq 0, \\ (\alpha - \|x_\star\|_2)^2 + \beta^2 = R^2}} \sup_{\gamma \geq 0} \inf_{u \in \mathbb{R}^n} \frac{\frac{1}{n} \sum_{i=1}^n f(a_i^\top x_\star, u_i) + \frac{\gamma}{7n} \left(\|Ax_\star \alpha / \|x_\star\|_2 - u + \beta \tilde{g}\|_2^2 - \beta^2 \omega(x_\star, X)^2 \right)}{(\alpha - \|x_\star\|_2)^2 + \beta^2}.$$

Note that the $\inf_{u \in \mathbb{R}^n}$ is separable over $i \in [n]$, and each $\inf_{u_i \in \mathbb{R}}$ has an explicit solution given by

$$\begin{aligned} & \inf_{u_i \in \mathbb{R}} f(a_i^\top x_\star, u_i) + \frac{\gamma}{7} \cdot \left(\alpha a_i^\top x_\star / \|x_\star\|_2 + \beta \tilde{g}_i - u_i \right)^2 \\ & \geq \begin{cases} \frac{1}{7} \frac{\gamma \cdot |h'(6a_i^\top x_\star)|}{\gamma + |h'(6a_i^\top x_\star)|} \cdot \left((\alpha / \|x_\star\|_2 - 1) a_i^\top x_\star + \beta \tilde{g}_i \right)^2, & \text{if } -6a_i^\top x_\star \leq \beta \tilde{g}_i + \frac{\alpha}{\|x_\star\|_2} a_i^\top x_\star \leq 6a_i^\top x_\star, a_i^\top x_\star > 0, \\ 0, & \text{otherwise.} \end{cases} \\ & = \mathcal{T}_i(\alpha, \beta, \gamma), \end{aligned} \tag{52}$$

where in the last step we use definition (34). We defer the explicit calculations of inequality (52) to the end of this section. Now, combining the two pieces in the display above, we obtain

$$\mathcal{Q}(Ax_\star, g, \tilde{g}) \geq \inf_{\substack{\alpha \in \mathbb{R}, \beta \geq 0, \\ (\alpha - \|x_\star\|_2)^2 + \beta^2 = R^2}} \sup_{\gamma \geq 0} \frac{\frac{1}{n} \sum_{i=1}^n \mathcal{T}_i(\alpha, \beta, \gamma) - \frac{\gamma}{7n} \beta^2 \omega(x_\star, X)^2}{(\alpha - \|x_\star\|_2)^2 + \beta^2} = \mathcal{D}(Ax_\star, \tilde{g}),$$

where in the last step we recall definition (33). This concludes the proof.

It remains to prove Claim (51) and to verify Eq. (52).

Proof of Claim (51). We consider three cases.

Case 1: $a_i^\top x_\star \leq 0$ or $u_i > 6a_i^\top x_\star$ or $u_i < -6a_i^\top x_\star$. Note that in this case, $f(a_i^\top x_\star, u_i) = 0$. Moreover, since $h(\cdot)$ is non-increasing, we obtain

$$[h(a_i^\top x_\star) - h(u_i)] \cdot [u_i - a_i^\top x_\star] \geq 0 \geq f(a_i^\top x_\star, u_i).$$

Case 2: $a_i^\top x_\star > 0$ and $0 \leq u_i \leq 6a_i^\top x_\star$. By the mean value theorem, we obtain

$$h(a_i^\top x_\star) - h(u_i) = h'(t_i)(a_i^\top x_\star - u_i) \text{ for some } \min(a_i^\top x_\star, u_i) \leq t_i \leq \max(a_i^\top x_\star, u_i).$$

Note that $h'(t_i) \leq 0$ and thus

$$[h(a_i^\top x_\star) - h(u_i)] \cdot [u_i - a_i^\top x_\star] = |h'(t_i)| \cdot [u_i - a_i^\top x_\star]^2 \geq |h'(6a_i^\top x_\star)| \cdot [u_i - a_i^\top x_\star]^2,$$

where in the last step we use the fact that $0 \leq t_i \leq 6a_i^\top x_\star$ and that $|h'(\cdot)|$ is strictly decreasing when its argument is positive.

Case 3: $a_i^\top x_\star > 0$ and $-6a_i^\top x_\star \leq u_i < 0$. Note that $h(u_i) = h(0)$ for $u_i < 0$ and thus

$$\begin{aligned} [h(a_i^\top x_\star) - h(u_i)] \cdot [u_i - a_i^\top x_\star] &= [h(a_i^\top x_\star) - h(0)] \cdot [u_i - a_i^\top x_\star] \\ &\stackrel{(i)}{\geq} |h'(6a_i^\top x_\star)| \cdot a_i^\top x_\star \cdot [a_i^\top x_\star - u_i] \\ &\geq \frac{1}{7} |h'(6a_i^\top x_\star)| \cdot [a_i^\top x_\star - u_i]^2, \end{aligned}$$

where in step (i) we use the mean value theorem once more, and in the last step we use $u_i \in [-6a_i^\top x_\star, 0]$. This concludes the proof of claim (51).

Verifying Eq. (52). We consider two cases:

Case 1: $a_i^\top x_\star \alpha / \|x_\star\|_2 + \beta \tilde{g}_i < -6a_i^\top x_\star$ **or** $a_i^\top x_\star \alpha / \|x_\star\|_2 + \beta \tilde{g}_i > 6a_i^\top x_\star$ **or** $a_i^\top x_\star \leq 0$. Note that in this case, by letting $u_i = a_i^\top x_\star \alpha / \|x_\star\|_2 + \beta \tilde{g}_i$ and recalling Eq. (50), we obtain

$$f(a_i^\top x_\star, u_i) + \gamma \cdot \left(\alpha a_i^\top x_\star / \|x_\star\|_2 + \beta \tilde{g}_i - u_i \right)^2 = 0.$$

Since $f(a_i^\top x_\star, u_i) \geq 0$ for all $u_i \in \mathbb{R}$, thus $u_i = a_i^\top x_\star \alpha / \|x_\star\|_2 + \beta \tilde{g}_i$ attains the minimum value 0.

Case 2: $-6a_i^\top x_\star \leq a_i^\top x_\star \alpha / \|x_\star\|_2 + \beta \tilde{g}_i \leq 6a_i^\top x_\star$ **and** $a_i^\top x_\star > 0$. Taking the derivative and setting it to zero yields

$$|h'(6a_i^\top x_\star)| \cdot (u_i - a_i^\top x_\star) \cdot \mathbb{1}\{a_i^\top x_\star > 0, -6a_i^\top x_\star \leq u_i \leq 6a_i^\top x_\star\} + \gamma \cdot (u_i - a_i^\top x_\star \alpha / \|x_\star\|_2 - \beta \tilde{g}_i) = 0$$

The above equation has a unique solution

$$u_i = \frac{|h'(6a_i^\top x_\star)| \cdot a_i^\top x_\star + \gamma \cdot (a_i^\top x_\star \alpha / \|x_\star\|_2 + \beta \tilde{g}_i)}{|h'(6a_i^\top x_\star)| + \gamma},$$

and its corresponding function value is

$$f(a_i^\top x_\star, u_i) + \frac{\gamma}{7} \cdot \left(\alpha a_i^\top x_\star / \|x_\star\|_2 + \beta \tilde{g}_i - u_i \right)^2 = \frac{1}{7} \frac{\gamma \cdot |h'(6a_i^\top x_\star)|}{\gamma + |h'(6a_i^\top x_\star)|} \cdot \left((\alpha / \|x_\star\|_2 - 1) a_i^\top x_\star + \beta \tilde{g}_i \right)^2.$$

This concludes the proof. \square

B.2 Proof of Lemma 2

To reduce the notational burden, we define $\ell_{\alpha, \beta} : \mathbb{R} \mapsto \mathbb{R}$ as

$$\ell_{\alpha, \beta}(\gamma) = \frac{1}{n} \sum_{i=1}^n \mathcal{T}_i(\alpha, \beta, \gamma) - \frac{\gamma \cdot \beta^2}{7n} \cdot \omega(X, x_\star)^2,$$

where $\mathcal{T}_i(\alpha, \beta, \gamma)$ is defined in Eq. (34). Computing the gradient (i.e. derivative) with respect to γ yields

$$\begin{aligned} \nabla \ell_{\alpha, \beta}(\gamma) &= \frac{1}{7n} \sum_{i=1}^n \frac{|h'(6a_i^\top x_\star)|^2 \left((\alpha / \|x_\star\|_2 - 1) a_i^\top x_\star + \beta \tilde{g}_i \right)^2}{(\gamma + |h'(6a_i^\top x_\star)|)^2} \times \\ &\quad \mathbb{1} \left\{ -6a_i^\top x_\star \leq \beta \tilde{g}_i + \frac{\alpha}{\|x_\star\|_2} a_i^\top x_\star \leq 6a_i^\top x_\star, a_i^\top x_\star > 0 \right\} - \frac{\beta^2 \omega(x_\star, X)^2}{7n} \end{aligned}$$

Proof of finiteness. We first prove that $\bar{\gamma}(\alpha, \beta) < \infty$. Note that

$$\lim_{\gamma \rightarrow +\infty} \nabla \ell_{\alpha, \beta}(\gamma) = -\frac{\beta^2 \omega(x_\star, X)^2}{7n} \stackrel{(i)}{\leq} -\frac{\beta^2 \bar{\omega}(x_\star, X)^2 / 4}{7n} < 0,$$

where step (i) holds with probability at least $1 - 2 \exp(-\bar{\omega}(x_\star, X)^2 / 8)$ by applying inequality (55d) in Lemma 4, and in the last step we use $\beta > 0$ and $\bar{\omega}(x_\star, X) > 0$. Since $\ell_{\alpha, \beta}(\gamma)$ is a concave function of γ , we obtain that with probability at least $1 - 2 \exp(-\bar{\omega}(x_\star, X)^2 / 8)$, the scalar $\bar{\gamma}(\alpha, \beta)$ is finite uniformly for all (α, β) satisfying Eq. (39).

Proof that $\bar{\gamma}(\alpha, \beta) \geq \gamma_*$. To show this, we will show that $\nabla \ell_{\alpha, \beta}(\gamma_*) > 0$ with the desired high-probability. Note that since $\nabla \ell_{\alpha, \beta}(\gamma)$ is non-increasing over γ , this implies $\nabla \ell_{\alpha, \beta}(\gamma) \geq \nabla \ell_{\alpha, \beta}(\gamma_*) > 0$ for all $0 \leq \gamma \leq \gamma_*$. This further implies $\ell_{\alpha, \beta}(\gamma)$ is strictly increasing when $0 \leq \gamma \leq \gamma_*$. Further by definition

$$\bar{\gamma}(\alpha, \beta) = \arg \sup_{\gamma > 0} \ell_{\alpha, \beta}(\gamma).$$

We thus conclude that $\bar{\gamma}(\alpha, \beta) \geq \gamma_*$.

We start with obtaining a lower bound on $\nabla \ell_{\alpha, \beta}(\gamma_*)$. We begin with a bound on the indicator terms in this expression. Substituting inequality (44) into the definition of the gradient, we have

$$\begin{aligned} \nabla \ell_{\alpha, \beta}(\gamma_*) &\geq \frac{1}{7n} \sum_{i=1}^n \frac{|h'(6a_i^\top x_*)|^2 ((\alpha / \|x_*\|_2 - 1)a_i^\top x_* + \beta \tilde{g}_i)^2}{(\gamma_* + |h'(6a_i^\top x_*)|)^2} \mathbb{1} \left\{ -\frac{a_i^\top x_*}{4\|x_*\|_2} \leq \tilde{g}_i \leq \frac{a_i^\top x_*}{\|x_*\|_2}, a_i^\top x_* > 0 \right\} - \frac{\beta^2 \omega(x_*, X)^2}{7n} \\ &= \frac{(\alpha - \|x_*\|_2)^2}{7} \frac{1}{n} \sum_{i=1}^n \delta_i + \frac{\beta^2}{7} \frac{1}{n} \sum_{i=1}^n \eta_i - \frac{2(\alpha - \|x_*\|_2)\beta}{7} \frac{1}{n} \sum_{i=1}^n \iota_i - \frac{\beta^2 \omega(x_*, X)^2}{7n}, \end{aligned} \quad (53)$$

where in the last step we use the shorthands

$$\delta_i = \frac{|h'(6a_i^\top x_*)|^2 (a_i^\top x_*)^2 / \|x_*\|_2^2}{(\gamma_* + |h'(6a_i^\top x_*)|)^2} \cdot \mathbb{1} \left\{ -\frac{a_i^\top x_*}{4\|x_*\|_2} \leq \tilde{g}_i \leq \frac{a_i^\top x_*}{\|x_*\|_2}, a_i^\top x_* > 0 \right\}, \quad (54a)$$

$$\eta_i = \frac{|h'(6a_i^\top x_*)|^2 \cdot \tilde{g}_i^2}{(\gamma_* + |h'(6a_i^\top x_*)|)^2} \cdot \mathbb{1} \left\{ -\frac{a_i^\top x_*}{4\|x_*\|_2} \leq \tilde{g}_i \leq \frac{a_i^\top x_*}{\|x_*\|_2}, a_i^\top x_* > 0 \right\}, \quad (54b)$$

$$\iota_i = \frac{|h'(6a_i^\top x_*)|^2 \cdot \tilde{g}_i a_i^\top x_* / \|x_*\|_2}{(\gamma_* + |h'(6a_i^\top x_*)|)^2} \cdot \mathbb{1} \left\{ -\frac{a_i^\top x_*}{4\|x_*\|_2} \leq \tilde{g}_i \leq \frac{a_i^\top x_*}{\|x_*\|_2}, a_i^\top x_* > 0 \right\}. \quad (54c)$$

We next invoke the following auxiliary lemma to control the random variables in the RHS of inequality (53).

Lemma 4. *Let $\{\delta_i, \eta_i, \iota_i\}_{i=1}^n$ be defined in Eq. (54a), Eq. (54b), and Eq. (54c), respectively. Let $\omega(x_*, X)$ and $\bar{\omega}(x_*, X)$ be defined in Eq. (32) and Eq. (17), respectively. Then the following holds for all $\epsilon \in (0, 1)$ and for all $t > 0$:*

$$\Pr \left\{ \left| \frac{1}{n} \sum_{i=1}^n \delta_i - \mathbb{E}\{\delta_i\} \right| \geq 8 \log(2/\epsilon)/n + \sqrt{64 \mathbb{E}\{\delta_i\} \log(2/\epsilon)/n} \right\} \leq \epsilon, \quad (55a)$$

$$\Pr \left\{ \left| \frac{1}{n} \sum_{i=1}^n \eta_i - \mathbb{E}\{\eta_i\} \right| \geq 8 \log(2/\epsilon)/n + \sqrt{64 \mathbb{E}\{\eta_i\} \log(2/\epsilon)/n} \right\} \leq \epsilon, \quad (55b)$$

$$\Pr \left\{ \left| \frac{1}{n} \sum_{i=1}^n \iota_i \right| \geq 8 \log(2/\epsilon)/n + \sqrt{64 \mathbb{E}\{\delta_i\} \log(2/\epsilon)/n} \right\} \leq \epsilon, \quad (55c)$$

$$\mathbb{E}[\delta_i] \wedge \mathbb{E}[\eta_i] \geq \frac{16 \bar{\omega}(x_*, X)^2}{n} \quad \text{and} \quad \Pr \left\{ |\omega(x_*, X) - \bar{\omega}(x_*, X)| \geq t \right\} \leq 2 \exp(-t^2/2). \quad (55d)$$

We provide the proof of Lemma 4 in Section B.2.1. By setting $\epsilon = 2 \exp(-\bar{\omega}(x_*, X)^2/64)$ and applying inequality (55) and the union bound, we obtain that with probability at least $1 - 6 \exp(-$

$\bar{\omega}(x_*, X)^2/64$), we have

$$\begin{aligned}\frac{1}{n} \sum_{i=1}^n \delta_i &\geq \mathbb{E}\{\delta_i\} - \frac{\bar{\omega}(x_*, X)^2}{8n} - \sqrt{\mathbb{E}\{\delta_i\}} \sqrt{\frac{\bar{\omega}(x_*, X)^2}{n}}, \\ \frac{1}{n} \sum_{i=1}^n \eta_i &\geq \mathbb{E}\{\eta_i\} - \frac{\bar{\omega}(x_*, X)^2}{8n} - \sqrt{\mathbb{E}\{\eta_i\}} \sqrt{\frac{\bar{\omega}(x_*, X)^2}{n}}, \\ \left| \frac{1}{n} \sum_{i=1}^n \iota_i \right| &\leq \frac{\bar{\omega}(x_*, X)^2}{8n} + \sqrt{\mathbb{E}\{\delta_i\}} \sqrt{\frac{\bar{\omega}(x_*, X)^2}{n}}.\end{aligned}$$

Substituting the inequalities in the display above into inequality (53) yields

$$\begin{aligned}7\nabla\ell_{\alpha,\beta}(\gamma_*) &\geq (\alpha - \|x_*\|_2)^2 \cdot \left(\mathbb{E}\{\delta_i\} - \frac{\bar{\omega}(x_*, X)^2}{8n} - \sqrt{\mathbb{E}\{\delta_i\}} \sqrt{\frac{\bar{\omega}(x_*, X)^2}{n}} \right) \\ &\quad + \beta^2 \cdot \left(\mathbb{E}\{\eta_i\} - \frac{\bar{\omega}(x_*, X)^2}{8n} - \sqrt{\mathbb{E}\{\eta_i\}} \sqrt{\frac{\bar{\omega}(x_*, X)^2}{n}} \right) \\ &\quad - 2|\alpha - \|x_*\|_2|\beta \cdot \left(\frac{\bar{\omega}(x_*, X)^2}{8n} + \sqrt{\mathbb{E}\{\delta_i\}} \sqrt{\frac{\bar{\omega}(x_*, X)^2}{n}} \right) - \beta^2 \frac{\omega(x_*, X)^2}{n}.\end{aligned}$$

Note that by the AM-GM inequality, we obtain $2|\alpha - \|x_*\|_2|\beta \leq (\alpha - \|x_*\|_2)^2 + \beta^2$ and

$$2|\alpha - \|x_*\|_2|\beta \sqrt{\mathbb{E}\{\delta_i\}} \sqrt{\frac{\bar{\omega}(x_*, X)^2}{n}} \leq 2\beta^2 \frac{\bar{\omega}(x_*, X)^2}{n} + \frac{(\alpha - \|x_*\|_2)^2}{2} \mathbb{E}\{\delta_i\}.$$

Substituting above, we have

$$\begin{aligned}7\nabla\ell_{\alpha,\beta}(\gamma_*) &\geq (\alpha - \|x_*\|_2)^2 \cdot \left(\frac{1}{2} \mathbb{E}\{\delta_i\} - \frac{\bar{\omega}(x_*, X)^2}{4n} - \sqrt{\mathbb{E}\{\delta_i\}} \sqrt{\frac{\bar{\omega}(x_*, X)^2}{n}} \right) \\ &\quad + \beta^2 \cdot \left(\mathbb{E}\{\eta_i\} - \frac{(2 + 1/8)\bar{\omega}(x_*, X)^2}{n} - \sqrt{\mathbb{E}\{\eta_i\}} \sqrt{\frac{\bar{\omega}(x_*, X)^2}{n}} \right) - \beta^2 \frac{\omega(x_*, X)^2}{n}.\end{aligned}$$

Continuing, using inequality (55d) yields

$$\begin{aligned}&\frac{1}{2} \mathbb{E}\{\delta_i\} - \frac{\bar{\omega}(x_*, X)^2}{4n} - \sqrt{\mathbb{E}\{\delta_i\}} \sqrt{\frac{\bar{\omega}(x_*, X)^2}{n}} \\ &= 0.25 \mathbb{E}\{\delta_i\} - \frac{\bar{\omega}(x_*, X)^2}{4n} + 0.25 \mathbb{E}\{\delta_i\} - \sqrt{\mathbb{E}\{\delta_i\}} \sqrt{\frac{\bar{\omega}(x_*, X)^2}{n}} \geq \frac{3\bar{\omega}(x_*, X)^2}{n},\end{aligned}$$

and similar calculation yields

$$\mathbb{E}\{\eta_i\} - \frac{(2 + 1/8)\bar{\omega}(x_*, X)^2}{n} - \sqrt{\mathbb{E}\{\eta_i\}} \sqrt{\frac{\bar{\omega}(x_*, X)^2}{n}} \geq \frac{3\bar{\omega}(x_*, X)^2}{n}.$$

Putting the pieces together yields

$$\begin{aligned}7\nabla\ell_{\alpha,\beta}(\gamma_*) &\geq ((\alpha - \|x_*\|_2)^2 + \beta^2) \cdot \frac{3\bar{\omega}(x_*, X)^2}{n} - \beta^2 \frac{\omega(x_*, X)^2}{n} \\ &\stackrel{(i)}{\geq} ((\alpha - \|x_*\|_2)^2 + \beta^2) \cdot \frac{(3 - 9/4)\bar{\omega}(x_*, X)^2}{n} > 0,\end{aligned}$$

where in step (i) we use inequality (55d) so that

$$\omega(x_*, X) \leq (3/2)\bar{\omega}(x_*, X) \quad \text{with probability } \geq 1 - 2 \exp(-\bar{\omega}(x_*, X)^2/8).$$

In conclusion, we obtain that $\nabla\ell_{\alpha,\beta}(\gamma_*) > 0$ with probability at least $1 - 8 \exp(-\bar{\omega}(x_*, X)^2/64)$, as desired. \square

B.2.1 Proof of Lemma 4

We prove each part separately.

Proof of inequality (55a). To reduce the notational burden, we define a function $\psi : \mathbb{R} \mapsto \mathbb{R}$

$$\psi(t) = \frac{|h'(6\|x_\star\|_2 t)|^2}{(\gamma_\star + |h'(6\|x_\star\|_2 t)|)^2}, \quad \text{for all } t \in \mathbb{R}. \quad (56)$$

We first claim that δ_i satisfies the Bernstein condition (see, e.g., [Wainwright \(2019, Eq. 2.16\)](#))

$$|\mathbb{E}\{(\delta_i - \mathbb{E}[\delta_i])^k\}| \leq \frac{1}{2} \cdot k! \cdot 4^{k-2} \cdot 16\mathbb{E}[\delta_i] \quad \text{for all } k = 2, 3, \dots \quad (57)$$

deferring its proof to the end of this section. Consequently, by [Wainwright \(2019, Proposition 2.3\)](#), we obtain that for $a = 32\mathbb{E}\{\delta_i\}/n$ and $b = 4/n$,

$$\mathbb{E}\left\{e^{\lambda \frac{1}{n} \sum_{i=1}^n (\delta_i - \mathbb{E}\{\delta_i\})}\right\} = \prod_{i=1}^n \mathbb{E}\left\{e^{\frac{\lambda}{n} (\delta_i - \mathbb{E}\{\delta_i\})}\right\} \leq \exp\left\{\frac{a\lambda^2/2}{1 - |\lambda|b}\right\} \quad \text{for all } \lambda \in (-1/b, 1/b),$$

and

$$\Pr\left\{\frac{1}{n} \sum_{i=1}^n \delta_i - \mathbb{E}\{\delta_i\} \geq t\right\} \leq e^{-\frac{t^2/2}{a+bt}}.$$

Thus, letting $t = 2b \log(2/\epsilon) + \sqrt{2a \log(2/\epsilon)}$, we obtain that for $\epsilon \in (0, 1]$,

$$\Pr\left\{\frac{1}{n} \sum_{i=1}^n \delta_i - \mathbb{E}\{\delta_i\} \geq 2b \log(2/\epsilon) + \sqrt{2a \log(2/\epsilon)}\right\} \leq \epsilon/2.$$

Following the same steps yields the lower tail bound

$$\Pr\left\{\frac{1}{n} \sum_{i=1}^n \delta_i - \mathbb{E}\{\delta_i\} \leq -2b \log(2/\epsilon) - \sqrt{2a \log(2/\epsilon)}\right\} \leq \epsilon/2.$$

Applying the union bound thus yields

$$\Pr\left\{\left|\frac{1}{n} \sum_{i=1}^n \delta_i - \mathbb{E}\{\delta_i\}\right| \geq 2b \log(2/\epsilon) + \sqrt{2a \log(2/\epsilon)}\right\} \leq \epsilon.$$

The desired result follows by noting $a = 32\mathbb{E}\{\delta_i\}/n$ and $b = 4/n$. It remains to prove the inequality (57).

Establishing the Bernstein condition (57). Let $\bar{\delta}_i = \delta_i - \mathbb{E}\{\delta_i\}$. Applying Jensen's inequality to the convex function $t \mapsto t^k$, we find that

$$|\bar{\delta}_i|^k \leq (|\delta_i| + \mathbb{E}\{|\delta_i|\})^k \leq 2^{k-1} (|\delta_i|^k + \mathbb{E}\{|\delta_i|^k\}).$$

Consequently,

$$\mathbb{E}\{|\bar{\delta}_i|^k\} \leq 2^{k-1} (\mathbb{E}\{|\delta_i|^k\} + \mathbb{E}\{|\delta_i|\}^k) \leq 2^k \mathbb{E}\{|\delta_i|^k\}.$$

Letting $G, \tilde{G} \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, 1)$, we compute

$$\begin{aligned} \mathbb{E}\{|\delta_i|^k\} &= \mathbb{E}\{\psi(G)^k G^{2k} \cdot \mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4, G > 0\}\} \\ &\stackrel{(i)}{\leq} \mathbb{E}\{\psi(G)G^{2k} \cdot \mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4, G > 0\}\} \\ &= \mathbb{E}\left\{\psi(G)G^{2k} \int_{-G/4}^{G/4} \phi(t)dt \cdot \mathbb{1}\{G > 0\}\right\}, \end{aligned} \quad (58)$$

where step (i) follows from the sandwich relation $0 \leq \psi(t) \leq 1$ for all $t \in \mathbb{R}$ and in the final step we have used the shorthand $\phi(t) = \exp\{-t^2/2\}/(\sqrt{2\pi})$. Integration by parts yields

$$\begin{aligned} \mathbb{E}\left\{\psi(G)G^{2k} \int_{-G/4}^{G/4} \phi(t) \cdot \mathbb{1}\{G > 0\}dt\right\} &= (2k-1)\mathbb{E}\left\{\psi(G)G^{2k-2} \int_{-G/4}^{G/4} \phi(t)dt \cdot \mathbb{1}\{G > 0\}\right\} \\ &+ \mathbb{E}\left\{\psi'(G)G^{2k-1} \int_{-G/4}^{G/4} \phi(t)dt \cdot \mathbb{1}\{G > 0\}\right\} + \mathbb{E}\left\{\psi(G)G^{2k-1} \frac{\phi(G/4)}{2} \cdot \mathbb{1}\{G > 0\}\right\}. \end{aligned} \quad (59)$$

We next bound the last two terms in the equality display above. Note that $\psi'(t) \leq 0$ for all $t > 0$ since $\psi(t)$ (56) is non-increasing for $t > 0$, and thus the second term is non-positive, i.e.,

$$\mathbb{E}\left\{\psi'(G)G^{2k-1} \int_{-G/4}^{G/4} \phi(t)dt \cdot \mathbb{1}\{G > 0\}\right\} \leq 0.$$

For the third term, note that

$$G \frac{\phi(G/4)}{2} \cdot \mathbb{1}\{G > 0\} \leq \int_{-G/4}^{G/4} \phi(t)dt \cdot \mathbb{1}\{G > 0\},$$

and thus we can bound the third term as

$$\mathbb{E}\left\{\psi(G)G^{2k-1} \frac{\phi(G/4)}{2} \cdot \mathbb{1}\{G > 0\}\right\} \leq \mathbb{E}\left\{\psi(G)G^{2k-2} \int_{-G/4}^{G/4} \phi(t)dt \cdot \mathbb{1}\{G > 0\}\right\}.$$

Substituting the bounds of the second and third term into Eq. (59) yields that for $k \geq 1$

$$\mathbb{E}\left\{\psi(G)G^{2k} \int_{-G/4}^{G/4} \phi(t) \cdot \mathbb{1}\{G > 0\}dt\right\} \leq 2k \cdot \mathbb{E}\left\{\psi(G)G^{2k-2} \int_{-G/4}^{G/4} \phi(t)dt \cdot \mathbb{1}\{G > 0\}\right\}.$$

Note that the above inequality holds for all natural numbers $k \geq 1$. Iteratively applying the inequality in the display above yields, for $k \geq 1$,

$$\mathbb{E}\left\{\psi(G)G^{2k} \int_{-G/4}^{G/4} \phi(t) \cdot \mathbb{1}\{G > 0\}dt\right\} \leq 2^{k-1}k! \cdot \mathbb{E}\left\{\psi(G)G^2 \int_{-G/4}^{G/4} \phi(t)dt \cdot \mathbb{1}\{G > 0\}\right\}. \quad (60)$$

Consequently, recalling inequality (58), we obtain the upper bound

$$\mathbb{E}\{|\delta_i|^k\} \leq 2^{k-1}k! \cdot \mathbb{E}\left\{\psi(G)G^2 \int_{-G/4}^{G/4} \phi(t)dt \cdot \mathbb{1}\{G > 0\}\right\} = 2^{k-1}k! \cdot \mathbb{E}\{\delta_i\},$$

where in the last step we use the definition of δ_i (54a). Putting the pieces together yields the desired result.

Proof of inequality (55b). We first claim that the random variable $\eta_i - \mathbb{E}\{\eta_i\}$ satisfies the following Bernstein condition, deferring its proof to the end,

$$|\mathbb{E}\{(\eta_i - \mathbb{E}\{\eta_i\})^k\}| \leq \frac{1}{2} \cdot k! \cdot 4^{k-2} \cdot 16\mathbb{E}\{\eta_i\} \quad \text{for all } k = 2, 3, 4, \dots \quad (61)$$

The rest of proof is identical to the proof of inequality (55a), so we omit it.

It remains to verify inequality (61). Let $\bar{\eta}_i = \eta_i - \mathbb{E}\{\eta_i\}$. Following identical steps as in the proof of inequality (57), we obtain

$$\mathbb{E}\{|\bar{\eta}_i|^k\} \leq 2^k \mathbb{E}\{|\eta_i|^k\}. \quad (62)$$

Continuing, by definition of η_i , we obtain that for $G, \tilde{G} \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, 1)$

$$\begin{aligned} \mathbb{E}\{|\eta_i|^k\} &= \mathbb{E}\left\{\psi(G)^k \tilde{G}^{2k} \cdot \mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4, G > 0\}\right\} \\ &\leq \mathbb{E}\left\{\psi(G) \tilde{G}^{2k} \cdot \mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4, G > 0\}\right\}, \end{aligned} \quad (63)$$

where in the last step we use $0 \leq \psi(t) \leq 1$ for all $t > 0$. Conditioning on G and for $G > 0$, we obtain

$$\begin{aligned} \mathbb{E}\left\{\tilde{G}^{2k} \cdot \mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4\}\right\} &= \int_{-G/4}^{G/4} t^{2k} \phi(t) dt \\ &\stackrel{(i)}{=} (2k-1) \cdot \int_{-G/4}^{G/4} t^{2k-2} \phi(t) dt - t^{2k-1} \phi(t) \Big|_{-G/4}^{G/4} \\ &\stackrel{(ii)}{\leq} (2k-1) \cdot \int_{-G/4}^{G/4} t^{2k-2} \phi(t) dt \\ &= (2k-1) \cdot \mathbb{E}\left\{\tilde{G}^{2k-2} \cdot \mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4\}\right\}, \end{aligned}$$

where in step (i) we use integration by parts and in step (ii) we use $t^{2k-1} \phi(t) \Big|_{-G/4}^{G/4} \geq 0$. The inequality in the display holds for any positive integer k and thus applying it sequentially yields that, conditionally on G and $G > 0$,

$$\begin{aligned} \mathbb{E}\left\{\tilde{G}^{2k} \cdot \mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4\}\right\} &\leq \prod_{i=1}^k (2i-1) \cdot \mathbb{E}\left\{\tilde{G}^2 \cdot \mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4\}\right\} \\ &\leq 2^k k! \cdot \mathbb{E}\left\{\tilde{G}^2 \cdot \mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4\}\right\}. \end{aligned} \quad (64)$$

Substituting the inequality in the display into inequality (63), we obtain

$$\mathbb{E}\{|\eta_i|^k\} \leq 2^k k! \cdot \mathbb{E}\left\{\psi(G) \tilde{G}^2 \cdot \mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4, G > 0\}\right\} = 2^k k! \cdot \mathbb{E}\{\eta_i\}$$

The desired result follows by substituting the inequality in the display into inequality (62).

Proof of inequality (55c). We again show that ι_i satisfies a Bernstein condition. First, note that for odd k ,

$$\mathbb{E}\{\iota_i^k\} = \mathbb{E}\left\{\psi(G)^k \tilde{G}^k G^k \mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4, G > 0\}\right\} = 0,$$

as $\tilde{G}^k \mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4, G > 0\}$ is an odd function of \tilde{G} and $\tilde{G}, G \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, 1)$. Continuing, using the fact $0 \leq \psi(t) \leq 1$ for all $t \in \mathbb{R}$, we obtain

$$\begin{aligned} \mathbb{E}\{\iota_i^{2k}\} &\leq \mathbb{E}\left\{\psi(G)G^{2k}\tilde{G}^{2k}\mathbb{1}\{G > 0\}\mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4\}\right\} \\ &\stackrel{(i)}{\leq} 2^k k! \cdot \mathbb{E}\left\{\psi(G)G^{2k}\mathbb{1}\{G > 0\}\mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4\}\right\} \\ &\stackrel{(ii)}{\leq} (2^k k!)^2 \cdot \mathbb{E}\left\{\psi(G)G^2\mathbb{1}\{G > 0\}\mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4\}\right\} = (2^k k!)^2 \cdot \mathbb{E}\{\delta_i\}, \end{aligned}$$

where in step (i) we use inequality (64) so that conditionally on G and $G > 0$,

$$\mathbb{E}\left\{\tilde{G}^{2k}\mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4\}\right\} \leq 2^k k! \cdot \mathbb{E}\left\{\mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4\}\right\},$$

and in step (ii) we use inequality (60). We thus have

$$\mathbb{E}\{\iota_i^{2k}\} \leq \frac{1}{2} \cdot (2k)! \cdot 4^{2k-2} \cdot 16\mathbb{E}\{\delta_i\}, \quad \text{for all } k = 2, 3, 4, \dots$$

Following the identical steps in the proof of inequality (55a), we obtain that for all $\epsilon \in (0, 1)$,

$$\Pr\left\{\left|\frac{1}{n}\sum_{i=1}^n \iota_i\right| \geq 8\log(2/\epsilon)/n + \sqrt{64\mathbb{E}\{\delta_i\}\log(2/\epsilon)/n}\right\} \leq \epsilon.$$

Proof of inequality (55d). We first lower bound $\mathbb{E}\{\delta_i\}$. Expanding the definition of δ_i (54a) yields

$$\mathbb{E}\{\delta_i\} = \mathbb{E}\left\{\frac{|h'(6\|x_\star\|_2 G)|^2 G^2 \mathbb{1}\{G > 0\}}{(|h'(6\|x_\star\|_2 G)|^2 + \gamma_\star)^2} \cdot \int_{-G/4}^{G/4} \frac{e^{-t^2/2}}{\sqrt{2\pi}} dt\right\}, \quad \text{where } G, \tilde{G} \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, 1).$$

Further note that for $G > 0$,

$$\int_{-G/4}^{G/4} \frac{t^2 e^{-t^2/2}}{\sqrt{2\pi}} dt \leq \left(\frac{G}{4}\right)^2 \int_{-G/4}^{G/4} \frac{e^{-t^2/2}}{\sqrt{2\pi}} dt.$$

Hence, invoking the definition of γ_\star as in Eq. (18), we find that

$$\mathbb{E}\{\delta_i\} \geq 16\mathbb{E}\left\{\frac{|h'(6\|x_\star\|_2 G)|^2 \mathbb{1}\{G > 0\}}{(|h'(6\|x_\star\|_2 G)|^2 + \gamma_\star)^2} \cdot \int_{-G/4}^{G/4} \frac{t^2 e^{-t^2/2}}{\sqrt{2\pi}} dt\right\} \geq \frac{16\bar{\omega}(x_\star, X)^2}{n}.$$

We next turn to lower bounding $\mathbb{E}\{\eta_i\}$. By definition of η_i (54b),

$$\mathbb{E}\{\eta_i\} = \mathbb{E}\left\{\frac{|h'(6\|x_\star\|_2 G)|^2 \tilde{G}^2 \mathbb{1}\{G > 0\}}{(|h'(6\|x_\star\|_2 G)|^2 + \gamma_\star)^2} \cdot \mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4\}\right\}.$$

Conditioning on G in tandem with the event $G > 0$, we have

$$\mathbb{E}\left\{\tilde{G}^2 \mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4\}\right\} = \int_{-G/4}^{G/4} \frac{t^2 e^{-t^2/2}}{\sqrt{2\pi}} dt.$$

Putting the two pieces together yields

$$\mathbb{E}\{\eta_i\} \geq \mathbb{E}\left\{\frac{|h'(6\|x_\star\|_2 G)|^2 \mathbb{1}\{G > 0\}}{(|h'(6\|x_\star\|_2 G)|^2 + \gamma_\star)^2} \cdot \int_{-G/4}^{G/4} \frac{t^2 e^{-t^2/2}}{\sqrt{2\pi}} dt\right\} \geq \frac{16\bar{\omega}(x_\star, X)^2}{n},$$

where the last step follows from the definition of γ_\star , i.e., Eq. (18). We next turn to prove the concentration of $\omega(x_\star, X)$ (32). To reduce the notational burden, we let $X^\perp = \{P_{x_\star}^\perp x : x \in X\}$ and define $\Psi : \mathbb{R}^d \mapsto \mathbb{R}$ as $\Psi(g) = \sup_{v \in X^\perp} \langle v/\|v\|_2, g \rangle$. Note that by definition (32), we have $\omega(x_\star, X) = \Psi(g)$ for $g \sim \mathcal{N}(0, I_d)$. It is equivalent to prove concentration of the random variable $\Psi(g)$ for $g \sim \mathcal{N}(0, I_d)$. Note that for all g_1, g_2 , we obtain

$$\Psi(g_1) - \Psi(g_2) = \sup_{v \in X^\perp} \left\langle \frac{v}{\|v\|_2}, g_1 \right\rangle - \sup_{v \in X^\perp} \left\langle \frac{v}{\|v\|_2}, g_2 \right\rangle \leq \sup_{v \in X^\perp} \left\langle \frac{v}{\|v\|_2}, g_1 - g_2 \right\rangle \leq \|g_1 - g_2\|_2.$$

Similar arguments show $\Psi(g_2) - \Psi(g_1) \leq \|g_1 - g_2\|_2$ for all $g_1, g_2 \in \mathbb{R}^d$. Therefore, $\Psi(\cdot)$ is a 1-Lipschitz function, so that applying [Wainwright](#) (e.g., 2019, Theorem 2.26) yields

$$\Pr\left\{|\Psi(g) - \mathbb{E}\{\Psi(g)\}| \geq t\right\} \leq 2 \exp(-t^2/2) \quad \text{for all } t > 0.$$

The desired result follows by noting $\omega(x_\star, X) = \Psi(g)$ and $\bar{\omega}(x_\star, X) = \mathbb{E}\{\Psi(g)\}$ for $g \sim \mathcal{N}(0, I_d)$. \square

B.3 Proof of Lemma 3

Note that by definition of $\{\theta_i, \vartheta_i, \xi_i\}$ in Eq. (46),

$$|\theta_i| \leq \sum_{j=1}^W s_j \mu_j \frac{(a_i^\top x_\star)^2}{\|x_\star\|_2^2}, \quad |\vartheta_i| \leq \sum_{j=1}^W s_j \mu_j \tilde{g}_i^2, \quad |\xi_i| \leq \sum_{j=1}^W s_j \mu_j \frac{a_i^\top x_\star}{\|x_\star\|_2} \tilde{g}_i.$$

Since $a_i \sim \mathcal{N}(0, I_d)$ and $\tilde{g}_i \sim \mathcal{N}(0, 1)$, it follows that θ_i , ϑ_i , and ξ_i are sub-exponential random variables, i.e., there exists a universal constant $C > 0$ such that

$$\|\theta_i\|_{\psi_1} \vee \|\vartheta_i\|_{\psi_1} \vee \|\xi_i\|_{\psi_1} \leq C \sum_{j=1}^W s_j \mu_j.$$

Applying Bernstein's inequality ([Vershynin, 2018](#), Theorem 2.8.1), we deduce that there exists a universal constant $C' > 0$ such that with probability at least $1 - 6n^{-10}$,

$$\left| \frac{1}{n} \sum_{i=1}^n \theta_i - \mathbb{E}\{\theta_i\} \right| \vee \left| \frac{1}{n} \sum_{i=1}^n \vartheta_i - \mathbb{E}\{\vartheta_i\} \right| \vee \left| \frac{1}{n} \sum_{i=1}^n \xi_i - \mathbb{E}\{\xi_i\} \right| \leq C' \sum_{j=1}^W s_j \mu_j \sqrt{\frac{\log(n)}{n}}.$$

We claim that there exists a universal and positive constant c such that,

$$\mathbb{E}\{\theta_i\} \geq c \sum_{j=1}^W \frac{s_j \mu_j}{(\mu_j \|x_\star\|_2)^4 \vee 1}, \quad \mathbb{E}\{\vartheta_i\} \geq c \sum_{j=1}^W \frac{s_j \mu_j}{(\mu_j \|x_\star\|_2)^4 \vee 1}, \quad \mathbb{E}\{\xi_i\} = 0,$$

deferring its proof to the end of this section. Combining the inequalities in the display with condition (21) and adjusting constants, we obtain that the deviation $\sqrt{\log(n)/n}$ is of smaller order term than both $\mathbb{E}\{\theta_i\}$ and $\mathbb{E}\{\vartheta_i\}$. We thus conclude that with probability at least $1 - 6n^{-10}$,

$$\frac{1}{n} \sum_{i=1}^n \theta_i \wedge \frac{1}{n} \sum_{i=1}^n \vartheta_i \gtrsim \sum_{j=1}^W \frac{s_j \mu_j}{(\mu_j \|x_\star\|_2)^4 \vee 1} \quad \text{and} \quad \left| \frac{1}{n} \sum_{i=1}^n \xi_i \right| \lesssim \sum_{j=1}^W s_j \mu_j \cdot \sqrt{\frac{\log(n)}{n}}.$$

This concludes the main proof. It remains to establish the claim on $\mathbb{E}\{\theta_i\}$, $\mathbb{E}\{\vartheta_i\}$, and $\mathbb{E}\{\xi_i\}$.

Lower bound on $\mathbb{E}\{\theta_i\}$. By definition (46), we obtain that for $G, \tilde{G} \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, 1)$,

$$\begin{aligned} \mathbb{E}\{\theta_i\} &= \mathbb{E}\left\{ |h'(6\|x_\star\|_2 G)| G^2 \mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4, G > 0\} \right\} \\ &= \mathbb{E}\left\{ |h'(6\|x_\star\|_2 G)| G^2 \mathbb{1}\{G > 0\} \int_{-G/4}^{G/4} \frac{e^{-t^2/2}}{\sqrt{2\pi}} dt \right\} \\ &\geq \mathbb{E}\left\{ |h'(6\|x_\star\|_2 G)| G^2 \mathbb{1}\{G > 0\} \frac{G}{2} \frac{e^{-G^2/32}}{\sqrt{2\pi}} \right\}. \end{aligned}$$

Note that by Eq. (6), $|h'(t)| = \sum_{j=1}^W s_j \mu_j \exp(-\mu_j t)$ for all $t > 0$. Thus, we obtain

$$\begin{aligned} \mathbb{E}\{\theta_i\} &\geq \frac{1}{2\sqrt{2\pi}} \sum_{j=1}^W s_j \mu_j \mathbb{E}\left\{ \exp(-6\mu_j \|x_\star\|_2 G - G^2/32) \mathbb{1}\{G > 0\} G^3 \right\} \\ &= \frac{1}{4\pi} \sum_{j=1}^W s_j \mu_j \int_0^{+\infty} e^{-6\mu_j \|x_\star\|_2 t - t^2/32} \cdot t^3 \cdot e^{-t^2/2} dt \\ &\geq \frac{1}{4\pi} \sum_{j=1}^W s_j \mu_j \int_0^{+\infty} e^{-6\mu_j \|x_\star\|_2 t - t^2} \cdot t^3 dt. \end{aligned} \tag{65}$$

Note that if $\mu_j \|x_\star\|_2 \leq 1$, then

$$\int_0^{+\infty} e^{-6\mu_j \|x_\star\|_2 t - t^2} \cdot t^3 dt \geq \int_0^{+\infty} e^{-6t - t^2} \cdot t^3 dt := c_0,$$

where c_0 is a universal and positive constant. If $\mu_j \|x_\star\|_2 \geq 1$, then

$$\begin{aligned} \int_0^{+\infty} e^{-6\mu_j \|x_\star\|_2 t - t^2} \cdot t^3 dt &\geq \int_0^{(3\mu_j \|x_\star\|_2)^{-1}} e^{-6\mu_j \|x_\star\|_2 t - t^2} \cdot t^3 dt \\ &\stackrel{(i)}{\geq} e^{-3} \int_0^{(3\mu_j \|x_\star\|_2)^{-1}} t^3 dt \gtrsim \frac{1}{\mu_j^4 \|x_\star\|_2^4}, \end{aligned}$$

where in step (i) we use

$$-6\mu_j \|x_\star\|_2 t - t^2 \geq -2 - (3\mu_j \|x_\star\|_2)^{-2} \geq -3, \quad \text{for all } 0 \leq t \leq (3\mu_j \|x_\star\|_2)^{-1} \text{ and } \mu_j \|x_\star\|_2 \geq 1.$$

Putting the two lower bounds together yields that there exists a universal and positive constant c such that

$$\int_0^{+\infty} e^{-6\mu_j \|x_\star\|_2 t - t^2} \cdot t^3 dt \geq \frac{c}{(\mu_j^4 \|x_\star\|_2^4) \vee 1}. \tag{66}$$

Substituting the inequality in the display above into inequality (65) yields

$$\mathbb{E}\{\theta_i\} \geq \frac{c}{4\pi} \sum_{j=1}^W \frac{s_j \mu_j}{(\mu_j \|x_\star\|_2)^4 \vee 1},$$

which concludes the proof.

Lower bound on $\mathbb{E}\{\vartheta_i\}$. By definition (46), we obtain that for $G, \tilde{G} \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, 1)$,

$$\begin{aligned} \mathbb{E}\{\varphi_i\} &= \mathbb{E}\left\{|h'(6\|x_\star\|_2 G)| \tilde{G}^2 \mathbb{1}\{-G/4 \leq \tilde{G} \leq G/4, G > 0\}\right\} \\ &= \mathbb{E}\left\{|h'(6\|x_\star\|_2 G)| \mathbb{1}\{G > 0\} \int_{-G/4}^{G/4} \frac{t^2 e^{-t^2/2}}{\sqrt{2\pi}} dt\right\} \\ &\gtrsim \mathbb{E}\left\{|h'(6\|x_\star\|_2 G)| \mathbb{1}\{G > 0\} e^{-G^2/32} G^3\right\}, \end{aligned}$$

where in the last step we use for $G > 0$

$$\int_{-G/4}^{G/4} \frac{t^2 e^{-t^2/2}}{\sqrt{2\pi}} dt \gtrsim e^{-G^2/32} \int_{-G/4}^{G/4} t^2 dt \gtrsim e^{-G^2/32} G^3.$$

Consequently, we obtain

$$\begin{aligned} \mathbb{E}\{\vartheta_i\} &\gtrsim \sum_{j=1}^W s_j \mu_j \mathbb{E}\left\{\exp\left(-6\mu_j \|x_\star\|_2 G - G^2/32\right) \mathbb{1}\{G > 0\} G^3\right\} \\ &= \sum_{j=1}^W s_j \mu_j \int_0^{+\infty} e^{-6\mu_j \|x_\star\|_2 t - t^2/32} \cdot t^3 \cdot e^{-t^2/2} dt \\ &\geq \sum_{j=1}^W s_j \mu_j \int_0^{+\infty} e^{-6\mu_j \|x_\star\|_2 t - t^2} \cdot t^3 dt \gtrsim \sum_{j=1}^W \frac{s_j \mu_j}{(\mu_j \|x_\star\|_2)^4 \vee 1}, \end{aligned}$$

where in the last step we use inequality (66). This concludes the proof.

Proof that $\mathbb{E}\{\xi_i\} = 0$. By definition (46), we obtain

$$\mathbb{E}\{\xi_i\} = \mathbb{E}\left\{|h'(6\|x_\star\|_2 G)| G \tilde{G} \cdot \mathbb{1}\{G > 0, -G/4 \leq \tilde{G} \leq G/4\}\right\}.$$

Since for any $G > 0$, $\tilde{G} \cdot \mathbb{1}\{G > 0, -G/4 \leq \tilde{G} \leq G/4\}$ is an odd function of \tilde{G} and $G, \tilde{G} \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, 1)$, then by condition on G ,

$$\mathbb{E}\left\{\tilde{G} \cdot \mathbb{1}\{G > 0, -G/4 \leq \tilde{G} \leq G/4\}\right\} = 0.$$

We thus obtain $\mathbb{E}\{\xi_i\} = 0$. □