

# Transforming semitotal colorings into total colorings

Italo J. Dejter  
 University of Puerto Rico  
 Rio Piedras, PR 00936-8377  
[italo.dejter@gmail.com](mailto:italo.dejter@gmail.com)

## Abstract

Let  $2 \leq k \in \mathbb{Z}$ . We focus on how to obtain total colorings of distance transitive cubic graphs and cage graphs by means of a semitotal-to-total coloring procedure transformation.

## 1 Introduction

We engage into the task of transforming semitotal colorings of distance transitive graphs and cage graphs into total colorings of those respective graphs. We pass to define and situate in the literature the necessary concepts of such engagement.

**Definition 1.** Given a simple connected graph  $\Gamma$ , a *total coloring* or *TC* of  $\Gamma$  is a color assignment to the vertices and edges of  $\Gamma$  such that no two incident or adjacent elements (vertices or edges) are assigned the same color.

A recent survey [18] contains an updated bibliography on TCs, with the TC Conjecture, posed independently by Behzad [1, 2] and by Vizing [27], that asserts that the total chromatic number of  $\Gamma$  (namely, the least number of colors required by a TC of  $\Gamma$ ) be either  $\Delta(\Gamma) + 1$  or  $\Delta(\Gamma) + 2$ , where  $\Delta$  is the largest degree of any vertex of  $\Gamma$ .

The TC Conjecture was established for cubic graphs [15, 7, 24, 26], meaning that the total chromatic number of cubic graphs is either 4 or 5. To decide whether a cubic graph  $\Gamma$  has total chromatic number  $\Delta(\Gamma) + 1$ , even for bipartite cubic graphs, is NP-hard [25].

Williams and Holroyd [28] released one of the conditions of a total coloring of a simple connected graph  $G$  with maximum degree  $\Delta = \Delta(G)$  by defining a *semitotal graph coloring* of  $G$  as a color assignment to the vertices and edges of  $G$  that uses  $\Delta + 1$  colors and has the properties of a total coloring, except that adjacent vertices need not have distinct colors.

**Definition 2.** A *semitotal coloring* of  $G$  is a function  $\mu : V(G) \cup E(G) \rightarrow \{1, 2, \dots, \Delta + 1\}$  such that any two adjacent edges  $e_1, e_2 \in E(G)$  have  $\mu(e_1) \neq \mu(e_2)$  and every  $v \in V(G)$  has  $\mu(v) \neq \mu(e), \forall e \in E(G)$  incident to  $v$ .

**Definition 3.** A  $\beta$ -edge of  $G$  with respect to  $\mu$  is an edge  $e = v_1v_2$  such that  $\mu(v_1) = \mu(v_2)$ .

**Definition 4.** Let  $\beta(\mu)$  be the number of  $\beta$ -edges with respect to  $\mu$ . Then, the *beta parameter*  $\beta(\mu)$  is defined as the minimum of the numbers  $\beta(\mu)$  such that  $\mu$  is a semitotal coloring of  $G$ .

From [29] we now that a cycle  $C_n$  is of type 1 if and only if its order  $n$  is a multiple of 3, while a complete graph  $K_n$  of order  $n$  is of type 1 if and only if  $n$  is odd. Furthermore, [28] shows that if  $n \equiv 1$  or  $2 \pmod{3}$ , then  $\beta(C_n) = 2$ . It further shows that if  $n$  is even, then  $\beta(K_n) = n/2$ .

## 2 Main result

In this paper, a selective transformation procedure is considered that takes a semitotal coloring of  $G$  into one or more singular steps consisting each into iteratively modifying maximum paths that consist of two alternate edge colors  $c_1$  and  $c_2$ . Being alternate in those two colors, each such maximum path has to have their initial and terminal end-vertices being in  $\{c_1, c_2\}$ , with either  $c_1 = c_2$  or  $c_1 \neq c_2$ . The way in which such transformation procedure is done may change a semitotal coloring that is not total into a total coloring and should be taken selectively, unless  $\beta(\mu) > 0$ .




Figure 1: Applications to the Heawood and Pappus graphs.

**Theorem 5.** Let  $G$  be a simple connected graph with maximum vertex degree  $\Delta = \Delta(G)$ . Let  $\mu : V(G) \cup E(G) \rightarrow \{1, 2, \dots, \Delta + 1\}$  be a semitotal coloring. Let  $v_1, v_2 \in V(G)$ ,  $e_1, \dots, e_k \in E(G)$  and let  $S = \{v_1, e_1, e_2, \dots, e_k, v_2\}$  be a maximum sequence of contiguous incident or adjacent elements (meaning  $v_1$  endvertex of  $e_1$ ,  $v_2$  endvertex of  $e_k$ , and  $e_i$  adjacent

to  $e_{i+1}, \forall i = 1, 2, \dots, k-1$ ) with  $\mu(S) = \{c_1, c_2\}$ ,  $c_1 = \mu(v_1)$ ,  $c_2 \neq c_1$ . Let  $\mu(e_{2j-1}) = c_2$  and  $\mu(e_{2j}) = c_1$  ( $1 \leq j \leq \lfloor k/2 \rfloor$ ), with  $\mu(v_2) \neq \mu(e_k)$ . Then there is a semitotal coloring  $\mu' : V(G) \cup E(G) \rightarrow \{1, 2, \dots, \Delta + 1\}$  with  $\mu(S) = \{c_1, c_2\}$ ,  $\mu'(v_1) = c_2$  and  $\mu'(v_1) = \mu'(e_\ell)$  if and only if  $\ell = 2j$  for some  $j$  such that  $1 \leq j \leq \lfloor k/2 \rfloor$ . By saying that  $\mu'$  is a  $\beta$ -reduction if  $v_1$  is an endvertex of a  $\beta$ -edge of  $\mu$  in  $G$  with  $\beta(\mu') < \beta(\mu)$ , a finite number of  $\beta$ -reductions  $\mu, \mu', \mu^{(2)}, \dots, \mu^{(t)}$  with  $\beta(u) > \beta(\mu') > \beta(\mu^{(2)}) > \dots > \beta(\mu^{(t)})$  leads to a total coloring of  $G$  if and only if  $\beta(\mu^{(t)}) = 0$ .

*Proof.* There are two cases. First, assume that  $k$  is odd. W.l.o.g., assume  $k = 5$ . Then:




Figure 2: Application to the Desargues graph.

$G$	$v_1$	$e_1$	$e_2$	$e_3,$	$e_4$	$e_5$	$v_2$	(1)
$\mu$	$c_1$	$c_2$	$c_1$	$c_2$	$c_1$	$c_2$	$c_1$	
$\mu'$	$c_2$	$c_1$	$c_2$	$c_1$	$c_2$	$c_1$	$c_2$	

Second, assume that  $k$  is even. Wl.o.g., assume  $k = 4$ . Then,

$G$	$v_1$	$e_1$	$e_2$	$e_3,$	$e_4$	$v_2$	(2)
$\mu$	$c_1$	$c_2$	$c_1$	$c_2$	$c_1$	$c_2$	
$\mu'$	$c_2$	$c_1$	$c_2$	$c_1$	$c_2$	$c_1$	

In either case, if  $v_1$  is the endvertex of a  $\beta$ -edge of  $\mu$ , then it is not an endvertex of a  $\beta$ -edge of  $\mu'$ . By a selective finite iteration as said in the statement, a total coloring maybe at our disposition. □




Figure 3: Foster 90-vertex graph and  $K_{3,3}$ .

### 3 Distance transitive graphs

Examples of applications of Theorem 5 to distance transitive graphs in this section are followed by additional examples for cage graphs in Section 5. Those cases of cubic hamiltonian graphs may use the LCF notation for their expression [6], [16]. In the cases of distance transitive graphs, their intersection arrays [4] will be mentioned.

**Example 6.** For a case of completeness, we precede examples of application of Theorem 5 with the case of total coloring the 3-cube  $Q_3$ , the starting point of the treatment in [8], namely,

$$\begin{array}{cccccc|cccc} 0 & \bar{3} & 1 & \bar{0} & 2 & \bar{1} & 3 & \bar{2} & 0 & & 0 & \bar{2} & 1 & \bar{3} & 2 & \bar{0} & 3 & \bar{1} & 0 \\ 1 & | & 2 & | & 3 & | & 0 & | & 1 & & 3 & | & 0 & | & 1 & | & 2 & | & 3 & | \\ 2 & \bar{0} & 3 & \bar{1} & 0 & \bar{2} & 1 & \bar{3} & 2 & & 2 & \bar{1} & 3 & \bar{2} & 0 & \bar{3} & 1 & \bar{0} & 2 \end{array} \neq \quad (3)$$

in two different ways via the two shown cutouts, with  $Q_3$  recoverable by gluing in parallel the leftmost and rightmost vertical segments in the cutouts; these separated by the inequality symbol  $\neq$ . The vertices and edges in either cutout are given via large and small (color) numbers, respectively, with the small numbers accompanying horizontal and vertical segments indicating the edges. The vertex colors are similar in both cutouts but the edge colors differ, so these edge colorings are orthogonal. Antipodal vertices of  $Q_3$ , differing 3 in distance, receive the same color number in either total coloring. In addition, we mention that the intersection array of  $Q_3$  is  $\{3, 2, 1; 1, 2, 3\}$  and that, being  $Q_3$  hamiltonian, its LCF notation is  $[-3, 3]^4$ .

**Example 7.** The distance transitive Heawood graph *Hea*, whose intersection array is  $\{3, 2, 2; 1, 1, 3\}$ , represented on the left of Figure 1 with LCF notation  $[5, -5]^7$  for an external Hamilton cycle  $H$  and an internal 1-factor  $F$ , has a total coloring obtained from semitotal colorings by a succession of transformations, each starting at a certain vertex  $v$ , then changing colors along an alternate path of edge colors and ending in another vertex  $w \neq v$ . In the figure, *Hea* is represented joining  $F$  with edges colored green = 3 with  $H$  colored clockwise from the upper-left vertex with colored cycle

$$C = (1_0 2_1 0_2 0_1 2_0 1_2 0_1 2_0 1_2 0_1 2_0 1_2 0_1 0_2), \quad (4)$$

where 0=hazel, 1=red and 2=blue. Then, with initial vertex being the one corresponding to the third vertex color (0 = hazel) in  $H$ , we have the alternatively colored maximum path  $P$  formed by initial and terminal vertex colors delimiting successive alternate edge colors being hazel = 0 and green = 3:

$$P = 0_3 1_0 2_3 1_0 2_3 0. \quad (5)$$

This path  $P$  is then is transformed into the alternate path:

$$P' = 3_0 1_3 2_0 1_3 2_0 3, \quad (6)$$

that makes the initial semitotal coloring into a total coloring.

**Definition 8.** A semitotal or total coloring  $\mu : V(G) \cup E(G) \rightarrow \{1, 2, \dots, \Delta + 1\}$  of a connected and simple graph  $G$  with largest degree  $\Delta$  is said to be *special* if there exists a color  $c \in \{1, 2, \dots, \Delta + 1\}$  such that no  $v \in V(G)$  has  $\mu(v) = c$ .

**Example 9.** The distance transitive Pappus graph  $Pap$ , whose intersection array is  $\{3, 2, 2, 1; 1, 2, 3\}$ , represented on the right of Figure 1 with LCF notation  $[5, 7, -7, 7, -7, -5]^3$  via an external Hamilton cycle  $H$  and an internal 1-factor  $F$ , shows  $Pap$  with a total coloring that uses the same color for all interior edges and so uses only the remaining colors for the vertices and edges of the external Hamilton cycle  $H$ , yielding the counterclockwise colored cycle starting at any vertex colored hazel = 0 (because of periodicity mod 3 since  $H$  has length 18 divisible by 3):

$$C = (0_1 2_0 1_2 0_1 2_0 1_2 0_1 2_0 1_2 0_1 2_0 1_2 0_1 2_0 1_2) = (0_1 2_0 1_2)^4. \quad (7)$$

Thus,  $Pap$  has a special total coloring in which one color is missing on vertices, namely color green = 3.




Figure 4: Coxeter graph.

**Example 10.** The 90-vertex Foster graph  $Fos$ , represented on the left of Figure 3 with LCF notation  $[17, -9, 37, -37, 9, -17]^{15}$  for an external Hamilton cycle  $H$  and an internal 1-factor  $F$ , has a special total coloring whose associated colored cycle is:

$$\mu(H) = (0_1 2_0 1_2)^{30}. \quad (8)$$

This is a distance transitive graph with intersection array  $3, 2, 2, 2, 2, 1, 1, 1; 1, 1, 1, 1, 2, 2, 2, 3$ ].

**Example 11.** The distance transitive Desargues graph  $Des$ , whose intersection array is  $\{3, 2, 2, 1, 1, 1; 1, 1, 2, 2, 3\}$ , with LCF notation  $[5, -5, 9, -9]^5$  for an external Hamilton cycle  $H$  and an internal 1-factor  $F$ , appears fourthfold in Figure 2, has effectively from left to right and then from top to bottom four semitotal colorings related via transformations as in Theorem 5 and as exemplified above Example 7 in the case of the Heawood graph  $Hea$ , with the lower-right final drawing showing effectively a total coloring. Here and elsewhere, dashed or thin edges are meant as edges with the same color on their respective endvertices, that is the  $\beta$ -edges. In fact, from semitotal to total coloring in  $Des$  is shown from its upper-left depiction in the figure to its lower-right one in three applications of Theorem 5: departing from the upper-left depiction a change of color at onr vertex carries a change of edge colors as shown. Second, the oriented vertex-and-edge path 0310230 in the upper-right depiction is transformed into the similar path 3013203 in the lower-left depiction with the oriented endvertex-and-edge colored path 20120 transformed into the similar path 02102.




Figure 5: Total colorings of  $Pet$  and its pull-back  $Dod$ .

Here is the presentation of the colored 6-cycles of these Heawood, Pappus and Desargues graphs in the three following successive tables:

$$\begin{array}{cccc}
 (1_2 0_1 3_2 1_0 2_1 3_0) & (1_0 2_3 0_1 2_0 1_2 0_3) & (3_2 1_0 2_1 3_2 0_1 2_0) & (3_2 0_3 1_2 0_1 2_3 1_0) \\
 (2_1 3_2 0_1 2_3 1_2 0_3) & (2_3 1_0 2_3 1_2 0_1 3_0) & (0_1 2_3 1_2 0_1 2_0 1_3) & (0_1 2_3 0_1 3_2 1_0 2_3) \\
 (2_0 1_2 0_1 2_3 1_2 0_3) & (1_2 0_3 1_0 2_1 3_2 0_3) & (0_1 2_3 1_2 0_1 3_2 1_3) & (2_3 1_0 3_2 0_1 2_3 1_0) \\
 (1_2 0_1 2_0 1_2 0_1 2_0) & (0_1 3_0 2_3 1_2 0_1 2_3) & & 
 \end{array} \tag{9}$$

$$\begin{array}{cccccc}
 (0_1 2_0 1_2 0_1 2_0 1_3) & (1_2 0_3 2_0 1_2 0_1 2_3) & (0_3 1_2 0_3 1_2 0_3 1_2) & (0_2 1_0 2_1 0_3 2_1 0_3) & (2_0 1_3 0_2 1_0 2_1 0_3) \\
 (0_1 2_0 1_2 0_1 2_1 0_3) & (1_2 0_3 2_0 1_2 0_1 2_3) & (1_0 2_3 1_0 2_3 1_0 2_3) & (1_0 2_1 0_2 1_3 0_1 2_3) & (2_0 1_3 0_2 1_3 2_0 1_3) \\
 (0_1 2_0 1_2 0_1 2_0 1_3) & (1_2 0_3 2_0 1_2 0_1 2_3) & (0_1 2_3 0_1 2_3 0_1 2_3) & (2_1 0_3 1_2 0_1 2_1 0_3) & (1_0 2_3 0_1 2_0 1_2 0_3)
 \end{array} \tag{10}$$

$$\begin{array}{ccccc}
(0_3 2_0 3_1 2_3 1_0 3_2) & (1_0 3_1 0_2 1_0 2_1 3_2) & (2_1 3_0 1_3 0_1 2_0 1_3) & (2_0 1_2 3 1 2_0 3_2 0_3) & (3_2 0_1 2_0 3_1 0_3 2_0) \\
(2_0 3_1 0_2 3_0 0_2 1_3) & (3_1 2_3 1_2 3_0 1_3 0_2) & (0_2 1_0 2_3 1_2 3_1 2_3) & (1_3 0_1 2_3 0_1 2_0 3_2) & (3_1 2_0 3_1 2_0 3_1 2_0) \\
(3_1 2_0 3_2 1_0 2_1 0_2) & (0_2 1_3 2_1 0_2 3_0 2_3) & (1_3 0_2 3_0 2_3 0_1 3_2) & (3_1 2_3 0_1 3_0 1_3 2_0) & (2_0 3_2 1_0 3_1 2_0 1_3) \\
(1_0 3_2 0_1 3_2 1_0 3_2) & (2_1 3_2 1_3 2_0 3_2 1_3) & (2_0 1_3 2_0 1_3 2_1 0_3) & (3_2 0_3 2_1 0_2 3_0 2_1) & (0_3 2_1 3_0 2_3 0_1 3_2)
\end{array} \tag{11}$$

**Example 12.** The complete bipartite graph  $K_{3,3}$  has a special semitotal coloring  $\mu$  depicted and colored as on the right side of Figure 3, with  $\beta(\mu) = 35602$ .




Figure 6: Biggs-Smith graph

**Example 13.** The 28-vertex Coxeter graph  $Cox$  is shown in Figure 4 with  $\beta(Cox) \leq 3$ , where the yellow areas are delimited by 9-cycles forming a 2-factor of  $Cox \setminus \{v\}$ , where  $v$  is the central vertex of this representation of  $Cox$ .



**Theorem 14.** Given a 2-fold covering graph map  $\phi : G \rightarrow G'$  of simple connected graphs  $G$  and  $G'$  with common maximum  $\Delta$  and given a semitotal or total coloring  $\mu' : G' \rightarrow \{1, 2, \dots, \Delta + 1\}$ , then there exists a corresponding semitotal or total coloring  $\mu : G \rightarrow \{1, 2, \dots, \Delta + 1\}$  behaving as a lifting of  $\mu'$ .

*Proof.* Consider the inverse image  $v$  or  $e$  of any vertex  $v'$  or edge  $e'$  of  $G'$  and assign the color  $\mu(v)$  or  $\mu(e)$  to  $v$  or  $e$ , respectively, expressed as  $\mu(v) = \mu(v')$  or  $\mu(e) = \mu(e')$ .  $\square$




Figure 7: Application to the Petersen and Robertson graphs.

**Example 15.** Figure 5 shows an example of Theorem 14 that produces a total coloring of the dodecahedral graph  $Dod$  as a pull-back of a total coloring of the Petersen graph  $Pet$ . See also example 20, below. The intersection array of  $Pet$  is  $\{3, 2; 1, 1\}$ . It is the only distance transitive cubic graph which is a snark. The intersection array of  $Dod$  is  $\{3, 2, 111; 111, 2, 3\}$ . Not used in the present application of Theorem 5 though, the LCF of  $Dod$  is  $[10, 7, 4, -4, -7, 10, -4, 7, -7, 4]$ .

**Example 16.** The Biggs-Smith graph  $BS$  is represented in Figure 6 as the union of an external Hamilton cycle  $H$  with the vertices of  $V(BS)$  denoted by the numeric order in which they are suggested to appear in  $H$  in the figure and an enclosed 1-factor  $F$ , where a special semitotal coloring  $\mu$  is indicated with the edges of  $F$  in green and the vertices

and edges of  $H$  alternatively colored red, blue and gray in such a way that all vertices in  $V(BG) = \{0, \dots, 101\}$  divisible by 3 have red color, expressed in the figure by citing their corresponding number. Then, the  $\beta$ -edges of  $\mu$  are those green edges the absolute value of the difference of whose endvertex numeric orders are divisible by 3. Thus,  $\beta(BS) \leq 17$ .

Note that Figure 6 is a representation of  $BS$  depending of its LCF notation:

[16, 24, -38, 17, 34, 48, -19, 41, -35, 47, -20, 34, -36, 21, 14, 48, -16, -36, -43, 28, -17, 21, 29, -43, 46, -24, 28, -38, -14, -50, -45, 21, 8, 27, -21, 20, -37, 39, -34, -44, -8, 38, -21, 25, 15, -34, 18, -28, -41, 36, 8, -29, -21, -48, -28, -20, -47, 14, -8, -15, -27, 38, 24, -48, -18, 25, 38, 31, -25, 24, -46, -14, 28, 11, 21, 35, -39, 43, 36, -38, 14, 50, 43, 36, -11, -36, -24, 45, 8, 19, -25, 38, 20, -24, -14, -21, -8, 44, -31, -38, -28, 37]

This in turn yield a list of the edges of 1-factor  $F$ , namely:

(0, 16) (1, 25) (2, 66) (3, 20) (4, 38) (5, 53) (6, 89) (7, 48) (8, 75) (9, 56) (10, 92)  
(11, 45) (12, 78) (13, 34) (14, 28) (15, 63) (16, 0) (17, 83) (18, 77) (19, 47) (20, 3) (21, 42)  
(22, 51) (23, 82) (24, 70) (25, 1) (26, 54) (27, 91) (28, 14) (29, 81) (30, 87) (31, 52) (32, 40)  
(33, 60) (34, 13) (35, 55) (36, 101) (37, 76) (38, 4) (39, 97) (40, 32) (41, 79) (42, 21) (43, 68)  
(44, 59) (45, 11) (46, 64) (47, 19) (48, 7) (49, 85) (50, 58) (51, 22) (52, 31) (53, 5) (54, 26)  
(55, 35) (56, 9) (57, 71) (58, 50) (59, 44) (60, 33) (61, 99) (62, 86) (63, 15) (64, 46) (65, 90)  
(66, 2) (67, 98) (68, 43) (69, 93) (70, 24) (71, 57) (72, 100) (73, 84) (74, 95) (75, 8) (76, 47)  
(77, 18) (78, 12) (79, 41) (80, 94) (81, 29) (82, 23) (83, 17) (84, 73) (85, 49) (86, 62) (87, 30)  
(88, 96] (89, 6) (90, 65) (91, 27) (92, 10) (93, 69) (94, 80) (95, 74) (96, 88) (97, 53) (98, 67)  
(99, 61) (100, 72) (101, 63)

which is represented in Figure 6. Of these green edges, those in thin trace (in contrast with those in thick trace), are the  $\beta$ -edges, namely:

*blue* : (1, 25), (13, 34), (22, 51), (31, 52), (37, 76), (46, 64), (49, 85);  
*gray* : ((5, 53), (17, 83), (62, 86), (65, 90), (74, 95);  
*red* : (12, 78), (15, 63), (30, 82), (33, 60), (69, 93).

## 4 Relation to efficient total colorings of regular graphs

In [8], we studied the relations between total colorings of  $k$ -regular graphs of girth  $k + 1$  ( $1 < k \in \mathbb{Z}$ ) and efficient dominating sets (EDSs) [10, 13, 11, 14, 17, 22]. In [12], we treated similar relations but for the girth  $2k$  in the place of the girth  $k + 1$  of [8]. The following definition combining both the concepts of total coloring and efficient domination, given in [8], is taken into consideration having being central both in [8] and [12]. None of the graphs considered in the present paper has an efficient total coloring, as can be seen in all those cases presented above.

**Definition 17.** [8] A coloring of a connected  $k$ -regular simple graph  $\Gamma$  ( $2 \leq k \in \mathbb{Z}$ ) is said to be an *efficient TC*, or *ETC*, (and  $\Gamma$  said to be *ETCed*), if:

- (a) (TC condition) each  $v \in V(\Gamma)$  together with its neighbors are assigned all the colors in  $[k + 1] = \{0, 1, \dots, k\}$  via a bijection  $N[v] = N(v) \cup \{v\} \leftrightarrow [k + 1]$ , where  $N[v]$  and  $N(v)$  are the *closed neighborhood* of  $v$  and the *open neighborhood* of  $v$ , respectively [11];




Figure 8: Semitotal colorings for the 30-vertex Tutte-Coxeter graph, and 2 for the 42-vertex cage; showing below 3  $\beta$ -induced Heawood and 7 non- $\beta$ -induced 6-subgraphs.

- (b) the TC in item (a) partitions  $V(\Gamma)$  into  $k + 1$  *efficient dominating sets* (*EDS*), also called *perfect codes*, namely independent (stable) subsets  $S_i \subseteq V(\Gamma)$  such that for any  $v \in V(\Gamma) \setminus S_i$ ,  $|N[v] \cap S_i| = 1$ , where  $i$  varies in  $[k + 1]$ , [10, 13, 11, 14, 17, 22].

Under conditions (a)-(b), it is seen that the total chromatic number of  $\Gamma$  is  $\Delta(\Gamma) + 1$ .

**Example 18.** For  $1 < j \in \mathbb{Z}$ , consider the  $6j$ -cycle graph  $C_{6j} = (v_1, e_1, v_2, e_2, \dots, e_{6j-1}, v_{6j}, e_{6j})$ , where  $e_i$  is the edge with end-vertices  $v_i$  and  $v_{i+1}$ , for  $1 \leq i \leq 6j - 1$ , and  $e_{6j}$  is the edge with end-vertices  $v_{6j}$  and  $v_1$ . Let the successive vertices and edges of  $C_{6j}$  be colored together in one fixed direction of  $C_{6j}$  in the order of the colors in the color cycle  $(0, 1, 2, 0, 1, 2, \dots, 0, 1, 2)$ . This clearly yields an ETC of  $C_{6j}$ .

In the rest of [8] and [12], respectively, total coloring of finite connected simple finite connected simple  $k$ -regular graphs  $\Gamma$  ( $k \geq 2$ ) of girths  $k + 1$  and  $2k$  were dealt with. Our purpose there was to determine ETCs of such graphs.

Some of these ETCs yield *edge-half-girth colorings*, (an adaptation of the edge-girth colorings of [8], see Definition 19, below) on the prism  $\Gamma \square K_2$ , where  $K_2 = P_2$  is the complete graph on two vertices, that is the path  $P_k$  consisting of  $k = 2$  vertices.

**Definition 19.** Let  $\Gamma$  be a finite connected  $k$ -regular simple graph of girth  $k + 1$ . An *edge-girth coloring*, or *EGC*, of  $\Gamma$  is a proper edge coloring via  $k + 1$  colors, each girth cycle colored with  $k$  colors, each color used precisely once.

With respect to the examples of application of Theorem 5 in the present work, we observe the following:

1. The Heawood graph *Hea* has a total coloring.
2. The Pappus graph *Pap* has a special total coloring.
3. The 90-vertex Foster graph *Fos* has a special total coloring.
4.  $K_{3,3}$  has a semitotal coloring  $\mu$  with  $\beta(\mu) = 3$ .
5. The Desargues graph *Des* has a total coloring.
6. The Tutte-Coxeter graph has semitotal coloring  $\mu$  with  $\beta(\mu) \leq 5$ , (in Section 5).
7. The 6-cage has a semitotal coloring  $\mu$  with  $\beta(\mu) \leq 21$ , (in Section 5).

## 5 Cage graphs

As for cage graphs, we already treated  $K_{3,3}$ , the Heawood graph and the Petersen graphs in Examples 12, 7, 9 and 20, namely the (3, 4)-, (3, 6)- and (3, 5)-cages on respective 6, 14 and 10 vertices. In addition, we consider the following cage graphs.

**Example 20.** We apply Theorem 5 to the Petersen graph  $Pet$  of 10 vertices and the Robertson graph  $Rob$  of 19 vertices, see Figure 7. These graphs are the  $(3, 5)$ -cage and the  $(4, 5)$ -cage, respectively. To exemplify Theorem 5 for  $Pet$ , let us use its representation as in [23, Figure 1] but colored semitotal as in the upper-left of the figure. This semitotal coloring of  $Pet$  on the upper-left of the figure has just one  $\beta$ -edge, which is in dashed green in contrast to the other edges of  $Pet$ , in thick trace. A single application of Theorem 5 yields the total coloring of  $Pet$  in the upper right of the figure, performed by exchanging the colors 0 (hazel) and 1 (red) along the endvertices of the alternately recoloring path and its edges, that produces the oriented vertex colored path 0323231; this is then transformed into the oriented vertex color path 1323230. These paths seen as vertex-and-edge colored paths 01302130212301 and 1031203120310, respectively. Similarly,  $Rob$  is shown in the lower-left of the figure with a semitotal coloring having just two  $\beta$ -edges. This is transformed into a total coloring of  $Rob$  by exchanging the colors 3 c(green) and 1 (red) along the vertex and edge path 132103413 and making it into 312301431.

**Example 21.** The Tutte 8-cage [19] or Tutte-Coxeter graph  $TuCo$  [5] is a cubic graph on 30 vertices and 45 edges which is the Levi graph of the Cremona-Richmond configuration. It is the only unique  $(3, 8)$ -cage graph and Moore graph.  $TuCo$  is hamiltonian with LCF notation  $[-13, -9, 7, -7, 9, 13]$  and is represented in the upper-left of Figure 8 figuring a semitotal coloring that shows that its  $\beta$ -parameter is equal to or less than 5. By the way, the intersection array of TuCo is  $\{3, 2, 2, 1; 1, 1, 1, 2\}$ .

**Example 22.** Consider the  $(5, 6)$ -cage  $\square_5^6$  on 42 vertices, depicted on the upper-right, lower-left and lower-right of Figure 8. Then,  $\square_5^6$  is hamiltonian with extended LCF notation  $[(-11, -15, 7), (11, 15, -7)]^7$ . In the upper-right of the figure there is a special semitotal coloring  $\mu_{\square_5^6}$  of  $\square_5^6$ , with vertex perodic colors being just 0,1,2, selected from the palette of six colors available in this case, namely 0 = black, 1 = red, 2 = blue, 3 = green, 4 = hazel and 5 = violet. The edges with color 3 = green are the only  $\beta$ -edges in this case and they say that  $\beta(\mu_{\square_5^6}) \leq 21$ .

Another interesting semitotal coloring  $\mu'_{\square_5^6}$  of  $\square_5^6$  is shown with the (thick) edge set of  $\square_5^6$  partitioned into those edges in the lower-left and those in the lower-right of the figure, where the vertices are maintained in their position as in the upper-right depiction of  $\square_5^6$ . In the lower left, there is the disjoint union of three copies of the Heawood graph which are formed from all the  $\beta$ -edges of  $\mu'_{\square_5^6}$ . In the lower right, there is the disjoint union of seven copies of the 6-cycle  $C_6$ .

**Question 23.** Are there any other semitotal colorings of regular graphs having the  $\beta$ -edges forming copies of interesting subgraphs, as in the second part of Example 22?

## References

- [1] M. Behzad, *Graphs and their chromatic numbers*, PhD thesis, Michigan State University, 1965.
- [2] , M. Behzad, *The total chromatic number*, Proc. Conf. Combin. Math. and Appl. (1969), 1–8.

- [3] N. L. Biggs and D. H., *On trivalent graphs*, Bulletin of the London Mathematical Society, **3** (1971), 155–158.
- [4] A. E. Brouwer, I. J. Dejter and C. Thomassen, *Highly symmetric subgraphs of hypercubes*, **2** (1993), 25–29.
- [5] A. E. Brouwer, A. M. Cohen and A. Neumaier, *Distance Regular Graphs*, Springer-Verlag, Berlin 1989.
- [6] H. S. M. Coxeter, R. Frucht and D. L. Powers, *Zero-symmetric graphs, trivalent graphical regular representations of groups*, Academic Press, New York, 1981.
- [7] S. Dantas, C. M. H. de Figueiredo, G. Mazzuocollo, M. Preissmann, V. F. dos Santos, D. Sasaki, *On the total coloring of generalized Petersen graphs*, Discrete Math., **339** (2016), 1471–1475.
- [8] I. J. Dejter, *Total coloring of regular graphs of girth = degree +1*, Ars Combinatoria.
- [9] I. J. Dejter, *Equitable factorizations of Hamming shells*, Discrete Mathematics, **261** (2003), no. 1-3, 177–187.
- [10] I. J. Dejter, *Worst-case efficient dominating sets in digraphs*, Discrete Appl. Math, **161** (2003), 944–952. .
- [11] I. J. Dejter and O. Serra, *Efficient dominating sets in Cayley graphs*, Discrete Appl. Math., **129** (2003), 319–328.
- [12] I. J. Dejter, *Total coloring of  $k$ -regular graphs of girth  $2k$* , preprint.
- [13] I. J. Dejter and O. Tomaiconza, *Nonexistence of efficient dominating sets in the Cayley graphs generated by transposition trees of diameter 3*, Discrete Appl. Math, **232** (2017), 116–124.
- [14] Y-P. Deng, Y.Q. Sun, Q. Liu and H-C. Wang, *Efficient dominating sets in circulant graphs*, Discrete Mathematics, **340** (2017), 1503–1507.
- [15] Y. Feng, W. Lin, *A concise proof for total coloring subcubic graphs*, Inform. Process. Lett., **113** (2013), 664–665.
- [16] R. Frucht, *A canonical representation of trivalent hamiltonian graphs*, J. Graph Th., **(1)** (1976), 45–60.
- [17] T. Haynes, S. T. Hedetniemi and M. A. Henning, *Efficient Domination in Graphs*, in *Domination in Graphs: Core Concepts*, Springer Monographs in Mathematics, Springer Cham [https://doi.org/10.1007/978-3-031-09496-5\\_9](https://doi.org/10.1007/978-3-031-09496-5_9).
- [18] J. Geetha, N. Narayanan and K. Somasundaram, *Total colorings-a survey*, AKCE Int. Jour. of Graphs and Combin., **20**, (2023), issue 3. 339–351.
- [19] C. Godsil and G. Royle, *Algebraic Graph Theory*, Springer-Verlag, 2001.

- [20] P. Gregor, A. Merino and T. Mütze, *Star transpositions Gray codes for multiset permutations*, J. of Graph Theory, **103(2)**, (2023), 212–270.
- [21] M. Klin, J. Lauri and M. Ziv-Av, *Links between two semisymmetric graphs on 112 vertices through the lens of association schemes*, Jour. Symbolic Comput., **47-10** (2012), 1175-1191.
- [22] M. Knor and P. Potočnik, *Efficient domination in vertex-transitive graphs*, Eur. Jour. Combin., **33** (2012), 1755–1764.
- [23] A. Orbanic, T. Pisanski, M. Randić and B. Servatius, *Blanusa double*, Mathematical Communications, **9(1)**, 91–103.
- [24] M. Rosenfeld, *On the total chromatic number of a graph*, Israel J. Math., **9** (1971), 396–402.
- [25] A. Sánchez-Arroyo, *Determining the total coloring number is NP-hard*, Discrete Math., **78** (1979), 315–319.
- [26] N. Vijayaditya, *On total chromatic number of a graph*, J. London Math. Soc., **2** (1971), 405–408.
- [27] V. G. Vizing, *On an estimate of the chromatic class of a  $p$ -graph*, Discret Analiz, **3** (1969), 25–30.
- [28] J. Williams, F. Holroyd, *semitotal graph colorings, the beta parametr and total chromatic number*, Discrete Mathematics, **(208)**(2008), 950–954.
- [29] H.-P. Yap, *Total colourings of graphs*, Lecture Notes in Mathematics, vol. 1623, Springer/Verlag, Berlin/Heidelberg, 1996.