

# SINGULAR SPDEs WITH THE CAUCHY-RIEMANN OPERATOR ON A TORUS

ZDZISŁAW BRZEŹNIAK, MIKHAIL NEKLYUDOV, AND EVELINA SHAMAROVA

**ABSTRACT.** We prove the existence of solution to the following  $\mathbb{C}^3$ -valued singular SPDE on the 2D torus  $\mathbb{T}^2$ :

$$(CR) \quad \partial_{\bar{z}} r = r \times \bar{r} + i\gamma \mathcal{W},$$

where  $\partial_{\bar{z}} := \frac{1}{2}(\partial_x + i\partial_y)$  is the Cauchy-Riemann operator on  $\mathbb{T}^2$ ,  $\mathcal{W} = (\mathcal{W}_1, \mathcal{W}_2, \mathcal{W}_3)$  is a real 3D white noise on  $\mathbb{T}^2$  whose component  $\mathcal{W}_3$  has zero mean over  $\mathbb{T}^2$ ,  $\gamma := (\gamma_1, \gamma_2, \gamma_3)$  is an  $\mathbb{R}^3$ -vector and  $\gamma \mathcal{W} := (\gamma_1 \mathcal{W}_1, \gamma_2 \mathcal{W}_2, \gamma_3 \mathcal{W}_3)$ .

**Keywords:** Singular SPDE; Cauchy-Riemann operator; White noise; 2D torus; Hölder-Zygmund space; Cross-product nonlinearity; Young product

**2020 MSC:** 60H15, 35J46

## 1. INTRODUCTION

**1.1. Motivation.** We start by noticing that the Cauchy-Riemann operator  $\partial_{\bar{z}} := \frac{1}{2}(\partial_x + i\partial_y)$  and its complex conjugate  $\partial_z := \frac{1}{2}(\partial_x - i\partial_y)$  are building blocks of the 2D free Dirac operator

$$(1.1) \quad \mathcal{D} := -2i \begin{pmatrix} 0 & \partial_z \\ \partial_{\bar{z}} & 0 \end{pmatrix} = -i\sigma_1 \partial_x - i\sigma_2 \partial_y,$$

where  $\sigma_1 := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  and  $\sigma_2 := \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$  are the Pauli matrices, see, e.g. [5]. Since equation (CR) can be rewritten as  $\partial_{\bar{z}} \bar{r} = -r \times \bar{r} - i\gamma \mathcal{W}$ , by using the free Dirac operator (1.1), we see that equation (CR) can be written in the following equivalent form

$$(1.2) \quad \mathcal{D} \begin{pmatrix} \bar{r} \\ r \end{pmatrix} = \begin{pmatrix} 2i \\ -2i \end{pmatrix} r \times \bar{r} + \begin{pmatrix} 2 \\ -2 \end{pmatrix} (\gamma \mathcal{W}).$$

Equation (1.2) can be regarded as a white-noise perturbation of a stationary Dirac equation. Dirac operators have wide-ranging applications in quantum mechanics, relativistic field theory, differential geometry, and Clifford algebras [13, 19]. By rewriting equation (CR) in terms of Dirac operators, we establish a connection to these areas, suggesting potential applications of our equation to these fields. We also remark that stationary deterministic Dirac equations have been studied in the literature [8, 12].

The quadratic term  $r \times \bar{r}$  in equation (1.2) represents a nonlinear self-interaction, where the solution  $r$  couples with its own complex conjugate, introducing an interesting mathematical structure. One possible interpretation of this cross-product term is that it describes a nonlinear coupling between spinor-like components in a

quantum mechanical system. Nonlinearities of quadratic type appear, for example, in Dirac-Klein-Gordon systems containing an interaction term that couples a Dirac field with a scalar field, as well as an interaction term representing a self-interaction of the Dirac field [18]. Quadratic nonlinearities appear also in quadratic Dirac equations, which describe self-interacting fermions [18]. From a mathematical perspective, the cross-product term plays a crucial role in balancing singularities introduced by the white noise  $\mathscr{W}$ . Specifically, this nonlinearity contains a self-correcting mechanism that cancels the most singular terms, ensuring the existence of solutions as limits in a Hölder space of negative order. This interplay between the structure of our nonlinearity and the solvability of the equation highlights an interesting and novel aspect of this work.

Another motivation for studying equation (CR) comes from the following stochastic equation

$$(1.3) \quad \Delta B = 2\partial_x B \times \partial_y B + 4\gamma\mathscr{W},$$

where  $B : \mathbb{T}^2 \rightarrow \mathbb{R}^3$ . The homogeneous version of equation (1.3) can be derived from the homogeneous Landau-Lifshitz-Gilbert (LLG) equation

$$u \times \Delta u = 0 \text{ with } |u| = 1$$

via the Bäcklund transformation  $\nabla B := (u \times \partial_y u, -u \times \partial_x u)$ , see e.g. [16, p. 50-52]. This transformation provides a mathematical connection between the LLG equation and equation (1.3), suggesting that (1.3) can be viewed as a stochastic generalization of the LLG equation.

It is straightforward to prove, on a heuristic level, that  $r = i\partial_z B$  is a solution to (CR) if and only if  $B$  is a solution to (1.3). This provides an additional motivation for studying equation (CR), as it establishes a link between our stochastic equation and a well-known physical model. In Section 6, we deduce the existence of a solution  $B$  to (1.3) from the existence of a solution  $r$  to (CR) under an additional assumption on the white noise  $\mathscr{W}$ .

Thus, in this work we are motivated to study equation (CR). To the best of our knowledge, the Cauchy-Riemann operators in the context of singular SPDEs have not been studied before. Moreover, the periodic setting introduces additional challenges, as the solvability of (CR) depends on the mean-zero condition of the right-hand side. More specifically, as we demonstrate in Lemma A.1, the equation  $\partial_{\bar{z}} r = f$  is solvable if and only if  $\int_{\mathbb{T}^2} f = 0$ .

We remark that, unlike most singular SPDEs considered in the literature, see e.g. [3, 4, 9, 14, 15], ours is of the first-order, where the kernel of the operator  $\partial_{\bar{z}}$  increases the regularity by 1. In this sense, and also since  $\partial_{\bar{z}}\partial_z = \frac{1}{2}\Delta$ , our work is related to recent studies of singular SPDEs involving the fractional Laplacian  $(-\Delta)^{\frac{1}{2}}$  [1, 7, 10, 11], which can also be regarded as a first-order differential operator due to its kernel's regularity-increasing properties. However, our focus on the Cauchy-Riemann operator introduces different challenges and requires new techniques, distinct from those used for fractional Laplacians. By developing these techniques, we aim to extend the existing framework to a broader class of first-order singular SPDEs.

**1.2. The method.** In this paper, we consider the mild formulation of equation (CR), see e.g. (3.2), with the complex-valued Green function  $G = 2\partial_{\bar{z}}K$  of the operator  $\partial_{\bar{z}}$  on  $\mathbb{T}^2$ , where  $K$  is the Green function for the Laplacian with zero mean on the torus  $\mathbb{T}^2$ . From this “mild” equation, we deduce that the expected regularity of a solution to equation (CR) is  $0^-$ . This implies that the quadratic term  $r \times \bar{r}$  is ill-defined. To overcome this difficulty, we apply the method used by Da Prato–Debussche in [9] which involves decomposing the solution  $r$  into a singular part and a smoother remainder. More specifically, we rewrite the mild formulation of equation (CR) with respect to a new variable  $R = r + \gamma\xi$ , where  $\xi := 2iG * \mathscr{W}$  is the singular term. This decomposition allows us to isolate and control the additive term in the equation for  $R$ , namely, the term

$$(1.4) \quad \zeta := 2G * (\xi \times \bar{\xi}).$$

Let us observe that the term  $\vartheta := \xi \times \bar{\xi}$  is ill-defined, since  $\xi$  has negative regularity  $0^-$ . We overcome this difficulty by employing a smooth approximation of  $\xi$ , redefining  $\vartheta$  as the corresponding limit, and showing that its components, we take the first component for certainty, are of the following form:

$$(1.5) \quad \vartheta_1 = 2\partial_1[(K * \mathscr{W}_2)\operatorname{Re} \xi_3] - 2\partial_2[(K * \mathscr{W}_2)\operatorname{Im} \xi_3].$$

Deducing the above expression involves a renormalization procedure via cancellation of divergent terms. We observe that (1.5) is well defined since the expression inside the square brackets can be understood in the Young product sense, see [2]. Moreover, this expression suggests that the expected regularity of  $\vartheta$  is  $-1^-$ . Had this been the case, the expected regularity of  $\zeta$ , in (1.4), would be  $0^-$  and therefore the expected regularity of  $R$  would also be  $0^-$ . Such a regularity is too low for the product  $R \times \bar{\xi}$  to be well-defined. Hence, a fixed-point argument cannot be realized in a Hölder space of a positive regularity. A natural way to remedy this would be to consider a lift of our equation to a space of modeled distributions, as in [2, 3, 14, 15].

However, we are able to avoid using this approach by proving, exploiting another reformulation of (1.5), that  $\zeta$  has regularity  $1^-$ , see Proposition 4.8 and Corollary 4.10. This subtle regularity analysis represents a significant technical innovation and is the cornerstone of our existence proof. It allows us to carry out the Banach fixed-point argument in the Hölder space of a positive regularity  $C^{1-\kappa}(\mathbb{T}^2)$ ,  $\kappa \in (0, \frac{1}{2})$ , by understanding the vector product  $R \times \bar{\xi}$ , and other similar, in the Young sense. The main ingredient of our regularity analysis is the following representation of the component  $\vartheta_1$ , as well as  $\vartheta_2$  and  $\vartheta_3$ :

$$\vartheta_1 = 2\partial_1[(K * \mathscr{W}_2 - (K * \mathscr{W}_2)(x))\operatorname{Re} \xi_3] - 2\partial_2[(K * \mathscr{W}_2 - (K * \mathscr{W}_2)(x))\operatorname{Im} \xi_3],$$

for an arbitrary  $x \in \mathbb{T}^2$ . As in (1.5), a renormalization occurs through a cancellation of certain divergent terms. We remark that including  $(K * \mathscr{W}_2)(x)$  in the expression (1.5) is critical in our analysis.

It remains to notice that to analyze the ill-defined product  $\xi \times \bar{\xi}$ , we employ a smooth white noise approximation by the standard mollifier and show the following convergence rate  $\|\xi \times \bar{\xi} - \xi^\varepsilon \times \bar{\xi}^\varepsilon\|_{C^{-\kappa}(\mathbb{T}^2)} \lesssim \varepsilon^{\frac{\kappa}{2}} \|\mathscr{W}\|_{C^{-1-\kappa}(\mathbb{T}^2)}$ . In [9], the authors addressed ill-defined products on a 2D torus by proving a convergence of Galerkin approximations in a negative-order Sobolev space.

### 1.3. Main result.

**Definition 1.1.** A 3D white noise on  $\mathbb{T}^d$  is a Gaussian random field

$$\mathscr{W} = (\mathscr{W}_1, \mathscr{W}_2, \mathscr{W}_3) : \mathbb{T}^d \times \Omega \rightarrow \mathbb{R}^3$$

with the following Fourier series representation:

$$(1.6) \quad \mathscr{W}_j = \eta_j + \operatorname{Re} \left[ \sum_{k \in \mathbb{Z}^d \setminus \{0\}} \eta_{kj} e^{i(k,x)} \right], \quad j = 1, 2, 3,$$

where  $\eta_j$ ,  $j = 1, 2, 3$ , are standard i.i.d. real Gaussian random variables and  $\eta_{kj}$ ,  $j = 1, 2, 3$ ,  $k \in \mathbb{Z}^d \setminus \{0\}$ , are standard i.i.d. complex Gaussian random variables. Furthermore, the series (1.6) converge in  $\mathscr{D}'(\mathbb{T}^d)$ .

**Remark 1.2.** We say that the white noise  $\mathscr{W}_j$  has zero mean over  $\mathbb{T}^d$  if, in representation (1.6),  $\eta_j = 0$ . We say that  $\mathscr{W}$  has zero mean over  $\mathbb{T}^d$  if  $\mathscr{W}_1$ ,  $\mathscr{W}_2$ , and  $\mathscr{W}_3$  have zero means over  $\mathbb{T}^d$ .

The following is the main result of our paper.

**Theorem 1.3.** *Let  $(\Omega, \mathscr{F}, \mathbb{P})$  be a probability space and let  $\mathscr{W} = (\mathscr{W}_1, \mathscr{W}_2, \mathscr{W}_3)$  be a 3D white noise on  $\mathbb{T}^2$  such that  $\mathscr{W}_3$  has zero mean over  $\mathbb{T}^2$ . Further let  $(\mathscr{W}_1^\varepsilon, \mathscr{W}_2^\varepsilon, \mathscr{W}_3^\varepsilon)$  be a mollification of  $\mathscr{W}$  by the standard mollifier  $\rho_\varepsilon$ , that is,  $\mathscr{W}_k^\varepsilon = \rho_\varepsilon * \mathscr{W}_k$ ,  $k = 1, 2, 3$ . Assume that  $\kappa \in (0, \frac{1}{2})$ . Then, for every  $\delta > 0$ , there exists a number  $\varsigma > 0$  and a set  $\Omega_\delta \subset \Omega$ , depending also on  $\kappa$ , with  $\mathbb{P}(\Omega_\delta) > 1 - \delta$ , such that for every  $\gamma = (\gamma_1, \gamma_2, \gamma_3)$  with  $|\gamma| < \varsigma$ , and for every  $\varepsilon > 0$ , the system*

$$(mCR) \quad \begin{cases} \partial_{\bar{z}} r_1^\varepsilon = r_2^\varepsilon \bar{r}_3^\varepsilon - \bar{r}_2^\varepsilon r_3^\varepsilon + \gamma_1 i \mathscr{W}_1^\varepsilon, \\ \partial_{\bar{z}} r_2^\varepsilon = \bar{r}_1^\varepsilon r_3^\varepsilon - r_1^\varepsilon \bar{r}_3^\varepsilon + \gamma_2 i \mathscr{W}_2^\varepsilon, \\ \partial_{\bar{z}} r_3^\varepsilon = r_1^\varepsilon \bar{r}_2^\varepsilon - \bar{r}_1^\varepsilon r_2^\varepsilon + \gamma_3 i \mathscr{W}_3^\varepsilon, \end{cases}$$

considered for  $\omega \in \Omega_\delta$ , has a solution  $r^\varepsilon = (r_1^\varepsilon, r_2^\varepsilon, r_3^\varepsilon) \in C^\infty(\mathbb{T}^2, \mathbb{C}^3)$  such that there exists a limit

$$r := \lim_{\varepsilon \rightarrow 0} r^\varepsilon \text{ in } C^{-\kappa}(\mathbb{T}^2) \quad \text{for every } \omega \in \Omega_\delta.$$

Moreover, the convergence holds with the following rate:

$$(1.7) \quad \|r^\varepsilon - r\|_{C^{-\kappa}(\mathbb{T}^2)} \lesssim \varepsilon^{\frac{\kappa}{2}} \quad \text{uniformly in } \omega \in \Omega_\delta.$$

**Definition 1.4.** By a solution  $r^\varepsilon = (r_1^\varepsilon, r_2^\varepsilon, r_3^\varepsilon)$  to system (mCR) we understand a classical solution. By a solution to (CR) we understand a limit of  $r^\varepsilon$ , as  $\varepsilon \rightarrow 0$ , in some sense.

In other words, Theorem 1.3 can be rephrased in the following way.

**Theorem 1.5.** *Let  $(\Omega, \mathscr{F}, \mathbb{P})$  be a probability space and let  $\mathscr{W} = (\mathscr{W}_1, \mathscr{W}_2, \mathscr{W}_3)$  be a 3D white noise on  $\mathbb{T}^2$  such that  $\mathscr{W}_3$  has zero mean over  $\mathbb{T}^2$ . Then, equation (CR) has a solution.*

**Remark 1.6.** Naturally, instead of assuming that the third component  $\mathscr{W}_3$  has zero mean over  $\mathbb{T}^2$ , we could alternatively assume that the first  $\mathscr{W}_1$  or the second component  $\mathscr{W}_2$  has zero mean over  $\mathbb{T}^2$ .

## 2. DEFINITIONS AND NOTATIONS

In what follows, we denote by  $\mathbb{T}^d := [-\pi, \pi]^d / \sim$  (where  $\sim$  is an equivalence relation that identifies opposite faces of the hypercube  $[-\pi, \pi]^d$ ), the  $d$ -dimensional torus which we will treat as a compact Riemannian manifold and a topological group. By 0 we will denote the corresponding zero element of  $\mathbb{T}^d$ .

By  $\mathcal{D}(\mathbb{T}^d)$  we denote the space of  $C^\infty$ -class functions on  $\mathbb{T}^d$  and by  $\mathcal{D}'(\mathbb{T}^d)$  the space of distributions on  $\mathbb{T}^d$ , i.e. the topological dual of  $\mathcal{D}(\mathbb{T}^d)$ . Note that the space  $\mathcal{D}(\mathbb{T}^d)$ , also called the space of test functions, is a Fréchet space endowed with the naturally defined countable family of seminorms. If  $U$  is an open subset of  $\mathbb{T}^d$ , then by  $\mathcal{D}(U)$  we denote the class of those test functions whose support is contained in  $U$ . Similarly, by  $\mathcal{D}'(U)$  we denote the topological dual of  $\mathcal{D}(U)$ .

For a test function  $\varphi \in \mathcal{D}(\mathbb{T}^d)$ ,  $\lambda > 0$  and  $x \in \mathbb{T}^d$ , we define the rescaled centered function  $\varphi_x^\lambda$  as follows:

$$\varphi_x^\lambda(y) := \frac{1}{\lambda^d} \varphi\left(\frac{y-x}{\lambda}\right).$$

Furthermore, we write  $\varphi_x$  instead of  $\varphi_x^1$ .

Let  $r \in \mathbb{N}_0$  and  $\alpha \in \mathbb{R}$ . Define the classes of test functions

$$(2.1) \quad \begin{aligned} \mathcal{B}^r &:= \{\varphi \in \mathcal{D}(\mathbb{T}^d) : \|\partial^k \varphi\|_\infty \leq 1, \text{ for all } 0 \leq |k| \leq r\}, \\ \mathcal{B}_\alpha &:= \{\varphi \in \mathcal{D}(\mathbb{T}^d) : \int_{\mathbb{T}^d} \varphi(z) z^k dz = 0, \text{ for all } 0 \leq |k| \leq \alpha\}, \text{ if } \alpha \geq 0, \\ \mathcal{B}_\alpha^r &:= \mathcal{B}^r \cap \mathcal{B}_\alpha, \text{ if } \alpha \geq 0, \end{aligned}$$

where  $k = (k_1, \dots, k_d)$  is a multiindex and if  $z = (z_1, \dots, z_d) \in \mathbb{T}^d$ ,  $z^k := z_1^{k_1} \dots z_d^{k_d}$ .

**Remark 2.1.** Note that our definitions of  $\mathcal{B}^r$ ,  $\mathcal{B}_\alpha$ , and  $\mathcal{B}_\alpha^r$  differ from those in [2, Definition A.9], as we have taken into account the periodicity of our problem.

Define the dyadic interval

$$[0, 1]_{\mathbb{D}} := \{2^{-n}, n \in \mathbb{N}_0\}.$$

**Definition 2.2** (Periodic Hölder-Zygmund spaces  $\mathcal{Z}^\alpha(\mathbb{T}^d)$ ). Assume that  $\alpha \in \mathbb{R}$ . If  $\alpha < 0$ , we denote by  $r = [-\alpha + 1]$  the smallest positive integer  $r > -\alpha$ . We define a “pseudo-norm”  $|f|_{\mathcal{Z}^\alpha}$  for  $f \in \mathcal{D}'(\mathbb{T}^d)$  as follows.

If  $\alpha \geq 0$ , then we put

$$(2.2) \quad |f|_{\mathcal{Z}^\alpha} := \sup_{\psi \in \mathcal{B}^0} |f(\psi)| + \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0, 1]_{\mathbb{D}}, \\ \varphi \in \mathcal{B}_\alpha^0}} \frac{|f(\varphi_x^\lambda)|}{\lambda^\alpha}.$$

If  $\alpha < 0$ , then, with  $r = [-\alpha + 1]$ , we put

$$(2.3) \quad |f|_{\mathcal{Z}^\alpha} := \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0, 1]_{\mathbb{D}}, \\ \varphi \in \mathcal{B}^r}} \frac{|f(\varphi_x^\lambda)|}{\lambda^\alpha}.$$

We define the periodic Hölder-Zygmund space  $\mathcal{Z}^\alpha := \mathcal{Z}^\alpha(\mathbb{T}^d)$  as the space of distributions  $f \in \mathcal{D}'(\mathbb{T}^d)$  for which  $|f|_{\mathcal{Z}^\alpha} < \infty$ .

**Remark 2.3.** By Lemma C.7, Definition 2.2 agrees with the classical definition of a Hölder-Zygmund space in the periodic setting (see Remark C.6) only if  $\alpha \notin \mathbb{N}$ . For  $\alpha \in \mathbb{N}$ ,  $\mathcal{Z}^\alpha(\mathbb{T}^d)$  defines a distinct space, which is not utilized in our analysis.

**Remark 2.4.** It is known that classical Hölder-Zygmund spaces are not separable and that the space  $C^\infty$  is not dense in them, see [17]. For this reason, we define so called little Hölder-Zygmund spaces  $\mathcal{X}^\alpha$ .

**Definition 2.5** (Periodic little Hölder-Zygmund spaces  $\mathcal{X}^\alpha(\mathbb{T}^d)$ ). We define a periodic little Hölder-Zygmund space  $\mathcal{X}^\alpha := \mathcal{X}^\alpha(\mathbb{T}^d)$  as the closure of  $C^\infty(\mathbb{T}^d)$  in  $\mathcal{Z}^\alpha(\mathbb{T}^d)$ . We furthermore denote by  $|\cdot|_\alpha$ ,  $\alpha \in \mathbb{R}$ , the restriction of the norm (2.2), for  $\alpha \geq 0$ , and (2.3), for  $\alpha < 0$ , to  $\mathcal{X}^\alpha(\mathbb{T}^d)$ .

**Remark 2.6.** It is important to notice that by Lemma B.3, a Gaussian white noise  $\mathscr{W}$  on  $\mathbb{T}^d$ , as well as its convolutions  $K * \mathscr{W}$  and  $G * \mathscr{W}$  (see Definition 2.9), are elements of little Hölder-Zygmund spaces. Furthermore, by Lemma E.1, Young products are well defined (specifically, uniquely defined) when they are restricted to little Hölder-Zygmund spaces. We also remark that Young products are not uniquely defined in classical Hölder-Zygmund spaces, see [6, Remark 14.2]. For these reasons, little Hölder-Zygmund spaces is a natural choice for our analysis.

**Remark 2.7.** For simplicity of notation, we often write  $\mathcal{X}^\alpha$ ,  $\mathcal{Z}^\alpha$ ,  $C^\alpha$ , and  $C^\infty$  instead of  $\mathcal{X}^\alpha(\mathbb{T}^d)$ ,  $\mathcal{Z}^\alpha(\mathbb{T}^d)$ ,  $C^\alpha(\mathbb{T}^d)$ , and  $C^\infty(\mathbb{T}^d)$ , respectively.

If  $f$  is  $\mathbb{C}$ -valued, then we set  $|f|_\alpha^2 := |\operatorname{Re} f|_\alpha^2 + |\operatorname{Im} f|_\alpha^2$ . The above definitions are for  $\mathbb{C}$ -valued distributions. If  $f$  is a  $Y$ -valued distribution, where  $Y$  is an  $n$ -dimensional Euclidean space over  $\mathbb{C}$ , i.e., if  $f = (f_1, \dots, f_n)$ , where  $f_k$  are  $\mathbb{C}$ -valued, then we define the norm

$$(2.4) \quad \|f\|_\alpha^2 := \sum_{k=1}^n |f_k|_\alpha^2.$$

Remark that for  $Y$ -valued distributions, we use the notation  $\|\cdot\|_\alpha$ , rather than  $|\cdot|_\alpha$ , for respective Hölder-Zygmund norms.

**Remark 2.8.** In [2, Definition A.9], the Hölder-Zygmund norms are defined with respect to sets of test functions supported on unit balls centered in the origin. In Lemma C.7, we show that both definitions are equivalent for  $\alpha \notin \mathbb{N}$ , that is, we can equivalently define the Hölder-Zygmund norms as in Definition C.3.

**Definition 2.9** (Green function for  $\partial_{\bar{z}}$ ). Let  $K$  be the Green function for the Laplacian on  $\mathbb{T}^2$ . We define

$$(2.5) \quad G := 2\partial_z K = \partial_1 K - i\partial_2 K.$$

We call  $G$  the Green function for the operator  $-2\partial_{\bar{z}}$ . Therefore,  $-2G$  is the Green function for the Cauchy-Riemann operator  $\partial_{\bar{z}}$ . We also put

$$G_1 := \partial_1 K \text{ and } G_2 := \partial_2 K.$$

## 3. SOLVING PROBLEM (mCR)

3.1. **Mild formulation of problem (mCR).** Let the function  $G$  be given by (2.5). We define

$$(3.1) \quad \xi_k^\varepsilon := 2iG * \mathscr{W}_k^\varepsilon = 2G_2 * \mathscr{W}_k^\varepsilon + 2iG_1 * \mathscr{W}_k^\varepsilon, \quad k = 1, 2, 3.$$

Note that we can write

$$\xi_k^\varepsilon := \xi_{2k}^\varepsilon + i\xi_{1k}^\varepsilon.$$

**Proposition 3.1.** *Let  $c > 0$ ,  $\varepsilon > 0$  and  $\gamma_k \in \mathbb{R}$ ,  $k = 1, 2, 3$ . There exists  $\mathbb{C}$ -valued Gaussian mean zero random variables  $a$  and  $b$  such that if  $(r_1^\varepsilon, r_2^\varepsilon, r_3^\varepsilon)$  is an  $C^\alpha(\mathbb{T}^2)$ -valued random variable, for some  $\alpha \in (0, 1)$ , which is a pathwise solution of the system*

$$(3.2) \quad \begin{cases} r_1^\varepsilon = 2G * (\overline{r_2^\varepsilon} r_3^\varepsilon - r_2^\varepsilon \overline{r_3^\varepsilon}) - \gamma_1 \xi_1^\varepsilon + a, \\ r_2^\varepsilon = 2G * (r_1^\varepsilon \overline{r_3^\varepsilon} - \overline{r_1^\varepsilon} r_3^\varepsilon) - \gamma_2 \xi_2^\varepsilon + b, \\ r_3^\varepsilon = 2G * (\overline{r_1^\varepsilon} r_2^\varepsilon - r_1^\varepsilon \overline{r_2^\varepsilon}) - \gamma_3 \xi_3^\varepsilon + c, \end{cases}$$

then  $r_k^\varepsilon \in C^\infty(\mathbb{T}^2)$ ,  $k = 1, 2, 3$ , and  $(r_1^\varepsilon, r_2^\varepsilon, r_3^\varepsilon)$  solves (mCR).

Before we embark on the proof of this proposition we need to state and prove the following auxiliary result.

**Lemma 3.2.** *Assume  $\alpha \in (0, 1)$ ,  $g_k \in C^\alpha(\mathbb{T}^2, \mathbb{R})$  and  $f_k = iG * g_k$ , for  $k = 1, 2$ . Then, in  $C^{\alpha+1}(\mathbb{T}^2, \mathbb{R})$ ,*

$$(3.3) \quad f_1 \overline{f_2} - \overline{f_1} f_2 = \partial_1(K * g_1 \partial_2 K * g_2) - \partial_2(K * g_1 \partial_1 K * g_2).$$

and

$$(3.4) \quad \int_{\mathbb{T}^2} \text{Im } f_1 \overline{f_2} = 0.$$

*Proof.* First of all, we note that for any complex numbers  $z_1$  and  $z_2$ ,

$$2i \text{Im } z_1 \overline{z_2} = z_1 \overline{z_2} - \overline{z_1} z_2.$$

Note that

$$iG = 2i\partial_z K = i(\partial_1 K - i\partial_2 K) = \partial_2 K + i\partial_1 K,$$

where  $K$  is the Green function of the Laplacian on  $\mathbb{T}^2$ , which is real-valued. Since  $g_k$  are  $C^\alpha$ , then  $K * g_k$  are  $C^{2+\alpha}$  and

$$f_k = iG * g_k = \partial_2 K * g_k + i\partial_1 K * g_k.$$

Thus, we have

$$\begin{aligned} f_1 \overline{f_2} - \overline{f_1} f_2 &= 2i \text{Im } f_1 \overline{f_2} = (\partial_1 K * g_1)(\partial_2 K * g_2) - (\partial_2 K * g_1)(\partial_1 K * g_2) \\ &= \partial_1(K * g_1 \partial_2 K * g_2) - (K * g_1) \partial_{12}^2 K * g_2 - \partial_2(K * g_1 \partial_1 K * g_2) + (K * g_1) \partial_{12}^2 K * g_2 \\ &= \partial_1(K * g_1 \partial_2 K * g_2) - \partial_2(K * g_1 \partial_1 K * g_2) \end{aligned}$$

which proves (3.3) and (3.4).  $\square$

*Proof of Proposition 3.1.* We start by noticing that since the solution  $r^\varepsilon := (r_1^\varepsilon, r_2^\varepsilon, r_3^\varepsilon)$  to (mCR) is  $C^\alpha(\mathbb{T}^2)$ -valued, it is automatically  $C^\infty(\mathbb{T}^2)$ -valued. Indeed,  $(\xi_1^\varepsilon, \xi_2^\varepsilon, \xi_3^\varepsilon)$  is in  $C^\infty(\mathbb{T}^2)$ . On the other hand, by Lemmata C.12 and D.3, a convolution with  $G$  increases the regularity of the expression in parenthesis, on the right-hand side of (mCR), by 1, inferring that  $r^\varepsilon$  is in  $C^{1+\alpha}(\mathbb{T}^2)$ . By induction, we conclude that  $r^\varepsilon$  is in  $C^\infty(\mathbb{T}^2)$ .

In what follows, with a slight abuse of notation, we omit the index  $\varepsilon$ , that is, we write  $\varkappa_1, \varkappa_2$  and  $\varkappa_3$  instead of  $\varkappa_1^\varepsilon, \varkappa_2^\varepsilon$  and  $\varkappa_3^\varepsilon$ . The same applies to other objects in (mCR).

Define  $a$  and  $b$  by the formulas

$$(3.5) \quad 2ac = i\gamma_2\eta_2, \quad 2bc = -i\gamma_1\eta_1,$$

where  $\eta_1$  and  $\eta_2$  are independent standard real Gaussian random variables as in Definition 1.1. From now on, we choose and fix  $\omega \in \Omega$  so that  $\eta_k = \eta_1(\omega)$  and  $\xi_k = \xi_k(\omega)$  are real numbers. Let us introduce  $\rho_k, k = 1, 2, 3$ , as follows:

$$(3.6) \quad r_1 = \varrho_1 + a, \quad r_2 = \varrho_2 + b, \quad r_3 = \varrho_3 + c.$$

Since  $(r_1, r_2, r_3) \in C^\alpha(\mathbb{T}^2)$ , we infer that  $(\varrho_1, \varrho_2, \varrho_3) \in C^\alpha(\mathbb{T}^2)$  and  $(\varrho_1, \varrho_2, \varrho_3)$  is a solution to the following system

$$(3.7) \quad \begin{cases} \varrho_1 = -4i G * \operatorname{Im} [(\varrho_2 + b)(\overline{\varrho_3} + c)] - \gamma_1\xi_1, \\ \varrho_2 = 4i G * \operatorname{Im} [(\varrho_1 + a)(\overline{\varrho_3} + c)] - \gamma_2\xi_2, \\ \varrho_3 = -4i G * \operatorname{Im} [(\varrho_1 + a)(\overline{\varrho_2} - b)] - \gamma_3\xi_3. \end{cases}$$

Then, it follows from (3.1) and (3.7) that we can find functions  $g_j \in C^\alpha(\mathbb{T}^2, \mathbb{R})$  satisfying the assumptions of Lemma 3.2 and

$$\varrho_j = iG * g_j, \quad j = 1, 2, 3.$$

Hence, by Lemma 3.2, for all  $k, j = 1, 2, 3, k \neq j$ ,

$$\int \operatorname{Im} \varrho_k \overline{\varrho_j} = 0.$$

In addition, we notice that  $\int_{\mathbb{T}^2} \varrho_j = 0$  for  $j = 1, 2, 3$ . By Lemma A.1,

$$(3.8) \quad \begin{aligned} \partial_{\bar{z}}\varrho_1 &= 2i \operatorname{Im} [(\varrho_2 + b)(\overline{\varrho_3} + c)] + \gamma_1i(\varkappa_1 - \eta_1) - 2bc \\ &= 2i \operatorname{Im} [(\varrho_2 + b)(\overline{\varrho_3} + c)] + \gamma_1i\varkappa_1, \\ -\partial_{\bar{z}}\varrho_2 &= 2i \operatorname{Im} [(\varrho_1 + a)(\overline{\varrho_3} + c)] - \gamma_2i(\varkappa_2 - \eta_2) - 2ac \\ &= 2i \operatorname{Im} [(\varrho_1 + a)(\overline{\varrho_3} + c)] - \gamma_2i\varkappa_2, \\ \partial_{\bar{z}}\varrho_3 &= 2i \operatorname{Im} [(\varrho_1 + a)(\overline{\varrho_2} - b)] + \gamma_3i\varkappa_3. \end{aligned}$$

By the change of variable (3.6),  $(r_1, r_2, r_3)$  solves (mCR).  $\square$

**Corollary 3.3.** *If  $\varkappa_1$  and  $\varkappa_2$  have zero means over  $\mathbb{T}^2$ , then the statement of Proposition 3.1 holds with  $a = b = c = 0$ .*

*Proof.* Recall that the fact that  $\varkappa_1$  and  $\varkappa_2$  have zero means over  $\mathbb{T}^2$  means that  $\eta_1 = \eta_2 = 0$ . Therefore, system (3.8) (and therefore system (mCR)) is satisfied with  $a = b = c = 0$ .  $\square$



### 3.2. The fixed-point argument.

**Lemma 3.4.** *Throughout this lemma, we choose and fix  $\gamma = (\gamma_1, \gamma_2, \gamma_3)$ . For  $\varepsilon > 0$ , let*

$$(3.9) \quad R^\varepsilon := r^\varepsilon + \gamma \xi^\varepsilon,$$

where  $\gamma \xi^\varepsilon := (\gamma_1 \xi_1^\varepsilon, \gamma_2 \xi_2^\varepsilon, \gamma_3 \xi_3^\varepsilon)$ . Then, system (3.2) is equivalent to

$$(3.10) \quad R^\varepsilon = -2G * (R^\varepsilon \times \overline{R^\varepsilon} - (\gamma \xi^\varepsilon) \times \overline{R^\varepsilon} - R^\varepsilon \times (\gamma \overline{\xi^\varepsilon})) - \tilde{\gamma} \zeta^\varepsilon + \begin{pmatrix} a \\ b \\ c \end{pmatrix},$$

where

$$(3.11) \quad \zeta^\varepsilon := 2G * (\xi^\varepsilon \times \overline{\xi^\varepsilon}), \quad \tilde{\gamma} := (\gamma_2 \gamma_3, \gamma_1 \gamma_3, \gamma_1 \gamma_2)$$

and by  $\tilde{\gamma} \zeta^\varepsilon$ , we mean  $(\tilde{\gamma}_1 \zeta_1^\varepsilon, \tilde{\gamma}_2 \zeta_2^\varepsilon, \tilde{\gamma}_3 \zeta_3^\varepsilon)$ .

**Remark 3.5.** Note that so far  $\zeta^\varepsilon$  makes sense only for  $\varepsilon > 0$  because the product  $\xi^\varepsilon \times \overline{\xi^\varepsilon}$  is ill-defined for  $\varepsilon = 0$ .

*Proof of Lemma 3.4.* One just needs to notice that (3.2) is equivalent to

$$r^\varepsilon = -2G * (r^\varepsilon \times \overline{r^\varepsilon}) + \begin{pmatrix} a \\ b \\ c \end{pmatrix}.$$

The substituting  $r^\varepsilon = R^\varepsilon - \gamma \xi^\varepsilon$ , we obtain (3.10). The proof is complete.  $\square$

For every  $\nu \in \mathbb{R}$ , introduce the Banach space

$$E_\nu = \{R = (R_1, R_2, R_3) : \mathbb{T}^2 \rightarrow \mathbb{C}^3 : R_k \in \mathcal{X}^\nu(\mathbb{T}^2), k = 1, 2, 3\}$$

with the norm  $\|\cdot\|_\nu$  defined by (2.4). For every  $\sigma > 0$ , we set

$$c = \frac{\sigma}{4} \quad \text{and define } a, b \text{ via } c \text{ by formulas (3.5).}$$

Given  $\varepsilon > 0$ ,  $\sigma > 0$ , and  $\kappa \in (0, \frac{1}{2})$ , define a map  $\Gamma_{\varepsilon, \sigma} : E_{1-\kappa} \rightarrow E_{1-\kappa}$ ,

$$(3.12) \quad \Gamma_{\varepsilon, \sigma} R := -2G * \left( R \times \overline{R} + (\gamma \xi^\varepsilon) \times \overline{R} + R \times (\gamma \overline{\xi^\varepsilon}) \right) - \tilde{\gamma} \zeta^\varepsilon + \begin{pmatrix} a \\ b \\ c \end{pmatrix}.$$

As we will see from Lemmata 3.7 and 3.8,  $\xi^\varepsilon$  and  $\zeta^\varepsilon$  are well defined also for  $\varepsilon = 0$ . Therefore, (3.12) also defines a map  $\Gamma_{0, \sigma}$  for every  $\sigma > 0$ . Our aim is to show that for every  $\delta > 0$ , there exists a set  $\Omega_\delta \subset \Omega$ ,  $\mathbb{P}(\Omega_\delta) > 1 - \delta$ , and  $\sigma > 0$  such that for every  $\varepsilon \geq 0$  and  $|\gamma| \lesssim \sigma^2$ , for every  $\omega \in \Omega_\delta$ , the map  $\Gamma_{\varepsilon, \sigma} : E_{1-\kappa} \rightarrow E_{1-\kappa}$  is well defined and has a unique fixed point.

For this aim, as well as to define the map  $\Gamma_{0, \sigma}$ , we need the following auxiliary Lemmata 3.7 and 3.8.

**Remark 3.6.** Note that by Lemma C.15,  $C^\infty(\mathbb{T}^2) = \bigcap_{\nu} \mathcal{X}^\nu(\mathbb{T}^2)$ . Therefore,

$$\mathcal{W}^\varepsilon, \xi^\varepsilon, \zeta^\varepsilon \in E_\nu \quad \text{for all } \nu \in \mathbb{R} \text{ and } \varepsilon > 0.$$

**Lemma 3.7.** *For all  $\kappa \in (0, 1)$ ,  $\xi \in E_{-\kappa}$ . Moreover, for all  $\varepsilon \geq 0$ ,*

$$\|\xi^\varepsilon\|_{-\kappa} \lesssim \|\mathscr{W}\|_{-1-\kappa},$$

where  $\mathscr{W}$  on the right-hand side is the non-mollified 3D white noise appearing in (CR),  $\xi^\varepsilon$  is defined by (3.1),  $\xi^0 := \xi$ , and the constant in the above inequality does not depend on  $\varepsilon \geq 0$ .

*Proof.* By Lemma B.3,  $\mathscr{W}_k$ ,  $k = 1, 2, 3$ , are in  $\mathscr{D}^{-1-\kappa}(\mathbb{T}^2)$ . Furthermore, by Lemma D.3,  $\xi_k$ ,  $k = 1, 2, 3$ , are in  $\mathscr{D}^{-\kappa}(\mathbb{T}^2)$ . Since  $\xi_{kj}^\varepsilon = G_k * \mathscr{W}_j^\varepsilon$ ,  $k = 1, 2$ ;  $j = 1, 2, 3$ , one has

$$|\xi_{kj}^\varepsilon|_{-\kappa} \lesssim |\mathscr{W}_j^\varepsilon|_{-1-\kappa} \lesssim |\mathscr{W}_j|_{-1-\kappa}.$$

In particular, we have used Lemma C.9.  $\square$

**Lemma 3.8.** *Let, for  $\varepsilon > 0$ ,  $\zeta^\varepsilon$  be defined by (3.11) and  $\kappa \in (0, \frac{1}{2})$ . Then  $\zeta^\varepsilon$  has a limit  $\zeta$  in  $E_{1-\kappa}$  as  $\varepsilon \rightarrow 0$ . Furthermore, for all  $\varepsilon \geq 0$  and  $\kappa \in (0, \frac{1}{2})$ ,*

$$(3.13) \quad \|\zeta^\varepsilon\|_{1-\kappa} \lesssim \|\mathscr{W}\|_{-1-\kappa}^2,$$

where  $\zeta^0 := \zeta$ , and the constant in the above inequality does not depend on  $\varepsilon \geq 0$ .

Lemma 3.8 is a fundamental result of this work, underpinning the proof of Theorem 1.3. Its proof is postponed to Subsection 4, where we give the explicit expression for  $\zeta$  and prove (3.13).

**Lemma 3.9.** *If  $\varepsilon \geq 0$ ,  $\sigma > 0$ , and  $\kappa \in (0, \frac{1}{2})$ , then the map  $\Gamma_{\varepsilon, \sigma} : E_{1-\kappa} \rightarrow E_{1-\kappa}$ , given by (3.12), is well-defined.*

*Proof of Lemma 3.9.* Let us choose and fix  $R \in E_{1-\kappa}$ . In view of Lemmata 3.8 and D.3, it is sufficient to show that  $R \times \bar{R} + (\gamma\xi^\varepsilon) \times \bar{R} + R \times (\gamma\bar{\xi}^\varepsilon) \in E_{-\kappa}$ .

Since,  $\bar{R} \in E_{1-\kappa}$  and the space  $E_{1-\kappa}$  is an algebra, we infer that  $R \times \bar{R} \in E_{1-\kappa} \subset E_{-\kappa}$ . Moreover, by Lemma E.1 and since  $\xi^\varepsilon \in E_{-\kappa}$ , we infer that both  $(\gamma\xi^\varepsilon) \times \bar{R}$  and  $R \times (\gamma\bar{\xi}^\varepsilon)$  belong to  $E_{-\kappa}$ . The proof is complete.  $\square$

Let us consider the closed ball in  $E_{1-\kappa}$  of radius  $\sigma > 0$  (to be fixed later)

$$M_{1-\kappa, \sigma} := \{R \in E_{1-\kappa} : \|R\|_{1-\kappa} \leq \sigma\}.$$

Obviously,  $M_{1-\kappa, \sigma}$  is a complete metric space.

**Proposition 3.10.** *For every  $\delta \in (0, 1)$  and  $\kappa \in (0, \frac{1}{2})$ , there exists a set  $\Omega_\delta \subset \Omega$ ,  $\mathbb{P}(\Omega_\delta) > 1 - \delta$ , and a number  $\sigma > 0$  such that for all  $\omega \in \Omega_\delta$ ,  $\varepsilon \geq 0$ , and  $\gamma = (\gamma_1, \gamma_2, \gamma_3)$  satisfying  $|\gamma| \lesssim \sigma^2$ ,  $\Gamma_{\varepsilon, \sigma}$  maps  $M_{1-\kappa, \sigma}$  into itself and the corresponding map  $\Gamma_{\varepsilon, \sigma} : M_{1-\kappa, \sigma} \rightarrow M_{1-\kappa, \sigma}$  is a strict contraction and hence has a unique fixed point in  $M_{1-\kappa, \sigma}$ . Moreover, the contraction constant of  $\Gamma_{\varepsilon, \sigma}$  does not depend on  $\varepsilon \geq 0$  and  $\omega \in \Omega_\delta$ .*

**Remark 3.11.** We reinforce that Proposition 3.10 is also valid for the map  $\Gamma_{0, \sigma}$ ;  $\xi^0$  and  $\zeta^0$  are understood as in Lemmata 3.7 and 3.8.

*Proof of Proposition 3.10.* Let  $\delta \in (0, 1)$  be fixed arbitrarily. Since the Gaussian random variables  $\eta_1$  and  $\eta_2$  are finite a.s., by Lemma B.2 we can find a  $\Lambda \in (0, +\infty)$  such that  $\mathbb{P}(\Omega_\delta) > 1 - \delta$ , where

$$(3.14) \quad \Omega_\delta := \left\{ \omega : \|\mathscr{W}\|_{-1-\kappa} + |\eta_1| + |\eta_2| < \Lambda \right\}.$$

Note that by Lemmata 3.7, 3.8 and C.9,  $\|\xi^\varepsilon\|_{-\kappa}$  and  $\|\zeta^\varepsilon\|_{1-\kappa}$  are bounded on  $\Omega_\delta$  uniformly in  $\varepsilon$  by a constant depending only on  $\Lambda$ . Take  $R \in M_{1-\kappa, \sigma}$ . Recall that in the definition of the map  $\Gamma_{\varepsilon, \sigma}$  by (3.12), we took  $c = \frac{\sigma}{4}$ . Choose  $\gamma = (\gamma_1, \gamma_2, \gamma_3)$  such that  $|\gamma| < \frac{\sigma^2}{16\Lambda}$ . Then,  $|a| + |b| + c < \frac{\sigma}{2}$ . By Lemmata 3.7, 3.8, and E.1,

$$\|\Gamma_{\varepsilon, \sigma} R\|_{1-\kappa} \leq C(\sigma^2 + |\gamma|\sigma\|\xi^\varepsilon\|_{-\kappa} + |\gamma|^2\|\zeta^\varepsilon\|_{1-\kappa}) + \frac{\sigma}{2} \leq C_\Lambda(\sigma^2 + \sigma^3 + \sigma^4) + \frac{\sigma}{2} \leq \sigma,$$

where  $C$  is a constant that depends only on  $\kappa$  and  $G$ , and  $C_\Lambda$  is another constant that depends only on  $C$  and  $\Lambda$ . The latter inequality is achieved by choosing  $\sigma$  such that

$$(3.15) \quad C_\Lambda(\sigma + \sigma^2 + \sigma^3) < \frac{1}{2}.$$

Therefore,  $\Gamma_{\varepsilon, \sigma}$  maps  $M_{1-\kappa, \sigma}$  to  $M_{1-\kappa, \sigma}$ . Further, take two points  $R, \hat{R} \in M_{1-\kappa, \sigma}$ . We obtain

$$(3.16) \quad \|\Gamma_{\varepsilon, \sigma} R - \Gamma_{\varepsilon, \sigma} \hat{R}\|_{1-\kappa} \leq 2C(\sigma + |\gamma|\|\xi^\varepsilon\|_{-\kappa})\|R - \hat{R}\|_{1-\kappa} \leq 2C_\Lambda(\sigma + \sigma^2)\|R - \hat{R}\|_{1-\kappa}.$$

Observe that by (3.15),  $2C_\Lambda(\sigma + \sigma^2) < 1$ . Therefore,  $\Gamma_{\varepsilon, \sigma}$  has a unique fixed point in  $M_{1-\kappa, \sigma}$ . We also remark that the fixed-point constant does not depend on  $\varepsilon \geq 0$  and  $\omega \in \Omega_\delta$ .  $\square$

**Remark 3.12.** We remark that the proof holds also for  $\varepsilon = 0$ . In this case,  $\xi^0$  and  $\zeta^0$  are the limits of  $\xi^\varepsilon$  and  $\zeta^\varepsilon$  in  $E_{-\kappa}$  and  $E_{1-\kappa}$ , respectively.

#### 4. RENORMALIZATION AND REGULARITY OF $\zeta^\varepsilon$ .

Recall that  $\zeta^\varepsilon$ , defined for  $\varepsilon > 0$  as

$$\zeta^\varepsilon := 2G * (\xi^\varepsilon \times \overline{\xi^\varepsilon}),$$

is a noise term in the fixed-point equation (3.4). We also recall that the proof of Lemma 3.8 was postponed to this section. According to Lemma 3.8,  $\zeta^\varepsilon$  has a limit  $\zeta$  in  $E_{1-\kappa}$ . This is a non-trivial and key result of this work since, as it will be clear from Lemma 4.2, the explicit expression for  $\xi^\varepsilon \times \overline{\xi^\varepsilon}$  suggests that  $\zeta \in E_{-\kappa}$ . The latter property, however, is not sufficient for the fixed-point argument to work. The first step in proving Lemma 3.8 is Lemma 4.2 in which, in particular, we prove that the  $\vartheta := \lim_{\varepsilon \rightarrow 0} \xi^\varepsilon \times \overline{\xi^\varepsilon}$  exists in  $\mathscr{D}'(\mathbb{T}^2)$ . The second step is Proposition 4.8, where we prove that  $\vartheta \in \cap_{\kappa \in (0, \frac{1}{2})} E_{-\kappa}$ . Corollary 4.10, which is a direct consequence of Proposition 4.8, implies Lemma 3.8.

**Lemma 4.1.** *Let  $A \in \mathbb{R}$  be a constant. Then, under the assumptions of Lemma 3.2, in  $C^{\alpha+1}(\mathbb{T}^2, \mathbb{R})$ , we have the identity*

$$f_1 \overline{f_2} - \overline{f_1} f_2 = \partial_1((K * g_1 - A) \partial_2 K * g_2) - \partial_2((K * g_1 - A) \partial_1 K * g_2).$$

*Proof.* Using the definition of  $G$  and  $f_k$ ,  $k = 1, 2$ , we obtain

$$\begin{aligned} f_1 \overline{f_2} - \overline{f_1} f_2 &= (\partial_1(K * g_1 - A))(\partial_2 K * g_2) - (\partial_2(K * g_1 - A))(\partial_1 K * g_2) \\ &= \partial_1((K * g_1 - A) \partial_2 K * g_2) - (K * g_1) \partial_{12}^2 K * g_2 - \partial_2((K * g_1 - A) \partial_1 K * g_2) + (K * g_1) \partial_{12}^2 K * g_2 \\ &= \partial_1((K * g_1 - A) \partial_2 K * g_2) - \partial_2((K * g_1 - A) \partial_1 K * g_2). \end{aligned}$$

□

**Lemma 4.2.** *The components of  $\vartheta^\varepsilon := \xi^\varepsilon \times \overline{\xi^\varepsilon}$  satisfy the following relations:*

$$(4.1) \quad \begin{aligned} \vartheta_1^\varepsilon &= 2(\partial_1[(K * \mathscr{W}_2^\varepsilon) \xi_{23}^\varepsilon] - \partial_2[(K * \mathscr{W}_2^\varepsilon) \xi_{13}^\varepsilon]), \\ \vartheta_2^\varepsilon &= 2(\partial_1[(K * \mathscr{W}_3^\varepsilon) \xi_{21}^\varepsilon] - \partial_2[(K * \mathscr{W}_3^\varepsilon) \xi_{11}^\varepsilon]), \\ \vartheta_3^\varepsilon &= 2(\partial_1[(K * \mathscr{W}_1^\varepsilon) \xi_{22}^\varepsilon] - \partial_2[(K * \mathscr{W}_1^\varepsilon) \xi_{12}^\varepsilon]), \end{aligned}$$

where  $\xi_{jk}^\varepsilon$  have been introduced in (3.1). Furthermore,  $\vartheta^\varepsilon$  satisfy, for any  $x \in \mathbb{T}^2$ , the additional relations

$$(4.2) \quad \begin{aligned} \vartheta_1^\varepsilon &= 2(\partial_1[(K * \mathscr{W}_2^\varepsilon - K * \mathscr{W}_2^\varepsilon(x)) \xi_{23}^\varepsilon] - \partial_2[(K * \mathscr{W}_2^\varepsilon - K * \mathscr{W}_2^\varepsilon(x)) \xi_{13}^\varepsilon]), \\ \vartheta_2^\varepsilon &= 2(\partial_1[(K * \mathscr{W}_3^\varepsilon - K * \mathscr{W}_3^\varepsilon(x)) \xi_{21}^\varepsilon] - \partial_2[(K * \mathscr{W}_3^\varepsilon - K * \mathscr{W}_3^\varepsilon(x)) \xi_{11}^\varepsilon]), \\ \vartheta_3^\varepsilon &= 2(\partial_1[(K * \mathscr{W}_1^\varepsilon - K * \mathscr{W}_1^\varepsilon(x)) \xi_{22}^\varepsilon] - \partial_2[(K * \mathscr{W}_1^\varepsilon - K * \mathscr{W}_1^\varepsilon(x)) \xi_{12}^\varepsilon]), \end{aligned}$$

where both (4.1) and (4.2) are valid in  $C^\infty(\mathbb{T}^2)$ . Moreover, there exists a limit

$$(4.3) \quad \vartheta := \lim_{\varepsilon \rightarrow 0} \vartheta^\varepsilon \text{ in } \mathscr{D}'(\mathbb{T}^2),$$

and the components  $\vartheta_k$  of  $\vartheta$ ,  $k = 1, 2, 3$ , are given by

$$(4.4) \quad \begin{aligned} \vartheta_1 &= 2(\partial_1[(K * \mathscr{W}_2) \xi_{23}] - \partial_2[(K * \mathscr{W}_2) \xi_{13}]), \\ \vartheta_2 &= 2(\partial_1[(K * \mathscr{W}_3) \xi_{21}] - \partial_2[(K * \mathscr{W}_3) \xi_{11}]), \\ \vartheta_3 &= 2(\partial_1[(K * \mathscr{W}_1) \xi_{22}] - \partial_2[(K * \mathscr{W}_1) \xi_{12}]), \end{aligned}$$

and also by

$$(4.5) \quad \begin{aligned} \vartheta_1 &= 2(\partial_1[(K * \mathscr{W}_2 - K * \mathscr{W}_2(x)) \xi_{23}] - \partial_2[(K * \mathscr{W}_2 - K * \mathscr{W}_2(x)) \xi_{13}]), \\ \vartheta_2 &= 2(\partial_1[(K * \mathscr{W}_3 - K * \mathscr{W}_3(x)) \xi_{21}] - \partial_2[(K * \mathscr{W}_3 - K * \mathscr{W}_3(x)) \xi_{11}]), \\ \vartheta_3 &= 2(\partial_1[(K * \mathscr{W}_1 - K * \mathscr{W}_1(x)) \xi_{22}] - \partial_2[(K * \mathscr{W}_1 - K * \mathscr{W}_1(x)) \xi_{12}]), \end{aligned}$$

where identities (4.5) are valid for any  $x \in \mathbb{T}^2$ , and the products in the square brackets, in both (4.4) and (4.5), are understood as Young products (in the sense of Lemma E.1).

*Proof.* We have

$$\vartheta_1^\varepsilon = \xi_2^\varepsilon \overline{\xi_3^\varepsilon} - \xi_3^\varepsilon \overline{\xi_2^\varepsilon}, \quad \xi_k^\varepsilon = iG * (2\mathscr{W}_k^\varepsilon), \quad k = 2, 3.$$

By Lemma 4.1, for every  $A \in \mathbb{R}$ ,

$$\xi_2^\varepsilon \overline{\xi_3^\varepsilon} - \xi_3^\varepsilon \overline{\xi_2^\varepsilon} = 4\partial_1[(K * \mathscr{W}_2^\varepsilon - A) \partial_2 K * \mathscr{W}_3^\varepsilon] - 4\partial_2[(K * \mathscr{W}_2^\varepsilon - A) \partial_1 K * \mathscr{W}_3^\varepsilon].$$

Recalling that  $2\partial_2 K * \mathscr{W}_3^\varepsilon = \xi_{23}^\varepsilon$ ,  $2\partial_1 K * \mathscr{W}_3^\varepsilon = \xi_{13}^\varepsilon$ , and setting  $A = 0$ , we obtain the expression for  $\vartheta_1^\varepsilon$  in (4.1). The identities for  $\vartheta_2^\varepsilon$  and  $\vartheta_3^\varepsilon$  in (4.1) can be obtained likewise.

Further, by choosing and fixing  $x \in \mathbb{T}^2$ , and then setting  $A = K * \mathcal{W}_2^\varepsilon(x)$ , we deduce the first identity in (4.2). The last two identities in (4.2) can be obtained likewise.

Finally, for any test function  $\varphi$ ,

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \vartheta_1^\varepsilon(\varphi) &= \lim_{\varepsilon \rightarrow 0} \int (K * \mathcal{W}_2^\varepsilon - A) \xi_{23}^\varepsilon \partial_1 \varphi + \lim_{\varepsilon \rightarrow 0} \int (K * \mathcal{W}_2^\varepsilon - A) \xi_{13}^\varepsilon \partial_2 \varphi \\ &= \int (K * \mathcal{W}_2 - A) \xi_{23} \partial_1 \varphi + \int (K * \mathcal{W}_2 - A) \xi_{13} \partial_2 \varphi. \end{aligned}$$

Setting  $A = 0$ , we obtain the first expression in (4.4). Next, setting  $A = K * \mathcal{W}_2^\varepsilon(x)$ , where  $x \in \mathbb{T}^2$  is arbitrary, we obtain the first expression in (4.5). We also remark that by Lemma B.2,  $(K * \mathcal{W}_2 - A) \in \mathcal{X}^{1-\kappa}$  and  $\xi_{13}, \xi_{23} \in \mathcal{X}^{-\kappa}$ ,  $\kappa \in (0, \frac{1}{2})$ . The last two expressions in (4.4) and (4.5) can be obtained likewise.  $\square$

**Remark 4.3.** Identities (4.2) and (4.5) are important to compute the regularity of  $\vartheta$ . Namely, using these identities, we will show that  $\vartheta \in E_{-\kappa}$  and  $\vartheta^\varepsilon \rightarrow \vartheta$  in  $E_{-\kappa}$ .

**Proposition 4.4.** *Let  $\tilde{\mathcal{B}}^0$  be the class of test functions introduced in Definition C.2. Assume that  $0 < \kappa < \varkappa < 1$ ,  $\eta \in \mathcal{X}^{-\kappa}(\mathbb{T}^d)$  and  $u \in \mathcal{X}^\varkappa(\mathbb{T}^d)$ . Then, for any  $\nu \in (\kappa, \varkappa)$ ,*

$$(4.6) \quad \sup_{x \in \mathbb{T}^d, \varphi \in \tilde{\mathcal{B}}^0, \lambda \in (0, 1]} \frac{|[(u - u(x))\eta](\varphi_x^\lambda)|}{\lambda^{\nu-\kappa}} \leq [u]_\nu |\eta|_{-\kappa} \lesssim |u|_\nu |\eta|_{-\kappa},$$

where

$$[u]_\nu := \sup_{\{x \in \mathbb{T}^d, |y| \leq 1\}} \frac{|u(x+y) - u(x)|}{|y|^\nu}$$

and the product  $(u - u(x))\eta$  is understood as the Young product in the sense of Lemma E.1.

**Remark 4.5.** By classical argument, it follows that (4.6) is fulfilled for every  $\eta \in \mathcal{X}^{-\kappa}(\mathbb{T}^d)$  and every  $u$  belonging to the closure of  $\mathcal{X}^\varkappa(\mathbb{T}^d)$  in  $\mathcal{X}^\nu(\mathbb{T}^d)$ .

**Remark 4.6.** We borrowed the type of expression  $(u - u(x))\eta$  from [3]. It has a connection with the idea of germs, see [2, 3].

*Proof of Proposition 4.4.* By Lemma C.16,  $u \in \mathcal{X}^\nu(\mathbb{T}^d)$  for all  $\nu \in (0, \varkappa)$  and

$$\lim_{\varepsilon \rightarrow 0} |u - u^\varepsilon|_\nu = 0,$$

where  $u^\varepsilon = u * \rho_\varepsilon$ . Therefore, it is enough to obtain the bound (4.6) for every  $u = \rho_\varepsilon * \tilde{u}$  which is a mollification of some  $\tilde{u} \in \mathcal{X}^\nu(\mathbb{T}^d)$ .

Let us choose and fix  $\nu \in (\kappa, \varkappa)$  and define the function  $\psi : \mathbb{T}^d \times (0, 1] \times \overline{B_1(0)} \rightarrow \mathbb{R}$ ,

$$\psi(x, \lambda, y) = (u(x + \lambda y) - u(x))\lambda^{-\nu}.$$

Since  $|\psi(x, \lambda, y)| \leq [u]_\nu$  uniformly in all arguments, we have

$$\begin{aligned}
& \sup_{x \in \mathbb{T}^d, \varphi \in \tilde{\mathcal{B}}^0, \lambda \in (0,1]} \frac{|[(u - u(x))\eta](\varphi_x^\lambda)|}{\lambda^{\nu-\kappa}} \\
&= \sup_{x \in \mathbb{T}^d, \varphi \in \tilde{\mathcal{B}}^0, \lambda \in (0,1]} \frac{|(\psi(x, \lambda, \cdot)\eta(x + \lambda \cdot))(\varphi)|}{\lambda^{-\kappa}} \\
&= \sup_{\lambda \in (0,1]} \sup_{x \in \mathbb{T}^d, \varphi \in \tilde{\mathcal{B}}^0} \frac{|\eta(x + \lambda \cdot)(\varphi\psi(x, \lambda, \cdot))|}{\lambda^{-\kappa}} \\
&\leq [u]_\nu \sup_{\lambda \in (0,1]} \sup_{x \in \mathbb{T}^d, \varphi \in \tilde{\mathcal{B}}^0} \frac{|\eta(x + \lambda \cdot)(\varphi)|}{\lambda^{-\kappa}} \\
&\leq [u]_\nu \sup_{x \in \mathbb{T}^d, \varphi \in \tilde{\mathcal{B}}^0, \lambda \in (0,1]} \frac{|\eta(\varphi_x^\lambda)|}{\lambda^{-\kappa}} = [u]_\nu |\eta|_{-\kappa} \lesssim |u|_\nu |\eta|_{-\kappa},
\end{aligned}$$

where the last inequality is true by the equivalence of the norms  $\|u\|_{C^\nu(\mathbb{T}^d)}$  and  $|u|_\nu$ , see Lemma C.12.  $\square$

**Corollary 4.7.** *Assume that  $\eta \in \mathcal{X}^{-\kappa}(\mathbb{T}^2)$ ,  $\kappa \in (0, \frac{1}{2})$ , and let  $\eta^\varepsilon = \eta * \rho_\varepsilon$ . Then, for all  $\varepsilon \geq 0$  and  $j = 1, 2, 3$ ,*

$$\begin{aligned}
(4.7) \quad \sup_{x \in \mathbb{T}^2, \varphi \in \tilde{\mathcal{B}}^0, \lambda \in (0,1]} \frac{|[(K * \mathscr{W}_j^\varepsilon - K * \mathscr{W}_j^\varepsilon(x))\eta^\varepsilon](\varphi_x^\lambda)|}{\lambda^{1-\kappa}} &\lesssim |K * \mathscr{W}_j^\varepsilon|_{1-\frac{\kappa}{2}} |\eta^\varepsilon|_{-\frac{\kappa}{2}} \\
&\lesssim |\mathscr{W}_j|_{-1-\frac{\kappa}{2}} |\eta|_{-\frac{\kappa}{2}},
\end{aligned}$$

where  $\mathscr{W}_j^0 := \mathscr{W}_j$ ,  $\eta^0 := \eta$ .

*Proof.* Since  $\mathscr{W}_j \in \mathcal{X}^{-1-\kappa}(\mathbb{T}^2)$  for all  $\kappa \in (0, \frac{1}{2})$ , by Lemma D.5,  $K * \mathscr{W}_j \in \mathcal{X}^{1-\kappa}(\mathbb{T}^2)$ . By Proposition 4.4, the left-hand side of (4.7) is not greater than  $|K * \mathscr{W}_j^\varepsilon|_{1-\frac{\kappa}{2}} |\eta^\varepsilon|_{-\frac{\kappa}{2}}$ . By Lemma C.9, the latter expression is not greater than  $|K * \mathscr{W}_j|_{1-\kappa} |\eta|_{-\kappa}$ .  $\square$

**Proposition 4.8.** *Let  $\vartheta$  be the limit of  $\vartheta^\varepsilon := \xi^\varepsilon \times \overline{\xi^\varepsilon}$  in  $\mathcal{D}'(\mathbb{T}^2)$  whose existence is obtained in Lemma 4.2. Then, for every  $\kappa \in (0, \frac{1}{2})$ ,*

$$\vartheta \in E_{-\kappa}.$$

Furthermore, for all  $\varepsilon \geq 0$ ,  $\kappa \in (0, \frac{1}{2})$ , and for  $j = 1, 2, 3$ ,

$$(4.8) \quad \|\vartheta_j^\varepsilon\|_{-\kappa} \lesssim |\mathscr{W}_{j_1}|_{-1-\frac{\kappa}{2}} |\mathscr{W}_{j_2}|_{-1-\frac{\kappa}{2}},$$

$$(4.9) \quad \|\vartheta_j^\varepsilon - \vartheta_j\|_{-\kappa} \lesssim |\mathscr{W}_{j_1}^\varepsilon - \mathscr{W}_{j_1}|_{-1-\frac{\kappa}{2}} |\mathscr{W}_{j_2}|_{-1-\frac{\kappa}{2}} + |\mathscr{W}_{j_2}^\varepsilon - \mathscr{W}_{j_2}|_{-1-\frac{\kappa}{2}} |\mathscr{W}_{j_1}|_{-1-\frac{\kappa}{2}},$$

where  $j_1 := j + 1 \pmod{3}$  and  $j_2 := j + 2 \pmod{3}$ ,  $\vartheta^0 := \vartheta$ .

**Remark 4.9.** Let us observe that (4.8) implies the bound

$$\|\vartheta^\varepsilon\|_{-\kappa} \lesssim \|\mathscr{W}\|_{-1-\kappa}^2.$$

*Proof of Proposition 4.8.* By Lemma 4.2,  $\vartheta^\varepsilon$  has the representation (4.2) valid for any  $x \in \mathbb{T}^2$  and its limit  $\vartheta$  has the representation (4.5).

In what follows, we will compute the norm  $|\vartheta_1|_{-\kappa}$ . The norms  $|\vartheta_2|_{-\kappa}$  and  $|\vartheta_3|_{-\kappa}$  can be computed likewise. We will use the equivalence of the norms  $|\cdot|_{-\kappa}$  and

$|\cdot|_{\tilde{\mathcal{F}}^{-\kappa}}$ , obtained in Appendix C, where the latter norm is introduced in Definition C.2. By Lemmata 4.2 and C.7, for any  $\kappa \in (0, \frac{1}{2})$  and  $\varepsilon \geq 0$ ,

$$\begin{aligned}
|\vartheta_1^\varepsilon|_{-\kappa} &\lesssim |\vartheta_1^\varepsilon|_{\tilde{\mathcal{F}}^{-\kappa}} = \sup_{\substack{x \in \mathbb{T}^2, \lambda \in (0,1], \\ \varphi \in \tilde{\mathcal{B}}^1}} \frac{|\vartheta_1^\varepsilon(\varphi_x^\lambda)|}{\lambda^{-\kappa}} \\
&= 2 \sup_{\substack{x \in \mathbb{T}^2, \lambda \in (0,1], \\ \varphi \in \tilde{\mathcal{B}}^1}} \lambda^\kappa \left| \partial_1 \left[ (K * \mathscr{W}_2^\varepsilon - K * \mathscr{W}_2^\varepsilon(x)) \xi_{23}^\varepsilon \right] (\varphi_x^\lambda) \right. \\
&\quad \left. - \partial_2 \left[ (K * \mathscr{W}_2^\varepsilon - K * \mathscr{W}_2^\varepsilon(x)) \xi_{13}^\varepsilon \right] (\varphi_x^\lambda) \right| \\
(4.10) \quad &\leq 2 \sup_{\substack{x \in \mathbb{T}^2, \lambda \in (0,1], \\ \varphi \in \tilde{\mathcal{B}}^1}} \frac{\left| \left[ (K * \mathscr{W}_2^\varepsilon - K * \mathscr{W}_2^\varepsilon(x)) \xi_{23}^\varepsilon \right] (\partial_1 \varphi)_x^\lambda \right|}{\lambda^{1-\kappa}} \\
&\quad + 2 \sup_{\substack{x \in \mathbb{T}^2, \lambda \in (0,1], \\ \varphi \in \tilde{\mathcal{B}}^1}} \frac{\left| \left[ (K * \mathscr{W}_2^\varepsilon - K * \mathscr{W}_2^\varepsilon(x)) \xi_{13}^\varepsilon \right] (\partial_2 \varphi)_x^\lambda \right|}{\lambda^{1-\kappa}}.
\end{aligned}$$

We evaluate the first term on the right-hand side of formula (4.10). The second term can be evaluated likewise. By Corollary 4.7,

$$\begin{aligned}
&\sup_{\substack{x \in \mathbb{T}^2, \lambda \in (0,1], \\ \varphi \in \tilde{\mathcal{B}}^1}} \frac{\left| \left[ (K * \mathscr{W}_2^\varepsilon - K * \mathscr{W}_2^\varepsilon(x)) \xi_{23}^\varepsilon \right] (\partial_1 \varphi)_x^\lambda \right|}{\lambda^{1-\kappa}} \\
&\leq \sup_{\substack{x \in \mathbb{T}^2, \lambda \in (0,1], \\ \psi \in \tilde{\mathcal{B}}^0}} \frac{\left| \left[ (K * \mathscr{W}_2^\varepsilon - K * \mathscr{W}_2^\varepsilon(x)) \xi_{23}^\varepsilon \right] \psi_x^\lambda \right|}{\lambda^{1-\kappa}} \\
&\lesssim |\mathscr{W}_2^\varepsilon|_{-1-\frac{\kappa}{2}} |\xi_{23}^\varepsilon|_{-\frac{\kappa}{2}} \lesssim |\mathscr{W}_2|_{-1-\frac{\kappa}{2}} |\mathscr{W}_3|_{-1-\frac{\kappa}{2}}.
\end{aligned}$$

This proves (4.8). To show (4.9), we note that by (4.2) and (4.5),

$$(4.11) \quad \vartheta_1^\varepsilon - \vartheta_1 = 2\partial_1 \left[ (K * (\mathscr{W}_2^\varepsilon - \mathscr{W}_2) - K * (\mathscr{W}_2^\varepsilon(x) - \mathscr{W}_2(x))) \xi_{23}^\varepsilon \right]$$

$$(4.12) \quad - 2\partial_2 \left[ (K * (\mathscr{W}_2^\varepsilon - \mathscr{W}_2) - K * (\mathscr{W}_2^\varepsilon(x) - \mathscr{W}_2(x))) \xi_{13}^\varepsilon \right]$$

$$(4.13) \quad + 2\partial_1 \left[ (K * \mathscr{W}_2 - K * \mathscr{W}_2(x)) (\xi_{23}^\varepsilon - \xi_{23}) \right]$$

$$(4.14) \quad - 2\partial_2 \left[ (K * \mathscr{W}_2 - K * \mathscr{W}_2(x)) (\xi_{13}^\varepsilon - \xi_{13}) \right]$$

$$=: \theta_1 + \theta_2,$$

where  $\theta_1$  is the sum of the term on the right-hand side of (4.11) and the term in (4.12);  $\theta_2$  is the sum of the terms in (4.13) and (4.14). From (4.11–4.14) it follows that the argument on obtaining the bound for  $|\vartheta_1^\varepsilon|_{-\kappa}$  works for both  $\theta_1$  and  $\theta_2$ . In the case of  $\theta_1$ , in the above argument, we substitute  $\mathscr{W}_2^\varepsilon$  with  $\mathscr{W}_2^\varepsilon - \mathscr{W}_2$  to obtain the bound

$$|\theta_1|_{-\kappa} \lesssim |\mathscr{W}_2^\varepsilon - \mathscr{W}_2|_{-1-\frac{\kappa}{2}} \left( |\xi_{13}^\varepsilon|_{-\frac{\kappa}{2}} + |\xi_{13}^\varepsilon|_{-\frac{\kappa}{2}} \right) \lesssim |\mathscr{W}_2^\varepsilon - \mathscr{W}_2|_{-1-\frac{\kappa}{2}} |\mathscr{W}_3|_{-1-\frac{\kappa}{2}}.$$

In the case of  $\theta_2$ , in the argument on obtaining the bound for  $|\vartheta_1^\varepsilon|_{-\kappa}$ , we substitute  $\xi_{23}^\varepsilon$  with  $\xi_{23}^\varepsilon - \xi_{23}$  and  $\xi_{13}^\varepsilon$  with  $\xi_{13}^\varepsilon - \xi_{13}$ . Thus, we obtain

$$|\theta_2|_{-\kappa} \lesssim |\mathscr{W}_2^\varepsilon|_{-1-\frac{\kappa}{2}} |\xi_{23}^\varepsilon - \xi_{23}|_{-\frac{\kappa}{2}} + |\mathscr{W}_2^\varepsilon|_{-1-\frac{\kappa}{2}} |\xi_{13}^\varepsilon - \xi_{13}|_{-\frac{\kappa}{2}} \lesssim |\mathscr{W}_2|_{-1-\frac{\kappa}{2}} |\mathscr{W}_3^\varepsilon - \mathscr{W}_3|_{-1-\frac{\kappa}{2}}.$$

This implies (4.9).  $\square$

We have the following corollary which implies Lemma 3.8.

**Corollary 4.10.** *Let  $\zeta$  be the limit of  $\zeta^\varepsilon := 2G * (\xi^\varepsilon \times \bar{\xi}^\varepsilon)$  in  $\mathscr{D}'(\mathbb{T}^2)$ . Then,*

$$\zeta \in E_{1-\kappa}.$$

Furthermore, for all  $\varepsilon \geq 0$  and  $\kappa \in (0, \frac{1}{2})$ ,

$$\begin{aligned} \|\zeta^\varepsilon\|_{1-\kappa} &\lesssim \|\mathscr{W}\|_{-1-\kappa}^2, \\ \|\zeta_j^\varepsilon - \zeta_j\|_{1-\kappa} &\lesssim |\mathscr{W}_{j_1}^\varepsilon - \mathscr{W}_{j_1}|_{-1-\frac{\kappa}{2}} |\mathscr{W}_{j_2}|_{-1-\frac{\kappa}{2}} + |\mathscr{W}_{j_2}^\varepsilon - \mathscr{W}_{j_2}|_{-1-\frac{\kappa}{2}} |\mathscr{W}_{j_1}|_{-1-\frac{\kappa}{2}}, \end{aligned}$$

where  $j = 1, 2, 3$ ,  $j_1 := j + 1 \pmod{3}$ ,  $j_2 := j + 2 \pmod{3}$ ,  $\zeta^0 := \zeta$ .

## 5. RATE OF CONVERGENCE AND PROOF OF THE MAIN RESULT

The aim of this section is to prove Theorem 1.3. This is done at the end of the section.

The next result is important for determining the rate of convergence (1.7). More specifically, it is used to prove the convergence  $R^\varepsilon \rightarrow R$  in  $E_{1-\kappa}$  and evaluate the rate of this convergence.

**Lemma 5.1.** *The following inequalities are satisfied for all  $\varepsilon > 0$  and  $\kappa \in (0, \frac{1}{2})$  :*

$$(5.1) \quad \|\xi^\varepsilon - \xi\|_{-\kappa} \lesssim \varepsilon^{\frac{\kappa}{2}} \|\mathscr{W}\|_{-1-\kappa}^2,$$

$$(5.2) \quad \|\vartheta^\varepsilon - \vartheta\|_{-\kappa} \lesssim \varepsilon^{\frac{\kappa}{2}} \|\mathscr{W}\|_{-1-\kappa}^2,$$

$$(5.3) \quad \|\zeta^\varepsilon - \zeta\|_{1-\kappa} \lesssim \varepsilon^{\frac{\kappa}{2}} \|\mathscr{W}\|_{-1-\kappa}^2.$$

*Proof.* By Lemma B.3,

$$|\mathscr{W}_j^\varepsilon - \mathscr{W}_j|_{-1-\kappa} \lesssim \varepsilon^{\frac{\kappa}{2}} |\mathscr{W}_j|_{-1-\frac{\kappa}{2}}, \quad j = 1, 2, 3, \quad \text{a.s.}$$

Since  $\xi_{kj} = 2G_k * \mathscr{W}_j$ , we have

$$|\xi_{kj}^\varepsilon - \xi_{kj}|_{-\kappa} = 2|G_k * \mathscr{W}_j^\varepsilon - G_k * \mathscr{W}_j|_{-\kappa} \lesssim |\mathscr{W}_j^\varepsilon - \mathscr{W}_j|_{-1-\kappa} \lesssim |\varepsilon|^{\frac{\kappa}{2}} |\mathscr{W}_j|_{-1-\frac{\kappa}{2}}.$$

Further, by Proposition 4.8,

$$\begin{aligned} |\vartheta_1^\varepsilon - \vartheta_1|_{-\kappa} &\lesssim |\mathscr{W}_2^\varepsilon - \mathscr{W}_2|_{-1-\frac{\kappa}{2}} |\mathscr{W}_3^\varepsilon|_{-1-\frac{\kappa}{2}} + |\mathscr{W}_3^\varepsilon - \mathscr{W}_3|_{-1-\frac{\kappa}{2}} |\mathscr{W}_2|_{-1-\frac{\kappa}{2}} \\ &\lesssim \varepsilon^{\frac{\kappa}{2}} |\mathscr{W}_2|_{-1-\frac{\kappa}{2}} |\mathscr{W}_3|_{-1-\frac{\kappa}{2}}. \end{aligned}$$

Likewise, we evaluate  $|\vartheta_j^\varepsilon - \vartheta_j|_{-\kappa}$  for  $j = 2, 3$ . This implies (5.1). Finally, recalling that  $\zeta^\varepsilon = 2G * \vartheta^\varepsilon$ , we obtain (5.3).  $\square$

**Lemma 5.2.** *Let  $\delta > 0$ ,  $\kappa \in (0, \frac{1}{2})$ , and let  $\Omega_\delta \subset \Omega$  be the set constructed in Proposition 3.10. Further let  $\sigma, \varsigma > 0$  be numbers, defined in the same proposition, for which the maps  $\Gamma_{\varepsilon, \sigma}$  and  $\Gamma_{0, \sigma}$  possess unique fixed points  $R^\varepsilon$  and, respectively,  $R$*



in  $M_{\sigma, 1-\kappa}$  for  $|\gamma| < \varsigma$ . Then, there exists a constant  $C_\delta > 0$ , depending only on  $\delta$ , such that for all  $\omega \in \Omega_\delta$ ,

$$(5.4) \quad \|R - R^\varepsilon\|_{1-\kappa} \leq C_\delta \varepsilon^{\frac{\kappa}{2}}.$$

*Proof.* Let us choose and fix  $\varepsilon > 0$ . Let us recall that  $R^\varepsilon$  is a solution of equation (3.10) and  $R$  is a solution of equation (3.10) with  $\varepsilon = 0$ . We write  $\xi$  and  $\zeta$  instead of  $\xi^0$  and  $\zeta^0$  when  $\varepsilon = 0$ . Similar to (3.16), keeping the same constants  $C$  and  $C_\Lambda$  as in (3.16), using equation (3.10) and recalling that  $|\gamma| < \frac{\sigma^2}{16\Lambda}$ , we infer that there exists  $\tilde{C} > 0$  such that

$$\begin{aligned} \|R - R^\varepsilon\|_{1-\kappa} &\leq 2C(\sigma + |\gamma| \|\xi^\varepsilon\|_{-\kappa}) \|R - R^\varepsilon\|_{1-\kappa} + 2C\sigma|\gamma| \|\xi - \xi^\varepsilon\|_{-\kappa} + |\gamma|^2 \|\zeta - \zeta^\varepsilon\|_{1-\kappa}, \\ &\leq 2C_\Lambda(\sigma + \sigma^2) \|R - R^\varepsilon\|_{1-\kappa} + \tilde{C}(\|\xi - \xi^\varepsilon\|_{-\kappa} + \|\zeta - \zeta^\varepsilon\|_{1-\kappa}). \end{aligned}$$

Recall that  $C_\Lambda$  satisfies (3.15) so that  $2C_\Lambda(\sigma + \sigma^2) < 1$ . By Lemma 5.1 and the choice (3.14) of the set  $\Omega_\delta$ , we obtain (5.4).  $\square$

In order to conclude the proof of Theorem 1.3 we finally need the following result on the rate of convergence of  $r^\varepsilon$ . It uses the change of variables (3.9).

**Proposition 5.3.** *Let the objects  $\delta > 0$ ,  $\kappa \in (0, \frac{1}{2})$ ,  $\Omega_\delta \subset \Omega$ ,  $\varsigma > 0$ ,  $R^\varepsilon$ , and  $R$  be as in Lemma 5.2. Then, for every  $\varepsilon > 0$  and  $\omega \in \Omega_\delta$ , the function*

$$r^\varepsilon = R^\varepsilon - \gamma \xi^\varepsilon,$$

where  $|\gamma| < \varsigma$ , solves (mCR). Moreover, there exists a limit  $r := \lim_{\varepsilon \rightarrow 0} r^\varepsilon$  in  $E_{-\kappa}$  and a constant  $\hat{C}_\delta > 0$ , depending only on  $\delta$ , such that for all  $\omega \in \Omega_\delta$ ,

$$(5.5) \quad \|r^\varepsilon - r\|_{-\kappa} \leq \hat{C}_\delta \varepsilon^{\frac{\kappa}{2}}.$$

*Proof.* By Lemma 3.4, (3.2) and (3.10) are equivalent for every  $\varepsilon > 0$  (provided that  $a, b, c$  are as in the aforementioned lemma). Since for all  $\varepsilon > 0$ ,  $R^\varepsilon \in E_{1-\kappa}$  and  $\xi^\varepsilon \in E_{1-\kappa}$  (see Remark 3.6), by Proposition 3.1,  $r^\varepsilon$  solves (mCR). Define

$$r := R - \gamma \xi.$$

Then, since  $|\mathscr{W}_j|_{-1-\kappa}$ ,  $j = 1, 2, 3$ , are bounded in  $\omega \in \Omega_\delta$ , (5.5) is fulfilled on  $\Omega_\delta$  by Lemmata 5.1 and 5.2.  $\square$

*Proof of Theorem 1.3.* Let the objects  $\delta > 0$ ,  $\Omega_\delta \subset \Omega$ ,  $\varsigma > 0$ ,  $r^\varepsilon$ , and  $r$  be as in Proposition 5.3 and  $|\gamma| < \varsigma$ . By Proposition 5.3,  $r^\varepsilon$  solves (mCR) and there exists a limit  $r := \lim_{\varepsilon \rightarrow 0} r^\varepsilon$  in  $E_{-\kappa}$ . By Lemma C.12,  $\mathcal{Z}^{-\kappa}(\mathbb{T}^2)$  and  $C^{-\kappa}(\mathbb{T}^2)$  coincide for  $\kappa \in (0, \frac{1}{2})$ . Therefore,  $r^\varepsilon \rightarrow r$  also in  $C^{-\kappa}(\mathbb{T}^2)$ . The bound (5.5) is therefore exactly what is required in (1.7). The theorem is proved.  $\square$

## 6. BÄCKLUND-TRANSFORMED, WHITE-NOISE PERTURBED LANDAU-LIFSHITZ-GILBERT EQUATION

In this section we prove the existence of solution to equation (1.3). Here, we additionally assume that  $\mathscr{W}_1$  and  $\mathscr{W}_2$  have zero means over  $\mathbb{T}^2$ .

**Theorem 6.1.** *Let  $\mathscr{W} = (\mathscr{W}_1, \mathscr{W}_2, \mathscr{W}_3)$  be a 3D white noise on  $\mathbb{T}^2$  with the zero mean over  $\mathbb{T}^2$  and let  $r$  be the solution to (CR) constructed in Theorem 1.3. Then,*

$$(6.1) \quad B = 2i\overline{G} * r$$

belongs to  $E_{1-\kappa}$  and is a solution to (1.3). Moreover,  $\text{Im } B = 0$ .

*Proof.* Note that since  $G = 2\partial_{\bar{z}}K$ , by (3.2),  $\int_{\mathbb{T}^2} r^\varepsilon = 0$ . It is straightforward to verify that if we set

$$(6.2) \quad r^\varepsilon = \frac{1}{2}(\partial_y B^\varepsilon + i\partial_x B^\varepsilon) = i\partial_{\bar{z}} B^\varepsilon,$$

then, since  $\partial_{\bar{z}} r^\varepsilon = \frac{i}{4}\Delta B^\varepsilon$  and  $r^\varepsilon \times \overline{r^\varepsilon} = \frac{i}{2}(\partial_x B^\varepsilon \times \partial_y B^\varepsilon)$ , we obtain that  $B^\varepsilon$  satisfies the equation

$$\Delta B^\varepsilon = 2\partial_x B^\varepsilon \times \partial_y B^\varepsilon + 4\gamma \mathscr{W}^\varepsilon,$$

where  $\mathscr{W}^\varepsilon = \mathscr{W} * \rho_\varepsilon$ . By (6.2) and Lemma A.1,  $B^\varepsilon = 2i\overline{G} * r^\varepsilon$  is a solution. Passing to the limit as  $\varepsilon \rightarrow 0$ , we obtain (6.1). Note that since  $r \in E_{-\kappa}$  and  $i\overline{G} = 2i\partial_{\bar{z}}K$ ,  $B \in E_{1-\kappa}$ .

Finally, by (mCR),  $\text{Re}(\partial_{\bar{z}} r^\varepsilon) = 0$ , which is equivalent to  $\partial_x p^\varepsilon = \partial_y q^\varepsilon$ , where  $p^\varepsilon = \text{Re } r^\varepsilon$  and  $q^\varepsilon = \text{Im } r^\varepsilon$ . On the other hand,

$$\text{Im}(i\overline{G} * r^\varepsilon) = \partial_x K * p^\varepsilon - \partial_y K * q^\varepsilon = K * (\partial_x p^\varepsilon - \partial_y q^\varepsilon) = 0$$

which proves that  $\text{Im } B = 0$ . □

In the appendices, we collect some important useful results.

#### APPENDIX A. ON THE CAUCHY-RIEMANN OPERATOR

**Lemma A.1.** *Let  $G$  be the Green function defined by (2.5). If  $f \in C^\alpha(\mathbb{T}^2)$ ,  $\alpha \in (0, 1)$ ,  $C \in \mathbb{C}$  is a constant and a function  $r : \mathbb{T}^2 \rightarrow \mathbb{C}$  satisfies*

$$(A.1) \quad r = G * f + C$$

then  $r \in C^{1+\alpha}(\mathbb{T}^2)$  and satisfies the identity

$$-2\partial_{\bar{z}} r = f - [f],$$

where  $[f] := (4\pi^2)^{-1} \int f$  denotes the average over  $\mathbb{T}^2$ .

*Proof.* By Lemmata C.12 and D.3,  $r \in C^{1+\alpha}(\mathbb{T}^2)$ . Note that

$$(A.2) \quad \partial_z \partial_{\bar{z}} = \frac{1}{4}\Delta \text{ on } C^{2+\alpha}(\mathbb{T}^2).$$

Thus, by applying the operator  $2\partial_{\bar{z}}$  to both sides of equality (A.1) we infer that

$$2\partial_{\bar{z}} r = 2\partial_{\bar{z}} G * f = 4\partial_{\bar{z}} \partial_z K * (f - [f]) = \Delta(K * (f - [f])) = [f] - f.$$

To justify the last step, we observe that if  $\int_{\mathbb{T}^2} g = 0$ , then

$$(A.3) \quad -\Delta u = g \quad \iff \quad u = K * g + C.$$

for some constant  $C \in \mathbb{R}$ . □

APPENDIX B. ON THE WHITE NOISE ON  $\mathbb{T}^d$ ,  $d \geq 2$ 

**Lemma B.1.** *Let  $\mathscr{W}$  be a 3D Gaussian white noise, as in Definition 1.1. If  $\mathscr{W}^\varepsilon := \mathscr{W}_j * \rho_\varepsilon$ ,  $j = 1, 2, 3$ , are mollified white noises, where  $\rho_\varepsilon$  is a standard mollifier, then,*

$$(B.1) \quad \mathscr{W}_j^\varepsilon = \eta_j + \rho_\varepsilon * \operatorname{Re} \left[ \sum_{k \in \mathbb{Z}^d \setminus \{0\}} \eta_{kj} e^{i(k,x)} \right].$$

*Proof.* Applying the convolution with  $\rho_\varepsilon$  in (1.6), we obtain (B.1).  $\square$

**Lemma B.2.** *Let  $\mathscr{W} : \Omega \rightarrow \mathscr{D}'(\mathbb{T}^d)$  be a Gaussian white noise. Then, for any  $\kappa > 0$ ,  $\mathscr{W} \in C^{-\frac{d}{2}-\kappa}(\mathbb{T}^d)$  a.s. Moreover, for any  $\delta > 0$ , there exists  $\Lambda > 0$  such that*

$$\mathbb{P}(\|\mathscr{W}\|_{C^{-\frac{d}{2}-\kappa}(\mathbb{T}^d)} < \Lambda) > 1 - \delta.$$

*Proof.* The first part of the Lemma is known, see e.g., [3] or [20], according to which, for any  $\kappa > 0$ ,

$$\mathbb{P}\{\mathscr{W} \in C^{-\frac{d}{2}-\kappa}(\mathbb{T}^d)\} = 1.$$

The second part trivially follows from the first but since we use it, we show a proof.

$$\{\mathscr{W} \in C^{-\frac{d}{2}-\kappa}(\mathbb{T}^d)\} = \bigcup_{N=1}^{\infty} A_N, \quad \text{where } A_N = \{\|\mathscr{W}\|_{C^{-\frac{d}{2}-\kappa}(\mathbb{T}^d)} < N\}.$$

Since  $A_N \subset A_{N+1}$ , by the continuity of the measure  $\mathbb{P}$ ,

$$\lim_{N \rightarrow \infty} \mathbb{P}\{\|\mathscr{W}\|_{C^{-\frac{d}{2}-\kappa}(\mathbb{T}^d)} < N\} = 1.$$

$\square$

**Lemma B.3.** *Let  $\mathscr{W} : \Omega \rightarrow \mathscr{D}'(\mathbb{T}^d)$  be a Gaussian white noise. Then, for any  $\kappa > 0$ ,  $\mathscr{W} \in \mathscr{L}^{-\frac{d}{2}-\kappa}(\mathbb{T}^d)$  a.s. Furthermore, for any  $\varepsilon > 0$  and  $\kappa > \varkappa > 0$ ,*

$$(B.2) \quad |\mathscr{W}^\varepsilon - \mathscr{W}|_{-\frac{d}{2}-\kappa} \lesssim \varepsilon^{\kappa-\varkappa} |\mathscr{W}|_{-\frac{d}{2}-\varkappa}, \quad \text{a.s.}$$

*Proof.* Let us choose and fix  $\kappa > \varkappa > 0$  and let  $\ell = [\frac{d}{2} + \kappa + 1]$ , where  $[\cdot]$  denotes the integer part. By Lemma B.2,  $\mathscr{W} \in \mathscr{L}^{-\frac{d}{2}-\varkappa} \cap \mathscr{L}^{-\frac{d}{2}-\kappa}$ . By Proposition C.11, one can compute  $|\mathscr{W}^\varepsilon - \mathscr{W}|_{-\frac{d}{2}-\kappa}$  by taking the supremum over  $\varphi \in \mathscr{B}^{\ell+1}$ . First, we note that

$$\mathscr{W}^\varepsilon(\varphi_x^\lambda) = \mathscr{W}(\rho_\varepsilon * \varphi_x^\lambda) = \mathscr{W} \left( \int \rho(z) \varphi_x^\lambda(\cdot - \varepsilon z) dz \right) = \int \rho(z) \mathscr{W}(\varphi_{x+\varepsilon z}^\lambda) dz,$$

where  $\rho$  is a standard mollifier. Define  $\psi : \mathbb{T}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ ,

$$\psi(y; w) := \frac{\varphi(y-w) - \varphi(y)}{|w|^{\kappa-\varkappa}}.$$

We have

$$\begin{aligned}
|\mathscr{W}^\varepsilon - \mathscr{W}|_{\mathscr{X}^{-\frac{d}{2}-\kappa}} &\lesssim \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \mathscr{B}^{\ell+1}}} \frac{|(\mathscr{W}^\varepsilon - \mathscr{W})(\varphi_x^\lambda)|}{\lambda^{-\frac{d}{2}-\kappa}} \\
&\leq \int \rho(z) dz \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \mathscr{B}^{\ell+1}}} \frac{|\mathscr{W}(\varphi_{x+\varepsilon z}^\lambda - \varphi_x^\lambda)|}{\lambda^{-\frac{d}{2}-\kappa}} \\
&\leq \varepsilon^{\kappa-\varkappa} \int \rho(z) |z|^{\kappa-\varkappa} dz \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \mathscr{B}^{\ell+1}}} \frac{|\mathscr{W}(\psi_x^\lambda(\cdot; \frac{\varepsilon z}{\lambda}))|}{\lambda^{-\frac{d}{2}-\varkappa}} \\
\text{(B.3)} \quad &\lesssim \varepsilon^{\kappa-\varkappa} \left[ \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \mathscr{B}^{\ell+1}, |w| \leq 1}} \frac{|\mathscr{W}(\psi_x^\lambda(\cdot; w))|}{\lambda^{-\frac{d}{2}-\varkappa}} + \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \mathscr{B}^{\ell+1}, |w| > 1}} \frac{|\mathscr{W}(\psi_x^\lambda(\cdot; w))|}{\lambda^{-\frac{d}{2}-\varkappa}} \right].
\end{aligned}$$

Since for  $\varphi \in \mathscr{B}^{\ell+1}$ ,  $\|D^l \varphi\|_\infty \leq 1$ ,  $|l| = 1, 2, \dots, \ell + 1$ , we have the following bound uniformly in  $|w| \leq 1$ :

$$\text{(B.4)} \quad \|\partial_1^k \psi(\cdot; w)\|_\infty \lesssim \|D^l \varphi\|_\infty \lesssim 1, \quad |l| = |k| + 1, \quad |k| = 0, 1, \dots, \ell,$$

where by  $\partial_1^k$ , we understand a partial derivative of order  $|k|$  with respect to the first argument of  $\psi$  and by  $D^l$  we understand a derivative of order  $|l|$  of  $\varphi$ . By (B.4), the first supremum in the line (B.3) is bounded by

$$\sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \psi \in \mathscr{B}^\ell}} \frac{|\mathscr{W}(\psi_x^\lambda)|}{\lambda^{-\frac{d}{2}-\varkappa}} \lesssim |\mathscr{W}|_{\mathscr{X}^{-\frac{d}{2}-\varkappa}},$$

where the last bound is implied by Proposition C.11, as  $\ell \geq [\frac{d}{2} + \varkappa + 1]$ . The second supremum in the line (B.3) is bounded by

$$\sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \mathscr{B}^{\ell+1}, |w| > 1}} \frac{|\mathscr{W}((\varphi-w)_x^\lambda)|}{\lambda^{-\frac{d}{2}-\varkappa}} + \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \mathscr{B}^{\ell+1}}} \frac{|\mathscr{W}(\varphi_x^\lambda)|}{\lambda^{-\frac{d}{2}-\varkappa}} \lesssim |\mathscr{W}|_{\mathscr{X}^{-\frac{d}{2}-\varkappa}}.$$

This immediately implies (B.2).  $\square$

## APPENDIX C. ON HÖLDER-ZYGMUND SPACES AND NORMS

In [2, Definition A.9], the sets  $\mathscr{B}^r$ ,  $\mathscr{B}_\alpha$ , and  $\mathscr{B}_\alpha^r$  were defined in a different manner. Here we would like to show the equivalence of the corresponding Hölder-Zygmund norms.

**Remark C.1.** Let  $B_1(0) \subset \mathbb{R}^d$  be an open ball of radius 1 centered at the origin. If  $\varphi \in \mathscr{D}(B_1(0))$ , then  $\varphi_0^\lambda \in \mathscr{D}(B_1(0))$  for  $\lambda \in (0, 1)$ . Since  $B_1(0) \subset [-\pi, \pi]^d$ , both  $\varphi$  and  $\varphi_0^\lambda$  can be extended  $2\pi$ -periodically, in both variables, from  $[-\pi, \pi]^d$  to  $\mathbb{R}^d$ . This way,  $\varphi_y^\lambda = \varphi_0^\lambda(\cdot - y)$ ,  $\lambda \in (0, 1]$ , can be regarded as a function from  $\mathscr{D}(\mathbb{T}^d)$ .

**Definition C.2.** For  $r \in \mathbb{N} \cup \{0\}$  and  $\alpha \in \mathbb{R}$ , we define

$$\begin{aligned}\tilde{\mathcal{B}}^r &:= \{\varphi \in \mathcal{D}(B_1(0)) : \|\partial^k \varphi\|_\infty \leq 1, \text{ for all } 0 \leq |k| \leq r\}, \\ \tilde{\mathcal{B}}_\alpha &:= \{\varphi \in \mathcal{D}(B_1(0)) : \int_{\mathbb{T}^d} \varphi(z) z^k dz = 0, \text{ for all } 0 \leq |k| \leq \alpha\}, \text{ if } \alpha \geq 0, \\ \tilde{\mathcal{B}}_\alpha^r &:= \tilde{\mathcal{B}}^r \cap \tilde{\mathcal{B}}_\alpha, \text{ if } \alpha \geq 0,\end{aligned}$$

provided that  $\mathcal{D}(B_1(0)) \subset \mathcal{D}(\mathbb{T}^d)$  in the sense of Remark C.1.

**Definition C.3.** Define the Hölder-Zygmund norms  $|\cdot|_{\mathcal{Z}^{\alpha,x}}$  as follows. If  $\alpha \geq 0$ , then

$$(C.1) \quad |f|_{\mathcal{Z}^{\alpha,x}} := \sup_{y \in \mathbb{T}^d, \psi_x \in \tilde{\mathcal{B}}^0} |f(\psi_y)| + \sup_{\substack{y \in \mathbb{T}^d, \lambda \in (0,1], \\ \varphi_x \in \tilde{\mathcal{B}}_\alpha^0}} \frac{|f(\varphi_y^\lambda)|}{\lambda^\alpha}.$$

If  $\alpha < 0$ , then, with  $r = \lceil -\alpha + 1 \rceil$ ,

$$|f|_{\mathcal{Z}^{\alpha,x}} := \sup_{\substack{y \in \mathbb{T}^d, \lambda \in (0,1], \\ \varphi_x \in \tilde{\mathcal{B}}^r}} \frac{|f(\varphi_y^\lambda)|}{\lambda^\alpha}.$$

We define the Hölder-Zygmund space  $\tilde{\mathcal{Z}}^\alpha(\mathbb{T}^d)$  as a space of distributions  $f \in \mathcal{D}'(\mathbb{T}^d)$  for which  $|f|_{\mathcal{Z}^{\alpha,0}} < \infty$ . For simplicity of notation, we set

$$|f|_{\tilde{\mathcal{Z}}^\alpha} := |f|_{\mathcal{Z}^{\alpha,0}}.$$

**Remark C.4.** Remark that  $|\cdot|_{\tilde{\mathcal{Z}}^\alpha}$  is the same norm as in [2, Definition A.9].

**Remark C.5.** For  $\alpha \in \mathbb{R}$ , the Hölder-Zygmund seminorm  $|\cdot|_{\alpha,\mathcal{K}}$  is defined in exactly the same way as the norm  $|\cdot|_{\tilde{\mathcal{Z}}^\alpha}$ , see Definition C.3, but with  $\mathbb{T}^d$  replaced with a compact  $\mathcal{K} \subset \mathbb{R}^d$ . For the precise definition of  $|\cdot|_{\alpha,\mathcal{K}}$  and the Hölder-Zygmund space  $\mathcal{Z}^\alpha(\mathbb{R}^d)$ , we refer to [2, Definition 2.1].

**Remark C.6.** Note that  $\tilde{\mathcal{Z}}^\alpha(\mathbb{T}^d) = \mathcal{Z}^\alpha(\mathbb{R}^d) \cap \mathcal{D}'(\mathbb{T}^d)$ . Indeed, the norm  $|\cdot|_{\tilde{\mathcal{Z}}^\alpha}$  is the restriction of the seminorm  $|\cdot|_{\alpha,\mathbb{T}^d}$  to periodic distributions. Conversely, each seminorm  $|\cdot|_{\alpha,\mathcal{K}}$ , where  $\mathcal{K} \subset \mathbb{R}^d$  is a compact, can be bounded by  $|\cdot|_{\alpha,\mathbb{T}^d}$  due to the periodicity of the distributions.

Let  $\alpha \notin \mathbb{N}$ . The following lemma is about the equivalence of the norms  $|\cdot|_{\tilde{\mathcal{Z}}^\alpha}$  and  $|\cdot|_{\mathcal{Z}^\alpha}$ , where the latter is the norm introduced in Definition 2.2.

**Lemma C.7.** *For every  $\alpha \in \mathbb{R}$ ,  $\alpha \notin \mathbb{N}$ ,  $\mathcal{Z}^\alpha = \tilde{\mathcal{Z}}^\alpha$  and the norms  $|\cdot|_{\tilde{\mathcal{Z}}^\alpha}$  and  $|\cdot|_{\mathcal{Z}^\alpha}$  are equivalent.*

*Proof.* First, we show that  $|\cdot|_{\tilde{\mathcal{Z}}^\alpha}$  and  $|\cdot|_{\mathcal{Z}^{\alpha,x}}$  is the same norm. Indeed, let  $\alpha < 0$  and  $r = \lceil -\alpha + 1 \rceil$ . We have

$$|f|_{\mathcal{Z}^{\alpha,x}} = \sup_{\substack{y \in \mathbb{T}^d, \lambda \in (0,1], \\ \varphi_x \in \tilde{\mathcal{B}}^r}} \frac{|f(\varphi_y^\lambda)|}{\lambda^\alpha} = \sup_{\substack{y \in \mathbb{T}^d, \lambda \in (0,1], \\ \varphi_x \in \tilde{\mathcal{B}}^r}} \frac{|f(\varphi_{y+x}^\lambda)|}{\lambda^\alpha} = |f|_{\tilde{\mathcal{Z}}^\alpha}.$$

The latter identity holds by the  $2\pi$ -periodicity of  $\varphi_{y+x}$  in both variables. For  $\alpha \geq 0$ , the proof is the same.

Next, let us choose and fix a finite cover of  $\mathbb{T}^d$  by open balls  $\{B_{x_k}(1)\}$ . Furthermore, we choose a finite partition of unity  $\{\chi_k\}$  of  $\mathbb{T}^d$  subordinated to the cover  $\{B_{x_k}(1)\}$ . Let  $f \in \mathcal{Z}^\alpha$ ,  $\alpha < 0$ , and  $r = \lceil -\alpha + 1 \rceil$ . Then

$$\begin{aligned} |f|_{\mathcal{Z}^\alpha} &= \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \tilde{\mathcal{B}}^r}} \frac{|f(\varphi_x^\lambda)|}{\lambda^\alpha} \leq \sum_k \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \tilde{\mathcal{B}}^r}} \frac{|f((\varphi\chi_k)_x^\lambda)|}{\lambda^\alpha} \\ (C.2) \quad &\leq \sum_k \sup_{\substack{x \in \mathbb{T}^d, \lambda \in (0,1], \\ \psi_{x_k} \in \tilde{\mathcal{B}}^r}} \frac{|f(\psi_x^\lambda)|}{\lambda^\alpha} = \sum_k |f|_{\mathcal{Z}^\alpha, x_k} \lesssim |f|_{\tilde{\mathcal{Z}}^\alpha}. \end{aligned}$$

Now let  $\alpha \geq 0$ ,  $\alpha \notin \mathbb{N}$ ,  $n = \lfloor \alpha \rfloor$ , and, for a fixed  $x \in \mathbb{R}^d$ , let  $T_{n,x}$  be the Taylor polynomial of degree  $n$ , based at 0, for the function  $f(x + \cdot)$ . Since, the test function  $\varphi$  in (2.2) annihilates monomials of order up to  $n$ , Taylor's formula (with Lagrange remainder) implies

$$\begin{aligned} \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \tilde{\mathcal{B}}_\alpha^0}} \frac{|f(\varphi_x^\lambda)|}{\lambda^\alpha} &= \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \tilde{\mathcal{B}}_\alpha^0}} \frac{|(f(x + \lambda \cdot) - T_{n,x}(\lambda \cdot))(\varphi)|}{\lambda^\alpha} \\ (C.3) \quad &\lesssim \sum_{|\beta|=n} [D^\beta f]_\alpha \lesssim \|f\|_{C^\alpha(\mathbb{T}^d)} \lesssim |f|_{\tilde{\mathcal{Z}}^\alpha}, \end{aligned}$$

where  $\beta$  is a multiindex and  $D^\beta$  is a partial derivative of order  $|\beta|$ . The evaluation of the first term in (2.2) can be done by the same argument as in (C.2). Thus, we have proved that

$$|f|_{\mathcal{Z}^\alpha} \lesssim |f|_{\tilde{\mathcal{Z}}^\alpha}.$$

Let us prove the inverse inequality. Let  $f \in \mathcal{Z}^\alpha$ ,  $\lambda \in (0, 1]$ , and let  $\lambda_{n+1} = 2^{-(n+1)}$ ,  $\lambda_n = 2^{-n}$ ,  $n \in \mathbb{N}_0$ , be two consecutive points from  $[0, 1]_{\mathbb{D}}$  with the property

$$\lambda_{n+1} < \lambda \leq \lambda_n.$$

Note that  $\frac{\lambda_n}{\lambda} < \frac{\lambda_n}{\lambda_{n+1}} = 2$ .

Let  $\alpha < 0$ ,  $r = \lceil -\alpha + 1 \rceil$ , and  $\varphi \in \tilde{\mathcal{B}}^r$ . We have

$$\frac{|f(\varphi_x^\lambda)|}{\lambda^\alpha} \leq \left(\frac{\lambda_n}{\lambda}\right)^d \frac{|f(\psi_x^{\lambda_n})|}{\lambda_n^\alpha} < 2^d \frac{|f(\psi_x^{\lambda_n})|}{\lambda_n^\alpha}, \quad \text{where } \psi(y) = \varphi\left(\frac{\lambda_n}{\lambda}y\right).$$

Note that since  $1 \leq \frac{\lambda_n}{\lambda} < 2$ ,

$$\text{supp } \psi \subset B_1(0) \subset [-\pi, \pi]^d \quad \text{and} \quad \|D^k \psi\|_\infty \leq 2^{|k|} \|D^k \varphi\|_\infty \leq 2^r.$$

Therefore,

$$(C.4) \quad |f|_{\tilde{\mathcal{Z}}^\alpha} \lesssim |f|_{\mathcal{Z}^\alpha}.$$

Now let  $\alpha \geq 0$  and  $\varphi \in \tilde{\mathcal{B}}_\alpha^0$ . We have

$$\frac{|f(\varphi_x^\lambda)|}{\lambda^\alpha} \leq \left(\frac{\lambda_{n+1}}{\lambda}\right)^d \frac{|f(\psi_x^{\lambda_{n+1}})|}{\lambda_{n+1}^\alpha} < \frac{|f(\psi_x^{\lambda_{n+1}})|}{\lambda_{n+1}^\alpha}, \quad \text{where } \psi(y) = \varphi\left(\frac{\lambda_{n+1}}{\lambda}y\right).$$

Note that since  $\frac{1}{2} < \frac{\lambda_{n+1}}{\lambda} \leq 1$ ,

$$\text{supp } \psi \subset B_2(0) \subset [-\pi, \pi]^d \quad \text{and} \quad \|\psi\|_\infty = \|\varphi\|_\infty \leq 1.$$

Furthermore,  $\psi$  annihilates monomials of order up to  $[\alpha]$ , as  $\varphi$  does. The evaluation of the first term in (C.1) (for  $x = 0$ ) is straightforward. Thus, we obtained (C.4) for  $\alpha \geq 0$ .  $\square$

The following corollary follows directly from the proof of Lemma C.7.

**Corollary C.8.** *It holds that for every  $n \in \mathbb{N}$*

$$\mathcal{Z}^n(\mathbb{T}^d) \subset \tilde{\mathcal{Z}}^n(\mathbb{T}^d) \quad \text{and} \quad |\cdot|_{\tilde{\mathcal{Z}}^n} \lesssim |\cdot|_{\mathcal{Z}^n}.$$

**Lemma C.9.** *Suppose  $\eta \in \mathcal{Z}^{-\kappa}$ ,  $\kappa > 0$ . Then,*

$$|\eta^\varepsilon|_{\mathcal{Z}^{-\kappa-1}} \leq |\eta|_{\mathcal{Z}^{-\kappa}}, \quad \text{where} \quad \eta^\varepsilon := \rho_\varepsilon * \eta.$$

*Proof.* For the function  $(\eta^\varepsilon)^\varepsilon := \rho_\varepsilon * \eta^\varepsilon$ , we have

$$\begin{aligned} (\eta^\varepsilon)^\varepsilon(\varphi_x^\lambda) &= \int \eta^\varepsilon(z-y) \rho_\varepsilon(y) \varphi_x^\lambda(z) dy dz = \int \eta^\varepsilon(z) \rho_\varepsilon(y) \varphi_x^\lambda(z+y) dy dz \\ &= \int \eta^\varepsilon(\varphi_{x-y}^\lambda) \rho_\varepsilon(y) dy. \end{aligned}$$

Passing to the limit as  $\varepsilon \rightarrow 0$ , we obtain

$$\eta^\varepsilon(\varphi_x^\lambda) = \int \eta(\varphi_{x-y}^\lambda) \rho_\varepsilon(y) dy.$$

We have

$$\sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \mathcal{B}^{[\kappa+1]}}} |\eta^\varepsilon(\varphi_x^\lambda)| \leq \int \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \mathcal{B}^{[\kappa+1]}}} |\eta(\varphi_{x-y}^\lambda)| \rho_\varepsilon(y) dy \leq \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \mathcal{B}^{[\kappa+1]}}} |\eta(\varphi_x^\lambda)|,$$

where the latter inequality is true by the periodicity of  $\varphi_x^\lambda$ .  $\square$

**Lemma C.10.** *Let  $\kappa > 0$  and  $\eta \in \mathcal{Z}^{-\kappa}$ . Then,*

$$|\partial_i \eta|_{\mathcal{Z}^{-\kappa-1}} \leq |\eta|_{\mathcal{Z}^{-\kappa}}.$$

*Proof.* We have

$$\begin{aligned} |\partial_i \eta|_{\mathcal{Z}^{-\kappa-1}} &= \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \mathcal{B}^r, r=[2+\kappa]}} \frac{|\partial_i \eta(\varphi_x^\lambda)|}{\lambda^{-1-\kappa}} = \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \mathcal{B}^r, r=[2+\kappa]}} \frac{|\eta(\partial_i \varphi_x^\lambda)|}{\lambda^{-\kappa}} \\ &\leq \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \psi \in \mathcal{B}^r, r=[1+\kappa]}} \frac{|\eta(\psi_x^\lambda)|}{\lambda^{-\kappa}} = |\eta|_{\mathcal{Z}^{-\kappa}}. \end{aligned}$$

$\square$

**Proposition C.11.** *Let  $\alpha < 0$ . The Hölder-Zygmund spaces  $\mathcal{Z}^\alpha(\mathbb{T}^d)$  can be equivalently defined with the use of any integer  $r > [-\alpha + 1]$ , that is, as the space of distributions  $f \in \mathcal{D}'(\mathbb{T}^d)$  with the finite norm*

$$(C.5) \quad |f|_{\mathcal{Z}^\alpha}^{(r)} := \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \mathcal{B}^r}} \frac{|f(\varphi_x^\lambda)|}{\lambda^\alpha}, \quad \text{where } r > [-\alpha + 1], r \in \mathbb{N}.$$

Furthermore, the norms  $|\cdot|_{\mathcal{Z}^\alpha}^{(r)}$  and  $|\cdot|_{\mathcal{Z}^\alpha}$  are equivalent; in particular, the norms  $|\cdot|_\alpha$  and  $|\cdot|_\alpha^{(r)}$  are equivalent, where the latter is the restriction of  $|\cdot|_{\mathcal{Z}^\alpha}^{(r)}$  to  $\mathcal{Z}^\alpha(\mathbb{T}^d)$ .

*Proof.* Let  $r > [-\alpha + 1]$ ,  $r \in \mathbb{N}$ . Define

$$(C.6) \quad |f|_{\tilde{\mathcal{Z}}^\alpha}^{(r)} := \sup_{\substack{x \in \mathbb{T}^d, \lambda \in (0,1], \\ \varphi \in \mathcal{B}^r}} \frac{|f(\varphi_x^\lambda)|}{\lambda^\alpha}.$$

The equivalence of the norms (C.5) and (C.6) follows by exactly the same argument as in the proof of Lemma C.7. Let us prove the equivalence of  $|\cdot|_{\mathcal{Z}^\alpha}^{(r)}$  and  $|\cdot|_{\tilde{\mathcal{Z}}^\alpha}$ .

Since  $\tilde{\mathcal{Z}}^\alpha(\mathbb{T}^d) = \mathcal{Z}^\alpha(\mathbb{R}^d) \cap \mathcal{D}'(\mathbb{T}^d)$ , see Remark C.6, by [14, Lemma 14.13],  $|\cdot|_{\tilde{\mathcal{Z}}^\alpha}^{(r)}$  and  $|\cdot|_{\tilde{\mathcal{Z}}^\alpha}$  define the same Banach space. Note that

$$|f|_{\tilde{\mathcal{Z}}^\alpha}^{(r)} \leq |f|_{\tilde{\mathcal{Z}}^\alpha}, \quad f \in \tilde{\mathcal{Z}}^\alpha,$$

since the supremum in the definition of  $|f|_{\tilde{\mathcal{Z}}^\alpha}^{(r)}$  is over a larger set. Since the norms define the same Banach space, their equivalence is a consequence of the open mapping theorem applied to the bounded linear operator

$$J: (\tilde{\mathcal{Z}}^\alpha, |\cdot|_{\tilde{\mathcal{Z}}^\alpha}) \rightarrow (\mathcal{Z}^\alpha, |\cdot|_{\mathcal{Z}^\alpha}^{(r)}), \quad f \mapsto f.$$

Since, by Lemma C.7, the norms  $|\cdot|_{\tilde{\mathcal{Z}}^\alpha}$  and  $|\cdot|_{\mathcal{Z}^\alpha}$  are equivalent, we conclude the equivalence of the norms  $|\cdot|_{\mathcal{Z}^\alpha}$  and  $|\cdot|_{\mathcal{Z}^\alpha}^{(r)}$ .  $\square$

**Lemma C.12.** *For  $\alpha \notin \mathbb{N}$ ,  $\mathcal{Z}^\alpha(\mathbb{T}^d) = C^\alpha(\mathbb{T}^d)$  and the norms in  $\mathcal{Z}^\alpha(\mathbb{T}^d)$  and  $C^\alpha(\mathbb{T}^d)$  are equivalent. Moreover, if  $n = [\alpha]$ , then*

$$C^{n+1} \subset \mathcal{Z}^\alpha \subset C^n.$$

*Proof.* By Lemma C.7,  $\mathcal{Z}^\alpha(\mathbb{T}^d) = \tilde{\mathcal{Z}}^\alpha(\mathbb{T}^d)$  and the norms  $|\cdot|_{\mathcal{Z}^\alpha}$  and  $|\cdot|_{\tilde{\mathcal{Z}}^\alpha}$  are equivalent. By Remark C.6,  $\tilde{\mathcal{Z}}^\alpha(\mathbb{T}^d) = \mathcal{Z}^\alpha(\mathbb{R}^d) \cap \mathcal{D}'(\mathbb{T}^d)$ , where  $\mathcal{Z}^\alpha(\mathbb{R}^d)$  is the classical Hölder-Zygmund space, see [2, 15]. Furthermore, by [14, Proposition 14.15],  $\mathcal{Z}^\alpha(\mathbb{R}^d) = C^\alpha(\mathbb{R}^d)$ . Therefore,  $\tilde{\mathcal{Z}}^\alpha(\mathbb{T}^d) = C^\alpha(\mathbb{T}^d)$ .

Let us show the equivalence of the norms in  $\tilde{\mathcal{Z}}^\alpha(\mathbb{T}^d)$  and  $C^\alpha(\mathbb{T}^d)$ . If  $\alpha < 0$ , then we have nothing to prove since  $C^\alpha(\mathbb{T}^d)$  is  $\tilde{\mathcal{Z}}^\alpha(\mathbb{T}^d)$  by definition, see [15, Definition 3.7, Section 3]. Let  $\alpha \geq 0$ ,  $\alpha \notin \mathbb{N}$ . By the same argument as in (C.3), we obtain

$$\sup_{\substack{x \in \mathbb{T}^d, \lambda \in (0,1], \\ \varphi \in \mathcal{B}_\alpha^0}} \frac{|f(\varphi_x^\lambda)|}{\lambda^\alpha} \lesssim \sum_{|\beta|=n} [D^\beta f]_\alpha,$$

where  $\beta$  is a multiindex and  $D^\beta$  is a partial derivative of order  $|\beta|$ . Hence,

$$|f|_{\tilde{\mathcal{Z}}^\alpha} \lesssim \|f\|_{C^\alpha}.$$



Finally, by the same argument as in the proof of Proposition C.11, involving the open mapping theorem, the norms  $|\cdot|_{\mathcal{Z}^\alpha}$  and  $|\cdot|_{C^\alpha}$  are equivalent. This implies the first statement of the lemma. The last statement of the lemma is an immediate consequence of the first statement.  $\square$

**Lemma C.13.** *Assume  $\kappa, \nu > 0$ ,  $\nu \notin \mathbb{N}$ . Then,  $\mathcal{Z}^\nu \subset \mathcal{Z}^{-\kappa}$  and*

$$(C.7) \quad |f|_{\mathcal{Z}^{-\kappa}} \lesssim |f|_{\mathcal{Z}^\nu}.$$

*Proof.* Take  $f \in \mathcal{Z}^\nu$  and let  $\ell = [\kappa + 1]$ . By Lemma C.12,

$$\begin{aligned} |f|_{\mathcal{Z}^{-\kappa}} &= \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \mathcal{B}^\ell}} \lambda^\kappa |f(\varphi_x^\lambda)| = \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \mathcal{B}^\ell}} \lambda^\kappa |f(x + \lambda \cdot)(\varphi)| \\ &\leq \sup_{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}} \lambda^\kappa |f(x)| \sup_{\varphi \in \mathcal{B}^\ell} \left| \int_{\mathbb{T}^d} \varphi(y) dy \right| \\ &\quad + \sup_{\substack{x \in \mathbb{T}^d, \lambda \in [0,1]_{\mathbb{D}}, \\ \varphi \in \mathcal{B}^\ell}} \lambda^{\kappa+\nu} \left| \int_{\mathbb{T}^d} |y|^\nu \frac{|f(x + \lambda y) - f(x)|}{|\lambda y|^\nu} \varphi(y) dy \right| \\ &\lesssim [\|f\|_\infty + [f]_\nu] \sup_{\varphi \in \mathcal{B}^\ell} \left| \int_{\mathbb{T}^d} \varphi(y) dy \right| \lesssim \|f\|_{C^\nu} \lesssim |f|_{\mathcal{Z}^\nu}. \end{aligned}$$

$\square$

The following corollary is an immediate consequence of Lemma C.13.

**Corollary C.14.** *Assume  $\kappa, \nu > 0$ ,  $\nu \notin \mathbb{N}$ . Then,  $\mathcal{Z}^\nu \subset \mathcal{Z}^{-\kappa}$  and*

$$(C.8) \quad |f|_{-\kappa} \lesssim |f|_\nu.$$

*Proof.* Inequality (C.7) immediately implies that  $\mathcal{Z}^\nu \subset \mathcal{Z}^{-\kappa}$  and (C.8) holds.  $\square$

**Lemma C.15.** *The following equality is true:*

$$C^\infty = \bigcap_{\alpha \in \mathbb{R}} \mathcal{Z}^\alpha = \bigcap_{\alpha \in \mathbb{R}} \mathcal{Z}^\alpha.$$

*Proof.* By Lemma C.12 and since  $C^\infty \subset \mathcal{Z}^n$ ,  $n \in \mathbb{N}$ , it follows that  $C^\infty \subset \bigcap_{\alpha \in \mathbb{R}} \mathcal{Z}^\alpha$ . On the other hand, by Lemmata C.13 and C.12,  $\bigcap_{\alpha \in \mathbb{R}} \mathcal{Z}^\alpha \subset \bigcap_{\alpha \notin \mathbb{N}, \alpha > 0} C^\alpha \subset \bigcap_{\alpha \notin \mathbb{N}, \alpha > 0} C^{[\alpha]} = C^\infty$ . It remains to notice that  $\bigcap_{\alpha \in \mathbb{R}} \mathcal{Z}^\alpha \subset \bigcap_{\alpha \in \mathbb{R}} \mathcal{Z}^\alpha$  and that  $C^\infty \subset \mathcal{Z}^\alpha$  for all  $\alpha \in \mathbb{R}$ .  $\square$

**Lemma C.16.** *Let  $0 < \varkappa < 1$ . Then, for every  $u \in \mathcal{Z}^\varkappa(\mathbb{T}^d)$  and every  $\nu \in (0, \varkappa)$ ,*

$$(C.9) \quad \lim_{\varepsilon \rightarrow 0} |u - u^\varepsilon|_\nu = 0,$$

where  $u^\varepsilon = u * \rho_\varepsilon$ . In particular,

$$\mathcal{Z}^\varkappa(\mathbb{T}^d) \subset \mathcal{Z}^\nu(\mathbb{T}^d).$$

*Proof.* The second part of the Lemma follows trivially from the first so it is sufficient to prove the first part.

For this purpose, let us choose and fix  $u \in \mathcal{Z}^\varkappa(\mathbb{T}^d)$ . By Lemma C.12,  $u \in C^\varkappa(\mathbb{T}^d)$  and it suffices to prove that

$$\lim_{\varepsilon \rightarrow 0} [u - u^\varepsilon]_\nu = 0,$$

where  $[u]_\nu := \sup_{x, y \in \mathbb{T}^d, x \neq y} \frac{|u(x) - u(y)|}{|x - y|^\nu}$ . Define the function  $F : \mathbb{T}^d \times \mathbb{T}^d \rightarrow \mathbb{R}$ ,

$$F(x, y) := \begin{cases} \frac{u(x) - u(y)}{|x - y|^\nu}, & \text{if } x \neq y, \\ 0, & \text{if } x = y. \end{cases}$$

Let  $\sigma > 0$  be fixed arbitrary. Since  $u \in C^\varkappa(\mathbb{T}^d)$  with  $\varkappa > \nu$ ,  $F$  is continuous. Hence, since  $\mathbb{T}^d \times \mathbb{T}^d$  is compact,  $F$  is uniformly continuous and therefore one can find  $\varepsilon_0 > 0$  such that whenever  $|z| < \varepsilon_0$ ,

$$|F(x - z, y - z) - F(x, y)| < \sigma \quad \text{for all } x, y \in \mathbb{T}^d.$$

Therefore, if  $0 < \varepsilon < \varepsilon_0$ , then

$$\begin{aligned} & \frac{|(u^\varepsilon - u)(x) - (u^\varepsilon - u)(y)|}{|x - y|^\nu} \\ &= \left| \int \rho_\varepsilon(z) \frac{u(x - z) - u(y - z)}{|x - y|^\nu} dz - \int \rho_\varepsilon(z) \frac{u(x) - u(y)}{|x - y|^\nu} dz \right| \\ &= \left| \int \rho_\varepsilon(z) (F(x - z, y - z) - F(x, y)) dz \right| < \sigma. \end{aligned}$$

Taking the supremum over  $x, y \in \mathbb{T}^d$  of the left-hand side, we infer that

$$[u - u^\varepsilon]_\nu \leq \sigma.$$

By Lemma C.12,  $u \in \mathcal{X}^\nu(\mathbb{T}^d)$  and (C.9) holds.  $\square$

#### APPENDIX D. SINGULAR KERNELS

**Lemma D.1.** *Let  $\beta > 0$ . For a  $\beta$ -regularizing kernel  $\mathcal{K}$  (see [2, Definition 2.3] or [15, Assumption 5.1]), the map  $f \mapsto \mathcal{K} * f$  is continuous from  $\mathcal{L}^\alpha(\mathbb{R}^d)$  to  $\mathcal{L}^{\alpha+\beta}(\mathbb{R}^d)$  for every  $\alpha \in \mathbb{R}$ . Moreover,*

$$(D.1) \quad |\mathcal{K} * f|_{\beta+\alpha, \mathcal{K}} \lesssim |f|_{\alpha, \tilde{\mathcal{K}}},$$

where  $\mathcal{K} \subset \mathbb{R}^d$  is a compact and  $\tilde{\mathcal{K}} \supset \mathcal{K}$  is another compact strictly containing  $\mathcal{K}$ .

*Proof.* For the proof see, e.g. [14, Theorem 14.17]. The bound (D.1) is obtained in the proof of Theorem 2.13 in [2].  $\square$

In the periodic case, the above statement implies the following result.

**Lemma D.2.** *Let  $\beta > 0$  and let  $\mathcal{K}$  be a  $\beta$ -regularizing kernel on a  $d$ -dimensional torus  $\mathbb{T}^d$ . Then, for every  $\alpha \in \mathbb{R}$ , the map*

$$(D.2) \quad \mathcal{L}^\alpha(\mathbb{T}^d) \rightarrow \mathcal{L}^{\alpha+\beta}(\mathbb{T}^d), \quad f \mapsto \mathcal{K} * f$$

*is linear and continuous. Moreover, for every  $\alpha \in \mathbb{R}$ , the map (D.2) maps  $\mathcal{L}^\alpha(\mathbb{T}^d)$  to  $\mathcal{L}^{\alpha+\beta}(\mathbb{T}^d)$ . In particular,*

$$|\mathcal{K} * f|_{\mathcal{L}^{\beta+\alpha}} \lesssim |f|_{\mathcal{L}^\alpha} \quad \text{and} \quad |\mathcal{K} * f|_{\beta+\alpha} \lesssim |f|_\alpha.$$

*Proof.* By Lemma C.7, the norms  $|\cdot|_{\mathcal{Z}^\alpha}$  (in the sense of Definition 2.2) and  $|\cdot|_{\alpha, \mathbb{T}^2}$  (in the sense of [2, Definition 2.1]) are equivalent. We remark that  $|\cdot|_{\alpha, \mathbb{T}^d}$  and  $|\cdot|_{\mathcal{Z}^\alpha}$  is the same norm, see Definition C.3 and Remark C.4. On the other hand, in Definition 2.2,  $\varphi_x^\lambda$  is a periodic function on  $\mathbb{T}^d$ . Therefore, the supremum in  $x$  running over an enlargement of  $\mathbb{T}^d$  equals to the supremum over  $\mathbb{T}^d$ .

Finally, since  $\mathcal{H} * f \in C^\infty$  for every  $f \in C^\infty$ , we infer that (D.2) maps  $\mathcal{Z}^\alpha(\mathbb{T}^d)$  to  $\mathcal{Z}^{\alpha+\beta}(\mathbb{T}^d)$ .  $\square$

**Lemma D.3.** *The Green function  $G$  for the operator  $-2\partial_{\bar{z}}$  is a 1-regularizing kernel on  $\mathbb{T}^2$ . Moreover, for every  $\alpha \in \mathbb{R}$ , the map*

$$(D.3) \quad \mathcal{Z}^\alpha(\mathbb{T}^2) \ni f \mapsto G * f \in \mathcal{Z}^{\alpha+1}(\mathbb{T}^2),$$

*is linear and continuous, and maps  $\mathcal{Z}^\alpha(\mathbb{T}^2)$  to  $\mathcal{Z}^{\alpha+1}(\mathbb{T}^2)$ . In particular,*

$$(D.4) \quad |G * f|_{\mathcal{Z}^{\alpha+1}} \lesssim |f|_{\mathcal{Z}^\alpha} \quad \text{and} \quad |G * f|_{\alpha+1} \lesssim |f|_\alpha.$$

For the proof, we need the following result.

**Lemma D.4.** *The Green function  $K$  for the Laplacian  $-\Delta$  on  $\mathbb{T}^2$  is in  $C^\infty(\mathbb{T}^2 \setminus \{0\})$ . For the derivatives of  $D^\beta K$ , where  $\beta$  is a multiindex, we have the bounds*

$$|D^\beta K(x)| \leq C_n |x|^{-n}, \quad |\beta| = n, \quad n = 1, 2, \dots,$$

where  $C_n > 0$  is a constant depending only on  $n$ .

*Proof.* For the proof see [4, Proposition B.1].  $\square$

*Proof of Lemma D.3.* Since  $G = 2\partial_{\bar{z}}K$ , by Lemma D.4,

$$|D^\beta G(x)| \leq C_{n+1} |x|^{-(n+1)}, \quad |\beta| = n, \quad n = 0, 1, 2, \dots$$

In particular,

$$|G(x)| \leq \frac{C_1}{|x|}.$$

By [2, Lemma 2.9],  $G(x - y)$  is a 1-regularizing kernel on  $\mathbb{T}^2$ . Therefore, the statement of the lemma follows from Lemma D.2.  $\square$

**Lemma D.5.** *Let  $K$  be the Green function of the Laplacian  $-\Delta$  on  $\mathbb{T}^2$ . Then, for every  $\alpha \in \mathbb{R}$ , the map*

$$(D.5) \quad \mathcal{Z}^\alpha(\mathbb{T}^2) \rightarrow \mathcal{Z}^{\alpha+2}(\mathbb{T}^2), \quad f \mapsto K * f$$

*is linear and continuous. Moreover, for every  $\alpha \in \mathbb{R}$ , the map (D.5) maps  $\mathcal{Z}^\alpha(\mathbb{T}^2)$  to  $\mathcal{Z}^{\alpha+2}(\mathbb{T}^2)$ . In particular,*

$$|K * f|_{\mathcal{Z}^{\alpha+2}} \lesssim |f|_{\mathcal{Z}^\alpha} \quad \text{and} \quad |K * f|_{\alpha+2} \lesssim |f|_\alpha.$$

*Proof.* Since  $G = 2\partial_{\bar{z}}K$ , we have

$$G * f = 2\partial_{\bar{z}}(K * f).$$

Since  $\int_{\mathbb{T}^2} G * f = 0$ , we have

$$K * f = -\bar{G} * (G * f).$$

By Lemma D.3,

$$|K * f|_{\mathcal{Z}^{\alpha+2}} = |\overline{G} * (G * f)|_{\mathcal{Z}^{\alpha+2}} \lesssim |(G * f)|_{\mathcal{Z}^{\alpha+1}} \lesssim |f|_{\mathcal{Z}^{\alpha}}.$$

Finally, we notice that the map (D.5) maps  $\mathcal{Z}^{\alpha}(\mathbb{T}^2)$  to  $\mathcal{Z}^{\alpha+2}(\mathbb{T}^2)$  by the same argument as in Lemma D.2.  $\square$

## APPENDIX E. YOUNG PRODUCTS IN LITTLE HÖLDER-ZYGMUND SPACES

**Lemma E.1.** *Fix  $\alpha > 0$  and  $\beta \leq 0$  such that  $\alpha + \beta > 0$ . Then, there exists a unique bilinear continuous map*

$$(E.1) \quad \mathcal{Z}^{\alpha}(\mathbb{T}^d) \times \mathcal{Z}^{\beta}(\mathbb{T}^d) \ni (f, g) \mapsto f \cdot g \in \mathcal{Z}^{\beta}(\mathbb{T}^d)$$

which extends the usual product

$$(E.2) \quad C^{\infty}(\mathbb{T}^d) \times C^{\infty}(\mathbb{T}^d) \ni (f, g) \mapsto f \cdot g \in C^{\infty}(\mathbb{T}^d).$$

Furthermore, for all  $f \in \mathcal{Z}^{\alpha}(\mathbb{T}^d)$  and  $g \in \mathcal{Z}^{\beta}(\mathbb{T}^d)$ ,

$$(E.3) \quad |f \cdot g|_{\beta} \lesssim |f|_{\alpha} |g|_{\beta}.$$

*Proof.* By the definition of the spaces  $\mathcal{Z}^{\alpha}(\mathbb{T}^d)$  and  $\mathcal{Z}^{\beta}(\mathbb{T}^d)$  it is sufficient to prove that (E.3) is fulfilled for all  $f, g \in C^{\infty}(\mathbb{T}^d)$ . Then, the product (E.2) will be automatically extended to the map (E.1), and moreover, (E.3) will be fulfilled.

In [6, Theorem 14.1], it was shown that the inequality

$$(E.4) \quad |f \cdot g|_{K, \beta} \lesssim |f|_{K_4, \alpha} |g|_{K_4, \beta},$$

is valid, in particular, for  $f, g \in C^{\infty}(\mathbb{R}^d)$ , where  $K \subset \mathbb{R}^d$  is a compact and  $K_4 := K + \overline{B_4(0)}$  is its enlargement with  $B_4(0)$  being a ball of radius 4 centered at the origin, see [6, Formula (14.5)]. In the periodic case, we take  $K = \mathbb{T}^d$  and note that the norms  $|\cdot|_{\mathcal{Z}^{\beta}}$  and  $|\cdot|_{\mathbb{T}^d, \beta}$ , and also  $|\cdot|_{\mathcal{Z}^{\alpha}}$  and  $|\cdot|_{\mathbb{T}^d, \alpha}$ ,  $\alpha \notin \mathbb{N}$ , are equivalent by Lemma C.7 and Remark C.4. Then, (E.4) implies (E.3) for  $C^{\infty}(\mathbb{T}^d)$ -functions by the same argument as in the proof of Lemma D.2 for  $\alpha \notin \mathbb{N}$ . If  $\alpha \in \mathbb{N}$ , we obtain (E.3) by Corollary C.8.  $\square$

**Remark E.2.** In [6, Theorem 14.1], has was proved that the Young product

$$\mathcal{Z}^{\alpha}(\mathbb{T}^d) \times \mathcal{Z}^{\beta}(\mathbb{T}^d) \rightarrow \mathcal{Z}^{\beta}(\mathbb{T}^d), \quad (f, g) \mapsto f \cdot g$$

exists and extends the product (E.2), but it is not unique, see [6, Remark 14.2]. By Lemma E.1, the extension of the product (E.2) to  $\mathcal{Z}^{\alpha}(\mathbb{T}^d) \times \mathcal{Z}^{\beta}(\mathbb{T}^d)$  is unique and  $\mathcal{Z}^{\beta}(\mathbb{T}^d)$ -valued.

**Acknowledgements.** We express our gratitude for the support received during the preparation of this work. Z.B. acknowledges partial support from CNPq-Brazil (grant 409764/2023-0). E.S. acknowledges partial support from Universidade Federal da Paraíba (PROPESQ/PRPG/UFPB) through grant PIA13631-2020 (public calls nos. 03/2020 and 06/2021), as well as from CNPq-Brazil (grants 409764/2023-0 and 443594/2023-6) and CAPES MATH AMSUD (grant 88887.878894/2023-00). M.N., who is a member of the CNPq-Brazil (grant 443594/2023-6) and CAPES MATH AMSUD grant teams, acknowledges the collaborative support of these projects. He also acknowledges partial support from Universidade Federal do Amazonas and

Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). Finally, E.S. extends thanks to the Mathematics Department of the University of York (UK) for their hospitality, as this paper was partially written during her visit to the department.

## REFERENCES

- [1] N. Berglund and C. Kuehn. Model spaces of regularity structures for space-fractional SPDEs. *J Stat Phys* 168, 331–368, 2017.
- [2] L. Broux, F. Caravenna and L. Zambotti, Hairer’s multilevel Schauder estimates without regularity structures, *Transactions of the American Mathematical Society* vol 377, N 10, 6981–7035, 2024.
- [3] L. Broux, F. Caravenna and L. Zambotti, An example of singular elliptic stochastic PDE, *Matemática Contemporânea*, vol. 58, pp 3–67, 2023
- [4] Z. Brzeźniak, F. Flandoli and M. Maurelli. Existence and uniqueness for stochastic 2D Euler flows with bounded vorticity. *Arch. Rational Mech. Anal.* 221, 107–142, 2016.
- [5] J. Behrndt, M. Holzmann and M. Tusek. Two-dimensional Dirac operators with general  $\delta$ -shell interactions supported on a straight line. *J. Phys. A: Math. Theor.* 56 045201, 2023.
- [6] F. Caravenna and L. Zambotti, Hairer’s reconstruction theorem without regularity structures, *EMS Surv. Math. Sci.* 7, pp 207–251, 2020
- [7] L. Chiarini and C. Landim. Schauder estimates for fractional laplacians and nonlocal, one-dimensional singular SPDEs. *SIAM J. Math. Anal.*, vol 54, issue 2, 2163–2215, 2022.
- [8] N. L. Chuprikov. The stationary Dirac equation as a generalized Pauli equation for two quasiparticles. *Found Phys* 45, 644–656, 2015.
- [9] G. da Prato and A. Debussche. Two-dimensional Navier–Stokes equations driven by a space–time White Noise. *J. Funct. Anal.* 196, 180–210, 2002.
- [10] P. Duch. Renormalization of singular elliptic stochastic PDEs using flow equation. *Probability and Mathematical Physics*, vol. 6, No. 1, 111–138, 2025.
- [11] P. Duch, M. Gubinelli and P. Rinaldi, Parabolic stochastic quantisation of the fractional  $\Phi_3^4$  model in the full subcritical regime, arXiv:2303.18112 [math.PR], 2023.
- [12] M. J. Esteban and E. Séré. Stationary states of the nonlinear Dirac equation: A variational approach. *Commun. Math. Phys.* 171, 323–350, 1995.
- [13] J. Gilbert and M. Murray, *CLIFFORD ALGEBRAS AND DIRAC OPERATORS IN HARMONIC ANALYSIS*, Cambridge University Press, Cambridge, 1991.
- [14] P. K. Friz and M. Hairer, *A COURSE ON ROUGH PATHS. WITH AN INTRODUCTION TO REGULARITY STRUCTURES*. Springer, 2020, pp. xvi+346.
- [15] M. Hairer. A theory of regularity structures. *Inventiones Mathematicae.* 198 (2): 269–504, 2014.
- [16] F. Helein, *HARMONIC MAPS, CONSERVATION LAWS AND MOVING FRAMES*. Cambridge Tracts in Mathematics, 150. Cambridge University Press, 2002.
- [17] T. Runst and W. Sickel, *SOBOLEV SPACES OF FRACTIONAL ORDER, NEMYTSKIJ OPERATORS, AND NONLINEAR PARTIAL DIFFERENTIAL EQUATIONS*. De Gruyter Series in Nonlinear Analysis and Applications, 3. Walter de Gruyter & Co., Berlin, 1996.
- [18] S. Selberg. Global existence in the critical space for the Thirring and Gross-Neveu models coupled with the electromagnetic field. *Discrete Continuous Dyn. Syst. Ser.*, vol 38, issue 5, 2555–2569, 2018.
- [19] B. Thaller. *THE DIRAC EQUATION*, Springer-Verlag, Berlin, 1992.
- [20] M. C. Veraar, Regularity of Gaussian white noise on the  $d$ -dimensional torus. *Banach center publications*, vol 95, 385–398, 2011.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, YORK, YO10 5DD, UK

*Email address:* `zdzislaw.brzezniak@york.ac.uk`

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO AMAZONAS, MANAUS, BRAZIL

*Email address:* `misha.neklyudov@gmail.com`

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DA PARAÍBA, JOÃO PESSOA,  
BRAZIL

*Email address:* `evelina.shamarova@academico.ufpb.br`