

THE SCHRÖDER-BERNSTEIN PROPERTY FOR OPERATORS ON HILBERT SPACES

NICOLÁS CUERVO OVALLE, ISAAC GOLDBRING, AND NETANEL LEVI

ABSTRACT. We establish that the complete theory of a Hilbert space equipped with a normal operator has the Schröder-Bernstein property for separable models. This is a partial answer to a question of Argoty, Berenstein, and the first-named author. We also prove an analogous statement for unbounded self-adjoint operators.

1. INTRODUCTION

The classical Schröder-Bernstein theorem in set theory states that when X and Y are sets for which there are injections $X \hookrightarrow Y$ and $Y \hookrightarrow X$, then in fact there is a bijection between X and Y . It is natural to ask if this property holds when X and Y are not mere sets but have additional structure on them. One appropriate setting for such a generalization is to consider structures (in the model-theoretic sense) \mathcal{M} and \mathcal{N} in the same language for which there exist *elementary* embeddings $\mathcal{M} \hookrightarrow \mathcal{N}$ and $\mathcal{N} \hookrightarrow \mathcal{M}$ and to ask if in this case \mathcal{M} and \mathcal{N} must be isomorphic. When this phenomena holds for all models of a first-order theory, we say that theory has the **Schröder-Bernstein property** or **SB-property** for short. This property has been considered in classical model theory (see, for example, [5] and [6]) and in the setting of continuous logic in [1].

In the latter article, the question was posed as to whether or not the complete theory of a structure of the form (\mathcal{H}, T) has the SB-property, where \mathcal{H} is a Hilbert space and T is a bounded, self-adjoint operator on \mathcal{H} ([1, Question 2.29]). Regarding this question, two partial results were proven in [1]: first, the complete

The first named author was partially supported by NSF grant DMS-2054477. He would also like to thank the UC Irvine Department of Mathematics for their hospitality.

The second-named author was partially supported by NSF grant DMS-2054477.

The third named author was supported by NSF DMS-2052899, DMS-2155211, and Simons 896624.

theory of any such pair (\mathcal{H}, T) has the SB-property **up to perturbations** ([1, Theorem 2.28]) and has the actual SB-property when T has countable spectrum ([1, Proposition 2.30]).

In this paper, we give another partial answer to the above question by establishing the SB-property for separable models; in fact, one does not even need to assume that the embeddings between these structures are elementary. In addition, one does not even need to require the operator to be self-adjoint, but instead can assume that the operator is merely normal. We rephrase this positive resolution of the question in more functional analytic terms:

Theorem. *Let $\mathcal{H}_1, \mathcal{H}_2$ be separable Hilbert spaces and let $T_i : \mathcal{H}_i \rightarrow \mathcal{H}_i$ be normal operators for $i = 1, 2$. Suppose that there exist isometries $U_1 : \mathcal{H}_1 \rightarrow \mathcal{H}_2$ and $U_2 : \mathcal{H}_2 \rightarrow \mathcal{H}_1$ such that $T_1 U_2 = U_2 T_2$ and $T_2 U_1 = U_1 T_1$. Then T_1 and T_2 are unitarily equivalent, that is, there is a unitary transformation $U : \mathcal{H}_1 \rightarrow \mathcal{H}_2$ such that $U T_1 = T_2 U$.*

By applying the Cayley transform, one can use the previous theorem to prove a version for unbounded self-adjoint operators; we do this in the last section.

We would like to thank Alex Berenstein for discovering an error in the first version of this paper.

2. THE MAIN THEOREM

Throughout this paper, given two Borel measures μ_1, μ_2 on \mathbb{R} , we say μ_1 and μ_2 are mutually absolutely continuous, denoted $\mu_1 \sim \mu_2$, if $\mu_1 \ll \mu_2$ and $\mu_2 \ll \mu_1$. In addition, given two bounded operators T_1, T_2 acting on Hilbert spaces $\mathcal{H}_1, \mathcal{H}_2$, we will write $(\mathcal{H}_1, T_1) \simeq (\mathcal{H}_2, T_2)$ if T_1 and T_2 are unitarily equivalent, that is, if there exists a unitary transformation $U : \mathcal{H}_1 \rightarrow \mathcal{H}_2$ such that $T_1 = U^{-1} T_2 U$.

We remind the reader of the direct integral version of the spectral theorem for normal operators [3]:

Fact 2.1. *Suppose that T is a normal operator on a separable Hilbert space \mathcal{H} . Then there is a Borel probability measure μ on the spectrum $\sigma(T)$ of T and a measurable family $(\mathcal{H}_\lambda)_{\lambda \in \mathbb{R}}$ of Hilbert spaces such that T is unitarily equivalent to the multiplication operator on the direct integral $\int^\oplus \mathcal{H}_\lambda d\mu(\lambda)$. Moreover, this direct integral representation of T is a unitary invariant of T in the following sense: Suppose that for $i = 1, 2$, T_i is a bounded normal operator acting on a Hilbert space \mathcal{H}_i . Suppose, in addition that we have Borel probability measures μ_1, μ_2 , a μ_1 -measurable family of Hilbert spaces $(\mathcal{H}_\lambda^1)_{\lambda \in \mathbb{R}}$ and a μ_2 -measurable family of Hilbert spaces $(\mathcal{H}_\lambda^2)_{\lambda \in \mathbb{R}}$ such that for*

$i = 1, 2$, T_i is unitarily equivalent to the multiplication operator on the direct integral $\int^\oplus \mathcal{H}_\lambda^i d\mu_i(\lambda)$. Then $(\mathcal{H}_1, T_1) \simeq (\mathcal{H}_2, T_2)$ if and only if: $\mu_1 \sim \mu_2$ and for μ_1 -almost every $\lambda \in \mathbb{R}$, $\dim \mathcal{H}_\lambda^1 = \dim \mathcal{H}_\lambda^2$.

Corollary 2.2. Let $T : \mathcal{H} \rightarrow \mathcal{H}$ be a bounded normal operator acting on a separable Hilbert space \mathcal{H} . Suppose that $W \subseteq \mathcal{H}$ is a closed subspace which is invariant under T and under T^* . Consider the direct integral representations of T and of $T|_W$ given by

$$\int^\oplus \mathcal{H}_\lambda d\mu(\lambda),$$

$$\int^\oplus W_\lambda d\rho(\lambda)$$

respectively. Then $\rho \ll \mu$, and for ρ -almost every $\lambda \in \mathbb{R}$, $\dim W_\lambda \leq \dim \mathcal{H}_\lambda$.

Proof. Let us also consider the direct integral representation of $T|_{W^\perp}$:

$$\int^\oplus E_\lambda d\sigma(\lambda)$$

By the invariance of W under both T and T^* , we have that $T = T|_W \oplus T|_{W^\perp}$, which implies that the following is another direct integral representation of T :

$$\int^\oplus (W_\lambda \oplus E_\lambda) d(\rho + \sigma)(\lambda).$$

By Fact 2.1, this implies that $\rho + \sigma$ and μ are mutually absolutely continuous, and that for μ -almost every $\lambda \in \mathbb{R}$, the dimensions of \mathcal{H}_λ and of $W_\lambda \oplus E_\lambda$ are equal. In particular we obtain that $\rho \ll \mu$ and that for ρ -almost every $\lambda \in \mathbb{R}$, $\dim W_\lambda \leq \dim \mathcal{H}_\lambda$, as required. \square

The following proposition is the key to the entire proof:

Proposition 2.3. Let $T : \mathcal{H} \rightarrow \mathcal{H}$ be a bounded normal operator acting on a separable Hilbert space \mathcal{H} . Suppose that $W \subseteq V \subseteq \mathcal{H}$ are closed subspaces which are invariant under both T and under T^* . Suppose in addition that $(\mathcal{H}, T) \simeq (W, T|_W)$. Then $(\mathcal{H}, T) \simeq (V, T|_V)$.

Proof. Let us consider the direct integral representations of T , $T|_V$, and $T|_W$:

$$\int^\oplus \mathcal{H}_\lambda d\mu(\lambda)$$

$$\int^\oplus V_\lambda d\rho(\lambda),$$

$$\int^\oplus W_\lambda d\sigma(\lambda).$$

By Corollary 2.2, we have that $\sigma \ll \rho \ll \mu$, $\dim W_\lambda \leq \dim V_\lambda$ for σ -almost every $\lambda \in \mathbb{R}$, and $\dim V_\lambda \leq \dim \mathcal{H}_\lambda$ for ρ -almost every $\lambda \in \mathbb{R}$. On the other hand, by the unitary equivalence of T and of $T|_W$, we have that μ and σ are mutually absolutely continuous and $\dim \mathcal{H}_\lambda = \dim W_\lambda$ for σ -almost every $\lambda \in \mathbb{R}$. This implies that $\mu \ll \rho$, whence μ and ρ are mutually absolutely continuous, and that for ρ -almost every $\lambda \in \mathbb{R}$, $\dim \mathcal{H}_\lambda = \dim W_\lambda \leq \dim V_\lambda \leq \dim \mathcal{H}_\lambda$, whence $\dim \mathcal{H}_\lambda = \dim V_\lambda$ for ρ -almost every $\lambda \in \mathbb{R}$. By Fact 2.1, this implies that $(T, \mathcal{H}) \simeq (T|_V, V)$, as required. \square

We will also need the Fuglede-Putnam theorem (see, for example, [2, Chapter IX]):

Fact 2.4. *Suppose that $T_i : \mathcal{H}_i \rightarrow \mathcal{H}_i$ are bounded operators for $i = 1, 2$. Further suppose that $S : \mathcal{H}_1 \rightarrow \mathcal{H}_2$ is a bounded operator that intertwines T_1 and T_2 , that is, such that $ST_1 = T_2S$. Then S also intertwines T_1^* and T_2^* , that is, $ST_1^* = T_2^*S$.*

We are now ready to prove our main theorem. We repeat the statement for the convenience of the reader.

Theorem 2.5. *Let $\mathcal{H}_1, \mathcal{H}_2$ be separable Hilbert spaces and let $T_i : \mathcal{H}_i \rightarrow \mathcal{H}_i$ be bounded normal operators. Suppose that there exist isometries $U_1 : \mathcal{H}_1 \rightarrow \mathcal{H}_2$ and $U_2 : \mathcal{H}_2 \rightarrow \mathcal{H}_1$ such that $T_1U_2 = U_2T_2$ and $T_2U_1 = U_1T_1$. Then T_1 and T_2 are unitarily equivalent.*

Proof. Set $V := U_1(\mathcal{H}_1)$ and $W := U_1(U_2(\mathcal{H}_2))$. We claim that V and W are closed subspaces which are invariant under T_2 and under T_2^* . Closedness follows immediately from unitarity of U_1 and U_2 . To see that these spaces are T_2 -invariant, let us consider $\varphi_1 \in V$, $\varphi_2 \in W$ and take $\psi_1 \in \mathcal{H}_1$, $\psi_2 \in \mathcal{H}_2$ such that $\varphi_1 = U_1\psi_1$, $\varphi_2 = U_1U_2\psi_2$. Then we have

$$T_2\varphi_1 = T_2U_1\psi_1 = U_1T_1\psi_1 \in U_1(\mathcal{H}_1) = V,$$

$$T_2\varphi_2 = T_2(U_1U_2\psi_2) = U_1T_1U_2\psi_2 = U_1U_2T_2\psi_2 \in U_1U_2(\mathcal{H}_2) = W,$$

as required. It is left to show that both spaces are T_2^* -invariant. Note that the only thing we used to prove the invariance under T_2 is the intertwining properties of T_1 and T_2 under U_1 and U_2 . By Fact 2.4, this property also holds for T_1^* and T_2^* . Consequently, one can proceed along the same lines and obtain the invariance of W and V under T_2^* .

We now claim that $(T_2, \mathcal{H}_2) \simeq (T_2, T_2|_W)$. This is true since U_1U_2 is a surjective isometry between \mathcal{H}_2 and W , which means it is unitary. In addition, by the above calculation it commutes with T_2 . An identical argument implies that

$(T_1, \mathcal{H}_1) \simeq (T_2, T_2|_V)$. Finally, by Proposition 2.3, we get that $(T_2, \mathcal{H}) \simeq (T_2, V)$. By transitivity of the relation of unitary equivalence, we obtain $(T_1, \mathcal{H}_1) \simeq (T_2, \mathcal{H}_2)$ as required. \square

3. UNBOUNDED SELF-ADJOINT OPERATORS

Theorem 3.1. *Let \mathcal{H}_1 and \mathcal{H}_2 be separable Hilbert spaces and let $T_i : \mathcal{H}_i \rightarrow \mathcal{H}_i$ be unbounded self-adjoint operators. Further suppose that there exist isometries $U_1 : \mathcal{H}_1 \rightarrow \mathcal{H}_2$ and $U_2 : \mathcal{H}_2 \rightarrow \mathcal{H}_1$ such that $U_1 T_1 \subseteq T_2 U_1$ and $U_2 T_2 \subseteq T_1 U_2$. Then $(\mathcal{H}_1, T_1) \simeq (\mathcal{H}_2, T_2)$ in the sense that there is a unitary transformation $U : \mathcal{H}_1 \rightarrow \mathcal{H}_1$ such that $U T_1 U^{-1} = T_2$.*

Proof. For each $i = 1, 2$, let $V_i := (T_i - i)(T_i + i)^{-1}$ be the Cayley transform of T_i , which is a unitary operator on \mathcal{H}_i . Note then that we have

$$\begin{aligned} U_1 V_1 &= U_1 (T_1 - i)(T_1 + i)^{-1} \\ &= (T_2 - i)U_1 (T_1 + i)^{-1} \\ &= (T_2 - i)(T_2 + i)^{-1}U_1 \\ &= V_2 U_1. \end{aligned}$$

In the same way, we have that $U_2 V_2 = V_1 U_2$. Since each V_i is unitary (and hence normal), the proof of Theorem 2.5, shows that U_1 is a surjective isometry. It follows then that $U_1 T_1 U_1^{-1} \subseteq T_2$ and thus $U_1 T_1 U_1^{-1} = T_2$ as self-adjoint operators have no proper symmetric extensions. \square

REFERENCES

- [1] C. Argoty, A. Berenstein, and N. Cuervo Ovalle, *The SB-property on metric structures*, Archive for Mathematical Logic, 2025, p. 1-29.
- [2] J.B. Conway, *A course in functional analysis*, Graduate Texts in Mathematics **96** (1990), Springer-Verlag.
- [3] J. Dixmier, *Les algèbres d'opérateurs dans l'espace Hilbertien*, Chapter II, Gauthier-Villars, Paris (1969).
- [4] J Goodrick, *When are elementarily bi-embeddable models isomorphic?*, Ph.D. thesis, University of California, Berkeley (2007).
- [5] J Goodrick and MC Laskowski, *The Schröder-Bernstein property for α -saturated models*, Proceedings of the American Mathematical Society **142** (2014), 1013-1023.
- [6] T. A. Nurmagambetov, *Characterization of ω -stable theories of bounded dimension*, Algebra and Logic **28** (1989), 388-396.

DEPARTMENT OF MATHEMATICS, LOS ANDES UNIVERSITY, BOGOTÁ, COLOMBIA, CRA. 1 #18A-12

Email address: `n.cuervo10@uniandes.edu.co`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, IRVINE, 340 ROWLAND HALL (BLDG.# 400), IRVINE, CA 92697-3875

Email address: `isaac@math.uci.edu`

URL: `http://www.math.uci.edu/~isaac`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, IRVINE, 340 ROWLAND HALL (BLDG.# 400), IRVINE, CA 92697-3875

Email address: `netanel1@uci.edu`