THE SCHRÖDER-BERNSTEIN PROPERTY FOR OPERATORS ON HILBERT SPACES

NICOLÁS CUERVO OVALLE, ISAAC GOLDBRING, AND NETANEL LEVI

ABSTRACT. We establish that the complete theory of a Hilbert space equipped with a normal operator has the Schröder-Bernstein property. This in particular answers a question of Argoty, Berenstein, and the first-named author. We also prove an analogous statement for unbounded self-adjoint operators.

1. INTRODUCTION

The classical Schröder-Bernstein theorem in set theory states that when X and Y are sets for which there are injections $X \hookrightarrow Y$ and $Y \hookrightarrow X$, then in fact there is a bijection between X and Y. It is natural to ask if this property holds when X and Y are not mere sets but have additional structure on them. One appropriate setting for such a generalization is to consider structures (in the model-theoretic sense) \mathcal{M} and \mathcal{N} in the same language for which there exist *elementary* embeddings $\mathcal{M} \hookrightarrow \mathcal{N}$ and $\mathcal{N} \hookrightarrow \mathcal{M}$ and to ask if in this case \mathcal{M} and \mathcal{N} must be isomorphic. When this phenomena holds for all models of a first-order theory, we say that theory has the **Schröder-Bernstein property** or **SB-property** for short. This property has been considered in classical model theory (see, for example, [5] and [6]) and in the setting of continuous logic in [1].

In the latter article, the question was posed as to whether or not the complete theory of a structure of the form $(\mathcal{H}, \mathsf{T})$ has the SB-property, where \mathcal{H} is a Hilbert space and T is a bounded, self-adjoint operator on \mathcal{H} ([1, Question 2.29]). Regarding this question, two partial results were proven in [1]: first, the complete theory of any such pair $(\mathcal{H}, \mathsf{T})$ has the SB-property **up to perturbations** ([1, Theorem 2.28]) and has the actual SB-property when T has countable spectrum ([1, Proposition 2.30]).

The first named author was partially supported by NSF grant DMS-2054477. He would also like to thank the UC Irvine Department of Mathematics for their hospitality.

The second-named author was partially supported by NSF grant DMS-2054477.

The third named author was supported by NSF DMS-2052899, DMS-2155211, and Simons 896624.

In this paper, we give a positive answer to the above question; in fact, one does not even need to assume that the embeddings between these structures are elementary. In addition, one does not even need to require the operator to be selfadjoint, but instead can assume that the operator is merely normal. We rephrase this positive resolution of the question in more functional analytic terms:

Theorem. Let $\mathfrak{H}_1, \mathfrak{H}_2$ be Hilbert spaces and let $T_i : \mathfrak{H}_i \to \mathfrak{H}_i$ be normal operators for i = 1, 2. Suppose that there exist isometries $U_1 : \mathfrak{H}_1 \to \mathfrak{H}_2$ and $U_2 : \mathfrak{H}_2 \to \mathfrak{H}_1$ such that $T_1 U_2 = U_2 T_2$ and $T_2 U_1 = U_1 T_1$. Then T_1 and T_2 are unitarily equivalent, that is, there is a unitary transformation $U : \mathfrak{H}_1 \to \mathfrak{H}_2$ such that $UT_1 = T_2 U$.

From our proof it will follow that the transformations U_1 and U_2 are unitary and so one can take $U = U_1$ in the above theorem.

By applying the Cayley transform, one can use the previous theorem to prove a version for unbounded self-adjoint operators; we do this in the last section.

2. The main theorem

We remind the reader of the direct integral version of the spectral theorem for normal operators [3]:

Fact 2.1. Suppose that T is a normal operator on a separable Hilbert space \mathcal{H} . Then there is a Borel probability measure μ on the spectrum $\sigma(T)$ and a measurable family $(\mathcal{H}_{\lambda})_{\lambda \in \sigma(T)}$ of Hilbert spaces such that T is unitarily equivalent to the multiplication operator on the direct integral $\int^{\oplus} \mathcal{H}_{\lambda} d\mu(\lambda)$. Moreover, this direct integral representation of T is a unitary invariant of T in the sense that if ν is another Borel probability measure on $\sigma(T)$ and $(\mathcal{H}'_{\lambda})_{\lambda \in \sigma(T)}$ is another measurable family of Hilbert spaces such that T is unitarily conjugate to the multiplication operator on $\int^{\oplus} \mathcal{H}'_{\lambda} d\nu(\lambda)$, then μ and ν are mutually absolutely continuous and dim $(\mathcal{H}_{\lambda}) = \dim(\mathcal{H}'_{\lambda})$ for μ -almost all $\lambda \in \sigma(T)$.

The following proposition is the key to the entire proof:

Proposition 2.2. Let $T : \mathcal{H} \to \mathcal{H}$ be a bounded normal operator. Suppose that $V \subseteq \mathcal{H}$ is a closed subspace invariant under both T and T^{*}. Suppose in addition that $(\mathcal{H}, T) \simeq (V, T|_V)$. Then $V = \mathcal{H}$.

Proof. We first suppose that \mathcal{H} is separable. Set $W := V^{\perp}$. By assumption, W is T-invariant. Consider the direct integral decompositions of T and $T|_W$:

$$(\mathfrak{H},T)\simeq \Bigl({\textstyle\int}^{\oplus}\mathfrak{H}_{\lambda}\,d\mu\left(\lambda\right),M\Bigr),$$

$$(W, \mathsf{T}|_W) \simeq \left(\int^{\oplus} W_{\lambda} \, \mathrm{d}\nu(\lambda), \mathsf{N}\right),$$

where M and N are the corresponding multiplication operators. Combining this with our assumption, we obtain

$$(\mathfrak{H},\mathsf{T})\simeq\left(\int^{\oplus}\mathfrak{H}_{\lambda}\,d\mu\left(\lambda\right)\oplus\int^{\oplus}W_{\lambda}\,d\nu\left(\lambda\right),\mathsf{F}\right)=\left(\int^{\oplus}\left(\mathfrak{H}_{\lambda}\oplus W_{\lambda}\right)\,d\left(\mu+\nu\right)\left(\lambda\right),\mathsf{F}\right)$$

where F is the corresponding multiplication operator. Thus, the measures μ and $\mu + \nu$ are mutually absolutely continuous and, since the multiplicity function is determined up to unitary equivalence, we have that for $(\mu + \nu)$ -almost every $\lambda \in \mathbb{R}$, dim $(W_{\lambda}) = 0$ and in particular this happens ν -almost everywhere. This implies that W = 0 and so $V = \mathcal{H}$, as required.

We now consider the general case. Consider the model-theoretic structure $\mathcal{M} := (\mathcal{H}, \mathsf{T}, \mathsf{P}, \mathsf{U})$, where \mathcal{H} and T are as in the statement of the proposition, $\mathsf{P} : \mathcal{H} \to \mathcal{H}$ is the orthogonal projection operator onto V , V is invariant under both T , and T^* , and $\mathsf{U} : \mathcal{H} \to \mathcal{H}$ is the unitary operator that witnesses $(\mathcal{H}, \mathsf{T}) \simeq (\mathsf{V}, \mathsf{T}|_{\mathsf{V}})$. Note that all of this is expressible in first-order logic as follows. Indeed, besides axioms stating that T , P , and U are linear, we add the following axioms:

(1) $\sup_{x} d(Px, P^{2}x) = 0$ (2) $\sup_{x,y} |\langle Px, y \rangle - \langle x, Py \rangle| = 0$ (3) $\sup_{x} d(PTPx, TPx) = 0$ (4) $\sup_{x,y} \langle Px, TPy \rangle - \langle Px, Ty \rangle| = 0$ (5) $\sup_{x,y} |\langle Ux, Uy \rangle - \langle x, y \rangle| = 0$ (6) $\sup_{x} d(PUx, Ux) = 0$ (7) $\sup_{x} \inf_{y} d(Uy, Px) = 0$ (8) $\sup_{x} d(UTx, TUx) = 0$

Note that the 7th item merely states that the image of U is dense in V, but since U is an isometry, it follows that it is in fact onto V.

Suppose now, towards a contradiction, that $V \neq \mathcal{H}$. Then there is $\varepsilon > 0$ such that, letting σ denote the sentence $\inf_x \sup_y (\varepsilon - d(Px, y))$, we have $\sigma^{\mathcal{M}} = 0$. Let $\mathcal{M}' := (\mathcal{H}', \mathsf{T}', \mathsf{P}', \mathsf{U}')$ be a separable elementary substructure of \mathcal{M} , which exists by Downward Löwenheim-Skolem. Note then that $V' := \mathsf{P}'(\mathcal{H}')$ is a closed subspace of \mathcal{H}' invariant under both T' and $(\mathsf{T}')^*$. Since $\sigma^{\mathcal{M}'} = 0$, we have that $V' \subsetneq \mathcal{H}'$, contradicting the first part of the proof.

We are now ready to prove our main theorem. We repeat the statement for the convenience of the reader.

Theorem 2.3. Let $\mathcal{H}_1, \mathcal{H}_2$ be Hilbert spaces and let $\mathsf{T}_i : \mathcal{H}_i \to \mathcal{H}_i$ be bounded normal operators. Suppose that there exist isometries $\mathsf{U}_1 : \mathcal{H}_1 \to \mathcal{H}_2$ and $\mathsf{U}_2 : \mathcal{H}_2 \to \mathcal{H}_1$ such that $\mathsf{T}_1\mathsf{U}_2 = \mathsf{U}_2\mathsf{T}_2$ and $\mathsf{T}_2\mathsf{U}_1 = \mathsf{U}_1\mathsf{T}_1$. Then T_1 and T_2 are unitarily equivalent.

Proof. Set $U := U_1U_2$, so $U : \mathcal{H}_2 \to \mathcal{H}_2$ is an isometry for which $T_2U = UT_2$. We claim that $\widetilde{\mathcal{H}}_2 := U(\mathcal{H}_2)$ is a T-invariant subspace. Indeed, given $\psi \in \widetilde{\mathcal{H}}_2$, take $\phi \in \mathcal{H}_2$ such that $\psi = U_1U_2\phi$. Then we have

$$T_2\psi = T_2U_1U_2\varphi = U_1T_1U_2\varphi = U_1U_2T_2\varphi \in \widetilde{\mathcal{H}}_2.$$

We next claim that $\widetilde{\mathcal{H}}_2$ is also invariant under T^{*}. Indeed, by the Fuglede-Putnam theorem [2, Chapter IX, Theorem 6.7], we have that T^{*}U = UT^{*}. Consequently, if $\psi \in \widetilde{\mathcal{H}}_2$, taking $\varphi \in \mathcal{H}_2$ such that $U\varphi = \psi$, we have $T^*\psi = T^*U\varphi = UT^*\varphi \in \widetilde{\mathcal{H}}_2$.

Let $\widetilde{T}_2 \coloneqq T_2|_{\widetilde{\mathcal{H}}_2}$. Since $T_2 U = UT_2$, we have that $\widetilde{T}_2 U = UT_2$. Consequently, we have the unitarily equivalence $(\mathcal{H}_2, T_2) \simeq (\widetilde{\mathcal{H}}_2, \widetilde{T}_2) = (\widetilde{\mathcal{H}}_2, T|_{\widetilde{\mathcal{H}}_2})$. By Lemma 2.2, we get that $\widetilde{\mathcal{H}}_2 = \mathcal{H}_2$. Finally, note that

$$\mathfrak{H}_{2} = \widetilde{\mathfrak{H}}_{2} = \mathfrak{U}_{1} \left(\mathfrak{U}_{2} \left(\mathfrak{H}_{2} \right) \right) \subseteq \mathfrak{U}_{1} \left(\mathfrak{H}_{1} \right) \subseteq \mathfrak{H}_{2},$$

which of course implies that $U_1(\mathcal{H}_1) = \mathcal{H}_2$. It follows that U_1 is a surjective isometry and thus a unitary transformation. Taking $U = U_1$ concludes the proof of the theorem.

Remark 2.4 (Remark for the model theorists). In classical logic, the SB property implies superstability [4, 5]. It seems likely that the same result holds true in the setting of continuous logic and will be the subject of a forthcoming paper by the first two authors. In that case, a consequence of our main theorem is that the complete theory of any structure of the form (\mathcal{H}, T) , where T is a normal operator on a Hilbert space \mathcal{H} , is superstable. It appears that this result might be folklore, but seems to have not yet appeared in the literature.

3. Unbounded self-adjoint operators

Theorem 3.1. Let \mathcal{H}_1 and \mathcal{H}_2 be Hilbert spaces and let $T_i : \mathcal{H}_i \to \mathcal{H}_i$ be unbounded self-adjoint operators. Further suppose that there exist isometries $U_1 : \mathcal{H}_1 \to \mathcal{H}_2$ and

 $U_2 : \mathcal{H}_2 \to \mathcal{H}_1$ such that $U_1T_1 \subseteq T_2U_1$ and $U_2T_2 \subseteq T_1U_2$. Then $(\mathcal{H}_1, T_1) \simeq (\mathcal{H}_2, T_2)$ in the sense that there is a unitary transformation $U : \mathcal{H}_1 \to \mathcal{H}_1$ such that $UT_1U^{-1} = T_2$.

Proof. For each i = 1, 2, let $V_i := (T_i - i)(T_i + i)^{-1}$ be the Cayley transform of T_i , which is a unitary operator on \mathcal{H}_i . Note then that we have

$$U_1V_1 = U_1(T_1 - i)(T_1 + i)^{-1}$$

= $(T_2 - i)U_1(T_1 + i)^{-1}$
= $(T_2 - i)(T_2 + i)^{-1}U_1$
= V_2U_1 .

In the same way, we have that $U_2V_2 = V_1U_2$. Since each V_i is unitary (and hence normal), the proof of Theorem 2.3, shows that U_1 is a surjective isometry. It follows then that $U_1T_1U_1^{-1} \subseteq T_2$ and thus $U_1T_1U_1^{-1} = T_2$ as self-adjoint operators have no proper symmetric extensions.

References

- [1] C. Argoty, A. Berenstein, and N. Cuervo Ovalle, *The SB-property on metric structures*, Archive for Mathematical Logic, 2025, p. 1-29.
- [2] J.B. Conway, *A course in functional analysis*, Graduate Texts in Mathematics **96** (1990), Springer-Verlag.
- [3] J. Dixmier, *Les algèbres d'opérateurs dans l'espace Hilbertien*, Chapter II, Gauthier-Villars, Paris (1969).
- [4] J Goodrick, When are elementarily bi-embeddable models isomorphic?, Ph.D. thesis, University of California, Berkeley (2007).
- [5] J Goodrick and MC Laskowski, *The Schröder-Bernstein property for a-saturated models*, Proceedings of the American Mathematical Society 142 (2014), 1013-1023.
- [6] T. A. Nurmagambetov, Characterization of ω-stable theories of bounded dimension, Algebra and Logic 28 (1989), 388-396.

DEPARTMENT OF MATHEMATICS, LOS ANDES UNIVERSITY, BOGOTÁ, COLOMBIA, CRA. 1 #18A-12

Email address: n.cuervo10@uniandes.edu.co

Department of Mathematics, University of California, Irvine, 340 Rowland Hall (Bldg.# 400), Irvine, CA 92697-3875

Email address: isaac@math.uci.edu

URL: http://www.math.uci.edu/~isaac

6 NICOLÁS CUERVO OVALLE, ISAAC GOLDBRING, AND NETANEL LEVI

Department of Mathematics, University of California, Irvine, 340 Rowland Hall (Bldg.# 400), Irvine, CA 92697-3875

Email address: netanell@uci.edu