TOPOLOGICAL ADELIC CURVES: ALGEBRAIC COVERINGS, GEOMETRY OF NUMBERS AND HEIGHTS OF CLOSED POINTS

ANTOINE SÉDILLOT

ABSTRACT. In this article, we introduce topological adelic curves. Roughly speaking, a topological adelic curve is a topological space of (generalised) absolute values on a given field satisfying a product formula. Topological adelic curves are topological counterparts to adelic curves introduced by Chen and Moriwaki. They aim at handling Arakelov geometry over possibly uncountable fields and give further ideas in the formalisation of the analogy between Diophantine approximation and Nevanlinna theory. Using the notion of pseudo-absolute values developed in [Séd24a], we prove several fundamental properties of topological adelic curves: algebraic coverings, Harder-Narasimhan formalism, existence of volume functions. We also define height of closed points and give a generalisation of Nevanlinna's first main theorem in this framework.

CONTENTS

Introduction	2
Conventions and notation	8
Part 1. Topological adelic curves: definition and algebraic coverings	9
1. Reminders on pseudo absolute values and adelic curves	9
2. Topological adelic curves	14
3. Families of topological adelic curves	17
4. Algebraic covering of topological adelic curves	20
Part 2. Adelic vector bundles and Harder-Narasimhan filtrations over	
topological adelic curves	25
5. Pseudo-norm families	25
6. Adelic vector bundles	36
7. Slopes of adelic vector bundles: proper case	42
Part 3. Arithmetic varieties over topological adelic curves: adelic line	
bundles and heights	47
8. Pseudo-metrics: local case	47
9. Pseudo-metric families	52
10. Height of closed point	68
Appendix A. Nevanlinna theory of complex functions	71

Date: March 27, 2025.

The author was partly supported by the collaborative research center SFB 1085 *Higher Invariants - Interactions between Arithmetic Geometry and Global Analysis* funded by the Deutsche Forschungsgemeinschaft. The main part of this research was conducted at Université Paris Cité.

References

INTRODUCTION

Motivations and background.

Arakelov geometry over number fields and arithmetic function fields. Arakelov theory stems from the analogy between number fields and function fields. Roughly speaking, one can formulate this analogy as follows: the geometry of schemes of finite type over $\text{Spec}(\mathbb{Z})$ should be similar to the geometry of schemes of finite type over a smooth projective curve. Unfortunately, schemes over $\text{Spec}(\mathbb{Z})$ are not "compact" and it is not quite clear how to "compactify" them within the world of schemes. To address this issue, Arakelov [Ara74] added analytic data to algebro-geometric objects. Arakelov's ideas have been used by Faltings [Fal91] in his proof of Mordell's conjecture and in the proof of Bogomolov's conjecture [Ull98, Zha98].

Studying the arithmetic of fields that are more general than global fields has also been developed to a great extent. Lang remarked that it was natural to study *arithmetic function fields*, namely finite type field extensions of \mathbb{Q} [Lan74, Lan86]. Indeed, Mordell-Weil's and Faltings' theorems both hold over such fields [Lan91]. Later, Moriwaki constructed a height theory over arithmetic function fields [Mor00] (see also [BGPS16]). Recently, Vojta proved a version of Roth's theorem over arithmetic function fields [Voj21].

The study of infinite algebraic extensions of number fields has also been a great inspiration for developing analogues of the tools of Diophantine geometry. Let us mention for instance a version of Siegel's lemma [RT96], the study of tensorial semistability [BC13] and the introduction of Siegel fields [GR17].

Arakelov geometry over adelic curves. In [CM19], Chen and Moriwaki introduced an Arakelov theory over arbitrary countable fields. The central object of the theory is called an *adelic curve*. Namely, an adelic curve is the data $S = (K, (\Omega, \nu), (| \cdot |_{\omega})_{\omega \in \Omega})$, where K is a field, (Ω, ν) is a measure space and $(| \cdot |_{\omega})_{\omega \in \Omega}$ is a family of absolute values on K. Moreover, an adelic curve $S = (K, (\Omega, \nu), (| \cdot |_{\omega})_{\omega \in \Omega})$ is called *proper* if the following *product formula* holds.

$$\forall a \in K^{\times}, \quad \int_{\Omega} \log |f|_{\omega} \nu(\mathrm{d}\omega) = 0.$$

Adelic curves arise naturally in various number theoretic situations. In particular, any global field can be naturally equipped with an adelic structure. More generally, any countable field can be endowed with an adelic structure. Furthermore, adelic curves allow to study global fields, trivially valued fields and arithmetic function fields uniformly.

Let us now introduce the counterpart of the usual tools of Arakelov geometry over adelic curves. Let $S = (K, (\Omega, \nu), (|\cdot|_{\omega})_{\omega \in \Omega})$ be an adelic curve. For any K-scheme X, for any $\omega \in \Omega$, we denote by K_{ω} the completion of K with respect to the absolute value $|\cdot|_{\omega}$ and $X_{\omega} := X \otimes_K K_{\omega}$. The avatar of a line bundle in algebraic geometry is called an *adelic line bundle*. Let X be a K-scheme, an adelic line bundle over X is the data $\overline{L} = (L, \varphi)$, where L is a line bundle over X and $\varphi = (\varphi_{\omega})_{\omega \in \Omega}$ is a family of continuous metrics (in the sense of Berkovich analytic spaces) over each $L_{\omega} := L \otimes_{\mathcal{O}_X} \mathcal{O}_{X_{\omega}}$. In addition, the metric family is subject to dominance and measurability conditions (cf. §6.1 in [CM19]). Among the results of the theory, let us first mention the development of the geometry of numbers via the slope theory of adelic vector bundles ([CM19], Chapter IV). Note that even in the case of number fields, Chen-Moriwaki's approach yields a new interpretation of known results. An arithmetic intersection theory for adelic line bundles is also constructed in [CM21]. The study of the positivity of adelic line bundles and a Hilbert-Samuel formalism are introduced in [CM22]. As an application, a generalisation of Bogomolov's conjecture over a (countable) field of characteristic zero is proven. Note also that a version of Roth's theorem over a particular class of adelic curves is established by Dolce and Zucconi in [DZ23]. This result generalises Vojta's aforementioned result [Voj21].

Nevanlinna theory and M-fields. Another, but yet not disconnected, motivation for our work is to study the analogy between Diophantine geometry and Nevanlinna theory. This analogy was spotted first by Osgood [Osg81] and further explored by Vojta in [Voj87]. Roughly speaking, Nevanlinna theory is the study of equations of the form f(z) = a, where f is meromorphic on \mathbb{C} and $a \in \mathbb{P}^1(\mathbb{C})$. It builds on two fundamental theorems. Through the analogy, the first one corresponds to Weil's theorem for heights (e.g. [BG06], Theorem 2.3.8). The second one is seen as an analogue of Roth's theorem [Rot55]. In Appendix A, we recalled the basic notions of Nevanlinna theory as well as the main ideas of the analogy.

In [Gub97], Gubler introduced the notion of *M*-fields, with the idea of including Nevanlinna theory in an Arakelov theoretic framework. Roughly speaking, a *M*-field *K* is a field *K* equipped with a measure space *M* such that any element *a* in *K* defines an integrable real function |a|. defined almost-everywhere on *M*. Moreover, these functions are assumed to satisfy the axioms of absolute values almost everywhere. The following example of *M*-field coming from Nevanlinna theory is fundamental for our purposes. Consider the field $\mathcal{M}(\mathbb{C})$ of meromorphic complex functions. Fix a real number R > 0 and set $M_R := \{z \in \mathbb{C} : |z|_{\infty} \leq R\}$ where the boundary $\{z \in \mathbb{C} : |z|_{\infty} = R\}$ is equipped with the Haar probability measure and the open disc $\{z \in \mathbb{C} : |z|_{\infty} < R\}$ is equipped with a counting measure. For any $f \in \mathcal{M}(\mathbb{C})$, consider the map

$$(z \in M_R) \mapsto \begin{cases} |f(z)|_{\infty} & \text{if } |z|_{\infty} = R, \\ e^{-\operatorname{ord}(f,z)} & \text{if } |z|_{\infty} = R, \end{cases}$$

which is well-defined everywhere except poles of f on the circle of radius R, hence almost everywhere. Then one can check that we have a M_R -field $\mathcal{M}(\mathbb{C})$.

Using M-fields, Gubler obtains a generalisation of Nevanlinna's first main theorem which includes notably the construction of a height function for fields of arithmetic nature. Nonetheless, it does not seem clear how one could hope to obtain further results, e.g. geometry of numbers, in the framework of M-fields.

Goal. The goal of this article is to introduce objects of Arakelov geometric nature allowing to handle uncountable fields and to formalise the analogy with Nevanlinna theory.

Hints from the above discussion. On the one hand, in the theory of adelic curves, the countability condition is imposed by the fact that the parameter space of absolute values is a measure space. This is due to the apparition of suprema and infima of measurable functions when considering operations on adelic vector bundles. Although this approach gives a lot of flexibility, we cannot expect the tools to transpose directly in the uncountable

setting. A natural idea is to consider a topological space as parameter space and replace the measurability conditions with (semi-)continuity ones.

On the other hand, the Nevanlinna theory example of M-field suggests that the space of possible arithmetic inputs should be larger than the space of usual absolute values. Note that even in the case of a classical adelic curve structure on $\mathbb{Q}(T)$, namely coming from classical Arakelov geometry over $\mathbb{P}^1_{\mathbb{Z}}$ ([CM19], §3.2.5), the natural topological space parametrising the Archimedean absolute values is the set of transcendental elements of the complex unit disc. From the point of view of measure theory, it is just the difference of the complex unit disc and the countable subset of algebraic numbers. However, this space equipped with its usual topology is very pathological.

Pseudo-absolute values and globally valued fields. The framework of globally valued fields, introduced by Ben Yaacov-Hrushovski [Hru16], gives another approach to handle arithmetic over fields. Roughly speaking, a *globally valued field* (GVF for short), is a field equipped with a family of heights satisfying the usual height compatibility axioms. This notion originates from model theory and there are several equivalent characterisations of GVFs ([BYDHS24], Theorem 7.7). The link with the above discussion is the following: a countable GVF is an equivalence class of proper adelic curves (*loc. cit.*, Corollary 7.11). This link with model theory yields another motivation for developing Arakelov geometry over uncountable fields: indeed, the ultraproduct construction is fundamental in model theory and in general, ultraproducts are uncountable.

Over a possibly uncountable field K, the GVF structures can be interpreted as a suitable measure on the space of pseudo-absolute values (or of pseudo-valuations) on K. More precisely, a pseudo-absolute value on a field K is a map $|\cdot|: K \to [0, +\infty]$ satisfying

- (i) |1| = 1 and |0| = 0;
- (ii) for all $a, b \in K$, $|a + b| \le |a| + |b|$;
- (iii) for all $a, b \in K$ such that $\{|a|, |b|\} \neq \{0, +\infty\}, |ab| = |a||b|.$

Moreover, $A_{|\cdot|} := \{a \in K : |a| < +\infty\}$ is a valuation ring of K with maximal ideal $m_{|\cdot|} := \{a \in K : |a| = 0\}$ and $|\cdot|$ induces an absolute value on the residue field $\kappa_{|\cdot|} := A_{|\cdot|}/\mathfrak{m}_{|\cdot|}$. In other terms, a pseudo-absolute value is an absolute value allowing "singularities". This notion was first introduced by Weil in [Wei51] and was developed independently by Ben Yaacov-Destic-Hrushovski-Szachniewicz [BYDHS24] and the author in [Séd24a].

Moreover, the space M_K of all pseudo-absolute values on a field K equipped with the topology of pointwise convergence is compact Hausdorff and can be interpreted as a Berkovich Zariski-Riemann space ([BYDHS24], Proposition 2.7 and [Séd24a], Theorems A-C). Considering the development of Berkovich's non-Archimedean analytic geometry [Ber90, CL06, CLD12, GK17, GK19], it is natural to expect a "global analytic" approach to Arakelov geometry [Pau09, YZ21, CG24].

Pseudo-absolute values are thus a natural candidate to encode the local aspects of the theoretical constructions that will follow later on. Their theory was developed in detail in [Séd24a] with this goal in mind.

Content of the article. This article consists of three parts. In the first one, we introduce topological adelic curves. A topological adelic curve is the data $S = (K, \phi : \Omega \to V, \nu)$, where K is a field, Ω is a locally compact Hausdorff topological space, $\phi : (\omega \in \Omega) \mapsto |\cdot|_{\omega} \in M_K$ is a continuous map between Ω and the set M_K of all pseudo-absolute values on K, and ν is a

Borel measure on Ω such that, for any $f \in K^{\times}$, the function $(\omega \in \Omega) \mapsto \log |f|_{\omega} \in [-\infty, +\infty]$ is ν -integrable (§2). We say that the topological adelic curve S is proper if the following product formula

$$\forall f \in K^{\times}, \quad \int_{\Omega} \log |f|_{\omega} \nu(\mathrm{d}\omega) = 0,$$

holds.

Let us give an example of a topological adelic curve arising in Nevanlinna theory. We fix R > 0. We define a topological adelic curve $S_R = (K_R, (\Omega_R, \nu_R), \phi_R)$, where

- K_R is the field of (germs of) meromorphic functions over the closed disc $\overline{D(R)} :=$
- $\{z \in \mathbb{C} : |z|_{\infty} \leq R\} \subset \mathbb{C};$ $\Omega_R = \{z \in \mathbb{C} : |z|_{\infty} < R\} \coprod \{z \in \mathbb{C} : |z|_{\infty} = R\}, \text{ where } \{z \in \mathbb{C} : |z|_{\infty} < R\} \text{ is }$ equipped with the discrete topology and $\{z \in \mathbb{C} : |z|_{\infty} = R\}$ is equipped with the usual topology;
- the map $\phi_R : \Omega_R \to M_{K_R}$ is defined by

$$\forall z \in \Omega_R, \quad \phi_R(z) := \begin{cases} (f \in K_R) \mapsto |f(z)|_{\infty} \in [0, +\infty] & \text{if } |z|_{\infty} = R\\ e^{-\operatorname{ord}(\cdot, z)} & \text{if } |z|_{\infty} < R \end{cases}$$

• the measure ν_R is defined by

$$\forall z \in \{z' \in \mathbb{C} : |z'|_{\infty} < R\}, \quad \nu_R(\{z\}) := \begin{cases} \log \frac{R}{|z|_{\infty}} & \text{if} \quad 0 < |z|_{\infty} < R, \\ \log R & \text{if} \quad z = 0, \end{cases}$$

and ν_R is the Lebesgue probability measure on $\{z \in \mathbb{C} : |z|_{\infty} = R\}$.

A major obstacle in the above construction lies in the fact that the product formula is not satisfied. Indeed, Jensen's formula yields

$$\forall f \in K_R^{\times}, \quad \int_{\Omega_R} \log |f|_{\omega} \nu_R(\mathrm{d}\omega) = \log |c(f,0)|,$$

where c(f,0) denotes the first non-zero coefficient in the Laurent series expansion of f in 0.

To address the above issue, we introduce the notion of *families of topological adelic curves*. Let us present the general idea. In the above example, the product formula does not hold. In the context of Diophantine geometry, the product formula is the ingredient that allows to define height functions relative to a metrised line bundle. The analogue of this construction in Nevanlinna theory is the first main theorem. The latter results from the fact that the defect in the product formula for the collection of topological adelic curves $(S_R)_{R>0}$ is a bounded function of R (in fact constant in the present case). Starting from this observation, we can define the (compatible) families of topological adelic curves and define an (asymptotic) notion of properness for these families. §3).

We conclude the first part by studying algebraic coverings of topological adelic curves (§4), namely the extension of topological adelic structures w.r.t. algebraic extensions of the base field. More precisely, we have the following result (which is a variant of Proposition 1.9 in [BYDHS24]).

Theorem A (Propositions 4.3.1, 4.4.1 and Remark 4.4.2). Let $S = (K, \phi : \Omega \to V, \nu)$ be a topological adelic curve. Let K'/K be an algebraic field extension.

(1) There exist a topological adelic curve $S' := S \otimes_K K' := (K', \phi' : \Omega' \to M_{K'}, \nu')$ and a morphism of topological adelic curves $S' \to S$. Moreover, S' is proper if so is S.

(2) Assume that K'/K is Galois and that, for any $\omega \in \Omega$, the residue field of the pseudo-absolute value $|\cdot|_{\omega}$ is perfect. Then we have a homeomorphism

 $\Omega' / \operatorname{Aut}(L/K) \cong \Omega.$

Moreover, if ν is Radon, then S' is the only topological adelic curve with base field K' extending S' with Galois-invariant measure.

The second part of the article is devoted to defining adelic vector bundles over a topological adelic curve and to constructing Harder-Narasimhan filtrations. Let $S = (K, \phi : \Omega \to V, \nu)$ be a topological adelic curve. A *pseudo-norm family* on a *K*-vector space of finite rank *E* is a family $\xi = (\|\cdot\|_{\omega})_{\omega\in\Omega}$, where $\|\cdot\|_{\omega}$ is a pseudo-norm on *E* over the pseudo-absolute value $\phi(\omega) \in M_K$ (the suitable generalisation of a norm over a valued field). An *adelic vector bundle* on *S* is then defined as a pair $\overline{E} = (E, \xi)$, where *E* is a *K*-vector space of finite rank and ξ is a pseudo-norm family on *E* satisfying suitable regularity and integrability conditions (cf. §5). Adelic vector bundles are "almost stable" with respect to the usual algebraic operations performed over vector spaces (cf. Proposition 6.1.4). Concerning the "almost stable" assertion, let us mention for now that the algebraic operations preserve the integrability property that will be needed in what follows. We also give the family counterpart of adelic vector bundles over a family of topological adelic curves and include an example coming from Nevanlinna theory (§6.4).

In the case where the base topological adelic curve is proper, the Arakelov degree of an adelic vector bundle $\overline{E} = (E, \xi)$ is defined as

$$\widehat{\operatorname{deg}}(\overline{E}) := -\int_{\Omega} \log \|\eta\|_{\omega, \det} \nu(\mathrm{d}\omega),$$

where $\eta \in \det(E) \setminus \{0\}$, is independent of the choice of η . Denote also by $\mu(\overline{E}) := \widehat{\deg}(\overline{E} =)/\dim_K(E)$ the *slope* of \overline{E} . Define as well the *maximal slope* $\mu_{\max}(\overline{E}) := \sup_{0 \neq F \subseteq E} \mu(\overline{F})$ and the *minimal slope* $\widehat{\mu}_{\min}(\overline{E}) := \inf_{E \to G \neq \{0\}} \widehat{\mu}(\overline{G})$.

In §7, we adapt the Harder-Narasimhan formalism for adelic vector bundles over a proper topological adelic curve.

Theorem B (Theorem 7.4.2). Let $\overline{E} = (E, \xi)$ be an adelic vector bundle on S. We assume that the pseudo-norm family ξ is ultrametric on Ω_{um} . Then there exists a unique flag

$$0 = E_0 \subsetneq E_1 \subsetneq \cdots \subsetneq E_n = E,$$

of \overline{E} , such that

- (1) for any i = 1, ..., n, $\overline{E_i/E_{i-1}}$ is semistable, i.e. for any non-zero vector subspace $F \subset E$, we have $\mu_{\min}(\overline{F}) \leq \mu_{\min}(\overline{E})$;
- (2) we have the inequalities

$$\widehat{\mu}(\overline{E_1/E_0}) > \cdots > \widehat{\mu}(\overline{E_n/E_{n-1}}).$$

In the third and last part, we introduce the higher dimensional counterpart of the previous part: namely we study algebraic varieties over a topological adelic curve. We fix a topological adelic curve $S = (K, \phi : \Omega \to V, \nu)$ and a projective K-scheme $\pi : X \to \text{Spec}(K)$.

As customary in Arakelov geometry, we want to consider analytic spaces attached to X at each $\omega \in \Omega$. Since elements of Ω correspond to pseudo-absolute values, we need some adjustments. Fix a family $\mathcal{X} = (\mathcal{X}_{\omega})_{\omega \in \Omega}$, where the \mathcal{X}_{ω} 's are projective schemes over

the finiteness ring $A_{\omega} = \{a \in K : |a|_{\omega} < +\infty\}$ with generic fibre X. The analytic space associated with X in ω (w.r.t. \mathcal{X}) is defined as $X_{\omega}^{\operatorname{an}} := (\mathcal{X}_{\omega} \otimes_{A_{\omega}} \widehat{\kappa_{\omega}})^{\operatorname{an}}$, where $\widehat{\kappa_{\omega}}$ denotes the completion of the residue field of A_{ω} and analytifications are considered in the sense of Berkovich. Let L be a line bundle on X and $\omega \in \Omega$. The usual notion of metric on Berkovich spaces can be transposed in this context and is called *pseudo-metric* (§8.1), it is the data $(\mathcal{L}_{\omega}, \varphi_{\omega})$ of a line bundle \mathcal{L}_{ω} on \mathcal{X}_{ω} with generic fibre L and a metric φ_{ω} on the pullback of \mathcal{L}_{ω} to $X_{\omega}^{\operatorname{an}}$. Therefore a *pseudo-metric family* on X is defined as a family $(\mathcal{L}, \varphi) = ((\mathcal{L}_{\omega}, \varphi_{\omega}))_{\omega \in \Omega}$ where the $(\mathcal{L}_{\omega}, \varphi_{\omega})$'s are pseudo-metrics on L in the ω 's. $(L, (\mathcal{L}, \varphi))$ is called *adelic line bundle* if the pseudo-metric family (\mathcal{L}, φ) satisfies certain regularity and integrability conditions (§9.1-9.5). We also extend this construction to the context of adelic line bundles, including an example in Nevanlinna theory (§9.6).

We also discuss the pushforward of adelic line bundles. More precisely, let $L = (L, (\mathcal{L}_{\omega}, \varphi_{\omega})_{\omega \in \Omega})$ be an adelic line bundle on X. Assume that X is geometrically reduced and that all the models \mathcal{X}_{ω} 's are flat and finitely presented. Then by taking supremum pseudo-norms, $(\varphi_{\omega})_{\omega \in \Omega}$ defines a pseudo-norm family $\pi_*\varphi = (\|\cdot\|_{\omega})_{\omega \in \Omega}$ on $\pi_*L = H^0(X, L)$ (Proposition-Definition8.1.1). Under technical conditions, we prove that $\pi_*\overline{L} := (\pi_*L, \pi_*\varphi)$ is an adelic vector bundle on S(Theorem 9.7.9). This result allows us to define arithmetic $(\chi$ -)volume functions on proper topological adelic curves (§9.8).

The last section introduces heights of closed points. In the proper case, our result is the following.

Theorem C (Theorem 10.1.2). Assume that the topological adelic curve S is proper. Let $\overline{L_1} = (L_1, \varphi_1), \overline{L_2} = (L_2, \varphi_2)$ be adelic line bundles on X. Then the following assertions hold. (1) For any closed point P of X, we have

$$h_{\overline{L_1}+\overline{L_2}}(P) = h_{\overline{L_1}}(P) + h_{\overline{L_2}}(P).$$

(2) Let P be a closed point of X. Assume that $L_1 = L_2$ and that φ_1, φ_2 have the same model family. Then we have

$$h_{\overline{L_1}}(P) = h_{\overline{L_2}}(P) + O(1),$$

where the bound does not depend on P (but depends on φ_1, φ_2).

If the topological adelic curve is not proper, e.g. in Nevanlinna theory, we have a family analogue. Let us describe it in the previous example. We consider the family $\mathbf{S} = (S_R = (K_R, \phi_R : \Omega_R \to V_R, \nu_R))_{R>0}$ of topological adelic curves arising in Nevanlinna theory. Denote $K_{\infty} := \varprojlim_{R>0} K_R = \mathcal{M}(\mathbb{C})$, namely the field of meromorphic functions on \mathbb{C} . Let $X_{\infty} \to \operatorname{Spec}(K_{\infty})$ be a geometrically integral projective variety. For any R > 0, denote $X_R := X_{\infty} \otimes_{K_{\infty}} K_R$. Any closed point $P_{\infty} \in X_{\infty}$ defines a family $\mathbf{P} := (P_R)_{R>0}$, where the P_R 's are closed points of the X_R 's.

Then we have the notion of adelic line bundles on the family $\mathbf{X} := (X_R)_{R>0}$ (§9.6). Let $\overline{\mathbf{L}} = (\overline{L_R})_{R>0}$ be an adelic line bundle on \mathbf{X} , in particular the L_R 's come from a line bundle L_{∞} on X_{∞} . An important example is the following. Assume that there exists a complex geometrically integral projective variety X such that $X_{\infty} = X \otimes_{\mathbb{C}} K_{\infty}$. Then any metrised line bundle on X defines an adelic line bundle on \mathbf{X} . Such adelic line bundles are called *induced by a complex metrised line bundle*.

Let $P_{\infty} \in X_{\infty}$ be a closed point and let s_{∞} be a meromorphic section of L_{∞} which is non-vanishing at P_{∞} . This defines a family $\mathbf{s} = (s_R)_{R>0}$ such that the s_R 's are meromorphic

sections of the L_R 's which are non vanishing at the P_R 's. We can now define the *height* $h_{\overline{\mathbf{L}},\mathbf{s}}(\mathbf{P}): (R > 0) \to \mathbb{R}$ of \mathbf{P} w.r.t. $\overline{\mathbf{L}}$ and \mathbf{s} (§10.2). Note that the choice of \mathbf{s} does not modify the height up to a bounded function: namely, if \mathbf{s}' is another suitable family of sections such that the height $h_{\overline{\mathbf{L}},\mathbf{s}'}(\mathbf{P})$ makes sense, we have

$$h_{\overline{\mathbf{L}},\mathbf{s}'}(\mathbf{P}) = h_{\overline{\mathbf{L}},\mathbf{s}}(\mathbf{P}) + O(1)$$

Our result is the following.

Theorem D (cf. Theorem 10.2.2 for a more general statement). Let $\overline{\mathbf{L}^{(1)}}, \overline{\mathbf{L}^{(2)}}$ be adelic line bundles on **X**. Let P_{∞} be a closed point of X_{∞} and $s_{\infty}^{(1)}, s_{\infty}^{(2)}$ be meromorphic sections of $L_{\infty}^{(1)}, L_{\infty}^{(2)}$ non vanishing at P_{∞} respectively, yielding families $\mathbf{s}^{(1)}, \mathbf{s}^{(2)}$. Then the following assertions hold.

(1) We have

$$h_{\overline{\mathbf{L}^{(1)}}+\overline{\mathbf{L}^{(2)}},\mathbf{s}^{(1)}+\mathbf{s}^{(2)}}(P) = h_{\overline{\mathbf{L}^{(1)}},\mathbf{s}^{(1)}}(P) + h_{\overline{\mathbf{L}^{(2)}},\mathbf{s}^{(2)}}(P).$$

(2) Assume that the underlying line bundles of $\overline{\mathbf{L}^{(1)}}$ and $\overline{\mathbf{L}^{(2)}}$ coincide and that they are both induced by a complex metrised line bundle. Then we have

$$h_{\overline{\mathbf{L}^{(1)}},\mathbf{s}^{(1)}}(P) = h_{\overline{\mathbf{L}^{(2)}},\mathbf{s}^{(2)}}(P) + O(1)$$

with bound independent on P.

This result gives a generalisation of Nevanlinna's first main theorem and of ([Gub97], Theorem 3.18) in the case of closed points.

Upcoming work. Let us mention open questions that should be addressed in the future.

- In view of the sketched results from ([Séd24b], Chapter IV), an arithmetic intersection theory of adelic line bundles should be defined over a proper topological adelic curve. This would allow to define heights of subvarieties and would give the full generalisation of ([Gub97], Theorem 3.18). Moreover, in view of the results in [DHS24], we expect the definability of this arithmetic intersection product over an arbitrary GVF, which could lead to a generalisation of the results in this article over more general fields.
- The definition of arithmetic volumes invites questions concerning the regularity of these volumes. In view of [Séd23], we expect differentiability properties to hold for these volumes. Note that this differentiability is a key ingredient for proving existential closedness for globally valued fields (cf. [Sza23]) and would achieve a major step towards the existence of a model companion for the language of GVF ([BYDHS24], Conjecture 12.7).

Acknowledgements. We would like to thank Huayi Chen for his support and discussions during the elaboration of this paper. We also thank Keita Goto, Walter Gubler, Klaus Künnemann and Jérôme Poineau for numerous remarks and suggestions.

CONVENTIONS AND NOTATION

- All rings considered in this article are commutative with unit.
- Let A be a ring. We denote by Spm(A) the set of maximal ideals of A.
- By a local ring (A, \mathfrak{m}) , we mean that A is a local ring and \mathfrak{m} is its maximal ideal. In general, if A is a local ring, the maximal ideal of A is denoted by \mathfrak{m}_A .

- Let A be a ring and let $X \to \operatorname{Spec}(A)$ be a scheme over A. Let $A \to B$ be an A-algebra. Then we denote $X \otimes_A B := X \times_{\operatorname{Spec}(A)} \operatorname{Spec}(B)$.
- Let K be a field and $X \to \operatorname{Spec}(K)$ be a K-scheme. For any domain A with fraction field K, we call model of X/A any A-scheme $\mathcal{X} \to \operatorname{Spec}(A)$ whose generic fibre is isomorphic to X. A model of $\pi : \mathcal{X} \to \operatorname{Spec}(A) X/A$ is respectively called projective, flat, coherent if π is projective, flat, finitely presented.
- Let k be a field. We denote by $|\cdot|_{\text{triv}}$ the trivial absolute value on k. If we have an embedding $k \hookrightarrow \mathbb{C}$, we denote by $|\cdot|_{\infty}$ the restriction of the usual Archimedean absolute value on \mathbb{C} .
- Let $(k, |\cdot|)$ be a valued field. Unless mentioned otherwise and when no confusion may arise, we will denote by \hat{k} the completion of k w.r.t. $|\cdot|$.
- Let (X, \mathcal{O}_X) be a locally ringed space. We make no distinction between line bundles and invertible \mathcal{O}_X -module and we use the additive notation for tensor products of line bundles.
- Let $(\Omega, \mathcal{A}, \nu)$ be a measure space. Denote by $\mathcal{L}^1(\Omega, \mathcal{A}, \nu)$ be the set of all ν -integrable functions $f: \Omega \to [-\infty, +\infty]$. Let $f: \Omega \to [-\infty, +\infty]$, we define

$$\int_{\Omega}^{\bar{}} f(\omega)\nu(\mathrm{d}\omega) := \inf\left\{\int_{\Omega} g(\omega)\nu(\mathrm{d}\omega) : g \in \mathcal{L}^{1}(\Omega, \mathcal{A}, \nu) \text{ and } f \leq g \quad \nu\text{-a.e.}\right\},$$

and

$$\underline{\int}_{\Omega} f(\omega)\nu(\mathrm{d}\omega) := \sup\left\{\int_{\Omega} g(\omega)\nu(\mathrm{d}\omega) : g \in \mathcal{L}^{1}(\Omega, \mathcal{A}, \nu) \text{ and } g \leq f \quad \nu\text{-a.e.}\right\}.$$

We say that f is ν -dominated if

$$\int_{\Omega} \overline{f}(\omega)\nu(\mathrm{d}\omega) < +\infty \quad \mathrm{and} \quad \underline{\int}_{\Omega} f(\omega)\nu(\mathrm{d}\omega) > -\infty.$$

Equivalently, f is ν -dominated iff there exists $g \in \mathcal{L}^1(\Omega, \mathcal{A}, \nu)$ such that $|f| \leq g \nu$ -a.e.

Part 1. Topological adelic curves: definition and algebraic coverings

1. Reminders on pseudo absolute values and adelic curves

1.1. Pseudo-absolute values.

1.1.1. Definition. Let K be a field. A pseudo-absolute value on K is a map $|\cdot|: K \to [0, +\infty]$ satisfying

- (i) |0| = 0 and |1| = 1;
- (ii) for all $a, b \in K$, $|a + b| \le |a| + |b|$;
- (iii) for all $a, b \in K$ such that $\{|a|, |b|\} \neq \{0, +\infty\}, |ab| = |a||b|.$

Recall that any pseudo-absolute value $|\cdot|$ on K determines a finiteness ring $A_{|\cdot|} = \{|\cdot| < +\infty\}$, which is a valuation ring of K with maximal ideal $\mathfrak{m}_{|\cdot|} = \{|\cdot| = 0\}$, called its *kernel*. Moreover, $|\cdot|$ induces an absolute value on the residue field $\kappa_{|\cdot|} := A_{|\cdot|}/\mathfrak{m}_{|\cdot|}$ called the residue absolute value. A pseudo-absolute value is called Archimedean, non-Archimedean, residually trivial if the associated residue absolute value is Archimedean, non-Archimedean, trivial.

We use the same notation as in [Séd24a]. Namely, by "let $(|\cdot|, A, \mathfrak{m}, \kappa)$ be a pseudo-absolute value", we mean that $|\cdot|$ is a pseudo-absolute value on K with finiteness ring A, kernel \mathfrak{m} , residue field κ . By default, when we write "let v be a pseudo absolute value", we mean

 $v = (| \cdot ||_v, A_v, \mathfrak{m}_v, \kappa_v)$. Moreover, if v is a pseudo-absolute value on K, we denote by $\widehat{\kappa_v}$ the completion of the residue field κ_v w.r.t. the residue absolute value induced by v.

1.1.2. Extension of pseudo-absolute values. Let L/K be a finite separable field extension. Let v be a pseudo-absolute value on K. Denote by A' the integral closure of A_v in L. This is a Prüfer domain, namely its prime localisations are valuation rings. Moreover, the extensions of A_v to L are in bijection with Spm(A'). For any $\mathfrak{m}_w \in \text{Spm}(A')$, we denote by κ_w the corresponding residue field, this is a finite field extension of κ_v .

- **Proposition 1.1.1** ([Séd24a], Proposition 3.1.2). (1) There is a bijective correspondence between the set of pseudo-absolute values on L above v and the set of extensions of the residue absolute value of v with respect to extensions of the form $\kappa_v \to \kappa_w$, where w runs over the set of maximal ideals of A'.
 - (2) Furthermore, we have the equality

$$\sum_{\mathbf{n}_w \in \operatorname{Spm}(A')} \frac{1}{|\operatorname{Spm}(A')|} \sum_{i|v} \frac{[\widehat{\kappa_{w,i}} : \widehat{\kappa_v}]_s}{[\kappa_w : \kappa_v]_s} = 1,$$
(1)

where, for all $\mathfrak{m}_w \in \operatorname{Spm}(A')$, *i* runs over the set of extensions of the residue absolute value of $|\cdot|_v$ to κ_w and $\widehat{\kappa_{w,i}}$ denotes the completion of κ_w for any such absolute value.

Now consider an arbitrary finite field extension L/K. Denote by K' the separable closure of K in L.

Proposition 1.1.2 ([Séd24a], Corollary 3.2.2). Let v be a pseudo-absolute value on K. Then the set of extensions of v on L is in bijection with the set of extensions of v on K'.

The following proposition characterises the action of the group of automorphisms on pseudo-absolute values.

Proposition 1.1.3 ([Séd24a], Propositions 3.3.1 and 3.3.4). Let L/K be an algebraic field extension.

(1) Aut(L/K) induces a right action on M_L as follows. For all $x \in M_L$, for all $\tau \in Aut(L/K)$, the map

$$|\cdot|_{\tau(x)} : \begin{vmatrix} L & \longrightarrow & [0, +\infty] \\ a & \longmapsto & |\tau(a)|_x \end{vmatrix}$$

defines a pseudo-absolute value on L denoted by $x \circ \tau$.

(2) Assume that L/K is Galois with Galois group G. Let $v \in M_K$ such that the residue field κ_v is perfect. Then G acts transitively on the set $M_{L,v}$ of extensions of v to L.

1.1.3. Space of pseudo-absolute values and integral structures. Let K be a field. Recall that the set M_K of all pseudo-absolute values on K equipped with the topology of point-wise convergence is a (non-empty) compact Hausdorff topological space (*loc. cit.*, Theorem 7.1.2). We denote by $M_{K,ar}$ and $M_{K,um}$ respectively the set of Archimedean and non-Archimedean pseudo-absolute values on K. Define a map $\epsilon : M_{K,ar} \to]0,1]$ by sending any $|\cdot| \in M_{K,ar}$ to the unique $\epsilon(|\cdot|) \in]0,1]$ such that the restriction of the residue absolute value of $|\cdot|$ to \mathbb{Q} is $|\cdot|_{\infty}^{\epsilon(|\cdot|)}$.

An integral structure for K is a Banach ring $(A, \|\cdot\|)$ such that A is a Prüfer domain with fraction field K. If $(A, \|\cdot\|)$ is an integral structure for K, then the Berkovich analytic spectrum $\mathcal{M}(A, \|\cdot\|)$ identifies as a closed subspace of M_K (*loc. cit.*, Proposition 9.1.5). Moreover, an integral structure $(A, \|\cdot\|)$ for K is called *tame* if

- (i) $\mathcal{M}(A, \|\cdot\|)$ contains the trivial absolute value on K;
- (ii) for any ultrametric element $|\cdot|_x \in \mathcal{M}(A, \|\cdot\|)$ and any $f \in A$, the inequality

 $|f|_x \le 1$

is satisfied;

(iii) $(A, \|\cdot\|)$ is a uniform Banach ring.

Proposition 1.1.4 ([Séd24a], Proposition 9.3.3). Let $(A, \|\cdot\|)$ be a tame integral structure for K. Let L/K be an algebraic extension. Then the integral closure B of A in L can be equipped with a norm $\|\cdot\|_B$ such that $(B, \|\cdot\|_B)$ is a tame integral structure for L. Moreover, $\mathcal{M}(B, \|\cdot\|_B)$ can be identified with the preimage of $\mathcal{M}(A, \|\cdot\|)$ via the restriction $M_L \to M_K$.

1.1.4. Examples of integral structure in Nevanlinna theory. Let R > 0 and let D(R) denote the complex closed disc of radius R. We denote respectively by $A_R = \mathcal{O}(\overline{D(R)})$ and $K_R = \mathcal{M}(\overline{D(R)})$ the ring of germs of holomorphic functions and the field of germs of meromorphic functions on $\overline{D(R)}$. Let $\|\cdot\|_R$ denote the supremum norm on $\overline{D(R)}$ and define $\|\cdot\|_{R,\text{hyb}} := \max\{\|\cdot\|_R, |\cdot|_{\text{triv}}\}$, where $\|\|\cdot\|_{\text{triv}}$ denotes the trivial norm on A_R . Then $(A_R, \|\cdot\|_{R,\text{hyb}})$ is a Banach ring and $(A_R, \|\cdot\|_{R,\text{hyb}})$ defines an integral structure for K([Séd24a], Example 9.2.1 (4)). Moreover, results from (*loc. cit.*, §9.4.3) imply that the integral structure $(A_R, \|\cdot\|_{R,\text{hyb}})$ is tame and the space $V_R := \mathcal{M}(A_R, \|\cdot\|_{R,\text{hyb}})$ has the following description.

Proposition 1.1.5 (*loc. cit.*, Proposition 9.4.6). (i) We have homeomorphisms

$$V_{R,\mathrm{ar}} \cong]0,1] \times \overline{D(R)}, \quad V_{R,\mathrm{um}} \cong \bigsqcup_{z \in \overline{D(R)}} [0,+\infty]/\sim,$$

where \sim denotes the equivalence relation which identifies the extremity 0 of each branch.

(ii) $V_{R,ar}$ is dense in V_R .

1.1.5. *Pseudo-norms.* Let K be a field and $v \in M_K$ be a pseudo-absolute value on K. Let E be a finite-dimensional vector space over K of dimension d. A *pseudo-norm* on E in v is a map $\|\cdot\|_v : E \to [0, +\infty]$ satisfying the following conditions:

- (i) $||0||_v = 0$ and there exists a basis $(e_1, ..., e_d)$ of E such that $||e_1||_v, \cdots, ||e_d||_v \in \mathbb{R}_{>0}$, such a basis is called *adapted* to $||\cdot||_v$;
- (ii) for any $(\lambda, x) \in K \times E$ such that $\{|\lambda|_v, ||x||_v\} \neq \{0, +\infty\}$, we have $\|\lambda x\|_v = |\lambda|_v ||x||_v$; (iii) for any $x, y \in E$, $\|x + y\|_v \le \|x\|_v + \|y\|_v$.

Under these assumptions, $(E, \|\cdot\|_v)$ is called a *pseudo-normed* vector space in v.

Proposition 1.1.6 ([Séd24a], Proposition 6.1.3). Let $(E, \|\cdot\|_v)$ be a pseudo-normed vector space in v. The finiteness module $\mathcal{E}_{\|\cdot\|_v} = \{\|\cdot\|_v < +\infty\}$ is a free A_v -module of rank dgenerated by any basis of E satisfying condition (i) above, the kernel $N_{\|\cdot\|_v} := \{\|\cdot\|_v = 0\}$ is equal to the A_v -submodule $\mathfrak{m}_v \mathcal{E}_{\|\cdot\|_v}$ of the finiteness module. Moreover, $\|\cdot\|_v$ induces a norm on the residue vector space $\widehat{E}_{\|\cdot\|_v} := \mathcal{E}_{\|\cdot\|_v} \otimes_{A_v} \widehat{\kappa_v}$ called the residue norm.

We say that a pseudo-norm $\|\cdot\|_v$ is *ultrametric*, resp. *Hermitian* if so is the residue norm. We also use the notation from ([Séd24a], §6). Namely, by "let $(\|\cdot\|, \mathcal{E}, N, \hat{E})$ be a pseudo-norm on the *K*-vector space *E* in *v*", we mean that $\|\cdot\|$ is a pseudo-norm on *K* in *v* with finiteness module \mathcal{E} , kernel *N* and residue vector space \hat{E} . Moreover, without additional specification, by "let $\|\cdot\|_v$ be a pseudo-norm on *E* in *v*", we mean the pseudo-norm $(\|\cdot\|_v, \mathcal{E}_v, N_v, \hat{E}_v)$. Moreover, if no confusion may arise, we omit "in *v*".

Recall that in (*loc. cit.*, §6.2), we have introduced the usual algebraic constructions for pseudo-normed vector spaces. More precisely, let $(E, \|\cdot\|_v)$ be a pseudo-normed vector space in $v \in M_K$.

- (1) Let F be a non-zero vector subspace of E. Then $\|\cdot\|_v$ induces a pseudo-norm on F denoted again by $\|\cdot\|_v$.
- (2) Let G be non-zero quotient of E. Then $\|\cdot\|_v$ induces a quotient pseudo-norm on G denoted by $\|\cdot\|_{v,G}$.
- (3) $\|\cdot\|_v$ induces a *dual* pseudo-norm on E^{\vee} denoted by $\|\cdot\|_{v,*}$.
- (4) Let $(E', \|\cdot\|'_v)$ be another pseudo-normed vector space. Then this data induces an ϵ -tensor product pseudo-norm on E in. Likewise, we have a π -tensor product pseudo-norm on E.
- (5) Let $i \ge 1$ be an integer. Then $\|\cdot\|_v$ induces the $i^{th}\epsilon$ -exterior power pseudo-norm and $i^{th}\pi$ -exterior power pseudo-norm on $\Lambda^i E$ denoted respectively by $\|\cdot\|_{v,\Lambda_{\epsilon}^i E}$ and $\|\cdot\|_{v,\Lambda_{\pi}^i E}$. In the particular, when i = d, the pseudo-norm $\|\cdot\|_{v,\Lambda_{\pi}^i E}$ is called the determinant pseudo norm on det(E) and is denoted by $\|\cdot\|_{v,\text{det}}$.

We now list generalisations of useful properties of norms in the context of pseudo-norms.

Proposition 1.1.7 ([Séd24a], Propositions 6.2.2-6.2.4). Let $(E, (\|\cdot\|_v, \mathcal{E}_v, N_v, \widehat{E}_v))$ be a pseudo-normed finite-dimensional K-vector space in $v \in M_K$.

- (1) Let G be a quotient of E. Then the dual pseudo-norm $\|\cdot\|_{v,G,*}$ on G^{\vee} identifies with the restriction of the pseudo-norm $\|\cdot\|_{v,*}$ on E^{\vee} to G^{\vee} .
- (2) The inequality

$$\|\cdot\|_{v,**} \le \|\cdot\|_v$$

holds, where $\|\cdot\|_{v,**}$ denotes the dual pseudo-norm of $\|\cdot\|_{v,*}$ on $E^{\vee\vee} \cong E$. Moreover, if either v is Archimedean, or if v is non-Archimedean and the pseudo-norm $\|\cdot\|_v$ is ultrametric, then we have

$$\|\cdot\|_{v,**} = \|\cdot\|_{v}.$$

(3) Let $(e_1, ..., e_r)$ be a basis of E which is adapted to $\|\cdot\|_v$. Then, for any $\eta \in \det(E)$, we have the equality

$$\|\eta\|_{v,\det} = \inf \{\|x_1\|_v \cdots \|x_r\|_v : x_1, ..., x_r \in \mathcal{E}_v \text{ and } \eta = x_1 \wedge \cdots \wedge x_r \}.$$

Definition 1.1.8. Let *E* be a finite-dimensional vector space over *K* and let $\mathbf{e} = (e_1, ..., e_r)$ be any basis of *E*. A pseudo-norm $\|\cdot\|$ on *E* in $v = (|\cdot|, A, \mathfrak{m}, \kappa)$ for which \mathbf{e} is an adapted basis is called *diagonalisable* if there exists a basis $\mathbf{e}' = (e'_1, ..., e'_r)$ of *E* such that the transition

matrix from **e** to \mathbf{e}' belongs to $\operatorname{GL}_r(A)$ and we have

$$\forall a_1, ..., a_r \in K^r, \quad \|a_1 e_1' + ... + a_r e_r'\| = \begin{cases} \sqrt{\sum_{i=1}^r |a_i|^2 \|e_i'\|^2} & \text{if } v \text{ is Archimedean}, \\ \max_{i=1,...,r} |a_i| \|e_i'\| & \text{if } v \text{ is ultrametric.} \end{cases}$$

In that case, we say that \mathbf{e}' is *orthogonal* for $\|\cdot\|$, it is moreover called *orthonormal* if it satisfies the additional condition: $\|e'_i\| = 1$ for all i = 1, ..., r.

1.2. Adelic curves.

1.2.1. Adelic structures over a field. An adelic curve is the data $S = (K, (\Omega, \mathcal{A}, \nu), (|\cdot|_{\omega})_{\omega \in \Omega})$ where K is a field, $(\Omega, \mathcal{A}, \nu)$ is a measure space and $(|\cdot|_{\omega})_{\omega \in \Omega}$ is a family of absolute values on K satisfying the following condition:

$$\forall a \in K^{\times}, \quad (\omega \in \Omega) \mapsto \log |a|_{\omega} \in \mathbb{R}$$

is \mathcal{A} -measurable and ν -integrable. The adelic curve S is called *proper* if the product formula

$$\forall a \in K^{\times}, \quad \int_{\Omega} \log |a|_{\omega} \nu(\mathrm{d}\omega) = 0$$

holds.

Let $S = (K, (\Omega, \mathcal{A}, \nu), (|\cdot|_{\omega})_{\omega \in \Omega})$ and $S' = (K', (\Omega', \mathcal{A}', \nu'), (|\cdot|_{\omega'})_{\omega' \in \Omega'})$ be two adelic curves. A morphism $\alpha : S' \to S$ of adelic curves is a triplet $\alpha = (\alpha^{\sharp}, \alpha_{\sharp}, I_{\alpha})$, where

- $\alpha^{\sharp}: K \to K'$ is a field extension;
- $\alpha_{\sharp}: (\Omega', \mathcal{A}') \to (\Omega, \mathcal{A})$ is a measurable map such that

$$\forall \omega' \in \Omega', \quad \forall a \in K, \quad |\alpha^{\sharp}(a)|_{\omega'} = |a|_{\alpha_{\sharp}(\omega')}.$$

Moreover, the direct image of ν' by α_{\sharp} is assumed to be equal to ν , namely, for any $f \in \mathcal{L}^1(\Omega, \nu)$, we have

$$\int_{\Omega} f \mathrm{d}\nu = \int_{\Omega'} f \circ \alpha_{\sharp} \mathrm{d}\nu';$$

• $I_{\alpha} : \mathcal{L}^1(\Omega', \mathcal{A}', \nu') \to \mathcal{L}^1(\Omega, \mathcal{A}, \nu)$ is a disintegration kernel of α_{\sharp} , namely I_{α} is a linear map such that, for all $g \in \mathcal{L}^1(\Omega', \mathcal{A}', \nu')$, we have

$$\int_{\Omega} I_{\alpha}(g) \mathrm{d}\nu = \int_{\Omega'} g \mathrm{d}\nu',$$

and which, for all $f \in \mathcal{L}^1(\Omega, \mathcal{A}, \nu)$, sends the equivalence class of $f \circ \alpha_{\sharp}$ to the class of f.

1.2.2. Algebraic coverings of adelic curves. Let $S = (K, (\Omega, \mathcal{A}, \nu), (|\cdot|_{\omega})_{\omega \in \Omega})$ be an adelic curve. Let K'/K be an algebraic extension. Then it is possible to define an adelic curve $S \otimes_K K'$ and a morphism $\alpha_{K'/K} : S \otimes_K K' \to S$. Moreover, if S is proper, then $S \otimes_K K'$ is also proper.

1.2.3. *Examples of adelic curves.* On a field, there are many possible adelic structures. We list some examples.

- Any global field (i.e. number field or function field) can be equipped with a natural adelic structure which corresponds to the classical way to perform arithmetic geometry over these fields ([CM19], §3.2.1-3.2.2).
- Given any field K and any measure space $(\Omega, \mathcal{A}, \nu)$, we can consider the proper adelic curve $S = (K, (\Omega, \mathcal{A}, \nu), (|\cdot|_{\mathrm{triv}})_{\omega \in \Omega})$, where the family $(|\cdot|_{\mathrm{triv}})_{\omega \in \Omega}$ consists of copies of the trivial absolute on K. This kind of adelic curve is of particular importance when considering the geometry of numbers and appears naturally when considering Harder-Narasimhan filtrations of adelic vector bundles ([CM24]).
- Function fields of polarised varieties and arithmetic varieties can be endowed with proper adelic structures ([CM19], §3.2.4-3.2.6). This gives a unified approach to higher dimensional of classical arithmetic geometry over global fields.
- Let K be a countable field of characteristic zero. In ([CM21], §2.7), it is proved that there exists an adelic curve $S = (K, (\Omega, \mathcal{A}, \nu), (|\cdot|_{\omega})_{\omega \in \Omega})$ satisfying the following properties:
 - (1) S is proper;
 - (2) for any $\omega \in \Omega$, $|\cdot|_{\omega}$ is a non-trivial absolute value on K;
 - (3) the set $\{\omega \in \Omega : |\cdot|_{\omega} \text{ is non-Archimedean}\}$ is infinite and countable;
 - (4) let \overline{K} be an algebraic closure of K, for any subfield $E_0 \subset \overline{K}$ which is finitely generated over \mathbb{Q} , then the set

$$\{a \in \overline{K} : h_{S \otimes_K \overline{K}}(1, a) \le C \text{ and } [K_0(a) : K_0] \le \delta\}$$

is finite for all $C \in \mathbb{R}_{>0}$ and $\delta \in \mathbb{Z}_{>1}$.

2. TOPOLOGICAL ADELIC CURVES

2.1. Definitions.

Definition 2.1.1. We define the category TAC of topological adelic curves as follows. An object of this category is the data $S = (K, \phi : \Omega \to M_K, \nu)$ where

- K is a field;
- Ω is a locally compact Hausdorff topological space called the *adelic space* of S and $\phi: \Omega \to M_K$ is a continuous map called the *structural morphism* of S;
- ν is a Borel measure on Ω satisfying the following condition: for all $f \in K^{\times}$, the map

$$\begin{array}{cccc} |f|_{\cdot} : & \Omega & \longrightarrow & \mathbb{R}_{\geq 0} \cup \{+\infty\} \\ \omega & \longmapsto & |f|_{\phi(\omega)} \end{array}$$

is such that the function $\log |f|$ is ν -integrable. Note that as $\phi : \Omega \to M_K$ is continuous, by definition of the topology of M_K , for all $f \in K^{\times}$, the map f| is continuous.

Then we define morphisms between topological adelic curves. Let $S_K = (K, \phi_K : \Omega_K \to M_K, \nu_K)$ and $S_L = (L, \phi_L : \Omega_L \to M_L, \nu_L)$ be two topological adelic curves. A morphism $\alpha : S_L \to S_K$ is the data $(\alpha^{\sharp}, \alpha_{\sharp}, I_{\alpha})$ where

- $\alpha^{\sharp}: K \to L$ is a field extension;
- $\alpha_{\sharp}: \Omega_L \to \Omega_K$ is a continuous map inducing a commutative diagram

$$\begin{array}{ccc} \Omega_L & \stackrel{\phi_L}{\longrightarrow} & M_L \\ & \downarrow^{\alpha_{\sharp}} & & \downarrow^{\pi_{L/K}} \\ \Omega_K & \stackrel{\phi_K}{\longrightarrow} & M_K \end{array}$$

Moreover, the direct image of ν_L by α_{\sharp} is assumed to be equal to ν_K . Namely, for any $f \in \mathcal{L}^1(\Omega_K, \nu_K)$, we have

$$\int_{\Omega_K} f \mathrm{d}\nu_K = \int_{\Omega_L} f \circ \alpha_\sharp \mathrm{d}\nu_L;$$

• $I_{\alpha}: L^1(\Omega_L, \nu_L) \to L^1(\Omega_K, \nu_K)$ is a disintegration kernel of α_{\sharp} , namely I_{α} is a linear map such that, for all $g \in L^1(\Omega_L, \nu_L)$, we have

$$\int_{\Omega_K} I_\alpha(g) \mathrm{d}\nu_K = \int_{\Omega_L} g \mathrm{d}\nu_L,$$

and which, for all $f \in L^1(\Omega_K, \nu_K)$, maps the equivalence class of $f \circ \alpha_{\sharp}$ to the class of f.

If $S = (K, \phi : \Omega \to M_K, \nu)$ is a topological adelic curve, for all $f \in K^{\times}$, we define the *defect* $d_S(f)$ by

$$d_S(f) := \int_{\Omega} \log |f|_{\omega} \nu(\mathrm{d}\omega).$$

The topological adelic curve S is called *proper* if, for all $f \in K^*$ we have $d_S(f) = 0$.

Definition 2.1.2. An integral topological adelic curve is a topological adelic curve $(K, \phi : \Omega \to M_K, \nu)$ such that there exists a tame integral structure $(A, \|\cdot\|_A)$ for K (cf. §1.1.3) such that the image of the structural morphism $\phi : \Omega \to M_K$ lies in the global space of pseudo-absolute values $V := \mathcal{M}(A, \|\cdot\|)$. The space V is called the *integral space* of S and the integral structure $(A, \|\cdot\|_A)$ is called the *underlying integral structure* of S.

Let $S_K = (K, \phi_K : \Omega_K \to M_K, \nu_K)$ and $S_L = (L, \phi_L : \Omega_L \to M_L, \nu_L)$ be two integral topological adelic curves with respective integral spaces V_K, V_L . A morphism $\alpha = (\alpha^{\sharp}, \alpha_{\sharp}, I_{\alpha})$: $S_L \to S_K$ of topological adelic curves is called *integral* if there exists a continuous map $\widetilde{\alpha_{\sharp}} : V_L \to V_K$ such that the diagram

$$\begin{array}{ccc} \Omega_L & \stackrel{\phi_L}{\longrightarrow} & V_L & \longrightarrow & M_L \\ & & & & \downarrow \widetilde{\alpha_{\sharp}} & & \downarrow \pi_{L/K} \\ \Omega_K & \stackrel{\phi_K}{\longrightarrow} & V_K & \longrightarrow & M_K \end{array}$$

is commutative.

We define the category ITAC of *integral topological adelic curves* as the subcategory of TAC curves whose objects are integral topological adelic curves and whose morphisms are integral morphisms of topological adelic curves.

- **Notation 2.1.3.** (1) Let $S = (K, \phi : \Omega \to M_K, \nu)$ be a topological adelic curve. Expect mentioned otherwise, for any $\omega \in \Omega$, we denote by A_{ω} the finiteness ring of the pseudo-absolute value $\varphi(\omega)$. Likewise, the kernel, resp. the residue field, resp. the underlying valuation of $\varphi(\omega)$, is denoted by $\mathfrak{m}_{\omega}, \kappa_{\omega}, v_{\omega}$.
 - (2) By "let $S = (K, \phi : \Omega \to V, \nu)$ be an integral topological adelic curve, we mean that $(K, \phi : \Omega, \to M_K, \nu)$ is a topological adelic curve with integral space V.

(3) We can use the notation from §1.1.3 in the context of topological adelic curves. Let $S = (K, \phi : \Omega \to M_K, \nu)$ be a topological adelic curve. Then we define Ω_{ar} , Ω_{um} as the respective preimages of $M_{K,ar}, M_{K,um}$ through the structural morphism $\phi : \Omega \to M_K$. We also have a map $\epsilon : (\omega \in \Omega_{ar}) \mapsto \epsilon(\phi(\omega)) =: \epsilon(\omega) \in]0, 1]$.

Proposition 2.1.4. Let $S = (K, \phi : \Omega \to M_K, \nu)$ be a topological adelic curve.

- (i) For all $f \in K$, let $\Omega_f := \{\omega \in \Omega : f \in A_\omega\}$. Then we have $\nu(\Omega \setminus \Omega_f) = 0$. Moreover, if $f \in K^{\times}$, then $\nu(\{\omega \in \Omega : |f|_{\omega} = 0\}) = 0$.
- (ii) Ω_{ar} , resp. Ω_{um} , is an open, resp. a closed subset of Ω .
- (iii) If ϵ is bounded from below on Ω_{ar} , then $\nu(\Omega_{ar}) < +\infty$ and Ω_{ar} is a closed subset of Ω .

Proof. Let $f \in K$. If f = 0, then $\Omega_f = \Omega$ and thus $\nu(\Omega \setminus \Omega_f) = 0$. Assume that $f \neq 0$. Then $\log |f|$. is ν -integrable and $\Omega_f = \{\omega \in \Omega : \log |f|_{\omega} \neq +\infty\}$. Hence $\nu(\Omega \setminus \Omega_f) = 0$. Finally, as $f \neq 0$, $\log |f^{-1}|$. is ν -integrable and $\{\omega \in \Omega : |f|_{\omega} = 0\} = \Omega \setminus \Omega_{f^{-1}}$. Thus $\nu(\{\omega \in \Omega : |f|_{\omega} = 0\}) = 0$. This concludes the proof of (i).

We now show (*ii*). $\Omega_{ar} = \varphi^{-1}(M_{K,ar})$ is open as φ is continuous. Likewise, Ω_{um} is closed. To show (iii), we may assume that $\Omega_{ar} \neq \emptyset$. Then $\operatorname{char}(K) = 0$ and $\log |2| \in L^1(\Omega, \nu)$. Therefore, the function $f := \max\{0, \log |2|\}$ is ν -integrable. Let 0 < m a lower bound for ϵ on Ω_{ar} . Then

$$m \log(2) \int_{\Omega_{\mathrm{ar}}} \nu(\mathrm{d}\omega) \le \int_{\Omega} f(\omega) \nu(\mathrm{d}\omega) \log < +\infty,$$

hence $\nu(\Omega_{\rm ar}) < +\infty$. Let $(\omega_{\alpha})_{\alpha \in I}$ be a generalised convergent sequence in $\Omega_{\rm ar}$ with limit $\omega \in \Omega$. Then the generalised sequence $(\epsilon(\omega_{\alpha}))_{\alpha \in I} \in [m, 1]^I$ is bounded. By Bolzano-Weierstrass, up to considering a subsequence, we may assume that $(\epsilon(\omega_{\alpha}))_{\alpha \in I}$ converges to some $\epsilon \in [m, 1]$. Then $|2|_{\omega} = \lim_{\alpha \in I} |2|_{\omega_{\alpha}} = \lim_{\alpha \in I} 2^{\epsilon(\omega_{\alpha})} = 2^{\epsilon} > 1$. Therefore, $\omega \in \Omega_{\rm ar}$ and it follows that $\Omega_{\rm ar}$ is closed.

2.2. Examples and constructions.

2.2.1. Number fields. Let K be a number field with ring of integers \mathcal{O}_K . Let Ω_{um} be the set of closed points of $\operatorname{Spec}(\mathcal{O}_K)$ and Ω_{ar} be the set of all fields embeddings $\sigma: K \to \mathbb{C}$, both equipped with the discrete topology. Then $\Omega := \Omega_{um} \sqcup \Omega_{ar}$ is a discrete topological space. For any $\omega \in \Omega_{ar}$, let $\phi(\omega) \in M_K$ denote the Archimedean absolute value on K corresponding to the complex embedding such that $\epsilon(\omega) = 1$. For any $\omega \in \Omega_{um}$ above a prime number p, let $\phi(\omega)$ denote the non-Archimedean extension of the p-adic absolute value such that the absolute value of p equals 1/p. Then the morphism $\phi: \Omega \to M_K$ is continuous. For any $\omega \in \Omega$, denote by K_{ω} , resp. \mathbb{Q}_{ω} the completion of K, resp. \mathbb{Q} , w.r.t. the absolute value $\phi(\omega)$, resp. the restriction of $\phi(\omega)$ to \mathbb{Q} . Now set $\nu(\{\omega\}) := [K_{\omega} : \mathbb{Q}_{\omega}]$. Then the usual product formula implies that $S = (K, \phi: \Omega \to M_K, \nu)$ is a proper topological adelic curve. Moreover, the homeomorphism $M_K \cong \mathcal{M}(\mathcal{O}_K, \max_{v \in \Omega_{ar}}\{|\cdot|_v\})$ from ([Séd24a], Proposition 7.2.1) implies that S is an integral topological adelic curve. Note that the discrete topology is the coarsest topology on Ω making the map $\phi: \Omega \to M_K$ continuous. This follows from the fact that, for any $\omega \in \Omega$, one can exhibit a neighbourhood U_{ω} of $\phi(\omega)$ in Ω such that $\phi^{-1}(U_{\omega}) \cap \Omega = \{\omega\}$.

2.2.2. Nevanlinna theory: analytic functions on a compact disc. We fix R > 0. We define a topological adelic curve $S_R = (K_R, \phi_R : \Omega_R \to M_{K_R}, \nu_R)$, where

- K_R is the field of meromorphic functions on the closed disc $D(R) := \{z \in \mathbb{C} : |z|_{\infty} \le R\} \subset \mathbb{C};$
- $\Omega_R = \{z \in \mathbb{C} : |z|_{\infty} < R\} \coprod \{z \in \mathbb{C} : |z|_{\infty} = R\}$, where $\{z \in \mathbb{C} : |z|_{\infty} < R\}$ is equipped with the discrete topology and $\{z \in \mathbb{C} : |z|_{\infty} = R\}$ is equipped with the usual topology;
- the map $\phi_R : \Omega_R \to M_{K_R}$ is defined by

$$\forall z \in \Omega, \quad \phi_R(z) := \begin{cases} v_{z,\mathrm{ar},1} = \left[(f \in K_R) \mapsto |f(z)|_{\infty} \in [0, +\infty] \right] & \text{if } |z|_{\infty} = R, \\ (f \in K_R) \mapsto e^{-\operatorname{ord}(f,z)} \in \mathbb{R}_{\ge 0} & \text{if } |z|_{\infty} < R; \end{cases}$$

• ν_R is defined by

$$\forall z \in \Omega_{R,\mathrm{um}}, \quad \nu_R(\{z\}) := \begin{cases} \log \frac{R}{|z|_{\infty}} & \text{if } 0 < |z|_{\infty} < R, \\ \log R & \text{if } z = 0, \end{cases}$$

and ν_R is the Lebesgue with total mass 1 on $\Omega_{R,ar}$.

Moreover, results in §1.1.4 imply that S_R is an integral topological adelic curve. Finally, the Jensen formula yields

$$\forall f \in K_R^{\times}, \quad d_R(f) := d_{S_R}(f) = \log |c(f,0)|_{\infty},$$

where c(f, 0) denotes the first non-zero coefficient in the Laurent series expansion of f in 0.

2.2.3. Restriction to a locally closed subset. Let $S = (K, \phi : \Omega \to M_K, \nu)$ be a topological adelic curve. Let $\Omega_0 \subset \Omega$ be a locally closed subset. Then Ω_0 is a locally compact Hausdorff M_K -topological space (cf. [Bou71], Chapitre I, §9.7, Proposition 13). Moreover, if ν_0 denotes the restriction of the measure ν to Ω_0 , then $(K, \phi : \Omega_0 \to M_K, \nu_0)$ is a topological adelic curve called the *restriction* of S to Ω_0 . Note that, in general, if S is proper, S_0 needs not be proper. Likewise, if S is an integral topological adelic curve with integral space V, Ω_0 is a locally compact Hausdorff V-topological space and $(K, \phi_0 : \Omega_0 \to V, \nu_0)$ is an integral topological adelic curve.

3. Families of topological adelic curves

In this section, we introduce to notion of families of topological adelic curves. It should be seen as a preliminary approach to including the analogy between Diophantine approximation and Nevanlinna theory in an Arakelov theoretic framework.

3.1. Definition and examples.

Definition 3.1.1. Let (I, \leq_I) be a poset and let Ω_0 be a locally compact Hausdorff topological space. A collection of topological adelic curves $(S_i = (K_i, \phi_i : \Omega_i \to M_{K_i}, \nu_i))_{i \in I}$ is called a *family of topological adelic curves* if the following conditions hold:

- (1) for all $i, i' \in I$, we have $\operatorname{char}(K_i) = \operatorname{char}(K_{i'})$;
- (2) $(K_i)_{i \in I}$ defines an inverse system in the category of fields, namely, for any $i, i' \in I$ such that $i \leq_I i'$, we have a field inclusion $K_{i'} \subset K_i$;

(3) there exists a family of continuous maps

$$\alpha_i: \Omega_i \to \Omega_0, \quad i \in I,$$

which satisfies the following condition:

$$\forall i, i' \in I, \quad \forall (\omega, \omega') \in \Omega_i \times \Omega_{i'}, \quad \alpha_i(\omega) = \alpha_{i'}(\omega') \Rightarrow |\cdot|_{\omega'|K_i \cap K_{i'}} = |\cdot|_{\omega|K_i \cap K_{i'}}.$$

In that case, the topological space Ω_0 is called the *base space* of the family $(S_i)_{i \in I}$ and the family $(\alpha_i)_{i \in I}$ is called the *family of structural morphisms* of the family $(S_i)_{i \in I}$.

- **Example 3.1.2.** (1) Let $S = (K, \phi : \Omega \to M_K, \nu)$ be a topological adelic curve. Let I be any set endowed with the trivial ordering. We consider the collection $(S)_{i \in \mathbb{N}}$. By considering the base space Ω and the family of structural morphisms consisting of the identity $\mathrm{Id} : \Omega \to \Omega$, we obtain a family of topological adelic curves. The case $I = \mathbb{N}$ should be seen as the counterpart of classical Diophantine approximation in our framework.
 - (2) We equip $\mathbb{R}_{>0}$ with the usual ordering. Consider the collection $(S_R)_{R>0}$, where for any R > 0, $S_R = (K_R, \phi_R : \Omega_R \to M_{K_R}, \nu_R)$ is the topological adelic curve defined in §2.2.2. Consider the topological space $\Omega_0 := \mathbb{C} \bigsqcup (\mathbb{C} \smallsetminus \{0\})$, where \mathbb{C} is equipped with the discrete topology and $\mathbb{C} \smallsetminus \{0\}$ is equipped with the usual topology. For any R > 0, let

$$\begin{array}{cccc} \alpha_R : & \Omega_R & \longrightarrow & \Omega_0 \\ & & & \\ \omega & \longmapsto & \begin{cases} \omega \in \mathbb{C} & \text{ if } \omega \in \Omega_{R, \text{um}}, \\ \omega \in \mathbb{C} \smallsetminus \{0\} & \text{ if } \omega \in \Omega_{R, \text{ar}}. \end{cases} \end{array}$$

Then, for any R > 0, α_R is a continuous injective map and $(S_R)_{R>0}$ is a family of topological adelic curves with base space Ω_0 and family of structural morphisms $(\alpha_R : \Omega_R \to \Omega_0)_{R>0}$.

3.2. Asymptotically proper family of topological adelic curves.

Definition 3.2.1. Let (I, \leq_I) be a poset. Let $\mathbf{S} = (S_i = (K_i, \phi_i : \Omega_i \to M_{K_i}, \nu_i))_{i \in I}$ be a family of topological adelic curves with base space Ω_0 and family of structural morphisms $(\alpha_i : \Omega_i \to \Omega_0)_{i \in I}$. Denote

$$K_{\infty} := \varprojlim_{i \in I} K_i.$$

For any $f \in K_{\infty}^{\times}$, define the function

$$d_{\cdot}(f): \begin{vmatrix} I & \longrightarrow & \mathbb{R} \\ i & \longmapsto & d_{S_i}(f) \end{vmatrix}$$

This function is called the *defect* of f w.r.t. the family **S**.

Let ~ be an equivalence relation on $\mathcal{F}(I,\mathbb{R})$ which is compatible with the additive group structure. We say that the family **S** is *asymptotically proper* w.r.t. ~ if

$$\forall f \in K_{\infty}^{\times}, \quad d_{\cdot}(f) \sim 0.$$

Example 3.2.2. (1) Let $S = (K, \phi : \Omega \to M_K, \nu)$ be a topological adelic curve. Let $I = \mathbb{N}$ endowed with the trivial ordering. Consider the family **S** of topological adelic curves defined in Example 3.1.2 (1). Let ~ be the equivalence relation defined by

$$\forall f, g \in \mathcal{F}(I, \mathbb{R}), \quad f \sim g \Leftrightarrow f - g = O(1).$$

Then the family **S** is asymptotically proper w.r.t. ~ iff S is proper. Indeed if S is not proper, then there exists $f \in K$ such that $d_S(f) \neq 0$. Then $\mathbf{f} := (f, f^2, f^3, ...) \in K_{\infty}$ satisfies $|d_i(\mathbf{f})| \to_{i \to +\infty} +\infty$.

(2) We equip $\mathbb{R}_{>0}$ with the usual ordering. Consider the collection $\mathbf{S} = (S_R)_{R>0}$ constructed in Example 3.1.2 (2). Let $\sim_{O(1)}$ be the equivalence relation on $\mathcal{F}(I,\mathbb{R})$ defined by

$$\forall f, g \in \mathcal{F}(I, \mathbb{R}), \quad f \sim_{O(1)} g \Leftrightarrow f - g = O(1).$$

Then K_{∞} identifies with the field of meromorphic functions on \mathbb{C} and, for any $f \in K_{\infty}$, for any R > 0, we have

$$d_R(f) = \log |c(f,0)|_{\infty} = O(1).$$

Hence the family **S** is asymptotically proper w.r.t. $\sim_{O(1)}$.

Note that we could also have chosen the equivalence relation $\sim_{\rm cst}$ defined by

$$\forall f,g \in \mathcal{F}(I,\mathbb{R}), \quad f \sim_{\mathrm{cst}} g \Leftrightarrow (\exists a \in \mathbb{R}, \quad f-g=a).$$

Then **S** is asymptotically proper w.r.t. \sim_{cst} .

Furthermore, let $h : \mathbb{R}_{>0} \to \mathbb{R}_{>0}$ be any function such that $h(R) \to_{R \to +\infty} +\infty$. For any R > 0, define another measure ν'_R on Ω_R by

$$\forall z \in \Omega_{R,\mathrm{um}}, \quad \nu_R'(\{z\}) := \begin{cases} \frac{1}{h(R)} \log \frac{R}{|z|_{\infty}} & \text{if } 0 < |z|_{\infty} < R, \\ \frac{1}{h(R)} \log R & \text{if } z = 0, \end{cases}$$

and ν'_R is the Lebesgue with total mass 1/h(R) on $\Omega_{R,ar}$. This construction yields a family of topological adelic curves $\mathbf{S}' := (S'_R := (K_R, \phi_R : \Omega_R \to M_{K_R}, \nu'_R))_{R>0}$ which is asymptotically proper w.r.t. the equivalence relation $\sim_{o(1)}$ defined by

 $\forall f,g \in \mathcal{F}(I,\mathbb{R}), \quad f \sim_{o(1)} g \Leftrightarrow f - g = o(1).$

Indeed, for any $f \in K_{\infty}^{\times}$, we have

$$d_{S'_R}(f) = \frac{\log |c(f,0)|_{\infty}}{h(R)} \to_{R \to +\infty} 0.$$

3.3. Adelic space attached to a family of topological adelic curves.

Definition 3.3.1. Let (I, \leq_I) be a poset equipped with the order topology. Let $\mathbf{S} = (S_i = (K_i, \phi_i : \Omega_i \to M_{K_i}, \nu_i))_{i \in I}$ be a family of topological adelic curves with base space Ω_0 and family of structural morphisms $(\alpha_i : \Omega_i \to \Omega_0)_{i \in I}$. Assume that there exists a family $\beta = (\beta_j)_{j \in J}$, where J is an arbitrary index set and, for any $j \in J$, $\beta_j : I \to \bigsqcup_{i \in I} \Omega_i$ is a map, such that

$$\forall i \in I, \quad \beta_j(i) \in \Omega_i,$$

and $\alpha_i \circ \beta_j : I \to \Omega_0$ is continuous.

We define the *adelic space* attached to the families \mathbf{S}, β as the set

$$\Omega_{\infty} := \left(\bigsqcup_{i \in I} \Omega_i\right) / \sim,$$

where \sim denotes the equivalence relation defined by

$$\forall (i,\omega), (i',\omega') \in \bigsqcup_{i \in I} \Omega_i, \quad (i,\omega) \sim (i',\omega') \Leftrightarrow \alpha_i(\omega) = \alpha_{i'}(\omega').$$

The set Ω_{∞} is equipped with the finest topology making the maps

$$\begin{split} p &: \bigsqcup_{i \in I} \Omega_i \to \Omega_{\infty}, \\ \overline{\beta_j} &:= p \circ \beta_j : I \to \Omega_{\infty}, \quad j \in J, \end{split}$$

continuous.

Remark 3.3.2. Using the same notation as above, we see that there exists a canonical injective continuous map $\Omega_{\infty} \to \Omega_0$ induced by the α_i 's. Therefore, up to shrinking the codomain of the α_i 's, it makes sense to assume that Ω_{∞} and Ω_0 are homeomorphic.

- **Example 3.3.3.** (1) Let $S = (K, \phi : \Omega \to M_K, \nu)$ be a topological adelic curve. Let I be any set endowed with the trivial ordering. Consider the family **S** of topological adelic curves defined in Example 3.1.2 (1). Let $J = \emptyset$. Then the adelic space attached to **S** can be identified with Ω .
 - (2) We equip $\mathbb{R}_{>0}$ with the usual ordering. Consider the collection $\mathbf{S} = (S_R)_{R>0}$ constructed in Example 3.1.2 (2). Consider the family $\beta = (\beta_z)_{z \in \mathbb{S}^1}$, where, for any $z \in \mathbb{S}^1$, for any R > 0, $\beta_z(R) = Rz \in \Omega_{R,ar}$. Then the adelic space attached to \mathbf{S} and β is homeomorphic to Ω_0 .

4. Algebraic covering of topological adelic curves

4.1. Finite separable extension. Throughout this subsection, we fix a topological adelic curve $S_K = (K, \phi_K : \Omega_K \to M_K, \nu_K)$ and a finite separable extension L/K. We will construct a canonically determined adelic structure on L whose corresponding adelic curve maps to S.

4.1.1. Case of a general topological adelic curve. Let $\Omega_L := \Omega_K \times_{V_K} V_L$ equipped with the fibre product topology. Denote by $\phi_L : \Omega_L \to M_L$ the pullback of ϕ_K . Then Ω_L is a locally compact Hausdorff space. By definition of the fibre product topology, for any $f \in L^{\times}$, the map $\omega \in \Omega_L \mapsto \log |f|_{\omega} \in [-\infty, +\infty]$ is continuous.

Proposition 4.1.1. The restriction morphism $\pi_{L/K} : M_L \to M_K$ is surjective, proper, with finite fibres.

Proof. $\pi_{L/K}$ is proper since it is continuous and both M_K and M_L are compact Hausdorff. $\pi_{L/K}$ is surjective by construction and has finite fibres by Proposition 1.1.1.

We now adapt the constructions in ([CM19], §3.3) to define a measure ν_L on Ω_L . For any $\omega \in \Omega_K$, let $\Omega_{L,\omega}$ denote the fibre of ω through the restriction map $\pi_{L/K} : \Omega_L \to \Omega_K$. Let $\omega \in \Omega$. Denote by I_{ω} the cardinality of the set of valuation rings on L extending A_{ω} .

Proposition 4.1.2. Using the same notation as in Proposition 1.1.1, the measure $\Omega_{L,\omega}$ defined by

$$\forall x \in \Omega_{L,\omega}, \quad \mathbb{P}_{L,\omega}(\{x\}) := \frac{1}{|I_{\omega}|} \frac{[\widehat{\kappa_x} : \widehat{\kappa_{\omega}}]}{[\kappa_x : \kappa_{\omega}]},$$

is a probability measure.

Proof. It is a direct consequence of (1).

Definition 4.1.3. Let $f: \Omega_L \to [-\infty, +\infty]$. Define $I_{L/K}(f): \Omega_K \to [-\infty, +\infty]$ by

$$I_{L/K}(f)(\omega) := \sum_{x \in \pi_{L/K}^{-1}(\omega)} \mathbb{P}_{L,\omega}(\{x\})f(x).$$

The σ -algebra defined in ([CM19], §3.3) coincides with the Borel σ -algebra on Ω_L . Therefore, we can reproduce *mutatis mutandis* the arguments of *ibid*. to obtain a measure ν_L such that the equality

$$\nu_L(A) = \int_{\Omega_K} \sum_{x \in \Omega_{L,\omega}} \mathbb{P}_{L,\omega}(\{x\}) \mathbf{1}_A(x) \mathrm{d}\nu_K, \tag{2}$$

holds for any Borel subset $A \subset \Omega_L$, where $\mathbf{1}_A$ denotes the characteristic function of A. Furthermore, ν_L satisfies all the conditions of ([CM19], Theorem 3.3.7). Thus we deduce the following proposition.

Proposition 4.1.4. We use the above notation. Then $S_L = S \otimes_K L := (L, \phi_L : \Omega_L \to M_L, \nu_L)$ is a topological adelic curve. We further have a morphism $\alpha_{L/K} : S_L \to S_K$. Moreover, for any $f \in L^{\times}$, we have

$$d_{S_L}(f) = \frac{1}{[L:K]} d_{S_K}(N_{L/K}(f)).$$
(3)

In particular, if S_K is proper, then S_L is proper.

Proof. We only need to justify that we have a morphism of topological adelic curves $\alpha_{L/K}$: $S_L \to S_K$. Denote by $\alpha_{L/K}^{\sharp}: K \to L$ the field extension. Define $\alpha_{L/K,\sharp}: \Omega_L \to \Omega_K$ as the pullback of $\pi_{L/K}: M_L \to M_K$ by $\varphi_K: \Omega_K \to M_K$. By construction, the diagram in Definition 2.1.1 is commutative. Moreover, ([CM19], Theorem 3.3.7 (3)) implies that $\alpha_{L/K,\sharp,*}\nu_L = \nu_K$. Finally, from ([CM19], Theorem 3.3.7 (1-2)), we get that $I_{L/K}$ (cf. Definition 4.1.3) is a disintegration kernel of $\alpha_{L/K,\sharp}$.

4.1.2. Case of an integral topological adelic curve. Assume that S_K is integral. Let $(A, \|\cdot\|_A)$ be the underlying integral structure and let $V_K = \mathcal{M}(A, \|\cdot\|_A)$ be the integral space. Denote by B the integral closure of A in L. As recalled in §1.1.3, there exists a norm $\|\cdot\|_B$ on B such that $(B, \|\cdot\|_B)$ is a tame integral structure for L and $V_L := \mathcal{M}(B, \|\cdot\|_B) = \pi_{L/K}^{-1}(\mathcal{M}(A, \|\cdot\|_A))$. This is a compact Hausdorff topological space and we denote by $\pi_{L/K} : V_L \to V_K$ the (continuous) restriction morphism. Now the following proposition implies that the image of structural morphism $\phi_L : \Omega_L \to M_L$ lies in V_L and $S \otimes_K L = (L, \phi_L : \Omega_L \to V_L, \nu_L)$ is an integral topological adelic curve.

Proposition 4.1.5. The restriction morphism $\pi_{L/K} : V_L \to V_K$ is surjective, proper, with finite fibres.

Proof. $\pi_{L/K}$ is proper since it is continuous and both V_L and V_K are compact Hausdorff. $\pi_{L/K}$ is surjective by construction and has finite fibres by Proposition 4.1.1.

4.2. Finite extension. Throughout this subsection, we fix a topological adelic curve $S_K = (K, \phi_K : \Omega_K \to M_K, \nu_K)$ and a finite extension L/K. Denote by K'/K the separable closure of K in L and let $S_{K'} := S \otimes_K K' := (K', \phi_{K'} : \Omega_{K'} \to M_{K'}, \nu_{K'})$ be the topological adelic curve constructed in §4.1.

Proposition 4.2.1. (1) The spaces M_L et $M_{K'}$ are homeomorphic.

- (2) There exist a topological adelic curve $S_L := (L, \phi_L : \Omega_L \to M_L, \nu_L)$ and a morphism $S_L \to S_{K'} \to S$ of topological adelic curves.
- (3) Assume that S_K is an integral topological adelic curve and denote by (A, || · ||_A) its underlying integral structure and by V_K its integral space. Let A' and A_L denote respectively the integral closures of A in K' and L. We respectively endow A' and A_L with the norm || · ||_{A'} and || · ||_{A_L} from Proposition 1.1.4. Then V_L := M(A_L) and V_{K'} = M(A') are homeomorphic and the topological adelic curve S_L is integral.

Proof. (1) Corollary 1.1.2 provides a continuous bijection $\pi_{L/K'}: M_L \to M_{K'}$. Since both M_L and $M_{K'}$ are compact Hausdorff, we obtain (1).

(2) The homeomorphism $\pi_{L/K'}: M_L \to M_{K'}$ induces by pullback a homeomorphism $\Omega_L := \pi_{L/K'}^{-1}(\Omega_{K'}) \to \Omega_{K'}$ and a continuous map $\phi_L: \Omega_L \to M_L$. Denote by ν_L the pushforward of $\nu_{K'}$ by the inverse of the latter homeomorphism. Then $S_L := (L, \phi_L: \Omega_L \to M_L, \nu_L)$ is a topological adelic curve.

(3) $(A_L, \|\cdot\|_{A_L})$ is an integral structure for L (cf. Proposition 1.1.4). Moreover, the restriction map $\pi_{L/K'} : \mathcal{M}(A_L) \to \mathcal{M}(A')$ is a restriction of the homeomorphism of (1) which is surjective. Hence it is a homeomorphism. The last part of the assertion follows directly. \Box

We now state a compatibility result for successive extensions of the base field.

Proposition 4.2.2. Let $K_2/K_1/K$ be successive finite extensions. Then we have a canonical isomorphism of topological adelic curves

$$(S_K \otimes_K K_1) \otimes_{K_1} K_2 \cong S_K \otimes_K K_2$$

making the below diagram commute.



Moreover, if S_K is integral, the above isomorphism and diagram lie in the category of integral topological adelic curves.

Proof. Let $(S_K \otimes_K K_1) \otimes_{K_1} K_2 = (K_2, \phi'_2 : \Omega'_2 \to M_{K_2}, \nu'_2)$ and $S_K \otimes_K K_2 = (K_2, \phi_2 : \Omega_2 \to M_{K_2}, \nu_2)$. The definition of Ω_2, Ω'_2 implies that there exists a canonical homeomorphism $\Omega_2 \cong \Omega'_2$. From ([CM19], (3.15)), we deduce that ν_2 can be identified with ν'_2 via the previous homeomorphism. Commutativity of the diagram follows from Proposition 4.2.1 (3).

In the integral case, we denote respectively by A_1 and A_2 the integral closures of A in K_1 and K_2 . Note that the construction of the extension of the norm $\|\cdot\|_A$ from Proposition 1.1.4 is compatible with successive extension. Since V'_2 is obtained by considering the integral closure of A_1 in L, which is equal to A_2 , we obtain $V_2 = V'_2$.

4.3. Algebraic extension. Throughout this subsection, we fix a topological adelic curve $S_K = (K, \phi_K : \Omega_K \to M_K, \nu_K)$ and an algebraic extension L/K. Let $\mathcal{E}_{L/K}$ be the set of all sub-extensions L/K'/K with K'/K finite. It is a directed set with respect to the inclusion relation and we have

$$L = \bigcup_{K'/K \in \mathcal{E}_{L/K}} K'.$$

Then Proposition 4.2.2 ensures that we have a cofiltered inverse system of topological adelic curves $(S_{K'})_{K' \in \mathcal{E}_{L/K}}$ whose arrows are denoted by $\pi_{K''/K'} : S_{K''} \to S_{K'}$. We will prove that the inverse limit of this inverse system exists and is a topological adelic curve with field L.

Proposition 4.2.2 yields an inverse system $(S_{K'} = (K', \phi_{K'} : \Omega_{K'} \to M_{K'}, \nu_{K'}))_{K' \in \mathcal{E}_{L/K}}$ in TAC which induces an inverse system of Cartesian diagrams of the form

$$\begin{array}{ccc} \Omega_{K'} & \stackrel{\phi_{K'}}{\longrightarrow} & M_{K'} \\ & & & \downarrow^{\pi_{K'/K}} \\ \Omega_K & \stackrel{\phi_K}{\longrightarrow} & M_K \end{array}$$

indexed by $\mathcal{E}_{L/K}$ and whose vertical arrows are proper. Hence its inverse limit can be written as

$$\begin{array}{ccc} \Omega_L & \stackrel{\phi_L}{\longrightarrow} & M_L \\ \downarrow & & \downarrow^{\pi_{L/K}} , \\ \Omega_K & \stackrel{\phi_K}{\longrightarrow} & M_K \end{array}$$

where $\Omega_L := \varprojlim_{K' \in \mathcal{E}_{L/K}} \Omega_{K'}$. Since the map $\pi_{L/K}$ is proper and the $\Omega_{K'}$ are locally compact Hausdorff topological spaces, Ω_L is locally compact Hausdorff.

We can now adapt the arguments from ([CM19], §3.4) to construct a Borel measure ν_L on Ω_L and a disintegration kernel for $\pi_{L/K} : \Omega_L \to \Omega_K$.

Hence we obtain the following proposition.

Proposition 4.3.1. We use the same notation as above. Then $S_L := S_K \otimes_K L := (L, \phi_L : \Omega_L \to M_L, \nu_L)$ is a topological adelic curve and we have an isomorphism

$$\lim_{K'\in\mathcal{E}_{L/K}}S_K\otimes_K K'.$$

Moreover, if S_K is proper, then S_L is proper. Finally, if S_K is integral, then S_L is integral.

Proof. Except for the assertion about integrality, the proposition follows from the above paragraph. We now assume that S_K is integral. Let $(A, \|\cdot\|_A)$ denote its underlying integral structure and let V_K be its integral space. Let B denote the integral closure of A in L. Then Proposition 1.1.4 and its proof yield a norm $\|\cdot\|_B$ as well as an isomorphism

$$V_L := \mathcal{M}(B, \|\cdot\|_B) \cong \lim_{K' \in \mathcal{E}_{L/K}} \mathcal{M}(A_{K'}, \|\cdot\|_{A_{K'}}) =: V_{K'},$$

where, for all $K' \in \mathcal{E}_{L/K}$, $(A_{K'}, \|\cdot\|_{A_{K'}})$ denotes the extension of $(A, \|\cdot\|_A)$ over K'. Therefore, we see that the image of the structural morphism $\phi_L : \Omega_L \to M_L$ lies in V_L and S_L is integral.

4.4. Action of the Galois group. Throughout this subsection, we fix a topological adelic curve $S = (K, \phi : \Omega \to M_K, \nu)$ and an algebraic extension L/K whose group of K-linear automorphisms is denoted by $\operatorname{Aut}(L/K)$. Let $S_L := (L, \phi_L : \Omega_L \to M_L, \nu_L)$ be the topological adelic curve $S \otimes_K L$ defined in §4.3.

Proposition 4.4.1. (i) The action of $\operatorname{Aut}(L/K)$ on M_L introduced in Proposition 1.1.3 (1) induces continuous and proper actions of $\operatorname{Aut}(L/K)$ on M_L and Ω_L . Moreover,

if S is integral and V_L denotes the integral space of S_L , the action of $\operatorname{Aut}(L/K)$ on M_L induces a continuous and proper action on V_L .

- (ii) We assume that L/K is Galois and that, for any $v \in M_K$, the residue field κ_v is perfect. Then for any $\omega \in \Omega$, the actions of $\operatorname{Aut}(L/K)$ on $\Omega_{L,\omega}$ and $M_{L,\phi(\omega)}$ are transitive. Moreover, if S is integral with underlying global space of pseudo-absolute values V. Then for any $v \in V$, $\operatorname{Aut}(L/K)$ acts transitively on $V_{L,v}$, where V_L denotes the underlying global space of pseudo-absolute values of S_L .
- (iii) We use the same assumptions as in (ii). Then we have homeomorphisms

 $\Omega_L / \operatorname{Aut}(L/K) \cong \Omega, \quad M_L/G \cong M_K.$

Moreover, if S is integral, we have a homeomorphism

$$V_L / \operatorname{Aut}(L/K) \cong V.$$

Proof. By considering the trivial actions on V and Ω , we see that $\operatorname{Aut}(L/K)$ induces actions on Ω_L and V_L . Let us show that these actions are continuous. First, assume that L/K is finite. Since $\operatorname{Aut}(L/K)$ is discrete, it is enough to prove that, for any $\tau \in \operatorname{Aut}(L/K)$, for any $a \in L$, the map $(|\cdot|_x \in M_L) \mapsto |a|_{\tau(x)}$ is continuous. This is clear by definition of the topology on M_L and since $|a|_{\tau(x)} = |\tau(a)|_x$. If now L/K is infinite, let us show that the map

$$\alpha_{L/K} : \left| \begin{array}{ccc} \operatorname{Aut}(L/K) \times M_L & \longrightarrow & M_L \\ (\tau, v) & \longmapsto & v \circ \tau \end{array} \right|$$

is continuous. By definition of the topology on M_L , it is enough to prove that for any intermediate extension L/K'/K with K'/K finite, the map $\alpha_{L/K} \circ \pi_{L/K'}$ is continuous. Let K' be such an intermediate extension. By definition of the topology on $\operatorname{Aut}(L/K)$ and M_L , the map $\beta : (\tau, v) \in \operatorname{Aut}(L/K) \times M_L \mapsto (\tau_{|K'}, v_{|K'}) \in \operatorname{Aut}(K'/K) \times V_{K'}$ is continuous. The finite case implies that $\alpha_{K'/K}$ is continuous. We conclude by using the fact that $\alpha_{L/K} \circ \pi_{L/K'} = \alpha_{K'/K} \circ \beta$. Since $\phi_L : \Omega_L \to M_L$ is continuous, we obtain the continuity of the action on Ω_L . These actions are proper since M_L and Ω_L are locally compact Hausdorff and $\operatorname{Aut}(L/K)$ is Hausdorff. In the case where S is integral, the action of $\operatorname{Aut}(L/K)$ on M_L induces a continuous and proper action on V_L as V_L is a compact subset of M_L which is stabilised by the action. This concludes the proof of (i).

(ii) is a direct consequence of Proposition 1.1.3 (2).

We now prove (iii). Denote $G := \operatorname{Aut}(L/K)$. Since $K = L^G$, the restriction maps $\pi_{L/K} : V_L \to V$ and $\pi_{L/K} : \Omega_L \to \Omega$ induce continuous maps $M_L/G \to M_K$ and $\Omega_L/G \to \Omega$ such that the diagram

$$\begin{array}{ccc} \Omega_L & \stackrel{\phi_L}{\longrightarrow} & M_L \\ \downarrow & & \downarrow \\ \Omega_L/G & \longrightarrow & M_L/G \\ \downarrow & & \downarrow \\ \Omega & \stackrel{\phi}{\longrightarrow} & M_K \end{array}$$

is Cartesian. (ii) ensures that the arrows $\Omega_L/G \to \Omega$ and $M_L/G \to M_K$ are injective, hence bijective. Since both M_L and M_K are compact Hausdorff, the arrow $M_L/G \to M_K$ is a homeomorphism. Therefore, the arrow $\Omega_L/G \to \Omega$ is a homeomorphism as it is the pullback of $M_L/G \to M_K$ by $\phi : \Omega \to M_K$. In case where S is integral we use that the map $V_L/G \to V$ is the pullback of $M_L/G \to M_K$ by the natural inclusion. Hence it is a homeomorphism. This concludes the proof of the proposition.

Remark 4.4.2. We assume that the hypotheses of Proposition 1.1.3 (ii) hold. Using ([BYDHS24], Lemma 10.4), if the measure ν is assumed to be Radon, we see that the measure ν_L is the only Aut(L/K)-invariant measure on Ω_L whose pushforward via $\pi_{L/K}$ is ν .

Part 2. Adelic vector bundles and Harder-Narasimhan filtrations over topological adelic curves

In this part, we study the intrinsic geometry of topological adelic curves. This is done by introducing what plays the role of a vector bundle on a curve in algebraic geometry. We introduce the counterpart of norm families in our context and introduce various regularity and dominance conditions (§5). After that, we define adelic vector bundles on a topological adelic curve (§6). Finally, we study slope theory for adelic vector bundles on a topological adelic curve (§7).

5. Pseudo-norm families

In this section, we globalise the constructions of §1.1.5. Throughout this section, we fix a topological adelic curve $S = (K, \phi : \Omega \to M_K, \nu)$. Recall that, for any $\omega \in \Omega$, we denote by A_{ω} and κ_{ω} , the finiteness ring and the residue field of ω respectively. From now on, we assume that for any $\omega \in \Omega_{\rm ar}$, we have $\epsilon(\omega) = 1$. In that case Proposition 2.1.4 implies that $\nu(\Omega_{\rm ar}) < +\infty$ and $\Omega_{\rm ar}, \Omega_{\rm um}$ are open subsets of Ω . Note that this assumption is not too harmful to the generality since we can replace, for any $\omega \in \Omega_{\rm ar}$, the pseudo-absolute value $|\cdot|_{\omega}$ by the pseudo-absolute value $|\cdot|_{\omega}^{1/\epsilon(\omega)}$ and the measure ν by the measure $\tilde{\nu} := (\mathbf{1}_{\Omega_{\rm um}} + \epsilon \mathbf{1}_{\Omega, {\rm ar}})\nu$.

5.1. **Definitions.** Let *E* be a finite-dimensional *K*-vector space. For any $\omega \in \Omega$, we call *pseudo-norm* in ω on *E* any map $\|\cdot\|_{\omega} : E \to [0, +\infty]$ such that $\|\cdot\|_{\omega}$ is a pseudo-norm in $\varphi(\omega)$ on *E* (cf. §1.1.5).

Definition 5.1.1. Let *E* be a finite-dimensional *K*-vector space. We call *pseudo-norm* family on *E* any family $\xi = (\|\cdot\|_{\omega})_{\omega \in \Omega}$ where, for any $\omega \in \Omega$, $\|\cdot\|_{\omega}$ is a pseudo-norm in ω on *E*. We assume that the additional following condition holds:

(*) for any $\omega \in \Omega$, there exist an open neighbourhood U of ω in Ω and a basis $(e_1, ..., e_d)$ of E such that, for any $i \in \{1, ..., d\}$, $||e_i||_{\omega} \in \mathbb{R}_{>0}$. Such a basis is called *adapted* to the pseudo-norm family ξ in ω .

Moreover, if there exists a basis $(e_1, ..., e_d)$ of E such that, for any $\omega \in \Omega$, for all i = 1, ..., r, we have $||e_i||_{\omega} \in \mathbb{R}_{>0}$, we say that $(e_1, ..., e_d)$ is globally adapted to ξ (on U). Finally, we say that the pseudo-norm family ξ is ultrametric, resp. Hermitian, if $|| \cdot ||_{\omega}$ is ultrametric for any $\omega \in \Omega_{\text{um}}$, resp. if $|| \cdot ||_{\omega}$ is Hermitian for any $\omega \in \Omega_{\text{ar}}$.

Notation 5.1.2. Let E be a finite-dimensional K-vector space.

(1) By "let $\xi = (\|\cdot\|_{\omega}, \mathcal{E}_{\omega}, N_{\omega}, \widehat{E}_{\omega})_{\omega \in \Omega}$ be a pseudo-norm family on E", we mean that, for any $\omega \in \Omega$, the pseudo-norm $\|\cdot\|_{\omega}$ has finiteness module \mathcal{E}_{ω} , kernel N_{ω} and residue vector space \widehat{E}_{ω} .

- (2) In case there is no explicit notation as above, for any pseudo-norm family $\xi = (\|\cdot\|_{\omega})_{\omega\in\Omega}$ on E, for any $\omega\in\Omega$, we denote by
 - \mathcal{E}_{ω} the finiteness module of $\|\cdot\|_{\omega}$;
 - N_{ω} the kernel of $\|\cdot\|_{\omega}$;
 - $-\widehat{E_{\omega}}$ the residue vector space of $\|\cdot\|_{\omega}$.

Example 5.1.3. Let *E* be a finite-dimensional *K*-vector space and fix a basis $\mathbf{e} = (e_1, ..., e_r)$ of *E*. We fix $\omega \in \Omega$ and we denote respectively by \mathcal{E}_{ω} and \widehat{E}_{ω} the restriction of scalars of *E* to A_{ω} and the corresponding residue vector space. Then \mathbf{e} defines compatible isomorphisms

$$\begin{array}{cccc} E & & & & & & & & \\ & \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\ K^n & & & & & & & \\ K^n & & & & & & & & \\ \hline \end{array}$$

For any $\lambda_1, ..., \lambda_r \in \widehat{\kappa_{\omega}}$, we set

$$\|\lambda_1 e_1 + \dots + \lambda_r e_r\|_{\mathbf{e},\omega} = \begin{cases} \max\{|\widetilde{\lambda_1}|_{\omega}, \dots, |\widetilde{\lambda_d}|_{\omega}\}, & \text{if } \omega \in \Omega_{\mathrm{um}}, \\ |\widetilde{\lambda_1}|_{\omega} + \dots + |\widetilde{\lambda_d}|_{\omega}, & \text{if } \omega \in \Omega_{\mathrm{ar}}, \end{cases}$$

where $|\cdot|$ denotes the residue absolute value on $\widehat{\kappa_{\omega}}$. Then $\|\cdot\|_{\mathbf{e},\omega}$ defines a norm on $\widehat{E_{\omega}}$. By lifting $\|\cdot\|_{\mathbf{e},\omega}$ to a pseudo-norm on E in ω , we obtain a pseudo-norm family $\xi_{\mathbf{e}} = (\xi_{\mathbf{e},\omega})_{\omega\in\Omega}$ on E called the *model pseudo-norm family* associated with the basis \mathbf{e} .

We also define a Hermitian pseudo-norm family $\xi_{\mathbf{e},2}$ as follows. For any $\lambda_1, ..., \lambda_r \in \widehat{\kappa_{\omega}}$, we set

$$\|\lambda_1 e_1 + \dots + \lambda_r e_r\|_{\mathbf{e},2,\omega} = \begin{cases} \max\{|\widetilde{\lambda_1}|_{\omega}, \dots, |\widetilde{\lambda_d}|_{\omega}\}, & \text{if } \omega \in \Omega_{\mathrm{um}}, \\ (|\widetilde{\lambda_1}|_{\omega}^2 + \dots + |\widetilde{\lambda_d}|_{\omega}^2)^{1/2}, & \text{if } \omega \in \Omega_{\mathrm{ar}}. \end{cases}$$

By the same arguments as above, we can lift the construction to obtain a pseudo-norm family $\xi_{\mathbf{e},2}$ on E. Note that \mathbf{e} is a basis of E which is globally adapted to both $\xi_{\mathbf{e}}$ and $\xi_{\mathbf{e},2}$.

The following lemma studies more closely condition (*) in Definition 5.1.1.

Lemma 5.1.4. Let E be a finite-dimensional K-vector space equipped with a pseudo-norm family $\xi = (\|\cdot\|_{\omega})_{\omega\in\Omega}$. Let $\omega \in \Omega$ and let $(e_1, ..., e_r)$ be a basis of E such that, for any i = 1, ..., r, we have $\|e_i\|_{\omega} \in \mathbb{R}_{>0}$. Then $(e_1, ..., e_r)$ is an adapted basis to ξ in ω .

Proof. By hypothesis, there exist a basis $(e'_1, ..., e'_r)$ of E together with an open neighbourhood U of ω in Ω such that,

$$\forall \omega' \in U, \quad \forall i = 1, \dots, r, \quad \|e'_i\|_{\omega'} \in \mathbb{R}_{>0}.$$

It suffices to prove that there exists an open neighbourhood V of ω in Ω such that, $||e_1||$. has value in $\mathbb{R}_{>0}$ on V. As $||e_1||_{\omega} < +\infty$, we can write $e_1 = a_1e'_1 + \cdots + a_re'_r$, where $a_1, \ldots, a_r \in A_{\omega}$. As $||e_1||_{\omega} > 0$, we may assume that $a_1 \in A_{\omega}^{\times}$ and $a_2, \ldots, a_r \in A_{\omega}$. Now, by continuity of the maps $|a_1|_{\ldots}, \ldots, |a_r|_{\ldots} : \Omega \to [0, +\infty]$, there exists an open neighbourhood $V \subset U$ of ω in Ω such that

$$\forall \omega' \in V, \quad a_1 \in A_{\omega'}^{\times}, \quad a_2, ..., a_r \in A_{\omega'}.$$

Thus Proposition 1.1.6 implies that $||e_1||$. has value in $\mathbb{R}_{>0}$ in V.

26

We end this subsection with two results concerning pseudo-norm families which possess a globally adapted basis.

Lemma 5.1.5. Let E be a finite-dimensional K-vector space equipped with a pseudo-norm family $\xi = (\|\cdot\|_{\omega})_{\omega \in \Omega}$. Assume that there exists a basis $(e_1, ..., e_r)$ of E which is globally adapted to ξ . Let $(e'_1, ..., e'_r)$ be another basis of E. Then there exists an open subset $\Omega' \subset \Omega$ such that, for all $\omega \in \Omega'$, the basis $(e'_1, ..., e'_r)$ is adapted to ξ in ω and $\nu(\Omega \setminus \Omega') = 0$.

Proof. For any i = 1, ..., r, we write $e'_i = a_1^{(i)}e_1 + \cdots + a_r^{(i)}e_r$, where $a_1^{(i)}, ..., a_r^{(i)} \in K$. Let $\omega \in \Omega$. Note that the basis $(e'_1, ..., e'_r)$ is adapted to ξ in ω iff, for any $i, j \in \{1, ..., r\}$, $a_j^{(i)} \in A_{\omega}^{\times}$. As the functions $\log |a_j^{(i)}| : \Omega \to [-\infty, +\infty]$ are continuous and ν -integrable for i, j = 1, ..., r, we obtain the desired assertion.

Lemma 5.1.6. Let E be a finite-dimensional K-vector space equipped with a pseudo-norm family $\xi = (\|\cdot\|_{\omega})_{\omega \in \Omega}$. Let $s \in E \setminus \{0\}$. Then the set

$$\{\omega \in \Omega : \|s\|_{\omega} \in \{0, +\infty\}\}$$

is a locally closed subset of Ω which has measure zero with respect to ν .

Proof. Let $s \in E \setminus \{0\}$. And denote $F_s := \{\omega \in \Omega : ||s||_{\omega} \in \{0, +\infty\}\}$. We may assume that F_s is non-empty. Let $\omega_0 \in F_s$. Let $(e_1, ..., e_r)$ be a basis of E which is globally adapted to ξ .

Write $s = s_1 e_1 + \cdots + s_r e_r$, where $s_1, ..., s_r \in K$. Then, for any $\omega \in \Omega$, ω belongs to F_s iff there exists $i \in \{1, ..., r\}$ such that $s_i \in K \setminus A_{\omega}^{\times}$. For any i = 1, ..., r, denote $F_i := \{\omega \in \Omega : s_i \in K \setminus A_{\omega}^{\times}\}$. As $|s_1|, ..., |s_r|$ are continuous, F_s is locally closed. Thus

$$F_s \subset F_1 \cup \cdots \cup F_r.$$

As the functions $\log |s_i| : \Omega \to [-\infty, +\infty]$ for i = 1, ..., r are ν -integrable, $\nu(F_1) = \cdots = \nu(F_r) = 0$ and therefore F_s has measure zero.

5.2. Algebraic constructions on pseudo-norm families.

Proposition-Definition 5.2.1. Let *E* be a finite-dimensional *K*-vector space equipped with a pseudo-norm family $\xi = (\|\cdot\|_{\omega})_{\omega \in \Omega}$.

- (1) Let $F \subset E$ be a vector subspace of E. Then the family of restrictions $\xi_{|F} := (\| \cdot \|_{\omega|F})_{\omega \in \Omega}$ is a pseudo-norm family on F called the restriction of ξ .
- (2) Let G be a quotient of E. Then the family of quotient pseudo-norms $\xi_G := (\|\cdot\|_{\omega,G})_{\omega \in \Omega}$ is a pseudo-norm family on G called the quotient of ξ .
- (3) The family of dual pseudo-norms $\xi^{\vee} := (\|\cdot\|_{\omega,*})_{\omega \in \Omega}$ is a pseudo-norm family on the dual vector space E^{\vee} called the dual of ξ .
- (4) Let E' be another finite-dimensional K-vector space equipped with a pseudo-norm family ξ' = (|| · ||'_ω)_{ω∈Ω}. For any ω ∈ Ω, let || · ||_{ω,π} and || · ||_{ω,ϵ} be respectively the π-tensor product and the ϵ-tensor product of the pseudo-norms || · ||_ω and || · ||'_ω. We denote by ξ ⊗_{ϵ,π} ξ' the pseudo-norm family on E ⊗ E' consisting of the pseudo-norms || · ||_{ω,ϵ} for ω ∈ Ω_{um} and || · ||_{ω,π} for ω ∈ Ω_{ar}. This family is called the ϵ, π-tensor product as well as the π-tensor product of ξ and ξ'.
- (5) Let $i \ge 1$ be an integer. We denote by $\Lambda^i \xi$ the pseudo-norm family on $\Lambda^i E$ as the quotient of the ϵ, π -tensor product pseudo-norm family on $E^{\otimes i}$. This family is called

the exterior power pseudo-norm family on $\Lambda^i E$. If $i = \dim_K(E)$, the pseudo-norm family $\Lambda^i \xi$ is called the determinant pseudo-norm family of ξ and is denoted by $\det(\xi)$.

(6) Let $\xi' = (\|\cdot\|'_{\omega}, \mathcal{E}'_{\omega}, N'_{\omega}, \widehat{E'_{\omega}})_{\omega \in \Omega}$ be another pseudo-norm family on E Assume that for any $\omega \in \Omega$, there exists a basis \mathbf{e} of E such that \mathbf{e} is both adapted to ξ and ξ' in ω . Then, for any $\omega \in \Omega$, we have equalities

$$\mathcal{E}_{\omega} = \mathcal{E}'_{\omega}, \quad N_{\omega} = N'_{\omega}, \quad \widehat{E_{\omega}} := \widehat{E'_{\omega}}.$$

We define the local distance function by

$$(\omega \in \Omega) \mapsto d_{\omega}(\xi, \xi') := \sup_{\overline{s} \in \widehat{E_{\omega}} \setminus \{0\}} \left| \log \|\overline{s}\|_{\omega} - \log \|\overline{s}\|_{\omega}' \right| = \sup_{s \in E} \left| \log \|s\|_{\omega} - \log \|s\|_{\omega}' \right|,$$

where we use the convention that, for any $\omega \in \Omega$, for any $s \in E$ such that $||s||_{\omega} \in \{0, +\infty\}$, we have $\log ||s||_{\omega} - \log ||s||'_{\omega} = 0$.

Proof. It suffices to prove that condition (*) from Definition 5.1.1 holds for the pseudo-norm families in (1)-(5). Fix $\omega \in \Omega$.

(1) By construction of ξ_F , there exists a basis $(e_1, ..., e_r)$ of F which can be enlarged in a basis $(e_1, ..., e_r, e_{r+1}, ..., e_d)$ such that $||e_1||_{\omega}, ..., ||e_d||_{\omega} > 0$. Then Lemma 5.1.4 implies that there exists an open neighbourhood U of ω in Ω such that

$$\forall \omega' \in U, \quad \forall i = 1, ..., d, \quad ||e_i||_{\omega'} \in \mathbb{R}_{>0}.$$

Thus $(e_1, ..., e_r)$ is a basis of F which is adapted to ξ_F in ω .

(2) Write G = E/F for some vector subspace $F \subset E$. By construction of ξ_G , there exists a basis $(e_1, ..., e_r, e_{r+1}, ..., e_d)$ of E such that $(e_1, ..., e_r)$ is a basis of F, the image of $(e_{r+1}, ..., e_d)$ in E/F is a basis and, for any i = r + 1, ..., d, $\|\overline{e_i}\|_{\omega} > 0$. Then one can use Lemma 5.1.4 to conclude as above.

(3) Let $(e_1, ..., e_d)$ be a basis of E which is adapted to ξ in ω , say on an open neighbourhood U of ω . Then it follows from the construction of ξ^{\vee} that, for any $\omega' \in U$, $(e_1^{\vee}, ..., e_d^{\vee})$ is a basis of E^{\vee} such that $\|e_1^{\vee}\|_{\omega',*}, ..., \|e_d^{\vee}\|_{\omega',*} > 0$. Hence the basis $(e_1^{\vee}, ..., e_d^{\vee})$ is adapted to ξ^{\vee} in ω .

(4) Let $(e_1, ..., e_d)$ and $(e'_1, ..., e'_{d'})$ respectively be basis of E and E' that are adapted to ξ and ξ' in ω on respective open neighbourhood U and U' of ω . The construction of π -tensor product and ϵ -tensor product implies that the tensor product basis $(e_1, ..., e_d) \otimes (e'_1, ..., e'_{d'})$ is an adapted basis of $E \otimes E'$ in ω on $U \cap U'$.

(5) This follows from (2) and (4).

5.3. Dominated pseudo-norm families. In this subsection, we introduce an integrability condition for pseudo-norm families. Most of its content is an adaptation of §4.1.2 of [CM19] in our context. Recall that a function $f: \Omega \to [-\infty, +\infty]$ is called ν -dominated iff there exists $g \in \mathcal{L}^1(\Omega, \mathcal{A}, \nu)$ such that $|f| \leq g \nu$ -a.e.

Definition 5.3.1. Let *E* be a finite-dimensional *K*-vector space equipped with a pseudonorm family $\xi = (\|\cdot\|_{\omega})_{\omega \in \Omega}$. The family is called *dominated* if the pseudo-norm families ξ and ξ^{\vee} are *upper dominated*, namely we have

$$\forall f \in E \smallsetminus \{0\}, \quad \int_{\Omega}^{\overline{}} \log \|f\|_{\omega} \nu(\mathrm{d}\omega) < +\infty,$$

and

$$\forall \varphi \in E^* \smallsetminus \{0\}, \quad \int_{\Omega} \log \|\varphi\|_{\omega,*} \nu(\mathrm{d}\omega) < +\infty.$$

 ξ is called *strongly dominated* if ξ is dominated and the distance function ($\omega \in \Omega$) $\mapsto d_{\omega}(\xi, \xi^{\vee\vee}) \in \mathbb{R}_{\geq 0}$ (cf. Definition 5.2.1 (6)) is ν -dominated.

Example 5.3.2. Using the notation of Example 5.1.3, we show that for any basis $\mathbf{e} = (e_1, ..., e_r)$ of a *K*-vector space *E*, the pseudo-norm families $\xi_{\mathbf{e}}, \xi_{\mathbf{e},2}$ are dominated. Indeed, for any $\omega \in \Omega$, we have

$$\forall \lambda_1, \dots, \lambda_d \in K, \quad \log \|\lambda_1 e_1 + \dots + \lambda_r e_r\|_{\mathbf{e},\omega} \le \max_{i=1,\dots,r} |\lambda_i|_\omega + \log(r) \mathbf{1}_{\Omega_{\mathrm{ar}}}(\omega),$$

$$\forall \lambda_1, \dots, \lambda_d \in K, \quad \log \|\lambda_1 e_1 + \dots + \lambda_r e_r\|_{\mathbf{e},2,\omega} \le \max_{i=1,\dots,r} |\lambda_i|_\omega + \frac{1}{2}\log(r) \mathbf{1}_{\Omega_{\mathrm{ar}}}(\omega).$$

where $\mathbf{1}_{\Omega_{ar}}$ denotes the characteristic function of Ω_{ar} . Moreover, we have

$$\forall \omega \in \Omega, \quad \forall \alpha_1, ..., \alpha_d \in K, \quad \|\alpha_1 e_1^{\vee} + \dots + \alpha_r e_r^{\vee}\|_{\mathbf{e}, \omega, *} = \max_{i=1, \dots, r} |\alpha_i|_{\omega},$$

$$\xi_{\mathbf{e},2}^{\vee} = \xi_{\mathbf{e}^{\vee},2},$$

where $\mathbf{e}^{\vee} := (e_1^{\vee}, ..., e_r^{\vee})$ denotes the dual basis attached to \mathbf{e} . Therefore, the pseudo-norm families $\xi_{\mathbf{e}}$ and $\xi_{\mathbf{e},2}$ are dominated.

Remark 5.3.3 ([CM19], Rem. 4.1.12). If a pseudo-norm family ξ consists of ultrametric pseudo-norms ξ_{ω} for any $\omega \in \Omega_{um}$, being dominated is equivalent to being strongly dominated.

Proposition 5.3.4. Let *E* be a finite-dimensional *K*-vector space equipped with a pseudonorm family $\xi = (\|\cdot\|_{\omega})_{\omega \in \Omega}$. Assume that $\dim_K(E) = 1$. Then the following assertions are equivalent.

- (i) ξ is dominated.
- (ii) For any $s \in E \setminus \{0\}$, the function $(\omega \in \Omega) \mapsto \log ||s||_{\omega}$ is ν -dominated.
- (iii) There exists $s \in E \setminus \{0\}$, the function $(\omega \in \Omega) \mapsto \log ||s||_{\omega}$ is ν -dominated.

Proof. The arguments of the proof of ([CM19], Proposition 4.1.16) can be adapted directly. \Box

- **Proposition 5.3.5.** (1) Let *E* be a finite-dimensional *K*-vector space equipped with a pseudo-norm family $\xi = (\|\cdot\|_{\omega})_{\omega \in \Omega}$. If ξ is dominated (resp. strongly dominated), then for any vector subspace $F \subset E$, the restriction of ξ to *F* is a dominated (resp. strongly dominated) pseudo-norm family on *F*.
 - (2) Let E be a finite-dimensional K-vector space equipped with a pseudo-norm family $\xi = (\|\cdot\|_{\omega})_{\omega \in \Omega}$. If ξ is dominated (resp. strongly dominated), then for any vector subspace $F \subset E$, the quotient pseudo-norm family $\xi_{E/F}$ is dominated (resp. strongly dominated).
 - (3) Let E be a finite-dimensional K-vector space equipped with a dominated pseudo-norm family $\xi = (\|\cdot\|_{\omega})_{\omega\in\Omega}$. Then the dual pseudo-norm family ξ^{\vee} on E^{\vee} is strongly dominated.
 - (4.a) Let E, F be finite-dimensional K-vector spaces equipped with dominated pseudo-norm families ξ, ξ' respectively. Then the ϵ, π -tensor product pseudo-norm family $\xi \otimes_{\epsilon,\pi} \xi'$ is a strongly dominated pseudo-norm family on $E \otimes_K F$.

- (4.b) Let E, F be finite-dimensional K-vector spaces equipped with dominated pseudo-norm families ξ, ξ' respectively. Then the ϵ -tensor product pseudo-norm family $\xi \otimes_{\epsilon,\pi} \xi'$ is a strongly dominated pseudo-norm family on $E \otimes_K F$.
 - (5) Let E be a finite-dimensional K-vector space equipped with a dominated pseudo-norm family ξ . Let $i \in \mathbb{N}$. Then the pseudo-norm family $\Lambda^i \xi$ is strongly dominated.
 - (6) Let E be a finite-dimensional K-vector space equipped with a pseudo-norm family $\xi = (\|\cdot\|_{\omega})_{\omega \in \Omega}$. If ξ is dominated, then the determinant pseudo-norm family $\det(\xi)$ is strongly dominated.
 - (7) Let K'/K be a finite extension of fields. Let $S' = (K', \phi' : \Omega' \to M_{K'}, \nu')$ be the topological adelic curve constructed in §4.2. Denote by $\pi_{K'/K} : \Omega' \to \Omega$ the projection. Let E be a finite-dimensional K-vector space and let $E' := E \otimes_K K'$. Let $\xi = (\|\cdot\|_{\omega})_{\omega \in \Omega}$ and $\xi' = (\|\cdot\|'_{\omega'})_{\omega' \in \Omega'}$ be pseudo-norm families on E and E'respectively such that

$$\forall \omega \in \Omega, \quad \forall \omega' \in \pi_{K'/K}^{-1}(\{\omega\}), \quad \forall s \in E, \quad \|s\|_{\omega'} = \|s\|_{\omega}.$$

If ξ' is dominated, then ξ is dominated.

Proof. (1) The upper domination of ξ implies that ξ_F is upper dominated. Let $\overline{\varphi} \in F^{\vee}$. Recall that Proposition 1.1.7 (1) implies that ξ_F^{\vee} identifies with the quotient pseudo-norm family of ξ^{\vee} via $\pi : E^{\vee} \to F^{\vee}$. Let $\varphi \in E^{\vee}$ be such that $\pi(\varphi) = \overline{\varphi}$. As ξ^{\vee} is upper dominated, we deduce that there exists an integrable function $g : \Omega \to [0, +\infty]$ such that $\log \|\varphi\|_{\cdot,*} \leq g$ ν -a.e.. Therefore, for almost all $\omega \in \Omega$, we have

$$\log \|\overline{\varphi}\|_{F,\omega,*} \le \log \|\varphi\|_{\omega,*} \le g(\omega),$$

hence ξ_F^{\vee} is upper dominated.

We now prove the strong domination property. Assume that $(\omega \in \Omega) \mapsto d_{\omega}(\xi, \xi^{\vee}) \in \mathbb{R}$ is ν -dominated. Proposition 1.1.7 (1) implies that ξ_F^{\vee} identifies with a quotient pseudo-norm family of ξ^{\vee} and ξ_F^{\vee} identifies with a restricted pseudo-norm family of ξ^{\vee} . Therefore, ([CM19], Proposition 1.1.42) yields

$$\forall \omega \in \Omega, \quad 0 \le d_{\omega}(\xi_F, \xi_F^{\lor \lor}) \le d_{\omega}(\xi, \xi^{\lor \lor}).$$

Hence we deduce the ν -domination of $(\omega \in \Omega) \mapsto d_{\omega}(\xi_F, \xi_F^{\vee})$.

Ρ

(2) Let $\overline{x} \in E/F$ and let x be a lift of \overline{x} in x. Then for any $\omega \in \Omega$, we have $\log \|\overline{x}\|_{E/F,\omega} \leq \log \|x\|_{\omega}$. Therefore, the upper domination of ξ implies that $\xi_{E/F}$ is itself upper dominated. Moreover, Proposition 1.1.7 (1) implies that $\xi_{E/F}^{\vee}$ identifies with the restriction of ξ^{\vee} to $(E/F)^{\vee} \subset E^{\vee}$. Thus (1) yields the upper domination of $\xi_{E/F}^{\vee}$.

The strong domination property is proven similarly to that of (1). $\xi_{E/F}^{\vee\vee}$ identifies with a quotient pseudo-norm family of $\xi^{\vee\vee}$ and therefore ([CM19], Proposition 1.1.42) yields

$$\forall \omega \in \Omega, \quad 0 \le d_{\omega}(\xi_{E/F}, \xi_{E/F}^{\vee \vee}) \le d_{\omega}(\xi, \xi^{\vee \vee})$$

(3) By definition, ξ^{\vee} is upper dominated. Moreover, Proposition 1.1.7 (2) implies that

$$\forall x \in E \smallsetminus \{0\}, \forall \omega \in \Omega, \quad \log \|x\|_{\omega,**} \le \log \|x\|_{\omega}.$$

From the upper domination of ξ we obtain that $\xi^{\vee\vee}$ is upper dominated. Finally, Remark 5.3.3 implies that ξ^{\vee} is strongly dominated.

(4.a) and (4.b) Let $\xi \otimes_{\pi} \xi' = (\|\cdot\|_{\omega,\pi})_{\omega \in \Omega}$ and $\xi \otimes_{\epsilon} \xi' = (\|\cdot\|_{\omega,\epsilon})_{\omega \in \Omega}$. We first remark that the upper domination of ξ and ξ' implies that $\xi \otimes_{\pi} \xi'$ is upper dominated. Indeed, for

any $\varphi \in E \otimes_K E' \setminus \{0\}$, for any expression $\varphi = \sum_{i=1}^N \varphi_i \otimes \varphi'_i$, where $N \ge 1$ is an integer and $(\varphi_i, \varphi'_i) \in E \times E'$ for i = 1, ..., N, we have

$$\log \|\varphi\|_{\omega,\pi} \le \log \left(\sum_{i=1}^N \|\varphi_i\|_{\omega} \|\varphi_i'\|_{\omega}'\right) \le \mathbf{1}_{\Omega_{\mathrm{ar}}}(\omega) \log N + \max_{i=1,\dots,N} \log \|\varphi\|_{\omega} + \max_{i=1,\dots,N} \log \|\varphi'\|_{\omega}'.$$

Since $\nu(\Omega_{\mathrm{ar}}) < +\infty$, the function $\mathbf{1}_{\Omega_{\mathrm{ar}}}(\cdot) \log N$ is ν -integrable. Moreover, since ξ and ξ' are upper dominated, the functions $\max_{i=1,\dots,N} \log \|\varphi\|$. and $\max_{i=1,\dots,N} \log \|\varphi'\|'$ are upper dominated. Therefore $\xi \otimes_{\pi} \xi'$ is upper dominated. As, for any $\omega \in \Omega$, we have $\|\cdot\|_{\omega,\epsilon} \leq \|\cdot\|_{\omega,\pi}$, we deduce that $\xi \otimes_{\epsilon} \xi'$ is upper dominated. Note that we also obtain that $\xi \otimes_{\epsilon,\pi} \xi'$ is upper dominated.

We now prove that $(\xi \otimes_{\epsilon} \xi')^{\vee}$ is upper dominated. Recall that ([CM19], Propositions 1.1.57 and 1.2.20) imply that $(\xi \otimes_{\epsilon} \xi')^{\vee}$ can be identified with $\xi^{\vee} \otimes_{\epsilon,\pi} \xi'^{\vee}$. As ξ^{\vee} and ξ'^{\vee} are upper dominated, the above paragraph implies that $\xi^{\vee} \otimes_{\epsilon,\pi} \xi'^{\vee}$ is upper dominated. Hence $(\xi \otimes_{\epsilon} \xi')^{\vee}$ is upper dominated and $\xi \otimes_{\epsilon} \xi'$ is dominated.

Combining (3) with (4.a), we obtain that $\xi^{\vee} \otimes_{\epsilon} \xi'^{\vee}$ is dominated. Since $\xi \otimes_{\epsilon,\pi} \xi'$ can be identified with $(\xi^{\vee} \otimes_{\epsilon} \xi'^{\vee})^{\vee}$, (3) furnishes the domination of $\xi \otimes_{\epsilon,\pi} \xi'$.

Finally $\xi \otimes_{\epsilon} \xi'$ and $\xi \otimes_{\epsilon,\pi} \xi'$ are strongly dominated since they are both ultrametric on Ω_{um} .

(5) It is a direct consequence of (4.b) and (2).

(6) Let $\eta \in \det(E) \setminus \{0\}$. Write $\eta = e_1 \wedge \cdots \wedge e_d$ for some basis $(e_1, ..., e_d)$ of E. Then Proposition 1.1.7 (3) implies that

$$\forall \omega \in \Omega, \quad |\log \|\eta\|_{\omega, \det} \leq |\log \|e_1\|_{\omega} + \dots + |\log \|e_d\|_{\omega} |.$$

Since ξ is dominated, for any i = 1, ..., d, the function $(\omega \in \Omega) \mapsto \log ||e_i||_{\omega} \in [-\infty, +\infty]$ is ν -dominated. Therefore, the function $(\omega \in \Omega) \mapsto \log ||\eta||_{\omega, \det} \in [-\infty, +\infty]$ is ν -dominated and Proposition 5.3.4 allows to conclude that $\det(\xi)$ is dominated. Since $\det(E)$ is a K-vector space of dimension 1, $\det(\xi)$ is ultrametric on $\Omega_{\rm um}$ and therefore $\det(\xi) = \det(\xi)^{\vee\vee}$ which implies the desired strong dominance.

(7) Let $s \in E \setminus \{0\}$. Combining the results of §4.2 with the hypothesis, we have

$$\bar{\int_{\Omega}} \log \|s\|_{\omega} \nu(\mathrm{d}\omega) = \bar{\int_{\Omega'}} \log \|s\|'_{\omega'} \nu'(\mathrm{d}\omega') < +\infty.$$

Thus ξ is upper dominated.

Let $\alpha \in E^{\vee} \setminus \{0\}$. Let $\omega \in \Omega$ and let $\omega' \in \pi_{K'/K}^{-1}(\{\omega\})$. Then we have

$$\|\alpha\|_{\omega,*} = \sup_{x \in \mathcal{E}_{\omega} \smallsetminus \mathfrak{m}_{\omega} \mathcal{E}_{\omega}} \frac{|\varphi(x)|_{\omega}}{\|x\|_{\omega}} = \sup_{x \in \mathcal{E}_{\omega} \smallsetminus \mathfrak{m}_{\omega} \mathcal{E}_{\omega}} \frac{|\varphi(x)|_{\omega}}{\|x\|'_{\omega'}} \le \|\alpha\|'_{\omega',*}.$$

As ξ'^{\vee} is upper dominated, ξ^{\vee} is upper dominated as well.

Proposition 5.3.6. Let *E* be a finite-dimensional *K*-vector space equipped with two pseudonorm families ξ_1, ξ_2 such that, for any $\omega \in \Omega \in \mathbb{R}_{\geq 0}$, there exists a basis \mathbf{e}_{ω} of *E* which is both adapted to ξ_1 and ξ_2 (so that the local distance function ($\omega \in \Omega$) $\mapsto d_{\omega}(\xi_1, \xi_2) \in \mathbb{R}_{\geq 0}$ is well-defined). Assume that ξ_1 is dominated and that the local distance function ($\omega \in \Omega$) $\mapsto d_{\omega}(\xi_1, \xi_2) \in \mathbb{R}_{\geq 0}$ is ν -dominated. Then ξ_2 is dominated.

Proof. For i = 1, 2, we denote $\xi_i = (\| \cdot \|_{i,\omega})_{\omega \in \Omega}$. Let $s \in E \setminus \{0\}$. With the conventions defining the local distance function, we have the inequalities

$$\int_{\Omega} \log \|s\|_{2,\omega} \nu(\mathrm{d}\omega) \leq \int_{\Omega} \log \|s\|_{1,\omega} \nu(\mathrm{d}\omega) + \int_{\Omega} d_{\omega}(\xi_1,\xi_2) \nu(\mathrm{d}\omega) < +\infty.$$

Hence ξ_2 is upper dominated.

Let $\alpha \in E^{\vee} \setminus \{0\}$. We have the inequalities

$$\begin{split} \int_{\Omega}^{\overline{}} \log \|\alpha\|_{2,\omega,*}\nu(\mathrm{d}\omega) &\leq \int_{\Omega}^{\overline{}} \log \|\alpha\|_{1,\omega,*}\nu(\mathrm{d}\omega) + \int_{\Omega}^{\overline{}} d_{\omega}(\xi_{1}^{\vee},\xi_{2}^{\vee})\nu(\mathrm{d}\omega) \\ &\leq \int_{\Omega}^{\overline{}} \log \|\alpha\|_{1,\omega,*}\nu(\mathrm{d}\omega) + \int_{\Omega}^{\overline{}} d_{\omega}(\xi_{1},\xi_{2})\nu(\mathrm{d}\omega) < +\infty, \end{split}$$

where the third inequality comes from ([CM19], Proposition 1.1.43). Hence ξ_2^{\vee} is upper dominated. Finally, we obtain that ξ_2 is dominated. \Box

The end of this subsection is devoted to relating the strong dominance of a pseudo-norm family to model pseudo-norm families (cf. Example 5.1.3) in the case where there exists a globally adapted basis.

Proposition 5.3.7. Let E be a finite-dimensional K-vector space equipped with a dominated pseudo-norm family ξ for which there exists a globally adapted basis. Then for any basis \mathbf{e} of E which is globally adapted to ξ , there exists a ν -integrable function $A_{\mathbf{e}}: \Omega \to [0, +\infty]$ such that, for any algebraic extension L/K, for any $x \in \Omega_L$, we have

$$d_x(\xi_L, \xi_{\mathbf{e},L}) \le A_{\mathbf{e}}(\pi_{L/K}(x)).$$

Proof. Using the assumption of being globally adapted, the measure theoretic arguments in the proof of ([CM19], Proposition 4.1.7) have counterparts in our setting. The proof goes exactly along the same lines. \Box

Corollary 5.3.8. Let *E* be a finite-dimensional *K*-vector space equipped with two pseudonorm families ξ_1, ξ_2 such that there exists a basis **e** which is both globally adapted to ξ_1 and ξ_2 . Assume that ξ_1 and ξ_2 are ultrametric on Ω_{um} . If ξ_1 and ξ_2 are both dominated, then the local distance function ($\omega \in \Omega$) $\mapsto d_{\omega}(\xi_1, \xi_2) \in \mathbb{R}_{>0}$ is ν -dominated.

Proof. Let **e** of *E* which is globally adapted to ξ_1, ξ_2 . Proposition 5.3.7 with L = K implies that the local distance functions

 $(\omega \in \Omega) \mapsto d_{\omega}(\xi_1^{\vee \vee}, \xi_{\mathbf{e}}) = d_{\omega}(\xi_1, \xi_{\mathbf{e}}) \in \mathbb{R}_{\geq 0}, \quad (\omega \in \Omega) \mapsto d_{\omega}(\xi_2^{\vee \vee}, \xi_{\mathbf{e}}) = d_{\omega}(\xi_2, \xi_{\mathbf{e}}) \in \mathbb{R}_{\geq 0},$ are ν -dominated. As

$$\forall \omega \in \Omega, \quad d_{\omega}(\xi_1, \xi_2) \le d_{\omega}(\xi_1, \xi_{\mathbf{e}}) + d_{\omega}(\xi_2, \xi_{\mathbf{e}}),$$

we obtain the desired ν -dominance.

The following proposition is an adaptation of ([CM19], Corollary 4.1.10) in our setting.

Corollary 5.3.9. Let E be a finite-dimensional K-vector space equipped with a pseudo-norm family ξ . Assume that there exists a basis of E which is globally adapted to ξ . Then the following assertions are equivalent:

(i) the pseudo-norm family ξ is dominated and the local distance function $(\omega \in \Omega) \mapsto d_{\omega}(\xi, \xi^{\vee \vee}) \in \mathbb{R}_{>0}$ is ν -dominated;

- (ii) for any basis **e** of *E* such that **e** is globally adapted to ξ , the local distance function $(\omega \in \Omega) \mapsto d_{\omega}(\xi, \xi_{\mathbf{e}}) \in \mathbb{R}_{>0}$ is ν -dominated;
- (iii) there exists a **e** basis of E such that **e** is globally adapted to ξ and the local distance function ($\omega \in \Omega$) $\mapsto d_{\omega}(\xi, \xi_{\mathbf{e}}) \in \mathbb{R}_{>0}$ is ν -dominated.

Proof. (i) \Rightarrow (ii) Let **e** of *E* which is globally adapted to ξ_1, ξ_2 . Proposition 5.3.7 implies that the local distance function ($\omega \in \Omega$) $\mapsto d_{\omega}(\xi^{\vee\vee}, \xi_{\mathbf{e}}) \in \mathbb{R}_{\geq 0}$ is ν -dominated. As $\xi_{\mathbf{e}}$ is dominated, Proposition 5.3.6 implies that $\xi^{\vee\vee}$ is dominated. As

$$\forall \omega \in \Omega, \quad d_{\omega}(\xi, \xi_{\mathbf{e}}) \le d_{\omega}(\xi, \xi^{\vee \vee}) + d_{\omega}(\xi^{\vee \vee}, \xi_{\mathbf{e}}),$$

we obtain the desired ν -dominance.

(ii) \Rightarrow (iii) This is trivial.

(iii) \Rightarrow (i) Let **e** be a basis of *E* such that **e** is globally adapted to ξ and the local distance function ($\omega \in \Omega$) $\mapsto d_{\omega}(\xi, \xi_{\mathbf{e},\omega}) \in \mathbb{R}_{\geq 0}$ is ν -dominated. Then Proposition 5.3.6 yields the dominance of ξ . Moreover, Proposition 5.3.7 implies that the local distance function ($\omega \in \Omega$) $\mapsto d_{\omega}(\xi^{\vee\vee}, \xi_{\mathbf{e}}) \in \mathbb{R}_{\geq 0}$ is ν -dominated. As

$$\forall \omega \in \Omega, \quad d_{\omega}(\xi, \xi^{\vee \vee}) \le d_{\omega}(\xi, \xi_{\mathbf{e}}) + d_{\omega}(\xi^{\vee \vee}, \xi_{\mathbf{e}}),$$

we obtain the desired ν -dominance.

Corollary 5.3.10. Let E be a finite-dimensional K-vector space equipped with a pseudonorm family ξ . Assume that there exists a basis of E which is globally adapted to ξ . If Eis dominated, then for any algebraic extension L/K, the extension of scalars ξ_L is strongly dominated.

Proof. Write $S \otimes_K L = (L, \phi_L : \Omega_L \to M_L, \nu_L)$. Let **e** be a basis of E which is globally adapted to ξ . Proposition 5.3.7 implies that the local distance function $(\omega \in \Omega_L) \mapsto d_{\omega}(\xi_L, \xi_{\mathbf{e},L}) \in \mathbb{R}_{\geq 0}$ is ν_L -dominated. As $\xi_{\mathbf{e},L}$ is dominated, Proposition 5.3.6 implies that ξ_L is dominated. \Box

5.4. **Regularity and measurability conditions for pseudo-norm families.** In this subsection, we introduce regularity conditions for pseudo-norm families. Here the theory differs from the classical theory of adelic curves, due to the topological nature of the adelic space.

Definition 5.4.1. Let *E* be a finite-dimensional *K*-vector space equipped with a pseudo-norm family $\xi = (\|\cdot\|_{\omega})_{\omega \in \Omega}$.

- (1) The family ξ is called *continuous*, respectively *usc*, *lsc* if, for any $s \in E$, the map $(\omega \in \Omega) \mapsto \log \|s\|_{\omega} \in [-\infty, +\infty]$ is continuous, respectively usc, *lsc*. We denote by $\mathcal{N}(E)^{\text{cont}}$, respectively $\mathcal{N}(E)^{\text{usc}}$, $\mathcal{N}(E)^{\text{lsc}}$ the set of continuous, respectively usc, *lsc* pseudo-norm families on E.
- (2) The family ξ is called ν -measurable if, for any $s \in E$, the map $(\omega \in \Omega) \mapsto \log ||s||_{\omega} \in [-\infty, +\infty]$ is ν -measurable. We denote by $\mathcal{N}(E)^{\nu}$ the set of ν -measurable pseudo-norm families on E.
- **Example 5.4.2.** (1) Let ξ be a pseudo-norm family on a finite-dimensional *K*-vector space *E*. Assume that ξ is either usc or lsc. Then ξ is a ν -measurable norm family on *E*. In particular, if ξ is continuous, then ξ is ν -measurable.
 - (2) Assume that Ω is discrete. Let E be a finite-dimensional K-vector space equipped with a pseudo-norm family $\xi = (\|\cdot\|_{\omega})_{\omega \in \Omega}$. Then ξ is continuous.

(3) Model pseudo-norm families constructed in Example 5.1.3 are continuous. Indeed, as Ω_{ar} and Ω_{um} are open, it follows from the definition of such pseudo-norm families on each of these open sets.

Proposition 5.4.3. Let *E* be a finite-dimensional *K*-vector space equipped with a pseudonorm family $\xi = (\|\cdot\|_{\omega})_{\omega \in \Omega}$.

- (1) Let $F \subset E$ be any vector subspace. Assume that ξ is lsc, resp. usc. Then the restriction pseudo-norm family $\xi_F = (\|\cdot\|_{F,\omega})_{\omega \in \Omega}$ on F induced by ξ is lsc, resp. usc.
- (2) Assume that ξ is usc. Then the dual pseudo-norm family $\xi^{\vee} = (\|\cdot\|_{\omega,*})_{\omega\in\Omega}$ on E^{\vee} is lsc, hence is ν -measurable.
- (3) Let $\pi : E \twoheadrightarrow G$ be a quotient vector space of E. Assume that ξ is usc. Then the quotient pseudo-norm family $\xi_G = (\|\cdot\|_{G,\omega})_{\omega \in \Omega}$ induced by ξ is usc, hence is ν -measurable.
- (4) Let $n \ge 1$ be an integer. For any i = 1, ..., n, let E_i be a finite-dimensional K-vector space equipped with a pseudo-norm family $\xi_i = (\|\cdot\|_{i,\omega})_{\omega \in \Omega}$.
 - (4.a) Assume that $\xi_1, ..., \xi_n$ are usc. Then the π -tensor product $\xi_1 \otimes_{\pi} \cdots \otimes_{\pi} \xi_n$ is a usc and ν -measurable pseudo-norm family on $E_1 \otimes \cdots \otimes E_n$.
 - (4.b) Assume that the dual pseudo-norm families $\xi_1^{\vee}, ..., \xi_n^{\vee}$ are continuous. Then the ϵ -tensor product $\xi_1 \otimes_{\epsilon} \cdots \otimes_{\epsilon} \xi_n$ is a lsc and ν -measurable pseudo-norm family on $E_1 \otimes \cdots \otimes E_n$.
 - (4.c) Assume that the dual pseudo-norm families $\xi_1^{\vee}, ..., \xi_n^{\vee}$ are continuous. Then the ϵ, π -tensor product $\xi_1 \otimes_{\epsilon,\pi} \cdots \otimes_{\epsilon,\pi} \xi_n$ is a ν -measurable pseudo-norm family on $E_1 \otimes \cdots \otimes E_n$.
- (5) Assume that ξ is usc. The determinant pseudo-norm family det (ξ) on det(E) is usc and ν -measurable.
- (6) Let L/K be an algebraic extension. Then the extension of scalars pseudo-norm family $\xi_L = (\|\cdot\|_x)_{x \in \Omega_L}$ on $E_L := E \otimes_K L$ induced by ξ is lsc, hence ν -measurable.

Proof. (1) It is immediate from the definition of a continuous pseudo-norm family.

(2) Let $\varphi \in E^{\vee} \setminus \{0\}$ and $a \in \mathbb{R}_{>0}$. Assume that $\|\varphi\|_{\omega,*} \in]a, +\infty]$. Let us prove that $\|\varphi\|_{\cdot,*}^{-1}(]a, +\infty]$ is open. Let $x \in \mathcal{E}_{\omega} \setminus \mathfrak{m}_{\omega}\mathcal{E}_{\omega}$. By upper semi-continuity of $\|x\|$ on Ω , there exists an open neighbourhood U of ω such that $\|x\|$ has value in $\mathbb{R}_{\geq 0}$ on U. Then the function $(\omega' \in U) \mapsto \frac{|\varphi(x)|_{\omega'}}{\|x\|_{\omega'}} \in \mathbb{R}_{>0}$ is lower semi-continuous. Hence up to shrinking U, we may assume that for any $\omega' \in U$, we have

$$a < \frac{|\varphi(x)|_{\omega'}}{\|x\|_{\omega'}} \le \|\varphi\|_{\omega',*}.$$

We deduce the desired lower semi-continuity.

(3) Let $\pi : E \to G$ be a quotient vector space of E. By definition, for any $\overline{x} \in E/F$, we have

$$\|\overline{x}\|_{\omega,E/F} = \inf_{x \in \pi^{-1}(\overline{x})} \|x\|_{\omega}.$$

Thus $\|\overline{x}\|_{\cdot,E/F}$ is the infimum of a family of usc functions and hence is usc. Since log is non-decreasing, $\log \|\overline{x}\|_{\cdot,E/F}$ is usc.

(4.a) Let $x \in E_1 \otimes \cdots \otimes E_n$ and $a \in \mathbb{R}_{>0}$. We prove that $||x||_{,\pi}^{-1}([0,a[)$ is open. Let $\omega \in ||x||_{\pi}^{-1}([0,a[))$. By definition of $||x||_{\omega,\pi}$, there exist a decomposition $x = \sum_{i=1}^{N} x_1^{(i)} \otimes \cdots \otimes x_n^{(i)}$,

with $\|x_j^{(i)}\|_{j,\omega} < +\infty$ for any i = 1, ..., N and j = 1, ..., n, such that

$$\sum_{i=1}^{N} \|x_1^{(i)}\|_{1,\omega} \cdots \|x_n^{(i)}\|_{n,\omega} < a.$$

Let U be an open neighbourhood of ω such that, for any $\omega' \in U$, for any i = 1, ..., N and j = 1, ..., n, we have $||x_j^{(i)}||_{j,\omega'} < +\infty$. Up to shrinking U, we may assume that, for any $\omega' \in U$, we have

$$||x||_{\omega',\pi} \le \sum_{i=1}^{N} ||x_1^{(i)}||_{1,\omega'} \cdots ||x_n^{(i)}||_{n,\omega'} < a.$$

Thus $||x||_{,\pi}$ and $\log ||x||_{,\pi}$ are use and ν -measurable.

(4.b) Let $x \in E_1 \otimes \cdots \otimes E_n$ and $a \in \mathbb{R}_{>0}$. We prove that $||x||_{,\epsilon}^{-1}(]a, +\infty]$) is open. Let $\omega \in ||x||_{\epsilon}^{-1}(]a, +\infty]$). By definition of $||x||_{\omega,\epsilon}$, there exists $(f_1, ..., f_n) \in E_1^{\vee} \times \cdots \times E_n^{\vee}$ such that $0 < ||f_1||_{1,\omega,*}, ..., ||f_n||_{n,\omega,*} < +\infty$ and

$$a < \frac{|x(f_1, ..., f_n)|_{\omega}}{\|f_1\|_{1,\omega,*} \cdots \|f_n\|_{n,\omega,*}}.$$

By continuity $|x(f_1, ..., f_n)|$ and Lemma 5.1.4, there exists an open neighbourhood U of ω such that, for any $\omega' \in U$, we have

$$0 < \|f_1\|_{1,\omega',*}, \dots, \|f_n\|_{n,\omega',*} < +\infty.$$

Up to shrinking U, by continuity of $||f_1||_{1,\cdot,*}, ..., ||f_n||_{n,\cdot,*}$, we may assume that

$$\forall \omega' \in U, \quad a < \frac{|x(f_1, ..., f_n)|_{\omega'}}{\|f_1\|_{1, \omega', *} \cdots \|f_n\|_{n, \omega', *}} \le \|x\|_{\omega', \epsilon}.$$

Hence $||x||_{\cdot,\epsilon}$ and $\log ||x||_{\cdot,\epsilon}$ are lsc and ν -measurable.

(4.c) Let $x \in E_1 \otimes \cdots \otimes E_n$ and $A \subset [-\infty, +\infty]$ be a measurable set. By (4.a) and (4.b), the sets

$$(\log \|x\|_{\cdot,\pi})^{-1}(A), \quad (\log \|x\|_{\cdot,\epsilon})^{-1}(A)$$

are ν -measurable. We can then conclude by using the equality

$$(\log \|x\|_{\cdot,\epsilon,\pi})^{-1}(A) = \left((\log \|x\|_{\cdot,\pi})^{-1}(A) \cap \Omega_{\mathrm{ar}} \right) \cup \left((\log \|x\|_{\cdot,\epsilon})^{-1}(A) \cap \Omega_{\mathrm{um}} \right)$$

combined with the fact that Ω_{ar} and Ω_{um} are both ν -measurable.

(5) Let $\eta \in \det(E)$ and $\omega \in \Omega$. Assume that $\|\eta\|_{\omega,\det} < +\infty$. Let $a > \|\eta\|_{\omega,\det}$. We show that $\|\eta\|_{\cdot,\det}^{-1}([0,a[)$ is an open neighbourhood of ω . By Proposition 1.1.7 (3), there exist $x_1, ..., x_r \in \mathcal{E}_{\omega}$ such that $\eta = x_1 \wedge \cdots \wedge x_r$ and

$$\|\eta\|_{\omega,\det} \le \|x_1\|_{\omega} \cdots \|x_r\|_{\omega} < a.$$

Let $\mathbf{e} = (e_1, ..., e_r)$ a basis of E which is adapted to ξ . Thus, for any i = 1, ..., r, $x_i = x_i^{(1)}e_1 + \cdots + x_i^{(r)}e_r \in \bigoplus_{i=1}^r A_{\omega}e_i$. Let U be an open neighbourhood of ω such that, for any $\omega' \in U$, for any $i, j \in \{1, ..., r\}$, $x_i^{(j)} \in A_{\omega'}$. By semi-continuity of $||x_1||_{\cdot}, ..., ||x_r||_{\cdot}$, up to shrinking U, we may assume that, for any $\omega' \in U$, we have

$$\|\eta\|_{\omega',\det} \le \|x_1\|_{\omega'} \cdots \|x_r\|_{\omega'} < a$$

Hence, for any a > 0, the set $\|\eta\|_{\cdot,\det}^{-1}([0,a[))$ is open and $\det(\xi)$ is use and ν -measurable.

6. Adelic vector bundles

Throughout this section, we fix a topological adelic curve $S = (K, \phi : \Omega \to M_K, \nu)$. Recall that, for any $\omega \in \Omega$, we denote by A_{ω} and κ_{ω} , the finiteness ring and the residue field of ω respectively.

6.1. Definitions.

Definition 6.1.1. Let ξ be a pseudo-norm family on a finite-dimensional K-vector space E. Then $\overline{E} = (E, \xi)$ is called

- a upper/lower semi-continuous adelic vector bundle (usc/lsc adelic vector bundle for short), if the pseudo-norm families ξ and ξ[∨] are upper/lower semi-continuous and ξ is dominated;
- (2) an *adelic vector bundle* if the pseudo-norm families ξ and ξ^{\vee} are continuous and ξ is dominated;
- (3) a measurable adelic vector bundle on S, if the pseudo-norm families ξ and ξ^{\vee} are ν -measurable and ξ is dominated.

Note that any (usc/lsc) adelic vector bundle is a measurable adelic vector bundle.

If $\overline{E} = (E, \xi)$ is a (usc/lsc/measurable) adelic vector bundle on S and if the pseudo-norm family ξ is strongly dominated, then \overline{E} is called a *strongly (usc/lsc/measurable) adelic vector* bundle on S.

Moreover, an adelic vector bundle $\overline{E} = (E, \xi)$ on S is called *Hermitian* if ξ is Hermitian.

Proposition 6.1.2. Let ξ be a pseudo-norm family on a finite-dimensional K-vector space E and assume that ξ is ultrametric on Ω_{um} . Then $\overline{E} = (E, \xi)$ is a usc adelic vector bundle on S iff $\overline{E} = (E, \xi)$ is an adelic vector bundle on S.

Proof. An adelic vector bundle is a use adelic vector bundle. We show the converse implication. Assume that \overline{E} is use. It suffices to prove that ξ and ξ^{\vee} are both lse. As ξ is use, Proposition 5.4.3 (2) implies that ξ^{\vee} is lse. Moreover, since ξ is ultrametric on $\Omega_{\rm um}$, $\xi = \xi^{\vee\vee}$ and, as ξ^{\vee} is use, we obtain that ξ is lse.

Proposition 6.1.3. Let $\xi = (\|\cdot\|_{\omega})_{\omega \in \Omega}$ be a pseudo-norm family on a K-vector space E of dimension 1. Then (E, ξ) is an adelic vector bundle on S iff ξ is continuous and dominated.

Proof. The direct implication being immediate, it suffices to show the converse. Assume that ξ is continuous and dominated. Let $\varphi \in E^{\vee} \setminus \{0\}$. Let $x \in E$ such that $\varphi(x) = 1$. Then, for any $\omega \in \Omega$, we have $\|\varphi\|_{\omega,*} = 1/\|x\|_{\omega}$, with the convention $1/0 = +\infty$ and $1/+\infty = 0$. Thus ξ^{\vee} is continuous.

Proposition 6.1.4. Let $\overline{E} = (E, \xi)$ be an adelic vector bundle on S.

- (1) Let $F \subset E$ be any vector subspace. Then (F, ξ_F) is an adelic vector bundle on S.
- (2) $\overline{E}^{\vee} = (E^{\vee}, \xi^{\vee})$ is a lsc adelic vector bundle on S. Moreover, if $\xi^{\vee\vee}$ is continuous, then \overline{E}^{\vee} is an adelic vector bundle on S. In particular, if ξ is ultrametric on $\Omega_{\rm um}$, then \overline{E}^{\vee} is an adelic vector bundle on S.

36
- (3) Let $\pi : E \to G$ be a quotient vector space of E. Let ξ_G denote the quotient pseudonorm family on G induced by ξ . Then (G, ξ_G) is a usc adelic vector bundle on S. Moreover, if ξ is ultrametric on Ω_{um} , then (G, ξ_G) is an adelic vector bundle on S.
- (4) Let $n \ge 1$ be an integer. For any i = 1, ..., n, let $\overline{E_i} = (E_i, \xi_i)$ be an adelic vector bundle on S. We assume that the pseudo-norm families $\xi_1, ..., \xi_n$ are ultrametric on Ω_{um} .
 - (4.a) The ϵ -tensor product $\overline{E_1} \otimes_{\epsilon} \cdots \otimes_{\epsilon} \overline{E_n}$ is a measurable adelic vector bundle on S.
 - (4.b) The (ϵ, π) -tensor product $\overline{E_1} \otimes_{\epsilon, \pi} \cdots \otimes_{\epsilon, \pi} \overline{E_n}$ is a measurable adelic vector bundle on S.
- (5) $(\det(E), \det(\xi))$ is a measurable adelic vector bundle on S. Moreover, if ξ is Hermitian, then $(\det(E), \det(\xi))$ is an adelic vector bundle on S.

Proof. (1) Propositions 5.3.5 (1) and 5.4.3 (1) imply that ξ_F is continuous and dominated. Now Proposition 5.4.3 (2) implies that ξ_F^{\vee} is lsc. Moreover, Proposition 1.1.7 (1) implies that ξ_F^{\vee} identifies with the a quotient pseudo-norm family of ξ^{\vee} , and is thus use by Proposition 5.4.3. Hence ξ_F^{\vee} is continuous.

(2) As ξ is dominated, ξ^{\vee} is dominated (cf. Proposition 5.3.5 (3)). Since ξ^{\vee} is continuous, Proposition 5.4.3 (2) implies that $\xi^{\vee\vee}$ is lsc. The last assertion comes from the fact that if ξ is ultrametric on $\Omega_{\rm um}$, $\xi = \xi^{\vee\vee}$.

(3) First note that Proposition 5.3.5 (2) implies that ξ_G is dominated. Now Proposition 1.1.7 (1) combined with (1) implies that ξ_G^{\vee} is continuous. Moreover, Proposition 5.4.3 (3) yields that ξ_G is usc. Thus (G, ξ_G) is a usc adelic vector bundle on S. If we further assume that ξ is ultrametric on $\Omega_{\rm um}$, ξ_G is ultrametric on $\Omega_{\rm um}$ as well and $\xi_G^{\vee\vee} = \xi_G$ and Proposition 5.4.3 (2) implies that ξ_G is lsc and therefore (G, ξ_G) is an adelic vector bundle on S.

(4.a) Proposition 5.3.5 (4.a) implies that $\xi_1 \otimes_{\epsilon} \cdots \otimes_{\epsilon} \xi_n$ is (strongly) dominated. Proposition 5.4.3 (4.a) implies that $\xi_1 \otimes_{\epsilon} \cdots \otimes_{\epsilon} \xi_n$ is lsc and hence ν -measurable. Combining the identification $(\xi_1 \otimes_{\epsilon} \cdots \otimes_{\epsilon} \xi_n)^{\vee} = \xi_1^{\vee} \otimes_{\epsilon,\pi} \cdots \otimes_{\epsilon,\pi} \xi_n^{\vee}$ with Proposition 5.4.3 (4.b), we obtain that $(\xi_1 \otimes_{\epsilon} \cdots \otimes_{\epsilon} \xi_n)^{\vee}$ is ν -measurable (cf. $\xi_i^{\vee\vee} = \xi_i$ is continuous for i = 1, ..., r).

(4.b) Proposition 5.3.5 (4.b) implies that $\xi_1 \otimes_{\epsilon,\pi} \cdots \otimes_{\epsilon,\pi} \xi_n$ is (strongly) dominated. Proposition 5.4.3 (4.b) implies that $\xi_1 \otimes_{\epsilon,\pi} \cdots \otimes_{\epsilon,\pi} \xi_n$ is ν -measurable. Moreover, as $\xi_i^{\vee\vee} = \xi$ for i = 1, ..., n, we have an identification $\xi_1 \otimes_{\epsilon,\pi} \cdots \otimes_{\epsilon,\pi} \xi_n = (\xi_1^{\vee} \otimes_{\epsilon} \cdots \otimes_{\epsilon} \xi_n^{\vee})^{\vee}$ and therefore $(\xi_1 \otimes_{\epsilon,\pi} \cdots \otimes_{\epsilon,\pi} \xi_n)^{\vee} = (\xi_1^{\vee} \otimes_{\epsilon} \cdots \otimes_{\epsilon} \xi_n^{\vee})^{\vee} = \xi_1^{\vee} \otimes_{\epsilon} \cdots \otimes_{\epsilon} \xi_n^{\vee}$, where the last identification comes from the fact that $\xi_1 \otimes_{\epsilon} \cdots \otimes_{\epsilon} \xi_n$ is ultrametric on $\Omega_{\rm um}$. Henceforth, the pseudo-norm family $(\xi_1 \otimes_{\epsilon,\pi} \cdots \otimes_{\epsilon,\pi} \xi_n)^{\vee}$ is lsc (and ν -measurable).

(5) Proposition 5.3.5 (6) implies that $\det(\xi)$ is dominated. Proposition 5.4.3 (5) implies that the pseudo-norm families $\det(\xi)$ and $\det(\xi^{\vee})$ are usc. Moreover, (3) implies that $(\det(\xi))^{\vee}$ is lsc. Thus $(\det(E), \det(\xi))$ is a measurable adelic vector bundle on S. If we further assume that ξ is Hermitian, Proposition 6.1.3 implies that it suffices to prove that $\det(\xi)$ is continuous. Then ([CM19], Proposition 1.2.47) implies that, for any $\omega \in \Omega$, we have identifications

$$\det(\xi) = \det(\xi)^{\vee \vee} = \det(\xi^{\vee})^{\vee}.$$

Therefore we can use Proposition 5.4.3 (3) to obtain that $det(\xi)$ is lsc.

6.2. Arakelov degree.

Definition 6.2.1. Let $\overline{E} := (E, \xi = (\|\cdot\|_{\omega})_{\omega \in \Omega})$ be a measurable adelic vector bundle on S. Let $s \in E \setminus \{0\}$, then the map $\log \|s\| : \Omega \to [-\infty, +\infty]$ is ν -integrable and we define the

Arakelov degree of s as

$$\widehat{\operatorname{deg}}_{\xi}(s) := -\int_{\Omega} \log \|s\|_{\omega} \nu(\mathrm{d}\omega).$$

Definition 6.2.2. We assume that the adelic curve S is proper. Let $\overline{E} := (E, \xi = (\|\cdot\|_{\omega})_{\omega \in \Omega})$ be an usc adelic vector bundle on S. Then the quantity

$$\widehat{\operatorname{deg}}(\overline{E}) := \widehat{\operatorname{deg}}_{\operatorname{det}(\xi)}(\eta) = -\int_{\Omega} \log \|\eta\|_{\omega, \operatorname{det}} \nu(\operatorname{d}\omega),$$

where $\eta \in \det(E) \setminus \{0\}$, is independent of the choice of η . We call it the Arakelov degree of \overline{E} .

6.3. Example in Nevanlinna theory. In this subsection, we give an example of adelic vector bundles in the context of Nevanlinna theory. Fix R > 0 and consider the topological adelic curve $S_R = (K_R, \phi_R : \Omega_R \to M_{K_R}, \nu_R)$ constructed in §2.2.2. For any $\omega \in \Omega_R$, we denote by $A_{R,\omega}$ the finiteness ring on ω . Let E be an arbitrary finite-dimensional \mathbb{C} -vector space equipped with a norm $\|\cdot\|$. Let us see that this data induces an adelic vector bundle on S_R . Let $E_R := E \otimes_{\mathbb{C}} K_R$.

Let $\omega \in \Omega_{R,\mathrm{ar}}$, we denote $\mathcal{E}_{R,\omega} := E \otimes_{\mathbb{C}} A_{R,\omega}$. It is a free $A_{R,\omega}$ of rank $\dim_{\mathbb{C}}(E)$ and $\widehat{E_{R,\omega}} := \mathcal{E}_{R,\omega} \otimes_{A_{R,\omega}} \mathbb{C}$ identifies with E. By lifting the norm $\|\cdot\|$ on E, we obtain a local-pseudo-norm on E_R in ω , which is denoted by $\|\cdot\|_{\omega}$.

Let $\omega \in \Omega_{R,\text{um}}$. Recall that $A_{R,\omega} = K_R$ as $|\cdot|_{\omega}$ is a usual absolute value. Then the completion $K_{R,\omega}$ of K_R w.r.t. $|\cdot|_{\omega}$ is isomorphic to $\mathbb{C}((T))$. Denote by $K_{R,\omega}^{\circ} \cong \mathbb{C}[[T]]$ the corresponding valuation ring of $K_{R,\omega}$. Then $K_{R,\omega}^{\circ}$ is the completion of the discrete valuation ring

$$\{f \in K_R : f(\omega) \in \mathbb{P}^1_{\mathbb{C}} \smallsetminus \{0, \infty\}\}.$$

Let $\mathcal{E}_{R,\omega} := E \otimes_{\mathbb{C}} K_{R,\omega}^{\circ}$, it is a free sub- $K_{R,\omega}^{\circ}$ -module of rank $\dim_{\mathbb{C}}(E)$ of $E_{R,\omega} := E \otimes_{K_R} K_{R,\omega}$, hence it is a (finitely generated) lattice of E_R . We consider the ultrametric lattice norm $\|\cdot\|_{\omega}$ induced by $\mathcal{E}_{R,\omega}$ on $E_{R,\omega}$ (cf. e.g. Definition 1.1.23 in [CM19]). Recall that, since $|\cdot|_{\omega}$ is discrete, $\mathcal{E}_{R,\omega}$ coincides with the unit ball of $E_{R,\omega}$ w.r.t. $\|\cdot\|_{\omega}$.

From the two above paragraphs, we have a collection $\xi = (\|\cdot\|_{\omega})_{\omega \in \Omega_R}$ of pseudo-norms on E_R .

Proposition 6.3.1. We use the same notation as above. Then $\overline{E_R} := (E_R, \xi_R)$ is an adelic vector bundle on S_R .

Proof. Let us first prove that ξ is a pseudo-norm family on E_R (in the sense of Definition 5.1.1). By construction of the pseudo-norms $\|\cdot\|_{\omega}$, any basis of E defines a basis of E_R which is adapted to $\|\cdot\|_{\omega}$ for all $\omega \in \Omega_R$. Thus ξ_R is a pseudo-norm family on E_R .

Let us now show that ξ_R is dominated. Fix an arbitrary basis $(e_1, ..., e_r)$ of E (which is globally adapted to ξ_R). Let $s = s_1e_1 + \cdots + s_re_r \in E_R \setminus \{0\}$, where $s_1, ..., s_r \in K_R$. Let $\omega \in \Omega_{R,\text{um}}$, then $||s||_{\omega} \neq 1$ iff ω is either a zero or a pole of s_i for some $i \in \{1, ..., r\}$. Since elements of K_R only have a finite number of zeroes and poles, this means that $||s||_{\omega} = 1$ for all but a finite number of $\omega \in \Omega_{R,\text{um}}$. As $\Omega_{R,\text{um}}$ is a discrete topological space equipped with a counting measure, $(\omega \in \Omega_{R,\text{um}}) \mapsto \log ||s||_{\omega} \in [-\infty, +\infty]$ is a ν_R -integrable function. Now let $\omega \in \Omega_{R,\text{ar}}$. Then we have

$$||s||_{\omega} \le \max_{1 \le i \le r} |s_i|_{\omega} \max_{1 \le i \le r} ||e_i||.$$

As $\nu_R(\Omega_{R,\mathrm{ar}}) = 1$, the function $(\omega \in \Omega_{R,\mathrm{ar}}) \mapsto \max_{1 \leq i \leq r} \log \|e_i\| \in [-\infty, +\infty]$ is ν_R -integrable. Moreover, for any i = 1, ..., r, the function $\log |s_i|$. is ν_R -integrable. Hence $(\omega \in \Omega_{R,\mathrm{ar}}) \mapsto \max_{1 \leq i \leq r} \log |s_i|_{\omega} \in [-\infty, +\infty]$ is ν_R -integrable. Finally, we obtain that ξ_R is upper dominated.

Let $\alpha = \alpha_1 e_1^{\vee} + \cdots + \alpha_r e_r^{\vee} \in E_R^{\vee} \setminus \{0\}$, where $\alpha_1, \dots, \alpha_r \in K_R$. Let $\omega \in \Omega_{R, \text{um}}$. Denote $\mathcal{E}_{R, \omega}^{\vee} := \text{Hom}_{K_{R, \omega}^{\circ}}(\mathcal{E}_{R, \omega}, K_{R, \omega}^{\circ})$. Then ([CM19], Proposition 1.1.34) yields the equality

$$\|\cdot\|_{\omega,*} = \|\cdot\|_{\mathcal{E}_{B,\omega}^{ee}}$$

of norms on $E_{R,\omega}$. Therefore the same argument used in the above paragraph for the lattice $\mathcal{E}_{R,\omega}^{\vee}$ instead of $\mathcal{E}_{R,\omega}$ implies that $\|\alpha\|_{\omega,*} = 1$ for almost all $\omega \in \Omega_{R,\mathrm{um}}$ and thus $[(\omega \in \Omega_{R,\mathrm{um}}) \mapsto \log \|\alpha\|_{\omega,*}]$ is a ν_R -integrable function. Now let $\omega \in \Omega_{R,\mathrm{ar}}$. Then we have

$$\|\alpha\|_{\omega,*} \le \max_{1 \le i \le r} |\alpha_i|_{\omega} \max_{1 \le i \le r} \|e_i^{\vee}\|_*,$$

and the same arguments of the above paragraph imply that ξ_R^{\vee} is upper dominated. Hence ξ_R is a dominated pseudo-norm family on E.

Finally, let us prove that ξ_R and ξ_R^{\vee} are continuous. We start by showing the continuity of ξ_R . Let $s \in E_R$. By discreteness of $\Omega_{R,\text{um}}$, it suffices to prove that ||s||. is continuous on $\Omega_{R,\text{ar}}$. Fix an arbitrary basis $(e_1, ..., e_r)$ of E (which is globally adapted to ξ_R). Write $s = s_1e_1 + \cdots + s_re_r \in E_R \setminus \{0\}$, where $s_1, ..., s_r \in K_R$. By definition, for any $\omega \in \Omega_{R,\text{ar}}$, we have

$$\|s\|_{\omega} = \begin{cases} \|s_1(\omega)e_1 + \cdots + s_r(\omega)e_r\| & \text{if } s_1(\omega), \dots, s_r(\omega) \in \mathbb{P}^1(\mathbb{C}) \smallsetminus \{\infty\}, \\ +\infty & \text{otherwise.} \end{cases}$$

Note that the set $U := \{\omega \in \Omega_{R,ar} : s_1(\omega), ..., s_r(\omega) \in \mathbb{P}^1(\mathbb{C}) \setminus \{\infty\}\}$ is the complement of a finite set, hence it is open. The continuity of ||s||. on U is equivalent to the continuity of the map

$$f: (\omega \in U) \mapsto s_1(\omega)e_1 + \cdots + s_r(\omega)e_r \in E,$$

where E is equipped with the topology induced by the norm $\|\cdot\|$. By continuity of the maps $|s_1|, ..., |s_r|$ on U, we see that f is continuous w.r.t. the topology on E induced by the infinite norm w.r.t. the basis $(e_1, ..., e_r)$, which is the same as the desired topology by equivalence of norms. Therefore, $\|s\|$ is continuous on U. It remains to prove that $\|s\|$ is continuous at any point of the finite set $\Omega_{R,ar} \setminus U$. Let $\omega_0 \in \Omega_{R,ar} \setminus U$. Since $\Omega_{R,ar} \setminus U$ is discrete, it suffices to prove the continuity of $\|s\|$ on a neighbourhood V of ω_0 such that $V \cap (\Omega_{R,ar} \setminus U) = \{\omega_0\}$. To show that $\|s\|$ is continuous at ω_0 , it suffices to show that

$$\lim_{V \ni \omega \to \omega_0} \|s\|_{\omega} = \|s\|_{\omega_0} = +\infty.$$

By symmetry, we may assume that s_1 has minimal valuation, i.e. has a pole of the greatest order in ω_0 , among $s_1, ..., s_r$. Denote this order by p < 0. We see that, up to shrinking V, we can write $s = (T - \omega_0)^p s'$, where s' satisfies

$$\forall \omega \in V, \quad \|s'\|_{\omega} < +\infty.$$

By the above case of continuity and compactness of $\Omega_{R,ar}$, we obtain that ||s'||. is bounded on V. As

$$\lim_{V\ni\omega\to\omega_0}(\omega-\omega_0)^p=+\infty,$$

we obtain the desired continuity.

To conclude the proof, we prove that ξ_R^{\vee} is continuous. As in the previous case, it suffices to prove that, for any $\alpha \in E_R^{\vee}$, the function $\|\alpha\|_{,\omega}$ is continuous on $\Omega_{R,\mathrm{ar}}$. Note that, by construction, for $\omega \in \Omega_{R,\mathrm{ar}}$, the pseudo-norm $\|\cdot\|_{\omega,*}$ is the pseudo-norm constructed on E_R^{\vee} by lifting the norm $\|\cdot\|_*$ on E^{\vee} . Therefore, the above case can be applied and implies that ξ_R^{\vee} is continuous.

Definition 6.3.2. Let E be a finite-dimensional complex vector space equipped with a norm $\|\cdot\|$. Let R > 0. Let $E_R := E_{\mathbb{C}} \otimes_{\mathbb{C}} K_R$ and $\xi_R = (\|\cdot\|_{\omega})_{\omega \in \Omega_R}$ be the pseudo-norm family on E_R constructed as above. Then the adelic vector bundle $\overline{E_R} := (E_R, \xi_R)$ is called the *induced adelic vector bundle on* S_R by the complex normed vector space $(E, \|\cdot\|)$. Note that any basis of E is globally adapted to ξ_R .

Remark 6.3.3. Let E be a finite-dimensional complex vector space equipped with a norm $\|\cdot\|$. Let R > 0. Consider a finite extension K'/K_R . Then one can use the same arguments as above to prove that $(E, \|\cdot\|)$ induces an adelic vector bundle on the topological adelic curve $S_R \otimes_{K_R} K'$.

6.4. Adelic vector bundles on families of topological adelic curves.

Definition 6.4.1. Let (I, \leq_I) be a (non-empty) totally ordered set. Let $\mathbf{S} = (S_i = (K_i, \phi_i : \Omega_i \to M_{K_i}, \nu_i))_{i \in I}$ be a family of topological adelic curves with base space Ω_0 and family of structural morphisms $(\alpha_i : \Omega_i \to \Omega_0)_{i \in I}$ (cf. Definition 3.1.1). By a *(usc/lsc/measurable)* adelic vector bundle on \mathbf{S} , we mean a collection $\overline{\mathbf{E}} = (\overline{E_i} = (E_i, \xi_i = (\|\cdot\|_{i,\omega})_{\omega \in \Omega_i}))_{i \in I}$ where

- (i) for any $i \in I$, $\overline{E_i}$ is an (usc/lsc/measurable) adelic vector bundle on S_i ;
- (ii) for any $i \leq_I i' \in I$, we have an isomorphism $E_{i'} \otimes_{K_{i'}} K_i \cong E_i$, in particular $E_{i'} \subset E_i$;
- (iii) for any $i \leq_I i' \in I$, for any $(\omega, \omega') \in \Omega_i \times \Omega_{i'}$ such that $\alpha_i(\omega) = \alpha_{i'}(\omega')$, we have

$$\forall s \in E_i, \quad \|f\|_{i,\omega} = \|f\|_{i',\omega'}.$$

Note that (ii) implies that for any $i \in I$, $\dim_{K_i}(E_i)$ does not depend on i. We denote this integer by $\dim(\overline{\mathbf{E}})$ and call it the *dimension* of \mathbf{E} .

Example 6.4.2. We consider the totally ordered set $\mathbb{R}_{>0}$ equipped with the usual total ordering. Consider the family of topological adelic curves $\mathbf{S} = (S_R)_{R>0}$ constructed in Example 3.1.2 (2). Let $\overline{E} = (E, \|\cdot\|)$ be a finite-dimensional complex normed vector space. Let R > 0. In §6.3, we have constructed the induced adelic vector bundle $\overline{E_R} = (E_R, \xi_R)$ on S_R by \overline{E} (cf. Definition 6.3.2). By construction, the collection $(\overline{E_R})_{R>0}$ is an adelic vector bundle on \mathbf{S} .

Proposition-Definition 6.4.3. Let (I, \leq_I) be a totally ordered set. Let $\mathbf{S} = (S_i = (K_i, \phi_i : \Omega_i \to M_{K_i}, \nu_i))_{i \in I}$ be a family of topological adelic curves with base space Ω_0 and family of structural morphisms $(\alpha_i : \Omega_i \to \Omega_0)_{i \in I}$. Let $\overline{\mathbf{E}} = (\overline{E_i} = (E_i, \xi_i = (\| \cdot \|_{i,\omega})_{\omega \in \Omega_i}))_{i \in I}$ be an adelic vector bundle on \mathbf{S} . Denote

$$K_{\infty} := \lim_{i \in I} K_i = \bigcap_{i \in I} K_i, \quad E_{\infty} := \lim_{i \in I} E_i = \bigcap_{i \in I} E_i$$

(1) Let $F_{\infty} \subset E_{\infty}$ be any vector subspace. For any $i \in I$, let $F_i := F_{\infty} \otimes_{K_{\infty}} K_i \subset E_i$ and denote by ξ_{i,F_i} the restriction of ξ_i to F_i . Then $\overline{\mathbf{F}} := (\overline{F_i} = (F_i, \xi_{i,F_i}))_{i \in I}$ is an adelic vector bundle on \mathbf{S} .

- (2) $\overline{\mathbf{E}}^{\vee} := (\overline{E_i}^{\vee} = (E_i^{\vee}, \xi_i^{\vee}))_{i \in I}$ is a lsc adelic vector bundle on **S**. Moreover, if for any $i \in I, \xi_i^{\vee \vee}$ is continuous, then $\overline{\mathbf{E}}^{\vee}$ is an adelic vector bundle on S. In particular, if for any $i \in I, \xi_i$ is ultrametric on $\Omega_{i,\text{um}}$, then $\overline{\mathbf{E}}^{\vee}$ is an adelic vector bundle on **S**.
- (3) Let $\pi : E_{\infty} \to G_{\infty}$ be a quotient vector space of E_{∞} . For any $i \in I$, let $G_i := G_{\infty} \otimes_{K_{\infty}} K_i$, it is a quotient vector space of E_i and let ξ_{i,G_i} denote the quotient pseudo-norm family on G_i induced by ξ_i . Then $\overline{\mathbf{G}} := (\overline{G_i} = (G_i, \xi_{i,G_i}))_{i \in I}$ is a usc adelic vector bundle on \mathbf{S} . Moreover, if for any $i \in I$, ξ_i is ultrametric on $\Omega_{i,\mathrm{um}}$, then $\overline{\mathbf{G}}$ is an adelic vector bundle on \mathbf{S} .
- (4) Let $n \ge 1$ be an integer. For any j = 1, ..., n, let $\mathbf{E}^{(\mathbf{j})} = (\overline{E_i^{(j)}} = (E_i^{(j)}, \xi_i^{(j)}))_{i \in I}$ be an adelic vector bundle on **S**. We assume that, for any $i \in I$ the pseudo-norm families $\xi_i^{(1)}, ..., \xi_i^{(n)}$ are ultrametric on $\Omega_{i, \text{um}}$.
 - (4.a) The ϵ -tensor product $\overline{\mathbf{E}^{(1)}} \otimes_{\epsilon} \cdots \otimes_{\epsilon} \overline{\mathbf{E}^{(n)}} := (\overline{E_i^{(1)}} \otimes_{\epsilon} \cdots \otimes_{\epsilon} \overline{E_i^{(n)}})_{i \in I}$ is a measurable adelic vector bundle on \mathbf{S} .
 - (4.b) The (ϵ, π) -tensor product $\overline{\mathbf{E}^{(1)}} \otimes_{\epsilon,\pi} \cdots \otimes_{\epsilon,\pi} \overline{\mathbf{E}^{(n)}} := (\overline{E_i^{(1)}} \otimes_{\epsilon,\pi} \cdots \otimes_{\epsilon,\pi} \overline{E_i^{(n)}})_{i \in I}$ is a measurable adelic vector bundle on \mathbf{S} .
- (5) $\det(\overline{\mathbf{E}}) := (\det(\overline{E_i}))_{i \in I}$ is a measurable adelic vector bundle on **S**. Moreover, if for any $i \in I$, ξ_i is Hermitian, then $\det(\overline{\mathbf{E}})$ is an adelic vector bundle on **S**.

Proof. It follows from Proposition 6.1.4 combined with the fact that all the involved algebraic constructions are compatible with the extension of scalars. \Box

Definition 6.4.4. Let (I, \leq_I) be a totally ordered set. Let $\mathbf{S} = (S_i = (K_i, \phi_i : \Omega_i \rightarrow M_{K_i}, \nu_i))_{i \in I}$ be a family of topological adelic curves with base space Ω_0 and family of structural morphisms $(\alpha_i : \Omega_i \rightarrow \Omega_0)_{i \in I}$. Let $\overline{\mathbf{E}} = (\overline{E_i} = (E_i, \xi_i = (\| \cdot \|_{i,\omega})_{\omega \in \Omega_i}))_{i \in I}$ be a measurable adelic vector bundle on \mathbf{S} . Denote

$$K_{\infty} := \lim_{i \in I} K_i = \bigcap_{i \in I} K_i, \quad E_{\infty} := \lim_{i \in I} E_i = \bigcap_{i \in I} E_i.$$

For any $s \in E_{\infty} \setminus \{0\}$, define

$$\widehat{\operatorname{deg}}_{\overline{\mathbf{E}}}(s) : \left| \begin{array}{ccc} I & \longrightarrow & \mathbb{R} \\ i & \longmapsto & \widehat{\operatorname{deg}}_{\overline{E_i}}(s) \end{array} \right|$$

The function $\overline{\deg}_{\overline{\mathbf{E}}}(s)$ is called the Arakelov degree of s w.r.t. $\overline{\mathbf{E}}$.

Let ~ be an equivalence relation on $\mathcal{F}(I, \mathbb{R})$ which is compatible with the additive group structure. Assume that **S** is asymptotically proper w.r.t. ~ (cf. Definition 3.2.1). Assume that $\overline{\mathbf{E}}$ is an adelic vector bundle on **S**. Let $\eta, \eta' \in \det(E_{\infty}) \setminus \{0\}$. Then

$$\widehat{\operatorname{deg}}_{\operatorname{det}(\overline{\mathbf{E}})}(\eta) - \widehat{\operatorname{deg}}_{\operatorname{det}(\overline{\mathbf{E}})}(\eta') \sim 0.$$

Therefore the class of $\widehat{\deg}_{\det(\overline{\mathbf{E}})}(\eta)$ in $\mathcal{F}(I,\mathbb{R})/\sim$ yields a well-defined element denoted by $\widehat{\deg}(\overline{\mathbf{E}})$ called the Arakelov degree of $\overline{\mathbf{E}}$.

Example 6.4.5. We consider the totally ordered set $\mathbb{R}_{>0}$ equipped with the usual total ordering. Consider the family of topological adelic curves $\mathbf{S} = (S_R)_{R>0}$ constructed in Example 3.1.2 (2). Let $\overline{E} = (E, \|\cdot\|)$ be a complex normed vector space of dimension 1.

We consider the adelic vector bundle $\overline{\mathbf{E}} = (E_R, \xi_R)_{R>0}$ constructed in Example 6.4.2. Let $e \in E \setminus \{0\}$ and let $f \in K_{\infty} = \mathcal{M}(\mathbb{C})^{\times}$. Then

$$\forall R > 0, \quad \widehat{\operatorname{deg}}_{\overline{\mathbf{E}}}(f \cdot e)(R) = T(R, f) - \ln \|e\|$$

(cf. §A for the notation T(R, f)).

7. Slopes of adelic vector bundles: proper case

In this section, we introduce slope theory for adelic vector bundles on a fixed proper topological adelic curve adelic curve $S = (K, \phi : \Omega \to M_K, \nu)$. Most of the results can be directly adapted from ([CM19], §4.3).

7.1. Degree, positive degree.

Proposition 7.1.1. Let $\overline{E} = (E, \xi)$ be an adelic vector bundle on S. The following assertions hold.

(1) Assume that \overline{E} is Hermitian. Then we have the equality

$$\widehat{\operatorname{deg}}(E,\xi) = -\widehat{\operatorname{deg}}(E^{\vee},\xi^{\vee}).$$

(2) In general, we have the inequality

$$0 \le \widehat{\deg}(E,\xi) + \widehat{\deg}(E^{\vee},\xi^{\vee}) \le \frac{1}{2} \dim_K(E) \log \dim_K(E) \nu(\Omega_{\mathrm{ar}}).$$

(3) Let $\overline{E'} = (E', \xi')$ be another adelic vector bundle on S. Assume that the double-dual pseudo-norm families $\xi^{\vee\vee}, \xi'^{\vee\vee}$ are continuous (e.g. if ξ and ξ' are ultrametric on $\Omega_{\rm um}$). Then we have the equality

$$\widehat{\operatorname{deg}}(\overline{E} \otimes_{\epsilon,\pi} \overline{E'}) = \dim_K(E')\widehat{\operatorname{deg}}(\overline{E}) + \dim_K(E)\widehat{\operatorname{deg}}(\overline{E'}).$$

(4) Let

$$0 = E_0 \subset E_1 \subset \cdots \subset E_n = E$$

be a flag of vector subspaces of E. For any i = 1, ..., n, denote by ξ_i the restriction of ξ to E_i and by η_i the quotient pseudo-norm family induced by ξ_i on E_i/E_{i-1} . Then we have the inequality

$$\sum_{i=1}^{n} \widehat{\deg}(E_i/E_{i-1},\eta_i) \le \widehat{\deg}(E,\xi).$$

Moreover, if ξ is Hermitian, then the above inequality is an equality.

Proof. (1) and (2) are respectively the counterpart of ([CM19], Propositions 4.3.8 and 4.3.10).

First remark that (3) is true if \overline{E} and $\overline{E'}$ are adelic line bundles on S. The general case follows from the fact that the canonical isomorphism

$$\det(E)^{\otimes \dim_K(E')} \otimes_K \det(E')^{\dim_K(E)} \cong \det(E \otimes_K E')$$

yields an identification

$$\det(\xi)^{\otimes \dim_K(E')} \otimes_{\epsilon,\pi} \det(\xi')^{\otimes \dim_K(E)} = \det(\xi \otimes_{\epsilon,\pi} \xi')$$

of pseudo-norm families.

(4) is an analogue of ([CM19], Proposition 4.1.13).

42

Definition 7.1.2. Let $\overline{E} = (E, \xi)$ be an adelic vector bundle on S. Let $F \subset E$ be any vector subspace and denote by ξ_F the restriction of ξ to F. Then Proposition 6.1.4 (1) implies that $\overline{F} := (F, \xi_F)$ is an adelic vector bundle on S. We define the *positive degree* of \overline{E} as

$$\widehat{\operatorname{deg}}_+(\overline{E}) := \sup_{F \subseteq E} \widehat{\operatorname{deg}}(\overline{F}),$$

where F runs over the set of all vector subspaces of E.

Remark 7.1.3. The positive degree of an adelic vector bundle plays the role of the number of "small sections" in the classical framework of Arakelov geometry over number fields.

7.2. Slopes.

Definition 7.2.1. Let $\overline{E} = (E, \xi)$ be a measurable adelic vector bundle on S. Assume that $E \neq \{0\}$.

(1) We define the *slope* of \overline{E} as

$$\widehat{\mu}(\overline{E}) := \frac{\widehat{\deg}(\overline{E})}{\dim_K(E)}.$$

(2) We define the maximal slope of \overline{E} as

$$\widehat{\mu}_{\max}(\overline{E}) := \sup_{\{0\} \neq F \subseteq E} \widehat{\mu}(\overline{F}),$$

where F runs over the set of non-zero vector subspaces of E.

(3) Assume that E is an adelic vector bundle on S. Let $E \twoheadrightarrow G$ be a quotient vector space of E. Denote by ξ_G the quotient pseudo-norm family on G induced by ξ and let $\overline{G} := (G, \xi_G)$. Proposition 6.1.4 (2) implies that \overline{G} is a lsc adelic vector bundle on S. We define the *minimal slope* of \overline{E} as

$$\widehat{\mu}_{\min}(\overline{E}) := \inf_{E \to G \neq \{0\}} \widehat{\mu}(\overline{G}),$$

where G runs over the set of non-zero quotient vector spaces of E.

7.3. Harder-Narasimhan filtration: Hermitian case.

Proposition 7.3.1. Let $\overline{E} = (E, \xi)$ be a non-zero Hermitian measurable adelic vector bundle. Let $\Theta(E)$ denote the set of all pairs of vector subspaces of E. For any $(E_1, E_2) \in \Theta(E)$, for i = 1, 2, we denote by $\overline{E_i} = (E_i, \xi_i)$, where ξ_i denotes the restriction of ξ to E_i , this is an adelic vector bundle on S.

(1) For any $(E_1, E_2) \in \Theta(E)$, we have the inequality

$$\widehat{\deg}(\overline{E_1 \cap E_2}) + \widehat{\deg}(\overline{E_1 + E_2}) \le \widehat{\deg}(\overline{E_1}) + \widehat{\deg}(\overline{E_2}).$$

(2) There exists a unique non-zero vector subspace $E_{des} \subset E$ such that

$$\widehat{\mu}(\overline{E_{\text{des}}}) = \widehat{\mu}_{\max}(\overline{E}).$$

(3) We have

$$\widehat{\mu}_{\max}(\overline{E}) < +\infty, \quad \widehat{\mu}_{\min}(\overline{E}) > -\infty.$$

Proof. (1) is a direct adaptation of ([CM19], Corollary 1.2.52).

(2) is a consequence of ([CM19], Proposition 4.3.28)

Finally, (2) implies that $\widehat{\mu}_{\max}(\overline{E}) < +\infty$. Moreover, The inequality $\widehat{\mu}_{\min}(\overline{E}) > -\infty$ follows from the inequality $0 \leq \widehat{\mu}_{\max}(\overline{E}) + \widehat{\mu}_{\min}(\overline{E})$.

Definition 7.3.2. Let $\overline{E} = (E, \xi)$ be a non-zero Hermitian measurable adelic vector bundle. The vector subspace $E_{\text{des}} \subset E$ constructed in Proposition 7.3.1 (2) is called the *destabilising* vector subspace of \overline{E} . If $E_{\text{des}} = E$, then \overline{E} is called *semistable*.

Let $\overline{E} = (E,\xi)$ be a non-zero Hermitian adelic vector bundle on S. Let $E_1 \subset E_2$ be vector subspaces of E. For i = 1, 2, we denote by ξ_i the restriction of ξ to E_i . We denote η by η the quotient pseudo-norm family induced by ξ_2 on E_2/E_1 and call it the *subquotient pseudo-norm* family induced by ξ on E_2/E_1 . We also denote $\overline{E_2/E_1} = (E_2/E_1, \eta)$. Note that $\overline{E_2/E_1}$ is a Hermitian lsc adelic vector bundle on S. Therefore, it makes sense to study the semistability of $\overline{E_2/E_1}$.

Using the construction of Proposition 7.3.1(2), we can construct inductively a flag

$$0 = E_0 \subsetneq E_1 \subsetneq \cdots \subsetneq E_n = E,$$

called the Harder-Narasimhan flag of \overline{E} , such that, for any i = 1, ..., n, $\overline{E_i/E_{i-1}}$ is semistable.

Proposition 7.3.3. We use the same notation as above. For any i = 1, ..., n, let $\mu_i := \hat{\mu}(\overline{E_i/E_{i-1}})$. Then we have

$$\mu_1 > \cdots > \mu_n.$$

Proof. This is an adaptation of ([CM19], Proposition 4.3.37) in our setting: all the ingredients of the proof have a counterpart.

7.4. Harder-Narasimhan filtration: general case. In the recent work [CJ23], Chen and Jeannin developed a very general framework for proving the existence of Harder-Narasimhan filtrations which is inspired by game theory.

7.4.1. Preliminaries. Recall that a poset $(\mathcal{L}, \leq_{\mathcal{L}})$ is a bounded lattice if \mathcal{L} possesses a greatest element \top together with a lowest element \perp and such that any finite subset of \mathcal{L} admits a supremum and an infimum w.r.t. $\leq_{\mathcal{L}}$. If $(\mathcal{L}, \leq_{\mathcal{L}})$ is a complete lattice, we define

$$\mathcal{P}_{<}(\mathcal{L}) := \{ (a, b) \in \mathcal{L}^2 : a <_{\mathcal{L}} b \}.$$

For any $(a, b) \in \mathcal{P}_{<}(\mathcal{L})$, we denote

$$\mathcal{L}_{[a,b]} := \{ c \in \mathcal{L} : a \leq_{\mathcal{L}} c \leq_{\mathcal{L}} b \}, \quad \mathcal{L}_{]a,b]} := \mathcal{L}_{[a,b]} \smallsetminus \{a\}, \quad \mathcal{L}_{[a,b[} := \mathcal{L}_{[a,b]} \smallsetminus \{b\}.$$

Recall that a bounded lattice (S, \leq) is called a *complete lattice* if every subset of S admits a supremum and an infimum w.r.t. \leq .

The setting we consider is as follows. We consider a bounded lattice $(\mathcal{L}, \leq_{\mathcal{L}})$, a complete lattice (S, \leq) together with a *pay-off* function

$$\mu: \mathcal{P}_{<}(\mathcal{L}) \to S.$$

This data is called a Harder-Narasimhan game on $(\mathcal{L}, \leq_{\mathcal{L}})$ with pay-off function μ . In that case, we define the optimal pay-off threshold

$$\mu_A^* := \inf_{x \in \mathcal{L} \setminus \{\top\}} \sup_{y \in \mathcal{L}_{]x,\top]}} \mu(x, y).$$

We say that the Harder-Narasimhan game is *semi-stable* if

 $\forall y \in \mathcal{L} \smallsetminus \{\bot\}, \quad \mu_A(\bot, y) \neq \mu_A(\bot, \top).$

Let $(x, y) \in \mathcal{P}_{\leq}(\mathcal{L})$. Then the restriction $\mu_{[x,y]}$ of μ to $\mathcal{P}_{\leq}(\mathcal{L}_{[x,y]})$ yields the pay-off function of a Harder-Narasimhan game called the *restriction* of the Narasimhan game to $\mathcal{L}_{[x,y]}$. We denote by $\mu_A(x, y)$ the optimal pay-off threshold of this restricted Harder-Narasimhan game. We now introduce Harder-Narasimhan filtrations in this general setting.

Theorem 7.4.1 ([CJ23], Theorem 1.1). Let $(\mathcal{L}, \leq_{\mathcal{L}})$ be a bounded lattice and let a (S, \leq) be a complete lattice together with a pay-off function

$$\mu: \mathcal{P}_{<}(\mathcal{L}) \to S.$$

We make the following assumptions.

- (i) (S, \leq) is totally ordered.
- (ii) The pay-off function μ is convex, namely for any $(x,y) \in \mathcal{L}^2$ such that $x \not\leq_{\mathcal{L}} y$ we have the inequality

$$\mu(x \wedge y, x) \le \mu(y, x \lor y).$$

(iii) $(\mathcal{L}, \leq_{\mathcal{L}})$ satisfies the ascending chain condition, namely, there does not exist a family $(x_n)_{n\in\mathbb{N}}$ in \mathcal{L} such that

$$x_0 <_{\mathcal{L}} x_1 <_{\mathcal{L}} \cdots <_{\mathcal{L}} x_n <_{\mathcal{L}} x_{n+1} <_{\mathcal{L}} \cdots$$

(iv) $(\mathcal{L}, \leq_{\mathcal{L}})$ satisfies the μ_A -descending chain condition, namely, for any $a \in \mathcal{L}$, there does not exist a family $(x_n)_{n\in\mathbb{N}}$ in \mathcal{L} such that

$$x_0 >_{\mathcal{L}} x_1 >_{\mathcal{L}} \cdots >_{\mathcal{L}} x_n >_{\mathcal{L}} x_{n+1} >_{\mathcal{L}} \cdots$$

and

$$\mu_A(a, x_0) < \mu_A(a, x_1) < \dots < \mu_A(a, x_n) < \mu_A(a, x_{n+1}) < \dots$$

Then there exists a unique increasing sequence

$$\perp = a_0 <_{\mathcal{L}} a_1 <_{\mathcal{L}} \cdots <_{\mathcal{L}} a_n = \top$$

such that

- (1) for any i = 1, ..., n the restricted Harder-Narasimhan game $\mathcal{L}_{[a_{i-1},a_i]}$ with pay-off function $\mu_{[x,y]}$ is semi-stable;
- (2) we have the inequalities

$$\mu_A(a_0, a_1) > \cdots > \mu_A(a_{n-1}, a_n).$$

7.4.2. Harder-Narasimhan filtration on a proper topological adelic curve. We now apply the results of the above paragraph to prove the existence of a Harder-Narasimhan filtration for (not necessarily Hermitian) adelic vector bundles on a proper topological adelic curve. Recall that we have fixed a proper topological adelic curve $S = (K, \phi : \Omega \to M_K, \nu)$.

Let $\overline{E} = (E, \xi)$ be an adelic line bundle on S. We assume that the pseudo-norm family ξ is ultrametric on Ω_{um} . We consider the set $\mathcal{L}(E)$ of vector subspaces of E, equipped with the ordering defined by the inclusion relation, it is a bounded lattice. We also consider the totally ordered set $[-\infty, +\infty]$ with the usual ordering. For any $(F', F) \in \mathcal{P}_{<}(\mathcal{L}(E))$, namely $F' \subsetneq F$ are vector subspaces of E, recall that the subquotient $\overline{F/F'}$ is an adelic vector bundle on S (cf. Proposition 6.1.4 (3)). Therefore we can define

$$\mu(F',F) := \widehat{\mu}(\overline{F/F'}) \in \mathbb{R}.$$

Then we obtain a Harder-Narasimhan game on $\mathcal{L}(E)$ with pay-off function μ . Note that, for any $(F', F) \in \mathcal{P}_{\leq}(\mathcal{L}(E))$, we have

$$\mu_A(F',F) = \widehat{\mu}_{\min}(\overline{F/F'}) \in [-\infty,+\infty].$$

Note that, for any $(F', F) \in \mathcal{P}_{<}(\mathcal{L}(E))$, the semi-stability of the Harder-Narasimhan game is equivalent to saying that, for any vector subspace $F \subset E$, we have the inequality

$$\widehat{\mu}_{\min}(\overline{F}) \le \widehat{\mu}_{\min}(\overline{E}).$$

Theorem 7.4.2. Let $\overline{E} = (E,\xi)$ be an adelic vector bundle on S. We assume that the pseudo-norm family ξ is ultrametric on Ω_{um} . Then there exists a unique flag

$$0 = E_0 \subsetneq E_1 \subsetneq \cdots \subsetneq E_n = E,$$

of \overline{E} , such that

- (1) for any i = 1, ..., n, $\overline{E_i/E_{i-1}}$ is semistable;
- (2) we have the inequalities

$$\widehat{\mu}(\overline{E_1/E_0}) > \cdots > \widehat{\mu}(\overline{E_n/E_{n-1}}).$$

Proof. We will apply Theorem 7.4.1 in our setting. Let us see that the hypotheses (i-iv) are satisfied. (i), (iii) and (iv) are trivially true. We only need to check that the pay-off function μ is convex, namely, for any vector subspaces $F, F' \subset E$ such that $F' \nsubseteq F$, we have the inequality

$$\widehat{\mu}(\overline{F/(F\cap F')}) \leq \widehat{\mu}(\overline{(F+F')/F})$$

of slopes of adelic vector bundles on S. The canonical isomorphism

$$f: F'/(F \cap F') \to (F + F')/F$$

is constructed as follows. An element $\alpha \in F'/(F \cap F')$, represented by some $x' \in F'$, is mapped to the class of x', viewed as an element of F + F', in (F + F')/F. This is independent of the choice of x'. Write $\xi = (\|\cdot\|_{\omega})_{\omega \in \Omega}$. For any $\omega \in \Omega$, denote respectively by $\|\cdot\|_{\omega,1}$ and $\|\cdot\|_{\omega,2}$ the subquotient pseudo-norm induced by $\|\cdot\|_{\omega}$ on $F'/(F \cap F')$ and (F + F')/F. Let $\alpha \in F'/(F \cap F')$. For any representative $x' \in F'$ of α , by construction of f, we have the inequality

$$\|f(\alpha)\|_{\omega,2} \le \|x'\|_{\omega}.$$

As x' is arbitrary, we obtain

$$\forall \alpha \in F'/(F \cap F'), \quad \|f(\alpha)\|_{\omega,2} \le \|\alpha\|_{\omega,1}.$$

Therefore, we have

$$\widehat{\operatorname{deg}}(\overline{F'/(F\cap F')}) \le \widehat{\operatorname{deg}}(\overline{(F+F')/F})$$

We can conclude by using the fact that

$$\dim_{K}(F'/(F \cap F')) = \dim_{K}(F') - \dim_{K}(F \cap F'))$$

=
$$\dim_{K}(F + F') - \dim_{K}(F) = \dim_{K}((F + F')/F).$$

Definition 7.4.3. Let $\overline{E} = (E, \xi)$ be an adelic line bundle on S. We assume that the pseudo-norm family ξ is ultrametric on Ω_{um} . Then the flag constructed in Theorem 7.4.2 is called the *Harder-Narasimhan flag* of \overline{E} .

Remark 7.4.4. In the classical theory of adelic curve, to obtain an analogue finiteness result to Proposition 7.3.1 (3), Chen and Moriwaki crucially use ([CM19], Theorem 4.1.26) to approximate general adelic norm families by Hermitian ones. Unfortunately, in our context, it is not quite clear how to adapt such a result. We insist on the fact that, for now, we do not have a finiteness result for maximal and minimal slopes for non-Hermitian adelic vector bundles on S.

Part 3. Arithmetic varieties over topological adelic curves: adelic line bundles and heights

In this final part, we study the higher dimensional analogue of the previous one. We start by studying the local aspects of the theory (\$8). Then we globalise these local ingredients (\$9). We finish by constructing global height functions (\$10).

8. PSEUDO-METRICS: LOCAL CASE

In this section, we introduce the higher dimensional analogue of pseudo-absolute values and pseudo-norms. Roughly speaking, this is done by considering models over the finiteness ring of a pseudo-valued field and by using the classical theory of Berkovich spaces on the special fibre of the model. We extend the usual notions of Fubini-Study metrics and semi-positive metrics to the pseudo-metric framework.

Throughout this section, we fix a field K equipped with a pseudo-absolute value $v = (|\cdot|, A, \mathfrak{m}, \kappa) \in M_K$. Recall that we denote by $\hat{\kappa}$ the completed residue field of v.

8.1. **Pseudo-metrics.** We first recall some notions from ([Séd24a], §8.1). Fix a field K equipped with a pseudo-absolute value $v = (|\cdot|, A, \mathfrak{m}, \kappa) \in M_K$. Recall that we denote by $\hat{\kappa}$ the completed residue field of v. Fix a projective K-scheme X and a projective model \mathcal{X} of X/A. The completed special fibre of \mathcal{X} is defined by $\widehat{\mathcal{X}}_s := \mathcal{X} \otimes_A \hat{\kappa}$. Then the (local) model analytic space associated with \mathcal{X} is the Berkovich analytification $\widehat{\mathcal{X}}_s^{\mathrm{an}}$.

Let *L* be a line bundle on *X*. A *(local) pseudo-metric* on *L* is the data $((\mathcal{X}, \mathcal{L}), \varphi)$, where $(\mathcal{X}, \mathcal{L})$ is a projective model of (X, L) over *A* and φ is a metric on the pullback of \mathcal{L} to $\widehat{\mathcal{X}}_s$. Such a pseudo-metric (\mathcal{L}, φ) is respectively called *lsc*, *usc*, *continuous* if the metric φ is lsc,

usc, continuous. When there is no possible confusion on the model \mathcal{X} , we allow ourselves to denote the pseudo-metric by (\mathcal{L}, φ) .

Proposition-Definition 8.1.1 ([Séd24a], §8.1). Assume that X is geometrically reduced if A = K. Let L be a line bundle on X and $((\mathcal{X}, \mathcal{L}), \varphi)$ be a pseudo-metric on L, where \mathcal{X} is a flat coherent model of X/A. Then φ induces a pseudo-norm $\|\cdot\|_{(\mathcal{L},\varphi)}$ on $H^0(X, L)$ with finiteness module $H^0(\mathcal{X}, \mathcal{L})$, kernel $\mathfrak{m}H^0(\mathcal{X}, \mathcal{L})$ and residue norm $\|\cdot\|_{\varphi}$, namely the supnorm over $\widehat{\mathcal{X}}_s^{\mathrm{an}}$ induced by the metric φ . When no confusion may arise, we simply denote $\|\cdot\|_{(\mathcal{L},\varphi)}$ by $\|\cdot\|_{\varphi}$.

From now on and until the end of this subsection, we fix a projective K-scheme X and a projective model \mathcal{X} of X over A. All pseudo-metrics are considered w.r.t. the model \mathcal{X} .

Notation 8.1.2. Let *L* be a line bundle on *X* and let $(\mathcal{X}, \mathcal{L})$ be a model of (X, L). Then we denote by L_v the pullback of \mathcal{L} to the completed special fibre $X_v := \mathcal{X} \otimes_A \hat{\kappa}$.

- **Proposition-Definition 8.1.3.** (1) Let *L* be a line bundle on *X*. Let (\mathcal{L}, φ) be a pseudometric on *L*. Then $(-\mathcal{L}, -\varphi)$ is a pseudo-metric on -L. Moreover, if (\mathcal{L}, φ) is respectively continuous/usc/lsc, then $(-\mathcal{L}, -\varphi)$ is continuous/lsc/usc.
 - (2) Let L, L' be two line bundles on X. Let $(\mathcal{L}, \varphi), (\mathcal{L}', \varphi')$ be pseudo-metrics on L, L'respectively. Then $(\mathcal{L}+\mathcal{L}', \varphi+\varphi')$. Moreover, if (\mathcal{L}, φ) and (\mathcal{L}', φ') are both respectively continuous/usc/lsc, then $(\mathcal{L}+\mathcal{L}', \varphi+\varphi')$ is continuous/usc/lsc.
 - (3) Let $f: Y \to X$ be a projective morphism of K-schemes. Let $g: \mathcal{Y}$ be a projective model of Y over A and let $g: \mathcal{Y} \to \mathcal{X}$ be a projective morphism such that g extends f. Then we have a commutative diagram with Cartesian squares



Let L be a line bundle on X. Let (\mathcal{L}, φ) be a pseudo-metric on L. Then $(g^*\mathcal{L}, \tilde{g}^*\varphi)$ is a pseudo-metric on L. By abuse of notation, we denote this pseudo-metric by $g^*(\mathcal{L}, \varphi)$ and $\tilde{g}^*\varphi$ by $g^*\varphi$. $g^*(\mathcal{L}, \varphi)$ is called the pullback of (\mathcal{L}, φ) w.r.t. the morphism $g: \mathcal{Y} \to \mathcal{X}$. Moreover, $g^*(\mathcal{L}, \varphi)$ is continuous/usc/lsc if φ is continuous/usc/lsc.

- (4) Let K'/K be a field extension and let $v' = (|\cdot|', A', \mathfrak{m}', \kappa')$ be a pseudo-absolute value on L such that v' extends v. Consider the fibre products $f : X' := X \otimes_K K' \to X$ and $g : \mathcal{X}' := \mathcal{X} \otimes_A A'$.
 - (i) Then \mathcal{X}' is a projective model of X' over A'. Moreover, \mathcal{X}' is flat/coherent if \mathcal{X} is flat/coherent. Moreover, we have a natural isomorphism $\mathcal{X}'_s \cong \mathcal{X} \otimes_{\kappa} \kappa'$.
 - (ii) Let (L, φ) be a pseudo-metric on L. Let L', L' denote the pullbacks of L, L to X', X' respectively. Then (X', L') is a model of (X', L') and denote by φ' the metric on L'_{v'} induced by φ on L_v. Then f*(L, φ) := (L', φ') defines a pseudo-metric on L' which is called the extension of (L, φ) w.r.t. the field extension K'/K.

Proof. (1) Since $(\mathcal{X}, \mathcal{L})$ is a model of (X, L), we deduce that $(\mathcal{X}, \mathcal{L}^{\vee})$ is a model of (X, L^{\vee}) . Moreover, φ^{\vee} defines a metric on L_v^{\vee} which is respectively continuous/lsc/usc if φ is continuous/usc/lsc.

(2) It is clear that $(\mathcal{X}, \mathcal{L} + \mathcal{L}')$ is a model of (X, L + L'). The assertion about the regularity of $(\mathcal{L} + \mathcal{L}', \varphi + \varphi')$ follows from the definition of the sum of two metrics on a Berkovich space.

(3) We first justify that the squares are Cartesian. For the first one, we have isomorphisms

$$\mathcal{Y} \times_{\mathcal{X}} X \cong \mathcal{Y} \times_{\mathcal{X}} (\mathcal{X} \otimes_A K) \cong \mathcal{Y} \otimes_A K \cong Y.$$

For the second one, we have isomorphisms

$$\mathcal{Y}_s \cong \mathcal{Y} \otimes_A \kappa \cong \mathcal{Y} \times_{\mathcal{X}} (\mathcal{X} \otimes_A \kappa) \cong \mathcal{Y} \times_{\mathcal{X}} \mathcal{X}_s.$$

Finally, the assertion about pseudo-metrics is clear from the properties of usual metrics on Berkovich spaces.

(4.i) We have isomorphisms

$$\mathcal{X}' \otimes_{A'} K' \cong (\mathcal{X} \otimes_A A') \otimes_{A'} K' \cong \mathcal{X} \otimes_A K' \cong (\mathcal{X} \otimes_A K) \otimes_K K' \cong X',$$

hence \mathcal{X}' is a model of X' over A. Moreover, \mathcal{X}' is a projective A'-scheme. Since being flat and of finite presentation is preserved by base change, \mathcal{X}' is flat/coherent if \mathcal{X} is flat/coherent. We now justify the assertion about the Cartesian squares. Finally, we have isomorphisms

$$\mathcal{X}_s \otimes_\kappa \kappa' = (\mathcal{X} \otimes_A \kappa) \otimes_\kappa \kappa' \cong \mathcal{X} \otimes_A \kappa' \cong (\mathcal{X} \otimes_A A') \otimes_{A'} \kappa' \cong \mathcal{X}' \otimes_{A'} \kappa' = \mathcal{X}'_s.$$

(4.ii) This is clear using (4.i) and the properties of usual metrics on Berkovich Spaces. \Box

Remark 8.1.4. Let $f: Y \to X$ be a projective morphism of K-schemes. Let L be a line bundle on X equipped with a pseudo-metric (\mathcal{L}, φ) . Then the extension of (\mathcal{L}, φ) to f^*L is subject to the choice of a model \mathcal{Y} of Y over A extending f. Let \mathcal{Y}' be an arbitrary model of Y over A. Then consider the schematic closure \mathcal{Y} of the graph of f in $\mathcal{Y}' \times_{\text{Spec}(A)} \mathcal{X}$. Then we see that \mathcal{Y} is a model of Y over A such that \mathcal{Y} dominates \mathcal{Y}' and $g: \mathcal{Y} \to \mathcal{X}$ extends f.

Definition 8.1.5. Let *L* be a line bundle on *X*. Let $(\mathcal{L}, \varphi), (\mathcal{L}, \varphi')$ be two continuous pseudo-metrics on *L*. Then $(\mathcal{O}_{\mathcal{X}}, \varphi - \varphi')$ is a pseudo-metric on $\mathcal{O}_{\mathcal{X}}$. Then we define the *distance* between φ and φ' as

$$d_v(\varphi,\varphi') := \sup_{x \in X_v^{\mathrm{an}}} |\log| \cdot |_{\varphi}(x) - \log| \cdot |_{\varphi'}(x)|,$$

where, for all $x \in X_v^{\text{an}}$, $\log |\cdot|_{\varphi}(x) - \log |\cdot|_{\varphi'}(x) := \log |\ell|_{\varphi}(x) - \log |\ell|_{\varphi'}(x)$ for some $\ell \in L(x)$ (this value does not depend on the choice of ℓ).

8.2. Fubini-Study pseudo-metric and Fubini-Study operator. The following construction is a crucial example of pseudo-metrics. Let E be a finite-dimensional K-vector space. Let $(\|\cdot\|, \mathcal{E}, N, \hat{E})$ be a pseudo-norm on E. Let $x \in X_v^{\text{an}}$, we denote by $|\cdot|_{\overline{E}}(x)$

- the ϵ -extension of scalars of the residue norm induced by $\|\cdot\|$ on $\widehat{E} \otimes_{\widehat{\kappa}} \widehat{\kappa}(x)$ if v is non-Archimedean;
- the π -extension of scalars of the residue norm induced by $\|\cdot\|$ on $\widehat{E} \otimes_{\widehat{\kappa}} \widehat{\kappa}(x)$ if v is Archimedean.

Consider the projective bundle $\mathbb{P}(\mathcal{E}) \to \operatorname{Spec}(A)$ associated with \mathcal{E} and denote by $\mathcal{O}_{\mathcal{E}}(1)$ the universal line bundle on $\mathbb{P}(\mathcal{E})$. Likewise, we have the projective bundles $\mathbb{P}(E) \to \operatorname{Spec}(K)$ and $\mathbb{P}(\widehat{E}) \to \operatorname{Spec}(\widehat{\kappa})$ for which we denote respectively by $\mathcal{O}_E(1)$ and $\mathcal{O}_{\widehat{E}}(1)$ the associated

universal line bundles. Then $(\mathbb{P}(\mathcal{E}), \mathcal{O}_{\mathcal{E}}(1))$ is a model of $(\mathbb{P}(E), \mathcal{O}_{E}(1))$ over A and we have a commutative diagram whose horizontal arrows are surjective



For any $x \in X_v^{\mathrm{an}}$, we denote by $|\cdot|_{\overline{E},\mathrm{FS}}(x)$ the quotient norm on $\mathcal{O}_{\widehat{E}}(1)(x) := \mathcal{O}_{\widehat{E}}(1) \otimes_{\widehat{\kappa}} \widehat{\kappa}(x)$ induced by the norm $|\cdot|_{\overline{E}}(x)$ on $\widehat{E} \otimes_{\widehat{\kappa}} \widehat{\kappa}(x)$ constructed above. Then the family $\varphi_{\overline{E},\mathrm{FS}} := (|\cdot|_{\overline{E},\mathrm{FS}}(x))_{x \in X_v^{\mathrm{an}}}$ defines a metric on $\mathcal{O}_{\widehat{E}}(1)$. Therefore, we obtain a pseudo-metric $(\mathcal{O}_{\mathcal{E}}(1), \varphi_{\overline{E},\mathrm{FS}})$ on $\mathcal{O}_{E}(1)$.

Definition 8.2.1. We use the same notation as above. The pseudo-metric $(\mathcal{O}_{\mathcal{E}}(1), \varphi_{\overline{E}, FS})$ on $\mathcal{O}_E(1)$ is called the *Fubini-Study* pseudo-metric on $\mathcal{O}_E(1)$ associated with the locally pseudo-normed vector space \overline{E} .

Proposition 8.2.2 ([CM19], Proposition 2.2.12). We use the same notation as above. Then the Fubini-Study pseudo-metric ($\mathcal{O}_{\mathcal{E}}(1), \varphi_{\overline{E}, FS}$) is continuous.

Example 8.2.3. Let *E* be a finite-dimensional *K*-vector space and fix a basis $(e_1, ..., e_d)$ of *E*. Fix $\lambda_1, ..., \lambda_r$. Let $\|\cdot\|$ denote the diagonalisable pseudo-norm on *E* such that $(e_1, ..., e_d)$ is an orthogonal basis of $\|\cdot\|$ such that, for any i = 1, ..., d, we have $\|e_i\| = e^{-\lambda_i}$. Namely, for any $x = x_1e_1 + \cdots + x_de_d \in E$, where $x_1, ..., x_d \in K$, we have

$$||x|| = \begin{cases} \max_{i=1,\dots,d} |x_i|e^{-\lambda_i} & \text{if } v \in M_{K,\text{um}}, \\ \sqrt{\sum_{i=1}^d |x_i|^2 e^{-2\lambda_i}} & \text{if } v \in M_{K,\text{ar}}. \end{cases}$$

Let $\overline{E} := (E, \|\cdot\|)$. Then we can see $e_1, ..., e_d$ as global sections of $\mathcal{O}_E(1)$ without common zeroes and $(e_1, ..., e_d)$ is adapted to $\|\cdot\|$. Moreover, the Fubini-Study pseudo-metric $(\mathcal{O}_{\mathcal{E}}(1), \varphi)$ is given as follows (cf. [BE21], Lemma 7.17). Let τ be a local trivialisation of $\mathcal{O}_{\widehat{E}}(1)$, then for any i = 1, ..., d we consider the function $f_i := s_i/\tau$ and we have

$$-\log|\tau|_{\varphi} = \begin{cases} \max_{i=1,\dots,d} \log|f_i| + \lambda_i & \text{if } v \in M_{K,\text{um}}, \\ \frac{1}{2}\log\sqrt{\sum_{i=1}^d |f_i|^2 e^{2\lambda_i}} & \text{if } v \in M_{K,\text{ar}}. \end{cases}$$

We now consider the general situation where we have a line bundle L on the projective K-variety X. Fix a model $(\mathcal{X}, \mathcal{L})$ of (X, L). Assume that there exist a finite-dimensional K-vector space E equipped with a pseudo-norm $(\|\cdot\|, \mathcal{E}, N, \widehat{E})$ and surjective morphism of sheaves $\beta : \mathcal{E} \otimes_A \mathcal{O}_{\mathcal{X}} \to \mathcal{L}$. Denote $\overline{E} := (E, \|\cdot\|)$. Then β yields a morphism of schemes $g : \mathcal{X} \to \mathbb{P}(\mathcal{E})$ such that \mathcal{L} is isomorphic to $g^*\mathcal{O}_{\mathcal{E}}(1)$. By considering generic fibres, g induces a morphism $f : X \to \mathbb{P}(E)$ which is extended by g by construction.

Definition 8.2.4. We use the same notation as above. The pullback of the Fubini-Study pseudo-metric on $\mathcal{O}_E(1)$ by g is a continuous pseudo-metric on L which is called the *quotient* pseudo-metric induced by \overline{E} and β .

Consider the particular case where $E \subset H^0(X, L)$ is a basepoint free vector subspace. Let $\|\cdot\|$ be a diagonalisable norm on E as in Example 8.2.3. Let $\overline{E} = (E, \|\cdot\|)$. Then the quotient pseudo-metric on L induced by \overline{E} and β is called a *Fubini-Study pseudo-metric* on L.

More generally, if L is a semi-ample line bundle on X. We say that a pseudo-metric (\mathcal{L}, φ) is *Fubini-Study* if there exists an integer $n \geq 1$ such that nL is globally generated and a Fubini-Study local pseudo metric $(n\mathcal{L}, \psi)$ on nL such that $\varphi = n^{-1}\psi$. The class of Fubini-Study pseudo-metrics on L is denoted by FS(L).

We now introduce the Fubini-Study operator for pseudo-metrics.

Definition 8.2.5. We assume that the model \mathcal{X} is flat and coherent. Let L be a line bundle on X. Let (\mathcal{L}, φ) be a pseudo-metric on L. We assume that \mathcal{L} is globally generated. Thus we have a surjection $\beta : H^0(\mathcal{X}, \mathcal{L}) \otimes_A \mathcal{O}_{\mathcal{X}} \to \mathcal{L}$. Moreover, the pseudo-metric (\mathcal{L}, φ) induces a pseudo-norm $\|\cdot\|_{\varphi}$ on $H^0(X, L)$ (cf. Definition 8.1.1). Then we can use the construction of Definition 8.2.4 to produce a continuous quotient pseudo-metric $(\mathcal{L}, \varphi_{\rm FS})$ on L, it is called the *Fubini-Study pseudo-metric* associated with (\mathcal{L}, φ) .

Proposition 8.2.6 ([CM19], Proposition 2.2.23). We assume that the model \mathcal{X} is flat and coherent. Let L be a line bundle on X and let (\mathcal{L}, φ) be a quotient pseudo-metric on L. Then, for any integer $n \geq 1$, we have $(n\mathcal{L}, n\varphi) = (n\mathcal{L}, (n\varphi)_{\text{FS}})$.

8.3. Semi-positive pseudo-metrics.

Definition 8.3.1. Let *L* be a line bundle on *X*. Fix a continuous pseudo-metric (\mathcal{L}, φ) on *L*. For any integer $n \ge 1$, we denote by $(n\mathcal{L}, \varphi_n)$ the pseudo-metric $(n\mathcal{L}, (n\varphi)_{\text{FS}})$.

(1) Assume that \mathcal{L} is semi-ample. Choose an integer $n \geq 1$ such that $n\mathcal{L}$ is globally generated. Then (\mathcal{L}, φ) is called *semi-positive* if the sequence

$$\frac{d_v(nk\varphi,\varphi_{nk})}{nk}, \quad k \ge 1$$

converges to 0. Note that this definition does not depend on the choice of n.

- (2) Assume that \mathcal{L} is semi-ample. Then (\mathcal{L}, φ) is called *plurisubharmonic* (*psh* for short) if there exists a sequence of $(\mathcal{L}, \varphi_i)_{i \geq 1}$ of Fubini-Study pseudo-metrics such that the sequence of function $(\varphi \varphi_i)_{i > 1}$ converges uniformly to the null function.
- (3) Let L be a line bundle on X. Then (\mathcal{L}, φ) is called *integrable* if there exist line bundles L_1, L_2 on X equipped respectively with plurisubharmonic pseudo-metrics $(\mathcal{L}_1, \varphi_1), (\mathcal{L}_2, \varphi_2)$ such that \mathcal{L}_1 and \mathcal{L}_2 are very ample, $L = L_1 L_2$ and $\varphi = \varphi_1 \varphi_2$.
- **Remark 8.3.2.** (1) A plurisubharmonic pseudo-metric is at first glance a special case of semi-positive pseudo-metric. If $\hat{\kappa} = \mathbb{C}$, then for any semi-positive pseudo-metric $(\mathcal{L}, \varphi), \varphi$ is a plurisubharmonic metric in the usual sense (cf. [Zha95], Theorem 3.5).
 - (2) The terminology and definition of plurisubharmonic pseudo-metrics is inspired from [BE21].

Proposition 8.3.3. Let L be a line bundle on X. Let (\mathcal{L}, φ) be a continuous pseudo-metric on L. Let K'/K be a field extension and let $v' = (|\cdot|', A', \mathfrak{m}', \kappa')$ be a pseudo absolute value on K' extending v. Consider the fibre product $f : X' := X \otimes_K K' \to X$ and denote by L' the

pullback of L to X'. If (\mathcal{L}, φ) is semi-positive, then $f^*(\mathcal{L}, \varphi)$ is a semi-positive pseudo-metric on L'.

Proof. Recall that the pseudo-metric $f^*(\mathcal{L}, \varphi)$ is equal to (\mathcal{L}', φ') , where \mathcal{L}' is the pullback of \mathcal{L} to $\mathcal{X}' := \mathcal{X} \otimes_A A'$ and φ' is the pullback of φ to $\mathcal{X}'_s \otimes_{\kappa'} \hat{\kappa'}$. ([CM21], Remark 3.2.7) implies that the metric φ' is semi-positive. Therefore, the local pseudo metric (\mathcal{L}', φ') is semi-positive.

Proposition 8.3.4. Assume that the special fibre \mathcal{X}_s is geometrically reduced. Let L be a line bundle on X. Let (\mathcal{L}, φ) be a continuous pseudo-metric on L with \mathcal{L} semi-ample. Then (\mathcal{L}, φ) is psh iff is is semi-positive.

Proof. First note that it suffices to prove that if (\mathcal{L}, φ) is semi-positive, then (\mathcal{L}, φ) is psh. Thus we assume that (\mathcal{L}, φ) is semi-positive. The $\hat{\kappa} = \mathbb{C}$ is treated in Remark 8.3.2. The $\hat{\kappa} = \mathbb{R}$ is dealt with by combining Proposition 8.3.3 with ([BE21], Theorem 7.31).

The non-Archimedean and non trivially valued case follows from ([CM21], Proposition 3.2.19). For the trivially valued case, we choose a transcendental extension K'/K and a pseudo-absolute value $v' = (|\cdot|', A', \mathfrak{m}', \kappa')$ extending v such that κ' is non-trivially valued. Consider the fibre product $f : X' := X \otimes_K K' \to X$ and the pullback L' of L to X'. Then the pullback of (\mathcal{L}, φ) to X' is semi-positive by Proposition 8.3.3. Then by the non-trivially valued case, $f^*(\mathcal{L}, \varphi) =: (\mathcal{L}', \varphi')$ is psh. By ([BE21], Theorem 7.31), φ' is continuous psh metric on $L'_{v'}$ and therefore φ is psh and (\mathcal{L}, φ) is a (continuous) psh pseudo-metric on L. \Box

9. PSEUDO-METRIC FAMILIES

In this section, we introduce the global counterpart of §8. We first introduce the global analogue of pseudo-metrics: pseudo-metric families (§9.1). Then we introduce integrability and regularity conditions for pseudo-metric families (§9.2-9.3). After that, we introduce the pushforward of a pseudo-metric family (§9.4). We are now able to define the notion of adelic line bundles (§9.5). Then we introduce adelic line bundles for families of topological adelic curves (§9.6. All the previously introduced notions are studied in the case of an integral topological adelic curve (§9.7). Finally, we define the arithmetic volume and the χ -volume of adelic line bundles subject to specific conditions (§9.8).

9.1. **Definitions.** In this subsection, we fix a topological adelic curve $S = (K, \phi : \Omega \to M_K, \nu)$ and a projective K-scheme X. Recall that, for any $\omega \in \Omega$, we denote by A_{ω} and κ_{ω} , the finiteness ring and the residue field of ω respectively.

Definition 9.1.1. Let *L* be an invertible \mathcal{O}_X -module.

- (1) A model family of (X, L) is a family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$ such that, for any $\omega \in \Omega$, $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})$ is a projective model of (X, L) over A_{ω} . A model family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$ of (X, L) is called *very ample*, resp. *flat*, resp. *coherent* if, for any $\omega \in \Omega$, \mathcal{L}_{ω} is very ample, resp. \mathcal{X}_{ω} is flat, resp. \mathcal{X}_{ω} is coherent.
- (2) A pseudo-metric family φ on L is a family $((\mathcal{X}_{\omega}, \mathcal{L}_{\omega}), \varphi_{\omega})_{\omega \in \Omega}$ where, for any $\omega \in \Omega$, (i) $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$ is a model family of (X, L), which is called the *model family* of φ ; (ii) $(\mathcal{L}_{\omega}, \varphi_{\omega})$ is a pseudo-metric in ω on L.

Notation 9.1.2. Let L be an invertible \mathcal{O}_X -module.

- (1) If $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$ is a model family of (X, L), by "let $\varphi = (\varphi_{\omega})_{\omega \in \Omega}$ be pseudo-metric family with model family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$ ", we mean that, φ is a pseudo-metric family on L with model family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$ such that, for any $\omega \in \Omega$, $(\mathcal{L}_{\omega}, \varphi_{\omega})$ satisfies (ii) in Definition 9.1.1.
- (2) If $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$ is a model family of (X, L), then for any $\omega \in \Omega$, we denote respectively by X_{ω}^{an} the Berkovich analytic space $(\mathcal{X}_{\omega} \otimes_{A_{\omega}} \widehat{\kappa_{\omega}})^{\mathrm{an}}$ and by L_{ω} the pullback of \mathcal{L} to $\mathcal{X}_{\omega} \otimes_{A_{\omega}} \widehat{\kappa_{\omega}}$.

Remark 9.1.3. We consider the case where $X = \operatorname{Spec}(K')$, where K'/K is a finite extension of fields. For any $\omega \in \Omega$, let A'_{ω} be the integral closure of A_{ω} in K', it is a semi-local Prüfer domain and we denote $\mathcal{X}_{\omega} := \operatorname{Spec}(A'_{\omega})$. Let L be a line bundle on X, i.e. a one-dimensional K'-vector space. Then a pseudo-metric family φ on L with model family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})$ is the same as a pseudo-norm family ξ_L on the one-dimensional K'-vector space L, relatively to the topological adelic curve $S' := S \otimes_K K'$.

Definition 9.1.4. Let *L* be a line bundle on *X* equipped with a pseudo-metric family φ with model family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$.

- (1) We have a pseudo-metric family $-\varphi$ with model family $(\mathcal{X}_{\omega}, -\mathcal{L}_{\omega})_{\omega \in \Omega}$ on -L.
- (2) Let L' be another line bundle on X equipped with a pseudo-metric family φ' with model family $(\mathcal{X}_{\omega}, \mathcal{L}'_{\omega})_{\omega \in \Omega}$. Then we have a pseudo-metric family $\varphi + \varphi'$ with model family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega} + \mathcal{L}'_{\omega})_{\omega \in \Omega}$.
- (3) Let $f: Y \to X$ be a projective morphism of K-schemes. Assume that, for any $\omega \in \Omega$, f extends to a A_{ω} -morphism $g_{\omega}: \mathcal{Y}_{\omega} \to \mathcal{X}_{\omega}$, where \mathcal{Y}_{ω} is a model of Y over A_{ω} . Then we have a pseudo-metric family $f^*\varphi$ with model family $(\mathcal{Y}_{\omega}, g_{\omega}^* \mathcal{L}_{\omega})_{\omega \in \Omega}$ on f^*L .
- (4) Let P: Spec $(K') \to X$ be a closed point. Denote by $S' = (K', \phi' : \Omega' \to M_{K'}, \nu')$ the topological adelic curve $S \otimes_K K'$. Let $\omega' \in \Omega'$ with finiteness ring $A'_{\omega'}$, denote $\omega := \pi_{K'/K}(\omega')$. Then we have a commutative diagram



and the valuative criterion of properness yields a unique morphism $\operatorname{Spec}(A'_{\omega'}) \to \mathcal{X}_{\omega}$. The image of the closed point of $\operatorname{Spec}(A'_{\omega'})$ together with the residue absolute value of ω' furnish a point $x_{\omega'} = (P_{\omega'}, |\cdot|_{\omega'}) \in X^{\operatorname{an}}_{\omega}$, where $P_{\omega'}$ is a closed point. Then the norm $|\cdot|_{\varphi_{\omega}}(x_{\omega'})$ can be lifted to a pseudo-norm $P^*\varphi_{\omega}$ on $E' := P^*L$. Thus the family $P^*\varphi := (P^*\varphi_{\omega})_{\omega\in\Omega}$ is a pseudo-norm family on P^*L (w.r.t. the topological adelic curve S').

Example 9.1.5. A crucial example of pseudo-metric families is the so-called "quotient pseudo-metric families" which are defined as follows. Let $\overline{E} = (E, \xi)$, where E is a finitedimensional K-vector space and $\xi = (\|\cdot\|_{\omega}, \mathcal{E}_{\omega}, N_{\omega}, \widehat{E}_{\omega})_{\omega \in \Omega}$ is a pseudo-norm family on E. Let L be a line bundle on X. Fix a model family $(\mathcal{X}, \mathcal{L}) := (\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$ of (X, L). Assume that, for any $\omega \in \Omega$, there exists a surjective homomorphism $\beta_{\omega} : \mathcal{E}_{\omega} \otimes_{A_{\omega}} \mathcal{O}_{\mathcal{X}_{\omega}} \to \mathcal{L}_{\omega}$. Then, for any $\omega \in \Omega$, we consider the quotient pseudo-metric $(\mathcal{L}_{\omega}, \varphi_{\omega})$ induced by $(E, \|\cdot\|_{\omega})$ and β_{ω} (cf. Definition 8.2.4). Then the family $((\mathcal{X}_{\omega}, \mathcal{L}_{\omega}), \varphi_{\omega})_{\omega \in \Omega}$ defines a pseudo-metric family on L which is called the *quotient pseudo-metric family* induced by \overline{E} and $\beta := (\beta_{\omega})_{\omega \in \Omega}$ on L. **Proposition 9.1.6.** Let *L* be a line bundle on *X*. Let $(\mathcal{X}, \mathcal{L}) := (\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$ be a model family for *L*. Assume that there exist a finite-dimensional *K*-vector space *E* equipped with two dominated pseudo-norm families $\xi = (\|\cdot\|_{\omega}, \mathcal{E}_{\omega}, N_{\omega}, \widehat{E}_{\omega})_{\omega \in \Omega}, \xi'$ which possess a common global adapted basis, together with a family of surjective homomorphisms $\beta := (\beta_{\omega} : \mathcal{E}_{\omega} \otimes_{A_{\omega}} \mathcal{O}_{\mathcal{X}_{\omega}} \to \mathcal{L}_{\omega})_{\omega \in \Omega}$. Let φ, φ' be respectively the quotient pseudo-metric families on *L* induced by $\beta, (E, \xi)$ and $\beta, (E, \xi')$. Then the local distance function $(\omega \in \Omega) \mapsto d_{\omega}(\varphi, \varphi')$ is ν -dominated.

Proof. By ([CM19], Proposition 2.2.20), for any $\omega \in \Omega$, we have the inequality

$$d_{\omega}(\varphi,\varphi') \le d_{\omega}(\xi^{\vee\vee},\xi'^{\vee\vee}).$$

Using Corollary 5.3.8, as $\xi_1^{\vee\vee}$ and $\xi_2^{\vee\vee}$ are both ultrametric on $\Omega_{\rm um}$, the local distance function $(\omega \in \Omega) \mapsto d_{\omega}(\xi^{\vee\vee}, \xi'^{\vee\vee})$ is ν -dominated. Hence so is $(\omega \in \Omega) \mapsto d_{\omega}(\varphi, \varphi')$.

9.2. Dominated pseudo-metric family. In this subsection, we fix a topological adelic curve $S = (K, \phi : \Omega \to M_K, \nu)$ and a projective K-scheme X. Recall that, for any $\omega \in \Omega$, we denote by A_{ω} and κ_{ω} , the finiteness ring and the residue field of ω respectively.

Definition 9.2.1. Let *L* be a line bundle on *X*. Let $\varphi := (\varphi_{\omega})_{\omega \in \Omega}$ be a pseudo-metric family on *L* with model family $(\mathcal{X}, \mathcal{L}) := (\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$.

- (1) Assume that the model family $(\mathcal{X}, \mathcal{L})$ is very ample. Then φ is called *dominated*, if there exist
 - (i) a finite-dimensional K-vector space E together with a dominated pseudo-norm family $\xi = (\|\cdot\|_{\omega}, \mathcal{E}_{\omega}, N_{\omega}, \widehat{E}_{\omega})_{\omega \in \Omega}$ on E;
 - (ii) a family of surjective homomorphisms $\beta := (\beta_{\omega} : \mathcal{E}_{\omega} \otimes_{A_{\omega}} \mathcal{O}_{\mathcal{X}_{\omega}} \to \mathcal{L}_{\omega})_{\omega \in \Omega}$ which induces a family of closed immersions $g := (g_{\omega} : \mathcal{X}_{\omega} \to \mathbb{P}(\mathcal{E}_{\omega}))_{\omega \in \Omega}$;

such that the quotient metric family φ' induced by (E,ξ) and β satisfies: the local distance function $(\omega \in \Omega) \mapsto d_{\omega}(\varphi, \varphi')$ is ν -dominated.

(2) In the general case, we say that φ is *dominated* if there exist two very ample line bundles L_1, L_2 on X, respectively equipped with two dominated pseudo-metric families φ_1, φ_2 , whose underlying model families are very ample, such that $L = L_1 - L_2$ and $\varphi = \varphi_1 - \varphi_2$.

Proposition 9.2.2. Let L_1, L_2 be two very ample line bundles on X equipped respectively with pseudo-metric families $\varphi_1 := (\varphi_{1,\omega})_{\omega \in \Omega}, \varphi_2 := (\varphi_{2,\omega})_{\omega \in \Omega}$ with respective very ample model families $(\mathcal{X}_{\omega}, \mathcal{L}_{1,\omega})_{\omega \in \Omega}, (\mathcal{X}_{\omega}, \mathcal{L}_{2,\omega})_{\omega \in \Omega}$. If φ_1 and φ_2 are dominated, then $\varphi_1 + \varphi_2$ is dominated.

Proof. As φ_1, φ_2 are dominated, there exist two finite-dimensional K-vector spaces E_1, E_2 equipped respectively with dominated pseudo-norm families ξ_1, ξ_2 , together with surjective homomorphisms $\beta_1 : E_1 \otimes_K \mathcal{O}_X \to L_1, \beta_2 : E_2 \otimes_K \mathcal{O}_X \to L_2$ inducing closed immersions $\iota_1 : X \hookrightarrow \mathbb{P}(E_1), \iota_2 : X \hookrightarrow \mathbb{P}(E_2)$ such that the respective induced quotient pseudo-metric families φ'_1, φ'_2 have respective model families $(\mathcal{X}_{\omega}, \mathcal{L}_{1,\omega})_{\omega \in \Omega}, (\mathcal{X}_{\omega}, \mathcal{L}_{2,\omega})_{\omega \in \Omega}$ and the local distance functions $(\omega \in \Omega) \mapsto d_{\omega}(\varphi_1, \varphi'_1), (\omega \in \Omega) \mapsto d_{\omega}(\varphi_2, \varphi'_2)$ are ν -dominated.

Then we have a closed immersion $X \hookrightarrow \mathbb{P}(E_1 \otimes_K E_2)$ by composing $(\iota_1, \iota_2) : X \to \mathbb{P}(E_1) \times \mathbb{P}(E_2)$ with the Segre embedding $\mathbb{P}(E_1) \times \mathbb{P}(E_2) \to \mathbb{P}(E_1 \otimes_K E_2)$. Moreover, by considering the dominated pseudo-norm family $\xi_1 \otimes_{\epsilon,\pi} \xi_2$ on $E_1 \otimes_K E_2$, we see that the induced quotient pseudo-metric family on $L_1 + L_2$ is $\varphi'_1 + \varphi'_2$. As we have

$$\forall \omega \in \Omega, \quad d_{\omega}(\varphi_1 + \varphi_2, \varphi_1' + \varphi_2') \le d_{\omega}(\varphi_1, \varphi_1') + d_{\omega}(\varphi_2, \varphi_2'),$$

we obtain the desired dominance.

Remark 9.2.3. We can adapt ([CM19], Remark 2.1.10) in our context to see that (1) and (2) in Definition 9.2.1 are equivalent when the model family is very ample. Let L be a line bundle on X and fix a very ample model family $(\mathcal{X}, \mathcal{L}) := (\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$. Assume that (1) holds. Then we can write L = 2L - L and $\varphi = 2\varphi - \varphi$. As $(\mathcal{X}_{\omega}, 2\mathcal{L}_{\omega})_{\omega \in \Omega}$ is a very ample model family for 2L and 2φ is dominated by Proposition 9.2.2, (2) holds. Conversely, write $L = L_1 - L_2$ and $\varphi = \varphi_1 - \varphi_2$, where L_1, L_2 are very ample line bundles on X equipped respectively with dominated pseudo-metric families φ_1, φ_2 whose underlying model families are very ample. Let φ' be a dominated pseudo-metric family in the sense of (1). Then again Proposition 9.2.2 implies that $\varphi' + \varphi_1$ is dominated in the sense of (1). Therefore, the local distance function

$$(\omega \in \Omega) \mapsto d_{\omega}(\varphi_2, \varphi' + \varphi_1) = d_{\omega}(\varphi, \varphi')$$

is ν -dominated (cf. Proposition 9.1.6). Hence (1) holds.

Proposition 9.2.4. Let L be a line bundle on X. Let $(\mathcal{X}, \mathcal{L}) := (\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$ be a model family of (X, L). Let E be a finite-dimensional K-vector space equipped with a pseudonorm family $\xi = (\|\cdot\|_{\omega}, \mathcal{E}_{\omega}, N_{\omega}, \widehat{E_{\omega}})_{\omega \in \Omega}$. Assume that there exists a family of surjective homomorphisms $\beta := (\beta_{\omega} : \mathcal{E}_{\omega} \otimes_{A_{\omega}} \mathcal{O}_{\mathcal{X}_{\omega}} \to \mathcal{L}_{\omega})$. If ξ is dominated, then the quotient pseudo-metric family φ induced by $\overline{E} = (E, \xi)$ and β (cf. Example 9.1.5) is dominated.

Proof. Since X is projective, there exists a very ample line bundle L' such that L + L' is also very ample. Fix a very ample model family $(\mathcal{X}_{\omega}, \mathcal{L}'_{\omega})_{\omega \in \Omega}$ of (X, L') such that $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega} + \mathcal{L}'_{\omega})_{\omega \in \Omega}$ is a very ample model family of L + L'. Fix a finite-dimensional K-vector space E' equipped with a dominated pseudo-norm family $\xi' = (\|\cdot\|'_{\omega}, \mathcal{E}'_{\omega}, N'_{\omega}, \widehat{E'_{\omega}})_{\omega \in \Omega}$ and a family of surjective homomorphisms $\beta' = (\beta'_{\omega} : \mathcal{E}'_{\omega} \otimes_{A_{\omega}} \mathcal{O}_{\mathcal{X}_{\omega}} \to \mathcal{L}'_{\omega})$ which induces a family of closed immersions $g' = (g'_{\omega} : \mathcal{X}_{\omega} \hookrightarrow \mathbb{P}(\mathcal{E}'_{\omega}))_{\omega \in \Omega}.$

Denote by $g = (g_{\omega} : \mathcal{X}_{\omega} \to \mathbb{P}(\mathcal{E}_{\omega}))_{\omega \in \Omega}$ the family of morphisms defined by β . Fix $\omega \in \Omega$. Then we have a morphism $(g_{\omega}, g'_{\omega}) : \mathcal{X}_{\omega} \to \mathbb{P}(\mathcal{E}_{\omega}) \times_{\operatorname{Spec}(A_{\omega})} \mathbb{P}(\mathcal{E}'_{\omega})$. As $\mathbb{P}(\mathcal{E}_{\omega}) \times_{\operatorname{Spec}(A_{\omega})} \mathbb{P}(\mathcal{E}'_{\omega}) \to \mathbb{P}(\mathcal{E}_{\omega})$ $\mathbb{P}(\mathcal{E}')$ is separated and g'_{ω} is a closed immersion, $(g_{\omega}, g'_{\omega})$ is closed immersion ([GW10], Remark 9.11). Moreover, we have the Segre embedding $\mathbb{P}(\mathcal{E}_{\omega}) \times_{\operatorname{Spec}(A_{\omega})} \mathbb{P}(\mathcal{E}'_{\omega}) \to \mathbb{P}(\mathcal{E}_{\omega} \otimes_{A_{\omega}} \mathcal{E}'_{\omega})$, which is a closed immersion ([GW10], Proposition 8.20). Therefore the composition with $(g_{\omega}, g'_{\omega})$ yields a closed immersion $g''_{\omega} : \mathcal{X}_{\omega} \hookrightarrow \mathbb{P}(\mathcal{E}_{\omega} \otimes_{A_{\omega}} \mathcal{E}'_{\omega}).$ Therefore, the family of surjective homomorphisms $\beta'' = (\beta_{\omega} \otimes \beta'_{\omega})_{\omega \in \Omega}$ induces the family

of closed immersions $g'' = (g''_{\omega})_{\omega \in \Omega}$.

Let φ' denote the quotient pseudo-metric family on L' induced by β' and (E',ξ') . φ' is dominated by definition. Moreover, the pseudo-metric family $\varphi + \varphi'$ is the quotient pseudometric family induced by $(E \otimes_K E', \xi \otimes_{\epsilon,\pi} \xi')$ and the family of surjective homomorphisms β'' . Using Definition 9.1.4, we see that $\xi \otimes_{\epsilon,\pi} \xi'$ is dominated. Therefore $\varphi + \varphi'$ is a dominated pseudo-metric family on L + L'. Writing $(L, \varphi) = (L + L', \varphi + \varphi') - (L', \varphi')$, we obtain that the pseudo-metric family φ is dominated.

Proposition 9.2.5. Let L be a line bundle on X. Let $\varphi := (\varphi_{\omega})_{\omega \in \Omega}$ be a pseudo-metric family on L with model family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$.

- (1) If φ is dominated, then $-\varphi$ is a dominated pseudo-metric family on -L.
- (2) Let L' be another line bundle on X equipped with a pseudo-metric family φ' . If φ and φ' are dominated, then $\varphi + \varphi'$ is a dominated pseudo-metric family on L + L'.

- (3) Let φ' be another pseudo-metric family on L with model family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$. If φ' is dominated and the local distance function $(\omega \in \Omega) \mapsto d_{\omega}(\varphi, \varphi')$ is ν -dominated, then φ is dominated.
- (4) Let f: Y → X be a projective morphism of K-schemes. Assume that, for any ω ∈ Ω, f extends to a A_ω-morphism g_ω: Y_ω → X_ω, where Y_ω is a projective model of Y over A_ω. If φ is dominated, then f^{*}φ is a dominated pseudo-metric family on f^{*}L.
- (5) Let φ' be another pseudo-metric family on L with model family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$. If φ and φ' are both dominated, then the local distance function $(\omega \in \Omega) \mapsto d_{\omega}(\varphi, \varphi')$ is ν -dominated.

Proof. (1) Write $(L, \varphi) = (L_1, \varphi_1) - (L_2, \varphi_2)$ with L_1, L_2 very ample line bundles on X and φ_1, φ_2 dominated pseudo-metric families on L_1, L_2 (whose underlying model families are very ample). Then $(-L, -\varphi) = (L_2, \varphi_2) - (L_1, \varphi_1)$ and therefore $-\varphi$ is dominated.

(2) Write $(L, \varphi) = (L_1, \varphi_1) - (L_2, \varphi_2)$ with L_1, L_2 very ample line bundles on X and φ_1, φ_2 dominated pseudo-metric families on L_1, L_2 and $(L', \varphi') = (L'_1, \varphi'_1) - (L'_2, \varphi'_2)$ with L'_1, L'_2 very ample line bundles on X and φ'_1, φ'_2 dominated pseudo-metric families on L'_1, L'_2 . Then $\varphi_1 + \varphi'_1$ and $\varphi_2 + \varphi'_2$ are respectively dominated pseudo-metric families on $L_1 + L'_1$ and $L_2 + L'_2$ by Proposition 9.2.2. Hence, by writing $(L+L', \varphi+\varphi') = (L_1+L'_1, \varphi_1+\varphi'_1) - (L_2+L'_2, \varphi_2+\varphi'_2)$, we obtain that $\varphi + \varphi'$ is dominated.

(3) We first consider the very ample case. By hypothesis, there exist a finite-dimensional K-vector space E together with a dominated pseudo-norm family ξ on E and a family of surjective homomorphisms $\beta := (\beta_{\omega} : \mathcal{E}_{\omega} \otimes_{A_{\omega}} \mathcal{O}_{\mathcal{X}_{\omega}} \to \mathcal{L}_{\omega})$ such that the quotient metric family ψ induced by (E, ξ) and β satisfies: the function $(\omega \in \Omega) \mapsto d_{\omega}(\varphi', \psi)$ is ν -dominated. Then we have

$$\forall \omega \in \Omega, \quad d_{\omega}(\varphi, \psi) \le d_{\omega}(\varphi, \varphi') + d_{\omega}(\varphi', \psi).$$

Hence φ is dominated.

In the general case, we write $(L, \varphi') = (L_1, \varphi'_1) - (L_2, \varphi'_2)$ with L_1, L_2 very ample line bundles on X and φ'_1, φ'_2 dominated pseudo-metric families on L_1, L_2 (whose underlying model families are very ample). Let $\varphi_1 := \varphi + \varphi'_2$ and $\varphi_2 := \varphi'_2$. Then $\varphi = \varphi_1 - \varphi_2$ and φ_1, φ_2 are respectively pseudo-metric families on the very ample line bundles $L + L_2 = L_1$ and L_2 . Moreover, we have

$$\forall \omega \in \Omega, \quad d_{\omega}(\varphi_1, \varphi_1') = d_{\omega}(\varphi + \varphi_2', \varphi_1') = d_{\omega}(\varphi, \varphi').$$

Therefore, the local distance function $(\omega \in \Omega) \mapsto d_{\omega}(\varphi_1, \varphi'_1)$ is dominated and by the very ample case, φ_1 is dominated. As φ_2 is dominated, φ is dominated.

(4) We start with the very ample case. By hypothesis, there exist a finite-dimensional *K*-vector space *E* together with a dominated pseudo-norm family ξ on *E* and a family of surjective homomorphisms $\beta := (\beta_{\omega} : \mathcal{E}_{\omega} \otimes_{A_{\omega}} \mathcal{O}_{\mathcal{X}_{\omega}} \to \mathcal{L}_{\omega})$ such that the quotient metric family φ' induced by (E,ξ) and β satisfies: the function $(\omega \in \Omega) \mapsto d_{\omega}(\varphi,\varphi')$ is ν -dominated. Then $f^*(\beta) : E \otimes_K \mathcal{O}_Y \to f^*L$ is a surjective homomorphism and $f^*\varphi'$ is the quotient pseudo-metric family induced by (E,ξ) and $f^*(\beta)$. Note that both $f^*\varphi$ and $f^*\varphi'$ have model family $(\mathcal{Y}_{\omega}, g^*_{\omega}\mathcal{L}_{\omega})_{\omega\in\Omega}$. Proposition 9.2.4 implies that $f^*\varphi'$ is dominated. Moreover, we have

$$\forall \omega \in \Omega, \quad d_{\omega}(f^*\varphi, f^*\varphi') \le d_{\omega}(\varphi, \varphi').$$

Thus $(\omega \in \Omega) \mapsto d_{\omega}(f^*\varphi, f^*\varphi')$ is ν -dominated and (3) implies that $f^*\varphi$ is dominated.

Let us now consider the general case. Write $(L, \varphi) = (L_1, \varphi_1) - (L_2, \varphi_2)$ with L_1, L_2 very ample line bundles on X and φ_1, φ_2 dominated pseudo-metric families on L_1, L_2 (whose underlying model families are very ample). Then $f^*\varphi_1, f^*\varphi_2$ are dominated by the very ample case. Since $f^*\varphi = f^*\varphi_1 - f^*\varphi_2$, by (1) and (2) we obtain that $f^*\varphi$ is dominated.

(5) Let L_1 be a very ample line bundle on X such that $L_2 = L + L_1$ is very ample. Fix a dominated pseudo-metric family φ_1 in L_1 whose underlying model family is very ample. We may assume that the induced model family of L_2 is also very ample. Set $\varphi_2 = \varphi + \varphi_1$ and $\varphi'_2 = \varphi' + \varphi_1$. (2) implies that φ_2 and φ'_2 are both dominated. Since L_2 is very ample with a very ample model family, Proposition 9.1.6 combined with the triangle inequality imply that the local distance function

$$(\omega \in \Omega) \mapsto d_{\omega}(\varphi_2, \varphi'_2) = d_{\omega}(\varphi, \varphi')$$

is ν -dominated.

9.3. Regularity conditions for pseudo-metric families. In this subsection, we fix a topological adelic curve $S = (K, \phi : \Omega \to M_K, \nu)$ and a projective K-scheme X. Recall that, for any $\omega \in \Omega$, we denote by A_{ω} and κ_{ω} , the finiteness ring and the residue field of ω respectively.

Definition 9.3.1. Let L be a line bundle on X equipped with a pseudo-metric family $\varphi = (\varphi_{\omega})_{\omega \in \Omega}$ with model family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$. We say that the pseudo-metric family φ is respectively usc, lsc, continuous if, for any closed point P of X, the pseudo-norm family $P^*\varphi$ (cf. Definition 9.1.4 (4)) on P^*L is usc, lsc, continuous.

- **Proposition 9.3.2.** (1) Let L be a line bundle on X equipped with a respectively usc, lsc, continuous pseudo-metric family φ . Then the pseudo-metric family $-\varphi$ on -L is respectively usc, lsc, continuous.
 - (2) Let L_1, L_2 be two line bundles on X equipped with respectively usc, lsc, continuous pseudo-metric families φ_1, φ_2 respectively. Then $\varphi_1 + \varphi_2$ is a respectively a usc, lsc, continuous pseudo-metric family on $L_1 + L_2$.
 - (3) Let f: Y → X be a projective morphism of K-schemes. Assume that, for any ω ∈ Ω, f extends to a A_ω-morphism g_ω: Y_ω → X_ω, where Y_ω is a projective model of Y over A_ω. Let L be a line bundle equipped with a respectively usc, lsc, continuous pseudo-metric family φ. Then the pseudo-metric family f*φ on f*L is respectively usc, lsc, continuous.

Proof. (1) Let P be a closed point of X. Then $P^*(-\varphi) = (P^*\varphi)^{\vee}$ and the result follows from Proposition 6.1.3.

(2) Let P be a closed point of X. Then $P^*(\varphi_1 + \varphi_2) = P^*\varphi_1 \otimes P^*\varphi_2$. Then the continuity follows from the explicit description of tensor product pseudo-norm families over vector spaces of dimension 1.

(3) It is seen directly from the construction of the pullback pseudo-metric family. \Box

Proposition 9.3.3. Let $\overline{E} = (E, \xi)$ where E is a finite-dimensional K-vector space and $\xi = (\|\cdot\|_{\omega})$ is a pseudo-norm family on E which is assumed to be ultrametric on Ω_{um} . Let L be a line bundle on X and let $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$ be a model family of (X, L). Assume that, for any $\omega \in \Omega$, there exists a surjective homomorphism $\beta_{\omega} : \mathcal{E}_{\omega} \otimes_{A_{\omega}} \mathcal{O}_{\mathcal{X}_{\omega}} \to \mathcal{L}_{\omega}$. Denote by φ the corresponding quotient pseudo-metric family on L (cf. Example 9.1.5). If the pseudo-norm family ξ is continuous, then φ is a usc pseudo-metric family on L.

Proof. Let P be a closed point of X. Using Remark 9.1.3, we see that $P^*\varphi$ identifies with the quotient pseudo-norm family of the pseudo-norm family $\xi \otimes_K \kappa(P)$ via the surjective homomorphism $[\cdot]_P : E \otimes_K \kappa(P) \to P^*L$. In general, the pseudo-norm family $\xi \otimes_K \kappa(P)$ is only known to be lsc, which is not sufficient to conclude.

Write $S \otimes_K \kappa(P) = (\kappa(P), \phi_P : \Omega_P \to M_{\kappa(P)}, \nu_P)$, $P^*\varphi = (\|\cdot\|_{P,\varphi,x})_{x\in\Omega_P}$ and $\xi_L = (\|\cdot\|_{P,x}, \mathcal{E}_P, N_P, \widehat{E_{P,x}})_{x\in\Omega_P}$. For any $x \in \Omega_P$, we denote by $(A_{P,x}, \mathfrak{m}_{P,x}, \kappa_{P,x})$ the finiteness ring of x. As P^*L is a quotient vector space of dimension 1 of $E \otimes_K \kappa(P)$, for any $x \in \Omega_P$, the residue vector space of P^* in x is a quotient vector space of dimension 1 of $\widehat{E_{P,x}}$. Hence we can use ([CM19], Proposition 1.3.26) to obtain

$$\forall \ell \in P^*L, \quad \forall x \in \Omega_P, \qquad \|\ell\|_{P,\varphi,x} = \inf_{(s,\lambda) \in E \times \kappa(P)^{\times}, [s]_P = \lambda \ell} |\lambda|_x^{-1} \|s\|_{\pi_{\kappa(P)/K}(x)}$$

As for any $\lambda \in \kappa(P)$, the map $(x \in \Omega_P) \mapsto |\lambda|_x$ is continuous and, for any $s \in E$, the map $(x \in \Omega_L) \mapsto ||s||_{\pi_{\kappa(P)/K}(x)}$ is continuous, we obtain that $P^*\varphi$ is usc.

9.4. Pushforward of pseudo-metric families. This subsection is devoted to studying the behaviour of the supremum pseudo-norm family determined by a pseudo-metric family. This notion is crucial in view of developing volume functions on topological adelic curves. We fix a topological adelic curve $S = (K, \phi : \Omega \to M_K, \nu)$ and a projective K-scheme X. Recall that, for any $\omega \in \Omega$, we denote by A_{ω} and κ_{ω} , the finiteness ring and the residue field of ω respectively.

Definition 9.4.1. Assume that X is geometrically reduced. Let L be a line bundle on X. Let $\varphi = (\varphi_{\omega})_{\omega}$ be a pseudo-metric family on L with a flat and coherent model family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$. For any $\omega \in \Omega$, φ_{ω} induces a pseudo-norm $\|\cdot\|_{\varphi_{\omega}}$ on $H^0(X, L)$ (cf. Definition 8.1.1). In the case where the family $\xi = (\|\cdot\|_{\varphi_{\omega}})_{\omega \in \Omega}$ is a pseudo-norm family in the sense of Definition 5.1.1, we call it the *pushforward pseudo-norm family* of φ . In that case, we say that the pushforward pseudo-norm family of φ is *well-defined*.

Remark 9.4.2. We use the notation of Definition 9.4.1. In general, it is not clear that the condition (*) in Definition 5.1.1 is satisfied for the family ξ of supremum pseudo-norms. We will give a criterion when the topological adelic curve is assumed to be integral.

Example 9.4.3. Let E be a finite-dimensional K-vector space and let $\xi = (\|\cdot\|_{\omega})_{\omega\in\Omega}$ be a pseudo-norm family on E. For any $\omega \in \Omega$, denote by \mathcal{E}_{ω} the finiteness module of $\|\cdot\|_{\omega}$. Assume that X is geometrically reduced. Let L be a line bundle on X and let $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega\in\Omega}$ be a model family of (X, L). Assume that we have a family $\beta = (\beta_{\omega} : \mathcal{E}_{\omega} \otimes_{A_{\omega}} \mathcal{O}_{\mathcal{X}_{\omega}} \to \mathcal{L}_{\omega})_{\omega\in\Omega}$ of surjective homomorphisms. Denote by φ the quotient pseudo-metric family defined by (E, ξ) and β . Then the pushforward pseudo-norm family of ξ is well-defined. Moreover, if ξ possesses a globally adapted basis, then the pushforward pseudo-norm family of φ possesses a globally adapted basis.

Proposition 9.4.4. Assume that X is geometrically reduced. Let L be a line bundle on X. Let $\varphi = (\varphi_{\omega})_{\omega}$ be a pseudo-metric family on L with a flat and coherent model family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$. Assume that the pushforward pseudo-norm family of φ is well-defined. We denote it by $\xi = (\|\cdot\|_{\varphi_{\omega}})_{\omega \in \Omega}$. Let K'/K be a finite field extension. Let $X_{K'} := X \otimes_K K'$, let $L' = L \otimes_K K'$ and let $S' := S \otimes_K K' = (K', \phi' : \Omega' \to M_{K'}, \nu')$ denote the topological adelic curve constructed in §4.2. Denote by φ' the pseudo-metric family on L' induced by extension of scalars by φ . Let $(\mathcal{X}'_{\omega'}, \mathcal{L}'_{\omega'})_{\omega' \in \Omega'}$ denote the model family of φ' . Then the following hold.

- (1) The pushforward pseudo-norm family ξ' of φ' is well-defined.
- (2) If ξ' is dominated, then ξ is dominated.

Proof. (1) follows directly from the fact that the ξ is well-defined. Moreover, the pseudo-norm families ξ, ξ' satisfy the hypotheses of Proposition 9.2.5 (7). Hence (2) holds.

We are now able to prove the analogue of ([CM19], Theorem 6.1.13) in our setting.

Theorem 9.4.5. Assume that X is geometrically integral. Let L be a line bundle on X. Let $\varphi = (\varphi_{\omega})_{\omega}$ be a pseudo-metric family on L with a flat and coherent model family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$. Assume that the pushforward pseudo-norm family of φ is well-defined and possesses a globally adapted basis. If φ is dominated, then ξ is strongly dominated.

Proof. As ξ is ultrametric on $\Omega_{\rm um}$, it suffices to prove that ξ is dominated.

Claim 9.4.6. Assume that the model family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$ is very ample, in particular L is very ample. Then ξ is dominated.

Proof. Let $E = H^0(X, L)$ and $r := \dim_K(E)$. As in the proof of ([CM19], Theorem 6.1.13), there exist a finite extension K'/K and closed points $P_1, ..., P_r$ of X such that $\kappa(P_i) \subset K'$ for all i = 1, ..., r. Moreover, we have a strictly decreasing sequence of K-vector spaces

$$\{0\} = E_r \subsetneq \cdots \subsetneq E_r = E \otimes_K K',$$

such that

$$\forall i \in \{1, ..., r\}, \quad E_i = \{s \in E_0 : s(P_1) = \cdots s(P_r) = 0\}.$$

By Proposition 9.4.4, we may assume that K = K'.

Let $\alpha_1, ..., \alpha_r$ denote respectively local bases of L around $P_1, ..., P_r$. Then we define a basis $(\theta_1, ..., \theta_r)$ of E^{\vee} as follows.

$$\forall i \in \{1, \dots, r\}, \quad \forall s \in E, \quad \theta_i(s) := f_s(P_i),$$

where, for any $s \in E$, $s = f_s \alpha_i$ around P_i . Denote by $(e_1, ..., e_r)$ the corresponding dual basis of E.

Let $\omega \in \Omega$. Define a pseudo-norm on E in ω as follows. Let $s \in E$ written as $s = s_1e_1 + \cdots + s_re_r$, where $s_1, \ldots, s_r \in K$. We set

$$||s||'_{\omega} := \max_{i=1,\dots,r} |s_i|_{\omega}.$$

Then $\xi' = (\|\cdot\|'_{\omega})_{\omega}$ is a pseudo-norm family on E and the basis $(e_1, ..., e_r)$ is globally adapted basis to ξ' . Note that ξ' is dominated. Moreover, Lemma 5.1.5 implies that, there exists an open subset $\Omega' \subset \Omega$ such that, for any $\omega \in \Omega'$, the finiteness modules of $\|\cdot\|_{\omega}$ and $\|\cdot\|'_{\omega}$ coincide and $\nu(\Omega') = 0$. From now on, we work on the topological adelic curve $S' = (K, \phi' : \Omega' \to M_K, \nu_{|\Omega'})$ obtained by restriction of the adelic structure on Ω' (cf. §2.2.3).

We have a family of surjective homomorphisms $\beta := (\beta_{\omega} : \mathcal{E}_{\omega} \otimes_{A_{\omega}} \mathcal{O}_{\mathcal{X}_{\omega}} \to \mathcal{L}_{\omega})_{\omega \in \Omega'}$. We consider the quotient pseudo-metric family $\varphi' = (\varphi'_{\omega})_{\omega \in \Omega'}$ induced by (E, ξ') and β .

Let us prove that, for any $\omega \in \Omega'$, we have $\|\cdot\|'_{\omega} = \|\cdot\|_{\varphi'_{\omega}}$. Let $\omega \in \Omega'$. Using ([CM19], Proposition 2.2.23), we have the inequality $\|\cdot\|_{\varphi'_{\omega}} \leq \|\cdot\|'_{\omega}$. We prove the converse inequality. By the valuative criterion of properness, for any i = 1, ..., r, the rational point P_i defines

a rational point $\widetilde{P_{i,\omega}}$ of the special fibre $\mathcal{X}_{\omega,s}$ (the unique closed point of $\overline{\{P_i\}} \cap \mathcal{X}_{\omega,s}$). Let $(i,j) \in \{1,...,r\}$. Then we have

$$|e_i|_{\varphi'_{\omega}}(\widetilde{P_{i,\omega}}) = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{cases}$$

Let $s = s_1 e_1 + \dots + s_r e_r \in \mathcal{E}_{\omega}$, where $s_1, \dots, s_r \in A_{\omega}$. Then

$$\forall i = 1, \dots, r, \quad |s|_{\varphi'_{\omega}}(\widetilde{P_{i,\omega}}) = \|s\|'_{\omega} \le \|s\|_{\varphi'_{\omega}}.$$

Therefore, the following holds.

$$\forall \omega \in \Omega', \quad d_{\omega}(\xi, \xi') \le d_{\omega}(\|\cdot\|_{\varphi_{\omega}}, \|\cdot\|_{\varphi_{\omega}}) \le d_{\omega}(\varphi, \varphi').$$

As φ is dominated, the local distance function $(\omega \in \Omega') \mapsto d_{\omega}(\varphi, \varphi')$ is $\nu_{|\Omega'}$ -dominated. Therefore using Proposition 5.3.6, we obtain that ξ is dominated. This concludes the proof of the claim.

Claim 9.4.7. For any $s \in H^0(X, L) \setminus \{0\}$, the function $(\omega \in \Omega) \mapsto \log \|s\|_{\varphi_\omega}$ is ν -dominated.

Proof. Let $s \in H^0(X, L) \setminus \{0\}$. Lemma 5.1.6 implies that there exists a locally closed subset $\Omega' \subset \Omega$ such that $\nu(\Omega \setminus \Omega') = 0$ and, for any $\omega \in \Omega'$, $||s||_{\varphi_{\omega}} \notin \{0, +\infty\}$. We denote by $S' = (K, \phi' : \Omega' \to M_K, \nu_{|\Omega'})$ the restriction of the adelic structure introduced in §2.2.3.

Choose a very ample line bundle L_1 on X with a very ample model family $(\mathcal{X}_{\omega}, \mathcal{L}_{1,\omega})_{\omega \in \Omega'}$ of L_1 , such that $L_2 := L + L_1$ is very ample and the model family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega} + \mathcal{L}_{1,\omega})_{\omega \in \Omega'}$ is very ample. For any $\omega \in \Omega'$, we denote $\mathcal{E}_{1,\omega} := H^0(\mathcal{X}_{\omega}, \mathcal{L}_{1,\omega})$ and $\mathcal{E}_{2,\omega} := H^0(\mathcal{X}_{\omega}, \mathcal{L}_{2,\omega})$. Note that multiplication by s yields a family of injective homomorphisms $(\mathcal{E}_{1,\omega} \hookrightarrow \mathcal{E}_{2,\omega})_{\omega \in \Omega'}$.

We consider an arbitrary dominated pseudo-norm family $\xi'_2 = (\|\cdot\|'_{2,\omega})_{\omega\in\Omega'}$ on $E_2 := H^0(X,L)$ such that, for any $\omega \in \Omega'$, the finiteness module of $\|\cdot\|'_{2,\omega}$ is $\mathcal{E}_{2,\omega}$. Denote by $\xi'_1 = (\|\cdot\|'_{1,\omega})_{\omega\in\Omega'}$ the restriction of ξ'_2 to $E_1 := H^0(X,L)$. Note that for any $\omega \in \Omega'$, the finiteness module of $\|\cdot\|'_{1,\omega}$ is $\mathcal{E}_{1,\omega}$. Note that Proposition 5.3.5 (1) implies that ξ'_1 is dominated. Let i = 1, 2. Denote by φ'_i the quotient pseudo-metric family induced by ξ'_i and the family of surjective homomorphisms $\beta'_i = (\beta'_{i,\omega} : \mathcal{E}_{i,\omega} \otimes_{A_\omega} \mathcal{O}_{\mathcal{X}_\omega} \to \mathcal{L}_{i,\omega})_{\omega\in\Omega'}$. Let $\varphi' := \varphi'_2 - \varphi'_1$, this is a pseudo-metric family on L with the same model family as φ . Hence Proposition 9.2.5 (5) implies that the local distance function $(\omega \in \Omega') \mapsto d_\omega(\varphi, \varphi')$ is $\nu_{|\Omega'}$ -dominated.

Arguing as in the proof of ([CM19], Claim 6.1.16), we obtain $||s||_{\varphi'_{\omega}} \leq 1$. Moreover, for any $u \in E$, which belongs to $\bigcap_{\omega \in \Omega'} \mathcal{E}_{1,\omega}$ (up to shrinking Ω'), we have

$$\|su\|_{\varphi'_{2,\omega}} \le \|s\|_{\varphi'_{\omega}} \|u\|_{\varphi'_{1,\omega}}.$$

Using the fact that ξ'_1 and ξ'_2 are dominated. we deduce that the function $(\omega \in \Omega') \mapsto \log ||s||_{\varphi'_{\omega}}$ is $\nu_{|\Omega'}$ -dominated. We can conclude using the fact that

$$\forall \omega \in \Omega', \quad d_{\omega}(\|\cdot\|_{\varphi_{\omega}}, \|\cdot\|_{\varphi'_{\omega}}) \le d_{\omega}(\varphi, \varphi').$$

We now prove Theorem 9.4.5. Choose a very ample line bundle L_1 on X with a very ample model family $(\mathcal{X}_{\omega}, \mathcal{L}_{1,\omega})_{\omega \in \Omega'}$ of L_1 , such that $L_2 := L + L_1$ is very ample and the model family $(\mathcal{X}_{\omega}, \mathcal{L}_{2,\omega} := \mathcal{L}_{\omega} + \mathcal{L}_{1,\omega})_{\omega \in \Omega'}$ is very ample. Fix a global section $t \in H^0(X, L_1)$. Let $\varphi_1 = (\varphi_{1,\omega})_{\omega \in \Omega}$ be a dominated pseudo-metric family such that the pushforward metric family of φ_1 is well-defined and possesses a globally adapted basis and, for any $\omega \in \Omega$, we have $||t||_{\varphi_{\omega}} \leq 1$ (it is possible to do so by choosing a suitable quotient pseudo-metric family

cf. Example 9.4.3). Let $\varphi_2 := \varphi + \varphi_1$ and denote by ξ_2 the pushforward pseudo-metric family of φ_2 . Proposition 9.2.5 (2) implies that φ_2 is dominated. As $(\mathcal{X}_{\omega}, \mathcal{L}_{2,\omega})_{\omega \in \Omega}$ is very ample, Claim 9.4.6 implies that ξ_2 is strongly dominated.

Note that by hypothesis, for any $\omega \in \Omega$, $t \in H^0(\mathcal{X}_{\omega}, \mathcal{L}_{1,\omega})$ and multiplication by t gives an injective homomorphism $H^0(\mathcal{X}_{\omega}, \mathcal{L}_{\omega}) \hookrightarrow H^{\mathcal{X}_{\omega}, \mathcal{L}_{2,\omega}}$. Fix a basis $(s_1, ..., s_r)$ of $H^0(X, L)$ which is globally adapted to ξ . For i = 1, ..., r, write $t_i := ts_i \in H^0(X, L_2)$. Enlarge $(t_1, ..., t_r)$ to a basis $\mathbf{t} := (t_1, ..., t_n)$ of $H^0(X, L_2)$ which is globally adapted to ξ_2 . Let $\xi_{\mathbf{t}, 2} = (\|\cdot\|_{\mathbf{t}, 2, \omega})_{\omega \in \Omega}$ denote the model pseudo-norm family on $H^0(X, L_2)$ defined by the basis \mathbf{t} (cf. Example 5.1.3). Corollary 5.3.9 implies that the local distance function $(\omega \in \Omega) \mapsto d_{\omega}(\xi_2, \xi_{\mathbf{t}, 2})$ is ν -dominated. Thus there exists a ν -dominated function $A : \Omega \to [-\infty, +\infty]$ such that, for almost all $\omega \in \Omega$, for any $(\lambda_1, ..., \lambda_r) \in K^r$, we have

$$\log \|\lambda_1 s_1 + \dots + \lambda_r s_r\|_{\varphi_{\omega}} \ge \log \|\lambda_1 t_1 + \dots + \lambda_r t_r\|_{\varphi_{2,\omega}}$$
$$\ge \log \|\lambda_1 t_1 + \dots + \lambda_r t_r\|_{\varphi_{t,2,\omega}} - A(\omega), \tag{4}$$

where the first inequality comes from the fact that $||t||_{\varphi_{1,\omega}} \leq 1$. Moreover, we have the inequality

$$\log \|\lambda_1 s_1 + \dots + \lambda_r s_r\|_{\varphi_{\omega}} \le \log \|\lambda_1 t_1 + \dots + \lambda_r t_r\|_{\varphi_{\mathbf{t},2,\omega}} + \max_{1 \le i \le r} \log \|s_i\|_{\varphi_{\omega}}.$$

Claim 9.4.7 implies that there exists a ν -dominated function $B: \Omega \to [-\infty, +\infty]$ such that the inequality

$$\log \|\lambda_1 s_1 + \dots + \lambda_r s_r\|_{\varphi_{\omega}} \le \log \|\lambda_1 t_1 + \dots + \lambda_r t_r\|_{\varphi_{\mathbf{t},2,\omega}} + B(\omega)$$
(5)

holds for almost all $\omega \in \Omega$.

Denote by ξ' the restriction of the pseudo-norm family $\xi_{t,2}$ to $H^0(X, L)$. Then the basis $(s_1, ..., s_r)$ is both globally adapted to ξ and ξ' . Moreover (4) and (5) imply that the local distance function ($\omega \in \Omega$) $\mapsto d_{\omega}(\xi, \xi')$ is ν -dominated. By Corollary 5.3.9, the pseudo-norm family ξ is strongly dominated.

Proposition 9.4.8. We consider the case where X = Spec(K'), where K'/K is a finite extension of fields. We denote $S' := S \otimes_K K' = (K', \phi' : \Omega' \to M_{K'}, \nu')$. For any $\omega \in \Omega$, let A'_{ω} be the integral closure of A_{ω} in K', we denote $\mathcal{X}_{\omega} := \text{Spec}(A'_{\omega})$. Let L be a line bundle on X equipped with a pseudo-metric family φ with model family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$. Denote by ξ_L the pseudo-norm family on L (w.r.t. the topological adelic curve S') introduced in Remark 9.1.3. Assume that ξ_L possesses a globally adapted basis. Then φ is dominated iff ξ_L is dominated.

Proof. We first assume that φ is dominated. Since the model family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$ is very ample, there exist a finite-dimensional K-vector space E together with a dominated pseudonorm family $\xi_E = (\|\cdot\|_{\omega}, \mathcal{E}_{\omega}, N_{\omega}, \widehat{E_{\omega}})_{\omega \in \Omega}$ on E which possesses a globally adapted basis and a family of surjective homomorphisms $\beta := (\beta_{\omega} : \mathcal{E}_{\omega} \otimes_{A_{\omega}} A'_{\omega} \to \mathcal{L}_{\omega})_{\omega \in \Omega}$ such that the quotient pseudo-metric family φ' induced by (E, ξ) and β satisfies: φ' has model family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$ and the function $(\omega \in \Omega) \mapsto d_{\omega}(\varphi, \varphi')$ is ν -dominated. We denote by ξ'_L the pseudo-norm family on L w.r.t. the topological adelic curve S determined by φ' . Then by definition, ξ'_L identifies with the quotient norm family on L (viewed as a K'-vector space) of the extension of scalars $\xi_{E,K'}$ via the surjection $E \otimes_K K' \to L$. Corollary 5.3.10 implies that $\xi_{E,K'}$ is dominated and Proposition 5.3.5 (2) implies that ξ'_L is dominated. As

$$\forall \omega \in \Omega', \quad d_{\omega}(\xi_L, \xi'_L) = d_{\omega}(\varphi, \varphi'),$$

Proposition 5.3.6 ensures that ξ_L is dominated.

Conversely, assume that ξ_L is dominated. Let $e \in L$ be a globally adapted basis to ξ_L . Then denote by $\xi_e = (\|\cdot\|_{e,\omega})_{\omega \in \Omega}$ the pseudo-norm family on $K \cdot e$ defined by

$$\forall \omega \in \Omega, \quad \forall \lambda \in K, \quad \|\lambda e\|_{e,\omega} = |\lambda|_{\omega}.$$

Then ξ_e is dominated and $\xi_{e,K'}$ is dominated by Corollary 5.3.10. Corollary 5.3.9 implies that the local distance function $(\omega \in \Omega') \mapsto d_{\omega}(\xi_{e,K'}, \xi_L)$ is ν' -dominated. Denote by φ' the pseudo-metric family induced by the family of surjective homomorphisms $\beta = (\beta_{\omega} : A_{\omega} \cdot e \otimes_{A_{\omega}} A'_{\omega} \cong \mathcal{L}_{\omega})_{\omega \in \Omega}$. Then, for any $\omega \in \Omega'$, we have $d_{\omega}(\xi_{e,K'}, \xi_L) = d_{\omega}(\varphi, \varphi')$. Hence Proposition 9.2.5 (3) implies that φ is dominated. \Box

9.5. Adelic line bundles. In this subsection, we fix a topological adelic curve $S = (K, \phi : \Omega \to M_K, \nu)$ and a projective K-scheme X. Recall that, for any $\omega \in \Omega$, we denote by A_{ω} and κ_{ω} , the finiteness ring and the residue field of ω respectively.

Definition 9.5.1. Let L be a line bundle on X equipped with a pseudo-metric family φ . We say that $\overline{L} = (L, \varphi)$ is respectively a *usc/lsc adelic line bundle* on X if φ is both usc/lsc and dominated. Moreover, we say that \overline{L} is a *(continuous) adelic line bundle* if φ is both continuous and dominated.

Moreover, an adelic line bundle $(L, \varphi = (\varphi_{\omega})_{\omega \in \Omega})$ is called *semi-positive*, resp. *integrable*, if, for any $\omega \in \Omega$, the locally pseudo-metrised line bundle (L, φ_{ω}) is semi-positive, resp. integrable.

Proposition 9.5.2. (1) Let $\overline{L} = (L, \varphi)$ be a usc/lsc/continuous adelic line bundle on X. Then $-\overline{L} := (-L, -\varphi)$ is respectively a usc/lsc/continuous adelic line bundle on X.

- (2) Let $\overline{L_1} = (L_1, \varphi_1), \overline{L_2} = (L_2, \varphi_2)$ be both usc/lsc/continuous adelic line bundles on X. Then $\overline{L_1} + \overline{L_2} := (L_1 + L_2, \varphi_1 + \varphi_2)$ is respectively a usc/lsc/continuous adelic line bundle on X.
- (3) Let $f: Y \to X$ be a projective morphism of K-schemes. Assume that, for any $\omega \in \Omega$, f extends to a A_{ω} -morphism $g_{\omega}: \mathcal{Y}_{\omega} \to \mathcal{X}_{\omega}$, where \mathcal{Y}_{ω} is a projective model of Yover A_{ω} . Let $\overline{L} = (L, \varphi)$ be an adelic line bundle on X. Then $f^*\overline{L}: (f^*L, f^*\varphi)$ is respectively a usc/lsc/continuous adelic line bundle on X.

Proof. This is a direct consequence of Propositions 9.2.5 and 9.3.2.

Proposition 9.5.3. Let $\overline{E} = (E, \xi)$ where E is a finite-dimensional K-vector space and $\xi = (\|\cdot\|_{\omega}, \mathcal{E}_{\omega}, N_{\omega}, \widehat{E}_{\omega})_{\omega \in \Omega}$ is a pseudo-norm family on E which is assumed to be ultrametric on Ω_{um} . Let L be a line bundle on X and let $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$ be a model family of (X, L). Assume that, for any $\omega \in \Omega$, there exists a surjective homomorphism $\beta_{\omega} : \mathcal{E}_{\omega} \otimes_{A_{\omega}} \mathcal{O}_{\mathcal{X}_{\omega}} \to \mathcal{L}_{\omega}$. Denote by φ the corresponding quotient pseudo-metric family on L (cf. Example 9.1.5). Assume that $\overline{E} = (E, \xi)$ is an adelic line bundle on S. Then $\overline{L} = (L, \varphi)$ is an adelic line bundle on X.

Proof. The fact that φ is dominated is Proposition 9.2.4. Proposition 9.3.3 implies that φ is use. As ξ^{\vee} is continuous and as $-\varphi$ identifies with the quotient pseudo-metric family induced by (E^{\vee}, ξ^{\vee}) , Proposition 9.3.3 implies that $-\varphi$ is use as well. This implies that φ is lsc, and therefore continuous.

Example 9.5.4 (Adelic line bundles in Nevanlinna theory). Let R > 0. Consider the topological adelic curve $S_R = (K_R, \phi_R : \Omega_R \to M_{K_R}, \nu_R)$ be the topological adelic curve

defined in §2.2.2. Let X be a reduced projective \mathbb{C} -scheme and denote $X_R := X \otimes_{\mathbb{C}} K_R$. For any $\omega \in \Omega_R$, denote by $A_{R,\omega}$ the finiteness ring of ω and let $\mathcal{X}_{R,\omega} := X \otimes_{\mathbb{C}} A_{R,\omega}$, it is a projective model of X_R over $A_{R,\omega}$. Let (L,φ) be a metrised line bundle on X, namely L is a line bundle on X and φ is a continuous metric on L. Further, assume that L is globally generated. For any $\omega \in \Omega_R$, we denote $\mathcal{L}_{R,\omega} := L \otimes_{\mathcal{O}_X} \mathcal{O}_{\mathcal{X}_{R,\omega}}$ and $L_R := L \otimes_{\mathcal{O}_X} \mathcal{O}_{\mathcal{X}_R}$. Note that L_R and the $\mathcal{L}_{R,\omega}$ for $\omega \in \Omega_R$ are globally generated as well.

Let us show that (L, φ) induces an adelic line bundle on X_R . Let $\omega \in \Omega_{R,ar}$. Then the special fibre of $\mathcal{X}_{R,\omega}$ identifies with X. Then $(\mathcal{L}_{R,\omega}, \varphi)$ defines a pseudo-metric on L_R . Let $\omega \in \Omega_{R,um}$. As $\phi_R(\omega)$ is a usual absolute value on K_R , we have $X_R = \mathcal{X}_{R,\omega}$. Then the completion $K_{R,\omega}$ of K_R w.r.t. $|\cdot|_{\omega}$ is isomorphic to $\mathbb{C}((T))$. Denote by $K_{R,\omega}^{\circ} \cong \mathbb{C}[[T]]$ the corresponding valuation ring of $K_{R,\omega}$. Let $\widetilde{\mathcal{X}_{R,\omega}} := X \otimes_{\mathbb{C}} K_{R,\omega}^{\circ}$ and $\widetilde{\mathcal{L}_{R,\omega}} := L \otimes_{\mathcal{O}_X} \mathcal{O}_{\widetilde{\mathcal{X}_{R,\omega}}}$. Then $(\widetilde{\mathcal{X}_{R,\omega}, \widetilde{\mathcal{L}_{R,\omega}})$ is a model of $X_R \otimes_{K_R} K_{R,\omega}$ over $K_{R,\omega}^{\circ}$ and the model metric $\varphi_{R,\omega}$ determined by this model yields a local pseudo metric on L_R in ω . These construction yield a pseudo-metric family $\varphi_R = (\varphi_{R,\omega})_{\omega \in \Omega_R}$ with model family $(\mathcal{X}_{R,\omega}, \mathcal{L}_{R,\omega})_{\omega \in \Omega_R}$. Let us prove that φ_R is dominated and continuous.

Denote $E := H^0(X, L)$ and

$$E_R := E \otimes_{\mathbb{C}} K_R, \quad \forall \omega \in \Omega_{R,\mathrm{ar}}, \quad \mathcal{E}_{R,\omega} := E \otimes_{\mathbb{C}} A_{R,\omega}, \quad \forall \omega \in \Omega_{R,\mathrm{um}}, \quad \widetilde{\mathcal{E}_{R,\omega}} := E \otimes_{\mathbb{C}} K_{R,\omega}^{\circ}.$$

Let $\|\cdot\|_{\varphi}$ denote the supremum norm on E induced by φ and denote by φ_{FS} the Fubini-Study (usual) metric on L associated with φ . Recall that φ_{FS} is the quotient metric associated with the complex normed vector space $(E, \|\cdot\|_{\varphi})$. Moreover, by using the results of §6.3, we have an adelic vector bundle (E_R, ξ_R) on S_R . As for any $\omega \in \Omega_R$, $\mathcal{X}_{R,\omega}$ is flat and coherent, we denote by $\varphi_{R,\text{FS}}$ Fubini-Study pseudo-metric family associated with (E_R, ξ_R) .

Let $\omega \in \Omega_{R,\text{um}}$. As $\mathbb{C} \to K_{R,\omega}^{\circ}$ is flat, for any $\omega \in \Omega_{R,\text{um}}$, flat base change yields $\widetilde{\mathcal{E}_{R,\omega}} = H^0(\widetilde{\mathcal{X}_{R,\omega}}, \widetilde{\mathcal{L}_{R,\omega}})$ and the evaluation morphism $\widetilde{\mathcal{E}_{R,\omega}} \otimes_{K_{R,\omega}^{\circ}} \mathcal{O}_{\widetilde{\mathcal{X}_{R,\omega}}} \to \widetilde{\mathcal{L}_{R,\omega}}$ is surjective. Using ([CM19], Proposition 2.3.12), we obtain that $\varphi_{R,\omega}$ identifies with the quotient metric induced by the lattice norm defined by the lattice $\widetilde{\mathcal{E}_{R,\omega}}$ inside $\widetilde{\mathcal{E}_{R,\omega}} := \widetilde{\mathcal{E}_{R,\omega}} \otimes_{K_{R,\omega}^{\circ}} K_{R,\omega}$. Note that from the description of ξ_R given in §6.3, we have $\varphi_{R,\text{FS},\omega} = \varphi_{\omega}$.

Moreover, we have

$$\forall \omega \in \Omega_R, \quad d_\omega(\varphi_R, \varphi_{R, \text{FS}}) = \mathbf{1}_{\Omega_{R, \text{ar}}}(\omega) d(\varphi, \varphi_{\text{FS}}),$$

where $d(\varphi, \varphi_{\rm FS})$ denotes the usual distance between the (complex) continuous metrics φ and $\varphi_{\rm FS}$. As $\nu_R(\Omega_{R,\rm ar}) < +\infty$, we obtain that the local distance function $[(\omega \in \Omega) \mapsto d_{\omega}(\varphi_R, \varphi_{R,\rm FS}]$ is ν_R -dominated. Using Proposition 6.3.1 and Proposition 9.2.4, we obtain that $\varphi_{R,\rm FS}$ is dominated. Therefore, by using Proposition 9.2.5 (3), we see that the pseudometric family φ_R is dominated.

We conclude by proving the continuity of φ_R . Let P be a closed point of X_R . Denote by $S_{R,P} = (\kappa(P), \phi_P : \Omega_{R,P} \to M_{\kappa(P)}, \nu_{R,P})$ the topological adelic curve $S_R \otimes_{K_R} \kappa(P)$. By definition of the pseudo-norm family $P^*\varphi_R$ on P^*L , we see that $P^*\varphi_R$ coincides with the pseudo-norm family determined by $(E, \|\cdot\|)$ on $\kappa(P)$. Using Remark 6.3.3, we obtain the continuity of $P^*\varphi_R$.

Definition 9.5.5. We use the same notation as in Example 9.5.4. Let R > 0. Then the adelic line bundle constructed in Example 9.5.4 is called the *adelic line bundle induced* by (L, φ) on X_R .

Proposition 9.5.6. We use the same notation as in Example 9.5.4. Assume that X is reduced. Let R > 0 and let (L, φ) be a metrised line bundle on X. We denote by $\overline{L_R} = (L_R, \varphi_R)$ the adelic line bundle on X_R induced by (L, φ) . Denote by ξ_R the collection of supremum pseudonorms on $E_R := H^0(X_R, L_R)$ induced by the pseudo-metric family φ . Then $\overline{E_R} := (E_R, \xi_R)$ is an adelic line bundle on S_R .

Proof. We first note that ξ_R is a well-defined pseudo-norm family on E_R . Indeed, any basis of the space of global sections $E := H^0(X, L)$ yields a globally adapted basis for ξ_R . Denote by $\|\cdot\|$ the supremum norm on $H^0(X, L)$ induced by the continuous metric φ .

Write $\xi_R = (\|\cdot\|_{\varphi_{\omega}})_{\omega \in \Omega_R}$. Let $\omega \in \Omega_{R,ar}$. Then $\|\cdot\|_{\varphi_{\omega}}$ coincides with the local pseudo-norm on $H^0(X, L)$ in ω whose residue vector space is E and whose residue norm is $\|\cdot\|$. Let $\omega \in \Omega_{R,um}$. As X is reduced, ([CM19], Proposition 2.3.16 (3)) implies that $\|\cdot\|_{\varphi_{\omega}}$ coincides with the lattice norm used in §6.3.

Using this description, we see that (E_R, ξ_R) coincides with the adelic vector bundle induced by the normed vector space $(E, \|\cdot\|)$ (cf. Definition 6.3.2). This allows us to conclude. \Box

9.6. Adelic line bundles on families of topological adelic curves. Throughout this subsection, we consider the following setting. Let (I, \leq_I) be a (non-empty) totally ordered set. Let $\mathbf{S} = (S_i = (K_i, \phi_i : \Omega_i \to M_{K_i}, \nu_i))_{i \in I}$ be a family of topological adelic curves with base space Ω_0 and family of structural morphisms $(\alpha_i : \Omega_i \to \Omega_0)_{i \in I}$ (cf. Definition 3.1.1). We denote

$$K_{\infty} := \varprojlim_{i \in I} K_i = \bigcap_{i \in I} K_i.$$

Let $\pi_{\infty} : X_{\infty} \to \operatorname{Spec}(K_{\infty})$ be a projective K_{∞} -scheme. For any $i \in I$, we denote $\pi_i : X_i := X_{\infty} \otimes_{K_{\infty}} K_i \to \operatorname{Spec}(K_i)$ and $\mathbf{X} := (X_i)_{i \in I}$. For any $i \in I$, for any $\omega \in \Omega_i$, we denote by $(A_{i,\omega}, \mathfrak{m}_{i,\omega}, \kappa_{i,\omega})$ the finiteness ring of ω .

Definition 9.6.1. By a *usc/lsc/continuous adelic line bundle* on **X**, we mean a collection $\overline{\mathbf{L}} = (\overline{L_i})_{i \in I}$, where

- (i) for any $i \in I$, $\overline{L_i} = (L_i, ((\mathcal{X}_{i,\omega}, \mathcal{L}_{i,\omega}), \varphi_{i,\omega})_{\omega \in \Omega_i})$ is respectively a usc/lsc/continuous adelic line bundle on X_i ;
- (ii) there exists a line bundle L_{∞} on X_{∞} such that, for any $i \in I$, $L_i = L_{\infty} \otimes_{\mathcal{O}_{X_{\infty}}} \mathcal{O}_{X_i}$;
- (iii) for any $i, i' \in I$ with $i \leq I$ i', for any $(\omega, \omega') \in \Omega_i \times \Omega_{i'}$ such that $\alpha_i(\omega) = \alpha_{\omega'}$, we have

$$\mathcal{X}_{i,\omega} = \mathcal{X}_{i',\omega'} \otimes_{A_{i,\omega}} A_{i',\omega'}, \quad \mathcal{L}_{i,\omega} = \mathcal{L}_{i',\omega'} \otimes_{\mathcal{O}_{\mathcal{X}_{i,\omega}}} \mathcal{O}_{\mathcal{X}_{i',\omega'}}$$

and the pseudo-metric $\varphi_{i,\omega}$ is the pullback of $\varphi_{i',\omega'}$ by the morphism $\varpi_{(i,\omega),(i',\omega')}$: $\mathcal{X}_{i,\omega} \to \mathcal{X}_{i',\omega'}$.

- **Proposition 9.6.2.** (1) Let $\overline{\mathbf{L}} = (\overline{L_i})_{i \in I}$ be a usc/lsc/continuous adelic vector bundle on \mathbf{X} . Then $-\overline{\mathbf{L}} := (-\overline{L_i})_{i \in I}$ is respectively a usc/lsc/continuous adelic vector bundle on \mathbf{X} .
 - (2) Let $\overline{\mathbf{L}^{(1)}} = (\overline{L_i^{(1)}})_{i \in I}, \overline{\mathbf{L}^{(2)}} = (\overline{L_i^{(2)}})_{i \in I}$ be both usc/lsc/continuous adelic line bundles on \mathbf{X} . Then $\overline{\mathbf{L}^{(1)}} + \overline{\mathbf{L}^{(2)}} := (\overline{L_i^{(1)}} + \overline{L_i^{(2)}})_{i \in I}$ is respectively a usc/lsc/continuous adelic line bundle on \mathbf{X} .

Proof. This is a consequence of Proposition 9.5.2 combined with the fact that condition (iii) in Definition 9.6.1 is compatible with taking dual and tensors products. \Box

Example 9.6.3. We consider the totally ordered set $\mathbb{R}_{>0}$ equipped with the usual total ordering. Consider the family of topological adelic curves $\mathbf{S} = (S_R)_{R>0}$ constructed in Example 3.1.2 (2). Let X be a reduced projective \mathbb{C} -scheme and let (L, φ) be a metrised line bundle on X, where L is assumed to be globally generated. Using the construction of Example 9.5.4, for any R > 0, we obtain an adelic line bundle $\overline{L_R} = (L_R, \varphi_R)$ on $X_R := X \otimes_{\mathbb{C}} K_R$. Then $\overline{\mathbf{L}} := (\overline{L_R})_{R>0}$ is an adelic line bundle on $\mathbf{X} := (X_R)_{R>0}$.

9.7. **Pseudo-metric families for integral topological adelic curves.** In this subsection, we study the case where the base topological adelic curve is integral. By fixing a model over the integral structure, we obtain a family of models and we can further make use of the results coming from the theory of global Berkovich spaces (cf. [LP24]).

We fix an integral topological adelic curve $S = (K, \phi : \Omega \to M_K, \nu)$ and a projective *K*-scheme *X*. We denote by $(A, \|\cdot\|)$ the underlying integral structure of *S* and we let $V := \mathcal{M}(A, \|\cdot\|)$. We also fix a flat and coherent model $\mathcal{X} \to \operatorname{Spec}(A)$ of *X* over *A*. We further assume that $(A, \|\cdot\|)$ is a geometric base ring ([LP24], Définition 3.3.8) and we denote by $\mathcal{X}^{\operatorname{an}}$ the Berkovich analytic space attached to \mathcal{X} (*loc. cit.*, Définition 4.1.5), this is a compact Hausdorff topological space by (*loc. cit.*, Proposition 6.5.3). Recall that, for any $\omega \in \Omega$, we denote by A_{ω} and κ_{ω} , the finiteness ring and the residue field of ω respectively.

9.7.1. Global adelic space.

Definition 9.7.1. We denote by $\mathcal{X}_{\Omega}^{an}$ the topological fibre product $\mathcal{X}^{an} \times_V \Omega$ and we call it the *global adelic space* attached to X w.r.t. the model \mathcal{X} . We have a Cartesian diagram of topological space



such that the map $p: \mathcal{X}_{\Omega}^{\mathrm{an}} \to \Omega$ is proper (as the pullback of the proper map $\mathcal{X}^{\mathrm{an}} \to V$).

Notation 9.7.2. (1) Let $\mathcal{U} \subset \mathcal{X}$ be a Zariski open subset. We denote by $\mathcal{U}_{\omega}^{an}$ the fibre product $\mathcal{U}^{an} \times_V \Omega$.

(2) Let \mathcal{L} be a line bundle on \mathcal{X} . For any $\omega \in \Omega$, we denote by \mathcal{L}_{ω} the pullback of \mathcal{L} to $\mathcal{X}_{\omega} = \mathcal{X} \otimes_A A_{\omega}$.

9.7.2. Global pseudo-metrics. Let \mathcal{L} be a line bundle on \mathcal{X} . By a metric φ on \mathcal{L} , we mean a family $\varphi = (|\cdot|_{\varphi_{\omega}}(x))_{(x,\omega)\in\mathcal{X}_{\Omega}^{an}}$, where, for any $(x,\omega)\in\mathcal{X}_{\Omega}^{an}$, $|\cdot||_{\varphi}(x)$ is a norm on the $\widehat{\kappa_{\omega}}$ -vector space $\mathcal{L}(x,\omega) := \mathcal{L} \otimes_{\mathcal{O}_{\mathcal{X}}} \widehat{\kappa_{\omega}}$. We say that a metric φ on \mathcal{L} is strongly continuous if, for any local section s of \mathcal{L} on a Zariski open subset $\mathcal{U} \subset \mathcal{X}$, the function

$$((x,\omega) \in \mathcal{U}_{\Omega}^{\mathrm{an}}) \mapsto |s|_{\varphi_{\omega}}(x) \in [0,+\infty[$$

is continuous. Likewise, we have the notion of strongly upper/lower semi-continuous metric.

Let L be a line bundle on X. By a model global pseudo-metric (\mathcal{L}, φ) on L over \mathcal{X} , we mean that \mathcal{L} is a line bundle on \mathcal{X} such that $(\mathcal{X}, \mathcal{L})$ is a model of (X, L) and that φ is a metric on \mathcal{L} . In this article, we will not consider any other kind of global pseudo-metrics. Therefore by "global pseudo-metric" we mean model global pseudo-metric.

Proposition 9.7.3. Let L be a line bundle on X and let (\mathcal{L}, φ) be a global pseudo-metric on L over \mathcal{X} .

- (1) Then the family $(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})_{\omega \in \Omega}$ is a flat and coherent model family of (X, L) in the sense of Definition 9.1.1.
- (2) For any $\omega \in \Omega$, we denote by φ_{ω} the family $(|\cdot|_{\varphi_{\omega}(x)})_{x \in X_{\omega}^{\mathrm{an}}}$. Then $((\mathcal{X}_{\omega}, \mathcal{L}_{\omega}), \varphi_{\omega})_{\omega \in \Omega}$ defines a pseudo-metric family on \mathcal{L} .

Proof. (1) holds by stability of flatness and finite presentation along base change and (2) follows from ([LP24], Proposition 4.5.3). \Box

Conversely, let $((\mathcal{X}_{\omega}, \mathcal{L}_{\omega}), \varphi_{\omega})_{\omega \in \Omega}$ be a pseudo-metric family of a line bundle L on X. Assume that there exists a model $(\mathcal{X}, \mathcal{L})$ of (X, L) such that, for any $\omega \in \Omega$, the pullback of \mathcal{L} to \mathcal{X}_{ω} is \mathcal{L}_{ω} . Then, by patching all the pseudo-metrics φ_{ω} , we obtain a global pseudo-metric on L. Thus global pseudo-metrics are special kinds of pseudo-metric families.

Notation 9.7.4. Let L be a line bundle on X and let (\mathcal{L}, φ) be a global pseudo-metric on L. By abuse of notation, we will identify the global pseudo-metric φ with the corresponding pseudo-metric family $((\mathcal{X}_{\omega}, \mathcal{L}_{\omega}), \varphi_{\omega})_{\omega \in \Omega}$. For instance, a global pseudo-metric is called *dominated* if the corresponding pseudo-metric family is so.

Remark 9.7.5. We consider the case where $X = \operatorname{Spec}(K')$, where K'/K is a finite extension of fields, and $\mathcal{X} = \operatorname{Spec}(A')$, where A' denotes the integral closure of A in K'. Let L be a line bundle on X, i.e. a one-dimensional K'-vector space. Then, for any $\omega \in \Omega$, the local analytic space $X_{\omega}^{\operatorname{an}}$ is a discrete finite set. Therefore, a global pseudo-metric (\mathcal{L}, φ) is the same as a pseudo-norm family ξ_L on the one-dimensional K'-vector space L, relatively to the (integral) topological adelic curve $S' := S \otimes_K K'$.

9.7.3. Pushforward of global pseudo-metrics. Throughout this paragraph, we assume that X is geometrically integral.

Let *L* be a line bundle on *X* and let (\mathcal{L}, φ) be a global pseudo metric on *L*. Then for any $\omega \in \Omega$, we can consider the sup-pseudo-norm $\|\cdot\|_{\varphi_{\omega}}$ (cf. 8.1.1). Let $\xi := (\|\cdot\|_{\varphi_{\omega}})_{\omega \in \Omega}$.

Proposition 9.7.6. Let L be a line bundle on X and let (\mathcal{L}, φ) be a global pseudo metric on L. Assume that the global pseudo-metric φ is strongly continuous. Then ξ is well defined and is an upper semi-continuous pseudo-norm family on $H^0(X, L)$.

Proof. Recall that, for any $\omega \in \Omega$, $\|\cdot\|_{\varphi_{\omega}}$ is a pseudo-norm on $H^0(X, L)$ with finiteness module $H^0(\mathcal{X}_{\omega}, \mathcal{L}_{\omega})$. Let $s \in H^0(X, L)$, let us prove that the function $(\omega \in \Omega) \mapsto \|s\|_{\varphi_{\omega}}$ is upper semi-continuous. We will use the following lemma.

Lemma 9.7.7 ([CM22], Proposition 2.2.4). Let $\pi : X \to Y$ be a closed continuous mapping of topological spaces and let $f : X \to [-\infty, +\infty]$ be an upper semi-continuous function. Assume that, for any $y \in Y$, the fibre $\pi^{-1}(y)$ is compact. Then the function

$$\phi_f: \left| \begin{array}{ccc} Y & \longrightarrow & [0,1] \\ y & \longmapsto & \sup_{x \in \pi^{-1}(y)} f(x) \end{array} \right|$$

is upper semi-continuous.

Let $s \in H^0(X, L)$. Apply the above lemma with $X = \mathcal{X}_{\Omega}^{\mathrm{an}}$, $Y = \Omega$ and $f : ((x, \omega) \in \mathcal{X}_{\Omega}^{\mathrm{an}}) \mapsto |s|_{\varphi_{\omega}}(x)$, which is continuous. Note that $p : \mathcal{X}_{\Omega}^{\mathrm{an}} \to \Omega$ is a proper map since Ω is locally compact and Hausdorff. Since, for any $\omega \in \Omega$, we have

$$\sup_{(x,\omega)\in p^{-1}(\omega)}|s|_{\varphi_{\omega}}(x)=\|s\|_{\varphi_{\omega}},$$

we deduce the desired upper semi-continuity.

Thus it suffices to prove that for any $\omega \in \Omega$, there exists an adapted basis for ξ in ω . We fix $\omega_0 \in \Omega$. Since $\|\cdot\|_{\varphi_{\omega_0}}$ is a pseudo-norm on E, there exists a basis $(s_1, ..., s_d)$ of E such that, for any i = 1, ..., d, we have $\|s_i\|_{\varphi_{\omega_0}} \in \mathbb{R}_{>0}$. As, for any i = 1, ..., d, the map $(\omega \in \Omega) \mapsto \|s_i\|_{\varphi_{\omega}} \in [0, +\infty]$ is upper semi-continuous, there exists an open neighbourhood U of ω_0 in Ω such that, for any $\omega \in U$, we have

$$\forall i = 1, \dots, d, \quad \|s_i\|_{\varphi_\omega} < +\infty.$$

Therefore, it remains to prove that, up to shrinking U, we have

$$\forall i = 1, \dots, d, \quad \|s_i\|_{\varphi_\omega} > 0.$$

For any closed point P of X and x and any element of $\Omega_{\kappa(P),\omega_0}$, by the valuative criterion of properness, we obtain a point in the local model analytic space X_{ω}^{an} which is denoted by (P, x) and whose image in X_{ω} is a closed point. Since closed points of X_{ω} are dense in X_{ω}^{an} , for any i = 1, ..., d, there exist a closed point P of X and an element $x \in \Omega_{\kappa(P),\omega_0}$ such that $|s_i|_{\varphi_{\omega_0}}(P, x) > 0$. By continuity of φ , up to shrinking U, we may assume that, for any $\omega \in U$, there exists a point $P_{i,\omega} \in X_{\omega}^{an}$ such that $|s|_{\varphi_{\omega}}(P_{i,\omega}) > 0$. In particular, we obtain

$$\forall i = 1, \dots, d, \quad \|s_i\|_{\varphi_\omega} > 0.$$

Hence ξ is a usc pseudo-norm family on $H^0(X, L)$

Definition 9.7.8. Let *L* be a line bundle on *X* and let (\mathcal{L}, φ) be a strongly continuous global pseudo metric on *L*. Then the usc pseudo-norm family $\xi = (\|\cdot\|_{\varphi_{\omega}})_{\omega \in \Omega}$ is called the *pushforward pseudo-norm family* induced by the continuous global pseudo-metric φ on *L*.

Theorem 9.7.9. Let L be a line bundle on X and let (\mathcal{L}, φ) be a strongly continuous and dominated global pseudo metric on L. Denote by ξ the pushforward pseudo-norm family of φ . Assume that ξ possesses a globally adapted basis. Then ξ is a usc and dominated pseudo-norm family on $H^0(X, L)$.

Proof. It is a consequence of Theorem 9.4.5 and Proposition 9.7.6.

9.8. Volume functions on a proper topological adelic curve. In this subsection, we fix a proper topological adelic curve $S = (K, \phi : \Omega \to M_K, \nu)$ and a geometrically integral projective K-scheme $\pi : X \to \text{Spec}(K)$ of dimension d.

Definition 9.8.1. Let $\overline{L} = (L, \varphi)$ be an adelic line bundle on X. For any integer $n \ge 1$, let $\pi_*(n\overline{L}) := (H^0(X, nL), \pi_*(n\varphi))$, where $\pi_*(n\varphi)$ denotes the collection of supremum pseudonorms on $H^0(X, nL)$ induced by $n\varphi$. Assume that, for any integer $n \ge 1$, $\pi_*(n\varphi)$ is well defined and is dominated and usc, so that the Arakelov degree $\widehat{\deg}(\pi_*(n\overline{L}))$ makes sense.

Then we define the χ -volume

$$\widehat{\mathrm{vol}}_{\chi}(\overline{L}) := \limsup_{n \to +\infty} \frac{\widehat{\mathrm{deg}}(\pi_*(n\overline{L}))}{n^{d+1}/(d+1)!},$$

and the arithmetic volume

$$\widehat{\mathrm{vol}}(\overline{L}) := \limsup_{n \to +\infty} \frac{\overline{\mathrm{deg}}_+(\pi_*(n\overline{L}))}{n^{d+1}/(d+1)!}$$

Example 9.8.2. Assume that the topological adelic curve is integral and fix a flat and coherent projective model \mathcal{X} over the integral structure. Let L be a line bundle on X equipped with a global pseudo-metric (\mathcal{L}, φ) w.r.t. the model \mathcal{X} . Denote $\overline{L} := (L, (\mathcal{L}, \varphi))$. Assume that (\mathcal{L}, φ) is strongly continuous and dominated and that, for any integer $n \geq 1$, the pushforward pseudo-norm family $\pi_*(n\varphi)$ possesses a globally adapted basis (e.g. when φ comes from a quotient of a pseudo-norm family which possesses a globally adapted basis as in Example 9.4.3). Then we can consider the χ -volume $\widehat{vol}_{\chi}(\overline{L})$ and the arithmetic volume $\widehat{vol}(\overline{L})$.

10. Height of closed point

In this section, we introduce global heights over topological adelic curves. We start by constructing height functions for closed points over a proper topological adelic curve (§10.1). Then we give the family counterpart for asymptotically proper families of topological adelic curves (§10.2).

10.1. Height of closed points on a proper topological adelic curve. In this subsection, we fix a proper topological adelic curve $S = (K, \phi : \Omega \to M_K, \nu)$ and a projective K-scheme X.

Definition 10.1.1. Let $\overline{L} = (L, \varphi)$ be an adelic line bundle on X. Let P be a closed point of X. The pseudo-norm family $P^*\varphi$ is continuous and dominated (cf. Proposition 9.4.8). As P^*L is a $\kappa(P)$ -vector space of dimension 1, Proposition 6.1.3 implies that $P^*\overline{L} := (P^*L, P^*\varphi)$ is an adelic line bundle on $S_P := S \otimes_K \kappa(P)$. As S is proper, S_P is proper as well (cf. Proposition 4.3.1). We define the *height* of P w.r.t. \overline{L} as

$$h_{\overline{L}}(P) := \widehat{\deg}_S(P^*\overline{L}).$$

Theorem 10.1.2. Let X be a projective K-scheme and let $\overline{L_1} = (L_1, \varphi_1), \overline{L_2} = (L_2, \varphi_2)$ be adelic line bundles on X. Then the following assertions hold.

(1) For any closed point P of X we have

$$h_{\overline{L_1}+\overline{L_2}}(P) = h_{\overline{L_1}}(P) + h_{\overline{L_2}}(P).$$

(2) Let P be a closed point of X. Assume that $L_1 = L_2$ and that φ_1, φ_2 have the same model family. Then we have

$$\left|h_{\overline{L_1}}(P) - h_{\overline{L_2}}(P)\right| \leq \int_{\Omega} d_{\omega}(\varphi_1, \varphi_2)\nu(\mathrm{d}\omega) < +\infty.$$

Proof. (1) This follows from Proposition 7.1.1 (3).

(2) Let P be a closed point of X. Write $S \otimes_K \kappa(P) = (\kappa(P), \phi_P : \Omega_P \to M_{\kappa(P)}, \nu_P)$. By definition, for any $\omega \in \Omega$, for any $x \in \pi_{\kappa(P)/K}^{-1}(\omega)$, we have

$$d_x(P^*\varphi_1, P^*\varphi_2) \le d_\omega(\varphi_1, \varphi_2)$$

By definition of $h_{\overline{L_1}}(P), h_{\overline{L_2}}(P)$, we obtain

$$\left|h_{\overline{L_1}}(P) - h_{\overline{L_2}}(P)\right| \le \int_{\Omega_P}^{\overline{}} d_x (P^* \varphi_1, P^* \varphi_2) \nu_P(\mathrm{d}x) \le \int_{\Omega}^{\overline{}} d_\omega(\varphi_1, \varphi_2) \nu(\mathrm{d}\omega).$$

The finiteness assertion follows from Proposition 9.2.5 (5).

Remark 10.1.3. Theorem 10.1.2 should be seen as the analogue of Nevanlinna's first main theorem in Arakelov geometry (cf. Theorem A.2.1).

10.2. Height of closed points on asymptotically proper families of topological adelic curves. In this subsection, we consider the following setting. Let (I, \leq_I) be a (non-empty) totally ordered set. Let $\mathbf{S} = (S_i = (K_i, \phi_i : \Omega_i \to M_{K_i}, \nu_i))_{i \in I}$ be a family of topological adelic curves with base space Ω_0 and family of structural morphisms $(\alpha_i : \Omega_i \to \Omega_0)_{i \in I}$ (cf. Definition 3.1.1). For any $i \in I$ and for any $\omega \in \Omega_i$, we denote by $A_{i,\omega}$ the finiteness ring of ω . Moreover, we denote

$$K_{\infty} := \varprojlim_{i \in I} K_i = \bigcap_{i \in I} K_i.$$

Let $\pi_{\infty} : X_{\infty} \to \operatorname{Spec}(K_{\infty})$ be a projective K_{∞} -scheme. For any $i \in I$, we denote $\pi_i : X_i := X_{\infty} \otimes_{K_{\infty}} K_i \to \operatorname{Spec}(K_i)$ and $\mathbf{X} := (X_i)_{i \in I}$. For any $i \in I$, for any $\omega \in \Omega_i$, we denote by $(A_{i,\omega}, \mathfrak{m}_{i,\omega}, \kappa_{i,\omega})$ the finiteness ring of ω . We also fix an equivalence relation \sim on $\mathcal{F}(I, \mathbb{R})$ which is compatible with the additive group structure and assume that the family \mathbf{S} is asymptotically proper w.r.t. \sim (cf. Definition 3.2.1).

Definition 10.2.1. Let $\overline{\mathbf{L}} = (\overline{L_i})_{i \in I}$ be an adelic line bundle on \mathbf{X} . For any $i \in I$, write $\overline{L_i} = (L_i, \varphi_i), L_{\infty} := \varprojlim_{i \in I} L_i$ and $\varphi := (\varphi_i)_{i \in I}$. Let P_{∞} be a closed point of X_{∞} . Then, for any $i \in I$, P_{∞} determines a closed point P_i of X_i and write $\mathbf{P} := (P_i)_{i \in I}$. We say that \mathbf{P} is the closed point of \mathbf{X} determined by P_{∞} .

Moreover, for any $s_{\infty} \in P_{\infty}^* L_{\infty}$, for any $i \in I$, we denote by s_i the image of s_{∞} in $P_i^* L_i$ and $\mathbf{s} := (s_i)_{i \in I}$. Then, for any $i \in I$, $P_i^* \overline{L_i} := (P_i^* L_i, P_i^* \varphi_i)$ defines an adelic vector bundle on **S** (condition (iii) in Definition 9.6.1 implies condition (iii) in Definition 6.4.1). Then define

$$\widehat{\deg}_{\mathbf{P}^*\varphi}(\mathbf{s}) : \left| \begin{array}{ccc} I & \longrightarrow & \mathbb{R} \\ i & \longmapsto & \widehat{\deg}_{P_i^*\varphi_i}(s_i) \end{array} \right.$$

As **S** is asymptotically proper, it follows that the class of $\overline{\deg}_{\varphi}(\mathbf{s})$ in $\mathcal{F}(I, \mathbb{R})/\sim$ is independent on the choice of s_{∞} . This class is denoted by $h_{\overline{\mathbf{L}}}(\mathbf{P})$ and called the *height* of **P** w.r.t. the adelic line bundle $\overline{\mathbf{L}}$.

Theorem 10.2.2. Let $\overline{\mathbf{L}^{(1)}}, \overline{\mathbf{L}^{(2)}}$ addic line bundles on \mathbf{X} . Let \mathbf{P} be the closed point of \mathbf{X} determined by a closed point P_{∞} of X_{∞} . Then the following assertions hold.

(1) We have the equality

$$h_{\overline{\mathbf{L}^{(1)}}+\overline{\mathbf{L}^{(2)}}}(\mathbf{P}) = h_{\overline{\mathbf{L}^{(1)}}}(\mathbf{P}) + h_{\overline{\mathbf{L}^{(2)}}}(\mathbf{P})$$

in $\mathcal{F}(I,\mathbb{R})/\sim$.

(2) Assume that the equivalence relation \sim is compatible with the partial ordering on $\mathcal{F}(I,\mathbb{R})$ induced by the usual ordering on \mathbb{R} . For j = 1, 2, write $\overline{\mathbf{L}^{(j)}} = (L_i^{(j)}, \varphi_i^{(j)})_{i \in I}$. Assume that, for any $i \in I$, $L_i^{(1)} = L_i^{(2)}$ and that $\varphi_i^{(1)}, \varphi_i^{(2)}$ have the same model family and that the function

$$d(\overline{\mathbf{L}^{(1)}}, \overline{\mathbf{L}^{(2)}}) : (i \in I) \mapsto \int_{\Omega_i}^{\overline{i}} d_\omega(\varphi_i^{(1)}, \varphi_i^{(2)}) \nu_i(\mathrm{d}\omega) \in \mathbb{R}$$

satisfies

$$d(\overline{\mathbf{L}^{(1)}}, \overline{\mathbf{L}^{(2)}}) \sim 0.$$

Then we have the equality

$$h_{\overline{\mathbf{L}^{(1)}}}(\mathbf{P}) = h_{\overline{\mathbf{L}^{(2)}}}(\mathbf{P})$$

in $\mathcal{F}(\mathbb{R}_{>0},\mathbb{R})/\sim$.

Proof. (1) For j = 1, 2, let $s_{\infty}^{(j)}$ be an element of $P^{\infty}L_{\infty}^{(j)}$ inducing a family $\mathbf{s}^{(j)} := (s_i^{(j)})i \in I$. Then $s_{\infty} := s_{\infty}^{(1)} \cdot s_{\infty}^{(2)}$ is an element of $P^{\infty}(L_{\infty}^{(1)} + L_{\infty}^{(2)})$ inducing a family $\mathbf{s} := (s_i^{(1)} \cdot s_i^{(2)})_{i \in i}$. As we have

$$\forall i \in I, \quad \widehat{\deg}_{\mathbf{P}^*(\varphi^{(1)} + \varphi^{(2)})}(\mathbf{s}) = \widehat{\deg}_{\mathbf{P}^*\varphi^{(1)}}(\mathbf{s}^{(1)}) + \widehat{\deg}_{\mathbf{P}^*\varphi^{(2)}}(\mathbf{s}^{(2)})$$

we can conclude using the fact the definition of the heights $h_{\overline{\mathbf{L}^{(1)}}+\overline{\mathbf{L}^{(2)}}}(\mathbf{P}), h_{\overline{\mathbf{L}^{(1)}}}(\mathbf{P}), h_{\overline{\mathbf{L}^{(2)}}}(\mathbf{P})$ is independent on the choice of $s_{\infty}^{(1)}, s_{\infty}^{(2)}$.

(2) Let $i \in I$. Denote $S_{i,P_i} := S_i \otimes_{K_i} \kappa(P_i) = (\kappa(P_i), \phi_{i,P_i} : \Omega_{i,P_i} \to M_{\kappa(P_i)}, \nu_{i,P_i})$. As in the proof of Theorem 10.1.2 (2), for any $\omega \in \Omega_i$, for any $x \in \pi_{\kappa(P_i)/K_i}^{-1}(\omega)$, we have

$$d_x(P_i^*\varphi_i^{(1)}, P_i^*\varphi_i^{(2)}) \le d_\omega(\varphi_i^{(1)}, \varphi_i^{(2)})$$

Let s_{∞} be an element of $P^{\infty}L_{\infty}$ inducing a family $\mathbf{s} := (s_i)i \in I$. Then we have

$$\forall i \in I, \quad |\widehat{\operatorname{deg}}_{\mathbf{P}^*\varphi^{(1)}}(\mathbf{s})(i) - \widehat{\operatorname{deg}}_{\mathbf{P}^*\varphi^{(2)}}(\mathbf{s})(i)| \le d(\overline{\mathbf{L}^{(1)}}, \overline{\mathbf{L}^{(2)}})(i)$$

Using the assumption on the equivalence relation \sim , we obtain the desired result.

Example 10.2.3. We consider the totally ordered set $\mathbb{R}_{>0}$ equipped with the usual total ordering. Consider the family of topological adelic curves $\mathbf{S} = (S_R)_{R>0}$ constructed in Example 3.1.2 (2). We consider the equivalence relation $\sim_{O(1)}$ on $\mathcal{F}(\mathbb{R}_{>0},\mathbb{R})$ with respect to which the family \mathbf{S} is asymptotically proper (cf. 3.2.2).

Let X be a reduced projective \mathbb{C} -scheme and let (L, φ) be a metrised line bundle on X, where L is assumed to be globally generated. For any $R \in \mathbb{R}_{>0} \cup \{\infty\}$, we denote $X_R := X \otimes_{\mathbb{C}} K_R$ (recall that $K_{\infty} = \mathcal{M}(\mathbb{C})$ the field of meromorphic functions on \mathbb{C}). In Example 9.6.3, we have constructed an adelic line bundle $\overline{\mathbf{L}} := (\mathbf{L}, \varphi) := (L_R, \varphi_R)_{R>0}$ on $\mathbf{X} := (X_R)_{R>0}$.

Let $\mathbf{P} = (P_R)_{R>0}$ be the closed point of \mathbf{X} determined by a rational point $P_{\infty} \in X_{\infty}(K_{\infty}) = X(\mathcal{M}(\mathbb{C}))$. From the point of view explained in §A.2, P_{∞} corresponds to a holomorphic curve $f : \mathbb{C} \to X$ and for any R > 0, P_R corresponds to the restriction of f to the closed disc of radius R in \mathbb{C} .

Let s be an regular meromorphic section of L such that $f(\mathbb{C}) \not\subseteq |\operatorname{div}(s)|$. For any $R \in \mathbb{R}_{>0} \cup \{\infty\}$, denote y s_i the regular meromorphic section of L_R determined by pullback of s and write $\mathbf{s} := (s_R)_{R>0}$. Then, for any R > 0, we have the equality

$$\deg_{\mathbf{P}^*\varphi}(\mathbf{s}) = T_{f,(L,\varphi,s)}(R),$$

(cf. §A.2 for the notation $T_{f,(L,\varphi,s)}(R)$).

Note that the equivalence relation $\sim_{O(1)}$ is compatible with the partial ordering on $\mathcal{F}(\mathbb{R}_{>0},\mathbb{R})$ and that for any other continuous metric φ' on L determining another adelic line bundle $\overline{\mathbf{L}'}$ on \mathbf{X} , we have

$$\forall R>0, \quad d(\overline{L},\overline{L'})(R) \leq \left|\max_{x\in X(\mathbb{C})}\log\frac{|s|_{\varphi'}(x)}{|s|_{\varphi}(x)}\right| < +\infty,$$

where s denotes an arbitrary regular meromorphic section of L such that $f(\mathbb{C}) \nsubseteq |\operatorname{div}(s)|$. Hence the function $d(\overline{L}, \overline{L'})$ is bounded.

Therefore, Theorem 10.2.2 gives a generalisation of Theorem A.2.1 in our context (see also Theorem 3.18 in [Gub97]).

APPENDIX A. NEVANLINNA THEORY OF COMPLEX FUNCTIONS

A.1. Classical Nevanlinna theory. We denote $K = \mathcal{M}(\mathbb{C})$, namely the field of meromorphic functions on \mathbb{C} , which is the fraction field of the ring of entire functions $A := \mathcal{O}(\mathbb{C})$. Let $E = \sum a_i[z_i]$ be a divisor on \mathbb{C} , with the z_i pairwise distinct. For any $k \in \mathbb{N} \cup \{\infty\}$, we define the truncated counting functions

$$\forall t > 0, \quad n_k(t, E) := \sum_{|z_i|_{\infty} < t} \min\{k, a_i\},$$

and

$$\forall r > 1, \quad N_k(r, E) := \int_1^r \frac{n_k(t, E)}{t} dt.$$

For ease of notation, we let $n(t, E) := n_{\infty}(t, E)$ and $N(r, E) := N_{\infty}(r, E)$.

Let $f \in K$, for any $a \in \mathbb{P}^1(\mathbb{C})$, we denote by $(f)_a$ the divisor of associated to (f-a), if $a \neq \infty$, and to 1/f if $a = \infty$. Then the *proximity function* in ∞ of f is defined by

$$\forall r>0, \quad m(r,f):=\frac{1}{2\pi}\int_0^{2\pi}\log^+|f(re^{i\theta})|_\infty d\theta.$$

The function $(r \in \mathbb{R}_{>0}) \mapsto m(r, 1/(f-a))$ is called the proximity function of f in $a \in \mathbb{C}$. Finally, the *height* of f (with respect to ∞) is defined by

$$\forall r > 1, \quad T(r, f) := m(r, f) + N(r, (f)_{\infty}).$$

Morally, the proximity function of f measures the mean approximation of f to ∞ on a circle of given radius. The counting function counts how many times f attains ∞ , i.e. has a pole, in an open disc of given radius. The philosophy of Nevanllina is that these two functions contain all the necessary information concerning the behaviour of f with respect to ∞ . The characteristic function, namely the sum of the two previous ones, behaves as a height function in Diophantine geometry:

$$\forall f_1, f_2 \in K, \quad T(r, f_1 + f_2) \le T(r, f_1) + T(r, f_2) + \log 2.$$

Theorem A.1.1 (Nevanlinna's first main theorem, [NW14], Theorem 1.1.17). Let $f \in K$ and $a \in \mathbb{C}$. Then

$$T(r, \frac{1}{f-a}) = T(r, f) + O(1),$$

where the bound O(1) is a bounded function of r, with bound depending only on f and a.

Morally, Nevanlinna's inequality, which is itself a consequence of Nevanlinna's first theorem, gives an upper bound for the counting function in terms of height.

In fact, other height functions may be defined. For our purposes, it is more convenient to work in a more geometric framework, namely, we will consider any $f \in K$ as a holomorphic

curve $f : \mathbb{C} \to \mathbb{P}^1(\mathbb{C})$. Let *L* be a holomorphic line bundle on $P^1(\mathbb{C})$, equipped with a hermitian metric φ and an invertible meromorphic section *s*. Denote $D := \operatorname{div}(s)$. We define

$$\begin{split} N_f(r,D) &:= \operatorname{ord}(f^*D,0) \log r + \sum_{0 < |z|_{\infty} < r} \operatorname{ord}(f*D,z) \log \left| \frac{r}{z} \right|_{\infty}, \\ m_f(r,D) &:= -\frac{1}{2\pi} \int_0^{2\pi} \log |s(f(re^{i\theta}))|_{\varphi} d\theta, \\ T_f(r,D) &:= N_f(r,D) + m_f(r,D). \end{split}$$

The previous constructions are now the special case $L = \mathcal{O}(1)$, φ the standard metric, and $D = [\infty]$. Then Nevanlinna's first main theorem is rephrased as follows. Consider L and f fixed. Then the height function $T_f(r, D)$ does not depend, up to a bounded function of r, neither of the choices of the metric φ nor of the meromorphic section s.

The Northcott property has a counterpart in Nevanlinna theory: this is the Liouville theorem, which states that an entire function is constant iff T(r, f) is a bounded function of r.

Let $f \in K$. Define the order of f as

$$\rho(f) := \limsup_{r \to \infty} \frac{T(r, f)}{\log r}.$$

Nevanlinna's first theorem shows that this quantity does not depend on the choice of height function.

As it was previously mentioned, Nevanlinna's first theorem yields an upper bound for counting functions of meromorphic functions. Nevanlinna's second theorem gives a result in the other direction: namely it yields a lower bound. To state it, we need consider multiple proximity functions.

Theorem A.1.2 (Nevanlinna's second theorem, [NW14], Theorem 1.2.5). Let $f \in K$. Let $q \ge 1$ be an integer and let $a_1, ..., a_q \in \mathbb{P}^1(\mathbb{C})$. Then

$$\sum_{i=1}^{q} m_f(r, a_i) \le_{exc} 2T_f(r) + O(\log^+ T_f(r)) + o(\log r),$$

or, equivalently,

$$\sum_{i=1}^{q} N_f(r, (f)_{a_i}) \ge_{exc} (q-2)T_f(r) - O(\log^+ T_f(r)) - o(\log r),$$

where \leq_{exc} means that the inequalities hold for all r > 0 except on a set of finite Lebesgue measure.

Roughly speaking, Nevanlinna's second main theorem says that a meromorphic function on \mathbb{C} cannot avoid "too many points", in that case, at most 2.

It is possible to measure to what extent the counting function at a point is significantly smaller than the height via the so-called defect. More precisely, let $f \in K$, and let $a \in \mathbb{P}^1(\mathbb{C})$. Then the *defect* of f at a is defined by

$$\delta_f(a) := \liminf_{r \to \infty} \frac{m_f(r, a)}{T_f(r)} = 1 - \limsup_{r \to \infty} \frac{N_f(r, (f)_a)}{T_f(r)}$$
where the second equality comes from Nevanlinna's first theorem. The same theorem implies that we have the inequality $0 \le \delta_f(a) \le 1$ and Nevanlinna's second theorem yields

$$\sum_{a \in \mathbb{P}^1(\mathbb{C})} \delta_f(a) \le 2$$

In the $\delta_f(a) > 0$ case, we say that a is a *deficient* value for f. With this notion at hand, Nevanlinna's second theorem implies that a given meromorphic function cannot have too many deficient values.

A.2. Holomorphic curves on a projective variety. In this subsection, we fix a complex projective variety X and we study holomorphic maps $f : \mathbb{C} \to X$. The latter are called *holomorphic curves* in X. Since X is projective, f induces a holomorphic map $f : \mathbb{C} \to \mathbb{P}^n(\mathbb{C})$ and Weierstrass factorisation theorem ([Ahl66], Chapter 5, Theorem 8) implies that there exist entire functions $f_0, ..., f_n$ without common zeroes such that $f(x) = [f_0(x) : \cdots : f_n(x)]$ for all $x \in \mathbb{C}$. This way we can see $f : \mathbb{C} \to X$ as a K-point $f \in X(K)$, where $K = \mathcal{M}(\mathbb{C})$. In this context, we can extend the definitions of counting, proximity and height functions.

Let D be a divsor on X. Assume that $f(\mathbb{C}) \not\subset |D|$, so that f^*D is a divisor on \mathbb{C} . Define the *counting function*

$$N_f(r, D) := \operatorname{ord}(f^*D, 0) \log r + \sum_{0 < |z|_{\infty} < r} \operatorname{ord}(f^*D, z) \log \frac{r}{|z|_{\infty}},$$

for all r > 0. Denote $L := \mathcal{O}_X(D)$ and let s be a regular meromorphic section of de L such that $D = \operatorname{div}(s)$. Let φ be a continuous Hermitian metric on L. Define the proximity function

$$m_f(r, (L, \varphi, s)) := -\frac{1}{2\pi} \int_0^{2\pi} \log |s|_{\varphi}(f(re^{i\theta})) d\theta,$$

for all r > 0. Finally, the *height function* is defined by

$$T_{f,(L,\varphi,s)}(r) := m_f(r,(L,\varphi,s)) + N_f(r,D),$$

for all r > 0.

We can now state the two main theorems of Nevanlinna theory for holomorphic curves.

Theorem A.2.1 (Nevanlinna's first theorem ([BG06], Theorem 13.2.9)). Let $f : \mathbb{C} \to X$ be a holomorphic curve, with X a projective complex variety. Let L be a line bundle on X, let s be a regular meromorphic section of L and let φ be a continuous Hermitian metric on L. Denote $D := \operatorname{div}(s)$ and assume that $f(\mathbb{C}) \not\subset |D|$. This data defines a height function $(r > 0) \mapsto T_{f,(L,\varphi,s)}(r) \in \mathbb{R}$.

(1) Let φ' be another continuous Hermitian metric on L. Then we have

$$T_{f,(L,\varphi,s)}(r) - T_{f,L,\varphi',s}(r) = \frac{1}{2\pi} \int_0^{2\pi} \log \frac{|s|_{\varphi'}}{|s|_{\varphi}} (f(re^{i\theta})) d\theta,$$

for all r > 0. In particular, the LHS in the above equality is a bounded function of r.

(2) Let s' be another regular meromorphic section of L such that $f(\mathbb{C}) \not\subset |\operatorname{div}(s')|$. Then we have

$$T_{f,(L,\varphi,s)}(r) - T_{f,L,\varphi,s'}(r) = \log |c((s'/s) \circ f, 0)|_{\infty},$$

ANTOINE SÉDILLOT

for all r > 0, where $c((s'/s) \circ f, 0)$ denotes the first non-zero coefficient in the Laurent series expansion of the meromorphic function $(s'/s) \circ f$.

As in the complex case, we study the asymptotic behaviour of the height function. The latter tends to $+\infty$ as $r \to +\infty$ if the holomorphic curve is not constant. Therefore, we consider height functions associated with different choices of metrics and regular meromorphic sections as equivalent. Properties of height functions are similar to the ones appearing in Diophantine geometry.

Proposition A.2.2. Let $f : \mathbb{C} \to X$ be a holomorphic curve, where X denotes a complex projective variety.

(1) Let (L_1, φ_1, s_1) , (L_2, φ_2, s_2) be continuous Hermitian metrised Cartier divisors on X. Let φ_{1+2} be a continuous Hermitian metric on $L_1 \otimes L_2$. Assume that $f(\mathbb{C}) \not\subset |\operatorname{div}(s_1)| \cup |\operatorname{div}(s_2)|$. Then

 $T_{f,(L_1\otimes L_2,\varphi_{1+2},s_1\otimes s_2)}(r) = T_{f,(L_1,\varphi_1,s_1)}(r) + T_{f,(L_2,\varphi_2,s_2)}(r) + O(1).$

(2) Let $\alpha : X \to Y$ be a morphism between complex projective varieties. Let (L, φ) be a continuous Hermitian line bundle Y and let s be a regular meromorphic section of L. Assume that $f(\mathbb{C}) \not\subset |\operatorname{div}(s)|$ and that $\alpha(X) \not\subset |\operatorname{div}(s)|$. Then

$$T_{f\circ\alpha,(L,\varphi,s)}(r) = T_{f,(f^*L,f^*\varphi,f^*s)}(r) + O(1).$$

(3) Let (L, φ) be a Hermitian line bundle L on X which is globally generated. Let s be a regular meromorphic section of L. Then the height function $T_{f,(L,\varphi,s)}$ is bounded from below.

In the context of holomorphic curves, Nevanlinna's second theorem is not known. Its conjectural statement is known as the Griffith conjecture.

Conjecture A.2.3 (Griffiths' Conjecture). Let A be an ample line bundle on a complex projective variety X. Denote $K_X := \wedge^{\dim(X)} T_X^*$ the canonical line bundle on X. Let D be a normal crossing divisor on X.

(1) Then, for any holomorphic curve $f : \mathbb{C} \to X$ with Zariski dense image, the inequality

$$m_{f,D}(r) + T_{f,K_X}(r) \leq_{\text{exc}} O(\log^+ T_{f,A}(r)) + o(\log(r))$$

holds.

(2) For any $\epsilon > 0$, there exists an algebraic subset $Z \nsubseteq X$ such that, for any holomorphic curve $f : \mathbb{C} \to X$ such that $f(\mathbb{C}) \not\subset Z$, we have

$$\forall C \in \mathbb{R}, \quad m_{f,D}(r) + T_{f,K_X}(r) \leq_{\text{exc}} \epsilon T_{f,A}(r) + C.$$

This conjecture is known in the case where X is a curve and in the $X = \mathbb{P}^n(\mathbb{C})$ case (cf. [Voj97] refining a result of Cartan). In the recent preprint [DH22], Dong and Hu announced a proof of the Griffiths conjecture. Unfortunately, there seems to be a gap in the paper.

A.3. Analogy with Diophantine approximation. We conclude this appendix by giving the idea of the (heuristic) analogy between Diophantine approximation and Nevanlinna theory. For more details, we refer to [Voj87, BG06, Voj10].

74

Diophantine approximation	Nevanlinna theory
Z	$\mathcal{O}(\mathbb{C})$
Q	$\mathcal{M}(\mathbb{C})$
$\{b_i: i \in I\} \subset \mathbb{Q}$ infinite	$f \in \mathcal{O}(\mathbb{C})$ non constant
$i \in I$	r > 0
$\{ \cdot _{\infty}\} \subset S$ finite set of absolute values on \mathbb{Q}	$\{\theta: \theta \in [0, 2\pi]\}$
$M_{\mathbb{Q}}\smallsetminus S$	$\{z \in \mathbb{R} : z _{\infty} < r\}$
$ b_i _v, v \in S$	$ f(re^{i\theta}) _{\infty}, \ \theta \in [0, 2\pi]$
$ b_i _v, v \notin S$	$\operatorname{ord}(f, z), z _{\infty} < r$
Height	Characteristic function
$h(b_i) = \sum_{v \in M_{\mathbb{Q}}} \log^+ b_i _v$	$T(f,\infty,r)$
Proximity function	Proximity function
$\forall a \in \mathbb{Q}, m_S(a, b_i) = \sum_{v \in S} \log^+ \frac{1}{b_i - a} _v$	$\forall a \in \mathbb{C}, m(f, a, r)$
Counting function	Counting function
$\forall a \in \mathbb{Q}, N_S(a, b_i) = \sum_{v \notin S} \log^+ \frac{1}{b_i - a} _v$	$\forall a \in \mathbb{C}, N(f, a, r)$
Product formula	Jensen's formula
$\sum_{v \in S} \log b_i _v = 0$	$T(f, \infty, r) - T(f, 0, r) = \log c(f, 0) _{\infty}$
Height theory	First main theorem
$\forall a \in \mathbb{Q}, \ m_S(a, b_i) + N_S(a, b_i) = h(b_i) + O(1)$	$\forall a \in \mathbb{C}, T(f, a, r) = T(f, \infty, r) + O(1)$
Roth's theorem	Second main theorem (weak form)
$\forall \epsilon > 0, \forall m \in \mathbb{Z} \ge 1, \forall a_1,, a_m \in \mathbb{Q},$	$\forall \epsilon > 0, \forall m \in \mathbb{Z}_{\geq 1}, \forall a_1,, a_m \in \mathbb{C},$
$\sum_{j=1}^{n} m(b_i, a_j) \leq_{\text{exc}} (2+\epsilon)h(b_i)$	$\left \sum_{j=1}^{n} m(f, a_j, r) \leq_{\text{exc}} (2+\epsilon) T(f, \infty, r)\right $
Defect	Defect
$\forall a \in \mathbb{Q}, \delta(a) := \liminf_{i \in I} \frac{m_S(a, b_i)}{h(b_i)}$	$\forall a \in \mathbb{C}, \delta(a) := \liminf_{r \to +\infty} \frac{m(f, a, r))}{T(f, \infty, r)}$

In the line concerning Roth's theorem, \leq_{exc} means that the inequality holds for all $i \in I$ except a finite number.

References

- [Ahl66] Lars Valerian Ahlfors. Complex Analysis. McGraw-Hill Book Company, 2nd edition, 1966.
- [Ara74] Souren Yu Arakelov. Intersection theory of divisors on an arithmetic surface. Math. USSR Izvestija, 8:1167–1180, 1974.
- [BC13] Jean-Benoît Bost and Huayi Chen. Concerning the semistability of tensor products in Arakelov geometry. Journal des Mathématiques Pures et Appliquées, 99(4), 2013.
- [BE21] Sébastien Boucksom and Dennis Eriksson. Spaces of norms, determinant of cohomology and Fekete points in non-Archimedean geometry. *Advances in Mathematics*, 378:107501, February 2021.
- [Ber90] Vladimir G. Berkovich. Spectral Theory and Analytic Geometry over Non-Archimedean Fields. American Mathematical Soc., 1990.
- [BG06] Enrico Bombieri and Walter Gubler. *Heights in Diophantine Geometry*. New Mathematical Monographs 4, Cambridge University Press, Cambridge, 2006.
- [BGPS16] José Ignacio Burgos Gil, Patrice Philippon, and Martín Sombra. Height of varieties over finitely generated fields. *Kyoto Journal of Mathematics*, 56(1), April 2016.
- [Bou71] Nicolas Bourbaki. Topologie Générale. Springer-Verlag Berlin Heidelberg, 1971.
- [BYDHS24] Itaï Ben Yaacov, Pablo Destic, Ehud Hrushovski, and Michał Szachniewicz. Globally valued fields: foundations, 2024. arXiv:2409.04570.
- [CG24] Yulin Cai and Walter Gubler. Abstract divisorial spaces and arithmetic intersection numbers, 2024. arXiv:2409.00611.

ANTOINE SÉDILLOT

- [CJ23] Huayi Chen and Marion Jeannin. Harder-Narasimhan games, 2023. arXiv:2306.08283.
 [CL06] Antoine Chambert-Loir. Mesures et équidistribution sur les espaces de Berkovich. Journal für
- die reine und angewandte Mathematik, 2006(595):215–235, 2006. [CLD12] Antoine Chambert-Loir and Antoine Ducros. Formes différentielles réelles et courants sur les
- [CLD12] Antoine Chambert-Loir and Antoine Ducros. Formes différentielles réelles et courants sur les espaces de Berkovich, 2012. arXiv:1204.6277.
- [CM19] Huayi Chen and Atushi Moriwaki. Arakelov geometry over adelic curves, volume 2258 of Lectures Notes in Mathematics. Springer Singapore, 2019.
- [CM21] Huayi Chen and Atushi Moriwaki. Arithmetic intersection theory over adelic curves, 2021. arXiv:2103.15646.
- [CM22] Huayi Chen and Atsushi Moriwaki. Hilbert-Samuel formula and positivity over adelic curves, 2022. arXiv:2207.02033.
- [CM24] Huayi Chen and Atsushi Moriwaki. *Positivity in Arakelov Geometry over Adelic Curves*. Progress in Mathematics. Birkhäuser Cham, 2024.
- [DH22] Xianjing Dong and Peichu Hu. On Griffiths conjecture, 2022. arXiv:2203.10985.
- [DHS24] Pablo Destic, Nuno Hultberg, and Michał Szachniewicz. Continuity of heights in families and complete intersections in toric varieties, 2024. arXiv:2412.15988.
- [DZ23] Paolo Dolce and Francesco Zucconi. On the generalisation of Roth's theorem, 2023. arXiv:2111.12409.
- [Fal91] Gerd Faltings. Diophantine Approximation on Abelian Varieties. Annals of Mathematics, 133(3):549–576, 1991.
- [GK17] Walter Gubler and Klaus Künnemann. A tropical approach to nonarchimedean Arakelov geometry. Algebra and Number Theory, 11(1):77–180, Jan 2017.
- [GK19] Walter Gubler and Klaus Künnemann. Positivity properties of metrics and delta-forms. Journal fur die Reine und Angewandte Mathematik, 2019(752), 2019.
- [GR17] Eric Gaudron and Gaël Rémond. Corps de Siegel. Journal für die reine und angewandte Mathematik, 2017(726):187–247, January 2017.
- [Gub97] Walter Gubler. Heights of subvarieties over M-fields. In Arithmetic geometry (Cortona 1994), pages p. 190–227. Cambridge University Press, 1997.
- [GW10] Ulrich Görtz and Torsten Wedhorn. Algebraic geometry. I, Schemes with examples and exercises. Springer Studium Mathematik - Master. Springer Spektrum Wiesbaden, 2nd ed. 2020 edition, 2010.
- [Hru16] Ehud Hrushovski. A logic for global fields. Séminaire d'Arithmétique et de Géométrie Algébrique d'Orsay, 2016.
- [Lan74] Serge Lang. Higher dimensional diophantine problems. Bulletin of the American Mathematical Society, 80(5):779–787, 1974.
- [Lan86] Serge Lang. Hyperbolic and Diophantine analysis. Bulletin (New Series) of the American Mathematical Society, 14(2):159–205, 1986.
- [Lan91] Serge Lang. Number Theory III: Diophantine geometry. (Encyclopedia of mathematical sciences 60). Springer, Berlin-Heidelberg, 1991.
- [LP24] Thibaud Lemanissier and Jérôme Poineau. Espaces de Berkovich Globaux. Progress in Mathematics. Birkhäuser Cham, 2024.
- [Mor00] Atushi Moriwaki. Arithmetic height functions over finitely generated fields. *Inventiones Mathematicae*, 2000.
- [NW14] Junjiro Noguchi and Jörg Winkelmann. Nevanlinna Theory in Several Complex Variables and Diophantine Approximation. Springer Japan, 2014.
- [Osg81] C.F. Osgood. A number-theoretic-differential equations approach to generalizing Nevanlinna theory. Indian Journal of Mathematics, 23(1-3):1–15, 1981.
- [Pau09] Frédéric Paugam. Global analytic geometry. Journal of Number Theory, 129(10):2295 2327, 2009.
- [Rot55] Klaus Friedrich Roth. Rational approximations to algebraic numbers. *Mathematika*, 1955.
- [RT96] Damien Roy and Jeffrey L Thunder. An absolute Siegel's Lemma. Journal f
 ür die reine und angewandte Mathematik, 1996(476):1–26, 1996.
- [Séd23] Antoine Sédillot. Differentiability of the χ -volume function over an adelic curve, 2023. arXiv:2303.03377.

76

- [Séd24a] Antoine Sédillot. Pseudo-absolute values: foundations, 2024. arXiv:2411.03905.
- [Séd24b] Antoine Sédillot. Study of projective varieties over adelic curves. Theses, Université Paris Cité, July 2024. https://theses.hal.science/tel-04990131v1/file/va_Sedillot_Antoine.pdf.
- [Sza23] Michał Szachniewicz. Existential closedness of $\bar{\mu}$ as a globally valued field via arakelov geometry, 2023. arXiv:2306.06275.
- [Ull98] Emmanuel Ullmo. Positivité et Discrétion des Points Algébriques des Courbes. Annals of Mathematics, 147(1):167–179, 1998.
- [Voj87] Paul Vojta. Diophantine Approximations and Value Distribution Theory. Lecture Notes in Mathematics 1239. Springer-Verlag Berlin Heidelberg, 1987.
- [Voj97] Paul Vojta. On Cartan's Theorem and Cartan's Conjecture. American Journal of Mathematics, 119(1):1–17, 1997.
- [Voj10] Paul Vojta. Diophantine Approximation and Nevanlinna Theory, pages 111–224. Springer Berlin Heidelberg, Berlin, Heidelberg, 2010.
- [Voj21] Paul Vojta. Roth's Theorem over arithmetic function fields. Algebra and Number Theory, 15(8):1943–2017, November 2021.
- [Wei51] André Weil. Arithmetic on algebraic varieties. Annals of Mathematics, 53(3):412–444, 1951.
- [YZ21] Xinyi Yuan and Shou-Wu Zhang. Adelic line bundles over quasi-projective varieties, 2021. arXiv:2105.13587.
- [Zha95] Shou-Wu Zhang. Positive line bundles on arithmetic vaireties. Journal of the American Mathematical Society, 1995.
- [Zha98] Shou-Wu Zhang. Equidistribution of Small Points on Abelian Varieties. Annals of Mathematics, 147(1):159–165, 1998.

A. SÉDILLOT, MATHEMATIK, UNIVERSITÄT REGENSBURG, 93040 REGENSBURG, GERMANY *Email address:* antoine.sedillot@mathematik.uni-regensburg.de