NONLINEAR ASYMPTOTIC STABILITY OF 2D TAYLOR-COUETTE FLOW IN THE EXTERIOR DISK

TE LI, PING ZHANG, AND YIBIN ZHANG

ABSTRACT. In this paper, we consider the asymptotic stability of the 2D Taylor-Couette flow in the exterior disk, with a small kinematic viscosity $\nu \ll 1$ and a large rotation coefficient |B|. Due to the degeneracy of the Taylor-Couette flow at infinity, we cannot expect the solution to decay exponentially in a space-time decoupled manner. As stated in previous work [20], even space-time coupled exponential decay can not be expected, and at most, we can obtain space-time coupled polynomial decay. To handle the space-time coupled decay multiplier, the previous time-independent resolvent estimate methods no longer work. Therefore, this paper introduces time-dependent resolvent estimates to deal with the spacetime coupled decay multiplier Λ_k . We remark that the choice of Λ_k is not unique, here we just provide one way to construct it. Finally, as an application, we derive a transition threshold bound of $\frac{1}{2}$, which is the same as that for the Taylor-Couette flow in the bounded region.

Contents

1. Introduction	2
1.1. Derivation of the perturbation equation	3
1.2. Key ideas	4
1.3. Main theorems	5
1.4. Notations	8
2. Fundamental energy estimates	9
3. Resolvent estimates without enhanced dissipation weight	12
4. Resolvent estimates with enhanced dissipation weight	20
5. Space-time estimates of the vorticity in nonzero modes	23
5.1. Space-time estimates of the inhomogeneous equation (5.2)	24
5.2. Space-time estimates of the homogeneous equation (5.1)	25
6. Proof of Theorem 1.3 and Theorem 1.4.	32
Appendix A. Local wellposedness of (1.3)	37
Appendix B. Basic Calculus	38
Appendix C. Basic properties of \mathcal{K}_k and Λ_k	40
Acknowledgement	42
Declarations	43
Conflict of interest	43
Data availability	43
References	43

Date: March 27, 2025.

1. INTRODUCTION

We consider 2D incompressible Navier-Stokes equations in the exterior disk $\Omega \stackrel{\text{def}}{=} \{x \in \mathbb{R}^2 : |x| \ge 1\}$:

(1.1)
$$\begin{cases} \partial_t V - \nu \Delta V + V \cdot \nabla V + \nabla P = 0, \quad (t, x) \in \mathbb{R}^+ \times \Omega, \\ \operatorname{div} V = 0, \\ V(0, x) = V_0(x), \end{cases}$$

where $V \in \mathbb{R}^2$ denotes the fluid velocity and P the scalar pressure function, which guarantees the divergence free condition of the velocity field, ν designates the kinematic viscosity of the fluid.

In this paper, we study the stability of a two-dimensional rotating flow in the exterior region Ω , which is a steady solution to (1.1). The corresponding velocity, vorticity, and scalar pressure are given as follows:

(1.2a)
$$V^{\star}(r,\theta) = \begin{pmatrix} -\sin\theta\\\cos\theta \end{pmatrix} \left(Ar + \frac{B}{r}\right) \stackrel{\text{def}}{=} \begin{pmatrix} v^{\star,1}\\v^{\star,2} \end{pmatrix}, \quad W^{\star}(r) \stackrel{\text{def}}{=} \partial_1 v^{\star,2} - \partial_2 v^{\star,1} = 2A,$$

(1.2b)
$$P^{\star}(r,\theta) = P^{\star}(r) \quad \text{with} \quad \partial_r P^{\star}(r) = \frac{1}{r} \left(Ar + \frac{B}{r} \right)^2$$

where A, B are two constants and we assume that $\nu \ll |B|$ throughout this paper. The steady flow of the form (1.2a) is well-known as Taylor-Couette flow.

As a classic example of rotating flow, Taylor-Couette (TC) flow has been extensively studied in both the physical and mathematical contexts. In physical experiments, TC flow always refers to the fluid flow between two concentric cylinders, where the inner cylinder rotates while the outer cylinder can be either stationary or rotating. This flow configuration was first studied by G. I. Taylor in the 1920s [25]. Despite its simple structure, TC flow exhibits complex and rich stability properties, and has become a classic problem in fluid dynamics due to its rich flow patterns, as referenced in such review books as [12, 10, 23, 24, 18]. These corresponding flow patterns are controlled by the Reynolds number and the radius ratio of the cylinders. For example, the TC apparatus is also an excellent prototype for studying the transport properties of most astrophysical or geophysical rotating shear flows [14]. Depending on the rotational speed of each cylinder, different flow states can be achieved, corresponding to increasing or decreasing angular velocity and angular momentum. Likewise, regarding the issue of asymptotic stability at high Reynolds numbers [6], TC flow, as a typical rotating shear flow, also exhibits rich decay properties.

This paper primarily investigates the fully nonlinear long-time asymptotic stability behavior near the TC flow in the exterior disk. Earlier, the well-posedness and stability of the two-dimensional stationary Navier-Stokes equations in exterior regions [5, 16] were widely studied and discussed. The following results are related to the fully nonlinear stability of the time-dependent Navier-Stokes equations near the 2D TC flow.

- Under the condition $|B| \ll 1$ and with the domain being the exterior disk, [21, 22] used perturbation theory to study the $L^p L^q$ estimate for the semigroup generated by the linearized operator, and further addressed the global well-posedness and nonlinear asymptotic stability for small initial data in the L^2 space.
- In [2], nonlinear enhanced dissipation and transition thresholds near the TC flow were obtained in the bounded annular region $(r, \theta) \in [1, R] \times \mathbb{T}$. Since the domain is compact, the shear effect of the TC flow is non-degenerate, i.e., $(A + \frac{B}{r^2})'$ is non-vanishing.

In the case when ν is sufficiently small and |B| is large enough, that is, when the Reynolds number is high and the rotational effects are strong, the fully nonlinear long-time asymptotic stability problem of the 2D TC flow in the exterior disk is both interesting and challenging.

1.1. Derivation of the perturbation equation. Let (V^*, P^*) be determined by (1.2). We denote $U \stackrel{\text{def}}{=} V - V^*$, $p \stackrel{\text{def}}{=} P - P^*$, and consider the following perturbation equations of the system (1.1) near the 2D TC flow:

(1.3)
$$\begin{cases} \partial_t U - \nu \Delta U + V^* \cdot \nabla U + U \cdot \nabla V^* + U \cdot \nabla U + \nabla p = 0, \\ \operatorname{div} U = 0, \quad (t, x) \in \mathbb{R}^+ \times \Omega, \\ U(t = 0) = U(0). \end{cases}$$

We shall present a local well-posedness result concerning the system (1.3) in Appendix A.

Since this paper aims at investigating the impact of degeneracy, which is caused by the external region, on the nonlinear stability of TC flow, we impose the impermeable and Navierslip boundary conditions for the velocity field:

(1.4)
$$U \cdot \mathbf{n}|_{\partial\Omega} = 0, \quad \left(\partial_1 U^2 - \partial_2 U^1\right)|_{\partial\Omega} = 0.$$

Let $W \stackrel{\text{def}}{=} \operatorname{curl} U \stackrel{\text{def}}{=} \partial_1 U^2 - \partial_2 U^1$ be the vorticity of U, and we rewrite (1.3-1.4) in the equivalent vorticity-stream formulation:

(1.5)
$$\begin{cases} \partial_t W - \nu \Delta W + V^* \cdot \nabla W + U \cdot \nabla W = 0, & (t, x) \in \mathbb{R}^+ \times \Omega, \\ \Delta \Phi = W, \quad U = \nabla^\perp \Phi^{\text{def}}_{=} \left(-\partial_2 \Phi, \partial_1 \Phi \right)^T, \\ W|_{\partial\Omega} = \Phi|_{\partial\Omega} = 0, \quad W(t = 0) = W(0). \end{cases}$$

Let $\mathbf{e_r} = (\cos \theta, \sin \theta)$ and $\mathbf{e}_{\theta} = (-\sin \theta, \cos \theta)$. In polar coordinates, (1.5) is equivalent to

$$(1.6) \qquad \begin{cases} \partial_t W - \nu \left(\partial_r^2 + \frac{1}{r} \partial_r + \frac{1}{r^2} \partial_\theta^2 \right) W + \left(A + \frac{B}{r^2} \right) \partial_\theta W + \frac{1}{r} (\partial_r \Phi \partial_\theta W - \partial_\theta \Phi \partial_r W) = 0, \\ \left(\partial_r^2 + \frac{1}{r} \partial_r + \frac{1}{r^2} \partial_\theta^2 \right) \Phi = W, \quad (t, r, \theta) \in \mathbb{R}^+ \times [1, \infty) \times \mathbb{T}, \\ U = U^r \mathbf{e_r} + U^\theta \mathbf{e_\theta} \quad \text{with} \quad U^r = -\frac{\partial_\theta \Phi}{r} \quad \text{and} \quad U^\theta = \partial_r \Phi, \\ W(t = 0) = W(0). \end{cases}$$

By expanding the functions $(W, \Phi, U^r, U^{\theta})$ in Fourier series, we write

(1.7)

$$W(t,r,\theta) = \sum_{k \in \mathbb{Z}} e^{ik\theta} \widehat{W}_k(t,r), \quad \Phi(t,r,\theta) = \sum_{k \in \mathbb{Z}} e^{ik\theta} \widehat{\Phi}_k(t,r),$$

$$U^r(t,r,\theta) = \sum_{k \in \mathbb{Z}} e^{ik\theta} \widehat{U}_k^r(t,r), \quad U^\theta(t,r,\theta) = \sum_{k \in \mathbb{Z}} e^{ik\theta} \widehat{U}_k^\theta(t,r),$$

Then by substituting (1.7) into (1.6) and comparing the Fourier coefficients of the resulting equations, we obtain

$$(1.8) \qquad \begin{cases} \partial_t \widehat{W}_k - \nu \Big(\partial_r^2 + \frac{1}{r} \partial_r - \frac{k^2}{r^2}\Big) \widehat{W}_k + \Big(A + \frac{B}{r^2}\Big) ik \widehat{W}_k \\ + \frac{i}{r} \sum_{\ell \in \mathbb{Z}} \Big((k-\ell) \partial_r \widehat{\Phi}_\ell \widehat{W}_{k-\ell} - \ell \widehat{\Phi}_\ell \partial_r \widehat{W}_{k-\ell}\Big) = 0, \quad (t,r,\theta) \in \mathbb{R}^+ \times [1,\infty) \times \mathbb{T} \\ \Big(\partial_r^2 + \frac{1}{r} \partial_r - \frac{k^2}{r^2}\Big) \widehat{\Phi}_k = \widehat{W}_k, \quad \widehat{U}_k^r = -\frac{ik}{r} \widehat{\Phi}_k, \quad \widehat{U}_k^\theta = \partial_r \widehat{\Phi}_k, \\ \widehat{W}_k(t=0) = \widehat{W}_k(0). \end{cases}$$

To get rid of the terms involving $\frac{1}{r}\partial_r$ and A in (1.8), we introduce

(1.9)
$$w_k \stackrel{\text{def}}{=} r^{\frac{1}{2}} e^{ikAt} \widehat{W}_k, \quad \varphi_k \stackrel{\text{def}}{=} r^{\frac{1}{2}} e^{ikAt} \widehat{\Phi}_k, \quad u_k^r \stackrel{\text{def}}{=} r^{\frac{1}{2}} e^{ikAt} \widehat{U}_k^r, \quad u_k^{\theta} \stackrel{\text{def}}{=} r^{\frac{1}{2}} e^{ikAt} \widehat{U}_k^{\theta}.$$

It is easy to observe that

$$||W||_{L^{2}(\Omega)}^{2} \approx \sum_{k \in \mathbb{Z}} ||\widehat{W}_{k}||_{L^{2}((1,\infty);rdr)}^{2} = \sum_{k \in \mathbb{Z}} ||w_{k}||_{L^{2}((1,\infty);dr)}^{2},$$
$$||\partial_{r}W||_{L^{2}(\Omega)}^{2} \approx \sum_{k \in \mathbb{Z}} ||\partial_{r}\widehat{W}_{k}||_{L^{2}((1,\infty);rdr)}^{2} = \sum_{k \in \mathbb{Z}} ||r^{\frac{1}{2}}\partial_{r}(r^{-\frac{1}{2}}w_{k})||_{L^{2}((1,\infty);dr)}^{2}$$

which will be frequently used throughout this paper. Moreover, we deduce from (1.8) that

(1.10a)
$$\partial_t w_0 + \mathcal{T}_0 w_0 = r^{-\frac{1}{2}} \sum_{\ell \in \mathbb{Z} \setminus \{0\}} i\ell \partial_r \left(r^{-1} \varphi_\ell w_{-\ell} \right), \quad (t,r) \in \mathbb{R}^+ \times [1,+\infty),$$

(1.10b)
$$\partial_t w_k + \mathcal{T}_k w_k = \sum_{\ell \in \mathbb{Z} \setminus \{0\}} i \left[r^{-\frac{1}{2}} \ell \partial_r \left(r^{-1} \varphi_\ell w_{k-\ell} \right) - k r^{-1} \partial_r \left(r^{-\frac{1}{2}} \varphi_\ell \right) w_{k-\ell} \right] - i k r^{-\frac{3}{2}} u_0^\theta w_k,$$

(1.10c)
$$\left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2}\right)\varphi_k = w_k,$$

(1.10d)
$$u_{k}^{r} = -i\frac{k}{r}\varphi_{k}, \quad u_{k}^{\theta} = r^{\frac{1}{2}}\partial_{r}(r^{-\frac{1}{2}}\varphi_{k}) = \partial_{r}\varphi_{k} - \frac{1}{2r}\varphi_{k},$$

(1.10e)
$$w_{k}|_{r=1} = \varphi_{k}|_{r=1} = 0, \quad \text{for any } k \in \mathbb{Z},$$

where \mathcal{T}_k is defined as

(1.11)
$$\mathcal{T}_{k}^{\mathrm{def}} = -\nu \Big(\partial_{r}^{2} - \frac{k^{2} - \frac{1}{4}}{r^{2}}\Big) + i\frac{kB}{r^{2}}, \quad D(\mathcal{T}_{k})^{\mathrm{def}} = H^{2}\big((1,\infty);dr\big) \cap H^{1}_{0}\big((1,\infty);dr\big).$$

So far, we have transformed the original problem into exploring the dynamic and long-time behavior of the system (1.10).

1.2. Key ideas. It's well known that, for transport-diffusion equations, the enhanced dissipation scale is closely related to the degree of degeneracy of the velocity field, see for instance [1, 4]. However, as pointed out in [20], due to the infinite degeneracy of TC flow in the exterior disk (i.e., $\partial_r^n(r^{-2})$ vanishes at infinity for any $n \ge 0$), the previous results are no longer applicable to $e^{-\mathcal{T}_k t}$, which motivates us to investigate the mapping properties of $e^{-\mathcal{T}_k t}$ in some space-time weighted spaces in [20]. In particular, we observe from Theorem 1.2 of [20] that due to the degeneracy at infinity, the non-homogeneous equation (1.10b) cannot exhibit exponential decay. Specifically, **Lemma 1.1** (Theorem 1.2, [20]). For any positive continuous functions $a_1(r)$, $a_2(r)$ and strictly decreasing function $\phi(r)$, there exists a sequence of functions, $\{(w_n, f_n)\}$, that solves

$$\partial_t w_n + \mathcal{T}_k w_n = f_n, \quad w_n|_{t=0} = 0, \quad w_n|_{r=1,\infty} = 0,$$

and satisfies for any $n \in \mathbb{N}$

$$\|e^{t\phi(r)}a_1(r)w_n\|_{L^2(\mathbb{R}^+;L^2_r)} + \|e^{t\phi(r)}a_2(r)f_n\|_{L^2(\mathbb{R}^+;L^2_r)} < +\infty.$$

However, there holds

$$\limsup_{n \to +\infty} \frac{\|e^{t\phi(r)}a_1(r)w_n\|_{L^2(\mathbb{R}^+;L^2_r)}}{\|e^{t\phi(r)}a_2(r)f_n\|_{L^2(\mathbb{R}^+;L^2_r)}} = +\infty.$$

Lemma 1.1 indicates that even if employing space-time coupling to mitigate the adverse effects of degeneracy at infinity in the Taylor-Couette flow, one still cannot obtain exponential nonlinear enhanced dissipation decay as in the Couette flow. Consequently, for the long-time behavior near the Taylor-Couette flow in the exterior region, even with space-time coupling, at most polynomial nonlinear decay can be expected.

In fact, Theorem 1.1 of [20] establishes that the solution to the linear homogeneous equation exhibits polynomial space-time coupled decay. Precisely:

Lemma 1.2 (Theorem 1.1, [20]). Let $k \in \mathbb{Z} \setminus \{0\}$, $q \in \mathbb{N}$, $\frac{\nu}{|kB|} \ll (1+q)^{-3}$ and let w be determined by

$$\partial_t w + \mathcal{T}_k w = 0, \quad w|_{t=0} = w(0), \quad w|_{r=1,\infty} = 0.$$

There exists constant C = C(q) > 0 independent of ν, k, B and $w_k(0)$, so that

$$\left\| (1 + \kappa_k t/r^2)^q w \right\|_{L^{\infty}(\mathbb{R}^+;L^2_r)} + \kappa_k^{1/2} \left\| (1 + \kappa_k t/r^2)^q r^{-1} w \right\|_{L^2(\mathbb{R}^+;L^2_r)} \le C \|w(0)\|_{L^2_r}.$$

The reasons for exploring the stability of the TC flow in the exterior disk list as follows:

- Firstly, due to the degeneracy of the Taylor-Couette flow at infinity, we cannot expect the solution to decay exponentially in a space-time decoupled manner. One may check [11, 7] for details.
- (2) Lemma 1.1 tells us that even the coupled space-time exponential decay is not valid.
- (3) Lemma 1.1 and Lemma 1.2 tell us that for the fully nonlinear equation (1.10b), one can at most expect polynomial space-time coupling decay.

At this point, a new issue arises: to close the *a priori* estimates for the nonlinear system (1.6), we need space-time coupling multipliers. The previous resolvent estimate methods [26, 9, 13, 2] are all aimed at space-time decoupled cases (essentially performing a Fourier transform in the time direction). Therefore, here we introduce time-dependent resolvent estimate in Section 4.

In the case of polar coordinates for the exterior disk, there is a critical issue: the weight of the stream function with respect to r differs from w by $1/r^2$. A simple basic fact is that the derivatives of $1/r^2$ are non-vanishing in [1, R] yet they may converge to zero at infinity. Therefore, for technical convenience, we shall choose Λ_k as the long-time space-time decay weight. The basic properties of Λ_k will be provided with a detailed explanation after we present the main theorem.

1.3. Main theorems. The following result is the main result of this paper, which concerns global well-posedness and the asymptotic behavior of the solution to the system (1.10).

Theorem 1.3. Let $0 < \varepsilon < 2$ and \hat{C} be a large constant, we denote $\Lambda_k \stackrel{\text{def}}{=} \log (\hat{C}r^2 + \nu^{\frac{1}{3}}|kB|^{\frac{2}{3}}t)$ to be the enhanced dissipation weight. Then there exist constants $C_{\varepsilon}, C'_{\varepsilon}, \hat{C}_{\varepsilon}$ depending only on ε , such that if $C_{\varepsilon} \geq \hat{C}_{\varepsilon}$ and

(1.12)
$$\mathcal{M}(0) \stackrel{\text{def}}{=} \| rw_0(0) \|_{L^2_r} + \| u_0^{\theta}(0) \|_{L^2_r} + \sum_{k \in \mathbb{Z} \setminus \{0\}} M_k(0) \le C_{\varepsilon} |\nu B|^{\frac{1}{2}}, \quad \text{with}$$

$$M_{k}(0) \stackrel{\text{def}}{=} \| r^{\varepsilon+2} w_{k}(0) \Lambda_{k}(0, \cdot) \|_{L_{r}^{2}} + \| r^{\varepsilon+3} \partial_{r} w_{k}(0) \Lambda_{k}(0, \cdot) \|_{L_{r}^{2}}, \quad \text{for } k \neq 0,$$

the system (1.10) has a unique global solution $(w_0, w_k, \varphi_k, u_k^r, u_k^{\theta})$ which satisfies

$$\mathcal{E}(T) \stackrel{\text{def}}{=} E_0(T) + 100 \sum_{k \in \mathbb{Z} \setminus \{0\}} E_k(T) \le C'_{\varepsilon} \mathcal{M}(0), \quad \text{for any } T > 0 \quad \text{with}$$

(1.13)
$$\begin{split} E_0(T) \stackrel{\text{def}}{=} & \|rw_0\|_{L^{\infty}_T(L^2_r)} + \|u_0^{\theta}\|_{L^{\infty}_T(L^2_r)} + \nu^{\frac{1}{2}} \|w_0\|_{L^2_T(L^2_r)}, \\ & E_k(T) \stackrel{\text{def}}{=} \|r^{\varepsilon+1} w_k \Lambda_k\|_{L^{\infty}_T(L^2_r)} + \mu^{\frac{1}{2}}_k \|r^{\varepsilon} w_k \Lambda_k\|_{L^2_T(L^2_r)} \\ & + \|kB|^{\frac{1}{2}} |k|^{\frac{1}{2}} \Big(\|r^{\varepsilon-1} \partial_r \varphi_k \Lambda_k\|_{L^2_T(L^2_r)} + \|k\| \|r^{\varepsilon-2} \varphi_k \Lambda_k\|_{L^2_T(L^2_r)} \Big), \quad \text{for } k \neq 0. \end{split}$$

It is easy to observe from Lemma B.3 and (1.10d) that

(1.14)
$$|kB|^{\frac{1}{2}}|k| \left\| r^{\varepsilon - \frac{3}{2}} \varphi_k \Lambda_k \right\|_{L^2_T(L^\infty_r)} \lesssim E_k(T)$$

As a corollary of Theorem 1.3, we obtain the pointwise logarithmic-type nonlinear enhanced dissipation estimate.

Theorem 1.4. Under the assumptions of Theorem 1.3, we have

(1.15)
$$||r^{1+\varepsilon}w_k(t)||_{L^2_r} \lesssim_{\varepsilon} \frac{1}{\log\left(\hat{C} + \nu^{\frac{1}{3}}|kB|^{\frac{2}{3}}t\right)} \mathcal{M}(0), \quad \text{for } k \neq 0, t > 0,$$

and

(1.16)
$$\lim_{t \to +\infty} t^{\frac{1}{2}} \|w_0(t)\|_{L^2_r} = 0.$$

We have the following remarks in order concerning Theorem 1.3: (a) The Choice of Λ_k :

In fact, throughout this paper, we only require the following properties for Λ_k :

A.
$$\mu_k^{-1} r^2 \left| \frac{\partial_t \Lambda_k}{\Lambda_k} \right| + r^2 \left| \frac{\partial_r^2 \Lambda_k}{\Lambda_k} \right| + r \left| \frac{\partial_r \Lambda_k}{\Lambda_k} \right| \ll 1$$
, where $\mu_k^{\text{def}} \max\left(\nu k^2, \nu^{\frac{1}{3}} |kB|^{\frac{2}{3}}\right)$;

- B. $\Lambda_k \leq \Lambda_\ell \Lambda_{k-\ell}$ for $k, \ell \in \mathbb{Z}$; C. $\frac{\Lambda_k(t,r)}{\Lambda_k(t,s)} \lesssim \max\left(\left(\frac{r}{s}\right)^{\varepsilon_1}, \left(\frac{s}{r}\right)^{\varepsilon_2}\right)$ for some $\varepsilon_1, \varepsilon_2 \ll 1$.

Rather than using the explicit expression for Λ_k ,

- (1) Property A will be used to absorb terms arising from commutators $[\partial_t, \Lambda_k]$ and $[\partial_r^2, \Lambda_k].$
- (2) Property B is designed to deal with the frequency interactions in the nonlinear terms.
- (3) Property C will be used to recover the elliptic estimate (see Lemma C.1 and the estimation of terms involving K_k).
 - Thus, if one prefers polynomially enhanced dissipation for w, we can

replace
$$\log\left(\hat{C}r^2 + \nu^{\frac{1}{3}}|kB|^{\frac{2}{3}}t\right)$$
 by $\left(e + \frac{\nu^{\frac{1}{3}}|kB|^{\frac{2}{3}}t}{r^2}\right)^o$,

where $\delta \ll_{\varepsilon} 1$. However, for simplicity, this paper considers only the logarithmic-type enhanced dissipation.

• Furthermore, we can replace $\nu^{\frac{1}{3}}|kB|^{\frac{2}{3}}$ by μ_k to provide better dissipation for high frequencies $(|k| \ge \nu^{-\frac{1}{2}}|B|^{\frac{1}{2}})$, and the proof is the same. We retain $\nu^{\frac{1}{3}}|kB|^{\frac{2}{3}}$ in Λ_k to emphasize the dissipation scale $\mathcal{O}(\nu^{-\frac{1}{3}}|B|^{-\frac{2}{3}})$.

(b) The Criticality of Theorem 1.3:

- (1) If we impose additional spatial weights r^{β} ($\beta > 0$) on w_0 , beyond those in $E_0(T)$, similar results as Theorem 1.3 hold only for $t \leq \nu^{-1}$. In this case, we cannot expect global stability results for general data, since Proposition 2.6 indicates that $e^{-\mathcal{T}_0 t}$ is linearly unstable in these weighted spaces. On the other hand, if we remove some spatial weights r^{β} from $E_0(T)$, we cannot close the energy estimate for $E_k(T)$. Therefore, up to this point, $E_0(T)$ is critical.
- (2) In the linear theory of TC flow in the exterior domain (see [20]), the linear evolution $e^{-\mathcal{T}_k t}$ can exhibit polynomial dissipation $(1 + \kappa_k t)^{-\delta}$ when $\frac{\nu}{|kB|} \ll (1 + \delta)^{-3}$, at the cost of losing equivalent spatial weights. For nonlinear evolution, we can only deal with $\delta \ll_{\varepsilon} 1$, because the elliptic estimate

 $\|r^{-2}\Lambda_k\varphi_k\|_{L^2_r} \le C\|\Lambda_k w_k\|_{L^2_r} \quad \text{for some } C \text{ depending only on } \delta,$

fails for $\Lambda_k = \left(e + \frac{\nu^{\frac{1}{3}} |kB|^{\frac{2}{3}} t}{r^2}\right)^{\delta}$ and large $|\delta|$. Therefore, up to this point, the enhanced dissipation weight $\left(e + \frac{\nu^{\frac{1}{3}} |kB|^{\frac{2}{3}} t}{r^2}\right)^{\delta}$ with $\delta \ll_{\varepsilon} 1$ may be optimal for the nonlinear problem.

(c) **Transition threshold:** The mathematic version of Transition threshold was formulated by Bedrossian-Gremain-Masmoudi [3] as follows:

Given a norm $\|\cdot\|_X$, find a $\beta = \beta(X)$ so that

$$\begin{split} \|u_{\rm in}\|_X \leqslant \ \nu^\beta \ \Rightarrow \ {\rm stability}, \\ \|u_{\rm in}\|_X \gg \ \nu^\beta \ \Rightarrow \ {\rm instability}, \end{split}$$

where the exponent $\beta = \beta(X) > 0$ is called the transition threshold. In the Λ_k -integrable space in this paper, Theorem 1.3 shows that we can obtain transition threshold estimates similar to those in bounded domain, as can be seen in [2], despite the infinite degeneracy of r^{-2} in exterior disk.

(d) **Biot-Savart Law:** The corresponding integral kernels are different for $k \neq 0$ and k = 0, specifically as follows:

• For the case of $k \neq 0$ in (1.10c), we consider $u_k^{\theta}, u_k^r, rw_k \in L^2((1,\infty), dr)$. In this case, φ_k can be explicitly expressed as

(1.17)
$$\varphi_k(r) = -\mathcal{K}_k[w_k](r) \stackrel{\text{def}}{=} -\int_1^\infty K_k(r,s)w_k(s)\,ds \quad \text{with} \\ K_k(r,s) \stackrel{\text{def}}{=} \frac{1}{2|k|}(rs)^{\frac{1}{2}} \Big(\min\left(\frac{r}{s},\frac{s}{r}\right)^{|k|} - \frac{1}{r^{|k|}s^{|k|}}\Big).$$

Here we list some basic properties of the integral kernel function $K_k(r, s)$:

(1.18a)
$$\partial_r K_k(r,s) = \frac{1}{2|k|} \left(\left(|k| + \frac{1}{2} \right) \left(\frac{r}{s} \right)^{|k| - \frac{1}{2}} \mathbb{1}_{r \le s} + \left(\frac{1}{2} - |k| \right) \left(\frac{s}{r} \right)^{|k| + \frac{1}{2}} \mathbb{1}_{s \le r} \right)^{|k| - \frac{1}{2}} \mathbb{1}_{s \le r}$$

(1.18b)
$$+ \left(|k| - \frac{1}{2} \right) \frac{1}{r^{|k| + \frac{1}{2}} s^{|k| - \frac{1}{2}}} ,$$
$$K_k(1, s) = K_k(r, 1) = \partial_r K_k(r, 1) = 0,$$

(1.18c)
$$\partial_r \partial_s K_k(r,s) = \delta(r-s) + \widetilde{K}_k(r,s)$$
 in the sense of distribution

 $(1.18d) |k|^{-1}rs|\widetilde{K}_k(r,s)| + s|\partial_s K_k(r,s)| + r|\partial_r K_k(r,s)| + |k||K_k(r,s)| \lesssim (rs)^{\frac{1}{2}}\min(\frac{r}{s},\frac{s}{r})^{|k|},$

where

$$\widetilde{K}_k(r,s) = -\frac{k^2 - \frac{1}{4}}{2|k|} \Big(\frac{r^{|k| - \frac{1}{2}}}{s^{|k| + \frac{1}{2}}} \mathbbm{1}_{r \le s} + \frac{s^{|k| - \frac{1}{2}}}{r^{|k| + \frac{1}{2}}} \mathbbm{1}_{s \le r} \Big) - \frac{(|k| - \frac{1}{2})^2}{2|k|} \frac{1}{r^{|k| + \frac{1}{2}}s^{|k| + \frac{1}{2}}}.$$

• For the case of k = 0, it's easy to observe that $u_0^r = 0$ and it follows from $W = r^{-1}\partial_r (rU^{\theta}) - r^{-1}\partial_{\theta}U^r$ that

(1.19)
$$w_0 = r^{-\frac{1}{2}} \partial_r \left(r^{\frac{1}{2}} u_0^{\theta} \right) = \partial_r u_0^{\theta} + \frac{1}{2r} u_0^{\theta}$$

It is natural to assume that $u_0^{\theta}, rw_0 \in L^2((1,\infty), dr)$. Then, according to Lemma B.3, we obtain

$$\lim_{r \to \infty} |r^{\frac{1}{2}} u_0^{\theta}| = 0.$$

Thus, in view of (1.19), we have

$$u_0^{\theta}(r) = -r^{-\frac{1}{2}} \int_r^{\infty} s^{\frac{1}{2}} w_0(s) ds.$$

Due to the logarithmic singularity of φ_0 at $r = \infty$, we directly use its explicit expression for the velocity field u_0^{θ} instead of recovering it through the stream function.

Finally, we sketch the organization of the paper:

- In Section 2, we present some fundamental energy estimates;
- In Sections 3 and 4, we discuss the resolvent estimates with or without the enhanced dissipation weight Λ_k ;
- In Section 5, we use the resolvent estimates to derive the estimates of $E_k(T)$ for $k \neq 0$;
- In Section 6, we provide the proofs of Theorem 1.3 and Theorem 1.4.

1.4. Notations. Throughout the whole paper, we denote

$$\Omega \stackrel{\text{def}}{=} \left\{ x \in \mathbb{R}^2 : |x| \ge 1 \right\} = \left\{ (r, \theta) : r \in [1, \infty), \theta \in \mathbb{T} \right\}.$$

For $1 \leq p \leq \infty$, we denote

$$L^{p}(\Omega) \stackrel{\text{def}}{=} L^{p}(\Omega; dx) = L^{p}((1, \infty) \times \mathbb{T}; r dr d\theta),$$

 $L_r^{p\stackrel{\text{def}}{=}}L^p\big((1,\infty);dr\big), \quad L_T^p\stackrel{\text{def}}{=}L^p\big((0,T);dt\big), \quad H_r^k\stackrel{\text{def}}{=}H^k\big((1,\infty);dr\big), \quad H_{r,0}^k\stackrel{\text{def}}{=}H_0^k\big((1,\infty);dr\big).$

By $\langle \cdot, \cdot \rangle_{L^2_r}$, we mean the natural inner product in the Hilbert space L^2_r , i.e.,

$$\langle f,g\rangle_{L^2_r} \stackrel{\mathrm{def}}{=} \int_1^\infty f\bar{g}dr,$$

and $\langle \cdot, \cdot \rangle_{L^2(\Omega)}$ is similarly defined.

By w', we mean $\partial_r w$. And $\mathbb{1}_E(r)$ designates the characteristic function on $E \subset [1, \infty)$. We denote $g \leq f$ if there exists a constant C > 0 so that $g \leq Cf$, moreover we denote $g \ll f$ if

(1.20)

this C is universal and small enough, and we use $g \lesssim_{\varepsilon} f$ to emphasize dependence of C on ε . Throughtout this paper, we denote

$$\kappa_k \stackrel{\text{def}}{=} \nu^{\frac{1}{3}} |kB|^{\frac{2}{3}}, \quad \mu_k \stackrel{\text{def}}{=} \max(\nu k^2, \kappa_k), \quad \mathcal{T}_k \stackrel{\text{def}}{=} -\nu \left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2}\right) + i\frac{kB}{r^2}, \quad D \stackrel{\text{def}}{=} H_r^2 \cap H_{r,0}^1.$$

2. Fundamental energy estimates

In this section, we first establish some fundamental energy estimates for the fully nonlinear equations (1.3) and (1.5).

Lemma 2.1. Let W be a smooth enough solution of (1.5) on [0,T]. Then, for any $T \in (0,T)$, one has

(2.1)
$$\|W(t)\|_{L^{2}(\Omega)}^{2} + 2\nu \|\nabla W\|_{L^{2}(L^{2}(\Omega))}^{2} = \|W(0)\|_{L^{2}(\Omega)}^{2}.$$

Proof. Due to div U = 0 and (1.4), we get, by taking $L^2(\Omega)$ inner product of the equation (1.5) with W and using integration by parts, that

$$0 = \frac{1}{2} \frac{d}{dt} \|W(t)\|_{L^{2}(\Omega)}^{2} + \nu \|\nabla W\|_{L^{2}(\Omega)}^{2} + \frac{1}{2} \int_{\partial\Omega} (V^{\star} + U) \cdot \mathbf{n} |W|^{2} dS - \nu \int_{\partial\Omega} \nabla W \cdot \mathbf{n} W dS$$

$$= \frac{1}{2} \frac{d}{dt} \|W(t)\|_{L^{2}(\Omega)}^{2} + \nu \|\nabla W\|_{L^{2}(\Omega)}^{2}.$$

By integrating the above inequality over [0, t], we obtain (2.1).

Lemma 2.2. Under the assumption that $U \in H^2(\Omega)$, div U = 0, $W = \operatorname{curl} U$, and U satisfies the boundary condition (1.4), we have

(2.2)
$$-\int_{\Omega} \Delta U \cdot U dx = \|W\|_{L^2(\Omega)}^2.$$

Proof. Let $\mathbf{n} = (\mathbf{n}^1, \mathbf{n}^2)$ be the outward normal vector to the domain Ω . We get, by using integration by parts, that

$$\begin{split} \|W\|_{L^{2}(\Omega)}^{2} &= \int_{\Omega} (\partial_{1}U^{2} - \partial_{2}U^{1})^{2} dx = \int_{\Omega} (\partial_{1}U^{2})^{2} dx + \int_{\Omega} (\partial_{2}U^{1})^{2} dx - 2 \int_{\Omega} \partial_{1}U^{2} \partial_{2}U^{1} dx \\ &= \int_{\partial\Omega} U^{2} \mathbf{n}^{1} \partial_{1}U^{2} dS - \int_{\Omega} U^{2} \partial_{1}^{2}U^{2} dx + \int_{\partial\Omega} U^{1} \mathbf{n}^{2} \partial_{2}U^{1} dS - \int_{\Omega} U^{1} \partial_{2}^{2}U^{1} dx \\ &- \int_{\partial\Omega} U^{2} \mathbf{n}^{1} \partial_{2}U^{1} dS - \int_{\partial\Omega} U^{1} \mathbf{n}^{2} \partial_{1}U^{2} dS + \int_{\Omega} U^{2} \partial_{2} \partial_{1}U^{1} + U^{1} \partial_{1} \partial_{2}U^{2} dx, \end{split}$$

from which, $\partial_1 U^1 + \partial_2 U^2 = 0$ and $(\partial_1 U^2 - \partial_2 U^1)|_{\partial\Omega} = 0$, we infer

$$||W||_{L^{2}(\Omega)}^{2} = \int_{\partial\Omega} (U^{2}\mathbf{n}^{1} - U^{1}\mathbf{n}^{2})(\partial_{1}U^{2} - \partial_{2}U^{1})dS - \int_{\Omega} \Delta U \cdot Udx$$
$$= -\int_{\Omega} \Delta U \cdot Udx.$$

This leads to (2.2).

Lemma 2.3. Let U be a smooth enough solution of (1.3) on [0,T]. Then for any $t \in (0,T)$, one has

(2.3)
$$\|U(t)\|_{L^{2}(\Omega)}^{2} + 2\nu \|W\|_{L^{2}_{t}(L^{2}(\Omega))}^{2} \\ \leq \|U(0)\|_{L^{2}(\Omega)}^{2} + 8\pi |B| \sum_{k \in \mathbb{Z} \setminus \{0\}} |k| \|\frac{\varphi_{k}}{r^{2}}\|_{L^{2}_{t}(L^{2}_{r})} \left\|\frac{\partial_{r}\varphi_{-k}}{r}\right\|_{L^{2}_{t}(L^{2}_{r})}.$$

Proof. Thanks to (1.4), $V^{\star} \cdot \mathbf{n}|_{\partial\Omega} = 0$ and (2.2), we get, by taking $L^2(\Omega)$ inner product of (1.3) with U, that

(2.4)
$$\frac{1}{2}\frac{d}{dt}\|U\|_{L^{2}(\Omega)}^{2} + \nu\|W\|_{L^{2}(\Omega)}^{2} = -\int_{\Omega} (U \cdot \nabla)V^{\star} \cdot U dx.$$

Yet by virtue of (1.2) and the fact: $\partial_r \mathbf{e}_{\theta} = 0, \partial_{\theta} \mathbf{e}_{\theta} = -\mathbf{e}_{\mathbf{r}}$, we write

$$\begin{split} \int_{\Omega} (U \cdot \nabla) V^{\star} \cdot U dx &= \int_{\Omega} \left(U^r \partial_r + U^{\theta} \frac{\partial_{\theta}}{r} \right) \left(A r \mathbf{e}_{\theta} + \frac{B}{r} \mathbf{e}_{\theta} \right) \cdot \left(U^r \mathbf{e}_{\mathbf{r}} + U^{\theta} \mathbf{e}_{\theta} \right) dx \\ &= \int_{\Omega} \left(\left(A - \frac{B}{r^2} \right) U^r \mathbf{e}_{\theta} - \left(A + \frac{B}{r^2} \right) U^{\theta} \mathbf{e}_{\mathbf{r}} \right) \cdot \left(U^r \mathbf{e}_{\mathbf{r}} + U^{\theta} \mathbf{e}_{\theta} \right) dx \\ &= -2B \int_{\Omega} \frac{U^r U^{\theta}}{r^2} dx, \end{split}$$

from which, (1.7), (1.9), (1.10d) and $\varphi_{-k} = \overline{\varphi_k}$ (since φ is real function), we infer

$$\begin{split} \int_{\Omega} (U \cdot \nabla) V^{\star} \cdot U dx &= -4\pi B \sum_{k \in \mathbb{Z}} \int_{1}^{\infty} \frac{\widehat{U}_{k}^{r} \widehat{U}_{-k}^{\theta}}{r^{2}} r dr = -4\pi B \sum_{k \in \mathbb{Z}} \int_{1}^{\infty} \frac{u_{k}^{r} u_{-k}^{\theta}}{r^{2}} dr \\ &= 4\pi B \sum_{k \in \mathbb{Z} \setminus \{0\}} ik \int_{1}^{\infty} \frac{\varphi_{k} \partial_{r} \varphi_{-k}}{r^{3}} dr - 4\pi B \sum_{k \in \mathbb{Z} \setminus \{0\}} ik \int_{1}^{\infty} \frac{\varphi_{k} \varphi_{-k}}{2r^{4}} dr \\ &= 4\pi B \Re \sum_{k \in \mathbb{Z} \setminus \{0\}} ik \int_{1}^{\infty} \frac{\varphi_{k} \partial_{r} \varphi_{-k}}{r^{3}} dr. \end{split}$$

By inserting the above inequality into (2.4), we find

$$\frac{1}{2}\frac{d}{dt}\|U\|_{L^{2}(\Omega)}^{2} + \nu\|W\|_{L^{2}(\Omega)}^{2} \le 4\pi|B|\sum_{k\in\mathbb{Z}\setminus\{0\}}|k|\|\frac{\varphi_{k}}{r^{2}}\|_{L^{2}_{r}}\left\|\frac{\partial_{r}\varphi_{-k}}{r}\right\|_{L^{2}_{r}}.$$

By integrating the above inequality over [0, t], we achieve (2.3).

Next, we study the mapping properties of the semigroup $e^{-\mathcal{T}_k t}$ in polynomial weighted spaces.

Lemma 2.4. Let $\alpha \in \mathbb{R}$ and $r^{\alpha}w \in D$. Then one has

(2.5a)
$$\Re \langle \mathcal{T}_k w, r^{2\alpha} w \rangle_{L^2_r} \ge \nu (k^2 - \alpha^2) \| r^{\alpha - 1} w \|_{L^2_r}^2.$$

(2.5b)
$$\Re \langle \mathcal{T}_k w, r^{2\alpha} w \rangle_{L^2_r} = \nu \| r^{\alpha} w' \|_{L^2_r}^2 + \nu \left(k^2 - \frac{1}{4} - 2\alpha^2 + \alpha \right) \| r^{\alpha - 1} w \|_{L^2_r}^2$$

Proof. We first get, by using Lemma B.1, that

$$\Re \langle \mathcal{T}_k w, r^{2\alpha} w \rangle_{L^2_r} = \nu \left(\| r^{\alpha} w' \|_{L^2_r}^2 - \alpha (2\alpha - 1) \| r^{\alpha - 1} w \|_{L^2_r}^2 \right) + \nu \left(k^2 - \frac{1}{4} \right) \| r^{\alpha - 1} w \|_{L^2_r}^2,$$

which implies (2.5b).

By applying (2.5b) and Lemma B.2, we obtain

$$\Re \langle \mathcal{T}_k w, r^{2\alpha} w \rangle_{L^2_r} \ge \nu \left((\alpha - 1/2)^2 \| r^{\alpha - 1} w \|_{L^2_r}^2 - \alpha (2\alpha - 1) \| r^{\alpha - 1} w \|_{L^2_r}^2 \right) + \nu \left(k^2 - \frac{1}{4} \right) \| r^{\alpha - 1} w \|_{L^2_r}^2$$
$$= \nu (k^2 - \alpha^2) \| r^{\alpha - 1} w \|_{L^2_r}^2.$$

This completes the proof of (2.5a).

Proposition 2.5. Let w be a smooth enough solution of the following linear equation:

(2.6)
$$\partial_t w - \nu \left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2} \right) w + iS(r)w = 0 \quad \text{with } w|_{r=1,\infty} = 0$$

where the shearing profile $S(r) \in C_b([1,\infty))$ is a real function. Then as long as $|\alpha| \leq |k|$, $||r^{\alpha}w(t)||_{L^2_r}$ is nonincreasing with respect to time $t \geq 0$.

Proof. By taking the real part of the L_r^2 inner product of (2.6) with $r^{2\alpha}w$ and then using Lemma B.2, we deduce that for $|\alpha| \leq |k|$

$$0 = \frac{1}{2} \frac{d}{dt} \|r^{\alpha} w(t)\|_{L_{r}^{2}}^{2} + \Re \Big\langle -\nu \Big(\partial_{r}^{2} - \frac{k^{2} - \frac{1}{4}}{r^{2}}\Big)w + iS(r)w, r^{2\alpha}w\Big\rangle_{L_{r}^{2}}$$

$$\geq \frac{1}{2} \frac{d}{dt} \|r^{\alpha} w(t)\|_{L_{r}^{2}}^{2} + \nu(k^{2} - \alpha^{2}) \|r^{\alpha - 1}w\|_{L_{r}^{2}}^{2},$$

which implies that $||r^{\alpha}w(t)||_{L^2_r}$ is nonincreasing with respect to time $t \ge 0$.

The above proposition provides a basic polynomially weighted energy estimate for the kmode equation with a general shearing profile S(r). However, the following proposition shows that even if S(r) = 0 and $|\alpha| > |k|$, we may not have a uniform bound in time for $||r^{\alpha}w||_{L^{2}_{r}}$.

Proposition 2.6. There exists C > 0, so that for any increasing function $f(r) \ge 0$ and initial data $0 \le w_g(r) \in C_c^{\infty}(2,4)$, if $\nu t \ge 6$, there holds

(2.7)
$$||rf(r)e^{-\mathcal{T}_0 t}w_g||_{L^2_r} \ge Cf(\sqrt{\nu t})||w_g||_{L^1_r}.$$

Proof. We first observe that $w \stackrel{\text{def}}{=} e^{-\mathcal{T}_0 t} w_q$ solves

$$\begin{cases} \partial_t w - \nu \Big(\partial_r^2 + \frac{1}{4r^2} \Big) w = 0, \quad (t,r) \in \mathbb{R}^+ \times [1,+\infty), \\ w(t=0) = w_g, \quad w|_{r=1,\infty} = 0. \end{cases}$$

Then $\eta(x) \stackrel{\text{def}}{=} |x|^{-\frac{1}{2}} w(|x|)$ verifies

$$\begin{cases} \partial_t \eta - \nu \Delta \eta = \partial_t \eta - \nu (\partial_r^2 + \frac{1}{r} \partial_r) \eta = 0, \quad (t, x) \in \mathbb{R}^+ \times \Omega, \\ \eta(0, x) = |x|^{-\frac{1}{2}} w_g(|x|), \quad \eta|_{\partial\Omega} = 0. \end{cases}$$

If we denote the heat kernel of Dirichlet Laplacian on Ω by $p_{\Omega}(t, x, y)$, then one has

(2.8)
$$\eta(t,x) = \int_{|y|\ge 1} p_{\Omega}(\nu t, x, y) |y|^{-\frac{1}{2}} w_g(|y|) \, dy = \int_{2\le |y|\le 4} p_{\Omega}(\nu t, x, y) |y|^{-\frac{1}{2}} w_g(|y|) \, dy.$$

By virtue of Theorem 2.7 below, we get for $|x| \ge 2, \nu t \ge 6$,

$$\eta(t,x) \gtrsim \int_{2 \le |y| \le 4} \frac{1}{\nu t} e^{-c_2 \frac{|x-y|^2}{\nu t}} w_g(|y|) \, dy \gtrsim \frac{1}{\nu t} e^{-c_3 \frac{|x|^2}{\nu t}} \|w_g\|_{L^1_r},$$

which implies

$$\begin{aligned} \left\| rf(r)e^{-\mathcal{T}_{0}t}w_{g} \right\|_{L_{r}^{2}}^{2} &= \int_{|x|\geq 1} |x|^{2}f^{2}(|x|)|\eta(t,x)|^{2} \, dx \geq f^{2}(\sqrt{\nu t}) \int_{|x|\geq \sqrt{\nu t}} |x|^{2}|\eta(t,x)|^{2} \, dx \\ &\gtrsim f^{2}(\sqrt{\nu t}) \int_{|x|\geq \sqrt{\nu t}} \frac{|x|^{2}}{(\nu t)^{2}} e^{-2c_{3}\frac{|x|^{2}}{\nu t}} \, dx \|w_{g}\|_{L_{r}^{1}}^{2} \gtrsim f^{2}(\sqrt{\nu t}) \|w_{g}\|_{L_{r}^{1}}^{2}. \end{aligned}$$

This completes the proof of (2.7).

Theorem 2.7 ([17, 27]). There exist some constants $c_1, c_2 > 0$, so that for $|x|, |y| \ge 2, t \ge 1$, there holds

(2.9)
$$\frac{c_1}{t}e^{-c_2\frac{|x-y|^2}{t}} \le p_{\Omega}(t,x,y) \le \frac{1}{c_1t}e^{-\frac{|x-y|^2}{c_2t}}.$$

However, in the following sections, we will show that, due to the special transport structure of Taylor-Couette flow, i.e., $S(r) = \frac{kB}{r^2}$, a uniform bound in time for $||r^{\alpha}w||_{L^2_r}$ can be obtained for $\nu/|kB| \ll (1+|\alpha|)^{-3}$. This is one of the key ideas of this paper.

3. Resolvent estimates without enhanced dissipation weight

In order to a uniform operator norm estimate for the semigroup $e^{-t\mathcal{T}_k}$, we first investigate the following fundamental resolvent equation:

(3.1)
$$(\mathcal{T}_k - ikB\lambda)w = -\nu \Big(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2}\Big)w + ikB(r^{-2} - \lambda)w = F$$

with $w \in H_r^2 \cap H_{r,0}^1$, $\lambda \in \mathbb{R}$ and $|k| \ge 1$. By density argument, we may always assume without loss of generality that $\varphi, w, F \in \mathcal{C}_c^{\infty}([1,\infty))$ in this section, unless otherwise specified. To proceed, we introduce:

$$(3.2) \quad \|w\|_{H^1_k}^2 \stackrel{\text{def}}{=} \|\partial_r w\|_{L^2_r}^2 + (k^2 - \frac{1}{4}) \left\|\frac{w}{r}\right\|_{L^2_r}^2 \quad \text{and} \quad \|w\|_{H^{-1}_k} \stackrel{\text{def}}{=} \inf_{w = \partial_r g_1 + \frac{k}{r} g_2} \left(\|g_1\|_{L^2_r} + \|g_2\|_{L^2_r}\right).$$

Then for $w \in H_{r,0}^1$, $F \in H_r^{-1}$, $k \in \mathbb{Z} \setminus \{0\}$ and $\alpha \in \mathbb{R}$, we have

(3.3)
$$\|\langle w, F \rangle_{L_r^2} \| \le \|w\|_{H_k^1} \|F\|_{H_k^{-1}}, \\ \|r^{1+\alpha}w(s)\|_{H_k^{-1}} \approx \inf_{w=\partial_r F_1 + \frac{k}{r}F_2} (\|r^{1+\alpha}F_1\|_{L_r^2} + \|r^{1+\alpha}F_2\|_{L_r^2}).$$

Below we fix some decreasing function $\rho \in \mathcal{C}^{\infty}(\mathbb{R})$ satisfying

$$\rho(z) = 1, \text{ for } z \le -1, \qquad \rho(z) = -1, \text{ for } z \ge 1.$$

Proposition 3.1. For any $k \in \mathbb{Z} \setminus \{0\}$, $\alpha \in \mathbb{R}$, $\lambda \in \mathbb{R}$, if we assume that $\frac{\nu}{|kB|} \ll (1+|\alpha|)^{-3}$, there exists constant C > 0 independent of $\nu, k, B, \lambda, \alpha$ so that

(3.4)
$$\nu \| r^{\alpha} w' \|_{L^{2}_{r}}^{2} + \mu_{k} \| r^{\alpha-1} w \|_{L^{2}_{r}}^{2} \leq C \left(\left| \langle F, r^{2\alpha} w \rangle_{L^{2}_{r}} \right| + \left| \langle F, r^{2\alpha} \tilde{\rho} w \rangle_{L^{2}_{r}} \right| \mathbb{1}_{(0,1)}(\lambda) \right),$$

where $\mu_k = \max\left(\nu k^2, \nu^{\frac{1}{3}} |kB|^{\frac{2}{3}}\right)$ and $\tilde{\rho} \stackrel{\text{def}}{=} \rho\left(\frac{r - \lambda^{-\frac{1}{2}}}{\nu^{\frac{1}{3}} |kB|^{-\frac{1}{3}} \lambda^{-\frac{1}{2}}}\right)$ for $\lambda \in (0, 1)$.

Moreover, we have

(3.5a)
$$\nu^{\frac{1}{2}} \mu_k^{\frac{1}{2}} \| r^{\alpha} w' \|_{L^2_r} + \mu_k \| r^{\alpha-1} w \|_{L^2_r} + |kB| \| r^{\alpha+1} (r^{-2} - \lambda) w \|_{L^2_r} \le C \| r^{\alpha+1} F \|_{L^2_r},$$

(3.5b)
$$\nu \|r^{\alpha}w\|_{H^{1}_{k}} + \nu^{\frac{1}{2}}\mu^{\frac{1}{2}}_{k}\|r^{\alpha-1}w\|_{L^{2}_{r}} + |kB|\|r^{\alpha}(r^{-2}-\lambda)w\|_{H^{-1}_{k}} \le C(1+|\alpha/k|)\|r^{\alpha}F\|_{H^{-1}_{k}}.$$

Proof. We first recall from [20] that (3.4) holds for $\alpha = 0$, i.e., for $\nu \ll |kB|$, there holds

(3.6)
$$\nu \|w'\|_{L^2_r}^2 + \mu_k \|w/r\|_{L^2_r}^2 \le C(|\langle F, w \rangle_{L^2_r}| + |\langle F, \tilde{\rho}w \rangle_{L^2_r} |\mathbb{1}_{(0,1)}(\lambda)).$$

For arbitrary $\alpha \in \mathbb{R}$, we observe from (3.1) that $\eta \stackrel{\text{def}}{=} r^{\alpha} w$ and $G \stackrel{\text{def}}{=} r^{\alpha} F$ satisfy

(3.7)
$$-\nu \Big(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2}\Big)\eta + ikB(r^{-2} - \lambda)\eta = G + \nu(\alpha + \alpha^2)r^{-2}\eta - 2\nu\alpha r^{-1}\partial_r\eta.$$

Then we get, by applying (3.6) and setting $\tilde{\alpha} \stackrel{\text{def}}{=} 1 + |\alpha|$, that

$$\begin{split} \nu \|\eta'\|_{L_{r}^{2}}^{2} + \mu_{k} \|\eta/r\|_{L_{r}^{2}}^{2} &\leq C\left(\left|\langle G,\eta\rangle_{L_{r}^{2}}\right| + \left|\langle G,\tilde{\rho}\eta\rangle_{L_{r}^{2}}\right|\mathbb{1}_{(0,1)}(\lambda)\right) \\ &+ C\nu \left|\langle(\alpha+\alpha^{2})r^{-2}\eta - 2\alpha r^{-1}\partial_{r}\eta,\eta\rangle_{L_{r}^{2}}\right| \\ &+ C\nu \left|\langle(\alpha+\alpha^{2})r^{-2}\eta - 2\alpha r^{-1}\partial_{r}\eta,\tilde{\rho}\eta\rangle_{L_{r}^{2}}\right|\mathbb{1}_{(0,1)}(\lambda) \\ &\leq C\left(\left|\langle G,\eta\rangle_{L_{r}^{2}}\right| + \left|\langle G,\tilde{\rho}\eta\rangle_{L_{r}^{2}}\right|\mathbb{1}_{(0,1)}(\lambda)\right) + C\nu\tilde{\alpha}^{2}\|\eta/r\|_{L_{r}^{2}}^{2} + \frac{1}{2}\nu\|\eta'\|_{L_{r}^{2}}^{2}. \end{split}$$

Notice that $\nu \tilde{\alpha}^2 \ll \mu_k$ if $\frac{\nu}{|kB|} \ll \tilde{\alpha}^{-3}$, so that we deduce from the above inequality that

$$\nu \|\eta'\|_{L^2_r}^2 + \mu_k \|\eta/r\|_{L^2_r}^2 \le C(|\langle F, r^{2\alpha}w\rangle_{L^2_r}| + |\langle F, r^{2\alpha}\tilde{\rho}w\rangle_{L^2_r}|\mathbb{1}_{(0,1)}(\lambda)),$$

which together with the fact:

$$\nu \|r^{\alpha}w\|_{H^{1}_{k}}^{2} + \mu_{k}\|r^{\alpha-1}w\|_{L^{2}_{r}}^{2} \approx \nu \|r^{\alpha}w'\|_{L^{2}_{r}}^{2} + \mu_{k}\|r^{\alpha-1}w\|_{L^{2}_{r}}^{2} \approx \nu \|\eta'\|_{L^{2}_{r}}^{2} + \mu_{k}\|\eta/r\|_{L^{2}_{r}}^{2}$$

ensures (3.4).

Next let us turn to the proof of (3.5a) and (3.5b). We first observe from (3.3) that (3.8) $|\langle F, r^{2\alpha}w\rangle_{L^2_r}| \leq \min(||r^{\alpha+1}F||_{L^2_r}||r^{\alpha-1}w||_{L^2_r}, ||r^{\alpha}F||_{H^{-1}_k}||r^{\alpha}w||_{H^1_k}),$

and for $\lambda \in (0, 1)$,

$$\begin{split} \left| \langle F, r^{2\alpha} \tilde{\rho} w \rangle_{L_{r}^{2}} \right| &\leq \min \left(\| r^{\alpha+1} F \|_{L_{r}^{2}} \| r^{\alpha-1} \tilde{\rho} w \|_{L_{r}^{2}}, \ \| r^{\alpha} F \|_{H_{k}^{-1}} \| r^{\alpha} \tilde{\rho} w \|_{H_{k}^{1}} \right) \\ &\lesssim \min \left(\| r^{\alpha+1} F \|_{L_{r}^{2}} \| r^{\alpha-1} w \|_{L_{r}^{2}}, \ \| r^{\alpha} F \|_{H_{k}^{-1}} (\| r^{\alpha} w \|_{H_{k}^{1}} + \| \partial_{r} \tilde{\rho} r^{\alpha} w \|_{L_{r}^{2}}) \right). \end{split}$$

Due to $\frac{\nu}{|kB|} \ll (1+|\alpha|)^{-3}, \nu^{\frac{1}{3}}|kB|^{-\frac{1}{3}}$ is sufficiently small, so that one has

$$\|\partial_r \tilde{\rho} r^{\alpha} w\|_{L^2_r} \lesssim \frac{\lambda^{-\frac{1}{2}}}{\nu^{\frac{1}{3}} |kB|^{-\frac{1}{3}} \lambda^{-\frac{1}{2}}} \|r^{\alpha-1} w\|_{L^2_r} \lesssim \nu^{-\frac{1}{3}} |kB|^{\frac{1}{3}} \|r^{\alpha-1} w\|_{L^2_r},$$

as a result, it comes out

(3.9)
$$|\langle F, r^{2\alpha} \tilde{\rho} w \rangle_{L^2_r}| \lesssim \min \left(\|r^{\alpha+1} F\|_{L^2_r} \|r^{\alpha-1} w\|_{L^2_r}, \\ \|r^{\alpha} F\|_{H^{-1}_k} \left(\|r^{\alpha} w\|_{H^1_k} + \nu^{-\frac{1}{3}} |kB|^{\frac{1}{3}} \|r^{\alpha-1} w\|_{L^2_r} \right) \right).$$

Then we deduce from (3.4), (3.8) and (3.9) that

$$\nu \|r^{\alpha}w'\|_{L^{2}_{r}}^{2} + \mu_{k}\|r^{\alpha-1}w\|_{L^{2}_{r}}^{2} \leq C\|r^{\alpha+1}F\|_{L^{2}_{r}}\|r^{\alpha-1}w\|_{L^{2}_{r}} \leq C\mu_{k}^{-1}\|r^{\alpha+1}F\|_{L^{2}_{r}}^{2} + \frac{\mu_{k}}{2}\|r^{\alpha-1}w\|_{L^{2}_{r}}^{2},$$

and

$$\begin{split} &\nu \|r^{\alpha}w'\|_{L_{r}^{2}}^{2}+\mu_{k}\|r^{\alpha-1}w\|_{L_{r}^{2}}^{2} \\ &\leq C\|r^{\alpha}F\|_{H_{k}^{-1}}\left(\|r^{\alpha}w\|_{H_{k}^{1}}+\nu^{-\frac{1}{3}}|kB|^{\frac{1}{3}}\|r^{\alpha-1}w\|_{L_{r}^{2}}\right) \\ &\leq C\left(\nu^{-1}+\nu^{-\frac{2}{3}}|kB|^{\frac{2}{3}}\mu_{k}^{-1}\right)\|r^{\alpha}F\|_{H_{k}^{-1}}^{2}+\frac{1}{2}\left(\nu\|r^{\alpha}w\|_{H_{k}^{1}}^{2}+\mu_{k}\|r^{\alpha-1}w\|_{L_{r}^{2}}^{2}\right), \end{split}$$

from which and the fact that $|kB|^{\frac{2}{3}}\mu_k^{-1} \leq \nu^{-\frac{1}{3}}$, we infer

(3.10a)
$$\nu^{\frac{1}{2}} \mu_k^{\frac{1}{2}} \| r^{\alpha} w' \|_{L^2_r} + \mu_k \| r^{\alpha-1} w \|_{L^2_r} \le C \| r^{\alpha+1} F \|_{L^2_r},$$

(3.10b)
$$\nu \|r^{\alpha}w\|_{H^{1}_{k}} + \nu^{\frac{1}{2}}\mu^{\frac{1}{2}}_{k}\|r^{\alpha-1}w\|_{L^{2}_{r}} \leq C\|r^{\alpha}F\|_{H^{-1}_{k}}.$$

Thanks to (3.10), it suffices to estimate $(r^{-2} - \lambda)w$ in order to complete the proof of (3.5). Notice that

-1

$$\begin{split} \|kBr^{\alpha}(r^{-2}-\lambda)w\|_{H_{k}^{-1}} &\leq \|r^{\alpha}F\|_{H_{k}^{-1}} + \nu \left\|r^{\alpha}\left(\partial_{r}^{2}-\frac{k^{2}-\frac{1}{4}}{r^{2}}\right)w\right\|_{H_{k}^{-1}} \\ &\lesssim \|r^{\alpha}F\|_{H_{k}^{-1}} + \nu \left(1+|\frac{\alpha}{k}|\right)\|r^{\alpha}\partial_{r}w\|_{L_{r}^{2}} + \nu|k|\|r^{\alpha-1}w\|_{L_{r}^{2}} \\ &\lesssim \left(1+|\frac{\alpha}{k}|+\frac{\nu^{\frac{1}{2}}|k|}{\mu_{k}^{\frac{1}{2}}}\right)\|r^{\alpha}F\|_{H_{k}^{-1}} \lesssim (1+|\alpha/k|)\|r^{\alpha}F\|_{H_{k}^{-1}}, \end{split}$$

from which and (3.10b), we obtain (3.5b).

On the other hand, we get, by using integration by parts, that

$$\begin{split} \langle F, r^{2\alpha+2}(r^{-2}-\lambda)w \rangle_{L^2_r} \\ &= \langle -\nu \left(\partial_r^2 - (k^2 - 1/4)r^{-2}\right)w + ikB \left(r^{-2} - \lambda\right)w, r^{2\alpha+2} \left(r^{-2} - \lambda\right)w \rangle_{L^2_r} \\ &= \nu \langle w', \left(r^{2\alpha+2}(r^{-2}-\lambda)w\right)' \rangle_{L^2_r} + \nu (k^2 - 1/4) \langle w, r^{2\alpha}(r^{-2}-\lambda)w \rangle_{L^2_r} + ikB \|r^{\alpha+1}(r^{-2}-\lambda)w\|_{L^2_r}^2 \\ &= \nu \langle w', r^{2\alpha+2}(r^{-2}-\lambda)w' \rangle_{L^2_r} + \nu \langle w', \left(2\alpha r^{2\alpha-1} - \lambda(2\alpha+2)r^{2\alpha+1}\right)w \rangle_{L^2_r} \\ &+ \nu (k^2 - 1/4) \langle w, r^{2\alpha}(r^{-2}-\lambda)w \rangle_{L^2_r} + ikB \|r^{\alpha+1}(r^{-2}-\lambda)w\|_{L^2_r}^2. \end{split}$$

By taking imaginary part of the above inequality and rewriting

$$2\alpha r^{2\alpha-1} - \lambda(2\alpha+2)r^{2\alpha+1} = 2(1+\alpha)r^{2\alpha+1}(r^{-2}-\lambda) - 2r^{2\alpha-1},$$

we obtain

$$\begin{aligned} &|kB| ||r^{\alpha+1}(r^{-2}-\lambda)w||_{L_{r}^{2}}^{2} \\ &\leq \left|\langle F, r^{2\alpha+2}(r^{-2}-\lambda)w\rangle_{L_{r}^{2}}\right| + 2\nu\tilde{\alpha}\left|\langle w', r^{2\alpha+1}(r^{-2}-\lambda)w\rangle_{L_{r}^{2}}\right| + 2\nu\left|\langle w', r^{2\alpha-1}w\rangle_{L_{r}^{2}}\right| \\ &\leq \left(||r^{\alpha+1}F||_{L_{r}^{2}} + 2\nu\tilde{\alpha}||r^{\alpha}w'||_{L_{r}^{2}}\right)||r^{\alpha+1}(r^{-2}-\lambda)w||_{L_{r}^{2}} + 2\nu||r^{\alpha}w'||_{L_{r}^{2}}||r^{\alpha-1}w||_{L_{r}^{2}}.\end{aligned}$$

Observing from (3.5a) that

$$\nu^{\frac{2}{3}} |kB|^{\frac{1}{3}} ||r^{\alpha}w'||_{L^{2}_{r}} + \nu^{\frac{1}{3}} |kB|^{\frac{2}{3}} ||r^{\alpha-1}w||_{L^{2}_{r}} \le C ||r^{\alpha+1}F||_{L^{2}_{r}}.$$

Then we get, by applying Young's inequality and $\left(\frac{\nu}{|kB|}\right)^{\frac{1}{3}} \ll \tilde{\alpha}^{-1}$, that

$$\begin{aligned} |kB| \|r^{\alpha+1}(r^{-2} - \lambda)w\|_{L^{2}_{r}}^{2} \lesssim &\frac{1}{|kB|} \|r^{\alpha+1}F\|_{L^{2}_{r}}^{2} + \frac{\nu^{2}\tilde{\alpha}^{2}}{|kB|} \|r^{\alpha}w'\|_{L^{2}_{r}}^{2} + \nu\|r^{\alpha}w'\|_{L^{2}_{r}}^{2} \|r^{\alpha-1}w\|_{L^{2}_{r}} \\ \lesssim &\left(\frac{1}{|kB|} + \frac{\nu^{2}\tilde{\alpha}^{2}}{|kB|}\nu^{-\frac{4}{3}}|kB|^{-\frac{2}{3}}\right) \|r^{\alpha+1}F\|_{L^{2}_{r}}^{2} \lesssim \frac{1}{|kB|} \|r^{\alpha+1}F\|_{L^{2}_{r}}^{2}, \end{aligned}$$

which together with (3.10b) ensures (3.5a). This completes the proof of Proposition 3.1.

Below we present the resolvent estimate of φ :

Proposition 3.2. Let $k \in \mathbb{Z} \setminus \{0\}, \varepsilon \in (0, 2), \lambda \in \mathbb{R}$, there exists constant $C_{\varepsilon} > 0$ independent of ν, k, B, λ , so that

(3.11)
$$\mu_k^{\frac{1}{2}} |kB|^{\frac{1}{2}} |k|^{\frac{1}{2}} \left(\left\| \frac{\varphi'}{r^{1-\varepsilon}} \right\|_{L^2_r} + |k| \left\| \frac{\varphi}{r^{2-\varepsilon}} \right\|_{L^2_r} \right) \le C_{\varepsilon} \|r^{2+\varepsilon}F\|_{L^2_r}.$$

Proof. If $\lambda \leq 0$, it's easy to observe from Proposition 3.1 that for $|\alpha| \leq 3$

$$\|r^{\alpha-1}w\|_{L^2_r} \le \|r^{\alpha+1}(r^{-2}-\lambda)w\|_{L^2_r} \le \frac{C}{|kB|} \|r^{\alpha+1}F\|_{L^2_r}$$

which together with Lemma C.1 ensures

$$\begin{split} \left\| \frac{\varphi'}{r^{1-\varepsilon}} \right\|_{L^{2}_{r}} + |k| \left\| \frac{\varphi}{r^{2-\varepsilon}} \right\|_{L^{2}_{r}} \lesssim_{\varepsilon} |k|^{-1} \|r^{\varepsilon}w\|_{L^{2}_{r}} \lesssim_{\varepsilon} |k|^{-1} \min\left(|kB|^{-1}, \mu_{k}^{-1}\right) \|r^{2+\varepsilon}F\|_{L^{2}_{r}} \\ \lesssim_{\varepsilon} \mu_{k}^{-\frac{1}{2}} |kB|^{-\frac{1}{2}} |k|^{-1} \|r^{2+\varepsilon}F\|_{L^{2}_{r}}. \end{split}$$

If $\lambda > 0$ and $\nu k^2 \ge |B|$, we deduce from Lemma C.1 that

$$\begin{split} \left\| \frac{\varphi'}{r^{1-\varepsilon}} \right\|_{L^{2}_{r}} + |k| \left\| \frac{\varphi}{r^{2-\varepsilon}} \right\|_{L^{2}_{r}} \lesssim_{\varepsilon} |k|^{-1} \| r^{\varepsilon} w \|_{L^{2}_{r}} \lesssim_{\varepsilon} |k|^{-1} \mu_{k}^{-1} \| r^{2+\varepsilon} F \|_{L^{2}_{r}} \\ \lesssim_{\varepsilon} \mu_{k}^{-\frac{1}{2}} (\nu k^{2})^{-\frac{1}{2}} |k|^{-1} \| r^{2+\varepsilon} F \|_{L^{2}_{r}} \\ \lesssim_{\varepsilon} \mu_{k}^{-\frac{1}{2}} |kB|^{-\frac{1}{2}} |k|^{-\frac{1}{2}} \| r^{2+\varepsilon} F \|_{L^{2}_{r}}. \end{split}$$

Therefore it remains to consider (3.11) for the case: $\lambda > 0$ and $\nu k^2 \leq |B|$ (i.e., $\mu_k = \nu^{\frac{1}{3}} |kB|^{\frac{2}{3}}$). Let us denote $\tilde{\delta} \stackrel{\text{def}}{=} (\frac{\nu}{|kB|})^{\frac{1}{3}} \ll 1$, and

(3.12)
$$E \stackrel{\text{def}}{=} \{ r \ge 1 : |1 - \sqrt{\lambda}r| \le \tilde{\delta} \},$$
$$E_m \stackrel{\text{def}}{=} \{ r \ge 1 : \tilde{\delta} \le |1 - \sqrt{\lambda}r| \le \frac{1}{2} \},$$
$$E_o \stackrel{\text{def}}{=} \{ r \ge 1 : \frac{1}{2} \le |1 - \sqrt{\lambda}r| \},$$

and split $w = w(\mathbb{1}_E + \mathbb{1}_{E_m} + \mathbb{1}_{E_o})$. Then we get, by applying Lemma C.1 that

$$\left\|\frac{\varphi'}{r^{1-\varepsilon}}\right\|_{L^{2}_{r}} + |k| \left\|\frac{\varphi}{r^{2-\varepsilon}}\right\|_{L^{2}_{r}} \lesssim_{\varepsilon} |k|^{-\frac{1}{2}} \left(\|r^{\varepsilon-\frac{1}{2}}w\mathbb{1}_{E}\|_{L^{1}_{r}} + \|r^{\varepsilon-\frac{1}{2}}w\mathbb{1}_{E_{m}}\|_{L^{1}_{r}}\right) + |k|^{-1} \|r^{\varepsilon}w\mathbb{1}_{E_{o}}\|_{L^{2}_{r}}.$$

It follows from Lemmas B.4 and B.5 that

$$\begin{split} \|r^{\varepsilon-\frac{1}{2}}w\mathbb{1}_{E}\|_{L^{1}_{r}} + \|r^{\varepsilon-\frac{1}{2}}w\mathbb{1}_{E_{m}}\|_{L^{1}_{r}} \lesssim \|r^{-\frac{1}{2}}\mathbb{1}_{E}\|_{L^{2}_{r}}\|r^{\varepsilon}w\|_{L^{2}_{r}} + \left\|r^{-\frac{1}{2}}\frac{\mathbb{1}_{E_{m}}}{1-\lambda r^{2}}\right\|_{L^{2}_{r}}\|r^{\varepsilon}(1-\lambda r^{2})w\|_{L^{2}_{r}} \\ \lesssim \tilde{\delta}^{\frac{1}{2}}\|r^{\varepsilon}w\|_{L^{2}_{r}} + \tilde{\delta}^{-\frac{1}{2}}\|r^{\varepsilon}(1-\lambda r^{2})w\|_{L^{2}_{r}}, \end{split}$$

from which and Proposition 3.1, we infer

$$\begin{split} & \left\|\frac{\varphi'}{r^{1-\varepsilon}}\right\|_{L^{2}_{r}} + |k| \left\|\frac{\varphi}{r^{2-\varepsilon}}\right\|_{L^{2}_{r}} \\ & \lesssim_{\varepsilon} |k|^{-\frac{1}{2}} \Big(\Big(\frac{\nu}{|kB|}\Big)^{\frac{1}{6}} \nu^{-\frac{1}{3}} |kB|^{-\frac{2}{3}} + \Big(\frac{\nu}{|kB|}\Big)^{-\frac{1}{6}} |kB|^{-1} + |k|^{-\frac{1}{2}} |kB|^{-1} \Big) \|r^{2+\varepsilon}F\|_{L^{2}_{r}} \\ & \lesssim_{\varepsilon} |k|^{-\frac{1}{2}} \nu^{-\frac{1}{6}} |kB|^{-\frac{5}{6}} \|r^{2+\varepsilon}F\|_{L^{2}_{r}} = |k|^{-\frac{1}{2}} \mu_{k}^{-\frac{1}{2}} |kB|^{-\frac{1}{2}} \|r^{2+\varepsilon}F\|_{L^{2}_{r}}. \end{split}$$

This completes the proof of Proposition 3.11.

To proceed further, we define sublinear operator $Q_{a,b}$, which will be frequently used in this paper, via

(3.13)
$$\mathcal{Q}_{a,b}[f](r) \stackrel{\text{def}}{=} \int_1^\infty (rs)^{-\frac{1}{2}} \min\left(\left(\frac{r}{s}\right)^a, \left(\frac{s}{r}\right)^b\right) |f|(s) ds.$$

Lemma 3.3. Let $\mathcal{Q}_{a,b}[f]$ be defined by (3.13). Then we have

(3.14a)
$$\|\mathcal{Q}_{a,b}[f]\|_{L^2_r} \le \left(\frac{1}{a} + \frac{1}{b}\right) \|f\|_{L^2_r} \quad \text{if } a, b > 0,$$

(3.14b)
$$||r^{\frac{1}{2}}\mathcal{Q}_{a,b}[f]||_{L^2_r} \le \left(\frac{1}{2a+1} + \frac{1}{2b-1}\right)^{\frac{1}{2}} ||f||_{L^1_r} \quad \text{if } a > -\frac{1}{2} \quad \text{and} \quad b > \frac{1}{2}.$$

Proof. In view of (3.13), we get, by using changes of variables: $\tilde{r} = \log r$, $\tilde{s} = \log s$, that

$$e^{\frac{\tilde{r}}{2}}\mathcal{Q}_{a,b}[f](e^{\tilde{r}}) = \int_{0}^{+\infty} \min\left(e^{a(\tilde{r}-\tilde{s})}, e^{b(\tilde{s}-\tilde{r})}\right) e^{\frac{\tilde{s}}{2}} |f|(e^{\tilde{s}}) d\tilde{s},$$
$$e^{\tilde{r}}\mathcal{Q}_{a,b}[f](e^{\tilde{r}}) = \int_{0}^{+\infty} \min\left(e^{(a+\frac{1}{2})(\tilde{r}-\tilde{s})}, e^{(b-\frac{1}{2})(\tilde{s}-\tilde{r})}\right) e^{\tilde{s}} |f|(e^{\tilde{s}}) d\tilde{s}$$

Applying Young Inequality gives rise to

$$\|\mathcal{Q}_{a,b}[f]\|_{L^{2}_{r}} = \|e^{\frac{\tilde{r}}{2}}\mathcal{Q}_{a,b}[f](e^{\tilde{r}})\|_{L^{2}(\mathbb{R}^{+};d\tilde{r})} \le \left(\frac{1}{a} + \frac{1}{b}\right)\|e^{\frac{\tilde{s}}{2}}|f|(e^{\tilde{s}})\|_{L^{2}(\mathbb{R}^{+};d\tilde{s})} \le \left(\frac{1}{a} + \frac{1}{b}\right)\|f\|_{L^{2}_{r}},$$
and

and

$$\begin{aligned} \|r^{\frac{1}{2}}\mathcal{Q}_{a,b}[f]\|_{L^{2}_{r}} &= \|e^{\tilde{r}}\mathcal{Q}_{a,b}[f](e^{\tilde{r}})\|_{L^{2}(\mathbb{R}^{+};d\tilde{r})} \leq \left(\frac{1}{2a+1} + \frac{1}{2b-1}\right)^{\frac{1}{2}} \|e^{\tilde{s}}|f|(e^{\tilde{s}})\|_{L^{1}(\mathbb{R}^{+};d\tilde{s})} \\ &\leq \left(\frac{1}{2a+1} + \frac{1}{2b-1}\right)^{\frac{1}{2}} \|f\|_{L^{1}_{r}}. \end{aligned}$$

This completes the proof of (3.14).

Proposition 3.4. Let $k \in \mathbb{Z} \setminus \{0\}, \varepsilon \in (0, 2), \lambda \in \mathbb{R}$, there exists constant $C_{\varepsilon} > 0$ independent of ν, k, B, λ , so that

(3.15)
$$\nu^{\frac{1}{2}} |kB|^{\frac{1}{2}} |k|^{\frac{1}{2}} \left(\left\| \frac{\varphi'}{r^{1-\varepsilon}} \right\|_{L^{2}_{r}} + |k| \left\| \frac{\varphi}{r^{2-\varepsilon}} \right\|_{L^{2}_{r}} \right) \le C_{\varepsilon} \|r^{1+\varepsilon}F\|_{H^{-1}_{k}}.$$

Proof. In the case of $\lambda \leq 0$, it's easy to observe from (3.1) that for $|\alpha| \leq 3$,

$$|kB||\langle (r^{-2} - \lambda)w, r^{2\alpha}w\rangle_{L^2_r}| \le |\Im\langle F, r^{2\alpha}w\rangle_{L^2_r}| + \nu|\Im\langle\partial_r w, \partial_r(r^{2\alpha})w\rangle_{L^2_r}|,$$

which implies in this case that

$$|kB| \|r^{\alpha-1}w\|_{L^{2}_{r}}^{2} \lesssim |\langle F, r^{2\alpha}w\rangle_{L^{2}_{r}}| + \nu \|r^{\alpha}\partial_{r}w\|_{L^{2}_{r}} \|r^{\alpha-1}w\|_{L^{2}_{r}}.$$

Applying Young's inequality and Proposition 3.1 gives

$$\begin{split} \|kB\| \|r^{\alpha-1}w\|_{L^2_r}^2 &\lesssim \|r^{\alpha}F\|_{H^{-1}_k} \|r^{\alpha}w\|_{H^1_k} + \nu \|r^{\alpha}\partial_r w\|_{L^2_r} \|r^{\alpha-1}w\|_{L^2_r} \\ &\lesssim (\nu^{-1} + \nu^{-\frac{1}{2}}\mu_k^{-\frac{1}{2}}) \|r^{\alpha}F\|_{H^{-1}_k}^2 \lesssim \nu^{-1} \|r^{\alpha}F\|_{H^{-1}_k}^2, \end{split}$$

from which and Lemma C.1, we infer

(3.16)
$$\left\|\frac{\varphi'}{r^{1-\varepsilon}}\right\|_{L^{2}_{r}} + |k| \left\|\frac{\varphi}{r^{2-\varepsilon}}\right\|_{L^{2}_{r}} \lesssim_{\varepsilon} |k|^{-1} \|r^{\varepsilon}w\|_{L^{2}_{r}} \lesssim_{\varepsilon} |k|^{-1} \nu^{-\frac{1}{2}} |kB|^{-\frac{1}{2}} \|r^{1+\varepsilon}F\|_{H^{-1}_{k}}.$$

In the case when $\lambda > 0$ and $\nu k^2 \ge |B|$, we deduce from Proposition 3.1 and Lemma C.1 that

(3.17)
$$\begin{aligned} \left\|\frac{\varphi'}{r^{1-\varepsilon}}\right\|_{L^{2}_{r}} + |k| \left\|\frac{\varphi}{r^{2-\varepsilon}}\right\|_{L^{2}_{r}} \lesssim_{\varepsilon} |k|^{-1} \|r^{\varepsilon}w\|_{L^{2}_{r}} \\ \lesssim_{\varepsilon} |k|^{-1} \nu^{-\frac{2}{3}} |kB|^{-\frac{1}{3}} \|r^{1+\varepsilon}F\|_{H^{-1}_{k}} \\ \lesssim_{\varepsilon} \nu^{-\frac{1}{2}} |kB|^{-\frac{1}{2}} |k|^{-\frac{1}{2}} \|r^{1+\varepsilon}F\|_{H^{-1}_{k}}.\end{aligned}$$

We claim that if $\lambda > 0$ and $\nu k^2 \leq |B|$, then

(3.18a)
$$\nu^{\frac{1}{2}} |kB|^{\frac{1}{2}} |k|^{\frac{3}{2}} \left\| \frac{\varphi}{r^{2-\varepsilon}} \right\|_{L^{2}_{r}} \lesssim_{\varepsilon} \|r^{1+\varepsilon}F\|_{H^{-1}_{k}},$$

(3.18b)
$$\nu^{\frac{1}{2}} |kB|^{\frac{1}{2}} |k|^{\frac{1}{2}} \left\| \frac{\varphi'}{r^{1-\varepsilon}} \right\|_{L^{2}_{r}} \leq C_{\varepsilon} \|r^{1+\varepsilon}F\|_{H^{-1}_{k}}$$

By summarizing (3.16), (3.17) and (3.18), we complete the proof of (3.15).

It remains to prove (3.18).

Proof of (3.18). In the case when $\lambda > 0$ and $\nu k^2 \leq |B|$, we denote $r_0 \stackrel{\text{def}}{=} \lambda^{-\frac{1}{2}}$, $\tilde{\delta} \stackrel{\text{def}}{=} (\frac{\nu}{|kB|})^{\frac{1}{3}} \ll 1$ and $\delta \stackrel{\text{def}}{=} \tilde{\delta}r_0$. Let E, E_m, E_o be defined as in (3.12). Since $\nu k^2 \leq |B|$, we have $|k|\tilde{\delta} \leq 1$. Let $\eta(r) \in \mathcal{C}_0^{\infty}(\mathbb{R})$ be supported in (-2, 2) and equal to 1 for $s \in [-1, 1]$, and we denote $\eta(\frac{r-r_0}{\delta})$ by η_{δ} . Then we write

(3.19)

$$-\frac{\varphi}{r^{2-\varepsilon}} = \int_{1}^{\infty} r^{-2+\varepsilon} K_{k}(r,s)w(s)ds$$

$$= \int_{1}^{\infty} r^{-2+\varepsilon} K_{k}(r,s)\frac{1-\eta_{\delta}(s)}{s^{-2}-\lambda}(s^{-2}-\lambda)w(s)ds$$

$$+ \int_{1}^{\infty} r^{-2+\varepsilon} K_{k}(r,s)w(s)\eta_{\delta}(s)ds \stackrel{\text{def}}{=} J_{1} + J_{2},$$

and

(3.20)

$$-\frac{\varphi'}{r^{1-\varepsilon}} = \int_{1}^{\infty} r^{-1+\varepsilon} \partial_{r} K_{k}(r,s) w(s) ds$$

$$= \int_{1}^{\infty} r^{-1+\varepsilon} \partial_{r} K_{k}(r,s) \frac{1-\eta_{\delta}(s)}{s^{-2}-\lambda} (s^{-2}-\lambda) w ds$$

$$+ \int_{1}^{\infty} r^{-1+\varepsilon} \partial_{r} K_{k}(r,s) w(s) \eta_{\delta}(s) ds \stackrel{\text{def}}{=} J_{3} + J_{4}.$$

By density argument, we only consider the case when $(s^{-2} - \lambda)w(s) = \partial_s F_1(s) + \frac{k}{s}F_2(s)$ with $F_1, F_2 \in \mathcal{C}^{\infty}_c([1,\infty))$. Then by virtue of (3.3), we have

(3.21)
$$\|r^{1+\varepsilon}(s^{-2}-\lambda)w(s)\|_{H_k^{-1}} \approx \inf_{F_1,F_2} \left(\|r^{1+\varepsilon}F_1\|_{L_r^2} + \|r^{1+\varepsilon}F_2\|_{L_r^2}\right).$$

Below we divide the proof into the following two steps: **Step 1.** The proof of (3.18a).

Due to K(r, 1) = 0, we get, by using integration by parts, that

$$J_{1} = -\int_{1}^{\infty} r^{-2+\varepsilon} \partial_{s} \Big(K_{k}(r,s) \frac{1-\eta_{\delta}(s)}{s^{-2}-\lambda} \Big) F_{1}(s) ds + \int_{1}^{\infty} r^{-2+\varepsilon} K_{k}(r,s) \frac{1-\eta_{\delta}(s)}{s^{-2}-\lambda} \frac{k}{s} F_{2}(s) ds$$

=: $J_{11} + J_{12} + J_{13} + J_{14}$,

where

$$J_{11} \stackrel{\text{def}}{=} -\int_{1}^{\infty} r^{-2+\varepsilon} \partial_s K_k(r,s) \frac{1-\eta_{\delta}(s)}{s^{-2}-\lambda} F_1(s) ds,$$

$$J_{12} \stackrel{\text{def}}{=} \delta^{-1} \int_{1}^{\infty} r^{-2+\varepsilon} K_k(r,s) \frac{\eta_{\delta}'(s)}{s^{-2}-\lambda} F_1(s) ds,$$

$$J_{13} \stackrel{\text{def}}{=} -2 \int_{1}^{\infty} r^{-2+\varepsilon} K_k(r,s) \frac{1-\eta_{\delta}(s)}{s^3(s^{-2}-\lambda)^2} F_1(s) ds,$$

$$J_{14} \stackrel{\text{def}}{=} \int_1^\infty r^{-2+\varepsilon} K_k(r,s) \frac{k(1-\eta_\delta(s))}{s(s^{-2}-\lambda)} F_2(s) ds.$$

We first get, by applying (1.18d) and $\frac{\mathbb{1}_{E_o}}{|1-\lambda r^2|} \lesssim 1$, that

$$\begin{split} |J_{11}| \lesssim \int_{1}^{\infty} r^{-2+\varepsilon} \left(\frac{r}{s}\right)^{\frac{1}{2}} \min\left(\frac{r}{s}, \frac{s}{r}\right)^{|k|} s^{2} \frac{\mathbb{1}_{E_{m}} + \mathbb{1}_{E_{o}}}{|1 - \lambda s^{2}|} |F_{1}(s)| ds \\ \leq r^{\frac{1}{2}} \int_{1}^{\infty} (rs)^{-\frac{1}{2}} \left(\frac{s}{r}\right)^{\frac{3}{2}-\varepsilon} \min\left(\frac{r}{s}, \frac{s}{r}\right)^{|k|} \frac{\mathbb{1}_{E_{m}}}{|1 - \lambda s^{2}|} s^{\frac{1}{2}+\varepsilon} |F_{1}(s)| ds \\ + \int_{1}^{\infty} (rs)^{-\frac{1}{2}} \left(\frac{s}{r}\right)^{1-\varepsilon} \min\left(\frac{r}{s}, \frac{s}{r}\right)^{|k|} \frac{\mathbb{1}_{E_{o}}}{|1 - \lambda s^{2}|} s^{1+\varepsilon} |F_{1}(s)| ds \\ \lesssim r^{\frac{1}{2}} \mathcal{Q}_{|k| - \frac{3}{2} + \varepsilon, |k| + \frac{3}{2} - \varepsilon} \left[r^{\frac{1}{2}+\varepsilon} \frac{\mathbb{1}_{E_{m}}}{1 - \lambda r^{2}} F_{1} \right] + \mathcal{Q}_{|k| - 1+\varepsilon, |k| + 1-\varepsilon} [r^{1+\varepsilon} F_{1}], \end{split}$$

from which, Lemmas 3.3 and B.5 and $|k| \leq \tilde{\delta}^{-1}$, we infer

$$\begin{split} \|J_{11}\|_{L^{2}_{r}} \lesssim_{\varepsilon} |k|^{-\frac{1}{2}} \left\| r^{-\frac{1}{2}} \frac{\mathbb{1}_{E_{m}}}{1-\lambda r^{2}} \right\|_{L^{2}_{r}} \|r^{1+\varepsilon}F_{1}\|_{L^{2}_{r}} + |k|^{-1} \|r^{1+\varepsilon}F_{1}\|_{L^{2}_{r}} \\ \lesssim_{\varepsilon} |k|^{-\frac{1}{2}} \tilde{\delta}^{-\frac{1}{2}} \|r^{1+\varepsilon}F_{1}\|_{L^{2}_{r}} \lesssim_{\varepsilon} |k|^{-\frac{3}{2}} \tilde{\delta}^{-\frac{3}{2}} \|r^{1+\varepsilon}F_{1}\|_{L^{2}_{r}}. \end{split}$$

Similarly, we get, by using Lemma C.1, that

$$\begin{split} \sum_{i=2}^{4} \|J_{1i}\|_{L_{r}^{2}} \lesssim \delta^{-1} \left\| r^{-2+\varepsilon} \mathcal{K}_{k} \left[\frac{\eta_{\delta}'}{s^{-2} - \lambda} F_{1} \right] \right\|_{L_{r}^{2}} + \left\| r^{-2+\varepsilon} \mathcal{K}_{k} \left[\frac{1 - \eta_{\delta}}{s^{3}(s^{-2} - \lambda)^{2}} F_{1} \right] \right\|_{L_{r}^{2}} \\ &+ \left\| r^{-2+\varepsilon} \mathcal{K}_{k} \left[\frac{k(1 - \eta_{\delta})}{s(s^{-2} - \lambda)} F_{2} \right] \right\|_{L_{r}^{2}} \\ \lesssim \delta^{-1} |k|^{-\frac{3}{2}} \left\| r^{-\frac{1}{2}+\varepsilon} \frac{\eta_{\delta}'}{r^{-2} - \lambda} F_{1} \right\|_{L_{r}^{1}} \\ &+ |k|^{-\frac{3}{2}} \left\| r^{-\frac{1}{2}+\varepsilon} \frac{\mathbb{1}_{E_{m}}}{r^{3}(r^{-2} - \lambda)^{2}} F_{1} \right\|_{L_{r}^{1}} + |k|^{-2} \left\| r^{\varepsilon} \frac{\mathbb{1}_{E_{o}}}{r^{3}(r^{-2} - \lambda)^{2}} F_{1} \right\|_{L_{r}^{2}} \\ &+ |k|^{-\frac{3}{2}} \left\| r^{-\frac{1}{2}+\varepsilon} \frac{k \mathbb{1}_{E_{m}}}{r(r^{-2} - \lambda)} F_{2} \right\|_{L_{r}^{1}} + |k|^{-2} \left\| r^{\varepsilon} \frac{k \mathbb{1}_{E_{o}}}{r(r^{-2} - \lambda)} F_{2} \right\|_{L_{r}^{2}}, \end{split}$$

which together with Lemma B.5 and $|k| \leq \tilde{\delta}^{-1}$ ensures that

$$\begin{split} \sum_{i=2}^{4} \|J_{1i}\|_{L_{r}^{2}} \lesssim_{\varepsilon} \delta^{-1} |k|^{-\frac{3}{2}} \left\| r^{\frac{1}{2}} \frac{\mathbb{1}_{\delta \leq |r-r_{0}| \leq 2\delta}}{1 - \lambda r^{2}} \right\|_{L_{r}^{2}} \|r^{1+\varepsilon} F_{1}\|_{L_{r}^{2}} \\ &+ |k|^{-\frac{3}{2}} \left\| r^{-\frac{1}{2}} \frac{\mathbb{1}_{E_{m}}}{(1 - \lambda r^{2})^{2}} \right\|_{L_{r}^{2}} \|r^{1+\varepsilon} F_{1}\|_{L_{r}^{2}} + |k|^{-2} \left\| r^{1+\varepsilon} \frac{\mathbb{1}_{E_{o}}}{(1 - \lambda r^{2})^{2}} F_{1} \right\|_{L_{r}^{2}} \\ &+ |k|^{-\frac{1}{2}} \left\| r^{-\frac{1}{2}} \frac{\mathbb{1}_{E_{m}}}{1 - \lambda r^{2}} \right\|_{L_{r}^{2}} \|r^{1+\varepsilon} F_{2}\|_{L_{r}^{2}} + |k|^{-1} \left\| r^{1+\varepsilon} \frac{\mathbb{1}_{E_{o}}}{1 - \lambda r^{2}} F_{2} \right\|_{L_{r}^{2}} \\ &\lesssim_{\varepsilon} \left(\delta^{-1} |k|^{-\frac{3}{2}} r_{0} \tilde{\delta}^{-\frac{1}{2}} + |k|^{-\frac{3}{2}} \tilde{\delta}^{-\frac{3}{2}} + |k|^{-2} \right) \|r^{1+\varepsilon} F_{1}\|_{L_{r}^{2}} \\ &+ \left(|k|^{-\frac{1}{2}} \tilde{\delta}^{-\frac{1}{2}} + |k|^{-1} \right) \|r^{1+\varepsilon} F_{2}\|_{L_{r}^{2}} \\ &\lesssim_{\varepsilon} |k|^{-\frac{3}{2}} \tilde{\delta}^{-\frac{3}{2}} \left(\|r^{1+\varepsilon} F_{1}\|_{L_{r}^{2}} + \|r^{1+\varepsilon} F_{2}\|_{L_{r}^{2}} \right). \end{split}$$

By summarizing the above estimates, we obtain

$$\|J_1\|_{L^2_r} \lesssim_{\varepsilon} \nu^{-\frac{1}{2}} |kB|^{\frac{1}{2}} |k|^{-\frac{3}{2}} (\|r^{1+\varepsilon}F_1\|_{L^2_r} + \|r^{1+\varepsilon}F_2\|_{L^2_r}),$$

from which, (3.21) and Proposition 3.1, we infer

4 A . F

$$(3.22) \|J_1\|_{L^2_r} \lesssim_{\varepsilon} \nu^{-\frac{1}{2}} |kB|^{\frac{1}{2}} |k|^{-\frac{3}{2}} \|r^{1+\varepsilon}(r^{-2}-\lambda)w\|_{H^{-1}_k} \lesssim_{\varepsilon} \nu^{-\frac{1}{2}} |kB|^{-\frac{1}{2}} |k|^{-\frac{3}{2}} \|r^{1+\varepsilon}F\|_{H^{-1}_k}.$$

While we get, by using Proposition 3.1 and Lemmas C.1 and B.4, that

(3.23)
$$\begin{aligned} \|J_2\|_{L^2_r} &= \left\|\frac{\mathcal{K}_k[w\eta_{\delta}]}{r^{2-\varepsilon}}\right\|_{L^2_r} \lesssim_{\varepsilon} |k|^{-\frac{3}{2}} \|r^{\varepsilon-\frac{1}{2}}w\eta_{\delta}\|_{L^1_r} \\ &\lesssim_{\varepsilon} |k|^{-\frac{3}{2}} \|r^{-\frac{1}{2}}\eta_{\delta}\|_{L^2_r} \|r^{\varepsilon}w\|_{L^2_r} \lesssim_{\varepsilon} |k|^{-\frac{3}{2}} \tilde{\delta}^{\frac{1}{2}}\nu^{-\frac{2}{3}} |kB|^{-\frac{1}{3}} \|r^{1+\varepsilon}F\|_{H^{-1}_k} \\ &\lesssim_{\varepsilon} \nu^{-\frac{1}{2}} |kB|^{-\frac{1}{2}} |k|^{-\frac{3}{2}} \|r^{1+\varepsilon}F\|_{H^{-1}_k}. \end{aligned}$$

In view of (3.19), by summing up (3.22) and (3.23), we achieve (3.18a). **Step 2.** The proof of (3.18b).

Similar to the estimation of J_2 , we get, by using Proposition 3.1 and Lemmas C.1 and B.4, that

(3.24)
$$\|J_{4}\|_{L_{r}^{2}} = \left\|\frac{\mathcal{K}_{k}^{\prime}[w\eta_{\delta}]}{r^{1-\varepsilon}}\right\|_{L_{r}^{2}} \lesssim_{\varepsilon} |k|^{-\frac{1}{2}} \|r^{\varepsilon-\frac{1}{2}}w\eta_{\delta}\|_{L_{r}^{1}} \\ \lesssim_{\varepsilon} |k|^{-\frac{1}{2}} \|r^{-\frac{1}{2}}\eta_{\delta}\|_{L_{r}^{2}} \|r^{\varepsilon}w\|_{L_{r}^{2}} \lesssim_{\varepsilon} |k|^{-\frac{1}{2}} \tilde{\delta}^{\frac{1}{2}}\nu^{-\frac{2}{3}} |kB|^{-\frac{1}{3}} \|r^{1+\varepsilon}F\|_{H_{k}^{-1}} \\ \lesssim_{\varepsilon} \nu^{-\frac{1}{2}} |kB|^{-\frac{1}{2}} |k|^{-\frac{1}{2}} \|r^{1+\varepsilon}F\|_{H_{k}^{-1}}.$$

To handle J_3 given by (3.20), we first get, by using integration by parts and (1.18a), that

$$J_3(r) = \int_1^\infty r^{-1+\varepsilon} \partial_r K_k(r,s) \frac{1-\eta_\delta(s)}{s^{-2}-\lambda} \partial_s F_1(s) ds$$
$$+ \int_1^\infty r^{-1+\varepsilon} \partial_r K_k(r,s) \frac{k(1-\eta_\delta(s))}{s(s^{-2}-\lambda)} F_2(s) ds$$
$$\stackrel{\text{def}}{=} J_{31} + J_{32} + J_{33} + J_{34} + J_{35},$$

where we define

$$\begin{split} J_{31} &\stackrel{\text{def}}{=} -\int_{1}^{\infty} r^{-1+\varepsilon} \widetilde{K}_{k}(r,s) \frac{1-\eta_{\delta}(s)}{s^{-2}-\lambda} F_{1}(s) ds, \\ J_{32} &\stackrel{\text{def}}{=} \delta^{-1} \int_{1}^{\infty} r^{-1+\varepsilon} \partial_{r} K_{k}(r,s) \frac{\eta_{\delta}'(s)}{s^{-2}-\lambda} F_{1}(s) ds, \\ J_{33} &\stackrel{\text{def}}{=} -2 \int_{1}^{\infty} r^{-1+\varepsilon} \partial_{r} K_{k}(r,s) \frac{1-\eta_{\delta}(s)}{s^{3}(s^{-2}-\lambda)^{2}} F_{1}(s) ds, \\ J_{34} &\stackrel{\text{def}}{=} \int_{1}^{\infty} r^{-1+\varepsilon} \partial_{r} K_{k}(r,s) \frac{1-\eta_{\delta}(s)}{s^{-2}-\lambda} \frac{k}{s} F_{2}(s) ds, \\ J_{35} &\stackrel{\text{def}}{=} -r^{1+\varepsilon} \frac{1-\eta_{\delta}(r)}{1-\lambda r^{2}} F_{1}(r). \end{split}$$

By virtue of (1.18d), we get, by applying Lemma 3.3, that

$$\begin{split} \|J_{31}\|_{L^{2}_{r}} &\lesssim |k| \left\| r^{\frac{1}{2}} \mathcal{Q}_{|k| - \frac{3}{2} + \varepsilon, |k| + \frac{3}{2} - \varepsilon} \left[r^{\frac{1}{2} + \varepsilon} \frac{\mathbb{1}_{E_{m}}}{1 - \lambda r^{2}} F_{1} \right] \right\|_{L^{2}_{r}} \\ &+ |k| \left\| \mathcal{Q}_{|k| - 1 + \varepsilon, |k| + 1 - \varepsilon} \left[r^{1 + \varepsilon} \frac{\mathbb{1}_{E_{o}}}{1 - \lambda r^{2}} F_{1} \right] \right\|_{L^{2}_{r}} \\ &\lesssim_{\varepsilon} |k|^{\frac{1}{2}} \left\| r^{-\frac{1}{2}} \frac{\mathbb{1}_{E_{m}}}{1 - \lambda r^{2}} \right\|_{L^{2}_{r}} \|r^{1 + \varepsilon} F_{1}\|_{L^{2}_{r}} + \left\| r^{1 + \varepsilon} \frac{\mathbb{1}_{E_{o}}}{1 - \lambda r^{2}} F_{1} \right\|_{L^{2}_{r}}, \end{split}$$

from which, $|k| \leq \tilde{\delta}^{-1}$ and Lemma B.5, we infer

$$\|J_{31}\|_{L^{2}_{r}} \lesssim_{\varepsilon} |k|^{-\frac{1}{2}} \tilde{\delta}^{-\frac{3}{2}} \|r^{1+\varepsilon}F_{1}\|_{L^{2}_{r}} \lesssim_{\varepsilon} \nu^{-\frac{1}{2}} |kB|^{-\frac{1}{2}} |k|^{-\frac{1}{2}} \|r^{1+\varepsilon}F_{1}\|_{L^{2}_{r}}.$$

Whereas we get, by a similar derivation of the estimate of $\sum_{i=2}^{4} \|J_{1i}\|_{L_r^2}$, that

$$\begin{split} \sum_{i=2}^{4} \|J_{3i}\|_{L_{r}^{2}} \lesssim \delta^{-1} \left\| r^{-1+\varepsilon} \mathcal{K}_{k}' \Big[\frac{\eta_{\delta}'}{s^{-2} - \lambda} F_{1} \Big] \right\|_{L_{r}^{2}} + \left\| r^{-1+\varepsilon} \mathcal{K}_{k}' \Big[\frac{1 - \eta_{\delta}}{s^{3}(s^{-2} - \lambda)^{2}} F_{1} \Big] \right\|_{L_{r}^{2}} \\ &+ \left\| r^{-1+\varepsilon} \mathcal{K}_{k}' \Big[\frac{k(1 - \eta_{\delta})}{s(s^{-2} - \lambda)} F_{2} \Big] \right\|_{L_{r}^{2}} \\ \lesssim \delta^{-1} |k|^{-\frac{1}{2}} \left\| r^{-\frac{1}{2}+\varepsilon} \frac{\eta_{\delta}'}{r^{-2} - \lambda} F_{1} \right\|_{L_{r}^{1}} \\ &+ |k|^{-\frac{1}{2}} \left\| r^{-\frac{1}{2}+\varepsilon} \frac{1}{r^{3}(r^{-2} - \lambda)^{2}} F_{1} \right\|_{L_{r}^{1}} + |k|^{-1} \left\| r^{\varepsilon} \frac{1 E_{o}}{r^{3}(r^{-2} - \lambda)^{2}} F_{1} \right\|_{L_{r}^{2}} \\ &+ |k|^{-\frac{1}{2}} \left\| r^{-\frac{1}{2}+\varepsilon} \frac{k \mathbbm_{E_{m}}}{r(r^{-2} - \lambda)^{2}} F_{2} \right\|_{L_{r}^{1}} + |k|^{-1} \left\| r^{\varepsilon} \frac{k \mathbbm_{E_{o}}}{r(r^{-2} - \lambda)} F_{2} \right\|_{L_{r}^{2}} \\ &\lesssim_{\varepsilon} \delta^{-1} |k|^{-\frac{1}{2}} \left\| r^{\frac{1}{2}} \frac{1 \frac{\delta \leq |r-r_{0}| \leq 2\delta}{1 - \lambda r^{2}} \right\|_{L_{r}^{2}} \|r^{1+\varepsilon} F_{1}\|_{L_{r}^{2}} \\ &+ |k|^{-\frac{1}{2}} \left\| r^{-\frac{1}{2}} \frac{1 \frac{\delta \leq |r-r_{0}| \leq 2\delta}{1 - \lambda r^{2}} \right\|_{L_{r}^{2}} \|r^{1+\varepsilon} F_{1}\|_{L_{r}^{2}} \\ &+ |k|^{-\frac{1}{2}} \left\| r^{-\frac{1}{2}} \frac{1 E_{m}}{(1 - \lambda r^{2})^{2}} \right\|_{L_{r}^{2}} \|r^{1+\varepsilon} F_{1}\|_{L_{r}^{2}} + |k|^{-1} \left\| \frac{1 E_{o}}{(1 - \lambda r^{2})^{2}} r^{1+\varepsilon} F_{1} \right\|_{L_{r}^{2}} \\ &+ |k|^{\frac{1}{2}} \left\| r^{-\frac{1}{2}} \frac{1 E_{m}}{1 - \lambda r^{2}} \right\|_{L_{r}^{2}} \|r^{1+\varepsilon} F_{2}\|_{L_{r}^{2}} + \|\frac{1 E_{o}}{1 - \lambda r^{2}} r^{1+\varepsilon} F_{2} \right\|_{L_{r}^{2}} \\ &\leq_{\varepsilon} \left(\delta^{-1} |k|^{-\frac{1}{2}} r_{0} \delta^{-\frac{1}{2}} + |k|^{-\frac{1}{2}} \delta^{-\frac{3}{2}} + |k|^{-1} \right) \|r^{1+\varepsilon} F_{1}\|_{L_{r}^{2}} + (|k|^{\frac{1}{2}} \delta^{-\frac{1}{2}} + 1) \|r^{1+\varepsilon} F_{2}\|_{L_{r}^{2}} \\ &\lesssim_{\varepsilon} |k|^{-\frac{1}{2}} \delta^{-\frac{3}{2}} \left(\|r^{1+\varepsilon} F_{1}\|_{L_{r}^{2}} + \|r^{1+\varepsilon} F_{2} \|_{L_{r}^{2}} \right). \end{split}$$

Finally for J_{35} , we have

$$\|J_{35}\|_{L^2_r} \lesssim \left\|\frac{1-\eta_{\delta}}{1-\lambda r^2}\right\|_{L^{\infty}_r} \|r^{1+\varepsilon}F_1\|_{L^2_r} \lesssim \tilde{\delta}^{-1}\|r^{1+\varepsilon}F_1\|_{L^2_r} \lesssim |k|^{-\frac{1}{2}}\tilde{\delta}^{-\frac{3}{2}}\|r^{1+\varepsilon}F_1\|_{L^2_r}.$$

In view of (3.20), by summarizing the above estimates, and then applying (3.21) and Proposition 3.1, we achieve (3.18b).

4. Resolvent estimates with enhanced dissipation weight

In this section, we establish pointwise enhanced dissipation for the following linearized equations with $k \neq 0$:

(4.1a)
$$\partial_t w_k + \mathcal{T}_k w_k = F_k, \quad w_k(t=0) = w_k(0), \quad w_k|_{r=1,\infty} = 0,$$

(4.1b)
$$\left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2}\right)\varphi_k = w_k, \quad \varphi_k|_{r=1,\infty} = 0.$$

Since we shall deal with L^2 type estimates, for simplicity, we always set $w_k(0) = 0$ and assume that $\varphi_k, w_k, F \in \mathcal{C}^{\infty}_c([0,\infty) \times [1,\infty))$. Recall that $\kappa_k \stackrel{\text{def}}{=} \nu^{\frac{1}{3}} |kB|^{\frac{2}{3}}$ and $\Lambda_k(t,r) \stackrel{\text{def}}{=} \log (\hat{C}r^2 + \kappa_k t)$, we denote

(4.2)
$$w_{\star} \stackrel{\text{def}}{=} \Lambda_k(t, r) w_k, \quad F_{\star} \stackrel{\text{def}}{=} \Lambda_k(t, r) F_k, \quad \varphi_{\star} \stackrel{\text{def}}{=} \Lambda_k(t, r) \varphi_k.$$

Then in view of (4.1), we find

$$(4.3) \quad \begin{cases} \partial_t w_\star - \nu \left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2}\right) w_\star + \frac{ikB}{r^2} w_\star \\ = F_\star + \frac{\partial_t \Lambda_k}{\Lambda_k} w_\star - \nu \frac{\partial_r^2 \Lambda_k}{\Lambda_k} w_\star + 2\nu \left(\frac{\partial_r \Lambda_k}{\Lambda_k}\right)^2 w_\star - 2\nu \frac{\partial_r \Lambda_k}{\Lambda_k} \partial_r w_\star \stackrel{\text{def}}{=} F_\star + R^{w_\star}, \\ \left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2}\right) \varphi_\star = w_\star + \frac{\partial_r^2 \Lambda_k}{\Lambda_k} \varphi_\star - 2\left(\frac{\partial_r \Lambda_k}{\Lambda_k}\right)^2 \varphi_\star + 2\frac{\partial_r \Lambda_k}{\Lambda_k} \partial_r \varphi_\star \stackrel{\text{def}}{=} w_\star + R^{\varphi_\star}, \\ w_\star(0) = 0, \quad w_\star|_{r=1,+\infty} = \varphi_\star|_{r=1,+\infty} = 0. \end{cases}$$

The key idea to handle the equations (4.3) is to view $R^{w_{\star}}$ and $R^{\varphi_{\star}}$ as small perturbations and using the resolvent estimates we obtained in Section 3.

Proposition 4.1. For any $k \in \mathbb{Z} \setminus \{0\}$, $\alpha \in [0, 4]$, T > 0, there exist constants C > 0 and \hat{C} sufficiently large, which are independent of ν, k, B, α, T , so that

(4.4)
$$\nu^{\frac{1}{2}} \mu_k^{\frac{1}{2}} \| r^{\alpha} \partial_r w_\star \|_{L^2_T(L^2_r)} + \mu_k \| r^{\alpha - 1} w_\star \|_{L^2_T(L^2_r)} \le C \| r^{\alpha + 1} F_\star \|_{L^2_T(L^2_r)}.$$

Proof. Without loss of generality, we set $T = \infty$ and define

$$\hat{f}(\lambda) \stackrel{\text{def}}{=} \int_0^\infty e^{-it\lambda} f(t) dt.$$

We may extend f(t) = 0 for $t \le 0$ in case f(0) = 0.

Then we get, by taking Fourier transformation to the w_{\star} equation in (4.3) with respect to the time variable, that

(4.5)
$$-\nu \left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2}\right) \widehat{w}_{\star}(kB\lambda, r) + ikB(r^{-2} - \lambda)\widehat{w}_{\star}(kB\lambda, r) \\ = \widehat{f}_{\star}(kB\lambda, r) + \widehat{R^{w_{\star}}}(kB\lambda, r),$$

from which and Proposition 3.1, we infer

$$\nu \mu_{k} \| r^{\alpha} \partial_{r} \widehat{w}_{\star}(kB\lambda, r) \|_{L^{2}_{r}}^{2} + \mu_{k}^{2} \| r^{\alpha-1} \widehat{w}_{\star}(kB\lambda, r) \|_{L^{2}_{r}}^{2} \\
\leq C \Big(\| r^{\alpha+1} \widehat{F}_{\star}(kB\lambda, r) \|_{L^{2}_{r}}^{2} + \| r^{\alpha+1} \widehat{R^{w_{\star}}}(kB\lambda, r) \|_{L^{2}_{r}}^{2} \Big).$$

By integrating the above inequality over \mathbb{R} with respect to λ variable, and then using Plancherel's equality, we obtain

$$(4.6) \quad \nu\mu_k \|r^{\alpha} \partial_r w_\star\|_{L^2_T(L^2_r)}^2 + \mu_k^2 \|r^{\alpha-1} w_\star\|_{L^2_T(L^2_r)}^2 \le C \Big(\|r^{\alpha+1} F_\star\|_{L^2_T(L^2_r)}^2 + \|r^{\alpha+1} R^{w_\star}\|_{L^2_T(L^2_r)}^2 \Big).$$

Yet it follows from Lemma C.2 that

$$\|r^{\alpha+1}R^{w_{\star}}\|_{L^{2}_{T}(L^{2}_{r})}^{2}$$

$$(4.7) \qquad \leq C\Big(\frac{\nu^{\frac{1}{3}}|kB|^{\frac{2}{3}}}{\hat{C}\log\hat{C}}\|r^{\alpha-1}w_{\star}\|_{L^{2}_{T}(L^{2}_{r})} + \frac{\nu}{\log\hat{C}}\|r^{\alpha}\partial_{r}w_{\star}\|_{L^{2}_{T}(L^{2}_{r})} + \frac{\nu}{\log\hat{C}}\|r^{\alpha-1}w_{\star}\|_{L^{2}_{T}(L^{2}_{r})}\Big)^{2}$$

$$\leq \frac{C}{\log^{2}\hat{C}}\Big(\nu^{2}\|r^{\alpha}\partial_{r}w_{\star}\|_{L^{2}_{T}(L^{2}_{r})}^{2} + (\nu^{\frac{1}{3}}|kB|^{\frac{2}{3}})^{2}\|r^{\alpha-1}w_{\star}\|_{L^{2}_{T}(L^{2}_{r})}^{2}\Big).$$

By inserting (4.7) into (4.6), and taking \hat{C} to be large enough, we arrive at (4.4).

Proposition 4.2. For any $k \in \mathbb{Z} \setminus \{0\}$, $\alpha \in [0, 4]$, T > 0, there exist constants C > 0 and \hat{C} sufficiently large, which are independent of ν, k, B, α, T , so that

(4.8)
$$\nu \| r^{\alpha} w_{\star} \|_{L^{2}_{T}(H^{1}_{k})} + \nu^{\frac{1}{2}} \mu^{\frac{1}{2}}_{k} \| r^{\alpha-1} w_{\star} \|_{L^{2}_{T}(L^{2}_{r})} \leq C \| r^{\alpha} F_{\star} \|_{L^{2}_{T}(H^{-1}_{k})}.$$

Proof. Thanks to (3.8-3.9) and Proposition 3.1, we deduce from (4.5) that

$$\begin{split} \nu \| r^{\alpha} \widehat{w}_{\star}(kB\lambda,r) \|_{L^{2}_{\lambda}(H^{1}_{k})}^{2} + \mu_{k} \| r^{\alpha-1} \widehat{w}_{\star}(kB\lambda,r) \|_{L^{2}_{\lambda}(L^{2}_{r})}^{2} \\ &\leq C \int_{\mathbb{R}} \Big| \left\langle r^{\alpha} \widehat{F}_{\star}, r^{\alpha} \widehat{w}_{\star} \right\rangle_{L^{2}_{r}} \Big| (kB\lambda) d\lambda + C \int_{\mathbb{R}} \Big| \left\langle r^{\alpha} \widehat{F}_{\star}, r^{\alpha} \tilde{\rho} \widehat{w}_{\star} \right\rangle_{L^{2}_{r}} \Big| (kB\lambda) \mathbb{1}_{(0,1)} (\lambda) d\lambda \\ &+ C \int_{\mathbb{R}} \Big| \left\langle r^{\alpha} \widehat{R^{w_{\star}}}, r^{\alpha} \widehat{w}_{\star} \right\rangle_{L^{2}_{r}} \Big| (kB\lambda) d\lambda + C \int_{\mathbb{R}} \Big| \left\langle r^{\alpha} \widehat{R^{w_{\star}}}, r^{\alpha} \tilde{\rho} \widehat{w}_{\star} \right\rangle_{L^{2}_{r}} \Big| (kB\lambda) \mathbb{1}_{(0,1)} (\lambda) d\lambda \\ &\leq C \nu^{-1} \| r^{\alpha} \widehat{F}_{\star} (kB\lambda,r) \|_{L^{2}_{\lambda}(H^{-1}_{k})} \\ &+ \frac{1}{4} \Big(\nu \| r^{\alpha} \widehat{w}_{\star} (kB\lambda,r) \|_{L^{2}_{\lambda}(H^{1}_{k})}^{2} + \nu^{\frac{1}{3}} |kB|^{\frac{2}{3}} \| r^{\alpha-1} \widehat{w}_{\star} (kB\lambda,r) \|_{L^{2}_{\lambda}(L^{2}_{r})}^{2} \Big) \\ &+ C \| r^{\alpha+1} \widehat{R^{w_{\star}}} (kB\lambda,r) \|_{L^{2}_{\lambda}(L^{2}_{r})}^{2} \| r^{\alpha-1} \widehat{w}_{\star} (kB\lambda,r) \|_{L^{2}_{\lambda}(L^{2}_{r})}^{2}, \end{split}$$

from which and and Plancherel's equality, we infer

(4.9)
$$\frac{3}{4} \left(\nu \| r^{\alpha} w_{\star} \|_{L^{2}_{T}(H^{1}_{k})}^{2} + \mu_{k} \| r^{\alpha-1} w_{\star} \|_{L^{2}_{T}(L^{2}_{r})}^{2} \right) \leq C \nu^{-1} \| r^{\alpha} F_{\star} \|_{L^{2}_{T}(H^{-1}_{k})}^{2} + C \| r^{\alpha+1} R^{w_{\star}} \|_{L^{2}_{T}(L^{2}_{r})}^{2} \| r^{\alpha-1} w_{\star} \|_{L^{2}_{T}(L^{2}_{r})}.$$

Whereas we get, by using (4.7) and taking \hat{C} to be large enough, that

$$C \|r^{\alpha+1}R^{w_{\star}}\|_{L^{2}_{T}(L^{2}_{r})} \|r^{\alpha-1}w_{\star}\|_{L^{2}_{T}(L^{2}_{r})}$$

$$\leq \frac{C}{\log \hat{C}} \|r^{\alpha-1}w_{\star}\|_{L^{2}_{T}(L^{2}_{r})} (\mu_{k}\|r^{\alpha-1}w_{\star}\|_{L^{2}_{T}(L^{2}_{r})} + \nu\|r^{\alpha}\partial_{r}w_{\star}\|_{L^{2}_{T}(L^{2}_{r})})$$

$$\leq \frac{1}{4} (\nu\|r^{\alpha}w_{\star}\|^{2}_{L^{2}_{T}(H^{1}_{k})} + \mu_{k}\|r^{\alpha-1}w_{\star}\|^{2}_{L^{2}_{T}(L^{2}_{r})}).$$

By inserting the above inequality into (4.9), we achieve (4.8).

Proposition 4.3. For any
$$k \in \mathbb{Z} \setminus \{0\}$$
, $\varepsilon \in (0, 2)$, $T > 0$, there exist constants $C_{\varepsilon} > 0$ and $\hat{C}_0 \gg 1$ depending only on ε , so that for $\hat{C} \ge \hat{C}_0$,

(4.10a)
$$\mu_k^{\frac{1}{2}} |kB|^{\frac{1}{2}} |k|^{\frac{1}{2}} \left(\left\| \frac{\partial_r \varphi_{\star}}{r^{1-\varepsilon}} \right\|_{L^2_T(L^2_r)} + |k| \left\| \frac{\varphi_{\star}}{r^{2-\varepsilon}} \right\|_{L^2_T(L^2_r)} \right) \le C_{\varepsilon} \|r^{2+\varepsilon} F_{\star}\|_{L^2_T(L^2_r)},$$

(4.10b)
$$\nu^{\frac{1}{2}} |kB|^{\frac{1}{2}} |k|^{\frac{1}{2}} \left(\left\| \frac{\partial_r \varphi_{\star}}{r^{1-\varepsilon}} \right\|_{L^2_T(L^2_r)} + |k| \left\| \frac{\varphi_{\star}}{r^{2-\varepsilon}} \right\|_{L^2_T(L^2_r)} \right) \le C_{\varepsilon} \|r^{1+\varepsilon} F_{\star}\|_{L^2_T(H^{-1}_k)}.$$

Proof. By virtue of (1.17) and φ_{\star} equation of (4.3), we write

$$-\varphi_{\star} = \mathcal{K}_k[w_{\star}] + \mathcal{K}_k[R^{\varphi_{\star}}],$$

which implies

$$(4.11) \qquad \begin{aligned} \left\| \frac{\partial_r \varphi_{\star}}{r^{1-\varepsilon}} \right\|_{L^2_T(L^2_r)} + |k| \left\| \frac{\varphi_{\star}}{r^{2-\varepsilon}} \right\|_{L^2_T(L^2_r)} &\leq \left\| \frac{\partial_r \mathcal{K}_k[w_{\star}]}{r^{1-\varepsilon}} \right\|_{L^2_T(L^2_r)} + |k| \left\| \frac{\mathcal{K}_k[w_{\star}]}{r^{2-\varepsilon}} \right\|_{L^2_T(L^2_r)} \\ &+ \left\| \frac{\partial_r \mathcal{K}_k[R^{\varphi_{\star}}]}{r^{1-\varepsilon}} \right\|_{L^2_T(L^2_r)} + |k| \left\| \frac{\mathcal{K}_k[R^{\varphi_{\star}}]}{r^{2-\varepsilon}} \right\|_{L^2_T(L^2_r)} \end{aligned}$$

Below for simplicity, we skip the standard transformation between $L^2_{\lambda}(L^2_r)$ and $L^2_T(L^2_r)$. We first, get, by using Proposition 3.2, (4.7) and Proposition 4.1, that

$$\begin{split} & \left\| \frac{\partial_r \mathcal{K}_k[w_\star]}{r^{1-\varepsilon}} \right\|_{L_T^2(L_r^2)} + |k| \left\| \frac{\mathcal{K}_k[w_\star]}{r^{2-\varepsilon}} \right\|_{L_T^2(L_r^2)} \\ \lesssim_{\varepsilon} \frac{1}{\mu_k^{\frac{1}{2}} |kB|^{\frac{1}{2}} |k|^{\frac{1}{2}}} \left(\|r^{2+\varepsilon} F_\star\|_{L_T^2(L_r^2)} + \|r^{2+\varepsilon} R^{w_\star}\|_{L_T^2(L_r^2)} \right) \\ \lesssim_{\varepsilon} \frac{1}{\mu_k^{\frac{1}{2}} |kB|^{\frac{1}{2}} |k|^{\frac{1}{2}}} \left(\|r^{2+\varepsilon} F_\star\|_{L_T^2(L_r^2)} + \nu^{\frac{1}{3}} |kB|^{\frac{2}{3}} \|r^{\varepsilon} w_\star\|_{L_T^2(L_r^2)} + \nu \|r^{1+\varepsilon} \partial_r w_\star\|_{L_T^2(L_r^2)} \right) \\ \lesssim_{\varepsilon} \frac{1}{\mu_k^{\frac{1}{2}} |kB|^{\frac{1}{2}} |k|^{\frac{1}{2}}} \|r^{2+\varepsilon} F_\star\|_{L_T^2(L_r^2)}. \end{split}$$

While we get, by applying Propositions 3.2, 3.4 and 4.2, that

$$\begin{split} & \left\| \frac{\partial_r \mathcal{K}_k[w_\star]}{r^{1-\varepsilon}} \right\|_{L^2_T(L^2_r)} + |k| \left\| \frac{\mathcal{K}_k[w_\star]}{r^{2-\varepsilon}} \right\|_{L^2_T(L^2_r)} \\ \lesssim_{\varepsilon} \frac{1}{\nu^{\frac{1}{2}} |kB|^{\frac{1}{2}} |k|^{\frac{1}{2}}} \|r^{1+\varepsilon} F_\star\|_{L^2_T(H^{-1}_k)} + \frac{1}{\mu^{\frac{1}{2}}_k |kB|^{\frac{1}{2}} |k|^{\frac{1}{2}}} \|r^{2+\varepsilon} R^{w_\star}\|_{L^2_T(L^2_r)} \\ \lesssim_{\varepsilon} \frac{1}{\nu^{\frac{1}{2}} |kB|^{\frac{1}{2}} |k|^{\frac{1}{2}}} \|r^{1+\varepsilon} F_\star\|_{L^2_T(H^{-1}_k)} \\ & + \frac{1}{\nu^{\frac{1}{6}} |kB|^{\frac{5}{6}} |k|^{\frac{1}{2}}} \left(\nu^{\frac{1}{3}} |kB|^{\frac{2}{3}} \|r^{\varepsilon} w_\star\|_{L^2_T(L^2_r)} + \nu \|r^{1+\varepsilon} \partial_r w_\star\|_{L^2_T(L^2_r)}\right) \\ \lesssim_{\varepsilon} \frac{1}{\nu^{\frac{1}{2}} |kB|^{\frac{1}{2}} |k|^{\frac{1}{2}}} \|r^{1+\varepsilon} F_\star\|_{L^2_T(H^{-1}_k)}. \end{split}$$

And it follows from Lemmas C.1 and C.2 that

$$\begin{split} \left\| \frac{\partial_r \mathcal{K}_k[R^{\varphi_\star}]}{r^{1-\varepsilon}} \right\|_{L^2_T(L^2_r)} + |k| \left\| \frac{\mathcal{K}_k[R^{\varphi_\star}]}{r^{2-\varepsilon}} \right\|_{L^2_T(L^2_r)} \lesssim_{\varepsilon} |k|^{-1} \|r^{\varepsilon} R^{\varphi_\star}\|_{L^2_T(L^2_r)} \\ \lesssim_{\varepsilon} \frac{|k|^{-1}}{\log \hat{C}} \Big(\left\| \frac{\partial_r \varphi_\star}{r^{1-\varepsilon}} \right\|_{L^2_T(L^2_r)} + |k| \left\| \frac{\varphi_\star}{r^{2-\varepsilon}} \right\|_{L^2_T(L^2_r)} \Big). \end{split}$$

By substituting the above inequalities into (4.11) and then taking \hat{C} to be sufficiently large, we achieve (4.10).

5. Space-time estimates of the vorticity in nonzero modes

In this section, we primarily investigate the enhanced dissipation and integrated inviscid damping estimates concerning (4.1). Below we split the solution of (4.1) $w_k = w_{k,H} + w_{k,I}$ and $\varphi_k = \varphi_{k,H} + \varphi_{k,I}$, where $(w_{k,H}, \varphi_{k,H})$ and $(w_{k,I}, \varphi_{k,I})$ satisfy respectively the following two equations:

(5.1)
$$\begin{cases} \partial_t w_{k,H} - \nu \left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2}\right) w_{k,H} + \frac{ikB}{r^2} w_{k,H} = 0, \quad (t,r) \in \mathbb{R}^+ \times [1, +\infty), \\ w_{k,H}(t=0) = w_k(0), \quad w_{k,H}|_{r=1,\infty} = 0, \\ \left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2}\right) \varphi_{k,H} = w_{k,H}, \quad \varphi_{k,H}|_{r=1,\infty} = 0, \end{cases}$$

and

(5.2)
$$\begin{cases} \partial_t w_{k,I} - \nu \left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2} \right) w_{k,I} + \frac{ikB}{r^2} w_{k,I} = f_1 + f_2, \quad (t,r) \in \mathbb{R}^+ \times [1, +\infty), \\ w_{k,I}(t=0) = 0, \quad w_{k,I}|_{r=1,\infty} = 0, \\ \left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2} \right) \varphi_{k,I} = w_{k,I}, \quad \varphi_{k,I}|_{r=1,\infty} = 0, \end{cases}$$

where we write $F_k = f_1 + f_2$.

The main result of Section 5 states as follows:

Proposition 5.1. Let $k \in \mathbb{Z} \setminus \{0\}$, $\varepsilon \in (0, 2)$ let w_k be the solution to (4.1) and the energy functionals $M_k(0)$ and $E_k(T)$ are given respectively by (1.12) and (1.13). Then there exist constants $C_{\varepsilon} > 0$, $\hat{C}_0 \gg 1$ depending only on ε , so that for $\hat{C} \geq \hat{C}_0$ there holds

(5.3)
$$E_k(T) \leq C_{\varepsilon} \Big(M_k(0) + \mu_k^{-\frac{1}{2}} \| r^{2+\varepsilon} f_1 \Lambda_k \|_{L^2_T(L^2_r)} + \nu^{-\frac{1}{2}} \| r^{1+\varepsilon} f_2 \Lambda_k \|_{L^2_T(H^{-1}_k)} \Big).$$

Proof. Proposition 5.1 follows Propositions 5.2 and 5.3 below.

5.1. Space-time estimates of the inhomogeneous equation (5.2).

Proposition 5.2. Let $k \in \mathbb{Z} \setminus \{0\}$, $\varepsilon \in (0, 2)$ and let $(w_{k,I}, \varphi_{k,I})$ solve (5.2). Then there exist constants $C_{\varepsilon} > 0$, $\hat{C}_0 \gg 1$ depending only on ε , so that for $\hat{C} \ge \hat{C}_0$,

(5.4)
$$\|r^{1+\varepsilon}w_{k,I}\Lambda_{k}\|_{L_{T}^{\infty}(L_{r}^{2})} + \nu^{\frac{1}{2}}\|r^{1+\varepsilon}w_{k,I}'\Lambda_{k}\|_{L_{T}^{2}(L_{r}^{2})} + \mu^{\frac{1}{2}}_{k}\|r^{\varepsilon}w_{k,I}\Lambda_{k}\|_{L_{T}^{2}(L_{r}^{2})} + |kB|^{\frac{1}{2}}|k|^{\frac{1}{2}}\left(\left\|\frac{\varphi_{k,I}'}{r^{1-\varepsilon}}\Lambda_{k}\right\|_{L_{T}^{2}(L_{r}^{2})} + |k|\left\|\frac{\varphi_{k,I}}{r^{2-\varepsilon}}\Lambda_{k}\right\|_{L_{T}^{2}(L_{r}^{2})}\right) \leq C_{\varepsilon}\left(\mu^{-\frac{1}{2}}_{k}\|r^{2+\varepsilon}f_{1}\Lambda_{k}\|_{L_{T}^{2}(L_{r}^{2})} + \nu^{-\frac{1}{2}}\|r^{1+\varepsilon}f_{2}\Lambda_{k}\|_{L_{T}^{2}(H_{k}^{-1})}\right).$$

Proof. We first get, by applying Propositions 4.1-4.3 and Lemma C.2, that

(5.5)

$$\begin{aligned}
\nu^{\frac{1}{2}} \| r^{1+\varepsilon} w_{k,I}' \Lambda_{k} \|_{L^{2}_{T}(L^{2}_{r})} + \mu^{\frac{1}{2}}_{k} \| r^{\varepsilon} w_{k,I} \Lambda_{k} \|_{L^{2}_{T}(L^{2}_{r})} \\
+ \| kB \|^{\frac{1}{2}} \| k \|^{\frac{1}{2}} \left(\left\| \frac{\varphi_{k,I}'}{r^{1-\varepsilon}} \Lambda_{k} \right\|_{L^{2}_{T}(L^{2}_{r})} + \| k \| \left\| \frac{\varphi_{k,I}}{r^{2-\varepsilon}} \Lambda_{k} \right\|_{L^{2}_{T}(L^{2}_{r})} \right) \\
\lesssim_{\varepsilon} \mu^{-\frac{1}{2}}_{k} \| r^{2+\varepsilon} f_{1} \Lambda_{k} \|_{L^{2}_{T}(L^{2}_{r})} + \nu^{-\frac{1}{2}} \| r^{1+\varepsilon} f_{2} \Lambda_{k} \|_{L^{2}_{T}(H^{-1}_{k})}.
\end{aligned}$$

Whereas we get, by taking the real part of the L_r^2 inner product of (5.2) with $r^{2+2\varepsilon}\Lambda_k^2 w_{k,I}$, and then using integration by parts and Lemma C.2, that

$$\frac{d}{dt} \|r^{1+\varepsilon} w_{k,I} \Lambda_k\|_{L_r^2}^2 \lesssim \left| \langle w_{k,I}, r^{2+2\varepsilon} w_{k,I} \partial_t \Lambda_k^2 \rangle_{L_r^2} \right| + \nu \left| \langle w_{k,I}, r^{2+2\varepsilon} w_{k,I} \partial_r^2 \Lambda_k^2 \rangle_{L_r^2} \right| \\
+ \|r^{2+\varepsilon} f_1 \Lambda_k\|_{L_r^2} \|r^{\varepsilon} w_{k,I} \Lambda_k\|_{L_r^2}^2 + \|r^{1+\varepsilon} f_2 \Lambda_k\|_{H_k^{-1}} \|r^{1+\varepsilon} w_{k,I} \Lambda_k\|_{H_k^1}^1 \\
\lesssim \nu \|r^{1+\varepsilon} w_{k,I}' \Lambda_k\|_{L_r^2}^2 + \mu_k \|r^{\varepsilon} w_{k,I} \Lambda_k\|_{L_r^2}^2 \\
+ \mu_k^{-1} \|r^{2+\varepsilon} f_1 \Lambda_k\|_{L_r^2}^2 + \nu^{-1} \|r^{1+\varepsilon} f_2 \Lambda_k\|_{H_k^{-1}}^2.$$

By integrating the above inequality over [0, T] and using (5.5), we achieve (5.4).

5.2. Space-time estimates of the homogeneous equation (5.1). In this subsection, we present the space-time estimates of the solution to the homogeneous linear equation (5.1). The main idea will be based on a decomposition of $w_{k,H}$, which was already used in [20] to derive linear enhanced dissipation around TC flow. For reader's convenience, we shall explain the motivation of the decomposition. Indeed if $\nu = 0$ in (5.1), the corresponding solution of (5.1) can be expressed as a rotation of the initial data, i.e. $e^{-i\frac{kB}{r^2}t}w_k(0)$. This motivates us to consider the solution $w_{k,H}$ may be in the vicinity of a solution similar to $e^{-i\frac{kB}{r^2}t}w_k(0)$. And we decompose the solution $w_{k,H}$ into

(5.6)
$$w_{k,H} = w_{k,H}^{(1)} + w_{k,H}^{(2)} \quad \text{with} \quad w_{k,H}^{(1)} \stackrel{\text{def}}{=} e^{-i\frac{kB}{r^2}t} w_k^{(1)}.$$

Then in view of (5.1), we find

$$e^{-i\frac{kB}{r^2}t}\partial_t w_k^{(1)} - \nu \left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2}\right) \left(e^{-i\frac{kB}{r^2}t}w_k^{(1)}\right)$$
$$= -\left(\partial_t w_{k,H}^{(2)} - \nu \left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2}\right)w_{k,H}^{(2)} + \frac{ikB}{r^2}w_{k,H}^{(2)}\right).$$

It is easy to observe that

$$-\nu \left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2}\right) \left(e^{-i\frac{kB}{r^2}t} w_k^{(1)}\right)$$

$$= -e^{-i\frac{kB}{r^2}t} \nu \left(\left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2}\right) w_k^{(1)} - \left|\frac{2kBt}{r^3}\right|^2 w_k^{(1)} + \frac{4ikBt}{r^3} \partial_r w_k^{(1)} - \frac{6ikBt}{r^4} w_k^{(1)}\right).$$

As a consequence, we obtain

$$e^{-i\frac{kB}{r^2}t} \left(\partial_t w_k^{(1)} - \nu \left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2} \right) w_k^{(1)} + \nu \left| \frac{2kBt}{r^3} \right|^2 w_k^{(1)} \right)$$

$$= - \left(\partial_t w_{k,H}^{(2)} - \nu \left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2} \right) w_{k,H}^{(2)} + \frac{ikB}{r^2} w_{k,H}^{(2)} \right)$$

$$+ e^{-i\frac{kB}{r^2}t} \nu \left(\frac{4ikBt}{r^3} \partial_r w_k^{(1)} - \frac{6ikBt}{r^4} w_k^{(1)} \right),$$

where the term

$$-\nu \left(\left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2} \right) w_k^{(1)} - \left| \frac{2kBt}{r^3} \right|^2 w_k^{(1)} \right)$$

is what we refer to as the enhanced dissipation term.

The computation motivates us to construct $w_k^{(1)}$ and $w_{k,H}^{(2)}$ in (5.6) through

(5.7)
$$\begin{cases} \partial_t w_k^{(1)} - \nu \Big(\partial_r^2 - \frac{k^2 + \Theta^2}{r^2} \Big) w_k^{(1)} + \nu \Big| \frac{2kBt}{r^3} \Big|^2 w_k^{(1)} = 0, \\ w_k^{(1)}|_{t=0} = w_k(0), \quad w_k^{(1)}|_{r=1,+\infty} = 0, \\ \Big(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2} \Big) \varphi_{k,H}^{(1)} = e^{-i\frac{kB}{r^2}t} w_k^{(1)}, \quad \varphi_{k,H}^{(1)}|_{r=1,+\infty} = 0, \end{cases}$$

and

(5.8)
$$\begin{cases} \partial_t w_{k,H}^{(2)} - \nu \left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2} \right) w_{k,H}^{(2)} + \frac{ikB}{r^2} w_{k,H}^{(2)} \\ = e^{-i\frac{kB}{r^2}t} \nu \left(i\frac{4kBt}{r^3} \partial_r w_k^{(1)} - i\frac{6kBt}{r^4} w_k^{(1)} + \frac{4 \cdot \Theta^2 + 1}{4r^2} w_k^{(1)} \right), \\ w_{k,H}^{(2)}|_{t=0} = 0, \quad w_{k,H}^{(2)}|_{r=1,+\infty} = 0, \\ \left(\partial_r^2 - \frac{k^2 - \frac{1}{4}}{r^2} \right) \varphi_{k,H}^{(2)} = w_{k,H}^{(2)}, \quad \varphi_{k,H}^{(2)}|_{r=1,+\infty} = 0. \end{cases}$$

Here $\Theta \gg 1$ is a fixed constant, which we shall take $\Theta \ge 10^9$ in the rest of this section.

Proposition 5.3. Let $(w_{k,H}, \varphi_{k,H})$ solve the homogeneous equation (5.1). Then for any $k \in \mathbb{Z} \setminus \{0\}, \varepsilon \in (0,2), T > 0$, there exist a constant $C_{\varepsilon} > 0$ depending only on ε , so that

(5.9)
$$\|r^{1+\varepsilon}w_{k,H}\Lambda_k\|_{L^{\infty}_T(L^2_r)} + \nu^{\frac{1}{2}}\|r^{1+\varepsilon}\partial_r w_{k,H}\Lambda_k\|_{L^2_T(L^2_r)} + \mu^{\frac{1}{2}}_k\|r^{\varepsilon}w_{k,H}\Lambda_k\|_{L^2_T(L^2_r)} + |kB|^{\frac{1}{2}}|k|^{\frac{1}{2}}\Big(\Big\|\frac{\partial_r\varphi_{k,H}}{r^{1-\varepsilon}}\Lambda_k\Big\|_{L^2_T(L^2_r)} + |k|\Big\|\frac{\varphi_{k,H}}{r^{2-\varepsilon}}\Lambda_k\Big\|_{L^2_T(L^2_r)}\Big) \le C_{\varepsilon}M_k(0).$$

Proof. In view of (5.8), we get, by applying Proposition 5.2 and Propositions 5.4 and 5.5 below, that

$$\begin{split} \|r^{1+\varepsilon}w_{k,H}^{(2)}\Lambda_{k}\|_{L_{T}^{\infty}(L_{r}^{2})} + \nu^{\frac{1}{2}}\|r^{1+\varepsilon}\partial_{r}w_{k,H}^{(2)}\Lambda_{k}\|_{L_{T}^{2}(L_{r}^{2})} + \mu^{\frac{1}{2}}_{k}\|r^{\varepsilon}w_{k,H}^{(2)}\Lambda_{k}\|_{L_{T}^{2}(L_{r}^{2})} \\ &+ |kB|^{\frac{1}{2}}|k|^{\frac{1}{2}}\Big(\Big\|\frac{\partial_{r}\varphi_{k,H}^{(2)}}{r^{1-\varepsilon}}\Lambda_{k}\Big\|_{L_{T}^{2}(L_{r}^{2})} + |k|\Big\|\frac{\varphi_{k,H}^{(2)}}{r^{2-\varepsilon}}\Lambda_{k}\Big\|_{L_{T}^{2}(L_{r}^{2})}\Big) \\ &\lesssim_{\varepsilon}\mu_{k}^{-\frac{1}{2}}\nu\Big(\Big\|\frac{kBt}{r^{1-\varepsilon}}\partial_{r}w_{k}^{(1)}\Lambda_{k}\Big\|_{L_{T}^{2}(L_{r}^{2})} + \Big\|\frac{kBt}{r^{2-\varepsilon}}w_{k}^{(1)}\Lambda_{k}\Big\|_{L_{T}^{2}(L_{r}^{2})} + \|r^{\varepsilon}w_{k}^{(1)}\Lambda_{k}\|_{L_{T}^{2}(L_{r}^{2})}\Big) \\ &\lesssim_{\varepsilon}\left((\nu/\mu_{k})^{\frac{1}{2}} + (\nu/\mu_{k})\right)\Big(\|r^{1+\varepsilon}w_{k}(0)\Lambda_{k}(0)\|_{L_{r}^{2}} + \|r^{2+\varepsilon}\partial_{r}w_{k}(0)\Lambda_{k}(0)\|_{L_{r}^{2}}\Big) \\ &\lesssim_{\varepsilon}\|r^{2+\varepsilon}w_{k}(0)\Lambda_{k}(0)\|_{L_{r}^{2}} + \|r^{3+\varepsilon}\partial_{r}w_{k}(0)\Lambda_{k}(0)\|_{L_{r}^{2}}, \end{split}$$

which together with (5.10), (5.20) and (5.22) ensures (5.13). We thus complete the proof of Proposition 5.3.

Proposition 5.4. Let $k \in \mathbb{Z} \setminus \{0\}$, $0 \le \alpha \le 4$, $\hat{C} \gg 1$ and $w_k^{(1)}$ solve (5.7). Then there exists a constant C > 0, which is independent of $\nu, k, B, \hat{C}, T, \alpha$, so that

(5.10)
$$\begin{aligned} \|r^{\alpha}w_{k}^{(1)}\Lambda_{k}\|_{L^{\infty}_{T}(L^{2}_{r})}^{2} + \nu\|r^{\alpha}\partial_{r}w_{k}^{(1)}\Lambda_{k}\|_{L^{2}_{T}(L^{2}_{r})}^{2} + \mu_{k}\|r^{\alpha-1}w_{k}^{(1)}\Lambda_{k}\|_{L^{2}_{T}(L^{2}_{r})}^{2} \\ + \kappa_{k}\|\frac{\kappa_{k}t}{r^{3-\alpha}}w_{k}^{(1)}\Lambda_{k}\|_{L^{2}_{T}(L^{2}_{r})}^{2} \leq C\|r^{\alpha}w_{k}(0)\Lambda_{k}(0)\|_{L^{2}_{r}}^{2}. \end{aligned}$$

Proof. For the brevity of notations, we only present the proof of (5.10) for the case $\alpha = 1$, and we divide our proof into the following six steps:

Step 1. The change of unknown: $\eta \stackrel{\text{def}}{=} w_k^{(1)} \Lambda_k$.

Let $\eta \stackrel{\text{def}}{=} w_k^{(1)} \Lambda_k$. Then in view of (5.7), we find

(5.11)
$$\begin{cases} \partial_t \eta - \nu \Big(\partial_r^2 - \frac{k^2 + \Theta^2}{r^2} \Big) \eta + \nu \Big| \frac{2kBt}{r^3} \Big|^2 \eta = R_1^{\eta} + R_2^{\eta}, \\ \eta|_{t=0} = \eta(0) = w_k(0)\Lambda_k(0), \quad \eta|_{r=1,+\infty} = 0, \end{cases}$$

where we denote $R_1^{\eta \text{def}} = -2\nu \frac{\partial_r \Lambda_k}{\Lambda_k} \partial_r \eta + (2\nu |\frac{\partial_r \Lambda_k}{\Lambda_k}|^2 - \nu \frac{\partial_r^2 \Lambda_k}{\Lambda_k})\eta$ and $R_2^{\eta \text{def}} = \frac{\partial_t \Lambda_k}{\Lambda_k}\eta$. It is easy to observe from Lemma C.2 that

$$\|r\partial_r w_k^{(1)} \Lambda_k\|_{L^2_T(L^2_r)} + \|w_k^{(1)} \Lambda_k\|_{L^2_T(L^2_r)} \approx \|r\partial_r \eta\|_{L^2_T(L^2_r)} + \|\eta\|_{L^2_T(L^2_r)}$$

So that in order prove (5.10), it remains to prove

(5.12)
$$\|r\eta\|_{L^{\infty}_{T}(L^{2}_{r})}^{2} + \nu \|r\partial_{r}\eta\|_{L^{2}_{T}(L^{2}_{r})}^{2} + \kappa_{k} \left\|\frac{\kappa_{k}t}{r^{2}}\eta\right\|_{L^{2}_{T}(L^{2}_{r})}^{2} + \mu_{k}\|\eta\|_{L^{2}_{T}(L^{2}_{r})}^{2} \le C \|r\eta(0)\|_{L^{2}_{r}}^{2}.$$

Step 2. Dyadic decomposition.

We first fix some function $\chi(r) \in \mathcal{C}_c^2[\frac{2}{3}, 2]$ (see the construction in [20]) and denote $\chi_j \stackrel{\text{def}}{=} \chi(\frac{r}{2^j})$ so that

(5.13)
$$\begin{cases} 0 \le \chi_j \le 1, \quad \chi_0(1) = 1, \quad \sum_{j \ge 0} \chi_j(r) = 1, \quad \frac{1}{2} \le \sum_{j \ge 0} \chi_j^2(r) \le 1, \ \forall \ r \ge 1, \\ \sup_r \chi_j \subset [2^{j+1}/3, 2^{j+1}], \quad \chi_j \chi_\ell = 0 \ \forall \ |j-\ell| \ge 2, \quad \|\chi\|_{C^2} \le 4 \cdot 10^6. \end{cases}$$

Then it is obvious to observe that

$$||r^{\alpha}\eta||_{L^{2}_{r}}^{2} \approx \sum_{j\geq 0} 2^{2\alpha j} ||\eta_{j}||_{L^{2}_{r}}^{2},$$

where $\eta_i \stackrel{\text{def}}{=} \chi_i \eta$, which satisfies (5.14)

$$\begin{cases} \partial_t \eta_j - \nu \Big(\partial_r^2 - \frac{k^2 + \Theta^2}{r^2} \Big) \eta_j + \nu \Big| \frac{2kBt}{r^3} \Big|^2 \eta_j = \chi_j (R_1^\eta + R_2^\eta) - \nu (2^{-2j} \chi_j'' \eta + 2^{1-j} \chi_j' \partial_r \eta), \\ \eta_j|_{t=0} = \eta_j(0), \quad \eta_j|_{r=\max(1,\frac{2j+1}{3})} = \eta_j|_{r=2^{j+1}} = 0, \quad (t,r) \in \mathbb{R}^+ \times [\max(1,2^{j+1}/3),2^{j+1}], \end{cases}$$

here and in what follows, we always denote $\chi'_{j} \stackrel{\text{def}}{=} \chi'(\frac{r}{2^{j}})$ and $\chi''_{j} \stackrel{\text{def}}{=} \chi''(\frac{r}{2^{j}})$. Noticing that $\eta(r) = \sum_{\ell \geq 0} \eta_{\ell}(r)$ for $r \geq 1$ and $\chi_{j}\chi_{\ell} = 0$ if $|j - \ell| \geq 2$, we also have

$$\chi''_j\eta = \sum_{|\ell-j| \le 1, \ell \ge 0} \chi''_j\eta_\ell, \qquad \chi'_j\partial_r\eta = \sum_{|\ell-j| \le 1, \ell \ge 0} \chi'_j\partial_r\eta_\ell.$$

Next we introduce the following energy functional:

$$\|\eta\|_{Y_T}^2 \stackrel{\text{def}}{=} \frac{1}{2} \sum_{j \ge 0} \|r\eta_j\|_{L_T^{\infty}(L_r^2)}^2 + \sum_{j \ge 0} \|\eta_j\|_{X_T}^2, \quad \text{with}$$
$$_j\|_{X_T}^2 \stackrel{\text{def}}{=} \frac{1}{2} \|r\eta_j(T)\|^2 + \nu \|r\partial_r\eta_j\|_{L^2(L^2)}^2 + \nu (k^2 + \Theta^2) \|\eta_j\|_{L^2(L^2)}^2 + 4\kappa_k^3 \|\frac{\tau}{2}\eta_j\|_L^2$$

(5.15)

$$\|\eta_j\|_{X_T}^2 \stackrel{\text{def}}{=} \frac{1}{2} \|r\eta_j(T)\|^2 + \nu \|r\partial_r\eta_j\|_{L^2_T(L^2_r)}^2 + \nu(k^2 + \Theta^2) \|\eta_j\|_{L^2_T(L^2_r)}^2 + 4\kappa_k^3 \|\frac{\tau}{r^2}\eta_j\|_{L^2_T(L^2_r)}^2$$

Step 3. The control of $\kappa_k \|\eta\|_{L^2_T(L^2_r)}^2$ by $\|\eta\|_{Y_T}^2$.

(1) $D_1 \stackrel{\text{def}}{=} \{ (\tau, r) \in (0, T) \times (1, +\infty) : r^2 \le \kappa_k \tau \};$ (2) $D_2 \stackrel{\text{def}}{=} \{ (\tau, r) \in (0, T) \times (1, +\infty) : 0 \le \kappa_k \tau \le r^2 \}.$

Here D_1 is the region where enhanced dissipation occurs. Then we decompose

$$\kappa_k \|\eta\|_{L^2_T(L^2_r)}^2 = \kappa_k \Big(\iint_{(\tau,r)\in D_1} + \iint_{(\tau,r)\in D_2} \Big) |\eta|^2 dr d\tau \stackrel{\text{def}}{=} \mathcal{I}_1 + \mathcal{I}_2.$$

Since $\sum_{j\geq 0} \chi_j \geq \frac{1}{2}$, one has

$$\mathcal{I}_{1} \leq \kappa_{k} \left\| \frac{\kappa_{k}\tau}{r^{2}} \eta \right\|_{L^{2}_{T}(L^{2}_{r})}^{2} \leq 2\kappa_{k} \sum_{j \geq 0} \left\| \frac{\kappa_{k}\tau}{r^{2}} \eta_{j} \right\|_{L^{2}_{T}(L^{2}_{r})}^{2} \leq \frac{1}{2} \|\eta\|_{Y_{T}}^{2}.$$

Whereas it follows from a direct computation that

$$\begin{aligned} \mathcal{I}_{2} &\leq 2\kappa_{k} \sum_{j \geq 0} \iint_{(\tau, r) \in D_{2}} |\eta_{j}|^{2} dr d\tau \\ &\leq 2\kappa_{k} \sum_{j \geq 0} \int_{0}^{\min(T, 2^{2j+2}/\kappa_{k})} \int_{1}^{\infty} \frac{9}{2^{2j+2}} |r\eta_{j}|^{2} dr d\tau \\ &\leq 18 \sum_{j \geq 0} \|r\eta_{j}\|_{L^{\infty}_{T}(L^{2}_{r})}^{2} \leq 36 \|\eta\|_{Y_{T}}^{2}. \end{aligned}$$

As a result, it comes out

(5.16)
$$\kappa_k \|\eta\|_{L^2_T(L^2_r)}^2 \lesssim \|\eta\|_{Y_T}^2.$$

Step 4. The estimation of $\|\eta\|_{Y_T}$. We get, by first taking the real part of L_r^2 inner product of (5.14) with $r^2\eta_j$ and then using integration by parts and (2), that

$$\begin{aligned} \frac{d}{dt} \|\eta_j(t)\|_{X_t}^2 &\leq \left| \langle r^2 \chi_j R_1^\eta, \eta_j \rangle_{L_r^2} \right| + \left| \langle r^2 \chi_j R_2^\eta, \eta_j \rangle_{L_r^2} \right| + \nu \|\eta_j\|_{L_r^2}^2 \\ &+ \nu \sum_{|\ell-j| \leq 1, \ell \geq 0} (2^{-2j} \|r^2 \chi_j''\|_{L_r^\infty} \|\eta_\ell\|_{L_r^2} + 2^{1-j} \|r\chi_j'\|_{L_r^\infty} \|r\partial_r \eta_\ell\|_{L_r^2}) \|\eta_j\|_{L_r^2}. \end{aligned}$$

Observing that $\text{supp}_r \chi_j \in [2^{j+1}/3, 2^{j+1}]$ and $\|\chi_j''\|_{L^{\infty}}, \|\chi_j'\|_{L^{\infty}} \leq 4 \cdot 10^6$, we find

$$\frac{d}{dt} \|\eta_j(t)\|_{X_t}^2 \le \left| \langle r^2 \chi_j R_1^\eta, \eta_j \rangle_{L_r^2} \right| + \left| \langle r^2 \chi_j R_2^\eta, \eta_j \rangle_{L_r^2} \right| + \nu \|\eta_j\|_{L_r^2}^2 + 16 \cdot 10^6 \nu \sum_{|\ell-j| \le 1, \ell \ge 0} (\|\eta_\ell\|_{L_r^2} + \|r\partial_r \eta_\ell\|_{L_r^2}) \|\eta_j\|_{L_r^2},$$

Then by integrating the above inequality over [0, T] and using the definition of X_T given by (5.15), we obtain

$$\begin{aligned} &\frac{1}{2} \|r\eta_j\|_{L^{\infty}_T(L^2_r)}^2 + \|\eta_j\|_{X_T}^2 \le 2 \times \left(\frac{1}{2} \|r\eta_j(0)\|_{L^2_r}^2 + \int_0^T \left|\langle r^2\chi_j R^\eta_1, \eta_j \rangle_{L^2_r}\right| d\tau \\ &+ \int_0^T \left|\langle r^2\chi_j R^\eta_2, \eta_j \rangle_{L^2_r} \left|d\tau + \frac{1}{\Theta^2} \|\eta_j\|_{X_T}^2 + 16 \cdot 10^6 (1/\Theta + 1/\Theta^2) \|\eta_j\|_{X_T} \sum_{|\ell-j|\le 1, \ell\ge 0} \|\eta_\ell\|_{X_T} \right). \end{aligned}$$

By summarizing the above inequalities for $j \ge 0$ and using Cauchy Inequality, we achieve

$$\begin{aligned} \|\eta\|_{Y_{T}}^{2} &\leq \sum_{j\geq 0} \|r\eta_{j}(0)\|_{L_{r}^{2}}^{2} + 2\int_{0}^{T} \sum_{j\geq 0} |\langle r^{2}\chi_{j}R_{1}^{\eta},\eta_{j}\rangle_{L_{r}^{2}} |d\tau \\ &+ 2\int_{0}^{T} \sum_{j\geq 0} |\langle r^{2}\chi_{j}R_{2}^{\eta},\eta_{j}\rangle_{L_{r}^{2}} |d\tau \\ &+ \frac{2}{\Theta^{2}} \|\eta\|_{Y_{T}}^{2} + 16 \cdot 10^{6} (1/\Theta + 1/\Theta^{2}) \big(\|\eta\|_{Y_{T}}^{2} + 3\sum_{j\geq 0} \sum_{|\ell-j|\leq 1,\ell\geq 0} \|\eta_{\ell}\|_{X_{T}}^{2}\big) \\ &= \sum_{j\geq 0} \|r\eta_{j}(0)\|_{L_{r}^{2}}^{2} + \big(2/\Theta^{2} + 16 \cdot 10^{7} (1/\Theta + 1/\Theta^{2})\big) \|\eta\|_{Y_{T}}^{2} \\ &+ 2\int_{0}^{T} \sum_{j\geq 0} |\langle r^{2}\chi_{j}R_{1}^{\eta},\eta_{j}\rangle_{L_{r}^{2}} |d\tau + 2\int_{0}^{T} \sum_{j\geq 0} |\langle r^{2}\chi_{j}R_{2}^{\eta},\eta_{j}\rangle_{L_{r}^{2}} |d\tau. \end{aligned}$$

Step 5. The estimation of terms in (5.17) involving R_i^{η} .

We first observe that

(5.18)
$$\nu \|r\partial_r\eta\|_{L^2_T(L^2_r)}^2 + \nu(k^2 + \Theta^2) \|\eta\|_{L^2_T(L^2_r)}^2 \lesssim \|\eta\|_{Y_T}^2$$

While it follows from Lemma C.2 that

9

$$\begin{split} &\sum_{i=1}^{2} \int_{0}^{T} \sum_{j\geq 0} \left| \langle r^{2}\chi_{j}R_{i}^{\eta}, \eta_{j} \rangle_{L_{r}^{2}} \left| d\tau \right. \\ &\lesssim \frac{\nu}{\log \hat{C}} (\|r\partial_{r}\eta\|_{L_{T}^{2}(L_{r}^{2})} + \|\eta\|_{L_{T}^{2}(L_{r}^{2})}) \|\eta\|_{L_{T}^{2}(L_{r}^{2})} + \frac{\kappa_{k}}{\hat{C}\log \hat{C}} \|\eta\|_{L_{T}^{2}(L_{r}^{2})}^{2} \\ &\lesssim \frac{\nu}{\log \hat{C}} \|r\partial_{r}\eta\|_{L_{T}^{2}(L_{r}^{2})}^{2} + \frac{\nu + \kappa_{k}}{\log \hat{C}} \|\eta\|_{L_{T}^{2}(L_{r}^{2})}^{2}. \end{split}$$

So that by taking $\hat{C} \gg 1$ and using (5.16) and (5.18), we obtain

(5.19)
$$\sum_{i=1}^{2} \int_{0}^{T} \sum_{j \ge 0} \left| \langle r^{2} \chi_{j} R_{i}^{\eta}, \eta_{j} \rangle_{L_{r}^{2}} \right| d\tau \le \frac{1}{8} \|\eta\|_{Y_{T}}^{2}$$

Step 6. The end of the proof.

Due to $\Theta \ge 10^9$, by inserting (5.19) into (5.17), we achieve

$$\begin{split} \|\eta\|_{Y_T}^2 &\leq \sum_{j\geq 0} \|r\eta_j(0)\|_{L_r^2}^2 + \left(1/4 + 2/\Theta^2 + 16\cdot 10^7 (1/\Theta + 1/\Theta^2)\right) \|\eta\|_{Y_T}^2 \\ &\leq \|r\eta(0)\|_{L_r^2}^2 + \frac{1}{2} \|\eta\|_{Y_T}^2, \end{split}$$

which together with (5.16) and (5.18) ensures (5.10). We thus complete the proof of Proposition 5.4.

Proposition 5.5. Let $k \in \mathbb{Z} \setminus \{0\}$, $\alpha \in [1, 5]$, $\hat{C} \gg 1$ and $w_k^{(1)}$ solve (5.7). Then there exists constant C > 0 independent of $\nu, k, B, \hat{C}, T, \alpha$, so that

(5.20)
$$\begin{aligned} & \left\| r^{\alpha} \partial_{r} w_{k}^{(1)} \Lambda_{k} \right\|_{L_{T}^{\infty}(L_{r}^{2})}^{2} + \nu \left\| r^{\alpha} \partial_{r}^{2} w_{k}^{(1)} \Lambda_{k} \right\|_{L_{T}^{2}(L_{r}^{2})}^{2} + \mu_{k} \left\| r^{\alpha-1} \partial_{r} w_{k}^{(1)} \Lambda_{k} \right\|_{L_{T}^{2}(L_{r}^{2})}^{2} \\ & + \kappa_{k} \left\| \frac{\kappa_{k} t}{r^{3-\alpha}} \partial_{r} w_{k}^{(1)} \Lambda_{k} \right\|_{L_{T}^{2}(L_{r}^{2})}^{2} \leq C \left(\left\| r^{\alpha} \partial_{r} w_{k}(0) \Lambda_{k}(0) \right\|_{L_{r}^{2}}^{2} + \left\| r^{\alpha-1} w_{k}(0) \Lambda_{k}(0) \right\|_{L_{r}^{2}}^{2} \right). \end{aligned}$$

Proof. We inherit the notations used in the proof of Proposition 5.4, and only present the detailed proof for the case $\alpha = 2$. We first get, by taking the real part of the L_r^2 inner product of (5.14) with $-\partial_r(r^4\partial_r\eta_j)$, that

(5.21)
$$\frac{1}{2} \frac{d}{dt} \|r^2 \partial_r \eta_j\|_{L^2_r}^2 + \nu \Re \Big\langle -\Big(\partial_r^2 - \frac{k^2 + \Theta^2}{r^2}\Big)\eta_j + \frac{4|kB|^2 t^2}{r^6} \eta_j, -\partial_r (r^4 \partial_r \eta_j)\Big\rangle_{L^2_r} \\ = \Re \left\langle \chi_j (R^\eta_1 + R^\eta_2) - \nu (2^{-2j} \chi''_j \eta + 2^{1-j} \chi'_j \partial_r \eta), -\partial_r (r^4 \partial_r \eta_j) \right\rangle_{L^2_r}.$$

By using integration by parts, one has

$$\begin{aligned} \Re \Big\langle -\nu \Big(\partial_r^2 - \frac{k^2 + \Theta^2}{r^2} \Big) \eta_j + \nu \frac{4|kB|^2 t^2}{r^6} \eta_j, \ -\partial_r (r^4 \partial_r \eta_j) \Big\rangle_{L^2_r} \\ &= \nu \|r^2 \partial_r^2 \eta_j\|_{L^2_r}^2 + \nu \Re \left\langle \partial_r^2 \eta_j, 4r^3 \partial_r \eta_j \right\rangle_{L^2_r} \\ &+ \nu (k^2 + \Theta^2) \|r \partial_r \eta_j\|_{L^2_r}^2 - 2\nu (k^2 + \Theta^2) \Re \left\langle r \eta_j, \partial_r \eta_j \right\rangle_{L^2_r} \\ &+ 4\nu |kBt|^2 \|r^{-1} \partial_r \eta_j\|_{L^2_r}^2 - 24\nu |kBt|^2 \Re \left\langle r^{-2} \eta_j, r^{-1} \partial_r \eta_j \right\rangle_{L^2_r} \\ &\geq \frac{\nu}{2} \Big(\|r^2 \partial_r^2 \eta_j\|_{L^2_r}^2 + (k^2 + \Theta^2) \|r \partial_r \eta_j\|_{L^2_r}^2 \Big) + 2\nu |kBt|^2 \|r^{-1} \partial_r \eta_j\|_{L^2_r}^2 \\ &- C \Big(\nu \|r \partial_r \eta_j\|_{L^2_r}^2 + \nu (k^2 + \Theta^2) \|\eta_j\|_{L^2_r}^2 + \nu |kBt|^2 \|r^{-2} \eta_j\|_{L^2_r}^2 \Big). \end{aligned}$$

By using the fact: $\left|\frac{\partial_t \Lambda_k}{\Lambda_k}\right| + r \left|\partial_r \frac{\partial_t \Lambda_k}{\Lambda_k}\right| \lesssim \frac{\kappa_k}{\log \hat{C}} \frac{1}{r^2}$, we obtain

$$\begin{aligned} \Re \left\langle \chi_{j} R_{2}^{\eta}, -\partial_{r} (r^{4} \partial_{r} \eta_{j}) \right\rangle_{L_{r}^{2}} \\ &= \left\langle \frac{\partial_{t} \Lambda_{k}}{\Lambda_{k}} \eta_{j}, -\partial_{r} (r^{4} \partial_{r} \eta_{j}) \right\rangle_{L_{r}^{2}} = \left\| \sqrt{\frac{\partial_{t} \Lambda_{k}}{\Lambda_{k}}} r^{2} \partial_{r} \eta_{j} \right\|_{L_{r}^{2}}^{2} + \left\langle \partial_{r} \left(\frac{\partial_{t} \Lambda_{k}}{\Lambda_{k}} \right) \eta_{j}, r^{4} \partial_{r} \eta_{j} \right\rangle \\ &\lesssim \frac{\kappa_{k}}{\log \hat{C}} \left(\| r \partial_{r} \eta_{j} \|_{L_{r}^{2}}^{2} + \| \eta_{j} \|_{L_{r}^{2}}^{2} \right). \end{aligned}$$

It follows from Lemma C.2 that

$$\begin{aligned} &\Re\left\langle\chi_{j}R_{1}^{\eta}-\nu(2^{-2j}\chi_{j}''\eta+2^{1-j}\chi_{j}'\partial_{r}\eta),-\partial_{r}(r^{4}\partial_{r}\eta_{j})\right\rangle_{L_{r}^{2}}\\ &\lesssim\left(\|\chi_{j}r^{2}R_{1}^{\eta}\|_{L_{r}^{2}}+\nu\sum_{|\ell-j|\leq 1,\ell\geq 0}\left(\|\eta_{\ell}\|_{L_{r}^{2}}+\|r\partial_{r}\eta_{\ell}\|_{L_{r}^{2}}\right)\right)\times\left(\|r^{2}\partial_{r}^{2}\eta_{j}\|_{L_{r}^{2}}+\|r\partial_{r}\eta_{j}\|_{L_{r}^{2}}\right)\\ &\leq C\nu\sum_{|\ell-j|\leq 1,\ell\geq 0}\left(\|\eta_{\ell}\|_{L_{r}^{2}}^{2}+\|r\partial_{r}\eta_{\ell}\|_{L_{r}^{2}}^{2}\right)+\frac{\nu}{4}\left(\|r^{2}\partial_{r}^{2}\eta_{j}\|_{L_{r}^{2}}^{2}+\|r\partial_{r}\eta_{j}\|_{L_{r}^{2}}^{2}\right).\end{aligned}$$

By substituting the above estimates into (5.21) and integrating the resulting inequality over [0, T], we find

$$\begin{split} \|r^{2}\partial_{r}\eta_{j}\|_{L_{T}^{\infty}(L_{r}^{2})}^{2} + \nu\|r^{2}\partial_{r}^{2}\eta_{j}\|_{L_{T}^{2}(L_{r}^{2})}^{2} + \nu(k^{2} + \Theta^{2})\|r\partial_{r}\eta_{j}\|_{L_{T}^{2}(L_{r}^{2})}^{2} + 4\kappa_{k}\left\|\frac{\kappa_{k}t}{r}\partial_{r}\eta_{j}\right\|_{L_{T}^{2}(L_{r}^{2})}^{2} \\ \lesssim \|r^{2}\partial_{r}\eta_{j}(0)\|_{L_{r}^{2}}^{2} + \frac{\kappa_{k}}{\log\hat{C}}\left(\|r\partial_{r}\eta_{j}\|_{L_{T}^{2}(L_{r}^{2})}^{2} + \|\eta_{j}\|_{L_{T}^{2}(L_{r}^{2})}^{2}\right) \\ + \sum_{|\ell-j|\leq 1,\ell\geq 0}\left(\nu\|r\partial_{r}\eta_{\ell}\|_{L_{T}^{2}(L_{r}^{2})}^{2} + \nu(k^{2} + \Theta^{2})\|\eta_{\ell}\|_{L_{T}^{2}(L_{r}^{2})}^{2} + \kappa_{k}\left\|\frac{\kappa_{k}t}{r^{2}}\eta_{\ell}\right\|_{L_{T}^{2}(L_{r}^{2})}^{2}\right). \end{split}$$

By summarizing the above inequalities for $j \ge 0$, and using (5.16) and (5.18), we achieve

$$\|r\partial_r\eta\|_{Y_T}^2 \lesssim \|r^2\partial_r\eta(0)\|_{L^2_r}^2 + \frac{\kappa_k}{\log\hat{C}} \left(\|r\partial_r\eta\|_{L^2_T(L^2_r)}^2 + \|\eta\|_{L^2_T(L^2_r)}^2\right) + \|\eta\|_{Y_T}^2$$

$$\lesssim \|r^2 \partial_r \eta(0)\|_{L^2_r}^2 + \frac{1}{\log \hat{C}} \|r \partial_r \eta\|_{Y_T}^2 + \|\eta\|_{Y_T}^2$$

So that by choosing $\hat{C} \gg 1$ and then applying Proposition 5.4, we obtain (5.20). This completes the proof of Propsition 5.5.

The next Proposition concerns the space-time estimates for the stream function $\varphi_{k,H}$, which is also referred to as the inviscid damping estimate.

Proposition 5.6. Let $k \in \mathbb{Z} \setminus \{0\}$ and $\varphi_{k,H}$ solve the homogeneous equation (5.1), we denote $\varphi_{k,H}^{(1)} \stackrel{\text{def}}{=} e^{-i\frac{kB}{r^2}t} \varphi_{k,H}$. Then there exist constants $C_{\varepsilon} > 0, \hat{C}_0 \gg 1$ depending only on ε , so that for $\hat{C} \geq \hat{C}_0$,

(5.22)
$$|kB|^{\frac{1}{2}} |k|^{\frac{1}{2}} \Big(\left\| r^{-1+\varepsilon} \partial_r \varphi_{k,H}^{(1)} \Lambda_k \right\|_{L^2_T(L^2_r)} + |k| \left\| r^{-2+\varepsilon} \varphi_{k,H}^{(1)} \Lambda_k \right\|_{L^2_T(L^2_r)} \Big) \le C_{\varepsilon} M_k(0).$$

Proof. Without loss of generality, we assume that $w_k^{(1)} \in \mathcal{C}_c^{\infty}([1,\infty))$. Observing that $\frac{d}{ds}e^{-i\frac{kB}{s^2}t} = \frac{2ikBt}{s^3}e^{-i\frac{kB}{s^2}t}$, we get, by using integration by parts, that

(5.23)
$$-r^{-2+\varepsilon}\Lambda_{k}(t,r)\varphi_{k,H}^{(1)} = \int_{1}^{\infty} r^{-2+\varepsilon}\Lambda_{k}(t,r)K_{k}(r,s)e^{-i\frac{kB}{s^{2}}t}w_{k}^{(1)}ds$$
$$= \frac{1}{2ikBt}\int_{1}^{\infty} r^{-2+\varepsilon}\Lambda_{k}(t,r)K_{k}(r,s)s^{3}w_{k}^{(1)}d\left(e^{-i\frac{kB}{s^{2}}t}\right)$$
$$= -\frac{1}{2ikBt}\int_{1}^{\infty} r^{-2+\varepsilon}\Lambda_{k}(t,r)\partial_{s}\left(K_{k}(r,s)s^{3}w_{k}^{(1)}\right)e^{-i\frac{kB}{s^{2}}t}ds,$$

and by (1.18a) that

$$-r^{-1+\varepsilon}\Lambda_{k}(t,r)\partial_{r}\varphi_{k,H}^{(1)} = \int_{1}^{\infty} r^{-1+\varepsilon}\Lambda_{k}(t,r)\partial_{r}K_{k}(r,s)e^{-i\frac{kB}{s^{2}}t}w_{k}^{(1)}ds$$

$$= \frac{1}{2ikBt}\int_{1}^{\infty} r^{-1+\varepsilon}\Lambda_{k}(t,r)\partial_{r}K_{k}(r,s)s^{3}w_{k}^{(1)}d\left(e^{-i\frac{kB}{s^{2}}t}\right)$$

$$= -\frac{1}{2ikBt}r^{-1+\varepsilon}\Lambda_{k}(t,r)r^{3}w_{k}^{(1)}(r)e^{-i\frac{kB}{r^{2}}t}$$

$$-\frac{1}{2ikBt}\int_{1}^{\infty} r^{-1+\varepsilon}\Lambda_{k}(t,r)\widetilde{K}_{k}(r,s)s^{3}w_{k}^{(1)}e^{-i\frac{kB}{s^{2}}t}ds$$

$$-\frac{1}{2ikBt}\int_{1}^{\infty} r^{-1+\varepsilon}\Lambda_{k}(t,r)\partial_{r}K_{k}(r,s)\partial_{s}\left(s^{3}w_{k}^{(1)}\right)e^{-i\frac{kB}{s^{2}}t}ds$$

Then we deduce from (C.6) and (1.18d), that

$$\begin{split} |r^{-2+\varepsilon}\Lambda_{k}(t,r)\varphi_{k,H}^{(1)}| + |k|^{-1}|r^{-1+\varepsilon}\Lambda_{k}(t,r)\partial_{r}\varphi_{k,H}^{(1)}| &- \frac{1}{2|k||kB|t}r^{2+\varepsilon}\Lambda_{k}(t,r)|w_{k}^{(1)}| \\ &\leq \frac{C}{|kB|t}\int_{1}^{\infty}r^{-2+\varepsilon}\Lambda_{k}(t,r)(\frac{r}{s})^{\frac{1}{2}}\min\left(\frac{r}{s},\frac{s}{r}\right)^{|k|}s^{3}\left(|w_{k}^{(1)}| + s|\partial_{s}w_{k}^{(1)}|\right)ds \\ &= \frac{C}{|kB|t}\int_{1}^{\infty}(rs)^{-\frac{1}{2}}(\frac{s}{r})^{1-\varepsilon}\frac{\Lambda_{k}(t,r)}{\Lambda_{k}(t,s)}\min\left(\frac{r}{s},\frac{s}{r}\right)^{|k|}s^{2+\varepsilon}\Lambda_{k}(t,s)\left(|w_{k}^{(1)}| + s|\partial_{s}w_{k}^{(1)}|\right)ds \\ &\leq \frac{C}{|kB|t}\int_{1}^{\infty}(rs)^{-\frac{1}{2}}(\frac{s}{r})^{1-\varepsilon}\max\left(1,(\frac{r}{s})^{\frac{2-\varepsilon}{2}}\right)\min\left(\frac{r}{s},\frac{s}{r}\right)^{|k|}s^{2+\varepsilon}\Lambda_{k}(t,s)\left(|w_{k}^{(1)}| + s|\partial_{s}w_{k}^{(1)}|\right)ds \\ &\leq \frac{C}{|kB|t}\mathcal{Q}_{|k|-1+\varepsilon,|k|-\frac{\varepsilon}{2}}\Big[s^{2+\varepsilon}\Lambda_{k}(t,s)\left(|w_{k}^{(1)}| + s|\partial_{s}w_{k}^{(1)}|\right)\Big], \end{split}$$

from which, (3.14a) and Propositions 5.4 and 5.5, we infer

$$\begin{split} |k|^{-1} \|r^{-1+\varepsilon} \partial_r \varphi_{k,H}^{(1)} \Lambda_k\|_{L^2_r} + \|r^{-2+\varepsilon} \varphi_{k,H}^{(1)} \Lambda_k\|_{L^2_r} \\ &\leq \frac{C_{\varepsilon}}{|kBt|} |k|^{-1} \Big(\|r^{2+\varepsilon} w_k^{(1)} \Lambda_k\|_{L^{\infty}_T(L^2_r)} + \|r^{3+\varepsilon} \partial_r w_k^{(1)} \Lambda_k\|_{L^{\infty}_T L^2_r} \Big) \\ &\leq \frac{C_{\varepsilon}}{|kBt|} |k|^{-1} \Big(\|r^{2+\varepsilon} w_k^{(1)}(0) \Lambda_k(0)\|_{L^2_r} + \|r^{3+\varepsilon} \partial_r w_k^{(1)}(0) \Lambda_k(0)\|_{L^2_r} \Big). \end{split}$$

Whereas we deduce from (C.3) and Proposition 5.4 that

$$\begin{aligned} |k| \|r^{-1+\varepsilon} \partial_r \varphi_{k,H}^{(1)} \Lambda_k\|_{L^2_r} + |k|^2 \|r^{-2+\varepsilon} \varphi_{k,H}^{(1)} \Lambda_k\|_{L^2_r} \\ &\leq C_{\varepsilon} \|r^{\varepsilon} w_k^{(1)} \Lambda_k\|_{L^{\infty}_T(L^2_r)} \leq C_{\varepsilon} \|r^{\varepsilon} w_k^{(1)}(0) \Lambda_k(0)\|_{L^2_r}. \end{aligned}$$

As a consequence, we arrive at

$$\left\| r^{-1+\varepsilon} \partial_r \varphi_{k,H}^{(1)} \Lambda_k(t) \right\|_{L^2_r}^2 + |k|^2 \left\| r^{-2+\varepsilon} \varphi_{k,H}^{(1)} \Lambda_k(t) \right\|_{L^2_r}^2$$

$$\leq C_{\varepsilon} \min\left(\frac{1}{|k|^2}, \frac{1}{|kBt|^2}\right) \left(\left\| r^{2+\varepsilon} w_k^{(1)}(0) \Lambda_k(0) \right\|_{L^2_r}^2 + \left\| r^{3+\varepsilon} \partial_r w_k^{(1)}(0) \Lambda_k(0) \right\|_{L^2_r}^2 \right).$$

By integrating the above inequality over [0, T], we complete the proof of Proposition 5.6. \Box

6. Proof of Theorem 1.3 and Theorem 1.4.

In this section, we present the proof of our main theorems. We begin by revisiting the fully nonlinear perturbation equations (1.10a) and (1.10b)

(6.1)
$$\begin{cases} \partial_t w_0 + \mathcal{T}_0 w_0 = r^{-\frac{1}{2}} \partial_r \Big(\sum_{\ell \in \mathbb{Z} \setminus \{0\}} i\ell r^{-1} \varphi_\ell w_{-\ell} \Big), \\ \partial_t w_k + \mathcal{T}_k w_k = f_1^{(1)} + f_1^{(2)} + f_1^{(3)} + f_2^{(1)} + f_2^{(2)}, \end{cases}$$

where

(6.2)
$$f_{1}^{(1)} \stackrel{\text{def}}{=} - ikr^{-1} \sum_{\ell \in \mathbb{Z} \setminus \{0,k\}} \partial_{r} \left(r^{-\frac{1}{2}} \varphi_{\ell}\right) w_{k-\ell}, \quad f_{1}^{(2)} \stackrel{\text{def}}{=} - ikr^{-\frac{3}{2}} u_{0}^{\theta} w_{k},$$
$$f_{1}^{(3)} \stackrel{\text{def}}{=} - ikr^{-1} \partial_{r} \left(r^{-\frac{1}{2}} \varphi_{k}\right) w_{0}, \quad f_{2}^{(1)} \stackrel{\text{def}}{=} ir^{-\frac{1}{2}} \sum_{\ell \in \mathbb{Z} \setminus \{0,k\}} \ell \partial_{r} \left(r^{-1} \varphi_{\ell} w_{k-\ell}\right),$$
$$f_{2}^{(2)} \stackrel{\text{def}}{=} ir^{-\frac{1}{2}} \partial_{r} \left(kr^{-1} \varphi_{k} w_{0}\right).$$

Lemma 6.1. Let w be a smooth enough solution to (1.10), and $\mathcal{M}(0), E_0(T)$ and $\mathcal{E}(T)$ be the energy functionals defined respectively by (1.12) and (1.13). Then there exists a constant C independent of ε, ν, B, T , such that

(6.3)
$$E_0(T) \le C\mathcal{M}(0) + \frac{1}{2}\mathcal{E}(T) + C|\nu B|^{-\frac{1}{2}}\mathcal{E}^2(T).$$

Proof. We first deduce from Lemma 2.3 that

$$\begin{split} &\|u_{0}^{\theta}\|_{L_{T}^{\infty}(L_{r}^{2})}^{2}+2\nu\|w_{0}\|_{L_{T}^{2}(L_{r}^{2})}^{2}\\ &\leq \|U\|_{L_{T}^{\infty}(L^{2}(\Omega))}^{2}+2\nu\|W\|_{L_{T}^{2}(L^{2}(\Omega))}^{2}\\ &\leq 2\Big(\|U(0)\|_{L^{2}(\Omega)}^{2}+8\pi|B|\sum_{k\in\mathbb{Z}\setminus\{0\}}|k|\Big\|\frac{\varphi_{k}}{r^{2}}\Big\|_{L_{T}^{2}(L_{r}^{2})}\Big\|\frac{\partial_{r}\varphi_{-k}}{r}\Big\|_{L_{T}^{2}(L_{r}^{2})}\Big)\\ &\leq 2\|U(0)\|_{L^{2}(\Omega)}^{2}+16\pi\sum_{k\in\mathbb{Z}\setminus\{0\}}E_{k}(T)E_{-k}(T), \end{split}$$

which implies

(6.4)
$$\|u_0^{\theta}\|_{L^{\infty}_T(L^2_r)} + \sqrt{2}\nu^{\frac{1}{2}} \|w_0\|_{L^2_T(L^2_r)} \le \sqrt{2} \times \left(\sqrt{2} \|U(0)\|_{L^2(\Omega)} + \sqrt{16\pi} \left(\sum_{k \neq 0} E_k(T) E_{-k}(T)\right)^{\frac{1}{2}}\right).$$

On the other hand, we get, by taking L_r^2 inner product of (1.10a) with r^2w_0 and using integration by parts, that

$$\begin{split} &\frac{1}{2} \frac{d}{dt} \| rw_0(t) \|_{L^2_r}^2 + \nu \| r\partial_r w_0 \|_{L^2_r}^2 - \frac{5\nu}{4} \| w_0 \|_{L^2_r}^2 \\ &\leq \left| \langle \sum_{k \in \mathbb{Z} \setminus \{0\}} ikr^{-\frac{1}{2}} \varphi_k w_{-k}, \ r^{-\frac{1}{2}} \partial_r \left(r^{\frac{3}{2}} w_0 \right) \rangle_{L^2_r} \right| \\ &\leq \left\| r^{-\frac{1}{2}} \sum_{k \in \mathbb{Z} \setminus \{0\}} k\varphi_k w_{-k} \right\|_{L^2_r} (\| r\partial_r w_0 \|_{L^2_r} + 3/2 \| w_0 \|_{L^2_r}) \\ &\leq \nu^{-1} \left\| r^{-\frac{1}{2}} \sum_{k \in \mathbb{Z} \setminus \{0\}} k\varphi_k w_{-k} \right\|_{L^2_r}^2 + \frac{\nu}{2} \| r\partial_r w_0 \|_{L^2_r}^2 + \frac{9\nu}{8} \| w_0 \|_{L^2_r}^2. \end{split}$$

By integrating the above inequality over [0, T], one has

$$\begin{aligned} & \left\| rw_0 \right\|_{L_T^{\infty}(L_r^2)}^2 + \nu \left\| r\partial_r w_0 \right\|_{L_T^2(L_r^2)}^2 \\ & \leq 2 \times \left(\left\| rw_0(0) \right\|_{L_r^2}^2 + 2\nu^{-1} \left\| r^{-\frac{1}{2}} \sum_{k \in \mathbb{Z} \setminus \{0\}} k\varphi_k w_{-k} \right\|_{L_T^2(L_r^2)}^2 + \frac{19\nu}{4} \left\| w_0 \right\|_{L_T^2(L_r^2)}^2 \right), \end{aligned}$$

from which and the fact:

$$x + y \le \sqrt{2}(a + b + c)$$
, if $x^2 + y^2 \le a^2 + b^2 + c^2$, for $x, y, a, b, c \ge 0$,

we infer

$$\begin{split} \|rw_0\|_{L^{\infty}_T(L^2_r)} &+ \nu^{\frac{1}{2}} \|r\partial_r w_0\|_{L^2_T(L^2_r)} \\ &\leq \sqrt{2} \times \left(\sqrt{2} \|rw_0(0)\|_{L^2_r} + 2\nu^{-\frac{1}{2}} \|r^{-\frac{1}{2}} \sum_{k \neq 0} k\varphi_k w_{-k} \|_{L^2_T(L^2_r)} + \sqrt{\frac{19}{2}} \nu^{\frac{1}{2}} \|w_0\|_{L^2_T(L^2_r)}\right) \\ &\leq 2 \|rw_0(0)\|_{L^2_r} + 2\sqrt{2}\nu^{-\frac{1}{2}} \sum_{k \neq 0} |k| \|r^{-\frac{3}{2}} \varphi_k\|_{L^2_T(L^\infty_r)} \|rw_{-k}\|_{L^{\infty}_T(L^2_r)} + \sqrt{19}\nu^{\frac{1}{2}} \|w_0\|_{L^2_T(L^2_r)}. \end{split}$$

Then along the same line to the derivation of (6.4), we deduce that

(6.5)
$$\|rw_0\|_{L^{\infty}_T(L^2_r)} + \nu^{\frac{1}{2}} \|r\partial_r w_0\|_{L^2_T(L^2_r)} \leq C \|rw_0(0)\|_{L^2_r} + C \|U(0)\|_{L^2(\Omega)} + C \|\nu B\|^{-\frac{1}{2}} \sum_{k \neq 0} E_k(T) E_{-k}(T) + \sqrt{19} \times \sqrt{16\pi} \Big(\sum_{k \neq 0} E_k(T) E_{-k}(T)\Big)^{\frac{1}{2}}.$$

In view of (1.12), we get, by summing up (6.4) with (6.5), that

$$\begin{split} E_{0}(T) &\leq C \Big(\|U(0)\|_{L^{2}(\Omega)} + \|rw_{0}(0)\|_{L^{2}_{r}} \Big) + C|\nu B|^{-\frac{1}{2}} \sum_{k \neq 0} E_{k}(T) E_{-k}(T) \\ &+ (\sqrt{2} + \sqrt{19})\sqrt{16\pi} \Big(\sum_{k \neq 0} E_{k}(T) E_{-k}(T) \Big)^{\frac{1}{2}} \\ &\leq C\mathcal{M}(0) + C|\nu B|^{-\frac{1}{2}} \mathcal{E}^{2}(T) + \frac{(\sqrt{2} + \sqrt{19})\sqrt{16\pi}}{100} \mathcal{E}(T) \\ &\leq C\mathcal{M}(0) + \frac{1}{2} \mathcal{E}(T) + C|\nu B|^{-\frac{1}{2}} \mathcal{E}^{2}(T), \end{split}$$

which leads to (6.3).

Lemma 6.2. For any $k, \ell \in \mathbb{Z}$ and any $\alpha \in (0, 1]$, there hold

(6.6)
$$|k|^{\alpha} \le |\ell|^{\alpha} + |k - \ell|^{\alpha}, \quad \log(|k| + |\ell| + e) \le \log(|k| + e)\log(|\ell| + e).$$

For any $\ell \in \mathbb{Z}$ and $\ell \neq 0, k$, there holds

$$(6.7) |k| \le 2|\ell||k-\ell|$$

Moreover, for any $\hat{C} \ge e$, one has

(6.8)
$$\Lambda_k \le \Lambda_\ell \Lambda_{k-\ell}.$$

Proof. (6.7) and the first inequality in (6.6) are obvious. Notice that

$$\log(a + x + y) \le \log(a + x)\log(a + y), \quad \text{for } x, y \ge 0, \ a \ge e,$$

which implies the second inequality in (6.6) and for $\hat{C} \ge e$,

$$\begin{split} \Lambda_k &= \log\left(\hat{C}r^2 + \nu^{\frac{1}{3}}|kB|^{\frac{2}{3}}t\right) \le \log\left(\hat{C}r^2 + \nu^{\frac{1}{3}}|B|^{\frac{2}{3}}|\ell|^{\frac{2}{3}}t + \nu^{\frac{1}{3}}|B|^{\frac{2}{3}}|k-\ell|^{\frac{2}{3}}t\right) \\ &\le \log\left(\hat{C}r^2 + \nu^{\frac{1}{3}}|B|^{\frac{2}{3}}|\ell|^{\frac{2}{3}}t\right)\log\left(\hat{C}r^2 + \nu^{\frac{1}{3}}|B|^{\frac{2}{3}}|k-\ell|^{\frac{2}{3}}t\right) = \Lambda_\ell\Lambda_{k-\ell}, \end{split}$$

which leads to (6.8). We thus complete the proof of Lemma 6.2.

Lemma 6.3. Let $k \in \mathbb{Z} \setminus \{0\}$ and $f_i^{(j)}, i = 1, 2, j = 1, 2, 3$, be given by (6.3). Then there exists a constant C_{ε} depending only on ε , such that

(6.9)
$$\mu_{k}^{-\frac{1}{2}} \Big(\|r^{2+\varepsilon} f_{1}^{(1)} \Lambda_{k}\|_{L_{T}^{2}(L_{r}^{2})} + \|r^{2+\varepsilon} f_{1}^{(2)} \Lambda_{k}\|_{L_{T}^{2}(L_{r}^{2})} + \|r^{2+\varepsilon} f_{1}^{(3)} \Lambda_{k}\|_{L_{T}^{2}(L_{r}^{2})} \Big) + \nu^{-\frac{1}{2}} \Big(\|r^{1+\varepsilon} f_{2}^{(1)} \Lambda_{k}\|_{L_{T}^{2}(H_{k}^{-1})} + \|r^{1+\varepsilon} f_{2}^{(2)} \Lambda_{k}\|_{L_{T}^{2}(H_{k}^{-1})} \Big) \leq C_{\varepsilon} |\nu B|^{-\frac{1}{2}} \sum_{\ell \in \mathbb{Z}} E_{\ell}(T) E_{k-\ell}(T).$$

Proof. Recall that $\mu_k = \max(\nu |k|^2, \nu^{\frac{1}{3}} |kB|^{\frac{2}{3}})$, we deduce from Lemma 6.2 that

 $\Lambda_k \lesssim \Lambda_\ell \Lambda_{k-\ell} \ (\forall k, \ell \in \mathbb{Z}), \quad |k|^{\frac{1}{2}} \lesssim |\ell|^{\frac{1}{2}} |k-\ell|^{\frac{1}{2}} \ (\text{for any } \ell \neq 0, k), \quad \mu_k^{-\frac{1}{2}} \leq |k|^{-\frac{1}{2}} |\nu B|^{-\frac{1}{4}},$ from which, (6.3) and (C.3), we infer

$$\begin{split} \mu_{k}^{-\frac{1}{2}} \| r^{2+\varepsilon} f_{1}^{(1)} \Lambda_{k} \|_{L_{T}^{2}(L_{r}^{2})} &\leq C |k| \mu_{k}^{-\frac{1}{2}} \sum_{\ell \in \mathbb{Z} \setminus \{0,k\}} \| r \partial_{r} (r^{-\frac{1}{2}} \varphi_{\ell}) \Lambda_{\ell} \|_{L_{T}^{\infty}(L_{r}^{\infty})} \| r^{\varepsilon} w_{k-\ell} \Lambda_{k-\ell} \|_{L_{T}^{2}(L_{r}^{2})} \\ &\leq C |k| \mu_{k}^{-\frac{1}{2}} \sum_{\ell \in \mathbb{Z} \setminus \{0,k\}} |\ell|^{-\frac{1}{2}} \| r w_{\ell} \Lambda_{\ell} \|_{L_{T}^{\infty}(L_{r}^{2})} \| r^{\varepsilon} w_{k-\ell} \Lambda_{k-\ell} \|_{L_{T}^{2}(L_{r}^{2})} \\ &\leq C |k| \mu_{k}^{-\frac{1}{2}} \sum_{\ell \neq k,0} |\ell|^{-\frac{1}{2}} \mu_{k-\ell}^{-\frac{1}{2}} E_{\ell}(T) E_{k-\ell}(T) \\ &\leq C |\nu B|^{-\frac{1}{2}} |k|^{\frac{1}{2}} \sum_{\ell \neq k,0} |\ell|^{-\frac{1}{2}} |k-\ell|^{-\frac{1}{2}} E_{\ell}(T) E_{k-\ell}(T) \\ &\leq C |\nu B|^{-\frac{1}{2}} \sum_{\ell \in \mathbb{Z} \setminus \{0,k\}} E_{\ell}(T) E_{k-\ell}(T). \end{split}$$

While it follows from (1.19) and Lemma B.3 that

 $(6.10) ||r^{\frac{1}{2}}u_0^{\theta}||_{L_T^{\infty}(L_r^{\infty})} \lesssim ||r\partial_r u_0^{\theta}||_{L_T^{\infty}(L_r^2)} + ||u_0^{\theta}||_{L_T^{\infty}(L_r^2)} \lesssim ||rw_0||_{L_T^{\infty}(L_r^2)} + ||u_0^{\theta}||_{L_T^{\infty}(L_r^2)}.$ Then we deduce from (6.3) that

$$\begin{aligned} \mu_k^{-\frac{1}{2}} \| r^{2+\varepsilon} f_1^{(2)} \Lambda_k \|_{L^2_T(L^2_r)} &\leq C |k| \mu_k^{-\frac{1}{2}} \| r^{\frac{1}{2}} u_0^{\theta} \|_{L^{\infty}_T(L^{\infty}_r)} \| r^{\varepsilon} w_k \Lambda_k \|_{L^2_T(L^2_r)} \\ &\leq C |k| \mu_k^{-\frac{1}{2}} E_0(T) \mu_k^{-\frac{1}{2}} E_k(T) \leq C |\nu B|^{-\frac{1}{2}} E_0(T) E_k(T). \end{aligned}$$

For $f_1^{(3)}$, we get, by using (C.3), that

$$\begin{split} \mu_{k}^{-\frac{1}{2}} \| r^{2+\varepsilon} f_{1}^{(3)} \Lambda_{k} \|_{L_{T}^{2}(L_{r}^{2})} &\lesssim |k| \mu_{k}^{-\frac{1}{2}} \| r^{\varepsilon} \partial_{r} (r^{-\frac{1}{2}} \varphi_{k}) \Lambda_{k} \|_{L_{T}^{2}(L_{r}^{\infty})} \| rw_{0} \|_{L_{T}^{\infty}(L_{r}^{2})} \\ &\lesssim_{\varepsilon} |k| \mu_{k}^{-\frac{1}{2}} |k|^{-\frac{1}{2}} \| r^{\varepsilon} w_{k} \Lambda_{k} \|_{L_{T}^{2}(L_{r}^{2})} \| rw_{0} \|_{L_{T}^{\infty}(L_{r}^{2})} \\ &\lesssim |k|^{\frac{1}{2}} \mu_{k}^{-1} E_{0}(T) E_{k}(T) \lesssim_{\varepsilon} |\nu B|^{-\frac{1}{2}} E_{0}(T) E_{k}(T). \end{split}$$

Notice that

$$r^{\frac{1}{2}+\varepsilon}\partial_r F\Lambda_k = \partial_r (r^{\frac{1}{2}+\varepsilon}F\Lambda_k) - F\partial_r (r^{\frac{1}{2}+\varepsilon}\Lambda_k), \quad \partial_r (r^{\frac{1}{2}+\varepsilon}\Lambda_k) \le Cr^{-\frac{1}{2}+\varepsilon}\Lambda_k,$$

where the constant C is independent of k. Then we get, by applying (1.14), that

$$\begin{split} \nu^{-\frac{1}{2}} \Big(\|r^{1+\varepsilon} f_{2}^{(1)} \Lambda_{k}\|_{L_{T}^{2}(H_{k}^{-1})} + \|r^{1+\varepsilon} f_{2}^{(2)} \Lambda_{k}\|_{L_{T}^{2}(H_{k}^{-1})} \Big) \\ \leq \nu^{-\frac{1}{2}} \sum_{\ell \in \mathbb{Z} \setminus \{0,k\}} |\ell| \Big(\|\partial_{r} (r^{-\frac{1}{2}+\varepsilon} \varphi_{\ell} w_{k-\ell} \Lambda_{k})\|_{L_{T}^{2}(H_{k}^{-1})} + \|\varphi_{\ell} w_{k-\ell} r^{-1} \partial_{r} (r^{\frac{1}{2}+\varepsilon} \Lambda_{k})\|_{L_{T}^{2}(H_{k}^{-1})} \Big) \\ + \nu^{-\frac{1}{2}} |k| \Big(\|\partial_{r} (r^{-\frac{1}{2}+\varepsilon} \varphi_{k} w_{0} \Lambda_{k})\|_{L_{T}^{2}(H_{k}^{-1})} + \|\varphi_{k} w_{0} r^{-1} \partial_{r} (r^{\frac{1}{2}+\varepsilon} \Lambda_{k})\|_{L_{T}^{2}(H_{k}^{-1})} \Big) \\ \lesssim \nu^{-\frac{1}{2}} \sum_{\ell \in \mathbb{Z} \setminus \{0,k\}} |\ell| \|r^{-\frac{1}{2}+\varepsilon} \varphi_{\ell} w_{k-\ell} \Lambda_{k}\|_{L_{T}^{2}(L_{r}^{2})} + \nu^{-\frac{1}{2}} |k| \|r^{-\frac{1}{2}+\varepsilon} \varphi_{k} w_{0} \Lambda_{k}\|_{L_{T}^{2}(L_{r}^{2})} \\ \lesssim \nu^{-\frac{1}{2}} \sum_{\ell \in \mathbb{Z} \setminus \{0,k\}} |\ell| \|r^{-\frac{3}{2}} \varphi_{\ell} \Lambda_{\ell}\|_{L_{T}^{2}(L_{r}^{\infty})} \|r^{1+\varepsilon} w_{k-\ell} \Lambda_{k-\ell}\|_{L_{T}^{\infty}(L_{r}^{2})} \end{split}$$

$$+ \nu^{-\frac{1}{2}} |k| \| r^{\varepsilon - \frac{3}{2}} \varphi_k \Lambda_k \|_{L^2_T(L^\infty_r)} \| r w_0 \|_{L^\infty_T(L^2_r)} \\ \lesssim |\nu B|^{-\frac{1}{2}} \sum_{\ell \in \mathbb{Z}} E_\ell(T) E_{k-\ell}(T).$$

This completes the proof of Lemma 6.3.

Now we are in a position to complete the proof of Theorem 1.3 and Theorem 1.4.

Proof of Theorem 1.3. Due to the local well-posedness established in Theorem A.2, here we focus on the *a priori* estimates. We first deduce from Proposition 5.1 and Lemma 6.3 that for $k \neq 0$,

$$\begin{split} E_k(T) &\lesssim_{\varepsilon} M_k(0) + \mu_k^{-\frac{1}{2}} \Big(\left\| r^{2+\varepsilon} f_1^{(1)} \Lambda_k \right\|_{L^2_T(L^2_r)} + \left\| r^{2+\varepsilon} f_1^{(2)} \Lambda_k \right\|_{L^2_T(L^2_r)} + \left\| r^{2+\varepsilon} f_1^{(3)} \Lambda_k \right\|_{L^2_T(L^2_r)} \Big) \\ &+ \nu^{-\frac{1}{2}} \Big(\left\| r^{1+\varepsilon} f_2^{(1)} \Lambda_k \right\|_{L^2_T(H^{-1}_k)} + \left\| r^{1+\varepsilon} f_2^{(2)} \Lambda_k \right\|_{L^2_T(H^{-1}_k)} \Big) \\ &\lesssim_{\varepsilon} M_k(0) + |\nu B|^{-\frac{1}{2}} \sum_{\ell \in \mathbb{Z}} E_\ell(T) E_{k-\ell}(T). \end{split}$$

While it follows from Lemma 6.1 that

$$E_0(T) \le C\mathcal{M}(0) + \frac{1}{2}\mathcal{E}(T) + C|\nu B|^{-\frac{1}{2}}\mathcal{E}^2(T).$$

By summing up the above inequalities for $k \in \mathbb{Z}$ and applying Young's inequality, we obtain

(6.11)
$$\mathcal{E}(T) \le C_{\varepsilon} \mathcal{M}(0) + \frac{1}{2} \mathcal{E}(T) + C_{\varepsilon} |\nu B|^{-\frac{1}{2}} \mathcal{E}^{2}(T).$$

Therefore, as long as

 $\mathcal{M}(0) \leq c_1(\varepsilon)\nu^{\frac{1}{2}}|B|^{\frac{1}{2}}$ for some sufficiently small $c_1(\varepsilon)$ depending only on ε ,

we deduce from (6.11) that

 $\mathcal{E}(T) \leq C_2(\varepsilon)\mathcal{M}(0)$ for any T > 0 and some $C_2(\varepsilon)$ depending only on ε .

This completes the proof of Theorem 1.3.

Proof of Theorem 1.4. By Theorem 1.3, we have

$$\log\left(\hat{C} + \nu^{\frac{1}{3}} |kB|^{\frac{2}{3}} t\right) \|r^{1+\varepsilon} w_k\|_{L^2_r} \le \|r^{1+\varepsilon} w_k \Lambda_k\|_{L^2_r} \lesssim_{\varepsilon} \mathcal{M}(0),$$

which leads to (1.15).

While we observe from Lemma 2.1 that $||W(t)||_{L^2(\Omega)}$ is decreasing with respect to time t. Furthermore, Theorem 1.3 implies $||W||_{L^2_T(L^2(\Omega))} < \infty$, so that we deduce that

$$\lim_{t \to \infty} \left(t^{\frac{1}{2}} \| W(t) \|_{L^{2}(\Omega)} \right) \le \sqrt{2} \lim_{t \to \infty} \left(\int_{t/2}^{t} \| W(s) \|_{L^{2}(\Omega)}^{2} \, ds \right)^{\frac{1}{2}} = 0.$$

This completes the proof of Theorem 1.4.

Appendix A. Local wellposedness of (1.3)

In this section, we present the result concerning the local well-posedness of the system (1.3).

Lemma A.1. Let $\varepsilon \in (0,2)$ and $T_0 \ge 0$. There holds

(A.1)
$$\|U\|_{H^1(\Omega)} \lesssim \|U\|_{\mathscr{M}(T_0)} \stackrel{def}{=} \|rw_0\|_{L^2_r} + \|u_0^\theta\|_{L^2_r} + \sum_{k \neq 0} \|r^{1+\varepsilon}w_k\Lambda_k(T_0, \cdot)\|_{L^2_r}$$

Proof. In view of (1.17) and $u_0^r = 0$, we get, by using Lemma C.1, that

$$\begin{aligned} \|U\|_{L^{2}(\Omega)}^{2} &\lesssim \|u_{0}^{\theta}\|_{L_{r}^{2}}^{2} + \sum_{k \neq 0} \left(\|u_{k}^{\theta}\|_{L_{r}^{2}}^{2} + \|u_{k}^{r}\|_{L_{r}^{2}}^{2} \right) \\ &\lesssim \|u_{0}^{\theta}\|_{L_{r}^{2}}^{2} + \sum_{k \neq 0} \left(\|\partial_{r}\varphi_{k}\|_{L_{r}^{2}}^{2} + \left\|\frac{k}{r}\varphi_{k}\right\|_{L_{r}^{2}}^{2} \right) \quad \text{[by using (1.10d)]} \\ &\lesssim \|u_{0}^{\theta}\|_{L_{r}^{2}}^{2} + \sum_{k \neq 0} |k|^{-2} \|rw_{k}\|_{L_{r}^{2}}^{2} \lesssim \|U\|_{\mathscr{M}(T_{0})}^{2}. \end{aligned}$$

Along the same line, we deduce that

$$\begin{split} \|\nabla U\|_{L^{2}(\Omega)}^{2} \lesssim \|\partial_{r}(U^{r}\mathbf{e}_{r} + U^{\theta}\mathbf{e}_{\theta})\|_{L^{2}(\Omega)}^{2} + \|r^{-1}\partial_{\theta}(U^{r}\mathbf{e}_{r} + U^{\theta}\mathbf{e}_{\theta})\|_{L^{2}(\Omega)}^{2} \\ \lesssim \|\partial_{r}U^{r}\|_{L^{2}(\Omega)}^{2} + \|\partial_{r}U^{\theta}\|_{L^{2}(\Omega)}^{2} \\ &+ \|r^{-1}\partial_{\theta}U^{r}\|_{L^{2}(\Omega)}^{2} + \|r^{-1}\partial_{\theta}U^{\theta}\|_{L^{2}(\Omega)}^{2} + \|r^{-1}U^{r}\|_{L^{2}(\Omega)}^{2} + \|r^{-1}U^{\theta}\|_{L^{2}(\Omega)}^{2} \\ \lesssim \|\partial_{r}u^{\theta}_{0}\|_{L^{2}_{r}}^{2} + \|r^{-1}u^{\theta}_{0}\|_{L^{2}_{r}}^{2} + \sum_{k\neq 0} \left(\|\partial_{r}u^{\theta}_{k}\|_{L^{2}_{r}}^{2} + \|kr^{-1}u^{\theta}_{k}\|_{L^{2}_{r}}^{2}\right) + \sum_{k\neq 0} \left(\|\partial_{r}u^{r}_{k}\|_{L^{2}_{r}}^{2} + \|kr^{-1}u^{r}_{k}\|_{L^{2}_{r}}^{2}\right) \\ \lesssim \|w_{0}\|_{L^{2}_{r}}^{2} + \|u^{\theta}_{0}\|_{L^{2}_{r}}^{2} + \sum_{k\neq 0} \left(\|\partial_{r}^{2}\varphi_{k}\|_{L^{2}_{r}}^{2} + \|kr^{-1}\partial_{r}\varphi_{k}\|_{L^{2}_{r}}^{2} + \|k^{2}r^{-2}\varphi_{k}\|_{L^{2}_{r}}^{2}\right) \\ \lesssim \|w_{0}\|_{L^{2}_{r}}^{2} + \|u^{\theta}_{0}\|_{L^{2}_{r}}^{2} + \sum_{k\neq 0} \left(\|w_{k}\|_{L^{2}_{r}}^{2} + \|k\partial_{r}\varphi_{k}\|_{L^{2}_{r}}^{2} + \|k^{2}r^{-1}\varphi_{k}\|_{L^{2}_{r}}^{2}\right) \\ \lesssim \|w_{0}\|_{L^{2}_{r}}^{2} + \|u^{\theta}_{0}\|_{L^{2}_{r}}^{2} + \sum_{k\neq 0} \left(\|w_{k}\|_{L^{2}_{r}}^{2} + \|k\partial_{r}\varphi_{k}\|_{L^{2}_{r}}^{2} + \|k^{2}r^{-1}\varphi_{k}\|_{L^{2}_{r}}^{2}\right) \\ \lesssim \|w_{0}\|_{L^{2}_{r}}^{2} + \|u^{\theta}_{0}\|_{L^{2}_{r}}^{2} + \sum_{k\neq 0} \left(\|w_{k}\|_{L^{2}_{r}}^{2} + \|rw_{k}\|_{L^{2}_{r}}^{2}\right) \lesssim \|U\|_{\mathscr{M}(T_{0})}^{2}, \end{split}$$

which completes the proof of (A.1).

Now the main result states as follows:

Theorem A.2. Let $0 < \varepsilon < 2$, $\hat{C} > \max\left(e, e^{\frac{4}{2-\varepsilon}}\right)$ and $T_0 \ge 0$. Then under the assumption that

(A.2)
$$\|U(0)\|_{\mathscr{M}(T_0)} = \|rw_0(0)\|_{L^2_r} + \|u_0^{\theta}(0)\|_{L^2_r} + \sum_{k \neq 0} \|r^{1+\varepsilon}w_k(0)\Lambda_k(T_0, \cdot)\|_{L^2_r} < \infty,$$

there exists T > 0 depending on ε , A, B, ν, T_0 and $||U(0)||_{\mathscr{M}(T_0)}$, so that the system (1.3) has a unique solution $U \in C([0, T]; \mathscr{M}(T_0))$. Moreover, we have

(A.3)
$$\begin{aligned} \|U\|_{N_{T}} \stackrel{\text{def}}{=} \|rw_{0}\|_{L_{T}^{\infty}(L_{r}^{2})} + \|u_{0}^{\theta}\|_{L_{T}^{\infty}(L_{r}^{2})} + \nu^{\frac{1}{2}}\|w_{0}\|_{L_{T}^{2}(L_{r}^{2})} + \sum_{k \neq 0} \|r^{1+\varepsilon}w_{k}\Lambda_{k}(T_{0},r)\|_{L_{T}^{\infty}(L_{r}^{2})} \\ + \nu^{\frac{1}{2}}\sum_{k \neq 0} \left(\|r^{1+\varepsilon}\partial_{r}w_{k}\Lambda_{k}(T_{0},r)\|_{L_{T}^{2}(L_{r}^{2})} + |k|\|r^{\varepsilon}w_{k}\Lambda_{k}(T_{0},r)\|_{L_{T}^{2}(L_{r}^{2})} \right) < \infty. \end{aligned}$$

The proof of the above theorem is through standard energy estimate and iteration scheme. We omit the details here.

APPENDIX B. BASIC CALCULUS

In this section, we present some inequalities which are based on calculus.

Lemma B.1. Let $k \in \mathbb{R}$, $w \in \mathcal{C}^{\infty}_{c}([1,\infty))$ with w(1) = 0, and $A(r) \in \mathcal{C}^{2}((1,\infty))$ be any real-valued function. Then, one has

$$\Re \langle \mathcal{T}_k w, A^2(r)w \rangle_{L^2_r} = \nu \|A(r)w'\|_{L^2_r}^2 - \frac{\nu}{2} \langle \left(A^2(r)\right)'', |w|^2 \rangle_{L^2_r} + \nu \left(k^2 - \frac{1}{4}\right) \|A(r)r^{-1}w\|_{L^2_r}^2.$$

Proof. Indeed we get, by using integration by parts, that

$$\Re \langle \mathcal{T}_k w, A^2(r)w \rangle_{L^2_r} = \nu \|A(r)w'\|_{L^2_r}^2 + \nu \Re \langle w', (A^2(r))'w \rangle_{L^2_r} + \nu \left(k^2 - \frac{1}{4}\right) \|A(r)r^{-1}w\|_{L^2_r}^2$$
$$= \nu \|A(r)w'\|_{L^2_r}^2 - \frac{\nu}{2} \langle \left(A^2(r)\right)'', |w|^2 \rangle_{L^2_r} + \nu \left(k^2 - \frac{1}{4}\right) \|A(r)r^{-1}w\|_{L^2_r}^2.$$

This completes the proof of Lemma B.1.

Lemma B.2. For any $\alpha \in \mathbb{R}$, $w \in \mathcal{C}^{\infty}_{c}([1,\infty))$ with w(1) = 0, we have

(B.1)
$$\|r^{\alpha}w'\|_{L^{2}_{r}}^{2} \ge (\alpha - 1/2)^{2} \|r^{\alpha - 1}w\|_{L^{2}_{r}}^{2}.$$

Proof. It is easy to observe that for any $\alpha, \beta \in \mathbb{R}$,

$$\begin{split} 0 &\leq \|r^{\beta}(r^{\alpha-\beta}w)'\|_{L^{2}_{r}}^{2} = \left\langle r^{\alpha}w' + (\alpha-\beta)r^{\alpha-1}w, r^{\alpha}w' + (\alpha-\beta)r^{\alpha-1}w\right\rangle_{L^{2}_{r}} \\ &= \|r^{\alpha}w'\|_{L^{2}_{r}}^{2} + (\alpha-\beta)^{2}\|r^{\alpha-1}w\|_{L^{2}_{r}}^{2} + (\alpha-\beta)\int_{1}^{\infty}r^{2\alpha-1}d|w|^{2} \\ &= \|r^{\alpha}w'\|_{L^{2}_{r}}^{2} + (\alpha-\beta)^{2}\|r^{\alpha-1}w\|_{L^{2}_{r}}^{2} - (2\alpha-1)(\alpha-\beta)\int_{1}^{\infty}r^{2\alpha-2}|w|^{2}dr \\ &= \|r^{\alpha}w'\|_{L^{2}_{r}}^{2} - \left((\alpha-1/2)^{2} - (\beta-1/2)^{2}\right)\|r^{\alpha-1}w\|_{L^{2}_{r}}^{2}. \end{split}$$

Taking $\beta = \frac{1}{2}$ in the above inequality leads to (B.1).

Lemma B.3. Let $\alpha \in \mathbb{R}$, $\hat{C}, a \geq 0$, $w \in \mathcal{C}^1([1,\infty))$, $A(r) = r^{\alpha}$ or $r^{\alpha} \log(\hat{C}r^2 + a)$. We assume that $||A(r)r^{-\frac{1}{2}}w||_{L^2_r} + ||A(r)r^{\frac{1}{2}}w'||_{L^2_r} < \infty$, then there exists a constant C > 0 independent of α, \hat{C}, a , such that

(B.2)
$$\|A(r)w\|_{L^{\infty}_{r}}^{2} \leq C \|A(r)r^{-\frac{1}{2}}w\|_{L^{2}_{r}} (\|A(r)r^{\frac{1}{2}}w'\|_{L^{2}_{r}} + (|\alpha|+1)\|A(r)r^{-\frac{1}{2}}w\|_{L^{2}_{r}}).$$

Moreover, there holds

(B.3)
$$\lim_{r \to +\infty} |A(r)w(r)| = 0.$$

Proof. Let $\chi(z) \in \mathcal{C}^{\infty}(\mathbb{R}_+)$ be a smooth function such that $\chi(z) = 1$ for $z \in [0,1]$ and $\chi(z) = 0$ for $z \ge 2$. We denote $\chi_R(r) \stackrel{\text{def}}{=} \chi\left(\frac{r-1}{R}\right) \in \mathcal{C}^{\infty}_c([1,\infty))$ for $R \ge 1$. Observing that $A'(r) \lesssim (|\alpha|+1)\frac{A(r)}{r}$ and $\partial_r \chi_R = \frac{1}{R} \chi'\left(\frac{r-1}{R}\right)$, we infer

$$\begin{split} |A(r)\chi_{R}w(r)|^{2} &= -\int_{r}^{\infty} \partial_{s} \left(A^{2}(s)\chi_{R}^{2}|w(s)|^{2} \right) ds \\ &= -\int_{r}^{\infty} A^{2}(s)\chi_{R}^{2} \left(w'(s)\overline{w}(s) + w(s)\overline{w}'(s) \right) ds \\ &- \int_{r}^{\infty} 2A(s)A'(s)\chi_{R}^{2}|w|^{2} ds - \int_{R+1}^{2R+1} 2A(s)^{2}\chi_{R}\partial_{s}\chi_{R}|w|^{2} ds \\ &\lesssim \|A(r)r^{-\frac{1}{2}}w\|_{L^{2}_{r}} \|A(r)r^{\frac{1}{2}}w'\|_{L^{2}_{r}} + (|\alpha|+1)\|A(r)r^{-\frac{1}{2}}w\|_{L^{2}_{r}}^{2}, \end{split}$$

which together with the fact

$$||A(r)w||_{L_r^{\infty}}^2 = \sup_{R \ge 1} ||A(r)\chi_R w||_{L_r^{\infty}}^2,$$

ensures (B.2).

While we get, by applying (B.2), that

$$\begin{split} \|A(r)(1-\chi_R)w\|_{L_r^\infty}^2 &\lesssim \|A(r)r^{-\frac{1}{2}}(1-\chi_R)w\|_{L_r^2} \left(\|A(r)r^{\frac{1}{2}}((1-\chi_R)w)'\|_{L_r^2} \\ &+ (|\alpha|+1)\|A(r)r^{-\frac{1}{2}}(1-\chi_R)w\|_{L_r^2} \right) \\ &\lesssim \|A(r)r^{-\frac{1}{2}}(1-\chi_R)w\|_{L_r^2} (\|A(r)r^{\frac{1}{2}}(1-\chi_R)w'\|_{L_r^2} \\ &+ \|A(r)r^{\frac{1}{2}}\partial_r\chi_Rw\|_{L_r^2} + (|\alpha|+1)\|A(r)r^{-\frac{1}{2}}(1-\chi_R)w\|_{L_r^2} \right) \\ &\lesssim \|A(r)r^{-\frac{1}{2}}(1-\chi_R)w\|_{L_r^2} (\|A(r)r^{\frac{1}{2}}(1-\chi_R)w'\|_{L_r^2} \\ &+ \|A(r)r^{-\frac{1}{2}}(1-\chi_{R/2})w\|_{L_r^2} + (|\alpha|+1)\|A(r)r^{-\frac{1}{2}}(1-\chi_R)w\|_{L_r^2} \right). \end{split}$$

Taking $R \to +\infty$ in the above inequality leads to (B.3). We thus complete the proof of Lemma B.3.

Lemma B.4. Let $\lambda > 0, 0 < \tilde{\delta} \ll 1$ and $E \stackrel{\text{def}}{=} \{r \ge 1 : |1 - \sqrt{\lambda}r| \le \tilde{\delta}\}$, then there holds $\|r^{-\frac{1}{2}}\mathbb{1}_E\|_{L^2_x} \lesssim \tilde{\delta}^{\frac{1}{2}}.$

Proof. Let $r_0 \stackrel{\text{def}}{=} \lambda^{-\frac{1}{2}}$, then we get, by a direct computation, that

$$\|r^{-\frac{1}{2}}\mathbb{1}_E\|_{L^2_r}^2 = \int_{E\cap[1,\infty)} \frac{dr}{r} \le \int_{r_0-\tilde{\delta}r_0}^{r_0+\delta r_0} \frac{dr}{r} = \log\frac{r_0+\tilde{\delta}r_0}{r_0-\tilde{\delta}r_0} = \log\left(1+\frac{2\tilde{\delta}}{1-\tilde{\delta}}\right) \le \frac{2\tilde{\delta}}{1-\tilde{\delta}} \lesssim \tilde{\delta}.$$

This finishes the proof of Lemma B.4.

Lemma B.5. Let $\tilde{\delta} \leq \frac{1}{4}$, $\lambda > 0$, $\beta = 1$ or 2, and let $E_m \stackrel{\text{def}}{=} \{r \geq 1 : \tilde{\delta} \leq |1 - \sqrt{\lambda}r| \leq \frac{1}{2}\}$. Then there exists a constant *C* independent of $\tilde{\delta}$ and λ such that

(B.4)
$$\left\| r^{-\frac{1}{2}} \frac{\mathbb{1}_{E_m}(r)}{(1-\lambda r^2)^{\beta}} \right\|_{L^2_r} \le C \tilde{\delta}^{-\beta+\frac{1}{2}},$$

Proof. By using the changes of variable: $\tilde{r} = \sqrt{\lambda}r$, we find for $\beta = 1$ or 2,

$$\begin{split} \left\| r^{-\frac{1}{2}} \frac{\mathbbm{1}_{E_m}}{(1-\lambda r^2)^{\beta}} \right\|_{L^2_r}^2 &= \int_1^\infty r^{-1} \frac{\mathbbm{1}_{E_m}}{(1-\lambda r^2)^{2\beta}} dr \le \int_{\tilde{\delta} \le |1-\tilde{r}| \le \frac{1}{2}} \tilde{r}^{-1} \frac{d\tilde{r}}{(1-\tilde{r}^2)^{2\beta}} \\ &\lesssim \int_{\tilde{\delta} \le |1-\tilde{r}| \le \frac{1}{2}} \frac{d\tilde{r}}{(1-\tilde{r})^{2\beta}} \lesssim \tilde{\delta}^{-2\beta+1}, \end{split}$$

which leads to (B.4), and we complete the proof of Lemma B.5.

Appendix C. Basic properties of \mathcal{K}_k and Λ_k

In this section, we shall present some basic properties of the operator: \mathcal{K}_k , which is given by (1.17), and the enhanced dissipation weight: $\Lambda_k = \log(\hat{C}r^2 + \kappa_k t)$.

Lemma C.1 (Basic properties of \mathcal{K}_k). For any $|k| \ge 1$, a > 0, $0 < \varepsilon < 2$ and $\hat{C} \ge e^{\frac{4}{2-\varepsilon}}$, there exists a constant $C_{\varepsilon} > 0$, which is independent of k, a, \hat{C} , such that

(C.1)
$$|k| \| r^{\varepsilon - 1} \mathcal{K}'_{k}[w] \|_{L^{2}_{r}} + |k|^{2} \| r^{\varepsilon - 2} \mathcal{K}_{k}[w] \|_{L^{2}_{r}} + |k|^{\frac{1}{2}} \| r^{\varepsilon - \frac{1}{2}} \mathcal{K}'_{k}[w] \|_{L^{\infty}_{r}}$$
$$+ |k|^{\frac{3}{2}} \| r^{\varepsilon - \frac{3}{2}} \mathcal{K}_{k}[w] \|_{L^{\infty}_{r}} \leq C_{\varepsilon} \| r^{\varepsilon} w \|_{L^{2}_{r}},$$

(C.2)
$$|k| \left\| r^{\varepsilon - 1} \mathcal{K}'_{k}[w] \right\|_{L^{2}_{r}} + |k|^{2} \left\| r^{\varepsilon - 2} \mathcal{K}_{k}[w] \right\|_{L^{2}_{r}} \le C_{\varepsilon} |k|^{\frac{1}{2}} \| r^{\varepsilon - \frac{1}{2}} w \|_{L^{1}_{r}},$$

$$\begin{aligned} \text{(C.3)} \qquad & |k| \left\| r^{\varepsilon-1} \log(\hat{C}r^2 + a) \mathcal{K}'_k[w] \right\|_{L^2_r} + |k|^2 \left\| r^{\varepsilon-2} \log(\hat{C}r^2 + a) \mathcal{K}_k[w] \right\|_{L^2_r} \\ & + |k|^{\frac{1}{2}} \left\| r^{\varepsilon-\frac{1}{2}} \log(\hat{C}r^2 + a) \mathcal{K}'_k[w] \right\|_{L^\infty_r} + |k|^{\frac{3}{2}} \left\| r^{\varepsilon-\frac{3}{2}} \log(\hat{C}r^2 + a) \mathcal{K}_k[w] \right\|_{L^\infty_r} \\ & \leq C_{\varepsilon} \| r^{\varepsilon} \log(\hat{C}r^2 + a) w \|_{L^2_r}, \\ \text{(C.4)} \qquad & |k| \left\| r^{\varepsilon-1} \log(\hat{C}r^2 + a) \mathcal{K}'_k[w] \right\|_{L^2_r} + |k|^2 \left\| r^{\varepsilon-2} \log(\hat{C}r^2 + a) \mathcal{K}_k[w] \right\|_{L^2_r} \\ & \leq C_{\varepsilon} |k|^{\frac{1}{2}} \| r^{\varepsilon-\frac{1}{2}} \log(\hat{C}r^2 + a) w \|_{L^1_r}. \end{aligned}$$

Proof. We first handle the estimates of the second term in (C.1) and (C.3). Recall from (1.17) that

$$\mathcal{K}_k[w] = \int_1^\infty K_k(r,s)w(s)ds, \quad \mathcal{K}'_k[w] = \int_1^\infty \partial_r K_k(r,s)w(s)ds.$$

Then for the operator $\mathcal{Q}_{a,b}[f]$ given by (3.13), we get, by using (1.18d), that

$$\begin{split} \left| \frac{\mathcal{K}_{k}[w]}{r^{2-\varepsilon}} \right| &\leq \int_{1}^{\infty} r^{\varepsilon-2} s^{-\varepsilon} |K_{k}(r,s)| s^{\varepsilon} |w|(s) ds \\ &\lesssim |k|^{-1} \int_{1}^{\infty} (rs)^{-\frac{1}{2}} \min\left(\left(\frac{r}{s}\right)^{|k|+\varepsilon-1}, \left(\frac{s}{r}\right)^{|k|+1-\varepsilon} \right) s^{\varepsilon} |w|(s) ds \\ &\lesssim |k|^{-1} \mathcal{Q}_{|k|+\varepsilon-1, |k|+1-\varepsilon} [s^{\varepsilon} w], \end{split}$$

which together with (3.14a) ensures that

(C.5)
$$\left\|\frac{\mathcal{K}_k[w]}{r^{2-\varepsilon}}\right\|_{L^2_r} \le \frac{C}{|k|} \left(\frac{1}{|k|+\varepsilon-1} + \frac{1}{|k|-\varepsilon+1}\right) \|r^{\varepsilon}w\|_{L^2_r} \le \frac{C_{\varepsilon}}{|k|^2} \|r^{\varepsilon}w\|_{L^2_r}.$$

While for $f(r) = \log(\hat{C}r^2 + a)r^{\frac{\varepsilon-2}{2}}$, it is straightforward to verify that for $\hat{C} \ge e^{\frac{4}{2-\varepsilon}}$,

$$f'(r) = \frac{2\hat{C}r}{\hat{C}r^2 + a}r^{\frac{\varepsilon - 2}{2}} - \frac{2 - \varepsilon}{2}r^{\frac{\varepsilon - 4}{2}}\log(\hat{C}r^2 + a) \le r^{\frac{\varepsilon - 4}{2}} \Big(2 - \frac{2 - \varepsilon}{2}\log\hat{C}\Big) \le 0,$$

then by separately considering the cases: $r \ge s$ and $r \le s$, to conclude that for any $a \ge 0$ and $\hat{C} \ge e^{\frac{4}{2-\varepsilon}}$, there holds

(C.6)
$$\frac{\log(\hat{C}r^2 + a)}{\log(\hat{C}s^2 + a)} \le \max\left(1, \left(\frac{r}{s}\right)^{\frac{2-\varepsilon}{2}}\right).$$

By virtue of (1.18d) and (C.6), one has

$$\begin{split} & \left| \frac{\log(\hat{C}r^2 + a)}{r^{2-\varepsilon}} \mathcal{K}_k[w] \right| \\ & \leq \frac{C}{|k|} \int_1^\infty r^{\varepsilon - 2} s^{-\varepsilon} (rs)^{\frac{1}{2}} \frac{\log(\hat{C}r^2 + a)}{\log(\hat{C}s^2 + a)} \min\left(\frac{r}{s}, \frac{s}{r}\right)^{|k|} s^{\varepsilon} \log(\hat{C}s^2 + a) |w|(s) ds \\ & \leq \frac{C}{|k|} \int_1^\infty (rs)^{-\frac{1}{2}} \left(\frac{r}{s}\right)^{\varepsilon - 1} \max\left(1, \left(\frac{r}{s}\right)^{\frac{2-\varepsilon}{2}}\right) \min\left(\frac{r}{s}, \frac{s}{r}\right)^{|k|} s^{\varepsilon} \log(\hat{C}s^2 + a) |w|(s) ds \\ & \leq \frac{C}{|k|} \int_1^\infty (rs)^{-\frac{1}{2}} \min\left(\left(\frac{r}{s}\right)^{|k| - 1+\varepsilon}, \left(\frac{s}{r}\right)^{|k| - \frac{\varepsilon}{2}}\right) s^{\varepsilon} \log(\hat{C}s^2 + a) |w|(s) ds \\ & \leq \frac{C}{|k|} \mathcal{Q}_{|k| - 1+\varepsilon, |k| - \frac{\varepsilon}{2}} [r^{\varepsilon} \log(\hat{C}r^2 + a) w], \end{split}$$

from which and (3.14a), we infer

(C.7)
$$\begin{aligned} \left\| \frac{\log(\hat{C}r^2 + a)}{r^{2-\varepsilon}} \mathcal{K}_k[w] \right\|_{L^2_r} &\leq \frac{C}{|k|} \Big(\frac{1}{|k| - \frac{\varepsilon}{2}} + \frac{1}{|k| + \varepsilon - 1} \Big) \|r^{\varepsilon} \log(\hat{C}r^2 + a)w\|_{L^2_r} \\ &\leq \frac{C_{\varepsilon}}{|k|^2} \|r^{\varepsilon} \log(\hat{C}r^2 + a)w\|_{L^2_r}. \end{aligned}$$

So far, we have completed the estimates of the second term in (C.1) and (C.3). By virtue of the estimates (1.18d), (C.6), and (3.14a), the estimates for the first term in (C.1) and (C.3) follows along the same line. For the third term in (C.1), we get, by applying Lemma B.3 with $A(r) = r^{\varepsilon - \frac{1}{2}}$ or $A(r) = r^{\varepsilon - \frac{1}{2}} \log(\hat{C}r^2 + a)$, that

$$\begin{split} \|A(r)\mathcal{K}'_{k}[w]\|_{L^{\infty}_{r}} &\lesssim \|A(r)r^{\frac{1}{2}}\mathcal{K}''_{k}[w]\|_{L^{2}_{r}}^{\frac{1}{2}}\|A(r)r^{-\frac{1}{2}}\mathcal{K}'_{k}[w]\|_{L^{2}_{r}}^{\frac{1}{2}} + \|A(r)r^{-\frac{1}{2}}\mathcal{K}'_{k}[w]\|_{L^{2}_{r}}^{\frac{1}{2}} \\ &\lesssim \left(\|A(r)r^{\frac{1}{2}}w\|_{L^{2}_{r}} + |k|^{2}\|A(r)r^{-\frac{3}{2}}\mathcal{K}_{k}[w]\|_{L^{2}_{r}}^{\frac{1}{2}}\right)^{\frac{1}{2}}\|A(r)r^{-\frac{1}{2}}\mathcal{K}'_{k}[w]\|_{L^{2}_{r}}^{\frac{1}{2}} + \|A(r)r^{-\frac{1}{2}}\mathcal{K}'_{k}[w]\|_{L^{2}_{r}}^{\frac{1}{2}} \\ &\lesssim_{\varepsilon} |k|^{-\frac{1}{2}}\|A(r)r^{\frac{1}{2}}w\|_{L^{2}_{r}}. \end{split}$$

The estimates of fourth term in (C.1) and (C.3) follow along the same lines.

Finally, for (C.2) and (C.4), we only present the estimate to typical term: $|k|^2 \left\| \frac{\mathcal{K}[w]}{r^{2-\varepsilon}} \right\|_{L^2}$. We first observe that

$$\begin{split} \frac{\mathcal{K}_{k}[w]}{r^{2-\varepsilon}} \Big| &\leq \int_{1}^{\infty} r^{\varepsilon-2} s^{-\varepsilon+\frac{1}{2}} |K_{k}(r,s)| s^{\varepsilon-\frac{1}{2}} |w|(s) ds \\ &\lesssim |k|^{-1} r^{\frac{1}{2}} \int_{1}^{\infty} r^{\varepsilon-\frac{5}{2}} s^{-\varepsilon+\frac{1}{2}} (rs)^{\frac{1}{2}} \min\left(\frac{r}{s}, \frac{s}{r}\right)^{|k|} s^{\varepsilon-\frac{1}{2}} |w|(s) ds \\ &= |k|^{-1} r^{\frac{1}{2}} \int_{1}^{\infty} (rs)^{-\frac{1}{2}} \left(\frac{s}{r}\right)^{\frac{3}{2}-\varepsilon} \min\left(\frac{r}{s}, \frac{s}{r}\right)^{|k|} s^{\varepsilon-\frac{1}{2}} |w|(s) ds \\ &= |k|^{-1} r^{\frac{1}{2}} \int_{1}^{\infty} (rs)^{-\frac{1}{2}} \min\left(\left(\frac{r}{s}\right)^{|k|-\frac{3}{2}+\varepsilon}, \left(\frac{s}{r}\right)^{|k|+\frac{3}{2}-\varepsilon}\right) s^{\varepsilon-\frac{1}{2}} |w(s)| ds \\ &= |k|^{-1} r^{\frac{1}{2}} \mathcal{Q}_{|k|-\frac{3}{2}+\varepsilon,|k|+\frac{3}{2}-\varepsilon} [r^{\varepsilon-\frac{1}{2}}w], \end{split}$$

which together with (3.14b) ensures that

$$\left\| \frac{\mathcal{K}_{k}[w]}{r^{2-\varepsilon}} \right\|_{L^{2}_{r}} \leq \frac{C}{|k|} \Big(\frac{1}{2(|k|+\varepsilon-\frac{3}{2})+1} + \frac{1}{2(|k|-\varepsilon+\frac{3}{2})-1} \Big)^{\frac{1}{2}} \|r^{\varepsilon-\frac{1}{2}}w\|_{L^{1}_{r}} \leq \frac{C_{\varepsilon}}{|k|^{\frac{3}{2}}} \|r^{\varepsilon-\frac{1}{2}}w\|_{L^{1}_{r}}.$$
 We thus complete the proof of Proposition C.1.

We thus complete the proof of Proposition C.1.

Lemma C.2. For $\beta \in \mathbb{R}$, $r \geq 1$, $\hat{C} \geq e$ and $\Lambda_k = \log(\hat{C}r^2 + \kappa_k t)$, we have

(C.8a)
$$\left|\frac{\partial_t \Lambda_k}{\Lambda_k}\right| = \frac{\kappa_k}{(\hat{C}r^2 + \kappa_k t)\log(\hat{C}r^2 + \kappa_k t)} \le \frac{\kappa_k}{\hat{C}\log\hat{C}}\frac{1}{r^2};$$

(C.8b)
$$\left|\frac{\partial_r \Lambda_k}{\Lambda_k}\right| \le \frac{2}{\log \hat{C}} \frac{1}{r}, \qquad \left|\frac{\partial_r^2 \Lambda_k}{\Lambda_k}\right| \le \frac{4}{\log \hat{C}} \frac{1}{r^2},$$

(C.8c)
$$\frac{\partial_r (r^{2\beta} \Lambda_k^2)}{r^{2\beta-1} \Lambda_k^2} \le 2|\beta| + \frac{4}{\log \hat{C}}, \quad \frac{\partial_r^2 (r^{2\beta} \Lambda_k^2)}{r^{2\beta-2} \Lambda_k^2} \le 2\beta(2\beta-1) + \frac{1}{\log \hat{C}}(16\beta+8) + \frac{8}{\log^2 \hat{C}}.$$

Proof. We first observe that

$$\partial_t \Lambda_k = \frac{\kappa_k}{\hat{C}r^2 + \kappa_k t}, \quad \partial_r \Lambda_k = \frac{2\hat{C}r}{\hat{C}r^2 + \kappa_k t}, \quad \partial_r^2 \Lambda_k = \frac{2\hat{C}\kappa_k t - 2\hat{C}^2 r^2}{(\hat{C}r^2 + \kappa_k t)^2},$$

which leads to (C.8a) and (C.8b).

Whereas observing that

$$\partial_r (r^{2\beta} \Lambda_k^2) = 2\beta r^{2\beta-1} \Lambda_k^2 + 2r^{2\beta} \Lambda_k \partial_r \Lambda_k,$$

and

$$\begin{aligned} \partial_r^2 (r^{2\beta} \Lambda_k^2) &= \partial_r^2 (r^{2\beta}) \Lambda_k^2 + 2 \partial_r r^{2\beta} \partial_r \Lambda_k^2 + r^{2\beta} \partial_r^2 \Lambda_k^2 \\ &= 2\beta (2\beta - 1) r^{2\beta - 2} \Lambda_k^2 + 8\beta r^{2\beta - 1} \Lambda_k \partial_r \Lambda_k + 2r^{2\beta} (\Lambda_k \partial_r^2 \Lambda_k + |\partial_r \Lambda_k|^2), \end{aligned}$$

we deduce (C.8c) from (C.8b).

Acknowledgement

T. Li is partially supported by National Natural Science Foundation of China under Grant 12421001. P. Zhang is partially supported by National Key R&D Program of China under grant 2021YFA1000800 and by National Natural Science Foundation of China under Grant 12421001, 12494542 and 12288201.

DECLARATIONS

Conflict of interest. The authors declare that there are no conflicts of interest.

Data availability. This article has no associated data.

References

- D. Albritton, R. Beekie and M. Novack, Enhanced dissipation and Hörmander's hypoellipticity, J. Funct. Anal., 283 (2022), Paper No. 109522.
- [2] X. An, T. He and T. Li, Nonlinear Asymptotic Stability and Transition Threshold for 2D Taylor-Couette Flows in Sobolev Spaces, Comm. Math. Phys., 405 (2024), Paper No. 146.
- [3] J. Bedrossian, P. Germain and N. Masmoudi, Stability of the Couette flow at high Reynolds number in 2D and 3D, Bull. Amer. Math. Soc. (N.S.), 56 (2019), 373-414.
- [4] J. Bedrossian and M. Coti Zelati, Enhanced dissipation, hypoellipticity, and anomalous small noise inviscid limits in shear flows, Arch. Ration. Mech. Anal., 224 (2017), 1161-1204.
- [5] W. Borchers and T. Miyakawa, On stability of exterior stationary Navier-Stokes flows, Acta Math., 174 (1995), 311-382.
- [6] D. Borrero-Echeverry, M. Schatz and R. Tagg, Transient turbulence in Taylor-Couette flow, *Phys. Rev. E*, 81 (2010), Paper No. 025301.
- [7] M. Coti Zelati, M. G. Delgadino and T. M. Elgindi, On the relation between enhanced dissipation timescales and mixing rates, *Comm. Pure Appl. Math.*, 73 (2020) 1205-1244.
- [8] M. Coti Zelati, M. Dolce and C.-C. Lo, Diffusion enhancement and Taylor dispersion for rotationally symmetric flows in discs and pipes, J. Math. Fluid Mech., 26 (2024), Paper No. 12.
- [9] Q. Chen, T. Li, D. Wei and Z. Zhang, Transition threshold for the 2-D Couette flow in a finite channel, Arch. Ration. Mech. Anal., 238 (2020), 125-183.
- [10] P. Chossat and G. Iooss, The Couette-Taylor Problem, Appl. Math. Sci., vol. 102, Springer-Verlag, New York, 1994.
- [11] P. Constantin, A. Kiselev, L. Ryzhik and A. Zlatoš, Diffusion and mixing in fluid flow, Ann. of Math, 168 (2008), 643-674.
- [12] C. David Andereck and F. Hayot, Ordered and Turbulent Patterns in Taylor-Couette Flow, NATO ASI Series B: Physics, vol. 297, 1991.
- [13] S. Ding and Z. Lin, Enhanced dissipation and transition threshold for the 2-D plane Poiseuille flow via resolvent estimate, J. Differential Equations, 332 (2022), 404-439.
- [14] B. Dubrulle, O. Dauchot, F. Daviaud, P.-Y. Longaretti, D. Richard and J.-P. Zahn, Stability and turbulent transport in Taylor-Couette flow from analysis of experimental data, *Phys. Fluids*, **17** (2005), Paper No. 095103.
- [15] Y. Feng, A. L. Mazzucato and C. Nobili, Enhanced dissipation by circularly symmetric and parallel pipe flows, *Phys. D*, 445 (2023), Paper No. 133640.
- [16] I. Gallagher, M. Higaki and Y. Maekawa, On stationary two-dimensional flows around a fast rotating disk, *Math. Nachr.*, **292** (2019), 273-308.
- [17] A. Grigorýan and L. Saloff-Coste, Dirichlet heat kernel in the exterior of a compact set, Comm. Pure Appl. Math., 55 (2002), 93-133.
- [18] S. Grossmann, D. Lohse and C. Sun, High-Reynolds Number Taylor-Couette Turbulence, Annu. Rev. Fluid Mech., 48 (2016), 53-80.
- [19] H. Kozono and T. Ogawa, On stability of Navier-Stokes flows in exterior domains, Arch. Ration. Mech. Anal., 128 (1994), 1-31.
- [20] T. Li, P. Zhang and Y. Zhang, Linear enhanced dissipation for the 2D Taylor-Couette flow in the exterior region: A supplementary example for Gearhart-Prüss type lemma, arXiv:2501.14187.
- [21] Y. Maekawa, On stability of steady circular flows in a two-dimensional exterior disk, Arch. Ration. Mech. Anal., 225 (2017), 287-374.
- [22] Y. Maekawa, Remark on stability of scale-critical stationary flows in a two-dimensional exterior disk, Springer Proceedings in Mathematics and Statistics, 215 (2017), 105-130.
- [23] T. Ma and S. Wang, Stability and Bifurcation of the Taylor Problem, Arch. Rational Mech. Anal., 181 (2006), 149-176.
- [24] G. Schneider, Nonlinear Stability of Taylor Vortices in Infinite Cylinders, Arch. Rational Mech. Anal., 144 (1998), 121-200.

- [25] G. I. Taylor, Stability of a viscous liquid contained between two rotating cylinders, Proc. R. Soc. Lond. A, 102 (1923), 541-542.
- [26] D. Wei, Diffusion and mixing in fluid flow via the resolvent estimate, Sci. China Math., 64 (2021), 507-518.
- [27] Q. S. Zhang, The global behavior of heat kernels in exterior domains, J. Funct. Anal., 200 (2003), 160-176.

(T. Li) STATE KEY LABORATORY OF MATHEMATICAL SCIENCES, ACADEMY OF MATHEMATICS & SYSTEMS SCIENCE, THE CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA. *Email address*: teli@amss.ac.cn

(P. Zhang) STATE KEY LABORATORY OF MATHEMATICAL SCIENCES, ACADEMY OF MATHEMATICS & SYS-TEMS SCIENCE, THE CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA, AND SCHOOL OF MATHE-MATICAL SCIENCES, UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, CHINA.

Email address: zp@amss.ac.cn

(Y. Zhang) ACADEMY OF MATHEMATICS & SYSTEMS SCIENCE, THE CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA.

Email address: zhangyibin22@mails.ucas.ac.cn