

ASSOCIATE PRIMES OF LOCAL COHOMOLOGY MODULES OVER CERTAIN QUOTIENTS OF REGULAR RINGS

TONY J. PUTHENPURAKAL

ABSTRACT. Let R be a regular ring containing a field k . Let $\mathbf{x} = x_1, \dots, x_r$ be a regular sequence in R such that $R/(\mathbf{x})$ is a regular ring. Fix $m \geq 1$. Set $A_m = R/(\mathbf{x})^m$. We show that for any ideal Q of A_m the set $\text{Ass} H_Q^i(A_m)$ is a finite set for $i \geq 0$, in the following cases:

- (1) $\text{char } k = p > 0$.
- (2) $\text{char } k = 0$, R is local or a smooth affine algebra over k .

1. INTRODUCTION

Let S be a Noetherian ring and let I be an ideal in S . Let $H_I^i(S)$ be the i^{th} -local cohomology module of S with respect to I . The question whether $\text{Ass} H_I^i(S)$ is a finite set was first raised by Huneke. If S is a regular ring containing a field k with $\text{char } k = p > 0$ then Huneke and Sharp proved that $\text{Ass} H_I^i(S)$ is a finite set for all $i \geq 0$, see [1]. If S is a regular ring containing a field k of characteristic zero then Lyubeznik proved that $\text{Ass} H_I^i(S)$ is a finite set or all $i \geq 0$ if either S is local or if S is a smooth affine algebra over k ; see [2]. In [4], it is proved that if $S \rightarrow R$ is a homomorphism of Noetherian rings that splits, then for every ideal I in S and every non-negative integer i , if $\text{Ass}_R H_{I_R}^i(R)$ is finite then $\text{Ass}_S H_I^i(S)$ is finite. The first example of a ring S such that $\text{Ass} H_I^i(S)$ is an infinite set was constructed by Singh [7]. In the paper [8], the authors constructed examples of rings with good properties such that $\text{Ass} H_I^i(S)$ is infinite. In this paper we construct large examples of singular rings S such that $\text{Ass} H_I^i(S)$ is finite for all $i \geq 0$.

1.1. Let R be a regular ring containing a field k . Let $\mathbf{x} = x_1, \dots, x_r$ be a regular sequence in R such that $R/(\mathbf{x})$ is a regular ring. Fix $m \geq 1$. Set $A_m = R/(\mathbf{x})^m$. We show the following result:

Theorem 1.2. *(with hypothesis as in 1.1. Let I be an ideal in R . Let P be a prime ideal in R containing \mathbf{x} . Fix $i \geq 0$ and $m \geq 1$. The following assertions are equivalent:*

- (i) $P \in \text{Ass}_R H_I^i(A_1)$.
- (ii) $P \in \text{Ass}_R H_I^i(A_m)$.

Note the result stated in the abstract easily follows from Theorem 1.2; see section 4. We note that $P \in \text{Ass}_R D$ if and only if $PA_P \in \text{Ass}_{R_P} D_P$ if and only if $P\widehat{A}_P \in \text{Ass}_{\widehat{R}_P}(D_P \otimes_{R_P} \widehat{R}_P)$. Thus we may assume $R = K[[X_1, \dots, X_d]]$ and $P = \mathfrak{m} = (X_1, \dots, X_d)$. As \mathbf{x} is a regular sequence in R with $R/(\mathbf{x})$ regular it

Date: March 27, 2025.

2020 Mathematics Subject Classification. Primary 13D45, 13A30; Secondary 13N10, 14B15.

Key words and phrases. local cohomology, graded local cohomology, Koszul cohomology, Rees Algebras, ring of differential operators.

follows that \mathbf{x} is part of regular system of parameters of R . So we may assume without loss of generality that $x_i = X_i$ for $1 \leq i \leq r$.

The proof of Theorem 1.2 splits into two cases depending on characteristic of k .

2. THE CASE WHEN $\text{char } k = p > 0$

2.1. Let K be a field of characteristic $p > 0$. Let $R = K[[X_1, \dots, X_d]]$ and let $\mathbf{x} = X_1, \dots, X_r$. Let $\mathfrak{m} = (X_1, \dots, X_d)$. Set $G = G_{\mathbf{x}}(R)$ be the associated graded ring of R with respect to \mathbf{x} . Note $G \cong R/(\mathbf{x})[X_1^*, \dots, X_r^*]$ is a regular ring. Let $\mathcal{S} = R[(\mathbf{x})t, t^{-1}]$ be the extended Rees algebra of the ideal (\mathbf{x}) . We note that \mathcal{S} is a regular ring. If T is a regular ring containing a field k of characteristic $p > 0$ then the notion of F_T modules and F_T -finite, F_T -modules was introduced by Lyubeznik, see [3]. We have \mathcal{S} is a graded $F_{\mathcal{S}}$ -finite, $F_{\mathcal{S}}$ -module. Also note that $\mathcal{S}_{t^{-1}} = R[t, t^{-1}]$ is also a graded $F_{\mathcal{S}}$ -finite, $F_{\mathcal{S}}$ -module. So $W = \mathcal{S}_{t^{-1}}/\mathcal{S} = \bigoplus_{n \geq 1} R/(\mathbf{x})^n$ is also a graded $F_{\mathcal{S}}$ -finite, $F_{\mathcal{S}}$ -module. Fix $m \geq 1$. Set $A_m = R/(\mathbf{x})^m$. We show

Theorem 2.2. (with hypotheses as in 2.1) *Let I be an ideal in R . The following assertions are equivalent:*

- (i) $\mathfrak{m} \in \text{Ass}_R H_I^i(A_1)$.
- (ii) $\mathfrak{m} \in \text{Ass}_R H_I^i(A_m)$.

Proof. We may assume K is infinite. We note that $H_{I\mathcal{S}}^i(W) = \bigoplus_{n \geq 1} H_I^i(R/(\mathbf{x})^n)$ is a $F_{\mathcal{S}}$ -finite, $F_{\mathcal{S}}$ -module. By [6, 4.1] it follows that

$$V = \ker(t^{-1}: H_{I\mathcal{S}}^i(W)(1) \rightarrow H_{I\mathcal{S}}^i(W))$$

is a graded F_G -finite F_G -module. For $n \geq 0$ we have exact sequences:

$$0 \rightarrow V_n \rightarrow H_I^i(R/(\mathbf{x})^{n+1}) \rightarrow H_I^i(R/(\mathbf{x})^n).$$

We also have $V_0 = H_I^i(R/(\mathbf{x}))$. An easy induction yields

$$\text{Ass } V_{n-1} \subseteq \text{Ass } H_I^i(R/(\mathbf{x})^n) \subseteq \bigcup_{m=0}^{n-1} \text{Ass } V_m.$$

Let $\mathfrak{n} = (X_{r+1}, \dots, X_d)$ be maximal ideal of G_0 . We note that $\mathfrak{m} \in \text{Ass}_R V_m$ if and only if the Koszul homology module $H_{d-r}(\mathfrak{n}, V_m) \neq 0$. Set $T = K[X_1^*, \dots, X_r^*]$. By [6, 1.1] we have that $H_{d-r}(\mathfrak{n}, V)$ is a graded F_T -finite, F_T -module. In particular we have by [6, 1.6(b)] that $H_{d-r}(\mathfrak{n}, V)_m = H_{d-r}(\mathfrak{n}, V_m)$ is non-zero for some $m \geq 0$ if and only if $H_{d-r}(\mathfrak{n}, V_j)$ is non-zero for ALL $j \geq 0$.

We now prove the result:

If $\mathfrak{m} \in \text{Ass}_R H_I^i(A_1)$ then as $V_0 = H_I^i(R/(\mathbf{x}))$ we have $\mathfrak{m} \in \text{Ass } V_0$. So $H_{d-r}(\mathfrak{n}, V)_0 \neq 0$. It follows that $H_{d-r}(\mathfrak{n}, V_{m-1})$ is non-zero. So $\mathfrak{m} \in \text{Ass } H_I^i(R/(\mathbf{x})^m)$.

Conversely if $\mathfrak{m} \in \text{Ass}_R H_I^i(A_m)$ then $\mathfrak{m} \in \bigcup_{j=0}^{m-1} \text{Ass } V_j$. Say $\mathfrak{m} \in \text{Ass } V_l$. Then by above argument $\mathfrak{m} \in \text{Ass } V_j$ for all $j \geq 0$. In particular $\mathfrak{m} \in \text{Ass } V_0$ and as $V_0 = H_I^i(R/(\mathbf{x}))$ the result follows. \square

3. THE CASE WHEN $\text{char } k = 0$

3.1. Let K be a field of characteristic 0. Let $R = K[[X_1, \dots, X_d]]$ and let $\mathbf{x} = X_1, \dots, X_r$. Let $\mathfrak{m} = (X_1, \dots, X_d)$. Set $G = G_{\mathbf{x}}(R)$ be the associated graded ring of R with respect to \mathbf{x} . Note $G \cong R/(\mathbf{x})[X_1^*, \dots, X_r^*]$ is a regular ring. Let $\mathcal{S} = R[(\mathbf{x})t, t^{-1}]$ be the extended Rees algebra of the ideal (\mathbf{x}) . We note that

\mathcal{S} is a regular ring. Let $D_0 =$ ring of K -linear differential operators on $G_0 = k[[X_{r+1}, \dots, X_d]]$. Let D be the r^{th} Weyl algebra over D_0 with the obvious grading. Then note that D is a ring of D -linear differential operators on G . We note that D is a ring of differential operators of G and \mathcal{S} . Also note that t^{-1} commutes with the action of differential operators. We have \mathcal{S} is a D -module. Also note that $\mathcal{S}_{t^{-1}} = R[t, t^{-1}]$ is also a D -module. So $W = \mathcal{S}_{t^{-1}}/\mathcal{S} = \bigoplus_{n \geq 1} R/(\mathbf{x})^n$ is also a D -module. Fix $m \geq 1$. Set $A_m = R/(\mathbf{x})^m$.

We show

Theorem 3.2. *(with hypotheses as in 3.1) Let I be an ideal in R . The following assertions are equivalent:*

- (i) $\mathfrak{m} \in \text{Ass}_R H_I^i(A_1)$.
- (ii) $\mathfrak{m} \in \text{Ass}_R H_I^i(A_m)$.

Proof. We note that

$$0 \rightarrow G \rightarrow W(1) \rightarrow W \rightarrow 0,$$

is a short exact sequence of D -modules. So the long exact sequence in cohomology $H_{I\mathcal{S}}^i(-)$ is a sequence graded of D -modules. As t^{-1} commutes with action of D it follows that

$$V = \ker(t^{-1}: H_{I\mathcal{S}}^i(W)(1) \rightarrow H_{I\mathcal{S}}^i(W))$$

is a graded D -module. As V is a graded quotient of $H_I^i(G)$ it follows that V is a graded holonomic generalized Eulerian D -module. For $n \geq 0$ we have exact sequences:

$$0 \rightarrow V_n \rightarrow H_I^i(R/(\mathbf{x})^{n+1}) \rightarrow H_I^i(R/(\mathbf{x})^n).$$

We also have $V_0 = H_I^i(R/(\mathbf{x}))$. An easy induction yields

$$\text{Ass } V_{n-1} \subseteq \text{Ass } H_I^i(R/(\mathbf{x})^n) \subseteq \bigcup_{m=0}^{n-1} \text{Ass } V_m.$$

Let $\mathfrak{n} = (X_{r+1}, \dots, X_d)$ be maximal ideal of G_0 . We note that $\mathfrak{m} \in \text{Ass}_R V_m$ if and only if the Koszul homology module $H_{d-r}(\mathfrak{n}, V_m) \neq 0$. Set $T = K[X_1^*, \dots, X_r^*]$. By [5, 1.20], we have that $H_{d-r}(\mathfrak{n}, V)$ is a graded generalized Eulerian holonomic module over the Weyl algebra in r -variables over K . In particular we have by [5, 6.1] that $H_{d-r}(\mathfrak{n}, V)_m = H_{d-r}(\mathfrak{n}, V_m)$ is non-zero for some $m \geq 0$ if and only if $H_{d-r}(\mathfrak{n}, V_j)$ is non-zero for ALL $j \geq 0$.

Rest of the argument is exactly similar as in 2.2. \square

4. PROOF OF RESULT STATED IN ABSTRACT

We now indicate a proof of the result stated in the abstract. Let I be the pre-image of Q in R . Let $\sharp E$ denote the number of elements in a set E . We have by Theorem 1.2;

$$\sharp \text{Ass}_{R/(\mathbf{x})^m} H_Q^i(R/(\mathbf{x})^m) = \sharp \text{Ass}_R H_I^i(R/(\mathbf{x})^m) = \sharp \text{Ass}_R H_I^i(R/(\mathbf{x})).$$

We note that if $\text{char } k = p > 0$ then as $R/(\mathbf{x})$ is regular, $\text{Ass}_R H_I^i(R/(\mathbf{x}))$ is a finite set.

If $\text{char } k = 0$, R is local or a smooth affine algebra over k , as $R/(\mathbf{x})$ is regular, $\text{Ass}_R H_I^i(R/(\mathbf{x}))$ is a finite set.

The result follows.

REFERENCES

- [1] C. Huneke and R. Sharp, *Bass Numbers of Local Cohomology Modules*, Transactions of the AMS 339, (1993), 765—779.
- [2] G. Lyubeznik, *Finiteness Properties of Local Cohomology Modules (an Application of D-modules to Commutative Algebra)*, Inv. Math. 113, (1993), 41—55.
- [3] ———, *F-modules: applications to local cohomology and D-modules in characteristic $p > 0$* , J. Reine Angew. Math. 491, (1997), 65—130.
- [4] L. Núñez-Betancourt, *Local cohomology properties of direct summands*, J. Pure Appl. Algebra 216 (2012), no. 10, 2137–2140.
- [5] T. J. Puthenpurakal, *Graded components of local cohomology modules*, Collect. Math. 73 (2022), no. 1, 135–171.
- [6] ———, *Koszul homology of F-finite module and applications*, eprint: arXiv:2307.04473
- [7] A. Singh, *p-torsion elements in local cohomology modules*, Math. Res. Lett. 7 (2000), no. 2-3, 165–176.
- [8] A. Singh and I. Swanson, *Associated primes of local cohomology modules and of Frobenius powers*, Int. Math. Res. Not. 2004, no. 33, 1703–1733.

DEPARTMENT OF MATHEMATICS, IIT BOMBAY, POWAI, MUMBAI 400 076, INDIA
Email address: tputhen@gmail.com