Optimal reinsurance in a competitive market

Lea Enzi^{*} & Stefan Thonhauser^{*}

Abstract

We study a stochastic differential game in a ruin theoretic environment. In our setting two insurers compete for market share, which is represented by a joint performance functional. Consequently, one of the insurers strives to maximize it, while the other seeks to minimize it. As a modelling basis we use classical surplus processes extended by dynamic reinsurance opportunities, which allows us to use techniques from the theory of piecewise deterministic Markov processes to analyze the resulting game. In this context we show that a dynamic programming principle for the upper and lower value of the game holds true and that these values are unique viscosity solutions to the associated Bellman-Isaacs equations. Finally, we provide some numerical illustrations.

1 Introduction

Stochastic differential games put players in a dynamic competition where both strategy and chance influence the outcome. Fleming and Souganidis (1989) laid the groundwork for the study of stochastic differential games. They considered two-player zero-sum stochastic differential games and established the key principles regarding value functions and associated Bellman-Isaacs equations. Numerous publications were ultimately based on this. Ferreira et al. (2019) for example extended this theory to a random time horizon. Buckdahn and Li (2008) considered a problem where the cost functionals are given by backward stochastic differential equations. An overview on different approaches together with a "weak formulation" of stochastic differential games was given by Possamaï et al. (2020).

This theory also proves to be suitable in the context of reinsurance, where insurers navigate a dynamic competition against each other and the uncertainties of claim arrivals. Modeling these interactions as a stochastic differential game provides a framework for investigating a range of problems. For example, Taksar and Zeng (2011) considered a reinsurance game where each of the players' wealth is represented by a surplus process which can be controlled individually by acquiring reinsurance. Assussen et al. (2019) on the other hand considered a game using the premium as control variate, accounting for the number of potential costumers.

^{*}Institute of Statistics, Graz University of Technology, Kopernikusgasse 24/III, 8010 Graz, Austria. lea.enzi@tugraz.at, stefan.thonhauser@tugraz.at

A considerable branch of actuarial literature deals with so-called reinsurance games based on Stackelberg equilibria, see for instance Chen and Shen (2018) or Cao et al. (2022). These games involve a leader and a follower with differing decision-making processes, resulting in an asymmetric situation.

The majority of existing literature relies on a diffusion approximation. However, we use a different modeling framework, which captures the specific characteristics of our problem better. We consider a similar game as in Taksar and Zeng (2011), where we use the "strategy against control" approach, introduced in Fleming and Souganidis (1989). The surplus process is given by the classical risk model, but we make use of the fact that this corresponds to a piecewise deterministic Markov process (PDMP). These types of processes were introduced by Davis (1984, 1993). We leverage the unique structure of these processes by exploiting the properties of the underlying jump filtration. A benefit of our approach is that we avoid the need for "restrictive strategies", due to the "simpler" filtration compared to diffusion processes. A comprehensive theoretical treatment of jumping Markov processes can be found in Jacod and Skorokhod (1996).

For the reinsurance control, we use the classical approach presented by Schmidli (2001, 2008) who aims at minimizing the probability of ruin. Further works related to this topic are for example by Azcue and Muler (2005, 2014) who consider a dividend maximization problem, or Eisenberg and Schmidli (2011) who minimize capital injections into an insurance portfolio.

The paper is structured as follows. In Section 2, we introduce the specific underlying insurance model and the stochastic differential game. In Section 3, we study the characteristics of the value functions and show that the dynamic programming principle holds. In Section 4, we prove that the value functions are viscosity solutions of the associated Bellman-Isaacs equations and that a comparison result holds true. In Section 5, we numerically solve these equations and compute approximations to the maximizing and minimizing controls.

2 Model setup

2.1 The underlying insurance model

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space. We consider two insurance portfolios with surplus processes of the form

$$X_t^i = x_i + c_i t - \sum_{j=1}^{N_t^i} Y_j^i, \quad i = 1, 2,$$
(2.1)

where $x_i \geq 0$ is the initial capital, $c_i > 0$ the premium rate and $N^i = (N_t^i)_{t\geq 0}$ is a homogeneous Poisson process with intensity $\lambda_i > 0$. Furthermore, $\{Y_j^i\}_{j\in\mathbb{N}}$ is a sequence of i.i.d. random variables with continuous distribution function F_Y^i , where $F_Y^i(0) = 0$. Moreover, $N^1, N^2, \{Y_j^1\}_{j\in\mathbb{N}}$ and $\{Y_j^2\}_{j\in\mathbb{N}}$ are mutually independent.

These processes are *piecewise deterministic Markov processes* (PDMPs), which were introduced by Davis (1984, 1993) and have a special structure that we will utilize later. Indeed, let $(\mathcal{Z}, \mathcal{A}, \ell)$ be the unit interval probability space, where $\mathcal{Z} = [0, 1]$, \mathcal{A} is the class of Lebesgue measurable sets and ℓ the Lebesgue

measure. Then, the random variable $U : \mathbb{Z} \to \mathbb{R}$ where U(z) = z is uniformly distributed on \mathbb{Z} . Now, let $\Omega = \prod_{i=1}^{\infty} \mathbb{Z}_i$, where $\{\mathbb{Z}_i\}_{i \in \mathbb{N}}$ are copies of \mathbb{Z} . Ω is called *Hilbert cube* and together with the product σ -algebra \mathcal{F} and product measure \mathbb{P} it builds the sample space for a countable sequence of i.i.d. random variables U_1, U_2, \ldots , where each $U_n \sim U(0, 1)$. For each $\omega = (\omega_1, \omega_2, \ldots) \in \Omega$, $U_n(\omega) = \omega_n$. Now, for fixed $i \in \{1, 2\}$, consider such a sequence of uniform random variables $\{U_i\}_{i \in \mathbb{N}}$ and define

$$S_{n}^{i}(\omega) = -\lambda_{i}^{-1} \ln(1 - U_{2n-1}(\omega)), \quad T_{n}^{i}(\omega) = \sum_{k=1}^{n} S_{k}^{i}(\omega), \quad (2.2)$$
$$X_{T_{n}^{i}}^{i}(\omega) = X_{T_{n-1}^{i}}^{i}(\omega) + c_{i}S_{n}^{i}(\omega) - (F_{Y}^{i})^{-1}(U_{2n}(\omega)),$$
$$N_{t}^{i}(\omega) = \sum_{k=1}^{\infty} \mathbb{1}_{\{T_{k}^{i}(\omega) \le t\}},$$

for $n \in \mathbb{N}$, where $T_0^i := 0$ and $X_0^i = x_i$. These definitions correspond to (2.1). For $i \in \{1, 2\}$ the generator of this PDMP is given by

$$\mathcal{A}^{i}f(x) = c_{i}f'(x) + \lambda_{i}\int_{0}^{\infty} f(x-y) - f(x) \,\mathrm{d}F_{Y}^{i}(y),$$

for $f \in \mathcal{D}(\mathcal{A}^i)$, which is the domain of the generator (see e.g. Davis (1993) or Rolski et al. (1999)).

2.2 A two players game

We consider a game similar to the one analyzed by Taksar and Zeng (2011). We assign the surplus processes (2.1) to two insurance companies, which compete on a common insurance market. We refer to the company corresponding to i = 1 as player 1 and the other as player 2. The players can observe each other's strategy. Therefore, it is necessary to employ a common filtration, i.e., the \mathbb{P} -completed filtration $\{\mathcal{F}_t\}_{t\geq 0}$ which is generated by the combined pure jump process. We define jump times similarly to (2.2), i.e., for each $\omega \in \Omega$ and i.i.d. uniform random variables U_1, U_2, \ldots with $U_n(\omega) = \omega_n$,

$$S_n(\omega) = -(\lambda_1 + \lambda_2)^{-1} \ln(1 - U_{2n-1}(\omega)), \quad T_n(\omega) = \sum_{k=1}^n S_k(\omega).$$
(2.3)

Now, we consider reinsurance controls of the following form:

Let $A_1, A_2 \subset \mathbb{R}$ be compact sets. Then, the reinsurance controls $u_1 = (u_t^1)_{t\geq 0}$ and $u_2 = (u_t^2)_{t\geq 0}$ are A_1 and A_2 -valued stochastic processes respectively, which are predictable on Ω in the sense of Definition (26.3) in (Davis, 1993, p. 67). This means, that there exists a measurable function $g_1^i : \mathbb{R}_0^+ \times \mathbb{R} \to A^i$ and measurable functions $g_k^i : \mathbb{R}_0^+ \times (\mathbb{R}_0^+ \times \mathbb{R})^{k-1} \to A^i, k = 1, 2, \ldots$, such that

$$u_{t}^{i}(\omega) = g_{1}^{i}(t,x) \mathbb{1}_{\{t \leq T_{1}(\omega)\}} + \sum_{k=2}^{\infty} g_{k}^{i}\left(t, S_{1}(\omega), X_{T_{1}}(\omega), \dots, S_{k-1}(\omega), X_{T_{k-1}}(\omega)\right) \mathbb{1}_{\{T_{k-1}(\omega) < t \leq T_{k}(\omega)\}},$$
(2.4)

for i = 1, 2. We call these controls *admissible* and denote the set of all such by \mathcal{U}_1 and \mathcal{U}_2 for i = 1, 2 respectively.

The part of the claim the first insurer has to pay is given by a retention or risk share function $r : \mathbb{R}_0^+ \times (A_1 \cup A_2) \to \mathbb{R}_0^+$. Thus, if a claim of size y occurs at time t, the insurer has to pay $r(y, u_t^i)$ and $y - r(y, u_t^i)$ is covered by the reinsurer. Consequently, it is natural to assume that r is monotone increasing in y and $r(y, \cdot) \in [0, y]$. Further, we assume that $r(y, \cdot) : (A_1 \cup A_2) \to \mathbb{R}_0^+$ is continuous for all $y \in \mathbb{R}_0^+$. The premium the first insurer has to pay for the reinsurance is described by some measurable and continuous function $p_i : A_i \to \mathbb{R}$, which reduces the premium incomes of the first insurer to

$$c_i(u_t^i) = c_i - p_i(u_t^i), \quad i = 1, 2.$$

If one assumes that reinsurance is expensive in the sense that a full cover results in a negative premium, then one would impose existence of a control $\hat{u}_i \in \mathcal{U}_i$ such that

$$c_i(\hat{u}_i) := \pi_i < 0, \quad i = 1, 2$$

In general, we will not assume this feature.

Remark 2.1. The theoretical framework remains unaffected by considering distinct retention levels, r_1 and r_2 , for the individual players. For simplicity of notation, we assume identical reinsurance types.

The controlled surplus processes are now given by

$$X_t^{i,u_i} = x_i + \int_0^t c_i(u_s^i) \,\mathrm{d}s - \sum_{j=1}^{N_t^i} r(Y_j^i, u_{T_j^i}^i), \quad i = 1, 2.$$

We define a new controlled process $X^{u_1u_2}$ as the difference of the two surplus processes

$$X_t^{u_1 u_2} = x + \int_0^t \left(c_1(u_s^1) - c_2(u_s^2) \right) \mathrm{d}s - \sum_{j=1}^{N_t^1} r(Y_j^1, u_{T_j^1}^1) + \sum_{k=1}^{N_t^2} r(Y_k^2, u_{T_k^2}^2), \quad (2.5)$$

where $x := x_1 - x_2$. We denote the common jump process by $N = N^1 + N^2$ with jump intensity $\lambda = \lambda_1 + \lambda_2$. Thus, $N_t \sim \text{Pois}(\lambda t)$ with jump times $\{T_j\}_{j \in \mathbb{N}}$. The definition of the jump times remains consistent with (2.3). Let a, b be real numbers with a < x < b and

$$\tau_x^{u_1 u_2} = \inf \left\{ t > 0 : X_t^{u_1 u_2} \notin [a, b] | X_0 = x \right\}.$$

We consider performance functionals of the form

$$J^{u_1 u_2}(x) = \mathbb{E}_x \left[\int_0^{\tau^{u_1 u_2}} e^{-\delta t} \zeta(X_t^{u_1 u_2}) \, \mathrm{d}t + e^{-\delta \tau^{u_1 u_2}} h(X_{\tau^{u_1 u_2}}^{u_1 u_2}) \right], \qquad (2.6)$$

where $\delta > 0$ and $\zeta : [a, b] \to \mathbb{R}_0^+$ is a measurable and Lipschitz continuous function. Further, $h : \mathbb{R} \to \mathbb{R}$ is measurable, Lipschitz continuous on $\mathbb{R} \setminus (a, b)$ and monotone increasing. Furthermore, it fulfills

$$\int_{\mathbb{R}} h(y)Q(x, \mathrm{d}y) = \int_{\mathbb{R}_{0}^{+}} h(x-y) \,\mathrm{d}F_{Y}^{1}(y) + \int_{\mathbb{R}_{0}^{+}} h(x+y) \,\mathrm{d}F_{Y}^{2}(y) < \infty$$

for all $x \in [a, b]$, where Q denotes the jump kernel of the process X.

Player one wants to maximize, and player two wants to minimize the functional (2.6), using their respective controls $u_1 \in \mathcal{U}_1$ and $u_2 \in \mathcal{U}_2$.

While in Taksar and Zeng (2011), the set of controls is immediately restricted to Markovian controls, we want to follow another - more general - approach based on Fleming and Souganidis (1989).

Definition 2.1. Let u and \tilde{u} be two controls. We say that $u \approx \tilde{u}$ if $u = \tilde{u} \ell \times \mathbb{P}$ -a.s. on compact time intervals.

An admissible strategy for player 1 is a mapping $\alpha : \mathcal{U}_2 \to \mathcal{U}_1$ for which $u_2 \approx \tilde{u}_2$ implies $\alpha[u_2] \approx \alpha[\tilde{u}_2]$. The set of all such strategies is denoted by Γ .

Equivalently, a mapping $\beta : \mathcal{U}_1 \to \mathcal{U}_2$ for which $u_1 \approx \tilde{u}_1$ implies $\beta[u_1] \approx \beta[\tilde{u}_1]$, is called an admissible strategy for player 2. The set of all such strategies is denoted by Δ .

With this strategy against control formulation, the lower value of the game is defined as

$$\underline{V}(x) = \inf_{\beta \in \Delta} \sup_{u_1 \in \mathcal{U}_1} J^{u_1 \beta(u_1)}(x), \qquad (2.7)$$

and the game's upper value is

$$\overline{V}(x) = \sup_{\alpha \in \Gamma} \inf_{u_2 \in \mathcal{U}_2} J^{\alpha(u_2)u_2}(x).$$
(2.8)

Remark 2.2. The precise definitions of the functions h and ζ leave room for interpretation of the game. For example, h correlates to what happens if the surplus of the first player is much larger than the one of the second player or vice versa. A possible interpretation is that if this event occurs, some restructuring of the "weaker" player occurs and the game restarts, or the two companies merge etc. With ζ , we can model for example running costs or gains but also provide a reason that player 1 might not want to exceed b immediately. All in all, the decisions only depend on a function of the difference of the surplus processes which describes the (possible) dominance of one player over the other. In our setting the two insurers are linked to each other by the shared performance functional and control interactions.

3 A-priori properties of the value functions

We start by showing that the performance functional $J^{u_1u_2}$ for two controls is quite regular. For $u_1 \in \mathcal{U}_1$ and $u_2 \in \mathcal{U}_2$ which have the form (2.4), we write $\phi^{u_1u_2}(\cdot, \cdot)$ for the function given by

$$\phi^{u_1 u_2}(t, x) = x + \int_0^t (c_1(g_k^1(s, w_{k-1})) - c_2(g_k^2(s, w_{k-1}))) \,\mathrm{d}s,$$

for $t \in [0, S_k)$, $x = X_{T_{k-1}}$ and $w_{k-1} = (S_1, X_{T_1}, \ldots, S_{k-1}, X_{T_{k-1}})$. The parameters S_j and X_{T_j} correspond to former inter-jump times and respective post-jump locations. The map $t \mapsto \phi^{u_1 u_2}(t, x)$ is the deterministic path of the controlled process between to consecutive jump times.

Lemma 3.1. The function $t \mapsto J^{u_1 u_2}(\phi^{u_1 u_2}(t, x))$ is bounded and continuous for any controls $u_1 \in \mathcal{U}_1$ and $u_2 \in \mathcal{U}_2$.

Proof. Let $x \in (a, b)$ and $u_1 \in \mathcal{U}_1, u_2 \in \mathcal{U}_2$. We start with the boundedness. Let $M_h := \sup_{x \in [a,b]} \int_0^\infty |h(y)| Q(x, \mathrm{d}y)$. Then,

$$\begin{aligned} |J^{u_1 u_2}(x)| &\leq \|\zeta\|_{\infty} \mathbb{E}_x \left[\int_0^{\tau^{u_1 u_2}} e^{-\delta t} \, \mathrm{d}t \right] + \mathbb{E}_x \left[|h(X^{u_1 u_2}_{\tau^{u_1 u_2}})| \right] \\ &\leq \frac{\|\zeta\|_{\infty}}{\delta} + M_h =: M. \end{aligned}$$

For the continuity, we need the form of a predictable process (2.4), which gives us the following: For some time $t \in \mathbb{R}_0^+$, there exist measurable functions g^1 and g^2 such that $u_t^1 = g^1(t, x)$ and $u_t^2 = g^2(t, x)$ until the first jump occurs. Let t^* be such that the process does not reach the boundary continuously before this time. Then,

$$J^{u_1 u_2}(x) = \mathbb{P}(T_1 > t^*) \left(\int_0^{t^*} e^{-\delta t} \zeta \left(x + \int_0^t c_1(g^1(s, x)) - c_2(g^2(s, x)) \, \mathrm{d}s \right) \mathrm{d}t \\ + \mathbb{E}_x \left[\int_{t^*}^{\tau^{u_1 u_2}} e^{-\delta t} \zeta(X_t^{u_1 u_2}) \, \mathrm{d}t + e^{-\delta \tau^{u_1 u_2}} h(X_{\tau^{u_1 u_2}}^{u_1 u_2}) \left| T_1 > t^* \right] \right) \\ + \mathbb{P}(T_1 \le t^*) \mathbb{E}_x \left[\int_0^{\tau^{u_1 u_2}} e^{-\delta t} \zeta(X_t^{u_1 u_2}) \, \mathrm{d}t + e^{-\delta \tau^{u_1 u_2}} h(X_{\tau^{u_1 u_2}}^{u_1 u_2}) \left| T_1 \le t^* \right] \right].$$

By using the tower and the Markov property, we get that

$$\begin{aligned} J^{u_1 u_2}(x) &= \mathbb{P}(T_1 > t^*) \left(\int_0^{t^*} e^{-\delta t} \zeta \left(x + \int_0^t c_1(g^1(s, x)) - c_2(g^2(s, x)) \, \mathrm{d}s \right) \mathrm{d}t \\ &+ e^{-\delta t^*} J^{\tilde{u}_1 \tilde{u}_2} \left(x + \int_0^{t^*} c_1(g^1(t, x)) - c_2(g^2(t, x)) \, \mathrm{d}t \right) \right) \\ &+ \mathbb{P}(T_1 \le t^*) \mathbb{E}_x \left[\int_0^{\tau^{u_1 u_2}} e^{-\delta t} \zeta(X_t^{u_1 u_2}) \, \mathrm{d}t + e^{-\delta \tau^{u_1 u_2}} h(X_{\tau^{u_1 u_2}}^{u_1 u_2}) \right| T_1 \le t^* \right], \end{aligned}$$

where $\tilde{u}_{t}^{i} = u_{t^{*}+t}^{i}, i = 1, 2$. Thus,

$$\left| J^{u_1 u_2}(x) - J^{\tilde{u}_1 \tilde{u}_2} \left(x + \int_0^{t^*} c_1(g^1(t,x)) - c_2(g^2(t,x)) \, \mathrm{d}t \right) \right| \\ \leq \left| \mathbb{P}(T_1 > t^*) e^{-\delta t^*} - 1 \right| M + \|\zeta\|_{\infty} t^* + \mathbb{P}(T_1 \le t^*) M \\ = M \left(2 - e^{-(\lambda_1 + \lambda_2)t^*} (e^{-\delta t^*} + 1) \right) + \|\zeta\|_{\infty} t^*.$$

For $t^* \to 0$ the r.h.s. converges to 0. Since u_i and \tilde{u}_i correspond to the same function g^i for i = 1, 2, we are done.

Corollary 3.1. The above proof also shows that $J^{u_1u_2}$ is absolutely continuous along the ODE path, independent of u_1 and u_2 . Therefore, it is in the domain of the generator of $X^{u_1u_2}$.

Lemma 3.2. \underline{V} and \overline{V} are bounded and continuous.

Proof. The boundedness follows directly from Lemma 3.1. For the continuity, we start with \underline{V} . Let x > y and consider the difference $\underline{V}(x) - \underline{V}(y)$,

$$\underline{V}(x) - \underline{V}(y) = \inf_{\beta \in \Delta} \sup_{u_1 \in \mathcal{U}_1} J^{u_1 \beta(u_1)}(x) - \inf_{\beta \in \Delta} \sup_{u_1 \in \mathcal{U}_1} J^{u_1 \beta(u_1)}(y).$$

For some $\varepsilon > 0$, let β^y be an admissible ε -optimal strategy corresponding to $\underline{V}(y)$, and u^x be an admissible ε -optimal control corresponding to $\underline{V}(x)$. We denote by X^x and X^y the controlled processes starting in x and y respectively, with corresponding exit times τ_x and τ_y . Notice, that u^x is also an admissible control for the process starting in y. From Definition 2.1, the answering strategy $\beta^y(u^x)$ is the same for both processes. Consequently, for a fixed ω the two processes move parallel until time $\tau_x \wedge \tau_y$. We get

$$\underline{V}(x) - \underline{V}(y) \leq \mathbb{E} \left[\int_{0}^{\tau_{x} \wedge \tau_{y}} e^{-\delta t} (\zeta(X_{t}^{x}) - \zeta(X_{t}^{y})) dt \right] \\ + \mathbb{E} \left[\int_{\tau_{y}}^{\tau_{x}} e^{-\delta t} \zeta(X_{t}^{x}) dt \, \mathbb{1}_{\{\tau_{y} < \tau_{x}\}} \right] - \mathbb{E} \left[\int_{\tau_{x}}^{\tau_{y}} e^{-\delta t} \zeta(X_{t}^{y}) dt \, \mathbb{1}_{\{\tau_{x} < \tau_{y}\}} \right] \\ + \mathbb{E} \left[e^{-\tau_{x}} h(X_{\tau_{x}}^{x}) - e^{-\tau_{y}} h(X_{\tau_{y}}^{y}) \right] + 2\varepsilon \\ \leq \int_{0}^{\infty} e^{-\delta t} L_{\zeta}(x-y) dt + L_{h} |x-y| + 2M \left(\mathbb{P}(\tau_{y} < \tau_{x}) + \mathbb{P}(\tau_{x} < \tau_{y}) \right) + 2\varepsilon.$$

Let $T_x := \inf\{t \ge 0 : X_t^x \le a + (x - y) \lor X_t^x \ge b + (x - y)\}$. Then, $\mathbb{P}(\tau_y < \tau_x) = \mathbb{P}(T_x < \tau_x)$, which goes to zero if $y \to x$. With a similar argument, $\mathbb{P}(\tau_x < \tau_y)$ goes to zero too.

The same holds for $\underline{V}(y) - \underline{V}(x)$ by choosing suitable controls and strategies. Therefore, $|\underline{V}(x) - \underline{V}(y)| \to 0$ as $|x - y| \to 0$. Now, we consider

$$\overline{V}(x) - \overline{V}(y) = \sup_{\alpha \in \Gamma} \inf_{u_2 \in U_2} J^{\alpha(u_2)u_2}(x) - \sup_{\alpha \in \Gamma} \inf_{u_2 \in U_2} J^{\alpha(u_2)u_2}(y)$$

By choosing an admissible ε -optimal strategy α^x corresponding to $\overline{V}(x)$ and an admissible ε -optimal control u^y corresponding to $\overline{V}(y)$ we get by similar arguments as before, that $|\overline{V}(x) - \overline{V}(y)| \to 0$ as $|x - y| \to 0$.

Remark 3.1. One may notice that in the following theorem we do not need to deal with so-called *r*-strategies as elaborated in Fleming and Souganidis (1989). This is because of the natural form of predictable processes with respect to the pure jump filtration, see (2.4).

Theorem 3.1 (Dynamic programming principle). Under our model assumptions, the following holds for any $\{\mathcal{F}_t\}_{t\geq 0}$ bounded stopping time T:

$$\underline{V}(x) = \inf_{\beta \in \Delta} \sup_{u_1 \in \mathcal{U}_1} \mathbb{E}_x \left[\int_0^{\tau^{u_1 \beta(u_1)} \wedge T} e^{-\delta s} \zeta \left(X_s^{u_1 \beta(u_1)} \right) \mathrm{d}s \right] \\
+ e^{-\delta T} \underline{V} \left(X_T^{u_1 \beta(u_1)} \right) \mathbb{1}_{\{T < \tau^{u_1 \beta(u_1)}\}} + e^{-\delta \tau^{u_1 \beta(u_1)}} h(X_{\tau^{u_1 \beta(u_1)}}^{u_1 \beta(u_1)}) \mathbb{1}_{\{T \ge \tau^{u_1 \beta(u_1)}\}} \right], \\
\overline{V}(x) = \sup_{\alpha \in \Gamma} \inf_{u_2 \in \mathcal{U}_2} \mathbb{E}_x \left[\int_0^{\tau^{\alpha(u_2)u_2} \wedge T} e^{-\delta s} \zeta \left(X_s^{\alpha(u_2)u_2} \right) \mathrm{d}s \right] , \quad (3.2) \\
+ e^{-\delta T} \overline{V} \left(X_T^{\alpha(u_2)u_2} \right) \mathbb{1}_{\{T < \tau^{\alpha(u_2)u_2}\}} + e^{-\delta \tau^{\alpha(u_2)u_2}} h(X_{\tau^{\alpha(u_2)u_2}}^{\alpha(u_2)u_2}) \mathbb{1}_{\{T \ge \tau^{\alpha(u_2)u_2}\}} \right].$$

Proof. We only prove (3.1), since (3.2) follows similarly. Let $T \ge 0$. We define

$$\begin{split} W(x) &:= \inf_{\beta \in \Delta} \sup_{u_1 \in \mathcal{U}_1} \mathbb{E}_x \left[\int_0^{\tau^{u_1 \beta(u_1)} \wedge T} e^{-\delta s} \zeta \left(X_s^{u_1 \beta(u_1)} \right) \mathrm{d}s \right. \\ &+ e^{-\delta T} \underline{V} \left(X_T^{u_1 \beta(u_1)} \right) \mathbbm{1}_{\{T < \tau^{u_1 \beta(u_1)}\}} + e^{-\delta \tau^{u_1 \beta(u_1)}} h(X_{\tau^{u_1 \beta(u_1)}}^{u_1 \beta(u_1)}) \mathbbm{1}_{\{T \ge \tau^{u_1 \beta(u_1)}\}} \right] \end{split}$$

Let $\varepsilon > 0$. There exists a strategy $\beta_{\varepsilon} \in \Delta$ such that

$$W(x) \geq \mathbb{E}_{x} \left[\int_{0}^{\tau^{u_{1}\beta_{\varepsilon}(u_{1})} \wedge T} e^{-\delta s} \zeta \left(X_{s}^{u_{1}\beta_{\varepsilon}(u_{1})} \right) \mathrm{d}s + e^{-\delta T} \underline{V} \left(X_{T}^{u_{1}\beta_{\varepsilon}(u_{1})} \right) \mathbb{1}_{\{T < \tau^{u_{1}\beta_{\varepsilon}(u_{1})}\}} \right. \\ \left. + e^{-\delta \tau^{u_{1}\beta_{\varepsilon}(u_{1})}} h(X_{\tau^{u_{1}\beta_{\varepsilon}(u_{1})}}^{u_{1}\beta_{\varepsilon}(u_{1})}) \mathbb{1}_{\{T \geq \tau^{u_{1}\beta_{\varepsilon}(u_{1})}\}} \right] - \varepsilon, \quad (3.3)$$

for all $u_1 \in \mathcal{U}_1$.

Furthermore, for each $\hat{x} \in [a, b]$ there exists a $\beta_{\hat{x}} \in \Delta$ s.t.,

$$\underline{V}(\hat{x}) > \sup_{u_1 \in \mathcal{U}_1} J^{u_1 \beta_{\hat{x}}(u_1)}(\hat{x}) - \varepsilon.$$
(3.4)

Since <u>V</u> is continuous and for fixed u_1 and β , $J^{u_1\beta(u_1)}$ is continuous too (for fixed $t \mapsto u_1(t) = (u_t^1, u_t^2)$), we can choose a grid $a = x_0 < x_1 < \cdots < x_{n-1} < x_n = b$, s.t. for $u_1 \in \mathcal{U}_1$ and $\beta \in \Delta$,

$$|J^{u_1\beta(u_1)}(x_i) - J^{u_1\beta(u_1)}(x_{i-1})| < \varepsilon,$$
(3.5)

and

$$|\underline{V}(x_i) - \underline{V}(x_{i-1})| < \varepsilon, \tag{3.6}$$

for $i \in \{1, \ldots, n\}$. For $u_1 \in \mathcal{U}_1$ and $s \ge 0$, we define

$$\beta(u_1)(\omega)_s = \begin{cases} \beta_{\varepsilon}(u_1)(\omega)_s & s \le T, \\ \sum_{i=1}^n \mathbb{1}_{[x_{i-1}, x_i)}(X_T^{u_1 \beta_{\varepsilon}(u_1)}) \beta_{x_{i-1}}(\tilde{u}_1)(\tilde{\omega})_{s-T} & s > T, \end{cases}$$
(3.7)

where $\tilde{u}_1(t) = u_1(T+t)$, $\omega = (U_1(\omega), U_2(\omega), \dots)$ and $\tilde{\omega} = (U_{2m-1}(\omega), U_{2m}(\omega), \dots)$ s.t. $T_{m-1}(\omega) \leq T < T_m(\omega)$, according to (2.3). From (3.7) we get:

$$\begin{split} J^{u_1\beta(u_1)}(x) &= \mathbb{E}_x \left[\int_0^{\tau^{u_1\beta(u_1)}} e^{-\delta s} \zeta(X_s^{u_1\beta(u_1)}) \,\mathrm{d}s + e^{-\delta \tau^{u_1\beta(u_1)}} h(X_{\tau^{u_1\beta(u_1)}}^{u_1\beta(u_1)}) \right] \\ &= \mathbb{E}_x \left[\int_0^{\tau^{u_1\beta(u_1)}\wedge T} e^{-\delta s} \zeta(X_s^{u_1\beta(u_1)}) \,\mathrm{d}s + e^{-\delta \tau^{u_1\beta(u_1)}} h(X_{\tau^{u_1\beta(u_1)}}^{u_1\beta(u_1)}) \right. \\ &+ \sum_{i=1}^n \mathbb{1}_{[x_{i-1},x_i)} (X_T^{u_1\beta_\varepsilon(u_1)}) e^{-\delta T} \left(\int_T^{\tau^{u_1\beta(u_1)}} e^{-\delta(s-T)} \zeta(X_s^{u_1\beta(u_1)}) \,\mathrm{d}s \right) \mathbb{1}_{\{T < \tau^{u_1\beta(u_1)}\}} \right]. \end{split}$$

Using the tower property of conditional expectations yields,

$$\begin{split} J^{u_{1}\beta(u_{1})}(x) &= \mathbb{E}_{x} \Bigg[\int_{0}^{\tau^{u_{1}\beta_{\varepsilon}(u_{1})}\wedge T} e^{-\delta s} \zeta(X_{s}^{u_{1}\beta_{\varepsilon}(u_{1})}) \,\mathrm{d}s \\ &+ e^{-\delta \tau^{u_{1}\beta_{\varepsilon}(u_{1})}} h(X_{\tau^{u_{1}\beta_{\varepsilon}(u_{1})}}^{u_{1}\beta_{\varepsilon}(u_{1})}) \mathbb{1}_{\{T \geq \tau^{u_{1}\beta_{\varepsilon}(u_{1})}\}} \\ &+ e^{-\delta T} \sum_{i=1}^{n} \mathbb{1}_{[x_{i-1},x_{i})}(X_{T}^{u_{1}\beta_{\varepsilon}(u_{1})}) \mathbb{E}_{x} \left[\left(\int_{T}^{\tau^{u_{1}\beta(u_{1})}} e^{-\delta(s-T)} \zeta(X_{s}^{u_{1}\beta(u_{1})}) \,\mathrm{d}s \right. \\ &+ e^{-\delta(\tau^{u_{1}\beta(u_{1})}-T)} h\left(X_{\tau^{u_{1}\beta(u_{1})}}^{u_{1}\beta(u_{1})}\right) \right) |\mathcal{F}_{T}] \,\mathbb{1}_{\{T < \tau^{u_{1}\beta_{\varepsilon}(u_{1})}\}} \right] \\ &= \mathbb{E}_{x} \left[\int_{0}^{\tau^{u_{1}\beta_{\varepsilon}(u_{1})}\wedge T} e^{-\delta s} \zeta(X_{s}^{u_{1}\beta_{\varepsilon}(u_{1})}) \,\mathrm{d}s + e^{-\delta\tau^{u_{1}\beta_{\varepsilon}(u_{1})}} h(X_{\tau^{u_{1}\beta_{\varepsilon}(u_{1})}}^{u_{1}\beta_{\varepsilon}(u_{1})}) \mathbb{1}_{\{T \geq \tau^{u_{1}\beta_{\varepsilon}(u_{1})\}}\}} \\ &+ \sum_{i=1}^{n} \mathbb{1}_{[x_{i-1},x_{i})}(X_{T}^{u_{1}\beta_{\varepsilon}(u_{1})}) e^{-\delta T} J^{\tilde{u}_{1}\beta_{x_{i-1}}(\tilde{u}_{1})}(X_{T}^{u_{1}\beta_{\varepsilon}(u_{1})}) \mathbb{1}_{\{T < \tau^{u_{1}\beta_{\varepsilon}(u_{1})\}}\}} \right]. \end{split}$$

By (3.4) and (3.5), we get that for $X_T^{u_1\beta_{\varepsilon}(u_1)} \in [x_i, x_{i+1}),$

$$\underline{V}(x_i) \ge J^{\tilde{u}_1 \beta_{x_i}(\tilde{u}_1)}(x_i) - \varepsilon \ge J^{\tilde{u}_1 \beta_{x_i}(\tilde{u}_1)}(X_T^{u_1 \beta_{\varepsilon}(u_1)}) - 2\varepsilon.$$

Note that this holds, because given the definition and purpose of \tilde{u}_1 , the same control (specified for x and its path) is applied in both J functionals above. Therefore, and by using (3.6),

$$\begin{aligned} J^{u_1\beta(u_1)}(x) \\ &\leq \mathbb{E}_x \left[\int_0^{\tau^{u_1\beta_{\varepsilon}(u_1)}\wedge T} e^{-\delta s} \zeta(X_s^{u_1\beta_{\varepsilon}(u_1)}) \,\mathrm{d}s + e^{-\delta \tau^{u_1\beta_{\varepsilon}(u_1)}} h(X_{\tau^{u_1\beta_{\varepsilon}(u_1)}}^{u_1\beta_{\varepsilon}(u_1)}) \mathbb{1}_{\{T \geq \tau^{u_1\beta_{\varepsilon}(u_1)}\}} \right] \\ &+ \sum_{i=1}^n \mathbb{1}_{[x_{i-1},x_i)} (X_T^{u_1\beta_{\varepsilon}(u_1)}) e^{-\delta T} \underline{V}(x_{i-1}) \mathbb{1}_{\{T < \tau^{u_1\beta_{\varepsilon}(u_1)}\}} \right] + 2\varepsilon \\ &\leq \mathbb{E}_x \left[\int_0^{\tau^{u_1\beta_{\varepsilon}(u_1)}\wedge T} e^{-\delta s} \zeta(X_s^{u_1\beta_{\varepsilon}(u_1)}) \,\mathrm{d}s + e^{-\delta \tau^{u_1\beta_{\varepsilon}(u_1)}} h(X_{\tau^{u_1\beta_{\varepsilon}(u_1)}}^{u_1\beta_{\varepsilon}(u_1)}) \mathbb{1}_{\{T \geq \tau^{u_1\beta_{\varepsilon}(u_1)}\}} \right] \\ &+ \sum_{i=1}^n \mathbb{1}_{[x_{i-1},x_i)} (X_T^{u_1\beta_{\varepsilon}(u_1)}) e^{-\delta T} \underline{V}(X_T^{u_1\beta_{\varepsilon}(u_1)}) \mathbb{1}_{\{T < \tau^{u_1\beta_{\varepsilon}(u_1)}\}} \right] + 3\varepsilon. \end{aligned}$$

Together with (3.3),

$$J^{u_1\beta(u_1)}(x) \le W(x) + 4\varepsilon,$$

and finally

$$\underline{V}(x) \le W(x) + 4\varepsilon. \tag{3.8}$$

For the other direction, we proceed in a similar way. We use that there exists some $\hat\beta_\varepsilon\in\Delta$ such that

$$\underline{V}(x) \ge \sup_{u_1 \in \mathcal{U}_1} \mathbb{E}_x \left[\int_0^{\tau^{u_1 \hat{\beta}_{\varepsilon}(u_1)}} e^{-\delta t} \zeta(X_t^{u_1 \hat{\beta}_{\varepsilon}(u_1)}) \, \mathrm{d}t + e^{-\delta \tau^{u_1 \hat{\beta}_{\varepsilon}(u_1)}} h(X_{\tau^{u_1 \hat{\beta}_{\varepsilon}(u_1)}}^{u_1 \hat{\beta}_{\varepsilon}(u_1)}) \right] - \varepsilon$$

$$(3.9)$$

For such a fixed $\hat{\beta}_{\varepsilon} \in \Delta$, there exists $u_{\varepsilon} \in \mathcal{U}_1$ such that

$$\begin{split} W(x) \leq \mathbb{E}_{x} \Biggl[\int_{0}^{\tau^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})} \wedge T} e^{-\delta s} \zeta \left(X_{s}^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})} \right) \mathrm{d}s + e^{-\delta T} \underline{V} \left(X_{T}^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})} \right) \mathbb{1}_{\{T < \tau^{u_{1}\hat{\beta}_{\varepsilon}(u_{1})}\}} \\ + e^{-\delta \tau^{u_{1}\hat{\beta}_{\varepsilon}(u_{1})}} h(X_{\tau^{u_{1}\hat{\beta}_{\varepsilon}(u_{1})}}^{u_{1}\hat{\beta}_{\varepsilon}(u_{\varepsilon})}) \mathbb{1}_{\{T \geq \tau^{u_{1}\hat{\beta}_{\varepsilon}(u_{1})}\}} \Biggr] + \varepsilon. \end{split}$$

$$(3.10)$$

Also, for each $\hat{x} \in [a, b]$, it holds that

$$\underline{V}(\hat{x}) \leq \sup_{u_1 \in \mathcal{U}_1} \mathbb{E}_{\hat{x}} \left[\int_0^{\tau^{u_1 \hat{\beta}_{\varepsilon}(u_1)}} e^{-\delta t} \zeta(X_t^{u_1 \hat{\beta}_{\varepsilon}(u_1)}) \, \mathrm{d}t + e^{-\delta \tau^{u_1 \hat{\beta}_{\varepsilon}(u_1)}} h\left(X_{\tau^{u_1 \hat{\beta}_{\varepsilon}(u_1)}}^{u_1 \hat{\beta}_{\varepsilon}(u_1)}\right) \right],$$

and therefore there exists $u_{\hat{x}} \in \mathcal{U}_1$ such that

$$\underline{V}(\hat{x}) \le J^{u_{\hat{x}}\hat{\beta}_{\varepsilon}(u_{\hat{x}})}(\hat{x}) + \varepsilon.$$
(3.11)

We define $u_1 \in \mathcal{U}_1$ by using the same grid as before, i.e., for $a = x_0 < x_1 < \cdots < x_{n-1} < x_n = b$:

$$u_t^1 = \begin{cases} u_t^{\varepsilon} & t \leq T, \\ \sum_{i=1}^n \mathbb{1}_{[x_{i-1}, x_i)} (X_T^{u_{\varepsilon} \hat{\beta}_{\varepsilon}(u_{\varepsilon})}) & u_{t-T}^{x_{i-1}} & t > T. \end{cases}$$

For $X_T^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})} \in [x_i, x_{i+1})$, we get by (3.5), (3.11) and (3.6),

$$J^{u_{x_i}\hat{\beta}_{\varepsilon}(u_{x_i})}(X_T^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})}) \ge J^{u_{x_i}\hat{\beta}_{\varepsilon}(u_{x_i})}(x_i) - \varepsilon \ge \underline{V}(x_i) - 2\varepsilon \ge \underline{V}(X_T^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})}) - 3\varepsilon.$$
(3.12)

By similar computations as before, together with (3.12) and (3.10),

$$\begin{split} J^{u_{1}\beta_{\varepsilon}(u_{1})}(x) \\ &= \mathbb{E}_{x} \left[\int_{0}^{\tau^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})} \wedge T} e^{-\delta s} \zeta(X_{s}^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})}) \,\mathrm{d}s + e^{-\delta \tau^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})}} h(X_{\tau^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})}}^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})}) \mathbb{1}_{\{T \geq \tau^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})}\}} \right] \\ &+ \sum_{i=1}^{n} \mathbb{1}_{[x_{i-1},x_{i})} (X_{T}^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})}) e^{-\delta T} J^{u_{x_{i-1}}\hat{\beta}_{\varepsilon}(u_{x_{i-1}})} (X_{T}^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})}) \mathbb{1}_{\{T < \tau^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})}\}} \right] \\ &\geq \mathbb{E}_{x} \left[\int_{0}^{\tau^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})} \wedge T} e^{-\delta s} \zeta(X_{s}^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})}) \,\mathrm{d}s + e^{-\delta \tau^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})}} h(X_{\tau^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})}}^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})}) \mathbb{1}_{\{T \geq \tau^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})}\}} \right] \\ &+ \sum_{i=1}^{n} \mathbb{1}_{[x_{i-1},x_{i})} (X_{T}^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})}) e^{-\delta T} \underline{V}(X_{T}^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})}) \mathbb{1}_{\{T < \tau^{u_{\varepsilon}\hat{\beta}_{\varepsilon}(u_{\varepsilon})}\}} \right] - 3\varepsilon \\ &\geq W(x) - 4\varepsilon. \end{split}$$

Finally, by (3.9),

$$W(x) \le \underline{V}(x) + 5\varepsilon. \tag{3.13}$$

Hence, by (3.8) and (3.13),

$$\underline{V}(x) = W(x).$$

From this result for a deterministic T one can derive the statement for bounded stopping times, using the particular form of $\{\mathcal{F}_t\}_{t\geq 0}$ stopping times (Davis, 1993, p. 261) and the standard procedure.

4 Characterization of the value functions

Since for the considered problem is not likely to allow for an explicit solution, we will have to rely on a numerical procedure. That is why we focus on existence and uniqueness of solutions to particular integro-differential equations.

4.1 Viscosity solutions to Bellmann-Isaacs equations

In this section we will show that \underline{V} and \overline{V} are viscosity solutions to their associated Bellman-Isaacs equations. For this purpose we define for v a smooth function, $u_1 \in A_1$ and $u_2 \in A_2$, the following:

$$\begin{split} \mathcal{L}(x,v(x),v'(x),v,u_1,u_2) &:= (c_1(u_1) - c_2(u_2)) \, v'(x) \\ &+ \lambda_1 \int_0^{\rho(x-a,u_1)} v(x-r(y,u_1)) \, \mathrm{d}F_Y^1(y) + \lambda_2 \int_0^{\rho(b-x,u_2)} v(x+r(y,u_2)) \, \mathrm{d}F_Y^2(y) \\ &+ \lambda_1 \int_{\rho(x-a,u_1)}^\infty h(x-r(y,u_1)) \, \mathrm{d}F_Y^1(y) + \lambda_2 \int_{\rho(b-x,u_2)}^\infty h(x+r(y,u_2)) \, \mathrm{d}F_Y^2(y) \\ &- (\delta + \lambda_1 + \lambda_2) v(x) + \zeta(x), \ x \in [a,b], \end{split}$$

where $\rho(z, u) := \inf \left\{ y \in \mathbb{R}_0^+ : r(y, u) \ge z \right\}.$

Theorem 4.1. (i) \underline{V} is a viscosity solution to the lower Bellman-Isaacs equation

$$\sup_{u_1 \in A_1} \inf_{u_2 \in A_2} \mathcal{L}(x, v(x), v'(x), v, u_1, u_2) = 0,$$
(4.1)

for $x \in (a, b)$.

(ii) \overline{V} is a viscosity solution to the upper Bellman-Isaacs equation

$$\inf_{u_2 \in A_2} \sup_{u_1 \in A_1} \mathcal{L}(x, v(x), v'(x), v, u_1, u_2) = 0.$$
(4.2)

for $x \in (a, b)$.

Proof. We only prove (ii), since (i) follows in a similar way. We start by showing that \overline{V} is a supersolution. Let $x \in (a, b)$ and let φ be a smooth function on [a, b], such that $\varphi(x) = \overline{V}(x)$ and $\overline{V} - \varphi \ge 0$. We have to show that

$$\inf_{u_2 \in A_2} \sup_{u_1 \in A_1} \mathcal{L}(x, \varphi(x), \varphi'(x), \varphi, u_1, u_2) \le 0.$$

$$(4.3)$$

To do so, we assume that there exists some $\theta > 0$ such that

$$\inf_{u_2 \in A_2} \sup_{u_1 \in A_1} \mathcal{L}(x, \varphi(x), \varphi'(x), \varphi, u_1, u_2) \ge \theta > 0.$$
(4.4)

We define,

$$L(x, u_1, u_2) := \mathcal{L}(x, \varphi(x), \varphi'(x), \varphi, u_1, u_2).$$

By (4.4), we have that

$$\inf_{u_2 \in A_2} \sup_{u_1 \in A_1} L(x, u_1, u_2) \ge \theta.$$

Thus, for each $u_2 \in A_2$, there exists an $u_1 \in A_1$ depending on u_2 , such that

$$L(x, u_1, u_2) \ge \theta.$$

Note that L is uniformly continuous in u_1 and u_2 , since c_1, c_2 and $r(y, \cdot)$ are continuous functions on a compact set. Therefore,

$$L(x, u_1, \xi) \ge \frac{3\theta}{4},$$

for all $\xi \in B(u_2, r) \cap A_2$ and some $r = r(u_2) > 0$. Because A_2 is compact, there exist finitely many distinct points $a_1, \ldots, a_n \in A_1$, $b_1, \ldots, b_n \in A_2$ and $r_1, \ldots, r_n > 0$ such that

$$A_2 \subset \bigcup_{i=1}^n B(b_i, r_i)$$

and

$$L(x, a_i, \xi) \ge \frac{3\theta}{4} \tag{4.5}$$

for $\xi \in B(b_i, r_i) \cap A_2$.

We define a function $f: A_2 \to A_1$ by setting

$$f(b) = \sum_{k=1}^{n} a_k \mathbb{1}_{\{b \in B(b_k, r_k) \setminus \bigcup_{i=1}^{k-1} B(b_i, r_i)\}}.$$

Defining

$$\alpha^*(u_2)(\omega)_t = f(u_2(\omega)_t),$$

leads to an admissible strategy $\alpha^* \in \Gamma$.

Since L is also uniformly continuous in x, there exists some h > 0 such that for \hat{x} with $|x - \hat{x}| < h$,

$$L(\hat{x}, f(u_2), u_2) \ge \frac{\theta}{2},$$

for all $u_2 \in A_2$.

Furthermore, there exists some $t^* > 0$ such that $\sup_{u_1 \in A_1, u_2 \in A_2} |(c_1(u_1) - c_2(u_2))|t^* \leq h$ and $x \pm \sup_{u_1 \in A_1, u_2 \in A_2} |(c_1(u_1) - c_2(u_2))|t^* \in (a, b)$. We set $T := T_1 \wedge t^*$ and $\varphi(x) = 0$ for $x \notin [a, b]$. By definition of φ and Theorem 3.1, it holds that

$$\begin{split} \varphi(x) &= \overline{V}(x) \\ &= \sup_{\alpha \in \Gamma} \inf_{u_2 \in \mathcal{U}_2} \mathbb{E}_x \left[\int_0^{\tau^{\alpha(u_2)u_2} \wedge T} e^{-\delta s} \zeta \left(X_s^{\alpha(u_2)u_2} \right) \mathrm{d}s + e^{-\delta T} \overline{V} \left(X_T^{\alpha(u_2)u_2} \right) \mathbb{1}_{\{T < \tau^{\alpha(u_2)u_2}\}} \right] \\ &\quad + e^{-\delta \tau^{\alpha(u_2)u_2}} h(X_{\tau^{\alpha(u_2)u_2}}^{\alpha(u_2)u_2}) \mathbb{1}_{\{T \ge \tau^{\alpha(u_2)u_2}\}} \right] \\ &\geq \inf_{u_2 \in \mathcal{U}_2} \mathbb{E}_x \left[\int_0^{\tau^{\alpha^*(u_2)u_2} \wedge T} e^{-\delta s} \zeta \left(X_s^{\alpha^*(u_2)u_2} \right) \mathrm{d}s + e^{-\delta T} \varphi \left(X_T^{\alpha^*(u_2)u_2} \right) \\ &\quad + e^{-\delta \tau^{\alpha^*(u_2)u_2}} h(X_{\tau^{\alpha^*(u_2)u_2}}^{\alpha^*(u_2)u_2}) \mathbb{1}_{\{T \ge \tau^{\alpha^*(u_2)u_2}\}} \right]. \end{split}$$

For some $\varepsilon > 0$, there exists $u_2^{\varepsilon} \in \mathcal{U}_2$ such that

$$\begin{split} \varphi(x) &\geq \mathbb{E}_x \left[\int_0^{\tau^{\alpha^*(u_2^{\varepsilon})u_2^{\varepsilon}} \wedge T} e^{-\delta s} \zeta \left(X_s^{\alpha^*(u_2^{\varepsilon})u_2^{\varepsilon}} \right) \mathrm{d}s + e^{-\delta T} \varphi \left(X_T^{\alpha^*(u_2^{\varepsilon})u_2^{\varepsilon}} \right) \\ &+ e^{-\delta \tau^{\alpha^*(u_2^{\varepsilon})u_2^{\varepsilon}}} h(X_{\tau^{\alpha^*(u_2^{\varepsilon})u_2^{\varepsilon}}}^{\alpha^*(u_2^{\varepsilon})u_2^{\varepsilon}}) \mathbbm{1}_{\{T \geq \tau^{\alpha^*(u_2^{\varepsilon})u_2^{\varepsilon}}\}} \right] - \varepsilon. \end{split}$$

Then, the Dynkin formula yields

$$\mathbb{E}_{x}\left[e^{-\delta T}\varphi(X_{T}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}})\right]$$
$$=\varphi(x)+\mathbb{E}_{x}\left[\int_{0}^{T}e^{-\delta s}\mathcal{A}\varphi(X_{s}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}})-\delta e^{-\delta s}\varphi(X_{s}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}})\,\mathrm{d}s\right].$$

Hence, we get that

$$\varepsilon \geq \mathbb{E}_{x} \left[\int_{0}^{\tau^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}\wedge T}} e^{-\delta s} \zeta \left(X_{s}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}} \right) \mathrm{d}s + \int_{0}^{T} e^{-\delta s} \mathcal{A}\varphi(X_{s}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}) - \delta e^{-\delta s}\varphi(X_{s}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}) \mathrm{d}s + e^{-\delta \tau^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}} h(X_{\tau^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}) \mathbb{1}_{\{T \geq \tau^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}\}} \right].$$

$$(4.6)$$

Further, note that

$$\mathbb{P}\left[T_1 \le t | T_1^i \le T_1^j\right] = \mathbb{P}\left[T_1 \le t\right], \quad i, j \in \{1, 2\}.$$

Therefore,

$$\begin{split} & \mathbb{E}_{x} \left[e^{-\delta \tau^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}} h(X_{\tau^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}) \mathbb{1}_{\{T \geq \tau^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}\}} \right] \\ &= \mathbb{E}_{x} \left[e^{-\delta T_{1}} h(X_{T_{1}}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}) \mathbb{1}_{\{\tau^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}=T_{1}\}} \mathbb{1}_{\{T_{1} \leq t^{*}\}} \right] \\ &= \mathbb{E}_{x} \left[e^{-\delta T_{1}} \left(\int_{\rho(X_{T_{1-}}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}-a,\alpha^{*}(u_{2}^{\varepsilon}))} h(X_{T_{1-}}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}-r(y,\alpha^{*}(u_{2}^{\varepsilon})_{T_{1}})) \, \mathrm{d}F_{Y}^{1}(y) \right. \\ &+ \int_{\rho(b-X_{T_{1-}}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}},u_{2}^{\varepsilon})} h(X_{T_{1-}}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}+r(y,u_{2}^{\varepsilon}(T_{1})) \, \mathrm{d}F_{Y}^{2}(y) \right) \mathbb{1}_{\{T_{1} \leq t^{*}\}} \right] \\ &= \frac{\lambda_{1}}{\lambda} \mathbb{E}_{x} \left[e^{-\delta T_{1}} \int_{\rho(X_{T_{1-}}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}-a,\alpha^{*}(u_{2}^{\varepsilon}))} h(X_{T_{1-}}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}-r(y,\alpha^{*}(u_{2}^{\varepsilon})_{T_{1}})) \, \mathrm{d}F_{Y}^{1}(y) \mathbb{1}_{\{T_{1} \leq t^{*}\}} \right] \\ &+ \frac{\lambda_{2}}{\lambda} \mathbb{E}_{x} \left[e^{-\delta T_{1}} \int_{\rho(b-X_{T_{1-}}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}},u_{2}^{\varepsilon})} h(X_{T_{1-}}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}+r(y,u_{2}^{\varepsilon}(T_{1})) \, \mathrm{d}F_{Y}^{2}(y) \mathbb{1}_{\{T_{1} \leq t^{*}\}} \right]. \end{split}$$

By compensation,

$$\begin{split} & \mathbb{E}_{x} \left[e^{-\delta T_{1}} \int_{\rho(X_{T_{1-}}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}} - a, \alpha^{*}(u_{2}^{\varepsilon}))} h(X_{T_{1-}}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}} - r(y, \alpha^{*}(u_{2}^{\varepsilon})_{T_{1}})) \, \mathrm{d}F_{Y}^{1}(y) \mathbb{1}_{\{T_{1} \leq t^{*}\}} \right] \\ &= \mathbb{E}_{x} \left[\int_{0}^{T_{1}} e^{-\delta t} \int_{\rho(X_{t-}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}} - a, \alpha^{*}(u_{2}^{\varepsilon}))} h(X_{t-}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}} - r(y, \alpha^{*}(u_{2}^{\varepsilon})_{t})) \, \mathrm{d}F_{Y}^{1}(y) \, \mathrm{d}N_{t} \mathbb{1}_{\{T_{1} \leq t^{*}\}} \right] \\ &= \mathbb{E}_{x} \left[\int_{0}^{T} e^{-\delta t} \int_{\rho(X_{t-}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}} - a, \alpha^{*}(u_{2}^{\varepsilon}))} h(X_{t-}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}} - r(y, \alpha^{*}(u_{2}^{\varepsilon})_{t})) \, \mathrm{d}F_{Y}^{1}(y) \, \mathrm{d}N_{t} \right] \\ &= \lambda \, \mathbb{E}_{x} \left[\int_{0}^{T} e^{-\delta t} \int_{\rho(X_{t-}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}} - a, \alpha^{*}(u_{2}^{\varepsilon}))} h(X_{t-}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}} - r(y, \alpha^{*}(u_{2}^{\varepsilon})_{t})) \, \mathrm{d}F_{Y}^{1}(y) \, \mathrm{d}I_{t} \right] . \end{split}$$

Similar calculations for the second summand and plugging this into (4.6), leads to

$$\begin{split} \varepsilon \geq \mathbb{E}_x \Bigg[\int_0^{\tau^{\alpha^*(u_2^\varepsilon)u_2^\varepsilon} \wedge T} e^{-\delta s} \zeta \left(X_s^{\alpha^*(u_2^\varepsilon)u_2^\varepsilon} \right) \mathrm{d}s \\ &+ \int_0^T e^{-\delta s} \mathcal{A}\varphi (X_s^{\alpha^*(u_2^\varepsilon)u_2^\varepsilon}) - \delta e^{-\delta s} \varphi (X_s^{\alpha^*(u_2^\varepsilon)u_2^\varepsilon}) \mathrm{d}s \\ &+ \lambda_1 \int_0^T e^{-\delta t} \int_{\rho(X_{t-}^{\alpha^*(u_2^\varepsilon)u_2^\varepsilon} - a, \alpha^*(u_2^\varepsilon))}^{\infty} h(X_{t-}^{\alpha^*(u_2^\varepsilon)u_2^\varepsilon} - r(y, \alpha^*(u_2^\varepsilon)_t)) \mathrm{d}F_Y^1(y) \mathrm{d}t \\ &+ \lambda_2 \int_0^T e^{-\delta t} \int_{\rho(b-X_{t-}^{\alpha^*(u_2^\varepsilon)u_2^\varepsilon}, u_2^\varepsilon)}^{\infty} h(X_{t-}^{\alpha^*(u_2^\varepsilon)u_2^\varepsilon} + r(y, u_2^\varepsilon(t)) \mathrm{d}F_Y^2(y) \mathrm{d}t \Bigg]. \end{split}$$

On the other hand, the last expectation equals,

$$\begin{split} \mathbb{E}_{x} \left[\int_{0}^{\tau^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}\wedge T}} e^{-\delta s} L(X_{s}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}, \alpha^{*}(u_{2}^{\varepsilon}), u_{2}^{\varepsilon}) \,\mathrm{d}s \right] \\ &= \mathbb{P}(T_{1} > t^{*}) \mathbb{E}_{x} \left[\int_{0}^{t^{*}} e^{-\delta s} L(X_{s}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}, \alpha^{*}(u_{2}^{\varepsilon}), u_{2}^{\varepsilon}) \,\mathrm{d}s \left| T_{1} > t^{*} \right] \\ &+ \mathbb{P}(T_{1} \le t^{*}) \mathbb{E}_{x} \left[\int_{0}^{T_{1}} e^{-\delta s} L(X_{s}^{\alpha^{*}(u_{2}^{\varepsilon})u_{2}^{\varepsilon}}, \alpha^{*}(u_{2}^{\varepsilon}), u_{2}^{\varepsilon}) \,\mathrm{d}s \left| T_{1} \le t^{*} \right] \\ &\geq \frac{\theta}{2\delta} \left(e^{-\lambda t^{*}} (1 - e^{-\delta t^{*}}) + \frac{1}{\delta + \lambda} \left(\delta + e^{-(\delta + \lambda)t^{*}} \lambda - e^{-\lambda t^{*}} (\delta + \lambda) \right) \right). \end{split}$$

If we now choose ε strictly smaller than the last line, we arrive at a contradiction, therefore (4.3) holds.

To prove that \overline{V} is a subsolution, we have to show that for some smooth function ψ on [a, b] with $\psi(x) = \overline{V}(x)$ and $\overline{V} - \psi \leq 0$,

$$\inf_{u_2 \in A_2} \sup_{u_1 \in A_1} \mathcal{L}(x, \psi(x), \psi'(x), \psi, u_1, u_2) \ge 0.$$
(4.7)

Suppose, there exists some $\eta > 0$ such that

$$\inf_{u_2 \in A_2} \sup_{u_1 \in A_1} \mathcal{L}(x, \psi(x), \psi'(x), \psi, u_1, u_2) \le -\eta < 0.$$
(4.8)

We define

$$K(x, u_1, u_2) := \mathcal{L}(x, \psi(x), \psi'(x), \psi, u_1, u_2)$$

Since A_2 is compact, we have by (4.8) that there exists some $\tilde{u}_2 \in A_2$ such that

$$\sup_{u_1} K(x, u_1, \tilde{u}_2) \le -\eta.$$

Since K is again uniformly continuous in x, there exists some $\tilde{h} > 0$ such that for \hat{x} with $|x - \hat{x}| < \tilde{h}$,

$$K(\hat{x}, \alpha(\tilde{u}_2), \tilde{u}_2) \le -\frac{\eta}{2},$$

for all $\alpha \in \Gamma$. Note that the constant control \tilde{u}_2 , given by $\tilde{u}_t^2 \equiv \tilde{u}_2$ is an admissible control, i.e., $\tilde{u}_2 \in \mathcal{U}_2$. Again, there exists some $\tilde{t} > 0$ such that $\sup_{u_1 \in A_1, u_2 \in A_2} |(c_1(u_1) - c_2(u_2))| \tilde{t} \leq \tilde{h}$ and $x \pm \sup_{u_1 \in A_1, u_2 \in A_2} |(c_1(u_1) - c_2(u_2))| \tilde{t} \in (a, b)$. We set $\tilde{T} := T_1 \wedge \tilde{t}$ and $\psi(x) = 0$ for $x \notin [a, b]$. Let $\varepsilon > 0$. Again by Theorem 3.1, the definition of ψ and the existence of some ε -optimal strategy α_{ε} , we get by similar calculations as before, that

$$-\varepsilon \leq \mathbb{E}_{x} \left[\int_{0}^{\tau^{\alpha_{\varepsilon}(\tilde{u}_{2})\tilde{u}_{2}} \wedge \tilde{T}} e^{-\delta s} \zeta \left(X_{s}^{\alpha_{\varepsilon}(\tilde{u}_{2})\tilde{u}_{2}} \right) \mathrm{d}s \right. \\ \left. + \int_{0}^{\tilde{T}} e^{-\delta s} \mathcal{A}\varphi(X_{s}^{\alpha_{\varepsilon}(\tilde{u}_{2})\tilde{u}_{2}}) - \delta e^{-\delta s}\varphi(X_{s}^{\alpha_{\varepsilon}(\tilde{u}_{2})\tilde{u}_{2}}) \mathrm{d}s \right. \\ \left. + \lambda_{1} \int_{0}^{\tilde{T}} e^{-\delta t} \int_{\rho(X_{t-}-a,\alpha_{\varepsilon}(\tilde{u}_{2}))}^{\infty} h(X_{t-} - r(y,\alpha_{\varepsilon}(\tilde{u}_{2})_{t})) \mathrm{d}F_{Y}^{1}(y) \mathrm{d}t \right. \\ \left. + \lambda_{2} \int_{0}^{\tilde{T}} e^{-\delta t} \int_{\rho(b-X_{t-},\tilde{u}_{2})}^{\infty} h(X_{t-} + r(y,\tilde{u}_{2}) \mathrm{d}F_{Y}^{2}(y) \mathrm{d}t \right].$$
(4.9)

As before the last expectation is,

$$\mathbb{E}_{x}\left[\int_{0}^{\tau^{\alpha_{\varepsilon}(\tilde{u}_{2})\tilde{u}_{2}}\wedge\tilde{T}}e^{-\delta s}K(X_{s}^{\alpha_{\varepsilon}(\tilde{u}_{2})\tilde{u}_{2}},\alpha_{\varepsilon}(\tilde{u}_{2}),\tilde{u}_{2})\,\mathrm{d}s\right]$$
$$\leq -\frac{\eta}{2\delta}\left(e^{-\lambda\tilde{t}}(1-e^{-\delta\tilde{t}})+\frac{1}{\delta+\lambda}\left(\delta+e^{-(\delta+\lambda)\tilde{t}}\lambda-e^{-\lambda\tilde{t}}(\delta+\lambda)\right)\right).$$

By choosing $\varepsilon < \frac{\eta}{2\delta} \left(e^{-\lambda \tilde{t}} (1 - e^{-\delta \tilde{t}}) + \frac{1}{\delta + \lambda} \left(\delta + e^{-(\delta + \lambda)\tilde{t}} \lambda - e^{-\lambda \tilde{t}} (\delta + \lambda) \right) \right)$ we obtain another contradiction. \Box

4.2 Uniqueness of solution to Bellman-Isaacs equation

Since \underline{V} and \overline{V} fulfill (3.1) and (3.2), it seems reasonable to use a fixed point argument to show uniqueness of the solution. Unfortunately, a direct link between a fixed point and (viscosity-)solutions can only be achieved if we have absolute continuity and the existence of Markovian optimal controls. Therefore, we are going to show a classical comparison theorem for the set of Bellman-Isaacs equations. The proof follows an idea of Azcue and Muler (2014).

Theorem 4.2. Let $\underline{v} : [a, b] \to \mathbb{R}$ and $\overline{v} : [a, b] \to \mathbb{R}$ be continuous and bounded viscosity sub- and supersolutions to (4.2) (or (4.1)). If it holds that $\underline{v}(a) \leq \overline{v}(a)$ and $\underline{v}(b) \leq \overline{v}(b)$, then $\underline{v}(x) \leq \overline{v}(x)$ for all $x \in [a, b]$.

Proof. Assume that there exists some $x_0 \in (a, b)$ where $\underline{v}(x_0) > \overline{v}(x_0)$. Then,

$$0 < \max_{x \in [a,b]} (\underline{v}(x) - \overline{v}(x)) = \widetilde{M}, \tag{4.10}$$

with maximizing argument x^* . We define for $\nu > 0$ and $(x, y) \in [a, b]^2$,

$$G_{\nu}(x,y) := \underline{v}(x) - \overline{v}(x) - g^{\nu}(x,y),$$

where

$$g^{\nu}(x,y) := \nu(x-y)^2,$$

and set

$$M_{\nu} = \max_{(x,y)\in[a,b]^2} G_{\nu}(x,y),$$

again with with maximizer (x_{ν}, y_{ν}) . We see, that

$$M_{\nu} \ge G_v(x^*, x^*) = \widetilde{M},$$

and therefore,

$$\liminf_{\nu \to \infty} M_{\nu} \ge \widetilde{M} > 0.$$
(4.11)

To exploit the differentiability of g^{ν} , we first need to show that (x_{ν}, y_{ν}) is not an element of the boundary of $[a, b]^2$, i.e., $(x_{\nu}, y_{\nu}) \in (a, b)^2$. Indeed, we immediately see

$$G_{\nu}(a,a) \le 0, \quad G_{\nu}(b,b) \le 0,$$

but we know that $M_{\nu} > 0$ from (4.11). Furthermore, for $(x, y) \in \partial [a, b]^2$ with $|x - y| \ge \sqrt{\varepsilon} > 0$ - the edges of the rectangle and the vertices $\{(a, b), (b, a)\},\$

$$G_{\nu}(x,y) = \underline{v}(x) - \overline{v}(y) - \nu(x-y)^2 \le 2M - \nu\varepsilon < 0,$$

for $\nu > 2M/\varepsilon$, where M is a bound of both \overline{v} and \underline{v} . Therefore, $(x_{\nu}, y_{\nu}) \notin \partial [a, b]^2$ for ν large enough.

Since G_{ν} attains a maximum at (x_{ν}, y_{ν}) , we have that for any $x \in [a, b]$, $G_{\nu}(x, y_{\nu}) \leq G_{\nu}(x_{\nu}, y_{\nu})$ which translates to

$$\underline{v}(x) - \underline{v}(x_{\nu}) \le g^{\nu}(x, y_{\nu}) - g^{\nu}(x_{\nu}, y_{\nu}).$$

Taylor's theorem then yields

$$\limsup_{x \to x_{\nu}} \frac{\underline{v}(x) - \underline{v}(x_{\nu}) - g_x^{\nu}(x_{\nu}, y_{\nu})(x - x_{\nu}) - o(|x - x_{\nu}|)}{|x - x_{\nu}|} \le 0$$

This implies that $g_x^{\nu}(x_{\nu}, y_{\nu}) \in D^+(\underline{v})(x_{\nu})$, which is the set of all super-differentials of \underline{v} in x_{ν} .

Equivalently, for any $y \in [a, b]$ we have $G_{\nu}(x_{\nu}, y) \leq G_{\nu}(x_{\nu}, y_{\nu})$ and

$$\overline{v}(y) - \overline{v}(y_{\nu}) \ge g^{\nu}(x_{\nu}, y_{\nu}) - g^{\nu}(x_{\nu}, y).$$

Again this leads to

$$\liminf_{y \to y_{\nu}} \frac{\overline{v}(y) - \overline{v}(y_{\nu}) - \left(-g_{y}^{\nu}(x_{\nu}, y_{\nu})(y - y_{\nu})\right) + o(|y - y_{\nu}|)}{|y - y_{\nu}|} \ge 0.$$

Thus, $-g_y^{\nu}(x_{\nu}, y_{\nu}) \in D^-(\overline{v})(y_{\nu})$, which is the set of all sub-differentials of \overline{v} in y_{ν} . Since \underline{v} and \overline{v} are sub- and supersolution of (4.2), the following holds:

$$\inf_{u_2 \in A_2} \sup_{u_1 \in A_1} \mathcal{L}(x_{\nu}, \underline{v}(x_{\nu}), g_x^{\nu}(x_{\nu}, y_{\nu}), \underline{v}(x_{\nu}), u_1, u_2) \ge 0,$$
(4.12)

$$\inf_{u_2 \in A_2} \sup_{u_1 \in A_1} \mathcal{L}(y_{\nu}, \overline{v}(y_{\nu}), -g_y^{\nu}(x_{\nu}, y_{\nu}), \overline{v}(y_{\nu}), u_1, u_2) \le 0.$$
(4.13)

By compactness, there exist $\underline{u}_1, \underline{u}_2$ and $\overline{u}_1, \overline{u}_2$ corresponding to optimizers in (4.12) and (4.13) respectively. Additionally, note that $g_x^{\nu}(x_{\nu}, y_{\nu}) = -g_y^{\nu}(x_{\nu}, y_{\nu})$.

Now, we subtract (4.12) from (4.13), choose $u_1 = \underline{u}_1$ and $u_2 = \overline{u}_2$ and get

$$\begin{aligned} &(\delta + \lambda_1 + \lambda_2) \left(\underline{v}(x_{\nu}) - \overline{v}(y_{\nu}) \right) \leq \zeta(x_{\nu}) - \zeta(y_{\nu}) \\ &+ \lambda_1 \left(\int_0^{\rho(x_{\nu} - a, \underline{u}_1)} \underline{v}(x_{\nu} - r(y, \underline{u}_1)) \, \mathrm{d}F_Y^1(y) - \int_0^{\rho(y_{\nu} - a, \underline{u}_1)} \overline{v}(y_{\nu} - r(y, \underline{u}_1)) \, \mathrm{d}F_Y^1(y) \right) \\ &+ \lambda_2 \left(\int_0^{\rho(b - x_{\nu}, \overline{u}_2)} \underline{v}(x_{\nu} + r(y, \overline{u}_2)) \, \mathrm{d}F_Y^2(y) - \int_0^{\rho(b - y_{\nu}, \overline{u}_2)} \overline{v}(y_{\nu} + r(y, \overline{u}_2)) \, \mathrm{d}F_Y^2(y) \right) \\ &+ \lambda_1 \left(\int_{\rho(x_{\nu} - a, \underline{u}_1)}^{\infty} h(x_{\nu} - r(y, \underline{u}_1)) \, \mathrm{d}F_Y^1(y) - \int_{\rho(y_{\nu} - a, \underline{u}_1)}^{\infty} h(y_{\nu} - r(y, \underline{u}_1)) \, \mathrm{d}F_Y^1(y) \right) \\ &+ \lambda_2 \left(\int_{\rho(b - x_{\nu}, \overline{u}_2)}^{\infty} h(x_{\nu} + r(y, \overline{u}_2)) \, \mathrm{d}F_Y^2(y) - \int_{\rho(b - y_{\nu}, \overline{u}_2)}^{\infty} h(y_{\nu} + r(y, \overline{u}_2)) \, \mathrm{d}F_Y^2(y) \right). \end{aligned}$$

$$(4.14)$$

To carry on, we need further properties of the maximizer (x_{ν}, y_{ν}) , especially for $\nu \to \infty$. At first we observe that

$$G_{\nu}(x_{\nu}, x_{\nu}) + G_{\nu}(y_{\nu}, y_{\nu}) \le 2G_{\nu}(x_{\nu}, y_{\nu})$$

which gives us the following estimate on $(x_{\nu} - y_{\nu})^2$:

$$\underline{v}(x_{\nu}) - \overline{v}(x_{\nu}) + \underline{v}(y_{\nu}) - \overline{v}(y_{\nu}) \le 2\left(\underline{u}(x_{\nu}) - \overline{v}(y_{\nu}) - \lambda(x_{\nu} - y_{\nu})^{2}\right)$$

$$\Leftrightarrow 2\nu(x_{\nu} - y_{\nu})^{2} \le \underline{v}(x_{\nu}) - \overline{v}(y_{\nu}) \le 2M$$

$$\Leftrightarrow (x_{\nu} - y_{\nu})^{2} \le \frac{M}{\nu}.$$
(4.15)

Using the compactness of $[a, b]^2$, we have existence of a sequence $(\nu_n)_{n \in \mathbb{N}}$ with $\nu_n \to \infty$, such that $\lim_{n\to\infty} (x_{\nu_n}, y_{\nu_n}) = (\overline{x}, \overline{y}) \in [a, b]^2$. From (4.15) we get $\overline{x} = \overline{y}$, together with (4.14) and (4.10), this gives us

$$\begin{split} (\delta + \lambda_1 + \lambda_2) \left(\underline{v}(\overline{x}) - \overline{v}(\overline{x}) \right)) \\ &\leq \lambda_1 \int_0^{\rho(\overline{x} - a, \underline{u}_1)} \underline{v}(\overline{x} - r(y, \underline{u}_1)) - \overline{v}(\overline{x} - r(y, \underline{u}_1)) \,\mathrm{d}F_Y^1(y) \\ &+ \lambda_2 \int_0^{\rho(b - \overline{x}, \overline{u}_2)} \underline{v}(\overline{x} + r(y, \overline{u}_2)) - \overline{v}(\overline{x} + r(y, \overline{u}_2)) \,\mathrm{d}F_Y^2(y) \leq (\lambda_1 + \lambda_2) \widetilde{M} \end{split}$$

Altogether, we arrive at the contradiction

$$\widetilde{M} \leq \liminf_{\nu \to \infty} M_{\nu} \leq \lim_{n \to \infty} M_{\nu_n} = \underline{v}(\overline{x}) - \overline{v}(\overline{x}) \leq \frac{\lambda_1 + \lambda_2}{\delta + \lambda_1 + \lambda_2} \widetilde{M}.$$

This proves the statement of the theorem.

Remark 4.1. Since the involved operators are non-local, the boundary values in a and b are not necessarily a-priori known, indeed they are part of the solution. Intuitively, this follows from controlling the distribution of the post-jump location and the sign of the drift close to the boundaries. \underline{V} and \overline{V} are both suband supersolutions to the respective equations, and therefore their uniqueness - as solutions to the Bellman-Isaacs equations - follows from the comparison theorem and the implicitly given boundary values.

5 Numerical examples

In the previous sections, we discussed that the value functions are viscosity solutions to Bellman-Isaacs equations. From the theory of viscosity solutions we know that if the functions \underline{V} and \overline{V} were continuously differentiable, they would be true solutions to these equations. Therefore, we are now trying to find numerical solutions and corresponding controls. Note that if we find controls u_1^* and u_2^* for which the upper and lower value coincide, a Nash equilibrium is immediately achieved.

For the numerical approach, we use a policy iteration algorithm: We divide the interval [a, b] into an equidistant grid $\{x_1, \ldots, x_n\}$, where $a = x_1 \le x_2 \le \cdots \le x_n = b$ and set $u_1^0 = u_2^0 = \overline{u}$ where \overline{u} corresponds to the constant control of no reinsurance. Then, we calculate $J^{u_1^0 u_2^0}$ by numerically solving

$$\mathcal{L}(x, J^{u_1^0 u_2^0}(x), (J^{u_1^0 u_2^0})'(x), J^{u_1^0 u_2^0}, u_1^0, u_2^0) = 0,$$
(5.1)

for every $x \in \{x_2, \ldots, x_{n-1}\}$. On the boundary, i.e., for $x_1 = a$ and $x_n = b$, we might have to either solve an equation or deal with a boundary value. We will discuss this later in a concrete example. Then, we update the controls one at a time, for example when starting with updating the control of player 2, we set for $x \in \{x_1, \ldots, x_n\}$,

$$u_{2}^{1}(x) = \operatorname*{arg\,min}_{u \in A_{2}} \mathcal{L}(x, J^{u_{1}^{0}u_{2}^{0}}(x), (J^{u_{1}^{0}u_{2}^{0}})'(x), J^{u_{1}^{0}u_{2}^{0}}, u_{1}^{0}, u).$$

Afterwards, we calculate $J^{u_1^0 u_2^1}$ like in (5.1) but with u_2^1 instead of u_2^0 . Then, we find u_1^1 by

$$u_1^1(x) = \operatorname*{arg\,max}_{u \in A_1} \mathcal{L}(x, J^{u_1^0 u_2^1}(x), (J^{u_1^0 u_2^1})'(x), J^{u_1^0 u_2^1}, u, u_2^1),$$

 $x \in \{x_1, \ldots, x_n\}$ and again solve the equation for $J^{u_1^1 u_2^1}$. We proceed in this way for u_2^2, u_1^2, \ldots until there are no more significant changes in the controls or value function. I.e., we fix a level $\varepsilon \approx 10^{-5}$ and stop the iteration if the maximal improvement on the grid is smaller than ε . Markov controls are inherently used by the nature of the schema. This procedure is also motivated by Fleming and Soner (2006), p. 379.

We consider the following (test-) functional:

$$J^{u_1 u_2}(x) = \mathbb{E}_x \left[e^{-\delta \tau^{u_1 u_2}} \mathbb{1}_{\{X^{u_1 u_2}_{\tau^{u_1 u_2}} \ge b\}} \right],$$
(5.2)

with corresponding value functions \underline{V} and \overline{V} . This results from setting $\zeta(x) = 0$ and $h(x) = \mathbb{1}_{\{x \ge b\}}$. If $\delta = 0$, this is just the probability that the process exits the interval [a, b] at the upper endpoint. It is intuitive that player one would aim to maximize this probability, as a larger probability corresponds to a greater expected surplus relative to player two. Conversely, player two wants to prevent the process from outgrowing b, therefore tries to find a minimizing strategy. Note that in this section, the terms "control" and "strategy" are used interchangeably.



Figure 1: Strategies of player 1 and 2 after 10 iterations. The controls of Player 1 and 2 correspond to the solid and dashed line respectively.

5.1 Proportional reinsurance, variance principle and exponential claims

We start by setting a = -4, b = 4 and $A = A_1 = A_2 = [0, 1]$. The risk share function $r : \mathbb{R}_0^+ \times A \to \mathbb{R}_0^+$ corresponding to proportional reinsurance is given by r(y, u) = yu. We assume that the claim heights are exponentially distributed with rate $1/\mu_i$, for $i \in \{1, 2\}$. For the first insurer's premium without acquiring reinsurance we apply the expectation principle, i.e., for $i \in \{1, 2\}$, $c_i = \lambda_i(1 + \eta_i)\mu_i$, with risk loading $\eta_i > 0$ and $\mu_i = \mathbb{E}(Y_1^i)$. For the reinsurer's premium, we apply the variance principle which leads us to the following premium function:

$$c_i(u) = c_i - \lambda_i(\mu_i(1-u) + \theta_i \mu_i^2(1-u)^2),$$

where $\theta_i > 0$ is the safety loading corresponding to the reinsurance premium. In Figure 1, we used the parameters $\lambda_1 = \lambda_2 = 1$, $\delta = 0.05$, $\mu_1 = 1$, $\mu_2 = 2$, $\eta_1 = 0.1$, $\eta_2 = 0.08$, $\theta_1 = \theta_2 = 0.12$. Here, we see that $c_2 = c_2(1) > c_1(u)$, for any $0 \le u \le 1$. Therefore, close to *a* player 2 will take the chance and produce a maximal negative drift, which results in the boundary value $\underline{V}(a) = \overline{V}(a) = 0$. In the middle of the interval, reinsurance is actively purchased, whereas close to *b* no reinsurance is taken in order to delay an upper exit. For player one, it is better not to acquire any reinsurance but to maximize the drift in order to exceed the upper bound of the interval as fast as possible. This also means that this player voluntarily accepts the full extent of potential claims.

5.2 Excess of loss reinsurance and expectation principle

In this framework, we set a = -2 and b = 2, as well as $A = A_1 = A_2 = [0, \infty]$. The risk share function for excess of loss reinsurance is given by r(y, M) = $\min(y, M)$. The claim amount which exceeds M is covered by the reinsurer. The first insurers premium is calculated like in Section 5.1.



Figure 2: Strategies of player 1 and 2 after 10 iterations with exponentially distributed claims. The solid line represents player one and the dashed player two. Technically, "no reinsurance" corresponds to an infinite value of the control, but for illustrative purposes it is shown in gray.



Figure 3: Strategies of player 1 and 2 after 10 iterations with Pareto II distributed claims. Again, the solid line shows the strategy of player one and the dashed one the strategy of player two, and the color gray stands for "no reinsurance".

5.2.1 Exponential claims

Let the claims be exponentially distributed with rate $1/\mu_i$, for $i \in \{1, 2\}$. The premium function is now calculated via the expectation principle, i.e.,

$$c_i(M) = c_i - \lambda_i (1 + \theta_i) \mu_i e^{-M/\mu_i}$$

With the same parameter values as in Section 5.1, we arrive at Figure 2. The plot suggests that in a certain range, the optimal control of player 2 is linear in x. Perhaps certain adjustments in the simulation procedure would lead to a straight line. However, in the interest of comparability and computational effort, the method remained unchanged across all examples - same discretization and termination criterion.

5.2.2 Pareto type II distributed claims

Now, we assume that the claims are Pareto II distributed with shape α_i and scale β_i . By applying the expectation principle, the premium function is given by

$$c_i(M) = c_i - \lambda_i (1+\theta_i) \left(\frac{\beta_i^{\alpha_i}}{\alpha_i - 1} (M+\beta_i)^{-\alpha_i + 1}\right).$$

In Figure 3, we chose $\alpha_1 = \alpha_2 = 3$, $\beta_1 = \beta_2 = 1$ and retain the remaining ones from Section 5.1. For these parameters, $\lim_{m\to\infty} c_1(m) > \lim_{m\to\infty} c_2(m)$, therefore $\underline{V}(b) = \overline{V}(b) = 1$.

Declarations

This research was funded in whole or in part by the Austrian Science Fund (FWF) [10.55776/P33317]. For open access purposes, the authors have applied a CC BY public copyright license to any author-accepted manuscript version arising from this submission.

References

- Søren Asmussen, Bent Jesper Christensen, and Julie Thøgersen. Nash equilibrium premium strategies for push-pull competition in a frictional nonlife insurance market. *Insurance Math. Econom.*, 87:92–100, 2019. doi: 10.1016/j.insmatheco.2019.02.002.
- Pablo Azcue and Nora Muler. Optimal reinsurance and dividend distribution policies in the Cramér-Lundberg model. *Math. Finance*, 15(2):261–308, 2005. doi: 10.1111/j.0960-1627.2005.00220.x.
- Pablo Azcue and Nora Muler. Stochastic optimization in insurance: A dynamic programming approach. SpringerBriefs in Quantitative Finance. Springer, New York, 2014.
- Rainer Buckdahn and Juan Li. Stochastic differential games and viscosity solutions of Hamilton-Jacobi-Bellman-Isaacs equations. SIAM J. Control Optim., 47(1):444–475, 2008. doi: 10.1137/060671954.

- Jingyi Cao, Dongchen Li, Virginia R. Young, and Bin Zou. Stackelberg differential game for insurance under model ambiguity. *Insurance Math. Econom.*, 106:128–145, 2022. doi: 10.1016/j.insmatheco.2022.06.003.
- Lv Chen and Yang Shen. On a new paradigm of optimal reinsurance: A stochastic stackelberg differential game between an insurer and a reinsurer. ASTIN Bulletin, 48(2):905–960, 2018. doi: 10.1017/asb.2018.3.
- Mark H. A. Davis. Piecewise-deterministic Markov processes: a general class of nondiffusion stochastic models. J. Roy. Statist. Soc. Ser. B, 46(3):353–388, 1984.
- Mark H. A. Davis. Markov models and optimization. Monographs on statistics and applied probability. Chapman & Hall, London, 1993.
- Julia Eisenberg and Hanspeter Schmidli. Minimising expected discounted capital injections by reinsurance in a classical risk model. Scand. Actuar. J., 2011 (3):155–176, 2011. doi: 10.1080/03461231003690747.
- M. Ferreira, D. Pinheiro, and S. Pinheiro. Two-player zero-sum stochastic differential games with random horizon. Adv. in Appl. Probab., 51(4):1209–1235, 2019. doi: 10.1017/apr.2019.47.
- Wendell H. Fleming and H. Mete Soner. Controlled Markov processes and viscosity solutions, volume 25 of Stochastic Modelling and Applied Probability. Springer, New York, 2nd edition, 2006.
- Wendell H. Fleming and Panagiotis E. Souganidis. On the existence of value functions of two-player, zero-sum stochastic differential games. *Indiana Univ. Math. J.*, 38(2):293–314, 1989. doi: 10.1512/iumj.1989.38.38015.
- Jean Jacod and Anatolii V. Skorokhod. Jumping Markov processes. Ann. Inst. H. Poincaré Probab. Statist., 32(1):11–67, 1996. ISSN 0246-0203.
- Dylan Possamaï, Nizar Touzi, and Jianfeng Zhang. Zero-sum path-dependent stochastic differential games in weak formulation. Ann. Appl. Probab., 30(3): 1415–1457, 2020. doi: 10.1214/19-AAP1533.
- Tomasz Rolski, Hanspeter Schmidli, Volker Schmidt, and Jozef Teugels. Stochastic processes for insurance and finance. Wiley Series in Probability and Statistics. John Wiley & Sons, Ltd., Chichester, 1999. doi: 10.1002/9780470317044.
- Hanspeter Schmidli. Optimal proportional reinsurance policies in a dynamic setting. Scand. Actuar. J., 2001(1):55–68, 2001. doi: 10.1080/ 034612301750077338.
- Hanspeter Schmidli. *Stochastic control in insurance*. Probability and its Applications. Springer-Verlag, London, 2008.
- Michael Taksar and Xudong Zeng. Optimal non-proportional reinsurance control and stochastic differential games. *Insurance Math. Econom.*, 48(1):64–71, 2011. doi: 10.1016/j.insmatheco.2010.09.006.