

Tilting objects in the extended heart of a t -structure

Alejandro Argudín-Monroy, Octavio Mendoza, Carlos E. Parra *

Abstract

Building on the recent work of Adachi, Enomoto and Tsukamoto on a generalization of the Happel-Reiten-Smalø tilting process, we study extended tilting objects in extriangulated categories with negative first extension. These objects coincide with the 1-tilting objects in abelian categories as in the work of Parra, Saorín and Virili. We will be particularly interested in the case where the extriangulated category in question is the heart $\mathcal{H}_{[t_1, t_2]}$ of an interval of t -structures $[t_1, t_2]$. Our main results consist of a characterization of the extended tilting objects of a heart $\mathcal{H}_{[t_1, t_2]}$ for the case when $t_2 \leq \Sigma^{-1}t_1$, and another one for the case when $\Sigma^{-2}t_1 < t_2$. In the first one, we give conditions for these tilting objects to coincide with the quasi-tilting objects of the abelian category $\mathcal{H}_{[t_1, \Sigma^{-1}t_1]}$. In the second one, it is given conditions for these to coincide with projective generators in the extriangulated category $\mathcal{H}_{[t_1, \Sigma t_2]}$.

1 Introduction

One of the fundamental tools to study a triangulated category is the notion of t -structure [8]. Among their properties, we can mention that any abelian category can be realized as the heart $\mathcal{H}_{\mathbf{x}} := \mathcal{X} \cap \mathcal{Y}$ of a t -structure $\mathbf{x} = (\mathcal{X}, \mathcal{Y})$ in a triangulated category. The Happel-Reiten-Smalø tilting (*right HRS-tilt*) process shows us how, starting from a t -structure $\mathbf{x} = (\mathcal{X}, \mathcal{Y})$, the torsion pairs of the heart $\mathcal{H}_{\mathbf{x}}$ parameterize a certain family of t -structures. Namely, they parameterize the t -structures $\mathbf{x}' = (\mathcal{X}', \mathcal{Y}')$ such that $\Sigma\mathcal{X} \subseteq \mathcal{X}' \subseteq \mathcal{X}$ [13]. Moreover, within the parameterized t -structures, we can highlight those whose heart is derived equivalent to $\mathcal{H}_{\mathbf{x}}$. These are the ones corresponding to the so-called *tilting torsion pairs*. It is of particular interest when such torsion pair can be chosen so that its heart has a projective generator. In which case, one has that the torsion class is of the form $\text{Gen}(V)$, where V is what is known as a *quasi-tilting object* [22, Proposition 3.8].

*The first named author was supported by a postdoctoral fellowship EPM(1) 2024 from SE-CIHTI. The first and second named authors were supported by the Project PAPIIT-IN100124 Universidad Nacional Autónoma de México. The third named author was supported by ANID+FONDECYT/REGULAR+1240253.

Recently, Adachi, Enomoto and Tsukamoto presented in [1] a generalization of the HRS-tilt process in the context of extriangulated categories with negative first extension, which can be referred as the *AET-tilt process*. To state the AET-tilt process in the context of a triangulated category \mathcal{D} , the following notions are presented. Given two t -structures $\mathbf{t}_1 = (\mathcal{X}_1, \mathcal{Y}_1)$ and $\mathbf{t}_2 = (\mathcal{X}_2, \mathcal{Y}_2)$ in \mathcal{D} such that $\mathbf{t}_1 \leq \mathbf{t}_2$ (that is $\mathcal{X}_1 \subseteq \mathcal{X}_2$), define the interval $[\mathbf{t}_1, \mathbf{t}_2]$ which is the class of all the t -structures $\mathbf{t}' = (\mathcal{X}', \mathcal{Y}')$ such that $\mathcal{X}_1 \subseteq \mathcal{X}' \subseteq \mathcal{X}_2$. The class $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} := \Sigma^{-1}\mathcal{Y}_1 \cap \mathcal{X}_2$ is known as the heart of the interval $[\mathbf{t}_1, \mathbf{t}_2]$. The AET-tilt process tells us that there exists a bijection between the t -structures in $[\mathbf{t}_1, \mathbf{t}_2]$ and the s -torsion pairs in the extriangulated category $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$.

The aim of this paper is to study the structure of the heart $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ as well as its tilting objects. To this end, we begin by pointing out that one can classify the hearts $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ into three different kinds. The first kind (*standard hearts*) consists of hearts $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ where $\mathbf{t}_1 = \Sigma \mathbf{t}_2$ (that is $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} = \mathcal{H}_{\mathbf{t}_2}$ which is an abelian category). The second kind (*restricted hearts*) consists of hearts $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ where $\Sigma \mathbf{t}_2 < \mathbf{t}_1 < \mathbf{t}_2$. Finally, the third kind (*extended hearts*) consists of hearts $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ where $\mathbf{t}_1 < \Sigma \mathbf{t}_2 < \mathbf{t}_2$.

Let us look with more precision at the content of the paper. In Section 2, we will look at some basic notions from the theory of extriangulated and triangulated categories. Specifically, it is recalled the notion of s -torsion pair in an extriangulated category (with negative first extension) and its relationship with t -structures (in case the extriangulated category in question is a triangulated one). Finally, it is stated the theorem of Adachi, Enomoto and Tsukamoto on the AET-tilt process.

The third section contains a deeper study on the structure of the heart $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ for an interval $[\mathbf{t}_1, \mathbf{t}_2]$ of s -torsion pairs in an extriangulated category. On the one hand, we will seek to study the s -torsion pairs of $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ as well as its closure properties. Specifically, we will see in Propositions 2.9 and 3.9 conditions for a s -torsion pair to be closed under extensions, (co)cones, direct summands, s -subobjects and s -quotients. On the other hand, we introduce the notion of *normal interval* (see Definition 3.5). We will see that, under certain conditions (see Theorem 3.13), if the interval $[\mathbf{t}_1, \mathbf{t}_2]$ is normal, then $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ is an exact and a semi-abelian category. If in addition, see Theorem 3.14, the ambient extriangulated category is triangulated, we have that the heart $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ is a quasi-abelian category. It is worth mentioning that similar results to the above ones have been proved (with different tools) by Rump, Schneiders, Tattar and Fiorot [26, 28, 29, 10]. By using the above, see Theorem 3.19, we will make use of the AET-tilt process to parameterize the torsion pairs in a quasi-abelian category.

In Section 4, we will apply the results obtained in Section 3 to study the t -structures of a triangulated category together with the extended hearts induced by them. In particular, we will review the right HRS-tilt and the left HRS-tilt processes (see Theorems 4.9 and 4.10).

Lastly, in Section 5, we introduce the notion of *extended tilting* object in an extriangulated category with negative first extension. This notion coincides with the definition of 1-tilting object in abelian categories introduced recently in [23].

We will see that the definition of extended tilting object in this more general context shares some properties with the abelian case. Specifically, we will show that an extended tilting object is \mathbb{E} -universal and that it also has projective dimension at most one, see Propositions 5.6 and 5.8. Having done so, we will seek to characterize extended tilting objects in restricted hearts and extended hearts. Namely, we will show that an object is extended tilting in a restricted heart $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ if and only if it is quasi-tilting in the standard heart $\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}$ (see Theorem 5.12). In contrast, given an interval $[\mathbf{t}_1, \mathbf{t}_2]$ with $\Sigma^{-2}\mathbf{t}_1 < \mathbf{t}_2$, an object V will be extended tilting in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ if and only if $\mathcal{H}_{[\mathbf{t}_1, \Sigma\mathbf{t}_2]}$ is cogenerating in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ and V is a projective generator in $\mathcal{H}_{[\mathbf{t}_1, \Sigma\mathbf{t}_2]}$. Using the above, we will exhibit an example of an extended tilting object in an extended heart that is a non-abelian category (see Example 5.17).

2 Preliminaries

Exact categories and triangulated categories are valuable tools of contemporary mathematics. Their main quality lies in the fact that they are suitable contexts for homological algebra. Recently, H. Nakaoka and Y. Palu presented in [18] the notion of extriangulated category: a concept that encompasses triangulated categories and exact categories. We will omit the precise definition of extriangulated category, see [18, Def. 2.12] for details, but we include below some essential notions for the convenience of the reader. To get an intuitive picture, the inexperienced reader may understand extriangulated categories as subcategories of triangulated categories which are closed under extensions. An **extriangulated category** consists of a triple $(\mathcal{D}, \mathbb{E}, \mathfrak{s})$, where \mathcal{D} is an additive category, $\mathbb{E} : \mathcal{D}^{op} \times \mathcal{D} \rightarrow \mathbf{Ab}$ is an additive bifunctor and \mathfrak{s} is a correspondence that associates an equivalence class $[A \xrightarrow{x} X \xrightarrow{y} B]$ to each element δ of $\mathbb{E}(B, A)$ satisfying a series of axioms as in [18, Def. 2.12]. To fix ideas, the reader can think of \mathcal{D} as an abelian category and $\mathbb{E} = \mathbf{Ext}_{\mathcal{D}}^1(-, -)$, or of \mathcal{D} as a triangulated category with $\mathbb{E} = \mathbf{Hom}_{\mathcal{D}}(-, -[1])$. Moreover, whenever we refer to an abelian category or a triangulated category as an extriangulated category, it will be by means of such a functor \mathbb{E} . Let $(\mathcal{D}, \mathbb{E}, \mathfrak{s})$ be an extriangulated category. By following [18], we have the following notions. For $A, B \in \mathcal{D}$, an element $\delta \in \mathbb{E}(B, A)$ is called an **\mathbb{E} -extension**, and the zero element $0 \in \mathbb{E}(B, A)$ is called the **split \mathbb{E} -extension**. If $\delta \in \mathbb{E}(B, A)$ and $\mathfrak{s}(\delta) = [A \xrightarrow{x} C \xrightarrow{y} B]$, we say that $A \xrightarrow{x} C \xrightarrow{y} B$ **realizes** δ or that it is an **\mathfrak{s} -conflation**, also known as an \mathbb{E} -triangle. In such case, we will use the notation $\delta : A \xrightarrow{x} C \xrightarrow{y} B$ or $A \xrightarrow{x} C \xrightarrow{y} B \xrightarrow{\delta}$. For every $\delta \in \mathbb{E}(B, A)$, $a \in \mathbf{Hom}_{\mathcal{D}}(A, A')$ and $b \in \mathbf{Hom}_{\mathcal{D}}(B', B)$, define $a \cdot \delta := \mathbb{E}(B, a)(\delta)$ and $\delta \cdot b := \mathbb{E}(b, A)(\delta)$. Observe that, each $\delta \in \mathbb{E}(B, A)$ induces the morphisms $\delta \cdot - : \mathbf{Hom}_{\mathcal{D}}(W, B) \rightarrow \mathbb{E}(W, A)$ and $- \cdot \delta : \mathbf{Hom}_{\mathcal{D}}(A, W) \rightarrow \mathbb{E}(B, W)$, for every $W \in \mathcal{D}$. Let $\delta \in \mathbb{E}(B, A)$ and $\delta' \in \mathbb{E}(B', A')$. A **morphism of extensions** $\delta \rightarrow \delta'$ is a pair $(a, b) \in \mathbf{Hom}_{\mathcal{D}}(A, A') \times \mathbf{Hom}_{\mathcal{D}}(B, B')$ such that $a \cdot \delta = \delta' \cdot b$. According to the definition of extriangulated category, we have that, for any morphism of \mathbb{E} -extensions $(a, b) : \delta \rightarrow \delta'$, with $\mathfrak{s}(\delta) = [A \xrightarrow{x} C \xrightarrow{y} B]$ and

$\mathfrak{s}(\delta') = [A' \xrightarrow{x'} C' \xrightarrow{y'} B']$, there exists a morphism $c : C \rightarrow C'$ such that $cx = x'a$ and $y'c = by$. In this case we say that (a, c, b) **realizes** the morphism $(a, b) : \delta \rightarrow \delta'$. Let $(\mathcal{D}, \mathbb{E}, \mathfrak{s})$ be an extriangulated category. Given two classes of objects $\mathcal{X}, \mathcal{Y} \subseteq \mathcal{D}$, we define the class $\mathcal{X} \star \mathcal{Y}$ as the subcategory of objects $C \in \mathcal{D}$ that admit an \mathfrak{s} -conflation $X \rightarrow C \rightarrow Y$ with $X \in \mathcal{X}$ and $Y \in \mathcal{Y}$. We say that a class of objects $\mathcal{X} \subseteq \mathcal{D}$ is **closed under extensions** if $\mathcal{X} \star \mathcal{X} \subseteq \mathcal{X}$. A class $\mathcal{Z} \subseteq \mathcal{D}$ is **closed under cones** if for any \mathfrak{s} -conflation $A \rightarrow B \rightarrow C$ with $A, B \in \mathcal{Z}$, we have that $C \in \mathcal{Z}$. Dually, \mathcal{Z} is **closed under cocones** if for any \mathfrak{s} -conflation $A \rightarrow B \rightarrow C$ with $B, C \in \mathcal{Z}$, we have that $A \in \mathcal{Z}$.

Remark 2.1. Let $(\mathcal{C}, \mathbb{E}, \mathfrak{s})$ be an extriangulated category and $\mathcal{X}, \mathcal{Y}, \mathcal{Z} \subseteq \mathcal{D}$ be classes of objects.

- (a) It is a known fact that $\mathcal{X} \star (\mathcal{Y} \star \mathcal{Z}) = (\mathcal{X} \star \mathcal{Y}) \star \mathcal{Z}$ (see for example [1, p.454]).
- (b) If \mathcal{X} is closed under extensions, then \mathcal{X} becomes an extriangulated category by restricting \mathbb{E} and \mathfrak{s} on \mathcal{X} , see [18, Rk. 2.18].
- (c) For the sake of simplicity, we write \mathcal{C} instead of $(\mathcal{C}, \mathbb{E}, \mathfrak{s})$ for referring to the extriangulated category $(\mathcal{C}, \mathbb{E}, \mathfrak{s})$.

The following long exact sequences that come from an \mathfrak{s} -conflation will be very useful throughout the paper.

Proposition 2.2. [18, Cor. 3.12] *For any \mathfrak{s} -conflation $\delta : A \xrightarrow{u} B \xrightarrow{v} C$ in an extriangulated category \mathcal{D} and any $W \in \mathcal{D}$, the sequences*

$$(W, A) \xrightarrow{(W, u)} (W, B) \xrightarrow{(W, v)} (W, C) \xrightarrow{\delta \cdot -} \mathbb{E}(W, A) \xrightarrow{\mathbb{E}(W, u)} \mathbb{E}(W, B) \xrightarrow{\mathbb{E}(W, v)} \mathbb{E}(W, C),$$

$$(C, W) \xrightarrow{(v, W)} (B, W) \xrightarrow{(u, W)} (A, W) \xrightarrow{-\delta} \mathbb{E}(C, W) \xrightarrow{\mathbb{E}(v, W)} \mathbb{E}(B, W) \xrightarrow{\mathbb{E}(u, W)} \mathbb{E}(A, W)$$

are exact, where $(W, -) := \mathcal{D}(W, -)$ and $(-, W) := \mathcal{D}(-, W)$.

Let \mathcal{C} be an additive category and $A \xrightarrow{u} B \xrightarrow{v} C$ be morphisms in \mathcal{C} . We recall that u is a **weak kernel** of v if $vu = 0$ and any morphism $a : W \rightarrow B$ in \mathcal{C} such that $va = 0$ factors through u . Dually, v is a **weak cokernel** of u if $vu = 0$ and any morphism $b : B \rightarrow W$ in \mathcal{C} such that $bu = 0$ factors through v .

Remark 2.3. Let \mathcal{D} be an extriangulated category and $\delta : A \xrightarrow{u} B \xrightarrow{v} C$ be any \mathfrak{s} -conflation. Then, as a consequence of Proposition 2.2, we have that u is a weak kernel of v and v is a weak cokernel of u .

2.1 Extriangulated categories with negative first extensions and \mathfrak{s} -torsion pairs

Let $\mathcal{D} = (\mathcal{D}, \mathbb{E}, \mathfrak{s})$ be an extriangulated category and $\delta : A \xrightarrow{u} B \xrightarrow{v} C$ be any \mathfrak{s} -conflation. By Proposition 2.2, we get that any $W \in \mathcal{D}$ gives the exact

sequences

$$\begin{aligned} (W, A) &\xrightarrow{(W, u)} (W, B) \xrightarrow{(W, v)} (W, C) \xrightarrow{\delta} \mathbb{E}(W, A) \xrightarrow{\mathbb{E}(W, u)} \mathbb{E}(W, B) \xrightarrow{\mathbb{E}(W, v)} \mathbb{E}(W, C), \\ (C, W) &\xrightarrow{(v, W)} (B, W) \xrightarrow{(u, W)} (A, W) \xrightarrow{-\delta} \mathbb{E}(C, W) \xrightarrow{\mathbb{E}(v, W)} \mathbb{E}(B, W) \xrightarrow{\mathbb{E}(u, W)} \mathbb{E}(A, W). \end{aligned}$$

It is well-known that the above exact sequences can be extended to the left for many extriangulated categories. For example, in the case of abelian categories, the sequence is extended with zeros; and for triangulated categories, it is extended with the functor $\mathcal{D}(W, -[-1])$. When an extriangulated category satisfies this, it is said to have negative first extension. In the following we recall such notion from [1], the reader is referred to [12] for additional information.

Definition 2.4. [1, Def. 2.3] Let $\mathcal{D} = (\mathcal{D}, \mathbb{E}, \mathfrak{s})$ be an extriangulated category. We say that \mathcal{D} has **negative first extension** if there is an additive bifunctor $\mathbb{E}^{-1} : \mathcal{D}^{op} \times \mathcal{D} \rightarrow \text{Ab}$, and for each $\delta \in \mathbb{E}(C, A)$ there exist two natural transformations $\delta^{-1} : \mathbb{E}^{-1}(-, C) \rightarrow \mathcal{D}(-, A)$ and $\delta_{-1} : \mathbb{E}^{-1}(A, -) \rightarrow \mathcal{D}(C, -)$ such that, for any \mathfrak{s} -conflation $\delta : A \xrightarrow{u} B \xrightarrow{v} C$ and each $W \in \mathcal{D}$, the sequences

$$\begin{aligned} \mathbb{E}^{-1}(W, A) &\xrightarrow{\mathbb{E}^{-1}(W, u)} \mathbb{E}^{-1}(W, B) \xrightarrow{\mathbb{E}^{-1}(W, v)} \mathbb{E}^{-1}(W, C) \xrightarrow{\delta^{-1}} (W, A) \xrightarrow{(W, u)} (W, B), \\ \mathbb{E}^{-1}(C, W) &\xrightarrow{\mathbb{E}^{-1}(v, W)} \mathbb{E}^{-1}(B, W) \xrightarrow{\mathbb{E}^{-1}(u, W)} \mathbb{E}^{-1}(A, W) \xrightarrow{\delta_{-1}} (C, W) \xrightarrow{(v, W)} (B, W) \end{aligned}$$

are exact. In this case, we say that $\mathcal{D} = (\mathcal{D}, \mathbb{E}, \mathfrak{s}, \mathbb{E}^{-1})$ is an extriangulated category with a negative first extension.

It can be seen that, by restricting the structure of an extriangulated category with negative first extension to a subcategory closed under extensions, we obtain a new extriangulated category with negative first extension, see [1, Ex. 2.4(iii)]. A particular case, in which we will be interested, is the one of subcategories of triangulated categories which are closed under extensions.

The structure of extriangulated categories allows us to develop concepts parallel to that of torsion pair in abelian categories. In this section we will present the notion of s -torsion for extriangulated categories with negative first extension and recall some basic properties.

We start with the following definition of torsion pair in extriangulated categories that comes naturally from the notion of torsion pair in abelian categories.

Definition 2.5. [14, Def. 3.1] Let \mathcal{D} be an extriangulated category. A **torsion pair** in \mathcal{D} is a pair $\mathbf{x} = (\mathcal{X}, \mathcal{Y})$ of full subcategories of \mathcal{D} which are closed under isomorphisms in \mathcal{D} and such that $\mathcal{D}(\mathcal{X}, \mathcal{Y}) = 0$ and $\mathcal{D} = \mathcal{X} \star \mathcal{Y}$.

In case the extriangulated category has negative first extensions, we have the notion of s -torsion pair. By following [1], we recall that the term “ s ” in s -torsion stands for “shift-closed” by means of [1, Lem. 3.3].

Definition 2.6. [1, Def. 3.1] Let \mathcal{D} be an extriangulated category with negative first extension. An **s -torsion pair** in \mathcal{D} is a torsion pair $\mathbf{x} = (\mathcal{X}, \mathcal{Y})$ such that $\mathbb{E}^{-1}(\mathcal{X}, \mathcal{Y}) = 0$.

Let \mathcal{D} be an extriangulated category with negative first extension. For a class $\mathcal{X} \subseteq \mathcal{D}$ and $i \in \{-1, 0, 1\}$, we define the right i -th orthogonal complement of \mathcal{X} by $\mathcal{X}^{\perp_i} := \{D \in \mathcal{D} : \mathbb{E}^i(\mathcal{X}, D) = 0\}$, where $\mathbb{E}^0(-, -) := \text{Hom}_{\mathcal{D}}(-, -)$. Dually, we have ${}^{\perp_i}\mathcal{X}$ which is the left i -th orthogonal complement of \mathcal{X} . In case the class \mathcal{X} has only one element, i.e. $\mathcal{X} = \{X\}$, we set $X^{\perp_i} := \mathcal{X}^{\perp_i}$ and ${}^{\perp_i}X := {}^{\perp_i}\mathcal{X}$. As a consequence of the above definitions, we get the following remark.

Remark 2.7. For an extriangulated category \mathcal{D} with negative first extension, a class $\mathcal{X} \subseteq \mathcal{D}$ and $i \in \{-1, 0, 1\}$, the classes \mathcal{X}^{\perp_i} and ${}^{\perp_i}\mathcal{X}$ are closed under extensions and direct summands.

Proposition 2.8. [1, Prop. 3.2] *Let $(\mathcal{T}, \mathcal{F})$ be an s -torsion pair in \mathcal{D} . Then, $\mathcal{T}^{\perp_0} = \mathcal{F}$ and ${}^{\perp_0}\mathcal{F} = \mathcal{T}$.*

Proposition 2.9. *For an s -torsion pair $(\mathcal{T}, \mathcal{F})$ in \mathcal{D} , the following statements hold true.*

- (a) \mathcal{T} is closed under cones, extensions and direct summands.
- (b) \mathcal{F} is closed under cocones, extensions and direct summands.
- (c) *Let $X \rightarrow Y \rightarrow Z$ be an \mathfrak{s} -conflation. Then:*
 - (c1) $X \in {}^{\perp_{-1}}\mathcal{F}$ if $Y, Z \in \mathcal{T}$;
 - (c2) $Z \in \mathcal{T}^{\perp_{-1}}$ if $X, Y \in \mathcal{F}$.
- (d) *For any s -torsion pair $(\mathcal{X}, \mathcal{Y})$ in \mathcal{D} , $\mathcal{X} \subseteq \mathcal{T} \Leftrightarrow \mathcal{F} \subseteq \mathcal{Y}$.*

Proof. The items (a), (b) and (c) follow from the long exact sequences appearing in Definition 2.6, Remark 2.7 and Proposition 2.8. Finally, (d) can be obtained from Proposition 2.8. \square

Remark 2.10. Let $\mathfrak{t} = (\mathcal{T}, \mathcal{F})$ be an s -torsion pair in \mathcal{D} . It follows that, for all $C \in \mathcal{D}$, there is an \mathfrak{s} -conflation $\delta_C : T_C \xrightarrow{u} C \xrightarrow{v} F_C$ with $T_C \in \mathcal{T}$ and $F_C \in \mathcal{F}$. An important property of s -torsion pairs is that the \mathfrak{s} -conflation δ_C is unique up to isomorphisms of \mathfrak{s} -conflations. Thus, we have a pair of functors: the torsion functor $\mathfrak{t} : \mathcal{D} \rightarrow \mathcal{T}$, $C \mapsto T_C$, and the torsion-free functor $(\mathbf{1} : \mathfrak{t}) : \mathcal{D} \rightarrow \mathcal{F}$, $C \mapsto F_C$, which are, respectively, the right and left adjoint functors of the inclusion functors $\mathcal{T} \rightarrow \mathcal{D}$ and $\mathcal{F} \rightarrow \mathcal{D}$. The reader is referred to [1, Prop. 3.7] for more details.

2.2 t -structures in triangulated categories

A particular case of an s -torsion pair in an extriangulated category with negative first extension is the notion of t -structure in a triangulated category. Since this will be an object of interest in the paper, we include below the definition and some remarks.

Recall that a triangulated category consists of a triple $(\mathcal{D}, \Sigma, \Delta)$, where \mathcal{D} is an additive category, $\Sigma : \mathcal{D} \rightarrow \mathcal{D}$ is an automorphism, and Δ is a class

of sequences of morphisms (called triangles) of the form $N \rightarrow M \rightarrow K \rightarrow \Sigma N$ satisfying a series of axioms. The reader is referred to [19] for a precise definition of triangulated category. The notion of t -structure was first introduced in [8]. Here we present an equivalent definition, see [24, Rk. 4.2] for details.

Definition 2.11. Let $(\mathcal{D}, \Sigma, \Delta)$ be a triangulated category. A pair $(\mathcal{U}, \mathcal{W})$ of full subcategories of \mathcal{D} is called a **t -structure** in \mathcal{D} if: \mathcal{U} and \mathcal{W} are closed under isomorphisms, $\Sigma\mathcal{U} \subseteq \mathcal{U}$ and $(\mathcal{U}, \Sigma^{-1}\mathcal{W})$ is a torsion pair in \mathcal{D} .

Remark 2.12. Let $(\mathcal{D}, \Sigma, \Delta)$ be a triangulated category and $(\mathcal{U}, \mathcal{W})$ a pair of full subcategories of \mathcal{D} .

- (a) [1, Lem. 3.3] The pair $(\mathcal{U}, \mathcal{W})$ is a t -structure in \mathcal{D} if and only if $(\mathcal{U}, \Sigma^{-1}\mathcal{W})$ is an s -torsion pair in \mathcal{D} .
- (b) [8] If $\mathbf{u} := (\mathcal{U}, \mathcal{W})$ is a t -structure in \mathcal{D} , then $\mathcal{H} = \mathcal{H}_{\mathbf{u}} := \mathcal{U} \cap \mathcal{W}$ is an abelian category and it is known as the **heart** of \mathbf{u} .
- (c) Let $t := (\mathcal{T}, \mathcal{F})$ be an s -torsion pair in \mathcal{D} . Then $\Sigma t := (\Sigma\mathcal{T}, \Sigma\mathcal{F})$ and $\Sigma^{-1}t := (\Sigma^{-1}\mathcal{T}, \Sigma^{-1}\mathcal{F})$ are s -torsion pairs in \mathcal{D} and $\Sigma\mathcal{T} \subseteq \mathcal{T} \subseteq \Sigma^{-1}\mathcal{T}$.

2.3 Intervals of s -torsion pairs

Let \mathcal{D} be an extriangulated category with negative first extension. Consider two s -torsion pairs $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ and $\mathbf{t}' = (\mathcal{T}', \mathcal{F}')$ in \mathcal{D} . By following [1], $\mathbf{t} \leq \mathbf{t}'$ if $\mathcal{T} \subseteq \mathcal{T}'$. By using this relation, we can consider the poset of s -torsion pairs in \mathcal{D} that is denoted by $\text{stors } \mathcal{D}$. Moreover, given $\mathbf{r} := (\mathcal{R}, \mathcal{M})$ and $\mathbf{s} := (\mathcal{S}, \mathcal{N})$ in $\text{stors } \mathcal{D}$ such that $\mathbf{r} \leq \mathbf{s}$, we have the interval $[\mathbf{r}, \mathbf{s}] := \{\mathbf{t} \in \text{stors } \mathcal{D} : \mathbf{r} \leq \mathbf{t} \leq \mathbf{s}\}$ and define the **heart** of $[\mathbf{r}, \mathbf{s}]$ as the class $\mathcal{H}_{[\mathbf{r}, \mathbf{s}]} := \mathcal{M} \cap \mathcal{S}$. Observe that $\mathcal{H}_{[\mathbf{r}, \mathbf{s}]}$ is closed under extensions, and thus $\mathcal{H}_{[\mathbf{r}, \mathbf{s}]}$ inherits an extriangulated structure with negative first extension from the triangulated category \mathcal{D} .

Example 2.13. Let \mathcal{D} be a triangulated category and $\mathbf{t} := (\mathcal{T}, \mathcal{F})$ be an s -torsion pair in \mathcal{D} . Then, by Remark 2.12(c), we have that $\Sigma\mathbf{t} := (\Sigma\mathcal{T}, \Sigma\mathcal{F})$ and $\Sigma^{-1}\mathbf{t} := (\Sigma^{-1}\mathcal{T}, \Sigma^{-1}\mathcal{F})$ are s -torsion pairs in \mathcal{D} and $\Sigma\mathbf{t} \leq \mathbf{t} \leq \Sigma^{-1}\mathbf{t}$. Now, consider the t -structures which correspond, respectively, to $\Sigma\mathbf{t}$ and $\Sigma^{-1}\mathbf{t}$; namely: $\mathbf{t}_* := (\Sigma\mathcal{T}, \Sigma^2\mathcal{F})$ and $\mathbf{t}^* := (\Sigma^{-1}\mathcal{T}, \mathcal{F})$. Therefore, it follows that $\mathcal{H}_{[\Sigma\mathbf{t}, \mathbf{t}]} = \Sigma\mathcal{F} \cap \mathcal{T} = \Sigma^{-1}\mathcal{H}_{\mathbf{t}_*}$ and $\mathcal{H}_{[\mathbf{t}, \Sigma^{-1}\mathbf{t}]} = \mathcal{F} \cap \Sigma^{-1}\mathcal{T} = \mathcal{H}_{\mathbf{t}^*}$ are abelian categories.

Example 2.14. [28, Sect. 1.2], [10, Thm. 2.15], [29, Sect. 5]. Let \mathcal{E} be a **quasi-abelian** category. That is, \mathcal{E} is an additive category such that: every morphism admits a kernel and a cokernel; and every push-out (pull-back) of a kernel (cokernel) is a kernel (cokernel). We can associate to \mathcal{E} a triangulated category \mathcal{D} . Moreover, \mathcal{D} is equipped with a pair of s -torsion pairs $\mathbf{t}_r, \mathbf{t}_\ell \in \text{stors } \mathcal{D}$ such that $\Sigma\mathbf{t}_r \leq \mathbf{t}_\ell \leq \mathbf{t}_r$ and $\mathcal{H}_{[\Sigma\mathbf{t}_r, \mathbf{t}_\ell]} \cong \mathcal{E}$. The category $\mathcal{R}_{\mathcal{E}} := \mathcal{H}_{[\Sigma\mathbf{t}_r, \mathbf{t}_r]}$ is known as the **right associated abelian category** of \mathcal{E} , and the category $\mathcal{L}_{\mathcal{E}} := \mathcal{H}_{[\Sigma\mathbf{t}_\ell, \mathbf{t}_\ell]}$ is known as the **left associated abelian category** of \mathcal{E} .

Remark 2.15. [1, Lem. 3.11] Let $\mathbf{r} := (\mathcal{R}, \mathcal{M}), \mathbf{s} := (\mathcal{S}, \mathcal{N}) \in \text{stors } \mathcal{D}$ be such that $\mathbf{r} \leq \mathbf{s}$. Then $\mathcal{S} = \mathcal{R} \star \mathcal{H}_{[\mathbf{r}, \mathbf{s}]}$ and $\mathcal{M} = \mathcal{H}_{[\mathbf{r}, \mathbf{s}]} \star \mathcal{N}$.

A classical result from [13] shows how to parameterize a certain family of t -structures by means of the torsion pairs of the heart of a given t -structure. The following theorem can be understood as a generalization of this result. It also extends similar results for quasi-abelian subcategories of abelian categories [29].

Theorem 2.16. [1, Thm. 3.9] *Let \mathcal{D} be an extriangulated category with negative first extension, and let $\mathbf{t}_1 = (\mathcal{T}_1, \mathcal{F}_1)$ and $\mathbf{t}_2 = (\mathcal{T}_2, \mathcal{F}_2)$ in $\text{stors } \mathcal{D}$ be such that $\mathbf{t}_1 \leq \mathbf{t}_2$. Then, there exist an isomorphism of posets*

$$\Phi : [\mathbf{t}_1, \mathbf{t}_2] \rightarrow \text{stors } \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}, (\mathcal{T}, \mathcal{F}) \mapsto (\mathcal{T} \cap \mathcal{F}_1, \mathcal{T}_2 \cap \mathcal{F})$$

with inverse

$$\Psi : \text{stors } \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} \rightarrow [\mathbf{t}_1, \mathbf{t}_2], (\mathcal{X}, \mathcal{Y}) \mapsto (\mathcal{T}_1 \star \mathcal{X}, \mathcal{Y} \star \mathcal{F}_2).$$

3 Extensions of hearts and their s -torsion pairs

Lemma 3.1. *Let \mathcal{D} be an extriangulated category with negative first extension, and let $\mathbf{t}_i = (\mathcal{T}_i, \mathcal{F}_i)$ in $\text{stors } \mathcal{D}$ for $i = 1, 2, 3$ be such that $\mathbf{t}_1 \leq \mathbf{t}_2 \leq \mathbf{t}_3$. Then*

$$\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]} = \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} \star \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}.$$

Moreover, for $C \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ and the \mathfrak{s} -conflation $T_2 \rightarrow C \rightarrow F_2$ given by \mathbf{t}_2 , we have that $F_2 \in \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$ and $T_2 \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$.

Proof. Since $\mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]} = \mathcal{T}_3 \cap \mathcal{F}_2 \subseteq \mathcal{T}_3 \cap \mathcal{F}_1 = \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$, $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} = \mathcal{T}_2 \cap \mathcal{F}_1 \subseteq \mathcal{T}_3 \cap \mathcal{F}_1 = \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ and $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ is closed under extensions, it follows that $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} \star \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]} \subseteq \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$. Let $C \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$. Consider the \mathfrak{s} -conflation $T_2 \rightarrow C \rightarrow F_2$ with $T_2 \in \mathcal{T}_2$ and $F_2 \in \mathcal{F}_2$ given by \mathbf{t}_2 . By Proposition 2.9(a), we obtain that $F_2 \in \mathcal{T}_3$ since $T_2 \in \mathcal{T}_2 \subseteq \mathcal{T}_3$ and $C \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]} \subseteq \mathcal{T}_3$. Similarly, it can be shown that $T_2 \in \mathcal{F}_1$. Therefore $F_2 \in \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$ and $T_2 \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$; proving that $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]} \subseteq \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} \star \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$. \square

Lemma 3.2. *For an extriangulated category \mathcal{D} with negative first extension, and $\mathbf{t}_i = (\mathcal{T}_i, \mathcal{F}_i)$ in $\text{stors } \mathcal{D}$ for $i = 1, 2, 3$ such that $\mathbf{t}_1 \leq \mathbf{t}_2 \leq \mathbf{t}_3$, the following statements hold true.*

- (a) *If $(\mathcal{T}, \mathcal{F})$ is an s -torsion pair in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$, then $(\mathcal{T}, \mathcal{F} \star \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]})$ is an s -torsion pair in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$.*
- (b) *If $(\mathcal{T}, \mathcal{F})$ is an s -torsion pair in $\mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$, then $(\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} \star \mathcal{T}, \mathcal{F})$ is an s -torsion pair in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$.*
- (c) *$(\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}, \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]})$ is an s -torsion pair in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$.*

- (d) If $(\mathcal{T}, \mathcal{F})$ is an s -torsion pair in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ with $\mathcal{T} \subseteq \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$, then $\mathcal{F}' := \mathcal{F} \cap \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ satisfies that $(\mathcal{T}, \mathcal{F}') \in \text{stors } \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ and $\mathcal{F} = \mathcal{F}' \star \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$.
- (e) If $(\mathcal{T}, \mathcal{F})$ is an s -torsion pair in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ with $\mathcal{F} \subseteq \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$, then $\mathcal{T}' := \mathcal{T} \cap \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$ satisfies that $(\mathcal{T}', \mathcal{F}) \in \text{stors } \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$ and $\mathcal{T} = \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} \star \mathcal{T}'$.
- (f) For every $(\mathcal{T}, \mathcal{F}) \in \text{stors } \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ there is $\mathbf{t} \in [\mathbf{t}_1, \mathbf{t}_2]$ such that $\mathcal{T} = \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}]}$ and $\mathcal{F} = \mathcal{H}_{[\mathbf{t}, \mathbf{t}_2]}$.

Proof. We only prove (a) and (d) since: (b) follows by dual arguments as (a), (c) is a particular case of (a), (e) follows by dual arguments as (d), and (f) is a consequence of Theorem 2.16.

(a) Let $(\mathcal{T}, \mathcal{F})$ be an s -torsion pair in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. Then $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} = \mathcal{T} \star \mathcal{F}$ and thus, by Lemma 3.1, we have that

$$\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]} = \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} \star \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]} = (\mathcal{T} \star \mathcal{F}) \star \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]} = \mathcal{T} \star (\mathcal{F} \star \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}).$$

It remains to prove that $\mathcal{D}(\mathcal{T}, \mathcal{F} \star \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}) = 0$ and that $\mathbb{E}^{-1}(\mathcal{T}, \mathcal{F} \star \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}) = 0$. Indeed, since $\mathcal{F} \subseteq \mathcal{T}^{\perp_0} \cap \mathcal{T}^{\perp_{-1}}$, $\mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]} \subseteq \mathcal{F}_2 \subseteq \mathcal{T}_2^{\perp_0} \cap \mathcal{T}_2^{\perp_{-1}} \subseteq \mathcal{T}^{\perp_0} \cap \mathcal{T}^{\perp_{-1}}$ and \mathcal{T}^{\perp_0} and $\mathcal{T}^{\perp_{-1}}$ are closed under extensions, we have that $\mathcal{D}(\mathcal{T}, \mathcal{F} \star \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}) = 0$ and $\mathbb{E}^{-1}(\mathcal{T}, \mathcal{F} \star \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}) = 0$.

(d) Let $\mathbf{t} = (\mathcal{T}, \mathcal{F}) \in \text{stors } \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ with $\mathcal{T} \subseteq \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. Consider the s -torsion pair $\mathbf{c} = (\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}, \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}) \in \text{stors } \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ showed in (c) and the following maps given by using Theorem 2.16:

$$\text{stors } \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]} \xrightarrow{\Psi_{13}} [\mathbf{t}_1, \mathbf{t}_3] \leftrightarrow [\mathbf{t}_1, \mathbf{t}_2] \xrightarrow{\Phi_{12}} \text{stors } \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}.$$

Since $\mathbf{t} \leq \mathbf{c}$ and $\mathbf{c} = \Phi_{13}(\mathbf{t}_2)$, we have that $\Psi_{13}(\mathbf{t}) \in [\mathbf{t}_1, \mathbf{t}_2]$. Therefore, $\Phi_{12}\Psi_{13}(\mathbf{t}) \in \text{stors } \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. Observe that

$$\Phi_{12}\Psi_{13}(\mathbf{t}) = ((\mathcal{T}_1 \star \mathcal{T}) \cap \mathcal{F}_1, \mathcal{T}_2 \cap (\mathcal{F} \star \mathcal{F}_3)) = (\mathcal{T}, \mathcal{F} \cap \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}).$$

Indeed, on the one hand, the contention $(\mathcal{T}_1 \star \mathcal{T}) \cap \mathcal{F}_1 \subseteq \mathcal{T}$ follows straightforward and the opposite contention follows from the fact that $\mathcal{T} \subseteq \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} = \mathcal{T}_2 \cap \mathcal{F}_1$. On the other hand, for the contention $\mathcal{T}_2 \cap (\mathcal{F} \star \mathcal{F}_3) \subseteq \mathcal{F} \cap \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ consider the facts that $\mathcal{F} \subseteq \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]} = \mathcal{F}_1 \cap \mathcal{T}_3$, $\mathcal{F}_3 \subseteq \mathcal{F}_1$ and $\mathcal{T}_2 \subseteq \mathcal{T}_3$; and for the opposite contention, observe that $\mathcal{F} \subseteq \mathcal{F} \star \mathcal{F}_3$ and $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} \subseteq \mathcal{T}_2$. Lastly, note that $\mathcal{F} = (\mathcal{F} \cap \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}) \star \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$ since $\mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]} \subseteq \mathcal{F}$ because $\mathcal{T} \subseteq \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. \square

As a application of Lemma 3.2, we can get the following example in a direct way. This was already known, from [28, Sect. 1.2], [10, Them. 2.15], [29, Sect. 5].

Example 3.3. Let \mathcal{E} be a quasi-abelian category (see Example 2.14). We have mentioned that there is a triangulated category \mathcal{D} equipped with a pair of s -torsion pairs $\mathbf{t}_r, \mathbf{t}_\ell \in \text{stors } \mathcal{D}$ such that $\Sigma \mathbf{t}_r \leq \mathbf{t}_\ell \leq \mathbf{t}_r$ and $\mathcal{H}_{[\Sigma \mathbf{t}_r, \mathbf{t}_\ell]} \cong \mathcal{E}$. It follows from Lemma 3.2 (c) that $(\mathcal{H}_{[\Sigma \mathbf{t}_r, \mathbf{t}_\ell]}, \mathcal{H}_{[\mathbf{t}_\ell, \mathbf{t}_r]})$ is a torsion pair in the abelian category $\mathcal{R}_{\mathcal{E}} := \mathcal{H}_{[\Sigma \mathbf{t}_r, \mathbf{t}_r]}$, and that $(\mathcal{H}_{[\Sigma \mathbf{t}_\ell, \Sigma \mathbf{t}_r]}, \mathcal{H}_{[\Sigma \mathbf{t}_r, \mathbf{t}_\ell]})$ is a torsion pair in the abelian category $\mathcal{L}_{\mathcal{E}} := \mathcal{H}_{[\Sigma \mathbf{t}_\ell, \mathbf{t}_\ell]}$. Therefore, \mathcal{E} can be realized as a torsion class in the abelian category $\mathcal{R}_{\mathcal{E}}$, or as torsion-free class in the abelian category $\mathcal{L}_{\mathcal{E}}$.

From the previous lemmas, we obtain the following result.

Corollary 3.4. *Let \mathcal{D} be an extriangulated category with negative first extension, and let $\mathbf{t}_i = (\mathcal{T}_i, \mathcal{F}_i)$ in $\text{stors } \mathcal{D}$, for $i = 1, \dots, n$, be such that $\mathbf{t}_1 \leq \dots \leq \mathbf{t}_n$. Then, the following statements hold true.*

- (a) $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_n]} = \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} \star \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]} \star \dots \star \mathcal{H}_{[\mathbf{t}_{n-1}, \mathbf{t}_n]}$.
- (b) If $(\mathcal{T}, \mathcal{F})$ is an s -torsion pair in $\mathcal{H}_{[\mathbf{t}_k, \mathbf{t}_{k+1}]}$ for some $1 \leq k < n$, then $(\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_k]} \star \mathcal{T}, \mathcal{F} \star \mathcal{H}_{[\mathbf{t}_{k+1}, \mathbf{t}_n]})$ is an s -torsion pair in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_n]}$.

Definition 3.5. Let \mathcal{D} be an extriangulated category with negative first extension, and let $\mathbf{t}_i = (\mathcal{T}_i, \mathcal{F}_i)$ in $\text{stors } \mathcal{D}$ for $i = 1, 2$ be such that $\mathbf{t}_1 \leq \mathbf{t}_2$. We say that $[\mathbf{t}_1, \mathbf{t}_2]$ is a **normal interval** if $\mathbb{E}^{-1}(\mathcal{T}_2, \mathcal{F}_1) = 0$.

Example 3.6. Let \mathcal{D} be an extriangulated category with negative first extension.

- (a) If $\mathbb{E}^{-1} = 0$ (e.g. in the case \mathcal{D} is exact), then any interval in $\text{stors } \mathcal{D}$ is normal.
- (b) Let \mathcal{D} be a triangulated category and $[\mathbf{t}_1, \mathbf{t}_2]$ be an interval in $\text{stors } \mathcal{D}$. Then $\mathbb{E}^{-1}(\mathcal{T}_2, \mathcal{F}_1) = \mathcal{D}(\mathcal{T}_2, \Sigma^{-1}\mathcal{F}_1)$. Therefore, if we fix \mathbf{t}_1 , the interval $[\mathbf{t}_1, \mathbf{t}_2]$ is normal if, and only if, \mathbf{t}_2 satisfies that $\Sigma^{-1}\mathcal{F}_1 \subseteq \mathcal{F}_2$, or equivalently that $\Sigma\mathcal{T}_2 \subseteq \mathcal{T}_1$. Note that these are precisely the s -torsion pairs (t -structures) parametrized by the HRS-tilting process [30, Prop. 2.1].

Lemma 3.7. *Let \mathcal{D} be an extriangulated category with negative first extension, and let $[\mathbf{t}_1, \mathbf{t}_2]$ be a normal interval in $\text{stors } \mathcal{D}$. Then $\mathbb{E}^{-1}(\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}, \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}) = 0$ and thus $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ is an exact category.*

Proof. Let $\mathbf{t}_i = (\mathcal{T}_i, \mathcal{F}_i)$, for $i = 1, 2$. Since $\mathbb{E}^{-1}(\mathcal{T}_2, \mathcal{F}_1) = 0$ and $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} = \mathcal{F}_1 \cap \mathcal{T}_2$, we get that $\mathbb{E}^{-1}(\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}, \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}) = 0$. Finally, by [1, Prop. 2.6], we conclude that $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ is an exact category. \square

Lemma 3.8. *For an extriangulated category \mathcal{D} with negative first extension and $\mathbf{t}_i = (\mathcal{T}_i, \mathcal{F}_i)$ in $\text{stors } \mathcal{D}$ for $i = 1, 2, 3$ such that $\mathbf{t}_1 \leq \mathbf{t}_2 \leq \mathbf{t}_3$, the following statements hold true.*

- (a) If $[\mathbf{t}_1, \mathbf{t}_3]$ is normal, then $[\mathbf{t}_1, \mathbf{t}_2]$ and $[\mathbf{t}_2, \mathbf{t}_3]$ are normal.
- (b) If $[\mathbf{t}_1, \mathbf{t}_2]$ and $[\mathbf{t}_2, \mathbf{t}_3]$ are normal and $\mathbb{E}^{-1}(\mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}, \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}) = 0$, then $[\mathbf{t}_1, \mathbf{t}_3]$ is normal.

Proof. The proof of (a) is straightforward since $\mathcal{F}_2 \subseteq \mathcal{F}_1$ and $\mathcal{T}_2 \subseteq \mathcal{T}_3$. Let $\mathbb{E}^{-1}(\mathcal{T}_3, \mathcal{F}_2) = 0$, $\mathbb{E}^{-1}(\mathcal{T}_2, \mathcal{F}_1) = 0$ and $\mathbb{E}^{-1}(\mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}, \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}) = 0$. In order to prove (b), we need to show that $\mathbb{E}^{-1}(\mathcal{T}_3, \mathcal{F}_1) = 0$. Indeed, by Remark 2.15 we get that $\mathcal{T}_3 = \mathcal{T}_1 \star \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ and $\mathcal{F}_1 = \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]} \star \mathcal{F}_3$. Hence, it is enough to prove that: (1) $\mathbb{E}^{-1}(\mathcal{T}_1, \mathcal{F}_3) = 0$, (2) $\mathbb{E}^{-1}(\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}, \mathcal{F}_3) = 0$, (3) $\mathbb{E}^{-1}(\mathcal{T}_1, \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}) = 0$, and (4) $\mathbb{E}^{-1}(\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}, \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}) = 0$. Notice that (1), (2) and (3) follow from

the equality $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]} = \mathcal{F}_1 \cap \mathcal{T}_3$ and the inclusion $\mathcal{F}_3 \subseteq \mathcal{F}_1$. In order to prove (4), we use that $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]} = \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} \star \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$ (see Lemma 3.1). Thus, we need to show that: (A) $\mathbb{E}^{-1}(\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}, \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}) = 0$, (B) $\mathbb{E}^{-1}(\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}, \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}) = 0$, (C) $\mathbb{E}^{-1}(\mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}, \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}) = 0$ and (D) $\mathbb{E}^{-1}(\mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}, \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}) = 0$. Notice that (A) and (D) follow from Lemma 3.7; the item (B) follows from the fact that $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} \subseteq \mathcal{T}_2$ and $\mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]} \subseteq \mathcal{F}_2$; and the item (C) is an hypothesis. \square

Proposition 3.9. *Let \mathcal{D} be an extriangulated category with negative first extension and $\mathbf{t}_1, \mathbf{t}_2, \mathbf{t}_3 \in \text{stors } \mathcal{D}$ such that $\mathbf{t}_1 \leq \mathbf{t}_2 \leq \mathbf{t}_3$. Then, for an \mathfrak{s} -conflation $X \rightarrow Y \rightarrow Z$ in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$, the following statements hold true,*

- (a) *If $[\mathbf{t}_2, \mathbf{t}_3]$ is normal and $Y \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$, then $Z \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$.*
- (b) *If $[\mathbf{t}_1, \mathbf{t}_2]$ is normal and $Y \in \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$, then $X \in \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$.*

Proof. We only prove (a) since the proof of (b) is similar. Consider an \mathfrak{s} -conflation $X \rightarrow Y \rightarrow Z$ in \mathcal{C} with $Y \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. Note that $X \in {}^{\perp -1} \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$ since $X \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]} = \mathcal{T}_3 \cap \mathcal{F}_1 \subseteq \mathcal{T}_3$, $\mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]} \subseteq \mathcal{F}_2$, and $[\mathbf{t}_2, \mathbf{t}_3]$ is normal. Then, it follows from the previous \mathfrak{s} -conflation that $Z \in {}^{\perp 0} \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$. Thus, by applying Proposition 2.8 to the s -torsion pair $(\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}, \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]})$ in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ (see Lemma 3.2 (c)), we have that $Z \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. \square

Definition 3.10. Let \mathcal{D} be an extriangulated category. We say that an \mathfrak{s} -conflation $N \xrightarrow{a} M \xrightarrow{b} K$ in \mathcal{D} is a **short exact sequence** if: a is the kernel of b in \mathcal{D} , and b is the cokernel of a in \mathcal{D} .

Proposition 3.11. *Let \mathcal{D} be an extriangulated category with negative first extension, and let $\mathbf{t}_i = (\mathcal{T}_i, \mathcal{F}_i)$ in $\text{stors } \mathcal{D}$ for $i = 1, 2, 3$ be such that $\mathbf{t}_1 \leq \mathbf{t}_2 \leq \mathbf{t}_3$. Then, for any $C \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$, the following statements hold true for the \mathfrak{s} -conflation $\eta : T_2 \xrightarrow{f_C} C \xrightarrow{g_C} F_2$ in \mathcal{D} given by \mathbf{t}_2 .*

- (a) *$T_2, F_2 \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$, and thus η is an \mathfrak{s} -conflation in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$.*
- (b) *If $[\mathbf{t}_2, \mathbf{t}_3]$ is normal, then f_C is a monomorphism in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ and f_C is the kernel of g_C in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$.*
- (c) *If $[\mathbf{t}_1, \mathbf{t}_2]$ is normal, then g_C is an epimorphism in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ and g_C is the cokernel of f_C in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$.*
- (d) *If $[\mathbf{t}_1, \mathbf{t}_2]$ and $[\mathbf{t}_2, \mathbf{t}_3]$ are normal, then η is a short exact sequence in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$.*

Proof. Notice that (a) follows from Lemma 3.1, (c) is dual to (b) and (d) follows from (b) and (c). Thus, it is enough to prove (b). (b) Let us show that f_C is monomorphism if $[\mathbf{t}_2, \mathbf{t}_3]$ is normal. Consider a morphism $h : W \rightarrow T_2$ with $W \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ and such that $f_C \circ h = 0$. By applying the functor $\mathcal{D}(W, -)$ to η , we get the exact sequence

$$\mathbb{E}^{-1}(W, F_2) \xrightarrow{\eta^{-1}} \mathcal{D}(W, T_2) \rightarrow \mathcal{D}(W, C).$$

Note that $h \in \text{Ker}(\mathcal{D}(f_C, C)) = \text{Im}(\eta^{-1})$. But $\mathbb{E}^{-1}(W, F_2) = 0$ since $W \in \mathcal{T}_3$, $F_2 \in \mathcal{F}_2$ and $[\mathbf{t}_2, \mathbf{t}_3]$ is normal. Therefore, $h = 0$ and thus f_C is a monomorphism. Finally, f_C is the kernel for g_C in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ since it is a monomorphism and a weak kernel for g_C . \square

Proposition 3.12. *For an extriangulated category \mathcal{D} with negative first extension, $\mathbf{t}_i = (\mathcal{T}_i, \mathcal{F}_i)$ in $\text{stors } \mathcal{D}$ for $i = 1, 2$ such that $[\mathbf{t}_1, \mathbf{t}_2]$ is a normal interval in $\text{stors } \mathcal{D}$ and $X, Y \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$, the following statements hold true.*

- (a) *Let $\chi : X \xrightarrow{f} Y \xrightarrow{c} Z$ be an \mathfrak{s} -conflation in \mathcal{D} . Then f has cokernel in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ and there are morphisms $X \xrightarrow{r} E \xrightarrow{h} Y$ in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ such that $f = hr$, r is an epimorphism in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ and $h = \text{Ker}(\text{Coker}(f)) =: \text{Im}(f)$ in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$.*
- (b) *Let $\kappa : K \xrightarrow{k} X \xrightarrow{f} Y$ be an \mathfrak{s} -conflation in \mathcal{D} . Then f has kernel in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ and there are morphisms $X \xrightarrow{r} E \xrightarrow{h} Y$ in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ such that $f = hr$, h is a monomorphism in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ and $r = \text{Coker}(\text{Ker}(f)) =: \text{CoIm}(f)$ in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$.*

Proof. We only prove (a) since the proof of (b) is dual. We have the \mathfrak{s} -conflation $\chi : X \xrightarrow{f} Y \xrightarrow{c} Z$ in \mathcal{D} . Since \mathcal{T}_2 is closed under cones (see Lemma 2.9) and $X, Y \in \mathcal{T}_2$, we have that $Z \in \mathcal{T}_2$. Consider the \mathfrak{s} -conflation $\eta_1 : T_1 \rightarrow Z \xrightarrow{c_1} F_1$ induced by \mathbf{t}_1 . Then $F_1 \in \mathcal{F}_1 \cap \mathcal{T}_2 = \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ since $T_1 \in \mathcal{T}_1 \subseteq \mathcal{T}_2$. Moreover, by [18, Rk. 2.22], we have the following commutative diagram in \mathcal{D}

$$\begin{array}{ccccc} X & \xrightarrow{r} & E & \longrightarrow & T_1 \\ \parallel & & \downarrow h & & \downarrow \\ X & \xrightarrow{f} & Y & \xrightarrow{c} & Z \\ & & \downarrow c_2 & & \downarrow c_1 \\ & & F_1 & \xlongequal{\quad} & F_1 \end{array}$$

where the rows and columns are \mathfrak{s} -conflations in \mathcal{D} . By Lemma 2.9 (a,b), we get that the \mathfrak{s} -conflation $\eta_2 : E \xrightarrow{h} Y \xrightarrow{c_2} F_1$ lies in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. Therefore (see Lemma 3.7) η_2 is a short exact sequence in the exact category $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$, and thus $h = \text{Ker}(c_2)$ and $c_2 = \text{Coker}(h)$. We assert that $r : X \rightarrow E$ is an epimorphism in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ and $c_2 : Y \rightarrow F_1$ is the cokernel of $f : X \rightarrow Y$ in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. Indeed, consider $W \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. Then, by applying the functor $\mathcal{D}(-, W)$ to the above diagram, we get the following exact and commutative diagram

$$\begin{array}{ccccc} \mathcal{D}(Z, W) & \xrightarrow{\mathcal{D}(c, W)} & \mathcal{D}(Y, W) & \xrightarrow{\mathcal{D}(f, W)} & \mathcal{D}(X, W) \\ \downarrow \mathcal{D}(h_1, W) & & \downarrow \mathcal{D}(h, W) & & \parallel \\ \mathcal{D}(T_1, W) & \longrightarrow & \mathcal{D}(E, W) & \xrightarrow{\mathcal{D}(r, W)} & \mathcal{D}(X, W) \end{array}$$

Since $W \in \mathcal{F}_1$, we get that $\mathcal{D}(T_1, W) = 0$ and thus $\mathcal{D}(r, W) : \mathcal{D}(E, W) \rightarrow \mathcal{D}(X, W)$ is a monomorphism. In particular, we conclude that $r : X \rightarrow E$ is an epimorphism in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. Let us show now that $c_2 = \text{Coker}(f)$ in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. Indeed, let $\lambda : Y \rightarrow W$ be such that $\lambda f = 0$. Then, by the above commutative diagram and since $\mathcal{D}(r, W) : \mathcal{D}(E, W) \rightarrow \mathcal{D}(X, W)$ is a monomorphism, we conclude that $\lambda h = 0$. Now, using that $c_2 = \text{Coker}(h)$, we get that there is a unique factorization of $\lambda : Y \rightarrow W$ through $c_2 : Y \rightarrow F_1$, proving that $c_2 = \text{Coker}(f)$ in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. \square

We recall that an additive category \mathcal{C} is **semi-abelian** if it has kernels, cokernels and the canonical morphism $\text{CoIm}(f) \rightarrow \text{Im}(f)$ is monic and epic.

Theorem 3.13. *Let \mathcal{D} be an extriangulated category with negative extension and $[\mathbf{t}_1, \mathbf{t}_2]$ be a normal interval in $\text{stors } \mathcal{D}$ such that every morphism $f : X \rightarrow Y$ in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ admits the following \mathfrak{s} -conflations in \mathcal{D} :*

$$\chi : X \xrightarrow{f} Y \xrightarrow{c} Z \text{ and } \kappa : K \xrightarrow{k} X \xrightarrow{f} Y.$$

Then, $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ is an exact and a semi-abelian category.

Proof. By Lemma 3.7, we know that the extriangulated category $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ is an exact category. By Proposition 3.12, we get that $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ has kernels and cokernels. Moreover, for $f : X \rightarrow Y$ in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$, we have the following commutative diagram in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$

$$\begin{array}{ccccc} & & E_1 & & \\ & \nearrow r_1 & & \searrow h_1 & \\ \text{Ker}(f) & \xrightarrow{k} & X & \xrightarrow{f} & Y \xrightarrow{c} \text{Coker}(f) \\ & \searrow r_2 & & \nearrow h_2 & \\ & & E_2 & & \end{array}$$

where $E_1 \xrightarrow{h_1} Y \xrightarrow{c} \text{Coker}(f)$ and $\text{Ker}(f) \xrightarrow{k} X \xrightarrow{r_2} E_2$ are short exact sequences in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$, r_1 is epic, and h_2 is monic. Therefore, since h_1 is monic and $h_1(r_1 k) = f k = 0$, we have that $r_1 k = 0$ and thus $\exists \lambda : E_2 \rightarrow E_1$ such that $\lambda r_2 = r_1$. In particular λ is epic since r_1 is epic. On the other hand, from the equality $h_1 \lambda r_2 = h_1 r_1 = f = h_2 r_2$, we get that $h_1 \lambda = h_2$ since r_2 is epic. But the equality $h_1 \lambda = h_2$ implies that λ is monic since h_2 is monic. \square

In general, an exact and semi-abelian category does not have to be quasi-abelian and neither abelian. In [27, Ex. 1] an example of this is given. However, in case the extriangulated category \mathcal{D} is triangulated, the theorem above can be strengthened as follows.

Theorem 3.14. *Let \mathcal{D} be a triangulated category and $[\mathbf{t}_1, \mathbf{t}_2]$ be a normal interval in $\text{stors } \mathcal{D}$. Then $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ is a quasi-abelian category.*

Proof. Since $[\mathbf{t}_1, \mathbf{t}_2]$ is normal, we have by Example 3.6(b) that $\Sigma \mathbf{t}_2 \leq \mathbf{t}_1 \leq \mathbf{t}_2$. Hence, by Lemma 3.2(c), it follows that $(\mathcal{H}_{[\Sigma \mathbf{t}_2, \mathbf{t}_1]}, \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]})$ is a torsion pair in the abelian category $\mathcal{H}_{[\Sigma \mathbf{t}_2, \mathbf{t}_2]}$. Now, by [26, Corollary, p.193], we have that a torsion-free class in an abelian category is quasi-abelian. Therefore, $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ is quasi-abelian. \square

3.1 Parameterizing subintervals

Let $[\mathbf{t}_1, \mathbf{t}_2]$ be an interval in $\text{stors } \mathcal{D}$. In Theorem 2.16, it is shown how the elements of $[\mathbf{t}_1, \mathbf{t}_2]$ are parameterized with the s -torsion pairs in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. The objective of this section will be to parameterize the objects of a subinterval of $[\mathbf{t}_1, \mathbf{t}_2]$ by means of the elements of s -torsion pairs in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$.

Lemma 3.15. *Let \mathcal{D} be an extriangulated category with negative first extension and let $\mathbf{t}_i = (\mathcal{T}_i, \mathcal{F}_i)$ in $\text{stors } \mathcal{D}$, for $i = 1, 2, 3$ be such that $\mathbf{t}_1 \leq \mathbf{t}_2 \leq \mathbf{t}_3$. Then, for $(\mathcal{X}, \mathcal{Y})$ in $\text{stors } \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$, the following statements hold true.*

- (a) $\mathbf{t}_2(\mathcal{Y}) \subseteq \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ and $(1 : \mathbf{t}_2)(\mathcal{Y}) \subseteq \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$.
- (b) $\mathbf{t}_2(\mathcal{X}) \subseteq \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ and $(1 : \mathbf{t}_2)(\mathcal{X}) \subseteq \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$.
- (c) If $[\mathbf{t}_2, \mathbf{t}_3]$ is normal, then $\mathbf{t}_2(\mathcal{Y}) \subseteq \mathcal{Y}$ and $(1 : \mathbf{t}_2)(\mathcal{Y}) \subseteq \mathcal{X}^{\perp-1}$.
- (d) If $[\mathbf{t}_1, \mathbf{t}_2]$ is normal, then $(1 : \mathbf{t}_2)(\mathcal{X}) \subseteq \mathcal{X}$ and $\mathbf{t}_2(\mathcal{X}) \subseteq {}^{\perp-1}\mathcal{Y}$.
- (e) the pair $\mathbf{t} = (\mathcal{T}_1 \star \mathcal{X}, \mathcal{Y} \star \mathcal{F}_3)$ is an s -torsion pair in \mathcal{D} such that $\mathbf{t}_1 \leq \mathbf{t} \leq \mathbf{t}_3$.

Proof. By Theorem 2.16, there is $(\mathcal{T}, \mathcal{F}) \in [\mathbf{t}_1, \mathbf{t}_3]$ such that $\mathcal{X} = \mathcal{T} \cap \mathcal{F}_1$ and $\mathcal{Y} = \mathcal{T}_3 \cap \mathcal{F}$. For $C \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$, consider the \mathfrak{s} -conflation $T_2 \rightarrow C \rightarrow F_2$ with $T_2 \in \mathcal{T}_2$ and $F_2 \in \mathcal{F}_2$ given by \mathbf{t}_2 . (a) If $C \in \mathcal{Y}$, then $T_2 \in \mathcal{F}_1$ and $F_2 \in \mathcal{T}_3$ by Lemma 2.9 since $C, F_2 \in \mathcal{F}_1$ and $C, T_2 \in \mathcal{T}_3$ respectively. Therefore, $T_2 \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ and $F_2 \in \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$. (b) It follows with similar arguments as (a). (c) Let $C \in \mathcal{Y}$. On the one hand, for any $X \in \mathcal{X}$, we have the exact sequence

$$\mathbb{E}^{-1}(X, F_2) \rightarrow \mathcal{D}(X, T_2) \rightarrow \mathcal{D}(X, C).$$

Here, observe that: $\mathcal{D}(X, C) = 0$ since $C \in \mathcal{Y}$; and $\mathbb{E}^{-1}(X, F_2) = 0$ since $X \in \mathcal{T} \subseteq \mathcal{T}_3$, $F_2 \in \mathcal{F}_2$ and $[\mathbf{t}_2, \mathbf{t}_3]$ is normal. And thus, $\mathbf{t}_2(C) = T_2 \in \mathcal{Y}$ by Proposition 2.8.

On the other hand, note that $F_2 \in \mathcal{X}^{\perp-1}$ by Lemma 2.9 since $F_2 \in \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]} \subseteq \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ (see (a) and Lemma 3.1), $C \in \mathcal{Y}$ and $T_2 \in \mathcal{Y}$. (d) Let $C \in \mathcal{X}$. On the one hand, for any $Y \in \mathcal{Y}$, we have the exact sequence

$$\mathbb{E}^{-1}(T_2, Y) \rightarrow \mathcal{D}(F_2, Y) \rightarrow \mathcal{D}(C, Y).$$

Here, observe that: $\mathcal{D}(C, Y) = 0$ since $C \in \mathcal{X}$; and $\mathbb{E}^{-1}(T_2, Y) = 0$ since $Y \in \mathcal{F} \subseteq \mathcal{F}_1$, $T_2 \in \mathcal{T}_2$ and $[\mathbf{t}_1, \mathbf{t}_2]$ is normal. And thus, $(1 : \mathbf{t}_2)(C) = F_2 \in \mathcal{X}$ by Proposition 2.8.

On the other hand, note that $T_2 \in {}^{\perp-1}\mathcal{Y}$ by Lemma 2.9 since $T_2 \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} \subseteq \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ (see (b) and Lemma 3.1), $C \in \mathcal{X}$ and $F_2 \in \mathcal{X}$. (e) It follows from Theorem 2.16. \square

Corollary 3.16. *Let \mathcal{D} be an extriangulated category with negative first extension and let $\mathbf{t}_i = (\mathcal{T}_i, \mathcal{F}_i)$ in $\text{stors } \mathcal{D}$, for $i = 1, 2, 3$ be such that $\mathbf{t}_1 \leq \mathbf{t}_2 \leq \mathbf{t}_3$. Then, for $(\mathcal{X}, \mathcal{Y})$ in $\text{stors } \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ and $\mathbf{t} = (\mathcal{T}, \mathcal{F}) := (\mathcal{T}_1 \star \mathcal{X}, \mathcal{Y} \star \mathcal{F}_3) \in [\mathbf{t}_1, \mathbf{t}_3]$, see Lemma 3.15 (e), the following statements hold true.*

- (a) $\mathbf{t} \in [\mathbf{t}_1, \mathbf{t}_2] \Leftrightarrow \mathcal{X} \subseteq \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} \Leftrightarrow \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]} \subseteq \mathcal{Y}$.
- (b) $\mathbf{t} \in [\mathbf{t}_2, \mathbf{t}_3] \Leftrightarrow \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} \subseteq \mathcal{X} \Leftrightarrow \mathcal{Y} \subseteq \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$.

Proof. (a) By Lemma 3.2 (c), we have that $(\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}, \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}) \in \text{stors } \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ and since $(\mathcal{X}, \mathcal{Y})$ in $\text{stors } \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$, we get by Proposition 2.9 (d) that $\mathcal{X} \subseteq \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} \Leftrightarrow \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]} \subseteq \mathcal{Y}$. Let now $\mathcal{T}_1 \subseteq \mathcal{T}_1 \star \mathcal{X} \subseteq \mathcal{T}_2$. Hence $\mathcal{X} \subseteq \mathcal{T}_2$. Moreover $\mathcal{X} \subseteq \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]} = \mathcal{F}_1 \cap \mathcal{T}_3 \subseteq \mathcal{F}_1$ and thus $\mathcal{X} \subseteq \mathcal{F}_1 \cap \mathcal{T}_2 = \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. Finally, assume that $\mathcal{X} \subseteq \mathcal{F}_1 \cap \mathcal{T}_2$. Hence $\mathcal{T}_1 \star \mathcal{X} \subseteq \mathcal{T}_2$, and so $\mathbf{t} \in [\mathbf{t}_1, \mathbf{t}_2]$. (b) The equivalence $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} \subseteq \mathcal{X} \Leftrightarrow \mathcal{Y} \subseteq \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$ follows from Lemma 3.2 (c) and Proposition 2.9 (d). Let $\mathcal{F}_2 \supseteq \mathcal{Y} \star \mathcal{F}_3 \supseteq \mathcal{F}_3$. Hence $\mathcal{Y} \subseteq \mathcal{F}_2$. Moreover $\mathcal{Y} \subseteq \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]} = \mathcal{F}_1 \cap \mathcal{T}_3 \subseteq \mathcal{T}_3$ and thus $\mathcal{Y} \subseteq \mathcal{F}_2 \cap \mathcal{T}_3 = \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$. Finally, let $\mathcal{Y} \subseteq \mathcal{F}_2 \cap \mathcal{T}_3$. Hence $\mathcal{F}_3 \subseteq \mathcal{Y} \star \mathcal{F}_3 \subseteq \mathcal{F}_2$, and so by Proposition 2.9 (d) we have $\mathbf{t} \in [\mathbf{t}_2, \mathbf{t}_3]$. \square

The following result is a generalization of Theorem 2.16 and it is very useful to parametrize torsion pairs in quasi-abelian categories, see Theorem 3.19. In order to do that, we start with the following notion.

Definition 3.17. Let \mathcal{D} be an extriangulated category with negative first extension and $\mathbf{x} \leq \mathbf{y} \leq \mathbf{z}$ in $\text{stors } \mathcal{D}$. We consider the classes of s -torsion pairs:

- (a) $\text{stors } \mathcal{H}_{[\mathbf{x}, \mathbf{y}, \mathbf{z}]}^+ := \{(\mathcal{T}, \mathcal{F}) \in \text{stors } \mathcal{H}_{[\mathbf{x}, \mathbf{z}]} : \mathcal{T} \subseteq \mathcal{H}_{[\mathbf{x}, \mathbf{y}]}\}$.
- (b) $\text{stors } \mathcal{H}_{[\mathbf{x}, \mathbf{y}, \mathbf{z}]}^- := \{(\mathcal{T}, \mathcal{F}) \in \text{stors } \mathcal{H}_{[\mathbf{x}, \mathbf{z}]} : \mathcal{H}_{[\mathbf{x}, \mathbf{y}]} \subseteq \mathcal{T}\}$.

Theorem 3.18. *For an extriangulated category \mathcal{D} with negative first extension and $\mathbf{t}_i = (\mathcal{T}_i, \mathcal{F}_i)$ in $\text{stors } \mathcal{D}$ for $i = 1, 2, 3, 4$ such that $\mathbf{t}_1 \leq \mathbf{t}_2 \leq \mathbf{t}_3 \leq \mathbf{t}_4$, the following statements hold.*

- (a) *The map $\Phi_r : [\mathbf{t}_2, \mathbf{t}_3] \rightarrow \text{stors } \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3, \mathbf{t}_4]}^+$, $(\mathcal{T}, \mathcal{F}) \mapsto (\mathcal{T} \cap \mathcal{F}_2, \mathcal{T}_4 \cap \mathcal{F})$, is an isomorphism of posets with inverse $(\mathcal{X}, \mathcal{Y}) \mapsto (\mathcal{T}_2 \star \mathcal{X}, \mathcal{Y} \star \mathcal{F}_4)$.*
- (b) *The map $\Phi_\ell : [\mathbf{t}_2, \mathbf{t}_3] \rightarrow \text{stors } \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2, \mathbf{t}_3]}^-$, $(\mathcal{T}, \mathcal{F}) \mapsto (\mathcal{T} \cap \mathcal{F}_1, \mathcal{T}_3 \cap \mathcal{F})$, is an isomorphism of posets with inverse $(\mathcal{X}, \mathcal{Y}) \mapsto (\mathcal{T}_1 \star \mathcal{X}, \mathcal{Y} \star \mathcal{F}_3)$.*
- (c) *The map $\varepsilon : \text{stors } \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3, \mathbf{t}_4]}^+ \rightarrow \text{stors } \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2, \mathbf{t}_3]}^-$ given by*

$$(\mathcal{X}, \mathcal{Y}) \mapsto ((\mathcal{T}_2 \star \mathcal{X}) \cap \mathcal{F}_1, (\mathcal{Y} \star \mathcal{F}_4) \cap \mathcal{T}_3),$$

is an isomorphism of posets with inverse

$$(\mathcal{T}, \mathcal{F}) \mapsto ((\mathcal{T}_1 \star \mathcal{T}) \cap \mathcal{F}_2, (\mathcal{F} \star \mathcal{F}_3) \cap \mathcal{T}_4).$$

Proof. We prove only (a) since (b) is similar and (c) follows from (a) and (b). From Theorem 2.16, we have the isomorphism of posets

$$\Phi : [\mathbf{t}_2, \mathbf{t}_4] \rightarrow \text{stors } \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_4]}, \quad (\mathcal{T}, \mathcal{F}) \mapsto (\mathcal{T} \cap \mathcal{F}_2, \mathcal{T}_4 \cap \mathcal{F}),$$

with inverse

$$\Psi : \text{stors } \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_4]} \rightarrow [\mathbf{t}_2, \mathbf{t}_4], \quad (\mathcal{X}, \mathcal{Y}) \mapsto (\mathcal{T}_2 \star \mathcal{X}, \mathcal{Y} \star \mathcal{F}_4).$$

Since $[\mathbf{t}_2, \mathbf{t}_3] \subseteq [\mathbf{t}_2, \mathbf{t}_4]$, by Corollary 3.16, we have that

$$\Psi^{-1}[\mathbf{t}_2, \mathbf{t}_3] = \{(\mathcal{X}, \mathcal{Y}) \in \text{stors } \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_4]} \mid \mathcal{X} \subseteq \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}\} = \text{stors } \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}^+.$$

Therefore, by restricting Φ and Ψ , we get the desired isomorphism. \square

As an application, we have the following result parametrizing the torsion pairs of a quasi-abelian category. We recall that in an exact category \mathcal{D} , see [1, Prop. 2.6], the class $\text{stors } \mathcal{D}$ is just the class of all the torsion pairs in \mathcal{D} .

Theorem 3.19. *Let \mathcal{E} be a quasi-abelian category, and let $\mathcal{L}_{\mathcal{E}}$ and $\mathcal{R}_{\mathcal{E}}$ be the associated abelian categories of \mathcal{E} (see Example 3.3). Then, the following posets are isomorphic.*

- (a) $\text{stors } \mathcal{E}$.
- (b) $\{(\mathcal{X}, \mathcal{Y}) \in \text{stors } \mathcal{L}_{\mathcal{E}} \mid \mathcal{Y} \subseteq \mathcal{E}\}$.
- (c) $\{(\mathcal{X}, \mathcal{Y}) \in \text{stors } \mathcal{R}_{\mathcal{E}} \mid \mathcal{X} \subseteq \mathcal{E}\}$.

Proof. Let \mathcal{D} be the triangulated category associated to \mathcal{E} . Recall that there are $\mathbf{t}_r, \mathbf{t}_\ell \in \text{stors } \mathcal{D}$ such that $\Sigma \mathbf{t}_r \leq \mathbf{t}_\ell \leq \mathbf{t}_r$ and $\mathcal{E} \cong \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$ (see Example 2.14). Then, the sought isomorphisms are given by Theorem 3.18 (under the setting: $\mathbf{t}_1 := \Sigma \mathbf{t}_\ell$, $\mathbf{t}_2 := \Sigma \mathbf{t}_r$, $\mathbf{t}_3 := \mathbf{t}_\ell$ and $\mathbf{t}_4 := \mathbf{t}_r$), together with the isomorphism $[\mathbf{t}_2, \mathbf{t}_3] \cong \text{stors } \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$ given by Theorem 2.16. \square

4 The extended heart of a t -structure

The objective of this section is to study a special kind of extensions of hearts in triangulated categories. The advantage that triangulated categories have over the rest of the extriangulated categories with negative first extensions is that the functor \mathbb{E}^{-1} coincides with $\mathcal{D}(-, \Sigma^{-1}-)$. So we have that $\mathcal{X}^{\perp_{-1}} = (\Sigma \mathcal{X})^{\perp_0}$ and ${}^{\perp_0}(\Sigma^{-1} \mathcal{X}) = {}^{\perp_{-1}} \mathcal{X}$. This fact is very useful when studying t -structures. We remind the reader that a pair of classes $(\mathcal{U}, \mathcal{W})$ is a t -structure if, and only if, $(\mathcal{U}, \Sigma^{-1} \mathcal{W})$ is an s -torsion pair, or equivalently $(\Sigma \mathcal{U}, \mathcal{W})$ is an s -torsion pair.

Definition 4.1. Let \mathcal{D} be a triangulated category and $\mathbf{u} = (\mathcal{U}, \mathcal{W})$ be a t -structure in \mathcal{D} . For $\mathbf{u}_1 := (\Sigma \mathcal{U}, \mathcal{W})$, $\mathbf{u}_2 := (\mathcal{U}, \Sigma^{-1} \mathcal{W})$ and $\mathbf{u}_3 := (\Sigma^{-1} \mathcal{U}, \Sigma^{-2} \mathcal{W})$, we have $\mathbf{u}_1 \leq \mathbf{u}_2 \leq \mathbf{u}_3$ in $\text{stors } \mathcal{D}$. Notice that $\mathcal{H} := \mathcal{H}_{[\mathbf{u}_1, \mathbf{u}_2]} = \mathcal{W} \cap \mathcal{U}$ is the heart $\mathcal{H}_{\mathbf{u}}$ of \mathbf{u} . The class $\mathcal{C} = \mathcal{C}_{\mathbf{u}} := \mathcal{H}_{[\mathbf{u}_1, \mathbf{u}_3]} = \mathcal{W} \cap \Sigma^{-1} \mathcal{U}$ is the **extended heart**

of \mathbf{u} . Recall that, for any $C \in \mathcal{D}$, we have the \mathfrak{s} -conflation $\tau_{\mathcal{U}}^{\leq}(C) \xrightarrow{f_{\mathcal{C}}} C \xrightarrow{g_{\mathcal{C}}} \tau_{\mathcal{U}}^{\geq}(C)$ given by \mathbf{u}_2 . Thus, the torsion pair \mathbf{u}_2 induces the torsion functor $\tau_{\mathcal{U}}^{\leq} : \mathcal{D} \rightarrow \mathcal{U}$ and the torsion-free functor $\tau_{\mathcal{U}}^{\geq} : \mathcal{D} \rightarrow \Sigma^{-1}\mathcal{W}$.

We recall that the heart $\mathcal{H} = \mathcal{H}_{\mathbf{u}}$ is an abelian category. On the other hand, since the extended heart $\mathcal{C} = \mathcal{C}_{\mathbf{u}}$ is closed under extensions in the triangulated category \mathcal{D} , we have that \mathcal{C} is an extriangulated category with negative first extension. In what follows, we show some basic properties of the extended heart \mathcal{C} .

Proposition 4.2. *For a triangulated category \mathcal{D} , a t -structure $\mathbf{u} = (\mathcal{U}, \mathcal{W})$ in \mathcal{D} , the heart $\mathcal{H} = \mathcal{H}_{\mathbf{u}}$ and the extended heart $\mathcal{C} = \mathcal{C}_{\mathbf{u}}$, the following statements hold true.*

- (a) $\mathcal{C} = \mathcal{H} \star \Sigma^{-1}\mathcal{H}$, $\tau_{\mathcal{U}}^{\leq}(\mathcal{C}) \subseteq \mathcal{H}$ and $\tau_{\mathcal{U}}^{\geq}(\mathcal{C}) \subseteq \Sigma^{-1}\mathcal{H}$.
- (b) If $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ is a torsion pair in \mathcal{H} , then $(\mathcal{T}, \mathcal{F} \star \Sigma^{-1}\mathcal{H})$, $(\mathcal{H} \star \Sigma^{-1}\mathcal{T}, \Sigma^{-1}\mathcal{F})$ and $(\mathcal{H}, \Sigma^{-1}\mathcal{H})$ are s -torsion pairs in \mathcal{C} .
- (c) For any $C \in \mathcal{C}$, the \mathfrak{s} -conflation $\tau_{\mathcal{U}}^{\leq}(C) \xrightarrow{f_{\mathcal{C}}} C \xrightarrow{g_{\mathcal{C}}} \tau_{\mathcal{U}}^{\geq}(C)$ given by \mathbf{u}_2 (see Definition 4.1) is a short exact sequence in \mathcal{C} .
- (d) If $\mathcal{H} \neq 0$, then the extriangulated category \mathcal{C} is not abelian.

Proof. Observe that $\mathcal{H}_{[\mathbf{u}_2, \mathbf{u}_3]} = \Sigma^{-1}\mathcal{H}$, $\mathcal{H}_{[\mathbf{u}_1, \mathbf{u}_2]} = \mathcal{H}$ and $\mathcal{C} = \mathcal{H}_{[\mathbf{u}_1, \mathbf{u}_3]}$. Then: (a) follows from Lemma 3.1, (b) from Lemma 3.2 and (c) from Proposition 3.11. (d) Let $0 \neq H \in \mathcal{H}$. Consider the canonical triangle $\Sigma^{-1}H \rightarrow 0 \rightarrow H \rightarrow H$ in the triangulated category \mathcal{D} . Then, by (a) we get the \mathfrak{s} -conflation $\eta : \Sigma^{-1}H \rightarrow 0 \rightarrow H$ in \mathcal{C} . Suppose that \mathcal{C} is abelian. Then η is a short exact sequence in \mathcal{C} and thus $H = 0$, which is a contradiction proving that \mathcal{C} is not abelian. \square

Definition 4.3. For a triangulated category \mathcal{D} , a t -structure $\mathbf{u} = (\mathcal{U}, \mathcal{W})$ in \mathcal{D} , and the extended heart $\mathcal{C} = \mathcal{C}_{\mathbf{u}}$ (see Definition 4.1), we introduce the following classes of t -structures in \mathcal{D} .

- (a) $\mathbb{T}_{\mathbf{u}}$ is the class of all the t -structures $(\mathcal{U}', \mathcal{W}')$ in \mathcal{D} such that $\Sigma\mathcal{U} \subseteq \mathcal{U}' \subseteq \Sigma^{-1}\mathcal{U}$.
- (b) $\mathbb{T}_{\mathbf{u}}^+ := \{(\mathcal{U}', \mathcal{W}') \in \mathbb{T}_{\mathbf{u}} : \mathcal{U}' \subseteq \mathcal{U}\}$.
- (c) $\mathbb{T}_{\mathbf{u}}^- := \{(\mathcal{U}', \mathcal{W}') \in \mathbb{T}_{\mathbf{u}} : \mathcal{U} \subseteq \mathcal{U}'\}$.

Lemma 4.4. *For a triangulated category \mathcal{D} , a t -structure $\mathbf{u} = (\mathcal{U}, \mathcal{W})$ in \mathcal{D} , the extended heart $\mathcal{C} = \mathcal{C}_{\mathbf{u}}$ (see Definition 4.1) and an s -torsion pair $(\mathcal{X}, \mathcal{Y})$ in \mathcal{C} , the following statements hold true.*

- (a) $\tau_{\mathcal{U}}^{\leq}(\mathcal{Y}) \subseteq \mathcal{Y}$, $\tau_{\mathcal{U}}^{\geq}(\mathcal{Y}) \subseteq \Sigma\mathcal{Y}$, $\tau_{\mathcal{U}}^{\geq}(\mathcal{X}) \subseteq \mathcal{X}$ and $\tau_{\mathcal{U}}^{\leq}(\mathcal{X}) \subseteq \Sigma^{-1}\mathcal{X}$.
- (b) $(\Sigma\mathcal{U} \star \mathcal{X}, \Sigma\mathcal{Y} \star \Sigma^{-1}\mathcal{W}) \in \mathbb{T}_{\mathbf{u}}$.

Proof. It follows from Lemma 3.15 since the intervals $[\mathbf{u}_1, \mathbf{u}_2]$ and $[\mathbf{u}_2, \mathbf{u}_3]$, considered in Definition 4.1, are normal. \square

Definition 4.5. Let $\mathbf{u} = (\mathcal{U}, \mathcal{W})$ be a t -structure in the triangulated category \mathcal{D} , and let $\mathcal{C} = \mathcal{C}_{\mathbf{u}}$ be the extended heart of \mathbf{u} (see Definition 4.1). For an s -torsion pair $\mathbf{x} = (\mathcal{X}, \mathcal{Y})$ in \mathcal{C} , we define the **extended tilt** $\mathbf{u}_{\mathbf{x}} = \mathbf{E}_{\mathbf{u}}(\mathbf{x})$ of \mathbf{u} with respect to \mathbf{x} as follows $\mathbf{u}_{\mathbf{x}} := (\mathcal{U}_{\mathbf{x}}, \mathcal{W}_{\mathbf{x}}) := (\Sigma\mathcal{U} \star \mathcal{X}, \Sigma\mathcal{Y} \star \Sigma^{-1}\mathcal{W})$. Thus, by Lemma 4.4(b), we get the map $\mathbf{E}_{\mathbf{u}} : \text{stors}\mathcal{C} \rightarrow \mathbb{T}_{\mathbf{u}}$, $(\mathcal{X}, \mathcal{Y}) \mapsto \mathbf{E}_{\mathbf{u}}(\mathcal{X}, \mathcal{Y})$ which is a morphism of posets.

Corollary 4.6. Let $\mathbf{u} = (\mathcal{U}, \mathcal{W})$ be a t -structure in the triangulated category \mathcal{D} , and let $\mathcal{C} = \mathcal{C}_{\mathbf{u}}$ and $\mathcal{H} = \mathcal{H}_{\mathbf{u}}$ be, respectively, the extended heart and the heart of \mathbf{u} . Then, for an s -torsion pair $\mathbf{x} = (\mathcal{X}, \mathcal{Y})$ in \mathcal{C} , the extended tilt $\mathbf{u}_{\mathbf{x}} := (\mathcal{U}_{\mathbf{x}}, \mathcal{W}_{\mathbf{x}})$, satisfies the following conditions.

- (a) $\Sigma\mathcal{U} \subseteq \mathcal{U}_{\mathbf{x}} \subseteq \mathcal{U} \Leftrightarrow \mathcal{X} \subseteq \mathcal{H} \Leftrightarrow \Sigma^{-1}\mathcal{H} \subseteq \mathcal{Y}$.
- (b) $\mathcal{U} \subseteq \mathcal{U}_{\mathbf{x}} \subseteq \Sigma^{-1}\mathcal{U} \Leftrightarrow \mathcal{H} \subseteq \mathcal{X} \Leftrightarrow \mathcal{Y} \subseteq \Sigma^{-1}\mathcal{H}$.

Proof. It follows from Corollary 3.16. \square

In the following result, we show that the extended tilt $\mathbf{E}_{\mathbf{u}}$ gives an isomorphism of posets between the s -torsion pairs in the extended heart $\mathcal{C} = \mathcal{C}_{\mathbf{u}}$ and some special class of t -structures. Moreover, such a bijection is well restricted on certain subposets described in the items (a) and (b) below.

Corollary 4.7. Let $\mathbf{u} = (\mathcal{U}, \mathcal{W})$ be a t -structure in the triangulated category \mathcal{D} with heart $\mathcal{H} = \mathcal{H}_{\mathbf{u}}$ and extended heart $\mathcal{C} = \mathcal{C}_{\mathbf{u}}$, and let $\mathbb{T}_{\mathbf{u}}^{\pm}$ be the classes described in Definition 4.3. Consider the following classes

- (a) $\text{stors}\mathcal{C}^+ := \{(\mathcal{X}, \mathcal{Y}) \in \text{stors}\mathcal{C} : \mathcal{X} \subseteq \mathcal{H}\}$;
- (b) $\text{stors}\mathcal{C}^- := \{(\mathcal{X}, \mathcal{Y}) \in \text{stors}\mathcal{C} : \mathcal{H} \subseteq \mathcal{X}\}$,

Then, the extended tilt $\mathbf{E}_{\mathbf{u}} : \text{stors}\mathcal{C} \rightarrow \mathbb{T}_{\mathbf{u}}$ is an isomorphism of posets whose inverse is given by $(\mathcal{U}', \mathcal{W}') \mapsto (\mathcal{U}' \cap \mathcal{W}, \Sigma^{-1}(\mathcal{W}' \cap \mathcal{U}))$. Moreover, $\mathbf{E}_{\mathbf{u}}(\text{stors}\mathcal{C}^+) = \mathbb{T}_{\mathbf{u}}^+$ and $\mathbf{E}_{\mathbf{u}}(\text{stors}\mathcal{C}^-) = \mathbb{T}_{\mathbf{u}}^-$.

Proof. It follows from Theorems 2.16 and 3.18. \square

Now, we want to relate the extended tilt $\mathbf{E}_{\mathbf{u}}$ with the right and left Happel-Reiten-Smalø tilt in the heart of a t -structure \mathbf{u} .

Definition 4.8. [13, 30] Let $\mathbf{u} = (\mathcal{U}, \mathcal{W})$ be a t -structure in the triangulated category \mathcal{D} , with heart $\mathcal{H} = \mathcal{H}_{\mathbf{u}} = \mathcal{U} \cap \mathcal{W}$.

- (a) The **right HRS-tilt** of $\mathbf{t} \in \text{stors}\mathcal{H}$, with respect to \mathbf{u} , is $\mathbf{R}_{\mathbf{u}}(\mathbf{t})$, where

$$\mathbf{R}_{\mathbf{u}} : \text{stors}\mathcal{H} \rightarrow \mathbb{T}_{\mathbf{u}}^+, \mathbf{t} = (\mathcal{T}, \mathcal{F}) \mapsto (\Sigma\mathcal{U} \star \mathcal{T}, \Sigma\mathcal{F} \star \mathcal{W}).$$

(b) The **left HRS-tilt** of $\mathbf{t} \in \text{stors } \mathcal{H}$, with respect to \mathbf{u} , is $L_{\mathbf{u}}(\mathbf{t})$, where

$$L_{\mathbf{u}} : \text{stors } \mathcal{H} \rightarrow \mathbb{T}_{\mathbf{u}}^-, \mathbf{t} = (\mathcal{T}, \mathcal{F}) \mapsto (\mathcal{U} \star \Sigma^{-1} \mathcal{T}, \mathcal{F} \star \Sigma^{-1} \mathcal{W}).$$

Notice that $\Sigma^{-1} : \mathbb{T}_{\mathbf{u}}^+ \rightarrow \mathbb{T}_{\mathbf{u}}^-$ is a bijection and $\Sigma^{-1} R_{\mathbf{u}} = L_{\mathbf{u}}$.

Now, for a given t -structure \mathbf{u} , we describe the relationship between the extended tilt $E_{\mathbf{u}}$ and the right HRS-tilt $R_{\mathbf{u}}$.

Theorem 4.9. *Let $\mathbf{u} = (\mathcal{U}, \mathcal{W})$ be a t -structure in \mathcal{D} with heart \mathcal{H} and extended heart \mathcal{C} , and let $E_{\mathbf{u}}^+ : \text{stors } \mathcal{C}^+ \rightarrow \mathbb{T}_{\mathbf{u}}^+$ be the isomorphism of posets given by the restriction of the extended tilt $E_{\mathbf{u}}$ on $\text{stors } \mathcal{C}^+$ (see Corollary 4.7). Then, the following statements hold true.*

(a) *The map $\mu^+ : \text{stors } \mathcal{H} \rightarrow \text{stors } \mathcal{C}^+$, $(\mathcal{T}, \mathcal{F}) \mapsto (\mathcal{T}, \mathcal{F} \star \Sigma^{-1} \mathcal{H})$, is an isomorphism of posets whose inverse is given by $\lambda^+ : \text{stors } \mathcal{C}^+ \rightarrow \text{stors } \mathcal{H}$, $(\mathcal{X}, \mathcal{Y}) \mapsto (\mathcal{X}, \mathcal{Y} \cap \mathcal{H})$. Furthermore $\mathcal{Y} = (\mathcal{Y} \cap \mathcal{H}) \star \Sigma^{-1} \mathcal{H}$ and $\mathcal{F} = (\mathcal{F} \star \Sigma^{-1} \mathcal{H}) \cap \mathcal{H}$.*

(b) $E_{\mathbf{u}}^+ \circ \mu^+ = R_{\mathbf{u}}$.

Proof. (a) Notice that the map μ^+ is well defined (see Proposition 4.2 (b)). Let us prove that λ^+ is well defined. Indeed, for $(\mathcal{X}, \mathcal{Y}) \in \text{stors } \mathcal{C}^+$, we need to show that $\mathcal{H} = \mathcal{X} \star (\mathcal{Y} \cap \mathcal{H})$. However, this equality can be obtained from Lemma 3.9 (a). Finally, the equalities in item (a) can be obtained from Proposition 2.8. (b) Let $(\mathcal{T}, \mathcal{F}) \in \text{stors } \mathcal{H}$. Then $E_{\mathbf{u}}^+ \mu^+(\mathcal{T}, \mathcal{F}) = (\Sigma \mathcal{U} \star \mathcal{T}, (\Sigma \mathcal{F} \star \mathcal{H}) \star \Sigma^{-1} \mathcal{W})$; and thus $E_{\mathbf{u}}^+ \mu^+(\mathcal{T}, \mathcal{F}) = R_{\mathbf{u}}(\mathcal{T}, \mathcal{F})$ since $(\Sigma \mathcal{F} \star \mathcal{H}) \star \Sigma^{-1} \mathcal{W} = \Sigma \mathcal{F} \star \mathcal{W}$. \square

Now, for a given t -structure \mathbf{u} , we describe the relationship between the extended tilt $E_{\mathbf{u}}$ and the left HRS-tilt $L_{\mathbf{u}}$.

Theorem 4.10. *Let $\mathbf{u} = (\mathcal{U}, \mathcal{W})$ be a t -structure in \mathcal{D} with heart \mathcal{H} and extended heart \mathcal{C} , and let $E_{\mathbf{u}}^- : \text{stors } \mathcal{C}^- \rightarrow \mathbb{T}_{\mathbf{u}}^-$ be the isomorphism of posets given by the restriction of the extended tilt $E_{\mathbf{u}}$ on $\text{stors } \mathcal{C}^-$ (see Corollary 4.7). Then, the following statements hold true.*

(a) *The map $\mu^- : \text{stors } \Sigma^{-1} \mathcal{H} \rightarrow \text{stors } \mathcal{C}^-$, $(\mathcal{T}, \mathcal{F}) \mapsto (\mathcal{H} \star \mathcal{T}, \mathcal{F})$, is an isomorphism of posets whose inverse is given by*

$$\lambda^- : \text{stors } \mathcal{C}^- \rightarrow \text{stors } \Sigma^{-1} \mathcal{H}, (\mathcal{X}, \mathcal{Y}) \mapsto (\mathcal{X} \cap \Sigma^{-1} \mathcal{H}, \mathcal{Y}).$$

Furthermore, $\mathcal{X} = \mathcal{H} \star (\mathcal{X} \cap \Sigma^{-1} \mathcal{H})$, $\mathcal{T} = (\mathcal{H} \star \mathcal{T}) \cap \Sigma^{-1} \mathcal{H}$ and $\text{stors } \mathcal{C}^- = \{(\mathcal{X}, \mathcal{Y}) \in \text{stors } \mathcal{C} : \mathcal{Y} \subseteq \Sigma^{-1} \mathcal{H}\}$.

(b) *We have the following commutative diagram of posets*

$$\begin{array}{ccc} \text{stors } \mathcal{C}^- & \xrightarrow{E_{\mathbf{u}}^-} & \mathbb{T}_{\mathbf{u}}^- \\ \mu^- \uparrow & & \uparrow L_{\mathbf{u}} \\ \text{stors } \Sigma^{-1} \mathcal{H} & \xleftarrow{\Sigma^{-1}} & \text{stors } \mathcal{H}, \end{array}$$

where each arrow is a poset isomorphism. That is, $E_{\mathbf{u}}^- \circ \mu^- \circ \Sigma^{-1} = L_{\mathbf{u}}$.

Proof. (a) From Corollary 4.6 (b), we get that

$$\mathbf{stors}\mathcal{C}^- = \{(\mathcal{X}, \mathcal{Y}) \in \mathbf{stors}\mathcal{C} : \mathcal{Y} \subseteq \Sigma^{-1}\mathcal{H}\}.$$

The above equality will be a strong point to prove the item (a). Indeed, from Lemma 3.2 (b), we have that the map μ^- is well defined. Let us prove that λ^- is also well defined. Indeed, for $(\mathcal{X}, \mathcal{Y}) \in \mathbf{stors}\mathcal{C}^-$, we need to show that $\Sigma^{-1}\mathcal{H} = (\mathcal{X} \cap \Sigma^{-1}\mathcal{H}) * \mathcal{Y}$. However, this equality can be obtained from Lemma 3.9 (b). Finally, the equalities in item (a) can be obtained from Proposition 2.8. (b) Let $(\mathcal{T}, \mathcal{F}) \in \mathbf{stors}\mathcal{H}$. Then

$$\mathbf{E}_{\mathbf{u}}^- \circ \mu^- \circ \Sigma^{-1}(\mathcal{T}, \mathcal{F}) = (\Sigma\mathcal{U} \star (\mathcal{H} * \Sigma^{-1}\mathcal{T}), \mathcal{F} * \Sigma^{-1}\mathcal{W});$$

and thus $\mathbf{E}_{\mathbf{u}}^- \circ \mu^- \circ \Sigma^{-1}(\mathcal{T}, \mathcal{F}) = \mathbf{L}_u(\mathcal{T}, \mathcal{F})$ since $\Sigma\mathcal{U} \star (\mathcal{H} * \Sigma^{-1}\mathcal{T}) = \mathcal{U} \star \Sigma^{-1}\mathcal{T}$. \square

5 Extended tilting objects

Let us begin by introducing the following notation inspired in the abelian case, see [20, 6].

Definition 5.1. Let \mathcal{D} be an extriangulated category. We say that $\mathcal{S} \subseteq \mathcal{D}$ is a **cogenerating class** in \mathcal{D} if, for each $X \in \mathcal{D}$, there is an \mathbb{E} -triangle $X \rightarrow S \rightarrow D$ with $S \in \mathcal{S}$.

Definition 5.2. Let \mathcal{D} be an extriangulated category and $V \in \mathcal{D}$.

- (a) We say that $X \in \mathcal{D}$ is **V -generated** if either $X = 0$ or there exists an \mathbb{E} -triangle $A \rightarrow \coprod_{\alpha} V \xrightarrow{f} X$ for some set α and $f \neq 0$. The class of all the V -generated objects in \mathcal{D} is denoted by $\text{Gen}_{\mathcal{D}}(V)$ or $\text{Gen}(V)$. If $\text{Gen}(V) = \mathcal{D}$ we say that V is a **generator** in \mathcal{D} .
- (b) We say that $X \in \mathcal{D}$ is **V -presented** if either $X = 0$ or there exists an \mathbb{E} -triangle $A \rightarrow \coprod_{\alpha} V \xrightarrow{f} X$ for some set α such that $f \neq 0$ and A is V -generated. The class of all the V -presented objects in \mathcal{D} is denoted by $\text{Pres}_{\mathcal{D}}(V)$ or $\text{Pres}(V)$.
- (c) $\text{Add}(V)$ is the class of all the objects in \mathcal{D} which are direct summands of coproducts of the form $\coprod_{\alpha} V$ where α is a set.

In the previous setting, we could have that $\text{Pres}(V) = \text{Gen}(V) = \mathcal{D}$ even if \mathcal{D} does not have arbitrary coproducts. For example, consider $\mathcal{D} = \text{mod}(\Lambda)$ the category of finitely generated left Λ -modules for some Artin algebra Λ . In this case $\text{Pres}(\Lambda) = \text{Gen}(\Lambda) = \text{mod}(\Lambda)$ but $\text{mod}(\Lambda)$ does not have arbitrary coproducts.

The following definition is inspired from [23, Def. 6.1].

Definition 5.3. Let \mathcal{D} be an extriangulated category with negative first extension. We say that $V \in \mathcal{D}$ is an **extended tilting object** if the following statements hold true.

- (T1) $\mathbf{t}(V) := (\text{Gen}(V), V^{\perp_0}) \in \mathbf{stors} \mathcal{D}$.
- (T2) $\text{Gen}(V) = \text{Pres}(V)$ and $\text{Gen}(V) = V^{\perp_1}$.
- (T3) $\text{Gen}(V)$ is a cogenerating class in \mathcal{D} .

In such case, we say that $\mathbf{t}(V)$ is the **extended tilting s -torsion pair** associated to V .

Let us show, in the following examples, that the definition of extended tilting object agrees with the usual notion of tilting object in those settings.

Example 5.4. Let \mathcal{D} be an extriangulated category with negative first extension.

- (a) If \mathcal{D} is abelian, then $V \in \mathcal{D}$ is a tilting object as in [23, Def. 6.1] if, and only if, V is extended tilting.
- (b) Let $\mathcal{D} = \text{Mod}(R)$ be the category of left R -modules, for a ring R . Then, by [23, Cor. 2.6 and 6.9] and [7, Thm. 3.11], we have that: $V \in \mathcal{D}$ is extended tilting if, and only if, V is 1-tilting in the sense of Angeleri-Coelho (see [2]).
- (c) Let \mathcal{D} be a triangulated category. Then, $V \in \mathcal{D}$ is extended tilting if, and only if, V is a zero object in \mathcal{D} . Indeed, let $V \in \mathcal{D}$ be extended tilting in \mathcal{D} . In this case, we have that $V^{\perp_1} = \Sigma^{-1}(V^{\perp_0})$. But $\Sigma^{-1}(V^{\perp_0}) \subseteq V^{\perp_0}$ since $(\text{Gen}(V), \Sigma(V^{\perp_0}))$ is a t -structure in \mathcal{D} (see Remark 2.12). Thus, $\text{Hom}_{\mathcal{D}}(V, V) = 0$ since $V \in V^{\perp_1} \subseteq V^{\perp_0}$.
- (d) Let \mathcal{D} be with enough projectives and injectives (i.e. every $D \in \mathcal{D}$ admits \mathfrak{s} -conflations $X \rightarrow I \rightarrow X'$ and $X'' \rightarrow P \rightarrow X$ such that $P \in {}^{\perp_1}\mathcal{D}$ and $I \in \mathcal{D}^{\perp_1}$). In [31], Zhu and Zhuang defined the notion of tilting class of projective dimension $\leq n$. In [31, Thm. 1] we can find conditions under which it is fulfilled that: if V is extended tilting, then $\text{Add}(V)$ is a tilting class of projective dimension ≤ 1 .

5.1 Projective dimension of extended tilting objects

We will show in this section that the projective dimension of a extended tilting object is ≤ 1 . For this, we will need to introduce the higher \mathbb{E} -extension groups of an extriangulated category \mathcal{D} .

As a precedent, let us recall that one can define higher \mathbb{E} -extension groups in exact categories by splicing short exact sequences [11, Chap. 6]. A similar thing has been done for extriangulated categories in [12, Sect. 3.2]. For the reader's convenience, and in order to have a language in the forthcoming proofs of the paper, we recall some ideas from [12, Sect. 3.2].

Let \mathcal{D} be a small extriangulated category. For $A, B \in \mathcal{D}$, we define recursively $\mathbb{E}^n(A, B)$ as follows. Set $\mathbb{E}^1(A, B) := \mathbb{E}(A, B)$. Also recall that, for $\alpha : A' \rightarrow A$, $\beta : B \rightarrow B'$ and $\delta \in \mathbb{E}(A, B)$, $\delta \cdot \alpha := \mathbb{E}(\alpha, B)(\delta)$ and

$$\beta \cdot \delta := \mathbb{E}(A, \beta)(\delta).$$

For $C \in \mathcal{D}$, denote by $(\delta_C \eta)$ the ordered pair $(\delta, \eta) \in \mathbb{E}(C, B) \times \mathbb{E}(A, C)$. Let $S^2(A, B) := (\coprod_{C \in \mathcal{D}} \mathbb{E}(C, B) \times \mathbb{E}(A, C)) / \sim$, where $(\delta_X \eta) \sim (\delta'_Y \eta')$ if:

- there is a morphism $f : X \rightarrow Y$ such that $\delta' \cdot f = \delta$ and $f \cdot \eta = \eta'$, or
- there is a morphism $f : Y \rightarrow X$ such that $\delta \cdot f = \delta'$ and $f \cdot \eta' = \eta$.

Note that \sim is reflexive and symmetric but not necessarily transitive. Hence, we consider the equivalence relation \approx generated by \sim . Lastly, define $\mathbb{E}^2(A, B) := (\coprod_{C \in \mathcal{D}} \mathbb{E}(C, B) \times \mathbb{E}(A, C)) / \approx$. It can be proved that $\mathbb{E}^2(A, B)$ is an abelian group with the Baer sum:

$$\overline{(\delta_X \eta)} + \overline{(\delta'_Y \eta')} := \overline{(\delta \cdot \pi_1 + \delta' \cdot \pi_2)_{X \amalg Y} (\mu_1 \cdot \eta + \mu_2 \cdot \eta')},$$

where $\pi_1 : X \amalg Y \rightarrow X$ and $\pi_2 : X \amalg Y \rightarrow Y$ are the canonical projections and $\mu_1 : X \rightarrow X \amalg Y$ and $\mu_2 : Y \rightarrow X \amalg Y$ are the canonical inclusions. Observe that the following equalities hold:

$$\overline{((\eta + \eta')_X \delta)} = \overline{((\eta \cdot \pi_1 + \eta' \cdot \pi_2)_{X \amalg X} (\mu_1 \cdot \delta + \mu_2 \cdot \delta))} = \overline{(\eta_X \delta)} + \overline{(\eta'_X \delta)}$$

where the first equality is induced by the morphism $\mu_1 + \mu_2 : X \rightarrow X \amalg X$;

$$\overline{(\eta_X (\delta + \delta'))} = \overline{((\eta \cdot \pi_1 + \eta \cdot \pi_2)_{X \amalg X} (\mu_1 \cdot \delta + \mu_2 \cdot \delta'))} = \overline{(\eta_X \delta)} + \overline{(\eta_X \delta')}$$

where the first equality is given by $\pi_1 + \pi_2 : X \amalg X \rightarrow X$;

$$\overline{(0_X \delta)} = \overline{((0 \cdot \pi_1 + 0 \cdot \pi_2)_{X \amalg X} (\mu_1 \cdot \delta + \mu_2 \cdot \delta))} = \overline{(0_X \delta)} + \overline{(0_X \delta)}$$

where the first equality follows from the morphism $\mu_1 + \mu_2 : X \rightarrow X \amalg X$; and similarly that

$$\overline{(\eta_X 0)} = \overline{((\eta \cdot \pi_1 + \eta \cdot \pi_2)_{X \amalg X} (\mu_1 \cdot 0 + \mu_2 \cdot 0))} = \overline{(\eta_X 0)} + \overline{(\eta_X 0)}.$$

By using this, one can prove that:

$$\overline{(\eta_X 0)} = 0 = \overline{(0_X \delta)} \text{ and } \overline{(-\eta_X \delta)} = -\overline{(\eta_X \delta)} = \overline{(\eta_X -\delta)}.$$

Lastly, for $\alpha : A' \rightarrow A$, $\beta : B \rightarrow B'$ and $\epsilon = \overline{(\delta_X \eta)} \in \mathbb{E}^2(A, B)$, define $\epsilon \cdot \alpha = \mathbb{E}^2(\alpha, B)(\epsilon) := \overline{(\delta_X (\eta \cdot \alpha))}$ and $\beta \cdot \epsilon := \mathbb{E}^2(A, \beta)(\epsilon) := \overline{((\beta \cdot \delta)_X \eta)}$.

Let $n > 1$. For $C \in \mathcal{D}$, denote by $(\delta_C \eta)$ an ordered pair $(\delta, \eta) \in \mathbb{E}(C, B) \times \mathbb{E}^n(A, C)$. Let $S^{n+1}(A, B) := (\coprod_{C \in \mathcal{D}} \mathbb{E}(C, B) \times \mathbb{E}^n(A, C)) / \sim$, where $(\delta_X \eta) \sim (\delta'_Y \eta')$ if

- there is a morphism $f : X \rightarrow Y$ such that $\delta' \cdot f = \delta$ and $f \cdot \eta = \eta'$, or
- there is a morphism $f : Y \rightarrow X$ such that $\delta \cdot f = \delta'$ and $f \cdot \eta' = \eta$.

Consider the equivalence relation \approx generated by \sim . Define $\mathbb{E}^{n+1}(A, B) := (\coprod_{C \in \mathcal{D}} \mathbb{E}(C, B) \times \mathbb{E}^n(A, C)) / \approx$. It can be proved that $\mathbb{E}^{n+1}(A, B)$ is an abelian group with the Baer sum:

$$\overline{(\delta \cdot_X \eta)} + \overline{(\delta' \cdot_Y \eta')} := \overline{(\delta \cdot \pi_1 + \delta' \cdot \pi_2) \cdot_{X \amalg Y} (\mu_1 \cdot \eta + \mu_2 \cdot \eta')},$$

where $\pi_1 : X \amalg Y \rightarrow X$ and $\pi_2 : X \amalg Y \rightarrow Y$ are the canonical projections and $\mu_1 : X \rightarrow X \amalg Y$ and $\mu_2 : Y \rightarrow X \amalg Y$ are the canonical inclusions. As before, we have that:

$$\overline{(\eta \cdot_X 0)} = 0 = \overline{(0 \cdot_X \delta)} \text{ and } \overline{(-\eta \cdot_X \delta)} = -\overline{(\eta \cdot_X \delta)} = \overline{(\eta \cdot_X -\delta)}.$$

Lastly, for $\alpha : A' \rightarrow A$, $\beta : B \rightarrow B'$ and $\epsilon = \overline{(\delta \cdot_X \eta)} \in \mathbb{E}^2(A, B)$, define $\epsilon \cdot \alpha := \mathbb{E}^{n+1}(\alpha, B)(\epsilon) := \overline{(\delta \cdot_X (\eta \cdot \alpha))}$ and $\beta \cdot \epsilon := \mathbb{E}^{n+1}(A, \beta)(\epsilon) := \overline{((\beta \cdot \delta) \cdot_X \eta)}$.

Definition 5.5. For a small extriangulated category \mathcal{D} and $V \in \mathcal{D}$, we say that the **projective dimension** of V is ≤ 1 , denoted by $\text{pd}(V) \leq 1$, if the equality $V^{\perp_1} = V^{\perp_{>0}} := \{X \in \mathcal{D} \mid \mathbb{E}^k(V, X) = 0 \forall k > 0\}$ holds true.

The following result is inspired from [23, Cor. 2.6].

Proposition 5.6. *Let \mathcal{D} be a small extriangulated category with negative first extension and $V \in \mathcal{D}$. If V is extended tilting, then $\text{pd}(V) \leq 1$.*

Proof. It is enough to show that $V^{\perp_1} \subseteq V^{\perp_{>0}}$. We proceed by induction on n to prove that $\mathbb{E}^n(V, V^{\perp_1}) = 0$. The case $n = 1$ is trivial.

Let $n > 1$ and assume that $\mathbb{E}^{n-1}(V, V^{\perp_1}) = 0$. For $X \in V^{\perp_1}$ and $\epsilon \in \mathbb{E}^n(V, X)$, we have that $\epsilon = \overline{(\eta \cdot_C \delta)}$ with $\eta \in \mathbb{E}(C, X)$ and $\delta \in \mathbb{E}^{n-1}(V, C)$. Let η be realized by $X \rightarrow E \rightarrow C$. Since $\text{Gen}(V)$ is cogenerating, there is an \mathfrak{s} -conflation $E \rightarrow T \rightarrow S$ with $T \in \text{Gen}(V)$. Then, by [18, Def. 2.12 (ET4)], there is a morphism $f : C \rightarrow C'$ and an \mathfrak{s} -conflation $\eta' : X \rightarrow T \rightarrow C'$ such that $\eta' \cdot f = \eta$ as can be seen in the following diagram

$$\begin{array}{ccccc} X & \longrightarrow & E & \longrightarrow & C \\ \parallel & & \downarrow & & \downarrow f \\ X & \longrightarrow & T & \longrightarrow & C' \\ & & \downarrow & & \downarrow \\ & & S & \xlongequal{\quad} & S. \end{array}$$

Observe that $C' \in V^{\perp_1}$ since $X, T \in V^{\perp_1}$ and $(V^{\perp_1}, V^{\perp_0})$ is an \mathfrak{s} -torsion pair (see Proposition 2.9 (a)). Hence $f \cdot \delta \in \mathbb{E}^{n-1}(V, C') = 0$ by induction hypothesis. Therefore $\epsilon = \overline{(\eta \cdot_C \delta)} = \overline{((\eta' \cdot f) \cdot_C \delta)} = \overline{(\eta' \cdot_{C'} (f \cdot \delta))} = \overline{(\eta' \cdot_{C'} 0)} = 0$. \square

5.2 Extended tilting objects are \mathbb{E} -universal

In this section we will show that extended tilting objects in extriangulated categories are \mathbb{E} -universal. Note that the notion of an \mathbb{E} -universal object given below is inspired from the notion of an **Ext**-universal object for abelian categories (see [23, Def. 5.6] or [5] for more details).

Definition 5.7. Let \mathcal{D} be an extriangulated category. An object V in \mathcal{D} is **\mathbb{E} -universal** if, for any $D \in \mathcal{D}$, there is an \mathbb{E} -triangle in \mathcal{D} of the form:

$$\varepsilon : D \rightarrow A \rightarrow \coprod_{\alpha} V$$

for some set $\alpha \neq \emptyset$, such that the canonical map $\text{Hom}_{\mathcal{D}}(V, \coprod_{\alpha} V) \rightarrow \mathbb{E}^1(V, D)$ is surjective. In such case, ε is called a **universal \mathbb{E} -triangle** of V by D .

We can adapt the proof of [23, Lem. 6.6] to our context as follows.

Proposition 5.8. *Let \mathcal{D} be an extriangulated category with negative first extension. Then, any extended tilting object $V \in \mathcal{D}$ is \mathbb{E} -universal.*

Proof. Let $V \in \mathcal{D}$ be an extended tilting object and $D \in \mathcal{D}$. Since $\mathcal{T} := \text{Gen}(V)$ is a cogenerating class in \mathcal{D} , there is an \mathfrak{s} -conflation $D \rightarrow T \rightarrow D'$ in \mathcal{D} with $T \in \mathcal{T}$. Now, from the \mathfrak{s} -conflation (see Remark 2.10) $\delta_{D'} : T' \rightarrow D' \rightarrow F$ associated to the s -torsion pair $\mathfrak{t}(V) := (\mathcal{T}, \mathcal{F}) = (\text{Gen}(V), V^{\perp_0})$ together with [18, Rk 2.22], we get the following commutative diagram in \mathcal{D}

$$\begin{array}{ccccc} D & \longrightarrow & Z & \longrightarrow & T' \\ \parallel & & \downarrow & & \downarrow \\ D & \longrightarrow & T & \longrightarrow & D' \\ & & \downarrow & & \downarrow \\ & & F & \xlongequal{\quad} & F, \end{array}$$

where $T' \in \mathcal{T}$, $F \in \mathcal{F}$ and all the rows and columns are \mathbb{E} -triangles. Applying the functor $\text{Hom}_{\mathcal{D}}(V, -)$ to the left column of the above diagram, we obtain the exact sequence:

$$0 = \text{Hom}_{\mathcal{D}}(V, F) \rightarrow \mathbb{E}^1(V, Z) \rightarrow \mathbb{E}^1(V, T) = 0$$

where the last equality hold, since $T \in \mathcal{T} = V^{\perp_1}$. Thus, $Z \in V^{\perp_1} = \mathcal{T}$. Now, let $T' \neq 0$. Then, from the equality $\mathcal{T} = \text{Pres}(V)$, [18, Prop. 3.15] together with the first row in the above diagram, we deduce the following commutative diagram in \mathcal{D} :

$$\begin{array}{ccccc} & & T'' \xlongequal{\quad} T'' & & \\ & & \downarrow & & \downarrow \\ D & \longrightarrow & D'' & \longrightarrow & \coprod_{\alpha} V \\ \parallel & & \downarrow & & \downarrow \\ D & \longrightarrow & Z & \longrightarrow & T' \end{array}$$

for some set $\alpha \neq \emptyset$ and $T'' \in \mathcal{T}$. In case $T' = 0$, we can use the same diagram by replacing the s -conflation $T'' \rightarrow \coprod_{\alpha} V \rightarrow T'$ with $V \rightarrow V \rightarrow 0$. In particular,

from this diagram, we get that $D'' \in \mathcal{T}$ since \mathcal{T} is closed under extensions in \mathcal{D} . Now, applying the functor $\text{Hom}_{\mathcal{D}}(V, -)$ to the upper row in the last diagram, we obtain the exact sequence:

$$\text{Hom}_{\mathcal{D}}(V, \coprod_{\alpha} V) \rightarrow \mathbb{E}^1(V, D) \rightarrow \mathbb{E}^1(V, D'') = 0$$

where the last equality is due to the orthogonality condition of V in \mathcal{T} . Therefore, such \mathbb{E} -triangle is a universal \mathbb{E} -triangle of V by C as desired. \square

5.3 Extended tilting objects in restricted hearts

In this section we will show that an extended tilting object in a restricted heart is nothing more than a quasi-tilting object in a standard heart. Let us recall the notion of quasi-tilting object from [23, Def. 6.1].

Definition 5.9. Let \mathcal{D} be an extriangulated category with negative first extension. We say that $V \in \mathcal{D}$ is a **quasi-tilting object** if the following statements hold true.

(T1) $\mathbf{t}(V) := (\text{Gen}(V), V^{\perp_0}) \in \text{stors } \mathcal{D}$.

(QT) $\text{Gen}(V) = \text{Pres}(V) = \overline{\text{Gen}}_{\mathcal{D}}(V) \cap V^{\perp_1}$, where $\overline{\text{Gen}}_{\mathcal{D}}(V)$ is the class of objects $X \in \mathcal{D}$ that admit an \mathfrak{s} -conflation $X \rightarrow V_0 \rightarrow X'$ in \mathcal{D} with $V_0 \in \text{Gen}(V)$.

Lemma 5.10. Let \mathcal{D} be an extriangulated category with negative first extension and $V \in \mathcal{D}$. If $(\mathcal{T}, \mathcal{F})$ is an s -torsion pair with $\mathcal{T} = \text{Gen}(V)$, then $\mathcal{F} = V^{\perp_0}$.

Proof. Since $V \in \mathcal{T}$, we have that $\mathcal{F} \subseteq V^{\perp_0}$. For the opposite contention, consider $X \in V^{\perp_0}$ and the canonical \mathfrak{s} -conflation $\eta : T \xrightarrow{u} X \rightarrow F$ with $T \in \mathcal{T}$ and $F \in \mathcal{F}$. Observe that, since $\mathcal{T} = \text{Gen}(V)$, $T = 0$ or there is an \mathfrak{s} -conflation $A \rightarrow V_0 \xrightarrow{f} X$ with $f \neq 0$ and $V_0 = \coprod_{\alpha} V$. But, the exact sequence

$$\mathbb{E}^{-1}(V_0, F) \xrightarrow{\eta^{-1}} \text{Hom}_{\mathcal{D}}(V_0, T) \rightarrow \text{Hom}_{\mathcal{D}}(V_0, X)$$

implies that $f \in \text{Ker}(\text{Hom}_{\mathcal{D}}(V_0, u)) = \text{Im}(\eta^{-1}) = 0$ because $V_0 \in \mathcal{T}$ and $F \in \mathcal{F}$. Therefore, $f = 0$ and thus $X \in \mathcal{F}$. \square

Lemma 5.11. Let \mathcal{D} be a triangulated category, $\mathbf{t}_1 \leq \mathbf{t}_2 \leq \mathbf{t}_3$ in $\text{stors } \mathcal{D}$, $V \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ and I be a set. Then, the coproduct $\coprod_I^{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}} V$ of I -copies of V in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ exists if, and only if, the coproduct $\coprod_I^{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}} V$ of I -copies of V in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ exists. In such case, we have that both coproducts are isomorphic.

Proof. By Lemma 3.2 (c), we have that $(\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}, \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}) \in \text{stors } \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$. Moreover, by Lemma 3.1 and Remark 2.10, we get that the inclusion functor $\iota : \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} \hookrightarrow \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ is a left adjoint of the functor $\mathbf{t}_2|_{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}} : \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]} \rightarrow \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. Using such adjoint, we get that $\coprod_I^{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}} V$ is a coproduct of I -copies of V in

$\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$. On the other hand, suppose that there exists $\coprod_I^{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}} V$, the coproduct of I -copies of V in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$. Using once again the Lemma 3.1, we have an \mathfrak{s} -conflation $\eta : T \rightarrow \coprod_I^{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}} V \xrightarrow{f} F$, where $T \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ and $F \in \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$. Note that $f \circ \mu_i = 0$ for all $i \in I$, where $\mu_i : V \hookrightarrow \coprod_I^{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}} V$ is the canonical inclusion. Hence, by the universal property of coproducts, we have that $f = 0$. Therefore, we have the distinguished triangle $\Sigma^{-1}F \rightarrow T \rightarrow \coprod_I^{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}} V \xrightarrow{f} F$, with $f = 0$, in the triangulated category \mathcal{D} , and thus $T \simeq \Sigma^{-1}F \amalg \coprod_I^{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}} V$. Hence $\coprod_I^{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}} V \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$, and thus, the existence of the coproduct $\coprod_I^{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}} V$ together with the isomorphism $\coprod_I^{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}} V \simeq \coprod_I^{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}} V$ is clear. \square

Theorem 5.12. *Let \mathcal{D} be a triangulated category and $\mathbf{t}_1 \leq \mathbf{t}_2$ in $\text{stors } \mathcal{D}$ be such that $\mathbf{t}_2 \leq \Sigma^{-1}\mathbf{t}_1$. If $V \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ satisfies that $\overline{\text{Gen}}_{\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}}(V) \subseteq \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$, then the following statements are equivalent.*

- (a) V is an extended tilting object in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$.
- (b) V is a quasi-tilting object in $\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}$ and $\overline{\text{Gen}}_{\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}}(V) = \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$.

Moreover, if one of the above statements holds true, then

$$V_{\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}}^{\perp_0} = V_{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}}^{\perp_0} \star \mathcal{H}_{[\mathbf{t}_2, \Sigma^{-1}\mathbf{t}_1]}.$$

Proof. Let $\mathbf{t}_1 = (\mathcal{T}_1, \mathcal{F}_1)$ and $\mathbf{t}_2 = (\mathcal{T}_2, \mathcal{F}_2)$. Observe that $\mathcal{T} := \text{Gen}_{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}}(V) = \text{Gen}_{\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}}(V)$ and $\text{Pres}_{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}}(V) = \text{Pres}_{\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}}(V)$ by Lemma 5.11 and Proposition 3.9. For simplicity, we denote $\overline{\text{Gen}}(V) := \overline{\text{Gen}}_{\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}}(V)$.

(b) \Rightarrow (a) Since V is quasi-tilting in $\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}$, we have the s -torsion pair $(\mathcal{T}, V_{\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}}^{\perp_0}) \in \text{stors } \mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}$. Hence, by Lemma 3.2(d) and Lemma 5.10, it follows that $(\mathcal{T}, V_{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}}^{\perp_0}) \in \text{stors } \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ and $V_{\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}}^{\perp_0} = V_{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}}^{\perp_0} \star \mathcal{H}_{[\mathbf{t}_2, \Sigma^{-1}\mathbf{t}_1]}$. We have proved that V satisfies (T1) in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. To prove (T2), it is enough to show that $\mathcal{T} = V_{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}}^{\perp_1}$. For this, observe that

$$\mathcal{T} = V_{\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}}^{\perp_1} \cap \overline{\text{Gen}}(V) = V_{\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}}^{\perp_1} \cap \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} = V_{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}}^{\perp_1}.$$

Lastly, to prove (T3), we observe that $\text{Gen}(V)$ is a cogenerating class in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ because $\overline{\text{Gen}}(V) = \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ in $\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}$ and \mathcal{T}_2 is closed under cones.

(a) \Rightarrow (b) We have $(\mathcal{T}, V_{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}}^{\perp_0}) \in \text{stors } \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. Thus, by Lemma 3.2(a) and Lemma 5.10, it follows that $(\mathcal{T}, V_{\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}}^{\perp_0}) \in \text{stors } \mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}$ and that $V_{\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}}^{\perp_0} = V_{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}}^{\perp_0} \star \mathcal{H}_{[\mathbf{t}_2, \Sigma^{-1}\mathbf{t}_1]}$. Therefore, V satisfies (T1) in $\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}$.

Let us prove that $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} = \overline{\text{Gen}}(V)$. Since V is extended tilting in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$, we have that $\text{Gen}(V)$ is a cogenerating class in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. Hence, $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} \subseteq \overline{\text{Gen}}(V)$. Therefore, we have proved the desired equality since the opposite contention is an hypothesis.

Lastly, we proceed to prove that $\overline{\text{Gen}}(V) \cap V_{\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}}^{\perp 1} = \mathcal{T}$. We have that $\mathcal{T} \subseteq \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} = \overline{\text{Gen}}(V)$ and thus

$$\mathcal{T} = \mathcal{T} \cap V_{\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}}^{\perp 1} \subseteq \overline{\text{Gen}}(V) \cap V_{\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}}^{\perp 1}$$

since $\mathcal{T} = V_{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}}^{\perp 1}$. For the inverse contention, consider $X \in \overline{\text{Gen}}(V) \cap V_{\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}}^{\perp 1}$. It follows that there is an \mathfrak{s} -conflation $X \rightarrow G \rightarrow X'$ with $G \in \mathcal{T}$. By Proposition 3.9(a), $X' \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. Note that $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} = \mathcal{T} \star V_{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}}^{\perp 0}$. Hence, there is an \mathfrak{s} -conflation $T \rightarrow X' \rightarrow W$ with $T \in \mathcal{T}$ and $W \in V_{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}}^{\perp 0}$. Then, by the Octahedral Axiom, we have the following commutative diagram with \mathfrak{s} -conflations as rows and columns:

$$\begin{array}{ccccc} X & \longrightarrow & M & \longrightarrow & T \\ \parallel & & \downarrow & & \downarrow \\ X & \longrightarrow & G & \longrightarrow & X' \\ & & \downarrow & & \downarrow \\ & & W & \xlongequal{\quad} & W \end{array}$$

Here, observe that the morphism $G \rightarrow W$ is equal to zero since $G \in \mathcal{T}$ and $W \in V_{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}}^{\perp 0}$, and thus, $M \cong G$. Now, since $T \in \mathcal{T} = \text{Pres}(V)$, there is an \mathfrak{s} -conflation $T' \rightarrow V_0 \rightarrow T$, where $V_0 \cong \coprod_{\alpha} V$ and $T' \in \mathcal{T}$. Lastly, from the \mathfrak{s} -conflations $T' \rightarrow V_0 \rightarrow T$ and $X \rightarrow M \rightarrow T$ and the Octahedral Axiom, we get the following commutative diagram with \mathfrak{s} -conflations as rows and columns:

$$\begin{array}{ccccc} & & T' & \xlongequal{\quad} & T' \\ & & \downarrow & & \downarrow \\ X & \longrightarrow & M' & \longrightarrow & V_0 \\ \parallel & & \downarrow & & \downarrow \\ X & \longrightarrow & M & \longrightarrow & T \end{array}$$

Here, we note that $M' \in \mathcal{T}$ since \mathcal{T} is closed under extensions; and that $X \rightarrow M' \rightarrow V_0$ splits since $X \in V_{\mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_1]}}^{\perp 1}$. Therefore, $X \in \mathcal{T}$. □

5.4 Extended tilting objects in extended hearts

Let \mathcal{D} be a triangulated category and $\mathbf{t}_1 \leq \mathbf{t}_2 \leq \mathbf{t}_3$ in $\text{stors } \mathcal{D}$. The aim of this section is to study extended tilting objects in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$. We will see that, under certain conditions, an object V in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ is extended tilting if, and only if, it is a projective generator in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. In particular, one could use this result to determine when the heart of a t -structure has a projective generator. Recall that this is of great interest for several reasons (see for example [9, 20]). In

particular, hearts with a projective generator are related to the theory of silting modules, see for example [3, 4].

Proposition 5.13. *For a triangulated category \mathcal{D} and $\mathbf{t}_1 \leq \mathbf{t}_2 \leq \mathbf{t}_3$ in $\text{stors } \mathcal{D}$, the following statements hold true for an extended tilting object $V \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$.*

- (a) *Let $\mathbf{t}_1 \leq \Sigma \mathbf{t}_2$. Then, $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ is a cogenerating class in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ and $\text{Gen}_{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}}(V) \subseteq \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$.*
- (b) *Let $\mathbf{t}_1 \leq \Sigma \mathbf{t}_2$ and $\mathbf{t}_3 = \Sigma^{-1} \mathbf{t}_2$. Then,*
 - (b1) *$\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} = \text{Gen}_{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}}(V)$ and V is a projective generator in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$;*
 - (b2) *if W is an extended tilting in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$, then $\text{Add}(V) = \text{Add}(W)$.*

Proof. Set $\mathbf{C} := \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ and $\mathbf{H} := \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$, and let $\mathbf{t} = (\mathcal{T}, \mathcal{F}) := (\text{Gen}_{\mathbf{C}}(V), V_{\mathbf{C}}^{\perp_0})$ be the associated extended tilting s -torsion pair in \mathbf{C} . Note that $\Sigma \mathbf{t} \leq \mathbf{t}$ for any $\mathbf{t} \in \text{stors } \mathcal{D}$, and that $[\mathbf{t}_2, \mathbf{t}_3]$ is normal if $\mathbf{t}_3 = \Sigma^{-1} \mathbf{t}_2$.

(a) It is enough to show that $\mathcal{T} = \text{Gen}_{\mathbf{C}}(V) \subseteq \mathbf{H}$ since $\text{Gen}_{\mathbf{C}}(V)$ is a cogenerating class in \mathbf{C} .

We start by proving that $V \in \mathbf{H}$. Indeed, from Lemma 3.1, we get an \mathfrak{s} -conflation $A \rightarrow V \xrightarrow{f} B$ in \mathbf{C} with $A \in \mathbf{H}$ and $B \in \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$. Suppose that $f \neq 0$. Then $B \in \text{Gen}_{\mathbf{C}}(V) = \mathcal{T}$. Now, since $B \in \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$, we have that $\Sigma B \in \Sigma \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]} = \mathcal{H}_{[\Sigma \mathbf{t}_2, \Sigma \mathbf{t}_3]} \subseteq \mathbf{C}$. Thus, we can consider the canonical \mathfrak{s} -conflation $\eta : T \rightarrow \Sigma B \rightarrow F$ with respect to the s -torsion pair \mathbf{t} , where $T \in \mathcal{T}$ and $F \in \mathcal{F}$. By applying the functor $\text{Hom}_{\mathbf{C}}(V, -)$ to η , we obtain the following exact sequence

$$\mathbb{E}^{-1}(V, F) \rightarrow \text{Hom}_{\mathbf{C}}(V, T) \rightarrow \text{Hom}_{\mathbf{C}}(V, \Sigma B).$$

Observe that $\text{Hom}_{\mathbf{C}}(V, T) = 0$ and $\Sigma B \in \mathcal{F}$ since $\mathbb{E}^{-1}(V, F) = 0$ and

$$\text{Hom}_{\mathbf{C}}(V, \Sigma B) = \text{Hom}_{\mathcal{D}}(V, \Sigma B) = \mathbb{E}_{\mathbf{C}}^1(V, B) = 0,$$

where the last equality is due to the fact that $B \in \text{Gen}_{\mathbf{C}}(V) = V^{\perp_1}$. Now, from the fact that $\Sigma B \in \mathcal{F}$, we obtain

$$0 = \mathbb{E}^{-1}(V, \Sigma B) = \text{Hom}_{\mathcal{D}}(V, B) = \text{Hom}_{\mathbf{C}}(V, B).$$

But this is a contradiction since $0 \neq f \in \text{Hom}_{\mathbf{C}}(V, B)$. Hence we conclude that $f = 0$. Thus, we get a distinguished triangle $\Sigma^{-1} B \rightarrow A \rightarrow V \xrightarrow{f} B$ in \mathcal{D} , with $f = 0$, and then $A = \Sigma^{-1} B \amalg V$. Therefore $V \in \mathbf{H}$ since $A \in \mathbf{H}$. Finally, by using that $V \in \mathbf{H}$ together with Lemma 5.11, we conclude that $\mathcal{T} = \text{Gen}_{\mathbf{C}}(V) \subseteq \mathbf{H}$.

(b1) By (a), we know that $\mathcal{T} = \text{Gen}_{\mathbf{C}}(V) \subseteq \mathbf{H}$. Let $\tilde{H} \in \mathbf{H}$. Consider the canonical \mathfrak{s} -conflation $T \rightarrow \tilde{H} \rightarrow F$ with respect to the s -torsion pair \mathbf{t} . From Proposition 3.9 (a), we have that $F \in \mathbf{H} = \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ and then $\Sigma^{-1} F \in \mathcal{H}_{[\Sigma^{-1} \mathbf{t}_1, \Sigma^{-1} \mathbf{t}_2]} \subseteq \mathbf{C}$. This implies, by (a), that $\Sigma^{-1} F \in V^{\perp_1} = \mathcal{T} \subseteq \mathbf{H}$ since

$$\mathbb{E}^1(V, \Sigma^{-1} F) = \text{Hom}_{\mathcal{D}}(V, \Sigma(\Sigma^{-1} F)) = \text{Hom}_{\mathbf{C}}(V, F) = 0.$$

Thus, we have the s -conflation $\Sigma^{-1}F \rightarrow T \rightarrow \tilde{H}$ in \mathcal{C} , with $\Sigma^{-1}F, T \in \mathcal{T}$. Hence by Proposition 2.9 (a), we get $\tilde{H} \in \mathcal{T}$. Now, in the s -conflation $T \rightarrow \tilde{H} \rightarrow F$ in \mathcal{C} , we have $\tilde{H}, T \in \mathcal{T}$. And then, by Proposition 2.9 (a), $F \in \mathcal{T} \cap \mathcal{F} = 0$. Therefore, $\tilde{H} \cong T$; and hence $\mathcal{H} \subseteq \text{Gen}_{\mathcal{C}}(V) = \mathcal{T}$, proving that $\mathcal{T} = \text{Gen}_{\mathcal{C}}(V) = \mathcal{H}$. Moreover, notice that V is a projective object of \mathcal{H} since $\mathcal{T} = V^{\perp_1}$. Lastly, we show that V is a generator of \mathcal{H} . For this, we take a non-zero object $X \in \mathcal{H}$, and recall that $\mathcal{H} = \mathcal{T} = \text{Gen}_{\mathcal{C}}(V) = \text{Pres}_{\mathcal{C}}(V)$. Thus, there is an \mathfrak{s} -conflation

$$\theta : Y \rightarrow \coprod_{\alpha}^{\mathcal{C}} V \rightarrow X,$$

in \mathcal{C} , for some set $\alpha \neq \emptyset$ and $Y \in \text{Gen}_{\mathcal{C}}(V) = \mathcal{H}$. From Lemma 5.11, we know that $\coprod_{\alpha}^{\mathcal{C}} V \simeq \coprod_{\alpha}^{\mathcal{H}} V$. Hence by the s -conflation θ , it follows that $X \in \text{Gen}_{\mathcal{H}}(V)$. Therefore $\text{Gen}_{\mathcal{H}}(V) = \mathcal{H}$ as desired.

(b2) Let us prove that $W \in \text{Add}(V)$. From the item (b1), we have that $\text{Pres}_{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}}(V) = \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]} = \text{Pres}_{\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}}(W)$. Hence, there is an \mathfrak{s} -conflation $\varepsilon : H \rightarrow \coprod_{\beta} V \rightarrow W$ in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. Note that ε splits since $W^{\perp_1} = \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ and thus $W \in \text{Add}(V)$. Similarly, we have that $V \in \text{Add}(W)$ and therefore $\text{Add}(V) = \text{Add}(W)$. \square

Theorem 5.14. *Let \mathcal{D} be a triangulated category and $\mathbf{t}_1 \leq \mathbf{t}_2 \leq \mathbf{t}_3$ in $\text{stors } \mathcal{D}$, be such that: $\mathbf{t}_1 \leq \Sigma \mathbf{t}_2$ and $\mathbf{t}_3 = \Sigma^{-1} \mathbf{t}_2$. Then, for $V \in \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$, the following statements are equivalent.*

- (a) V is an extended tilting object in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$.
- (b) $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$ is a cogenerating class in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ and V is a projective generator in $\mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$.

Proof. Set the notation $\mathcal{C} := \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_3]}$ and $\mathcal{H} := \mathcal{H}_{[\mathbf{t}_1, \mathbf{t}_2]}$. We also recall that $\mathbf{t} := (\mathcal{H}, \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]})$ is an s -torsion pair in \mathcal{C} , see Lemma 3.2. Note that $[\mathbf{t}_2, \mathbf{t}_3]$ is normal.

The implication (a) \Rightarrow (b) can be obtained from Proposition 5.13. Let us show that (b) implies (a). Indeed, let V be a projective generator in \mathcal{H} which is a cogenerating class in \mathcal{C} . Thus $\mathcal{H} = \text{Gen}_{\mathcal{H}}(V)$ and by Lemma 5.11 and Proposition 3.9 (a), it follows that $\mathcal{H} = \text{Gen}_{\mathcal{H}}(V) = \text{Gen}_{\mathcal{C}}(V)$. Now, using that $\text{Pres}_{\mathcal{H}}(V) = \text{Gen}_{\mathcal{H}}(V) = \mathcal{H}$ and $\text{Pres}_{\mathcal{H}}(V) \subseteq \text{Pres}_{\mathcal{C}}(V)$, we get that $\text{Pres}_{\mathcal{C}}(V) = \text{Gen}_{\mathcal{C}}(V) = \mathcal{H}$.

We assert now that $\mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]} = V^{\perp_0}$. Indeed, the inclusion $\mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]} \subseteq V^{\perp_0}$ follows since $V \in \mathcal{H}$ and $\mathbf{t} \in \text{stors } \mathcal{C}$. Now, let $Z \in V^{\perp_0}$. Consider the \mathfrak{s} -conflation $\delta_Z : H \rightarrow Z \rightarrow H'$ in \mathcal{C} with $H \in \mathcal{H}$ and $H' \in \mathcal{H}_{[\mathbf{t}_2, \mathbf{t}_3]}$ given by \mathbf{t} . Applying the functor $(V, -) := \text{Hom}_{\mathcal{D}}(V, -)$ in δ_Z , we get the exact sequence

$$(V, \Sigma^{-1}H') \rightarrow (V, H) \rightarrow (V, Z)$$

where: $(V, Z) = 0$ since $Z \in V^{\perp_0}$; and $(V, \Sigma^{-1}H') = 0$ since $V \in \mathcal{H}$, $\Sigma^{-1}H' \in \mathcal{H}_{[\mathbf{t}_3, \Sigma^{-1}\mathbf{t}_3]} \subseteq \mathcal{H}_{[\mathbf{t}_2, \Sigma^{-1}\mathbf{t}_3]}$ and $(\mathcal{H}, \mathcal{H}_{[\mathbf{t}_2, \Sigma^{-1}\mathbf{t}_3]}) \in \text{stors } \mathcal{H}_{[\mathbf{t}_1, \Sigma^{-1}\mathbf{t}_3]}$ (see Lemma 3.2 (c)). Hence $0 = \text{Hom}_{\mathcal{D}}(V, H) = \text{Hom}_{\mathcal{H}}(V, H)$ and thus $H = 0$ since V is a

generator in \mathbf{H} . This shows that $V^{\perp_0} = \mathcal{H}_{[t_2, t_3]}$. Therefore, $(\text{Gen}_{\mathbf{C}}(V), V^{\perp_0}) = (\mathbf{H}, \mathcal{H}_{[t_2, t_3]})$ which is an s -torsion pair in \mathbf{C} .

It remains to prove that $\text{Gen}_{\mathbf{C}}(V) = V^{\perp_1}$. Observe that this task is reduced to check that $V^{\perp_1} \subseteq \text{Gen}_{\mathbf{C}}(V)$ since V is a projective object in \mathbf{H} and $\text{Gen}_{\mathbf{C}}(V) = \mathbf{H}$. Let $C \in V^{\perp_1}$. Since \mathbf{H} is a cogenerating class in \mathbf{C} , there is an \mathfrak{s} -conflation $\eta : C \rightarrow H \rightarrow C'$ in \mathbf{C} , with $H \in \mathbf{H}$ and $C' \in \mathbf{C}$. Hence by Proposition 3.9 (a), we get that $C' \in \mathbf{H}$. If $C' = 0$, then $C \simeq H$ and thus $C \in \mathbf{H}$ as desired. Let $C' \neq 0$. Now, by using the fact that V is a generator in \mathbf{H} , we have an \mathfrak{s} -conflation $\eta' : H' \rightarrow \coprod_{\alpha} V \rightarrow C'$ in \mathbf{H} , where $\alpha \neq \emptyset$. Then, by [18, Prop. 3.15], we get the following commutative diagram in \mathbf{C}

$$\begin{array}{ccccc}
& & C & \xlongequal{\quad} & C \\
& & \downarrow & & \downarrow \\
H' & \longrightarrow & E & \longrightarrow & H \\
\parallel & & \downarrow & & \downarrow \\
H' & \longrightarrow & \coprod_{\alpha} V & \longrightarrow & C'
\end{array}$$

where all the rows and columns are s -conflations. Now, from the middle row, we get that $E \in \mathbf{H}$. Moreover, since $\text{Hom}_{\mathcal{D}}(\coprod_{\alpha} V, \Sigma C) \simeq \prod_{\alpha} \text{Hom}_{\mathcal{D}}(V, \Sigma C) = 0$, it follows that the middle column splits and thus $C \amalg \coprod_{\alpha} V = E \in \mathbf{H}$; and therefore $C \in \mathbf{H}$ as desired. \square

In what follows we will see some situations where Theorem 5.14 can be applied.

Corollary 5.15. *For a triangulated category \mathcal{D} , and a t -structure $\mathbf{u} = (\mathcal{U}, \mathcal{W})$ in \mathcal{D} with heart $\mathcal{H} = \mathcal{U} \cap \mathcal{W}$ and extended heart $\mathcal{C} = \mathcal{W} \cap \Sigma^{-1}\mathcal{U}$, the following statements hold true.*

- (a) $(\mathcal{H}, \Sigma^{-1}\mathcal{H}) \in \text{stors } \mathcal{C}$ and the extriangulated category \mathcal{C} is not abelian if $\mathcal{H} \neq 0$.
- (b) For any $V \in \mathcal{C}$, the following statements are equivalent:
 - (b1) V is an extended tilting object in \mathcal{C} ;
 - (b2) \mathcal{H} is a cogenerating class in \mathcal{C} and V is a projective generator in \mathcal{H} .
- (c) \mathcal{H} is a cogenerating class in \mathcal{C} if, and only if, $\mathcal{C} = \Sigma^{-1}\mathcal{H} * \mathcal{H}$.

Proof. Consider $\mathbf{u}_1 := (\Sigma\mathcal{U}, \mathcal{W})$, $\mathbf{u}_2 := (\mathcal{U}, \Sigma^{-1}\mathcal{W})$ and $\mathbf{u}_3 := (\Sigma^{-1}\mathcal{U}, \Sigma^{-2}\mathcal{W})$. Then we have that $\mathbf{u}_1 \leq \mathbf{u}_2 \leq \mathbf{u}_3$ in $\text{stors } \mathcal{D}$. Thus, the item (a) follows from Lemma 3.2 (c) and Proposition 4.2 (d); and the item (b) can be obtained from Theorem 5.14. Let us show (c). Indeed, it is clear that \mathcal{H} is a cogenerating class in \mathcal{C} , whenever $\mathcal{C} = \Sigma^{-1}\mathcal{H} * \mathcal{H}$. Assume now that \mathcal{H} is a cogenerating class in \mathcal{C} . Then, by Proposition 3.9 (a), we get that $\mathcal{C} = \Sigma^{-1}\mathcal{H} * \mathcal{H}$. \square

Let R be a ring and $\text{Mod-}R$ be the category of right R -modules. We recall that a torsion pair $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ in the abelian category $\text{Mod-}R$ splits if, for any R -module A , the canonical exact sequence (associated to the torsion pair \mathbf{t}) $0 \rightarrow T \rightarrow A \rightarrow F \rightarrow 0$ splits. Now consider the triangulated category $\mathcal{D}(R)$ which is the unbounded derived category of $\text{Mod-}R$. For a torsion pair $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ in $\text{Mod-}R$, we have (see [13, Prop. 2.1]) the Happel-Reiten-Smalø t -structure $\mathbf{u}(\mathbf{t}) := (\mathcal{U}_{\mathbf{t}}, \mathcal{W}_{\mathbf{t}})$ in $\mathcal{D}(R)$, where

$$\mathcal{U}_{\mathbf{t}} := \{X \in \mathcal{D}(R) : H^i(X) = 0 \text{ for } i > 0, H^0(X) \in \mathcal{T}\}$$

and

$$\mathcal{W}_{\mathbf{t}} := \{X \in \mathcal{D}(R) : H^i(X) = 0 \text{ for } i < -1, H^{-1}(X) \in \mathcal{F}\}.$$

Corollary 5.16. *Let R be a non trivial (i.e. $R \neq 0$) right hereditary ring and $\mathbf{t} = (\mathcal{T}, \mathcal{F})$ be a splitting torsion pair in $\text{Mod-}R$. Then, for the heart $\mathcal{H}_{\mathbf{t}} = \mathcal{U}_{\mathbf{t}} \cap \mathcal{W}_{\mathbf{t}}$ and the extended heart $\mathcal{C}_{\mathbf{t}} = \mathcal{W}_{\mathbf{t}} \cap \Sigma^{-1}\mathcal{U}_{\mathbf{t}}$ of the Happel-Reiten-Smalø t -structure $\mathbf{u}(\mathbf{t}) = (\mathcal{U}_{\mathbf{t}}, \mathcal{W}_{\mathbf{t}})$, the following statements hold true.*

- (a) $(\mathcal{H}_{\mathbf{t}}, \Sigma^{-1}\mathcal{H}_{\mathbf{t}}) \in \text{stors}\mathcal{C}_{\mathbf{t}}$, the extriangulated category $\mathcal{C}_{\mathbf{t}}$ is not abelian and $\mathcal{C}_{\mathbf{t}} = \Sigma^{-1}\mathcal{H}_{\mathbf{t}} * \mathcal{H}_{\mathbf{t}}$.
- (b) For any $V \in \mathcal{C}_{\mathbf{t}}$, the following statements are equivalent:
 - (b1) V is an extended tilting object in $\mathcal{C}_{\mathbf{t}}$;
 - (b2) V is a projective generator in $\mathcal{H}_{\mathbf{t}}$.

Proof. Since $R \neq 0$ and the torsion pair \mathbf{t} splits, we have that $\mathcal{H}_{\mathbf{t}} \neq 0$. Thus, by Corollary 5.15, it is enough to show that $\mathcal{C}_{\mathbf{t}} = \Sigma^{-1}\mathcal{H}_{\mathbf{t}} * \mathcal{H}_{\mathbf{t}}$.

Let $C \in \mathcal{C}_{\mathbf{t}}$. Since $(\mathcal{H}_{\mathbf{t}}, \Sigma^{-1}\mathcal{H}_{\mathbf{t}}) \in \text{stors}\mathcal{C}_{\mathbf{t}}$, there is an \mathfrak{s} -conflation $A \rightarrow C \rightarrow \Sigma^{-1}B$ in $\mathcal{D}(R)$, with $A, B \in \mathcal{H}_{\mathbf{t}}$. By using the exact sequence of cohomologies associated to the distinguished triangle $A \rightarrow C \rightarrow \Sigma^{-1}B \rightarrow \Sigma A$, we obtain the following information:

- (1) $H^k(C) = 0$ for all $k \neq -1, 0, 1$;
- (2) $H^{-1}(C) \cong H^{-1}(A) \in \mathcal{F}$;
- (3) there is a short exact sequence in $\text{Mod-}R$

$$\varepsilon : 0 \rightarrow H^0(A) \xrightarrow{f} H^0(C) \xrightarrow{g} H^{-1}(B) \rightarrow 0;$$

- (4) $H^1(C) \cong H^0(B) \in \mathcal{T}$.

From the short exact sequence ε together with the fact that $H^0(A) \in \mathcal{T}$ and $H^{-1}(B) \in \mathcal{F}$, we deduce that $H^0(A) = \mathbf{t}(H^0(C))$ and $H^{-1}(B) = (1 : \mathbf{t})(H^0(C))$. Hence ε splits since \mathbf{t} splits. Therefore, we can build the \mathfrak{s} -conflation

$$H^{-1}(B) \xrightarrow{\begin{bmatrix} g' \\ 0 \end{bmatrix}} H^0(C) \amalg \Sigma(H^{-1}(C)) \xrightarrow{\begin{bmatrix} f' & 0 \\ 0 & 1 \end{bmatrix}} H^0(A) \amalg \Sigma(H^{-1}(C))$$

in $\mathcal{D}(R)$, where f' is a left inverse of f and g' is a right inverse of g . Recall that C is quasi-isomorphic to $\coprod_{k \in \mathbb{Z}} \Sigma^{-k}(H^k(C)) = \Sigma^{-1}(H^1(C)) \amalg H^0(C) \amalg \Sigma(H^{-1}(C))$

(e.g. see [16, Lem. 4.4.15]). From this fact, together with item (1), we get a splitting s -conflation $\Sigma^{-1}(H^1(C)) \rightarrow C \rightarrow H^0(C) \amalg \Sigma(H^{-1}(C))$ in $\mathcal{D}(R)$. Now, by [18, Rk. 2.22], we get the following commutative diagram in $\mathcal{D}(R)$

$$\begin{array}{ccccc}
\Sigma^{-1}(H^1(C)) & \xrightarrow{\quad} & Z & \xrightarrow{\quad} & H^{-1}(B) \\
\parallel & & \downarrow & & \downarrow \\
\Sigma^{-1}(H^1(C)) & \xrightarrow{\quad} & C & \xrightarrow{\quad} & H^0(C) \amalg \Sigma(H^{-1}(C)) \\
& & \downarrow & & \downarrow \\
& & H^0(A) \amalg \Sigma(H^{-1}(C)) & \xlongequal{\quad} & H^0(A) \amalg \Sigma(H^{-1}(C)),
\end{array}$$

where all the rows and columns are s -conflations. Observe that $Z \in \Sigma^{-1}\mathcal{H}_{\mathbf{t}}$ since $\Sigma^{-1}H^1(C) \in \Sigma^{-1}\mathcal{T}$ by item (4) and $H^{-1}(B) \in \mathcal{F}$. Moreover, we have that $H^0(A) \amalg \Sigma(H^{-1}(C)) \in \mathcal{H}_{\mathbf{t}}$ since $H^0(A) \in \mathcal{T}$ and $\Sigma H^{-1}(C) \in \Sigma\mathcal{F}$ by item (2). From the middle column in the previous diagram, we deduce that $C \in \Sigma^{-1}\mathcal{H}_{\mathbf{t}} \star \mathcal{H}_{\mathbf{t}}$. This shows that $\mathcal{C}_{\mathbf{t}} = \Sigma^{-1}\mathcal{H}_{\mathbf{t}} \star \mathcal{H}_{\mathbf{t}}$. \square

Example 5.17. Let K be a field and $R := K \times K$. Observe that every torsion pair in $\text{Mod-}R$ splits since R is semi-simple. Furthermore, R is also a non trivial hereditary ring. Note that $I := K \times 0$ is a finitely generated idempotent two-sided ideal of R . Consider the torsion pair $\mathbf{t} := (\text{Mod-}R/I, R/I^{\perp_0})$ in $\text{Mod-}R$, and let $\mathcal{H}_{\mathbf{t}}$ be the heart and $\mathcal{C}_{\mathbf{t}}$ be the extended heart of the Happel-Reiten-Smalø t -structure $\mathbf{u}(\mathbf{t})$. Then, by [21, Ex. 5.8], we know that $\mathcal{H}_{\mathbf{t}}$ has a projective generator V . Thus, from Corollary 5.16, we conclude that V is an extended tilting object in the extriangulated category $\mathcal{C}_{\mathbf{t}}$ which is not abelian.

Now consider the following situation. Let \mathcal{D} be a triangulated category, $\mathbf{u} = (\mathcal{U}, \mathcal{W})$ a t -structure in \mathcal{D} and $k \geq 1$. Consider the s -torsion pairs in \mathcal{D} , namely: $\mathbf{u}_1^k := (\Sigma\mathcal{U}, \mathcal{W})$, $\mathbf{u}_2^k := (\Sigma^{-k}\mathcal{U}, \Sigma^{-k-1}\mathcal{W})$ and $\mathbf{u}_3^k := \Sigma^{-1}\mathbf{u}_2^k$. Notice that $\Sigma\mathbf{u}_1^k \leq \mathbf{u}_1^k \leq \mathbf{u}_2^k \leq \mathbf{u}_3^k$ and $\mathbf{u}_1^k \leq \Sigma\mathbf{u}_2^k$. Moreover the interval $[\mathbf{u}_2^k, \mathbf{u}_3^k]$ is normal and we can not assure that for the other intervals. We introduce the classes

$$\mathbf{H}^{(k)} := \mathcal{H}_{[\mathbf{u}_1^k, \mathbf{u}_2^k]} = \mathcal{W} \cap \Sigma^{-k}\mathcal{U} \quad \text{and} \quad \mathbf{C}^{(k)} := \mathcal{H}_{[\mathbf{u}_1^k, \mathbf{u}_3^k]} = \mathcal{W} \cap \Sigma^{-k-1}\mathcal{U}$$

of the t -structure \mathbf{u} . Observe that

$$\mathcal{H}_{[\mathbf{u}_2^k, \mathbf{u}_3^k]} = \Sigma^{-k-1}\mathcal{W} \cap \Sigma^{-k-1}\mathcal{U} = \Sigma^{-k-1}\mathcal{H},$$

where $\mathcal{H} = \mathcal{W} \cap \mathcal{U}$ is the heart of \mathbf{u} , and thus $\mathcal{H}_{[\mathbf{u}_2^k, \mathbf{u}_3^k]}$ is an abelian category. For $\mathbf{H}^{(k)}$ and $\mathbf{C}^{(k)}$ we only assure the structure of extriangulated categories with negative first extension. Hence, we can apply Theorem 5.14 and Lemma 3.2 (c) to the discussion above and thus we get the following result.

Corollary 5.18. *For $k \geq 1$, a triangulated category \mathcal{D} , a t -structure $\mathbf{u} = (\mathcal{U}, \mathcal{W})$ in \mathcal{D} with heart $\mathcal{H} = \mathcal{U} \cap \mathcal{W}$, and the extriangulated categories with negative first extension $\mathbf{H}^{(k)} := \mathcal{W} \cap \Sigma^{-k} \mathcal{U}$ and $\mathbf{C}^{(k)} := \mathcal{W} \cap \Sigma^{-k-1} \mathcal{U}$, the following statements hold true.*

- (a) $(\mathbf{H}^{(k)}, \Sigma^{-k-1} \mathcal{H}) \in \text{stors } \mathbf{C}^{(k)}$.
- (b) *For any $V \in \mathbf{C}^{(k)}$, the following statements are equivalent:*
 - (b1) V is an extended tilting object in $\mathbf{C}^{(k)}$;
 - (b2) $\mathbf{H}^{(k)}$ is a cogenerating class in $\mathbf{C}^{(k)}$ and V is a projective generator in $\mathbf{H}^{(k)}$.

Finally, we consider the following situation. Let \mathcal{D} be a triangulated category, $\mathbf{u} = (\mathcal{U}, \mathcal{W})$ a t -structure in \mathcal{D} and $k \geq 1$. Consider the s -torsion pairs in \mathcal{D} , namely: $\mathbf{v}_1^k := (\Sigma^{k+2} \mathcal{U}, \Sigma^{k+1} \mathcal{W})$, $\mathbf{v}_2^k := (\Sigma^2 \mathcal{U}, \Sigma \mathcal{W})$ and $\mathbf{v}_3^k := (\Sigma \mathcal{U}, \mathcal{W})$. Notice that $\mathbf{v}_3^k := \Sigma^{-1} \mathbf{v}_2^k$, $\Sigma \mathbf{v}_1^k \leq \mathbf{v}_1^k \leq \mathbf{v}_2^k \leq \mathbf{v}_3^k$ and $\mathbf{v}_1^k \leq \Sigma \mathbf{v}_2^k$. Moreover the interval $[\mathbf{v}_2^k, \mathbf{v}_3^k]$ is normal and we can not assure that for the other intervals. We introduce the classes

$$\mathbf{H}_{(k)} := \mathcal{H}_{[\mathbf{v}_1^k, \mathbf{v}_2^k]} = \Sigma^{k+1} \mathcal{W} \cap \Sigma^2 \mathcal{U} \quad \text{and} \quad \mathbf{C}_{(k)} := \mathcal{H}_{[\mathbf{v}_1^k, \mathbf{v}_3^k]} = \Sigma^{k+1} \mathcal{W} \cap \Sigma \mathcal{U}$$

of the t -structure \mathbf{u} . Observe that

$$\mathcal{H}_{[\mathbf{v}_2^k, \mathbf{v}_3^k]} = \Sigma \mathcal{W} \cap \Sigma \mathcal{U} = \Sigma \mathcal{H},$$

where $\mathcal{H} = \mathcal{W} \cap \mathcal{U}$ is the heart of \mathbf{u} , and thus $\mathcal{H}_{[\mathbf{v}_2^k, \mathbf{v}_3^k]}$ is an abelian category. For $\mathbf{H}_{(k)}$ and $\mathbf{C}_{(k)}$ we only assure the structure of extriangulated categories with negative first extension. Hence, we can apply Theorem 5.14 and Lemma 3.2 (c) to the discussion above and thus we get the following result.

Corollary 5.19. *For $k \geq 1$, a triangulated category \mathcal{D} , a t -structure $\mathbf{u} = (\mathcal{U}, \mathcal{W})$ in \mathcal{D} with heart $\mathcal{H} = \mathcal{U} \cap \mathcal{W}$, and the extriangulated categories with negative first extension $\mathbf{H}_{(k)} := \Sigma^{k+1} \mathcal{W} \cap \Sigma^2 \mathcal{U}$ and $\mathbf{C}_{(k)} := \Sigma^{k+1} \mathcal{W} \cap \Sigma \mathcal{U}$, the following statements hold true.*

- (a) $(\mathbf{H}_{(k)}, \Sigma \mathcal{H}) \in \text{stors } \mathbf{C}_{(k)}$.
- (b) *For any $V \in \mathbf{C}_{(k)}$, the following statements are equivalent:*
 - (b1) V is an extended tilting object in $\mathbf{C}_{(k)}$;
 - (b2) $\mathbf{H}_{(k)}$ is a cogenerating class in $\mathbf{C}_{(k)}$ and V is a projective generator in $\mathbf{H}_{(k)}$.

References

- [1] T. Adachi, H. Enomoto, M. Tsukamoto. Intervals of s -torsion pairs in extriangulated categories with negative first extensions. *Math. Proc. Camb. Philos. Soc.*, 174(3):451–469, 2023.

- [2] L. Angeleri Hügel, F. U. Coelho. Infinitely generated tilting modules of finite projective dimension. *Forum Math.*, 13(2):239–250, 2001.
- [3] L. Angeleri Hügel, F. Marks, J. Vitória. Silting modules. *Int. Math. Res. Not.* 2016(4):1251–1284, 2016.
- [4] A. Argudín-Monroy, D. Bravo, C.E. Parra. TTF classes generated by silting modules. *Preprint*, arXiv:2411.17581, 2024.
- [5] A. Argudín-Monroy, C. E. Parra. Universal co-extensions of torsion abelian groups. *J. Algebra*, 647:1–27, 2024.
- [6] M. Auslander, R.-O. Buchweitz. The homological theory of maximal Cohen-Macaulay approximations. *Mém. Soc. Math. Fr., Nouv. Sér.* 38:5–37, 1987.
- [7] S. Bazzoni, A characterization of n -cotilting and n -tilting modules. *J. Algebra*, 273(1):359–372, 2004.
- [8] A. Beilinson, J. Bernstein, P. Deligne. Faisceaux pervers. *Asterisque*, 100, 172 p. 1982.
- [9] A. Cipriani and J. Woolf. When are there enough projective perverse sheaves? *Glasg. Math. J.* 64(1):185–196, 2022.
- [10] L. Fiorot. n -quasi-abelian categories vs n -tilting torsion pairs. *Doc Math.* 26:149–197, 2021.
- [11] L. Frerick, D. Sieg. Exact categories in functional analysis. *Preprint*. 2010
- [12] M. Gorsky, H. Nakaoka, Y. Palu. Positive and negative extensions in extriangulated categories. *Preprint*, arXiv:2103.12482 [math.CT], 2021.
- [13] D. Happel, I. Reiten, S.O. Smalø. *Tilting in abelian categories and quasi-tilted algebras*. Providence, RI: American Mathematical Society (AMS), 1996.
- [14] J. He, Y. Hu, P. Zhou. Torsion pairs and recollements of extriangulated categories. *Comm. Algebra*, 50(5):2018–2036, 2022. DOI: 10.1080/00927872.2021.1996585
- [15] F. Kong, K. Song, P. Zhang. Decomposition of torsion pairs on module categories. *J. Algebra*, 388:248–267, 2013.
- [16] H. Krause, *Homological theory of representations*. Cambridge: Cambridge University Press, 2022
- [17] B. Mitchell, *Theory of Categories*. Academic Press, 1965.
- [18] H. Nakaoka, Y. Palu. Extriangulated categories, Hovey twin cotorsion pairs and model structures, *Cah. Topol. Geom. Differ. Categ.*, 60(2):117–193, 2019.

- [19] A. Neeman. *Triangulated categories*. Princeton, NJ: Princeton University Press, 2001.
- [20] C. E. Parra, M. Saorín. Direct limits in the heart of t-structure: the case of a torsion pair, *J. Pure Appl. Algebra*, 219(9):4117–4143, 2015.
- [21] C. E. Parra, M. Saorín, On hearts which are module categories. *J. Math. Soc. Japan*, 68(4):1421–1460, 2016.
- [22] C. E. Parra, M. Saorín, The HRS tilting process and Grothendieck hearts of t-structures. *Representations of algebras, geometry and physics, Maurice Auslander distinguished lectures and international conference, Woods Hole Oceanographic Institute, Woods Hole, MA, USA, April 25–30, 2018*. Providence, RI: American Mathematical Society (AMS):209–241, 2021.
- [23] C. E. Parra, M. Saorín, S. Virili. Tilting preenvelopes and cotilting precovers in general abelian categories. *Algebr. Represent. Theory*. 26(4):1087–1140, 2023.
- [24] C. E. Parra, M. Saorín, S. Virili. Locally finitely presented and coherent hearts. *Rev. Mat. Iberoam.* 39(1):201–268, 2023.
- [25] W. Rump. *-modules, tilting, and almost abelian categories. *Comm. Algebra*, 29(8): 3293–3325, 2001.
- [26] W. Rump. Almost abelian categories. *Cah. Topologie Géom. Différ. Catégoriques*, 42(3):163–225, 2001.
- [27] W. Rump. A counterexample to Raikov’s conjecture. *Bull. Lond. Math. Soc.*, 40(6):985–994, 2008.
- [28] J.P. Schneiders. *Quasi-abelian categories and sheaves*. SMF, Paris: Mém. Soc. Math. Fr., Nouv. Sér, 1998.
- [29] A. Tattar. Torsion pairs and quasi-abelian categories. *Algebr. Represent. Theory*, 24(6):1557–1581, 2021.
- [30] J. Woolf. Stability conditions, torsion theories and tilting, *J. Lond. Math. Soc.* 82, 3 (2010), pp. 663–682
- [31] B. Zhu, X. Zhuang. Tilting subcategories in extriangulated categories. *Front. Math. China*. 15(1):225–253, 2020.

Alejandro Argudín-Monroy
 Departamento de Matemáticas, Facultad de Ciencias,
 Universidad Nacional Autónoma de México,
 Circuito Exterior, Ciudad Universitaria,
 CDMX 04510, MÉXICO.
 argudin@ciencias.unam.mx

Octavio Mendoza Hernández
 Instituto de Matemáticas

Universidad Nacional Autónoma de México,
Circuito Exterior, Ciudad Universitaria,
CDMX 04510, MÉXICO.
`omendoza@matem.unam.mx`

Carlos Parra
Instituto de Ciencias Físicas y Matemáticas
Edificio Emilio Pugin, Campus Isla Teja
Universidad Austral de Chile
5090000 Valdivia, CHILE
`carlos.parra@uach.cl`