

THE DIRICHLET PROBLEM FOR THE PRESCRIBED CURVATURE EQUATIONS IN MINKOWSKI SPACE

BIN WANG

ABSTRACT. We study the Dirichlet problem for functions whose graphs are spacelike hypersurfaces with prescribed curvature in the Minkowski space and we obtain some new interior second order estimates for admissible solutions to the corresponding fully nonlinear elliptic partial differential equations.

CONTENTS

1. Introduction	1
2. Preliminaries	4
3. Global gradient estimates	7
4. The scalar curvature equation	10
5. The k -curvature equation	14
Acknowledgement	20
Data Availability	20
References	20

1. INTRODUCTION

The Dirichlet problem for the prescribed curvature equations in the Minkowski space is a physically motivated problem due to applications in the theory of relativity. However, the problem has not been fully explored in the literature compared to its Euclidean counterparts because the geometric nature of the ambient space causes substantial difficulties for deriving a priori estimates of admissible solutions. Hence, major advancements for this problem have stopped since the work of P. Bayard [2] and J. Urbas [46] in 2003, and only until very recent years C. Ren and Z. Wang [36, 37, 49] have made some important progress. In this note, we continue the investigation from the limited literature and obtain some new interior curvature estimates which also improve those already remarkable ones due to J. Urbas [46] and Z. Wang [49]. A more detailed literature review will soon be given below and our main results are stated in Theorem 1.2 and Theorem 1.3.

Remark 1.1. For literature in the Euclidean case, the reader is referred to the modern representative work of Guan-Ren-Wang [26] and Sheng-Urbas-Wang [42]; see also the very delicate work [24] of Guan-Spruck in hyperbolic space.

Let $\sigma_k : \mathbb{R}^n \rightarrow \mathbb{R}$ denote the k -th elementary symmetric polynomial, which is defined as

$$\sigma_k(x_1, \dots, x_n) = \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} x_{i_1} \cdots x_{i_k}.$$

2020 *Mathematics Subject Classification.* Primary 53C50, 53C21; Secondary 35B45, 35J60.

Key words and phrases. Spacelike hypersurfaces in the Minkowski space, prescribed curvature equations, curvature estimates, fully nonlinear elliptic PDEs.

We consider a fully nonlinear elliptic equation of the form

$$F[u] = f(\kappa_1(x), \dots, \kappa_n(x)) = \psi(x, u, Du), \quad x \in \Omega,$$

in a smooth bounded domain $\Omega \subseteq \mathbb{R}^n$, where f is a smooth symmetric function of n variables given by σ_k , ψ is a prescribed positive function and $\kappa = (\kappa_1, \dots, \kappa_n)$ denotes the principal curvatures of the graph of u over Ω .

The study of this class of fully nonlinear elliptic equations were initiated by Caffarelli-Nirenberg-Spruck [10–12] and Ivochkina [29–31], and further developed by Trudinger-Wang [43–45]. Indeed, the equation operator σ_k includes a large class of notable examples. For a smooth hypersurface Σ with principal curvatures $\kappa[\Sigma] = (\kappa_1, \dots, \kappa_n)$, the quantity $\sigma_k(\kappa[\Sigma])$ will be called the k -th mean curvature (or just k -curvature) of Σ . In particular, $\sigma_1(\kappa[\Sigma]) = \sum_{i=1}^n \kappa_i$ is the usual mean curvature, $\sigma_2(\kappa[\Sigma]) = \sum_{i < j} \kappa_i \kappa_j$ is the scalar curvature, and $\sigma_n(\kappa[\Sigma]) = \kappa_1 \cdots \kappa_n$ is the Gauss curvature. The other values of k also pertain to some important geometric problems in the sense that they can be reduced to solving some particular σ_k type equations; see e.g. [23] and [25, 55].

The aim of this note is to seek an admissible function u such that its graph $\Sigma = (x, u(x))$ over $\bar{\Omega}$ is a spacelike hypersurface in the Minkowski space and solves the following Dirichlet problem

$$(1.1) \quad \begin{aligned} F[u] &= \psi(x, u, Du) && \text{in } \Omega, \\ u &= \varphi && \text{on } \partial\Omega, \end{aligned}$$

where the meaning of being spacelike and admissible are specified in section 2.

The problem was first studied by Robert Bartnik and Leon Simon in their influential paper [1] for the case of prescribing mean curvature i.e. when $k = 1$ in (1.1); see also the extension by Gerhardt [18] to a general Lorentzian product manifold. The motivations mainly came from Einstein's theory of relativity, as first emphasized in the fundamental paper of Lichnerowicz [34]. Roughly speaking, spacelike hypersurfaces of constant mean curvature in the Minkowski space are important because they provide Riemannian submanifolds with properties which reflect those of the spacetime. In particular, they played a role in the initial proof [41] of the positive mass conjecture by Schoen and Yau. On the other hand, the study of such hypersurfaces was already posed by Calabi [13] at an earlier time for the quest of understanding its Bernstein type property; see the resolution of this problem by Cheng-Yau [14] and references citing their papers.

Later, (1.1) was solved by Delanoë [16] for $k = n$ i.e. the prescribed Gauss curvature equation; see also the work of Guan [22] in which the result was proved under a subsolution condition. The next interesting case would naturally be to solve the prescribed scalar curvature equation i.e. when $k = 2$ in (1.1). Bayard [2] was the first to tackle the problem and proved the solvability in dimensions three and four. Soon after that, Urbas [46] extended this result to all dimensions, however, Urbas' proof relied crucially on the additional assumption that both the boundary data φ and the domain are uniformly convex in the sense that $D^2\varphi \geq c_0 I$ in $\bar{\Omega}$ for some uniform constant $c_0 > 0$ and the principal curvatures of $\partial\Omega$ are bounded from below by a positive uniform constant. Our first main result is the removal of this assumption and the following improved existence theorem is obtained.

Theorem 1.2. *Let $\Omega \subseteq \mathbb{R}^n$ be a convex admissible bounded domain with a smooth boundary. Suppose $\psi(x, u, Du)$ is a smooth positive function satisfying $\psi_u \geq 0$ and $\varphi \in C^4(\bar{\Omega})$ is spacelike. Assume the existence of an admissible subsolution \underline{u} such that*

$$\begin{aligned} \sigma_2[\underline{u}] &\geq \psi(x, \underline{u}, D\underline{u}) && \text{in } \Omega, \\ \underline{u} &= \varphi && \text{on } \partial\Omega. \end{aligned}$$

Then there exists a unique admissible solution u to

$$(1.2) \quad \begin{aligned} \sigma_2[u] &= \psi(x, u, Du) && \text{in } \Omega, \\ u &= \varphi && \text{on } \partial\Omega, \end{aligned}$$

belonging to $C^{3,\alpha}(\bar{\Omega})$ for any $\alpha \in (0, 1)$.

The next goal is to continue the investigation for the remaining cases $3 \leq k \leq n - 1$ and we shall proceed by the standard continuity method along with the regularity theorem due to Evans-Krylov [17, 32]; this requires us to establish a priori estimates for admissible solutions up to the second order. In fact, the C^0 estimate follows directly from the comparison principle and the C^1 estimate has been successfully obtained by Bayard [2, Proposition 3.1] for all k . For the second order derivatives on the boundary $\partial\Omega$, Bayard [2, Section 4.2] proved the bound for all k if $\varphi = \text{const}$ and for a general boundary data, the condition $k = 2$ had to be imposed. Although the boundary C^2 estimate has not been obtained in the most general case, it is good enough for the moment. The real issue is the second order estimate in Ω which was not known for $3 \leq k \leq n - 1$.

The major difficulties come from two aspects. First, the operator σ_k is a lot more structurally complicated when $3 \leq k \leq n - 1$. As a comparison, when $k = 1$ the equation is quasilinear and when $k = n$ the equation is of Monge-Ampère type, both of which are extensively studied and many techniques could be adapted in our setting. Even when $k = 2$, there are structural advantages that were utilized by Bayard [2] and Urbas [46] but failed to hold for $k \geq 3$. The second major obstacle occurs in the interchanging formula (5.6) which gives rise to a negative curvature term. In contrast, the formula yields a positive curvature term for the Euclidean case which is rather crucial as demonstrated in [42].

In two recent papers [36, 37], C. Ren and Z. Wang proved two powerful concavity inequalities for the σ_{n-1} and σ_{n-2} operators by exploiting their algebraic structures in depth, and then they were able to overcome the difficulties and solve (1.1) for $k = n - 1$ and $k = n - 2$. However, it is not feasible to generalize their method for other k 's. In [28], Huang assumed $\psi = \psi(x, u, Du)$ has a special dependence on the gradient terms so that an extra positive curvature term could be extracted from the twice differentiation of the equation. Huang's method is inspiring but the assumptions are not applicable when e.g. $\psi = \psi(x, u)$ does not contain the gradient terms at all.

Now the ultimate goal is reduced to derive second order estimates in Ω for $3 \leq k \leq n - 3$ and general ψ . Z. Wang [49] has achieved the goal for admissible solutions whose graphs are $(k + 1)$ -convex. Our second main result establishes the estimates for admissible solutions whose graphs are semi-convex, which may attract more attention than theorem 1.2.

Theorem 1.3. *Let $2 \leq k \leq n - 1$ and let $\Omega \subseteq \mathbb{R}^n$ be a convex admissible domain with a smooth boundary. Assume $\psi(x, u, Du) \in C^2(\bar{\Omega} \times \mathbb{R} \times \mathbb{R}^n)$ is positive and $\varphi \in C^2(\bar{\Omega})$ is spacelike. Suppose $u \in C^4(\Omega) \cap C^2(\bar{\Omega})$ is an admissible solution to (1.1) that has a semi-convex graph Σ i.e. the principal curvatures $\kappa = (\kappa_1, \dots, \kappa_n)$ of Σ satisfy*

$$(1.3) \quad \kappa_i(x) \geq -K \quad \text{for all } x \in \Omega \text{ and all } 1 \leq i \leq n$$

for some $K > 0$. Then

$$\max_{\Omega} \kappa_{\max}(x) \leq C \left(1 + \max_{\partial\Omega} \kappa_{\max}(x) \right)$$

for some $C > 0$ depending on $n, k, K, \|u\|_{C^1(\bar{\Omega})}, \|\psi\|_{C^2(\mathcal{D})}$ and $\|\varphi\|_{C^1(\mathcal{D})}$, where

$$\mathcal{D} := \bar{\Omega} \times \left[\inf_{\Omega} u, \sup_{\Omega} u \right] \times \mathbb{R}^n.$$

Remark 1.4. The new contribution here is that our estimate improves that of Z. Wang [49] in the sense that semi-convexity is weaker than $(k + 1)$ -convexity; for a proof of this fact, the reader is referred to [33, Lemma 7] or [48, Lemma 2.13].

Remark 1.5. If one could obtain the curvature estimate without the additional semi-convexity assumption (1.3), then we would be able to conclude the solvability of the Dirichlet problem (1.1) for all k , at least for a constant boundary data.

In [37], C. Ren and Z. Wang had conjectured that their concavity inequality should hold for all k with $2k > n$. Hence, the desired curvature estimate is expected to hold for all k with $2k > n$, although a verification for their conjecture is still absent.

Conjecture 1.6. *Let $2 \leq k \leq n - 1$ satisfy $2k > n$ and let $\Omega \subseteq \mathbb{R}^n$ be a convex admissible bounded domain with a smooth boundary. Suppose $\psi(x, u, Du)$ is a smooth positive function satisfying $\psi_u \geq 0$ and $\varphi \in C^4(\overline{\Omega})$ is spacelike. Assume the existence of an admissible subsolution \underline{u} such that*

$$\begin{aligned} F[\underline{u}] &\geq \psi(x, \underline{u}, D\underline{u}) && \text{in } \Omega, \\ \underline{u} &= \varphi && \text{on } \partial\Omega. \end{aligned}$$

Then there exists a unique admissible solution u to (1.1) belonging to $C^{3,\alpha}(\overline{\Omega})$ for any $\alpha \in (0, 1)$.

To conclude the introduction, we mention some related research. In [40], Schnürer considered the Dirichlet problem in a general Lorentzian product manifold for a class of curvature functions which excludes the operator $\sigma_k(\kappa)$ concerned here. On the other hand, Gerhardt [19–21] studied closed hypersurfaces of prescribed curvature in Lorentzian manifolds as well. Indeed, much of our proof remains valid in a more generic setting, however, since the problem has not been fully solved even in some very simple cases, it might not be worthwhile to create more complications at this moment. For literature on entire spacelike hypersurfaces, the reader may be referred to the series of work [3–8, 38, 39, 50–54] by P. Bayard, Ph. Delanoë, C. Ren, A. Seppi, O. C. Schnürer, Z. Wang, L. Xiao. We also wish to call attention to a recent paper [27] of Guo-Jiao in which they studied the same Dirichlet problem for a different class of fully nonlinear equations.

The rest of this note is organized as follows. In section 2, we review preliminary concepts and fix notations. In section 3, we derive global gradient estimates for admissible solutions which extend the ones obtained by Bayard [2] in the sense that our estimates hold for a right-hand side $\psi = \psi(x, u, Du)$ that may contain gradient terms and we have replaced the assumption that both Ω and φ are strictly convex by the subsolution condition. In section 4, we prove curvature estimates for admissible solutions to the scalar curvature equation in all dimensions without assuming uniform convexity on Ω and φ , hence improving Urbas' result [46] and theorem 1.2 follows accordingly. In section 5, we prove theorem 1.3 which extends the estimate due to Z. Wang [49]. It might be noteworthy that in both section 4 and section 5, instead of ordinary curvature estimates, we actually have proved Pogorelov type interior curvature estimates assuming the boundary data is affine.

2. PRELIMINARIES

In this section, we review some basic concepts and fix notations which will be used throughout the subsequent sections but without directly quoting every time.

We shall always assume Ω is an admissible domain.

Definition 2.1. A smooth bounded domain $\Omega \subseteq \mathbb{R}^n$ is said to be admissible if at least $k - 1$ principal curvatures of $\partial\Omega$ are positive at each boundary point.

Lemma 2.2. *Every affine spacelike boundary data φ on $\partial\Omega$ has an admissible extension to $\overline{\Omega}$ if and only if the smooth bounded domain Ω is convex and admissible.*

Proof. See [2, Lemma 2.1]. □

Now recall that the Minkowski space $\mathbb{R}^{n,1}$ is the Euclidean space \mathbb{R}^{n+1} equipped with the metric

$$ds^2 = dx_1^2 + \cdots + dx_n^2 - dx_{n+1}^2.$$

Definition 2.3. A smooth hypersurface Σ in $\mathbb{R}^{n,1}$ is said to be k -convex if its principal curvatures

$$\kappa[\Sigma] \in \Gamma_k = \{\kappa \in \mathbb{R}^n : \sigma_j(\kappa) > 0 \quad \forall 1 \leq j \leq k\}$$

at every point; it is said to be spacelike if for every $p \in \Sigma$, the induced metric $\langle \cdot, \cdot \rangle_{T_p \Sigma}$ on the tangent space is positive definite.

Let Σ be the hypersurface in $\mathbb{R}^{n,1}$ given as the graph of a smooth function $u : \bar{\Omega} \rightarrow \mathbb{R}$. The induced metric and the second fundamental form of Σ are then

$$g_{ij} = \delta_{ij} - u_i u_j, \quad h_{ij} = \frac{u_{ij}}{\sqrt{1 - |Du|^2}}.$$

Remark 2.4. The graph of u over $\bar{\Omega}$ is a spacelike hypersurface in $\mathbb{R}^{n,1}$ if and only if

$$(2.1) \quad \sup_{\bar{\Omega}} |Du| < 1.$$

Thus, we may as well say a function u is spacelike in $\bar{\Omega}$ if (2.1) holds.

The unit normal vector field to Σ is

$$\nu = \frac{(Du, 1)}{\sqrt{1 - |Du|^2}}.$$

Note that we shall use $Du = (u_1, \dots, u_n)$ and $D^2u = (u_{ij})$ to denote the ordinary gradient vector and the ordinary Hessian matrix. On the other hand, for a chosen local orthonormal frame $\{e_1, \dots, e_n\}$ on $T\Sigma$, the symbol ∇ will denote the induced Levi-Civita connection on Σ . For a smooth function u on Σ , we set $\nabla_i u = \nabla_{e_i} u$ and $\nabla_{ij} u = \nabla^2 u(e_i, e_j)$. The norm of ∇u with respect to g_{ij} is then

$$|\nabla u| = \sqrt{g^{ij} u_i u_j} = \frac{|Du|}{\sqrt{1 - |Du|^2}},$$

where

$$g^{ij} = \delta_{ij} + \frac{u_i u_j}{1 - |Du|^2}$$

is the inverse of g_{ij} .

We also recall the following fundamental formulas for hypersurfaces in $\mathbb{R}^{n,1}$.

(Gauss formula)	$\nabla_{ij} X = h_{ij} \nu,$
(Weingarten formula)	$\nabla_i \nu = h_{ij} e_j,$
(Codazzi equation)	$\nabla_k h_{ij} = \nabla_j h_{ik},$
(Gauss equation)	$R_{ijst} = -(h_{is} h_{jt} - h_{it} h_{js}).$

For a symmetric matrix $A = (a_{ij})$ and an operator

$$F : \{\text{symmetric matrices}\} \rightarrow \mathbb{R},$$

we define

$$F^{ij}(A) = \frac{\partial F}{\partial a_{ij}}, \quad F^{ij,rs} = \frac{\partial^2 F}{\partial a_{ij}(A) \partial a_{rs}}.$$

When $F(A) = f(\lambda(A))$ depends only on the eigenvalues and when A is diagonal, we have $F^{ij} = f_i \delta_{ij}$ where

$$f_i = \frac{\partial f}{\partial \lambda_i}.$$

Moreover, we have

$$\sum_{i,j} F^{ij} a_{ij} = \sum_{i=1}^n f_i(\lambda(A)) \lambda_i, \quad \sum_{i,j,k} F^{ij} a_{ik} a_{jk} = \sum_{i=1}^n f_i(\lambda(A)) \lambda_i^2.$$

In this article, we are considering an equation of the form

$$F(A) = f(\lambda(A)) = \psi(x, u, Du),$$

where $f = \sigma_k$ and $A = (a_{ij})$ is given by

$$a_{ij} = \frac{1}{w} \gamma^{ik} u_{kl} \gamma^{lj}, \quad \gamma^{ik} = \delta_{ik} + \frac{u_i u_k}{w(1+w)},$$

since the principal curvatures $\kappa = (\kappa_1, \dots, \kappa_n)$ of Σ are eigenvalues of the following matrix

$$\frac{1}{w} \left(I + \frac{Du \otimes Du}{w^2} \right) D^2 u, \quad w = \sqrt{1 - |Du|^2}.$$

To solve our equation, we need also to define the notion of admissible solutions.

Definition 2.5. A function $u \in C^\infty(\Omega) \cap C(\overline{\Omega})$ is said to be admissible if its graph $\Sigma = (x, u(x))$ over $\overline{\Omega}$ is a k -convex spacelike hypersurface in $\mathbb{R}^{n,1}$. Equivalently, u is admissible if (2.1) holds and

$$\sigma_j[u] > 0 \quad \text{for all } x \in \overline{\Omega} \text{ and all } 1 \leq j \leq k.$$

Remark 2.6. The homogenized equation operator $(F[u])^{1/k}$ is concave with respect to second derivatives for admissible solutions. This is a key condition to invoke the Evan-Krylov device [17, 32].

Finally, we state some commonly used properties of the σ_k operator.

Notation 2.7. Observe that

$$\frac{\partial}{\partial x_i} \sigma_k(x) = \sigma_{k-1}(x) \Big|_{x_i=0} = \sigma_{k-1}(x_1, \dots, x_{i-1}, 0, x_{i+1}, \dots, x_n).$$

Throughout the article, we will use $\sigma_{k-1}(x|i)$ or $\sigma_k^{ii}(x)$ interchangeably to denote the first order derivatives. The notations $\sigma_{k-2}(x|ij)$ or $\sigma_k^{ii,jj}(x)$ are defined in a similar way for second order derivatives.

Lemma 2.8. For all $1 \leq k \leq n$ and $x \in \mathbb{R}^n$, we have

$$\begin{aligned} \sigma_k(x) &= x_i \sigma_{k-1}(x|i) + \sigma_k(x|i), \\ \sum_{i=1}^n x_i \sigma_{k-1}(x|i) &= k \sigma_k(x), \\ \sum_{i=1}^n \sigma_{k-1}(x|i) &= (n - k + 1) \sigma_{k-1}(x). \end{aligned}$$

Moreover, if $x_1 \geq \dots \geq x_n$ and $x \in \Gamma_k$, then

$$\sigma_k^{11}(x) \cdot x_1 \geq \frac{k}{n} \sigma_k(x).$$

Lemma 2.9 (Maclaurin's inequality). Let $2 \leq k \leq n$ and suppose $x = (x_1, \dots, x_n) \in \Gamma_k$. Denote by

$$H_k := \binom{n}{k}^{-1} \sigma_k.$$

Then we have

$$H_k^{1/k}(x) \leq \dots \leq H_2^{1/2}(x) \leq H_1(x).$$

Lemma 2.10. Let $\kappa = (\kappa_1, \dots, \kappa_n) \in \Gamma_k$. Suppose $\kappa_j \leq 0$ for some $1 \leq j \leq n$. Then

$$\sigma_k^{jj}(\kappa) \geq C(n, k) \sum_{i=1}^n \sigma_k^{ii}(\kappa)$$

and

$$\kappa_j \geq -\frac{n-k}{k} \kappa_1.$$

Proof. The first inequality follows from Lemma 2.8 by looking at

$$\sigma_{k-1} = \kappa_j \sigma_{k-1}^{jj}(\kappa) + \sigma_{k-1}(\kappa|j).$$

For the second inequality, see [37, Lemma 11]. \square

3. GLOBAL GRADIENT ESTIMATES

In this section, we obtain global gradient estimates when the prescribed function $\psi = \psi(x, u, Du)$ is allowed to contain gradient terms. Previously, this result was achieved by Bayard [2, Proposition 3.1] for $\psi = \psi(x)$. We first recall the standard comparison principle.

Lemma 3.1. *Let $u, v \in C^2(\Omega) \cap C(\bar{\Omega})$ be spacelike in Ω . Assume u is moreover admissible in Ω and the positive prescribed function $\psi(x, z, p) \in C^2(\Omega \times \mathbb{R} \times \mathbb{R}^n)$ satisfies $\psi_z \geq 0$. If $u \leq v$ on $\partial\Omega$ and*

$$\sigma_k[u] \geq \psi(x, u, Du), \quad \sigma_k[v] \leq \psi(x, v, Dv) \quad \text{in } \Omega,$$

then $u \leq v$ in Ω .

Proof. See e.g. [2, Theorem 5.1], [12, Lemma A], or [30, Theorem 3.1]. \square

By the work of Bartnik-Simon [1], there exists an admissible function \bar{u} such that

$$\begin{aligned} \sigma_1[\bar{u}] &= \left[\binom{n}{k}^{-1} \psi(x, \bar{u}, D\bar{u}) \right]^{1/k} \quad \text{in } \Omega, \\ \bar{u} &= \varphi \quad \text{on } \partial\Omega. \end{aligned}$$

Hence, it follows from the Maclaurin's inequality

$$\sigma_k[\bar{u}] \leq \binom{n}{k} \cdot (\sigma_1[\bar{u}])^k = \psi(x, \bar{u}, D\bar{u})$$

and so

$$u \leq \bar{u} \quad \text{in } \Omega$$

by the comparison principle.

Similarly, by assuming the existence of an admissible subsolution i.e. some admissible function \underline{u} such that

$$\begin{aligned} \sigma_k[\underline{u}] &\geq \psi(x, \underline{u}, D\underline{u}) \quad \text{in } \Omega, \\ \underline{u} &= \varphi \quad \text{on } \partial\Omega, \end{aligned}$$

we have

$$(3.1) \quad \underline{u} \leq u \leq \bar{u} \quad \text{in } \Omega.$$

Consequently, by the Hopf lemma, for the interior normal derivative at any point on $\partial\Omega$, we have

$$\frac{\partial \underline{u}}{\partial \gamma} \leq \frac{\partial u}{\partial \gamma} \leq \frac{\partial \bar{u}}{\partial \gamma}.$$

Thus, we conclude that

$$(3.2) \quad \sup_{\partial\Omega} |Du| \leq \max\{\sup_{\partial\Omega} |D\underline{u}|, \max_{\partial\Omega} |D\bar{u}|\} \leq 1 - \theta_0$$

for some $\theta_0 \in (0, 1)$ depending on \underline{u} , \bar{u} and $\text{diam}(\Omega)$.

Theorem 3.2. *Let $2 \leq k \leq n - 1$. Suppose $\psi(x, z, p) \in C^\infty(\bar{\Omega} \times \mathbb{R} \times \mathbb{R}^n)$ is positive and $\psi_z \geq 0$. Assume further the existence of an admissible subsolution. If $u \in C^3(\Omega) \cap C^1(\bar{\Omega})$ is an admissible solution of (1.1), then*

$$(3.3) \quad \sup_{\bar{\Omega}} |Du| \leq 1 - \theta$$

for some $\theta \in (0, 1)$ depending on $n, k, \theta_0, \sup_{\bar{\Omega}} |u|, \sup_{\mathcal{D}} |D\psi|$ and $\inf_{\mathcal{D}} \psi$ where $\mathcal{D} := \bar{\Omega} \times [\inf_{\Omega} u, \sup_{\Omega} u] \times \mathbb{R}^n$ and θ_0 is the constant in (3.2).

Proof. According to (3.2), it is sufficient to estimate the quantity

$$\tilde{w} := \frac{1}{w} = \frac{1}{\sqrt{1 - |Du|^2}}$$

and prove that

$$\sup_{\bar{\Omega}} \tilde{w} \leq C \left(1 + \sup_{\partial\Omega} \tilde{w} \right)$$

for some $C > 0$ depending on the known constants. As in [2, 27], we may consider the function

$$\tilde{Q} = \tilde{w} e^{Bu},$$

where B is a positive constant to be determined later. If \tilde{Q} achieves its maximum on $\partial\Omega$ then we have the bound

$$\sup_{\bar{\Omega}} w \leq \exp \left[B \cdot 2 \sup_{\bar{\Omega}} |u| \right] \sup_{\partial\Omega} w$$

and we are through. Suppose this is not the case and \tilde{Q} attains its maximum at some interior point $x_0 \in \Omega$. By rotating the standard coordinates $\{\epsilon_1, \dots, \epsilon_{n+1}\}$ of \mathbb{R}^{n+1} if necessary, we may assume that

$$u_1(x_0) = |Du(x_0)| > 0 \quad \text{and} \quad u_j(x_0) = 0 \quad \text{for } j \geq 2.$$

By further rotating $\{\epsilon_2, \dots, \epsilon_n\}$, we may also assume $\{u_{ij}(x_0)\}$ is diagonal for $i, j \geq 2$. Pick an orthonormal frame $\{e_1, \dots, e_n\}$ around the point $X_0 = (x_0, u(x_0))$ e.g.

$$e_i = \gamma^{is} \tilde{\partial}_s, \quad \gamma^{is} = \delta_{is} + \frac{u_i u_s}{w(1+w)}, \quad \tilde{\partial}_s = \epsilon_s + u_s \epsilon_{n+1},$$

so that

$$\nabla_1 u = \frac{|Du|}{w} = |\nabla u|, \quad \nabla_i u = u_i = 0 \quad \text{for } i \geq 2.$$

Then, by the Weingarten formula, we have

$$\nabla_i \tilde{w} = -\nabla_i \langle \nu, \epsilon_{n+1} \rangle = -\langle h_{ij} e_j, \epsilon_{n+1} \rangle = -h_{ij} \langle \gamma^{js} \tilde{\partial}_s, \epsilon_{n+1} \rangle = h_{ij} \nabla_j u.$$

Taking the logarithm, the function

$$Q := \log \tilde{Q} = \log \tilde{w} + Bu$$

also attains its maximum at x_0 . Consequently, at x_0 , we have

$$(3.4) \quad 0 = \nabla_i Q = \frac{\nabla_i \tilde{w}}{\tilde{w}} + B \nabla_i u = \frac{h_{1i} \nabla_1 u}{\tilde{w}} + B \nabla_i u$$

from which it follows that at x_0 ,

$$(3.5) \quad h_{11} = -B \tilde{w} \quad \text{and} \quad h_{1i} = 0 \quad \text{for } i \geq 2.$$

In the remaining part of the proof, all subsequent calculations are carried out at the point X_0 without explicitly saying so.

Since

$$h_{11} = \frac{u_{11}}{w^3} \quad \text{and} \quad h_{ij} = \frac{u_{ij}}{w} \quad \text{for } i, j \geq 2,$$

the matrix $\{h_{ij}\}$ is diagonal. Hence

$$F^{ij} := \frac{\partial F}{\partial h_{ij}} = \frac{\partial \sigma_k}{\partial \kappa_i} \delta_{ij}$$

and we calculate

$$\begin{aligned} (3.6) \quad 0 &\geq F^{ii} \nabla_{ii} Q \\ &= F^{ii} \frac{\nabla_i h_{i1} \nabla_1 u + h_{i1} \nabla_{i1} u}{\tilde{w}} - F^{ii} \frac{(h_{i1} \nabla_1 u)^2}{\tilde{w}^2} + B F^{ii} \nabla_{ii} u \\ &= F^{ii} h_{i1i} \frac{\nabla_1 u}{\tilde{w}} + F^{11} h_{11}^2 - B^2 F^{11} |Du|^2 + Bk\psi\tilde{w}, \end{aligned}$$

where we have used (3.5) and the Gauss formula

$$\nabla_{ij} u = -\nabla_{ij} \langle X, \epsilon_{n+1} \rangle = -\langle \nabla_{ij} X, \epsilon_{n+1} \rangle = -h_{ij} \langle \nu, \epsilon_{n+1} \rangle = \tilde{w} h_{ij}.$$

For the first term in (3.6), we differentiate $F(h_{ij}) = \psi$ and invoke the Codazzi equation and the Weingarten formula,

$$\begin{aligned} F^{ii} h_{i1i} &= F^{ii} h_{ii1} = \nabla_1 \psi = \psi_{x_j} \nabla_1 x_j + \psi_z \nabla_1 u + \partial_{p_k} \psi \nabla_1 \nu_k \\ &= \frac{\psi_{x_1}}{w} + \psi_z \frac{|Du|}{w} + \partial_{p_1} \psi \frac{h_{11}}{w} \\ &= \frac{\psi_{x_1}}{w} + \psi_z \frac{|Du|}{w} - \frac{B}{w^2} \partial_{p_1} \psi \\ &\geq \tilde{w} \psi_{x_1} + \tilde{w} \psi_z |Du| - B \tilde{w}^2 \partial_{p_1} \psi. \end{aligned}$$

Substituting this back into (3.6), we have

$$0 \geq \tilde{w} \psi_{x_1} |Du| + \tilde{w} \psi_z |Du|^2 - B \tilde{w}^2 |Du| \partial_{p_1} \psi + B^2 \tilde{w}^2 F^{11} - B^2 |Du|^2 F^{11} + Bk\psi\tilde{w}.$$

Since u is spacelike i.e. $|Du| < 1$, it follows that

$$\begin{aligned} &\tilde{w} \psi_{x_1} |Du| + \tilde{w} \psi_z |Du|^2 - B \tilde{w}^2 |Du| \partial_{p_1} \psi \\ &\geq -CB \tilde{w}^2 \end{aligned}$$

for some $C > 0$ depending on $\sup_{\mathcal{D}} |D\psi|$. Moreover, since $h_{11} < 0$ due to (3.5), we have from lemma 2.10 and the Maclaurin's inequality that

$$F^{11} \geq C(n, k) \sum_{i=1}^n F^{ii} \geq C(n, k, \inf \psi),$$

which leads to (if e.g. $\tilde{w} \geq 2$)

$$\begin{aligned} 0 &\geq -CB \tilde{w}^2 + \frac{B^2}{2} \tilde{w}^2 F^{11} \\ &\geq CB \tilde{w}^2 (B - C) \end{aligned}$$

for some $C > 0$ depending on $n, k, \inf_{\mathcal{D}} \psi$, and $\sup_{\mathcal{D}} |D\psi|$.

Now, if $B > 0$ was fixed to be a large number, then either we reach a contradiction here and the maximum is attained on the boundary, or $\tilde{w}(x_0) \leq C$ somewhere in the derivation above when we assumed $\tilde{w} \geq C$ is sufficiently large; in that case, we would have

$$\sup_{\bar{\Omega}} \tilde{w} \leq \tilde{w}(x_0) \exp \left[B \cdot 2 \sup_{\bar{\Omega}} |u| \right].$$

The proof is now complete. \square

4. THE SCALAR CURVATURE EQUATION

The curvature bound for the scalar curvature equation was first obtained by Bayard [2] in dimension three for $\psi = \psi(x)$ and in dimension four for $\psi = \text{const}$. It was then extended by Urbas [46] to all dimensions for $\psi = \psi(x, u)$. We remark that Urbas' proof in fact works for the case $\psi = \psi(x, u, Du)$ as well. However, as Urbas commented in [46], the proof relied crucially on the additional assumption that the boundary data φ is uniformly convex i.e. $D^2\varphi \geq c_0 I$ uniformly for some positive constant c_0 . The cruciality of this condition is illustrated in our Remark 4.4 below.

In this section, we improve the curvature estimates for the scalar curvature equation by removing the assumption of uniform convexity.

Theorem 4.1. *Let $\varphi \in C^2(\Omega) \cap C(\bar{\Omega})$ be spacelike and affine. Suppose $\psi(x, z, p) \in C^\infty(\bar{\Omega} \times \mathbb{R} \times \mathbb{R}^n)$ is positive. If $u \in C^4(\Omega) \cap C^2(\bar{\Omega})$ is an admissible solution to (1.2), then the maximum principal curvature*

$$\kappa_{\max}(x) := \max_{1 \leq i \leq n} \kappa_i(x)$$

of its graph Σ satisfies

$$\sup_{\Omega'} \kappa_{\max} \leq C(\Omega')$$

for any $\Omega' \subset\subset \Omega$, where $C(\Omega') > 0$ depends on $n, \theta, \|\psi\|_{C^2(\mathcal{D})}, \inf_{\mathcal{D}} \psi$, and $\|\varphi\|_{C^1(\bar{\Omega})}$; here θ is the constant in (3.3) and

$$\mathcal{D} := \bar{\Omega} \times [\inf_{\Omega} u, \sup_{\Omega} u] \times \mathbb{R}^n.$$

Remark 4.2. It is sufficient to assume φ is spacelike and satisfies $\sigma_2[\varphi] < \sigma_2[u]$ in Ω .

Proof. Denote by

$$\tilde{w} := \frac{1}{w} = \frac{1}{\sqrt{1 - |Du|^2}}.$$

We consider the quantity

$$\tilde{W}(X, \xi) = \eta^\beta h_{\xi\xi} \exp\left(\frac{\alpha}{2}|X|^2\right)$$

for $X \in \Sigma$ and $\xi \in T_X \Sigma$, where $\eta = \varphi - u$, and $\alpha, \beta > 0$ are some large constants to be chosen later.

Suppose the maximum of \tilde{W} is attained at some interior point $X_0 = (x_0, u(x_0)) \in \Sigma$ and some $\xi_0 \in T_{X_0} \Sigma$. We choose a local orthonormal frame $\{e_1, \dots, e_n\}$ around X_0 such that

$$\xi_0 = e_1, \quad \nabla_{e_i} e_j = 0 \quad \text{at } X_0.$$

We may also assume the second fundamental form $h_{ij} = \kappa_i \delta_{ij}$ is diagonal at X_0 with principal curvatures ordered as

$$\kappa_1 \geq \kappa_2 \geq \dots \geq \kappa_n.$$

Let $\zeta = e_1$. Then the function

$$W(X) = \eta^\beta h_{ab} \zeta_a \zeta_b$$

is defined near X_0 and attains an interior maximum also at X_0 . By our special choice of the frame, we find that $Z := h_{ab} \zeta_a \zeta_b$ satisfies

$$\nabla_i Z = \nabla_i h_{11} \quad \text{and} \quad \nabla_i \nabla_j Z = \nabla_i \nabla_j h_{11} \quad \text{at } X_0.$$

Therefore, by working with $\log W$, we obtain at X_0

$$(4.1) \quad 0 = \beta \frac{\nabla_i \eta}{\eta} + \frac{h_{11i}}{h_{11}} + \alpha \langle X, e_i \rangle,$$

$$(4.2) \quad 0 \geq \beta \frac{\nabla_{ii} \eta}{\eta} - \beta \left(\frac{\nabla_i \eta}{\eta} \right)^2 + \frac{h_{11ii}}{\kappa_1} - \frac{h_{11i}^2}{\kappa_1^2} + \alpha (1 + h_{ii} \langle X, \nu \rangle).$$

In what follows, we will carry out all calculations at the point X_0 without explicitly indicating so. Now, we shall contract (4.2) with

$$F^{ij} := \frac{\partial \sigma_k^{1/k}}{\partial h_{ij}}$$

and estimate term by term.

Remark 4.3. It is here and only here that we use the homogenized operator in order to invoke its concavity property more conveniently.

Recall the interchanging formula [46, Lemma 2.1], we have

$$\begin{aligned} F^{ij} \nabla_i \nabla_j h_{11} &= -F^{ij,kl} \nabla_1 h_{ij} \nabla_1 h_{kl} - F^{ij} h_{ij} \kappa_1^2 \\ &\quad + F^{ij} h_{ik} h_{jk} \kappa_1 + \nabla_1 \nabla_1 \psi \\ (4.3) \quad &= -F^{ij,kl} \nabla_1 h_{ij} \nabla_1 h_{kl} - F \kappa_1^2 \\ &\quad + \kappa_1 \sum_{i=1}^n F^{ii} \kappa_i^2 + \nabla_1 \nabla_1 \psi. \end{aligned}$$

For the first term, we apply [46, Lemma 2.2] in addition to concavity of $\sigma_k^{1/k}$ to obtain that

$$-F^{ij,kl} h_{ij1} h_{kl1} = \sum_{i \neq j} \frac{F^{ii} - F^{jj}}{\kappa_i - \kappa_j} h_{ij1}^2 \geq 2 \sum_{i \neq 1} \frac{F^{ii} - F^{11}}{\kappa_1 - \kappa_i} h_{11i}^2.$$

For the last term in (4.3), since ψ contains the gradient, we have

$$(4.4) \quad \nabla_1 \nabla_1 \psi \geq -C(1 + \kappa_1^2).$$

Next, we consider the term $F^{ii} \nabla_{ii} \eta$. To compute this we may extend φ to be constant in the ϵ_{n+1} direction. Then, since $u = X_{n+1}$ on Σ , we have

$$\begin{aligned} \nabla_i \nabla_j \eta &= \sum_{\alpha=1}^n \frac{\partial \varphi}{\partial X_\alpha} \nabla_i \nabla_j X_\alpha + \sum_{\alpha, \gamma=1}^n \frac{\partial^2 \varphi}{\partial X_\alpha \partial X_\gamma} \nabla_i X_\alpha \nabla_j X_\gamma - \nabla_i \nabla_j X_{n+1} \\ &= \sum_{\alpha=1}^n \frac{\partial \varphi}{\partial X_\alpha} h_{ij} \nu_\alpha + \sum_{\alpha, \gamma=1}^n \frac{\partial^2 \varphi}{\partial X_\alpha \partial X_\gamma} \nabla_i X_\alpha \nabla_j X_\gamma - h_{ij} \nu_{n+1} \end{aligned}$$

from which it follows that

$$(4.5) \quad F^{ii} \nabla_{ii} \eta \geq \left(\sum_{\alpha=1}^n \frac{\partial \varphi}{\partial X_\alpha} \nu_\alpha - \nu_{n+1} \right) F^{ii} h_{ii} \geq -C.$$

Substituting all these back into (4.2), we get

$$\begin{aligned} (4.6) \quad 0 &\geq 2 \sum_{i \neq 1} \frac{F^{ii} - F^{11}}{\kappa_1 - \kappa_i} \frac{h_{11i}^2}{\kappa_1} - \sum_{i=1}^n F^{ii} \frac{h_{11i}^2}{\kappa_1^2} + \sum_{i=1}^n F^{ii} \kappa_i^2 + \alpha \sum_{i=1}^n F^{ii} \\ &\quad - \beta \sum_{i=1}^n F^{ii} \frac{|\nabla_i \eta|^2}{\eta^2} - \frac{C\beta}{\eta} - C\kappa_1 - C. \end{aligned}$$

Let $\delta > 0$ be a number whose value is to be determined. We proceed by considering two cases.

Case 1: $\kappa_n \leq -\delta \kappa_1$.

In this case, we may use the fact that $|\nabla\eta| \leq C$ and the first order critical condition (4.1) to estimate

$$\begin{aligned} & \sum_{i=1}^n F^{ii} \frac{h_{11i}^2}{\kappa_1^2} + \beta \sum_{i=1}^n F^{ii} \frac{|\nabla_i \eta|^2}{\eta^2} \\ & \leq C\alpha^2 \sum_{i=1}^n F^{ii} + \frac{C\beta(1+\beta)}{\eta^2} \sum_{i=1}^n F^{ii}. \end{aligned}$$

On the other hand, since $\kappa_n \leq -\delta\kappa_1$, we have

$$\sum_{i=1}^n F^{ii} \kappa_i^2 \geq F^{nn} \kappa_n^2 \geq C(n, k) \delta^2 \kappa_1^2 \sum_{i=1}^n F^{ii}$$

by lemma 2.10. Hence, the inequality (4.6) becomes

$$(4.7) \quad 0 \geq \left(C\delta^2 \kappa_1^2 - \frac{C\beta(1+\beta)}{\eta^2} - C\alpha^2 \right) \sum_{i=1}^n F^{ii} - \frac{C\beta}{\eta} - C\kappa_1 - C$$

and a bound for $\eta^b \kappa_1$ at X_0 follows from this.

Case 2: $\kappa_n \geq -\delta\kappa_1$.

This time, in order to handle the negative third order terms, we partition the indices $\{1, \dots, n\}$ into

$$I = \{j : F^{jj} \leq \theta^{-1} F^{11}\}, \quad J = \{j : F^{jj} > \theta^{-1} F^{11}\}$$

for some $\theta > 0$ to be determined. Again, we use the first critical condition (4.1) and the fact that $|\nabla\eta| \leq C$ to estimate

$$\begin{aligned} & \sum_{i=1}^n F^{ii} \frac{h_{11i}^2}{\kappa_1^2} + \beta \sum_{i=1}^n F^{ii} \frac{|\nabla_i \eta|^2}{\eta^2} \\ & = \left(\sum_I + \sum_J \right) F^{ii} \frac{h_{11i}^2}{\kappa_1^2} + \beta \left(\sum_I + \sum_J \right) F^{ii} \frac{|\nabla_i \eta|^2}{\eta^2} \\ & \leq \left(\sum_I C\beta^3 F^{ii} \frac{|\nabla_i \eta|^2}{\eta^2} + \frac{C\alpha^2}{\beta} F^{ii} \right) + \sum_J F^{ii} \frac{h_{11i}^2}{\kappa_1^2} \\ (4.8) \quad & + \beta \sum_I F^{ii} \frac{|\nabla_i \eta|^2}{\eta^2} + \left(\sum_J \frac{C}{\beta} F^{ii} \frac{h_{11i}^2}{\kappa_1^2} + \frac{C\alpha^2}{\beta} F^{ii} \right) \\ & \leq C(\beta + \beta^3) \sum_I F^{ii} \frac{|\nabla_i \eta|^2}{\eta^2} + \left(1 + \frac{C}{\beta} \right) \sum_J F^{ii} \frac{h_{11i}^2}{\kappa_1^2} + \frac{C\alpha^2}{\beta} \sum_{i=1}^n F^{ii} \\ & \leq C(\beta + \beta^3) \frac{F^{11}}{\eta^2} + \left(1 + \frac{C}{\beta} \right) \sum_J F^{ii} \frac{h_{11i}^2}{\kappa_1^2} + \frac{C\alpha^2}{\beta} \sum_{i=1}^n F^{ii}, \end{aligned}$$

where we have used the Cauchy's inequality

$$\frac{h_{11i}^2}{\kappa_1^2} \leq C\epsilon\alpha^2 + \frac{\beta^2}{C\epsilon} \frac{|\nabla_i \eta|^2}{\eta^2}$$

with $\epsilon = 1/\beta$.

With (4.8) at hand, the inequality (4.6) becomes

$$(4.9) \quad \begin{aligned} 0 &\geq 2 \sum_{i \neq 1} \frac{F^{ii} - F^{11}}{\kappa_1 - \kappa_i} \frac{h_{11i}^2}{\kappa_1} - \left(1 + \frac{C}{\beta}\right) \sum_J F^{ii} \frac{h_{11i}^2}{\kappa_1^2} + \left(\alpha - \frac{C\alpha^2}{\beta}\right) \sum_{i=1}^n F^{ii} \\ &\quad + F^{11} \kappa_1^2 - \frac{C(\beta + \beta^3)}{\eta^2} F^{11} - \frac{C\beta}{\eta} - C\kappa_1 - C. \end{aligned}$$

Note that

$$\begin{aligned} &2 \sum_{i \neq 1} \frac{F^{ii} - F^{11}}{\kappa_1 - \kappa_i} \frac{h_{11i}^2}{\kappa_1} - \left(1 + \frac{C}{\beta}\right) \sum_J F^{ii} \frac{h_{11i}^2}{\kappa_1^2} \\ &\geq 2 \sum_J \frac{F^{ii} - F^{11}}{\kappa_1 - \kappa_i} \frac{h_{11i}^2}{\kappa_1} - \left(1 + \frac{C}{\beta}\right) \sum_J F^{ii} \frac{h_{11i}^2}{\kappa_1^2} \\ &\geq \sum_J \left[2 \cdot \frac{1 - \theta}{1 + \delta} - \left(1 + \frac{C}{\beta}\right)\right] F^{ii} \frac{h_{11i}^2}{\kappa_1^2} \end{aligned}$$

can be made non-negative if we choose appropriate values for the parameters δ, θ, β . Indeed, let

$$\frac{C}{\beta} = \varepsilon, \quad \frac{1 - \theta}{1 + \delta} \geq 1 - \varepsilon,$$

and for $\beta \geq 1$ large enough so that $0 < \varepsilon \leq 1/3$, we have

$$2 \cdot \frac{1 - \theta}{1 + \delta} - \left(1 + \frac{C}{\beta}\right) \geq 1 - 3\varepsilon \geq 0.$$

Consequently, we are left with

$$0 \geq \left(\alpha - \frac{C\alpha^2}{\beta}\right) \sum_{i=1}^n F^{ii} + F^{11} \kappa_1^2 - \frac{C(\beta + \beta^3)}{\eta^2} F^{11} - \frac{C\beta}{\eta} - C\kappa_1 - C.$$

Finally, there is only one troublesome term i.e. $-C\kappa_1$ to be handled, which arose from the interchanging formula (4.3) and the twice differentiation (4.4) of ψ . Indeed, by fixing $\beta \geq C\alpha^2$ is large enough, we have

$$\left(\alpha - \frac{C\alpha^2}{\beta}\right) \sum_{i=1}^n F^{ii} \geq C\alpha \sum_{i=1}^n F^{ii} \geq C\alpha\kappa_1$$

and a bound for $\eta^\beta \kappa_1$ at X_0 is achieved by choosing a large enough α .

The proof is now complete. \square

Several remarks are in order.

Remark 4.4. In [46], Urbas used the additional assumption that the boundary data φ is uniformly convex to obtain that

$$(4.10) \quad \sum_{i,j} F^{ij} \nabla_i \nabla_j \eta \geq c_0 \sum_{i=1}^n F^{ii} - C$$

instead of (4.5). Then, by further invoking the property that

$$(4.11) \quad \sum_{i=1}^n F^{ii} \geq C\kappa_1,$$

Urbas was able to handle the term $-C\kappa_1$ with the term

$$\beta \frac{F^{ij} \nabla_i \nabla_j \eta}{\eta} \geq \frac{C\beta}{\eta} \kappa_1$$

by fixing $\beta \geq 1$ large.

Remark 4.5. Both our proof and the proof of Urbas cannot be generalized to $k \geq 3$ because the property (4.11) holds only when $k = 2$; see the demonstration in [46, Remark at the end of page 315].

By an almost identical proof, we obtain the following and hence theorem 1.2.

Theorem 4.6. *Let $n \geq 3$ and let $\Omega \subseteq \mathbb{R}^n$ be a convex admissible domain with a smooth boundary. Assume $\psi(x, u, Du) \in C^2(\overline{\Omega} \times \mathbb{R} \times \mathbb{R}^n)$ is positive and $\varphi \in C^2(\overline{\Omega})$ is spacelike. Suppose $u \in C^4(\Omega) \cap C^2(\overline{\Omega})$ is an admissible solution to (1.2). Then the maximum principal curvature*

$$\kappa_{\max}(x) := \max_{1 \leq i \leq n} \kappa_i(x)$$

of its graph satisfies

$$\max_{\Omega} \kappa_{\max}(x) \leq C \left(1 + \max_{\partial\Omega} \kappa_{\max}(x) \right)$$

for some $C > 0$ depending on $n, \|u\|_{C^1(\overline{\Omega})}, \|\psi\|_{C^2(\mathcal{D})}$ and $\|\varphi\|_{C^1(\mathcal{D})}$, where

$$\mathcal{D} := \overline{\Omega} \times [\inf_{\Omega} u, \sup_{\Omega} u] \times \mathbb{R}^n.$$

5. THE k -CURVATURE EQUATION

Although the curvature estimates are now known for $k = 1, 2, n$ due to the work of Bartnik-Simon [1], Bayard [2], Delanoe [16], Urbas [46], and most recently for $k = n - 1, n - 2$ due to the work of Ren-Wang [36, 37], the question of whether the bound remains valid for $3 \leq k \leq n - 3$ is still open to this date. For this direction of research, some partial progress has been obtained: In [28], Huang proved the curvature bound for the k -curvature equation $\sigma_k[u] = \psi(X, \tilde{w})$ when the right-hand side is convex in \tilde{w} and

$$(5.1) \quad \frac{\partial \psi^{1/k}(X, \tilde{w})}{\partial \tilde{w}} \cdot \tilde{w} \geq \psi^{1/k}(X, \tilde{w}) \quad \text{for fixed } X = (x, u(x)),$$

where

$$\tilde{w} := \frac{1}{w} = \frac{1}{\sqrt{1 - |Du|^2}}.$$

The approach of Huang may be appealing for some particular functions ψ , but the assumption excludes the very desirable case when e.g. $\psi = \psi(X)$ does not contain gradient terms at all. In [49], Z. Wang obtained the bound for a general right-hand side $\psi = \psi(x, u, Du)$ and all k if the solution graph is additionally $(k + 1)$ -convex.

In this section, we establish the curvature bound for admissible solutions to the k -curvature equation which additionally have semi-convex graphs. Since semi-convexity is weaker than $(k + 1)$ -convexity [48, Lemma 2.13], our result generalizes the estimate due to Z. Wang. Our proof adapts the arguments from a work of Lu [35], where the core ideas are inspired by the novel paper of Guan-Ren-Wang [26].

Theorem 5.1. *Let $\varphi \in C^2(\Omega) \cap C(\overline{\Omega})$ be spacelike and affine. Suppose $\psi(x, z, p) \in C^\infty(\overline{\Omega} \times \mathbb{R} \times \mathbb{R}^n)$ is positive. If $u \in C^4(\Omega) \cap C^2(\overline{\Omega})$ is an admissible solution to (1.2) and its graph Σ is semi-convex i.e. principal curvatures $\kappa = (\kappa_1, \dots, \kappa_n)$ of Σ satisfy*

$$\kappa_i(x) \geq -K \quad \text{for all } x \in \overline{\Omega} \text{ and all } 1 \leq i \leq n,$$

then the maximum principal curvature

$$\kappa_{\max}(x) := \max_{1 \leq i \leq n} \kappa_i(x)$$

of Σ satisfies

$$\sup_{\Omega'} \kappa_{\max} \leq C(\Omega')$$

for any $\Omega' \subset\subset \Omega$, where $C(\Omega') > 0$ depends on $n, k, K, \theta, \|\psi\|_{C^2(\mathcal{D})}$, $\inf_{\mathcal{D}} \psi$, and $\|\varphi\|_{C^1(\bar{\Omega})}$; here θ is the constant in (3.3) and

$$\mathcal{D} := \bar{\Omega} \times \left[\inf_{\Omega} u, \sup_{\Omega} u \right] \times \mathbb{R}^n.$$

Proof. Denote by

$$\tilde{w} := \frac{1}{w} = \frac{1}{\sqrt{1 - |Du|^2}}$$

and let

$$\eta = \varphi - u.$$

The setting is exactly the same as in section 4 and we shall dive directly into the calculations. This time we use the following test function

$$W = \beta \log \eta + \log \kappa_{\max} + N\tilde{w} + \frac{\alpha}{2}|X|^2,$$

where $\beta, N, \alpha > 0$ are possibly large constants to be chosen later. The key difference here is that we are going to utilize almost all available positive terms which were plausibly omitted in section 4.

Suppose W attains its maximum at some interior point $X_0 = (x_0, u(x_0))$. We may choose a local orthonormal frame $\{e_1, \dots, e_n\}$ around X_0 such that the second fundamental form $h_{ij} = \kappa_i \delta_{ij}$ is diagonalized and

$$\kappa_{\max} = \kappa_1 \geq \kappa_2 \geq \dots \geq \kappa_n.$$

In case κ_1 has multiplicity $m > 1$ i.e.

$$\kappa_1 = \kappa_2 = \dots = \kappa_m > \kappa_{m+1} \geq \dots \geq \kappa_n,$$

we may apply a smooth approximation lemma due to Brendle-Choi-Daskalopoulos [9] to obtain that

$$(5.2) \quad \begin{aligned} \delta_{kl} \cdot (\kappa_1)_i &= h_{kli}, \quad 1 \leq k, l \leq m, \\ (\kappa_1)_{ii} &\geq h_{11ii} + 2 \sum_{p>m} \frac{h_{1pi}^2}{\kappa_1 - \kappa_p}, \end{aligned}$$

in the viscosity sense. Then at x_0 , we have

$$(5.3) \quad \begin{aligned} 0 &= \beta \frac{\nabla_i \eta}{\eta} + \frac{(\kappa_1)_i}{\kappa_1} + N \nabla_i \tilde{w} + \alpha \langle X, e_i \rangle = \beta \frac{\nabla_i \eta}{\eta} + \frac{h_{11i}}{\kappa_1} + N \nabla_i \tilde{w} + \alpha \langle X, \hat{e}_i \rangle, \\ 0 &\geq \beta \frac{\nabla_{ii} \eta}{\eta} - \beta \left(\frac{\nabla_i \eta}{\eta} \right)^2 + \frac{(\kappa_1)_{ii}}{\kappa_1} - \frac{(\kappa_1)_i^2}{\kappa_1^2} + N \nabla_{ii} \tilde{w} + \alpha (1 + h_{ii} \langle X, \nu \rangle) \\ (5.4) \quad &\geq \beta \frac{\nabla_{ii} \eta}{\eta} - \beta \left(\frac{\nabla_i \eta}{\eta} \right)^2 + \frac{h_{11ii}}{\kappa_1} + 2 \sum_{p>m} \frac{h_{1pi}^2}{\kappa_1(\kappa_1 - \kappa_p)} - \frac{h_{11i}^2}{\kappa_1^2} + N \nabla_{ii} \tilde{w} + \alpha (1 + h_{ii} \langle X, \nu \rangle). \end{aligned}$$

Contracting (5.4) with $F = \sigma_k$, we have

$$(5.5) \quad \begin{aligned} 0 &\geq \sum_{i=1}^n \frac{F^{ii} h_{11ii}}{\kappa_1} + 2 \sum_{i=1}^n \sum_{p>m} \frac{F^{ii} h_{1pi}^2}{\kappa_1(\kappa_1 - \kappa_i)} - \sum_{i=1}^n \frac{F^{ii} h_{11i}^2}{\kappa_1^2} \\ &\quad + \frac{\beta}{\eta} \sum_{i=1}^n F^{ii} \nabla_{ii} \eta - \beta \sum_{i=1}^n F^{ii} \left(\frac{\nabla_i \eta}{\eta} \right)^2 + N \sum_{i=1}^n F^{ii} w_{ii} + \alpha \sum_{i=1}^n F^{ii}. \end{aligned}$$

Now, in the Minkowski space, the interchanging formula reads

$$(5.6) \quad h_{11ii} = h_{ii11} + h_{11} h_{ii}^2 - h_{11}^2 h_{ii},$$

and so

$$F^{ii}h_{11ii} = F^{ii}h_{ii11} + \kappa_1 F^{ii}\kappa_i^2 - \kappa_1^2 F^{ii}\kappa_i.$$

Differentiating the equation $F = \psi$ twice yields

$$(5.7) \quad \sum_{i=1}^n F^{ii}h_{ii1} = \nabla_1 \psi$$

and

$$(5.8) \quad \sum_{p,q,r,s} F^{pq,rs}h_{pq1}h_{rs1} + \sum_{i=1}^n F^{ii}h_{ii11} = \nabla_1 \nabla_1 \psi \geq -C(1 + \kappa_1^2)$$

The inequality (5.5) becomes

$$(5.9) \quad \begin{aligned} 0 \geq & - \sum_{p,q,r,s} \frac{F^{pq,rs}h_{pq1}h_{rs1}}{\kappa_1} + \sum_{i=1}^n F^{ii}\kappa_i^2 - C\kappa_1 - C \\ & + 2 \sum_{i=1}^n \sum_{p>m} \frac{F^{ii}h_{1pi}^2}{\kappa_1(\kappa_1 - \kappa_i)} - \sum_{i=1}^n \frac{F^{ii}h_{11i}^2}{\kappa_1^2} \\ & + \frac{\beta}{\eta} \sum_{i=1}^n F^{ii}\nabla_{ii}\eta - \beta \sum_{i=1}^n F^{ii} \left(\frac{\nabla_i \eta}{\eta} \right)^2 + N \sum_{i=1}^n F^{ii}w_{ii} + \alpha \sum_{i=1}^n F^{ii}. \end{aligned}$$

We continue to expand the terms in (5.9). First, we have

$$\begin{aligned} - \sum_{p,q,r,s} \frac{F^{pq,rs}h_{pq1}h_{rsq}}{\kappa_1} &= - \sum_{p,q} \frac{F^{pp,qq}h_{pp1}h_{qq1}}{\kappa_1} + \sum_{p,q} \frac{F^{pp,qq}h_{pq1}^2}{\kappa_1} \\ &\geq - \sum_{p,q} \frac{F^{pp,qq}h_{pp1}h_{qq1}}{\kappa_1} + 2 \sum_{i>m} \frac{F^{11,ii}h_{11i}^2}{\kappa_1} \\ &= - \sum_{p,q} \frac{F^{pp,qq}h_{pp1}h_{qq1}}{\kappa_1} + 2 \sum_{i>m} \frac{F^{ii} - F^{11}}{\kappa_1(\kappa_1 - \kappa_i)} h_{11i}^2. \end{aligned}$$

On the other hand, we have

$$\begin{aligned} 2 \sum_{i=1}^n \sum_{p>m} \frac{F^{ii}h_{1pi}^2}{\kappa_1(\kappa_1 - \kappa_p)} &\geq 2 \sum_{p>m} \frac{F^{pp}h_{1pp}^2}{\kappa_1(\kappa_1 - \kappa_p)} + 2 \sum_{p>m} \frac{F^{11}h_{1p1}^2}{\kappa_1(\kappa_1 - \kappa_p)} \\ &= 2 \sum_{i>m} \frac{F^{ii}h_{ii1}^2}{\kappa_1(\kappa_1 - \kappa_i)} + 2 \sum_{i>m} \frac{F^{11}h_{11i}^2}{\kappa_1(\kappa_1 - \kappa_i)}. \end{aligned}$$

Substituting these back into (5.9), we have

$$(5.10) \quad \begin{aligned} 0 \geq & - \sum_{p,q} \frac{F^{pp,qq}h_{pp1}h_{qq1}}{\kappa_1} - \frac{F^{11}h_{111}^2}{\kappa_1^2} + 2 \sum_{i>m} \frac{F^{ii}h_{ii1}^2}{\kappa_1(\kappa_1 - \kappa_i)} - C\kappa_1 - C \\ & + 2 \sum_{i>m} \frac{F^{ii} - F^{11}}{\kappa_1(\kappa_1 - \kappa_i)} h_{11i}^2 + 2 \sum_{i>m} \frac{F^{11}h_{11i}^2}{\kappa_1(\kappa_1 - \kappa_i)} - \sum_{i \neq 1} \frac{F^{ii}h_{11i}^2}{\kappa_1^2} \\ & + \frac{\beta}{\eta} \sum_{i=1}^n F^{ii}\nabla_{ii}\eta - \beta \sum_{i=1}^n F^{ii} \left(\frac{\nabla_i \eta}{\eta} \right)^2 + \sum_{i=1}^n F^{ii}\kappa_i^2 + N \sum_{i=1}^n F^{ii}\tilde{w}_{ii} + \alpha \sum_{i=1}^n F^{ii}. \end{aligned}$$

We now look at the terms involving η . By the same calculation as in (4.5), we have

$$\sum_{i=1}^n F^{ii}\nabla_{ii}\eta \geq -C.$$

Also, invoking $|\nabla\eta| \leq C$ and the first critical condition (5.3), we have

$$\begin{aligned} \beta \sum_{i=1}^n F^{ii} \left(\frac{\nabla_i \eta}{\eta} \right)^2 &= \beta F^{11} \left(\frac{\nabla_1 \eta}{\eta} \right)^2 + \beta \sum_{i \neq 1} F^{ii} \left(\frac{\nabla_i \eta}{\eta} \right)^2 \\ &\leq \frac{C\beta}{\eta^2} F^{11} + \sum_{i \neq 1} \left(\frac{C}{\beta} F^{ii} \frac{h_{11i}^2}{\kappa_1^2} + \frac{CN^2}{\beta} F^{ii} (\nabla_i \tilde{w})^2 + \frac{C\alpha^2}{\beta} F^{ii} \right). \end{aligned}$$

That is, (5.10) reduces to

$$\begin{aligned} (5.11) \quad 0 &\geq - \sum_{p,q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} - \frac{F^{11} h_{111}^2}{\kappa_1^2} + 2 \sum_{i>m} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \kappa_i)} \\ &\quad + 2 \sum_{i>m} \frac{F^{ii} - F^{11}}{\kappa_1(\kappa_1 - \kappa_i)} h_{11i}^2 + 2 \sum_{i>m} \frac{F^{11} h_{11i}^2}{\kappa_1(\kappa_1 - \kappa_i)} - \left(1 + \frac{C}{\beta}\right) \sum_{i \neq 1} \frac{F^{ii} h_{11i}^2}{\kappa_1^2} \\ &\quad + \left(N \sum_{i=1}^n F^{ii} \tilde{w}_{ii} - \frac{CN^2}{\beta} \sum_{i=1}^n F^{ii} \tilde{w}_i^2 \right) + \left(\alpha - \frac{C\alpha^2}{\beta} \right) \sum_{i=1}^n F^{ii} \\ &\quad + \left(F^{11} \kappa_1^2 - \frac{C\beta}{\eta^2} F^{11} \right) - \frac{C\beta}{\eta} - C\kappa_1 - C. \end{aligned}$$

By choosing $\beta > 0$ large enough, we may assume the number

$$\varepsilon := \frac{C}{\beta}$$

is sufficiently small. Also, recall the following formulas [47, (3.1) and (3.13)],

$$\begin{aligned} F^{ij} \nabla_i \tilde{w} \nabla_j \tilde{w} &\leq |Du|^2 \tilde{w}^2 F^{ij} h_{ik} h_{jk}, \\ F^{ij} \nabla_i \nabla_j \tilde{w} &= \tilde{w} F^{ij} h_{im} h_{jm} + \langle \nabla \psi, \epsilon_{n+1} \rangle, \end{aligned}$$

where ϵ_{n+1} is the $(n+1)$ -th standard coordinate in \mathbb{R}^{n+1} . We are left with

$$\begin{aligned} (5.12) \quad 0 &\geq - \sum_{p,q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} - \frac{F^{11} h_{111}^2}{\kappa_1^2} + 2 \sum_{i>m} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \kappa_i)} \\ &\quad + 2 \sum_{i>m} \frac{F^{ii} - F^{11}}{\kappa_1(\kappa_1 - \kappa_i)} h_{11i}^2 + 2 \sum_{i>m} \frac{F^{11} h_{11i}^2}{\kappa_1(\kappa_1 - \kappa_i)} - (1 + \varepsilon) \sum_{i \neq 1} \frac{F^{ii} h_{11i}^2}{\kappa_1^2} \\ &\quad + CN \sum_{i=1}^n F^{ii} \kappa_i^2 + C\alpha \sum_{i=1}^n F^{ii} - \frac{C\beta}{\eta} - C\kappa_1 - CN - C, \end{aligned}$$

where we have also assumed $\eta^2 \kappa_1^2 \geq C\beta$ is sufficiently large. It remains only to handle the first two lines. Indeed, the second line could be easily handled as

$$h_{11i} = h_{i11} = \delta_{1i} \cdot (\kappa_1)_1 = 0, \quad 1 < i \leq m \quad \text{by (5.2)}$$

and so

$$\begin{aligned}
& 2 \sum_{i>m} \frac{F^{ii} - F^{11}}{\kappa_1(\kappa_1 - \kappa_i)} h_{11i}^2 + 2 \sum_{i>m} \frac{F^{11} h_{11i}^2}{\kappa_1(\kappa_1 - \kappa_i)} - (1 + \varepsilon) \sum_{i \neq 1} \frac{F^{ii} h_{11i}^2}{\kappa_1^2} \\
(5.13) \quad & = 2 \sum_{i>m} \frac{F^{ii} h_{11i}^2}{\kappa_1(\kappa_1 - \kappa_i)} - (1 + \varepsilon) \sum_{i>m} \frac{F^{ii} h_{11i}^2}{\kappa_1^2} \\
& = \sum_{i>m} \frac{F^{ii} h_{11i}^2}{\kappa_1^2(\kappa_1 - \kappa_i)} [2\kappa_1 - (1 + \varepsilon)(\kappa_1 - \kappa_i)] \\
& = \sum_{i>m} \frac{F^{ii} h_{11i}^2}{\kappa_1^2(\kappa_1 - \kappa_i)} [(1 - \varepsilon)\kappa_1 + (1 + \varepsilon)\kappa_i]
\end{aligned}$$

is non-negative by semi-convexity $\kappa_i \geq -K$. Hence,

$$\begin{aligned}
(5.14) \quad 0 & \geq - \sum_{p,q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} - \frac{F^{11} h_{111}^2}{\kappa_1^2} + 2 \sum_{i>m} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \kappa_i)} \\
& \quad + CN \sum_{i=1}^n F^{ii} \kappa_i^2 + C\alpha \sum_{i=1}^n F^{ii} - \frac{C\beta}{\eta} - C\kappa_1 - CN - C.
\end{aligned}$$

However, the major difficulty is to deal with the first line in (5.12) and for that purpose, we are going to apply some novel ideas from the work of Guan-Ren-Wang [26]. First, we recall a concavity lemma for the operator σ_k and a proof of which can be found in [35].

Lemma 5.2. *Let $\kappa = (\kappa_1, \dots, \kappa_n) \in \Gamma_k$ be ordered as $\kappa_1 \geq \dots \geq \kappa_n$ and $1 \leq l < k$. For any $\varepsilon, \delta, \delta_0 \in (0, 1)$, there exists some $\delta' > 0$ such that if $\kappa_l \geq \delta\kappa_1$ and $\kappa_{l+1} \leq \delta'\kappa_1$, then we have*

$$(5.15) \quad - \sum_{p,q} \frac{\sigma_k^{pp,qq} \xi_p \xi_q}{\sigma_k} + \frac{1}{\sigma_k^2} \left(\sum_{i=1}^n \sigma_k^{ii} \xi_i \right)^2 \geq (1 - \varepsilon) \frac{\xi_1^2}{\kappa_1^2} - \delta_0 \sum_{i>l} \frac{\sigma_k^{ii} \xi_i}{\kappa_1 \sigma_k}$$

for an arbitrary vector $\xi = (\xi_1, \dots, \xi_n) \in \mathbb{R}^n$.

Then, we proceed by an iteration argument that is very delicate; one could see applications of this new technique in some other settings [15, 48, 55]. We may still let $\varepsilon > 0$ denote a very small number without causing any confusion, and let $\delta_0 = 1/2$. Pick an arbitrary $\delta_1 \in (0, 1)$, say $\delta_1 = 1/3$, we would then trivially have $\kappa_1 \geq \delta_1 \kappa_1$. Now, by the lemma, there exists some $\delta_2 > 0$ such that if $\kappa_2 \leq \delta_2 \kappa_1$, then the inequality (5.15) holds. If we had $\kappa_2 \geq \delta_2 \kappa_1$, then we may continue to pick some δ_3 and see if we would have $\kappa_3 \leq \delta_3 \kappa_1$. The key argument is that this process either halts at some $1 \leq l < k$, or it goes on and we have $\kappa_k > \delta_k \kappa_1$. We now analyze the two cases.

Case 1: For each $1 \leq i \leq k$ there exists some $\delta_i > 0$ such that $\kappa_i \geq \delta_i \kappa_1$.

In this case, we immediately have

$$\begin{aligned}
\sigma_k & = \sum_{1 \leq i_1 < \dots < i_k \leq n} \kappa_{i_1} \cdots \kappa_{i_k} \\
& \geq \kappa_1 \cdots \kappa_k - C(n, k) \kappa_1 \cdots \kappa_{k-1} \cdot K \\
& \geq \kappa_1 \cdots \kappa_{k-1} (\kappa_k - CK) \\
& \geq C\delta_2 \cdots \delta_k \kappa_1^k
\end{aligned}$$

and the desired estimate follows.

Case 2: The process halts at some $1 \leq l < k$ and we have (5.15).

In this case, taking $\xi = (h_{ii1})$ and applying (5.7) and (5.2), the inequality (5.15) yields

$$\begin{aligned}
& - \sum_{p,q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} - \frac{F^{11} h_{111}^2}{\kappa_1^2} + 2 \sum_{i>m} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \kappa_i)} \\
\geq & - \frac{(\nabla_1 \psi)^2}{\kappa_1 \psi} + (1 - \varepsilon) \sigma_k \frac{h_{111}^2}{\kappa_1^3} - \frac{F^{11} h_{111}^2}{\kappa_1^2} + 2 \sum_{i>m} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \kappa_i)} - \frac{1}{2} \sum_{i>l} \frac{F^{ii} h_{ii1}^2}{\kappa_1^2} \\
\geq & - C \kappa_1 + (1 - \varepsilon) [F^{11} \kappa_1 + \sigma_k(\kappa|1)] \frac{h_{111}^2}{\kappa_1^3} - \frac{F^{11} h_{111}^2}{\kappa_1^2} + \sum_{i>m} \frac{F^{ii} h_{ii1}^2}{\kappa_1^2(\kappa_1 - \kappa_i)} (3\kappa_1 + \kappa_i) \\
\geq & - C \kappa_1 - \varepsilon \frac{F^{11} h_{111}^2}{\kappa_1^2} + (1 - \varepsilon) \sigma_k(\kappa|1) \frac{h_{111}^2}{\kappa_1^3}.
\end{aligned}$$

Therefore,

$$\begin{aligned}
(5.16) \quad 0 & \geq \frac{CN}{2} F^{11} \kappa_1^2 - \varepsilon \frac{F^{11} h_{111}^2}{\kappa_1^2} \\
& + C \alpha \sum_{i=1}^n F^{ii} + (1 - \varepsilon) \sigma_k(\kappa|1) \frac{h_{111}^2}{\kappa_1^3} \\
& + \frac{CN}{2} F^{11} \kappa_1^2 - \frac{C\beta}{\eta} - C \kappa_1 - CN - C.
\end{aligned}$$

Still, using the first order critical condition (5.3) and the fact that $\kappa_i \geq -K$, we can estimate

$$\begin{aligned}
(1 - \varepsilon) \sigma_k(\kappa|1) \frac{h_{111}^2}{\kappa_1^3} & \geq -C \kappa_2 \cdots \kappa_k \cdot K \cdot \frac{1}{\kappa_1} \cdot \left(\frac{C\beta^2}{\eta^2} + CN^2 \kappa_1^2 + C\alpha^2 \right) \\
& \geq -CKN^2 \kappa_1 \cdots \kappa_k
\end{aligned}$$

and

$$\begin{aligned}
-\varepsilon \frac{F^{11} h_{111}^2}{\kappa_1^2} & \geq -\varepsilon F^{11} \left(\frac{C\beta^2}{\eta^2} + CN^2 \kappa_1^2 + C\alpha^2 \right) \\
& \geq -\varepsilon CN^2 F^{11} \kappa_1^2
\end{aligned}$$

by assuming $\eta^2 \kappa_1^2$ is large. Thus,

$$\frac{CN}{2} F^{11} \kappa_1^2 - \varepsilon \frac{F^{11} h_{111}^2}{\kappa_1^2} \geq (CN - \varepsilon CN^2) F^{11} \kappa_1^2 \geq 0$$

by taking ε small enough e.g. of order C/N^2 . On the other hand,

$$\begin{aligned}
C \alpha \sum_{i=1}^n F^{ii} + (1 - \varepsilon) \sigma_k(\kappa|1) \frac{h_{111}^2}{\kappa_1^3} & \geq C \alpha \sigma_{k-1} - CKN^2 \kappa_1 \cdots \kappa_k \\
& \geq (C \alpha - CKN^2 \kappa_k) \kappa_1 \cdots \kappa_{k-1}.
\end{aligned}$$

If $C \alpha - CKN^2 \kappa_k \geq 0$, then we are done. Otherwise, taking α to be at least of order N^3 , we would have $\kappa_k \geq \frac{C\alpha}{KN^2} = CN$ and so

$$\sigma_k \geq \kappa_1 \cdots \kappa_k - C \kappa_1 \cdots \kappa_{k-1} \cdot K \geq \kappa_1 \cdots \kappa_{k-1} (CN - C) \geq CN \kappa_1$$

by choosing a large N .

Finally, we can remove the first two lines from (5.16) and since

$$F^{11} \kappa_1^2 = \sigma_{k-1}(\kappa|1) \kappa_1^2 \geq \frac{k}{n} \sigma_k \kappa_1,$$

the estimate follows by choosing N large.

□

By an almost identical proof, theorem 1.3 follows. Moreover, we could also provide an alternative proof for Ren-Wang's curvature estimates when $k = n - 1$ [36, Theorem 5] and $k = n - 2$ [37, Theorem 7].

Theorem 5.3. *Let $k = n - 1, n \geq 3$, or $k = n - 2, n \geq 5$, and let $\Omega \subseteq \mathbb{R}^n$ be a convex admissible domain with a smooth boundary. Assume $\psi(x, u, Du) \in C^2(\bar{\Omega} \times \mathbb{R} \times \mathbb{R}^n)$ is positive and $\varphi \in C^2(\bar{\Omega})$ is spacelike. Suppose $u \in C^4(\Omega) \cap C^2(\bar{\Omega})$ is an admissible solution to (1.1). Then the maximum principal curvature*

$$\kappa_{\max}(x) := \max_{1 \leq i \leq n} \kappa_i(x)$$

of its graph satisfies

$$\max_{\Omega} \kappa_{\max}(x) \leq C \left(1 + \max_{\partial\Omega} \kappa_{\max}(x) \right)$$

for some $C > 0$ depending on $n, \|u\|_{C^1(\bar{\Omega})}, \|\psi\|_{C^2(\mathcal{D})}$ and $\|\varphi\|_{C^1(\mathcal{D})}$, where

$$\mathcal{D} := \bar{\Omega} \times [\inf_{\Omega} u, \sup_{\Omega} u] \times \mathbb{R}^n.$$

Proof. In this case, the first line in (5.12) can be easily handled because we have the very powerful concavity inequalities [48, Lemma 2.11] due to Ren-Wang [36, 37],

$$-\sum_{p,q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} - \frac{F^{11} h_{111}^2}{\kappa_1^2} + 2 \sum_{i>m} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \kappa_i)} \geq -C \frac{(\nabla_1 \psi)^2}{\kappa_1} \geq -C \kappa_1.$$

Moreover, the second line (5.13) is still non-negative by lemma 2.10:

$$(1 - \varepsilon)\kappa_1 + (1 + \varepsilon)\kappa_i \geq \left(\frac{2k - n}{k} - \frac{n}{k}\varepsilon \right) \kappa_1 \geq 0$$

if $0 < \varepsilon < (2k - n)/n$ and $2k > n$. Since all the other parts of the calculations remain exactly the same, the proof is complete. □

Remark 5.4. Instead of $\log \kappa_{\max}$, Ren-Wang used $\log \log \sum_{j=1}^n e^{\kappa_j}$ in their test function which may cause the calculations to be a lot more involved.

ACKNOWLEDGEMENT

We would like to thank Professor Zhizhang Wang for kindly sharing a preprint of his report [49] with us, which cannot be easily found online. We are also greatly indebted to our Ph.D. advisor, Professor Man-Chun Lee, for much support in various aspects academically.

DATA AVAILABILITY

No data was used for the research described in the article.

REFERENCES

- [1] Robert Bartnik and Leon Simon, *Spacelike hypersurfaces with prescribed boundary values and mean curvature*, Comm. Math. Phys. **87** (1982/83), no. 1, 131–152. MR680653
- [2] Pierre Bayard, *Dirichlet problem for space-like hypersurfaces with prescribed scalar curvature in $\mathbb{R}^{n,1}$* , Calc. Var. Partial Differential Equations **18** (2003), no. 1, 1–30. MR2001880
- [3] ———, *Entire spacelike hypersurfaces of prescribed scalar curvature in Minkowski space*, Calc. Var. Partial Differential Equations **26** (2006), no. 2, 245–264. MR2222246
- [4] ———, *Entire scalar curvature flow and hypersurfaces of constant scalar curvature in Minkowski space*, Methods Appl. Anal. **16** (2009), no. 1, 87–118. MR2556831
- [5] ———, *Entire downward solitons to the scalar curvature flow in Minkowski space*, Ann. Inst. H. Poincaré C Anal. Non Linéaire **40** (2023), no. 6, 1353–1383. MR4656418

- [6] Pierre Bayard and Philippe Delanoë, *Entire spacelike radial graphs in the Minkowski space, asymptotic to the light-cone, with prescribed scalar curvature*, Ann. Inst. H. Poincaré C Anal. Non Linéaire **26** (2009), no. 3, 903–915. MR2526408
- [7] Pierre Bayard and Oliver C. Schnürer, *Entire spacelike hypersurfaces of constant Gauß curvature in Minkowski space*, J. Reine Angew. Math. **627** (2009), 1–29. MR2494911
- [8] Pierre Bayard and Andrea Seppi, *Entire hypersurfaces of constant scalar curvature in minkowski space*, preprint on arXiv, <https://arxiv.org/abs/2408.10042>.
- [9] Simon Brendle, Kyeongsu Choi, and Panagiota Daskalopoulos, *Asymptotic behavior of flows by powers of the Gaussian curvature*, Acta Math. **219** (2017), no. 1, 1–16. MR3765656
- [10] L. Caffarelli, L. Nirenberg, and J. Spruck, *The Dirichlet problem for nonlinear second-order elliptic equations. III. Functions of the eigenvalues of the Hessian*, Acta Math. **155** (1985), no. 3-4, 261–301. MR806416
- [11] ———, *Nonlinear second order elliptic equations. IV. Starshaped compact Weingarten hypersurfaces*, Current topics in partial differential equations, 1986, pp. 1–26. MR1112140
- [12] Luis Caffarelli, Louis Nirenberg, and Joel Spruck, *Nonlinear second-order elliptic equations. V. The Dirichlet problem for Weingarten hypersurfaces*, Comm. Pure Appl. Math. **41** (1988), no. 1, 47–70. MR917124
- [13] Eugenio Calabi, *Examples of Bernstein problems for some nonlinear equations*, Global Analysis (Proc. Sympos. Pure Math., Vols. XIV, XV, XVI, Berkeley, Calif., 1968), 1970, pp. 223–230. MR264210
- [14] Shiu Yuen Cheng and Shing Tung Yau, *Maximal space-like hypersurfaces in the Lorentz-Minkowski spaces*, Ann. of Math. (2) **104** (1976), no. 3, 407–419. MR431061
- [15] Jianchun Chu, *A simple proof of curvature estimate for convex solution of k -Hessian equation*, Proc. Amer. Math. Soc. **149** (2021), no. 8, 3541–3552. MR4273155
- [16] F. Delanoë, *The Dirichlet problem for an equation of given Lorentz-Gaussian curvature*, Ukrain. Mat. Zh. **42** (1990), no. 12, 1704–1710. MR1098472
- [17] Lawrence C. Evans, *Classical solutions of fully nonlinear, convex, second-order elliptic equations*, Comm. Pure Appl. Math. **35** (1982), no. 3, 333–363. MR649348
- [18] Claus Gerhardt, *H -surfaces in Lorentzian manifolds*, Comm. Math. Phys. **89** (1983), no. 4, 523–553. MR713684
- [19] ———, *Hypersurfaces of prescribed curvature in Lorentzian manifolds*, Indiana Univ. Math. J. **49** (2000), no. 3, 1125–1153. MR1803223
- [20] ———, *Hypersurfaces of prescribed mean curvature in Lorentzian manifolds*, Math. Z. **235** (2000), no. 1, 83–97. MR1785072
- [21] ———, *Hypersurfaces of prescribed scalar curvature in Lorentzian manifolds*, J. Reine Angew. Math. **554** (2003), 157–199. MR1952172
- [22] Bo Guan, *The Dirichlet problem for Monge-Ampère equations in non-convex domains and spacelike hypersurfaces of constant Gauss curvature*, Trans. Amer. Math. Soc. **350** (1998), no. 12, 4955–4971. MR1451602
- [23] Bo Guan and Pengfei Guan, *Convex hypersurfaces of prescribed curvatures*, Ann. of Math. (2) **156** (2002), no. 2, 655–673. MR1933079
- [24] Bo Guan and Joel Spruck, *Hypersurfaces of constant curvature in hyperbolic space. II*, J. Eur. Math. Soc. (JEMS) **12** (2010), no. 3, 797–817. MR2639319
- [25] Pengfei Guan, Junfang Li, and Yanyan Li, *Hypersurfaces of prescribed curvature measure*, Duke Math. J. **161** (2012), no. 10, 1927–1942. MR2954620
- [26] Pengfei Guan, Changyu Ren, and Zhizhang Wang, *Global C^2 -estimates for convex solutions of curvature equations*, Comm. Pure Appl. Math. **68** (2015), no. 8, 1287–1325. MR3366747
- [27] Mengru Guo and Heming Jiao, *The Dirichlet problem for a class of curvature equations in Minkowski space*, J. Differential Equations **425** (2025), 129–156. MR4851893
- [28] Yong Huang, *Curvature estimates of hypersurfaces in the Minkowski space*, Chinese Ann. Math. Ser. B **34** (2013), no. 5, 753–764. MR3079808
- [29] N. M. Ivochkina, *Solution of the Dirichlet problem for certain equations of Monge-Ampère type*, Mat. Sb. (N.S.) **128(170)** (1985), no. 3, 403–415, 447. MR815272
- [30] ———, *Solution of the Dirichlet problem for equations of m th order curvature*, Mat. Sb. **180** (1989), no. 7, 867–887, 991. MR1014618
- [31] ———, *The Dirichlet problem for the curvature equation of order m* , Algebra i Analiz **2** (1990), no. 3, 192–217. MR1073214
- [32] N. V. Krylov, *Boundedly inhomogeneous elliptic and parabolic equations in a domain*, Izv. Akad. Nauk SSSR Ser. Mat. **47** (1983), no. 1, 75–108. MR688919
- [33] Ming Li, Changyu Ren, and Zhizhang Wang, *An interior estimate for convex solutions and a rigidity theorem*, J. Funct. Anal. **270** (2016), no. 7, 2691–2714. MR3464054
- [34] André Lichnerowicz, *L’intégration des équations de la gravitation relativiste et le problème des n corps*, J. Math. Pures Appl. (9) **23** (1944), 37–63. MR14298

- [35] Siyuan Lu, *Curvature estimates for semi-convex solutions of Hessian equations in hyperbolic space*, Calc. Var. Partial Differential Equations **62** (2023), no. 9, Paper No. 257, 23. MR4659885
- [36] Changyu Ren and Zhizhang Wang, *On the curvature estimates for Hessian equations*, Amer. J. Math. **141** (2019), no. 5, 1281–1315. MR4011801
- [37] ———, *The global curvature estimate for the $n - 2$ Hessian equation*, Calc. Var. Partial Differential Equations **62** (2023), no. 9, Paper No. 239, 50. MR4646879
- [38] Changyu Ren, Zhizhang Wang, and Ling Xiao, *The convexity of entire spacelike hypersurfaces with constant σ_{n-1} curvature in Minkowski space*, J. Geom. Anal. **34** (2024), no. 6, Paper No. 189, 38. MR4735355
- [39] ———, *The prescribed curvature problem for entire hypersurfaces in Minkowski space*, Anal. PDE **17** (2024), no. 1, 1–40. MR4702314
- [40] Oliver C. Schnürer, *The Dirichlet problem for Weingarten hypersurfaces in Lorentz manifolds*, Math. Z. **242** (2002), no. 1, 159–181. MR1985454
- [41] Richard Schoen and Shing Tung Yau, *On the proof of the positive mass conjecture in general relativity*, Comm. Math. Phys. **65** (1979), no. 1, 45–76. MR526976
- [42] Weimin Sheng, John Urbas, and Xu-Jia Wang, *Interior curvature bounds for a class of curvature equations*, Duke Math. J. **123** (2004), no. 2, 235–264. MR2066938
- [43] Neil S. Trudinger and Xu-Jia Wang, *Hessian measures. I*, 1997, pp. 225–239. Dedicated to Olga Ladyzhenskaya. MR1634570
- [44] ———, *Hessian measures. II*, Ann. of Math. (2) **150** (1999), no. 2, 579–604. MR1726702
- [45] ———, *Hessian measures. III*, J. Funct. Anal. **193** (2002), no. 1, 1–23. MR1923626
- [46] John Urbas, *The Dirichlet problem for the equation of prescribed scalar curvature in Minkowski space*, Calc. Var. Partial Differential Equations **18** (2003), no. 3, 307–316. MR2018670
- [47] ———, *Interior curvature bounds for spacelike hypersurfaces of prescribed k -th mean curvature*, Comm. Anal. Geom. **11** (2003), no. 2, 235–261. MR2014878
- [48] Bin Wang, *Starshaped compact hypersurfaces in warped product manifolds I: prescribed curvature equations*, Calc. Var. Partial Differential Equations **63** (2024), no. 9, Paper No. 241, 35. MR4821890
- [49] Zhizhang Wang, *The global curvature estimates for Hessian equations*, Proceedings of the International Consortium of Chinese Mathematicians 2018, [2020] ©2020, pp. 367–384. MR4251152
- [50] Zhizhang Wang and Ling Xiao, *Entire spacelike hypersurfaces with constant σ_k curvature in Minkowski space*, Math. Ann. **382** (2022), no. 3-4, 1279–1322. MR4403224
- [51] ———, *Entire convex curvature flow in Minkowski space*, Calc. Var. Partial Differential Equations **62** (2023), no. 9, Paper No. 252, 27. MR4659387
- [52] ———, *Entire self-expanders for power of σ_k curvature flow in Minkowski space*, J. Funct. Anal. **284** (2023), no. 8, Paper No. 109866, 27. MR4543575
- [53] ———, *Entire spacelike constant σ_k curvature hypersurfaces with prescribed boundary data at infinity*, preprint on arXiv, <https://arxiv.org/abs/2107.03514>.
- [54] ———, *Entire σ_k curvature flow in Minkowski space*, preprint on arXiv, <https://arxiv.org/abs/2207.04552>.
- [55] Fengrui Yang, *Prescribed curvature measure problem in hyperbolic space*, Comm. Pure Appl. Math. **77** (2024), no. 1, 863–898. MR4666637

DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, NEW TERRITORIES, THE HONG KONG SPECIAL ADMINISTRATIVE REGION, THE PEOPLE’S REPUBLIC OF CHINA.

Email address: bwang@math.cuhk.edu.hk