

A Szegő limit theorem for a class of Toeplitz operators on the Bergman space of the unit ball with singular symbols

Daniel Ivan Ramirez Montaña

We obtain a Szegő limit theorem for a family of Toeplitz operators defined on the weighted Bergman space of the unit ball \mathbb{B}_n . The symbols of these operators are supported on some isotropic or co-isotropic submanifold $\Gamma \subseteq \mathbb{B}_n$ and can be seen, in general, as measures that are singular with respect to the Lebesgue measure on \mathbb{C}^n .

Contents

1	Introduction	2
1.1	Definitions	3
1.2	Statement of the main result	7
2	The Riemannian metric on Γ	8
3	The trace of a multiple composition	11
4	Asymptotic of the trace of a multiple composition	16
5	Section with the main theorem	22
5.1	The Szegő limit theorem for polynomials	22
5.2	The Szegő limit theorem	25
6	Corollaries and follow up results	30
6.1	Distribution of eigenvalues	30
6.2	Schatten norm estimates	32
6.3	An example	35
6.3.1	Operator norm	35
6.3.2	Eigenvalue distribution comparison	36
A	Proof of Lemma 5.9	38
B	The determinant of the Hessian	43
B.1	The determinant as a block matrix	43
B.2	Some symplectic geometry	48
	References	52

1 Introduction

In general, a Szegö limit theorem is a result that describes the asymptotic behavior of the spectral measures of an indexed family of operators as the parameter that indexes the family tends to a limit. This description is often made in terms of operator traces. Originally, G. Szegö delivered expressions for the limit of determinants of truncations of infinite Toeplitz matrices as the truncation order tended to infinity ([13]). These relations were later used to obtain information on the asymptotic distribution of the eigenvalues of the said matrices, expressing the limit of their traces through integrals. In particular, in [4] it is shown that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{j=1}^N f(\lambda_{j,N}) = \frac{1}{2\pi} \int_0^{2\pi} f(a(x)) dx ,$$

where $\lambda_{j,N}$ are the eigenvalues of the N -th order truncation of the Toeplitz matrix whose diagonals are the Fourier coefficients of a real valued $a \in L^\infty(\mathbb{S}^1)$ and f is a continuous function on $[\text{ess inf } a, \text{ess sup } a]$. Later works focused on reproducing this result with weaker conditions for a and applying to different families of functions f (e.g. [14], [16]).

Some of these ideas were generalized in [3]. For a bounded self-adjoint operator A on a Hilbert space H , we want a measure μ_A on \mathbb{R} such that

$$\text{Tr}(f(A)) = \int_{\mathbb{R}} f d\mu_A, \tag{1}$$

for any $f \in C(\mathbb{R})$. If H has finite dimension, this can be done using the Riesz-Markov-Kakutani theorem, since the assignation $f \mapsto \text{Tr}(f(A))$ results in a continuous functional on $C(\mathbb{R})$. In fact, μ_A is the sum of the Dirac measures centered on each eigenvalue of A . In general, this cannot be done when H is not of finite dimension, so an alternative construction is proposed. Taking an increasing sequence of finite-rank orthogonal projections P_n on H such that P_n converges to the identity in H , the correspondent sequence of measures μ_n satisfying (1) for the operator $P_n A|_{P_n(H)}$ is constructed. Therefore, a natural substitute for μ_A would be the weak limit of the μ_n measures. In this situation a Szegö limit theorem translates to finding a measure whose integral reproduces the limit of the traces of continuous images of a sequence of operators. Particular attention has been paid to this problem when A is replaced by a pseudo-differential operator (see [3], [8], [15]).

Recently in [2] results of this type were obtained considering various function spaces. In particular, the weighted Bergman space of the disc \mathbb{D} and the weighted Fock space on \mathbb{C} are treated. In [11] the problem was studied in the context of the Fock space of \mathbb{C}^n and considering symbols supported on some manifold, thus in general being measures that are singular with respect to the Lebesgue measure. In both of these works, the family of operators whose limiting behavior is considered is indexed by the parameter defining the associated weighted space. We will also follow this approach, obtaining an analogue result to the main one of [11] for the case of the weighted Bergman space of the unit ball \mathbb{B}_n .

1.1 Definitions

For $n \in \mathbb{N}$, let $\mathbb{B}_n := \{z \in \mathbb{C}^n : |z| < 1\}$ denote the unit ball in \mathbb{C}^n . For $\alpha > -1$, the Bergman space with standard weights $A_\alpha^2(\mathbb{B}_n)$ is the Hilbert space of all holomorphic functions in $L^2(\mathbb{B}_n, dv_\alpha)$, where

$$dv_\alpha(w) = c_\alpha(1 - |w|^2)^\alpha dv(w) ,$$

with $c_\alpha = \frac{\Gamma(\alpha+1+n)}{n!\Gamma(\alpha+1)}$ a normalizing constant and dv standing for the normalized Lebesgue measure on \mathbb{B}_n . Moreover, $A_\alpha^2(\mathbb{B}_n)$ is a reproducing kernel Hilbert space with the inner product inherited from $L^2(\mathbb{B}_n, dv_\alpha)$ and with kernel

$$K^\alpha(z, w) = \frac{1}{(1 - \langle z, w \rangle)^{n+1+\alpha}} , \quad z, w \in \mathbb{B}_n .$$

When $\alpha = 0$, $dv_0 = dv$ and $A^2(\mathbb{B}_n)$ and $K(z, w)$ are used for $A_0^2(\mathbb{B}_n)$ and $K^0(z, w)$, respectively. These are the standard (unweighted) Bergman spaces. The function K^α is known as the (weighted) Bergman kernel and it satisfies the reproducing property

$$\begin{aligned} f(z) &= \langle f, K^\alpha(\cdot, z) \rangle_{A_\alpha^2(\mathbb{B}_n)} \\ &= \int_{\mathbb{B}_n} K^\alpha(z, w) f(w) dv_\alpha(w) , \end{aligned}$$

for any $f \in A_\alpha^2(\mathbb{B}_n)$, $z \in \mathbb{B}_n$. This last integral expression makes sense even for more general classes of functions. In fact, it represents the orthogonal projection from $L^2(\mathbb{B}_n, dv_\alpha)$ to $A^2(\mathbb{B}_n)$, also called the Bergman projection. For a bounded Lebesgue measurable function a or a regular Borel measure μ , both on \mathbb{B}_n , their Toeplitz operators \mathcal{T}_a and \mathcal{T}_μ are defined by

$$\mathcal{T}_a f(z) = \int_{\mathbb{B}_n} K^\alpha(z, w) a(w) f(w) dv_\alpha(w) , \quad (2)$$

and

$$\mathcal{T}_\mu f(z) = c_\alpha \int_{\mathbb{B}_n} K^\alpha(z, w) f(w) (1 - |w|^2)^\alpha d\mu(w) , \quad (3)$$

for any $f \in A^2(\mathbb{B}_n)$, $z \in \mathbb{B}_n$. In each case, a or μ is called the symbol of the respective Toeplitz operator. The operator \mathcal{T}_μ is a natural generalization of \mathcal{T}_a , which in turn is also defined as being the compression to $A^2(\mathbb{B}_n)$ of the multiplying operator $M_a f = af$.

We present the Toeplitz operators we will work with.

Definition 1.1. Let $\Gamma \subset \mathbb{B}_n$ be a Riemannian submanifold of dimension d , $d\sigma$ the Riemannian volume element on Γ , and $a : \Gamma \rightarrow \mathbb{C}$ a function such that $a \in C_0^2(\Gamma)$. We define the Toeplitz operator $T_{ad\sigma}$ by

$$T_{ad\sigma} f(z) = c_\alpha \int_{\Gamma} K^\alpha(z, w) f(w) a(w) (1 - |w|^2)^\alpha d\sigma(w) , \quad (4)$$

for every $f \in A_\alpha^2(\mathbb{B}_n)$, $z \in \mathbb{B}_n$.

Remark 1.2. We would like to recall that $\Gamma \subset \mathbb{B}_n$ being a Riemannian submanifold means that the inclusion $\iota : \Gamma \rightarrow \mathbb{B}_n$ is a smooth embedding (an immersion which is also a topological embedding) and that the Riemannian metric on Γ is the pullback of the Riemannian metric on \mathbb{B}_n (See for example [9], p.15,132). The Riemannian metric on \mathbb{B}_n will always be taken to be the one obtained from the Bergman Hermitian metric (see further below), which makes \mathbb{B}_n a model for hyperbolic geometry.

Remark 1.3. • Although not made explicit in the notation, $T_{ad\sigma}$ depends on α . We are interested in the behavior of $T_{ad\sigma}$ as α tends to infinity.

- If $d < 2n$, the resulting symbol measure $\mu = ad\sigma$ is singular with respect to the Lebesgue measure on \mathbb{B}_n .
- If $d = 2n$, Γ is an open subset of \mathbb{B}_n and the measure $d\sigma(w) = \frac{\pi^n}{n!} \frac{dv(w)}{(1-|w|^2)^{n+1}}$ is a multiple of the invariant measure on \mathbb{B}_n . Therefore $T_a := T_{ad\sigma}$ is the “classical” Toeplitz operator (2) with symbol $\frac{\pi^n}{n!} \frac{a(w)}{(1-|w|^2)^{n+1}}$. Explicitly

$$\mathcal{T}_a = T_{\frac{n!}{\pi^n} a(\cdot) (1-|\cdot|^2)^{n+1}} .$$

- As a is taken to have compact support, $T_{ad\sigma}$ is bounded, compact, and belongs to each Schatten class $S_p(A_\alpha^2(\mathbb{B}_n))$, $p > 0$.
- The condition $a \in C^2(\Gamma)$ can be changed to $a \in L^\infty(\Gamma)$ without affecting the definition of $T_{ad\sigma}$. In fact, it is not needed for a substantial part of the results. However, it is included in the definition since it is necessary for the main theorem.

Prior to stating the Szegő limit theorem we need to introduce some geometric definitions (see, for example, [6], [7], [10]).

By writing the local holomorphic coordinates as $z_\ell = x_\ell + iy_\ell$, $\ell = 1, \dots, n$, \mathbb{B}_n can be naturally regarded both as a complex manifold of dimension n and as a smooth manifold of dimension $2n$. For any $p \in \mathbb{B}_n$ the respective real and complex tangent spaces are given in terms of basic tangent vectors by

$$\mathbb{T}_p \mathbb{B}_n = \text{span} \left\{ \frac{\partial}{\partial x_\ell} \Big|_p, \frac{\partial}{\partial y_\ell} \Big|_p \right\}_{\ell=1, \dots, n} \quad \text{and} \quad \mathbb{T}_p \mathbb{B}_n^{\mathbb{C}} = \text{span} \left\{ \frac{\partial}{\partial z_\ell} \Big|_p, \frac{\partial}{\partial \bar{z}_\ell} \Big|_p \right\}_{\ell=1, \dots, n},$$

with the relation between the elements of these bases being given by the Wirtinger derivatives

$$\frac{\partial}{\partial z_\ell} = \frac{1}{2} \left(\frac{\partial}{\partial x_\ell} - i \frac{\partial}{\partial y_\ell} \right), \quad \frac{\partial}{\partial \bar{z}_\ell} = \frac{1}{2} \left(\frac{\partial}{\partial x_\ell} + i \frac{\partial}{\partial y_\ell} \right).$$

The Hermitian metric b defined by

$$b_{jk}(p) := \frac{1}{n+1} \frac{\partial^2}{\partial z_j \partial \bar{z}_k} \log K(z, z) \Big|_{z=p},$$

$j, k = 1, \dots, n$ on $\mathbb{T}_p \mathbb{B}_n^{\mathbb{C}}$ for each $p \in \mathbb{B}_n$ is called the Bergman Hermitian metric and it makes \mathbb{B}_n a Hermitian manifold. It induces a Riemannian metric \mathring{b} on the underlying smooth manifold, thus giving it the structure of a Riemannian manifold. The metric \mathring{b} is described in terms of b by the matrix

$$\mathring{B}(p) := \left(\mathring{b}_{jk}(p) \right) = \begin{pmatrix} \text{Re } b_{jk}(p) & \text{Im } b_{jk}(p) \\ -\text{Im } b_{jk}(p) & \text{Re } b_{jk}(p) \end{pmatrix}.$$

The endomorphism J defined by

$$J \left(\frac{\partial}{\partial x_\ell} \Big|_p \right) = \frac{\partial}{\partial y_\ell} \Big|_p, \quad J \left(\frac{\partial}{\partial y_\ell} \Big|_p \right) = -\frac{\partial}{\partial x_\ell} \Big|_p$$

is a complex structure on $T_p\mathbb{B}_n$ that is compatible with \mathring{b} in the sense that \mathring{b} is invariant under J . This complex structure is in turn induced by the natural one of \mathbb{R}^{2n} , $(x_1, \dots, x_n, y_1, \dots, y_n) \mapsto (-y_1, \dots, -y_n, x_1, \dots, x_n)$. We also consider the symplectic forms defined by

$$\omega_p(\xi, \nu) := \mathring{b}_p(\xi, J\nu)$$

for $\xi, \nu \in T_p\mathbb{B}_n$.

Let $V \subseteq T_p\mathbb{B}_n$ be a subspace and denote by

$$V^\omega := \{\xi \in T_p\mathbb{B}_n : \omega_p(\xi, v) = 0 \quad \forall v \in V\} \quad (5)$$

its symplectic complement.

Definition 1.4. We say that V is isotropic (respectively co-isotropic, respectively Lagrangian) if $V \subseteq V^\omega$ (respectively $V \supseteq V^\omega$, respectively $V = V^\omega$).

Example 1.5. From the skew-symmetry of ω_p , it is straightforward that any subspace V of dimension one is isotropic. On the other hand, if V has codimension one then it is co-isotropic, as in this case V^ω is isotropic and so $V^\omega \subseteq (V^\omega)^\omega = V$. In addition, the total subspace $T_p\mathbb{B}_n$ is trivially co-isotropic.

For $\Gamma \subseteq \mathbb{B}_n$ a submanifold, let $D_p\iota : T_p\Gamma \rightarrow T_p\mathbb{B}_n$ be the differential of the inclusion map $\iota : \Gamma \rightarrow \mathbb{B}_n$ at the point $p \in \Gamma$. Then $D_p\iota(T_p\Gamma)$ is a subspace of $T_p\mathbb{B}_n$ of dimension d .

Definition 1.6. Let $\Gamma \subseteq \mathbb{B}_n$ be a submanifold. We say that Γ is isotropic (respectively co-isotropic, Lagrangian) if for every $p \in \Gamma$, the subspace $D_p\iota(T_p\Gamma) \subseteq T_p\mathbb{B}_n$ is isotropic (respectively co-isotropic, Lagrangian).

Example 1.7. It follows from Example 1.5 that any curve is isotropic and any hypersurface is co-isotropic. Also, \mathbb{B}_n or any open subset of \mathbb{B}_n is co-isotropic.

Our Szegő limit theorem will apply to submanifolds that are isotropic or co-isotropic. In order to include both cases in the same statement, we introduce the notation

$$d' = \begin{cases} d & \Gamma \text{ isotropic} \\ 2n - d & \Gamma \text{ co-isotropic} \end{cases} .$$

We also normalize $T_{ad\sigma}$ by means of

$$\widehat{T}_{ad\sigma} := \frac{n!}{2^{\frac{d'}{2}} \pi^{\frac{d}{2}}} \alpha^{-n+\frac{d}{2}} T_{ad\sigma} . \quad (6)$$

We will prove in Corollary 5.2 that $\|\widehat{T}_{ad\sigma}\|_{\mathcal{L}(A_\alpha^2(\mathbb{B}_n))} = O(1)$ as α tends to infinity, thus justifying the usage of the term normalization.

1.2 Statement of the main result

We can now state the main result of this paper. It is a Bergman space analogue of the main result of [11].

Theorem 1.8. [Szegő limit theorem] *Let $\Gamma \subseteq \mathbb{B}_n$ be a isotropic or co-isotropic submanifold, $a \in C_0^2(\Gamma)$ a positive function, and $R > 0$ such that $\sup_{\xi \in \Gamma} \frac{a(\xi)}{(1-|\xi|^2)^{n+1}} < R$. If φ is a function such that $\frac{\varphi(t)}{t^p} \in C[0, R]$ for some $p > 0$, then*

$$\lim_{\alpha \rightarrow \infty} \left(\frac{\pi}{\alpha}\right)^{\frac{d}{2}} \text{Tr} \left(\varphi \left(\widehat{T}_{ad\sigma} \right) \right) = \frac{1}{2^{\frac{d'}{2}}} \int_{\Gamma} \mathcal{Q}_{\frac{d'}{2}}(\varphi) \left(\frac{a(\xi)}{(1-|\xi|^2)^{n+1}} \right) d\sigma(\xi) , \quad (7)$$

where \mathcal{Q}_ε is the operator

$$\mathcal{Q}_\varepsilon(\varphi)(t) := \frac{1}{\Gamma(\varepsilon)} \int_0^t \varphi(s) \left(\ln \frac{t}{s} \right)^{\varepsilon-1} \frac{ds}{s}$$

when $\varepsilon > 0$ ($\Gamma(\varepsilon)$ is the Gamma function evaluated at ε) and the identity when $\varepsilon = 0$.

We single out the following particular situation. A form of case $n = 1$ appears as Theorem 1.1 in [2].

Example 1.9 (The absolutely continuous case). The case $\Gamma = \mathbb{B}_n$ is trivially co-isotropic and $d' = 0$. We thus get

$$\lim_{\alpha \rightarrow \infty} \left(\frac{\pi}{\alpha}\right)^n \text{Tr} \left(\varphi \left(\widehat{T}_a \right) \right) = \frac{\pi^n}{n!} \int_{\mathbb{B}_n} \varphi \left(\frac{a(w)}{(1-|w|^2)^{n+1}} \right) \frac{dv(w)}{(1-|w|^2)^{n+1}} .$$

In terms of the “classical” operator \mathcal{T}_a (2), we have $\mathcal{T}_a = \widehat{T}_{a(\cdot)(1-|\cdot|^2)^{n+1}}$ and so

$$\lim_{\alpha \rightarrow \infty} \frac{n!}{\alpha^n} \text{Tr}(\varphi(\mathcal{T}_a)) = \int_{\mathbb{B}_n} \varphi(a(w)) \frac{dv(w)}{(1-|w|^2)^{n+1}} .$$

The article has the following structure. In the following section, we describe some aspects of the intrinsic geometry of Γ . We obtain a relation involving the Riemannian metric and the symplectic form in Γ that will be needed later. In Section 3 we derive an integral representation for the trace of a composition of multiple Toeplitz operators with symbols sharing the same support, and then in Section 4 we make use of the stationary phase method to study the asymptotic behavior of this integral as α grows to infinity. The expression we obtain relies on computing the determinant of a given Hessian matrix, which we leave for Appendix B. The proof for the Szegö limit theorem is delivered in Section 5, first for polynomials and then for a wider family of continuous functions. A key step in the proof involves bounding the traces for small powers (exponents $0 < p < 1$) of the Toeplitz operators. This is done in Appendix A. Section 6 is reserved for establishing some corollaries to the main theorem, involving results on eigenvalue distribution and Schatten norm estimations.

The line of reasoning used here follows that in [11], with some of the proofs presented being adaptations to the Bergman setting of the ones therein. In particular, the proofs in Section 6 and Appendix B.1 remain virtually unmodified but are regardless included for the sake of completeness. On the other hand, parts of appendices A and B.2 generalize their analogues there.

Throughout the article, we will use the lone capital letter C to represent an arbitrary constant that may change from line to line. When deemed appropriate, we will highlight its independence from other variables.

2 The Riemannian metric on Γ

Let $\gamma : B(0, 1) \rightarrow \Gamma$ be a parametrization. For each $p = \gamma(t) \in \Gamma$, $t = (t_1, \dots, t_d) \in B(0, 1)$, the tangent space $T_p\Gamma$ is spanned by the tangent vectors $\frac{\partial}{\partial \gamma_j} \Big|_t$, $j = 1, \dots, d$. These tangent vectors are defined by

$$\begin{aligned} \frac{\partial}{\partial \gamma_j} \Big|_t : C^\infty(\Gamma) &\rightarrow \mathbb{R} \\ f &\mapsto \frac{d}{ds} \left[f(\gamma(t_1, \dots, t_j + s, \dots, t_d)) \right] \Big|_{s=0} . \end{aligned}$$

Also recall that for each $v \in T_p\Gamma$, $D_p\iota(v)$ is the tangent vector to \mathbb{B}_n at $p = \iota(p)$ determined by

$$D_p\iota(v)(f) = v(f \circ \iota) , \quad f \in C^\infty(\mathbb{B}_n) .$$

As a submanifold $\Gamma \subseteq \mathbb{B}_n$, the Riemannian metric g in Γ is given by the pullback of the metric \mathring{b} in \mathbb{B}_n . More precisely

$$g_p(u, v) = \mathring{b}_p(D_p\iota(u), D_p\iota(v)) ,$$

$\forall u, v \in T_p\Gamma$. We also define a skew-symmetric, nondegenerate bilinear form by

$$h_p(u, v) = \omega_p(D_p\iota(u), (D_p\iota(v))) ,$$

$\forall u, v \in T_p\Gamma$.

The following lemma relating g_p and h_p looks rather technical, but will be key during Section 4.

Lemma 2.1. *When taking into account the basis $\left\{ \frac{\partial}{\partial \gamma_j} \Big|_t \right\}_{j=1, \dots, d}$ of $T_p\Gamma$, the components of the forms g_p and h_p satisfy the relations*

$$g_{jk}(p) + ih_{jk}(p) = \sum_{\ell, r=1}^n b_{\ell r}(p) \cdot \partial_j \gamma_\ell(t) \cdot \partial_k \overline{\gamma_r}(t) , \quad (8)$$

$$g_{jk}(p) - ih_{jk}(p) = \sum_{\ell, r=1}^n \overline{b_{\ell r}(p)} \cdot \partial_j \overline{\gamma_\ell}(t) \cdot \partial_k \gamma_r(t) , \quad (9)$$

for $j, k = 1, \dots, d$.

Proof. For convenience, we will write $\frac{\partial}{\partial x_{n+\ell}} \Big|_p = \frac{\partial}{\partial y_\ell} \Big|_p$. First we note that for each $j = 1, \dots, d$

$$D_p\iota \left(\frac{\partial}{\partial \gamma_j} \Big|_t \right) = \sum_{\ell=1}^{2n} \partial_j (\iota \circ \gamma)_\ell(t) \frac{\partial}{\partial x_\ell} \Big|_p ,$$

where $(\iota \circ \gamma)_\ell$ denotes the ℓ -th component function of $\iota \circ \gamma : B(0, 1) \subset \mathbb{R}^d \rightarrow \mathbb{B}_n \subseteq \mathbb{R}^{2n}$, and $\partial_j (\iota \circ \gamma)_\ell(t)$ its partial derivative respect to the j -th variable valued at t . This is obtained straight from the definitions and using the chain rule. We may now compute

$$\begin{aligned} g_{jk}(p) &= \mathring{b}_p \left(D_p\iota \left(\frac{\partial}{\partial \gamma_j} \Big|_t \right), D_p\iota \left(\frac{\partial}{\partial \gamma_k} \Big|_t \right) \right) \\ &= \sum_{\ell, r=1}^{2n} \partial_j (\iota \circ \gamma)_\ell(t) \partial_k (\iota \circ \gamma)_r(t) \mathring{b}_p \left(\frac{\partial}{\partial x_\ell} \Big|_p, \frac{\partial}{\partial x_r} \Big|_p \right) \end{aligned}$$

$$\begin{aligned}
&= \sum_{\ell,r=1}^{2n} \partial_j(\iota \circ \gamma)_\ell(t) \partial_k(\iota \circ \gamma)_r(t) \mathring{b}_{\ell r}(\gamma(t)) \\
&= \sum_{\ell,r=1}^n \left[\left(\partial_j(\iota \circ \gamma)_\ell(t) \partial_k(\iota \circ \gamma)_r(t) + \partial_j(\iota \circ \gamma)_{n+\ell}(t) \partial_k(\iota \circ \gamma)_{n+r}(t) \right) \right. \\
&\quad \cdot \text{Re}b_{\ell r}(\gamma(t)) + \left(\partial_j(\iota \circ \gamma)_\ell(t) \partial_k(\iota \circ \gamma)_{n+r}(t) \right. \\
&\quad \left. \left. - \partial_j(\iota \circ \gamma)_{n+\ell}(t) \partial_k(\iota \circ \gamma)_r(t) \right) \text{Im}b_{\ell r}(\gamma(t)) \right].
\end{aligned}$$

In a similar fashion, we get

$$\begin{aligned}
h_{jk}(p) &= \mathring{b}_p \left(\mathbb{D}_p \iota \left(\frac{\partial}{\partial \gamma_j} \Big|_t \right), J \left(\mathbb{D}_p \iota \left(\frac{\partial}{\partial \gamma_k} \Big|_t \right) \right) \right) \\
&= \sum_{\ell,r=1}^n \left[\left(\partial_j(\iota \circ \gamma)_\ell(t) \partial_k(\iota \circ \gamma)_r(t) + \partial_j(\iota \circ \gamma)_{n+\ell}(t) \partial_k(\iota \circ \gamma)_{n+r}(t) \right) \right. \\
&\quad \cdot \text{Im}b_{\ell r}(\gamma(t)) + \left(\partial_j(\iota \circ \gamma)_{n+\ell}(t) \partial_k(\iota \circ \gamma)_r(t) \right. \\
&\quad \left. \left. - \partial_j(\iota \circ \gamma)_\ell(t) \partial_k(\iota \circ \gamma)_{n+r}(t) \right) \text{Re}b_{\ell r}(\gamma(t)) \right].
\end{aligned}$$

Using these two relations

$$\begin{aligned}
g_{jk}(p) + ih_{jk}(p) &= \sum_{\ell,r=1}^n \left[\left(\partial_j(\iota \circ \gamma)_\ell \partial_k(\iota \circ \gamma)_r + \partial_j(\iota \circ \gamma)_{n+\ell} \partial_k(\iota \circ \gamma)_{n+r} \right) \right. \\
&\quad \cdot \left(\text{Re}b_{\ell r}(p) + i \text{Im}b_{\ell r}(p) \right) + \left(\partial_j(\iota \circ \gamma)_{n+\ell} \partial_k(\iota \circ \gamma)_r \right. \\
&\quad \left. \left. - \partial_j(\iota \circ \gamma)_\ell \partial_k(\iota \circ \gamma)_{n+r} \right) \left(i \text{Re}b_{\ell r}(p) - \text{Im}b_{\ell r}(p) \right) \right] \\
&= \sum_{\ell,r=1}^n b_{\ell r}(p) \left[\left(\partial_j(\iota \circ \gamma)_\ell \partial_k(\iota \circ \gamma)_r + \partial_j(\iota \circ \gamma)_{n+\ell} \partial_k(\iota \circ \gamma)_{n+r} \right) \right. \\
&\quad \left. + i \left(\partial_j(\iota \circ \gamma)_{n+\ell} \partial_k(\iota \circ \gamma)_r - \partial_j(\iota \circ \gamma)_\ell \partial_k(\iota \circ \gamma)_{n+r} \right) \right] \\
&= \sum_{\ell,r=1}^n b_{\ell r}(p) \left[\partial_j \left((\iota \circ \gamma)_\ell + i(\iota \circ \gamma)_{n+\ell} \right) \right.
\end{aligned}$$

$$\begin{aligned}
& \cdot \partial_k \left((\iota \circ \gamma)_r - i(\iota \circ \gamma)_{n+r} \right) \Big] \\
& = \sum_{\ell, r=1}^n b_{\ell r}(p) \cdot \partial_j \gamma_\ell(t) \cdot \partial_k \overline{\gamma_r}(t) ,
\end{aligned}$$

from which (9) is also deduced.

Notice that, as we have done when writing the local holomorphic coordinates of \mathbb{B}_n , we have used $(\iota \circ \gamma)_\ell + i(\iota \circ \gamma)_{n+\ell} = \gamma_\ell$. \square

3 The trace of a multiple composition

The first step towards a proof of Theorem 1.8 relies on representing in terms of an integral the composition of multiple Toeplitz operators with symbols sharing the same support.

We recall some notions from the classic theory of Berezin symbols and kernels ([1]). Let $A \in \mathcal{L}(A_\alpha^2(\mathbb{B}_n))$. The associated kernel K_A^α is the function

$$K_A^\alpha(z, w) := Ak_w(z) = \langle Ak_w, k_z \rangle_{A_\alpha^2(\mathbb{B}_n)} ,$$

where $k_z = \frac{K^\alpha(\cdot, z)}{\sqrt{K^\alpha(z, z)}} = \left(\frac{\sqrt{1-|z|^2}}{1-\langle \cdot, z \rangle} \right)^{n+1+\alpha}$ denotes the normalized reproducing kernel at z . In the case when A belongs to the trace class, then K_A^α satisfies

$$\mathrm{Tr}(A) = \int_{\mathbb{B}_n} K_A^\alpha(z, z) \, dv_\alpha(z) , \quad (10)$$

and whenever B is another bounded operator we have

$$K_{AB}^\alpha(z, w) = \int_{\mathbb{B}_n} K_A^\alpha(z, y) K_B^\alpha(y, w) \, dv_\alpha(y) . \quad (11)$$

For the special case when $A = T_{ad\sigma}$ is a Toeplitz operator, Fubini's theorem and the reproducing property of the Bergman kernels yield in the expression

$$K_{T_{ad\sigma}}^\alpha(z, w) = c_\alpha \int_{\Gamma} K^\alpha(z, \xi) K^\alpha(\xi, w) a(\xi) (1 - |\xi|^2)^\alpha \, d\sigma(\xi) . \quad (12)$$

We present a formula for the trace of a composition of m Toeplitz operators associated with the same manifold $\Gamma \subset \mathbb{B}_n$.

Lemma 3.1. *Let $m \geq 2$ be a natural number, $a_1, \dots, a_m \in L^\infty(\Gamma)$ with compact support, and $S := T_{a_1 d\sigma} \circ \dots \circ T_{a_m d\sigma}$. Then*

$$\begin{aligned} \text{Tr}(S) = c_\alpha^m \int_{\Gamma^m} \prod_{j(\bmod m)} \left[\left(\frac{1 - |\xi_j|^2}{1 - \langle \xi_j, \xi_{j+1} \rangle} \right)^\alpha \frac{a_j(\xi_j)}{(1 - \langle \xi_j, \xi_{j+1} \rangle)^{n+1}} \right] \\ \cdot d\sigma(\xi_1) \cdots d\sigma(\xi_m) . \quad (13) \end{aligned}$$

Proof. By induction on the relation (11) one can prove that

$$\begin{aligned} K_S^\alpha(z, w) = \int_{\mathbb{B}_n^{m-1}} K_{T_{a_1 d\sigma}}^\alpha(z, y_1) \left[\prod_{j=2}^{m-1} K_{T_{a_j d\sigma}}^\alpha(y_{j-1}, y_j) \right] K_{T_{a_m d\sigma}}^\alpha(y_{m-1}, w) \\ \cdot dv_\alpha(y_{m-1}) \cdots dv_\alpha(y_1) . \end{aligned}$$

We write $y_0 = z$ and $y_m = w$. Replacing all $K_{T_{a_j d\sigma}}^\alpha$ for their equivalent form in (12), and using Fubini's theorem together with the reproducing property of the Bergman kernel, we get

$$\begin{aligned} K_S^\alpha(z, w) &= c_\alpha^m \int_{\mathbb{B}_n^{m-1}} \prod_{j=1}^m \left[\int_{\Gamma} K^\alpha(y_{j-1}, \xi_j) K^\alpha(\xi_j, y_j) a_j(\xi_j) (1 - |\xi_j|^2)^\alpha d\sigma(\xi_j) \right] \\ &\quad \cdot dv_\alpha(y_{m-1}) \cdots dv_\alpha(y_1) \\ &= c_\alpha^m \int_{\Gamma^m} \int_{\mathbb{B}_n^{m-1}} \prod_{j=1}^m \left[K^\alpha(y_{j-1}, \xi_j) K^\alpha(\xi_j, y_j) \right] dv_\alpha(y_1) \cdots dv_\alpha(y_{m-1}) \\ &\quad \cdot \prod_{\ell=1}^m [a_\ell(\xi_\ell) (1 - |\xi_\ell|^2)^\alpha] d\sigma(\xi_1) \cdots d\sigma(\xi_m) \\ &= c_\alpha^m \int_{\Gamma^m} K^\alpha(y_0, \xi_1) K^\alpha(\xi_m, y_m) \prod_{j=1}^{m-1} \left[\int_{\mathbb{B}_n} K^\alpha(\xi_j, y_j) k_{\xi_{j+1}}(y_j) dv_\alpha(y_j) \right] \\ &\quad \cdot \prod_{\ell=1}^m [a_\ell(\xi_\ell) (1 - |\xi_\ell|^2)^\alpha] d\sigma(\xi_1) \cdots d\sigma(\xi_m) \\ &= c_\alpha^m \int_{\Gamma^m} K^\alpha(z, \xi_1) K^\alpha(\xi_m, w) \prod_{j=1}^{m-1} \left[K^\alpha(\xi_j, \xi_{j+1}) \right] \end{aligned}$$

$$\cdot \prod_{\ell=1}^m [a_\ell(\xi_\ell)(1 - |\xi_\ell|^2)^\alpha] d\sigma(\xi_1) \cdots d\sigma(\xi_m) .$$

Using the trace formula in (10),

$$\begin{aligned} \mathrm{Tr}(S) &= \int_{\mathbb{B}_n} K_S^\alpha(z, z) dv_\alpha(z) \\ &= c_\alpha^m \int_{\mathbb{B}_n} \int_{\Gamma^m} K^\alpha(z, \xi_1) K^\alpha(\xi_m, z) \prod_{j=1}^{m-1} [K^\alpha(\xi_j, \xi_{j+1})] \\ &\quad \cdot \prod_{\ell=1}^m [a_\ell(\xi_\ell)(1 - |\xi_\ell|^2)^\alpha] d\sigma(\xi_1) \cdots d\sigma(\xi_m) dv_\alpha(z) \\ &= c_\alpha^m \int_{\Gamma^m} \int_{\mathbb{B}_n} K^\alpha(\xi_m, z) k_{\xi_1}(z) dv_\alpha(z) \prod_{j=1}^{m-1} [K^\alpha(\xi_j, \xi_{j+1})] \\ &\quad \cdot \prod_{\ell=1}^m [a_\ell(\xi_\ell)(1 - |\xi_\ell|^2)^\alpha] d\sigma(\xi_1) \cdots d\sigma(\xi_m) \\ &= c_\alpha^m \int_{\Gamma^m} \prod_{j \pmod{m}} [K^\alpha(\xi_j, \xi_{j+1}) a_j(\xi_j)(1 - |\xi_j|^2)^\alpha] d\sigma(\xi_1) \cdots d\sigma(\xi_m) , \end{aligned}$$

from where the statement follows. \square

We rewrite the expression (13) in the form

$$\begin{aligned} \mathrm{Tr}(S) &= c_\alpha^m \int_{\Gamma^m} \exp \left[i\alpha \left(i \mathrm{Log} \prod_{j=1}^m \frac{1 - \langle \xi_j, \xi_{j+1} \rangle}{1 - |\xi_j|^2} \right) \right] \left(\prod_{j=1}^m \frac{a_j(\xi_j)}{(1 - \langle \xi_j, \xi_{j+1} \rangle)^{n+1}} \right) \\ &\quad \cdot d\sigma(\xi_1) \cdots d\sigma(\xi_m) , \quad (14) \end{aligned}$$

with the understanding that the subindexes are always taken mod m (i.e. $\xi_{m+1} = \xi_1$). We also define

$$\begin{aligned} \Phi(\xi_1, \dots, \xi_m) &:= i \mathrm{Log} \prod_{j=1}^m \frac{1 - \langle \xi_j, \xi_{j+1} \rangle}{1 - |\xi_j|^2} \\ &= i \sum_{j=1}^m \mathrm{Log} \frac{1 - \langle \xi_j, \xi_{j+1} \rangle}{1 - |\xi_j|^2} \end{aligned} \quad (15)$$

$$= \frac{i}{n+1} \sum_{j=1}^m [\text{Log } K(\xi_j, \xi_j) - \text{Log } K(\xi_j, \xi_{j+1})] . \quad (16)$$

This notation will be useful when using stationary phase method later.

Remark 3.2. Notice that $\frac{1-\langle \xi_j, \xi_{j+1} \rangle}{1-|\xi_j|^2} \in (-\infty, 0]$ if and only if $\langle \xi_j, \xi_{j+1} \rangle \in [1, \infty)$. However, this does not occur since $\xi_j, \xi_{j+1} \in \mathbb{B}_n$. It follows from (15) that Φ is well defined and continuous on Γ^m . Moreover, from (14) we find that all the ξ_j involved in the integration are the ones in the support of a_j . Since each a_j has compact support in Γ , we conclude that there is an open neighborhood of $(-\infty, 0]$ such that only values outside this neighborhood are used to evaluate Log in the expression for Φ when computing (14).

We describe the behavior of $\text{Im}(\Phi(\xi_1, \dots, \xi_m))$ with the following result.

Lemma 3.3. $\text{Im}(\Phi(\xi_1, \dots, \xi_m)) \geq 0$ for any $(\xi_1, \dots, \xi_m) \in \Gamma^m$, with equality if and only if $\xi_1 = \xi_2 = \dots = \xi_m$.

We will make use of the following claim.

Claim 3.4. Let $m \in \mathbb{N}$, $m \geq 2$. Then for any $\{d_1, \dots, d_m\} \subset (0, 1)$,

$$\prod_{j=1}^m \frac{1 - d_j d_{j+1}}{1 - d_j^2} \geq 1 , \quad (17)$$

where $d_{m+1} := d_1$. Moreover, the equality holds if and only if $d_1 = d_2 = \dots = d_m$.

Proof. The proof is straightforward when $d_1 = d_2 = \dots = d_m$. Suppose that not all d_j are the same and let $d_0 := d_m$. We must show that the inequality in (17) is strict. Notice that if k' is such that $d_{k'+1} = d_{k'}$ then

$$\frac{1 - d_{k'-1} d_{k'}}{1 - d_{k'-1}^2} \cdot \frac{1 - d_{k'} d_{k'+1}}{1 - d_{k'}^2} = \frac{1 - d_{k'-1} d_{k'+1}}{1 - d_{k'-1}^2} ,$$

and so

$$\prod_{j=1}^m \frac{1 - d_j d_{j+1}}{1 - d_j^2} = \frac{1 - d_{k'-1} d_{k'+1}}{1 - d_{k'-1}^2} \prod_{\substack{j=1 \\ j \neq k'-1, k'}}^m \frac{1 - d_j d_{j+1}}{1 - d_j^2} .$$

The rightmost term is a product of the form of (17) over the set of $m - 1$ elements $\{d_1, \dots, d_m\} \setminus \{d_{k'}\}$. This shows that we can omit repeated consecutive elements from the list d_1, \dots, d_m and still get the same result. Hence, without loss of generality, we can assume that d_1, \dots, d_m are all different from their neighbors modulo m .

We proceed by induction over m . If $m = 2$ then

$$\frac{(1 - d_1 d_2)^2}{(1 - d_1^2)(1 - d_2^2)} = \frac{(1 - d_1 d_2)^2}{(1 - d_1 d_2)^2 - (d_1 - d_2)^2} > 1 .$$

Now assume the statement holds for $m - 1$. First, we make an observation: There exists $k \in \{1, \dots, m\}$ such that $d_k > d_{k-1}, d_{k+1}$ or $d_k < d_{k-1}, d_{k+1}$. To see this, suppose $d_2 > d_1$. If $d_3 < d_2$, then $k = 2$ satisfies the property, and if not, then $d_1 < d_2 < d_3$. Similarly if $d_4 < d_3$ we have finished, otherwise $d_1 < d_2 < d_3 < d_4$. In the worst scenario $d_1 < \dots < d_m$, which implies that $d_m > d_{m-1}, d_1$. The reasoning for when $d_2 < d_1$ is analogous.

Let k be such an index and define the quadratic polynomial $p(x) = (x - d_{k-1})(x - d_{k+1})$. Notice that for $|x| \neq 1$

$$p(x) > 0 \Leftrightarrow \frac{(1 - d_{k-1}x)(1 - d_{k+1}x)}{1 - x^2} > 1 - d_{k-1}d_{k+1} .$$

The roots of p are d_{k-1} and d_{k+1} , and so $p(x) > 0$ for all $x > d_{k-1}, d_{k+1}$ or $x < d_{k-1}, d_{k+1}$. In particular $p(d_k) > 0$, therefore

$$\frac{(1 - d_{k-1}d_k)(1 - d_{k+1}d_k)}{1 - d_k^2} > 1 - d_{k-1}d_{k+1} ,$$

and

$$\frac{1 - d_{k-1}d_k}{1 - d_{k-1}^2} \cdot \frac{1 - d_k d_{k+1}}{1 - d_k^2} > \frac{1 - d_{k-1}d_{k+1}}{1 - d_{k-1}^2} .$$

Then

$$\prod_{j=1}^m \frac{1 - d_j d_{j+1}}{1 - d_j^2} > \frac{1 - d_{k-1}d_{k+1}}{1 - d_{k-1}^2} \prod_{\substack{j=1 \\ j \neq k-1, k}}^m \frac{1 - d_j d_{j+1}}{1 - d_j^2} ,$$

which by the induction hypothesis is greater than 1 because, as before, the rightmost term is a product of the form of (17) over a set of $m - 1$ elements. \square

Proof of Lemma 3.3. For each $j = 1, \dots, m$ we have

$$|1 - \langle \xi_j, \xi_{j+1} \rangle| \geq 1 - |\langle \xi_j, \xi_{j+1} \rangle| \geq 1 - |\xi_j| |\xi_{j+1}|, \quad (18)$$

which we use for

$$\begin{aligned} \operatorname{Im}(\Phi(\xi_1, \dots, \xi_m)) &= \operatorname{Im} \left(i \operatorname{Log} \prod_{j=1}^m \frac{1 - \langle \xi_j, \xi_{j+1} \rangle}{1 - |\xi_j|^2} \right) \\ &= \ln \left| \prod_{j=1}^m \frac{1 - \langle \xi_j, \xi_{j+1} \rangle}{1 - |\xi_j|^2} \right| \\ &\geq \ln \prod_{j=1}^m \frac{1 - |\xi_j| |\xi_{j+1}|}{1 - |\xi_j|^2}. \end{aligned}$$

The last product is nonnegative by (17), thus proving the inequality part. The if part of the equality characterization is straightforward. On the other hand, if the equality holds, then by Claim 3.4 $|\xi_1| = \dots = |\xi_m|$. Inequality (18) involves the Cauchy-Schwarz inequality, so the equality being attained implies that ξ_j and ξ_{j+1} are linearly dependent, and thus $\xi_j = \pm \xi_{j+1}$. Finally, if $\xi_j = -\xi_{j+1}$, then

$$|1 - \langle \xi_j, \xi_{j+1} \rangle| = 1 + |\xi_j|^2 > 1 - |\xi_j| |\xi_{j+1}|,$$

which contradicts the assumption of equality, hence $\xi_j = \xi_{j+1}$ and $\xi_1 = \dots = \xi_m$. \square

4 Asymptotic of the trace of a multiple composition

This section is devoted to finding the asymptotic behavior of the integral expression in (13) when α tends to infinity. Throughout the section, we will keep $m \geq 2$ and $a_1, \dots, a_m \in C_0^2(\Gamma)$ fixed and denote $S := T_{a_1 d\sigma} \circ \dots \circ T_{a_m d\sigma}$.

Consider $\gamma : B(0, 1) \subset \mathbb{R}^d \rightarrow \Gamma$, $t = (t_1, \dots, t_d) \mapsto \gamma(t)$ a parametrization of an open set in Γ . Let $t, s_1, \dots, s_{m-1}, s_q = (s_1^{(q)}, \dots, s_d^{(q)})$ denote m variables in \mathbb{R}^d and $\mathbf{s} = (s_1, \dots, s_{m-1}) \in \mathbb{R}^{d(m-1)}$. We define

$$\begin{aligned} \Xi &: B(0, \varepsilon)^m \subset \mathbb{R}^{dm} \rightarrow \Gamma^m \\ \Xi(t, \mathbf{s}) &= (\gamma(t), \gamma(t + s_1), \dots, \gamma(t + s_{m-1})). \end{aligned}$$

Choosing ε small enough Ξ is a parametrization for an open set $\mathcal{V} \subseteq \Gamma^m$, which intersects the diagonal of Γ^m whenever $\mathbf{s} = \mathbf{0}$, i.e., $s_1 = \dots = s_{m-1} = 0$. Let $\chi \in C_0^\infty(\mathcal{V})$. We wish to compute

$$I_\chi(\alpha) = c_\alpha^m \int_{\mathcal{V}} e^{i\alpha\Phi(\xi_1, \dots, \xi_m)} \left(\prod_{j=1}^m \frac{a_j(\xi_j)}{(1 - \langle \xi_j, \xi_{j+1} \rangle)^{n+1}} \right) \chi(\xi_1, \dots, \xi_m) \cdot d\sigma(\xi_1) \cdots d\sigma(\xi_m), \quad (19)$$

asymptotically as $\alpha \rightarrow \infty$ using stationary phase method through the parametrization Ξ . The stationary phase will be made over \mathbf{s} with t fixed, and then we will integrate over t . Regarding \mathcal{V} as a subset of the product manifold Γ^m , its Riemannian metric G_Ξ satisfies $\det G_\Xi(t, \mathbf{s}) = \det G(t) \cdot \det G(t + s_1) \cdots \det G(t + s_{m-1})$. Make $\varphi(t, \mathbf{s}) = \Phi \circ \Xi = \Phi(\gamma(t), \gamma(t + s_1), \dots, \gamma(t + s_{m-1}))$ so that we have

$$\begin{aligned} \frac{I_\chi(\alpha)}{c_\alpha^m} &= \int_{B(0, \varepsilon)^m} e^{i\alpha\Phi(\Xi(t, \mathbf{s}))} \left(\prod_{j=1}^m \frac{a_j(\gamma(t + s_{j-1}))}{(1 - \langle \gamma(t + s_{j-1}), \gamma(t + s_j) \rangle)^{n+1}} \right) \chi(\Xi(t, \mathbf{s})) \\ &\quad \cdot \det G_\Xi(t, \mathbf{s})^{\frac{1}{2}} d(t, \mathbf{s}) \\ &= \int_{B(0, \varepsilon)} \left[\int_{B(0, \varepsilon)^{m-1}} e^{i\alpha\varphi(t, \mathbf{s})} \left(\prod_{j=1}^m \frac{a_j(\gamma(t + s_{j-1})) \det G(t + s_j)^{\frac{1}{2}}}{(1 - \langle \gamma(t + s_{j-1}), \gamma(t + s_j) \rangle)^{n+1}} \right) \right. \\ &\quad \left. \cdot \chi(\Xi(t, \mathbf{s})) d\mathbf{s} \right] dt \end{aligned} \quad (20)$$

(For simplicity of the expression, we have adopted the convention that $s_0 := 0 =: s_m$). We claim that $\varphi(t, \cdot)$ has a critic point at $\mathbf{s} = \mathbf{0}$. Let $1 \leq q \leq m - 1$ and $1 \leq j \leq d$. We differentiate $\varphi(t, \cdot)$ with respect to $s_j^{(q)}$ using (16) and that $\text{Log } K(z, w)$ is holomorphic in z and antiholomorphic in w . We obtain

$$-i \frac{\partial}{\partial s_j^{(q)}} \varphi(t, \mathbf{s}) = \frac{1}{n+1} \sum_{\ell=1}^n \left[\frac{\partial}{\partial z_\ell} \text{Log } K(\gamma(t + s_q), \gamma(t + s_q)) \cdot \partial_j \gamma_\ell(t + s_q) \right]$$

$$\begin{aligned}
& + \frac{\partial}{\partial \bar{w}_\ell} \text{Log } K(\gamma(t + s_q), \gamma(t + s_q)) \cdot \partial_j \bar{\gamma}_\ell(t + s_q) \\
& - \frac{\partial}{\partial z_\ell} \text{Log } K(\gamma(t + s_q), \gamma(t + s_{q+1})) \cdot \partial_j \gamma_\ell(t + s_q) \\
& - \frac{\partial}{\partial \bar{w}_\ell} \text{Log } K(\gamma(t + s_{q-1}), \gamma(t + s_q)) \cdot \partial_j \bar{\gamma}_\ell(t + s_q) \Big] \\
& \hspace{15em} (21)
\end{aligned}$$

from which it follows that $\nabla_{\mathbf{s}} \varphi(t, \mathbf{0}) = 0$.

We now use (21) to compute the Hessian of φ at $\mathbf{s} = \mathbf{0}$. Let $1 \leq q' \leq m-1$ and $1 \leq k \leq d$. Notice that if $q' \notin \{q-1, q, q+1\}$, then (21) does not depend on $s_k^{(q')}$, and therefore $\frac{\partial^2}{\partial s_k^{(q')} \partial s_j^{(q)}} \varphi(t, \mathbf{s}) = 0$. For the remaining three no null cases we have

$$\begin{aligned}
\frac{\partial^2}{\partial s_k^{(q-1)} \partial s_j^{(q)}} \varphi(t, \mathbf{0}) &= \frac{-i}{n+1} \sum_{\ell, r=1}^n \left[\frac{\partial^2}{\partial z_r \partial \bar{w}_\ell} \text{Log } K(\gamma(t), \gamma(t)) \partial_j \bar{\gamma}_\ell(t) \partial_k \gamma_r(t) \right], \\
\frac{\partial^2}{\partial s_k^{(q+1)} \partial s_j^{(q)}} \varphi(t, \mathbf{0}) &= \frac{-i}{n+1} \sum_{\ell, r=1}^n \left[\frac{\partial^2}{\partial z_\ell \partial \bar{w}_r} \text{Log } K(\gamma(t), \gamma(t)) \partial_j \gamma_\ell(t) \partial_k \bar{\gamma}_r(t) \right], \\
i \frac{\partial^2}{\partial s_k^{(q)} \partial s_j^{(q)}} \varphi(t, \mathbf{0}) &= \frac{1}{n+1} \sum_{\ell, r=1}^n \left[\frac{\partial^2}{\partial z_\ell \partial \bar{w}_r} \text{Log } K(\gamma(t), \gamma(t)) \partial_j \gamma_\ell(t) \partial_k \bar{\gamma}_r \right. \\
& \quad \left. + \frac{\partial^2}{\partial z_r \partial \bar{w}_\ell} \text{Log } K(\gamma(t), \gamma(t)) \partial_j \bar{\gamma}_\ell(t) \partial_k \gamma_r \right].
\end{aligned}$$

Recall that

$$\frac{1}{n+1} \frac{\partial^2}{\partial z_\ell \partial \bar{z}_r} \log K(z, z) \Big|_{z=\gamma(t)}$$

are the components $b_{\ell r}$ of the Bergman Hermitian metric of \mathbb{B}_n at $\gamma(t)$. Comparing with (8) and (9) of Lemma 2.1 we get

$$\begin{aligned}
-i \frac{\partial^2}{\partial s_k^{(q-1)} \partial s_j^{(q)}} \varphi(t, \mathbf{0}) &= -g_{jk}(p) + ih_{jk}(p), \\
-i \frac{\partial^2}{\partial s_k^{(q+1)} \partial s_j^{(q)}} \varphi(t, \mathbf{0}) &= -g_{jk}(p) - ih_{jk}(p),
\end{aligned}$$

$$-i \frac{\partial^2}{\partial s_k^{(g)} \partial s_j^{(g)}} \varphi(t, \mathbf{0}) = 2g_{jk}(p),$$

where in accordance with Section 2, $g_{jk}(p)$ and $h_{jk}(p)$ are the components at $p = \gamma(t)$ of the Riemannian metric g and the skew-symmetric form h in Γ . If we define the matrices $G = (g_{jk}(p))$ and $H = (h_{jk}(p))$, then the Hessian matrix of φ at $\mathbf{s} = \mathbf{0}$ can be expressed as a $(m-1) \times (m-1)$ block matrix with blocks of size $d \times d$ in the form

$$-i \text{Hess} \varphi(t, \mathbf{0}) = \begin{pmatrix} 2G & -G - iH & 0 & \dots & 0 \\ -G + iH & 2G & -G - iH & \dots & 0 \\ 0 & -G + iH & 2G & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & \dots & -G + iH & 2G & -G - iH \\ 0 & \dots & 0 & -G + iH & 2G \end{pmatrix}.$$

To express the determinant of the Hessian, make $W := G^{-1}H$ to rewrite

$$-i \text{Hess} \varphi(t, \mathbf{0}) = \begin{pmatrix} G & 0 & \dots & 0 \\ 0 & G & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & \dots & G & 0 \\ 0 & \dots & 0 & G \end{pmatrix} \begin{pmatrix} 2I & -I - iW & \dots & 0 \\ -I + iW & 2I & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & \dots & 2I & -I - iW \\ 0 & \dots & -I + iW & 2I \end{pmatrix}, \quad (22)$$

and let \mathcal{O}_{m-1} be the rightmost matrix of the factorization. We then have

$$\det(-i \text{Hess} \varphi(t, \mathbf{0})) = (\det G)^{m-1} \det \mathcal{O}_{m-1}.$$

In Appendix B it is shown that $\det \mathcal{O}_{m-1}$ is positive, while so is $\det G$ because G is positive-definite. This implies that $\det(-i \text{Hess} \varphi(t, \mathbf{0}))$ is also positive and so the application of the stationary phase method is justified. Apply Theorem 7.7.5 [5] to estimate the integral inside the brackets in (20), to obtain

$$\int_{B(0, \varepsilon)^{m-1}} e^{i\alpha \varphi(t, \mathbf{s})} \left(\prod_{j=1}^m \frac{a_j(\gamma(t + s_{j-1})) \det G(t + s_j)^{\frac{1}{2}}}{(1 - \langle \gamma(t + s_{j-1}), \gamma(t + s_j) \rangle)^{n+1}} \right) \chi(\Xi(t, \mathbf{s})) \, d\mathbf{s} =$$

$$\det \left(\frac{\alpha}{2\pi i} \text{Hess} \varphi(t, \mathbf{0}) \right)^{-\frac{1}{2}} \left[\prod_{j=1}^m \frac{a_j(\gamma(t)) \det G(t)^{\frac{1}{2}}}{(1 - |\gamma(t)|^2)^{n+1}} \right] \chi(\gamma(t), \dots, \gamma(t))$$

$$\cdot (1 + O(\alpha^{-1})).$$

Substituting this on (20) we get

$$\begin{aligned}
\frac{I_\chi(\alpha)}{c_\alpha^m} &= \int_{B(0,\varepsilon)} \left[\det \left(\frac{\alpha}{2\pi i} \text{Hess}\varphi(t,0) \right)^{-\frac{1}{2}} \left(\prod_{j=1}^m \frac{a_j(\gamma(t)) \det G(t)^{\frac{1}{2}}}{(1-|\gamma(t)|^2)^{n+1}} \right) \right. \\
&\quad \left. \cdot \chi(\gamma(t), \dots, \gamma(t)) (1 + O(\alpha^{-1})) \right] dt \\
&= \left(\frac{2\pi}{\alpha} \right)^{\frac{d(m-1)}{2}} \int_{B(0,\varepsilon)} \left(\prod_{j=1}^m a_j(\gamma(t)) \right) \cdot \frac{\chi(\gamma(t), \dots, \gamma(t))}{(1-|\gamma(t)|^2)^{m(n+1)}} \\
&\quad \cdot \frac{\det G(t)^{\frac{1}{2}}}{(\det \mathcal{O}_{m-1})^{\frac{1}{2}}} dt (1 + O(\alpha^{-1})) \\
&= \left(\frac{2\pi}{\alpha} \right)^{\frac{d(m-1)}{2}} \int_{\gamma(B(0,\varepsilon))} \left(\prod_{j=1}^m a_j(\xi) \right) \frac{\chi(\xi, \dots, \xi)}{(1-|\xi|^2)^{m(n+1)}} \frac{d\sigma(\xi)}{(\det \mathcal{O}_{m-1})^{\frac{1}{2}}} \\
&\quad \cdot (1 + O(\alpha^{-1})) . \tag{23}
\end{aligned}$$

In this last expression, the integration can be changed to be done on all Γ instead of just $\gamma(B(0,\varepsilon))$ without affecting the outcome. To see this, note that if $\xi \notin \gamma(B(0,\varepsilon))$ then $(\xi, \dots, \xi) \notin \Xi(B(0,\varepsilon)^m) = \mathcal{V} \supset \text{supp}\chi$.

Let $\gamma_1, \dots, \gamma_L : B(0,1) \subset \mathbb{R}^d \rightarrow \Gamma$ be parametrizations of Γ such that $\cup_{\ell=1}^L \gamma_\ell(B(0,1)) = \Gamma$. Without loss of generality, we have taken this as a finite union since, even if Γ is not compact, the integral in which we are interested is done on the supports of the functions a_j which have compact support. Take ε small enough and define

$$\Xi_\ell : B(0,\varepsilon)^m \rightarrow \Gamma^m$$

$$\Xi_\ell(t, \mathbf{s}) = (\gamma_\ell(t), \gamma_\ell(t + s_1), \dots, \gamma_\ell(t + s_{m-1})) ,$$

so that each $\Xi_\ell(B(0,\varepsilon)^m)$ is, as described in the previous construction, an open set in Γ^m that intersects the diagonal when $\mathbf{s} = \mathbf{0}$. Then

$$\mathcal{U} := \bigcup_{\ell=1}^L \Xi_\ell(B(0,\varepsilon)^m)$$

is an open neighborhood of the diagonal in Γ^m and $\{\Xi_\ell, B(0, \varepsilon)^m\}_{\ell=1, \dots, L}$ is an atlas for \mathcal{U} . It follows from (14) and Lemma 3.3 that $\text{Tr}(S)$ decays exponentially fast outside \mathcal{U} when α grows. Let $\{\chi_\ell\}_{\ell=1, \dots, L}$ be a partition of unity subordinated to the aforementioned atlas. Applying what we obtained in (23) to each χ_ℓ and substituting it into (14) we get

$$\begin{aligned}
\text{Tr}(S) &= c_\alpha^m \int_{\mathcal{U}} e^{i\alpha\Phi(\xi_1, \dots, \xi_m)} \left(\prod_{j=1}^m \frac{a_j(\xi_j)}{(1 - \langle \xi_j, \xi_{j+1} \rangle)^{n+1}} \right) d\sigma(\xi_1) \cdots d\sigma(\xi_m) \\
&\quad + O(\alpha^{-\infty}) \\
&= c_\alpha^m \sum_{\ell=1}^L \int_{\Xi_\ell(B(0, \varepsilon)^m)} e^{i\alpha\Phi(\xi_1, \dots, \xi_m)} \left(\prod_{j=1}^m \frac{a_j(\xi_j)}{(1 - \langle \xi_j, \xi_{j+1} \rangle)^{n+1}} \right) \chi_\ell(\xi_1, \dots, \xi_m) \\
&\quad \cdot d\sigma(\xi_1) \cdots d\sigma(\xi_m) + O(\alpha^{-\infty}) \\
&= \sum_{\ell=1}^L I_{\chi_\ell}(\alpha) + O(\alpha^{-\infty}) \\
&= c_\alpha^m \left(\frac{2\pi}{\alpha} \right)^{\frac{d(m-1)}{2}} \sum_{\ell=1}^L \int_{\Gamma} \left(\prod_{j=1}^m a_j(\xi) \right) \frac{\chi_\ell(\xi, \dots, \xi)}{(1 - |\xi|^2)^{m(n+1)}} \frac{d\sigma(\xi)}{(\det \mathcal{O}_{m-1})^{\frac{1}{2}}} \\
&\quad \cdot (1 + O(\alpha^{-1})) \\
&= c_\alpha^m \left(\frac{2\pi}{\alpha} \right)^{\frac{d(m-1)}{2}} \int_{\Gamma} \left(\prod_{j=1}^m a_j(\xi) \right) \frac{\sum_{\ell=1}^L \chi_\ell(\xi, \dots, \xi)}{(1 - |\xi|^2)^{m(n+1)}} \frac{d\sigma(\xi)}{(\det \mathcal{O}_{m-1})^{\frac{1}{2}}} \\
&\quad \cdot (1 + O(\alpha^{-1})) \\
&= c_\alpha^m \left(\frac{2\pi}{\alpha} \right)^{\frac{d(m-1)}{2}} \int_{\Gamma} \left(\prod_{j=1}^m \frac{a_j(\xi)}{(1 - |\xi|^2)^{n+1}} \right) \frac{d\sigma(\xi)}{(\det \mathcal{O}_{m-1})^{\frac{1}{2}}} (1 + O(\alpha^{-1})), \tag{24}
\end{aligned}$$

since $\sum_{\ell} \chi_\ell = 1$ in Γ .

In order to obtain an explicit formula from (24), we would like $\det \mathcal{O}_{m-1}$ to be computable, while it would be ideal for it to be constant for all $\xi = \gamma(t) \in \Gamma$. This does not need to be true in general, but in Appendix B we show that this is the case when Γ is isotropic or co-isotropic. Namely, in

Proposition B.11 we show that

$$\sqrt{\det \mathcal{O}_{m-1}} = \begin{cases} m^{\frac{d}{2}} & \Gamma \text{ isotropic} \\ 2^{(d-n)(m-1)} m^{n-\frac{d}{2}} & \Gamma \text{ co-isotropic} \end{cases}, \quad (25)$$

which, rewriting it in terms of d' , leads to

$$2^{\frac{d(m-1)}{2}} (\det \mathcal{O}_{m-1})^{-\frac{1}{2}} = 2^{\frac{d'(m-1)}{2}} m^{-\frac{d'}{2}} \quad (26)$$

for both Γ isotropic or co-isotropic. Also, recall that $\widehat{T}_{ad\sigma} = \frac{n!}{2^{\frac{d'}{2}} \pi^{\frac{d}{2}}} \alpha^{-n+\frac{d}{2}} T_{ad\sigma}$. We conclude this section by presenting a statement that describes the resulting estimation when adding this hypothesis.

Lemma 4.1. [Asymptotics for the trace of a composition] *Let Γ be isotropic or co-isotropic and $a_1, \dots, a_m \in C_0^2(\Gamma)$. Then*

$$\text{Tr} \left(\widehat{T}_{a_1 d\sigma} \circ \dots \circ \widehat{T}_{a_m d\sigma} \right) = \frac{\left(\frac{\alpha}{\pi}\right)^{\frac{d}{2}}}{(2m)^{\frac{d'}{2}}} \int_{\Gamma} \prod_{j=1}^m \frac{a_j(\xi)}{(1-|\xi|^2)^{n+1}} d\sigma(\xi) \left(1 + O\left(\frac{1}{\alpha}\right) \right).$$

Proof. It can be verified using Stirling's formula that

$$c_{\alpha} = \frac{\Gamma(\alpha + 1 + n)}{n! \Gamma(\alpha + 1)} = \frac{\alpha^n}{n!} \left(1 + O\left(\frac{1}{\alpha}\right) \right).$$

By the linearity of the trace, the result will follow from (24) by substituting the last identity for c_{α} and the expression for $2^{\frac{d(m-1)}{2}} (\det \mathcal{O}_{m-1})^{-\frac{1}{2}}$ given in (26). \square

5 Section with the main theorem

In this section, we demonstrate Theorem 1.8. First, we state the result for monomial functions and then extend it to polynomials by linearity. Next, we will use a Weierstrass argument to derive the theorem for continuous functions.

5.1 The Szegő limit theorem for polynomials

As a consequence of Lemma 4.1 we obtain the following proposition, which can be seen as a first form of our Szegő limit theorem applying to monomial functions $\varphi(t) = t^m$.

Proposition 5.1. *Let Γ be isotropic or co-isotropic and $a \in C_0^2(\Gamma)$. Then*

$$\mathrm{Tr} \left(\left(\widehat{T}_{ad\sigma} \right)^m \right) = \frac{\left(\frac{\alpha}{\pi} \right)^{\frac{d}{2}}}{(2m)^{\frac{d'}{2}}} \int_{\Gamma} \left(\frac{a(\xi)}{(1-|\xi|^2)^{n+1}} \right)^m d\sigma(\xi) \left(1 + O \left(\frac{1}{\alpha} \right) \right). \quad (27)$$

Proof. Take $a_1 = \dots = a_m = a$ in Lemma 4.1. \square

As a first application, we obtain an estimation of the growth rate of the operator norm of the Toeplitz operators. This result shows that the operators defined in (6) are indeed normalizations of the Toeplitz operators defined in Definition 1.1.

Corollary 5.2. *Let Γ be isotropic or co-isotropic and $a \in C_0^2(\Gamma)$. There exists a constant $C > 0$ such that*

$$\| \widehat{T}_{ad\sigma} \|_{\mathcal{L}(A_\alpha^2(\mathbb{B}_n))} < C$$

for all $\alpha > 0$.

Proof. Let $a = a_1 + ia_2$ with a_1 and a_2 real valued. We can write

$$\widehat{T}_{ad\sigma} = \widehat{T}_{(a_1+|a|\infty)d\sigma} + i\widehat{T}_{(a_2+|a|\infty)d\sigma} - (1+i)\widehat{T}_{|a|\infty d\sigma},$$

and so

$$\begin{aligned} \| \widehat{T}_{ad\sigma} \|_{\mathcal{L}(A_\alpha^2(\mathbb{B}_n))} &\leq \| \widehat{T}_{(a_1+|a|\infty)d\sigma} \|_{\mathcal{L}(A_\alpha^2(\mathbb{B}_n))} + \| \widehat{T}_{(a_2+|a|\infty)d\sigma} \|_{\mathcal{L}(A_\alpha^2(\mathbb{B}_n))} \\ &\quad + \sqrt{2} \| \widehat{T}_{|a|\infty d\sigma} \|_{\mathcal{L}(A_\alpha^2(\mathbb{B}_n))}, \end{aligned}$$

where the three operators on the right-hand side have positive symbols. Therefore, it suffices to prove the proposition for $a \geq 0$, which in turn implies that $\widehat{T}_{ad\sigma}$ is self-adjoint and positive.

Let $a \geq 0$ and ℓ be a positive integer. Then

$$\| \widehat{T}_{ad\sigma} \|_{\mathcal{L}(A_\alpha^2(\mathbb{B}_n))}^\ell = \| (\widehat{T}_{ad\sigma})^\ell \|_{\mathcal{L}(A_\alpha^2(\mathbb{B}_n))} \leq \mathrm{Tr} \left((\widehat{T}_{ad\sigma})^\ell \right). \quad (28)$$

On the other hand, using the result for the trace of the power of a Toeplitz operator in Proposition 5.1,

$$\mathrm{Tr} \left((\widehat{T}_{ad\sigma})^\ell \right) = \frac{\left(\frac{\alpha}{\pi} \right)^{\frac{d}{2}}}{(2\ell)^{\frac{d'}{2}}} \int_{\Gamma} \left(\frac{a(\xi)}{(1-|\xi|^2)^{n+1}} \right)^\ell d\sigma(\xi) \left(1 + O \left(\frac{1}{\alpha} \right) \right)$$

$$\leq C \left[\sup_{\xi \in \Gamma} \frac{a(\xi)}{(1 - |\xi|^2)^{n+1}} \right]^\ell \frac{\alpha^{\frac{d}{2}}}{\ell^{\frac{d'}{2}}},$$

with the constant C not depending on α nor ℓ . Substitute this into (28) and take ℓ -th roots to get

$$\|\widehat{T}_{ad\sigma}\|_{\mathcal{L}(A_\alpha^2(\mathbb{B}_n))} \leq C^{\frac{1}{\ell}} \left[\sup_{\xi \in \Gamma} \frac{a(\xi)}{(1 - |\xi|^2)^{n+1}} \right]^{\frac{d}{2\ell}} \frac{\alpha^{\frac{d}{2\ell}}}{\ell^{\frac{d'}{2\ell}}} \leq C \left(\alpha^{\frac{1}{\ell}} \right)^{\frac{d}{2}}, \quad (29)$$

again with the last constant C not depending on α nor ℓ , and where we have used that $\left(\frac{1}{\ell}\right)^{\frac{1}{\ell}} \leq 1$ for all ℓ . This is valid for ℓ any positive integer, particularly for any $\ell > \alpha$. Then

$$\|\widehat{T}_{ad\sigma}\|_{\mathcal{L}(A_\alpha^2(\mathbb{B}_n))} \leq C \inf_{\ell \in \mathbb{N}} \left(\alpha^{\frac{1}{\ell}} \right)^{\frac{d}{2}} \leq C \left(\alpha^{\frac{1}{\alpha}} \right)^{\frac{d}{2}} \leq C.$$

Here we have used that $\lim_{\alpha \rightarrow \infty} \alpha^{\frac{1}{\alpha}} = 1$. Since the constant C is independent of α , the statement is proved. \square

Remark 5.3. From (29) and the arguments given thereafter we find that if $a > 0$, $\sup_{\xi \in \Gamma} \frac{a(\xi)}{(1 - |\xi|^2)^{n+1}}$ is a bound for both the operator norm of $\widehat{T}_{ad\sigma}$ and its largest eigenvalue.

We rely on the operator \mathcal{Q}_ε to extend the result in Proposition 5.1, first to encompass polynomials. Recall that \mathcal{Q}_ε is defined by

$$\mathcal{Q}_\varepsilon(\varphi)(t) = \frac{1}{\Gamma(\varepsilon)} \int_0^t \varphi(s) \left(\ln \frac{t}{s} \right)^{\varepsilon-1} \frac{ds}{s}$$

when $\varepsilon > 0$ and the identity when $\varepsilon = 0$.

Remark 5.4. We summarize some important features of this operator.

- \mathcal{Q}_ε is linear in the first entry.
- For $p > 0$, $\mathcal{Q}_\varepsilon(s^p)(t) = \frac{t^p}{p^\varepsilon}$. This follows from the change of variables $x = p \ln \left(\frac{t}{s}\right)$. In particular $\mathcal{Q}_{\frac{d'}{2}}(s^m)(t) = \frac{t^m}{m^{\frac{d'}{2}}}$. This remains valid even for $t \in \mathbb{C}$. In this case, the integration domain is taken to be the open line segment joining 0 and t .

- Equation (27) in Proposition 5.1 can be rewritten as

$$\mathrm{Tr} \left(\left(\widehat{T}_{ad\sigma} \right)^m \right) = \frac{\left(\frac{\alpha}{\pi} \right)^{\frac{d}{2}}}{2^{\frac{d'}{2}}} \int_{\Gamma} \mathcal{Q}_{\frac{d'}{2}}(s^m) \left(a(\xi)(1 - |\xi|^2)^{-(n+1)} \right) d\sigma(\xi) \left(1 + O(\alpha^{-1}) \right) . \quad (30)$$

We now state the result for the polynomial case.

Proposition 5.5. [Szegö limit theorem for polynomials] *Let Γ be isotropic or co-isotropic, $a \in C_0^2(\Gamma)$, and P be a polynomial without constant term. Then*

$$\mathrm{Tr} \left(P \left(\widehat{T}_{ad\sigma} \right) \right) = \frac{\left(\frac{\alpha}{\pi} \right)^{\frac{d}{2}}}{2^{\frac{d'}{2}}} \int_{\Gamma} \mathcal{Q}_{\frac{d'}{2}}(P) \left(\frac{a(\xi)}{(1 - |\xi|^2)^{n+1}} \right) d\sigma(\xi) \left(1 + O\left(\frac{1}{\alpha} \right) \right) .$$

Proof. The proof follows from (30) and the linearity of both the trace and $\mathcal{Q}_{\varepsilon}$. \square

5.2 The Szegö limit theorem

We now wish to extend our result to apply to more general continuous functions. From now on, we will consider $a > 0$. Fix $R > 0$ so that

$$\sup_{\xi \in \Gamma} \frac{a(\xi)}{(1 - |\xi|^2)^{n+1}} \leq R .$$

From Remark 5.3 we infer that

$$\bigcup_{\alpha > 0} \sigma \left(\widehat{T}_{ad\sigma} \right) \subseteq [0, R] .$$

This is important as $\varphi(\widehat{T}_{ad\sigma})$ is only well defined if φ is continuous on the spectrum of $\widehat{T}_{ad\sigma}$. Because we seek to analyze the asymptotic behavior of $\varphi(\widehat{T}_{ad\sigma})$ when α grows, we require φ to be continuous on the spectrum of $\widehat{T}_{ad\sigma}$ for all α .

We define the family of functions for which we will extend the past results.

Definition 5.6.

$$\mathcal{C} = \left\{ \varphi \in C[0, R] : \exists p > 0, \frac{\varphi(t)}{t^p} \in C[0, R] \right\} .$$

Remark 5.7. Without loss of generality, we can restrict the definition to consider only $p \in (0, 1)$.

Proposition 5.8. *The family \mathcal{C} satisfies:*

- a) $\mathcal{C} \supset \{P : P \text{ is a polynomial without constant term}\}$.
- b) $\varphi\left(\widehat{T}_{ad\sigma}\right)$ is trace class for every $\varphi \in \mathcal{C}$.
- c) $\mathcal{Q}_\varepsilon(\varphi)(t)$ is well defined $\forall \varepsilon \geq 0, t \in [0, R]$ and $\varphi \in \mathcal{C}$. Moreover, if $p > 0$ such that $\frac{\varphi(s)}{s^p} \in C[0, R]$, then

$$|\mathcal{Q}_\varepsilon(\varphi)(t)| \leq \left\| \frac{\varphi}{s^p} \right\|_{L^\infty[0, R]} \frac{t^p}{p^\varepsilon}.$$

- d) For every $\varphi \in \mathcal{C}$

$$\int_{\Gamma} \mathcal{Q}_{\frac{d'}{2}}(\varphi) \left(\frac{a(\xi)}{(1 - |\xi|^2)^{n+1}} \right) d\sigma(\xi) < \infty.$$

Proof. a) is obvious. For b) let λ_j be the eigenvalues of $\widehat{T}_{ad\sigma}$. Recall that as a is taken to be positive, so is $\widehat{T}_{ad\sigma}$. Then for $\varphi \in \mathcal{C}$ the norm of $\varphi\left(\widehat{T}_{ad\sigma}\right)$ in the trace class $S_1(A_\alpha^2(\mathbb{B}_n))$ satisfies

$$\begin{aligned} \left\| \varphi\left(\widehat{T}_{ad\sigma}\right) \right\|_{S_1(A_\alpha^2(\mathbb{B}_n))} &= \sum_{j=1}^{\infty} |\varphi(\lambda_j)| \\ &= \sum_{j=1}^{\infty} \frac{|\varphi(\lambda_j)|}{|\lambda_j|^p} |\lambda_j|^p \\ &\leq \sup_{t \in [0, R]} \left| \frac{\varphi(t)}{t^p} \right| \sum_{j=1}^{\infty} |\lambda_j|^p \\ &= \left\| \frac{\varphi}{s^p} \right\|_{L^\infty[0, R]} \left\| \widehat{T}_{ad\sigma} \right\|_{S_p(A_\alpha^2(\mathbb{B}_n))}^p, \end{aligned}$$

which is finite because $\widehat{T}_{ad\sigma}$ belongs to the Schatten class $S_p(A_\alpha^2(\mathbb{B}_n))$. For c) we compute

$$|\mathcal{Q}_\varepsilon(\varphi)(t)| = \left| \frac{1}{\Gamma(\varepsilon)} \int_0^t \varphi(s) \left(\ln \frac{t}{s} \right)^{\varepsilon-1} \frac{ds}{s} \right|$$

$$\begin{aligned}
&\leq \frac{1}{\Gamma(\varepsilon)} \int_0^t \left| \frac{\varphi(s)}{s^p} \right| s^p \left(\ln \frac{t}{s} \right)^{\varepsilon-1} \frac{ds}{s} \\
&\leq \frac{\left\| \frac{\varphi}{s^p} \right\|_{L^\infty[0,R]}}{\Gamma(\varepsilon)} \int_0^t s^p \left(\ln \frac{t}{s} \right)^{\varepsilon-1} \frac{ds}{s} = \left\| \frac{\varphi}{s^p} \right\|_{L^\infty[0,R]} \frac{t^p}{p^\varepsilon}.
\end{aligned}$$

Finally, for d) we use c) to get

$$\begin{aligned}
\left| \int_{\Gamma} \mathcal{Q}_{\frac{d'}{2}}(\varphi) \left(\frac{a(\xi)}{(1-|\xi|^2)^{n+1}} \right) d\sigma(\xi) \right| &\leq \int_{\Gamma} \left| \mathcal{Q}_{\frac{d'}{2}}(\varphi) \left(\frac{a(\xi)}{(1-|\xi|^2)^{n+1}} \right) \right| d\sigma(\xi) \\
&\leq \left\| \frac{\varphi}{s^p} \right\|_{L^\infty[0,R]} \frac{1}{p^{\frac{d'}{2}}} \int_{\Gamma} \left(\frac{a(\xi)}{1-|\xi|^{n+1}} \right)^p d\sigma(\xi) \\
&\leq \left\| \frac{\varphi}{s^p} \right\|_{L^\infty[0,R]} \frac{\sigma(\text{supp}(a))}{p^{\frac{d'}{2}}} \left(\sup_{\xi \in \Gamma} \frac{a(\xi)}{1-|\xi|^{n+1}} \right)^p < \infty,
\end{aligned} \tag{31}$$

where $\sigma(\text{supp}(a)) = \int_{\text{supp}(a)} d\sigma(\xi)$. \square

For the proof of the Szegő limit theorem we will use the following two results. The proof of the first one relies strongly on the hyperbolic geometry of \mathbb{B}_n and is left for Appendix A.

Lemma 5.9. *Let $\Gamma \subset \mathbb{B}_n$ be a submanifold and $a \in C_0^2(\Gamma)$ a positive function. For $p \in (0, 1)$*

$$\left(\frac{\pi}{\alpha} \right)^{\frac{d}{2}} \text{Tr} \left(\left(\widehat{T}_{ad\sigma} \right)^p \right)$$

is bounded as $\alpha \rightarrow \infty$.

Lemma 5.10. *Let Γ be isotropic or co-isotropic, $a \in C_0^2(\Gamma)$ a positive function, and P a polynomial without constant term. For $p \in (0, 1)$,*

$$\begin{aligned}
\lim_{\alpha \rightarrow \infty} \left(\frac{\pi}{\alpha} \right)^{\frac{d}{2}} \text{Tr} \left[\left(\widehat{T}_{ad\sigma} \right)^p P \left(\widehat{T}_{ad\sigma} \right) \right] &= \\
&= \frac{1}{2^{\frac{d'}{2}}} \int_{\Gamma} \mathcal{Q}_{\frac{d'}{2}}(s^p P(s)) \left(\frac{a(\xi)}{(1-|\xi|^2)^{n+1}} \right) d\sigma(\xi).
\end{aligned}$$

Proof. (Proof of Lemma 5.10)

In order to ease the notation, we will write

$$\mathring{Q}(\varphi) := \frac{1}{2^{\frac{d'}{2}}} \int_{\Gamma} \mathcal{Q}_{\frac{d'}{2}}(\varphi) \left(\frac{a(\xi)}{(1-|\xi|^2)^{n+1}} \right) d\sigma(\xi).$$

and $S := \widehat{T}_{ad\sigma}$. Let $\epsilon > 0$ and $\{g_j\}_{j \in \mathbb{N}}$ be a sequence of polynomials without constant terms that converges uniformly to the function s^p in $[0, R]$. For each $j \in \mathbb{N}$ we have

$$\begin{aligned} & \left| \left(\frac{\pi}{\alpha} \right)^{\frac{d}{2}} \text{Tr}[(S^p)P(S)] - \mathring{Q}(s^p P(s)) \right| \leq \\ & \left(\frac{\pi}{\alpha} \right)^{\frac{d}{2}} \left| \text{Tr}[(S^p - g_j(S))P(S)] \right| + \left| \left(\frac{\pi}{\alpha} \right)^{\frac{d}{2}} \text{Tr}[g_j(S)P(S)] - \mathring{Q}(g_j P) \right| \\ & + \left| \mathring{Q}((g_j - s^p)P) \right| =: \mathring{I}_1 + \mathring{I}_2 + \mathring{I}_3. \end{aligned} \quad (32)$$

We first bound \mathring{I}_1 . Let \tilde{P} be the polynomial whose coefficients are the absolute values of the coefficients of P . In this way $|P(s)| \leq \tilde{P}(s)$ for all $s \in [0, R]$ and hence $(\tilde{P} - |P|)(S)$ is a positive operator, which in turn implies $\text{Tr}(|P(S)|) \leq \text{Tr}(\tilde{P}(S))$. As \tilde{P} is a polynomial without constant term, applying Proposition 5.5 we can find a constant C such that $\left(\frac{\pi}{\alpha} \right)^{\frac{d}{2}} \text{Tr}(\tilde{P}(S)) < C$. We can now get

$$\begin{aligned} \mathring{I}_1 &= \left(\frac{\pi}{\alpha} \right)^{\frac{d}{2}} \left| \text{Tr}[(S^p - g_j(S))P(S)] \right| \\ &\leq \left(\frac{\pi}{\alpha} \right)^{\frac{d}{2}} \|P(S)\|_{S_1(A_\alpha^2(\mathbb{B}_n))} \|S^p - g_j(S)\|_{\mathcal{L}(A_\alpha^2(\mathbb{B}_n))} \\ &= \left(\frac{\pi}{\alpha} \right)^{\frac{d}{2}} \text{Tr}(|P(S)|) \|s^p - g_j\|_{L^\infty[0, R]} \\ &\leq \left(\frac{\pi}{\alpha} \right)^{\frac{d}{2}} \text{Tr}(\tilde{P}(S)) \|s^p - g_j\|_{L^\infty[0, R]} \leq C \|s^p - g_j\|_{L^\infty[0, R]}. \end{aligned} \quad (33)$$

To estimate \mathring{I}_3 we first notice that

$$\left\| \frac{(g_j - s^p)P}{s^p} \right\|_{L^\infty[0, R]} \leq \|g_j - s^p\|_{L^\infty[0, R]} \left\| \frac{P}{s^p} \right\|_{L^\infty[0, R]},$$

with $\left\| \frac{P}{s^p} \right\|_{L^\infty[0,R]}$ bounded as P has no constant term and $p \in (0, 1)$. We then apply inequality (31) from d) in Proposition 5.8 to get

$$\begin{aligned} \mathring{I}_3 &\leq \frac{\sigma(\text{supp}(a))}{p^{\frac{d}{2}}} \left(\sup_{\xi \in \Gamma} \frac{a(\xi)}{1 - |\xi|^{n+1}} \right)^p \left\| \frac{(g_j - s^p)P}{s^p} \right\|_{L^\infty[0,R]} \\ &\leq C \|g_j - s^p\|_{L^\infty[0,R]} . \end{aligned} \quad (34)$$

Both (33) and (34) are valid for all $j \in \mathbb{N}$, and the constants involved in each of them do not depend on j or α . Since $\|g_j - s^p\|_{L^\infty[0,R]}$ tends to zero, we can choose $j^* \in \mathbb{N}$ large enough so that $\mathring{I}_1 < \frac{\epsilon}{3}$ and $\mathring{I}_3 < \frac{\epsilon}{3}$. Finally, for this fixed j^* , $g_{j^*}P$ is a polynomial without constant term, so by Proposition 5.5

$$\mathring{I}_2 = \left| \left(\frac{\pi}{\alpha} \right)^{\frac{d}{2}} \text{Tr}[g_{j^*}(S)P(S)] - \mathring{Q}(g_{j^*}P) \right| < \frac{\epsilon}{3}$$

for all α large enough, which together with (32) proves the claim. \square

We now give a proof of the main result of the paper.

Proof of Theorem 1.8 [Szegő limit theorem]. Because $\varphi \in \mathcal{C}$, $\varphi(0) = 0$ and we can find $p \in (0, 1)$ such that $\frac{\varphi(s)}{s^p} \in C[0, R]$ and $\frac{\varphi(s)}{s^p} = 0$ at $s = 0$. Let $\{p_j\}_{j \in \mathbb{N}}$ be a sequence of polynomials without constant terms that converges uniformly to $\frac{\varphi(s)}{s^p}$ in $[0, R]$. This in turn implies that $s^p p_j$ converges uniformly to φ in $[0, R]$. We use once more the notation for $\mathring{Q}(\varphi)$ and S that we defined in the proof of Lemma 5.10 and follow a similar approach. Let $\epsilon > 0$. For each $j \in \mathbb{N}$ we have

$$\begin{aligned} &\left| \left(\frac{\pi}{\alpha} \right)^{\frac{d}{2}} \text{Tr}(\varphi(S)) - \mathring{Q}(\varphi) \right| \leq \\ &\quad \left(\frac{\pi}{\alpha} \right)^{\frac{d}{2}} \left| \text{Tr}(\varphi(S) - S^p p_j(S)) \right| + \left| \left(\frac{\pi}{\alpha} \right)^{\frac{d}{2}} \text{Tr}(S^p p_j(S)) - \mathring{Q}(s^p p_j) \right| \\ &\quad + \left| \mathring{Q}(s^p p_j - \varphi) \right| =: \mathring{I}_4 + \mathring{I}_5 + \mathring{I}_6 . \end{aligned} \quad (35)$$

For \mathring{I}_4 :

$$\mathring{I}_4 = \left(\frac{\pi}{\alpha} \right)^{\frac{d}{2}} \left| \text{Tr}((\varphi - s^p p_j)(S)) \right|$$

$$\begin{aligned}
&= \left(\frac{\pi}{\alpha}\right)^{\frac{d}{2}} \left| \text{Tr} \left(S^p \left(\frac{\varphi}{s^p} - p_j \right) (S) \right) \right| \\
&\leq \left(\frac{\pi}{\alpha}\right)^{\frac{d}{2}} \|S^p\|_{S_1(A_\alpha^2(\mathbb{B}_n))} \left\| \left(\frac{\varphi}{s^p} - p_j \right) (S) \right\|_{\mathcal{L}(A_\alpha^2(\mathbb{B}_n))} \\
&= \left(\frac{\pi}{\alpha}\right)^{\frac{d}{2}} \text{Tr}(S^p) \left\| \frac{\varphi}{s^p} - p_j \right\|_{L^\infty[0,R]} \leq C \left\| \frac{\varphi}{s^p} - p_j \right\|_{L^\infty[0,R]}, \quad (36)
\end{aligned}$$

where the last inequality is obtained applying Lemma 5.9.

For \mathring{I}_6 , we employ inequality (31) from d) in Proposition 5.8 to obtain

$$\mathring{I}_6 \leq C \left\| \frac{s^p p_j - \varphi}{s^p} \right\|_{L^\infty[0,R]} = C \left\| \frac{\varphi}{s^p} - p_j \right\|_{L^\infty[0,R]}. \quad (37)$$

Both (36) and (37) hold for any j and are independent of α , so we can pick $j^* \in \mathbb{N}$ such that $\mathring{I}_4 < \frac{\epsilon}{3}$ and $\mathring{I}_6 < \frac{\epsilon}{3}$ for all $\alpha > 0$. Finally, for j^* fixed, Lemma 5.10 asserts $\mathring{I}_5 < \frac{\epsilon}{3}$ for all α large enough. Applying all these inequalities in (35) concludes the proof of the theorem. \square

6 Corollaries and follow up results

In this section we present some results that follow as applications of the Szegő limit theorem 1.8. In particular, we obtain results describing the distribution of the eigenvalues and the Schatten norms of $\widehat{T}_{ad\sigma}$.

6.1 Distribution of eigenvalues

We begin by stating the equality in Theorem 1.8 in an alternate form. Addressing the definition of \mathcal{Q} , the right hand side of (7) equals

$$\frac{1}{2^{\frac{d'}{2}} \Gamma(\frac{d'}{2})} \int_{\Gamma} \int_0^{\frac{a(\xi)}{(1-|\xi|^2)^{n+1}}} \varphi(s) \left(\ln \frac{a(\xi)}{s(1-|\xi|^2)^{n+1}} \right)^{\frac{d'}{2}-1} \frac{ds}{s} d\sigma(\xi),$$

and interchanging the integrals order by Fubini's theorem, it can be rewritten as

$$\begin{aligned}
&\frac{1}{2^{\frac{d'}{2}}} \int_0^{\max_{\xi \in \Gamma} \frac{a(\xi)}{(1-|\xi|^2)^{n+1}}} \varphi(s) \\
&\cdot \left[\frac{1}{\Gamma(\frac{d'}{2}) s} \int_{\left\{ \xi \in \Gamma : \frac{a(\xi)}{(1-|\xi|^2)^{n+1}} \geq s \right\}} \left(\ln \frac{a(\xi)}{s(1-|\xi|^2)^{n+1}} \right)^{\frac{d'}{2}-1} d\sigma(\xi) \right] ds.
\end{aligned}$$

If we define $\mathcal{D}_a(s)$ to be the function defined by what is inside the brackets in the last expression, then (7) can be restated as

$$\lim_{\alpha \rightarrow \infty} \left(\frac{\pi}{\alpha} \right)^{\frac{d}{2}} \text{Tr} \left(\varphi \left(\widehat{T}_{ad\sigma} \right) \right) = \frac{1}{2^{\frac{d'}{2}}} \int_0^{\max_{\xi \in \Gamma} \frac{\alpha(\xi)}{(1-|\xi|^2)^{n+1}}} \varphi(s) \mathcal{D}_a(s) \, ds . \quad (38)$$

Notice that by doing this, we are now interpreting the asymptotics given in the Szegő limit theorem as an integral with respect to a measure $\mathcal{D}_a(s) \, ds$ that is absolutely continuous with respect to the Lebesgue measure in \mathbb{R} .

Corollary 6.1. *Let $I \subset (0, \max_{\xi \in \Gamma} \frac{\alpha(\xi)}{(1-|\xi|^2)^{n+1}}]$ be a closed interval and denote by $\mathcal{N}_I(\widehat{T}_{ad\sigma})$ the number of eigenvalues of $\widehat{T}_{ad\sigma}$ in I . Then*

$$\lim_{\alpha \rightarrow \infty} \left(\frac{\pi}{\alpha} \right)^{\frac{d}{2}} \mathcal{N}_I(\widehat{T}_{ad\sigma}) = \frac{1}{2^{\frac{d'}{2}}} \int_I \mathcal{D}_a(s) \, ds .$$

Proof. Let \mathbb{I}_I be the indicator function of I , which is continuous on each spectrum $\sigma(\widehat{T}_{ad\sigma})$, $\alpha > 0$. If $\{\lambda_j\}_{j \in \mathbb{N}}$ are the singular values of $\widehat{T}_{ad\sigma}$ we notice that

$$\mathcal{N}_I(\widehat{T}_{ad\sigma}) = \sum_{\{j : \lambda_j \in I\}} 1 = \sum_{j=1}^{\infty} \mathbb{I}_I(\lambda_j) = \text{Tr} \left(\mathbb{I}_I \left(\widehat{T}_{ad\sigma} \right) \right) .$$

Let $I = [c, d]$ and consider the two sequences of trapezoidal functions $\{\psi_j\}_{j \in \mathbb{N}}$, $\{\phi_j\}_{j \in \mathbb{N}}$ defined by

$$\psi_j(x) = \begin{cases} 0 & x \in [0, c] \\ j(x-c) & x \in (c, c + \frac{1}{j}) \\ 1 & x \in [c + \frac{1}{j}, d - \frac{1}{j}] \\ -j(x-d) & x \in (d - \frac{1}{j}, d) \\ 0 & x \in [d, \infty) \end{cases} ,$$

$$\phi_j(x) = \begin{cases} 0 & x \in [0, c - \frac{1}{j}] \\ j(x-c) + 1 & x \in (c - \frac{1}{j}, c) \\ 1 & x \in [c, d] \\ -j(x-d) + 1 & x \in (d, d + \frac{1}{j}) \\ 0 & x \in [d + \frac{1}{j}, \infty) \end{cases} .$$

These functions satisfy

$$\lim_{j \rightarrow \infty} \psi_j(x) = \mathbb{I}_{(c,d)}(x) \quad , \quad \lim_{j \rightarrow \infty} \phi_j(x) = \mathbb{I}_I(x)$$

for all $x > 0$ and $\psi_j(x) \leq \mathbb{I}_I(x) \leq \phi_j(x)$ for all $j \in \mathbb{N}$ and $x > 0$. It follows that

$$\mathrm{Tr} \left(\psi_j(\widehat{T}_{ad\sigma}) \right) \leq \mathcal{N}_I(\widehat{T}_{ad\sigma}) \leq \mathrm{Tr} \left(\phi_j(\widehat{T}_{ad\sigma}) \right) ,$$

and by (38)

$$\begin{aligned} \int_0^{\max_{\xi \in \Gamma} \frac{a(\xi)}{(1-|\xi|^2)^{n+1}}} \psi_j(s) \mathcal{D}_a(s) \, ds &\leq \frac{1}{2^{\frac{d'}{2}}} \lim_{\alpha \rightarrow \infty} \left(\frac{\pi}{\alpha} \right)^{\frac{d'}{2}} \mathcal{N}_I(\widehat{T}_{ad\sigma}) \\ &\leq \int_0^{\max_{\xi \in \Gamma} \frac{a(\xi)}{(1-|\xi|^2)^{n+1}}} \phi_j(s) \mathcal{D}_a(s) \, ds \end{aligned} \quad (39)$$

for all $j \in \mathbb{N}$. Now note that $|\psi_j(x)| \leq |\phi_j(x)| \leq \phi_1(x)$ for all $x > 0$, $j \in \mathbb{N}$ and $\int_0^{\max_{\xi \in \Gamma} \frac{a(\xi)}{(1-|\xi|^2)^{n+1}}} \phi_1(s) \mathcal{D}_a(s) \, ds < \infty$. Then by the dominated convergence theorem

$$\lim_{j \rightarrow \infty} \int_0^{\max_{\xi \in \Gamma} \frac{a(\xi)}{(1-|\xi|^2)^{n+1}}} \psi_j(s) \mathcal{D}_a(s) \, ds = \int_0^{\max_{\xi \in \Gamma} \frac{a(\xi)}{(1-|\xi|^2)^{n+1}}} \mathbb{I}_{(c,d)}(x) \mathcal{D}_a(s) \, ds$$

and

$$\lim_{j \rightarrow \infty} \int_0^{\max_{\xi \in \Gamma} \frac{a(\xi)}{(1-|\xi|^2)^{n+1}}} \phi_j(s) \mathcal{D}_a(s) \, ds = \int_0^{\max_{\xi \in \Gamma} \frac{a(\xi)}{(1-|\xi|^2)^{n+1}}} \mathbb{I}_{[c,d]}(x) \mathcal{D}_a(s) \, ds ,$$

both of them being equal to $\int_I \mathcal{D}_a(s) \, ds$, which together with (39) concludes the proof. \square

6.2 Schatten norm estimates

Let $p > 0$. Recall that for A a compact operator in $A_\alpha^2(\mathbb{B}_n)$, A is a member of the Schatten class $S_p(A_\alpha^2(\mathbb{B}_n))$ if and only if $|A|^p$ is of trace class. This is equivalent to having the quantity

$$\|A\|_{S_p(A_\alpha^2(\mathbb{B}_n))} := \mathrm{Tr}((A^*A)^{\frac{p}{2}})^{\frac{1}{p}}$$

being finite. For $p \geq 1$, $\|\cdot\|_{S_p(A_\alpha^2(\mathbb{B}_n))}$ is a norm in $S_p(A_\alpha^2(\mathbb{B}_n))$, while when $0 < p < 1$ it is a quasi-norm satisfying

$$\|A + B\|_{S_p(A_\alpha^2(\mathbb{B}_n))}^p \leq 2 \left(\|A\|_{S_p(A_\alpha^2(\mathbb{B}_n))}^p + \|B\|_{S_p(A_\alpha^2(\mathbb{B}_n))}^p \right) .$$

Let $a \in C_0^2(\Gamma)$. If $\widehat{T}_{ad\sigma}$ is a Toeplitz operator, it can be expressed as the sum of three Toeplitz operators with positive symbols (see the beginning of

the proof of 5.2). Then, as $\|\cdot\|_p$ is at least a quasi-norm, we will have by Lemma 5.9 that

$$\left(\frac{\pi}{\alpha}\right)^{\frac{d}{2}} \operatorname{Tr}((\widehat{T}_{ad\sigma}^* \widehat{T}_{ad\sigma})^{\frac{p}{2}})$$

is bounded when $\alpha \rightarrow \infty$.

Lemma 6.2. *Let Γ be isotropic or co-isotropic and $a \in C_0^2(\Gamma)$. For any $p > 0$,*

$$\lim_{\alpha \rightarrow \infty} \left(\frac{\pi}{\alpha}\right)^{\frac{d}{2}} \operatorname{Tr}((\widehat{T}_{ad\sigma}^* \widehat{T}_{ad\sigma})^{\frac{p}{2}}) = \frac{1}{(2p)^{\frac{d'}{2}}} \int_{\Gamma} \left(\frac{|a(\xi)|}{(1-|\xi|^2)^{n+1}}\right)^p d\sigma.$$

Proof. For any Toeplitz operator, $\widehat{T}_{ad\sigma}^* = \widehat{T}_{\bar{a}d\sigma}$ and so we have for $m \in \mathbb{N}$

$$\begin{aligned} \operatorname{Tr}((\widehat{T}_{ad\sigma}^* \widehat{T}_{ad\sigma})^m) &= \frac{\left(\frac{\alpha}{\pi}\right)^{\frac{d}{2}}}{(4m)^{\frac{d'}{2}}} \int_{\Gamma} \frac{|a(\xi)|^{2m}}{(1-|\xi|^2)^{2m(n+1)}} d\sigma(\xi) \left(1 + O\left(\frac{1}{\alpha}\right)\right) \\ &= \frac{\left(\frac{\alpha}{\pi}\right)^{\frac{d}{2}}}{2^{d'}} \int_{\Gamma} \mathcal{Q}_{\frac{d'}{2}}(s^m) \left(\frac{|a(\xi)|^2}{(1-|\xi|^2)^{2(n+1)}}\right) d\sigma(\xi) \left(1 + O\left(\frac{1}{\alpha}\right)\right). \end{aligned}$$

Make $A := \widehat{T}_{ad\sigma}^* \widehat{T}_{ad\sigma}$. Using the same arguments as in Corollary 5.2 and the discussion at the beginning of Subsection 5.2 we can find a constant R such that $\sigma(A) \subseteq [0, R^2]$ for all $\alpha > 0$.

By the linearity of $\mathcal{Q}_{\frac{d'}{2}}$ we get

$$\lim_{\alpha \rightarrow \infty} \left(\frac{\pi}{\alpha}\right)^{\frac{d}{2}} \operatorname{Tr}(P(A)) = \frac{1}{2^{d'}} \int_{\Gamma} \mathcal{Q}_{\frac{d'}{2}}(P) \left(\frac{|a(\xi)|^2}{(1-|\xi|^2)^{2(n+1)}}\right) d\sigma(\xi) \quad (40)$$

for any P polynomial without constant term. Let $\epsilon > 0$. We compute

$$\begin{aligned} &\left| \left(\frac{\pi}{\alpha}\right)^{\frac{d}{2}} \operatorname{Tr}(A^{\frac{p}{2}}) - \frac{1}{(2p)^{\frac{d'}{2}}} \int_{\Gamma} \left(\frac{|a(\xi)|}{(1-|\xi|^2)^{n+1}}\right)^p d\sigma \right| \leq \left(\frac{\pi}{\alpha}\right)^{\frac{d}{2}} \left| \operatorname{Tr}(A^{\frac{p}{2}} - P(A)) \right| \\ &+ \left| \left(\frac{\pi}{\alpha}\right)^{\frac{d}{2}} \operatorname{Tr}(P(A)) - \frac{1}{2^{d'}} \int_{\Gamma} \mathcal{Q}_{\frac{d'}{2}}(P) \left(\frac{|a(\xi)|^2}{(1-|\xi|^2)^{2(n+1)}}\right) d\sigma \right| \\ &+ \frac{1}{2^{d'}} \left| \int_{\Gamma} \mathcal{Q}_{\frac{d'}{2}}(s^{\frac{p}{2}} - P) \left(\frac{|a(\xi)|^2}{(1-|\xi|^2)^{2(n+1)}}\right) d\sigma \right| =: \mathring{I}_7 + \mathring{I}_8 + \mathring{I}_9 \end{aligned} \quad (41)$$

for any such polynomial. We will consider a P of the form $P = P_1 P_2$ with P_1 and P_2 two other polynomials without constant term. By doing this we have

$$|s^{\frac{p}{2}} - P| \leq |s^{\frac{p}{4}}| |s^{\frac{p}{4}} - P_1(s)| + |P_1(s)| |s^{\frac{p}{4}} - P_2(s)|.$$

Using this, we estimate \mathring{I}_7 by

$$\begin{aligned} \mathring{I}_7 &\leq \left(\frac{\pi}{\alpha}\right)^{\frac{d}{2}} \left| \text{Tr}(A^{\frac{p}{4}}(A^{\frac{p}{4}} - P_1(A))) \right| + \left(\frac{\pi}{\alpha}\right)^{\frac{d}{2}} \left| \text{Tr}(P_1(A)(A^{\frac{p}{4}} - P_2(A))) \right| \\ &\leq \left(\frac{\pi}{\alpha}\right)^{\frac{d}{2}} \text{Tr}(A^{\frac{p}{4}}) \|s^{\frac{p}{4}} - P_1\|_{L^\infty[0, R^2]} + \left(\frac{\pi}{\alpha}\right)^{\frac{d}{2}} \text{Tr}(P_1(A)) \|s^{\frac{p}{4}} - P_2\|_{L^\infty[0, R^2]}, \end{aligned}$$

and, applying $c)$ of Proposition 5.8, \mathring{I}_9 by

$$\begin{aligned} \mathring{I}_9 &\leq \frac{1}{2^{\frac{d'}{2}} \left(\frac{p}{4}\right)^{\frac{d'}{2}}} \left\| \frac{s^{\frac{p}{2}} - P}{s^{\frac{p}{4}}} \right\|_{L^\infty[0, R^2]} \int_{\Gamma} \left(\frac{|a(\xi)|}{(1 - |\xi|^2)^{n+1}} \right)^{\frac{p}{2}} d\sigma(\xi) \\ &\leq \left(\frac{2}{p}\right)^{\frac{d'}{2}} \int_{\Gamma} \left(\frac{|a(\xi)|}{(1 - |\xi|^2)^{n+1}} \right)^{\frac{p}{2}} d\sigma(\xi) \\ &\quad \cdot \left(\|s^{\frac{p}{4}} - P_1\|_{L^\infty[0, R^2]} + \|P_1 s^{-\frac{p}{4}}\|_{L^\infty[0, R^2]} \|s^{\frac{p}{4}} - P_2\|_{L^\infty[0, R^2]} \right) \end{aligned}$$

for all $\alpha > 0$. We can now choose P_1 such that

$$\|s^{\frac{p}{4}} - P_1\|_{L^\infty[0, R^2]} < \frac{\epsilon}{6} \min \left\{ \left(\sup_{\alpha > 0} \left(\frac{\pi}{\alpha}\right)^{\frac{d}{2}} \text{Tr}(A^{\frac{p}{4}}) \right)^{-1}, \left(\left(\frac{2}{p}\right)^{\frac{d'}{2}} \int_{\Gamma} \left(\frac{|a(\xi)|}{(1 - |\xi|^2)^{n+1}} \right)^{\frac{p}{2}} d\sigma(\xi) \right)^{-1} \right\}$$

and P_2 such that

$$\begin{aligned} \|s^{\frac{p}{4}} - P_2\|_{L^\infty[0, R^2]} &< \frac{\epsilon}{6} \min \left\{ \left(\sup_{\alpha > 0} \left(\frac{\pi}{\alpha}\right)^{\frac{d}{2}} \text{Tr}(P_1(A)) \right)^{-1}, \right. \\ &\quad \left. \left(\left(\frac{2}{p}\right)^{\frac{d'}{2}} \|s^{-\frac{p}{4}} P_1\|_{L^\infty[0, R^2]} \int_{\Gamma} \left(\frac{|a(\xi)|}{(1 - |\xi|^2)^{n+1}} \right)^{\frac{p}{2}} d\sigma(\xi) \right)^{-1} \right\}. \end{aligned}$$

This ensures that $\mathring{I}_7 < \frac{\epsilon}{3}$ and $\mathring{I}_9 < \frac{\epsilon}{3}$. Finally for these fixed P_1 and P_2 we know from (40) that we can take α large enough so $\mathring{I}_8 < \frac{\epsilon}{3}$, which together with (41) concludes the proof. \square

The desired result describing the Schatten norms follows as a consequence.

Corollary 6.3. *Let Γ be isotropic or co-isotropic and $a \in C_0^2(\Gamma)$. For any $p > 0$,*

$$\lim_{\alpha \rightarrow \infty} \left(\frac{\pi}{\alpha}\right)^{\frac{d}{2p}} \|\widehat{T}_{ad\sigma}\|_{S_p(A_\alpha^2(\mathbb{B}_n))} = \left(\frac{1}{(2p)^{\frac{d'}{2}}} \int_{\Gamma} \left(\frac{|a(\xi)|}{(1 - |\xi|^2)^{n+1}} \right)^p d\sigma(\xi) \right)^{\frac{1}{p}}.$$

6.3 An example

We illustrate some of the results given by presenting a concrete example.

For $r \in (0, 1)$ let $\Gamma = r\mathbb{S}^1 = \{z \in \mathbb{C} : |z| = r\}$. If we use the parametrization $\gamma : (0, 1) \rightarrow \Gamma$, $\gamma(\theta) = re^{2\pi i\theta}$, then $d\sigma = 2\pi \frac{r}{1-r^2} d\theta$. Notice that in this case $d = 1$ and $n = 1$, thus $d' = 1$. Let $a \equiv 1$ in Γ and $T_r := T_{1d\sigma}$ be the associated Toeplitz operator. Then

$$T_r f(z) = 2\pi r(1-r^2)^{\alpha-1}(\alpha+1) \int_0^1 K^\alpha(z, re^{2\pi i\theta}) f(re^{2\pi i\theta}) d\theta$$

for any $f \in A_\alpha^2(\mathbb{D})$.

Recall that $e_m(z) = \delta_m z^m$ with $\delta_m := \sqrt{\frac{\Gamma(m+\alpha+2)}{m! \Gamma(\alpha+2)}}$, $m \in \mathbb{N}_0$, is an orthonormal basis for $A_\alpha^2(\mathbb{D})$, and that the Bergman kernel can be expressed as an hypergeometric series via

$$K^\alpha(z, w) = \frac{1}{(1-z\bar{w})^{\alpha+2}} = \sum_{j=0}^{\infty} \frac{\Gamma(\alpha+j+2)}{\Gamma(\alpha+2) j!} (z\bar{w})^j .$$

Using this we compute

$$\begin{aligned} \frac{T_r e_m(z)}{2\pi r(1-r^2)^{-\alpha+1}} &= (\alpha+1)\delta_m \int_0^1 \left(\sum_{j=0}^{\infty} \frac{\Gamma(\alpha+j+2)}{\Gamma(\alpha+2) j!} z^j r^j e^{-2\pi j i\theta} \right) r^m e^{2\pi m i\theta} d\theta \\ &= (\alpha+1)\delta_m \sum_{j=0}^{\infty} \frac{\Gamma(\alpha+j+2)}{\Gamma(\alpha+2) j!} r^{m+j} z^j \int_0^1 e^{2\pi(m-j)i\theta} d\theta \\ &= (\alpha+1)r^{2m} \frac{\Gamma(\alpha+m+2)}{\Gamma(\alpha+2) m!} e_m(z) , \end{aligned}$$

from where it follows that the eigenvalues of $\widehat{T}_r = \frac{1}{\sqrt{2\pi\alpha}} T_r$ are

$$\lambda_m = \sqrt{\frac{2\pi}{\alpha}} (1-r^2)^{\alpha-1} \frac{\Gamma(\alpha+m+2)}{\Gamma(\alpha+1)m!} r^{2m+1} , \quad (42)$$

$m \in \mathbb{N}_0$.

6.3.1 Operator norm

We seek to analyze the behavior of the operator norm of \widehat{T}_r by estimating its largest eigenvalue. For any fixed α , we notice that the sequence of quotients

$$\frac{\lambda_m}{\lambda_{m-1}} = \frac{(\alpha+1)r^2}{m} + r^2$$

is monotonically decreasing to $r^2 < 1$. Thus, the largest eigenvalue of \widehat{T}_r for this α is λ_{m^*} , where m^* is the largest natural number such that $\frac{(\alpha+1)r^2}{m^*} + r^2 \geq 1$. This is obtained when $m^* = \left\lfloor (\alpha+1) \frac{r^2}{1-r^2} \right\rfloor$. Assume α is such that $(\alpha+1) \frac{r^2}{1-r^2} \in \mathbb{N}$. Then in terms of the Beta function

$$\frac{\Gamma(\alpha + m^* + 2)}{\Gamma(\alpha + 1)m^*!} = \mathbf{B} \left(\alpha + 1, (\alpha + 1) \frac{r^2}{1 - r^2} \right)^{-1} r^{-2}.$$

If α grows, so does $m^* = (\alpha + 1) \frac{r^2}{1-r^2}$, and we can derive that

$$\mathbf{B} \left(\alpha + 1, (\alpha + 1) \frac{r^2}{1 - r^2} \right) \sim \sqrt{\frac{2\pi}{\alpha + 1}} (1 - r^2)^{\alpha + 1} r^{2m^* - 1}$$

from the known asymptotic formula $\mathbf{B}(x, y) \sim \sqrt{2\pi} \frac{x^{x-\frac{1}{2}} y^{y-\frac{1}{2}}}{(x+y)^{x+y-\frac{1}{2}}}$ for large x and y , which is in turn an application of Stirling's formula. Substituting this in the expression for λ_{m^*} in (42) we get

$$\lambda_{m^*} \sim \sqrt{\frac{\alpha + 1}{\alpha}} \frac{1}{(1 - r^2)^2} \sim \frac{1}{(1 - r^2)^2}$$

when α grows. This shows that the normalization proposed in (6) and Corollary 5.2 is optimal in terms of the growth rate of α , while also exhibiting the sharpness of the bound given in Remark 5.3.

6.3.2 Eigenvalue distribution comparison

Keeping with \widehat{T}_r as defined above, given an interval $[t_1, t_2] \subset \left(0, \frac{1}{(1-r^2)^2}\right)$ Corollary 6.1 simplifies to

$$\lim_{\alpha \rightarrow \infty} \sqrt{\frac{\pi}{\alpha}} \mathcal{N}_{[t_1, t_2]}(\widehat{T}_r) = \frac{\sqrt{8\pi r}}{1 - r^2} \left[\sqrt{\ln \left(\frac{1}{(1 - r^2)^{n+1} t_1} \right)} - \sqrt{\ln \left(\frac{1}{(1 - r^2)^{n+1} t_2} \right)} \right], \quad (43)$$

which is also written as an asymptotic equivalence in the form

$$\mathcal{N}_{[t_1, t_2]}(\widehat{T}_r) \sim \frac{\sqrt{8\alpha r}}{1 - r^2} \left[\sqrt{\ln \left(\frac{1}{(1 - r^2)^{n+1} t_1} \right)} - \sqrt{\ln \left(\frac{1}{(1 - r^2)^{n+1} t_2} \right)} \right]. \quad (44)$$

To test this result, for fixed values of $r \in (0, 1)$ and $t_1 < t_2 \leq \frac{1}{(1-r^2)^2}$, and distinct values of α , we have numerically counted the number of eigenvalues of \widehat{T}_r in $[t_1, t_2]$ using (42). For a pair of selected values, the comparisons between these numerical outputs and the expected results given by (43) and (44) are presented in the next two tables.

α	$\mathcal{N}_{[\frac{16}{15}, \frac{16}{9}]}(\widehat{T}_{\frac{1}{2}})$	R.H.S. of (44)	$\sqrt{\frac{\pi}{\alpha}} \mathcal{N}_{[\frac{16}{15}, \frac{16}{9}]}(\widehat{T}_{\frac{1}{2}})$
100	14	13.47	2.4814
500	30	30.13	2.378
10^3	42	42.61	2.3541
$5 \cdot 10^3$	95	95.29	2.3813
10^4	134	134.76	2.3750
$5 \cdot 10^4$	301	301.35	2.3859
10^5	426	426.17	2.3877

Table 1: Comparison for $r = \frac{1}{2}$, $t_1 = \frac{16}{15}$, $t_2 = \frac{16}{9}$. The values in the second column were obtained by counting numerically for how many values of m the eigenvalue λ_m as given in (42) lies in $[\frac{16}{15}, \frac{16}{9}]$. As expected, the numbers on the rightmost column seem to be tending to the value of the R.H.S. of (43), approximately 2.3887 in this case.

α	$\mathcal{N}_{[\frac{2}{5}, \frac{3}{5}]}(\widehat{T}_{\frac{1}{\sqrt{2}}})$	R.H.S. of (44)	$\sqrt{\frac{\pi}{\alpha}} \mathcal{N}_{[\frac{2}{5}, \frac{3}{5}]}(\widehat{T}_{\frac{1}{\sqrt{2}}})$
100	5	5.60	0.8862
500	12	12.52	0.9511
10^3	18	17.71	1.0088
$5 \cdot 10^3$	39	39.61	0.9775
10^4	56	56.02	0.9925
$5 \cdot 10^4$	125	125.28	0.9908
10^5	177	177.17	0.9920

Table 2: Comparison for $r = \frac{1}{\sqrt{2}}$, $t_1 = \frac{2}{5}$, $t_2 = \frac{3}{5}$. The values in the second column were obtained by counting numerically for how many values of m the eigenvalue λ_m as given in (42) lies in $[\frac{2}{5}, \frac{3}{5}]$. As expected, the numbers on the rightmost column seem to be tending to the value of the R.H.S. of (43), approximately 0.9930 in this case.

A Proof of Lemma 5.9

Throughout this section a will be a positive symbol and $0 < p < 1$. Lemma 5.9 is equivalent to there being some constant $C > 0$ such that

$$\mathrm{Tr}(T_{ad\sigma}^p) \leq C\alpha^{p(n-\frac{d}{2})+\frac{d}{2}}, \quad (45)$$

for all α large enough. We will prove this inequality.

Let $\widetilde{T}_{ad\sigma}$ be the Berezin symbol of $T_{ad\sigma}$, namely

$$\widetilde{T}_{ad\sigma}(z) = \langle T_{ad\sigma}k_z, k_z \rangle .$$

For any Hilbert space H and any positive operator $A \in \mathcal{L}(H)$, the relationship $\langle A^p x, x \rangle \leq \langle Ax, x \rangle^p$ is true for all x unitary in H whenever $0 < p < 1$ (Proposition 1.31 [17]). Combining this with the definition of $\widetilde{T}_{ad\sigma}$ yields in

$$\widetilde{T_{ad\sigma}^p}(z) \leq \left(\widetilde{T}_{ad\sigma}(z) \right)^p$$

for all $z \in \mathbb{B}_n$. Using the formula for the trace of an operator in terms of its Berezin symbol, we get

$$\begin{aligned} \mathrm{Tr}(T_{ad\sigma}^p) &= c_\alpha \int_{\mathbb{B}_n} \widetilde{T_{ad\sigma}^p}(z) \, d\tau(z) \\ &\leq c_\alpha \int_{\mathbb{B}_n} \left(\widetilde{T}_{ad\sigma}(z) \right)^p \, d\tau(z) \\ &= c_\alpha^{1+p} \int_{\mathbb{B}_n} \left[\int_{\Gamma} \left(\frac{(1-|z|^2)(1-|w|^2)}{|1-\langle z, w \rangle|^2} \right)^{n+1+\alpha} \frac{a(w) d\sigma(w)}{(1-|w|^2)^{n+1}} \right]^p \, d\tau(z) \end{aligned} \quad (46)$$

where $d\tau(z) = \frac{dv(z)}{(1-|z|^2)^{n+1}}$ is the invariant measure on \mathbb{B}_n .

Remark A.1. If φ_z denotes the involutive automorphism in \mathbb{B}_n that interchanges z with the origin, then

$$\frac{(1-|z|^2)(1-|w|^2)}{|1-\langle z, w \rangle|^2} = 1 - |\varphi_z(w)|^2 .$$

Moreover $|\varphi_z(w)|$ is the pseudo-hyperbolic distance between z and w , while the hyperbolic distance is $\beta(z, w) = \tanh^{-1} |\varphi_z(w)|$. This then implies

$$\frac{(1-|z|^2)(1-|w|^2)}{|1-\langle z, w \rangle|^2} \leq 1 ,$$

with equality if and only if $\beta(z, w) = 0$, if and only if $z = w$ (see [18] Sections 1.2 and 1.5, particularly Lemma 1.2 and Proposition 1.21).

For $\epsilon > 0$ we now define

$$\mathcal{V} = \{z \in \mathbb{B}_n : \exists w \in \Gamma, \beta(z, w) < \epsilon\}$$

a hyperbolic tubular neighborhood of Γ in \mathbb{B}_n .

Proposition A.2. *The neighborhood \mathcal{V} satisfies*

$$c_\alpha^{1+p} \int_{\mathbb{B}_n \setminus \mathcal{V}} \left[\int_{\Gamma} \left(\frac{(1-|z|^2)(1-|w|^2)}{|1-\langle z, w \rangle|^2} \right)^{n+1+\alpha} \frac{a(w) d\sigma(w)}{(1-|w|^2)^{n+1}} \right]^p d\tau(z) = O(\alpha^{-\infty}). \quad (47)$$

Proof. Let $r(a) > 0$ be such that $\beta(w, 0) < r(a)$ for all $w \in \text{supp}(a)$. There exists some constant $C_{r(a)} > 0$ that only depends on $r(a)$ such that

$$C_{r(a)}^{-1} \leq \frac{1-|u|^2}{1-|v|^2} \leq C_{r(a)}$$

for any $u, v \in \mathbb{B}_n$ with $\beta(u, v) < r(a)$ (Lemma 2.20 in [18]). Let $\rho < 1$ be such that $C_{r(a)}(1-\rho^2) < 1/2$. It suffices to prove that

$$\mathcal{J}_1 := \int_{(\mathbb{B}_n \setminus \mathcal{V}) \cap \{z \in \mathbb{B}_n : |z| \geq \rho\}} \left[* \right]^p d\tau(z) \quad \text{and} \quad \mathcal{J}_2 := \int_{(\mathbb{B}_n \setminus \mathcal{V}) \cap \{z \in \mathbb{B}_n : |z| < \rho\}} \left[* \right]^p d\tau(z)$$

are both $O(\alpha^{-\infty})$, where $*$ represents what is inside the brackets in (47).

If $z \in \mathbb{B}_n$, $w \in \text{supp}(a)$, the invariance under automorphisms of the Bergman metric leads to $\beta(\varphi_z(w), z) = \beta(w, 0) < r(a)$, and so

$$\begin{aligned} \frac{(1-|z|^2)(1-|w|^2)}{|1-\langle z, w \rangle|^2} &= \frac{1-|\varphi_z(w)|^2}{1-|z|^2} (1-|z|^2) \\ &\leq C_{r(a)}(1-|z|^2). \end{aligned}$$

Using this we compute

$$\mathcal{J}_1 \leq C_{r(a)}^{(n+1+\alpha)p} \left[\int_{\Gamma} \frac{a(w) d\sigma(w)}{(1-|w|^2)^{n+1}} \right]^p \int_{\{z \in \mathbb{B}_n : |z| \geq \rho\}} (1-|z|^2)^{\alpha p - (n+1)(1-p)} dv(z)$$

$$\begin{aligned} &\leq C (C_{r(a)}(1 - \rho^2))^{\alpha p} \\ &< C \frac{1}{2^{\alpha p}} = O\left(\frac{1}{\alpha^\infty}\right) \end{aligned}$$

for all α large enough so that $\alpha p - (n + 1)(1 - p) > 0$.

If $z \notin \mathcal{V}$ then $\beta(z, w) \geq \epsilon$ for all $w \in \Gamma$ and hence

$$1 - |\varphi_z(w)|^2 = 1 - \tanh^2 \beta(z, w) \leq 1 - \tanh^2 \epsilon ,$$

which we use to conclude that

$$\begin{aligned} \mathcal{J}_2 &\leq (1 - \tanh^2 \epsilon)^{(n+1+\alpha)p} \left[\int_{\Gamma} \frac{a(w) d\sigma(w)}{(1 - |w|^2)^{n+1}} \right]^p \int_{\{z \in \mathbb{B}_n : |z| < \rho\}} d\tau(z) \\ &\leq C(1 - \tanh^2 \epsilon)^{\alpha p} = O\left(\frac{1}{\alpha^\infty}\right) . \end{aligned}$$

□

We take a moment to focus on the case $d = 2n$, so Γ is an open subset of \mathbb{B}_n and $T_{ad\sigma} = T_a$. Considering the result in last statement and using it together with the first inequality in (46) we can get

$$\mathrm{Tr}(T_a) \leq c_\alpha \int_{\delta \mathbb{B}_n} \left(\widetilde{T}_a(z)\right)^p d\tau(z) + O\left(\frac{1}{\alpha^\infty}\right) ,$$

where $\delta < 1$ is so that $\delta \mathbb{B}_n \supset \mathrm{supp}(a)$. In this case, the exponent of α in (45) simplifies to n , which is the growth order of $c_\alpha = \frac{\alpha^n}{n!}(1 + O(\alpha^{-1}))$.

Therefore, it will suffice to show that $\int_{\delta \mathbb{B}_n} \left(\widetilde{T}_a(z)\right)^p d\tau(z)$ remains bounded when α grows. For this we recall that the Berezin transform is a contraction from $\mathcal{L}(A_\alpha^2(\mathbb{B}_n))$ to $L^\infty(\mathbb{B}_n)$, so

$$\begin{aligned} \int_{\delta \mathbb{B}_n} \left(\widetilde{T}_a(z)\right)^p d\tau(z) &\leq \|\widetilde{T}_a\|_{L^\infty(\mathbb{B}_n)}^p \int_{\delta \mathbb{B}_n} d\tau(z) \\ &\leq C \|T_a\|_{\mathcal{L}(A_\alpha^2(\mathbb{B}_n))}^p . \end{aligned}$$

Since $d = 2n$ implies Γ being co-isotropic and $d' = 0$, $\|T_a\|_{\mathcal{L}(A_\alpha^2(\mathbb{B}_n))} = \frac{\pi^n}{n!} \|\widehat{T}_a\|_{\mathcal{L}(A_\alpha^2(\mathbb{B}_n))}$ is bounded for all $\alpha > 0$ by Corollary 5.2, thus proving the claim for $d = 2n$.

Let $d \leq 2n - 1$. From Proposition A.2, and rewriting the integral expression in (46) we conclude that proving our claim relies on bounding the integral

$$\mathcal{I}_p(\alpha) := c_\alpha^{1+p} \int_{\mathcal{V}} \left[\int_{\Gamma} e^{\alpha \ln(1-|\varphi_z(w)|^2)} \frac{a(w) \, d\sigma(w)}{|1 - \langle z, w \rangle|^{2(n+1)}} \right]^p \frac{dv(z)}{(1 - |z|^2)^{(n+1)(1-p)}}.$$

We now give a parametrization for \mathcal{V} . By means of using a partition of unity subordinated to an open cover of the support of a if necessary, we can assume without loss of generality that the support of a is contained on the image of a single parametrization $\gamma : B(0, 1) \rightarrow \Gamma$ of Γ . For each $p = \gamma(t) \in \Gamma$ let $\{E_1(t), \dots, E_{2n-d}(t)\}$ be an orthonormal basis for the normal space $D_{p^\iota}(\mathbb{T}_p\Gamma)^\perp \subset \mathbb{T}_p\mathbb{B}_n$ at p , so the ambient tangent space $\mathbb{T}_p\mathbb{B}_n$ splits as a direct sum

$$\mathbb{T}_p\mathbb{B}_n = D_{p^\iota}(\mathbb{T}_p\Gamma) \oplus \text{span}\{E_1(t), \dots, E_{2n-d}(t)\}.$$

Also let $\exp_{\gamma(t)} : \mathbb{T}_p\mathbb{B}_n \rightarrow \mathbb{B}_n$ denote the exponential map. As the Bergman ball is geodesically complete, $\exp_{\gamma(t)}$ has all of $\mathbb{T}_p\mathbb{B}_n$ as its domain. Denote by $B'(0, \epsilon) \subset \mathbb{R}^{2n-d}$ the ball of radius ϵ centered at the origin. We define

$$\begin{aligned} \eta : B(0, 1) \times B'(0, \epsilon) \subset \mathbb{R}^d \times \mathbb{R}^{2n-d} &\rightarrow \mathbb{B}_n \\ \eta(t, x) &= \exp_{\gamma(t)} \left(\sum_{j=1}^{2n-d} x_j E_j(t) \right). \end{aligned}$$

By choosing ϵ small enough, we can guarantee that:

- i) $\eta(t, x)$ is in a geodesic ball centered at $\gamma(t)$ and therefore the Bergman and radial distances between $\eta(t, x)$ and $\gamma(t)$ coincide, this is

$$\beta(\gamma(t), \eta(t, x)) = \left(\sum_{j=1}^{2n-d} x_j^2 \right)^{\frac{1}{2}}.$$

- ii) The Bergman distance from $\eta(t, x)$ to Γ is attained at $\gamma(t)$, this is

$$\beta(\gamma(t), \eta(t, x)) \leq \beta(\gamma(s), \eta(t, x))$$

for all $\gamma(s) \in \Gamma$. Moreover t is the only critical point of the function $\beta(\gamma(\cdot), \eta(t, x))$ in $B(0, 1)$, therefore the last inequality is strict whenever $s \neq t$.

It follows that $\mathcal{V} = \eta(B(0, 1) \times B'(0, \epsilon))$ and

$$\mathcal{I}_p(\alpha) := c_\alpha^{1+p} \int_{B(0,1) \times B'(0,\epsilon)} \left[\int_{B(0,1)} e^{i\alpha\Psi((t,x),s)} \frac{a(\gamma(s)) \mathbf{m}(s) ds}{|1 - \langle \eta(t, x), \gamma(s) \rangle|^{2(n+1)}} \right]^p \cdot \frac{\mathbf{n}(t, x) dt dx}{(1 - |\eta(t, x)|^2)^{(n+1)(1-p)}}, \quad (48)$$

where $\mathbf{m}(s)ds = d\sigma(w)$, $\mathbf{n}(t, x)dtdx = dv(z)$, and we have introduced the phase function $\Psi((t, x), s) = -i \ln(1 - |\varphi_{\eta(t,x)}(\gamma(s))|^2)$. The observations made in ii) and in remark A.1 translate to that for each $(t, x) \in B(0, 1) \times B'(0, \epsilon)$, the function $\Psi((t, x), \cdot)$ has its only critical point at $s = t$.

Applying the method of stationary phase (Theorem 7.7.5 in [5]) yields in

$$\begin{aligned} & \int_{B(0,1)} e^{i\alpha\Psi((t,x),s)} \frac{a(\gamma(s)) \mathbf{m}(s) ds}{|1 - \langle \eta(t, x), \gamma(s) \rangle|^{2(n+1)}} \\ &= \left(\frac{2\pi}{\alpha} \right)^{\frac{d}{2}} e^{i\alpha\Psi((t,x),t)} \frac{a(\gamma(t)) \mathbf{m}(t)}{|1 - \langle \eta(t, x), \gamma(t) \rangle|^{2(n+1)}} \cdot \frac{1 + O\left(\frac{1}{\alpha}\right)}{\det(-i\text{Hess}\Psi((t, x), t))^{\frac{1}{2}}}. \end{aligned} \quad (49)$$

The continuity of the functions involved and the compactness of the domain also imply that

$$\frac{a(\gamma(t)) \mathbf{m}(t)}{|1 - \langle \eta(t, x), \gamma(t) \rangle|^{2(n+1)}} \cdot \frac{\det(-i\text{Hess}\Psi((t, x), t))^{-\frac{1}{2}} \mathbf{n}(t, x)}{(1 - |\eta(t, x)|^2)^{(n+1)(1-p)}}$$

is uniformly bounded for $(t, x) \in B(0, 1) \times B'(0, \epsilon)$. Implementing this and (49) in the expression for $\mathcal{I}_p(\alpha)$ in (48) we find that there exists some constant $C > 0$ such that for all α large enough

$$\begin{aligned} \frac{1}{c_\alpha^{1+p}} \mathcal{I}_p(\alpha) &\leq C \left(\frac{2\pi}{\alpha} \right)^{\frac{dp}{2}} \int_{B'(0,\epsilon)} e^{ip\alpha\Psi((t,x),t)} dx \\ &= C \left(\frac{2\pi}{\alpha} \right)^{\frac{dp}{2}} \int_{B'(0,\epsilon)} \left(1 - \tanh^2 \beta(\gamma(t), \eta(t, x)) \right)^{p\alpha} dx \\ &= C \left(\frac{2\pi}{\alpha} \right)^{\frac{dp}{2}} \int_{B'(0,\epsilon)} \left(1 - \tanh^2 |x| \right)^{p\alpha} dx. \end{aligned} \quad (50)$$

We can find some constant $C > 0$ such that $Cr^{2n-d-1} \leq (\tanh r)^{2n-d-1}$ for all $r \in (0, \epsilon)$. This can be done because $\frac{\tanh r}{r}$ tends to 1 as r approaches 0. Integrating in polar coordinates:

$$\begin{aligned}
\int_{B'(0, \epsilon)} \left(1 - \tanh^2 |x|\right)^{p\alpha} dx &= (2n-d) \int_0^\epsilon r^{2n-d-1} (1 - \tanh^2 r)^{p\alpha} dr \\
&\leq \frac{2n-d}{C} \int_0^\epsilon \tanh^{2n-d-1} r (1 - \tanh^2 r)^{p\alpha} dr \\
&= \frac{n - \frac{d}{2}}{C} \int_0^{\tanh^2 \epsilon} \rho^{n - \frac{d}{2} - 1} (1 - \rho)^{p\alpha - 1} d\rho \\
&\leq \frac{n - \frac{d}{2}}{C} \mathbf{B}\left(n - \frac{d}{2}, p\alpha\right), \tag{51}
\end{aligned}$$

where we have used the change of variables $\rho = \tanh^2 r$ and $\mathbf{B}(z, w)$ is the Beta function. As a consequence of the Stirling's formula the Beta function has the asymptotic approximation

$$\mathbf{B}\left(n - \frac{d}{2}, p\alpha\right) \sim \Gamma\left(n - \frac{d}{2}\right) (p\alpha)^{-n + \frac{d}{2}}$$

as $\alpha \rightarrow \infty$. Combining this and (51) with (50) and the fact that $c_\alpha \sim \alpha^n$ we get

$$\mathcal{I}_p(\alpha) \leq C\alpha^{n(1+p) - \frac{dp}{2} - n + \frac{d}{2}} = C\alpha^{p(n - \frac{d}{2}) + \frac{d}{2}}$$

which by Proposition A.2 implies (45), concluding the proof of Lemma 5.9.

B The determinant of the Hessian

B.1 The determinant as a block matrix

Recall from (22) that computing the determinant of the Hessian of $\varphi(t, \mathbf{0})$ relies on computing that of \mathcal{O}_{m-1} , which we have defined as the $(m-1)d \times (m-1)d$ matrix that expressed as a $(m-1) \times (m-1)$ block matrix with blocks of size $d \times d$ has the form

$$\mathcal{O}_{m-1} = \begin{pmatrix} 2I & -I - iW & \dots & 0 \\ -I + iW & 2I & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & \dots & 2I & -I - iW \\ 0 & \dots & -I + iW & 2I \end{pmatrix},$$

where $W := G^{-1}H$.

Let \mathcal{R} stand for the ring of $d \times d$ complex matrices generated by I and W . Then \mathcal{O}_{m-1} can also be regarded as a $(m-1) \times (m-1)$ matrix with entries in \mathcal{R} . As \mathcal{R} is commutative, any such matrix possesses a well-defined determinant in \mathcal{R} , which we will denote $\det_{\mathcal{R}}$, and that shares all the usual properties with determinants of square matrices with complex entries. The relation between this determinant and the usual one $\det \mathcal{O}_{m-1}$ is given in Theorem 1 in [12]. From that theorem it follows that

$$\det \mathcal{O}_{m-1} = \det \left(\det_{\mathcal{R}} \mathcal{O}_{m-1} \right).$$

Notice that the right-hand side of the equation makes sense since $\det_{\mathcal{R}} \mathcal{O}_{m-1}$ is an element of \mathcal{R} , and thus a $d \times d$ matrix with complex entries. We will use this fact to compute $\det \mathcal{O}_{m-1}$. This demonstration is the same as the one in [11].

Proposition B.1. *Let $q \in \mathbb{N}$. Then*

$$\det_{\mathcal{R}} \mathcal{O}_1 = 2I,$$

$$\det_{\mathcal{R}} \mathcal{O}_2 = 3I - W^2,$$

and

$$\det_{\mathcal{R}} \mathcal{O}_{q+2} = 2I \det_{\mathcal{R}} \mathcal{O}_{q+1} - (I + W^2) \det_{\mathcal{R}} \mathcal{O}_q. \quad (52)$$

Proof. The first two equalities for $\det_{\mathcal{R}} \mathcal{O}_1$ and $\det_{\mathcal{R}} \mathcal{O}_2$ are straightforward. For the one for $\det_{\mathcal{R}} \mathcal{O}_{q+2}$, expand the determinant along the top row. \square

In the following, we seek to obtain a closed form for $\det_{\mathcal{R}} \mathcal{O}_q$. The recursive relation in (52) can be used to get

$$\begin{aligned} \begin{pmatrix} \det_{\mathcal{R}} \mathcal{O}_{q+1} \\ \det_{\mathcal{R}} \mathcal{O}_{q+2} \end{pmatrix} &= \begin{pmatrix} 0 & I \\ -(I + W^2) & 2I \end{pmatrix} \begin{pmatrix} \det_{\mathcal{R}} \mathcal{O}_q \\ \det_{\mathcal{R}} \mathcal{O}_{q+1} \end{pmatrix} \\ &= \begin{pmatrix} 0 & I \\ -(I + W^2) & 2I \end{pmatrix}^q \begin{pmatrix} \det_{\mathcal{R}} \mathcal{O}_1 \\ \det_{\mathcal{R}} \mathcal{O}_2 \end{pmatrix}. \end{aligned} \quad (53)$$

Suppose W is invertible in \mathcal{R} . The eigenvalues in \mathcal{R} of $\begin{pmatrix} 0 & I \\ -(I+W^2) & 2I \end{pmatrix}$ are $I \pm iW$, and thus through diagonalization we can write

$$\begin{pmatrix} 0 & I \\ -(I + W^2) & 2I \end{pmatrix} = \begin{pmatrix} I & I \\ I + iW & I - iW \end{pmatrix} \begin{pmatrix} I + iW & 0 \\ 0 & I - iW \end{pmatrix}$$

$$\cdot \begin{pmatrix} \frac{1}{2}(I + iW^{-1}) & -\frac{i}{2}W^{-1} \\ \frac{1}{2}(I - iW^{-1}) & \frac{i}{2}W^{-1} \end{pmatrix},$$

with $\begin{pmatrix} I & I \\ I+iW & I-iW \end{pmatrix}$ and $\frac{1}{2}\begin{pmatrix} I+iW^{-1} & -iW^{-1} \\ I-iW^{-1} & iW^{-1} \end{pmatrix}$ being inverses in \mathcal{R} . We use this to compute the q -th power of $\begin{pmatrix} 0 & I \\ -(I+W^2) & 2I \end{pmatrix}$, which equals

$$\begin{pmatrix} & * \\ (I+W^2)((I-iW)^q - (I+iW)^q) & (I+iW)^{q+1} - (I-iW)^{q+1} \end{pmatrix},$$

and where the entries in the top row are not displayed since they are not needed for our purposes. Using this to compute the bottom row of (53), we get

$$\begin{aligned} 2iW \det_{\mathcal{R}} \mathcal{O}_{q+2} &= \left[(I+W^2) \left((I-iW)^q - (I+iW)^q \right) \right] \det_{\mathcal{R}} \mathcal{O}_1 \\ &\quad + \left[(I+iW)^{q+1} - (I-iW)^{q+1} \right] \det_{\mathcal{R}} \mathcal{O}_2 \\ &= \left[(I+W^2) \left((I-iW)^q - (I+iW)^q \right) \right] 2I \\ &\quad + \left[(I+iW)^{q+1} - (I-iW)^{q+1} \right] (3I - W^2) \\ &= (I-iW)^q \left[(I+W^2) 2I - (I-iW)(3I - W^2) \right] \\ &\quad + (I+iW)^q \left[-(I+W^2) 2I + (I+iW)(3I - W^2) \right] \\ &= (I-iW)^q \left[-(I-iW)^3 \right] + (I+iW)^q \left[(I+iW)^3 \right] \\ &= -(I-iW)^{q+3} + (I+iW)^{q+3}. \end{aligned}$$

Shift the values of q by making $m-1 = q+2$, and notice that

$$\begin{aligned} (I+iW)^m - (I-iW)^m &= \sum_{\ell=0}^m \binom{m}{\ell} I^{m-\ell} (iW)^\ell - \sum_{\ell=0}^m \binom{m}{\ell} I^{m-\ell} (-iW)^\ell \\ &= \sum_{\ell=0}^m \binom{m}{\ell} (i^\ell - (-i)^\ell) W^\ell \end{aligned}$$

$$= 2iW \sum_{\ell=0}^{\lfloor \frac{m-1}{2} \rfloor} \binom{m}{2\ell+1} (-1)^\ell W^{2\ell},$$

which is a polynomial in W . We now state the following.

Proposition B.2. *Let $m \geq 2$ and $W = G^{-1}H$. Then*

$$2iW \det_{\mathcal{R}} \mathcal{O}_{m-1} = (I + iW)^m - (I - iW)^m,$$

or equivalently

$$\det_{\mathcal{R}} \mathcal{O}_{m-1} = \sum_{\ell=0}^{\lfloor \frac{m-1}{2} \rfloor} \binom{m}{2\ell+1} (-1)^\ell W^{2\ell}. \quad (54)$$

Proof. It only remains to justify the case in which W is not invertible in \mathcal{R} . By the above arguments, for any invertible matrix, the value of the determinant of the corresponding block matrix will be in terms of the polynomial expression in (54), and hence the same expression will remain valid for W by continuity and the density of the invertible matrices in $M_{d \times d}(\mathbb{C})$. \square

We interpret the geometric meaning of the matrix W . For every $p \in \Gamma$ define the endomorphism $K_p : D_{p^\iota}(\mathbb{T}_p \Gamma) \rightarrow D_{p^\iota}(\mathbb{T}_p \Gamma)$ by

$$K_p = \Pi_p \circ J|_{D_{p^\iota}(\mathbb{T}_p \Gamma)}, \quad (55)$$

where $\Pi_p : \mathbb{T}_p \mathbb{B}_n \rightarrow D_{p^\iota}(\mathbb{T}_p \Gamma)$ is the orthogonal projection and $J|_{D_{p^\iota}(\mathbb{T}_p \Gamma)}$ is the restriction of J to $D_{p^\iota}(\mathbb{T}_p \Gamma)$.

Proposition B.3. *Consider $\gamma : B(0, 1) \rightarrow \Gamma$ a parametrization of Γ around p . The matrix W is the matrix representation of the endomorphism K_p with the basis for $D_{p^\iota}(\mathbb{T}_p \Gamma)$ given by $\left\{ D_{p^\iota} \left(\frac{\partial}{\partial \gamma_j} \Big|_t \right) \right\}_{j=1, \dots, d}$.*

Proof. The definition of W is equivalent to $GW = H$, which in turn implies

$$\begin{aligned} h_{jk} &= \sum_{\ell=1}^d g_{j\ell} w_{\ell k} \\ &= \sum_{\ell=1}^d g_p \left(\frac{\partial}{\partial \gamma_j} \Big|_t, \frac{\partial}{\partial \gamma_\ell} \Big|_t \right) w_{\ell k} \\ &= \sum_{\ell=1}^d \mathring{b}_p \left(D_{p^\iota} \left(\frac{\partial}{\partial \gamma_j} \Big|_t \right), D_{p^\iota} \left(\frac{\partial}{\partial \gamma_\ell} \Big|_t \right) \right) w_{\ell k} \end{aligned}$$

$$= \mathring{b}_p \left(D_{p^\ell} \left(\frac{\partial}{\partial \gamma_j} \Big|_t \right), \sum_{\ell=1}^d w_{\ell k} D_{p^\ell} \left(\frac{\partial}{\partial \gamma_\ell} \Big|_t \right) \right) \quad (56)$$

for any $1 \leq j, k \leq d$. On the other hand we know from the definition that

$$h_{jk} = \mathring{b}_p \left(D_{p^\ell} \left(\frac{\partial}{\partial \gamma_j} \Big|_t \right), J D_{p^\ell} \left(\frac{\partial}{\partial \gamma_k} \Big|_t \right) \right) .$$

This together with (56) leads to

$$\mathring{b}_p \left(D_{p^\ell} \left(\frac{\partial}{\partial \gamma_j} \Big|_t \right), J D_{p^\ell} \left(\frac{\partial}{\partial \gamma_k} \Big|_t \right) - \sum_{\ell=1}^d w_{\ell k} D_{p^\ell} \left(\frac{\partial}{\partial \gamma_\ell} \Big|_t \right) \right) = 0$$

for any $1 \leq j, k \leq d$. Since $\left\{ D_{p^\ell} \left(\frac{\partial}{\partial \gamma_j} \Big|_t \right) \right\}_{j=1, \dots, d}$ is a basis for $D_{p^\ell}(\mathbb{T}_p \Gamma)$, we get

$$J D_{p^\ell} \left(\frac{\partial}{\partial \gamma_k} \Big|_t \right) - \sum_{\ell=1}^d w_{\ell k} D_{p^\ell} \left(\frac{\partial}{\partial \gamma_\ell} \Big|_t \right) \in D_{p^\ell}(\mathbb{T}_p \Gamma)^\perp ,$$

and consequently

$$\begin{aligned} \Pi_p \circ J \left(D_{p^\ell} \left(\frac{\partial}{\partial \gamma_k} \Big|_t \right) \right) &= \sum_{\ell=1}^d w_{\ell k} \Pi_p \left(D_{p^\ell} \left(\frac{\partial}{\partial \gamma_\ell} \Big|_t \right) \right) \\ &= \sum_{\ell=1}^d w_{\ell k} D_{p^\ell} \left(\frac{\partial}{\partial \gamma_\ell} \Big|_t \right) \end{aligned}$$

for each $1 \leq k \leq d$, which proves the statement. \square

It can be verified from its definition in terms of Π_p and J that K_p is skew-adjoint with respect to \mathring{b}_p . This implies that W is skew-adjoint and hence all its nonzero eigenvalues are pure imaginary and come in conjugate pairs. Let $\pm i\lambda_k$, $k = 1, \dots, r$ stand for said eigenvalues with possible repetitions due to multiplicity and satisfying

$$0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_r . \quad (57)$$

Let P_{m-1} be the polynomial defined by the right-hand side of (54), this is

$$P_{m-1}(x) = \sum_{\ell=0}^{\lfloor \frac{m-1}{2} \rfloor} \binom{m}{2\ell+1} (-1)^\ell x^{2\ell} .$$

By the spectral mapping theorem the eigenvalues of $P_{m-1}(W)$ are

$$\begin{aligned} P_{m-1}(\pm i\lambda_k) &= \sum_{\ell=0}^{\lfloor \frac{m-1}{2} \rfloor} \binom{m}{2\ell+1} (-1)^\ell (\pm i\lambda_k)^{2\ell} \\ &= \sum_{\ell=0}^{\lfloor \frac{m-1}{2} \rfloor} \binom{m}{2\ell+1} \lambda_k^{2\ell} = \frac{(1+\lambda_k)^m - (1-\lambda_k)^m}{2\lambda_k}, \end{aligned}$$

each with double the multiplicity as in W , together with the eigenvalue $P_{m-1}(0) = m$ with multiplicity $d-2r$ corresponding to the kernel of W . We present a first statement giving an expression for $\det \mathcal{O}_{m-1}$ in terms of the eigenvalues of W .

Proposition B.4. *Let $\pm i\lambda_k$, $k = 1, \dots, r$ be the nonzero eigenvalues of W . Then*

$$\sqrt{\det \mathcal{O}_{m-1}} = m^{\frac{d}{2}-r} \prod_{k=1}^r \frac{(1+\lambda_k)^m - (1-\lambda_k)^m}{2\lambda_k}. \quad (58)$$

Proof.

$$\begin{aligned} \det \mathcal{O}_{m-1} &= \det \left(\det_{\mathcal{R}} \mathcal{O}_{m-1} \right) \\ &= \det P_{m-1}(W) \\ &= \prod_{\{\lambda : \lambda \text{ eigenvalue of } P_{m-1}(W)\}} \lambda \\ &= m^{d-2r} \prod_{k=1}^r \left(\frac{(1+\lambda_k)^m - (1-\lambda_k)^m}{2\lambda_k} \right)^2. \end{aligned}$$

□

B.2 Some symplectic geometry

In this section we describe the behavior of K_p when Γ is isotropic or co-isotropic in order to express the eigenvalues $\pm i\lambda_k$ of W needed to compute the determinant of \mathcal{O}_{m-1} .

Let (X, ω) be a symplectic vector space of dimension $2n$, where the symplectic form is constructed using the formula $\omega(\cdot, \cdot) = g(\cdot, J\cdot)$ with J a

complex structure on X and g an inner product on X compatible with J . Let $V \subseteq X$ be a subspace of dimension d and denote by V^ω its symplectic complement as defined in (5) and by V^\perp its orthogonal complement with respect to g . These sets are also subspaces of X and one can verify directly from the definitions that

$$V^\perp = J(V^\omega) \quad \text{and} \quad J(V^\perp) = V^\omega . \quad (59)$$

Notice that this will imply $\dim V^\omega = \dim V^\perp = 2n - d$.

According to the definition of K_p , we define the endomorphism $K_V : V \rightarrow V$ by

$$K_V = \Pi_V \circ J|_V ,$$

where $\Pi_V : X \rightarrow V$ is the orthogonal projection and $J|_V$ is the restriction of J to V . We will describe these transformations, obtaining results that we will then interpret to the particular case of K_p . These calculations appear in [11] with $T_p\Gamma$ instead of V .

Proposition B.5.

$$\ker K_V = V^\omega \cap V .$$

Proof. Using the relation in (59),

$$\begin{aligned} \ker K_V &= \ker (\Pi_V \circ J|_V) \\ &= J^{-1}(\ker \Pi_V) \cap V \\ &= J^{-1}(V^\perp) \cap V \\ &= V^\omega \cap V . \end{aligned}$$

□

Recall from Definition 1.4 that V is isotropic if $V \subseteq V^\omega$ and co-isotropic if $V \supseteq V^\omega$. The next corollary follows straightforwardly.

Corollary B.6. *If V is isotropic, then K_V is identically zero.*

This fully describes the isotropic case. We turn our attention to the co-isotropic case.

Proposition B.7. *If V is co-isotropic, then the image of K_V satisfies*

$$K_V(V) = J(V) \cap V .$$

Proof. Assume $\xi = K_V v$ for some $v \in V$. Then $Jv = \xi + u$ for some $u \in V^\perp = J(V^\omega)$. Next we can find some $w \in V^\omega$ such that $u = Jw$ and hence $\xi = J(v - w) \in J(V)$, since $w \in V^\omega \subseteq V$ because V is co-isotropic.

For the reverse inclusion, if $\xi \in J(V) \cap V$, then $\xi = Jv$ for some $v \in V$, and therefore $\xi = \Pi_V \xi = \Pi_V Jv = K_V v$. \square

Corollary B.8. *If V is co-isotropic then it splits as a direct sum*

$$V = K_V(V) \oplus V^\omega . \quad (60)$$

Proof. By Proposition B.5 $\ker K_V = V^\omega$, and thus

$$\dim(\ker K_V) = \dim V^\omega = 2n - d .$$

On the other hand, by the dimension theorem we have $\dim(\ker K_V) = d - \dim K_V(V)$. Comparing both expressions yields $\dim K_V(V) = 2d - 2n$. It follows that

$$\dim V^\omega = 2n - d \quad \text{and} \quad \dim K_V(V) = 2d - 2n ,$$

which together add up to $d = \dim V$. Finally, by the previous proposition $K_V(V) \subseteq J(V)$, and therefore

$$K_V(V) \cap V^\omega \subseteq J(V) \cap J(V^\perp) = \emptyset .$$

\square

Remark B.9. We have so far shown that

$$\dim K_V(V) = \begin{cases} 0 & V \text{ isotropic} \\ 2(d - n) & V \text{ co-isotropic} \end{cases} .$$

In general, the rank of K_V is always pair regardless of V being isotropic or co-isotropic because K_V is skew-adjoint. In order to maintain consistency with the definition we gave for the eigenvalues of K_p in (57), we will use r to denote half the rank of K_V .

Keeping up with V co-isotropic, consider the action of K_V on the splitting (60) of V : If $\xi \in V^\omega$, then $J\xi \in JV^\omega = V^\perp$ and hence $K_V \xi = \Pi_V J\xi = 0$. Alternatively, if $\xi \in K_V(V) \subseteq J(V)$, then $J\xi \in V$, thus in this scenario $K_V \xi = \Pi_V J\xi = J\xi$. From this we conclude that in the co-isotropic case, K_V

behaves like J on $K_V(V)$ and like the zero function on V^ω . We can therefore find bases for $K_V(V)$ and V^ω such that K_V has the matrix representation

$$(K_V) = \begin{pmatrix} 0 & -I_{r \times r} & 0 \\ I_{r \times r} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

where $I_{r \times r}$ is the identity matrix of size r . To obtain the eigenvalues of K_V , we now compute the zeros of

$$\begin{aligned} \det [(K_V) - \lambda I_{d \times d}] &= \det \begin{pmatrix} -\lambda I_{r \times r} & -I_{r \times r} & 0 \\ I_{r \times r} & -\lambda I_{r \times r} & 0 \\ 0 & 0 & -\lambda I_{(d-2r) \times (d-2r)} \end{pmatrix} \\ &= \det \begin{pmatrix} -\lambda I_{r \times r} & -I_{r \times r} \\ I_{r \times r} & -\lambda I_{r \times r} \end{pmatrix} \det (-\lambda I_{(d-2r) \times (d-2r)}) \\ &= (\lambda^2 + 1)^r (-\lambda)^{d-2r}. \end{aligned}$$

It follows that the eigenvalues of K_V are $\pm i$ each with multiplicity r and the zero eigenvalue with multiplicity $d - 2r$.

We return to the special case of the symplectic space $(\mathbb{T}_p \mathbb{B}_n, \omega_p)$, with $D_p \iota(\mathbb{T}_p \Gamma) \subset \mathbb{T}_p \mathbb{B}_n$ and K_p as defined in (55). The results exposed in this subsection can be summarized in the next statement.

Proposition B.10. *If Γ is isotropic, then K_p has no nonzero eigenvalues, this is, $r = 0$. If Γ is co-isotropic, then the eigenvalues of K_p are $\pm i$ each with multiplicity $r = d - n$ and the zero eigenvalue with multiplicity $2n - d$.*

We conclude the section with the result we use in Section 4 expressing the value of $\det \mathcal{O}_{m-1}$.

Proposition B.11. *Let Γ be isotropic or co-isotropic. Then*

$$\sqrt{\det \mathcal{O}_{m-1}} = \begin{cases} m^{\frac{d}{2}} & \Gamma \text{ isotropic} \\ 2^{(d-n)(m-1)} m^{n-\frac{d}{2}} & \Gamma \text{ co-isotropic} \end{cases}.$$

Proof. In both cases, we conclude by applying (58) from Proposition B.4. By Proposition B.10 we know that when Γ isotropic $r = 0$; while when Γ co-isotropic $r = d - n$ and $i\lambda_k = i$ for $k = 1, \dots, r$ and thus $\lambda_k = 1$ for $k = 1, \dots, r$.

□

References

- [1] Berezin, F.A., *Covariant and contravariant symbols of operators*, Math. USSR-Izv. 6:5, pp. 1117-1151, 1972.
- [2] Camper, T., Mitkovski, M., *A semi-classical Szegő-type theorem for Toeplitz operators*, arXiv:2411.19298, 2024
- [3] Guillemin, V., *Some classical theorems in spectral theory revisited*. In Seminar on Singularities of Solutions of Linear Partial Differential Equations, Ann. of Math. Stud. 91, pp. 219 -259 Princeton Univ. Press, Princeton, N.J., 1979.
- [4] Grenander, U., Szegő, G., *Toeplitz Forms and Their Applications*, University of California Press, 1958.
- [5] Hörmander, L., *The Analysis of Linear Partial Differential Operators I*, Second Edition, Grundlehren der mathematischen Wissenschaften 256, Springer-Verlag, 1990.
- [6] Huybrechts, D., *Complex Geometry: An Introduction*, Univesitext, Springer, 2005.
- [7] Krantz, S.G., *Function Theory of Several Complex Variables*, AMS Chelsea Publishing Series, American Mathematical Society, 2001.
- [8] Laptev, A., Safarov, Y., *Szegő type limit theorems*, Journal of Functional Analysis, 138, pp. 544-559, 1996.
- [9] Lee, J.M., *Riemannian Manifolds: An Introduction to Curvature*, Springer-Verlag, 1997.
- [10] Libermann, P., Marle, C.M., *Symplectic Geometry and Analytical Mechanics*, Mathematics and Its Applications, Springer Netherlands, 1987.
- [11] Pérez-Esteve, S., Uribe, A., *Szegő limit theorems for singular Berezin-Toeplitz operators*, Journal of Functional Analysis, vol. 278, No. 1, 2020.
- [12] Sylvester, J. R., *Determinants of block matrices*, The Mathematical Gazette, Vol. 84, No. 501, pp. 460-467, 2000.
- [13] Szegő, G., *Ein Grenzwertsatz über die Toeplitzschen Determinanten einer reellen positiven Funktion* Math. Ann., 76, pp. 490–503, 1915.

- [14] Widom, H., *Asymptotic behavior of block Toeplitz matrices and determinants. II* Adv. Math., 21, pp. 1–29, 1976.
- [15] Widom, H., *Eigenvalue distribution theorems for certain homogeneous spaces*, Journal of Functional Analysis, 32, pp. 139-147, 1979.
- [16] Zamarashkin, N. L., Tyrtshnikov, E. E., *Distribution of eigenvalues and singular values of Toeplitz matrices under weakened conditions on the generating function*, Mat. Sb., 188:8, pp. 83–92, 1997.
- [17] Zhu, K., *Operator Theory in Function Spaces*, Second Edition, Mathematical Surveys and Monographs, vol. 138, AMS, 2007.
- [18] Zhu, K., *Spaces of Holomorphic Functions in the Unit Ball*, Graduate Texts in Mathematics, vol. 226, Springer, New York, 2005.