# Orthosymplectic Donaldson-Thomas theory

Chenjing Bu

#### Abstract

We construct and study Donaldson–Thomas invariants counting orthogonal and symplectic objects in linear categories, which are a generalization of the usual Donaldson–Thomas invariants from the structure groups GL(n) to the groups O(n) and Sp(2n), and a special case of the intrinsic Donaldson–Thomas theory developed by the author, Halpern-Leistner, Ibáñez Núñez, and Kinjo [14–16]. Our invariants are defined using the motivic Hall algebra and its orthosymplectic analogue, the motivic Hall module. We prove wall-crossing formulae for our invariants, which relate the invariants with respect to different stability conditions.

As examples, we define Donaldson–Thomas invariants counting orthogonal and symplectic perfect complexes on a Calabi–Yau threefold, and Donaldson– Thomas invariants counting self-dual representations of a self-dual quiver with potential. In the case of quivers, we compute the invariants explicitly in some cases. We also define a motivic version of Vafa–Witten invariants counting orthogonal and symplectic Higgs complexes on a class of algebraic surfaces.

### Contents

1	Intro	duction	3
2	Categories and stacks		8
	2.1	Self-dual categories	8
	2.2	Moduli stacks	10
	2.3	Stability conditions	13
3	Rings of motives 1		
	3.1	Definition	16
	3.2	Motivic Hall algebras and modules	18
4	Invariants 2		
	4.1	Epsilon motives	21
	4.2	DT invariants	24
	4.3	Motivic DT invariants	25
5	Wall-crossing 2		
	5.1	Wall-crossing for epsilon motives	27
	5.2	An anti-symmetrized version	30
	5.3	Wall-crossing for DT invariants	32
6	Applications		37
	6.1	Self-dual quivers	37
	6.2	Orthosymplectic complexes	41
	6.3	DT invariants for curves	45
	6.4	DT invariants for threefolds	47
	6.5	Vafa–Witten type invariants for surfaces	49
Ref	References		

#### 52

### 1 Introduction

**1.1.1. Overview**. The theory of *Donaldson–Thomas* (DT) *invariants* has been a central topic in enumerative geometry, initiated in the works of Donaldson and Thomas [22] and Thomas [47], and further developed by Joyce and Song [35], Kontsevich and Soibelman [37], and many others.

The usual DT invariants are, roughly speaking, virtual counts of semistable objects in 3-Calabi–Yau linear categories, such as the category of coherent sheaves on a Calabi–Yau threefold, or the category of representations of a quiver with potential. These were first constructed by Joyce and Song [35], based on a formalism of Joyce [28–32] for constructing motivic enumerative invariants for abelian categories, and a similar approach was taken by Kontsevich and Soibelman [37].

However, until recently, it was not known how to extend this theory beyond the linear case. A step in this direction was taken by the author [10], who developed a theory of motivic enumerative invariants for moduli stacks of orthosymplectic objects, parallel to Joyce's formalism, and partially inspired by the works of Young [53–55] on self-dual quivers.

This theory was later generalized to *intrinsic DT theory* by the author, Halpern-Leistner, Ibáñez Núñez, and Kinjo [14–16], which is a new framework for enumerative geometry that is intrinsic to the moduli stack, and applies to all algebraic stacks satisfying mild assumptions. The new framework also makes it possible to significantly simplify the original work [10].

The present paper supersedes the author's work [10] by updating and simplifying the theory using the new framework of [14–16].

The main goal of this paper is to construct DT invariants counting *self-dual objects*, or orthosymplectic objects, in self-dual 3-Calabi–Yau linear categories. Self-dual objects typically do not form linear categories, so the usual linear DT theory does not apply.

As applications, we construct DT invariants counting orthosymplectic representations of self-dual quivers with potential, DT invariants counting orthosymplectic perfect complexes on curves and Calabi–Yau threefolds, and also a motivic version of Vafa–Witten invariants counting orthosymplectic Higgs complexes on a class of algebraic surfaces.

Orthosymplectic DT invariants are related to counting D-branes in string theories on Calabi–Yau 3-*orientifolds*, discussed in, for example, Witten [52, §5.2], Diaconescu, Garcia-Raboso, Karp, and Sinha [21], and Hori and Walcher [26].

**1.1.2.** The setting. The basic setting of our construction is as follows. We start with a *self-dual linear category*  $\mathcal{A}$ , that is a linear category equipped with a contravariant involution

$$(-)^{\vee} \colon \mathscr{A} \xrightarrow{\sim} \mathscr{A}^{\mathrm{op}}$$
.

For example,  $\mathscr{A}$  could be the category of vector bundles on a smooth projective curve, where the involution is given by taking the dual bundle. See §2.1 for more details and examples.

We then consider the fixed locus of the involution, which is a groupoid  $\mathscr{A}^{sd}$ , called the groupoid of *self-dual objects*. Explicitly, an object of  $\mathscr{A}^{sd}$  is a pair  $(x, \phi)$ , where x is an object of  $\mathscr{A}$ , and  $\phi : x \xrightarrow{\sim} x^{\vee}$  is an isomorphism satisfying  $\phi = \phi^{\vee}$ . For example, in the case of vector bundles,  $\mathscr{A}^{sd}$  consists of either orthogonal or symplectic vector bundles, depending on the choice of a sign  $\varepsilon = \pm 1$  when identifying a vector bundle with its double dual, which is a part of the data of the involution.

We also assume that  $\mathscr{A}$  is equipped with a moduli stack  $\mathscr{X}$  of objects, so the involution  $(-)^{\vee}$  defines an involution on  $\mathscr{X}$ . Its fixed locus  $\mathscr{X}^{sd}$  is the moduli stack of objects in  $\mathscr{A}^{sd}$ .

In fact, our theory does not essentially use the categories  $\mathscr{A}$  and  $\mathscr{A}^{sd}$ , and only depends on the stack  $\mathscr{X}$ , equipped with its linear structure and involution.

**1.1.3.** Harder–Narasimhan filtrations. Recall that given a *stability condition*  $\tau$  on an abelian category  $\mathscr{A}$ , which, for simplicity, we assume is given by a slope  $\tau(x) \in \mathbb{R}$  for each non-zero object  $x \in \mathscr{A}$ , satisfying certain conditions, there is the notion of  $\tau$ -semistable objects, and every object x has a unique Harder–Narasimhan (HN) filtration

with each quotient  $y_i = x_i/x_{i-1}$  non-zero and semistable, such that  $\tau(y_1) > \cdots > \tau(y_k)$ .

Now, suppose that  $\mathscr{A}$  is self-dual, and that  $\tau$  is compatible with the self-dual structure, in that  $\tau(x^{\vee}) = -\tau(x)$  for all non-zero objects x. Then, for a non-zero self-dual object  $(x, \phi)$ , we necessarily have  $\tau(x) = 0$ , and the self-dual structure  $\phi$  induces isomorphisms  $y_i \xrightarrow{\sim} y_{k+1-i}^{\vee}$  of its HN factors. In particular, if k is odd, the middle piece  $y_{(k+1)/2}$  admits an induced self-dual structure. For convenience, when k is even, we sometimes think of it as having the zero self-dual object as the middle piece.

Therefore, heuristically speaking, if we think of objects of  $\mathscr{A}$  as composed of semistable objects via HN filtrations, we should think of an object of  $\mathscr{A}^{sd}$  as composed of a series of semistable objects of  $\mathscr{A}$ , which are those in the left half of the HN filtration, together with a single semistable self-dual object in  $\mathscr{A}^{sd}$  in the middle; the factors in the right half are dual to those on the left, and do not contain new information.

**1.1.4. Orthosymplectic modules**. The above phenomenon suggests that the collection of orthosymplectic objects can be viewed as a module for the collection of linear objects in some sense. This can be made precise in several different ways.

For example, if we consider the symmetric monoidal groupoid  $(\mathscr{A}^{\approx}, \oplus, 0)$ , where  $\mathscr{A}^{\approx}$  is the underlying groupoid of  $\mathscr{A}$ , then the operation  $\oplus^{\mathrm{sd}} \colon \mathscr{A}^{\approx} \times \mathscr{A}^{\mathrm{sd}} \to \mathscr{A}^{\mathrm{sd}}$  sending (x, y) to  $x \oplus y \oplus x^{\vee}$ , equipped with the obvious self-dual structure, establishes  $\mathscr{A}^{\mathrm{sd}}$  as a module for  $\mathscr{A}^{\approx}$ . Similarly, the corresponding operation  $\oplus^{\mathrm{sd}} \colon \mathscr{X} \times \mathscr{X}^{\mathrm{sd}} \to \mathscr{X}^{\mathrm{sd}}$  on moduli stacks establishes  $\mathscr{X}^{\mathrm{sd}}$ as a module for the commutative monoid stack  $(\mathscr{X}, \oplus, 0)$ . As another example, consider the *ring of motives*  $\mathbb{M}(\mathcal{X})$  over  $\mathcal{X}$ , which we define in §3.1. It has the structure of the *motivic Hall algebra*, introduced by Joyce [29], whose multiplication \*is roughly given by parametrizing all possible extensions of given objects. Note that this is a different multiplication from the one given by the ring structure on the ring of motives. In the orthosymplectic case, the ring of motives  $\mathbb{M}(\mathcal{X}^{sd})$  is a module for the motivic Hall algebra  $\mathbb{M}(\mathcal{X})$ , which we call the *motivic Hall module*. The module structure  $\diamond$ , which we define in §3.2, is roughly given by parametrizing three-step self-dual filtrations, with given graded pieces  $x, y, x^{\vee}$ , with x an ordinary object and y a self-dual object.

Other similar constructions include *cohomological Hall modules* considered by Young [55], and twisted modules for Joyce vertex algebras introduced by the author [11].

**1.1.5.** Epsilon motives. The first main construction of this paper is that of *epsilon motives* for the moduli stack  $\mathcal{X}^{sd}$ , which we present in §4.1, parallel to the construction of Joyce [28–32] in the linear case.

Given a stability condition  $\tau$ , for each non-zero connected component  $\mathscr{X}_{\alpha} \subset \mathscr{X}$ , corresponding to an element  $\alpha \in \pi_0(\mathscr{X}) \setminus \{0\}$ , Joyce defined the *epsilon motive* 

$$\epsilon_{\alpha}(\tau) \in \mathbb{M}(\mathcal{X})$$

living in the ring of motives over  $\mathscr{X}$ , and supported on the semistable locus  $\mathscr{X}^{ss}_{\alpha}(\tau) \subset \mathscr{X}_{\alpha}$ . It can be thought of as an interpolation between the stable locus and the semistable locus.

The epsilon motive  $\epsilon_{\alpha}(\tau)$  satisfies an important property called the *no-pole theorem*, originally proved by Joyce [30, Theorem 8.7], which ensures that it has a well-defined Euler characteristic, which can be used to define DT invariants. In general, Euler characteristics of stacks are ill-behaved, as for example, any reasonable definition would give  $\chi(*/\mathbb{G}_m) = 1/0 = \infty$ . The no-pole theorem ensures that for epsilon motives, this type of divergence does not occur after multiplying by the motive  $[\mathbb{G}_m] = \mathbb{L} - 1$ , which roughly corresponds to removing the copy of  $\mathbb{G}_m$  from the stabilizer groups of all non-zero points of  $\mathcal{X}$ , corresponding to scalar automorphisms of objects in  $\mathcal{A}$ .

In the orthosymplectic case, we present a parallel construction. Given a self-dual stability condition  $\tau$ , for each connected component  $\theta \in \pi_0(\mathcal{X}^{sd})$ , we define the *epsilon motive* 

$$\epsilon^{\mathrm{sd}}_{ heta}( au) \in \mathbb{M}(\mathscr{X}^{\mathrm{sd}})$$
 ,

supported on the semistable locus  $\mathscr{X}^{\mathrm{sd},\mathrm{ss}}_{\theta}(\tau) \subset \mathscr{X}^{\mathrm{sd}}_{\theta}$ . It satisfies an analogous no-pole theorem, and will be used to define orthosymplectic DT invariants.

The construction of the epsilon motives  $\epsilon_{\theta}^{\text{sd}}(\tau)$  is a special case of a general construction in *intrinsic DT theory* by the author, Ibáñez Núñez, and Kinjo [15, §5.2], and we build upon the general framework by specifying explicit combinatorial data, called *stability measures* there, used to define the epsilon motives. We also use the motivic Hall module structure, described in §1.1.4, to make the construction more explicit in our setting. **1.1.6.** DT invariants. We now consider the case when  $\mathscr{X}$  is either a smooth stack, or a (-1)-shifted symplectic stack in the sense of Pantev, Toën, Vaquié, and Vezzosi [40]. The latter case often occurs when  $\mathscr{A}$  is a 3-Calabi–Yau category, such as the category of coherent sheaves on a Calabi–Yau threefold, or more precisely, a modification of this category that is self-dual, which we describe in §6.2.

In this case, following Joyce and Song [35], we define the *DT* invariant  $DT_{\alpha}(\tau) \in \mathbb{Q}$  for  $\mathcal{X}$  and a class  $\alpha \in \pi_0(\mathcal{X}) \setminus \{0\}$  as a weighted Euler characteristic

$$\mathrm{DT}_{\alpha}(\tau) = \int_{\mathcal{X}} (1 - \mathbb{L}) \cdot \epsilon_{\alpha}(\tau) \cdot v_{\mathcal{X}} \, d\chi \,,$$

where  $v_{\mathcal{X}}$  is the *Behrend function* of  $\mathcal{X}$ , a constructible function on  $\mathcal{X}$  introduced by Behrend [5] to capture the virtual geometry of  $\mathcal{X}$ . Multiplying by the factor  $(1 - \mathbb{L})$  corresponds to removing a copy of  $\mathbb{G}_m$  from all stabilizer groups in  $\mathcal{X}$ , as we mentioned in §1.1.5.

Using our self-dual epsilon motives, in §4.2, we define *self-dual DT invariants*  $DT_{\theta}^{sd}(\tau) \in \mathbb{Q}$  for  $\mathcal{X}^{sd}$  and a class  $\theta \in \pi_0(\mathcal{X}^{sd})$  by a similar weighted Euler characteristic

$$\mathrm{DT}^{\mathrm{sd}}_{\theta}(\tau) = \int_{\mathcal{X}^{\mathrm{sd}}} \epsilon^{\mathrm{sd}}_{\theta}(\tau) \cdot v_{\mathcal{X}^{\mathrm{sd}}} \, d\chi \,,$$

where we no longer need the factor  $(1 - \mathbb{L})$ , since a general point in  $\mathscr{X}^{sd}$  does not have  $\mathbb{G}_m$  in its stabilizer, as scaling a self-dual object does not preserve its self-dual structure.

We also define motivic enhancements of these invariants, called *motivic DT invariants*, and denoted by  $DT_{\alpha}^{mot}(\tau)$  and  $DT_{\theta}^{mot,sd}(\tau)$ . They live in a ring of monodromic motives over the base field, and in the (-1)-shifted symplectic case, they also depend on choices of orientations of the stacks  $\mathscr{X}$  and  $\mathscr{X}^{sd}$ . The linear case was due to Kontsevich and Soibelman [37].

**1.1.7. Wall-crossing formulae**. In §5, we prove *wall-crossing formulae* for our DT invariants, which relate the DT invariants of  $\mathcal{X}^{sd}$  for different stability conditions  $\tau$ , generalizing the wall-crossing formulae of Joyce and Song [35]. They are of the form

$$DT_{\theta}^{sd}(\tau_{-}) = \sum_{\substack{n \ge 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathcal{X}) \setminus \{0\}; \ \rho \in \pi_{0}(\mathcal{X}^{sd}):\\ \theta = \alpha_{1} + \alpha_{1}^{\vee} + \dots + \alpha_{n} + \alpha_{n}^{\vee} + \rho}} C(\alpha_{1}, \dots, \alpha_{n}, \rho; \tau_{+}, \tau_{-}) \cdot DT_{\alpha_{1}}(\tau_{+}) \cdots DT_{\alpha_{n}}(\tau_{+}) \cdot DT_{\rho}^{sd}(\tau_{+}) ,$$

$$(1.1.7.1)$$

where  $\tau_{\pm}$  are self-dual stability conditions satisfying certain conditions,  $C(\alpha_1, \ldots, \alpha_n, \rho; \tau_+, \tau_-)$  is a rational combinatorial coefficient, and the sum has only finitely many non-zero terms. A similar formula holds for the motivic DT invariants.

A key ingredient in proving the wall-crossing formula is the *motivic integral identity* for Behrend functions. In the linear case, the identity was proved by Joyce and Song [35, Theorem 5.11] in a numerical form, and conjectured by Kontsevich and Soibelman [37, Conjecture 4] in a stronger motivic form, which was proved by Lê [38]. In the general case, the identity was formulated and proved by the author [12, Theorem 4.2.2].

Wall-crossing formulae govern the structure of the DT invariants, and can be used to com-

pute them explicitly in some cases. The same wall-crossing structure is also satisfied by other types of enumerative invariants in the orthosymplectic case, such as Joyce's [34] homological enumerative invariants, extended by the author [11] to the case of orthosymplectic quivers, where the wall-crossing formula (1.1.7.1) plays a central role.

**1.1.8.** Application to quivers. In §6.1, we define DT invariants for *self-dual quivers with potential*, which are an orthosymplectic analogue of the usual DT theory for quivers with potential, studied in Joyce and Song [35, Ch. 7] and Kontsevich and Soibelman [37, §8].

Self-dual quivers were first introduced by Derksen and Weyman [19], and studied by Young [53–55] in the context of DT theory. These works were a main early source of inspiration for our work.

We also provide an algorithm for computing DT invariants for self-dual quivers where the potential is zero, and present some numerical results. We mention a relation between self-dual quivers and orthosymplectic coherent sheaves in Example 6.3.5.

**1.1.9.** Application to sheaves. In §6.4, we define DT invariants counting orthosymplectic perfect complexes on Calabi–Yau threefolds.

More precisely, we consider Bridgeland stability conditions on the threefold that are compatible with the self-dual structure, so the abelian category of semistable objects of slope 0 is self-dual. We then apply our theory to this category to define DT invariants. We also prove wall-crossing formulae relating the invariants for different Bridgeland stability conditions.

As mentioned in §1.1.1, we expect that these invariants are related to counting D-branes in orientifold string theories.

In §§6.3 and 6.5, we discuss two variants of this theory, DT invariants for curves and motivic Vafa–Witten type invariants for del Pezzo, K3, and abelian surfaces. These Vafa–Witten invariants are a motivic version and an orthosymplectic analogue of the theory of usual Vafa– Witten invariants developed by Tanaka and Thomas [45; 46].

**1.1.10.** Acknowledgements. The author thanks Dominic Joyce for his continuous support throughout this project, and for his many valuable comments and suggestions on the paper. The author also thanks Andrés Ibáñez Núñez and Tasuki Kinjo for helpful discussions.

The author was supported by the Mathematical Institute, University of Oxford.

1.1.11. Conventions. Throughout this paper, we use the following conventions:

- We work over a base field *K*.
- All *schemes*, *algebraic spaces*, and *algebraic stacks* over *K* are assumed to be locally of finite type over *K* and have affine diagonal.
- A *derived algebraic stack* over *K* is a derived stack over *K* that has an open cover by *geometric stacks* in the sense of Toën and Vezzosi [51, §2.2.3], and is assumed locally almost of finite presentation.

### 2 Categories and stacks

#### 2.1 Self-dual categories

**2.1.1.** We begin by introducing a notion of *self-dual linear categories*, as described in §1.1.2. This notion will not be essentially used in our main constructions, since we primarily work with moduli stacks of objects in such categories, which we will discuss in §2.2. However, they will be useful in providing motivations, as well as in studying examples and applications.

2.1.2. Self-dual linear categories. A self-dual K-linear category consists of the following data:

- A *K*-linear category  $\mathscr{A}$ .
- An equivalence of *K*-linear categories  $(-)^{\vee} : \mathscr{A} \xrightarrow{\sim} \mathscr{A}^{\mathrm{op}}$ , called the *dual functor*.
- A natural isomorphism  $\eta: (-)^{\vee\vee} \cong \operatorname{id}_{\mathscr{A}}$ , such that for any object  $x \in \mathscr{A}$ , we have  $\eta_{x^{\vee}} = (\eta_x^{\vee})^{-1} \colon x^{\vee\vee\vee} \cong x^{\vee}$ .

Given such a category  $\mathscr{A}$ , a *self-dual object* in  $\mathscr{A}$  is a pair  $(x, \phi)$ , where  $x \in \mathscr{A}$  and  $\phi \colon x \xrightarrow{\sim} x^{\vee}$  is an isomorphism, such that  $\phi^{\vee} = \phi \circ \eta_x$ :



We denote by  $\mathscr{A}^{sd}$  the groupoid of self-dual objects in  $\mathscr{A}$ , whose morphisms are isomorphisms compatible with the self-dual structures.

More conceptually, ignoring size issues, a self-dual *K*-linear category is a fixed point of the  $\mathbb{Z}_2$ -action on the 2-category of *K*-linear categories, given by taking the opposite category. A self-dual object in  $\mathscr{A}$  is a fixed point of the  $\mathbb{Z}_2$ -action on the underlying groupoid of  $\mathscr{A}$ , given by the dual functor  $(-)^{\vee}$ .

From this viewpoint, we can analogously define self-dual categories and objects in the context of higher categories, although more coherence data is needed if we were to write down the axioms explicitly.

**2.1.3. Example. Vector bundles.** Let *X* be a *K*-scheme, and let  $\mathscr{A} = \text{Vect}(X)$  be the *K*-linear exact category of vector bundles on *X* of finite rank.

For each choice of a sign  $\varepsilon \in \{\pm 1\}$ , there is a self-dual structure  $(-)^{\vee} \colon \mathscr{A} \xrightarrow{\sim} \mathscr{A}^{\text{op}}$  sending a vector bundle to its dual vector bundle, with the natural isomorphism  $\eta \colon (-)^{\vee \vee} \xrightarrow{\sim} \text{id}_{\mathscr{A}}$  given by  $\varepsilon$  times the usual identification.

A self-dual object in  $\mathscr{A}$  is a pair  $(E, \phi)$ , where *E* is a vector bundle on *X*, and  $\phi : E \xrightarrow{\sim} E^{\vee}$  is an isomorphism, satisfying  $\phi^{\vee} = \phi \circ \eta_E$ . Equivalently,  $\phi$  is a non-degenerate symmetric (or antisymmetric) bilinear form on *E* when  $\varepsilon = +1$  (or -1). In particular, if *K* is algebraically closed of characteristic  $\neq 2$ , then self-dual objects of  $\mathscr{A}$  can be identified with principal O(*n*)-bundles (or Sp(*n*)-bundles) on *X*.

**2.1.4. Example. Self-dual quivers.** Let *Q* be a *self-dual quiver*, that is, a quiver with an involution  $\sigma: Q \xrightarrow{\sim} Q^{\text{op}}$ , where  $Q^{\text{op}}$  is the opposite quiver of *Q*. See §6.1 for details.

Let  $\mathscr{A} = \operatorname{Mod}(KQ)$  be the *K*-linear abelian category of finite-dimensional representations of *Q* over *K*. There is a self-dual structure  $(-)^{\vee} : \mathscr{A} \xrightarrow{\sim} \mathscr{A}^{\operatorname{op}}$  sending a representation to the representation with the dual vector spaces and dual linear maps. This also involves choosing signs when defining  $\eta : (-)^{\vee\vee} \xrightarrow{\sim} \operatorname{id}_{\mathscr{A}}$ , as in the previous example. Again, see §6.1 for details.

Self-dual objects in  $\mathcal{A}$  are called *self-dual representations* of Q, which we think of as analogues of orthogonal or symplectic bundles in the quiver setting.

**2.1.5.** Non-example. Coherent sheaves. Let *X* be a connected, smooth, projective *K*-variety of positive dimension, and let  $\mathscr{A} = \operatorname{Coh}(X)$  be the abelian category of coherent sheaves on *X*.

Then  $\mathscr{A}$  does not admit a self-dual structure. This is because  $\mathscr{A}$  is *noetherian*, meaning that every ascending chain of subobjects of a given object stabilizes, while it is not *artinian*, in that there exists an infinite descending chain of subobjects  $\mathscr{O}_X \supset \mathscr{O}_X(-1) \supset \mathscr{O}_X(-2) \supset \cdots$ . Since taking the opposite category exchanges the properties of being noetherian and artinian, the category  $\mathscr{A}$  is not equivalent to  $\mathscr{A}^{\text{op}}$ .

This problem can be fixed, however, by considering the derived category  $\mathscr{D} = D^b \operatorname{Coh}(X)$ , and taking an alternative heart  $\mathscr{A}' \subset \mathscr{D}$  that is compatible with derived duality, which can be constructed from Bridgeland stability conditions. See §6.4 for details.

#### 2.1.6. Self-dual filtrations. We now discuss a useful construction in self-dual linear categories.

Let  $\mathscr{A}$  be a *self-dual exact category* over K, meaning a K-linear exact category with a selfdual structure, such that the dual functor  $(-)^{\vee}$  sends short exact sequences  $y \hookrightarrow x \twoheadrightarrow z$  to short exact sequences  $z^{\vee} \hookrightarrow x^{\vee} \twoheadrightarrow y^{\vee}$ .

For an integer  $n \ge 0$ , define the *K*-linear category  $\mathscr{A}^{(n)}$  of *n*-step filtrations in  $\mathscr{A}$  where objects are diagrams

with each sequence  $x_{i-1} \hookrightarrow x_i \twoheadrightarrow y_i$  short exact in  $\mathscr{A}$ , and morphisms are usual morphisms of diagrams. Define the *dual filtration* of (2.1.6.1) to be the *n*-step filtration

where  $x/x_i$  denotes the cokernel of the inclusion  $x_i \hookrightarrow x$ , which exists by the axioms of an exact category. We have the short exact sequence  $y_i \hookrightarrow x/x_{i-1} \twoheadrightarrow x/x_i$  by the third isomorphism theorem, which holds in any exact category.

This defines a self-dual structure on  $\mathscr{A}^{(n)}$ . Its self-dual objects are called *n*-step self-dual filtrations in  $\mathscr{A}$ , and will be an important idea in our subsequent constructions.

#### 2.2 Moduli stacks

**2.2.1.** We describe a set of axioms for algebraic stacks that behave like moduli stacks of objects in linear categories and self-dual linear categories, based on the notion of *linear moduli stacks* introduced by the author et al. [14, §7.1], which we call *linear stacks* here.

For the main constructions of this paper, it is enough to work only with the moduli stacks, without needing to refer to the original categories. This is also a benefit of the intrinsic framework for enumerative geometry developed in [14–16].

**2.2.2. Graded and filtered points.** Let  $\mathcal{X}$  be an algebraic stack over K. Following Halpern-Leistner [24], define the *stack of graded points* and the *stack of filtered points* of  $\mathcal{X}$  as the mapping stacks

$$\begin{aligned} \operatorname{Grad}(\mathcal{X}) &= \operatorname{Map}(*/\mathbb{G}_{\mathrm{m}}, \mathcal{X}) ,\\ \operatorname{Filt}(\mathcal{X}) &= \operatorname{Map}(\mathbb{A}^{1}/\mathbb{G}_{\mathrm{m}}, \mathcal{X}) , \end{aligned}$$

where we use the scaling action of  $\mathbb{G}_m$  on  $\mathbb{A}^1$ . These are again algebraic stacks over *K*.

Consider the morphisms

$$*/\mathbb{G}_{\mathrm{m}} \xleftarrow[pr]{0} \mathbb{A}^{1}/\mathbb{G}_{\mathrm{m}} \xleftarrow[1]{0} *$$

where pr is induced by the projection  $\mathbb{A}^1 \to *$ . These induce morphisms of stacks

$$\operatorname{Grad}(\mathscr{X}) \xrightarrow[\operatorname{sf}]{\operatorname{cond}} \operatorname{Filt}(\mathscr{X}) \xrightarrow[\operatorname{ev_0}]{\operatorname{ev_0}} \mathscr{X},$$

where the notations 'gr', 'sf', and 'tot' stand for the *associated graded point*, the *split filtration*, and the *total point*, respectively. The morphism gr is an  $\mathbb{A}^1$ -deformation retract, and the morphisms tot and ev<sub>1</sub> are representable, under our assumptions in §1.1.11.

**2.2.3.** Linear stacks. Following the author et al. [14, §7.1], define a *linear stack* over *K* to be the following data:

- An algebraic stack  $\mathcal{X}$  over K.
- A commutative monoid structure  $\oplus : \mathcal{X} \times \mathcal{X} \to \mathcal{X}$ , with unit  $0 \in \mathcal{X}(K)$ .
- $A * / \mathbb{G}_m$ -action  $\odot : * / \mathbb{G}_m \times \mathcal{X} \to \mathcal{X}$  respecting the monoid structure.

Note that these structures come with extra coherence data.

In this case, the set  $\pi_0(\mathcal{X})$  of connected components of  $\mathcal{X}$  carries the structure of a commutative monoid. We denote its operation by +, and its unit by 0.

We require the following additional property:

• There is an isomorphism

$$\coprod_{\gamma: \mathbb{Z} \to \pi_0(\mathcal{X})} \prod_{n \in \text{supp}(\gamma)} \mathcal{X}_{\gamma(n)} \xrightarrow{\sim} \text{Grad}(\mathcal{X}) , \qquad (2.2.3.1)$$

where  $\gamma$  runs through maps of sets  $\mathbb{Z} \to \pi_0(\mathcal{X})$  such that  $\operatorname{supp}(\gamma) = \mathbb{Z} \setminus \gamma^{-1}(0)$  is finite, and the morphism is defined by the composition

$$*/\mathbb{G}_{\mathrm{m}} \times \prod_{n \in \mathrm{supp}(\gamma)} \mathscr{X}_{\gamma(n)} \xrightarrow{(-)^{n}} \prod_{n \in \mathrm{supp}(\gamma)} (*/\mathbb{G}_{\mathrm{m}} \times \mathscr{X}_{\gamma(n)}) \xrightarrow{\odot} \prod_{n \in \mathrm{supp}(\gamma)} \mathscr{X}_{\gamma(n)} \xrightarrow{\oplus} \mathscr{X}$$

on the component corresponding to  $\gamma$ , where the first morphism is given by the *n*-th power map  $(-)^n \colon */\mathbb{G}_m \to */\mathbb{G}_m$  on the factor corresponding to  $\mathscr{X}_{\gamma(n)}$ .

We can think of (2.2.3.1) roughly as an isomorphism  $\operatorname{Grad}(\mathcal{X}) \simeq \mathcal{X}^{\mathbb{Z}}$ , where we only consider components of  $\mathcal{X}^{\mathbb{Z}}$  involving finitely many non-zero classes in  $\pi_0(\mathcal{X})$ .

Most examples of moduli stacks of objects in abelian categories are linear stacks. See [14, §7.1.3] for details.

#### **2.2.4.** Stacks of filtrations. For a linear stack $\mathcal{X}$ , we have canonical isomorphisms

$$\pi_0(\operatorname{Filt}(\mathscr{X})) \simeq \pi_0(\operatorname{Grad}(\mathscr{X})) \simeq \{\gamma \colon \mathbb{Z} \to \pi_0(\mathscr{X}) \mid \operatorname{supp}(\gamma) \text{ finite} \}$$

where the first isomorphism is induced by the morphism gr, and the second is given by (2.2.3.1).

For classes  $\alpha_1, \ldots, \alpha_n \in \pi_0(\mathcal{X})$ , there is a *stack of filtrations* 

$$\mathscr{X}^+_{\alpha_1,\ldots,\alpha_n} \subset \operatorname{Filt}(\mathscr{X})$$

defined as a component corresponding to a map  $\gamma$  as above whose non-zero values agree with the non-zero elements in  $\alpha_n, \ldots, \alpha_1$ , preserving order. We think of this as the stack parametrizing *n*-step filtrations with stepwise quotients of classes  $\alpha_1, \ldots, \alpha_n$ . The isomorphism type of this stack does not depend on the choice of  $\gamma$ , as in [14, §7.1].

The morphisms defined in §2.2.2 restrict to canonical morphisms gr:  $\mathscr{X}_{\alpha_1,...,\alpha_n}^+ \to \mathscr{X}_{\alpha_1} \times \cdots \times \mathscr{X}_{\alpha_n}$  and ev<sub>1</sub>:  $\mathscr{X}_{\alpha_1,...,\alpha_n}^+ \to \mathscr{X}_{\alpha_1+...+\alpha_n}$ , sending a filtration to its associated graded object and total object, respectively. These do not depend on the choice of  $\gamma$ .

We say that  $\mathscr{X}$  has *quasi-compact filtrations*, if for any  $\alpha_1, \ldots, \alpha_n \in \pi_0(\mathscr{X})$ , the morphism  $\operatorname{ev}_1: \mathscr{X}^+_{\alpha_1,\ldots,\alpha_n} \to \mathscr{X}_{\alpha_1+\ldots+\alpha_n}$  is quasi-compact. See also Halpern-Leistner [24, Definition 3.8.1]. This is a very mild condition, and is satisfied by all examples of our interest.

**2.2.5.** Self-dual linear stacks. We now introduce a notion of *self-dual linear stacks*, which describe moduli stacks of objects in self-dual linear categories.

Let  $\mathcal{X}$  be a linear stack over K. A *self-dual structure* on  $\mathcal{X}$  is a  $\mathbb{Z}_2$ -action on  $\mathcal{X}$ , given by an involution

$$(-)^{\vee} \colon \mathscr{X} \xrightarrow{\sim} \mathscr{X},$$

together with a 2-isomorphism  $\eta \colon (-)^{\vee\vee} \cong \operatorname{id}_{\mathscr{X}}$  with  $\eta_{(-)^{\vee}} = (\eta_{(-)}^{\vee})^{-1}$  similarly to §2.1.2, such that the involution respects the monoid structure  $\oplus$  on  $\mathscr{X}$ , and inverts the  $*/\mathbb{G}_{\mathrm{m}}$ -action  $\odot$ , meaning that it is equivariant with respect to the involution  $(-)^{-1} \colon */\mathbb{G}_{\mathrm{m}} \to */\mathbb{G}_{\mathrm{m}}$ . Note that extra coherence data is needed for these compatibility conditions as well.

In this case, we call  $\mathcal{X}$  a *self-dual linear stack*. Define the *stack of self-dual points* of  $\mathcal{X}$  as the fixed locus

$$\mathscr{X}^{\mathrm{sd}} = \mathscr{X}^{\mathbb{Z}_2}$$
 .

It has affine diagonal by Lemma 2.2.9 below. Note that this is different from the fixed locus of the automorphism  $(-)^{\vee}$  of  $\mathscr{X}$ , which would give the fixed locus of the corresponding  $\mathbb{Z}$ -action on  $\mathscr{X}$ , rather than that of the  $\mathbb{Z}_2$ -action.

There is a monoid action

$$\oplus^{\mathrm{sd}} \colon \mathscr{X} \times \mathscr{X}^{\mathrm{sd}} \longrightarrow \mathscr{X}^{\mathrm{sd}}$$
,

given by  $(x, y) \mapsto x \oplus y \oplus x^{\vee}$ . This induces a monoid action  $\pi_0(\mathcal{X}) \times \pi_0(\mathcal{X}^{sd}) \to \pi_0(\mathcal{X}^{sd})$ , which we often denote by  $(\alpha, \theta) \mapsto \alpha + \theta + \alpha^{\vee}$ , where  $\alpha + \alpha^{\vee}$  can also be seen as a class in  $\pi_0(\mathcal{X}^{sd})$ , corresponding to the case when  $\theta = 0$ .

2.2.6. Example. Let  $\mathscr{X} = \coprod_{n \in \mathbb{N}} */\mathrm{GL}(n)$  be the moduli stack of vector spaces over K, which is a linear moduli stack. Consider the involution  $(-)^{\vee} : */\mathrm{GL}(n) \to */\mathrm{GL}(n)$  sending a vector space to its dual, or equivalently, sending a matrix to its inverse transpose. Then  $\mathscr{X}^{\mathrm{sd}}$  is the classifying stack of non-degenerate symmetric (or anti-symmetric) bilinear forms, depending on the choice of the 2-morphism  $\eta$ , similarly to Example 2.1.3. In particular, if K is algebraically closed of characteristic  $\neq 2$ , then  $\mathscr{X}^{\mathrm{sd}}$  is either  $\coprod_{n \in \mathbb{N}} */\mathrm{O}(n)$  or  $\coprod_{n \in \mathbb{N}} */\mathrm{Sp}(2n)$ .

2.2.7. Self-dual graded points. The involution on  $\mathscr{X}$  induces an involution on  $\operatorname{Grad}(\mathscr{X})$ , and we may identify  $\operatorname{Grad}(\mathscr{X}^{\operatorname{sd}}) \simeq \operatorname{Grad}(\mathscr{X})^{\mathbb{Z}_2}$ . This gives an isomorphism

$$\operatorname{Grad}(\mathscr{X}^{\operatorname{sd}}) \simeq \coprod_{\substack{\gamma : \mathbb{Z} \setminus \{0\} \to \pi_0(\mathscr{X}) \\ \text{involutive,} \\ \gamma(0) \in \pi_0(\mathscr{X}^{\operatorname{sd}})}} \left( \mathscr{X}_{\gamma(0)}^{\operatorname{sd}} \times \prod_{n > 0 : \gamma(n) \neq 0} \mathscr{X}_{\gamma(n)} \right),$$
(2.2.7.1)

where  $\gamma$  runs through finitely supported maps that are *involutive*, meaning that  $\gamma(-n) = \gamma(n)^{\vee}$  for all n > 0, and  $\gamma(0)$  is a convenient notation which is independent of the map  $\gamma$ , and  $\mathcal{X}_{\gamma(0)}^{sd} \subset \mathcal{X}^{sd}$  denotes the component corresponding to  $\gamma(0)$ .

**2.2.8. Self-dual filtrations.** For classes  $\alpha_1, \ldots, \alpha_n \in \pi_0(\mathcal{X})$  and  $\theta \in \pi_0(\mathcal{X}^{sd})$ , define the *stack* of *self-dual filtrations* 

$$\mathscr{X}^{\mathrm{sd},+}_{\alpha_1,\ldots,\alpha_n,\theta} \subset \mathrm{Filt}(\mathscr{X}^{\mathrm{sd}})$$

as a component such that under the isomorphism  $\pi_0(\text{Filt}(\mathcal{X}^{\text{sd}})) \simeq \pi_0(\text{Grad}(\mathcal{X}^{\text{sd}}))$ , its corresponding map  $\gamma$  as above has  $\gamma(0) = \theta$ , and its non-zero values at positive integers agree with

the non-zero elements in  $\alpha_n, \ldots, \alpha_1$ , preserving order. This does not depend on the choice of  $\gamma$  by the constancy theorem of the author et al. [14, Theorem 6.1.2].

The stack  $\mathscr{X}_{\alpha_1,\ldots,\alpha_n,\theta}^{\mathrm{sd},+}$  can be thought of as parametrizing *self-dual filtrations* in the sense of §2.1.6, with stepwise quotients of classes  $\alpha_1,\ldots,\alpha_n,\theta,\alpha_n^{\vee},\ldots,\alpha_1^{\vee}$ .

The morphisms defined in §2.2.2 restrict to morphisms gr:  $\mathscr{X}_{\alpha_1,\dots,\alpha_n,\theta}^{\mathrm{sd},+} \to \mathscr{X}_{\alpha_1} \times \cdots \times \mathscr{X}_{\alpha_n} \times \mathscr{X}_{\theta}^{\mathrm{sd}}$ and  $\mathrm{ev}_1 : \mathscr{X}_{\alpha_1,\dots,\alpha_n,\theta}^{\mathrm{sd},+} \to \mathscr{X}_{\alpha_1+\dots+\alpha_n+\theta+\alpha_n^{\vee}+\dots+\alpha_1^{\vee}}^{\mathrm{sd}}$ . If  $\mathscr{X}$  has quasi-compact filtrations as in §2.2.4, then the morphism  $\mathrm{ev}_1$  described above is always quasi-compact.

**2.2.9. Lemma.** Let  $\mathscr{X}$  be an algebraic stack with affine diagonal, acted on by  $\mathbb{Z}_2$ . Then the forgetful morphism  $\mathscr{X}^{\mathbb{Z}_2} \to \mathscr{X}$  is affine.

**Proof.** Let  $\mathscr{J} = \mathscr{X} \times_{j_0, \mathscr{X} \times \mathscr{X}, j_1} \mathscr{X}$ , where  $j_0$  is the diagonal morphism, and  $j_1 = (\mathrm{id}, i)$ , where i is the involution. Let  $\pi : \mathscr{J} \to \mathscr{X}$  be the projection to the first factor, which is affine as  $\mathscr{X}$  has affine diagonal. Let  $\mathbb{Z}_2$  act on  $\mathscr{J}$  by the involution on the second factor, so we may identify  $\mathscr{J}^{\mathbb{Z}_2} \simeq \mathscr{X}^{\mathbb{Z}_2}$ . Then  $\pi$  is equivariant with respect to the trivial  $\mathbb{Z}_2$ -action on  $\mathscr{X}$ , so the forgetful morphism  $\mathscr{J}^{\mathbb{Z}_2} \to \mathscr{J}$  is a closed immersion, which can be seen by base changing along morphisms from affine schemes to  $\mathscr{X}$ . The composition  $\mathscr{X}^{\mathbb{Z}_2} \simeq \mathscr{J}^{\mathbb{Z}_2} \to \mathscr{J} \to \mathscr{X}$  is thus affine.

Note that in this lemma, we may allow the base to be any algebraic stack, where we assume that the relative diagonal of  $\mathcal{X}$  is affine, and that the action respects the structure morphism.

### 2.3 Stability conditions

**2.3.1.** We define a notion of *stability conditions* on linear stacks, based on ideas in the works of Rudakov [44], Joyce [30], Bridgeland [8], as well as Halpern-Leistner [24]. Such a stability condition will determine a *semistable locus* in the stack, and enumerative invariants will count the points in the semistable locus.

**2.3.2.**  $\Theta$ -stratifications. We first define  $\Theta$ -stratifications following Halpern-Leistner [24]. This is a geometric formulation of the existence and uniqueness of HN filtrations. We slightly weaken the original definition by discarding the ordering on the set of strata, which we will not need.

A  $\Theta$ -stratification of a stack  $\mathcal{X}$  is the following data:

• Open substacks  $\mathcal{S} \subset \operatorname{Filt}(\mathcal{X})$  and  $\mathcal{Z} \subset \operatorname{Grad}(\mathcal{X})$ , with  $\mathcal{S} = \operatorname{gr}^{-1}(\mathcal{Z})$ ,

such that for each  $\lambda \in \pi_0(\operatorname{Grad}(\mathcal{X})) \simeq \pi_0(\operatorname{Filt}(\mathcal{X}))$ , if we write  $\mathcal{S}_{\lambda} \subset \mathcal{S}$  and  $\mathcal{Z}_{\lambda} \subset \mathcal{Z}$  for the parts lying in the components  $\mathcal{X}_{\lambda}^+ \subset \operatorname{Filt}(\mathcal{X})$  and  $\mathcal{X}_{\lambda} \subset \operatorname{Grad}(\mathcal{X})$ , respectively, then:

For each λ, the morphism ev<sub>1</sub>: S<sub>λ</sub> → X is a locally closed immersion, and the family (S<sub>λ</sub>)<sub>λ</sub> defines a locally closed stratification of X.

In this case, each  $S_{\lambda}$  is called a *stratum*, and each  $\mathcal{Z}_{\lambda}$  is called the *centre* of the stratum  $S_{\lambda}$ .

**2.3.3. Stability conditions.** Let  $\mathcal{X}$  be a linear stack. A *stability condition* on  $\mathcal{X}$  is a map

$$\tau\colon \pi_0(\mathscr{X})\setminus\{0\}\longrightarrow T$$

to a totally ordered set *T*, satisfying the following conditions:

- (i) If  $\alpha_1, \alpha_2 \in \pi_0(\mathcal{X}) \setminus \{0\}$  and  $\tau(\alpha_1) \leq \tau(\alpha_2)$ , then  $\tau(\alpha_1) \leq \tau(\alpha_1 + \alpha_2) \leq \tau(\alpha_2)$ .
- (ii) For any class  $\alpha \in \pi_0(\mathcal{X})$ , the semistable locus

$$\mathscr{X}_{\alpha}^{ss}(\tau) = \mathscr{X}_{\alpha} \setminus \bigcup_{\substack{\alpha = \alpha_1 + \alpha_2 \\ \tau(\alpha_1) > \tau(\alpha_2)}} \operatorname{ev}_1(\mathscr{X}_{\alpha_1, \alpha_2}^+)$$
(2.3.3.1)

is open in  $\mathscr{X}_{\alpha}$ , where  $\alpha_1, \alpha_2$  are assumed non-zero. Moreover, for any  $t \in T$ , the union of all  $\mathscr{X}_{\alpha}^{ss}(\tau)$  with either  $\alpha = 0$  or  $\tau(\alpha) = t$  is an open linear substack of  $\mathscr{X}$ .

(iii) The open substacks

$$\mathcal{Z}_{\alpha_1,\dots,\alpha_n}(\tau) = \mathcal{X}_{\alpha_1}^{\mathrm{ss}}(\tau) \times \dots \times \mathcal{X}_{\alpha_n}^{\mathrm{ss}}(\tau) \subset \mathcal{X}_{\alpha_1} \times \dots \times \mathcal{X}_{\alpha_n}$$
  
$$\mathcal{S}_{\alpha_1,\dots,\alpha_n}(\tau) = \mathrm{gr}^{-1}(\mathcal{Z}_{\alpha_1,\dots,\alpha_n}(\tau)) \qquad \subset \mathcal{X}_{\alpha_1,\dots,\alpha_n}^+$$

for all  $n \ge 0$  and classes  $\alpha_1, \ldots, \alpha_n \in \pi_0(\mathcal{X}) \setminus \{0\}$  with  $\tau(\alpha_1) > \cdots > \tau(\alpha_n)$  define a  $\Theta$ -stratification of  $\mathcal{X}$  in the sense of §2.3.2.

More precisely, the last condition means that for each choice of  $\alpha_1, \ldots, \alpha_n$  as above, we choose an element  $\lambda$  as in §2.3.2 such that  $\mathscr{X}_{\lambda} \simeq \mathscr{X}_{\alpha_1} \times \cdots \times \mathscr{X}_{\alpha_n}$  and  $\mathscr{X}^+_{\lambda} \simeq \mathscr{X}^+_{\alpha_1,\ldots,\alpha_n}$ , and we set  $\mathscr{X}_{\lambda}$ and  $\mathscr{S}_{\lambda}$  as above; for all other  $\lambda$ , we set them to be empty.

2.3.4. Examples. Here are some examples of stability conditions on linear stacks.

- (i) Let  $\mathscr{X}$  be any linear stack. The constant map  $\tau \colon \pi_0(\mathscr{X}) \setminus \{0\} \to \{0\}$  is called the *trivial stability condition*, where  $\mathscr{X}^{ss}_{\alpha}(\tau) = \mathscr{X}_{\alpha}$  for all  $\alpha$ .
- (ii) Let  $\mathscr{X}$  be the moduli stack of representations of a quiver Q. Then any *slope function*  $\mu: Q_0 \to \mathbb{Q}$  induces a stability condition on  $\mathscr{X}$  given by

$$\tau(d) = \frac{\sum_{i \in Q_0} d_i \cdot \mu(i)}{\sum_{i \in Q_0} d_i}$$

for non-zero dimension vectors  $d \in \pi_0(\mathcal{X}) \setminus \{0\}$ , where the  $\Theta$ -stratification can be constructed from Ibáñez Núñez [27, Theorem 2.6.3]. See §6.1 for more details.

(iii) Let X be the moduli stack of coherent sheaves on a projective scheme Y over a field K of characteristic zero. Then *Gieseker stability* is a stability condition on X, where the choice of τ is described in Joyce [30, Example 4.16], and the Θ-stratification exists by Alper, Halpern-Leistner, and Heinloth [2, Example 7.28].

**2.3.5.** For self-dual linear stacks. Let  $\mathscr{X}$  be a self-dual linear stack over *K*, and let  $\tau$  be a stability condition on  $\mathscr{X}$ . We say that  $\tau$  is *self-dual*, if the following condition holds:

• For any  $\alpha, \beta \in \pi_0(\mathcal{X}) \setminus \{0\}$ , we have  $\tau(\alpha) \leq \tau(\beta)$  if and only if  $\tau(\alpha^{\vee}) \geq \tau(\beta^{\vee})$ .

In this case, for each  $\theta \in \pi_0(\mathcal{X}^{sd})$ , writing  $\alpha = j(\theta)$  for the corresponding class in  $\pi_0(\mathcal{X})$ , we have the *semistable locus* 

$$\mathscr{X}_{\theta}^{\mathrm{sd},\mathrm{ss}}(\tau) = \mathscr{X}_{\alpha}^{\mathrm{ss}}(\tau)^{\mathbb{Z}_2} \cap \mathscr{X}_{\theta}^{\mathrm{sd}} \subset \mathscr{X}_{\theta}^{\mathrm{sd}} , \qquad (2.3.5.1)$$

where  $\mathscr{X}^{\mathrm{sd}}_{\theta} \subset (\mathscr{X}_{\alpha})^{\mathbb{Z}_2}$  as an open and closed substack.

We have an induced  $\Theta\text{-stratification of }\mathcal{X}^{\mathrm{sd}}$  given by the open substacks

$$\begin{aligned} \mathcal{Z}_{\alpha_{1},\dots,\alpha_{n},\theta}^{\mathrm{sd}}(\tau) &= \mathcal{X}_{\alpha_{1}}^{\mathrm{ss}}(\tau) \times \dots \times \mathcal{X}_{\alpha_{n}}^{\mathrm{ss}}(\tau) \times \mathcal{X}_{\theta}^{\mathrm{sd},\mathrm{ss}}(\tau) \subset \mathcal{X}_{\alpha_{1}} \times \dots \times \mathcal{X}_{\alpha_{n}} \times \mathcal{X}_{\theta}^{\mathrm{sd}} ,\\ \mathcal{S}_{\alpha_{1},\dots,\alpha_{n},\theta}^{\mathrm{sd}}(\tau) &= \mathrm{gr}^{-1}(\mathcal{Z}_{\alpha_{1},\dots,\alpha_{n},\theta}^{\mathrm{sd}}(\tau)) \qquad \qquad \subset \mathcal{X}_{\alpha_{1},\dots,\alpha_{n},\theta}^{\mathrm{sd},+} ,\end{aligned}$$

where  $\alpha_1, \ldots, \alpha_n \in \pi_0(\mathcal{X}) \setminus \{0\}$  and  $\theta \in \pi_0(\mathcal{X}^{sd})$  are classes such that  $\tau(\alpha_1) > \cdots > \tau(\alpha_n) > 0$ . These strata and centres can also be realized as  $\mathbb{Z}_2$ -fixed loci in the strata and centres of the  $\Theta$ -stratification of  $\mathcal{X}$  given by  $\tau$ .

**2.3.6. Permissibility**. Let  $\mathscr{X}$  be a linear stack over *K*, and let  $\tau$  be a stability condition on  $\mathscr{X}$ . We say that  $\tau$  is *permissible*, if the following condition holds:

• For any  $\alpha \in \pi_0(\mathcal{X})$ , the semistable locus  $\mathcal{X}^{ss}_{\alpha}(\tau) \subset \mathcal{X}_{\alpha}$  is quasi-compact.

This is similar to the notion of permissible weak stability conditions in Joyce [30, Definition 4.7] and Joyce and Song [35, Definition 3.7].

We also explain in §4.1.5 below that this notion of permissibility implies the corresponding *stability measures* being permissible as in [15, §4.1.4]. The following lemma can be seen as a shadow of this result, which we prove directly.

**2.3.7. Lemma.** Let  $\mathcal{X}$  be a linear stack over K, with quasi-compact filtrations as in §2.2.4, and let  $\tau$  be a permissible stability condition on  $\mathcal{X}$ .

Then for any  $\alpha \in \pi_0(\mathcal{X}) \setminus \{0\}$ , there are only finitely many decompositions  $\alpha = \alpha_1 + \cdots + \alpha_n$ into classes  $\alpha_i \in \pi_0(\mathcal{X}) \setminus \{0\}$ , such that  $\tau(\alpha_i) = \tau(\alpha)$  and  $\mathcal{X}_{\alpha_i}^{ss}(\tau) \neq \emptyset$  for all *i*.

**Proof.** Let  $t = \tau(\alpha)$ . Then the open substack

$$\mathscr{X}( au;t) = \{0\} \cup \coprod_{\substack{eta \in \pi_0(\mathscr{X}) \setminus \{0\}: \ au(eta) = t}} \mathscr{X}^{ ext{ss}}_eta( au) \subset \mathscr{X}$$

is again a linear stack. Replacing  $\mathscr{X}$  by  $\mathscr{X}(\tau; t)$ , we may assume that  $\mathscr{X}$  has quasi-compact connected components, and that  $\tau$  is trivial.

By the finiteness theorem of the author et al. [14, Theorem 6.2.3], each connected component of  $\mathscr{X}$  has finitely many *special faces*. In this case, this is the statement that for any  $\alpha \in \pi_0(\mathscr{X}) \setminus \{0\}$ , there are finitely many decompositions  $\alpha = \alpha_1 + \cdots + \alpha_n$ , such that all other decompositions can be obtained from combining terms in these decompositions, and hence the total number of decompositions is finite.

## 3 Rings of motives

#### 3.1 Definition

**3.1.1.** We provide background material on *rings of motives* over an algebraic stack, following the author, Ibáñez Núñez, and Kinjo [15, §3], based on the theory of Joyce [31]. These rings will be used to construct DT invariants later.

Recall from §1.1.11 our running assumptions on algebraic stacks.

**3.1.2. The ring of motives.** Let  $\mathscr{X}$  be an algebraic stack over *K*, and let *A* be a commutative ring. The *ring of motives* over  $\mathscr{X}$  with coefficients in *A* is the *A*-module

$$\mathbb{M}(\mathcal{X};A) = \bigoplus_{\mathcal{X} \to \mathcal{X}} A \cdot [\mathcal{X}] / \sim,$$

where we run through isomorphism classes of representable morphisms  $\mathscr{Z} \to \mathscr{X}$  of finite type, with  $\mathscr{Z}$  quasi-compact, and  $\hat{\oplus}$  indicates that we take the set of *locally finite sums*, that is, possibly infinite sums  $\sum_{\mathscr{Z}\to\mathscr{X}} a_{\mathscr{Z}} \cdot [\mathscr{Z}]$ , such that for each quasi-compact open substack  $\mathscr{U} \subset \mathscr{X}$ , there are only finitely many  $\mathscr{Z}$  such that  $a_{\mathscr{X}} \neq 0$  and  $\mathscr{Z} \times_{\mathscr{X}} \mathscr{U} \neq \emptyset$ . The relation ~ is generated by locally finite sums of elements of the form

$$a \cdot ([\mathcal{Z}] - [\mathcal{Z}'] - [\mathcal{Z} \setminus \mathcal{Z}']),$$

where  $a \in A$ ,  $\mathscr{Z}$  is as above, and  $\mathscr{Z}' \subset \mathscr{Z}$  is a closed substack. The class  $[\mathscr{Z}] \in \mathbb{M}(\mathscr{X}; A)$  is called the *motive* of  $\mathscr{Z}$ .

For a representable morphism  $\mathscr{Z} \to \mathscr{X}$  of finite type, where  $\mathscr{Z}$  is not necessarily quasicompact, we can still define its motive  $[\mathscr{Z}] \in \mathbb{M}(\mathscr{X}; A)$ , by stratifying  $\mathscr{Z}$  into quasi-compact locally closed substacks,  $\mathscr{Z} = \bigcup_{i \in I} \mathscr{Z}_i$ , and defining  $[\mathscr{Z}] = \sum_{i \in I} [\mathscr{Z}_i]$  as a locally finite sum.

The ring structure on  $\mathbb{M}(\mathcal{X}; A)$  is given by  $[\mathcal{Z}] \cdot [\mathcal{Z}'] = [\mathcal{Z} \times_{\mathcal{X}} \mathcal{Z}']$  on generators, with unit element  $[\mathcal{X}]$ .

We also write  $\mathbb{M}(\mathcal{X})$  for  $\mathbb{M}(\mathcal{X};\mathbb{Z})$ , and  $\mathbb{M}(K;A)$  for  $\mathbb{M}(\text{Spec}(K);A)$ .

3.1.3. Properties. We list some basic properties of rings of motives; see [15, §3] for details.

(i) For a morphism  $f: \mathscr{Y} \to \mathscr{X}$ , there is a *pullback map* 

$$f^* \colon \mathbb{M}(\mathscr{X}; A) \longrightarrow \mathbb{M}(\mathscr{Y}; A)$$
,

given by  $[\mathscr{X}] \mapsto [\mathscr{X} \times_{\mathscr{X}} \mathscr{Y}]$  on generators, which is a ring homomorphism.

(ii) For a representable quasi-compact morphism  $f: \mathscr{Y} \to \mathscr{X}$ , there is a *pushforward map* 

$$f_!: \mathbb{M}(\mathcal{Y}; A) \longrightarrow \mathbb{M}(\mathcal{X}; A)$$
,

given by  $[\mathcal{Z}] \mapsto [\mathcal{Z}]$  on generators. This is not a ring homomorphism in general.

(iii) For stacks  $\mathcal{X}$  and  $\mathcal{Y}$ , there is an *external product* 

$$\boxtimes : \mathbb{M}(\mathcal{X}; A) \otimes \mathbb{M}(\mathcal{Y}; A) \longrightarrow \mathbb{M}(\mathcal{X} \times \mathcal{Y}; A) ,$$

given by  $[\mathscr{Z}] \otimes [\mathscr{Z}'] \mapsto [\mathscr{Z} \times \mathscr{Z}']$  on generators. The multiplication on  $\mathbb{M}(\mathscr{X}; A)$  can be realized as the external product for  $\mathscr{Y} = \mathscr{X}$  followed by pulling back along the diagonal.

(iv) For a representable quasi-compact morphism  $f: \mathcal{Y} \to \mathcal{X}$ , we have the *projection formula* 

$$f_!(a \cdot f^*(b)) = f_!(a) \cdot b \tag{3.1.3.1}$$

for  $a \in \mathbb{M}(\mathcal{Y}; A)$  and  $b \in \mathbb{M}(\mathcal{X}; A)$ , which can be verified on generators.

(v) For a pullback diagram

$$\begin{array}{ccc} \mathcal{Y}' & \xrightarrow{g'} & \mathcal{Y} \\ f' & & & \downarrow f \\ \mathcal{X}' & \xrightarrow{g} & \mathcal{X} \end{array}, \end{array}$$

where f is representable and quasi-compact, we have the *base change formula* 

$$g^* \circ f_! = f'_! \circ (g')^* \colon \mathbb{M}(\mathcal{Y}; A) \longrightarrow \mathbb{M}(\mathcal{X}'; A) . \tag{3.1.3.2}$$

(vi) For a generator  $[\mathcal{X}] \in \mathbb{M}(\mathcal{X}; A)$  and a vector bundle  $\mathcal{E} \to \mathcal{X}$  of rank *n*, we have

$$[\mathscr{E}] = \mathbb{L}^n \cdot [\mathscr{Z}] . \tag{3.1.3.3}$$

#### 3.1.4. Motivic integration. We also consider the localization

$$\widehat{\mathbb{M}}(\mathscr{X};A) = \mathbb{M}(\mathscr{X};A) \underset{A[\mathbb{L}]}{\otimes} A[\mathbb{L}^{\pm 1}, (\mathbb{L}^{k}-1)^{-1}:k > 0],$$

where  $\mathbb{L} = [\mathbb{A}^1]$  is the motive of the affine line, and  $\hat{\otimes}$  denotes the completed tensor product with respect to locally finite sums. We call this the *completed ring of motives* over  $\mathcal{X}$ .

For a quasi-compact algebraic stack  $\mathcal{X}$  over K, there is a *motivic integration map* 

$$\int_{\mathcal{X}} (-) \colon \mathbb{M}(\mathcal{X}; A) \longrightarrow \hat{\mathbb{M}}(K; A) ,$$

sending a generator of the form [Z/GL(n)] to the element  $[Z] \cdot [GL(n)]^{-1}$ , where *Z* is a quasicompact algebraic space over *K*. Such elements [Z/GL(n)] generate  $\mathbb{M}(\mathcal{X}; A)$ . See [15, §3] for details.

#### 3.1.5. Euler characteristics. Let

$$\widehat{\mathbb{M}}^{\mathrm{reg}}(K;A) = \mathbb{M}(K;A) \underset{A[\mathbb{L}]}{\otimes} A[\mathbb{L}^{\pm 1}, (1 + \mathbb{L} + \dots + \mathbb{L}^{k-1})^{-1} : k > 0] / (\mathbb{L} - 1) \text{-torsion}$$

be the subring of  $\hat{\mathbb{M}}(K; A)$  consisting of motives 'with no poles at  $\mathbb{L} = 1$ ', and suppose that *A* 

is a Q-algebra. Then there is an Euler characteristic map  $\chi: \widehat{\mathbb{M}}^{\operatorname{reg}}(K; A) \to A$ , sending  $\mathbb{L}$  to 1, and  $(1 + \mathbb{L} + \cdots + \mathbb{L}^{k-1})^{-1}$  to 1/k. See [15, §3] for details.

**3.1.6.** The virtual rank decomposition. Let  $\mathscr{X}$  be a stack over *K*, and let *A* be a commutative  $\mathbb{Q}$ -algebra. As in Joyce [31, §5] and the author, Ibáñez Núñez, and Kinjo [15, §5.1], there is a *virtual rank decomposition* 

$$\mathbb{M}(\mathcal{X};A) = \bigoplus_{k \ge 0} \mathbb{M}^{(k)}(\mathcal{X};A),$$

where  $\hat{\oplus}$  means taking locally finite sums as in §3.1.2, and each  $\mathbb{M}^{(k)}(\mathcal{X};A) \subset \mathbb{M}(\mathcal{X};A)$  is the submodule of motives of *pure virtual rank* k. Roughly speaking, having virtual rank k means having a pole of order at most k at  $\mathbb{L} = 1$  after motivic integration. We omit the precise definition here, which is complicated.

When  $\mathscr{X}$  is quasi-compact, the image of the map  $\int_{\mathscr{X}} (-) : \mathbb{M}^{(\leq k)}(\mathscr{X}; A) \to \hat{\mathbb{M}}(K; A)$  lies in the subspace  $(\mathbb{L}-1)^{-k} \cdot \hat{\mathbb{M}}^{\operatorname{reg}}(K; A) \subset \hat{\mathbb{M}}(K; A)$ , where  $\mathbb{M}^{(\leq k)} = \mathbb{M}^{(0)} \oplus \cdots \oplus \mathbb{M}^{(k)}$ . In particular, there is an Euler characteristic integration map

$$\int_{\mathcal{X}} \left(\mathbb{L}-1\right)^k \cdot (-) \, d\chi = \chi \circ \int_{\mathcal{X}} \left(\mathbb{L}-1\right)^k \cdot (-) \colon \, \mathbb{M}^{(\leq k)}(\mathcal{X};A) \longrightarrow A \, .$$

#### 3.2 Motivic Hall algebras and modules

**3.2.1.** We introduce the *motivic Hall algebra* for a linear stack, originally defined by Joyce [29], which is an associative algebra structure on the ring of motives  $\mathbb{M}(\mathcal{X})$ .

For self-dual linear stacks, we show that the ring of motives  $\mathbb{M}(\mathcal{X}^{sd})$  is a module for the motivic Hall algebra  $\mathbb{M}(\mathcal{X})$ , which we call the *motivic Hall module*.

Hall modules have been constructed and studied for other flavours of Hall algebras, such as by Young [53–55] in the context of Ringel's [42; 43] Hall algebras and that of cohomological Hall algebras. A similar construction in the context of Joyce's [33; 34] vertex algebras is obtained by the author [11]. Another closely related work is DeHority and Latyntsev [18], who studied the relation between the cohomological version and the vertex algebra version.

**3.2.2. The motivic Hall algebra**. Let  $\mathscr{X}$  be a linear stack over *K*, with quasi-compact filtrations as in §2.2.4. Define an operation

$$*: \mathbb{M}(\mathcal{X}) \otimes \mathbb{M}(\mathcal{X}) \longrightarrow \mathbb{M}(\mathcal{X})$$

by the composition

$$\mathbb{M}(\mathcal{X}) \otimes \mathbb{M}(\mathcal{X}) \xrightarrow{\boxtimes} \mathbb{M}(\mathcal{X} \times \mathcal{X}) \xrightarrow{\operatorname{gr}^*} \mathbb{M}(\mathcal{X}^+) \xrightarrow{\operatorname{(ev_1)}_!} \mathbb{M}(\mathcal{X}) ,$$

where  $\mathscr{X}^+$  denotes the disjoint union of the stacks of filtrations  $\mathscr{X}^+_{\alpha_1,\alpha_2}$  for all  $\alpha_1, \alpha_2 \in \pi_0(\mathscr{X})$ . Roughly speaking, for motives  $a, b \in \mathbb{M}(\mathscr{X})$ , the product  $a * b \in \mathbb{M}(\mathscr{X})$  parametrizes all possible extensions of objects parametrized by *a* and *b*, respectively.

We will see in Theorem 3.2.4 that the product \* is associative, and that it has a unit element  $[\{0\}] \in \mathbb{M}(\mathcal{X})$ , which is the motive of the component  $\{0\} \subset \mathcal{X}$ . This defines an associative algebra structure on  $\mathbb{M}(\mathcal{X})$ , called the *motivic Hall algebra* of  $\mathcal{X}$ .

**3.2.3. The motivic Hall module**. Now, let  $\mathscr{X}$  be a self-dual linear stack over *K*, with quasicompact filtrations as in §2.2.4. Define an operation

$$\diamond \colon \mathbb{M}(\mathscr{X}) \otimes \mathbb{M}(\mathscr{X}^{\mathrm{sd}}) \longrightarrow \mathbb{M}(\mathscr{X}^{\mathrm{sd}})$$

by the composition

$$\mathbb{M}(\mathscr{X}) \otimes \mathbb{M}(\mathscr{X}^{\mathrm{sd}}) \xrightarrow{\boxtimes} \mathbb{M}(\mathscr{X} \times \mathscr{X}^{\mathrm{sd}}) \xrightarrow{\mathrm{gr}^*} \mathbb{M}(\mathscr{X}^{\mathrm{sd},+}) \xrightarrow{(\mathrm{ev}_1)_!} \mathbb{M}(\mathscr{X}^{\mathrm{sd}}),$$

where  $\mathscr{X}^{\mathrm{sd},+}$  denotes the disjoint union of the stacks of filtrations  $\mathscr{X}^{+}_{\alpha,\theta}$  for all  $\alpha \in \pi_{0}(\mathscr{X})$  and  $\theta \in \pi_{0}(\mathscr{X}^{\mathrm{sd}})$ .

Again, roughly speaking, for motives  $a \in \mathbb{M}(\mathcal{X})$  and  $b \in \mathbb{M}(\mathcal{X}^{sd})$ , the product  $a \diamond b \in \mathbb{M}(\mathcal{X}^{sd})$  parametrizes the total objects of all possible three-step self-dual filtrations, as in §2.1.6, whose graded pieces are parametrized by a, b, and  $a^{\vee}$ , respectively,

We will prove in Theorem 3.2.4 that the product  $\diamond$  establishes  $\mathbb{M}(\mathcal{X}^{sd})$  as a left module for the motivic Hall algebra  $\mathbb{M}(\mathcal{X})$ . This is called the *motivic Hall module* of  $\mathcal{X}$ .

#### **3.2.4. Theorem.** Let $\mathcal{X}$ be a linear stack over K, with quasi-compact filtrations.

(i) Recall the operation \* defined in §3.2.2. Then for any  $a, b, c \in \mathbb{M}(\mathcal{X})$ , we have

$$[\{0\}] * a = a = a * [\{0\}], \qquad (3.2.4.1)$$

$$(a * b) * c = a * (b * c), \qquad (3.2.4.2)$$

where  $[\{0\}] \in \mathbb{M}(\mathcal{X})$  is the motive of the component  $\{0\} \subset \mathcal{X}$ .

(ii) Suppose that X is equipped with a self-dual structure. Consider the involution (−)<sup>∨</sup> on M(X) induced by the involution of X, and the operation ◊ defined in §3.2.3. Then for any a, b ∈ M(X) and c ∈ M(X<sup>sd</sup>), we have

$$a^{\vee} * b^{\vee} = (b * a)^{\vee},$$
 (3.2.4.3)

$$\left[ \left\{ 0 \right\} \right] \diamond c = c , \qquad (3.2.4.4)$$

$$a \diamond (b \diamond c) = (a \ast b) \diamond c . \tag{3.2.4.5}$$

**Proof.** For (3.2.4.1), it is enough to show that for any  $\alpha \in \pi_0(\mathcal{X})$ , the morphisms  $\mathcal{X}^+_{0,\alpha} \to \mathcal{X}_{\alpha}$  and  $\mathcal{X}^+_{\alpha,0} \to \mathcal{X}_{\alpha}$  are isomorphisms, which follows from the descriptions in §2.2.4.

For (3.2.4.2), we may assume that  $a \in \mathbb{M}(\mathcal{X}_{\alpha_1})$ ,  $b \in \mathbb{M}(\mathcal{X}_{\alpha_2})$ , and  $c \in \mathbb{M}(\mathcal{X}_{\alpha_3})$ , for some  $\alpha_1, \alpha_2, \alpha_3 \in \pi_0(\mathcal{X})$ . Applying the base change formula (3.1.3.2) to the pullback squares in the

diagrams

we see that both sides of (3.2.4.2) are equal to  $(ev_1)_! \circ gr^*(a \boxtimes b \boxtimes c)$ , where gr and  $ev_1$  are the outer compositions in both diagrams in (3.2.4.6). These diagrams are special cases of the *associativity theorem* of the author et al. [14, §6.3], as explained in [14, §7.1.7].

The relation (3.2.4.3) follows from the commutativity of the diagram

$$\begin{aligned}
\mathcal{X}_{\alpha_{1}} \times \mathcal{X}_{\alpha_{2}} & \stackrel{\mathrm{gr}}{\longleftarrow} & \mathcal{X}_{\alpha_{1},\alpha_{2}}^{+} & \stackrel{\mathrm{ev}_{1}}{\longrightarrow} & \mathcal{X}_{\alpha_{1}+\alpha_{2}} \\
\stackrel{(-)^{\vee}}{\longrightarrow} & \stackrel{(-)^{\vee}}{\searrow} & \stackrel{(-)^{\vee}}{\longrightarrow} & \stackrel{(-)^{\vee}}{\searrow} \\
\mathcal{X}_{\alpha_{2}^{\vee}} \times \mathcal{X}_{\alpha_{1}^{\vee}} & \stackrel{\mathrm{gr}}{\longleftarrow} & \mathcal{X}_{\alpha_{2}^{\vee},\alpha_{1}^{\vee}}^{+} & \stackrel{\mathrm{ev}_{1}}{\longrightarrow} & \mathcal{X}_{\alpha_{2}^{\vee}+\alpha_{1}^{\vee}},
\end{aligned}$$
(3.2.4.7)

where  $\alpha_1, \alpha_2 \in \pi_0(\mathcal{X})$ , and the middle vertical isomorphism is given by the  $\mathbb{Z}_2$ -action on Filt( $\mathcal{X}$ ).

The relation (3.2.4.4) follows from the isomorphism  $\mathscr{X}_{0,\theta}^{\mathrm{sd},+} \xrightarrow{\sim} \mathscr{X}_{\theta}^{\mathrm{sd}}$  for  $\theta \in \pi_0(\mathscr{X}^{\mathrm{sd}})$ . For (3.2.4.5), we have similar diagrams

where the pullback squares follow from the associativity theorem of the author et al. [14, §6.3]. Alternatively, these diagrams can be obtained by taking  $\mathbb{Z}_2$ -fixed loci in pullback diagrams analogous to (3.2.4.6) for 5-step filtrations. The relation (3.2.4.5) then follows from applying the base change formula (3.1.3.2) to these diagrams.

### 4 Invariants

In this section, we present the definition of *orthosymplectic DT invariants*, as a special case of the *intrinsic DT invariants* of the author, Ibáñez Núñez, and Kinjo [15], which is the main construction of this paper. The new input here, compared to the cited work, is the choice of coefficients, or *stability measures* in the sense of [15], when defining the epsilon motives in the orthosymplectic setting. We also use the motivic Hall algebra and module to make the construction more explicit in our setting.

#### 4.1 Epsilon motives

4.1.1. We define *epsilon motives* for linear and self-dual linear stacks, following Joyce [32] in the linear case and the construction of the author, Ibáñez Núñez, and Kinjo [15] for general algebraic stacks. These are elements of the rings of motives  $\mathbb{M}(\mathcal{X};\mathbb{Q})$  and  $\mathbb{M}(\mathcal{X}^{sd};\mathbb{Q})$ , depending on a stability condition  $\tau$ , and are obtained from motives of semistable loci,  $[\mathcal{X}^{ss}_{\alpha}(\tau)]$  and  $[\mathcal{X}^{sd,ss}_{\theta}(\tau)]$ , by removing certain parts of the strictly semistable locus. The purpose of doing this step is so that the *no-pole theorem*, Theorem 4.1.6, holds, allowing us to take the Euler characteristics of epsilon motives, which will then be used to define DT invariants.

Throughout, we assume that  $\mathcal{X}$  is a linear stack over *K*, with quasi-compact filtrations as in §2.2.4.

**4.1.2.** The linear case. Let  $\tau$  be a permissible stability condition on  $\mathscr{X}$ . Following Joyce [32], for each class  $\alpha \in \pi_0(\mathscr{X}) \setminus \{0\}$ , define the *epsilon motive*  $\epsilon_{\alpha}(\tau) \in \mathbb{M}(\mathscr{X}_{\alpha}; \mathbb{Q})$  by the formula

$$\epsilon_{\alpha}(\tau) = \sum_{\substack{n > 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathcal{X}) \setminus \{0\}:\\ \alpha = \alpha_{1} + \dots + \alpha_{n},\\ \tau(\alpha_{1}) = \dots = \tau(\alpha_{n})}} \frac{(-1)^{n-1}}{n} \cdot \left[\mathcal{X}_{\alpha_{1}}^{ss}(\tau)\right] * \dots * \left[\mathcal{X}_{\alpha_{n}}^{ss}(\tau)\right], \qquad (4.1.2.1)$$

where \* denotes multiplication in the motivic Hall algebra  $\mathbb{M}(\mathcal{X}; \mathbb{Q})$ . By Lemma 2.3.7, only finitely many terms in the sum are non-zero. Note that  $\epsilon_{\alpha}(\tau)$  is supported on  $\mathcal{X}_{\alpha}^{ss}(\tau)$ .

Formally inverting the formula (4.1.2.1), we obtain the relation

$$\left[\mathscr{X}_{\alpha}^{\mathrm{ss}}(\tau)\right] = \sum_{\substack{n > 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathscr{X}) \setminus \{0\}:\\ \alpha = \alpha_{1} + \dots + \alpha_{n},\\ \tau(\alpha_{1}) = \dots = \tau(\alpha_{n})}} \frac{1}{n!} \cdot \epsilon_{\alpha_{1}}(\tau) * \dots * \epsilon_{\alpha_{n}}(\tau) \ . \tag{4.1.2.2}$$

The relation between the coefficients  $(-1)^{n-1}/n$  and 1/n! are explained in §4.1.4 below.

One can also combine (4.1.2.2) with the relation

$$\begin{bmatrix} \mathscr{X}_{\alpha} \end{bmatrix} = \sum_{\substack{n > 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathscr{X}) \setminus \{0\}:\\ \alpha = \alpha_{1} + \dots + \alpha_{n},\\ \tau(\alpha_{1}) > \dots > \tau(\alpha_{n})}} \begin{bmatrix} \mathscr{X}_{\alpha_{1}}^{ss}(\tau) \end{bmatrix} * \dots * \begin{bmatrix} \mathscr{X}_{\alpha_{n}}^{ss}(\tau) \end{bmatrix},$$
(4.1.2.3)

which comes from the  $\Theta$ -stratification of  $\mathcal{X}$ , and can be an infinite but locally finite sum, giving the formula

$$\left[\mathscr{X}_{\alpha}\right] = \sum_{\substack{n > 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathscr{X}) \setminus \{0\}:\\ \alpha = \alpha_{1} + \dots + \alpha_{n},\\ \tau(\alpha_{1}) \ge \dots \ge \tau(\alpha_{n})}} \frac{1}{|W_{\alpha_{1}, \dots, \alpha_{n}}(\tau)|} \cdot \epsilon_{\alpha_{1}}(\tau) \ast \dots \ast \epsilon_{\alpha_{n}}(\tau) , \qquad (4.1.2.4)$$

where  $W_{\alpha_1,...,\alpha_n}(\tau)$  denotes the group of permutations  $\sigma$  of  $\{1,...,n\}$  such that  $\tau(\alpha_{\sigma(1)}) \ge \cdots \ge \tau(\alpha_{\sigma(n)})$ . This can be taken as an alternative definition of the invariants  $\epsilon_{\alpha}(\tau)$ , that is, they are

the unique set of motives such that (4.1.2.4) holds for all  $\alpha$ .

One can interpret (4.1.2.4) as considering a generalized version of HN filtrations, where the slopes of the quotients are non-increasing rather than strictly decreasing, and the sum is averaged over all possible orderings satisfying the non-increasing condition.

The motive  $\epsilon_{\alpha}(\tau)$  agrees with the motive denoted by  $\epsilon_{\mathcal{X}_{\alpha}}^{(1)}(\mu_{\tau})$  in [15, §5.2], where  $\mu_{\tau}$  is the *stability measure* associated to the stability condition  $\tau$ , as explained in [15, Example 4.1.7].

**4.1.3.** The self-dual case. Suppose that  $\mathcal{X}$  is equipped with a self-dual structure, and let  $\tau$  be a permissible self-dual stability condition on  $\mathcal{X}$ .

For each class  $\theta \in \pi_0(\mathcal{X}^{sd})$ , define the *epsilon motive*  $\epsilon_{\theta}^{sd}(\tau) \in \mathbb{M}(\mathcal{X}_{\theta}^{sd}; \mathbb{Q})$  by the formula

$$\epsilon_{\theta}^{\mathrm{sd}}(\tau) = \sum_{\substack{n \ge 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathcal{X}) \setminus \{0\}, \ \rho \in \pi_{0}(\mathcal{X}^{\mathrm{ss}}) \\ \theta = \alpha_{1} + \alpha_{1}^{\vee} + \dots + \alpha_{n} + \alpha_{n}^{\vee} + \rho, \\ \tau(\alpha_{1}) = \dots = \tau(\alpha_{n}) = 0}} \left( \frac{-1/2}{n} \right) \cdot \left[ \mathcal{X}_{\alpha_{1}}^{\mathrm{ss}}(\tau) \right] \diamond \cdots \diamond \left[ \mathcal{X}_{\alpha_{n}}^{\mathrm{ss}}(\tau) \right] \diamond \left[ \mathcal{X}_{\rho}^{\mathrm{sd},\mathrm{ss}}(\tau) \right],$$
(4.1.3.1)

where  $\diamond$  denotes the multiplication for the motivic Hall module, the notation  $\alpha_i + \alpha_i^{\vee}$  is from §2.2.5, and  $\binom{-1/2}{n}$  is the binomial coefficient. The sum only contains finitely many non-zero terms, and  $\epsilon_{\theta}^{\text{sd}}(\tau)$  is supported on the semistable locus  $\mathcal{X}_{\theta}^{\text{sd,ss}}(\tau) \subset \mathcal{X}_{\theta}^{\text{sd}}$ .

Formally inverting the formula (4.1.3.1), we obtain the relation

$$\left[\mathscr{X}_{\theta}^{\mathrm{sd},\mathrm{ss}}(\tau)\right] = \sum_{\substack{n \ge 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathscr{X}) \setminus \{0\}, \ \rho \in \pi_{0}(\mathscr{X}^{\mathrm{sd}}):\\ \theta = \alpha_{1} + \alpha_{1}^{\vee} + \dots + \alpha_{n} + \alpha_{n}^{\vee} + \rho,\\ \tau(\alpha_{1}) = \dots = \tau(\alpha_{n}) = 0}} \frac{1}{2^{n} n!} \cdot \epsilon_{\alpha_{1}}(\tau) \diamond \cdots \diamond \epsilon_{\alpha_{n}}(\tau) \diamond \epsilon_{\rho}^{\mathrm{sd}}(\tau) , \qquad (4.1.3.2)$$

which we explain further in §4.1.4. This can be combined with the relation

$$\begin{bmatrix} \mathcal{X}_{\theta}^{\mathrm{sd}} \end{bmatrix} = \sum_{\substack{n \ge 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathcal{X}) \setminus \{0\}, \ \rho \in \pi_{0}(\mathcal{X}^{\mathrm{sd}}): \\ \theta = \alpha_{1} + \alpha_{1}^{\vee} + \dots + \alpha_{n} + \alpha_{n}^{\vee} + \rho, \\ \tau(\alpha_{1}) > \dots > \tau(\alpha_{n}) > 0} \begin{bmatrix} \mathcal{X}_{\alpha_{1}}^{\mathrm{ss}}(\tau) \end{bmatrix} \diamond \begin{bmatrix} \mathcal{X}_{\rho}^{\mathrm{sd}}(\tau) \end{bmatrix}$$
(4.1.3.3)

from the  $\Theta$ -stratification of  $\mathscr{X}^{sd}$ , together with (4.1.2.2), to obtain the formula

$$\left[\mathcal{X}_{\theta}^{\mathrm{sd}}\right] = \sum_{\substack{n \ge 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathcal{X}) \setminus \{0\}, \ \rho \in \pi_{0}(\mathcal{X}^{\mathrm{sd}}):\\ \theta = \alpha_{1} + \alpha_{1}^{\vee} + \dots + \alpha_{n} + \alpha_{n}^{\vee} + \rho, \\ \tau(\alpha_{1}) \ge \dots \ge \tau(\alpha_{n}) \ge 0} \frac{1}{|W_{\alpha_{1},\dots,\alpha_{n}}^{\mathrm{sd}}(\tau)|} \cdot \epsilon_{\alpha_{1}}(\tau) \diamond \dots \diamond \epsilon_{\alpha_{n}}(\tau) \diamond \epsilon_{\rho}^{\mathrm{sd}}(\tau) , \qquad (4.1.3.4)$$

where  $W^{\text{sd}}_{\alpha_1,...,\alpha_n}(\tau)$  is the group of permutations  $\sigma$  of  $\{1,...,n,n^{\vee},...,1^{\vee}\}$ , such that  $\sigma(i)^{\vee} = \sigma(i^{\vee})$  for all *i*, where we set  $(i^{\vee})^{\vee} = i$ , satisfying the non-increasing condition  $\tau(\alpha_{\sigma(1)}) \ge \cdots \ge \tau(\alpha_{\sigma(n)}) \ge 0$ , where we set  $\alpha_{i^{\vee}} = \alpha_{i}^{\vee}$ . For example, we have  $|W^{\text{sd}}_{\alpha_1,...,\alpha_n}(\tau)| = 2^n n!$  if  $\tau(\alpha_1) = \cdots = \tau(\alpha_n) = 0$ .

The coefficients  $1/|W^{\rm sd}_{\alpha_1,\ldots,\alpha_n}(\tau)|$  in (4.1.3.4) can be seen as defining a *stability measure*  $\mu^{\rm sd}_{\tau}$  on  $\mathcal{X}^{\rm sd}_{\theta}$ , in the sense of [15]. The motive  $\epsilon^{\rm sd}_{\theta}(\tau)$  agrees with the motive  $\epsilon^{(0)}_{\mathcal{X}^{\rm sd}_{\theta}}(\mu^{\rm sd}_{\tau})$  in [15, §5.2].

**4.1.4. Explanations of the coefficients**. The relations between the coefficients in (4.1.2.1), (4.1.2.2), (4.1.3.1), and (4.1.3.2), can be seen more directly by setting

$$\begin{split} &\delta(\tau;t) = \left[\{0\}\right] + \sum_{\substack{\alpha \in \pi_0(\mathcal{X}) \setminus \{0\}:\\ \tau(\alpha) = t}} \left[\mathcal{X}_{\alpha}^{\rm ss}(\tau)\right], \qquad \delta^{\rm sd}(\tau) = \sum_{\substack{\theta \in \pi_0(\mathcal{X}^{\rm sd})\\ \theta \in \pi_0(\mathcal{X}^{\rm sd})}} \left[\mathcal{X}_{\theta}^{\rm sd,ss}(\tau)\right] \\ &\epsilon(\tau;t) = \sum_{\substack{\alpha \in \pi_0(\mathcal{X}) \setminus \{0\}:\\ \tau(\alpha) = t}} \epsilon_{\alpha}(\tau), \qquad \qquad \epsilon^{\rm sd}(\tau) = \sum_{\substack{\theta \in \pi_0(\mathcal{X}^{\rm sd})\\ \theta \in \pi_0(\mathcal{X}^{\rm sd})}} \epsilon_{\theta}^{\rm sd}(\tau), \end{split}$$

as motives on  $\mathcal{X}$  or  $\mathcal{X}^{sd}$ , where  $t \in T$ , so that these relations can be rewritten as

$$\begin{split} \epsilon(\tau;t) &= \log \delta(\tau;t) , \qquad \epsilon^{\rm sd}(\tau) = \delta(\tau;0)^{-1/2} \diamond \delta^{\rm sd}(\tau) , \\ \delta(\tau;t) &= \exp \epsilon(\tau;t) , \qquad \delta^{\rm sd}(\tau) = \exp \Bigl( \frac{1}{2} \epsilon(\tau;0) \Bigr) \diamond \epsilon^{\rm sd}(\tau) , \end{split}$$

where we take formal power series using the product in the motivic Hall algebra.

The coefficients  $(-1)^{n-1}/n$  and  $\binom{-1/2}{n}$  in (4.1.2.1) and (4.1.3.1) are determined by the coefficients 1/n! and  $1/(2^n n!)$  in (4.1.2.2) and (4.1.3.2) in this way. They are the unique choice of coefficients only depending on n, such that the *no-pole theorem*, Theorem 4.1.6, holds for the epsilon motives. The rough reason for this is that they ensure the combinatorial descriptions of the coefficients  $1/|W_{\alpha_1,...,\alpha_n}(\tau)|$  and  $1/|W_{\alpha_1,...,\alpha_n}^{sd}(\tau)|$  in (4.1.2.4) and (4.1.3.4), and from the viewpoint of [15], the no-pole theorem corresponds to the property that these coefficients sum up to 1 for all permutations  $\sigma$  as described for each of them, for fixed classes  $\alpha_i$ .

4.1.5. Remark on permissibility. In the situations above, the permissibility of the stability condition  $\tau$  implies that the stability measures  $\mu_{\tau}$  and  $\mu_{\tau}^{sd}$  are permissible in the sense of [15, §4.1.4], which follows from [15, Lemma 5.4.8].

**4.1.6.** The no-pole theorem. A key property of the epsilon motives is the *no-pole theorem*, which states that they have pure virtual ranks in the sense of §3.1.6. This will allow us to define numerical invariants, including DT invariants, by taking their Euler characteristics.

**Theorem**. Let  $\mathcal{X}$  be a linear stack over K, with quasi-compact filtrations.

- (i) For any permissible stability condition  $\tau$  on  $\mathcal{X}$ , and any  $\alpha \in \pi_0(\mathcal{X}) \setminus \{0\}$ , the motive  $\epsilon_{\alpha}(\tau)$  has pure virtual rank 1.
- (ii) If  $\mathscr{X}$  is equipped with a self-dual structure, then for any permissible self-dual stability condition  $\tau$  on  $\mathscr{X}$ , and any  $\theta \in \pi_0(\mathscr{X}^{sd})$ , the motive  $\epsilon_{\theta}^{sd}(\tau)$  has pure virtual rank 0.

These are special cases of the general no-pole theorem for intrinsic DT invariants in [15, Theorem 5.3.7], and we refer to the cited work for the proof. The linear case (i) was originally proved by Joyce [30, Theorem 8.7], under a slightly different setting. The self-dual case (ii) was originally proved in an earlier version of this paper, [10, Appendix E], under another slightly different setting.

#### 4.2 DT invariants

**4.2.1.** We now turn to the definition of *DT invariants* for linear and self-dual linear stacks, the main construction of this paper. The linear case was first due to Joyce and Song [35] and Kontsevich and Soibelman [37], and the self-dual case was first constructed by the author in an earlier version of this paper [10]. Here, we continue to follow the general construction of the author, Ibáñez Núñez, and Kinjo [15], specializing it to the self-dual linear case.

Throughout this section, we assume that the base field *K* is algebraically closed and has characteristic zero. We work with (-1)-*shifted symplectic stacks* over *K* in the sense of Pantev, Toën, Vaquié, and Vezzosi [40], which are derived algebraic stacks locally finitely presented over *K*, equipped with a (-1)-shifted symplectic form  $\omega$ .

**4.2.2. Local structure.** Following the author [12, §2.2.4], we introduce the following local conditions on algebraic stacks.

A stack is *étale* (or *Nisnevich*) *locally a quotient stack*, if it admits a representable étale (or Nisnevich) cover by quotient stacks of the form U/GL(n), with U an algebraic space.

A stack is *étale* (or *Nisnevich*) *locally fundamental*, if it admits a representable étale (or Nisnevich) cover by quotient stacks of the form U/GL(n), with U an affine scheme.

These conditions are preserved by taking  $\mathbb{Z}_2$ -fixed points under the assumptions in §1.1.11, by Lemma 2.2.9.

4.2.3. Derived linear stacks. Following the author et al. [13, §2.4.6], define a *derived linear stack* over *K* to be a derived algebraic stack  $\mathcal{X}$ , locally finitely presented over *K*, equipped with a monoid structure  $\oplus$  and a compatible  $*/\mathbb{G}_m$ -action  $\odot$ , such that the isomorphism (2.2.3.1) holds without taking the classical truncations, where we use the derived stack of graded points  $\operatorname{Grad}(\mathcal{X})$ , defined as the derived mapping stack from  $*/\mathbb{G}_m$  to  $\mathcal{X}$ .

As in [13, §3.1.7], define a (-1)-shifted symplectic linear stack over K to be a derived linear stack  $\mathcal{X}$  as above, equipped with a (-1)-shifted symplectic form  $\omega$ , such that there exists an equivalence  $\oplus^*(\omega) \simeq \omega \boxplus \omega$  on  $\mathcal{X} \times \mathcal{X}$ , where we do not require extra coherence conditions.

We further assume that the classical truncation  $\mathscr{X}_{cl}$  of  $\mathscr{X}$  satisfies the conditions in §1.1.11, has quasi-compact filtrations, and is étale locally a quotient stack.

We will often denote  $\mathscr{X}_{cl}$  simply by  $\mathscr{X}$ .

**4.2.4.** The linear case. Let  $\tau$  be a permissible stability condition on  $\mathscr{X}$ . Following the construction of Joyce and Song [35, Definition 5.15], but adapting it to our setting of linear stacks, for a class  $\alpha \in \pi_0(\mathscr{X}) \setminus \{0\}$ , define the *DT invariant*  $DT_{\alpha}(\tau) \in \mathbb{Q}$  by the formula

$$DT_{\alpha}(\tau) = \int_{\mathcal{X}_{\alpha}} (1 - \mathbb{L}) \cdot \epsilon_{\alpha}(\tau) \cdot v_{\mathcal{X}} d\chi, \qquad (4.2.4.1)$$

where the notation  $\int (-) d\chi$  is defined in §3.1.6, and  $v_{\mathcal{X}}$  is the *Behrend function* of  $\mathcal{X}$ , which is a constructible function on  $\mathcal{X}$  only depending on  $\mathcal{X}_{cl}$ , as in Joyce and Song [35, §4.1] or the

author [12, §2.5.6], and originally due to Behrend [5] for Deligne-Mumford stacks.

This integral is well-defined since  $\epsilon_{\alpha}(\tau)$  is supported on the semistable locus  $\mathscr{X}_{\alpha}^{ss}(\tau)$ , which is quasi-compact, and by the no-pole theorem, Theorem 4.1.6 (i).

**4.2.5. The self-dual case**. Assume further that  $\mathscr{X}$  is equipped with a  $\mathbb{Z}_2$ -action, preserving the (-1)-shifted symplectic form, such that the induced  $\mathbb{Z}_2$ -action on  $\mathscr{X}$  establishes it as a self-dual linear stack.

Let  $\tau$  be a permissible self-dual stability condition on  $\mathscr{X}$ . For a class  $\theta \in \pi_0(\mathscr{X}^{sd})$ , define the *self-dual DT invariant*  $DT^{sd}_{\theta}(\tau) \in \mathbb{Q}$  by the formula

$$DT_{\theta}^{\rm sd}(\tau) = \int_{\mathcal{X}_{\theta}^{\rm sd}} \epsilon_{\theta}^{\rm sd}(\tau) \cdot v_{\mathcal{X}^{\rm sd}} \, d\chi \;. \tag{4.2.5.1}$$

Again, this is well-defined by the fact that  $\epsilon_{\theta}^{sd}(\tau)$  is supported on  $\mathscr{X}_{\theta}^{sd,ss}(\tau)$ , which is quasicompact, and by the no-pole theorem, Theorem 4.1.6 (ii).

This is one of the main constructions of this paper, and is a special case of the intrinsic DT invariants in [15, §6.1] for the stability measure  $\mu_{\tau}^{sd}$  described in §4.1.3.

**4.2.6.** For smooth stacks. Let  $\mathscr{X}$  be a classical smooth linear stack which is étale locally a quotient stack, and consider its (-1)-shifted cotangent stack  $T^*[-1]\mathscr{X}$ , which has a canonical (-1)-shifted symplectic structure, making it a (-1)-shifted symplectic linear stack. We have  $(T^*[-1]\mathscr{X})_{cl} \simeq \mathscr{X}$ . If  $\mathscr{X}$  is equipped with a self-dual structure, then the fixed locus  $\mathscr{X}^{sd}$  is also smooth, and  $(T^*[-1]\mathscr{X})^{sd} \simeq T^*[-1]\mathscr{X}^{sd}$ .

In this case, we have  $v_{\mathcal{X}} = (-1)^{\dim \mathcal{X}}$  and  $v_{\mathcal{X}^{sd}} = (-1)^{\dim \mathcal{X}^{sd}}$ , and (4.2.4.1) - (4.2.5.1) become

$$DT_{\alpha}(\tau) = (-1)^{\dim \mathscr{X}_{\alpha}} \cdot \int_{\mathscr{X}_{\alpha}} (1 - \mathbb{L}) \cdot \epsilon_{\alpha}(\tau) \, d\chi \,, \qquad (4.2.6.1)$$

$$\mathrm{DT}_{\theta}^{\mathrm{sd}}(\tau) = (-1)^{\dim \mathcal{X}_{\theta}^{\mathrm{sd}}} \cdot \int_{\mathcal{X}_{\theta}^{\mathrm{sd}}} \epsilon_{\theta}^{\mathrm{sd}}(\tau) \, d\chi \,. \tag{4.2.6.2}$$

The invariants  $DT_{\alpha}(\tau)$  are essentially the same as those defined by Joyce [32, §6.2], denoted by  $J^{\alpha}(\tau)^{\Omega}$  there, while the invariants  $DT_{\theta}^{sd}(\tau)$  are new.

Note that the formulae (4.2.6.1)–(4.2.6.2) also make sense for smooth (self-dual) linear stacks over an arbitrary base field *K*, allowing us to also define DT invariants in this case.

#### 4.3 Motivic DT invariants

**4.3.1.** We also introduce motivic enhancements of the linear and self-dual DT invariants defined above, following the formalism of the author, Ibáñez Núñez, and Kinjo [15, §6.2]. These generalize the construction of Kontsevich and Soibelman [37] in the linear case.

**4.3.2. Monodromic motives.** For a stack  $\mathscr{X}$  over K and a commutative ring A, we have the ring of *monodromic motives*, denoted by  $\hat{\mathbb{M}}^{\text{mon}}(\mathscr{X}; A)$ . It is similar to  $\hat{\mathbb{M}}(\mathscr{X}; A)$  defined in §3.1.4, but its elements have the additional structure of a monodromy action. See [15, §6.2.2] for details.

When *A* contains  $\mathbb{Q}$ , there is an Euler characteristic map  $\chi : \hat{\mathbb{M}}^{\text{mon,reg}}(K;A) \to A$ , where  $\hat{\mathbb{M}}^{\text{mon,reg}}(K;A) \subset \hat{\mathbb{M}}^{\text{mon}}(K;A)$  is the subspace of elements that are regular at  $\mathbb{L} = 1$ , defined similarly to §3.1.5.

There is an element  $\mathbb{L}^{1/2} \in \widehat{\mathbb{M}}^{\mathrm{mon,reg}}(K;A)$  satisfying  $(\mathbb{L}^{1/2})^2 = \mathbb{L}$  and  $\chi(\mathbb{L}^{1/2}) = -1$ .

**4.3.3.** Orientations. For a (-1)-shifted symplectic stack  $\mathscr{X}$  over K, the *canonical bundle* of  $\mathscr{X}$  is the determinant line bundle of its cotangent complex,  $K_{\mathscr{X}} = \det \mathbb{L}_{\mathscr{X}}$ .

An *orientation* of  $\mathscr{X}$  is a line bundle  $K_{\mathscr{X}}^{1/2}$  on  $\mathscr{X}$ , with an isomorphism  $o_{\mathscr{X}} : (K_{\mathscr{X}}^{1/2})^{\otimes 2} \xrightarrow{\sim} K_{\mathscr{X}}$ . We sometimes abbreviate the pair  $(K_{\mathscr{X}}^{1/2}, o_{\mathscr{X}})$  as  $o_{\mathscr{X}}$ .

Given such an orientation, if the classical truncation  $\mathscr{X}_{cl}$  is Nisnevich locally a quotient stack as in §4.2.2, then there is an element  $v_{\mathscr{X}}^{mot} \in \widehat{\mathbb{M}}^{mot}(\mathscr{X};\mathbb{Z})$ , as in the author [12, §2.5.4], called the *motivic Behrend function*, originally constructed by Bussi, Joyce, and Meinhardt [17] and Ben-Bassat et al. [6].

**4.3.4.** Orientation data. By the author [12, Theorem 3.1.6] or [13, §6.1.6], an orientation  $o_{\mathcal{X}}$  induces an orientation  $o_{\text{Grad}(\mathcal{X})}$  of  $\text{Grad}(\mathcal{X})$ . An orientation  $o_{\mathcal{X}}$  is called an *orientation data*, if it satisfies the following compatibility condition:

• Under the isomorphism (2.2.3.1), the induced orientation  $o_{\text{Grad}(\mathcal{X})}$  of  $\text{Grad}(\mathcal{X})$  agrees with the product orientations on the left-hand side.

By Joyce and Upmeier [36, Theorem 3.6], such an orientation data exists canonically on moduli stacks of coherent sheaves on Calabi–Yau threefolds.

4.3.5. Self-dual orientation data. Now, let  $\mathscr{X}$  be a *self-dual* (-1)-*shifted symplectic linear stack*, that is, a stack  $\mathscr{X}$  as in §4.3.4, equipped with a  $\mathbb{Z}_2$ -action preserving the symplectic form  $\omega$ , compatible with the monoid structure  $\oplus$  and inverting the  $*/\mathbb{G}_m$ -action  $\odot$ .

In this case, the fixed locus  $\mathscr{X}^{sd} = \mathscr{X}^{\mathbb{Z}_2}$  carries an induced (-1)-shifted symplectic structure. However, an orientation of  $\mathscr{X}$  does not naturally induce one on  $\mathscr{X}^{sd}$ .

We define a *self-dual orientation data* on  $\mathcal{X}$  to be a pair  $(o_{\mathcal{X}}, o_{\mathcal{X}^{sd}})$  of orientations of  $\mathcal{X}$  and  $\mathcal{X}^{sd}$ , respectively, satisfying the following conditions:

- (i)  $o_{\mathcal{X}}$  is an orientation data.
- (ii) Under the isomorphism (2.2.7.1), the induced orientation of  $\operatorname{Grad}(\mathscr{X}^{\operatorname{sd}})$  agrees with the product orientations on the right-hand side.

The author does not know if such a self-dual orientation data, or even an orientation, exists in the case of coherent sheaves on Calabi–Yau threefolds, which we will discuss in §6.4 below.

**4.3.6.** Motivic DT invariants. Let  $\mathscr{X}$  be a (-1)-shifted symplectic linear stack over K, equipped with an orientation data as in §4.3.4. Assume that its classical truncation  $\mathscr{X}_{cl}$  is Nisnevich locally a quotient stack, as in §4.2.2.

For a permissible stability condition  $\tau$  on  $\mathscr{X}$ , and a class  $\alpha \in \pi_0(\mathscr{X}) \setminus \{0\}$ , following the construction of Kontsevich and Soibelman [37], define the *motivic* DT *invariant*  $DT^{mot}_{\alpha}(\tau) \in \widehat{\mathbb{M}}^{mot}(K;\mathbb{Q})$  by the formula

$$DT^{mot}_{\alpha}(\tau) = \int_{\mathscr{X}_{\alpha}} \left( \mathbb{L}^{1/2} - \mathbb{L}^{-1/2} \right) \cdot \epsilon_{\alpha}(\tau) \cdot v_{\mathscr{X}}^{mot} , \qquad (4.3.6.1)$$

where  $v_{\mathcal{X}}^{\text{mot}}$  is the motivic Behrend function of  $\mathcal{X}$  defined in §4.3.3.

Now, suppose further that  $\mathscr{X}$  is equipped with a self-dual structure as in §4.3.5, together with a self-dual orientation data.

For a self-dual permissible stability condition  $\tau$  and a class  $\theta \in \pi_0(\mathcal{X}^{sd})$ , define the *self-dual motivic* DT *invariant*  $DT_{\theta}^{mot,sd}(\tau) \in \hat{\mathbb{M}}^{mot}(K; \mathbb{Q})$  by

$$DT_{\theta}^{\text{mot,sd}}(\tau) = \int_{\mathcal{X}_{\theta}^{\text{sd}}} \epsilon_{\theta}^{\text{sd}}(\tau) \cdot v_{\mathcal{X}^{\text{sd}}}^{\text{mot}} .$$
(4.3.6.2)

This is also a main construction of this paper, and is a special case of the intrinsic motivic DT invariants in [15, §6.2] for the stability measure  $\mu_{\tau}^{\text{sd}}$  described in §4.1.3.

4.3.7. For smooth stacks. Let  $\mathscr{X}$  be a linear stack which is smooth and Nisnevich locally a quotient stack, and consider its (-1)-shifted cotangent stack  $T^*[-1]\mathscr{X}$ , as in §4.2.6. It has a canonical (-1)-shifted symplectic linear structure and orientation data, and in the self-dual case, also a canonical self-dual orientation data.

The motivic Behrend function of  $\mathscr{X}$  is  $\nu_{\mathscr{X}}^{\text{mot}} = \mathbb{L}^{-\dim \mathscr{X}/2}$  by the author [12, Theorem 2.5.5], where dim  $\mathscr{X}$  refers to the dimension of the classical smooth stack  $\mathscr{X}$ . The formulae (4.3.6.1)–(4.3.6.2) can be simplified to

$$DT_{\alpha}^{\text{mot}}(\tau) = \frac{\mathbb{L}^{1/2} - \mathbb{L}^{-1/2}}{\mathbb{L}^{\dim \mathcal{X}_{\alpha}/2}} \cdot \int_{\mathcal{X}_{\alpha}} \epsilon_{\alpha}(\tau) , \qquad (4.3.7.1)$$

$$\mathrm{DT}_{\theta}^{\mathrm{mot,sd}}(\tau) = \mathbb{L}^{-\dim \mathcal{X}_{\theta}^{\mathrm{sd}}/2} \cdot \int_{\mathcal{X}_{\theta}^{\mathrm{sd}}} \epsilon_{\theta}^{\mathrm{sd}}(\tau) .$$
(4.3.7.2)

## 5 Wall-crossing

#### 5.1 Wall-crossing for epsilon motives

5.1.1. We now discuss how to relate the epsilon motives and DT invariants defined in §§4.1–4.3 when we change the stability condition  $\tau$ . These relations are called *wall-crossing formulae*. We first prove wall-crossing formulae for epsilon motives in Theorem 5.1.3, which we then use in §5.3 to obtain wall-crossing formulae for DT invariants.

Throughout, let  $\mathscr{X}$  be a self-dual linear stack with quasi-compact filtrations as in §2.2.4. Results in the linear case will not need the self-dual structure on  $\mathscr{X}$ , and we will indicate this when it is the case.

**5.1.2.** Dominance of stability conditions. For stability conditions  $\tau_0$ ,  $\tau$  on  $\mathcal{X}$ , following Joyce [30, Definition 4.10], we say that  $\tau_0$  dominates  $\tau$ , if  $\tau(\alpha_1) \leq \tau(\alpha_2)$  implies  $\tau_0(\alpha_1) \leq \tau_0(\alpha_2)$  for all  $\alpha_1, \alpha_2 \in \pi_0(\mathcal{X}) \setminus \{0\}.$ 

In this case, the  $\Theta$ -stratification of  $\mathcal X$  given by  $\tau$  refines the one given by  $\tau_0$ , and in particular, we have  $\mathscr{X}^{ss}_{\alpha}(\tau) \subset \mathscr{X}^{ss}_{\alpha}(\tau_0)$  for all  $\alpha \in \pi_0(\mathscr{X}) \setminus \{0\}$ .

For example, every stability condition is dominated by the trivial stability condition.

**5.1.3. Theorem.** Let  $\tau_+, \tau_-, \tau_0$  be permissible self-dual stability conditions on  $\mathcal{X}$ , with  $\tau_0$  dominating both  $\tau_+$  and  $\tau_-$ . Then for any  $\alpha \in \pi_0(\mathcal{X})$  and  $\theta \in \pi_0(\mathcal{X}^{sd})$ , we have the relations

$$\left[\mathcal{X}_{\alpha}^{\mathrm{ss}}(\tau_{-})\right] = \sum_{\substack{n \ge 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathcal{X}) \setminus \{0\}:\\ \alpha = \alpha_{1} + \dots + \alpha_{n}}} S(\alpha_{1}, \dots, \alpha_{n}; \tau_{+}, \tau_{-}) \cdot \left[\mathcal{X}_{\alpha_{1}}^{\mathrm{ss}}(\tau_{+})\right] * \dots * \left[\mathcal{X}_{\alpha_{n}}^{\mathrm{ss}}(\tau_{+})\right],$$
(5.1.3.1)

$$\left[\mathcal{X}_{\theta}^{\mathrm{sd},\mathrm{ss}}(\tau_{-})\right] = \sum_{\substack{n \ge 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathcal{X}) \setminus \{0\}, \ \rho \in \pi_{0}(\mathcal{X}^{\mathrm{sd}}):\\ \theta = \alpha_{1} + \alpha_{1}^{\vee} + \dots + \alpha_{n} + \alpha_{n}^{\vee} + \rho}} S^{\mathrm{sd}}(\alpha_{1}, \dots, \alpha_{n}; \tau_{+}, \tau_{-}) \cdot \left[\mathcal{X}_{\alpha_{1}}^{\mathrm{ss}}(\tau_{+})\right] \diamond \cdots \diamond \left[\mathcal{X}_{\alpha_{n}}^{\mathrm{sd}}(\tau_{+})\right] \diamond \left[\mathcal{X}_{\rho}^{\mathrm{sd},\mathrm{ss}}(\tau_{+})\right],$$
(5.1.3.2)

$$\epsilon_{\alpha}(\tau_{-}) = \sum_{\substack{n \ge 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathcal{X}) \setminus \{0\}:\\ \alpha = \alpha_{1} + \dots + \alpha_{n}}} U(\alpha_{1}, \dots, \alpha_{n}; \tau_{+}, \tau_{-}) \cdot \epsilon_{\alpha_{1}}(\tau_{+}) * \dots * \epsilon_{\alpha_{n}}(\tau_{+}) , \qquad (5.1.3.3)$$

$$\epsilon_{\theta}^{\mathrm{sd}}(\tau_{-}) = \sum_{\substack{n \ge 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathcal{X}) \setminus \{0\}, \ \rho \in \pi_{0}(\mathcal{X}^{\mathrm{sd}}):\\ \theta = \alpha_{1} + \alpha_{1}^{\vee} + \dots + \alpha_{n} + \alpha_{n}^{\vee} + \rho}} U^{\mathrm{sd}}(\alpha_{1}, \dots, \alpha_{n}; \tau_{+}, \tau_{-}) \cdot \epsilon_{\alpha_{1}}(\tau_{+}) \diamond \dots \diamond \epsilon_{\alpha_{n}}(\tau_{+}) \diamond \epsilon_{\rho}^{\mathrm{sd}}(\tau_{+}), \quad (5.1.3.4)$$

in  $\mathbb{M}(\mathscr{X}_{\alpha}; \mathbb{Q})$  and  $\mathbb{M}(\mathscr{X}_{\theta}^{sd}; \mathbb{Q})$ , where the sums are finite, and

$$S(\alpha_{1}, \dots, \alpha_{n}; \tau_{+}, \tau_{-}) = \prod_{i=1}^{n-1} \left\{ \begin{array}{ll} 1, & \tau_{+}(\alpha_{i}) > \tau_{+}(\alpha_{i+1}) \text{ and} \\ & \tau_{-}(\alpha_{1} + \dots + \alpha_{i}) \leqslant \tau_{-}(\alpha_{i+1} + \dots + \alpha_{n}) \\ -1, & \tau_{+}(\alpha_{i}) \leqslant \tau_{+}(\alpha_{i+1}) \text{ and} \\ & \tau_{-}(\alpha_{1} + \dots + \alpha_{i}) > \tau_{-}(\alpha_{i+1} + \dots + \alpha_{n}) \\ 0, & \text{otherwise} \end{array} \right\} , \quad (5.1.3.5)$$
$$S^{\text{sd}}(\alpha_{1}, \dots, \alpha_{n}; \tau_{+}, \tau_{-}) = \prod_{i=1}^{n} \left\{ \begin{array}{ll} 1, & \tau_{+}(\alpha_{i}) > \tau_{+}(\alpha_{i+1}) \text{ and } \tau_{-}(\alpha_{1} + \dots + \alpha_{i}) \leqslant 0 \\ -1, & \tau_{+}(\alpha_{i}) \leqslant \tau_{+}(\alpha_{i+1}) \text{ and } \tau_{-}(\alpha_{1} + \dots + \alpha_{i}) > 0 \\ 0, & \text{otherwise} \end{array} \right\} , \quad (5.1.3.6)$$

$$U(\alpha_{1}, \dots, \alpha_{n}; \tau_{+}, \tau_{-}) = \sum_{\substack{0 = a_{0} < \dots < a_{m} = n, \ 0 = b_{0} < \dots < b_{\ell} = m: \\ \text{Writing } \beta_{i} = \alpha_{a_{i-1}+1} + \dots + \alpha_{a_{i}} \text{ for } i = 1, \dots, m, \\ \text{and } \gamma_{i} = \beta_{b_{i-1}+1} + \dots + \beta_{b_{i}} \text{ for } i = 1, \dots, \ell, \\ \text{we have } \tau_{+}(\alpha_{j}) = \tau_{+}(\beta_{i}) \text{ for all } a_{i-1} < j \leq a_{i}, \end{cases}$$
(5.1.3.7)

we have  $\tau_+(\alpha_j) = \tau_+(\beta_i)$  for all  $a_{i-1} < j \le a_i$ , and  $\tau_{-}(\gamma_{i}) = \tau_{-}(\alpha_{1} + \cdots + \alpha_{n})$  for all  $i = 1, \dots, \ell$ 

$$U^{\rm sd}(\alpha_{1},\ldots,\alpha_{n};\tau_{+},\tau_{-}) = \sum_{\substack{0 = a_{0} < \cdots < a_{m} \leq n, \ 0 = b_{0} < \cdots < b_{\ell} \leq m: \\ \text{Writing } \beta_{i} = \alpha_{a_{i-1}+1} + \cdots + \alpha_{a_{i}} \text{ for } i = 1,\ldots,m, \\ \text{and } \gamma_{i} = \beta_{b_{i-1}+1} + \cdots + \beta_{b_{i}} \text{ for } i = 1,\ldots,m, \\ \text{we have } \tau_{+}(\alpha_{j}) = \tau_{+}(\beta_{i}) \text{ for all } a_{i-1} < j \leq a_{i}, \\ \tau_{+}(\alpha_{j}) = 0 \text{ for all } j > a_{m}, \\ \text{and } \tau_{-}(\gamma_{i}) = 0 \text{ for all } i = 1,\ldots,\ell \end{cases} \begin{pmatrix} m \\ \prod_{i=1}^{m} \frac{1}{(a_{i} - a_{i-1})!} \end{pmatrix} \cdot \frac{1}{2^{n-a_{m}} (n - a_{m})!} , \quad (5.1.3.8)$$

where we set  $\tau_{+}(\alpha_{n+1}) = 0$  in (5.1.3.6).

For (5.1.3.1) and (5.1.3.3), we do not need  $\mathcal{X}$  or  $\tau_+, \tau_-, \tau_0$  to be self-dual.

The formulae (5.1.3.1) and (5.1.3.3) were originally due to Joyce [32, Theorem 5.2], under a slightly different setting. The self-dual versions (5.1.3.2) and (5.1.3.4) are new.

The coefficients (5.1.3.5)–(5.1.3.8) are combinatorial, and are defined whenever  $\tau_{\pm}$  are maps from the set  $C = \{\alpha_i + \cdots + \alpha_j \mid 1 \leq i \leq j \leq n\}$  of symbolic sums to totally ordered sets  $T_{\pm}$ , such that  $\tau_+(\gamma_1) \leq \tau_+(\gamma_2)$  implies  $\tau_+(\gamma_1) \leq \tau_+(\gamma_1 + \gamma_2) \leq \tau_+(\gamma_2)$  whenever  $\gamma_1, \gamma_2, \gamma_1 + \gamma_2 \in C$ , and similarly for  $\tau_-$ . For (5.1.3.6) and (5.1.3.8), we also require distinguished elements  $0 \in T_{\pm}$ .

**Proof.** The  $\Theta$ -stratifications of  $\mathscr{X}^{ss}_{\alpha}(\tau_0)$  and  $\mathscr{X}^{sd,ss}_{\theta}(\tau_0)$  defined by  $\tau_+$  and  $\tau_-$  give the relations

$$\begin{bmatrix} \mathscr{X}_{\alpha}^{\mathrm{ss}}(\tau_{0}) \end{bmatrix} = \sum_{\substack{n > 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathscr{X}) \setminus \{0\}:\\ \alpha = \alpha_{1} + \dots + \alpha_{n},\\ \tau_{0}(\alpha_{1}) = \dots = \tau_{0}(\alpha_{n}),\\ \tau_{\pm}(\alpha_{1}) > \dots > \tau_{\pm}(\alpha_{n})} \qquad (5.1.3.9)$$

$$\begin{bmatrix} \mathscr{X}_{\theta}^{\mathrm{sd},\mathrm{ss}}(\tau_{0}) \end{bmatrix} = \sum_{\substack{n \ge 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathscr{X}) \setminus \{0\},\\ \theta = \alpha_{1} + \alpha_{1}^{\vee} + \dots + \alpha_{n} + \alpha_{n}^{\vee} + \rho,\\ \tau_{0}(\alpha_{1}) = \dots = \tau_{0}(\alpha_{n}) = 0,\\ \tau_{\pm}(\alpha_{1}) > \dots > \tau_{\pm}(\alpha_{n}) > 0 \end{bmatrix}$$

where the '±' signs mean that we have a relation for  $\tau_+$ , and another for  $\tau_-$ . These are finite sums by Lemma 2.3.7, and agree with (5.1.3.1)–(5.1.3.2) with  $\tau_\pm$ ,  $\tau_0$  in place of  $\tau_+$ ,  $\tau_-$ .

These relations then imply the relations

$$\begin{bmatrix} \mathscr{X}_{\alpha}^{\mathrm{ss}}(\tau_{\pm}) \end{bmatrix} = \sum_{\substack{n > 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathscr{X}) \setminus \{0\}:\\ \alpha = \alpha_{1} + \dots + \alpha_{n},\\ \tau_{0}(\alpha_{1}) = \dots = \tau_{0}(\alpha_{n}),\\ \tau_{\pm}(\alpha_{1} + \dots + \alpha_{i}) > \tau_{\pm}(\alpha_{i+1} + \dots + \alpha_{n}) \text{ for } i = 1, \dots, n - 1}$$

$$\begin{bmatrix} \mathscr{X}_{\theta}^{\mathrm{sd},\mathrm{ss}}(\tau_{\pm}) \end{bmatrix} = \sum_{\substack{n \ge 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathscr{X}) \setminus \{0\}:\\ \theta = \alpha_{1} + \alpha_{1}^{\vee} + \dots + \alpha_{n} + \alpha_{n}^{\vee} + \rho,\\ \tau_{0}(\alpha_{1}) = \dots = \tau_{0}(\alpha_{n}) = 0,\\ \tau_{\pm}(\alpha_{1} + \dots + \alpha_{i}) > 0 \text{ for } i = 1, \dots, n \end{bmatrix} \diamond \cdots \diamond \begin{bmatrix} \mathscr{X}_{\alpha_{n}}^{\mathrm{ss}}(\tau_{0}) \end{bmatrix} \diamond \begin{bmatrix} \mathscr{X}_{\rho}^{\mathrm{sd},\mathrm{ss}}(\tau_{0}) \end{bmatrix}, \quad (5.1.3.12)$$

which agree with (5.1.3.1)-(5.1.3.2) with  $\tau_0$ ,  $\tau_{\pm}$  in place of  $\tau_+$ ,  $\tau_-$ . Indeed, these can be verified by expanding the right-hand sides of (5.1.3.11)-(5.1.3.12) using (5.1.3.9)-(5.1.3.10), then applying Lemma 5.1.4 below to see that the results are equal to the left-hand sides.

Now, expanding the right-hand sides of (5.1.3.11)-(5.1.3.12) for  $\tau_+$  using (5.1.3.9)-(5.1.3.10) for  $\tau_-$ , then applying Lemma 5.1.4 below, gives the general case of (5.1.3.1)-(5.1.3.2).

To verify the relations (5.1.3.3)-(5.1.3.4), we first substitute the relations (5.1.3.1)-(5.1.3.2), in (4.1.2.1), (4.1.3.1) for  $\tau_{-}$ , then substitute in (4.1.2.2), (4.1.3.2) for  $\tau_{+}$ . Keeping track of the coefficients gives the desired relations.

**5.1.4. Lemma.** For symbols  $\alpha_1, \ldots, \alpha_n$  and maps  $\tau_1, \tau_2, \tau_3$  from  $\{\alpha_i + \cdots + \alpha_j \mid 1 \le i \le j \le n\}$  to totally ordered sets with distinguished elements 0, we have the identities

$$S(\alpha_{1}, \dots, \alpha_{n}; \tau_{1}, \tau_{3}) = \sum_{(\beta_{1}, \dots, \beta_{m}) \in Q} S(\beta_{1}, \dots, \beta_{m}; \tau_{2}, \tau_{3}) \cdot \prod_{i=1}^{m} S(\alpha_{a_{i-1}+1}, \dots, \alpha_{a_{i}}; \tau_{1}, \tau_{2}), \quad (5.1.4.1)$$

$$S^{\text{sd}}(\alpha_{1}, \dots, \alpha_{n}; \tau_{1}, \tau_{3}) = \sum_{(\beta_{1}, \dots, \beta_{m}) \in Q^{\text{sd}}} S^{\text{sd}}(\beta_{1}, \dots, \beta_{m}; \tau_{2}, \tau_{3}) \cdot \left(\prod_{i=1}^{m} S(\alpha_{a_{i-1}+1}, \dots, \alpha_{a_{i}}; \tau_{1}, \tau_{2})\right) \cdot S^{\text{sd}}(\alpha_{a_{m}+1}, \dots, \alpha_{n}; \tau_{1}, \tau_{2}), \quad (5.1.4.2)$$

where

$$Q = \left\{ (\beta_1, \dots, \beta_m) \middle| \begin{array}{l} m \ge 1, \ 0 = a_0 < \dots < a_m = n, \\ \beta_i = \alpha_{a_{i-1}+1} + \dots + \alpha_{a_i} \text{ for all } i \end{array} \right\},$$
$$Q^{\text{sd}} = \left\{ (\beta_1, \dots, \beta_m) \middle| \begin{array}{l} m \ge 0, \ 0 = a_0 < \dots < a_m \le n, \\ \beta_i = \alpha_{a_{i-1}+1} + \dots + \alpha_{a_i} \text{ for all } i \end{array} \right\}.$$

**Proof.** The identity (5.1.4.1) was proved in Joyce [32, Theorem 4.5]. The identity (5.1.4.2) follows from (5.1.4.1) and the fact that  $S^{sd}(\alpha_1, \ldots, \alpha_n; \tau_i, \tau_j) = S(\alpha_1, \ldots, \alpha_n, \infty; \tau_i, \tau_j)$ , where we set  $\tau_i(\alpha_j + \cdots + \alpha_n + \infty) = 0$  for all *i* and all  $1 \le j \le n + 1$ .

5.1.5. Weakening the assumptions. In Theorem 5.1.3, we can slightly weaken the assumptions by allowing  $\tau_0$  to be non-permissible, so that  $\mathscr{X}^{ss}_{\alpha}(\tau_0)$  can be non-quasi-compact, and we add the extra assumption that the sums (5.1.3.11)–(5.1.3.12) are locally finite for all classes  $\alpha$ ,  $\theta$ . In this case, the relations (5.1.3.9)–(5.1.3.10) are always valid as locally finite sums, and the proof shows that the relations (5.1.3.1)–(5.1.3.4) still hold as locally finite sums.

#### 5.2 An anti-symmetrized version

**5.2.1.** In this section, we rewrite the relations (5.1.3.3)-(5.1.3.4) in terms of anti-symmetrized product operations, instead of the operations \* and  $\diamond$ . This will be useful in writing down wall-crossing formulae for DT invariants in §5.3 below.

As in §5.1.1, let  $\mathcal{X}$  be a self-dual linear stack with quasi-compact filtrations.

5.2.2. Lie algebras and twisted modules. The motivic Hall algebra  $\mathbb{M}(\mathcal{X})$  can be seen as a Lie algebra using the commutator

$$[a,b] = a * b - b * a . (5.2.2.1)$$

This was considered in Joyce [29, §5.2]. It is equipped with a contravariant involution  $(-)^{\vee}$ , meaning that  $[a^{\vee}, b^{\vee}] = [b, a]^{\vee}$  for  $a, b \in \mathbb{M}(\mathcal{X})$ , which follows from Theorem 3.2.4.

We define a similar anti-symmetrized operation  $\heartsuit: \mathbb{M}(\mathcal{X}) \otimes \mathbb{M}(\mathcal{X}^{sd}) \to \mathbb{M}(\mathcal{X}^{sd})$  by

$$a \heartsuit m = a \diamond m - a^{\vee} \diamond m , \qquad (5.2.2.2)$$

where  $\diamond$  is the multiplication in the motivic Hall module. This does not define a Lie algebra module, but a *twisted module*, in that it satisfies the relations

$$a \heartsuit m = -a^{\vee} \heartsuit m \,, \tag{5.2.2.3}$$

$$a \heartsuit (b \heartsuit m) - b \heartsuit (a \heartsuit m) = [a, b] \heartsuit m - [a^{\lor}, b] \heartsuit m .$$
(5.2.2.4)

We see (5.2.2.4) as a Jacobi identity twisted by the contravariant involution of the Lie algebra, giving the extra term  $[a^{\vee}, b] \heartsuit m$ .

Note that over  $\mathbb{Q}$ , a twisted module in this sense is equivalent to a usual module for the Lie subalgebra consisting of elements *a* with  $a^{\vee} = -a$ , with the action  $a \cdot m = (1/2)(a \heartsuit m)$ .

**5.2.3. Theorem.** The relations (5.1.3.3)-(5.1.3.4) can be written only using the Lie bracket [-, -] and the operation  $\heartsuit$ , without using the products \* or  $\diamond$ .

More precisely, using the notations of Theorem 5.1.3, we have the relations

$$\begin{aligned}
\epsilon_{\alpha}(\tau_{-}) &= \sum_{\substack{n \ge 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathcal{X}) \setminus \{0\}:\\ \alpha = \alpha_{1} + \dots + \alpha_{n}}} \tilde{U}(\alpha_{1}, \dots, \alpha_{n}; \tau_{+}, \tau_{-}) \cdot \left[ \dots \left[ \left[ \epsilon_{\alpha_{1}}(\tau_{+}), \epsilon_{\alpha_{2}}(\tau_{+}) \right], \dots \right], \epsilon_{\alpha_{n}}(\tau_{+}) \right], \quad (5.2.3.1) \\
\epsilon_{\theta}^{sd}(\tau_{-}) &= \sum_{\substack{n \ge 0; \ m_{1}, \dots, m_{n} > 0;\\ \alpha_{1,1}, \dots, \alpha_{1,m_{1}}; \dots; \alpha_{n,m_{n}} \in \pi_{0}(\mathcal{X}) \setminus \{0\}; \ \rho \in \pi_{0}(\mathcal{X}^{sd}):\\ \theta = (\alpha_{1,1} + \alpha_{1,1}^{\vee} + \dots + \alpha_{1,m_{1}} + \alpha_{1,m_{1}}^{\vee}) + \dots + (\alpha_{n,1} + \alpha_{n,1}^{\vee} + \dots + \alpha_{n,m_{n}} + \alpha_{n,m_{n}}^{\vee}) + \rho \\
&= \left[ \left[ \epsilon_{\alpha_{1,1}}(\tau_{+}), \dots \right], \epsilon_{\alpha_{1,m_{1}}}(\tau_{+}) \right] \heartsuit \dots \heartsuit \left[ \left[ \epsilon_{\alpha_{n,1}}(\tau_{+}), \dots \right], \epsilon_{\alpha_{n,m_{n}}}(\tau_{+}) \right] \heartsuit \epsilon_{\beta}^{sd}(\tau_{+}) , \quad (5.2.3.2) \end{aligned}$$

where  $\tilde{U}(\ldots)$  and  $\tilde{U}^{sd}(\ldots)$  are certain combinatorial coefficients, whose choices are not unique.

Here, the formulae (5.2.3.1)-(5.2.3.2) are just (5.1.3.3)-(5.1.3.4) with the terms grouped differently, and this theorem is essentially a combinatorial property of the coefficients U(...) and  $U^{sd}(...)$  stating that such regrouping is always possible. The non-uniqueness of the coefficients is due to relations in the Lie brackets and the twisted module operation, such as the Jacobi identity and (5.2.2.3)-(5.2.2.4).

**Proof.** The relation (5.2.3.1) was shown in Joyce [32, Theorem 5.4]. The relation (5.2.3.2) will follow from a more general result to appear in [16]; an earlier version of this paper, [10, Appendix D], contains a direct but rather complicated proof of this combinatorial property.  $\Box$ 

#### 5.3 Wall-crossing for DT invariants

**5.3.1.** In this section, we prove wall-crossing formulae for our self-dual DT invariants defined in §§4.2 and 4.3, using the wall-crossing formulae for epsilon motives established in Theorems 5.1.3 and 5.2.3. A key ingredient is the *motivic integral identity* for Behrend functions proved by the author [12], generalizing the integral identities in the linear case of Kontsevich and Soibelman [37, Conjecture 4], proved by Lê [38], and Joyce and Song [35, Theorem 5.11].

Throughout, let *K* be an algebraically closed field of characteristic 0, and let  $\mathscr{X}$  be a selfdual (-1)-shifted symplectic linear stack over *K*, as in §4.2.3. We further assume that the classical truncation  $\mathscr{X}_{cl}$  is Nisnevich locally fundamental, as in §4.2.2.

5.3.2. Theorem. Let  $\tau_+, \tau_-, \tau_0$  be permissible self-dual stability conditions on  $\mathscr{X}$ , with  $\tau_0$  dominating both  $\tau_+$  and  $\tau_-$ . Then for any  $\alpha \in \pi_0(\mathscr{X})$  and  $\theta \in \pi_0(\mathscr{X}^{sd})$ , we have the wall-crossing formulae

$$DT_{\alpha}(\tau_{-}) = \sum_{\substack{n \ge 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathcal{X}) \setminus \{0\}:\\ \alpha = \alpha_{1} + \dots + \alpha_{n}}} \tilde{U}(\alpha_{1}, \dots, \alpha_{n}; \tau_{+}, \tau_{-}) \cdot \ell(\alpha_{1}, \dots, \alpha_{n}) \cdot DT_{\alpha_{1}}(\tau_{+}) \cdots DT_{\alpha_{n}}(\tau_{+}) , \quad (5.3.2.1)$$

$$DT_{\theta}^{sd}(\tau_{-}) = \sum_{\substack{n \ge 0; \ m_{1}, \dots, m_{n} > 0;\\ \alpha_{1,1}, \dots, \alpha_{1,m_{1}}; \dots; \alpha_{n,m_{n}} \in \pi_{0}(\mathcal{X}) \setminus \{0\}; \ \rho \in \pi_{0}(\mathcal{X}^{sd}):\\ \theta = (\alpha_{1,1} + \alpha_{1,1}^{\vee} + \dots + \alpha_{1,m_{1}} + \alpha_{1,m_{1}}^{\vee}) + \dots + (\alpha_{n,1} + \alpha_{n,1}^{\vee} + \dots + \alpha_{n,m_{n}} + \alpha_{n,m_{n}}^{\vee}) + \rho} \ell^{sd}(\alpha_{1,1}, \dots, \alpha_{1,m_{1}}; \dots; \alpha_{n,1}, \dots, \alpha_{n,m_{n}}; \rho) :$$

$$\left(\mathrm{DT}_{\alpha_{1,1}}(\tau_{+})\cdots\mathrm{DT}_{\alpha_{1,m_{1}}}(\tau_{+})\right)\cdots\left(\mathrm{DT}_{\alpha_{n,1}}(\tau_{+})\cdots\mathrm{DT}_{\alpha_{n,m_{n}}}(\tau_{+})\right)\cdots\mathrm{DT}_{\beta}^{\mathrm{sd}}(\tau_{+})$$

where the sums contain finitely many non-zero terms, the coefficients  $\tilde{U}(...), \tilde{U}^{sd}(...) \in \mathbb{Q}$  are defined in Theorem 5.2.3, and the coefficients  $\ell(...), \ell^{sd}(...) \in \mathbb{Z}$  are defined in §5.3.6 below.

If, moreover,  $\mathcal{X}$  is equipped with an orientation data  $o_{\mathcal{X}}$  or a self-dual orientation data  $(o_{\mathcal{X}}, o_{\mathcal{X}^{sd}})$ , then we have the wall-crossing formulae

$$DT_{\alpha}^{mot}(\tau_{-}) = \sum_{\substack{n \ge 0; \ \alpha_{1}, \dots, \alpha_{n} \in \pi_{0}(\mathcal{X}) \setminus \{0\}:\\ \alpha = \alpha_{1} + \dots + \alpha_{n}}} \tilde{U}(\alpha_{1}, \dots, \alpha_{n}; \tau_{+}, \tau_{-}) \cdot L(\alpha_{1}, \dots, \alpha_{n}) \cdot DT_{\alpha_{1}}^{mot}(\tau_{+}) \cdots DT_{\alpha_{n}}^{mot}(\tau_{+}) ,$$
(5.3.2.3)

$$DT_{\theta}^{\text{mot,sd}}(\tau_{-}) = \sum_{\substack{n \ge 0; \ m_{1}, \dots, m_{n} \ge 0; \\ \alpha_{1,1}, \dots, \alpha_{1,m_{1}}; \dots; \alpha_{n,1}, \dots, \alpha_{n,m_{n}} \in \pi_{0}(\mathcal{X}) \setminus \{0\}; \ \rho \in \pi_{0}(\mathcal{X}^{\text{sd}}): \\ \theta = (\alpha_{1,1} + \alpha_{1,1}^{\vee} + \dots + \alpha_{1,m_{1}} + \alpha_{1,m_{1}}^{\vee}) + \dots + (\alpha_{n,1} + \alpha_{n,1}^{\vee} + \dots + \alpha_{n,m_{n}} + \alpha_{n,m_{n}}^{\vee}) + \rho \\ L^{\text{sd}}(\alpha_{1,1}, \dots, \alpha_{1,m_{1}}; \dots; \alpha_{n,1}, \dots, \alpha_{n,m_{n}}; \rho) \cdot \\ \left(DT_{\alpha_{1,1}}^{\text{mot}}(\tau_{+}) \cdots DT_{\alpha_{1,m_{1}}}^{\text{mot}}(\tau_{+})\right) \cdots \left(DT_{\alpha_{n,1}}^{\text{mot}}(\tau_{+}) \cdots DT_{\alpha_{n,m_{n}}}^{\text{mot}}(\tau_{+})\right) \cdots DT_{\rho}^{\text{mot,sd}}(\tau_{+}),$$
(5.3.2.4)

respectively, where the coefficients  $L(...), L^{sd}(...) \in \mathbb{Z}[\mathbb{L}^{\pm 1/2}]$  are defined in §5.3.6 below.

The proof of the theorem will be given in §5.3.9.

**5.3.3. Symmetric stacks**. The wall-crossing formulae in Theorem 5.3.2 provide a condition for the DT invariants to be independent of the choice of the stability condition.

We say that a (-1)-shifted symplectic stack  $\mathscr{X}$  is *numerically symmetric*, if vdim Filt $(\mathscr{X}) = 0$ , meaning that this holds on every connected component of Filt $(\mathscr{X})$ . See the author et al. [13, §4.3] for examples of stacks satisfying this condition.

For example, if  $\mathscr{X}$  is a self-dual (-1)-shifted symplectic linear stack, then  $\mathscr{X}$  is numerically symmetric if and only if vdim  $\mathscr{X}^+_{\alpha,\beta} = 0$  for all  $\alpha, \beta \in \pi_0(\mathscr{X})$ , and  $\mathscr{X}^{sd}$  is numerically symmetric if and only if vdim  $\mathscr{X}^{sd,+}_{\alpha,\theta} = 0$  for all  $\alpha \in \pi_0(\mathscr{X})$  and  $\theta \in \pi_0(\mathscr{X}^{sd})$ .

When  $\mathscr{X}$  and  $\mathscr{X}^{sd}$  are numerically symmetric, the coefficients L(...),  $\ell(...)$  are zero unless  $n \leq 1$ , and the coefficients  $L^{sd}(...)$ ,  $\ell^{sd}(...)$  are zero unless n = 0, which follow from their definitions. This immediately implies the following:

**5.3.4. Corollary.** In the situation of Theorem 5.3.2, assume that  $\mathcal{X}$  and  $\mathcal{X}^{sd}$  are numerically symmetric. Then the relations (5.3.2.1)–(5.3.2.4) simplify to

$$DT_{\alpha}(\tau_{-}) = DT_{\alpha}(\tau_{+}), \qquad DT_{\theta}^{sd}(\tau_{-}) = DT_{\theta}^{sd}(\tau_{+}), \qquad (5.3.4.1)$$

$$DT^{mot}_{\alpha}(\tau_{-}) = DT^{mot}_{\alpha}(\tau_{+}), \qquad DT^{mot,sd}_{\theta}(\tau_{-}) = DT^{mot,sd}_{\theta}(\tau_{+}).$$
(5.3.4.2)

In particular, if  $\mathcal{X}$  has quasi-compact connected components, then all the above invariants are independent of the choice of the stability condition.

Here, the final claim follows from taking  $\tau_0$  and  $\tau_+$  to be the trivial stability condition, which is permissible when  $\mathcal{X}$  has quasi-compact connected components.

The remaining part of this section is devoted to the proof of Theorem 5.3.2.

5.3.5. Lattice algebras and modules. Define

$$\Lambda_{\mathscr{X}} = \bigoplus_{\alpha \in \pi_0(\mathscr{X})} \hat{\mathbb{M}}^{\mathrm{mon}}(K; \mathbb{Q}) \cdot \lambda_{\alpha} , \qquad \Lambda_{\mathscr{X}}^{\mathrm{sd}} = \bigoplus_{\theta \in \pi_0(\mathscr{X}^{\mathrm{sd}})} \hat{\mathbb{M}}^{\mathrm{mon}}(K; \mathbb{Q}) \cdot \lambda_{\theta}^{\mathrm{sd}} ,$$

where  $\hat{\mathbb{M}}^{\text{mon}}(K;\mathbb{Q})$  is the ring of monodromic motives defined in §4.3.2. We define a product \* on  $\Lambda_{\mathcal{X}}$ , and a  $\Lambda_{\mathcal{X}}$ -module structure  $\diamond$  on  $\Lambda_{\mathcal{X}}^{\text{sd}}$ , by setting

$$\lambda_{\alpha} * \lambda_{\beta} = \frac{\mathbb{L}^{\operatorname{vdim} \mathscr{X}_{\alpha,\beta}^{+}/2}}{\mathbb{L}^{1/2} - \mathbb{L}^{-1/2}} \cdot \lambda_{\alpha+\beta}, \qquad \lambda_{\alpha} \diamond \lambda_{\theta}^{\operatorname{sd}} = \frac{\mathbb{L}^{\operatorname{vdim} \mathscr{X}_{\alpha,\theta}^{\operatorname{sd}}/2}}{\mathbb{L}^{1/2} - \mathbb{L}^{-1/2}} \cdot \lambda_{\alpha+\theta+\alpha^{\vee}}^{\operatorname{sd}}$$
(5.3.5.1)

for  $\alpha, \beta \in \pi_0(\mathcal{X})$  and  $\theta \in \pi_0(\mathcal{X}^{sd})$ . The associativity of these operations follow from the relations

$$\begin{aligned} & \operatorname{vdim} \mathscr{X}_{\alpha,\beta}^{+} + \operatorname{vdim} \mathscr{X}_{\alpha+\beta,\gamma}^{+} = \operatorname{vdim} \mathscr{X}_{\alpha,\beta,\gamma}^{+} = \operatorname{vdim} \mathscr{X}_{\alpha,\beta+\gamma}^{+} + \operatorname{vdim} \mathscr{X}_{\beta,\gamma}^{+} \,, \\ & \operatorname{vdim} \mathscr{X}_{\alpha,\beta}^{+} + \operatorname{vdim} \mathscr{X}_{\alpha+\beta,\theta}^{\operatorname{sd},+} = \operatorname{vdim} \mathscr{X}_{\alpha,\beta+\theta}^{\operatorname{sd},+} = \operatorname{vdim} \mathscr{X}_{\alpha,\beta+\theta+\beta^{\vee}}^{\operatorname{sd},+} + \operatorname{vdim} \mathscr{X}_{\beta,\theta}^{\operatorname{sd},+} \,, \end{aligned}$$

which follow from the derived versions of the associativity diagrams (3.2.4.6) and (3.2.4.8). The algebra  $\Lambda_{\mathcal{X}}$  is often called the *quantum torus* in the literature, such as in Kontsevich and

Soibelman [37, §6.2].

The map  $\lambda_{\alpha} \mapsto \lambda_{\alpha^{\vee}}$  defines a contravariant involution  $(-)^{\vee}$  of  $\Lambda_{\mathcal{X}}$ . We also write  $a \heartsuit m = a \diamond m - a^{\vee} \diamond m$  for  $a \in \Lambda_{\mathcal{X}}$  and  $m \in \Lambda_{\mathcal{X}}^{sd}$ , as in §5.2.2, which gives  $\Lambda_{\mathcal{X}}^{sd}$  the structure of a twisted module over the involutive Lie algebra  $\Lambda_{\mathcal{X}}$ , with the commutator Lie bracket.

We also define the numerical versions

$$\bar{\Lambda}_{\mathscr{X}} = \bigoplus_{\alpha \in \pi_0(\mathscr{X})} \mathbb{Q} \cdot \bar{\lambda}_{\alpha} , \qquad \bar{\Lambda}_{\mathscr{X}}^{\mathrm{sd}} = \bigoplus_{\theta \in \pi_0(\mathscr{X}^{\mathrm{sd}})} \mathbb{Q} \cdot \bar{\lambda}_{\theta}^{\mathrm{sd}} ,$$

which are no longer equipped with algebra structures, but have a Lie bracket and a twisted module operation  $\heartsuit$ , respectively, given by

$$\left[\bar{\lambda}_{\alpha}, \bar{\lambda}_{\beta}\right] = (-1)^{1 + \operatorname{vdim} \mathscr{X}_{\alpha,\beta}^{+}} \cdot \operatorname{vdim} \mathscr{X}_{\alpha,\beta}^{+} \cdot \bar{\lambda}_{\alpha+\beta}, \qquad (5.3.5.2)$$

$$\bar{\lambda}_{\alpha} \heartsuit \bar{\lambda}_{\theta}^{\mathrm{sd}} = (-1)^{1 + \mathrm{vdim}\,\mathcal{X}_{\alpha,\theta}^{\mathrm{sd},+}} \cdot \mathrm{vdim}\,\mathcal{X}_{\alpha,\theta}^{\mathrm{sd},+} \cdot \bar{\lambda}_{\alpha+\theta+\alpha^{\vee}}^{\mathrm{sd}} \,. \tag{5.3.5.3}$$

By [12, Lemma 3.1.7], we have vdim  $\mathscr{X}^+_{\beta,\alpha} = -\text{vdim} \, \mathscr{X}^+_{\alpha,\beta}$  and vdim  $\mathscr{X}^{\text{sd},+}_{\alpha^{\vee},\theta} = -\text{vdim} \, \mathscr{X}^{\text{sd},+}_{\alpha,\theta}$ , establishing (5.3.5.2) and (5.3.5.3) as limits of (5.3.5.1) as  $\mathbb{L}^{1/2} \to -1$ .

**5.3.6.** Coefficients. We can now define the coefficients  $L(...), L^{sd}(...)$ , etc., which appear in (5.3.2.1).

For  $\alpha_1, \ldots, \alpha_n \in \pi_0(\mathcal{X})$ , we record the coefficients of the Lie brackets in  $\Lambda_{\mathcal{X}}$  and  $\bar{\Lambda}_{\mathcal{X}}$  as

$$\left[\left[\ldots\left[\lambda_{\alpha_{1}},\lambda_{\alpha_{2}}\right],\ldots\right],\lambda_{\alpha_{n}}\right]=L(\alpha_{1},\ldots,\alpha_{n})\cdot\lambda_{\alpha_{1}+\cdots+\alpha_{n}},$$
(5.3.6.1)

$$\left[\left[\dots\left[\bar{\lambda}_{\alpha_1},\bar{\lambda}_{\alpha_2}\right],\dots\right],\bar{\lambda}_{\alpha_n}\right] = \ell(\alpha_1,\dots,\alpha_n) \cdot \bar{\lambda}_{\alpha_1+\dots+\alpha_n}, \qquad (5.3.6.2)$$

where  $L(\alpha_1, \ldots, \alpha_n) \in \mathbb{Z}[\mathbb{L}^{\pm 1/2}]$  and  $\ell(\alpha_1, \ldots, \alpha_n) \in \mathbb{Z}$ .

Similarly, for  $\alpha_{1,1}, \ldots, \alpha_{1,m_1}; \ldots; \alpha_{n,1}, \ldots, \alpha_{n,m_n} \in \pi_0(\mathcal{X})$  and  $\rho \in \pi_0(\mathcal{X}^{sd})$ , we also record the coefficients in

$$\begin{bmatrix} \dots [\lambda_{\alpha_{1,1}}, \lambda_{\alpha_{1,2}}], \dots ], \lambda_{\alpha_{1,m_1}} \end{bmatrix} \otimes \dots \otimes \begin{bmatrix} \dots [\lambda_{\alpha_{n,1}}, \lambda_{\alpha_{n,2}}], \dots ], \lambda_{\alpha_{n,m_n}} \end{bmatrix} \otimes \lambda_{\rho}^{\mathrm{sd}}$$
$$= L^{\mathrm{sd}}(\alpha_{1,1}, \dots, \alpha_{1,m_1}; \dots; \alpha_{n,1}, \dots, \alpha_{n,m_n}; \rho) \cdot \lambda_{\alpha_{1,1}+\alpha_{1,1}^{\vee}+\dots+\alpha_{n,m_n}+\alpha_{n,m_n}^{\vee}+\rho}, \qquad (5.3.6.3)$$

$$\begin{bmatrix} [\dots [\bar{\lambda}_{\alpha_{1,1}}, \bar{\lambda}_{\alpha_{1,2}}], \dots], \bar{\lambda}_{\alpha_{1,m_1}} \end{bmatrix} \otimes \cdots \otimes \begin{bmatrix} [\dots [\bar{\lambda}_{\alpha_{n,1}}, \bar{\lambda}_{\alpha_{n,2}}], \dots], \bar{\lambda}_{\alpha_{n,m_n}} \end{bmatrix} \otimes \bar{\lambda}_{\rho}^{\mathrm{sd}}$$
$$= \ell^{\mathrm{sd}}(\alpha_{1,1}, \dots, \alpha_{1,m_1}; \dots; \alpha_{n,1}, \dots, \alpha_{n,m_n}; \rho) \cdot \bar{\lambda}_{\alpha_{1,1}+\alpha_{1,1}^{\vee}+\dots+\alpha_{n,m_n}+\alpha_{n,m_n}^{\vee}+\rho}, \qquad (5.3.6.4)$$

where  $L^{\mathrm{sd}}(\ldots) \in \mathbb{Z}[\mathbb{L}^{\pm 1/2}]$  and  $\ell^{\mathrm{sd}}(\ldots) \in \mathbb{Z}$ .

These coefficients only depend on the numbers vdim  $\mathscr{X}^+_{\alpha,\beta}$  and vdim  $\mathscr{X}^{\mathrm{sd},+}_{\alpha,\theta}$  for  $\alpha, \beta \in \pi_0(\mathscr{X})$ and  $\theta \in \pi_0(\mathscr{X}^{\mathrm{sd}})$ . They have straightforward explicit expressions, which we omit.

We have the relations  $\ell(...) = L(...)|_{\mathbb{L}^{1/2}=-1}$  and  $\ell^{sd}(...) = L^{sd}(...)|_{\mathbb{L}^{1/2}=-1}$ . Also, L(...) and  $L^{sd}(...)$  are symmetric Laurent polynomials in  $\mathbb{L}^{1/2}$ , in that they are invariant under the transformation  $\mathbb{L}^{1/2} \mapsto \mathbb{L}^{-1/2}$ .

#### 5.3.7. The motivic integral identity. A crucial ingredient in proving wall-crossing formulae

for DT invariants is the *motivic integral identity* for the motivic Behrend function, first conjectured by Kontsevich and Soibelman [37, Conjecture 4] in the linear case, proved by Lê [38] in that case, and proved by the author [12, Theorem 4.2.2] in general.

Suppose that we are given a self-dual orientation data  $(o_{\mathcal{X}}, o_{\mathcal{X}^{sd}})$  on  $\mathcal{X}$ . The motivic integral identity states, in this case, that we have

$$\nu_{\mathcal{X}}^{\text{mot}} \boxtimes \nu_{\mathcal{X}}^{\text{mot}} = \mathbb{L}^{-\text{vdim}\,\mathcal{X}_{\alpha,\beta}^{+}/2} \cdot \operatorname{gr}_{!} \circ \operatorname{ev}_{1}^{*}(\nu_{\mathcal{X}}^{\text{mot}}) \qquad \text{in } \hat{\mathbb{M}}^{\text{mon}}(\mathcal{X}_{\alpha} \times \mathcal{X}_{\beta}) , \qquad (5.3.7.1)$$

$$\nu_{\mathcal{X}}^{\text{mot}} \boxtimes \nu_{\mathcal{X}^{\text{sd}}}^{\text{mot}} = \mathbb{L}^{-\text{vdim}\,\mathcal{X}_{\alpha,\theta}^{\text{sd}+/2}} \cdot \operatorname{gr}_{!} \circ \operatorname{ev}_{1}^{*}(\nu_{\mathcal{X}^{\text{sd}}}^{\text{mot}}) \quad \text{in } \hat{\mathbb{M}}^{\text{mon}}(\mathcal{X}_{\alpha} \times \mathcal{X}_{\theta}^{\text{sd}}) , \qquad (5.3.7.2)$$

where  $\alpha, \beta \in \pi_0(\mathcal{X})$  and  $\theta \in \pi_0(\mathcal{X}^{sd})$ , and the compositions are through  $\hat{\mathbb{M}}^{\text{mon}}(\mathcal{X}^{*}_{\alpha,\beta})$  and  $\hat{\mathbb{M}}^{\text{mon}}(\mathcal{X}^{sd,+}_{\alpha,\theta})$ , respectively. These identities imply the relations

$$\left(\int_{\mathcal{X}_{\alpha}} a \cdot v_{\mathcal{X}}^{\mathrm{mot}}\right) \cdot \left(\int_{\mathcal{X}_{\beta}} b \cdot v_{\mathcal{X}}^{\mathrm{mot}}\right) = \mathbb{L}^{-\mathrm{vdim}\,\mathcal{X}_{\alpha,\beta}^{+}/2} \cdot \int_{\mathcal{X}_{\alpha+\beta}} (a * b) \cdot v_{\mathcal{X}}^{\mathrm{mot}}, \qquad (5.3.7.3)$$

$$\left(\int_{\mathcal{X}_{\alpha}} a \cdot v_{\mathcal{X}^{\mathrm{sd}}}^{\mathrm{mot}}\right) \cdot \left(\int_{\mathcal{X}_{\theta}^{\mathrm{sd}}} m \cdot v_{\mathcal{X}^{\mathrm{sd}}}^{\mathrm{mot}}\right) = \mathbb{L}^{-\mathrm{vdim}\,\mathcal{X}_{\alpha,\theta}^{\mathrm{sd},+/2}} \cdot \int_{\mathcal{X}_{\alpha+\theta+\alpha}^{\mathrm{sd}}} (a \diamond m) \cdot v_{\mathcal{X}^{\mathrm{sd}}}^{\mathrm{mot}}, \qquad (5.3.7.4)$$

where  $a \in \mathbb{M}_{qc}(\mathscr{X}_{\alpha}; \mathbb{Q}), b \in \mathbb{M}_{qc}(\mathscr{X}_{\beta}; \mathbb{Q})$ , and  $m \in \mathbb{M}_{qc}(\mathscr{X}_{\theta}^{sd}; \mathbb{Q})$ . These follow from identifying both sides of each relation with the integrals

$$\mathbb{L}^{-\operatorname{vdim} \mathscr{X}^{+}_{\alpha,\beta}/2} \cdot \int_{\mathscr{X}^{+}_{\alpha,\beta}} \operatorname{gr}^{*}(a \boxtimes b) \cdot \operatorname{ev}_{1}^{*}(v_{\mathscr{X}}^{\operatorname{mot}}), \qquad \mathbb{L}^{-\operatorname{vdim} \mathscr{X}^{\operatorname{sd},+}_{\alpha,\theta}/2} \cdot \int_{\mathscr{X}^{\operatorname{sd},+}_{\alpha,\theta}} \operatorname{gr}^{*}(a \boxtimes m) \cdot \operatorname{ev}_{1}^{*}(v_{\mathscr{X}^{\operatorname{sd}}}^{\operatorname{mot}}),$$

respectively, using the projection formula (3.1.3.1). The relation (5.3.7.3) was first described by Kontsevich and Soibelman [37, Theorem 8].

**5.3.8. The numeric integral identity**. Using the numeric version of the motivic integral identity, proved by the author [12, Theorem 4.3.3], we can also obtain numerical versions of the integral relations (5.3.7.3)-(5.3.7.4),

$$\int_{\mathcal{X}_{\alpha+\beta}} (1-\mathbb{L}) \cdot [a,b] \cdot v_{\mathcal{X}} \, d\chi = (-1)^{1+\operatorname{vdim}\mathcal{X}_{\alpha,\beta}^+} \cdot \operatorname{vdim}\mathcal{X}_{\alpha,\beta}^+ \cdot \left(\int_{\mathcal{X}_{\alpha}} (1-\mathbb{L}) \cdot a \cdot v_{\mathcal{X}} \, d\chi\right) \cdot \left(\int_{\mathcal{X}_{\beta}} (1-\mathbb{L}) \cdot b \cdot v_{\mathcal{X}} \, d\chi\right), \tag{5.3.8.1}$$

$$\int_{\mathcal{X}_{\alpha+\theta+\alpha^{\vee}}^{\mathrm{sd}}} (a \heartsuit m) \cdot v_{\mathcal{X}^{\mathrm{sd}}} d\chi = (-1)^{1+\mathrm{vdim}\,\mathcal{X}_{\alpha,\theta}^{\mathrm{sd},+}} \cdot \mathrm{vdim}\,\mathcal{X}_{\alpha,\theta}^{\mathrm{sd},+} \cdot \left(\int_{\mathcal{X}_{\alpha}} (1-\mathbb{L}) \cdot a \cdot v_{\mathcal{X}} d\chi\right) \cdot \left(\int_{\mathcal{X}_{\theta}^{\mathrm{sd}}} m \cdot v_{\mathcal{X}^{\mathrm{sd}}} d\chi\right),$$
(5.3.8.2)

provided that the motives *a*, *b*, *m* are chosen so that the integrals on the right-hand sides are finite, that is, they lie in  $\hat{\mathbb{M}}^{\text{mon,reg}}(K; A)$  as in §4.3.2 before taking the Euler characteristics. These identities do not require orientations on  $\mathscr{X}$  or  $\mathscr{X}^{\text{sd}}$ . The identity (5.3.8.1) was proved by Joyce and Song [35, Theorem 5.14] in the setting of Calabi–Yau threefolds.

To prove them, we use a similar argument as in 5.3.7. Namely, for (5.3.8.1), we identify

the left-hand side with

$$\chi \left( (1 - \mathbb{L})^2 \cdot \left( -\int_{\mathbb{P}(\mathcal{X}^+_{\alpha,\beta})} \operatorname{gr}^*(a \boxtimes b) \cdot \operatorname{ev}_1^*(v_{\mathcal{X}}) + \int_{\mathbb{P}(\mathcal{X}^+_{\beta,\alpha})} \operatorname{\bar{gr}}^*(a \boxtimes b) \cdot \operatorname{\bar{ev}}_1^*(v_{\mathcal{X}}) \right) \right) + \int_{\mathcal{X}_{\alpha} \times \mathcal{X}_{\beta}} (1 - \mathbb{L}) \cdot (a \boxtimes b) \cdot \left( \mathbb{L}^{-h^1(\mathbb{L}_{\mathrm{gr}})} - \mathbb{L}^{-h^1(\mathbb{L}_{\mathrm{gr}})} \right) \cdot \oplus^*(v_{\mathcal{X}}) d\chi , \qquad (5.3.8.3)$$

where  $\mathbb{P}(\mathscr{X}_{\alpha,\beta}^+) = (\mathscr{X}_{\alpha,\beta}^+ \setminus \mathrm{sf}(\mathscr{X}_{\alpha} \times \mathscr{X}_{\beta}))/\mathbb{G}_{\mathrm{m}}$ , with the  $\mathbb{G}_{\mathrm{m}}$ -action given by choosing an identification of  $\mathscr{X}_{\alpha,\beta}^+$  with a component of Filt( $\mathscr{X}$ ), and  $\mathbb{P}(\mathscr{X}_{\beta,\alpha}^+)$  is defined similarly, using the opposite component. We denote by  $\overline{\mathrm{gr}}$ ,  $\overline{\mathrm{ev}}_1$  the maps  $\mathrm{gr}$ ,  $\mathrm{ev}_1$  for  $\mathscr{X}_{\beta,\alpha}^+$ , and by  $\mathbb{L}_{\mathrm{gr}}$  the relative cotangent complex of  $\mathscr{X}_{\alpha,\beta}^+$  over  $\mathscr{X}_{\alpha} \times \mathscr{X}_{\beta}$ . We regard  $h^1(\mathbb{L}_{\mathrm{gr}}) = \dim \mathrm{H}^1(\mathbb{L}_{\mathrm{gr}})$  as a constructible function on  $\mathscr{X}_{\alpha,\beta}^+$ , which can be pulled back to  $\mathscr{X}_{\alpha} \times \mathscr{X}_{\beta}$ . The factors  $\mathbb{L}^{-h^1(\mathbb{L}_{\mathrm{gr}})}$  and  $\mathbb{L}^{-h^1(\mathbb{L}_{\mathrm{gr}})}$  are due to the difference of stabilizer groups in  $\mathscr{X}_{\alpha,\beta}^+$  and  $\mathscr{X}_{\alpha} \times \mathscr{X}_{\beta}$ ; see [12, §4.3.4] for details. Applying [12, (4.3.3.2)], (5.3.8.3) becomes

$$\int_{\mathcal{X}_{\alpha} \times \mathcal{X}_{\beta}} (1 - \mathbb{L})^2 \cdot (a \boxtimes b) \cdot \left(h^1(\mathbb{L}_{\mathrm{gr}}) - h^0(\mathbb{L}_{\mathrm{gr}}) + h^0(\mathbb{L}_{\mathrm{gr}}) - h^1(\mathbb{L}_{\mathrm{gr}})\right) \cdot \oplus^*(\nu_{\mathcal{X}}) \, d\chi \,, \quad (5.3.8.4)$$

where we also replaced  $\mathbb{L}^{-h^1(\mathbb{L}_{gr})} - \mathbb{L}^{-h^1(\mathbb{L}_{gr})}$  by  $(1 - \mathbb{L}) \cdot (h^1(\mathbb{L}_{gr}) - h^1(\mathbb{L}_{gr}))$ , as they are equal modulo  $(1 - \mathbb{L})^2$ , so this will not affect the integral. By [12, Lemma 3.1.7], the alternating sum in (5.3.8.4) is equal to  $-\text{vdim } \mathcal{X}^+_{\alpha,\beta}$ . Finally, by [12, (4.3.3.1)], we have  $\oplus^*(v_{\mathcal{X}}) = (-1)^{\text{vdim } \mathcal{X}^+_{\alpha,\beta}} \cdot (v_{\mathcal{X}} \boxtimes v_{\mathcal{X}})$ , which identifies (5.3.8.4) with the right-hand side of (5.3.8.1).

The identity (5.3.8.2) can be proved analogously.

#### 5.3.9. Proof of Theorem 5.3.2. Consider the integration maps

$$\begin{split} (\mathbb{L}^{1/2} - \mathbb{L}^{-1/2}) \cdot \int_{\mathcal{X}} (-) \cdot v_{\mathcal{X}}^{\mathrm{mot}} \colon & \mathbb{M}_{\mathrm{qc}}(\mathcal{X}; \mathbb{Q}) \longrightarrow \Lambda_{\mathcal{X}} ,\\ & \int_{\mathcal{X}^{\mathrm{sd}}} (-) \cdot v_{\mathcal{X}^{\mathrm{sd}}}^{\mathrm{mot}} \colon & \mathbb{M}_{\mathrm{qc}}(\mathcal{X}^{\mathrm{sd}}; \mathbb{Q}) \longrightarrow \Lambda_{\mathcal{X}}^{\mathrm{sd}} , \end{split}$$

where the generators  $\lambda_{\alpha}$  and  $\lambda_{\theta}^{sd}$  record which components the motives are supported on. The relations (5.3.7.3)–(5.3.7.4) imply that these maps are algebra and module homomorphisms.

Similarly, the relations (5.3.8.1)–(5.3.8.2) imply that the integration maps

$$\int_{\mathcal{X}} (1 - \mathbb{L}) \cdot (-) \cdot v_{\mathcal{X}} d\chi \colon \mathbb{M}^*_{qc}(\mathcal{X}; \mathbb{Q}) \longrightarrow \bar{\Lambda}_{\mathcal{X}},$$
$$\int_{\mathcal{X}^{sd}} (-) \cdot v_{\mathcal{X}^{sd}} \colon \mathbb{M}^*_{qc}(\mathcal{X}^{sd}; \mathbb{Q}) \longrightarrow \bar{\Lambda}^{sd}_{\mathcal{X}},$$

are Lie algebra and twisted module homomorphisms, where the superscripts \* indicate subspaces of motives for which the integrals are finite. It follows from (5.3.8.1)–(5.3.8.2) that these subspaces are a Lie subalgebra and a sub-twisted module for this subalgebra, respectively.

The theorem is now a direct consequence of Theorem 5.2.3, by applying the above integration homomorphisms to the relations (5.2.3.1)-(5.2.3.2).

## 6 Applications

#### 6.1 Self-dual quivers

**6.1.1.** We apply our theory to study DT invariants counting *self-dual representations* of a *self-dual quiver*. These are an analogue of orthogonal and symplectic principal bundles on a variety, similar to how the usual quiver representations are analogous to vector bundles or coherent sheaves on varieties.

Self-dual quivers were first introduced by Derksen and Weyman [19] as a special case of *G*-quivers for G = O(n) or Sp(2n), and studied by Young [53–55] in the context of DT theory.

Throughout, we fix an algebraically closed field *K* of characteristic zero.

**6.1.2. Self-dual quivers.** To fix notations, recall that a *quiver* is a quadruple  $Q = (Q_0, Q_1, s, t)$ , where  $Q_0$  and  $Q_1$  are finite sets, thought of as the sets of vertices and edges, and  $s, t: Q_1 \rightarrow Q_0$  are the source and target maps.

For a quiver *Q*, a *self-dual structure* on *Q* consists of the following data:

(i) A contravariant involution

$$(-)^{\vee} \colon Q \xrightarrow{\sim} Q^{\operatorname{op}}$$

where  $Q^{\text{op}} = (Q_0, Q_1, t, s)$  is the opposite quiver of Q, such that  $(-)^{\vee\vee} = \text{id}$ .

(ii) Choices of signs

$$u: Q_0 \longrightarrow \{\pm 1\}, \qquad v: Q_1 \longrightarrow \{\pm 1\},$$

such that  $u(i) = u(i^{\vee})$  for all  $i \in Q_0$ , and  $v(a) v(a^{\vee}) = u(s(a)) u(t(a))$  for all  $a \in Q_1$ .

In this case, the *K*-linear abelian category Mod(KQ) of finite-dimensional representations of Q admits a self-dual structure, defined as follows. For a representation E of Q, write  $E_i$  for the vector space at  $i \in Q_0$  and  $e_a \colon E_{s(a)} \to E_{t(a)}$  the linear map for the edge  $a \in Q_1$ . Define the *dual representation*  $E^{\vee}$  by assigning the vector space  $(E_i^{\vee})^{\vee}$  to the vertex *i*, and the linear map  $v(a) \cdot (e_a^{\vee})^{\vee}$  to the edge *a*. Then, identify  $E^{\vee\vee}$  with *E* using the sign u(i) at each vertex *i*. As in §2.1.2, we have the groupoid  $Mod(KQ)^{sd}$  of *self-dual representations* of Q.

**6.1.3. Moduli stacks**. Let Q be a quiver, and let  $\mathcal{X}_Q$  be the moduli stack of representations of Q over K. Explicitly, we have

$$\mathscr{X}_{Q} = \coprod_{\alpha \in \mathbb{N}^{Q_{0}}} V_{\alpha}/G_{\alpha} , \qquad (6.1.3.1)$$

where  $V_{\alpha} = \bigoplus_{a \in Q_1} \text{Hom}(K^{\alpha_{s(a)}}, K^{\alpha_{t(a)}})$ , and  $G_{\alpha} = \prod_{i \in Q_0} \text{GL}(\alpha_i)$ .

If Q is equipped with a self-dual structure, the self-dual structure on Mod(KQ) extends to an involution of  $\mathcal{X}_O$ , establishing it as a self-dual linear stack. The homotopy fixed locus  $\mathcal{X}_O^{sd}$  can be seen as the moduli stack of self-dual representations of Q. Explicitly, we have

$$\mathcal{X}_{Q}^{\mathrm{sd}} \simeq \coprod_{\theta \in (\mathbb{N}^{Q_0})^{\mathrm{sd}}} V_{\theta}^{\mathrm{sd}} / G_{\theta}^{\mathrm{sd}} , \qquad (6.1.3.2)$$

where  $(\mathbb{N}^{Q_0})^{\mathrm{sd}} \subset \mathbb{N}^{Q_0}$  is the subset of dimension vectors  $\theta$  such that  $\theta_i = \theta_{i^{\vee}}$  for all  $i \in Q_0$ , and  $\theta_i$  is even if  $i = i^{\vee}$  and u(i) = -1. The vector space  $V_{\theta}^{\mathrm{sd}}$  and the group  $G_{\theta}^{\mathrm{sd}}$  are given by

$$V_{\theta}^{\mathrm{sd}} = \prod_{a \in Q_1^{\circ}/\mathbb{Z}_2} \operatorname{Hom}(K^{\theta_{s(a)}}, K^{\theta_{t(a)}}) \times \prod_{a \in Q_1^{+}} \operatorname{Sym}^2(K^{\theta_{t(a)}}) \times \prod_{a \in Q_1^{-}} \wedge^2(K^{\theta_{t(a)}}), \qquad (6.1.3.3)$$

$$G_{\theta}^{\mathrm{sd}} = \prod_{i \in Q_0^{\circ}/\mathbb{Z}_2} \mathrm{GL}(\theta_i) \times \prod_{i \in Q_0^{+}} \mathrm{O}(\theta_i) \times \prod_{i \in Q_0^{-}} \mathrm{Sp}(\theta_i) , \qquad (6.1.3.4)$$

where  $Q_0^{\circ}$  is the set of vertices *i* with  $i \neq i^{\vee}$ , and  $Q_0^{\pm}$  the sets of vertices *i* with  $i = i^{\vee}$  and  $u(i) = \pm 1$ . Similarly,  $Q_1^{\circ}$  is the set of edges *a* with  $a \neq a^{\vee}$ , and  $Q_1^{\pm}$  the sets of edges *a* with  $a = a^{\vee}$  and  $v(a) u(t(a)) = \pm 1$ .

6.1.4. Potentials. Recall that a *potential* on a quiver Q is an element  $W \in KQ/[KQ, KQ]$ , where KQ is the path algebra of Q, and  $[KQ, KQ] \subset KQ$  is the K-linear subspace spanned by commutators. Such an element can be seen as a formal linear combination of cyclic paths in Q, and there is a trace function  $\varphi_W = \operatorname{tr}(W): \mathscr{X}_Q \to \mathbb{A}^1$  defined by taking traces along cyclic paths in a representation. The derived critical locus

$$\mathscr{X}_{Q,W} = \operatorname{Crit}(\varphi_W) \subset \mathscr{X}_Q$$

admits a natural (-1)-shifted symplectic structure, and is a (-1)-shifted symplectic linear stack, equipped with a canonical orientation data.

When Q is equipped with a self-dual structure, the potential W is said to be *self-dual* if it is invariant under the involution of KQ sending a path to its dual path, multiplied by the product of the signs assigned to the edges in the path. In this case, the function  $\varphi_W$  is  $\mathbb{Z}_2$ -invariant, so  $\mathscr{X}_{Q,W}$  is a self-dual linear stack, and the fixed locus  $\mathscr{X}_{Q,W}^{sd}$  admits a natural (-1)-shifted symplectic derived structure and a canonical self-dual orientation data.

When the potential *W* is zero,  $\mathscr{X}_{Q,0} \simeq T^*[-1] \mathscr{X}_Q$  is the (-1)-shifted cotangent stack of the smooth stack  $\mathscr{X}_Q$ , as in §4.2.6, and in particular, its classical truncation coincides with  $\mathscr{X}_Q$ .

**6.1.5. Stability conditions.** A *slope function* on a quiver Q is a map  $\mu: Q_0 \to \mathbb{Q}$ . Given such a map, the *slope* of a dimension vector  $\alpha \in \mathbb{N}^{Q_0} \setminus \{0\}$  is the number

$$\tau(\alpha) = \frac{\sum_{i \in Q_0} \alpha_i \mu(i)}{\sum_{i \in Q_0} \alpha_i}$$

This defines a stability condition on the linear stack  $\mathscr{X}_Q$  in the sense of §2.3.3, where the  $\Theta$ -stratification can be constructed from Ibáñez Núñez [27, Theorem 2.6.3].

If *Q* is equipped with a self-dual structure, then a slope function  $\mu$  is said to be *self-dual* if  $\mu(i^{\vee}) = -\mu(i)$  for all  $i \in Q_0$ . In this case, the corresponding stability condition on Mod(*KQ*)

is self-dual, and the corresponding stability condition on  $\mathcal{X}_O$  is also self-dual.

The above also applies to quivers with potentials. For a potential W on a quiver Q, any slope function  $\tau$  on Q defines a stability condition on  $\mathscr{X}_{Q,W}$ , where the existence of a  $\Theta$ -stratification follows from [27, Theorem 2.6.3]. For a self-dual potential W on a self-dual quiver Q, a self-dual slope function  $\tau$  on Q defines a self-dual stability condition on  $\mathscr{X}_{Q,W}$ .

**6.1.6. DT invariants.** For a quiver Q, a potential W, a slope function  $\tau$  on Q, and a dimension vector  $\alpha \in \mathbb{N}^{Q_0} \setminus \{0\}$ , we have the DT invariants

$$DT_{\alpha}(\tau) \in \mathbb{Q}$$
,  $DT_{\alpha}^{mot}(\tau) \in \widehat{\mathbb{M}}^{mon}(K; \mathbb{Q})$ ,

defined as in §§4.2.4 and 4.3.6, for the (-1)-shifted symplectic linear stack  $\mathcal{X}_{Q,W}$ . These invariants were studied by Joyce and Song [35], Kontsevich and Soibelman [37], and others.

When Q is equipped with a self-dual structure and W,  $\tau$  are self-dual, we have the self-dual DT invariants

$$\mathrm{DT}_{\theta}(\tau) \in \mathbb{Q}, \qquad \mathrm{DT}_{\theta}^{\mathrm{mot}}(\tau) \in \widehat{\mathbb{M}}^{\mathrm{mon}}(K; \mathbb{Q}),$$

defined as in §§4.2.5 and 4.3.6 for the self-dual (-1)-shifted symplectic linear stack  $\mathscr{X}_{Q,W}$ . These are new constructions in this paper.

When the potential *W* is zero, we have  $\mathscr{X}_{Q,0} \simeq T^*[-1] \mathscr{X}_Q$  as in §6.1.4, and the discussions in §§4.2.6 and 4.3.7 apply, which provide more straightforward formulae for the DT invariants.

6.1.7. Wall-crossing formulae. For a self-dual quiver Q with a self-dual potential W, Theorem 5.3.2 applies to the self-dual (-1)-shifted symplectic linear stack  $\mathcal{X}_{Q,W}$ , proving wallcrossing formulae for the DT invariants defined in §6.1.6. We are allowed to take  $\tau_+$ ,  $\tau_-$  in the theorem to be any two self-dual slope functions, since we can take  $\tau_0$  in the theorem to be the trivial stability condition, which is permissible.

6.1.8. An algorithm for computing DT invariants. For a self-dual quiver Q, in the case when the potential W is zero, we describe an algorithm for computing all the invariants  $DT_{\alpha}(\tau)$ ,  $DT_{\alpha}^{mot}(\tau)$ ,  $DT_{\theta}(\tau)$ , and  $DT_{\theta}^{mot}(\tau)$ , for any self-dual slope function  $\tau$ .

First, we compute the motives of  $\mathscr{X}_{\alpha} = V_{\alpha}/G_{\alpha}$  and  $\mathscr{X}_{\theta}^{sd} = V_{\theta}^{sd}/G_{\theta}^{sd}$ , as in (6.1.3.1)–(6.1.3.2), in  $\mathbb{M}(K)$ . We use the relation (3.1.3.3) for the vector bundles  $\mathscr{X}_{\alpha} \to */G_{\alpha}$  and  $\mathscr{X}_{\theta}^{sd} \to */G_{\theta}^{sd}$ , and the motives

$$[*/\mathrm{GL}(n)] = \prod_{i=0}^{n-1} \frac{1}{\mathbb{L}^n - \mathbb{L}^i},$$
(6.1.8.1)

$$[*/\mathcal{O}(2n)] = \mathbb{L}^{n} \cdot \prod_{i=0}^{n-1} \frac{1}{\mathbb{L}^{2n} - \mathbb{L}^{2i}}, \qquad (6.1.8.2)$$

$$[*/O(2n+1)] = [*/Sp(2n)] = \mathbb{L}^{-n} \cdot \prod_{i=0}^{n-1} \frac{1}{\mathbb{L}^{2n} - \mathbb{L}^{2i}}, \qquad (6.1.8.3)$$

where the linear and symplectic cases follow from Joyce [31, Theorem 4.10], as these are *special groups* in the sense there, while the orthogonal cases are due to Dhillon and Young [20, Theorem 3.7]. We then have

$$\int_{\mathcal{X}_{\alpha}} v_{\mathcal{X}}^{\text{mot}} = \mathbb{L}^{-(\dim V_{\alpha} - \dim G_{\alpha})/2} \cdot [\mathcal{X}_{\alpha}] = \mathbb{L}^{(\dim V_{\alpha} + \dim G_{\alpha})/2} \cdot [*/G_{\alpha}], \quad (6.1.8.4)$$

$$\int_{\mathcal{X}_{\theta}^{\mathrm{sd}}} v_{\mathcal{X}^{\mathrm{sd}}}^{\mathrm{mot}} = \mathbb{L}^{-(\dim V_{\theta}^{\mathrm{sd}} - \dim G_{\theta}^{\mathrm{sd}})/2} \cdot [\mathcal{X}_{\theta}^{\mathrm{sd}}] = \mathbb{L}^{(\dim V_{\theta}^{\mathrm{sd}} + \dim G_{\theta}^{\mathrm{sd}})/2} \cdot [*/G_{\theta}^{\mathrm{sd}}], \qquad (6.1.8.5)$$

where  $[*/G_{\alpha}]$  and  $[*/G_{\theta}^{sd}]$  are products of the rational functions in (6.1.8.1)–(6.1.8.3).

Next, we compute the invariants  $DT_{\alpha}^{mot}(0)$  and  $DT_{\theta}^{sd,mot}(0)$  for the trivial slope function 0. These can be obtained from (4.3.7.1)–(4.3.7.2) by substituting in (4.1.2.1) and (4.1.3.1), then using the relations (5.3.7.3)–(5.3.7.4) to reduce to the known integrals (6.1.8.4)–(6.1.8.5). This process also shows that  $DT_{\alpha}^{mot}(0)$  and  $DT_{\theta}^{sd,mot}(0)$  are rational functions in  $\mathbb{L}^{1/2}$ , and evaluating them at  $\mathbb{L}^{1/2} = -1$  gives the numerical invariants  $DT_{\alpha}(0)$  and  $DT_{\theta}^{sd}(0)$ .

Finally, for a general self-dual slope function  $\tau$ , we may apply the wall-crossing formulae (5.3.2.1)–(5.3.2.4) to compute the invariants  $DT_{\alpha}(\tau)$ ,  $DT_{\theta}^{sd}(\tau)$ ,  $DT_{\alpha}^{mot}(\tau)$ , and  $DT_{\theta}^{sd,mot}(\tau)$  from the case when  $\tau = 0$ , which is already known.

As an alternative to the final step, we may first compute the integrals  $\int_{\mathcal{X}_{\alpha}^{ss}(\tau)} v_{\mathcal{X}}^{mot}$  and  $\int_{\mathcal{X}_{\theta}^{sd,ss}(\tau)} v_{\mathcal{X}^{sd}}^{mot}$  using the relations (5.1.3.11)–(5.1.3.12), together with (5.3.7.3)–(5.3.7.4) to reduce to the known integrals (6.1.8.4)–(6.1.8.5), then repeat the process above to obtain the invariants  $DT_{\alpha}^{mot}(\tau)$  and  $DT_{\theta}^{sd,mot}(\tau)$ , which are rational functions in  $\mathbb{L}^{1/2}$ . We then evaluate them at  $\mathbb{L}^{1/2} = -1$  to obtain the numerical invariants  $DT_{\alpha}^{sd}(\tau)$  and  $DT_{\theta}^{sd}(\tau)$ .

The author has implemented the above algorithm using a computer program, and some numerical results are presented below.

**6.1.9. Example. The point quiver.** Consider the point quiver Q with a single vertex and no edges, with the trivial slope function  $\tau = 0$ . There are two self-dual structures on Q, with the signs +1 and -1 assigned to the vertex, respectively.

We have the moduli stacks  $\mathscr{X}_Q = \coprod_{n \ge 0} */\mathrm{GL}(n)$  and  $\mathscr{X}_Q^{\mathrm{sd}} = \coprod_{n \ge 0} */\mathrm{O}(n)$  or  $\coprod_{n \ge 0} */\mathrm{Sp}(2n)$ , depending on the sign of the vertex. As in Joyce and Song [35, Example 7.19], the usual DT invariants of Q are given by

$$\mathrm{DT}_{\mathsf{A}_{n-1}} = \frac{1}{n^2}$$

for all  $n \ge 1$ , where the subscript  $A_{n-1}$  refers to the Dynkin type of GL(n).

Based on explicit computation following the algorithm in §6.1.8, we conjecture that

$$DT_{B_n}^{sd} = DT_{C_n}^{sd} = (-1)^n {\binom{-1/4}{n}}, \qquad DT_{D_n}^{sd} = (-1)^n {\binom{1/4}{n}},$$

where the subscripts  $B_n$ ,  $C_n$ , and  $D_n$  refer to the Dynkin types of O(2n+1), Sp(2n), and O(2n),

respectively. Equivalently, we have the generating series

$$\sum_{n \ge 0} q^n \cdot \mathrm{DT}_{\mathsf{B}_n}^{\mathsf{sd}} = \sum_{n \ge 0} q^n \cdot \mathrm{DT}_{\mathsf{C}_n}^{\mathsf{sd}} = (1-q)^{-1/4}, \qquad \sum_{n \ge 0} q^n \cdot \mathrm{DT}_{\mathsf{D}_n}^{\mathsf{sd}} = (1-q)^{1/4}$$

We expect to prove this conjecture in a future paper [16], and we expect that the coincidence of the type B and C invariants here should be related to the fact that these groups are Langlands dual to each other.

6.1.10. Example. The  $\tilde{A}_1$  quiver. Let  $Q = (\bullet \Rightarrow \bullet)$  be the quiver with two vertices and two arrows pointing in the same direction, called the  $\tilde{A}_1$  quiver. Consider the contravariant involution of Q that exchanges the two vertices but fixes the edges. We use the simplified notation  $\tilde{A}_1^{u,v}$ , where u, v are the signs in the self-dual structure. For example,  $\tilde{A}_1^{+,++}$  means that we take the sign +1 on all vertices and edges. Note that both vertices must have the same sign. We use the slope function  $\tau = (1, -1)$ .

Based on numerical evidence from applying the algorithm in §6.1.8, we conjecture that we have the generating series

$$\int_{-\infty}^{\infty} n/2 = -\operatorname{sd} \operatorname{mot}\left(\sum_{i=1}^{\infty} (1-q^{1/2} \mathbb{L}^{-1/2})(1-q^{1/2} \mathbb{L}^{1/2})\right) \text{ for } \tilde{A}_{1}^{+,++} \text{ and } \tilde{A}_{1}^{-,--}, \quad (6.1.10.1)$$

$$\sum_{n=0}^{\infty} q^{n/2} \cdot \mathrm{DT}_{(n,n)}^{\mathrm{sd},\mathrm{mot}}(\tau) = \begin{cases} \left(\frac{1+q^{1/2}}{1-q^{1/2}}\right)^{1/2} & \text{for } \tilde{A}_1^{+,+-} \text{ and } \tilde{A}_1^{-,+-} , \quad (6.1.10.2) \\ (1-q)^{1/2} & \text{for } \tilde{A}_1^{+,--} \text{ and } \tilde{A}_1^{-,++} . \quad (6.1.10.3) \end{cases}$$

This example is related to coherent sheaves on  $\mathbb{P}^1$ , as we will discuss in Example 6.3.5.

#### 6.2 Orthosymplectic complexes

**6.2.1.** In the following sections, §§6.3–6.5, we will apply our theory to study orthosymplectic DT invariants for certain smooth projective varieties over  $\mathbb{C}$ , in three variants which apply to curves, surfaces, and threefolds, respectively. This section will provide background material that is common to these settings.

Our DT invariants will count *orthogonal* or *symplectic complexes* on a variety, which are perfect complexes equipped with isomorphisms to their dual complexes.

We note that this approach of defining a coherent-sheaf-like version of principal bundles is different from the related construction of Gómez, Fernandez Herrero, and Zamora [23].

**6.2.2.** The setting. Throughout, we work over the complex number field  $\mathbb{C}$ , and we fix a connected, smooth, projective  $\mathbb{C}$ -variety *Y* of dimension *n*.

Let  $\bar{\mathcal{X}}$  be the derived moduli stack of perfect complexes on Y, as in Toën and Vaquié [50]. It is a derived algebraic stack locally of finite presentation over  $\mathbb{C}$ . By Pantev, Toën, Vaquié, and Vezzosi [40], if Y is a *Calabi–Yau n-fold*, meaning that its canonical bundle  $K_Y$  is trivial, then  $\bar{\mathcal{X}}$  has a (2 - n)-shifted symplectic structure. We fix the data  $(I, L, s, \varepsilon)$ , where  $I: Y \xrightarrow{\sim} Y$  is an involution,  $L \to Y$  is a line bundle,  $s \in \mathbb{Z}$ , and  $\varepsilon: L \xrightarrow{\sim} I^*(L)$  is an isomorphism such that  $I^*(\varepsilon) \circ \varepsilon = \mathrm{id}_L$ . Define a self-dual structure on  $\mathrm{Perf}(Y)$  by the dual functor

$$\mathbb{D} = \mathbb{R}\mathscr{H}om(I^*(-), L)[s] \colon \operatorname{Perf}(Y) \xrightarrow{\sim} \operatorname{Perf}(Y)^{\operatorname{op}}, \qquad (6.2.2.1)$$

and identify  $\mathbb{D}(\mathbb{D}(E))$  with *E* using the isomorphism  $\varepsilon$ , for all objects  $E \in \text{Perf}(Y)$ .

This self-dual structure induces a  $\mathbb{Z}_2$ -action on  $\bar{\mathcal{X}}$ , and the fixed locus  $\bar{\mathcal{X}}^{sd}$  is the moduli stack of self-dual perfect complexes on Y. In particular, when  $I = id_Y$  and  $\varepsilon = \pm id_L$ , the stack  $\bar{\mathcal{X}}^{sd}$  parametrizes L[s]-twisted orthogonal or symplectic complexes on Y, respectively. When  $L = \mathcal{O}_Y$  and s = 0, they are simply called orthogonal or symplectic complexes.

6.2.3. Bridgeland stability conditions. Consider the free abelian group

$$K(Y) = \{ \operatorname{ch}(E) \mid E \in \operatorname{Perf}(Y) \} \subset \operatorname{H}^{2\bullet}(Y; \mathbb{Q}) .$$
(6.2.3.1)

It has an involution  $(-)^{\vee}$  given by  $ch(E) \mapsto ch(\mathbb{D}(E))$ . Let  $K^{sd}(Y) \subset K(Y)$  be the fixed locus.

Define a *Bridgeland stability condition* on *Y* as in Bridgeland [8, Definition 5.1], which is a pair  $\tau = (Z, \mathcal{P})$ , where  $Z \colon K(Y) \to \mathbb{C}$  is a group homomorphism, and  $\mathcal{P}$  is a slicing of Perf(*Y*).

Let Stab(Y) be the set of Bridgeland stability conditions on *Y*, which has a topology given by a *generalized metric d*, that is, a metric allowing infinite distance, defined as in [8, §8] by

$$d(\tau, \tilde{\tau}) = \sup\left\{ |\phi^+(E) - \tilde{\phi}^+(E)|, |\phi^-(E) - \tilde{\phi}^-(E)|, |\log m(E) - \log \tilde{m}(E)| \mid E \neq 0 \right\}, \quad (6.2.3.2)$$

where *E* runs through all non-zero objects of Perf(*Y*), and  $\phi^+(E)$ ,  $\phi^-(E)$ , m(E) are the maximal phase, the minimal phase, and the sum of lengths of central charges of the  $\tau$ -HN factors of *E*, respectively, and similarly for  $\tilde{\tau}$ . The projection

$$\operatorname{Stab}(Y) \longrightarrow \operatorname{Hom}(K(Y), \mathbb{C})$$
 (6.2.3.3)

given by  $(Z, \mathscr{P}) \mapsto Z$  is a local homeomorphism, and equips Stab(Y) with the structure of a complex manifold.

The self-dual structure on Perf(*Y*) defined in §6.2.2 gives an anti-holomorphic involution  $(-)^{\vee}$ : Stab(*Y*)  $\rightarrow$  Stab(*Y*), given by  $(Z, \mathscr{P}) \mapsto (Z^{\vee}, \mathscr{P}^{\vee})$ , where  $Z^{\vee}(\alpha) = \overline{Z(\alpha^{\vee})}$  and  $\mathscr{P}^{\vee}(t) = \mathscr{P}(-t)^{\vee}$ . The fixed locus Stab<sup>sd</sup>(*Y*)  $\subset$  Stab(*Y*) is the set of *self-dual stability conditions*, and we have a local homeomorphism

$$\operatorname{Stab}^{\operatorname{sd}}(Y) \longrightarrow \operatorname{Hom}(K(Y), \mathbb{C})^{\mathbb{Z}_2},$$
 (6.2.3.4)

given by  $(Z, \mathscr{P}) \mapsto Z$ , where  $\mathbb{Z}_2$  acts on  $\text{Hom}(K(Y), \mathbb{C})$  via the anti-holomorphic involution  $Z \mapsto Z^{\vee}$ . This equips  $\text{Stab}^{\text{sd}}(Y)$  with the structure of a real analytic manifold.

6.2.4. Permissibility. We define subspaces of permissible Bridgeland stability conditions,

$$\operatorname{Stab}^{\circ}(Y) \subset \operatorname{Stab}(Y)$$
,  $\operatorname{Stab}^{\circ,\operatorname{sd}}(Y) \subset \operatorname{Stab}^{\operatorname{sd}}(Y)$ 

as maximal open subsets such that every element  $\tau = (Z, \mathscr{P})$  with  $Z(K(Y)) \subset \mathbb{Q} + i\mathbb{Q}$  satisfies the following conditions:

- (i) *Support property*. For any r > 0, there are only finitely many classes  $\alpha \in K(Y)$  admitting a semistable object, such that  $|Z(\alpha)| \leq r$ .
- (ii) *Generic flatness.* See Abramovich and Polishchuk [1, Problem 3.5.1], Halpern-Leistner [24, Definition 6.2.4], or Piyaratne and Toda [41, Definition 4.4] for the formulation.
- (iii) Boundedness. For any  $t \in \mathbb{R}$  and  $\alpha \in K(Y)$  with  $Z(\alpha) \in \mathbb{R}_{\geq 0} \cdot e^{\pi i t}$ , there is a quasi-compact open substack  $\mathscr{X}(\tau; t)_{\alpha} \subset \overline{\mathscr{X}}$  whose  $\mathbb{C}$ -points are the objects of  $\mathscr{P}(t)$  of class  $\alpha$ .

By Piyaratne and Toda [41, Proposition 4.12], if a stability condition  $\tau$  satisfies these conditions and has rational central charge, then a neighbourhood of  $\tau$  lies in Stab<sup>°</sup>(*Y*).

For  $\tau \in \text{Stab}^{\circ}(Y)$  and an interval  $J \subset \mathbb{R}$  of length |J| < 1, there is an open substack

$$\mathscr{X}(\tau;J) \subset \bar{\mathscr{X}}$$

whose  $\mathbb{C}$ -points are the objects of  $\mathscr{P}(J)$ , which we construct in §6.2.6 below. It is a derived linear stack in the sense of §4.2.3, and  $\tau$  defines a permissible stability condition on its classical truncation in the sense of §2.3, where the  $\Theta$ -stratification is constructed in §6.2.6 below.

In particular, if  $\tau \in \text{Stab}^{\circ,\text{sd}}(Y)$  and J = -J, then  $\mathscr{X}(\tau; J)$  is a self-dual derived linear stack, and the induced stability condition on  $\mathscr{X}(\tau; J)$  is self-dual. The stack  $\mathscr{X}(\tau; 0)^{\text{sd}}$  is the moduli stack of  $\tau$ -semistable self-dual complexes, which our orthosymplectic DT invariants will count.

**6.2.5.** Example. Let *Y* be either a curve, a surface, or a threefold satisfying the conjectural Bogomolov–Gieseker inequality of Bayer, Macrì, and Toda [4, Conjecture 3.2.7], and fix the data (*I*, *L*, *s*,  $\varepsilon$ ) as in §4.3.4. Suppose we are given an ample class  $\omega \in H^{1,1}(Y; \mathbb{Q})$  with  $I^*(\omega) = \omega$ .

In this case, we give an example of a permissible self-dual Bridgeland stability condition, with central charge valued in  $\mathbb{Q} + i\mathbb{Q}$ .

Let  $\beta = c_1(L)/2 \in \mathrm{H}^2(Y; \mathbb{Q})$ . Consider the group homomorphism  $Z_\omega \colon K(Y) \to \mathbb{C}$  given by

$$Z_{\omega}(\alpha) = i^{n-s} \cdot \int_{Y} \exp(-\beta - i\omega) \cdot \alpha$$
(6.2.5.1)

for  $\alpha \in K(Y)$ , where  $n = \dim Y$ . This is compatible with the self-dual structure, in the sense that we have  $Z_{\omega}(\mathbb{D}(\alpha)) = \overline{Z_{\omega}(\alpha)}$  for all  $\alpha \in K(Y)$ . Here, our coefficient  $i^{n-s}$  is only inserted to make  $Z_{\omega}$  self-dual, and does not essentially affect the stability condition.

There is a Bridgeland stability condition  $\tau_{\omega} = (Z_{\omega}, \mathscr{P}_{\omega}) \in \operatorname{Stab}^{\circ}(Y)$  with central charge  $Z_{\omega}$ , by the works of Toda [49] and Piyaratne and Toda [41]. See also the earlier works of Bridgeland [9] and Arcara and Bertram [3] in the case of surfaces. In fact, we can also choose  $\mathscr{P}_{\omega}$  so that  $\tau_{\omega}$  is self-dual, or equivalently, the slicing  $\mathscr{P}_{\omega}$  coincides with its dual slicing  $\mathscr{P}_{\omega}^{\vee}$  given by

$$\mathscr{P}^{\vee}_{\omega}(t) = \mathbb{D}(\mathscr{P}_{\omega}(-t))$$
.

This follows from Bayer, Macrì, and Toda [4, Remark 4.4.3], which is essentially the same statement in the case when s = 1, and the general case is constructed from this case by simply shifting the phase by (1 - s)/2.

**6.2.6.** In the situation of §6.2.4, for  $\tau \in \text{Stab}^{\circ}(Y)$  and an interval  $J \subset \mathbb{R}$  with |J| < 1, we construct the open substack  $\mathcal{X}(\tau; J) \subset \overline{\mathcal{X}}$  and its  $\Theta$ -stratification by  $\tau$ -HN types as follows.

Applying Piyaratne and Toda [41, Proposition 4.12], we may apply a phase shift and assume that  $J \subset ]\varepsilon, 1 - \varepsilon[$  for some  $\varepsilon > 0$ . Fix  $\alpha \in K(Y)$  of slope within J, and then choose a perturbation  $\tau' = (Z', \mathscr{P}')$  of  $\tau$  satisfying the above properties, with  $d(\tau', \tau) < \varepsilon$  and  $Z'(K(Y)) \subset \mathbb{Q}+i\mathbb{Q}$ . Then if  $\beta \in K(Y)$  is the class of a  $\tau$ -HN factor of an object of  $\mathscr{P}(J)$  of class  $\alpha$ , then  $Z'(\beta)$  must lie in the bounded region

$$\left\{r\mathrm{e}^{\pi\mathrm{i}t}\mid r\geq 0,\ t\in J_{\varepsilon}\right\}\cap\left\{Z'(\alpha)-r\mathrm{e}^{\pi\mathrm{i}t}\mid r\geq 0,\ t\in J_{\varepsilon}\right\}\subset\mathbb{C}\,,$$

where  $J_{\varepsilon}$  is the  $\varepsilon$ -neighbourhood of J, so the set B of such classes  $\beta$  is finite. We then choose  $\varepsilon$ small enough, possibly changing  $\tau'$ , so that for any  $\beta, \beta' \in B$ ,  $\arg Z(\beta) < \arg Z(\beta')$  implies  $\arg Z'(\beta) < \arg Z'(\beta')$ , where we take phases within  $J_{\varepsilon}$ . Now, Halpern-Leistner [24, Theorem 6.5.3] gives the open substack  $\mathcal{X}(\tau'; ]0, 1[)$  with a  $\Theta$ -stratification by  $\tau'$ -HN types. The part of  $\mathcal{X}(\tau; J)$  lying in  $\mathcal{X}_{\alpha}$  can be defined as a finite open union of strata.

To construct the  $\Theta$ -stratification on  $\mathcal{X}(\tau; J)$ , we follow the proof of [24, Theorem 6.5.3], with the following modifications. Instead of using rational weights for HN filtrations, we use *real-weighted filtrations* in the sense of [14, §§7.2–7.3]. As a result, we obtain real-weighted  $\Theta$ -stratifications, which non-canonically give usual  $\Theta$ -stratifications by [14, Proposition 7.2.12]. The key ingredients of the proof in [24] are the conditions (R), (S), and (B) there. The rationality condition (R) is no longer needed as we use real weights. The condition (S) needs to be modified to incorporate real weights, but the argument still works to prove it. The condition (B) follows from the quasi-compactness of  $\mathcal{X}(\tau; J)$ .

This also shows that any  $\tau \in \operatorname{Stab}^{\circ}(Y)$  satisfies the support property and the boundedness property in §6.2.4, where for the support property, fixing r > 0 and choosing  $\tau'$  rational with  $d(\tau', \tau) < \varepsilon$  with  $\varepsilon < 1/2$ , for any class  $\alpha$  with  $|Z(\alpha)| \leq r$  admitting a  $\tau$ -semistable object E, by considering the  $\tau'$ -HN filtration of E, we see that  $\alpha$  is a finite sum of classes  $\beta$  with  $|Z'(\beta)| < re^{\varepsilon}$ admitting  $\tau'$ -semistable objects, and these classes lie on the same side of a line in  $\mathbb{C}$ , so there are only finitely many choices.

#### 6.3 DT invariants for curves

**6.3.1.** We define DT invariants counting orthogonal and symplectic bundles on a curve. These are orthosymplectic versions of Joyce's motivic invariants counting vector bundles on a curve, as in [32, §6.3].

**6.3.2.** Let *C* be a connected, smooth, projective curve over  $\mathbb{C}$ , and fix the data (*I*, *L*, *s*,  $\varepsilon$ ) as in §6.2.2. This defines a self-dual structure on Perf(*C*).

Let  $\tau = (Z, \mathscr{P})$  be the Bridgeland stability condition defined in Example 6.2.5, where we choose the unique element  $\omega \in H^2(C; \mathbb{Q})$  with  $\int_C \omega = 1$ . Explicitly, we have

$$Z(E) = \mathbf{i}^{-s} \cdot \left( \left( 1 - \mathbf{i} \, \frac{\deg L}{2} \right) r + d \right) \tag{6.3.2.1}$$

for  $E \in \text{Perf}(C)$  with rank r and degree d, so that  $ch(E) = r + d\omega$ . Note that the choices of L and s do not affect which objects are semistable, although they affect which objects are self-dual. The subcategory  $\text{Vect}(C) \subset \text{Perf}(C)$  of vector bundles on C satisfies  $\text{Vect}(C) = \mathscr{P}(](-1-s)/2, (1-s)/2[)$ .

6.3.3. The even case. When *s* is even, the abelian category  $\mathscr{P}(0)$  consists of objects E[s/2] for semistable vector bundles *E* on *C* in the usual sense, whose rank *r* and degree *d* satisfy  $d = r \deg L/2$ . The self-dual objects are such *E* with isomorphisms  $\phi: E \xrightarrow{\sim} \mathscr{H}om(I^*(E), L)$  with  $I^*(\phi)^{\vee} \circ \phi = (-1)^{s/2} \cdot \varepsilon$ .

In particular, when  $L = \mathcal{O}_C$ , semistable self-dual complexes can be identified, up to a shift, with orthogonal or symplectic bundles on *C*, depending on whether  $(-1)^{s/2} \cdot \varepsilon = 1$  or -1, whose underlying vector bundles are semistable in the usual sense.

For each rank r > 0, we have the self-dual DT invariants

$$\mathrm{DT}_r^{\mathrm{sd}} \in \mathbb{Q}$$
,  $\mathrm{DT}_r^{\mathrm{sd},\mathrm{mot}} \in \widehat{\mathbb{M}}^{\mathrm{mon}}(K;\mathbb{Q})$ ,

counting semistable self-dual vector bundles of rank r as above, defined as in §§4.2.6 and 4.3.7 using the self-dual linear stack  $\mathscr{X}(\tau; 0)$  defined in §6.2.4 with the trivial stability condition.

6.3.4. The odd case. When *s* is odd,  $\mathscr{P}(0)$  consists of objects E[(s-1)/2] for torsion sheaves *E* on *C*, and the semistable self-dual objects are such *E* with isomorphisms  $\phi : E \xrightarrow{\sim} \mathbb{R}\mathscr{H}om(I^*(E), L[1])$  with  $I^*(\phi)^{\vee} \circ \phi = (-1)^{(s-1)/2} \cdot \varepsilon$ . For each degree d > 0, we have the self-dual DT invariants

$$\mathrm{DT}_{0,d}^{\mathrm{sd}} \in \mathbb{Q}$$
,  $\mathrm{DT}_{0,d}^{\mathrm{sd},\mathrm{mot}} \in \widehat{\mathbb{M}}^{\mathrm{mon}}(K;\mathbb{Q})$ ,

counting these self-dual torsion sheaves, defined similarly as above.

In fact, these invariants do not depend on the choice of L, since choosing a suitable Iinvariant open cover of C trivializing L, torsion sheaves supported on the open sets give an open cover of the moduli stacks, where pieces and intersections do not depend on L. It then



Figure 1. An equivalence of categories

follows from [15, Theorem 5.2.10 (i)] that the invariants do not depend on *L*.

6.3.5. Example. Invariants for  $\mathbb{P}^1$ . Consider the case when  $C = \mathbb{P}^1$  and  $I = id_{\mathbb{P}^1}$ . We describe the invariants in two situations.

When s = 0 and  $L = \mathcal{O}_{\mathbb{P}^1}$ , since every vector bundle on  $\mathbb{P}^1$  splits as a direct sum of line bundles, all semistable vector bundles of slope 0 are trivial bundles. The self-dual abelian category  $\mathscr{P}(0)$  is thus equivalent to the category of finite-dimensional  $\mathbb{C}$ -vector spaces, with one of the two self-dual structures described in Example 6.1.9, depending on the sign  $\varepsilon$ . The DT invariants agree with the ones given there.

When s = 1, invariants for  $\mathbb{P}^1$  are related to DT invariants for self-dual quivers. Indeed, as a special case of Bondal [7, Theorem 6.2], we have an equivalence

$$\Phi \colon \mathsf{Perf}(\mathbb{P}^1) \xrightarrow{\sim} \mathsf{D}^{\mathsf{b}}\mathsf{Mod}(\mathbb{C}Q) , \qquad (6.3.5.1)$$

where *Q* is the  $\tilde{A}^1$  quiver in Example 6.1.10, and  $\Phi(E) = (\mathbb{R}\Gamma(\mathbb{P}^1, E(-1)) \rightrightarrows \mathbb{R}\Gamma(\mathbb{P}^1, E))$ , with the two maps given by multiplying with the coordinate functions  $x_0, x_1 \in \Gamma(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(1))$ .

In fact, under the isomorphism  $\Phi$ , the self-dual structure on  $\operatorname{Perf}(\mathbb{P}^1)$  given by  $(I = \operatorname{id}_{\mathbb{P}^1}, L = \mathcal{O}_{\mathbb{P}^1}(-1), s = 1, \varepsilon)$  corresponds to the self-dual structure on  $\operatorname{D^bMod}(\mathbb{C}Q)$  given by the signs  $(\varepsilon, ++)$  in the notation of Example 6.1.10, as shown in Figure 1. Here, r and d denote the rank and degree of a complex on  $\mathbb{P}^1$ , and  $(d_0, d_1)$  is the dimension vector of a representation of Q. The two-way arrows indicate the dual operation, and the self-dual objects lie on the vertical axis on the left-hand side, or the horizontal axis on the right-hand side. The shaded regions indicate the usual heart of  $\operatorname{D^bMod}(\mathbb{C}Q)$  and the corresponding heart of  $\operatorname{Perf}(\mathbb{P}^1)$ . The right-hand side can also be viewed either as the central charge of  $\tau$ , or that of the stability condition on Q given by the slope function (1, -1).

In particular, the DT invariants in this case coincide with those in Example 6.1.10, and should be given by the conjectural formulae (6.1.10.1) and (6.1.10.3).

#### 6.4 DT invariants for threefolds

**6.4.1.** We define DT invariants counting orthogonal or symplectic complexes on a Calabi–Yau threefold. These invariants are one of the main applications of our theory, and are an extension of the usual DT invariants studied by Thomas [47], Joyce and Song [35], Kontsevich and Soibelman [37], and many others. We expect our invariants to be related to counting D-branes on Calabi–Yau 3-orientifolds, as discussed in Witten [52, §5.2], Diaconescu, Garcia-Raboso, Karp, and Sinha [21], and Hori and Walcher [26].

We also prove wall-crossing formulae for these invariants in Theorem 6.4.3, which relate the invariants for different Bridgeland stability conditions.

**6.4.2. Invariants.** Let *Y* be a Calabi–Yau threefold over  $\mathbb{C}$ , and fix the data (*I*, *L*, *s*,  $\varepsilon$ ) as in §6.2.2.

In this case, the derived stack  $\bar{\mathcal{X}}$  and the fixed locus  $\bar{\mathcal{X}}^{sd}$  are (-1)-shifted symplectic stacks. The stack  $\bar{\mathcal{X}}$  has an orientation data in the sense of §4.3.4, by Joyce and Upmeier [36, Theorem 3.6]. However, we do not know if the stack  $\bar{\mathcal{X}}^{sd}$  has an orientation in general.

Let  $\tau = (Z, \mathscr{P}) \in \operatorname{Stab}^{\circ, \operatorname{sd}}(Y)$  be a self-dual Bridgeland stability condition on Y, which is guaranteed to exist in the situation of Example 6.2.5. For each phase  $t \in \mathbb{R}$ , let  $\mathscr{X}(\tau; t) \subset \overline{\mathscr{X}}$  be the open substack as in §6.2.4, which is a (-1)-shifted symplectic linear stack, and has a self-dual structure when t = 0.

Given a class  $\alpha \in K(Y)$  with  $Z_{\omega}(\alpha) \in \mathbb{R}_{>0} \cdot e^{\pi i t}$ , define the numerical and motivic DT invariants

$$DT_{\alpha}(\tau) \in \mathbb{Q}$$
,  $DT_{\alpha}^{mot}(\tau) \in \hat{\mathbb{M}}^{mon}(\mathbb{C};\mathbb{Q})$ ,

as in §§4.2.4 and 4.3.6 for the stack  $\mathscr{X}(\tau;t)$  with the trivial stability condition, where we take the sum of DT invariants of connected components of the open and closed substack  $\mathscr{X}(\tau;t)_{\alpha} \subset \mathscr{X}(\tau;t)$ , and we use the orientation of Joyce and Upmeier [36] for the motivic version. These invariants are not new, and can be constructed from the formalisms of Joyce and Song [35] and Kontsevich and Soibelman [37].

When t = 0, for each  $\theta \in K^{sd}(Y)$  with  $Z_{\omega}(\theta) \in \mathbb{R}_{>0}$ , we have the numerical self-dual DT invariant

$$\mathrm{DT}^{\mathrm{sd}}_{\theta}(\tau) \in \mathbb{Q}$$
,

defined as in §4.2.5 for the self-dual linear stack  $\mathscr{X}(\tau; 0)$  with the trivial stability condition, where we sum over connected components of  $\mathscr{X}(\tau; 0)^{sd}_{\theta}$ . These are new invariants for Calabi–Yau threefolds, and are one of the main constructions of this paper.

If one can construct a self-dual orientation data on  $\mathscr{X}(\tau; 0)$  in the sense of §4.3.5, then the motivic self-dual DT invariant  $DT_{\theta}^{\text{mot,sd}}(\tau)$  will also be defined, as in §4.3.6.

**6.4.3. Theorem.** Let Y be a Calabi–Yau threefold over  $\mathbb{C}$ . Choose the data  $(I, L, s, \varepsilon)$  as in §6.2.2. Let  $\tau = (Z, \mathcal{P}), \tilde{\tau} = (\tilde{Z}, \tilde{\mathcal{P}}) \in \operatorname{Stab}^{\circ}(Y)$  be Bridgeland stability conditions.

- (i) If  $\tau$ ,  $\tilde{\tau}$  can be connected by a path of length < 1/4 in Stab<sup>°</sup>(Y), then for any class  $\alpha \in K(Y)$  with  $Z(\alpha) \neq 0$ , the wall-crossing formula (5.3.2.1) holds.
- (ii) If  $\tau, \tilde{\tau} \in \text{Stab}^{\circ, \text{sd}}(Y)$ , and they can be connected by a path of length < 1/4 in  $\text{Stab}^{\circ, \text{sd}}(Y)$ , then for any class  $\theta \in K^{\text{sd}}(Y)$  with  $Z(\theta) \in \mathbb{R}_{>0}$ , the wall-crossing formula (5.3.2.2) holds.

Here, the length of a path is defined as the supremum of sums of distances over all subdivisions. In (5.3.2.1)–(5.3.2.2), we use  $\tau, \tilde{\tau}$  in place of  $\tau_+, \tau_-$ . The sets  $\pi_0(\mathcal{X}), \pi_0(\mathcal{X}^{sd})$  in the formulae are defined using  $\mathcal{X} = \mathcal{X}(\tau; ]t - 1/4, t + 1/4[)$ , where t is a phase of  $Z(\alpha)$  in (i) or t = 0 in (ii). The coefficients  $\tilde{U}(\ldots), \tilde{U}^{sd}(\ldots)$  are defined using the total order on phases in ]t - 1/2, t + 1/2[.

Moreover, if we are given an orientation data on  $\mathcal{X}(\tau; ]t - 1/2, t + 1/2[)$ , or a self-dual orientation data on  $\mathcal{X}(\tau; ]-1/2, 1/2[)$ , respectively, then (i)–(ii) also hold for the motivic versions (5.3.2.3)–(5.3.2.4), where  $\alpha$  has phase t.

**Proof.** To avoid repetition, we prove (i)–(ii) using a common argument, where we write  $\alpha$  for  $\theta$  for (ii).

We first prove the following claim: For fixed  $\tau$  and a fixed class  $\alpha$  or  $\theta$ , there exists  $\delta > 0$ such that the wall-crossing formulae hold whenever  $d(\tau, \tilde{\tau}) < \delta$ , with the sets  $\pi_0(\mathcal{X}), \pi_0(\mathcal{X}^{sd})$ defined using  $\mathcal{X} = \mathcal{X}(\tau; t)$ , and we may take  $\tau_+, \tau_-$  in the formulae to be either  $\tau, \tilde{\tau}$  or  $\tilde{\tau}, \tau$ .

Write  $\mathscr{A} = \mathscr{P}(t)$ . Let  $K \subset K(Y)$  be the set of Chern characters of  $\tau$ -semistable objects in Perf(*Y*), and  $C \subset K$  the set of classes realized by objects in  $\mathscr{A}$ .

We choose  $0 < \delta < 1/8$  such that  $K \cap Z^{-1}(V_{4\delta}(e^{2\delta} \cdot Z(\alpha))) \subset C$ , where

$$V_u(z) = \{ r e^{\pi i \phi} \mid 0 \leq r \leq |z|, \ |\phi| \leq u \} \subset \mathbb{C} .$$

If  $\beta \in K(Y)$  is the class of a  $\tau$ -HN factor of a  $\tilde{\tau}$ -semistable object of class  $\alpha$ , then  $Z(\beta)$  must lie in  $V_{2\delta}(Z(\alpha))$ . By the choice of  $\delta$ , all such classes  $\beta$  have phase t, and are hence equal to  $\alpha$ . This implies that all  $\tilde{\tau}$ -semistable objects of class  $\alpha$  are  $\tau$ -semistable and are in  $\mathcal{A}$ .

Similarly, we may assume that all  $\tilde{\tau}$ -semistable objects with phase in  $[t - \delta, t + \delta]$  and norm  $\leq e^{\delta} \cdot |Z(\alpha)|$  are in  $\mathscr{A}$ . Indeed, such objects have  $\tau$ -phase in  $[t - 2\delta, t + 2\delta]$  and  $\tau$ -norm  $\leq e^{2\delta} \cdot |Z(\alpha)|$ , and this property holds by the choice of  $\delta$ .

It follows that for any object in  $\mathscr{A}$  of class  $\alpha$ , its  $\tilde{\tau}$ -HN factors also belong to  $\mathscr{A}$ . In other words,  $\tilde{\tau}$  almost defines a stability condition on  $\mathscr{X}(\tau; t)$  in the sense of §2.3.3, except that the  $\Theta$ -stratification is only defined on  $\mathscr{X}(\tau; t)_{\beta}$  for classes  $\beta \in C$  with  $|Z(\beta)| \leq |Z(\alpha)|$ . However, this is enough to prove wall-crossing for  $\alpha$ , as the other classes are irrelevant in the argument. The claim thus follows from Theorem 5.3.2, where  $\tau$  corresponds to trivial stability on  $\mathscr{X}(\tau; t)$ .

We now turn to the original statement of the theorem. Choose a path  $(\tau_s = (Z_s, \mathcal{P}_s))_{s \in [0,1]}$ of length  $\ell < 1/4$ , with  $\tau_0 = \tau$  and  $\tau_1 = \tilde{\tau}$ . By the compactness of [0, 1], our claim implies that we can choose  $0 = s_0 < \cdots < s_n = 1$  such that there are wall-crossing formulae between each  $\tau_{s_i}$  and  $\tau_{s_{i+1}}$ . We may thus apply (5.3.2.1), etc., to express  $DT_{\alpha}(\tau_{s_0})$ , etc., in terms of invariants for  $\tau_{s_1}$ , and so on, finally in terms of invariants for  $\tau_{s_n} = \tilde{\tau}$ . In each step, the involved invariants  $\mathrm{DT}_\beta(\tau_{\mathbf{s}_i})$  must satisfy that  $Z(\beta)$  lies in the bounded region

$$\{re^{\pi i\phi} \mid r \ge 0, \ |\phi - t| \le \ell\} \cap \{Z(\alpha) - re^{\pi i\phi} \mid r \ge 0, \ |\phi - t| \le \ell\} \subset \mathbb{C},$$

so that the sums (5.3.2.1), etc., can not only be written using some  $\pi_0(\mathcal{X}(\tau_{s_i}; t_i))$  and its self-dual version, as in the argument above, but also using the larger set  $\pi_0(\mathcal{X}(\tau; ]t - 1/2, t + 1/2[))$  and its self-dual version, where the coefficients  $\tilde{U}(\ldots)$ ,  $\tilde{U}^{sd}(\ldots)$  are zero for the newly introduced terms. The support property of  $\tau$  ensures that only finitely many non-zero terms appear in each step.

It remains to prove that the coefficients  $\tilde{U}(...)$ ,  $\tilde{U}^{sd}(...)$  respect composition of wallcrossing formulae, so that the wall-crossing formulae obtained from the above process are equivalent to (5.3.2.1), etc., from  $\tau$  directly to  $\tilde{\tau}$ . This follows from the fact that the coefficients S(...),  $S^{sd}(...)$  respect composition, which was proved in Lemma 5.1.4.

6.4.4. Generic stability conditions. Following Joyce and Song [35, Conjecture 6.12], we say that a stability condition  $\tau$  as above is *generic*, if for any  $\alpha, \beta \in K(Y)$  with  $Z(\alpha) = \lambda Z(\beta) \neq 0$  for some  $\lambda \in \mathbb{R}_{>0}$ , we have the numerical condition vdim  $\overline{\mathcal{X}}^+_{\alpha,\beta} = 0$ .

Similarly, when  $\tau$  is self-dual, we say that it is *generic* as a self-dual stability condition, if it is generic as above, and for any  $\alpha \in K(Y)$  of phase 0 and  $\theta \in K^{sd}(Y)$ , we have vdim  $\bar{\mathcal{X}}_{\alpha,\theta}^{sd,+} = 0$ .

By the first part in the proof of Theorem 6.4.3, combined with Corollary 5.3.4, we see that if  $\tau \in \text{Stab}^{\circ,\text{sd}}(Y)$  is generic, then for each class  $\alpha$  or  $\theta$  as in Theorem 6.4.3, there exists  $\delta > 0$ , such that the invariant  $\text{DT}_{\alpha}(\tau)$  or  $\text{DT}_{\theta}^{\text{sd}}(\tau)$  does not change if we move  $\tau$  inside its  $\delta$ -neighbourhood. Moreover, this also holds for the motivic versions  $\text{DT}_{\alpha}^{\text{mot}}(\tau)$  or  $\text{DT}_{\theta}^{\text{mot},\text{sd}}(\tau)$ , where the self-dual version requires a self-dual orientation data.

6.4.5. Expectations on deformation invariance. We expect that the numeric version of the orthosymplectic DT invariants,  $DT_{\theta}^{sd}(\tau)$ , should satisfy *deformation invariance*, analogously to Joyce and Song [35, Corollary 5.28] in the linear case, that is, they should stay constant under deformations of the complex structure of the threefold *Y*. However, we have not yet been able to prove this, as it does not seem straightforward to adapt the strategy of [35] using Joyce–Song pairs to our case, and further work is needed.

We do not expect the motivic version,  $DT_{\theta}^{sd,mot}(\tau)$ , to satisfy deformation invariance.

#### 6.5 Vafa-Witten type invariants for surfaces

**6.5.1.** We construct a motivic version of orthosymplectic analogues of *Vafa–Witten invariants* for algebraic surfaces, studied by Tanaka and Thomas [45; 46], Maulik and Thomas [39], and Thomas [48]. We define our invariants for surfaces *S* with  $K_S \leq 0$ .

Our invariants count *self-dual Higgs complexes* on a surface, which are orthosymplectic complexes defined in §6.2, equipped with Higgs fields. They are a generalization of *G*-Higgs *bundles*, introduced by Hitchin [25], for G = O(n) or Sp(2n).

Via the spectral construction, these invariants can be seen as a version of orthosymplectic DT invariants in §6.4, for the non-compact Calabi–Yau threefold  $K_S$ , the total space of the canonical bundle of the surface *S*, with an involution that reverses the fibre direction.

**6.5.2. Higgs complexes.** Let *S* be a connected, smooth, projective algebraic surface over  $\mathbb{C}$ , and fix the data (*I*, *L*, *s*,  $\varepsilon$ ) as in §6.2.2 defining a self-dual structure  $\mathbb{D}$  on Perf(*S*).

For an object  $E \in Perf(S)$ , a *Higgs field* on *E* is a morphism

$$\psi\colon E\longrightarrow E\otimes K_S$$

in Perf(*S*). We call such a pair  $(E, \psi)$  a *Higgs complex* on *S*.

A self-dual Higgs complex is then defined as a fixed point of the involution

$$(E, \psi) \longmapsto (\mathbb{D}(E), -\mathbb{D}(\psi) \otimes K_S)$$

on the  $\infty$ -groupoid of Higgs complexes, where  $\mathbb{D}(\psi) \colon \mathbb{D}(E) \otimes K_S^{-1} \to \mathbb{D}(E)$ .

More concretely, for a self-dual object  $(E, \phi) \in \text{Perf}(S)^{\text{sd}}$  with  $\text{Ext}^i(E, E \otimes K_S) = 0$  for all i < 0, where  $\phi \colon E \xrightarrow{\sim} \mathbb{D}(E)$ , a self-dual Higgs field on  $(E, \phi)$  is the same data as a Higgs field  $\psi \colon E \to E \otimes K_S$  such that  $(\phi \otimes K_S) \circ \psi = -(\mathbb{D}(\psi) \otimes K_S) \circ \phi$  as morphisms  $E \to \mathbb{D}(E) \otimes K_S$ .

**6.5.3. Moduli stacks**. Let  $\hat{\mathcal{Y}}$  be the derived moduli stack of perfect complexes on *S*. Let  $\hat{\mathcal{X}} = T^*[-1]\hat{\mathcal{Y}}$  be the (-1)-shifted cotangent stack of  $\hat{\mathcal{Y}}$ , equipped with the canonical (-1)-shifted symplectic structure.

Then  $\bar{\mathcal{X}}$  is a moduli stack of Higgs complexes on *S*, since at a  $\mathbb{C}$ -point  $E \in \bar{\mathcal{Y}}(\mathbb{C})$ , we have

$$\mathbb{L}_{\tilde{\mathscr{Q}}}[-1]|_{E} \simeq \mathbb{R}\mathrm{Hom}_{S}(E, E)^{\vee}[-2] \simeq \mathbb{R}\mathrm{Hom}_{S}(E, E \otimes K_{S}),$$

parametrizing Higgs fields on E.

The self-dual structure on Perf(*S*) determines a  $\mathbb{Z}_2$ -action on  $\overline{\mathscr{Y}}$ , which induces a  $\mathbb{Z}_2$ -action on  $\overline{\mathscr{X}}$ . We have  $\overline{\mathscr{X}}^{sd} \simeq T^*[-1] \overline{\mathscr{Y}}^{sd}$ , giving  $\overline{\mathscr{X}}^{sd}$  a canonical (-1)-shifted symplectic structure.

We regard  $\bar{\mathcal{X}}^{sd}$  as a moduli stack of self-dual Higgs complexes on *S*. This description agrees with the definition of a self-dual Higgs field, as the (-1)-shifted tangent map of the involution  $\mathbb{D}$ , as a map  $\mathbb{R}\text{Hom}(E, E) \xrightarrow{\sim} \mathbb{R}\text{Hom}(\mathbb{D}(E), \mathbb{D}(E))$ , is given by  $\psi \mapsto -\mathbb{D}(\psi)$ .

**6.5.4.** Stability conditions. We now restrict to the case when the anti-canonical bundle  $K_S^{-1}$  of *S* is ample or trivial, so that *S* is either a del Pezzo surface, a K3 surface, or an abelian surface. We abbreviate this condition as  $K_S \leq 0$ .

In this case, for any  $\tau \in \operatorname{Stab}^{\circ}(S)$  and any  $E \in \operatorname{Perf}(S)$ , every Higgs field  $\psi : E \to E \otimes K_S$ respects the  $\tau$ -HN filtration of E, since choosing a non-zero map  $\xi : K_S \to \mathcal{O}_S$ , the composition  $\xi \circ \psi : E \to E \otimes K_S \to E$  must preserve the HN filtration. Therefore, heuristically, a Higgs complex  $(E, \psi)$  is  $\tau$ -semistable if and only if E is  $\tau$ -semistable. This justifies the following series of definitions: For each interval  $J \subset \mathbb{R}$  of length |J| < 1, let  $\mathscr{Y}(\tau; J) \subset \overline{\mathscr{Y}}$  be the open substack of objects in  $\mathscr{P}(J)$ , and let  $\mathscr{X}(\tau; J) = \mathrm{T}^*[-1] \mathscr{Y}(\tau; J) \subset \overline{\mathscr{X}}$  be the corresponding open substack. The stacks  $\mathscr{X}(\tau; J)$  and  $\mathscr{Y}(\tau; J)$  are derived linear stacks. When J = -J, they are also self-dual derived linear stacks, and we have  $\mathscr{X}(\tau; J)^{\mathrm{sd}} \simeq \mathrm{T}^*[-1] \mathscr{Y}(\tau; J)^{\mathrm{sd}}$ .

Moreover,  $\tau$  defines permissible stability conditions on  $\mathscr{X}(\tau; J)$  and  $\mathscr{Y}(\tau; J)$ , in the sense of §2.3.5. Here, the  $\Theta$ -stratification on  $\mathscr{X}(\tau; J)$  can be obtained by following the proof of Halpern-Leistner [24, Theorem 6.5.3], similarly to §6.2.6, where the conditions (S) and (B) follow from the respective properties of  $\mathscr{Y}(\tau; J)$ .

**6.5.5. Invariants.** Suppose  $K_S \leq 0$  and  $\tau \in \text{Stab}^{\circ,\text{sd}}(S)$ . For a class  $\alpha \in K(S)$  with  $Z(\alpha) \neq 0$  or  $\theta \in K^{\text{sd}}(S)$  of phase 0, define the *Vafa–Witten type invariants* 

$$\operatorname{vw}_{\alpha}(\tau) \in \mathbb{Q}, \qquad \operatorname{vw}_{\theta}^{\operatorname{sd}}(\tau) \in \mathbb{Q}$$

counting semistable Higgs complexes of class  $\alpha$  or semistable self-dual Higgs complexes of class  $\theta$ , as the DT invariants in §4.2 for the (-1)-shifted symplectic linear stack  $\mathcal{X}(\tau; t)$  with the trivial stability condition, where  $t \in \mathbb{R}$  is a phase of  $Z(\alpha)$  or t = 0 for  $\theta$ .

Moreover, since  $\mathscr{X}(\tau; t)$  and  $\mathscr{X}(\tau; 0)^{sd}$  are (-1)-shifted cotangent stacks, they come with canonical orientations, which define an orientation data on  $\mathscr{X}(\tau; t)$  and a self-dual orientation data on  $\mathscr{X}(\tau; 0)$ . We use them to define *motivic Vafa–Witten type invariants* 

$$\operatorname{vw}_{\alpha}^{\operatorname{mot}}(\tau)$$
,  $\operatorname{vw}_{\theta}^{\operatorname{mot},\operatorname{sd}}(\tau) \in \widehat{\mathbb{M}}^{\operatorname{mon}}(\mathbb{C};\mathbb{Q})$ .

**6.5.6. Wall-crossing**. We have the following theorem stating the wall-crossing formulae for our Vafa–Witten invariants, which can be proved analogously to Theorem 6.4.3.

**Theorem.** Let S be a surface with  $K_S \leq 0$ , and choose the data  $(I, L, s, \varepsilon)$  as in §6.5.2. Let  $\tau, \tilde{\tau} \in$  Stab<sup>°</sup>(S) be Bridgeland stability conditions.

- (i) If τ, τ̃ can be connected by a path of length < 1/4 in Stab°(S), then for any class α ∈ K(S) with Z(α) ≠ 0, the wall-crossing formulae (5.3.2.1) and (5.3.2.3) hold for the invariants vw<sub>α</sub>(-), vw<sub>α</sub><sup>mot</sup>(-) when changing between τ and τ̃.
- (ii) If  $\tau, \tilde{\tau} \in \operatorname{Stab}^{\circ, \operatorname{sd}}(S)$ , and they can be connected by a path of length < 1/4 in  $\operatorname{Stab}^{\circ, \operatorname{sd}}(S)$ , then for any class  $\theta \in K^{\operatorname{sd}}(S)$  with  $Z(\theta) \in \mathbb{R}_{>0}$ , the wall-crossing formulae (5.3.2.2) and (5.3.2.4) hold for the invariants  $\operatorname{vw}_{\theta}^{\operatorname{sd}}(-)$ ,  $\operatorname{vw}_{\theta}^{\operatorname{mot,sd}}(-)$  when changing between  $\tau$  and  $\tilde{\tau}$ .

Here, the precise formulations of the wall-crossing formulae are as in Theorem 6.4.3.

**6.5.7. The case of K3 surfaces**. Suppose that *S* is a K3 surface or an abelian surface. Then for any  $E, F \in Perf(S)$ , we have the numerical relations

$$\operatorname{rk}\operatorname{Ext}^{\bullet}_{S}(E,F) = \operatorname{rk}\operatorname{Ext}^{\bullet}_{S}(F,E), \qquad \operatorname{rk}\operatorname{Ext}^{\bullet}_{S}(E,\mathbb{D}(E))^{\mathbb{Z}_{2}} = \operatorname{rk}\operatorname{Ext}^{\bullet}_{S}(\mathbb{D}(E),E)^{\mathbb{Z}_{2}},$$

where 'rk' denotes the alternating sum of dimensions, and  $(-)^{\mathbb{Z}_2}$  denotes the fixed part of the involution  $\phi \mapsto \mathbb{D}(\phi)$ . These relations imply that  $\bar{\mathcal{X}}$  and  $\bar{\mathcal{X}}^{sd}$  are numerically symmetric in the sense of §5.3.3. By Corollary 5.3.4 and Theorem 6.5.6, the invariants  $vw_{\alpha}(-)$ ,  $vw_{\theta}^{sd}(-)$ ,  $vw_{\alpha}^{mot}(-)$ , and  $vw_{\theta}^{mot,sd}(-)$  are locally constant functions on Stab<sup>°</sup>(S) or Stab<sup>°,sd</sup>(S).

## References

- [1] Abramovich, D., and Polishchuk, A., 'Sheaves of t-structures and valuative criteria for stable complexes', *J. Reine Angew. Math.* 2006/590 (2006). doi: 10.1515/crelle.2006.005, arXiv: math/0309435.
- [2] Alper, J., Halpern-Leistner, D., and Heinloth, J., 'Existence of moduli spaces for algebraic stacks', *Invent. Math.* 234/3 (2023), 949–1038. doi: 10.1007/s00222-023-01214-4, arXiv: 1812.01128.
- [3] Arcara, D., and Bertram, A., 'Bridgeland-stable moduli spaces for K-trivial surfaces', with an appendix by M. Lieblich, *J. Eur. Math. Soc.* 15/1 (2012), 1–38. doi: 10.4171/jems/354, arXiv: 0708.2247.
- Bayer, A., Macrì, E., and Toda, Y., 'Bridgeland stability conditions on threefolds. I: Bogomo-lov-Gieseker type inequalities', *J. Algebr. Geom.* 23/1 (2014), 117–163. doi: 10.1090/s1056-3911 -2013-00617-7, arXiv: 1103.5010.
- [5] Behrend, K., 'Donaldson-Thomas type invariants via microlocal geometry', Ann. Math. 170/3 (2009), 1307–1338. doi: 10.4007/annals.2009.170.1307, arXiv: math/0507523.
- [6] Ben-Bassat, O., Brav, C., Bussi, V., and Joyce, D., 'A "Darboux theorem" for shifted symplectic structures on derived Artin stacks, with applications', *Geom. Topol.* 19/3 (2015), 1287–1359. doi: 10.2140/gt.2015.19.1287, arXiv: 1312.0090.
- Bondal, A. I., 'Representation of associative algebras and coherent sheaves', *Math. USSR Izv.* 34/1 (1990), 23–42. doi: 10.1070/im1990v034n01abeh000583.
- [8] Bridgeland, T., 'Stability conditions on triangulated categories', Ann. Math. 166/2 (2007), 317–345. doi: 10.4007/annals.2007.166.317, arXiv: math/0212237.
- [9] Bridgeland, T., 'Stability conditions on K3 surfaces', Duke Math. J. 141/2 (2008), 241-291. doi: 10.1215/s0012-7094-08-14122-5, arXiv: math/0307164.
- Bu, C., 'Enumerative invariants in self-dual categories. I: Motivic invariants', preprint v3 (22 Jan. 2024), arXiv: 2302.00038v3.
- Bu, C., 'Enumerative invariants in self-dual categories. II: Homological invariants', preprint v2 (11 Sept. 2023), arXiv: 2309.00056v2.
- [12] Bu, C., 'A motivic integral identity for (-1)-shifted symplectic stacks', *Moduli*, to appear, arXiv: 2405.10092.
- [13] Bu, C., Davison, B., Ibáñez Núñez, A., Kinjo, T., and Pădurariu, T., 'Cohomology of symmetric stacks', preprint v1 (6 Feb. 2025), arXiv: 2502.04253v1.
- [14] Bu, C., Halpern-Leistner, D., Ibáñez Núñez, A., and Kinjo, T., 'Intrinsic Donaldson–Thomas theory. I: Component lattices of stacks', preprint v1 (19 Feb. 2025), arXiv: 2502.13892v1.
- [15] Bu, C., Ibáñez Núñez, A., and Kinjo, T., 'Intrinsic Donaldson–Thomas theory. II: Stability measures and invariants', preprint v1 (27 Feb. 2025), arXiv: 2502.20515v1.

- [16] Bu, C., Ibáñez Núñez, A., and Kinjo, T., 'Intrinsic Donaldson–Thomas theory. III: Wall-crossing and applications', in preparation.
- [17] Bussi, V., Joyce, D., and Meinhardt, S., 'On motivic vanishing cycles of critical loci', *J. Algebr. Geom.* 28/3 (2019), 405–438. doi: 10.1090/jag/737, arXiv: 1305.6428.
- [18] DeHority, S., and Latyntsev, A., 'Orthosymplectic modules of cohomological Hall algebras', preprint v1 (11 Jan. 2025), arXiv: 2501.06643v1.
- [19] Derksen, H., and Weyman, J., 'Generalized quivers associated to reductive groups', *Colloq. Math.* 94/2 (2002), 151–173. doi: 10.4064/cm94-2-1.
- [20] Dhillon, A., and Young, M. B., 'The motive of the classifying stack of the orthogonal group', *Michigan Math. J.* 65/1 (2016). doi: 10.1307/mmj/1457101817, arXiv: 1411.2710.
- [21] Diaconescu, D.-E., Garcia-Raboso, A., Karp, R. L., and Sinha, K., 'D-brane superpotentials in Calabi–Yau orientifolds', *Adv. Theor. Math. Phys.* 11/3 (2007), 471–516. doi: 10.4310/atmp.2007.v11.n3.a5, arXiv: hep-th/0606180.
- [22] Donaldson, S. K., and Thomas, R. P., 'Gauge theory in higher dimensions', in S. A. Huggett et al. (eds.), *The geometric universe: Science, geometry, and the work of Roger Penrose* (Oxford University Press, 1998), 31–47. doi: 10.1093/0s0/9780198500599.003.0003.
- [23] Gómez, T. L., Fernandez Herrero, A., and Zamora, A., 'The moduli stack of principal ρ-sheaves and Gieseker–Harder–Narasimhan filtrations', *Math. Z.* 307/3 (2024), 51, 67 pp. doi: 10.1007/ s00209-024-03497-6, arXiv: 2107.03918.
- [24] Halpern-Leistner, D., 'On the structure of instability in moduli theory', preprint v5 (4 Feb. 2022), arXiv: 1411.0627v5.
- [25] Hitchin, N. J., 'The self-duality equations on a Riemann surface', *Proc. Lond. Math. Soc. (3)*, 55/1 (1987), 59–126. doi: 10.1112/plms/s3-55.1.59.
- [26] Hori, K., and Walcher, J., 'D-brane categories for orientifolds the Landau–Ginzburg case', *J. High Energy Phys.* 2008/4 (2008), 030, 36 pp. doi: 10.1088/1126-6708/2008/04/030, arXiv: hep-th/0606179.
- [27] Ibáñez Núñez, A., 'Refined Harder–Narasimhan filtrations in moduli theory', preprint v1 (29 Nov. 2023), arXiv: 2311.18050v1.
- [28] Joyce, D., 'Configurations in abelian categories. I: Basic properties and moduli stacks', *Adv. Math.* 203/1 (2006), 194–255. doi: 10.1016/j.aim.2005.04.008, arXiv: math/0312190.
- [29] Joyce, D., 'Configurations in abelian categories. II: Ringel–Hall algebras', Adv. Math. 210/2 (2007), 635–706. doi: 10.1016/j.aim.2006.07.006, arXiv: math/0503029.
- [30] Joyce, D., 'Configurations in abelian categories. III: Stability conditions and identities', Adv. Math. 215/1 (2007), 153-219. doi: 10.1016/j.aim.2007.04.002, arXiv: math/0410267.
- [31] Joyce, D., 'Motivic invariants of Artin stacks and "stack functions", Q. J. Math. 58/3 (2007), 345–392. doi: 10.1093/qmath/ham019, arXiv: math/0509722.
- [32] Joyce, D., 'Configurations in abelian categories. IV: Invariants and changing stability conditions', *Adv. Math.* 217/1 (2008), 125–204. doi: 10.1016/j.aim.2007.06.011, arXiv: math/0410268.
- [33] Joyce, D., 'Ringel-Hall style vertex algebra and Lie algebra structures on the homology of moduli spaces', preliminary version (Mar. 2018), https://people.maths.ox.ac.uk/joyce/hall.pdf.

- [34] Joyce, D., 'Enumerative invariants and wall-crossing formulae in abelian categories', preprint v1 (8 Nov. 2021), arXiv: 2111.04694v1.
- [35] Joyce, D., and Song, Y., A theory of generalized Donaldson-Thomas invariants (Mem. Am. Math. Soc. 1020; American Mathematical Society, 2012). doi: 10.1090/s0065-9266-2011-00630-1, arXiv: 0810.5645.
- [36] Joyce, D., and Upmeier, M., 'Orientation data for moduli spaces of coherent sheaves over Calabi-Yau 3-folds', Adv. Math. 381 (2021), 107627, 47 pp. doi: 10.1016/j.aim.2021.107627, arXiv: 2001.00113.
- [37] Kontsevich, M., and Soibelman, Y., 'Stability structures, motivic Donaldson–Thomas invariants and cluster transformations', preprint v1 (16 Nov. 2008), arXiv: 0811.2435v1.
- [38] Lê, Q. T., 'Proofs of the integral identity conjecture over algebraically closed fields', *Duke Math. J.* 164/1 (2015), 157–194. doi: 10.1215/00127094-2869138, arXiv: 1206.5334.
- [39] Maulik, D., and Thomas, R. P., 'Sheaf counting on local K3 surfaces', *Pure Appl. Math. Q.* 14/3 (2018), 419–441. doi: 10.4310/pamq.2018.v14.n3.a1, arXiv: 1806.02657.
- [40] Pantev, T., Toën, B., Vaquié, M., and Vezzosi, G., 'Shifted symplectic structures', *Publ. Math., Inst. Hautes Étud. Sci.* 117 (2013), 271–328. doi: 10.1007/s10240-013-0054-1, arXiv: 1111.3209.
- [41] Piyaratne, D., and Toda, Y., 'Moduli of Bridgeland semistable objects on 3-folds and Donald-son-Thomas invariants', *J. Reine Angew. Math.* 747 (2019), 175–219. doi: 10.1515/crelle-2016-0006, arXiv: 1504.01177.
- [42] Ringel, C. M., 'Hall algebras', in *Topics in algebra. 1: Rings and representations of algebras* (Banach Cent. Publ. 26, part 1; 1990), 433–447.
- [43] Ringel, C. M., 'Hall algebras and quantum groups', *Invent. Math.* 101/1 (1990), 583–591. doi: 10
   .1007/bf01231516.
- [44] Rudakov, A., 'Stability for an abelian category', J. Algebra, 197/1 (1997), 231–245. doi: 10.1006/ jabr.1997.7093.
- [45] Tanaka, Y., and Thomas, R. P., 'Vafa–Witten invariants for projective surfaces. I: Stable case', J. Algebr. Geom. 29/4 (2020), 603–668. doi: 10.1090/jag/738, arXiv: 1702.08487.
- [46] Tanaka, Y., and Thomas, R. P., 'Vafa-Witten invariants for projective surfaces. II: Semistable case', *Pure Appl. Math. Q.* 13/3 (2018), 517–562. doi: 10.4310/pamq.2017.v13.n3.a6, arXiv: 1702.08488.
- [47] Thomas, R. P., 'A holomorphic Casson invariant for Calabi–Yau 3-folds, and bundles on K3 fibrations', J. Differ. Geom. 54/2 (2000), 367–438. doi: 10.4310/jdg/1214341649, arXiv: math/9806111.
- [48] Thomas, R. P., 'Equivariant K-theory and refined Vafa–Witten invariants', Commun. Math. Phys. 378/2 (2020), 1451–1500. doi: 10.1007/s00220-020-03821-1, arXiv: 1810.00078.
- [49] Toda, Y., 'Moduli stacks and invariants of semistable objects on K3 surfaces', Adv. Math. 217/6 (2008), 2736-2781. doi: 10.1016/j.aim.2007.11.010, arXiv: math/0703590.
- [50] Toën, B., and Vaquié, M., 'Moduli of objects in dg-categories', Ann. Sci. Éc. Norm. Supér. 40/3 (2007), 387-444. doi: 10.1016/j.ansens.2007.05.001, arXiv: math/0503269.
- [51] Toën, B., and Vezzosi, G., Homotopical algebraic geometry II: Geometric stacks and applications (Mem. Am. Math. Soc. 902; American Mathematical Society, 2008). doi: 10.1090/memo/0902, arXiv: math/0404373.

- [52] Witten, E., 'D-branes and K-theory', J. High Energy Phys. 1998/12 (1998), 019. doi: 10.1088/ 1126-6708/1998/12/019, arXiv: hep-th/9810188.
- [53] Young, M. B., 'The Hall module of an exact category with duality', *J. Algebra*, 446 (2016), 291–322. doi: 10.1016/j.jalgebra.2015.09.013, arXiv: 1212.0531.
- [54] Young, M. B., 'Self-dual quiver moduli and orientifold Donaldson–Thomas invariants', *Commun. Number Theory Phys.* 9/3 (2015), 437–475. doi: 10.4310/cntp.2015.v9.n3.a1, arXiv: 1408.4888.
- [55] Young, M. B., 'Representations of cohomological Hall algebras and Donaldson-Thomas theory with classical structure groups', *Commun. Math. Phys.* 380/1 (2020), 273–322. doi: 10.1007/ s00220-020-03877-z, arXiv: 1603.05401.

Chenjing Bu bu@maths.ox.ac.uk

Mathematical Institute, University of Oxford, Oxford OX2 6GG, United Kingdom.