

Decoherence Time Maximization and Partial Isolation for Open Quantum Harmonic Oscillator Memory Networks*

Igor G. Vladimirov¹, Ian R. Petersen², Guodong Shi^{2,3}

Abstract

This paper considers a network of open quantum harmonic oscillators which interact with their neighbours through direct energy and field-mediated couplings and also with external quantum fields. The position-momentum dynamic variables of the network are governed by linear quantum stochastic differential equations associated with the nodes of a graph whose edges specify the interconnection of the component oscillators. Such systems can be employed as Heisenberg picture quantum memories with an engineered ability to approximately retain initial conditions over a bounded time interval. We use the quantum memory decoherence time defined previously in terms of a fidelity threshold on a weighted mean-square deviation for a subset (or linear combinations) of network variables from their initial values. This approach is applied to maximizing a high-fidelity asymptotic approximation of the decoherence time over the direct energy coupling parameters of the network. The resulting optimality condition is a set of linear equations for blocks of a sparse matrix associated with the edges of the direct energy coupling graph of the network. We also discuss a setting where the quantum network has a subset of dynamic variables which are affected by the external fields only indirectly, through a complementary “shielding” system. This holds under a rank condition on the network-field coupling matrix and can be achieved through an appropriate field-mediated coupling between the component oscillators. The partially isolated subnetwork has a longer decoherence time in the high-fidelity limit, thus providing a particularly relevant candidate for a quantum memory.

I. INTRODUCTION

Due to the operator-valued nature of quantum mechanics, which describes the dynamics and statistical properties of the physical world on atomic and subatomic scales [12], [20], the quantum variables are of the power of the continuum even in the finite-level case. For example, the qubit, which is the simplest quantum mechanical system, involves the Hilbert space \mathbb{C}^2 as opposed to the two-point state space of classical binary systems (such as flip-flop circuits in digital computer electronics) or Bernoulli random variables. In line with the qualitatively different noncommutative structure of quantum variables and quantum states, their dynamics and quantum probabilistic properties [7] follow the nonclassical Heisenberg or Schrödinger pictures, including the postulate of unitary evolution for isolated systems completely specified by the Hamiltonian.

The fragility of quantum dynamics with respect to external fields and measurements complicates the manipulation of quantum systems needed for the initialization and preservation of particular quantum states or dynamic variables over the course of time. The latter tasks correspond to the stages of “writing” and storing the quantum information (followed by the “reading” phase) and can be implemented using photonics and light-matter interaction ([2], [6], [30], [31], [32] and references therein) or other platforms. These operations involve a controlled isolation of the quantum system from its environment, similarly to the existing paradigms of quantum computation [14], where the closed unitary evolution alternates with measurements.

The reversibility of unitary dynamics (especially in the zero Hamiltonian case where any quantum state and all the system variables are preserved in time) is particularly useful for quantum information storage. However, in contrast to this idealised scenario, the ability to retain quantum states or dynamic variables is corrupted by the unavoidable interaction of the quantum system with its environment. The system-field interaction gives rise to quantum noise [3] which forces the system to drift away from its initial condition in a dissipative (and hence, irreversible) fashion. The resulting open quantum dynamics are modelled by the Hudson-Parthasarathy quantum stochastic differential equations (QSDEs) [10], [16] driven by quantum Wiener processes. They are linear for open quantum harmonic oscillators (OQHOs) [11], [15], [18], [33] and quasi-linear for finite-level open quantum systems [1], [24]. Although the finite-level systems are particularly suitable for modelling multi-qubit registers, continuous variables systems (including the OQHOs with their position-momentum variables) are also used in the Gaussian channel models of quantum information theory [8].

For both classes of open quantum systems in the quantum stochastic calculus framework, their performance as a Heisenberg picture quantum memory in the storage phase has recently been described in terms of a decoherence time [26], [27]. The latter is defined as the time horizon at which a weighted mean-square deviation of the system variables from their initial values reaches a given fidelity threshold (see [25] and references therein for different yet related decoherence measures). The maximization of the memory decoherence time (or its asymptotic approximation in the high-fidelity or short-horizon limit) over the energy and coupling parameters of the quantum system improves its ability to approximately retain the initial conditions over a bounded time interval. The approximate decoherence time maximization for OQHOs has also been

*This work is supported by the Australian Research Council grant DP240101494.

^{1,2}School of Engineering, Australian National University, Canberra, ACT, Australia: igor.g.vladimirov@gmail.com, i.r.petersen@gmail.com.

³Australian Centre for Robotics, University of Sydney, Camperdown, Sydney, NSW, Australia: guodong.shi@sydney.edu.au.

considered in [28] for a partially isolated subsystem which is affected by the external fields only indirectly through another subsystem of the oscillator, thus leading to a qualitatively longer memory decoherence time. However, these optimization problems were considered in [26], [28] with respect to a single direct energy coupling matrix for a feedback interconnection of two OQHOs with direct energy and field-mediated couplings [34]. At the same time, the development of systematic methods for solving such problems for multicomponent quantum networks with arbitrary interconnection architecture is of theoretical and practical interest.

In the present paper, we extend the approach of [26], [28] to a quantum memory system organised as a network of OQHOs which interact with their neighbours through direct energy and field-mediated couplings and also with external quantum fields. The position-momentum dynamic variables of the network are governed by a set of cross-coupled linear QSDEs associated with the nodes of a graph whose edges specify the interconnection of the constituent oscillators. This allows the network to be represented as an augmented OQHO by making advantage of the linearity of the QSDEs (instead of the quantum feedback network formalism [5] in its full generality). We use the quantum memory decoherence time defined previously in terms of a fidelity threshold on a weighted mean-square deviation for a subset (or linear combinations) of network variables from their initial values. This approach is applied to maximizing the high-fidelity asymptotic approximation of the decoherence time over the direct energy coupling matrices of the network. The resulting necessary and sufficient condition of optimality is a set of linear algebraic equations (resembling the Sylvester equations [4], [22]) for blocks of a sparse matrix associated with the edges of the direct energy coupling graph of the network. We also discuss a setting where the quantum network has a subset of dynamic variables which are affected by the external fields only indirectly, through a complementary “shielding” system. This partial subnetwork isolation from the external fields and the related network decomposition hold under a certain rank condition on the network-field coupling matrix and can be achieved through an appropriate field-mediated coupling between the component oscillators. The partially isolated subnetwork acquires a longer decoherence time in the high-fidelity limit, thus providing a particularly relevant candidate for a quantum memory.

The paper is organised as follows. Section II describes the internal dynamic and external field variables of the network of OQHOs being considered. Section III specifies the direct energy and field-mediated couplings between the component OQHOs (including the neighbourhoods and Hamiltonians) and their coupling to the external fields. Section IV provides the set of QSDEs for the Heisenberg dynamics of the quantum network and computes the energy and system-field coupling matrices for the augmented OQHO which represents it. Section V quantifies the performance of the network as a temporary memory system in terms of weighted deviations of its dynamic variables from their initial conditions and the memory decoherence time. Section VI formulates and solves the problem of maximizing the approximate decoherence time in the high-fidelity limit over the direct energy coupling parameters of the network. Section VII specifies the above results (including the approximate decoherence time maximization) in application to the partial subnetwork isolation setting. Section VIII provides concluding remarks.

II. OPEN QUANTUM NETWORK AND EXTERNAL FIELDS

We consider a network of linear quantum stochastic systems at the vertex set V of a finite graph specified below. For any $j \in V$, the j th component system is a multi-mode open quantum harmonic oscillator (OQHO) with an even number n_j of internal dynamic variables which are time-varying self-adjoint operators of position-momentum type on (a dense domain of) a Hilbert space \mathfrak{H} . These n_j quantum variables are assembled into a column-vector $X_j(t)$ (the time argument $t \geq 0$ will often be omitted for brevity) and act initially (at $t = 0$) on a Hilbert space \mathfrak{H}_j . It is assumed that they satisfy the canonical commutation relations (CCRs)

$$[X_j(t), X_j(t)^T] = 2i\Theta_j, \quad [X_j(t), X_k(t)^T] = 0 \quad (1)$$

for all $t \geq 0$ and $j, k \in V$ such that $j \neq k$, where the transpose $(\cdot)^T$ applies to matrices and vectors of operators as if the latter were scalars, $i := \sqrt{-1}$ is the imaginary unit, and

$$\Theta_j := \frac{1}{2}I_{n_j/2} \otimes \mathbf{J} \quad (2)$$

is a nonsingular real antisymmetric matrix of order n_j . Here, $[\alpha, \beta^T] := ([\alpha_a, \beta_b])_{1 \leq a \leq r, 1 \leq b \leq s} = \alpha\beta^T - (\beta\alpha^T)^T$ is the matrix of commutators $[\alpha_a, \beta_b] = \alpha_a\beta_b - \beta_b\alpha_a$ for vectors $\alpha := (\alpha_a)_{1 \leq a \leq r}$, $\beta := (\beta_b)_{1 \leq b \leq s}$ of linear operators. Also, I_r is the identity matrix of order r , and \otimes is the tensor product of spaces or operators, including the Kronecker product of matrices as in (2), where

$$\mathbf{J} := \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \quad (3)$$

spans the subspace of antisymmetric matrices of order 2. In view of (1), (2), an augmented n -dimensional vector X of network variables, given by

$$X := (X_j)_{j \in V}, \quad n := \sum_{j \in V} n_j, \quad (4)$$

has a block-diagonal CCR matrix:

$$[X, X^T] = 2i\Theta, \quad \Theta := \text{diag}(\Theta_j) = \frac{1}{2}I_{n/2} \otimes \mathbf{J}. \quad (5)$$

The relation (5) holds, for example, if the network variables are implemented as the pairs $q_1, p_1, \dots, q_{n/2}, p_{n/2}$ of quantum mechanical positions q_r and momenta $p_r := -i\partial_{q_r}$, with $r = 1, \dots, n/2$, on the Schwartz space [29] of rapidly decreasing functions on $\mathbb{R}^{n/2}$.

In addition to the internal variables, the j th OQHO has multichannel input and output bosonic fields W_j, Y_{jk} (see Fig. 1 for an informal illustration), which consist of m_j and r_{jk} time-varying self-adjoint quantum variables, respectively (the




Fig. 1. An illustration of a network of OQHOs (depicted as numbered circles) with external quantum Wiener processes W_j and output fields Y_{jk} which mediate the coupling between the component OQHOs. This field-mediated coupling is represented by arrows, while dashed lines indicate the direct energy coupling.

dimensions m_j, r_{jk} are also even). The input field W_j is a quantum Wiener process on a symmetric Fock space \mathfrak{F}_j . The network-field space has the tensor-product structure

$$\mathfrak{H} := \mathfrak{H}^0 \otimes \mathfrak{F}, \quad \mathfrak{H}^0 := \otimes_{j \in V} \mathfrak{H}_j, \quad \mathfrak{F} := \otimes_{j \in V} \mathfrak{F}_j, \quad (6)$$

where the composite Fock space \mathfrak{F} accommodates the external fields, and the initial network space \mathfrak{H}^0 provides a domain for the initial network variables. Accordingly, the internal network variables and the output fields act on the network-field space \mathfrak{H} . Now, the input fields satisfy the two-point CCRs

$$[W_j(s), W_j(t)^T] = 2i \min(s, t) J_j, \quad [W_j(s), W_k(t)^T] = 0 \quad (7)$$

for all $s, t \geq 0$ and $j, k \in V$ such that $j \neq k$, where

$$J_j := I_{m_j/2} \otimes \mathbf{J} \quad (8)$$

is an orthogonal real antisymmetric matrix of order m defined in terms of (3), so that $J_j^2 = -I_{m_j}$. The right-hand side of (7) vanishes at $s = 0$ or $t = 0$ since the initial input field operators act as the identity operator on \mathfrak{F} , which commutes with any operator. Due to the continuous tensor-product structure [17] of the Fock space filtration, the relation (7) is equivalent to its fulfillment for all $s = t \geq 0$, whose incremental form is given by

$$d[W_j, W_k^T] = [dW_j, W_k^T] + [W_j, dW_k^T] + [dW_j, dW_k^T] = [dW_j, dW_k^T] = \begin{cases} 2iJ_j dt & \text{if } j = k \\ 0 & \text{if } j \neq k \end{cases}. \quad (9)$$

Here, the quantum Ito lemma [10] is used along with the property of the future-pointing Ito increments of the input quantum Wiener processes to commute with adapted processes (in the sense of the network-field space filtration). In particular,

$$[dW_k(t), [W_j(s)^T \ X_j(s)^T \ Y_j(s)^T]] = 0 \quad (10)$$

for all $t \geq s \geq 0$ and $j, k \in V$. We assume that the input fields are in the tensor-product vacuum state [16] (in particular, they are statistically independent for different component OQHOs)

$$\mathbf{v} := \otimes_{j \in V} \mathbf{v}_j \quad (11)$$

on the composite Fock space \mathfrak{F} , where \mathbf{v}_j is the vacuum state on the Fock space \mathfrak{F}_j . Accordingly, the CCRs (9) correspond to the imaginary parts of the quantum Ito relations

$$dW_j dW_k^T = \begin{cases} \Omega_j dt & \text{if } j = k \\ 0 & \text{if } j \neq k \end{cases}, \quad \Omega_j := I_{m_j} + iJ_j. \quad (12)$$

In what follows, it will be convenient to use (similarly to (4)) an augmented m -dimensional vector W of external fields given by

$$W := (W_j)_{j \in V}, \quad m := \sum_{j \in V} m_j. \quad (13)$$

From (8), (12), the quantum Ito relations for W take the form

$$dWdW^T = \Omega dt, \quad \Omega = I_m + iJ, \quad (14)$$

where

$$J := \text{diag}(J_j) = I_{m/2} \otimes \mathbf{J}. \quad (15)$$

The above described commutation and statistical structure of the external fields is essential for the dynamical and quantum probabilistic properties of the network. The latter also depend on the network-field interaction and the coupling between the component OQHOs considered in the next section.

III. DIRECT ENERGY AND FIELD-MEDIATED COUPLINGS

With every node $j \in V$ of the quantum network, we associate three subsets of the vertex set V of the graph which do not contain j :

$$N_j^0, N_j^+, N_j^- \subset V \setminus \{j\}, \quad j \in V. \quad (16)$$

Each of these subsets specifies a neighbourhood with which the j th OQHO interacts through a direct energy or field-mediated coupling.

More precisely, N_j^0 pertains to the direct energy coupling (for example, as between the OQHOs 1 and 2 and also between 2 and 3 in Fig. 1) and satisfies a symmetry property: for all $j, k \in V$ such that $j \neq k$, the inclusion $k \in N_j^0$ holds if and only if so does $j \in N_k^0$. The set N_j^0 is therefore a neighbourhood of the node j in an undirected graph (whose edges are represented by dashed lines in Fig. 1).

The set N_j^+ in (16) specifies a neighbourhood whose component OQHOs $k \in N_j^+$ “receive” the output fields Y_{jk} from the j th OQHO. In a similar fashion, the set N_j^- describes another neighbourhood whose component OQHOs $k \in N_j^-$ “send” their output fields Y_{kj} to the j th OQHO. The sets N_j^\pm are specified by the outgoing and incoming edges of the node j in a directed graph, respectively. Accordingly, for all $j, k \in V$ such that $j \neq k$, the inclusions $k \in N_j^+$ and $j \in N_k^-$ are equivalent; see Fig. 2. The output fields Y_{jk} form an r_j -dimensional “total” output field Y_j of the j th OQHO as




Fig. 2. Two component OQHOs j and k (with $j, k \in V$ such that $j \neq k$) belong to the neighbourhoods N_k^- and N_j^+ (shown as dashed boxes), respectively, if and only if they are coupled through the output field Y_{jk} from j to k .

$$Y_j := (Y_{jk})_{k \in N_j^+}, \quad r_j := \sum_{k \in N_j^+} r_{jk} \leq m_j \quad (17)$$

(the inequality is clarified below).

Ignoring the field-mediated coupling (to be considered later), the direct energy coupling is described in terms of the neighbourhoods N_j^0 and the following Hamiltonian of the OQHO network:

$$H^0 := \frac{1}{2} X^T R^0 X = \sum_{j \in V} (H_j + \sum_{k \in N_j^0} H_{jk}), \quad (18)$$

where $R^0 := (R_{jk}^0)_{j, k \in V} \in \mathbb{S}_n$ is the energy matrix of the network (where \mathbb{S}_n denotes the subspace of real symmetric matrices of order n), with $R_{jk}^0 := 0$ if $k \in V \setminus (\{j\} \cup N_j^0)$, and use is also made of (4). In (18),

$$H_j := \frac{1}{2} X_j^T R_j X_j \quad (19)$$

is the individual Hamiltonian of the j th OQHO specified by the corresponding diagonal block $R_j := R_{jj}^0 \in \mathbb{S}_{n_j}$ of R^0 , while the matrix $R_{jk}^0 = R_{kj}^{0T} \in \mathbb{R}^{n_j \times n_k}$ parameterises the direct energy coupling Hamiltonian for the j th and k th OQHOs:

$$H_{jk} := \frac{1}{2} X_j^T R_{jk}^0 X_k = H_{kj}, \quad j \in V, \quad k \in N_j^0, \quad (20)$$

where the equality uses the commutativity between X_j and X_k from (1) (for the same reason, H_{jk} are also self-adjoint operators). Therefore, the direct energy coupling between the OQHOs $j \in V$ and $k \in N_j^0$ (indicated by an edge of the undirected graph mentioned above), contributes $\frac{1}{2}(H_{jk} + H_{kj}) = H_{jk}$ to the total energy of the system.

In addition to the direct energy coupling between the component OQHOs through the symmetric interaction Hamiltonian terms (20), the coupling of the j th OQHO to the external field W_j is quantified by a matrix $M_j \in \mathbb{R}^{m_j \times n_j}$. Similarly, the coupling of the j th OQHO to the output field Y_{kj} (such as quantized electromagnetic radiation in the case of quantum optical [30] implementation of the network) from the k th OQHO, with $k \in N_j^-$, is specified by a matrix $N_{jk} \in \mathbb{R}^{r_{kj} \times n_j}$. In comparison with the direct energy coupling, the effect of the field-mediated coupling on the energetics of the network is less straightforward and will be recovered from the network dynamics discussed in the next section.

IV. HEISENBERG NETWORK DYNAMICS

In accordance with the direct energy and field-mediated coupling, the Heisenberg evolution of the network is governed by a set of linear quantum stochastic differential equations (QSDEs)

$$dX_j = \left(A_j X_j + \overbrace{\sum_{k \in N_j^0} A_{jk} X_k}^{\text{direct energy coupling}} \right) dt + B_j dW_j + \overbrace{\sum_{k \in N_j^-} E_{jk} dY_{kj}}^{\text{field-mediated coupling}}, \quad (21)$$

$$dY_j = C_j X_j dt + D_j dW_j, \quad j \in V. \quad (22)$$

Here, the matrices

$$A_j \in \mathbb{R}^{n_j \times n_j}, \quad A_{jk} \in \mathbb{R}^{n_j \times n_k}, \quad B_j \in \mathbb{R}^{n_j \times m_j}, \quad C_j \in \mathbb{R}^{r_j \times n_j}, \quad D_j \in \mathbb{R}^{r_j \times m_j}, \quad E_{jk} \in \mathbb{R}^{n_j \times r_{kj}}$$

are expressed in terms of the CCR matrices Θ_j from (1), (2), the individual energy matrices R_j of the component OQHOs from (19), the direct energy coupling matrices R_{jk}^0 from (20), the matrices M_j of coupling of the OQHOs to their external input fields and the field-mediated coupling matrices N_{jk} (see the last paragraph of the previous section) as

$$A_j = 2\Theta_j(R_j + M_j^T J_j M_j + \sum_{\ell \in N_j^-} N_{j\ell}^T \tilde{J}_{\ell j} N_{j\ell}), \quad (23)$$

$$A_{jk} = 2\Theta_j R_{jk}^0, \quad B_j = 2\Theta_j M_j^T, \quad (24)$$

$$C_j = 2D_j J_j M_j, \quad E_{jk} = 2\Theta_j N_{jk}^T, \quad (25)$$

where

$$\tilde{J}_{\ell j} := D_{\ell j} J_{\ell} D_{\ell j}^T. \quad (26)$$

This parameterization (including the quadratic functions of the coupling matrices in (23)) reflects the physical realizability (PR) conditions [11], [21] for the coefficients of linear QSDEs. Furthermore, in accordance with (17), (22), the matrices

$$C_j := (C_{jk})_{k \in N_j^+}, \quad D_j := (D_{jk})_{k \in N_j^+} \quad (27)$$

consist of blocks $C_{jk} \in \mathbb{R}^{r_{jk} \times n_j}$ and $D_{jk} \in \mathbb{R}^{r_{jk} \times m_j}$ with even numbers of rows r_{jk} , which, similarly to the first equality in (25), are related by

$$C_{jk} = 2D_{jk} J_j M_j. \quad (28)$$

The blocks D_{jk} of the matrix D_j describe the selection of the output field channels from the j th OQHO and their ‘‘distribution’’ over the receiving OQHOs $k \in N_j^+$. Accordingly, the matrix D_j is formed from r_j conjugate rows of a permutation matrix of order m_j (which clarifies the inequality in (17)), so that, without loss of generality, the quantum Ito matrix of the output Y_j of the j th OQHO in the relations

$$dY_j dY_k^T = \begin{cases} D_j \Omega_j D_j^T dt & \text{if } j = k \\ 0 & \text{if } j \neq k \end{cases}, \quad j, k \in V \quad (29)$$

has a form similar to that in (12):

$$D_j \Omega_j D_j^T = I_{r_j} + iI_{r_j/2} \otimes \mathbf{J}. \quad (30)$$

In particular, it follows from (12), (26), (27), (30) that for all $j, k \in V$,

$$D_{jk} J_j D_{j\ell}^T = \begin{cases} \tilde{J}_{jk} (= I_{r_{jk}/2} \otimes \mathbf{J}) & \text{if } k = \ell \in N_j^+ \\ 0 & \text{otherwise} \end{cases}. \quad (31)$$

The block-diagonal structure of the matrix $D_j J_j D_j^T$ in (30), (31) means the commutativity between the Ito increments (at the same moment of time) of the output fields from the j th OQHO towards different component OQHOs in the network:

$$[dY_{jk}, dY_{j\ell}^T] = 0, \quad k, \ell \in N_j^+, \quad k \neq \ell. \quad (32)$$

In accordance with the partitioning in (17), (27), the output Y_{jk} of the j th OQHO towards the k th OQHO satisfies the QSDE

$$dY_{jk} = C_{jk} X_j dt + D_{jk} dW_j, \quad j \in V, \quad k \in N_j^+. \quad (33)$$

As the k th diagonal block of (29), the quantum Ito relations

$$dY_{jk} dY_{jk}^T = D_{jk} \Omega_j D_{jk}^T dt$$

clarify the meaning of the matrix \tilde{J}_{jk} (cf. (23), (26), (31)) as the CCR matrix for the output field Y_{jk} in the sense that $[dY_{jk}, dY_{jk}^T] = 2i\tilde{J}_{jk} dt$, which complements the commutativity (32).

The following theorem represents the dynamics (21) of the quantum network as that of a composite OQHO with the augmented vectors X, W of network variables and input fields from (4), (13).

Theorem 1: The augmented vector X of the network variables (4) satisfies a linear QSDE

$$dX = AX dt + B dW \quad (34)$$

driven by the augmented external field vector W in (13), where the matrices $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{n \times m}$ are given by

$$A := 2\Theta(R + M^T J M), \quad B := 2\Theta M^T. \quad (35)$$

Here, Θ, J are the augmented CCR matrices of the network and external field variables from (5), (14), (15). Furthermore, $M := (M_{jk})_{j,k \in V} \in \mathbb{R}^{m \times n}$ is the network-field coupling matrix whose blocks $M_{jk} \in \mathbb{R}^{m_j \times n_k}$ are computed as

$$M_{jk} = \begin{cases} M_j & \text{if } j = k \\ D_{jk}^T N_{kj} & \text{if } k \in N_j^+ \\ 0 & \text{if } k \notin \{j\} \cup N_j^+ \end{cases}. \quad (36)$$

Also, $R := (R_{jk})_{j,k \in V} \in \mathbb{S}_n$ is the network energy matrix with blocks $R_{jk} \in \mathbb{R}^{n_j \times n_k}$ given by

$$R_{jj} = R_j, \quad (37)$$

$$R_{jk} = \chi_{N_j^0}(k) R_{jk}^0 + \chi_{N_j^-}(k) M_{kj}^T J_k M_k - \chi_{N_j^+}(k) M_j^T J_j M_{jk} \quad (38)$$

for all $j, k \in V$ such that $j \neq k$, where χ_S is the indicator function of a set S (that is, $\chi_S(z) = 1$ if $z \in S$ and $\chi_S(z) = 0$ if $z \notin S$). \square

Proof: Substitution of (23)–(28), (33) into (21) leads to

$$\begin{aligned} dX_j &= 2\Theta_j \left(\left((R_j + M_j^T J_j M_j + \sum_{\ell \in N_j^-} N_{j\ell}^T \tilde{J}_{\ell j} N_{j\ell}) X_j + \sum_{k \in N_j^0} R_{jk}^0 X_k \right) dt + M_j^T dW_j + \sum_{k \in N_j^-} N_{jk}^T D_{kj} (2J_k M_k X_k dt + dW_k) \right), \\ &= 2\Theta_j \left(\left(\sum_{k \in V} R_{jk}^0 X_k + \overbrace{(M_j^T J_j M_j + \sum_{\ell \in N_j^-} N_{j\ell}^T \tilde{J}_{\ell j} N_{j\ell})}^{\gamma_{jj}} \right) X_j + \sum_{k \in N_j^-} \overbrace{2N_{jk}^T D_{kj} J_k M_k X_k}^{\gamma_{jk}} \right) dt + \sum_{k \in V} M_{kj}^T dW_k \\ &= 2\Theta_j \sum_{k \in V} ((R_{jk}^0 + \gamma_{jk}) X_k dt + M_{kj}^T dW_k), \quad j \in V, \end{aligned} \quad (39)$$

where use is also made of (26), (36) and the blocks R_{jk}^0 of the energy matrix R^0 from (18)–(20) which does not take into account the field-mediated coupling between the component OQHOs. Here, $\Gamma := (\gamma_{jk})_{j,k \in V} \in \mathbb{R}^{n \times n}$ is an auxiliary matrix whose blocks $\gamma_{jk} \in \mathbb{R}^{n_j \times n_k}$ are defined on the right-hand side of (39):

$$\gamma_{jk} := \begin{cases} \sum_{\ell \in \{j\} \cup N_j^-} M_{\ell j}^T J_\ell M_{\ell j} & \text{if } j = k \\ 2M_{kj}^T J_k M_k & \text{if } k \in N_j^- \\ 0 & \text{if } k \notin \{j\} \cup N_j^- \end{cases} \quad (40)$$

for all $j, k \in V$. This allows the QSDEs (39) to be assembled into one QSDE

$$dX = 2\Theta(R^0 + \Gamma)X dt + \underbrace{2\Theta M^T}_B dW, \quad (41)$$

where the second notation from (35) is also used. Now note that

$$\mathbf{A}(\Gamma) = M^T J M, \quad (42)$$

where \mathbf{A} is the antisymmetrizer of square matrices: $\mathbf{A}(\Gamma) := \frac{1}{2}(\Gamma - \Gamma^T)$. Indeed,

$$(\mathbf{A}(\Gamma))_{jj} = \mathbf{A}(\gamma_{jj}) = \gamma_{jj} = (M^T J M)_{jj}, \quad j \in V \quad (43)$$

in view of the antisymmetry of the diagonal blocks γ_{jj} of Γ in (40) inherited from the matrices J_j in (8). Furthermore,

$$(\mathbf{A}(\Gamma))_{jk} = \frac{1}{2}(\gamma_{jk} - \gamma_{kj}^T) = \chi_{N_j^-}(k) M_{kj}^T J_k M_k + \chi_{N_j^+}(k) M_j^T J_j M_{jk} \quad (44)$$

for any $j, k \in V$ such that $j \neq k$. On the other hand, (36) yields

$$(M^T J M)_{jk} = \sum_{\ell \in V} M_{\ell j}^T J_\ell M_{\ell k} = \chi_{N_j^+}(k) M_j^T J_j M_{jk} + \chi_{N_j^-}(k) M_{kj}^T J_k M_k + \sum_{\ell \in (N_j^- \cap N_k^-) \setminus \{j, k\}} \underbrace{N_{j\ell}^T D_{\ell j} J_\ell D_{\ell k}^T N_{k\ell}}_0 \quad (45)$$

for all $j, k \in V$ such that $j \neq k$, with (31) making the rightmost sum in (45) vanish. A comparison of (44) with (45) leads to

$$(\mathbf{A}(\Gamma))_{jk} = (M^T J M)_{jk}, \quad j, k \in V, \quad j \neq k,$$

which, in combination with (43), establishes (42). Due to (42), there exists a unique matrix $R = R^T \in \mathbb{R}^{n \times n}$ satisfying

$$R^0 + \Gamma = R + M^T J M. \quad (46)$$

Indeed, since the energy matrix R^0 in (18) is symmetric while $M^T J M$ is antisymmetric, then, in view of (42), such a matrix R is found by taking the symmetric part on both sides of (46):

$$R = R^0 + \mathbf{S}(\Gamma), \quad (47)$$

where \mathbf{S} is the symmetrizer of square matrices: $\mathbf{S}(\Gamma) := \frac{1}{2}(\Gamma + \Gamma^T)$, with the equality of the antisymmetric parts of (46) being secured by (42). In view of (46), the QSDE (41) takes the form (34) with the matrices (35). It now remains to compute the blocks of the matrix R in (47). By the antisymmetry of γ_{jj} in (43), the diagonal blocks of R coincide with those of R^0 :

$$R_{jj} = R_j + \mathbf{S}(\gamma_{jj}) = R_j, \quad j \in V,$$

thus establishing (37). The off-diagonal blocks of R are found by using those of $\mathbf{S}(\Gamma)$:

$$(\mathbf{S}(\Gamma))_{jk} = \frac{1}{2}(\gamma_{jk} + \gamma_{kj}^T) = \chi_{N_j^-}(k) M_{kj}^T J_k M_k - \chi_{N_j^+}(k) M_j^T J_j M_{jk} \quad (48)$$

for all $j, k \in V$ such that $j \neq k$, where (40) is used similarly to (44). A combination of (47) with (48) establishes (37), (38), thus completing the proof. \blacksquare

The above proof (which uses the linearity of the component QSDEs instead of the general quantum feedback network formalism [5]) clarifies the role of the quadratic terms in (23) for the property (42) (including (43)). The latter is crucial for the existence and uniqueness of the symmetric energy matrix R in the representation (35) leading to the PR property

$$A\Theta + \Theta A^T + BJB^T = 0,$$

which corresponds to the preservation of the CCRs (5) for the internal network variables over the course of time [11], [21] and also substantially uses the commutativity (10).

Also note that the individual energy matrices R_j of the component OQHOs enter the network energy matrix R only through its diagonal blocks (37). Furthermore, in addition to the direct energy coupling between the OQHOs, the network-field coupling and the field-mediated coupling contribute to R only through its off-diagonal blocks (38). The resulting network Hamiltonian H is related to H^0 in (18) as

$$H = \frac{1}{2} X^T R X = H^0 + \frac{1}{2} \sum_{j, k \in V} X_j^T (\chi_{N_j^-}(k) M_{kj}^T J_k M_k - \chi_{N_j^+}(k) M_j^T J_j M_{jk}) X_k = H^0 - \sum_{j \in V} X_j^T M_j^T J_j \sum_{k \in N_j^+} D_{jk}^T N_{kj} X_k,$$

where use is also made of the blocks (36) of the network-field coupling matrix M which does not depend on the matrix R^0 .

The QSDE (34), coming from the internal direct energy and field-mediated coupling in the quantum network and its interaction with the external fields, makes the network variables evolve in time $t \geq 0$ as

$$X(t) = e^{tA} X(0) + Z(t), \quad Z(t) := \int_0^t e^{(t-s)A} B dW(s), \quad (49)$$

with the matrices A, B from (35). The fact that e^{tA} is a usual matrix exponential makes the quantum trajectory X in (49) (despite the operator-valued nature of its component processes) similar to the solution of classical linear SDEs. This resemblance is due to the linearity of the QSDE (34).

V. QUANTUM NETWORK AS TEMPORARY MEMORY SYSTEM

The linear network response $e^{tA}X(0)$ to the initial network variables in (49) can be regarded as a “useful signal” carrying information about $X(0)$. It is particularly so on time scales $t \ll \frac{1}{\|A\|}$, where $\|\cdot\|$ is the operator norm of a matrix. At such time horizons, the deviation of $e^{tA}X(0)$ from the initial condition $X(0)$ is relatively small in any suitable sense. This allows the network to be used as a temporary memory (in the Heisenberg picture) for storing (at least approximately) $X(0)$. However, the network response to the initial condition in (49) is “corrupted” by the zero-mean Gaussian quantum Ito process [23, Section 3] $Z := (Z_j)_{j \in V}$ adapted to the filtration of the Fock space \mathfrak{F} in (6) and driven by the quantum Wiener process W in the vacuum state (11), with $Z(0) = 0$. Note that the process Z commutes with the components of $X(0)$ (since the latter act on the different initial network space \mathfrak{H}_0 in (6)):

$$[X(0), Z(t)^T] = 0, \quad t \geq 0. \quad (50)$$

Furthermore, if the network-field quantum state on the space \mathfrak{H} in (6) also has a tensor-product structure given by

$$\rho := \rho_0 \otimes \nu,$$

with ρ_0 the initial network state on \mathfrak{H}_0 , then, in addition to (50), the process Z is statistically independent of $X(0)$. In particular,

$$\mathbf{E}(X(0)Z(t)^T) = \mathbf{E}X(0) \underbrace{\mathbf{E}Z(t)^T}_0 = 0, \quad (51)$$

where $\mathbf{E}\zeta := \text{Tr}(\rho\zeta)$ is the expectation of an operator ζ on the network-field space \mathfrak{H} .

The above commutation and statistical properties can be used in order to quantify the ability of the quantum network as a memory system to retain the initial condition of a quantum process

$$\varphi(t) := FX(t) = \sum_{j \in V} F_j X_j(t), \quad t \geq 0. \quad (52)$$

Here, $F \in \mathbb{R}^{s \times n}$ is a given constant matrix which has full row rank

$$\text{rank} F = s \leq n \quad (53)$$

and is partitioned into blocks $F_j \in \mathbb{R}^{s \times n_j}$ in accordance with (4):

$$F := (F_j^T)_{j \in V}^T. \quad (54)$$

The rows of F specify the coefficients of s independent linear combinations (in particular, a subset) of the network variables which need (or have a better chance) to be preserved over a bounded time interval. Note that, the process φ in (52), in general, involves the internal variables of different OQHOs in the network and thus corresponds to a spatially distributed storage of quantum information. The deviation of φ from its initial value is related by

$$\eta(t) := \varphi(t) - \varphi(0) = F\xi(t) \quad (55)$$

to a similar deviation for the underlying network process X :

$$\xi(t) := X(t) - X(0) = \alpha_t X(0) + Z(t). \quad (56)$$

Here, the last equality follows from (49) and employs an auxiliary matrix

$$\alpha_t := e^{tA} - I_n. \quad (57)$$

Following [26], [27], we will describe the quantum network memory performance in terms of a quadratic form

$$Q(t) := \eta(t)^T \eta(t) = \xi(t)^T \Sigma \xi(t), \quad (58)$$

where, in accordance with (55), the weighting matrix $0 \preceq \Sigma \in \mathbb{S}_n$ is factorised as

$$\Sigma := F^T F, \quad (59)$$

so that $\text{rank} \Sigma = s$ due to (53). Note that $\xi(0) = 0$, $\eta(0) = 0$, $\alpha_0 = 0$ and $Q(0) = 0$ in view of (55)–(58). At an arbitrary moment of time $t \geq 0$, the “size” of the deviation $\eta(t)$ in (55) can be quantified by a mean-square functional

$$\Delta(t) := \mathbf{E}Q(t) = \langle \Sigma, \text{Re} \Upsilon(t) \rangle = \|F\alpha_t \sqrt{P}\|^2 + \langle \Sigma, \text{Re} \Lambda(t) \rangle \quad (60)$$

using (58), as proposed in [26], [27], where $\|\cdot\|$, $\langle \cdot, \cdot \rangle$ are the Frobenius norm and inner product of matrices [9], respectively. Here,

$$\Upsilon(t) := \mathbf{E}(\xi(t)\xi(t)^T) = \alpha_t \Pi \alpha_t^T + \Lambda(t)$$

is the one-point second-moment matrix of the process ξ found in [26] by using (50), (51), (56), (57) and the second-moment matrix

$$\Pi := \mathbf{E}(X(0)X(0)^T) = P + i\Theta, \quad P := \text{Re}\Pi \quad (61)$$

of the initial network variables in $X(0)$, including their augmented CCR matrix Θ from (5), and the quantum covariance matrix

$$\Lambda(t) := \mathbf{E}(Z(t)Z(t)^T) = \int_0^t e^{sA} B \Omega B^T e^{sA^T} ds, \quad t \geq 0 \quad (62)$$

of the process Z in (49), with Ω the quantum Ito matrix from (14). The matrix $\Lambda(t)$ in (62) coincides with the controllability Gramian of the pair $(A, B\sqrt{\Omega})$ over the time interval $[0, t]$ and satisfies the initial value problem for the Lyapunov ODE

$$\dot{\Lambda}(t) = A\Lambda(t) + \Lambda(t)A^T + \mathcal{U}, \quad t \geq 0, \quad \Lambda(0) = 0, \quad (63)$$

where

$$\mathcal{U} := B\Omega B^T. \quad (64)$$

From (63), the time derivatives $\Lambda^{(k)} := d^k \Lambda / dt^k$ can be computed recursively as $\Lambda^{(k+1)} = A\Lambda^{(k)} + \Lambda^{(k)}A^T + \delta_{k0}\mathcal{U}$ for any $k \geq 0$, where δ_{jk} is the Kronecker delta. In particular, the first three derivatives at $t = 0$ are

$$\dot{\Lambda}(0) = \mathcal{U}, \quad (65)$$

$$\ddot{\Lambda}(0) = A\mathcal{U} + \mathcal{U}A^T, \quad (66)$$

$$\dddot{\Lambda}(0) = A^2\mathcal{U} + \mathcal{U}(A^T)^2 + 2A\mathcal{U}A^T. \quad (67)$$

In [26], the relations (64)–(67) were used in order to compute the following time derivatives of $\Delta(t)$ in (60) at $t = 0$:

$$\dot{\Delta}(0) = \|FB\|^2, \quad (68)$$

$$\ddot{\Delta}(0) = \langle \Sigma, ABB^T + BB^T A^T + 2APA^T \rangle, \quad (69)$$

thus leading to the short-horizon asymptotic behaviour of the function

$$\Delta(t) = \dot{\Delta}(0)t + \frac{1}{2}\ddot{\Delta}(0)t^2 + \mathcal{O}(t^3), \quad \text{as } t \rightarrow 0+. \quad (70)$$

For improving the performance of the network as a quantum memory system for storing the initial condition $\varphi(0)$ of the process (52), it is beneficial to minimize the mean-square deviation functional $\Delta(t)$ in (60) at a suitable time $t > 0$. Such minimization can be carried out over the energy and coupling parameters of the network. This problem can also be approached through maximizing the memory decoherence time [26], [27]

$$\tau(\varepsilon) := \inf\{t \geq 0 : \Delta(t) \geq \varepsilon\Delta_*\} \quad (71)$$

as a horizon by which the subnetwork variables do not deviate from their initial values “too far”. The threshold value $\varepsilon\Delta_*$ in (71) for the mean-square deviation functional (60) involves

$$\Delta_* := \mathbf{E}(\varphi(0)^T \varphi(0)) = \langle \Sigma, \Pi \rangle = \|F\sqrt{P}\|^2 \quad (72)$$

as a reference scale (obtained from (52), (59), (61)) and a dimensionless fidelity parameter $\varepsilon > 0$. The latter specifies a relative error threshold for $\varphi(t)$ to approximately reproduce $\varphi(0)$. While $\tau(0) = 0$ by $\Delta(0) = 0$ in (60), we eliminate from consideration the trivial case $\Delta_* = 0$ when $\tau(\varepsilon) = 0$ for any $\varepsilon > 0$ by assuming that

$$F\sqrt{P} \neq 0. \quad (73)$$

Under this condition, the decoherence time $\tau(\varepsilon)$ in (71), which is a nondecreasing function of the fidelity parameter ε , satisfies $\tau(\varepsilon) > 0$ for any $\varepsilon > 0$.

The condition (73) is relevant if $s < n$ when, in view of (53), the columns of the matrix F are linearly dependent. Since

$$[\varphi(t), \varphi(t)^T] = 2iF\Theta F^T \quad (74)$$

by (5), (52), the fulfillment of $F\sqrt{P} = 0$ in (72) implies that $iF\Theta F^T = F\Pi F^T \not\geq 0$ and hence, $F\Theta F^T = 0$, in which case, the components of $\varphi(t)$ commute with each other at any time $t \geq 0$. Here, use is made of the fact that a purely imaginary antisymmetric matrix is positive semi-definite if and only if it is zero. Therefore, if the vector $\varphi(0)$, to be retained by the network, is essentially quantum in the sense that it contains at least one pair of noncommuting variables, that is,

$$F\Theta F^T = \sum_{j \in V} F_j \Theta_j F_j^T \neq 0 \quad (75)$$

in (74), then the condition (73) is necessarily satisfied, which makes (75) sufficient for (73).

VI. APPROXIMATE MEMORY DECOHERENCE TIME MAXIMIZATION

By using the infinite differentiability of $\Delta(t)$ in (60) with respect to $t \geq 0$ along with its time derivatives (68), (69), it was established in [26] that the decoherence time $\tau(\varepsilon)$ in (71) has the following first and second right derivatives at $\varepsilon = 0$:

$$\tau'(0) = \frac{\|F\sqrt{P}\|^2}{\dot{\Delta}(0)} = \frac{\|F\sqrt{P}\|^2}{\|FB\|^2} > 0, \quad (76)$$

$$\tau''(0) = -\frac{\ddot{\Delta}(0)\tau'(0)^2}{\dot{\Delta}(0)} = -\frac{\langle \Sigma, ABB^T + BB^T A^T + 2APA^T \rangle \|F\sqrt{P}\|^4}{\|FB\|^6}. \quad (77)$$

Here, in addition to (73) (or its stronger version (75)), it is assumed that

$$FB \neq 0. \quad (78)$$

Despite having the physical dimension of time, both quantities $\tau'(0)$ and $\tau''(0)$ in (76), (77) resemble the signal-to-noise ratio since their numerators involve FPF^T pertaining to the initial condition $\varphi(0)$ to be stored by the network (see (52), (61)), while $FB B^T F^T = FB \text{Re} \Omega B^T F^T$ in the denominators is associated with the quantum diffusion matrix in

$$d\varphi d\varphi^T = FB \Omega B^T F^T dt$$

coming from the network-field coupling in view of (14), (35). Therefore, under the condition (78), the asymptotic behaviour of $\tau(\varepsilon)$ in the high-fidelity limit (of small values of $\varepsilon > 0$) can be described by an appropriately truncated Taylor series approximation

$$\tau(\varepsilon) = \widehat{\tau}(\varepsilon) + O(\varepsilon^3), \quad \widehat{\tau}(\varepsilon) := \tau'(0)\varepsilon + \frac{1}{2}\tau''(0)\varepsilon^2, \quad (79)$$

as $\varepsilon \rightarrow 0+$. Note that the quantity $\tau'(0)$ in (76) is independent of the matrix A , whereas $\tau''(0)$ in (77) depends on both matrices A and B , with its dependence on A being concave quadratic. As observed in [26], these properties are beneficial for an effective solution of the approximate decoherence time maximization problem

$$\widehat{\tau}(\varepsilon) \longrightarrow \sup \quad (80)$$

(for a given small value of ε) over a suitable set of those energy and coupling parameters of the network that can be varied. More precisely, for fixed network-field and field-mediated coupling, when the matrices M_j , D_{jk} , N_{kj} are fixed for all $j \in V$ and $k \in N_j^+$, and hence, so is the composite network-field coupling matrix M in (36), then the matrix B in (35) is also fixed. If, in addition, all of the individual energy matrices R_j of the component OQHOs are also fixed, then the matrix A in (35) is an affine function of the network energy matrix R and thus of the direct energy coupling matrices R_{jk}^0 in view of (38). This allows the maximization problem (80) over the matrices R_{jk}^0 to be reduced to a convex quadratic minimization problem in the following theorem which is a network extension of [26, Theorems 1, 2]. In order to formulate it, we define, for any $j \in V$ and $k \in N_j^0$, a linear operator g_{jk} on $\mathbb{R}^{n_j \times n_k}$ by

$$g_{jk}(N) := \Theta_j \Sigma_{jj} \Theta_j N P_{kk} + P_{jj} N \Theta_k \Sigma_{kk} \Theta_k + \Theta_j \Sigma_{jk} \Theta_k N^T P_{jk} + P_{jk} N^T \Theta_j \Sigma_{jk} \Theta_k = (g_{kj}(N^T))^T, \quad N \in \mathbb{R}^{n_j \times n_k}, \quad (81)$$

where Θ_j are the CCR matrices from (1), (2), and use is made of the appropriate blocks $(\cdot)_{jk}$ of the matrices Σ , P from (59), (61). Also, $K_{jk} : \mathbb{S}_n \rightarrow \mathbb{R}^{n_j \times n_k}$ is an affine map given by

$$K_{jk}(R) := L_{jk} + 2(\mathbf{S}(\Theta \Sigma \Theta R P))_{jk}, \quad (82)$$

with the matrix

$$L_{jk} := \frac{1}{2}(\mathbf{S}(\Theta \Sigma B(B^T + 2JMP)))_{jk} \quad (83)$$

being a constant term (which does not depend on R) in (82).

Theorem 2: Suppose the network-field and field-mediated coupling matrices M_j , N_{jk} are fixed along with the matrices D_{jk} and the individual energy matrices R_j of the component OQHOs in the network. Also, suppose the fidelity level ε in (71) and the matrix F in (52), (59) are fixed together with the matrix P in (61), and the conditions (73), (78) are satisfied. Then the direct energy coupling matrices R_{jk}^0 deliver a solution to the problem

$$\widehat{\tau}(\varepsilon) \longrightarrow \sup, \quad R_{jk}^0 = R_{kj}^{0T} \in \mathbb{R}^{n_j \times n_k}, \quad j \in V, \quad k \in N_j^0 \quad (84)$$

of maximizing the approximate decoherence time from (79) if and only if

$$g_{jk}(R_{jk}^0) + K_{jk}(\check{R}_{jk}) = 0, \quad (85)$$

where the maps g_{jk} , K_{jk} are given by (81)–(83). Here, the matrix $\check{R}_{jk} \in \mathbb{S}_n$ is obtained by letting $R_{jk}^0 = R_{kj}^{0T} := 0$ in (that is, removing R_{jk}^0 and R_{kj}^0 from) the network energy matrix R in (37), (38), so that $K_{jk}(\check{R}_{jk})$ does not depend on R_{jk}^0 . \square

Proof: Since $\tau'(0)$ in (76) does not depend on the network energy matrix R , the maximization (84) of $\widehat{\tau}(\varepsilon)$ from (79) reduces to that of $\tau''(0)$:

$$\sup_{R_{jk}^0} \widehat{\tau}(\varepsilon) = \tau'(0)\varepsilon + \frac{1}{2}\varepsilon^2 \sup_{R_{jk}^0} \tau''(0). \quad (86)$$

In turn, (77) allows the rightmost supremum in (86) to be represented as

$$\sup_{R_{jk}^0} \tau''(0) = -\frac{\|F\sqrt{P}\|^4}{\|FB\|^6} \inf_{R_{jk}^0} \ddot{\Delta}(0). \quad (87)$$

By (86) and (87), the optimization problem (84) is equivalent to

$$\ddot{\Delta}(0) \longrightarrow \inf, \quad R_{jk}^0 = R_{kj}^{0T} \in \mathbb{R}^{n_j \times n_k}, \quad j \in V, \quad k \in N_j^0, \quad (88)$$

which is a convex quadratic minimization problem in view of (69) and the affine dependence of A in (35) on R , with R being an affine function of the direct energy coupling matrices R_{jk}^0 . Therefore, the first-order optimality condition

$$\partial_{R_{jk}^0} \ddot{\Delta}(0) = 0, \quad j \in V, \quad k \in N_j^0, \quad (89)$$

for the partial Frechet derivatives [19] of $\ddot{\Delta}(0)$ with respect to the matrices R_{jk}^0 is both necessary and sufficient for them to deliver a global minimum to $\ddot{\Delta}(0)$. As obtained in the proof of [26, Theorem 1], the Frechet derivative of $\ddot{\Delta}(0)$ with respect to the whole network energy matrix R is given by

$$\partial_R \ddot{\Delta}(0) = -4\mathbf{S}(\Theta\Sigma(BB^T + 2AP)). \quad (90)$$

The first variation of the composite function $R_{jk}^0 \mapsto R \mapsto \ddot{\Delta}(0)$ can now be computed by applying the chain rule as

$$\delta \ddot{\Delta}(0) = \langle \partial_R \ddot{\Delta}(0), \delta R \rangle = \langle (\partial_R \ddot{\Delta}(0))_{jk}, \delta R_{jk}^0 \rangle + \langle (\partial_R \ddot{\Delta}(0))_{kj}, \delta R_{jk}^{0T} \rangle = 2\langle (\partial_R \ddot{\Delta}(0))_{jk}, \delta R_{jk}^0 \rangle. \quad (91)$$

Here, use is also made of the symmetry of the matrix $\partial_R \ddot{\Delta}(0)$ in (90) along with the relation $\langle \alpha^T, \beta^T \rangle = \langle \alpha, \beta \rangle$ for identically dimensioned matrices. A combination of (91) with (90) yields the partial Frechet derivatives in (89):

$$\partial_{R_{jk}^0} \ddot{\Delta}(0) = 2(\partial_R \ddot{\Delta}(0))_{jk} = -8(\mathbf{S}(\Theta\Sigma(BB^T + 2AP)))_{jk}. \quad (92)$$

The right-hand side of (92) is an affine function of R , which can be split into R_{jk}^0 -independent and R_{jk}^0 -dependent parts as

$$-\frac{1}{16} \partial_{R_{jk}^0} \ddot{\Delta}(0) = \frac{1}{2}(\mathbf{S}(\Theta\Sigma(BB^T + 2AP)))_{jk} = \frac{1}{2}(\mathbf{S}(\Theta\Sigma(BB^T + 2\check{A}_{jk}P)))_{jk} + g_{jk}(R_{jk}^0) = K_{jk}(\check{R}_{jk}) + g_{jk}(R_{jk}^0). \quad (93)$$

Here, the linear map g_{jk} from (81) is evaluated at the matrix R_{jk}^0 , and

$$K_{jk}(\check{R}_{jk}) := \frac{1}{2}(\mathbf{S}(\Theta\Sigma(BB^T + 2\check{A}_{jk}P)))_{jk}, \quad (94)$$

with

$$\check{A}_{jk} := 2\Theta(\check{R}_{jk} + M^T J M) = 2\Theta\check{R}_{jk} + B J M \quad (95)$$

in accordance with (35) and with the matrix \check{R}_{jk} being obtained by removing R_{jk}^0 (and also $R_{kj}^0 = R_{jk}^{0T}$) from R in (38). Note that the relations (94), (95) are in agreement with (82), (83). By (93), the optimality condition (89) takes the form (85), thus establishing the latter as a necessary and sufficient condition of optimality for the problem (88) and its equivalent (84). ■

The linear operator g_{jk} , defined by (81) on the Hilbert space $\mathbb{R}^{n_j \times n_k}$ (with the Frobenius inner product $\langle \cdot, \cdot \rangle$), is self-adjoint and negative semi-definite. Indeed, in view of (93),

$$g_{jk} = -\frac{1}{16} \partial_{R_{jk}^0}^2 \ddot{\Delta}(0) = g_{jk}^\dagger \preceq 0 \quad (96)$$

since the second Frechet derivative $\partial_{R_{jk}^0}^2 \ddot{\Delta}(0)$ of the convex quadratic function $R \mapsto \ddot{\Delta}(0)$ in (69) is a positive semi-definite self-adjoint operator. Therefore, if the operators g_{jk} are all negative definite (and hence, have well-defined inverses g_{jk}^{-1}), then the optimality condition (85) can be represented as

$$R_{jk}^0 = -g_{jk}^{-1}(K_{jk}(\check{R}_{jk})), \quad j \in V, \quad k \in N_j^0. \quad (97)$$

By using the vectorization $\vec{\cdot}$ (or $\text{vec}(\cdot)$) of matrices [13], the action of the linear operator g_{jk} in (81) can be expressed as

$$\text{vec}(g_{jk}(N)) = Q_{jk} \vec{N} \quad (98)$$

in terms of an auxiliary matrix $0 \succ Q_{jk} \in \mathbb{S}_{n_j n_k}$ given by

$$Q_{jk} := P_{kk} \otimes (\Theta_j \Sigma_{jj} \Theta_j) + (\Theta_k \Sigma_{kk} \Theta_k) \otimes P_{jj} + (P_{kj} \otimes (\Theta_j \Sigma_{jk} \Theta_k) + (\Theta_k \Sigma_{kj} \Theta_j) \otimes P_{jk}) \mathbf{T}_{jk}. \quad (99)$$

Here, \mathbf{T}_{jk} is a permutation matrix of order $n_j n_k$ which represents the matrix transpose in $\mathbb{R}^{n_j \times n_k}$ in vectorized form: $\text{vec}(N^T) = \mathbf{T}_{jk} \tilde{N}$ for any $N \in \mathbb{R}^{n_j \times n_k}$. Hence, in view of (96), the negative definiteness of g_{jk} is equivalent to $\det Q_{jk} \neq 0$. In this case, (98), (99) allow (97) to be vectorized as

$$\tilde{R}_{jk}^0 = -Q_{jk}^{-1} \text{vec}(K_{jk}(\check{R}_{jk})), \quad j \in V, k \in N_j^0. \quad (100)$$

Note that (85) describes a set of linear algebraic equations (with a resemblance to the Sylvester equations [4], [22]) for blocks R_{jk}^0 of a sparse symmetric matrix whose sparsity structure is specified by the edges of the direct energy coupling graph of the quantum network. In the case of $g_{jk} < 0$, the alternative form (97) of this set of equations (including its vectorized representation (100)) allows it to be approached as a fixed-point problem whose practical solution can take into account the specific ‘‘spatially distributed’’ structure. Also note that the network-field and field-mediated coupling matrices M_j, N_{jk} enter the set of equations (85) through the matrices (83) (which depend on M from (36) in a quadratic fashion in view of the second equality in (35)) and also through the matrices \check{R}_{jk} in view of (38).

VII. PARTIAL SUBNETWORK ISOLATION

As obtained in [28], under a certain rank condition, the matrix F in (52) can be found so as to select those linear combinations (or a subset) of the network variables which pertain to a partially isolated subnetwork whose dynamics are affected by the external fields only indirectly. This is specified by the following lemma which is an appropriate adaptation (with an almost verbatim proof) of [28, Lemma 1].

Lemma 1: Suppose the network-field coupling matrix M in (36) satisfies

$$d := n - \text{rank} M > 0.$$

Then for any $s \leq d$, there exists a full row rank matrix $F \in \mathbb{R}^{s \times n}$ such that

$$FB = 0 \quad (101)$$

for the matrix B from (35), and the process φ in (52) satisfies

$$\dot{\varphi}(t) = GX(t). \quad (102)$$

Here, the matrix $G \in \mathbb{R}^{s \times n}$ is given by

$$G := 2F\Theta R, \quad (103)$$

where Θ is the CCR matrix for the network variables from (5), and R is the network energy matrix computed in (37), (38). ■

In view of the partitioning (54) of the matrix F and the parameterization of the matrix B in (35), the condition (101) is equivalent to

$$\sum_{j \in V} F_j \Theta_j M_{kj}^T = F_k \Theta_k M_k^T + \sum_{j \in N_k^+} F_j \Theta_j N_{jk}^T D_{kj} = 0, \quad k \in V, \quad (104)$$

where the block-diagonal structure of the CCR matrix Θ in (5) is used together with the blocks (36) of the network-field coupling matrix M . The representation (104) of (101) shows that an appropriate ‘‘tuning’’ of the field-mediated coupling between the constituent QHOs in the network can provide an additional freedom in choosing the matrix F subject to (101).

The absence of the quantum diffusion term (involving W) on the right-hand side of (102) makes it an ODE and allows φ to be interpreted as a vector of subnetwork variables, partially isolated from the external fields. Compared to the completely autonomous subsystems arising in the quantum Kalman decomposition [35], this isolation is partial because φ is indirectly affected by W through ‘‘complementary’’ network variables which form a quantum process

$$\psi(t) := TX(t). \quad (105)$$

Here, $T \in \mathbb{R}^{(n-s) \times n}$ is any matrix whose rows complement those of the matrix F in Lemma 1 to a nonsingular matrix:

$$S := \begin{bmatrix} F \\ T \end{bmatrix} \in \mathbb{R}^{n \times n}, \quad \det S \neq 0.$$

Therefore, an appropriate similarity transformation allows the network QSDE (34) to be represented as an ODE for φ and a QSDE for ψ as

$$\dot{\varphi} = a_{11}\varphi + a_{12}\psi, \quad d\psi = (a_{21}\varphi + a_{22}\psi)dt + bdW, \quad (106)$$

where the matrices $a_{11} \in \mathbb{R}^{s \times s}$, $a_{12} \in \mathbb{R}^{s \times (n-s)}$, $a_{21} \in \mathbb{R}^{(n-s) \times s}$, $a_{22} \in \mathbb{R}^{(n-s) \times (n-s)}$ and $b \in \mathbb{R}^{(n-s) \times m}$ are given by

$$a := \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = SAS^{-1}, \quad b := TB.$$

The decomposition (106) of the network dynamics can be viewed as an interconnection of a partially isolated subnetwork Φ with the vector φ of internal variables in (52) interacting with a complementary subnetwork Ψ with the vector ψ of internal variables in (105), with only Ψ being directly affected by the external quantum Wiener process W in (13). In this sense, Ψ “shields” Φ from W ; see Fig. 3.




Fig. 3. A schematic representation of (106) as an interconnection of quantum subnetworks Φ , Ψ , where Φ is affected by the external fields W only through Ψ which interacts with W .

The special choice of the matrix F in Lemma 1 subject to (101) makes the coefficient $\dot{\Delta}(0)$ in (68) vanish. Hence, the quadratic term becomes dominant in (70) which acquires the form [28, Lemma 2]

$$\Delta(t) = \frac{1}{2} \ddot{\Delta}(0) t^2 + O(t^3), \quad \ddot{\Delta}(0) = 2 \|G\sqrt{P}\|^2, \quad (107)$$

as $t \rightarrow 0+$, with the matrices G from (103) and P from (61). The fact that the short-horizon asymptotic behaviour (107) is quadratic (rather than linear) in time is a consequence of the partial isolation of the subnetwork Φ from the external fields (see Fig. 3). This is a qualitatively slower growth of the function Δ compared to the case of matrices F satisfying (78). Accordingly, the partial subnetwork isolation prolongs the memory decoherence time in the high-fidelity limit as the following adaptation of [28, Theorem 1] shows.

Lemma 2: Suppose the conditions of Lemma 1 are satisfied together with (73) and

$$G\sqrt{P} \neq 0.$$

Then the memory decoherence time (71) behaves asymptotically as

$$\tau(\varepsilon) \sim \frac{\|F\sqrt{P}\|}{\|G\sqrt{P}\|} \sqrt{\varepsilon} =: \hat{\tau}(\varepsilon), \quad \text{as } \varepsilon \rightarrow 0+. \quad (108)$$

As a result of the partial subnetwork isolation, the asymptotic behaviour (108) in the high-fidelity limit yields a longer decoherence time compared to the case of (78), where $\tau(\varepsilon)$ is asymptotically linear with respect to ε in view of (76). Indeed, $\varepsilon = o(\sqrt{\varepsilon})$, as $\varepsilon \rightarrow 0+$. In the framework of (108), the maximization of $\tau(\varepsilon)$ at a given small value of ε can also be replaced with its approximate version (80). Assuming the matrices F and P to be fixed, the problem (80), with the approximate decoherence time $\hat{\tau}(\varepsilon)$ from (108), is equivalent to the minimization of the denominator

$$\|G\sqrt{P}\| = 2 \|F\Theta R\sqrt{P}\| \quad (109)$$

(see (103)), which is a convex function of the network energy matrix R . The following theorem is a counterpart of Theorem 2 in the partial isolation setting. Accordingly, (78) is replaced with (101), and (79) is replaced with (108).

Theorem 3: Suppose the network-field and field-mediated coupling matrices M_j, N_{jk} are fixed along with the matrices D_{jk} and the individual energy matrices R_j of the component OQHOs in the network. Also, suppose the fidelity level ε in (71) and the matrix F in (52), (59) are fixed together with the matrix P in (61), and the conditions (73), (101) are satisfied. Then the direct energy coupling matrices R_{jk}^0 deliver a solution to the problem (84) of maximizing the approximate decoherence time $\hat{\tau}(\varepsilon)$ from (108) if and only if they satisfy (85) with the maps g_{jk} from (81), and the maps K_{jk} from (82) being replaced with

$$K_{jk}(R) := 2(\mathbf{S}(\Theta\Sigma\Theta RP))_{jk}. \quad (110)$$

□

Proof: Although (110) can be obtained from (82) by noting that, in view of (59), $\Sigma B = F^T F B = 0$ in the partial subnetwork isolation case (101) and hence, $L_{jk} = 0$ in (83), we will provide an independent proof of the present theorem. As mentioned above, the problem (84), with the approximate decoherence time $\hat{\tau}(\varepsilon)$ from (108), reduces to the minimization of the quantity (109). The latter is equivalent to minimizing

$$\phi := \frac{1}{2} \|F \Theta R \sqrt{P}\|^2 = \frac{1}{16} \ddot{\Delta}(0) \quad (111)$$

(cf. (103), (107)) as a convex quadratic function of the direct energy coupling matrices $R_{jk}^0 = R_{kj}^{0T} \in \mathbb{R}^{n_j \times n_k}$, with $j \in V$, $k \in N_j^0$, and the $\frac{1}{2}$ -factor introduced for convenience. Therefore, (111) achieves a global minimum if and only if the partial Frechet derivatives vanish:

$$\partial_{R_{jk}^0} \phi = 0, \quad j \in V, k \in N_j^0. \quad (112)$$

By using the variational identity $\delta(\|z\|^2) = 2\langle z, \delta z \rangle$ which holds for a real matrix z , the first variation of ϕ in (111) with respect to the matrix R_{jk}^0 can be computed, similarly to (91), as

$$\delta \phi = \langle F \Theta R \sqrt{P}, F \Theta (\delta R) \sqrt{P} \rangle = -\langle \Theta F^T F \Theta R P, \delta R \rangle = -\langle \mathbf{S}(\Theta \Sigma \Theta R P), \delta R \rangle = -2\langle (\mathbf{S}(\Theta \Sigma \Theta R P))_{jk}, \delta R_{jk}^0 \rangle. \quad (113)$$

Here, use is also made of the antisymmetry of the CCR matrix Θ in (5) and the symmetry of P in (61) along with (59). The relation (113) leads to the partial Frechet derivative

$$-\partial_{R_{jk}^0} \phi = 2(\mathbf{S}(\Theta \Sigma \Theta R P))_{jk} = g_{jk}(R_{jk}^0) + K_{jk}(\check{R}_{jk}) \quad (114)$$

represented in terms of (81), (110). By (112), (114), the condition (85), with the maps K_{jk} from (110), is necessary and sufficient for optimality in the problem (84) with the approximate decoherence time from (108) in the partial subnetwork isolation case (101). \blacksquare

Note that Theorem 3 is essentially an adaptation of Theorem 2 to the partial subnetwork isolation setting, with both of them reducing to the minimization of $\ddot{\Delta}(0)$ from (69) which takes a specific form (107) in this case. Accordingly, the previous remarks on the general structure and numerical solution of the equations (85) for an optimal direct energy coupling remain applicable in this case too. At the same time, in the partial subnetwork isolation case, the maps K_{jk} in (110) do not involve the network-field and field-mediated coupling matrices. Therefore, other than affecting the choice of the matrix F subject to (101), these matrices enter the equations (85) only through the matrices \check{R}_{jk} .

VIII. CONCLUSION

For a network of OQHOs with direct energy and field-mediated couplings, driven by external quantum fields, we have considered the problem of maximizing the high-fidelity approximation of the memory decoherence time (associated with the weighted mean-square deviation of the network variables from their initial conditions) over the direct energy coupling matrices. Using the convex quadratic structure of this problem, we have obtained a necessary and sufficient condition of optimality in the form of a set of Sylvester-like linear algebraic equations for blocks of a matrix whose sparsity is specified by the edges of the direct energy coupling graph of the network. The approximate decoherence time maximization problem and its solution have also been discussed for a partially isolated subnetwork with a qualitatively longer decoherence time and thus with an enhanced memory system performance. The development of an algorithm for numerical solution of the resulting sets of equations can benefit from a combination of vectorization techniques with advanced graph theoretic considerations. Other possible directions of research on this topic include the extension of the quantum memory optimization approach beyond the high-fidelity (that is, short-horizon) approximations and to finite-level open quantum networks with multipartite interactions.

REFERENCES

- [1] L.A.D.Espinosa, Z.Miao, I.R.Petersen, V.Ugrinovskii, and M.R.James, Physical realizability and preservation of commutation and anticommutation relations for n -level quantum systems, *SIAM J. Control Optim.*, vol. 54, no. 2, 2016, pp. 632–661.
- [2] S.Fu, A.R.R.Carvalho, M.R.Hush, and M.R.James, Cross-phase modulation and entanglement in a compound gradient echo memory, *Phys. Rev. A*, vol. 93, 2016, 023809.
- [3] C.W.Gardiner, and P.Zoller, *Quantum Noise*, Springer, Berlin, 2004.
- [4] J.D.Gardiner, A.J.Laub, J.J.Amato, and C.B.Moler, Solution of the Sylvester Matrix Equation $AXB^T + CXD^T = E$, *ACM Trans. Math. Soft.*, vol. 18, no. 2, 1992, pp. 223–231.
- [5] J.Gough, and M.R.James, Quantum feedback networks: Hamiltonian formulation, *Comm. Math. Phys.*, vol. 287, 2009, pp. 1109–1132.
- [6] K.Heshami, D.G.England, P.C.Humphreys, P.J.Bustard, V.M.Acosta, J.Nunn, B.J.Sussman, Quantum memories: emerging applications and recent advances, *J. Mod. Opt.*, vol. 63, no. 20, 2016, pp. 2005–2028.
- [7] A.S.Holevo, *Statistical Structure of Quantum Theory*, Springer, Berlin, 2001.
- [8] A.S.Holevo, *Quantum Systems, Channels, Information: A Mathematical Introduction*, De Gruyter, Berlin, 2019.
- [9] R.A.Horn, and C.R.Johnson, *Matrix Analysis*, Cambridge University Press, New York, 2007.
- [10] R.L.Hudson, and K.R.Parthasarathy, Quantum Ito's formula and stochastic evolutions, *Commun. Math. Phys.*, vol. 93, 1984, pp. 301–323.
- [11] M.R.James, H.I.Nurdin, and I.R.Petersen, H^∞ control of linear quantum stochastic systems, *IEEE Trans. Automat. Contr.*, vol. 53, no. 8, 2008, pp. 1787–1803.

- [12] L.D.Landau, and E.M.Lifshitz, *Quantum Mechanics: Non-relativistic Theory*, 3rd Ed., Butterworth-Heinemann, Amsterdam, 1981.
- [13] J.R.Magnus, *Linear Structures*, Oxford University Press, New York, 1988.
- [14] M.A.Nielsen, and I.L.Chuang, *Quantum Computation and Quantum Information*, Cambridge University Press, Cambridge, 2000.
- [15] H.I.Nurdin, and N.Yamamoto, *Linear Dynamical Quantum Systems*, Springer, Netherlands, 2017.
- [16] K.R.Parthasarathy, *An Introduction to Quantum Stochastic Calculus*, Birkhäuser, Basel, 1992.
- [17] K.R.Parthasarathy, and K.Schmidt, *Positive Definite Kernels, Continuous Tensor Products, and Central Limit Theorems of Probability Theory*, Springer-Verlag, Berlin, 1972.
- [18] I.R.Petersen, Quantum linear systems theory, *Open Automat. Contr. Syst. J.*, vol. 8, 2017, pp. 67–93.
- [19] M.Reed, and B.Simon, *Methods of Modern Mathematical Physics: Functional Analysis*, vol. 1, 2nd Ed., Academic Press, San Diego, 1980.
- [20] J.J.Sakurai, *Modern Quantum Mechanics*, Addison-Wesley, Reading, Mass., 1994.
- [21] A.J.Shaiju, and I.R.Petersen, A frequency domain condition for the physical realizability of linear quantum systems, *IEEE Trans. Automat. Contr.*, vol. 57, no. 8, 2012, pp. 2033–2044.
- [22] R.E.Skelton, T.Iwasaki, and K.M.Grigoriadis, *A Unified Algebraic Approach to Linear Control Design*, Taylor & Francis, London, 1998.
- [23] I.G.Vladimirov, I.R.Petersen, and M.R.James, Multi-point Gaussian states, quadratic-exponential cost functionals, and large deviations estimates for linear quantum stochastic systems, *Appl. Math. Optim.*, vol. 83, 2021, pp. 83–137 (published 24 July 2018).
- [24] I.G.Vladimirov, and I.R.Petersen, Moment dynamics and observer design for a class of quasilinear quantum stochastic systems, *SIAM J. Control Optim.*, vol. 60, no. 3, 2022, pp. 1223–1249.
- [25] I.G.Vladimirov, and I.R.Petersen, Decoherence quantification through commutation relations decay for open quantum harmonic oscillators, *Systems & Control Letters*, vol. 178, 2023, 105585.
- [26] I.G.Vladimirov, and I.R.Petersen, Decoherence time in quantum harmonic oscillators as quantum memory systems, 2024 Australian and New Zealand Control Conference (ANZCC), Gold Coast, Australia, 2024, pp. 211–216.
- [27] I.G.Vladimirov, and I.R.Petersen, Decoherence time control by interconnection for finite-level quantum memory systems, *European Journal of Control*, 2024, 101054, published online 15 June 2024, <https://doi.org/10.1016/j.ejcon.2024.101054>.
- [28] I.G.Vladimirov, and I.R.Petersen, Optimization of partially isolated quantum harmonic oscillator memory systems by mean square decoherence time criteria, Australian and New Zealand Control Conference (ANZCC 2025), Gold Coast, 30-31 January 2025, Australia, pp. 64–69, preprint <https://arxiv.org/abs/2409.15720>, 24 September 2024.
- [29] V.S.Vladimirov, *Methods of the Theory of Generalized Functions*, Taylor & Francis, London, 2002.
- [30] D.F.Walls, and G.J.Milburn, *Quantum Optics*, Springer, Berlin, 2008.
- [31] Y.Wang, J.Li, S.Zhang, et al, Efficient quantum memory for single-photon polarization qubits, *Nat. Photonics*, vol. 13, 2019, pp. 346–351.
- [32] N.Yamamoto, and M.R.James, Zero-dynamics principle for perfect quantum memory in linear networks, *New J. Phys.*, vol. 16, 2014, 073032.
- [33] G.Zhang, and Z.Dong, Linear quantum systems: a tutorial, *Annual Reviews in Control*, vol. 54, 2022, pp. 274–294.
- [34] G.Zhang, and M.R.James, Direct and indirect couplings in coherent feedback control of linear quantum systems, *IEEE Trans. Automat. Contr.*, vol. 56, no. 7, 2011, 1535–1550.
- [35] G.Zhang, S.Grivopoulos, I.R.Petersen, and J.E.Gough The Kalman Decomposition for Linear Quantum Systems, *IEEE Transactions on Automatic Control*, vol. 63, no. 2, 2018, pp. 331–346.