

Surface Subgroups for Cocompact Lattices of Isometries of \mathbb{H}^{2n}

Jeremy Kahn and Zhenghao Rao

Abstract

We prove the existence of surface subgroups within any cocompact lattice Γ in $\mathrm{SO}(2n, 1)$ for $n \geq 2$. This result addresses the cases missing from the work of Hamenstädt in 2015, who constructed surface subgroups in cocompact lattices for all other rank-one semisimple Lie groups of non-compact type.

1 Introduction

A *surface group*, for our purposes, is the fundamental group of a closed hyperbolic surface. A surface *subgroup* of a group is a subgroup that is a surface group.

In [KM12], Kahn and Markovic construct surface subgroups of cocompact lattices in $\mathrm{Isom}(\mathbb{H}^3)$, while Kahn and Wright [KW21] later proved the existence of surface subgroups in non-uniform lattices in $\mathrm{Isom}(\mathbb{H}^3)$. In [Ham15], Hamenstädt constructs surface subgroups of cocompact lattices in any rank one simple Lie group *except* for $\mathrm{Isom}(\mathbb{H}^{2n})$ for $n \geq 2$. Kahn, Labourie and Mozes [KLM24] built surface subgroups in cocompact lattices in center-free, complex semisimple Lie group of non-compact type. In this paper, we show that for all $n \geq 2$, every cocompact lattice of isometries of \mathbb{H}^{2n} has a surface subgroup. Here is our main result.

Theorem 1.1. *Suppose $\Gamma < \mathrm{SO}^+(2n, 1)$ is a cocompact lattice with $n > 1$. Then Γ contains surface subgroups.*

Although we do not state a quantitative result, the surface subgroups we construct here could be K -quasi-Fuchsian for any $K > 1$, as mentioned by Hamenstädt [Ham15].

In this introduction, we describe the basic idea used in [KM12] (and [Ham15], [KW21], [KLM24], [Rao25]), then explain the difficulty of applying it to \mathbb{H}^{2n} , and finally outline our approach to resolve the issue.

In [KM12], the authors consider “good pants” in $M = \Gamma \backslash \text{Isom}(\mathbb{H}^3)$, which are ϵ -nearly isometric immersions of the “ R -perfect pants” into M ; the R -perfect pants is the unique hyperbolic pair of pants with three geodesic boundaries of length R . Each good pants in M has three *short orthogeodesics*, each connecting two of the boundary cuffs; each short orthogeodesic has two *feet*, which are unit normal vectors to the boundary geodesics and point toward the orthogeodesic. We then consider the distribution of the feet of good pants around each “good” closed geodesic γ in M . We see that these are evenly distributed in the unit normal bundle $N^1(\gamma)$, and this implies that there is a permutation $\sigma: \mathcal{F}_\gamma \rightarrow \mathcal{F}_\gamma$ (where \mathcal{F}_γ is the set of feet of good pants on γ) that closely approximates the translation $\tau: N^1(\gamma) \rightarrow N^1(\gamma)$ that sends each unit normal vector to the vector obtained by parallel transport (positive) distance one along γ , and then negating the normal vector. We use this permutation (and a doubling trick) to pair off the good pants, and thereby make a “good panted surface”. Finally, we verify that any good panted surface is π_1 -injective.

We can outline this approach as follows:

1. Define a notion of good pants (depending on parameters ϵ and R), which should be some precise version of “the image of the R -perfect pants by an immersion that is ϵ -close to an isometry”. Take one of each good pants (as in [KW21]), or define some other distribution on the finite set of (R, ϵ) -good pants.
2. For each good pants Π and boundary geodesic γ , describe the (short) foot of Π on γ .
3. For each closed geodesic γ in M that is good (in the sense of being the boundary of at least one good pants), show that the feet of good pants are *(approximately) evenly distributed*.
4. Conclude from this even distribution that there is a permutation of the feet that closely approximates the “negate and twist by one” translation τ on $N^1(\gamma)$.
5. Use this permutation and the “doubling trick” to match good pants along their common boundaries and form a good panted surface.

6. Show that every good panted surface is π_1 -injective.

In [Ham15], the author (implicitly) follows the six-step outline above; the greatest novelty lies in a completely different approach to Step 6, using the differential geometry of rank-one symmetric spaces (of non-compact type). To see why \mathbb{H}^{2n} presents a difficulty, we should properly understand the meaning of “evenly distributed” in Step 3. Let $\tilde{\gamma}$ be a component of the preimage of γ in the universal cover (the symmetric space), and let T_γ be a generator of the deck group preserving $\tilde{\gamma}$, so $\gamma = \tilde{\gamma}/T_\gamma$. We let Z_γ be the *centralizer* of T_γ , and Z'_γ the component of Z_γ containing the identity.

Here is an explicit description of Z'_γ and Z_γ in \mathbb{H}^{2n} . Since $T_\gamma \in Z'_\gamma \subset Z_\gamma$, we can think of Z'_γ and Z_γ as acting on \mathbb{H}^{2n}/T_γ , and in fact on $N^1(\gamma)$. For a generic γ , the monodromy (of parallel transport) at the fiber of $N^1(\gamma)$ at a single point is a rotation of that $(2n - 2)$ -sphere; we can then think of it as a rotation around the *North and South poles*. Indeed there is a North pole and a South pole for each fiber, and these poles (and all latitudes) are preserved by parallel transport. An element of Z'_γ shifts the fibers by some amount along γ , and then rotates each sphere by some amount around the poles, thus preserving latitude. For Z_γ we add one more generator, that negates latitude and interchanges the North and South poles.




Figure 1: One possible obstruction: the “gap” between the two components

In all cases, one can prove (as Step 3) that the good feet are approximately evenly distributed *with respect to* Z_γ . When τ (defined in Step 4) lies in Z'_γ , this even distribution implies the existence of the permutation required in

Step 4. This holds in every rank one case but \mathbb{H}^{2n} . In the case of \mathbb{H}^{2n} , this τ lies in $Z_\gamma \setminus Z'_\gamma$. In Figure 1, we illustrate a possible distribution of feet that is nearly invariant under Z_γ : in each fiber, the feet are clustered near the north and south poles, with a large gap in the lower latitudes. The problem is that there may be a few more feet in the northern “ice cap” than the southern one, and this would prevent the existence of a bijection between the two sets of feet. Moreover, there could in principle be even more troublesome cases, where the feet lie in bands on the sphere, as in Figure 2.




Figure 2: A worse case: the “bands”

To solve the above problem for \mathbb{H}^{2n} with $n \geq 2$, we first observe that in Step 3, the distribution ν_a of the feet of good pants on γ is approximated by the *estimated foot measure* μ_a , which can be defined analytically through third connections. The use of third connections and this observation are inspired by [KW21]. We then show that Radon–Nikodym derivative of μ_a with respect to the Lebesgue measure on $N^1(\gamma)$ is bounded above and below by constants that depend only on ϵ and the manifold M . This means that there cannot be any “gaps” in μ_a (or ν_a) that would be an obstruction to forming the permutation that approximates τ . We can then directly verify that ν_a satisfies the conditions of Hall’s Marriage Theorem, and finally produce the permutation in Step 4. We expect that this technique can be generalized to higher rank non-compact Lie groups with a “weak flip” [KLM24].

Section 2 introduces preliminaries on the geometry of pants, especially in $2n$ -dimensional hyperbolic space and manifolds. After introducing some properties of $\mathrm{SO}(2n, 1)$ (which also hold for general $\mathrm{SO}(n, 1)$), and basic

definitions for pants, we introduce the notion of *cuff-good* pants, which are pants whose cuffs are good curves (as in [KW21]). Using the *Steiner graph* (which we define) for pants, we then show that cuff-good pants are either good (in the sense of [Ham15] and [KM12], where we generate pants from connections between tripods), or *bad*, in that they are close to a geodesic immersion of a one-holed torus. This then allows us to determine when a pair of pants is good from one cuff and a *third connection* that generates the other two cuffs.

Section 3 contains the core estimate of this paper, where we approximate the distribution of cuff-good pants through counting the third connections. Given a closed geodesic γ , for any framed third connection, we compute the monodromy of the other two cuffs of the pants using matrix multiplication, and then derive an (almost) necessary and sufficient condition of generating cuff-good pants. Thus we can determine the *good region* in the space of third connections, which is a fiber bundle over the unit normal bundle $N^1(\gamma)$ of γ . The projection of the good region yields a measure on $N^1(\gamma)$, which is called the *estimated foot measure*. By evaluating the volume of the good region in each fiber, we show that the estimated foot measure is bounded above and below by a multiple of the Lebesgue measure; this rules out the “ice cap” or “band” obstruction described above.

In Section 4, we first show that our matching pattern satisfies the incompressibility criterion established by Hamenstädt [Ham15]. We then apply Hall’s Marriage Theorem (using our measure bounds and estimates of the Cheeger constant) to produce the permutation on the set of all good pants, which is used to prove the main result of this paper, the existence of π_1 -injective immersed surface in closed hyperbolic $2n$ -manifolds.

Acknowledgments. This work arose out of a conversation in 2014, in which Ursula Hamenstädt explained to the first author the problem that can arise when trying to build a surface out of pants in a quotient of \mathbb{H}^4 .

The second author would like to thank Fernando Al Assal for inviting him to visit MPI MIS in 2023. In a conversation during this visit, Al Assal believed that pants with good cuffs should be good. Although the claim is now known to be not entirely true, it inspired the second author to think about the monodromy of cuffs. Finally, the first and second authors managed to show that a pants in \mathbb{H}^{2n} with good cuffs is either good or bad. The second author would also like to thank Ursula Hamenstädt and Yi Liu for helpful conversations.

2 Some geometry of good pants

2.1 The structure of $\mathrm{SO}^+(2n, 1)$ and $\mathfrak{so}(2n, 1)$

Let $\mathcal{G} = \mathrm{SO}^+(2n, 1)$, which is the identity component of $\mathrm{SO}(2n, 1)$ and acts on \mathbb{H}^{2n} by orientation preserving isometries. We choose $p_0 = (0, \dots, 0, 1) \in \mathbb{H}^{2n}$ to be the base point, and $\mathbf{u}_0 = (0, \dots, 0, 1)$ a unit tangent vector at p_0 . Let \mathcal{K} be the stabilizer of p_0 , then \mathcal{K} , which is isomorphic to $\mathrm{SO}(2n)$, is a maximal compact subgroup of \mathcal{G} . The stabilizer \mathcal{M} of \mathbf{u}_0 is a subgroup of \mathcal{K} which is isomorphic to $\mathrm{SO}(2n - 1)$. Hence \mathcal{G}/\mathcal{K} can be identified with \mathbb{H}^{2n} , \mathcal{G}/\mathcal{M} can be identified with the unit tangent bundle $T^1(\mathbb{H}^{2n})$ of \mathbb{H}^{2n} , and \mathcal{G} itself can be identified with the frame bundle $\mathcal{F}(\mathbb{H}^{2n})$ of \mathbb{H}^{2n} . We always denote by \mathbf{e} the identity element of a group, and $[\cdot]$ the conjugacy class of a group element.

The left action of \mathcal{G} on \mathbb{H}^{2n} as isometry induces a left action of \mathcal{G} on $\mathcal{F}(\mathbb{H}^{2n})$. Now we define the right action of \mathcal{G} on $\mathcal{F}(\mathbb{H}^{2n})$ based on the left action, generalizing the definition in Section 2 of [Rao25] for dimension 2 and 3. Fix a base frame $\Psi_0 = \langle p_0, F_0 \rangle \in \mathcal{F}(\mathbb{H}^{2n})$. Then the right action of \mathcal{G} on $\mathcal{F}(\mathbb{H}^{2n})$ is defined as follows:

- (1) For any $g \in \mathcal{G}$, $\Psi_0 \cdot g = g \cdot \Psi_0$.
- (2) For any $\Psi \in \mathcal{F}(\mathbb{H}^{2n})$ and $g, h \in \mathcal{G}$, $g \cdot (\Psi \cdot h) = (g \cdot \Psi) \cdot h$.

Since the left action of \mathcal{G} on $\mathcal{F}(\mathbb{H}^{2n})$ is transitive and free, then for any Ψ , there exists unique $g \in \mathcal{G}$ such that $\Psi = g \cdot \Psi_0$. Thus for any $h \in \mathcal{G}$,

$$\Psi \cdot h = (g \cdot \Psi_0) \cdot h = g \cdot (\Psi_0 \cdot h),$$

which means the right action on Ψ_0 will tell us the right action on any other frames. Some common examples are frames flows and rotations.

There is a one parameter group $\mathcal{A} = \{a_t : t \in \mathbb{R}\}$ such that the right action of a_t on \mathcal{G} is the unit-speed frame flow, and such that \mathcal{A} commutes with \mathcal{M} . Let $\mathcal{B} = \mathcal{A}\mathcal{M} = \mathcal{M}\mathcal{A} < \mathcal{G}$. We can define the unstable and stable horospherical subgroups \mathcal{N}^+ and \mathcal{N}^- of \mathcal{G} , respectively, as follows:

$$\mathcal{N}^\pm = \{g \in \mathcal{G} : a_t g a_{-t} \rightarrow \mathbf{e}, \text{ as } t \rightarrow \pm\infty\}$$

Let $\mathfrak{a}, \mathfrak{m}, \mathfrak{n}^\pm$ be the corresponding Lie subalgebra in $\mathfrak{so}(2n, 1)$ of $\mathcal{A}, \mathcal{M}, \mathcal{N}^\pm$, respectively. Thus

$$\mathfrak{so}(2n, 1) = \mathfrak{a} \oplus \mathfrak{m} \oplus \mathfrak{n}^+ \oplus \mathfrak{n}^-,$$

as real vector spaces.

By hyperbolic geometry, we have the following facts:

1. For any $g \in \mathcal{G} - \{\mathcal{AN}^+ - \{e\}\}$, there exist unique $n^+ \in \mathcal{N}^+, n^- \in \mathcal{N}^-$ and $t \in \mathbb{R}$ such that

$$g = n^+ a_t n^-.$$

2. For any $X^+ \in \mathfrak{n}^+$ and $t > 0$,

$$a_t \exp(X^+) = \exp(e^{-t} X^+) a_t.$$

3. For any $X^- \in \mathfrak{n}^-$ and $t > 0$,

$$\exp(X^-) a_t = a_t \exp(e^{-t} X^-).$$

Lemma 2.1. *There exist $C, \epsilon_0 > 0$ such that for any $\epsilon_0 > \epsilon > 0$: Suppose that $u \in \mathcal{G}$ is ϵ -close to \mathbf{e} , then u can be uniquely written as*

$$u = n^+ b n^-,$$

for some $n^\pm \in \mathcal{N}^\pm$ and $b \in \mathcal{B}$. Moreover, n^\pm and b are $C\epsilon$ -close to \mathbf{e} .

Let $\|\cdot\|$ be the norm on $\mathfrak{so}(2n, 1)$ by its Killing form, then this norm is preserved by Ad_k for any $k \in \mathcal{K}$.

Lemma 2.2. *For any $X^+ \in \mathfrak{n}^+$, $t > 0$ and $m \in \mathcal{M}$, there exists X_1^+ satisfying the following inequality:*

$$\|X_1^+\| = e^{-t} \|X^+\|,$$

such that

$$a_t m \exp(X^+) = \exp(X_1^+) a_t m.$$

Proof. Let $X_1^+ = \text{Ad}_m(e^{-t} X^+)$, then $\|X_1^+\| = e^{-t} \|X^+\|$, and we have

$$\begin{aligned} a_t m \exp(X^+) (a_t m)^{-1} &= m a_t \exp(X^+) a_t^{-1} m^{-1} \\ &= m \exp(e^{-t} X^+) m^{-1} \\ &= \exp(\text{Ad}_m(e^{-t} X^+)) = \exp(X_1^+). \quad \square \end{aligned}$$

Similarly, we have another lemma.

Lemma 2.3. *For any $X^- \in \mathfrak{n}^-$, $t > 0$ and $m \in \mathcal{M}$, there exists X_1^- satisfying the following inequality:*

$$\|X_1^-\| = e^{-t}\|X^-\|,$$

such that

$$\exp(X^-)a_t m = a_t m \exp(X_1^-).$$

The next two lemmas are actually from the inefficiency theory in \mathcal{G} . The first one tells us that small perturbation can be absorbed into long frame flows and rotations after conjugacy, and the second involves more elements and perturbations which generalize the first one.

Lemma 2.4. *There exist $\epsilon_0, R, K > 0$ such that when $\epsilon_0 > \epsilon > 0$, the following holds: Suppose that $u \in \mathcal{G}$ is ϵ -close to \mathbf{e} , $t > R$ and $m \in \mathcal{M}$. Then there exists $t' > 0$ with $|t - t'| < K\epsilon$ and $m' \in \mathcal{M}$ that is $K\epsilon$ -close to m , such that $a_t m u$ is conjugate to $a_{t'} m'$.*

Proof. Suppose ϵ_0 and C are the constants in Lemma 2.1, and $R > 10000$ large enough, which will be determined later. Next we will find K satisfying the requirements.

By Lemma 2.1, there exist $n_1^\pm \in \mathcal{N}^\pm$, $a_{t_1} \in \mathcal{A}$ and $m_1 \in \mathcal{M}$ such that

$$u = n_1^+ a_{t_1} \hat{m}_1 n_1^-$$

and $n_1^\pm, \hat{m}_1, a_{t_1}$ are all $C\epsilon$ -close to \mathbf{e} . Then by Lemma 2.2 and Lemma 2.3, we have

$$\begin{aligned} [a_t m u] &= [a_t m n_1^+ a_{t_1} \hat{m}_1 n_1^-] = [a_{t/2} m n_1^+ a_{t_1} \hat{m}_1 n_1^- a_{t/2}] \\ &= [\hat{n}_1^+ a_{t/2} m a_{t_1} \hat{m}_1 a_{t/2} \hat{n}_1^-] \\ &= [a_{t+t_1} m \hat{m}_1 \hat{n}_1^- \hat{n}_1^+], \end{aligned}$$

where \hat{n}_1^- and \hat{n}_1^+ are $e^{-t}C\epsilon$ -close to \mathbf{e} . Now we let $t_1 = t + t_1$, $m_1 = m \hat{m}_1$ and $u_1 = \hat{n}_1^- \hat{n}_1^+$. Thus $|t - t_1| < C\epsilon$, m_1 is $C\epsilon$ -close to m , u_1 is $2e^{-t}C\epsilon$ -close to \mathbf{e} , and

$$[a_t m u] = [a_{t_1} m_1 u_1].$$

Let $K = 2C$, and R large enough such that $2e^{-R/2}C < 1/2$ and $t_i > R/2$ for all $i \in \mathbb{Z}_+$. Thus we can have sequences $\{t_i\}_{i=1}^\infty \subset \mathbb{R}_+$, $\{m_i\}_{i=1}^\infty \subset \mathcal{M}$

and $\{u_i\}_{i=1}^\infty \subset \mathcal{G}$ such that for any $i \in \mathbb{Z}_+$, $|t_i - t_{i+1}| < 2^{1-i}C\epsilon$, m_{i+1} is $2^{1-i}C\epsilon$ -close to m_i , u_i is $2^{-i}\epsilon$ -close to \mathbf{e} , and

$$[a_{t_i}m_iu_i] = [a_{t_{i+1}}m_{i+1}u_{i+1}].$$

Therefore the limits of these three sequences exist, and moreover, the limit of $\{u_i\}$ is \mathbf{e} . Let $t_i \rightarrow t'$ and $m_i \rightarrow m'$, as $i \rightarrow \infty$. Hence $[a_t m u] = [a_{t'} m']$, $|t - t'| < 2C\epsilon$, and m' is $2C\epsilon$ -close to m , which completes the proof. \square

Lemma 2.5. *There exist $R_0, \epsilon_0 > 0$ and $K > 1$ such that for any $\epsilon_0 > \epsilon > 0$ and $R > R_0$, the following holds: Suppose $u_1, u_2, v_1, v_2 \in \mathcal{G}$ that are ϵ -close to \mathbf{e} , $t_1, t_2 > R$, and $m_1, m_2 \in \mathcal{M}$. Then there exists $t \in \mathbb{R}$ and $m \in \mathcal{M}$ such that:*

1. $|t - (t_1 + t_2)| < K\epsilon$;
2. m is $K\epsilon$ -close to $m_1 m_2$;
3. $[a_{t_1} u_1 m_1 v_1 a_{t_2} u_2 m_2 v_2] = [a_t m]$.

Proof. Let R_0 be the constant R in the previous lemma, ϵ_1 be the constant in Lemma 2.1 (which is the same as in Lemma 2.4), C be the larger constant in Lemma 2.2 and Lemma 2.3, and $\epsilon_0 < \epsilon_1$ which will be determined later. Then as in the proof of Lemma 2.4, we have the decomposition $u_i = n_i^+ a_{\hat{t}_i} \hat{m}_i n_i^-$ for $i = 1, 2$. Then by Lemma 2.2 and Lemma 2.3, we have

$$\begin{aligned} [a_{t_1} u_1 m_1 v_1 a_{t_2} u_2 m_2 v_2] &= [a_{t_1} n_1^+ a_{\hat{t}_1} \hat{m}_1 n_1^- m_1 v_1 a_{t_2} n_2^+ a_{\hat{t}_2} \hat{m}_2 n_2^- m_2 v_2] \\ &= [\hat{n}_1^+ a_{t_1} a_{\hat{t}_1} \hat{m}_1 m_1 \hat{n}_1^- v_1 \hat{n}_2^+ a_{t_2} a_{\hat{t}_2} \hat{m}_2 m_2 \hat{n}_2^- v_2] \\ &= [a_{t_1 + \hat{t}_1} \hat{m}_1 m_1 u_3 a_{t_2 + \hat{t}_2} \hat{m}_2 m_2 u_4], \end{aligned}$$

where \hat{n}_i^+ 's are $Ce^{-R}\epsilon$ -close to \mathbf{e} , \hat{n}_i^- 's are $C\epsilon$ -close to \mathbf{e} , and

$$u_3 = \hat{n}_1^- v_1 \hat{n}_2^+, u_4 = \hat{n}_2^- v_2 \hat{n}_1^+$$

are $(1 + C + Ce^{-R})\epsilon$ -close to \mathbf{e} . Then we can repeat the above process to decompose u_3 as $u_3 = n_3^+ a_{\hat{t}_3} \hat{m}_3 n_3^-$, and we will have

$$[a_{t_1} u_1 m_1 v_1 a_{t_2} u_2 m_2 v_2] = [a_{t_1 + \hat{t}_1 + \hat{t}_3 + t_2 + \hat{t}_2} \hat{m}_1 m_1 \hat{m}_3 \hat{m}_2 m_2 \hat{n}_3^- u_4 \hat{n}_3^+],$$

where \hat{n}_3^\pm are $Ce^{-R}(1 + C + Ce^{-R})\epsilon$ -close to \mathbf{e} . Now let $t_5 = t_1 + \hat{t}_1 + \hat{t}_3 + t_2 + \hat{t}_2$, $m_5 = \hat{m}_1 m_1 \hat{m}_3 \hat{m}_2 m_2$ and $u_5 = \hat{n}_3^- u_4 \hat{n}_3^+$, and we know the distance between u_5 and \mathbf{e} is at most

$$(1 + C + Ce^{-R})\epsilon + 2Ce^{-R}(1 + C + Ce^{-R})\epsilon.$$

Thus we choose $\epsilon_0 < \epsilon_1$ such that

$$(1 + C + Ce^{-R})\epsilon_0 + 2Ce^{-R}(1 + C + Ce^{-R})\epsilon_0 = \epsilon_1,$$

and then we have u_5 is ϵ_1 -close to \mathbf{e} . By Lemma 2.4, we can find $t \in \mathbb{R}_+$, $m \in \mathcal{M}$ that are $2\epsilon_1$ -close to t_5, m_5 , respectively, such that $[a_{t_5}m_5u_5] = [a_tm]$. \square

2.2 Preliminary definitions

We want to introduce some important geometric definitions related with pants and surfaces in hyperbolic $2n$ -manifolds. Some are recalled from [KM12], [Ham15] and [Rao25].

2.2.1 Monodromy, good curves and normal bundles

Suppose M is a hyperbolic $2n$ -manifold and $\gamma \subset M$ is a closed geodesic. Then γ is the quotient of a geodesic in \mathbb{H}^{2n} under some isometry of \mathbb{H}^{2n} . Such an isometry is uniquely determined up to conjugacy by its translation length, which is the length of γ , and its rotation part, which is an element in $SO(2n - 1)$. The length of γ , denoted by $l(\gamma)$, is called ϵ -close to R , if

$$R - \epsilon < l(\gamma) < R + \epsilon,$$

for some $R, \epsilon > 0$.

The rotation part is called the *monodromy* of γ . The monodromy of γ is called ϵ -close to \mathbf{e} , if distance between the monodromy and \mathbf{e} is bounded by ϵ in $SO(2n - 1)$. We then say that a closed geodesic γ is (R, ϵ) -good if it has real length 2ϵ -close to $2R$ and monodromy ϵ -close to \mathbf{e} .

Remark 2.6. For $g \in SO(m)$, we can regard that g is ϵ -close to \mathbf{e} for some $\epsilon > 0$, if the angle between v and gv is no greater than ϵ for any unit vector v . For $g, g' \in SO(m)$, we say g is ϵ -close to g' , if $g^{-1}g'$ is ϵ -close to \mathbf{e} .

Suppose $\gamma: T \rightarrow M$ is a geodesic in M with unit speed and constant velocity, where T is a Riemannian 1-manifold. The unit normal bundle of γ is defined as

$$N^1(\gamma) = \{(u, v) : u \in U, v \in T_{\gamma(u)}M, \langle v, \gamma'(u) \rangle = 0, \|v\| = 1\}.$$

If γ is not closed, $N^1(\gamma)$ is isometric to $[0, l(\gamma)] \times S^{2n-2}$. Otherwise $N^1(\gamma)$ is the quotient of $\mathbb{R} \times S^{2n-2}$ by the translation along γ . In both cases, $N^1(\gamma)$

can be regarded as a metric space where the metric comes from the L^2 -metric on $\mathbb{R} \times S^{2n-2}$. Moreover, we define the normal frame bundle of γ as follows:

$$\mathcal{F}^\perp(\gamma) = \{(u, E) : u \in U, (\gamma'(u), E) \text{ is a positive-oriented frame of } T_u M\}.$$

2.2.2 Orthogeodesics, connections and feet

Suppose there are two oriented closed geodesics α_i , $i = 1, 2$, in M with unit speed and constant velocity. Suppose that $\eta: [0, l] \rightarrow M$ is a geodesic segment satisfying:

- η has unit speed and constant velocity;
- $\eta(0) \in \alpha_1, \eta(l) \in \alpha_2$;
- $\eta(0, l)$ meets with α 's orthogonally, $i = 1, 2$.

We then call η an *orthogeodesic* or a *connection* between α_1 and α_2 . In the case where α_1 and α_2 coincide, we will call η a *third connection*.

If α_1 and α_2 intersect, η_1 is the constant function; otherwise we let $\text{foot}_{\alpha_1}(\eta) = \eta'(0) \in N^1(\alpha_1)$ and $\text{foot}_{\alpha_2}(\eta) = -\eta'(l) \in N^1(\alpha_2)$ be the *feet* of η on α_0 and α_1 , respectively. The base point of a foot is called a *footpoint*.

Similarly, we can define the orthogeodesic and feet for two distinct oriented geodesics in \mathbb{H}^{2n} , where the orthogeodesic between them is always unique.

2.2.3 Pants and feet

Suppose Π_0 is a topological pair of pants considered as a manifold with boundary. We say a *pants* in M is an injective homomorphism $\rho: \pi_1(\Pi_0) \rightarrow \pi_1(M)$, up to conjugacy, which determines and is determined by a continuous map $f: \Pi \rightarrow M$, up to homotopy. So we can also call f or $f(\Pi_0)$ a pants. Let C_i 's be the boundary component of Π_0 , and we orient C_i such that Π_0 is on the left of C_i , for $i \in \mathbb{Z}/3\mathbb{Z}$. Then there is a unique oriented closed geodesic γ_i in M that is freely homotopic to $f(C_i)$. We can homotope f such that f maps C_i to γ_i , and we call such f a *nice pants*. From now on, when talking about pants in M , we will always assume that they are nice.

For any given simple non-separating arc α_i in Π_0 connecting C_{i-1} and C_{i+1} , we then can homotope f such that f maps α_i to an orthogeodesic η_i between γ_{i-1} and γ_{i+1} . Here η_i does not depend on the choice of α_i and f .

We call γ_i 's and η_i 's the *cuffs* and the *short orthogeodesic* of the pants $f(\Pi_0)$, respectively. The feet of η_{j-1} and η_{j+1} on γ_i are called the *short feet* of f on γ_i . When we talk about a cuff γ , the information of the associated pants is also carried by γ .

Moreover, for any simple arc β_i in Π_0 connecting C_i and itself such that C_{i-1} and C_{i+1} are in different component of $\Pi_0 - \beta_i$ (which always has two connected components), we can homotope f such that f sends β_i to an orthogeodesic ζ_i from γ_i to itself. ζ_i is independent of the choice of β_i and f , and we also call ζ_i is the third connection for γ_i in Π_0 . The two feet of ζ_i on γ_i are called the *long feet* of f on γ_i .

2.2.4 Tripods and framed tripods

We define a *tripod* in $T\mathbb{H}^{2n}$ or TM to be an ordered triple (v_0, v_1, v_2) of unit tangent vectors at the same base point x such that the three vectors mutually enclose an angle of $2\pi/3$. Then the vectors of a triple are contained in a real plane V in $T_x\mathbb{H}^{2n}$ or T_xM . The orientation of V is determined by the cyclic order of the triple.

A *framed tripod* in $T\mathbb{H}^{2n}$ or TM is a pair $((v_0, v_1, v_2), E)$ where (v_0, v_1, v_2) is a tripod contained in a plane V and E is a positive orthonormal frame in the orthogonal complement of V . We then can define how two framed tripods are related to each other.

The following definition is combined from Section 4 of [KM12] and Section 4 of [Ham15], which plays an essential role in the definition of good pants.

Definition 2.7. For $R > 0, \epsilon > 0$, we say two framed tripods $((v_0, v_1, v_2), E)$ at p and $((u_0, u_1, u_2), F)$ at q in TM are (R, ϵ) -*well-connected*, if there are geodesic segments $\gamma_i, i \in \mathbb{Z}/3\mathbb{Z}$, connecting p and q , such that:

- (1) The real length $l(\gamma_i)$ of γ_i satisfies that

$$R + \log \frac{4}{3} - \epsilon < l(\gamma_i) < R + \log \frac{4}{3} + \epsilon;$$

- (2) The angle between v_i and γ_i' at p is bounded by $e^{-R/4}$;
- (3) The angle between u_{-i} and $-\gamma_i'$ at q is bounded by $e^{-R/4}$;
- (4) Let $E_i = (v_i, v_i^\perp, E)$ be the positive frame at p determined by v_i and E , where v_i^\perp is orthogonal to v_i in the tangent plane determined by the

tripod, and the same for F_i at q . Let \hat{E}_i be the frame from the parallel transport of E_i along γ_i to q by negating the first two vectors. Then the element in $\text{SO}(2n)$ which transforms \hat{E}_i to F_i is ϵ -close to \mathbf{e} .

The geodesic segments γ_i are called *good connections* of the tripods, and the transformation defined in (4) is called the monodromy of γ_i .

We also want to single out another family of pairs of tripods, where the monodromy of each connection behaves more like an involution, rather than being close to \mathbf{e} .

Definition 2.8. For $R > 0, \epsilon > 0 > 0$, we say two framed tripods $((v_0, v_1, v_2), E)$ at p and $((u_0, u_1, u_2), F)$ at q in TM are (R, ϵ) -*badly-connected*, if there are geodesic segments $\gamma_i, i \in \mathbb{Z}/3\mathbb{Z}$, connecting p and q , such that items (1)–(3) in Definition 2.7 hold, and:

- (4') With \hat{E}_i and F_i defined as in (4) of Definition 2.7, the element in $\text{SO}(2n)$ which transforms \hat{E}_i to F_i is ϵ -close to $\mathbf{diag}\{1, -1, -1, 1, \dots, 1\}$.

We then say that the γ_i are *bad connections*.

2.2.5 Good and bad pants

Essentially, a pants group is (R, ϵ) -good (or bad) if there are (R, ϵ) -well-connected (or badly-connected) tripods that generate the group.

Definition 2.9. We say a pair of pants $f: \Pi_0 \rightarrow M$ is (R, ϵ) -*good*, if there are two (R, ϵ) -well-connected framed tripods in M with good connections γ_i , such that $\gamma_{i+1}\gamma_{i-1}^{-1}$ is freely homotopic to $f(C_i)$ for $i \in \mathbb{Z}/3\mathbb{Z}$. The (R, ϵ) -*bad pants* are defined the same way, but with bad connections. For convenience, we call a pants (R, ϵ) -*cuff-good*, if each cuff of this pants is an (R, ϵ) -good curve.

By Proposition 4.4 in [Ham15], we know that good pants have good boundary curves. Actually the same property holds for bad pants, which is not hard to verify, so we have the following proposition.

Proposition 2.10. *There exists a universal constant $\chi > 0$ such that any (R, ϵ) -good or bad pants is $(R, \chi\epsilon)$ -cuff-good.*

Remark 2.11. Although Hamenstädt ruled out the case of $\text{SO}(2n, 1)$ from the main theorem of [Ham15], those results before Section 7 in that paper also work well when the Lie group is some $\text{SO}(2n, 1)$ as indicated in Section 7 of that paper.

2.2.6 The doubling trick

This subsection follows Section 2.5 in [KW21]. One can also refer to Section 4.7 in [Rao25].

Suppose now that we have a family of pants $\mathcal{F} = \{f_i: P_i \rightarrow M\}$, where P_i 's are orientable. We take two copies of each $f_i: P_i \rightarrow M$ with opposite orientation and denote them by $f_i^+: P_i^+ \rightarrow M$ and $f_i^-: P_i^- \rightarrow M$. We then obtain a family of oriented pants $2\mathcal{F}$.

For each geodesic $\gamma \in M$, let $\mathcal{F}(\gamma)$ be the submultiset of $\cup_i \partial P_i$ consisting of those elements that are mapped to the unoriented γ in M . $\mathcal{F}(\gamma)$ could be empty if γ is not the image of any boundary component. We define $2\mathcal{F}(\gamma)$ in a similar way with $2\mathcal{F}$, by taking two copies of each $\alpha \in \mathcal{F}(\gamma)$ with opposite orientation, which is a submultiset of $(\cup_i \partial P_i^+) \cup (\cup_i \partial P_i^-)$. For $\alpha \in \mathcal{F}(\gamma)$, let $\alpha^+ \in 2\mathcal{F}(\gamma)$ that is mapped to γ and $\alpha^- \in 2\mathcal{F}(\gamma)$ that is mapped to γ^{-1} . Now suppose that there is a self-bijection σ_γ on $\mathcal{F}(\gamma)$. We then can define an involution τ_γ on $2\mathcal{F}(\gamma)$ by

$$\tau_\gamma(\alpha^+) = (\sigma_\gamma(\alpha))_-$$

and

$$\tau_\gamma(\alpha^-) = (\sigma_\gamma^{-1}(\alpha))_+.$$

By combining each involution for all γ , we get an involution τ on $(\cup_i \partial P_i^+) \cup (\cup_i \partial P_i^-)$. Then we identify α and $\tau(\alpha)$ for each $\alpha \in (\cup_i \partial P_i^+) \cup (\cup_i \partial P_i^-)$, and we will get a closed surface \mathcal{S} and a map $f: \mathcal{S} \rightarrow M$ by joining P_i^\pm 's and f_i^\pm 's via τ .

2.3 Steiner graph of pants

We first recall the result of Fermat point for hyperbolic triangles from Section 5 in [TX18]. Suppose $\triangle ABC$ is a hyperbolic triangle with three angles $\angle A, \angle B, \angle C$. When $\angle A, \angle B, \angle C < 2\pi/3$, then there is a unique Fermat point P inside the triangle such that $|PA| + |PB| + |PC|$ is minimal. When one of three angles is no less than $2\pi/3$, then the vertex corresponding to this angle is the unique Fermat point. In the former case,

$$\angle APB = \angle BPC = \angle CPA = 2\pi/3.$$

For $A, B \in M$, we let $\pi_1(M, A, B)$ denote the set of homotopy classes (rel endpoints) of paths in M that connect A and B . Then $\pi_1(M, A, A)$

can be identified with $\pi_1(M, A)$, and we can form $\alpha^{-1} \in \pi_1(M, B, A)$ and $\alpha\beta \in \pi_1(M, A, C)$ when $\alpha \in \pi_1(M, A, B)$ and $\beta \in \pi_1(M, B, C)$.

We can now be more precise with the definition of a pants Π in M . We can take a base point $*$ for our model pants Π_0 and draw three paths γ_i from the base point to itself, such that each γ_i is freely homotopic to a distinct cuff. Then the γ_i 's generate $\pi_1(\Pi_0)$, and $\gamma_0\gamma_1\gamma_2 = 1$. Given any $P \in M$, a pair of pants Π in M is determined by a representation $\rho: \pi_1(\Pi_0, *) \rightarrow \pi_1(M, P)$; we say ρ and $\rho': \pi_1(\Pi_0, *) \rightarrow \pi_1(M, P')$ determine the same pants if there is $\eta \in \pi_1(M, P, P')$ such that $\rho'(\gamma_i) = \eta^{-1}\rho(\gamma_i)\eta$ for each i .

Let $G(\Pi, M)$ denote the space of triples $(P, Q, (\eta_i))$, where $P, Q \in M$, each η_i is a geodesic segment from P to Q , and the representation from $\pi_1(\Pi_0, *)$ to $\pi_1(M, P)$ given by $\gamma_{i+1} \mapsto \eta_i\eta_{i+1}^{-1}$ determines Π . We define a function $l: G(\Pi, M) \rightarrow [0, \infty)$ by

$$l(P, Q, (\eta_i)) = |\eta_1| + |\eta_2| + |\eta_3|$$

of $P, Q \in M$. Since $G(\Pi, M)$ is locally compact and l is continuous and proper, the function l has a minimum at some $(P, Q, (\eta_i)) \in G(\Pi, M)$.

Theorem 2.12. *The function l has a unique local minimum (which is then the global minimum).*

Proof. Let us fix a base point $(P_*, Q_*, (\eta_{i,*}))$ in $G(\Pi, M)$, and let \tilde{M} be the universal cover of M . We can think of \tilde{M} as the space of all paths starting at P_* up to homotopy rel endpoints; we can do the same with P_* replaced by Q_* . Therefore we have

$$\tilde{M} \times \tilde{M} \cong \{(P, \alpha, Q, \beta) : P, Q \in M, \alpha \in \pi_1(M, P_*, P), \beta \in \pi_1(M, Q_*, Q)\},$$

We then can define the following map

$$\begin{aligned} \iota: \tilde{M} \times \tilde{M} &\rightarrow G(\Pi, M) \\ (P, \alpha, Q, \beta) &\mapsto (P, Q, ([\alpha^{-1}\eta_{i,*}\beta])), \end{aligned}$$

where $[\gamma]$ is the geodesic segment that is path-homotopic to γ . We claim that ι is a bijection.

If $(P, \alpha_1, Q, \beta_1)$ and $(P, \alpha_2, Q, \beta_2)$ have the same image, then

$$\alpha_1^{-1}\eta_{i,*}\beta_1 = \alpha_2^{-1}\eta_{i,*}\beta_2 \in \pi_1(M, P, Q),$$

for $i \in \mathbb{Z}/3\mathbb{Z}$. Thus $\eta_{i,*}^{-1}\alpha_2\alpha_1^{-1}\eta_{i,*}\beta_1\beta_2^{-1}$ is null-homotopic. Applying this for i and $i + 1$ and cancelling $\beta_1\beta_2^{-1}$, we obtain

$$\eta_{i,*}^{-1}\alpha_2\alpha_1^{-1}\eta_{i,*} = \eta_{i+1,*}^{-1}\alpha_2\alpha_1^{-1}\eta_{i+1,*},$$

as elements in $\pi_1(M, Q_*)$. We then know that $\alpha_2\alpha_1^{-1}$ commutes with $\eta_{i,*}\eta_{i+1,*}^{-1}$ up to homotopy, as elements in $\pi_1(M, P_*)$. Therefore $\alpha_2\alpha_1^{-1}$ lies in the centralizer of a pants group that determines Π , and is therefore trivial. Hence $\alpha_1 = \alpha_2$. Similarly, $\beta_1 = \beta_2$. Hence ι is injective.

On the other hand, for any $(P, Q, (\eta_i)) \in G(\Pi, M)$, the representation $\gamma_i \mapsto \eta_i\eta_{i+1}^{-1}$ determines the same pants Π as the representation $\gamma_i \mapsto \eta_{i,*}\eta_{i+1,*}^{-1}$ which is given by $(P_*, Q_*, (\eta_{i,*}))$. Thus there exists $\alpha \in \pi_1(M, P_*, P)$ such that

$$\eta_i\eta_{i+1}^{-1} = \alpha^{-1}\eta_{i,*}\eta_{i+1,*}^{-1}\alpha, \quad (2.13)$$

for $i \in \mathbb{Z}/3\mathbb{Z}$. Let $\beta = \eta_{i,*}^{-1}\alpha\eta_i \in \pi_1(M, Q_*, Q)$. Then by (2.13), we know β does not depend on the choice of i . Moreover, we have $\eta_i = \alpha^{-1}\eta_{i,*}\beta$ for each $i \in \mathbb{Z}/3\mathbb{Z}$, which implies $\iota(P, \alpha, Q, \beta) = (P, Q, (\eta_i))$. Hence ι is also surjective, and then the claim is proved.

We then can pull l back to $\tilde{M} \times \tilde{M}$, and the unique local minimum is implied from the observation that l is strictly convex (with respect to the product metric): this is because l is the sum of three distances, and distance function is strictly convex on $\tilde{M} \times \tilde{M}$ (except on the diagonal, where it is identically 0). \square

We observe that one of the η_i may have length 0 if P and Q are the same point, but this can hold for at most one η_i . The minimum length graph with vertices P, Q and connections η_i is called the *Steiner graph* of Π , and it is called *degenerate* if one of the η_i has length 0. We then prove the following claim.

Proposition 2.14. *There exists $R_0 > 0$ such that for all $R > R_0$ and $0 \leq \epsilon < 1$, the following holds: Suppose Π is an (R, ϵ) -cuff-good pants in a closed hyperbolic n -manifold M . Then the Steiner graph of Π is not degenerate. Moreover, the angle between any two connections is $2\pi/3$ at each vertex of the Steiner graph.*

Proof. Let P, Q be the two vertices and $\eta_i, i \in \mathbb{Z}/3\mathbb{Z}$ be the edges of the Steiner graph of Π . We let $\tilde{Q} \in \mathbb{H}^n$ be a lift of Q and $\tilde{\eta}_i$ be the lift of η_i such that $\tilde{\eta}_i$ connects \tilde{Q} and \tilde{P}_i , where P_i are three different lifts of P . Since

l reaches its minimal value at $(P, Q, (\eta_i))$, then by the above result, \tilde{Q} is the Fermat point of $\Delta\tilde{P}_1\tilde{P}_2\tilde{P}_3$.

If $(P, Q, (\eta_i))$ is degenerate, then \tilde{Q} coincides with some \tilde{P}_i , and

$$\begin{aligned} l(P, Q, (\eta_i)) &= \|\tilde{P}_i\tilde{P}_{i+1}\| + \|\tilde{P}_i\tilde{P}_{i+2}\| \geq l(\gamma_{i+2}) + l(\gamma_{i+1}) \\ &\geq 2(2R - 2\epsilon) = 4R - 4\epsilon. \end{aligned}$$

Here, $\|XY\|$ denotes the hyperbolic distance between X and Y for $X, Y \in \mathbb{H}^n$.

However, the function l can actually attain smaller values. Let α_i be the short orthogeodesic between γ_{i+1} and γ_{i+2} , then by hyperbolic trigonometry, there exists universal constant C such that

$$C^{-1}e^{-R/2} < l(\alpha_i) < Ce^{-R/2}.$$

Suppose $P' = \alpha_0 \cap \gamma_1$, and we let $\tilde{P}'_1, \tilde{P}'_2, \tilde{P}'_3$ be three lifts of P' such that the projection of $\tilde{P}'_i\tilde{P}'_{i+1}$ is freely homotopic to γ_{i+2} . We then let \tilde{Q}' be the Fermat point of the hyperbolic triangle $\Delta\tilde{P}'_1\tilde{P}'_2\tilde{P}'_3$, and Q' be the projection of \tilde{Q}' . Since $P' \in \gamma_1$, then the projection of $\tilde{P}'_2\tilde{P}'_0$ in M is γ_1 , so $\|\tilde{P}'_2\tilde{P}'_0\| = l(\gamma_1)$. By hyperbolic geometry, we know that there exists a universal constant $C > 0$ such that

$$\begin{aligned} 2R - 2\epsilon &\leq l(\gamma_2) \leq \|\tilde{P}'_0\tilde{P}'_1\| \leq 2R + C, \\ 2R - 2\epsilon &\leq l(\gamma_0) \leq \|\tilde{P}'_1\tilde{P}'_2\| \leq 3R + C. \end{aligned} \tag{2.15}$$

Therefore when R is large enough, each angle of $\Delta\tilde{P}'_1\tilde{P}'_2\tilde{P}'_3$ is less than $2\pi/3$. Hence, the Fermat point \tilde{Q}' is in the interior. Furthermore, there is a universal constant C' such that

$$\|\tilde{Q}'\tilde{P}'_0\| + \|\tilde{Q}'\tilde{P}'_1\| + \|\tilde{Q}'\tilde{P}'_2\| \leq \frac{1}{2} \left(\|\tilde{P}'_0\tilde{P}'_1\| + \|\tilde{P}'_1\tilde{P}'_2\| + \|\tilde{P}'_2\tilde{P}'_0\| \right) + C' \tag{2.16}$$

Thus when R is sufficiently large, by (2.15) and (2.16), we have

$$\begin{aligned} l(P', Q, (\eta'_i)) &= \|\tilde{Q}'\tilde{P}'_0\| + \|\tilde{Q}'\tilde{P}'_1\| + \|\tilde{Q}'\tilde{P}'_2\| \\ &\leq \frac{1}{2} ((2R + 2\epsilon) + (2R + C) + (3R + C)) + C' \\ &= 3.5R + \epsilon + C + C' < 4R - 4\epsilon, \end{aligned}$$

where η'_i is the projection of $\tilde{P}'_{i+1}\tilde{P}'_{i+2}$. That is contradictory to our assumption that $(P, Q, (\eta_i))$ was the Steiner graph. Therefore the Steiner graph of Π is not degenerate.




Figure 3: Lifts of P' in the universal cover

Furthermore, we know that \tilde{Q} , which is the Fermat point of $\Delta\tilde{P}_1\tilde{P}_2\tilde{P}_3$, is in the interior of $\Delta\tilde{P}_1\tilde{P}_2\tilde{P}_3$. Thus $\angle\tilde{P}_i\tilde{Q}\tilde{P}_{i+1} = 2\pi/3$, for $i \in \mathbb{Z}/3\mathbb{Z}$. Thus the angle between any two connections at Q is $2\pi/3$. We then know that the result holds at P as well by the symmetry of P and Q . \square

Remark 2.17. There are actually three (reduced) graphs with the homotopy type of a pants: the “theta”, which is what we are considering, the “barbell”, where there are segments from each point P, Q to itself, and a segment connecting them, and the “figure eight”, which can be thought of as either a degenerate theta or a degenerate barbell, and which therefore forms the boundary between them. One can then prove that there is a unique local minimum on the total space of graphs.

2.4 Good cuffs implies good or bad pants

In this section, we prove a surprising fact, that the converse of Proposition 2.10 is also true, up to an adjustment of constant.

For $A, B \in SO(2n-1)$, we say A is ϵ -close to B if AB^{-1} is ϵ -close to \mathbf{e} . Let $Z = \mathbf{diag}\{-1, 1, \dots, 1\} \in O(2n-1)$. The conjugacy by Z can be restricted to a self-map of $SO(2n-1)$, and we denote it by $\phi: SO(2n-1) \rightarrow SO(2n-1)$.

Let

$$\mathcal{Q} = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & X \end{pmatrix} : X \in SO(2n-2) \right\}$$

be a subgroup of $SO(2n-1)$, which is isomorphic to $SO(2n-2)$. Then the elements in \mathcal{Q} are fixed by ϕ .

Lemma 2.18. *There exists $\epsilon_0 > 0$ such that when $\epsilon_0 > \epsilon > 0$: Suppose $A \in SO(2n-1)$ is ϵ -close to $\phi(A)$, then there exists $Q \in \mathcal{Q}$, such that AQ is $\epsilon/2$ -close to either \mathbf{e} or $\mathbf{diag}\{-1, -1, 1, \dots, 1\}$.*

Proof. Let $\epsilon_0 = \pi/10$ and $F = (v_0, v_1, \dots, v_{2n-2})$ be a basis of \mathbb{R}^{2n-1} , where v_0 points towards the North Pole of S^{2n-2} . We first claim that for any $A \in SO(2n-1)$, we can always find a $Q \in \mathcal{Q}$ such that the left action of AQ is a rotation in a 2-dimensional subspace W containing v_0 , and the orthogonal complement W^\perp is pointwise-preserved by AQ . Actually, we consider the right action of A on F , and let

$$(v'_0, v'_1, \dots, v'_{2n-2}) = (v_0, v_1, \dots, v_{2n-2}) \cdot A,$$

and the claim is trivial when $v'_0 = v_0$. If not, let W be the subspace spanned by v_0 and v'_0 . Let P be the rotation in W that sends v_0 to v'_0 and preserves W^\perp , and let

$$(v'_0, v''_1, \dots, v''_{2n-2}) = (v_0, v_1, \dots, v_{2n-2}) \cdot P.$$

Since the right action of elements in \mathcal{Q} are those transformations preserving the first vector of the frame, so we can find $Q \in \mathcal{Q}$ such that

$$(v'_0, v''_1, \dots, v''_{2n-2}) = (v'_0, v'_1, \dots, v'_{2n-2}) \cdot Q.$$

Therefore

$$\begin{aligned} (v_0, v_1, \dots, v_{2n-2}) \cdot P &= (v'_0, v''_1, \dots, v''_{2n-2}) \\ &= (v'_0, v'_1, \dots, v'_{2n-2}) \cdot Q \\ &= ((v_0, v_1, \dots, v_{2n-2}) \cdot A) \cdot Q \\ &= (v_0, v_1, \dots, v_{2n-2}) \cdot (AQ), \end{aligned}$$

which proves the claim.

Now we only need to prove that the rotation angle of AQ is either close to 0 or π , since when the rotation angle is close to π , we can post-compose some $Q' \in \mathcal{Q}$ such that AQQ' is close to $\mathbf{diag}\{-1, -1, 1, \dots, 1\}$. We have

$$\phi(AQ) = ZAQZ = ZAZQ,$$

since Q commutes with Z . Thus AQ is ϵ -close to $\phi(AQ)$ by the bi-invariance of the metric on $SO(2n-1)$. On the other hand, the left action of Z is the reflection of the equator, and hence $\phi(X) = X^{-1}$ whenever the left action of $X \in SO(2n-1)$ is a rotation in a 2-dimensional subspace containing v_0 . So AQ is ϵ -close to $(AQ)^{-1}$, which means the rotation angle of AQ is $\epsilon/2$ -close to either 0 or π . \square

Let

$$R(\theta) = \begin{pmatrix} \cos \theta & \sin \theta & & & \\ -\sin \theta & \cos \theta & & & \\ & & 1 & & \\ & & & \ddots & \\ & & & & 1 \end{pmatrix} \in SO(2n), \text{ for } \theta \in [0, 2\pi],$$

then the right action of $R(\theta)$ on a $2n$ -frame is the rotation by θ in the plane spanned by the first two vectors, and Z can be viewed as the restriction of $R(\pi)$ on the space spanned by the last $2n-1$ vectors of the frame. For any $t \in \mathbb{R}$, let $G(t)$ be the right action of the frame flow along the first vector of the frame by distance t . We notice that $R(\pi)^2 = \text{Id}$, and $R(\pi)$ conjugates $G(t)$ to $G(-t)$, for any t .

The following lemma generalizes the inefficiency theory of hyperbolic geometry to dimension $2n$. We will skip the proof and one can refer to [KM15], [LM15], or [Rao25] for more details.

Lemma 2.19 (Inefficiency of Broken geodesic segments). *For any $0 < \epsilon < \pi$, there exist $C, L, m > 0$ such that for any $\theta \in (\epsilon - \pi, \pi - \epsilon)$: Suppose $t_1, t_2 > L$, then there exist $Y_1, Y_2 \in SO(2n)$ that are Ce^{-L} -close to Id , and $t_1 + t_2 > t > t_1 + t_2 - m$ such that*

$$G(t_1)R(\theta)G(t_2) = Y_1G(t)Y_2.$$

The next theorem is the reverse statement of Proposition 2.10. It shows that the length and monodromy of cuffs can provide enough information to tell whether the pants is good or not. We can then consider the set of all (R, ϵ) -cuff-good (and good) pants and use our estimates in Section 3 to prove in Section 4 that we can match these pants to form an incompressible surface.

Theorem 2.20. *There is a universal constant C such that if Π is a pants group that is (R, ϵ) -cuff-good, then Π is either $(R, C\epsilon)$ -good or $(R, C\epsilon)$ -bad.*

Proof. Suppose the three cuffs of an oriented pants Π in M are γ_0, γ_1 and γ_2 that are (R, ϵ) -good. We proceed by the following steps:

1. **Use the Steiner graph of Π to get two (triply) connected tripods.**

Let G be the Steiner graph of Π with vertices p, q and edges η_0, η_1, η_2 connecting p and q , such that $\eta_{i+1}\eta_{i-1}^{-1}$ is freely homotopic to γ_i , for $i \in \mathbb{Z}/3\mathbb{Z}$, and we denote by l_i the length of η_i . Without loss of generality, we assume for each $i \in \mathbb{Z}/3\mathbb{Z}$, η_i is parameterized by arc length and oriented from p to q . By Prop 2.14, the angle between η'_i and η'_j is $2\pi/3$ at p if $i \neq j \in \mathbb{Z}/3\mathbb{Z}$, and the same is also true for $-\eta'_i$'s at q . Thus $P = (p, \eta'_0, \eta'_1, \eta'_2)$ and $Q = (q, -\eta'_0, -\eta'_{-1}, -\eta'_{-2})$ are two tripods generating Π . Then we only need to find frames at p and q such that the monodromy along each η_i is $C\epsilon$ -close to \mathbf{e} , for some constant C .

2. **Choose frames for the orthogonal plane at each tripod, to get complete frames for \mathbb{R}^{2n} .**

We first choose arbitrary positive frames E and F at p and q in the orthogonal complement of the real plane α_p and α_q , respectively. Then $P_0 = (P, E)$ and $Q_0 = (Q, F)$ are two framed tripods that generate Π . For each $i \in \mathbb{Z}/3\mathbb{Z}$, there are vectors u_i at p and v_i at q such that (η'_i, u_i) and $(-\eta'_i, v_i)$ are positive frames of α_p and α_q , respectively. Let $E_i = (\eta'_i, u_i, E)$ and $F_i = (-\eta'_i, v_i, F)$ be the frames at p and q . Then $R(2\pi/3)$ sends E_i and F_i to E_{i+1} and F_{i-1} , respectively.

3. **For each connection, push these two frames to the midpoint of the connection and obtain a transformation X_i between the two new frames. Then write the monodromy of each cuff as a word in these X_i .**

Let x_i be the midpoint of η_i , for $i \in \mathbb{Z}/3\mathbb{Z}$. We then push forward E_i and F_i to x_i by parallel transport, and the frames we get at x_i are denoted by $E_i(x_i)$ and $F_i(x_i)$. Then there is a unique $X_i \in SO(2n)$ such that the right action of $R(\pi)X_i$ sends $E_i(x_i)$ to $F_i(x_i)$. We notice that the right action of $R(\pi)$ on $E_i(x_i)$ will send η'_i to $-\eta'_i$, therefore X_i preserves $-\eta'_i$.

Now we consider the conjugacy class of the element in \mathcal{G} of the broken geodesic $\eta_{i+1}\eta_{i+2}^{-1}$, which can be expressed as composition of right

actions on \mathcal{FM} as follows, up to conjugacy:

$$\begin{aligned}
& [R(\pi)X_{i+1}G\left(-\frac{l_{i+1}}{2}\right)R\left(\frac{-2\pi}{3}\right)G\left(\frac{l_{i+2}}{2}\right)X_{i+2}^{-1}R(\pi) \\
& G\left(-\frac{l_{i+2}}{2}\right)R\left(\frac{-2\pi}{3}\right)G\left(\frac{l_{i+1}}{2}\right)] \\
& = [R(\pi)X_{i+1}R(\pi)G\left(\frac{l_{i+1}}{2}\right)R(\pi)R\left(\frac{-2\pi}{3}\right) \\
& G\left(\frac{l_{i+2}}{2}\right)X_{i+2}^{-1}G\left(\frac{l_{i+2}}{2}\right)R(\pi)R\left(\frac{-2\pi}{3}\right)G\left(\frac{l_{i+1}}{2}\right)] \\
& = [R(\pi)X_{i+1}R(\pi)G\left(\frac{l_{i+1}}{2}\right)R\left(\frac{\pi}{3}\right) \\
& G\left(\frac{l_{i+2}}{2}\right)X_{i+2}^{-1}G\left(\frac{l_{i+2}}{2}\right)R\left(\frac{\pi}{3}\right)G\left(\frac{l_{i+1}}{2}\right)]
\end{aligned}$$

When R is large enough, by Lemma 2.19, we can find $Y_i^\pm \in \text{SO}(2n)$ and $l_i^\pm \in \mathbb{R}_+$, $i \in \mathbb{Z}/3\mathbb{Z}$ such that

$$\begin{aligned}
G\left(\frac{l_{i+1}}{2}\right)R\left(\frac{\pi}{3}\right)G\left(\frac{l_{i+2}}{2}\right) &= Y_{i+1}^+G(l_i^+)Y_{i+2}^+, \\
G\left(\frac{l_{i+2}}{2}\right)R\left(\frac{\pi}{3}\right)G\left(\frac{l_{i+1}}{2}\right) &= Y_{i+2}^-G(l_i^-)Y_{i+1}^-.
\end{aligned}$$

Then the above conjugacy class can be written as

$$\begin{aligned}
& [R(\pi)X_{i+1}R(\pi)Y_{i+1}^+G(l_i^+)Y_{i+2}^+X_{i+2}^{-1}Y_{i+2}^-G(l_i^-)Y_{i+1}^-] \\
& = [G(l_i^+)Y_{i+2}^+X_{i+2}^{-1}Y_{i+2}^-G(l_i^-)Y_{i+1}^-R(\pi)X_{i+1}R(\pi)Y_{i+1}^+].
\end{aligned}$$

By the definition of X_i and $R(\pi)$, we know that X_i and $R(\pi)X_iR(\pi)$ both preserve the first vector of frames. Therefore by Lemma 2.4, the length of γ_i can be approximated by $l_i^+ + l_i^-$, and the monodromy of γ_i can be estimated by $X_{i+2}^{-1}R(\pi)X_{i+1}R(\pi)$, which also preserves the tangent vector of γ_i , up to conjugacy, where the error terms are bounded by e^{-R} up to a universal constant.

4. **These transformations are right actions acting on the frames, but we can write the right actions as matrices and apply Lemma 2.18.**

Therefore $X_{i+2}^{-1}R(\pi)X_{i+1}R(\pi) = X_{i+2}^{-1}\phi(X_{i+1})$, which we can think of as an element in $\text{SO}(2n-1)$, is ϵ -close to \mathbf{e} , since γ_i is (R, ϵ) -good, $i \in \mathbb{Z}/3\mathbb{Z}$. Hence X_{i+2} is ϵ -close to $\phi(X_{i+1})$, $i \in \mathbb{Z}/3\mathbb{Z}$, and we then know X_i is 3ϵ -close to $\phi(X_i)$, and X_i is 2ϵ -close to X_j , for $i \in \mathbb{Z}/3\mathbb{Z}$ and $j \neq i$. When ϵ is small enough, by Lemma 2.18, we can find $Q \in \mathcal{Q}$ such that X_0Q is $3\epsilon/2$ -close to either \mathbf{e} or $\mathbf{diag}\{-1, -1, 1\}$. In the former case, we have X_iQ is $7\epsilon/2$ -close to \mathbf{e} , for all $i \in \mathbb{Z}/3\mathbb{Z}$. Since the right action of Q commutes with the frame flow and the rotations $R(\theta)$, we can replace X_i with X_iQ through replacing F with $F \cdot Q$. We then have a pair of well-connected framed tripods.

For the latter case, we can find a pair of badly-connected framed tripods for a similar reason. \square

3 Counting third connections

Now, suppose that γ_0 is a given (R, ϵ) -good curve. We want to find all third connections η such that the pants generated by each η is an (R, ϵ) -cuff-good pants, but also a *good* pants (i.e. not a *bad* one). More specifically, we want to know the distribution of the (long) feet of these third connections, and roughly what measure we get on the average of a pair of feet.

Given two unit normal vectors on γ_0 as two feet of a predicted third connection, we can estimate the volume of the space of monodromies for this third connection for which the two other cuffs will be good. This volume is zero unless the two unit normal vectors are close (when compared in either direction) and basically just depends on how close they are. We will make this much more explicit in this section.

In this way we obtain an effective estimate of the measure on the feet of the good third connections (first on the long feet, and then on their midpoint). With this, we will prove in the future that, for each good curve γ_0 , we can match the pants around γ_0 such that the feet for matching pants satisfy the usual condition of being on opposite sides of γ_0 (nearly antipodal points of the sphere) along with a shear by 1. We can then apply Hamenstädt's sufficient condition to conclude that the resulting surface is π_1 -injective.

3.1 Length of third connections

We start with a fundamental result in hyperbolic geometry, which is actually a special case of Lemma 2.19. This lemma helps us estimate the length of a third connection in terms of the length of cuffs in good pants. The same estimate in the two and three dimensional cases were used in [KM15, Section 3.3] and [KW21, Section 3.3], respectively.

Lemma 3.1. *There exists constant $C, L_0 > 0$ such that for all $L > L_0$, the following holds: Suppose γ^+, γ^- are two oriented geodesic segments in M which meet orthogonally at both endpoints satisfying that $\gamma^+\gamma^-$ is an oriented broken geodesic. If $l(\gamma^i) > L$, $i = +, -$, then*

$$|l(\gamma) - l(\gamma^+) - l(\gamma^-) + 2 \ln 2| < Ce^{-L},$$

where γ is the closed geodesic homotopic to $\gamma^+\gamma^-$.

Remark 3.2. The monodromy of geodesics in hyperbolic 3-manifolds can be regarded as the imaginary part of the complex length, which cannot be defined properly in higher-dimensional cases. Therefore we need to study the length and the monodromy of third connections separately.

Let η be a third connection for γ_0 of an (R, ϵ) -good pants, and suppose η subdivides γ_0 into two arcs α_1 and α_2 . We let u_i be the length of α_i for $i = 1, 2$, and w be the length of η . Then $u_1 + u_2 = l(\gamma_0)$. Let $L_{R,\epsilon}(\gamma_0)$ be the set of all pairs (x, y) such that

$$|x + y - 2 \ln 2 - 2R| < 2\epsilon$$

and

$$|(l(\gamma_0) - x) + y - 2 \ln 2 - 2R| < 2\epsilon.$$

Then the square region $L_{R,\epsilon}(\gamma_0) \subset \mathbb{R}^2$ can be used to approximate the set of (u_1, w) for all third connections of good pants, up to an exponentially small error term. Suppose that the measure on $\mathbb{R} \times \mathbb{R}$ is given by $e^{2y} dx dy$, then the area of $L_{R,\epsilon}(\gamma_0)$ is

$$\text{Vol}(L_{R,\epsilon}(\gamma_0)) = 8e^{4R-l(\gamma_0)}(e^{2\epsilon} - e^{-2\epsilon})^2 \sim 128\epsilon^2 e^{4R-l(\gamma_0)}, \quad (3.3)$$

as $\epsilon \rightarrow 0$.

3.2 Monodromy estimate and feet distance

In this subsection, we first compute the monodromies of the other two cuffs γ_1, γ_2 from the frames at footpoints of η , and then determine the good region of monodromies for the third connection that generates cuff-good pants in terms of the distance between two feet.

Let η be a third connection for γ_0 , where the endpoints of η are roughly R apart and the length of η is about R . We want to determine when the remaining two cuffs of the pants determined by γ_0 and η are (R, ϵ) -good curves. Let A, B be the points where γ_0 meets η such that η is oriented from A to B , and let S_i be the unit normal sphere of γ_0 at i for $i = A, B$. Let γ_0 be divided by η into α_1, α_2 such that $\gamma_0 = \alpha_1 \alpha_2^{-1}$, where α_1 and α_2 are both oriented from A to B . Let γ_1 be the oriented cuff homotopic to $\eta \alpha_1^{-1}$ and γ_2 be the oriented cuff homotopic to $\eta^{-1} \alpha_2$. Let y be the midpoint of η , and x_i be the midpoint of α_i for $i = 1, 2$.

We now want to set up a system of frames that will allow us to write the monodromies for the γ_i as a product of right actions. At point A , the unit tangent vectors $(\gamma'_0)_A$ of γ_0 and η'_A of η form a 2-plane P_A in the tangent space $T_A M$, and we temporarily choose an arbitrary frame E of the orthogonal complementary subspace of P_A in $T_A M$. Let $E_0 = (\eta'_A, -(\gamma'_0)_A, E)$, $E_1 = ((\gamma'_0)_A, \eta'_A, E)$ and $E_2 = (-(\gamma'_0)_A, -\eta'_A, E)$ be three frames at A . Similarly, we choose an arbitrary frame F of the orthogonal complementary subspace of P_B in $T_B M$, where P_B is the 2-plane spanned by $(\gamma'_0)_B$ and η'_B . Then we have three frames $F_0 = (-\eta'_B, -(\gamma'_0)_B, F)$, $F_1 = (-(\gamma'_0)_B, \eta'_B, F)$ and $F_2 = ((\gamma'_0)_B, -\eta'_B, F)$ at B .

We apply the frame flow to E_0 and F_0 until they arrive at y , and denote the resulting frames by $E_0(y)$ and $F_0(y)$. There then exists a unique $Y \in \text{SO}(2n)$ such that the right action by $R(\pi)Y$ sends $E_0(y)$ to $F_0(y)$. Similarly, we apply the frame flow to E_i, F_i , for $i = 1, 2$, until they arrive at x_i ; we again have $X_i \in \text{SO}(2n)$ such that $E_i(x_i) \cdot R(\pi)X_i = F_i(x_i)$. From Y and the X_i , we can compute the conjugacy class of the element in \mathcal{G} for each γ_i , $i \in \mathbb{Z}/3\mathbb{Z}$; it can be expressed as a composition of right actions as in the proof of Theorem 2.20. Then as in that proof, the monodromies for γ_0, γ_1 and γ_2 are given, up to conjugacy and an e^{-R} error term, by $X_2^{-1} \phi(X_1)$, $X_1^{-1} \phi(Y)$ and $Y^{-1} \phi(X_2)$, where $Y^{-1} \phi(X_2)$ is conjugate to $(X_2^{-1} \phi(Y))^{-1}$; from here we will consider the X_i and Y as elements in $\text{SO}(2n - 1)$ since they all preserve the first vector in the frame. This means (ignoring for the moment the very small error term), the resulting pants is cuff-good if and only if

the latter two transformations are ϵ -close to \mathbf{e} . So our goal becomes to **estimate the volume of the intersection of ϵ -neighborhoods of X_1 and X_2 (essentially determined by the distance between X_1 and X_2 in $\text{SO}(2n-1)$) whenever $X_2^{-1}\phi(X_1)$ is ϵ -close to \mathbf{e} .**

Let Λ be the monodromy of γ_0 . We define the spherical distance (or the angle) between η'_A and $-\eta'_B$ along α_1 , denoted by $d_{S^{2n-2}}^1(\eta'_A, -\eta'_B)$, as the angle between these two unit normal vectors when we bring them to x_1 by parallel transport. If we bring them to x_2 , the distance will equal the distance between $\Lambda^{-1}(\eta'_A)$ and $-\eta'_B$ along α_1 .

Proposition 3.4. *For any $\pi/30 > \epsilon > 0$, there exists $R_0 > 0$ such that when $R > R_0$, the following holds: Suppose η is a third connection for γ_0 such that the generated pants is (R, ϵ) -cuff-good and good. Then $d_{S^{2n-2}}^1(\eta'_A, -\eta'_B) < 3\epsilon/2$.*

Proof. The idea of proving this statement is similar to Part 4 of the proof of Theorem 2.20. Suppose X_1, X_2 and Y are defined as above; then we know that $X_i^{-1}\phi(Y)$ is ϵ -close to \mathbf{e} , and $X_2^{-1}\phi(X_1)$ is conjugate to Λ which is also ϵ -close to \mathbf{e} . Thus X_1 is 2ϵ -close to X_2 , and X_2 is ϵ -close to $\phi(X_1)$; hence X_1 is 3ϵ -close to $\phi(X_1)$. Then by Lemma 2.18, there exists $Q \in \mathcal{Q}$ such that X_1Q is $3\epsilon/2$ -close to \mathbf{e} or $\mathbf{diag}\{-1, -1, 1, \dots, 1\}$. Since the pants generated by γ_0 and η is good, so we can rule out the second possibility, which means X_1Q is $3\epsilon/2$ -close to \mathbf{e} . We can assume that X_1 is $3\epsilon/2$ -close to \mathbf{e} by replacing F with $F \cdot Q^{-1}$; here Q is considered as an element in $\text{SO}(2n-2)$. Therefore $\phi(X_1)$ is also $3\epsilon/2$ -close to \mathbf{e} , since $\phi(X_1)$ is conjugate to X_1 in $\text{O}(2n)$.

On the other hand, we notice that the second vector of $E_1(x_1)$ is the parallel transport of η'_A to x_1 , and the second vector of $F_1(x_1) \cdot R(\pi)$ is the parallel transport of $-\eta'_B$ to x_1 . Therefore by $\phi(X_1)$ sending $E_1(x_1)$ to $F_1(x_1) \cdot R(\pi)$ as $2n$ -frames and $d(\phi(X_1), \mathbf{e}) < 3\epsilon/2$, we have

$$d_{S^{2n-2}}^1(\eta'_A, -\eta'_B) < 3\epsilon/2. \quad \square$$

On the other hand, we want to show that when the two feet are sufficiently close, i.e. $d_{S^{2n-2}}^1(\eta'_A, -\eta'_B)$ is sufficiently small, the distance between X_1 and X_2 is bounded by $3\epsilon/2$; this will ensure that we have enough third connections in the good region.

Proposition 3.5. *For any $\pi/30 > \epsilon > 0$, there exists $R_0 > 0$ such that when $R > R_0$, the following holds: Suppose $\eta'_A, -\eta'_B$ are unit normal vectors at A*

and B respectively (where we keep the notation even though we are not given η) such that $d_{S^{2n-2}}^1(\eta'_A, -\eta'_B) < \epsilon/4$. Then

$$d(X_1, X_2) < \epsilon + 2d_{S^{2n-2}}^1(\eta'_A, -\eta'_B) < 3\epsilon/2.$$

Remark 3.6. The change of the frames E or F will not affect the distance between X_1 and X_2 ; therefore, $d(X_1, X_2)$ is well-defined without choosing E and F and can be calculated from any choice of E and F .

Proof. Let $\delta = d_{S^{2n-2}}^1(\eta'_A, -\eta'_B) < \epsilon/4$. After bringing η'_A and $-\eta'_B$ to x_1 , let P be a 2-plane in the normal space at x_1 to γ_0 that contains η'_A and $-\eta'_B$. Then we can choose frames E at A and F at B such that X_1 is the rotation in P that maps η'_A to $-\eta'_B$ and has rotation angle δ . So $d(X_1, \mathbf{e}) = \delta$, and hence $d(\phi(X_1), \mathbf{e}) = \delta$. Then by $d(X_2^{-1}\phi(X_1), \mathbf{e}) < \epsilon$, we know that

$$d(X_2, \mathbf{e}) \leq d(X_2, \phi(X_1)) + d(\phi(X_1), \mathbf{e}) = d(X_2^{-1}\phi(X_1), \mathbf{e}) + d(\phi(X_1), \mathbf{e}) < \epsilon + \delta.$$

Thus we have

$$d(X_1, X_2) \leq d(X_1, \mathbf{e}) + d(X_2, \mathbf{e}) < \epsilon + 2\delta < 3\epsilon/2. \quad \square$$

3.3 The average feet of good pants

For a pants P in a hyperbolic 2 or 3-manifold, a short foot on one cuff γ is always a midpoint of the long feet on γ in the unit normal bundle of γ , and P is divided into two isometric right-angled hexagons. But the same properties do not hold in hyperbolic manifolds of dimension greater than 3. In this section, we will define the average of long feet for good pants in M and estimate the difference between the new defined feet and the short feet in $N^1(\gamma)$.

Suppose that Π is an oriented (R, ϵ) -cuff-good and good pants with a boundary curve γ_0 , for $\epsilon > 0$ sufficiently small such that it satisfies all the above propositions, and R sufficiently large which will be determined later. Let η be the third connection for γ_0 with two long feet $u_1, u_2 \in N^1(\gamma_0)$, and n_1, n_2 be the two short feet on γ_0 such that u_i, n_i, u_{i+1} are ordered along the orientation of γ_0 for $i \in \mathbb{Z}/2\mathbb{Z}$.

Let us now define the *average feet* a_1, a_2 on γ_0 of Π . There are two midpoints b_1, b_2 of the base points for u_1 and u_2 , and we label them so that u_i, b_i, u_{i+1} are ordered along the orientation of γ_0 (as with the n_i). Fixing i ,

we parallel transport u_1 and u_2 to b_i (transporting each about one-quarter the length of γ). By Proposition 3.4, we know that the resulting unit normal vectors are $3\epsilon/2$ -close in the unit normal sphere at b_i . We then let a_i be the midpoint of the short geodesic (in this unit sphere at b_i) connecting them.

Then we have a quick result for average feet.

Proposition 3.7. *When R is sufficiently large, a_1 is 3ϵ -close to $\Lambda^{1/2}a_2$, where Λ is the monodromy of γ_0 .*

Proof. This directly follows from Proposition 3.4. □

Before stating the theorem, we state a simple result in hyperbolic geometry, and its proof is left to the reader.

Lemma 3.8. *There exists a universal constant $K_0, d_0 > 0$ such that when $d > d_0$ the following holds: Suppose γ_1, γ_2 are two geodesics in \mathbb{H}^n such that the real distance between γ_1 and γ_2 is d . For any two points $X_1, X_2 \in \gamma_2$, let $n_i \in N^1(\gamma_1)$ be the foot of the perpendicular geodesic from X_i to γ_1 , $i = 1, 2$. Then n_1 and n_2 are within distance $K_0 e^{-d}$ in $N^1(\gamma_1)$.*

Now we show that the average feet are close to the short feet for good pants.

Proposition 3.9. *There exists a universal constant $K > 0$ such that when R is large enough, a_i is Ke^{-R} -close to n_i , $i \in \mathbb{Z}/2\mathbb{Z}$.*

Proof. Let γ_1, γ_2 be the other two cuffs of the pants Π and η_i be the short orthogeodesic connecting γ_0 and γ_i for $i = 1, 2$, and n_i is one foot of η_i . Let β be the third connection for γ_0 with two long feet u_1, u_2 . We then focus on the lifts of the cuffs and feet to \mathbb{H}^{2n} and $T\mathbb{H}^{2n}$, and want to estimate the difference of lifts of a_i and n_i .

Let $\tilde{\gamma}_0$ be a lift of γ , and $\tilde{u}_1, \tilde{a}_1, \tilde{n}_1, \tilde{u}_2$ be the lifts of u_1, a_1, n_1, u_2 respectively such that \tilde{a}_1 and \tilde{n}_1 are the only lifts of a_1 and n_1 , respectively, between \tilde{u}_1 and \tilde{u}_2 . Without loss of generality, we then only need to show that \tilde{a}_1 is Ke^{-R} -close to \tilde{n}_1 for some constant $K > 0$.

Let $\tilde{\beta}$ be the lift of β such that \tilde{u}_1 is the foot for $\tilde{\beta}$, and $\tilde{\gamma}'_0$ be another lift of γ such that $\tilde{\beta}$ connects $\tilde{\gamma}_0$ and $\tilde{\gamma}'_0$. Let $\tilde{\eta}_1$ and $\tilde{\gamma}_1$ be the lifts of η_1 and γ_1 such that \tilde{n}_1 is the foot for $\tilde{\eta}_1$ at $\tilde{\gamma}_0 \cap \tilde{\eta}_1$ and $\tilde{\eta}_1$ is the orthogeodesic between $\tilde{\gamma}_0$ and $\tilde{\gamma}_1$. We then let $\tilde{\eta}'_1$ be the lift of η connecting $\tilde{\gamma}_1$ and $\tilde{\gamma}'_0$, and $Q = \tilde{\eta}'_1 \cap \tilde{\gamma}'_0$ and $Q_1 = \tilde{\eta}'_1 \cap \tilde{\gamma}_1$. We denote by $\partial\mathbb{H}^{2n}$ the boundary at infinity of

\mathbb{H}^{2n} , and let X_1, X_2 be the endpoints of $\tilde{\gamma}_1$ in $\partial\mathbb{H}^{2n}$ such that Q_1 is between X_1 and $\tilde{\gamma}_1 \cap \tilde{\eta}_1$. We drop the perpendicular geodesics from Q, Q_1, X_1 and X_2 to $\tilde{\gamma}_0$ with foot q, q_1, x_1 and x_2 on $\tilde{\gamma}_0$, respectively. Then we know that \tilde{n}_1 is the midpoint of x_1 and x_2 in $N^1(\tilde{\gamma}_0)$. On the other hand, we know that \tilde{a}_1 is the midpoint of \tilde{u}_1 and the foot of another lift of β on $\tilde{\gamma}_0$. Hence it is sufficient to prove that \tilde{u}_1 is $O(e^{-R})$ -close to x_1 .




Figure 4: Lifts of cuffs and feet in \mathbb{H}^{2n}

We let $R > d_0$ be large enough, where d_0 is from Lemma 3.8. Since γ_1 is a good curve, we have

$$2R - 2\epsilon < d_{\mathbb{H}^{2n}}(Q_1, \tilde{\eta}_1) = l(\gamma_1) < 2R + 2\epsilon.$$

We note that the distance between $\tilde{\gamma}_0$ and $\tilde{\gamma}_1$, which is the length of η_1 , is bounded by a constant multiple of $e^{-R/2}$. Therefore by elementary hyperbolic geometry, $d_{\mathbb{H}^{2n}}(Q_1, \tilde{\gamma}_0) > R$ and q_1 is e^{-R} -close to x_1 . We notice that

$d_{\mathbb{H}^{2n}}(Q_1, Q) = l(\tilde{\eta}'_1)$ is also equal to the length of η_1 , which is exponentially small. So by elementary hyperbolic geometry again and $d_{\mathbb{H}^{2n}}(Q_1, \tilde{\gamma}_0) > R$, we know that q is e^{-R} -close to q_1 . Finally, by Lemma 3.8, \tilde{u}_1 is $K_0 e^{-R}$ -close to q . Hence \tilde{u}_1 is $(K_0 + 2)e^{-R}$ -close to x_1 . \square

One may notice that the average feet can actually be defined without the given orientations of Π and γ , and the above results all holds as well. Here the orientations are provided just for convenience.

3.4 The volume of the intersection and the measure of feet of good pants

In this final subsection (of Section 3) we show that the volume of intersection is bounded above and below, for all pairs of long feet that are close (after being brought together by parallel transport). This is our fundamental estimate, which we will use to prove the matching result in Section 4.2.

3.4.1 Estimated foot measures

For each oriented closed geodesic γ_0 , a framed oriented third connection η can be described as a point $((u, (f_u, E)), (v, (f_v, F)), l, \Lambda)$ in

$$\mathbb{F}(\gamma_0) \equiv \mathcal{F}^\perp(\gamma_0) \times \mathcal{F}^\perp(\gamma_0) \times [0, \infty) \times \text{SO}(2n - 1),$$

where u, v are two footpoints of η , f_u is the tangent vector of η at u , f_v is the tangent vector of η at v , E, F are the corresponding orthogonal complement in the normal spaces, l is the length of η , and Λ is the monodromy along η (which lies in $\text{SO}(2n - 1)$ because it preserves the tangent vector to η). We then have an action of $\text{SO}(2n - 2)$ on each $\mathcal{F}^\perp(\gamma_0)$ factor of $\mathbb{F}(\gamma_0)$ as the rotation around the first vector of frame (with the resulting left or right multiplication on $\text{SO}(2n - 1)$), and therefore we have an action of $\text{SO}(2n - 2) \times \text{SO}(2n - 2)$ on $\mathbb{F}(\gamma_0)$. Let

$$\mathbb{I}(\gamma_0) = \mathbb{F}(\gamma_0) / \text{SO}(2n - 2) \times \text{SO}(2n - 2),$$

which is a $\text{SO}(2n - 1)$ -fiber bundle over $N^1(\gamma_0) \times N^1(\gamma_0) \times [0, \infty)$; then $\mathbb{I}(\gamma_0)$ is the space of third connections for γ_0 . We take the Riemannian metric on $\mathbb{F}(\gamma_0)$ to be the L^2 -metric over each coordinate and push it down to $\mathbb{I}(\gamma_0)$. Locally, we let the measure on $\mathbb{I}(\gamma_0)$ be the product of the Lebesgue (or

Haar) measures on $N^1(\gamma_0)$, $N^1(\gamma_0)$, and $\text{SO}(2n-1)$, and multiplied by $e^{2t} dt$ on $[0, \infty)$.

Let $\Pi_{R,\epsilon}(\gamma_0)$ be the set of pairs (P, η) where P is good and (R, ϵ) -cuff-good in M of which γ_0 is a boundary geodesic and η is an oriented third connection of P for γ_0 , and define $\pi_0: \Pi(\gamma_0) \rightarrow \mathbb{I}(\gamma_0)$ by $\pi_0(P, \eta) = Q(\eta)$, where $Q(\eta)$ is the image of η in $\mathbb{F}(\gamma_0)$, pushed down to $\mathbb{I}(\gamma_0)$. Then $\Pi_{R,\epsilon}(\gamma_0)$ is a finite set and the counting measure on $\Pi_{R,\epsilon}(\gamma_0)$ generates a point measure $\tilde{\nu}$ on $\mathbb{I}(\gamma_0)$.

Now we want to define two *good* regions to estimate $\tilde{\nu}$. Let $\mathcal{R}(\gamma_0) \subset \mathbb{I}(\gamma_0)$ be the region such that (P, η) *would* be good and (R, ϵ) cuff-good if $Q(\eta) \in \mathcal{R}(\gamma_0) \subset \mathbb{I}(\gamma_0)$. More formally, we let T_{γ_0} be the translation (defined up to conjugacy) in $\text{Isom}(\mathbb{H}^{2n})$ such that $\mathbb{H}^{2n}/T_{\gamma_0}$ is the cover of M associated to γ_0 . For any $\alpha \in \mathbb{I}(\gamma_0)$, there is a unique pants group $P(\alpha)$ (which has T_{γ_0} as one generator) such that $\mathbb{H}^{2n}/P(\alpha)$ has a third connection η (from γ_0 to itself), such that $Q(\eta) = \alpha$. We then say that $\alpha \in \mathcal{R}(\gamma_0) \subset \mathbb{I}(\gamma_0)$ if and only if $P(\alpha)$ is good and (R, ϵ) -cuff-good.

For any $\delta > 0$, we want to define $\mathcal{R}_\delta(\gamma_0) \subset \mathbb{I}(\gamma_0)$ to be the region of third connections that satisfy our previous estimate in Section 3.2. Formally, a point $\alpha \in \mathbb{I}(\gamma_0)$ is contained in $\mathcal{R}_\delta(\gamma_0)$ if and only if $P(\alpha)$ is good and there is a lift $((u, (f_u, E)), (v, (f_v, F)), l, \Lambda)$ of α in $\mathbb{F}(\gamma_0)$ such that:

1. if η is a third connection connecting a and b with length l , the lengths of two new cuffs are within $(2R - 2\epsilon, 2R + 2\epsilon)$;
2. suppose X_1, X_2 are the matrices of right actions defined as in Section 3.2 for frames at u and v , where f_u, f_v are regarded as $\eta'_A, -\eta'_B$; then $\phi(Y)$ lies in the intersection of δ -neighborhoods centered at X_1 and X_2 .

By definition, we have $\text{supp}(\tilde{\nu}) \subset \mathcal{R}(\gamma_0)$. By our discussion in Section 3.2, there exists a universal constant $C_0 > 0$, such that

$$\mathcal{R}_{\epsilon - C_0 e^{-R}}(\gamma_0) \subset \mathcal{R}(\gamma_0) \subset \mathcal{R}_{\epsilon + C_0 e^{-R}}(\gamma_0). \quad (3.10)$$

By Theorem 4.9 in [KW19], we have the following proposition.

Proposition 3.11. *There exists $q = q(M) \leq 1/2$ and a constant $C > 0$ such that for any $\epsilon > 0$, the following holds when R is sufficiently large: Let γ_0 be an (R, ϵ) -good curve and $\zeta = e^{-qR}$. Then for any $A \subset \mathcal{R}(\gamma_0) \subset \mathbb{I}(\gamma_0)$ we have*

$$(1 - \zeta)|\mathcal{N}_{-\zeta}(A)| \leq C \cdot \tilde{\nu}(A) \text{Vol}(M) \leq (1 + \zeta)|\mathcal{N}_\zeta(A)|,$$

here $\mathcal{N}_{-\zeta}(A) = \mathbb{I}(\gamma_0) - \mathcal{N}_\zeta(\mathbb{I}(\gamma_0) - A)$.

Let $\rho: N^1(\gamma_0) \times N^1(\gamma_0) \rightarrow N^1(\gamma_0) \times N^1(\gamma_0)$ be defined as $\rho(a, b) = (c, b)$ if a is a midpoint of c and b ; then every regular value of ρ has 4 preimages. We extend ρ to a map from $N^1(\gamma_0) \times N^1(\gamma_0) \times [0, \infty)$ to itself by preserving the third coordinate, and let $\mathbb{J}(\gamma_0) = \rho^*(\mathbb{I}(\gamma_0))$ be the pullback bundle. For each point (c, b, l, X) in our good region $\mathcal{R}(\gamma_0)$, there is only one preimage (a, b, l, X) under ρ_* such that:

1. The oriented distance from a to b lies in $(0, l(\gamma_0)/2)$.
2. After parallel transport to the same point on γ_0 , the distance between a and b (on a unit normal sphere) is bounded by $\pi/4$.

Here we need to assume that ϵ is no greater than some universal constant (by Proposition 3.4) to satisfy the second requirement. Let $\mathcal{S}(\gamma_0)$ and $\mathcal{S}_\delta(\gamma_0)$ be the corresponding component of $(\rho_*)^{-1}(\mathcal{R}(\gamma_0))$ and $(\rho_*)^{-1}(\mathcal{R}_\delta(\gamma_0))$, respectively, that satisfies the above two conditions, and let $\hat{\nu}$ be the pullback measure of $\tilde{\nu}$ on $\mathbb{J}(\gamma_0)$. By (3.10), we have

$$\mathcal{S}_{\epsilon - C_0 e^{-R}}(\gamma_0) \subset \mathcal{S}(\gamma_0) \subset \mathcal{S}_{\epsilon + C_0 e^{-R}}(\gamma_0). \quad (3.12)$$

Moreover, since the Jacobian of ρ is a constant, the result in Proposition 3.11 holds for $\hat{\nu}$ and $\mathcal{S}(\gamma_0) \subset \mathbb{J}(\gamma_0)$ as well.

Proposition 3.13. *There exists $q = q(M) \leq 1/2$ and a constant $C > 0$ such that for any $\epsilon > 0$, the following holds when R is sufficiently large: Let γ_0 be an (R, ϵ) -good curve and $\zeta = e^{-qR}$. Then for any $A \subset \mathcal{S}(\gamma_0) \subset \mathbb{J}(\gamma_0)$ we have*

$$(1 - \zeta)|\mathcal{N}_{-\zeta}(A)| \leq C \cdot \hat{\nu}(A) \text{Vol}(M) \leq (1 + \zeta)|\mathcal{N}_\zeta(A)|,$$

here $\mathcal{N}_{-\zeta}(A) = \mathbb{J}(\gamma_0) - \mathcal{N}_\zeta(\mathbb{J}(\gamma_0) - A)$.

Let $\pi: \mathbb{J}(\gamma_0) \rightarrow N^1(\gamma_0)$ be the projection to the first $N^1(\gamma_0)$ -coordinate. Let $\nu_a = \pi_*(\hat{\nu})$ be the *actual average foot measure*, and $\mu_a = \pi_*(\mathbb{1}_{\mathcal{S}(\gamma_0)})$ be the *estimated average foot measure*; then ν_a and μ_a are measures on $N^1(\gamma_0)$. The measures $\hat{\nu}, \nu_a$ and μ_a actually depend on the good curve γ_0 , and we will keep using the notations when there is no ambiguity. Clearly, the estimated average foot measure μ_a is invariant by the centralizer for the element of $\text{SO}(2n, 1)$ associated to γ_0 : translation along γ_0 , rotation of all the unit normal spheres by the same angle (around the north and south poles), and reflection through the equator of each sphere. We will prove a similar result to Proposition 3.13 for ν_a and μ_a on $N^1(\gamma_0)$ at the end of Section 3, through

fiberwise estimates for $\tilde{\nu}$ for $\mathbb{1}_{\mathcal{S}(\gamma_0)}$. For any $t \in N^1(\gamma_0)$, let $F^t = \pi^{-1}(t)$ be the preimage of t in $\mathbb{J}(\gamma_0)$; then F^t is a $\text{SO}(2n-1)$ -bundle over $N^1(\gamma_0) \times [0, \infty)$. We have the following lemma.

Lemma 3.14. *There exists $C_1 > 0$ such that when R is large enough: for any $t \in N^1(\gamma_0)$,*

$$F^t \cap N_\zeta(\mathcal{S}(\gamma_0)) \subset F^t \cap \mathcal{S}_{\epsilon+C_1\zeta}(\gamma_0),$$

where $\zeta = e^{-qR}$ as in Proposition 3.13.

Proof. Fix $t \in N^1(\gamma_0)$. By (3.12), we have

$$F^t \cap N_\zeta(\mathcal{S}(\gamma_0)) \subset F^t \cap N_\zeta(\mathcal{S}_{\epsilon+C_0e^{-R}}(\gamma_0)). \quad (3.15)$$

Suppose $\alpha = (t, y, l, Y) \in F^t \cap N_\zeta(\mathcal{S}_{\epsilon+C_0e^{-R}}(\gamma_0))$ for some $y \in N^1(\gamma_0)$, $l > 0$ and Y in the $\text{SO}(2n-1)$ -fiber at (t, y) . Then there exists $\beta = (t', y', l', Y') \in \mathcal{S}_{\epsilon+C_0e^{-R}}(\gamma_0)$ such that $d(\alpha, \beta) \leq \zeta$. Let $x \in N^1(\gamma_0)$ such that t is the midpoint of x and y , and x' the same for t' and y' . Since ρ is a Lipschitz map, there exists a universal constant C_2 , such that (x, y, l, Y) is $C_2\zeta$ -close to (x', y', l', Y') in $\mathbb{I}(\gamma_0)$. Therefore we can find the lifts (E_1, E_2, l, Y) and (E'_1, E'_2, l', Y') of (x, y, l, Y) and (x', y', l', Y') respectively, such that

1. (E_1, E_2, l, Y) is $C_2\zeta$ -close to (E'_1, E'_2, l', Y') in $\mathbb{F}(\gamma_0)$;
2. $\phi(Y')$ is in the $(\epsilon + C_0e^{-R})$ -neighborhood of X'_1 and X'_2 .

Here Z'_i (and Z_i) are the matrices for (E'_1, E'_2, l', X') (and (E_1, E_2, l, X)) as described in Condition 2 in the definition of $\mathcal{R}_\delta(\gamma_0)$. Now since $d(E_i, E'_i) < C_2\zeta$, so

$$d(X_1, X'_1), d(X_2, X'_2) < 2C_2\zeta.$$

Then by $d(Y, Y') < C_2\zeta$, we know Y is in the $(\epsilon + C_0e^{-R} + 3C_2\zeta)$ -neighborhood of X_1 and X_2 . Let $C_1 = C_0 + 3C_2$; then by $q \leq 1$, we have $\alpha \in F^t \cap \mathcal{S}_{\epsilon+C_0e^{-R}+3C_2\zeta}(\gamma_0) \subset F^t \cap \mathcal{S}_{\epsilon+C_1\zeta}(\gamma_0)$, and by (3.15),

$$F^t \cap N_\zeta(\mathcal{S}(\gamma_0)) \subset F^t \cap N_\zeta(\mathcal{S}_{\epsilon+C_0e^{-R}}(\gamma_0)) \subset F^t \cap \mathcal{S}_{\epsilon+C_1\zeta}(\gamma_0). \quad \square$$

By Lemma 3.1, we know that if $(t, y, l, X) \in F^t \cap \mathcal{S}_\delta(\gamma_0)$, then $l \in (R + 2 \ln 2 - 4\epsilon, R + 2 \ln 2 + 4\epsilon)$ and the distance from t to y is 4ϵ -close to $R/2$ (here the distance from t to y is half of the distance between two footpoints

of a *predicted* third connection). Since t is fixed, by parallel transport along γ_0 , we can think of $F^t \cap \mathcal{S}_\delta(\gamma_0)$ as a subset of a $\mathrm{SO}(2n-1)$ -bundle over

$$S^{2n-2} \times (R/2 - 1, R/2 + 1) \times [0, \infty).$$

Moreover, by our previous discussion in Section 3.2, the fiber of this bundle only depends on its projection on S^{2n-2} . Thus this bundle is a product space of D^t and $(R/2 - 1, R/2 + 1) \times [0, \infty)$, where D^t is a $\mathrm{SO}(2n-1)$ -bundle over S^{2n-2} . Let $\hat{\mathcal{S}}_\delta^t$ be the projection of $F^t \cap \mathcal{S}_\delta(\gamma_0)$ on D^t . Then by Lemma 3.1,

$$\hat{\mathcal{S}}_\delta^t \times \hat{L}_{R, \epsilon - Ce^{-R}} \subset F^t \cap \mathcal{S}_\delta(\gamma_0) \subset \hat{\mathcal{S}}_\delta^t \times \hat{L}_{R, \epsilon + Ce^{-R}}. \quad (3.16)$$

Here $\hat{L}_{R, \delta} \subset \mathbb{R} \times \mathbb{R}$ with measure $e^{2y} dx dy$ is the set of all pairs (x, y) satisfying:

1. $|2x + y - 2 \ln 2 - 2R| < 2\delta$;
2. $|(l(\gamma_0) - 2x) + y - 2 \ln 2 - 2R| < 2\delta$.

Similar to (3.3), we have

$$\mathrm{Vol}(\hat{L}_{R, \delta}) = 4e^{4R - l(\gamma_0)}(e^{2\delta} - e^{-2\delta})^2 \sim 64\delta^2 e^{4R - l(\gamma_0)}, \quad (3.17)$$

when $\delta \rightarrow 0$.

3.4.2 The volume of the space of good monodromies

In this section, we give an estimate of the volume of $\hat{\mathcal{S}}_\delta^t$ for any δ that is close to ϵ and any fixed $t \in N^1(\gamma_0)$. We first present a simple and rough estimate of the volume of the intersection of two ϵ -balls in $\mathrm{SO}(2n-1)$ in this subsection.

For $0 \leq d \leq 2r \leq \pi$, let $V(d, r, n)$ denote the volume of the intersection of two balls of radius r in $\mathrm{SO}(2n-1)$, whose centers lie at distance d , and $\hat{V}(d, r, n)$ be the analogous version in \mathbb{R}^{2n-1} . Then there exists a universal constant $M \equiv M(n) > 1$ such that

$$\frac{1}{M} \hat{V}(d, r, n) < V(d, r, n) < M \hat{V}(d, r, n),$$

and

$$\hat{V}(d, r, n) = 2 \int_0^{r/2-d} |B_{2n-2}(\sqrt{r^2 - (r-x)^2})| dx,$$

where $|B_{2n-2}(s)| = \pi^{n-1}s^{2n-2}/(n-1)!$ is the volume of the $2n-2$ dimensional ball of radius s . Letting $t = x/r$ and simplifying, we obtain

$$\widehat{V}(d, r, n) = \frac{2\pi^{n-1}r^{2n-1}}{(n-1)!} \int_0^{1-\frac{d}{2r}} (1 - (1-t)^2)^{n-1} dt$$

For fixed n , when $0 \leq d \leq 2r$, $\widehat{V}(d, r, n)$ is decreasing in d . Therefore,

$$\widehat{V}(d, r, n) \leq \widehat{V}(0, r, n) = |B_{2n-1}(r)| = \frac{2^n \pi^{n-1} r^{2n-1}}{(2n-1)!}.$$

When $d \leq 3r/2$, we have

$$\widehat{V}(d, r, n) \geq \widehat{V}(3r/2, r, n) = \frac{2\pi^{n-1}r^{2n-1}}{(n-1)!} \int_0^{1/4} (1 - (1-t)^2)^{n-1} dt.$$

And when $d \geq 2r - \eta$, we have

$$\begin{aligned} \widehat{V}(d, r, n) &\leq \widehat{V}(2r - \eta, r, n) = \frac{2\pi^{n-1}r^{2n-1}}{(n-1)!} \int_0^{\eta/2r} (1 - (1-t)^2)^{n-1} dt \\ &\leq \frac{2\pi^{n-1}r^{2n-1}}{(n-1)!} \frac{\eta}{2r} \left(\frac{2\eta}{2r}\right)^{n-1} \\ &= \frac{\pi^{n-1}r^{n-1}\eta^n}{(n-1)!}. \end{aligned}$$

Thus there exists a constant, denoted by M_n , such that

$$V(d, r, n) < M_n r^{2n-1}, \quad (3.18)$$

and, when $d \leq 3r/2$,

$$V(d, r, n) > r^{2n-1}/M_n. \quad (3.19)$$

and when $d \geq 2r - \eta$,

$$V(d, r, n) < M_n r^{n-1} \eta^n. \quad (3.20)$$

Now for any two points $x, y \in N^1(\gamma_0)$, let $m_x(y) \in N^1(\gamma_0)$ be the unique point such that x is a midpoint of y and $m_x(y)$. For simplicity, we will slightly abuse notation by using m to denote the same kind of maps for two points on S^{2n-2} . For $x_1, x_2 \in S^{2n-2}$, arbitrarily choose two collection of unit

vectors E, F such that (x_1, E) and x_2, F are orthonormal frames of R^{2n-1} with opposite orientations. Let $Z_1, Z_2 \in \text{SO}(2n-1)$ such that

$$(-x_1, E) \cdot Z_1 = (x_2, F)$$

and

$$(x_1, E) \cdot \Lambda Z_2 = (-x_2, F),$$

where Λ is the monodromy pf γ_0 ; we then have Λ is conjugate to $Z_2^{-1}\phi(Z_1)$. We define $f(x_1, x_2) = d(Z_1, Z_2)$; then f is well defined, since the distance is independent of the choices of frames E and F .

By our definition of $\hat{S}_\delta^t \subset D^t$, the projection of \hat{S}_δ^t on S^{2n-2} (as the base space of D^t) consists of $y \in S^{2n-2}$ such that $f(y, m_t(y)) < 2\delta$. Here t and y are considered as points in the same S^{2n-2} by parallel transport, since t is fixed. Therefore for we have,

$$\text{Vol}(\hat{S}_\delta^t) = \int_{S^{2n-2}} \mathbb{1}_{\{y: f(y, m_t(y)) < 2\delta\}} V(f(y, m_t(y)), \delta) d\sigma.$$

We now can present an upper and lower bound of this volume for δ that is close to ϵ .

Proposition 3.21. *There exists a universal constant $M_1 > 0$ such that for any $t \in N^1(\gamma_0)$ and $\delta > 8\epsilon/9$, we have*

$$\delta^{4n-3}/M_1 < \text{Vol}(\hat{S}_\delta^t) < M_1 \delta^{4n-3}.$$

Proof. We first establish the upper bound. By Proposition 3.4 and (3.18), we have for any $t \in N^1(\gamma_0)$,

$$\begin{aligned} \text{Vol}(\hat{S}_\delta^t) &= \int_{S^{2n-2}} \mathbb{1}_{\{y: f(y, m_t(y)) < 2\delta\}} V(f(y, m_t(y)), \delta) d\sigma \\ &\leq \int_{S^{2n-2}} \mathbb{1}_{\{y: d_{S^{2n-2}}(y, m_t(y)) < 3\delta/2\}} M \delta^{2n-1} d\sigma \\ &= \int_{S^{2n-2}} \mathbb{1}_{\{y: d_{S^{2n-2}}(y, t) < 3\delta/4\}} M \delta^{2n-1} d\sigma \\ &= \text{Vol}(\{y : d_{S^{2n-2}}(y, t) < 3\delta/4\}) \cdot M \delta^{2n-1} \\ &\leq C_1 \delta^{2n-2} \cdot M \delta^{2n-1} = C_1 M \delta^{4n-3}, \end{aligned}$$

for some universal constant $C_1 > 0$.

By Proposition 3.5, for $x, y \in S^{2n-2}$, if $d_{S^{2n-2}}(x, y) < 3\delta/16$, then

$$f(x, y) < \epsilon + 2d_{S^{2n-2}}(x, y) \leq 9\delta/8 + 3\delta/8 = 3\delta/2.$$

Together with (3.19), we have

$$\begin{aligned} \text{Vol}(\hat{S}_\delta^t) &= \int_{S^{2n-2}} \mathbb{1}_{\{y: f(y, m_t(y)) < 2\delta\}} V(f(y, m_t(y)), \delta) d\sigma \\ &\geq \int_{S^{2n-2}} \mathbb{1}_{\{y: f(y, m_t(y)) < 3\delta/2\}} \cdot \delta^{2n-1}/M d\sigma \\ &\geq \int_{S^{2n-2}} \mathbb{1}_{\{y: d_{S^{2n-2}}(y, m_t(y)) < 3\delta/16\}} \cdot \delta^{2n-1}/M d\sigma \\ &= \int_{S^{2n-2}} \mathbb{1}_{\{y: d_{S^{2n-2}}(y, t) < 3\delta/32\}} \cdot \delta^{2n-1}/M d\sigma \\ &= \text{Vol}(\{y : d_{S^{2n-2}}(y, t) < 3\delta/32\}) \cdot \delta^{2n-1}/M \\ &\geq C_2 \delta^{2n-2} \cdot \delta^{2n-1}/M = \frac{C_2}{M} \delta^{4n-3}, \end{aligned}$$

for some universal constant $C_2 > 0$.

Now we let $M_1 = \max\{C_1 M, M/C_2\}$, and then the result is proved. \square

With the above proposition we can prove the main result of this section, that the the estimated average foot measure μ_a is bounded above and below by Lebesgue measure.

Proposition 3.22. *There is a universal constant $B > 1$ such that for any $\epsilon > 0$, the following holds when R is sufficiently large: Suppose γ_0 is an (R, ϵ) -good curve, and λ is the Lebesgue measure on $N^1(\gamma_0)$. Then for any $t \in N^1(\gamma_0)$,*

$$B^{-1} e^{2R} \epsilon^{4n-1} < \left. \frac{d\mu_a}{d\lambda} \right|_t < B e^{2R} \epsilon^{4n-1}.$$

Proof. By definition, we know that for any $t \in N^1(\gamma_0)$,

$$\left. \frac{d\mu_a}{d\lambda} \right|_t = |F^t \cap \mathcal{S}(\gamma_0)|.$$

Then by (3.12), (3.16), (3.17), and Proposition 3.21, we have

$$\begin{aligned} |F^t \cap \mathcal{S}(\gamma_0)| &\leq |F^t \cap \mathcal{S}_{\epsilon+C_0 e^{-R}}(\gamma_0)| \\ &\leq |\hat{\mathcal{S}}_{\epsilon+C_0 e^{-R}}^t \times \hat{L}_{R, \epsilon+C_0 e^{-R}}| \\ &= \text{Vol}(\hat{\mathcal{S}}_{\epsilon+C_0 e^{-R}}^t) \times \text{Vol}(\hat{L}_{R, \epsilon+C_0 e^{-R}}) \\ &\leq M_1 (2\epsilon)^{4n-3} \times 800 e^{2R} \epsilon^2 = B_1 e^{2R} \epsilon^{4n-1}, \end{aligned}$$

for some universal constant B_1 . Similarly, there is another universal constant B_2 such that

$$|F^t \cap \mathcal{S}(\gamma_0)| \geq B_2 e^{2R} \epsilon^{4n-1}.$$

Let $B = \max\{B_1, B_2^{-1}\}$, and then the result is proved. \square

3.4.3 Counting good connections

We now compare the actual average foot measure and the estimated average foot measure on $N^1(\gamma_0)$.

Proposition 3.23. *There exists $q = q(M) \leq 1/2$ and constant $C, L > 0$ such that for any $\epsilon > 0$, the following holds when R is sufficiently large: Let γ_0 be an (R, ϵ) -good curve and $\zeta = e^{-qR}$. Then for any $B \subset N^1(\gamma_0)$, we have*

$$(1 - L\zeta/\epsilon)\mu_a(\mathcal{N}_{-\zeta}(B)) \leq C \cdot \nu_a(B) \text{Vol}(M) \leq (1 + L\zeta/\epsilon)\mu_a(\mathcal{N}_{\zeta}(B)),$$

here $\mathcal{N}_{-\zeta}(B) = N^1(\gamma_0) - \mathcal{N}_{\zeta}(N^1(\gamma_0) - B)$.

Proof. For any $B \subset N^1(\gamma_0)$, we have $\nu_a(B) = \hat{\nu}(\pi^{-1}(B))$. Then by Proposition 3.13, we have

$$\begin{aligned} & (1 - \zeta)|\mathcal{N}_{-\zeta}(\pi^{-1}(B) \cap \mathcal{S}(\gamma_0))| \\ & \leq C_0 \cdot \hat{\nu}(\pi^{-1}(B)) \cdot \text{Vol}(M) = C_0 \cdot \nu_a(B) \cdot \text{Vol}(M) \\ & \leq (1 + \zeta)|\mathcal{N}_{\zeta}(\pi^{-1}(B) \cap \mathcal{S}(\gamma_0))|, \end{aligned}$$

here C_0 is the constant C in Proposition 3.13. Thus we only need to show that there exists $L > 0$ such that

$$|\mathcal{N}_{\zeta}(\pi^{-1}(B) \cap \mathcal{S}(\gamma_0))| < (1 + L\zeta/\epsilon)\mu_a(\mathcal{N}_{\zeta}(B)),$$

and the proof of the other side of the inequality will be the same.

We notice that

$$\mathcal{N}_{\zeta}(\pi^{-1}(B) \cap \mathcal{S}(\gamma_0)) \subset \mathcal{N}_{\zeta}(\pi^{-1}(B)) \cap \mathcal{N}_{\zeta}(\mathcal{S}(\gamma_0)) = \pi^{-1}(\mathcal{N}_{\zeta}(B)) \cap \mathcal{N}_{\zeta}(\mathcal{S}(\gamma_0)),$$

and

$$\mu_a(\mathcal{N}_{\zeta}(B)) = |\pi^{-1}(\mathcal{N}_{\zeta}(B)) \cap \mathcal{S}(\gamma_0)|.$$

Therefore it suffices to show that there exists $L > 0$ such that

$$|\pi^{-1}(\mathcal{N}_{\zeta}(B)) \cap \mathcal{N}_{\zeta}(\mathcal{S}(\gamma_0))| \leq (1 + L\zeta/\epsilon)|\pi^{-1}(\mathcal{N}_{\zeta}(B)) \cap \mathcal{S}(\gamma_0)|.$$

We will prove this inequality fiberwise, i.e. we want to prove that there exists $L > 0$ such that for any $t \in N^1(\gamma_0)$,

$$|F^t \cap \mathcal{N}_\zeta(\mathcal{S}(\gamma_0))| \leq (1 + L\zeta/\epsilon) |F^t \cap \mathcal{S}(\gamma_0)|. \quad (3.24)$$

By Lemma 3.14 and (3.16), we have

$$F^t \cap \mathcal{N}_\zeta(\mathcal{S}(\gamma_0)) \subset F^t \cap \mathcal{S}_{\epsilon+C_1\zeta}(\gamma_0) \subset \hat{\mathcal{S}}_{\epsilon+C_1\zeta}^t \times \hat{L}_{R,\epsilon+C_1\zeta e^{-R}}. \quad (3.25)$$

By (3.12) and (3.16), we have

$$F^t \cap \mathcal{S}(\gamma_0) \supset F^t \cap \mathcal{S}_{\epsilon-C_0e^{-R}} \supset \hat{\mathcal{S}}_{\epsilon-C_0e^{-R}}^t \times \hat{L}_{R,\epsilon-C_0e^{-R}}. \quad (3.26)$$

By (3.17) and $\zeta = e^{-qR}$ for $q \leq 1/2$, we know that when R is sufficiently large,

$$\text{Vol}(\hat{L}_{R,\epsilon+C_1\zeta e^{-R}}) \leq (1 + \zeta) \text{Vol}(\hat{L}_{R,\epsilon-C_0e^{-R}}). \quad (3.27)$$

Hence now we only need to compare $\text{Vol}(\hat{\mathcal{S}}_{\epsilon+C_1\zeta}^t)$ and $\text{Vol}(\hat{\mathcal{S}}_{\epsilon-C_0e^{-R}}^t)$.

Let $b: D^t \rightarrow S^{2n-2}$ be the projection map, and for any $s \in S^{2n-2}$, let $G^s = b^{-1}(s) \cong \text{SO}(2n-1)$. By definition, we have $\hat{\mathcal{S}}_{\epsilon-C_0e^{-R}}^t \subset \hat{\mathcal{S}}_{\epsilon+C_1\zeta}^t$. Suppose $u \in b(\hat{\mathcal{S}}_{\epsilon+C_1\zeta}^t) \subset S^{2n-2}$; then $f(u, m_t(u)) \leq 2(\epsilon + C_1\zeta)$.

We now want to obtain an upper bound on $\text{Vol}(G^u \cap (\hat{\mathcal{S}}_{\epsilon+C_1\zeta}^t - \hat{\mathcal{S}}_{\epsilon-C_0e^{-R}}^t))$. We have two cases, depending on $f(u, m_t(u))$:

- If $2(\epsilon - C_0e^{-R}) < f(u, m_t(u)) \leq 2(\epsilon + C_1\zeta)$, then $u \notin b(\hat{\mathcal{S}}_{\epsilon-C_0e^{-R}}^t)$. Thus by (3.20), when R is sufficiently large, we have

$$\begin{aligned} \text{Vol}(G^u \cap (\hat{\mathcal{S}}_{\epsilon+C_1\zeta}^t - \hat{\mathcal{S}}_{\epsilon-C_0e^{-R}}^t)) &= \text{Vol}(G^u \cap \hat{\mathcal{S}}_{\epsilon+C_1\zeta}^t) \\ &= V(f(u, m_t(u)), \epsilon + C_1\zeta, n) \\ &< M_n(\epsilon + C_1\zeta)^{n-1} (2\epsilon + C_1\zeta - f(u, m_t(u)))^n \\ &< M_n(2\epsilon)^{n-1} (C_1\zeta + C_0e^{-R})^n \\ &< M_n 2^{2n-1} C_1^n \epsilon^{n-1} \zeta^n. \end{aligned} \quad (3.28)$$

- If $f(u, m_t(u)) \leq 2(\epsilon - C_0e^{-R})$, then $u \in b(\hat{\mathcal{S}}_{\epsilon-C_0e^{-R}}^t)$. Thus

$$\text{Vol}(G^u \cap (\hat{\mathcal{S}}_{\epsilon+C_1\zeta}^t - \hat{\mathcal{S}}_{\epsilon-C_0e^{-R}}^t)) = V(f(u, m_t(u)), \epsilon + C_1\zeta) - V(f(u, m_t(u)), \epsilon - C_0e^{-R}).$$

The right hand side of the above equation reaches its maximum when $f(u, m_t(u)) = 0$, i.e. the distance between two centers is 0; then the

maximum is the difference of the volume of two concentric balls. Hence there is a universal constant $M_2 > 0$ such that when R is sufficiently large, we have

$$\begin{aligned} \text{Vol}(G^u \cap (\hat{\mathcal{S}}_{\epsilon+C_1\zeta}^t - \hat{\mathcal{S}}_{\epsilon-C_0e^{-R}}^t)) &< M_2((\epsilon + C_1\zeta)^{2n-1} - (\epsilon - C_0e^{-R})^{2n-1}) \\ &< (8n - 4)M_2C_1\epsilon^{2n-2}\zeta. \end{aligned} \quad (3.29)$$

Hence by Propositions 3.4 and 3.21, (3.28), and (3.29), when R is sufficiently large, we have

$$\begin{aligned} \text{Vol}(\hat{\mathcal{S}}_{\epsilon+C_1\zeta}^t - \hat{\mathcal{S}}_{\epsilon-C_0e^{-R}}^t) &\leq (\text{Vol}(b(\hat{\mathcal{S}}_{\epsilon+C_1\zeta}^t)) - \text{Vol}(b(\hat{\mathcal{S}}_{\epsilon-C_0e^{-R}}^t))) \cdot M_n 2^{2n-1} C_1^n \epsilon^{n-1} \zeta^n \\ &\quad + \text{Vol}(b(\hat{\mathcal{S}}_{\epsilon-C_0e^{-R}}^t)) \cdot (8n - 4)M_2C_1\epsilon^{2n-2}\zeta \\ &\leq \text{Vol}(S^{2n-2}) \cdot M_n 2^{2n-1} C_1^n \epsilon^{n-1} \zeta^n + C_1\epsilon^{2n-2} \cdot (8n - 4)M_2C_1\epsilon^{2n-2}\zeta \\ &\leq M_3\epsilon^{4n-4}\zeta \leq M_4(\zeta/\epsilon) \cdot \text{Vol}(\hat{\mathcal{S}}_{\epsilon-C_0e^{-R}}^t), \end{aligned}$$

for some universal constants $M_3, M_4 > 0$. So we have

$$\text{Vol}(\hat{\mathcal{S}}_{\epsilon+C_1\zeta}^t) \leq (1 + M_4\zeta/\epsilon) \text{Vol}(\hat{\mathcal{S}}_{\epsilon-C_0e^{-R}}^t). \quad (3.30)$$

By (3.25), (3.26), (3.27), and (3.30), let $L = 2M_4 + 2$, and then (3.24) is proved. \square

4 Matching and the main theorem

In this last section, we will match each pair of pants to well-matched pants along each boundary component to obtain an incompressible surface in M .

4.1 Matching pattern and the incompressibility

We first provide some preliminaries on matching that we will use to complete the proof of the main theorem.

We start to introduce how we match pants together, which is a little bit different from the pattern used in [KM12] and [Ham15]. We then will show that if a closed surface in M is made by good pants in this pattern, it is incompressible by applying the definition and the theorem in [Ham15].

For an oriented closed geodesic γ in M , we introduce two actions on $N^1(\gamma)$. Let T_γ be the parallel transport along the orientation of γ by distance

1 and A_γ be given by the antipodal map on the unit normal sphere at each point of γ . We notice that T_γ commutes with A_γ , and let

$$\tau_\gamma = T_\gamma \circ A_\gamma = A_\gamma \circ T_\gamma: N^1(\gamma) \rightarrow N^1(\gamma).$$

Suppose that P_1, P_2 are two oriented (R, ϵ) -cuff-good and good pants in M that share a common cuff γ , for $\epsilon > 0$ sufficiently small and R sufficiently large. We then can define:

Definition 4.1. For $\sigma > 0$, P_1 and P_2 are σ -well-matched, if the following are satisfied:

1. The induced orientations of γ as a cuff of P_1 and P_2 are opposite.
2. Let a_1 be an average foot of P_1 , then there is an average foot a_2 of P_2 such that the distance between $\tau_\gamma(a_1)$ and a_2 in $N^1(\gamma)$ is less than σ .

Remark 4.2. We need to clarify that when considering the action of T_γ on a_1 , the orientation of γ is induced by the orientation of P_1 . Then this definition is symmetric in P_1 and P_2 , and γ is not necessarily oriented.

We then recall Hamenstädt's definition and result.

Definition 4.3 ([Ham15], Definition 6.1). For $\sigma > 0$, two pants P_1, P_2 are σ -well attached along a common boundary geodesic γ if the following holds true:

1. The orientation of γ as a boundary geodesic of P_1 and P_2 are opposite.
2. Let x be an endpoint of a short orthogeodesic of P_1 on γ . Then there is an endpoint y of a short orthogeodesic of P_2 on γ whose oriented distance to x is contained in the interval $[1 - \sigma, 1 + \sigma]$.
3. Let v_1 be the direction of the short orthogeodesic of P_1 at x and let v_2 be the direction of a short orthogeodesic of P_2 at y ; then the angle between v_2 and the image of $-v_1$ under parallel transport along the oriented subarc of γ connecting x to y is at most σ .

Proposition 4.4 ([Ham15], Proposition 6.2). *For any $b > 1$, there are $\delta \in (0, \pi/4]$ and $R_0 > 10$ with the following property. Let $R > R_0$ and let $S \subset M$ be a piecewise immersed closed surface composed of finitely many (R, δ) -good pants which are R^{-b} -well attached along their common boundary geodesics. Then S is incompressible.*

Remark 4.5. In [Ham15], for a closed oriented surface S of genus $g \geq 2$ and a closed Riemannian manifold M of non-positive sectional curvature, a continuous map $f: S \rightarrow M$ is called *incompressible* if $f_*: \pi_1(S) \rightarrow \pi_1(M)$ is injective.

Now we can state our matching result and prove it.

Proposition 4.6. *There are $\delta \in (0, \pi/4]$ and $R_0 > 10$ such that when $R > R_0$, the following holds: Suppose $S \subset M$ is a piecewise immersed closed surface composed of finitely many (R, δ) -good pants which are R^{-2} -well-matched along their common boundary geodesics. Then S is a π_1 -injective surface.*

Proof. We choose $b = 3/2$ in Proposition 4.4, then it is sufficient to show that when R_0 is large enough, if two (R, δ) -good pants are R^{-2} -well-matched, then they are $R^{-3/2}$ -well attached. And this is a quick result from the estimate of the difference between average feet and short feet in Proposition 3.9. \square

4.2 Matching pants

We follow the idea in [KW21] to use Hall's Marriage Theorem to match pants along each good curve. To be more specific, fix an arbitrary good curve γ_0 , and for each oriented good pants Π with γ_0 as a cuff, we want to match Π with a well-matched good pants that induces the opposite orientation on γ_0 . We then form a bipartite graph, whose vertices are oriented good pants P with an assigned average foot a . There is an edge that joins two vertices if and only if they are well-matched by γ_0 .

We let $\hat{\Pi}_{R,\epsilon}$ be the set of pairs (P, a) where P is an unoriented good and (R, ϵ) -cuff-good pants in M and a is an average foot of P , and $\hat{\Pi}_{R,\epsilon}^*$ is the set of all oriented (P, a) . Since each good pants has two average feet, so the projection $(P, a) \mapsto P$ is a 2-1 mapping. Moreover, $\hat{\Pi}_{R,\epsilon}$ and $\hat{\Pi}_{R,\epsilon}^*$ are finite sets, and $\hat{\Pi}_{R,\epsilon}^*$ is a double cover of $\hat{\Pi}_{R,\epsilon}$ by discarding the orientation.

For any (R, ϵ) -good curve γ_0 , let $\hat{\Pi}_{R,\epsilon}(\gamma_0) \subset \hat{\Pi}_{R,\epsilon}$ consist of (P, a) where γ_0 is a cuff of P , and $\hat{\Pi}_{R,\epsilon}^*(\gamma_0) \subset \hat{\Pi}_{R,\epsilon}^*$ be the set of oriented (P, a) where γ_0 is a cuff of P . Let $p: \hat{\Pi}_{R,\epsilon}(\gamma_0) \rightarrow N^1(\gamma_0)$ by $p(P, a) = a$, then the point measure induced by p on $N^1(\gamma_0)$ is the actual average foot measure ν_a .

In this subsection, we will prove the following theorem, which is analogous to Theorem 3.1 in [KM12], Theorem 5.1 in [KW21], and Theorem 6.6 in [Rao25].

Theorem 4.7. *For all $\epsilon > 0$, there exists $R_0 > 0$ such that when $R > R_0$, the following holds: Suppose γ_0 is an (R, ϵ) -good curve in M . Then there exists a permutation $\sigma_{\gamma_0}: \hat{\Pi}_{R,\epsilon}(\gamma_0) \rightarrow \hat{\Pi}_{R,\epsilon}(\gamma_0)$, such that for any $\pi \in \hat{\Pi}_{R,\epsilon}(\gamma_0)$,*

$$d(p(\sigma_{\gamma_0}(\pi)), \tau_{\gamma_0}(p(\pi))) < R^{-2}.$$

We recall that the Cheeger constant $h(M)$ for a smooth closed m -dimensional Riemannian manifold M is defined as

$$h(M) = \inf \frac{|\partial A|}{\min\{|A|, |M - A|\}},$$

among all m -dimensional submanifold $A \subset M$. Then for all these M , we have the following result.

Theorem 4.8 (Theorem 5.3 in [KW21]). *Suppose $A \subset M$ and $|\mathcal{N}_\eta(A)| \leq |M|/2$ for some $\eta > 0$, then*

$$|\mathcal{N}_\eta(A)| \geq (1 + \eta h(M))|A|.$$

We have another general result.

Proposition 4.9. *There exist constants $R_0 > 0$ such that when $R > R_0$: Suppose M is a S^{2n-2} -fiber bundle over a round circle with a circumference of $2R$, where S^{2n-2} is the standard unit sphere. Then $h(M) \geq 1/4R$.*

Proof. Let S_R^1 be the circle with a circumference of $2R$. Without loss of generality, we rescale the size of S^{2n-2} so that its volume equals 1. Suppose A is a smooth $(2n-1)$ -dimensional submanifold of M with $|A| \leq |M|/2$. For any $x \in S_R^1$, let S_x^{2n-2} be the fiber over x in M , and $q_x(A) = |A \cap S_x^{2n-2}|$ be the $(2n-2)$ -dimensional volume of the intersection. Let h be the Cheeger constant of S^{2n-2} , Then we have

$$|\partial A| \geq \int_{x \in S_R^1} h \cdot \min\{q_x(A), |S^{2n-2}| - q_x(A)\} dx.$$

Since M is a fiber bundle over a circle, so we also have

$$|\partial A| \geq \sup q_x(A) - \inf q_x(A).$$

If $\sup q_x(A) - \inf q_x(A) \geq 1/4$, then

$$\frac{|\partial A|}{|A|} \geq \frac{1/4}{|A|} \geq \frac{1/2}{|M|} = \frac{1}{4R}. \quad (4.10)$$

If $\sup q_x(A) - \inf q_x(A) \leq 1/4$, then by $|A| \leq |M|/2$, we have $\inf q_x(A) \leq |S^{2n-2}|/2 = 1/2$. Thus $q_x(A) \leq \sup q_x(A) \leq 3/4$, and then $|S^{2n-2}| - q_x(A) \geq 1/4$. Therefore

$$\begin{aligned} |\partial A| &\geq \int_{x \in S_R^1} h \cdot \min\{q_x(A), |S^{2n-2}| - q_x(A)\} dx \\ &\geq \int_{x \in S_R^1} h \cdot \min\{q_x(A), 1/4\} dx \\ &\geq \int_{x \in S_R^1} h \cdot \min\{q_x(A), q_x(A)/3\} dx \\ &= \frac{h}{3} \int_{x \in S_R^1} q_x(A) dx = \frac{h|A|}{3}. \end{aligned}$$

Hence $|\partial A|/|A| \geq h/3$.

Together with (4.10), when R is sufficiently large, we have

$$h(M) \geq \min\left\{\frac{1}{4R}, \frac{h}{3}\right\} = \frac{1}{4R}. \quad \square$$

For any (R, ϵ) -good curve γ_0 , $N^1(\gamma_0)$ is a mapping torus with length $l(\gamma_0) \leq 2R$. Then by Theorem 4.8, we have the following corollary.

Corollary 4.11. *If $A \subset N^1(\gamma_0)$ and $|\mathcal{N}_\eta(A)| \leq |N^1(\gamma_0)|/2$ for some $\eta > 0$, then*

$$|\mathcal{N}_\eta(A)| \geq \left(1 + \frac{\eta}{4R}\right)|A|.$$

Moreover, we have a similar result for μ_a .

Proposition 4.12. *If $A \subset N^1(\gamma_0)$ and $|\mathcal{N}_\eta(A)| \leq |N^1(\gamma_0)|/2$ for some $\eta > 0$, then*

$$\mu_a(\mathcal{N}_\eta(A)) \geq \left(1 + \frac{\eta}{4B^2R}\right)\mu_a(A),$$

where B is the constant in Proposition 3.22.

Proof. By Proposition 3.22 and Corollary 4.11, we have:

$$\begin{aligned} \mu_a(\mathcal{N}_\eta(A) - A) &\geq B^{-1}e^{2R}\epsilon^{4n-1}|\mathcal{N}_\eta(A) - A| \\ &\geq B^{-1}e^{2R}\epsilon^{4n-1}\frac{\eta}{4R}|A| \\ &\geq \frac{\eta}{4B^2R}\mu_a(A). \end{aligned} \quad \square$$

Now we can present the proof of Theorem 4.7.

Proof of Theorem 4.7. Let $\xi = R^{-2}$. By definition, we know that for any $A \subset N^1(\gamma_0)$,

$$\nu_a(A) = \#\{\pi \in \hat{\Pi}_{R,\epsilon}(\gamma_0) : p(\pi) \in A\}.$$

Then by Hall's Marriage Theorem, the result follows from the claim that

$$\nu_a(\mathcal{N}_\xi(A)) \geq \nu_a(\tau_{\gamma_0}(A)),$$

for any $A \subset N^1(\gamma_0)$. Now we prove this claim.

If $|\mathcal{N}_{\xi/2}(A)| \leq |N^1(\gamma_0)|/2$, by Proposition 3.23, Proposition 4.12, and the invariance of μ_a under τ_{γ_0} , we have when R is sufficiently large,

$$\begin{aligned} \nu_a(\mathcal{N}_\xi(A)) &\geq \frac{1}{C \text{Vol}(M)} (1 - L\zeta/\epsilon) \mu_a(\mathcal{N}_{\xi-\zeta}(A)) \\ &\geq \frac{1}{C \text{Vol}(M)} (1 - L\zeta/\epsilon) \mu_a(\mathcal{N}_{\xi/2}(A)) \\ &\geq \frac{1}{C \text{Vol}(M)} (1 - L\zeta/\epsilon) \left(1 + \frac{\xi/2 - \zeta}{4B^2R}\right) \mu_a(\mathcal{N}_\zeta(A)) \\ &\geq \frac{1}{C \text{Vol}(M)} (1 + L\zeta/\epsilon) \mu_a(\mathcal{N}_\zeta(A)) \\ &= \frac{1}{C \text{Vol}(M)} (1 + L\zeta/\epsilon) \mu_a(\mathcal{N}_\zeta(\tau_{\gamma_0}(A))) \\ &\geq \nu(\tau_{\gamma_0}(A)). \end{aligned}$$

Otherwise, let $A' = N^1(\gamma_0) - \mathcal{N}_\xi(A)$, then $|\mathcal{N}_{\xi/2}(A')| \leq |N^1(\gamma_0)|/2$. Then by the same reasoning, we have $\nu_a(\mathcal{N}_\xi(\tau_{\gamma_0}(A'))) \geq \nu_a(A')$. Therefore

$$\begin{aligned} \nu_a(N^1(\gamma_0)) - \nu_a(\tau_{\gamma_0}(A)) &= \nu_a(\tau_{\gamma_0}(N^1(\gamma_0) - A)) \\ &\geq \nu_a(\tau_{\gamma_0}(\mathcal{N}_\xi(A'))) \\ &\geq \nu_a(N^1(\gamma_0) - \mathcal{N}_\xi(A)). \end{aligned}$$

Hence $\nu_a(\mathcal{N}_\xi(A)) \geq \nu_a(\tau_{\gamma_0}(A))$. □

4.3 Proof of the main theorem

Proof of Theorem 1.1. We want to prove the theorem by showing that any closed hyperbolic $2n$ -manifold M has an immersed incompressible closed surface.

For any given $\epsilon > 0$, we choose R to be sufficiently large such that all previous statements hold for R and ϵ .

For any (R, ϵ) -good curve γ_0 , let τ_{γ_0} be a permutation of $\hat{\Pi}_{R, \epsilon}(\gamma_0)$ in Theorem 4.7. Then we apply the doubling trick on $\hat{\Pi}_{R, \epsilon}(\gamma_0)$ and τ_{γ_0} . We notice that $2\hat{\Pi}_{R, \epsilon}(\gamma_0) = \hat{\Pi}_{R, \epsilon}^*(\gamma_0)$. Therefore we find an involution τ^* of $\hat{\Pi}_{R, \epsilon}^*(\gamma_0)$ and match each pants with its image under τ^* . Then we know that P and P' are glued along γ_0 only if they are R^{-2} -well-matched.

Since the number of (R, ϵ) -good curve in M is finite (by Margulis argument), we can repeat the above step for each (R, ϵ) -good curve. Finally, we glue all pants in $\hat{\Pi}_{R, \epsilon}^*$ together and get a closed surface. We notice that this surface may have many connected components, but we can just choose an arbitrary one get a connected closed surface. Then by Proposition 4.6, this surface will be incompressible in M . □

References

- [Ham15] Ursula Hamenstädt. *Incompressible Surfaces in Rank One Locally Symmetric Spaces*. *Geometric and Functional Analysis*, 25(3):815–859, 2015.
- [KLM24] Jeremy Kahn, François Labourie, and Shahar Mozes. *Surface groups in uniform lattices of some semi-simple groups*. *Acta Mathematica*, 232(1):79–220, 2024.
- [KM12] Jeremy Kahn and Vladimir Marković. *Immersing almost geodesic surfaces in a closed hyperbolic three manifold*. *Annals of Mathematics*, pages 1127–1190, 2012.
- [KM15] Jeremy Kahn and Vladimir Marković. *The good pants homology and the Ehrenpreis conjecture*. *Annals of Mathematics*, pages 1–72, 2015.
- [KW19] Jeremy Kahn and Alex Wright. *Counting connections in a locally symmetric space*. <https://www.math.brown.edu/jk17/Connections.pdf>, 2019.

- [KW21] Jeremy Kahn and Alex Wright. *Nearly Fuchsian surface subgroups of finite covolume Kleinian groups*. *Duke Mathematical Journal*, 170(3):503–573, 2021.
- [LM15] Yi Liu and Vladimir Marković. *Homology of curves and surfaces in closed hyperbolic 3-manifolds*. *Duke Mathematical Journal*, 164(14):2723–2808, 2015.
- [Rao25] Zhenghao Rao. *Subgroups of genus-2 quasi-Fuchsian groups and cocompact Kleinian groups*. *Geometry & Topology*, 29(1):495–548, 2025.
- [TX18] Ser Peow Tan and Binbin Xu. *Carrier graphs for representations of the rank two free group into isometries of hyperbolic three space*. *arXiv preprint arXiv:1807.07277*, 2018.