

Variable selection via thresholding

Ka Long Keith Ho¹ and Hien Duy Nguyen^{1,2}

¹Joint Graduate School of Mathematics for Innovation, Kyushu University,
Fukuoka, Japan.

²Department of Mathematics and Physical Science, La Trobe University,
Melbourne, Australia.

March 28, 2025

Abstract

Variable selection comprises an important step in many modern statistical inference procedures. In the regression setting, when estimators cannot shrink irrelevant signals to zero, covariates without relationships to the response often manifest small but non-zero regression coefficients. The ad hoc procedure of discarding variables whose coefficients are smaller than some threshold is often employed in practice. We formally analyze a version of such thresholding procedures and develop a simple thresholding method that consistently estimates the set of relevant variables under mild regularity assumptions. Using this thresholding procedure, we propose a sparse, \sqrt{n} -consistent and asymptotically normal estimator whose non-zero elements do not exhibit shrinkage. The performance and applicability of our approach are examined via numerical studies of simulated and real data.

Keywords: Hard thresholding; Variable selection consistency; Sparse estimation; Asymptotic normality; Regression analysis.

1 Introduction

Regression analysis is a fundamental tool of statistical inference. In the modern context, the successful conduct of regression analysis typically requires a solution to the variable selection problem, which has been described and characterized in the works of George [2000], Fan and Lv [2010], and Heinze et al. [2018], among many others.

Variable selection arises as a necessary task in all regression settings, including generalized linear models (GLMs), nonlinear regression, mixed-effects models, functional models, and so forth.

However, due to the breadth of the subject, we will restrict our exposition to the linear regression setting, although we note that our methodology can be extended with minor modifications to most regression modeling tasks.

Let

$$\mathbf{Y} = \mathbf{X}\boldsymbol{\beta}_0 + \boldsymbol{\epsilon},$$

where $\mathbf{Y} = (Y_1, \dots, Y_n)^\top$, $\boldsymbol{\epsilon} = (\epsilon_1, \dots, \epsilon_n)^\top \in \mathbb{R}^n$ are the response and error vectors, and $\mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_n)^\top \in \mathbb{R}^{n \times p}$ is a design matrix, for $n, p \in \mathbb{N}$. Here, $\boldsymbol{\beta}_0 = (\beta_{0,1}, \dots, \beta_{0,p})^\top \in \mathbb{R}^p$ is the vector of true signals which are unknown to the analyst, who only observes \mathbf{X} and \mathbf{Y} . We shall further restrict ourselves to the random design scenario, where \mathbf{X}_i and ϵ_i are random quantities that are independent and identically distributed (IID) replicates of \mathbf{X} and ϵ , respectively, for each $i \in \{1, \dots, n\} = [n]$. We denote the common probability space on which all of our random objects are supported on $(\Omega, \mathcal{F}, \mathbb{P})$ with expectation operator \mathbb{E} .

We assume that p is large but finite and that $\boldsymbol{\beta}_0$ is sparse in the sense that $|\mathcal{S}_0| \gg 0$, where

$$\mathcal{S}_0 = \{j \in [p] : \beta_{0,j} = 0\}$$

is the set of irrelevant coordinates. The problem of variable selection under sparsity can be rephrased as the problem of estimating the set \mathcal{S}_0 . Traditional methods for addressing this problem, such as step-wise and best subset selection methods (see e.g., Hastie, 2009, Sec. 3.3), can be computationally prohibitive and slow, which has led to the large volume of research in sparse shrinkage estimators, such as the LASSO, bridge, and elastic net estimators, stemming from the works of Tibshirani [1996], Fu [1998], Fan and Li [2001], and Zou and Hastie [2005], among other early pioneers. Detailed accounts of the shrinkage estimation literature can be found in the texts of Bühlmann and van de Geer [2011], Hastie et al. [2015], Rish and Grabarnik [2014], and van de Geer [2016].

Typically, a shrinkage estimator $\hat{\boldsymbol{\beta}}_n = (\hat{\beta}_{n,1}, \dots, \hat{\beta}_{n,p})^\top$ will be such that the set of coordinates that are estimated to be irrelevant,

$$\hat{\mathcal{S}}_{0,n} = \{j \in [p] : \hat{\beta}_{n,j} = 0\}$$

is non-empty; i.e., $|\hat{\mathcal{S}}_{0,n}| > 0$. However, due to numerical implementations, it often occurs that the computed elements of $\hat{\boldsymbol{\beta}}_n$ will yield small but non-zero signals. This is the case, for example, when implementing the majorization–minimization algorithms for sparse penalties, such as in Hunter and Li [2005], Lloyd-Jones et al. [2018], and Abergel et al. [2024], or when using the “merit function” penalties of Zhao [2018, Ch. 7]. Furthermore, many shrinkage methods, such as the ridge estimator [Hoerl and Kennard, 1970] and its adaptive variants [Frommlet and Nuel, 2016, Dai et al., 2018, Abergel et al., 2024] output elements of $\hat{\boldsymbol{\beta}}_n$ that are sufficiently small as to be

considered negligible.

As per the references, an ad-hoc approach to handling such small estimates is to simply declare some small threshold $\delta > 0$ for which all estimated signals with absolute size smaller than δ are placed in the set of irrelevant coordinates. However, the question arises as to how small should the threshold δ be chosen. Practically, one may try a set of candidate thresholds and make a choice based on some empirical criterion.

In the following text, we shall consider a formalization of the described thresholding scheme whereupon one estimates the set of irrelevant signals \mathcal{S}_0 by the set $\hat{\mathcal{S}}_n$ of coordinates corresponding to elements of the estimated signal $\hat{\beta}_n$ whose size is less than some choice of a small threshold. That is, we produce an algorithm that takes some estimator $\hat{\beta}_n$ of β_0 as an input along with a sequence of threshold values $\delta_1 > \delta_2 > \dots > \delta_K > 0$, for some $K \in \mathbb{N}$. The algorithm then outputs an estimate $\hat{\mathcal{S}}_n$ of the set of irrelevant coordinates \mathcal{S}_0 , by eliminating all signals smaller than $\delta_{\hat{K}_n}$, where $\hat{K}_n \in [K]$ is chosen using an information criterion dependent on the sequence of threshold values along with their corresponding thresholded empirical risks. Under some regularity assumptions regarding the moments of \mathbf{X} and ϵ , and the minimum threshold size δ_K , we show that our algorithm is consistent in the sense that

$$\lim_{n \rightarrow \infty} \mathbb{P} \left(\hat{\mathcal{S}}_n = \mathcal{S}_0 \right) = 1. \quad (1)$$

To accompany our selection result, we also propose an estimator $\bar{\beta}_n$, whose j th coordinate is zero, if $j \in \hat{\mathcal{S}}_n$, and equal to the j th coordinate of the initial estimator of the regression model $\hat{\beta}_n$, otherwise (the so-called ‘‘hard threshold’’ estimator). Via the consistency of $\hat{\mathcal{S}}_n$, if the \sqrt{n} -blowup of $\hat{\beta}_n$ converges in distribution, then we can show that $\bar{\beta}_n$ satisfies an oracle result. Namely, this estimator is \sqrt{n} -consistent in its relevant coordinates, and the irrelevant signals are equal to zero with probability approaching one. Furthermore, if $\hat{\beta}_n$ is asymptotically normal, then the relevant coordinates of $\bar{\beta}_n$ are asymptotically normal as well.

Our technical results follow from recent work regarding the asymptotic distribution of the minimum value of an empirical risk minimization program by Westerhout et al. [2024] along with connections between such asymptotics and information criteria, reported in Nguyen [2024]. The proofs are further made possible using the random empirical processes theory of van der Vaart and Wellner [2007], as presented in van der Vaart and Wellner [2023]. Along with our theory, we also provide some illustrative numerical studies to demonstrate our methodology.

We note that our approaches are different and complementary to those of previous works that seek to use thresholding for variable selection and screening. For instance, Zhou [2009], van de Geer et al. [2011], Slawski and Hein [2013], Pokarowski and Mielniczuk [2015], Zheng et al. [2014], and Sun et al. [2019] consider thresholding already regularized estimators, in the high-dimensional setting. The regularized estimators are required to be well-behaved, via enforcement of strong moment and tail assumptions on the error ϵ and covariate \mathbf{X} and eigenvalue assumptions

on the Grammian, in order to establish variable selection consistency via finite sample bounds. In contrast, we take a wholly asymptotic approach, which provides qualitative instead of quantitative guarantees under weaker and different assumptions. Furthermore, our asymptotic normality result is obtained under different demands than those available in the literature. For instance, with normal noise and diminishing thresholds, Pötscher and Leeb [2009], Pötscher and Schneider [2011] and Schneider [2016] obtain marginal asymptotic normality results for the hard-thresholded least squares estimators. This contrasts with our results, which guarantee multivariate asymptotic normality without making assumptions on the noise distribution and for thresholds strictly bounded away from zero.

The remainder of the manuscript proceeds as follows. In Section 2, we describe the construction of our thresholding algorithm. Our main theoretical results are presented in Section 3, along with required technical preliminaries. Numerical studies are conducted in Section 4 and concluding remarks are provided in Section 5. Further proofs and auxiliary results are provided in the Appendix.

2 The thresholding algorithm

We retain notation from the introduction, and for each $k \in [K]$, we define a thresholding function $t_k : \mathbb{R} \rightarrow [0, 1]$, which serves as a smooth approximation to the step function threshold at δ_k :

$$\text{step}(b; \delta_k) = \begin{cases} 0 & \text{if } |b| \leq \delta_k, \\ 1 & \text{if } |b| > \delta_k. \end{cases} \quad (2)$$

We shall elaborate on the required properties of these functions in the sequel.

Given our estimate $\hat{\beta}_n$ of β_0 , we can apply the k th thresholding transformation

$$\mathbf{T}_k(\beta; \hat{\beta}_n) = \left(\beta_1 t_k(\hat{\beta}_{n,1}), \dots, \beta_p t_k(\hat{\beta}_{n,p}) \right)^\top, \quad (3)$$

which intuitively sets elements of β whose corresponding estimated signals are smaller than δ_k to zero. For brevity, we will sometimes write (3) as $\mathbf{T}_{n,k}(\beta)$. Furthermore, with the notation

$$\mathbf{D}_k(\beta) = \text{diag}(t_k(\beta_1), \dots, t_k(\beta_p)),$$

the diagonal matrix whose diagonal comprises the elements of $t_k(\beta_1), \dots, t_k(\beta_p)$, we further have the expression

$$\mathbf{T}_{n,k}(\beta) = \mathbf{D}_k(\hat{\beta}_n) \beta.$$

For each $k \in [K]$, we can now define the thresholded empirical risk

$$R_{n,k}(\boldsymbol{\beta}) = \frac{1}{n} \sum_{i=1}^n \{Y_i - \mathbf{X}_i^\top \mathbf{T}_{n,k}(\boldsymbol{\beta})\}^2,$$

where $R_{n,k} : \mathbb{R}^p \rightarrow \mathbb{R}$ can be understood as a random empirical process in the sense of van der Vaart and Wellner [2023, Sec. 3.13], since it depends not only on the averaging process, but also on the random vector $\hat{\boldsymbol{\beta}}_n$. If $\hat{\boldsymbol{\beta}}_n$ is a consistent estimator of $\boldsymbol{\beta}_0$, then we may anticipate that $R_{n,k}$ will converge to some limiting risk functional r_k , defined for each $\boldsymbol{\beta} \in \mathbb{R}^p$ by

$$r_k(\boldsymbol{\beta}) = \mathbb{E} \left[\{Y - \mathbf{X}^\top \mathbf{T}_k(\boldsymbol{\beta}, \boldsymbol{\beta}_0)\}^2 \right],$$

whose minimum value we can write as

$$\psi_k = \min_{\boldsymbol{\beta} \in \mathbb{R}^p} r_k(\boldsymbol{\beta}).$$

We note that this minimum and its empirical variant exist because both $R_{n,k}$ and r_k are quadratic, for each k and n . This is easier to see when writing:

$$R_{n,k}(\boldsymbol{\beta}) = \frac{1}{n} \sum_{i=1}^n \left\{ Y_i - \mathbf{X}_i^\top \mathbf{D}_k(\hat{\boldsymbol{\beta}}_n) \boldsymbol{\beta} \right\}^2 \quad \text{and} \quad r_k(\boldsymbol{\beta}) = \mathbb{E} \left[\{Y - \mathbf{X}^\top \mathbf{D}_k(\boldsymbol{\beta}_0) \boldsymbol{\beta}\}^2 \right].$$

Intuitively, the j th coordinate of $\mathbf{T}_k(\boldsymbol{\beta}; \boldsymbol{\beta}_0)$ is 0, when $|\beta_{0,j}| \leq \delta_k$ and β_j , otherwise. Thus, the minimum risk ψ_k is exactly the expected least squares value from fitting the linear regression model which excludes all variables in the k th irrelevant set

$$\mathcal{S}_k = \{j \in [p] : |\beta_{0,j}| \leq \delta_k\}.$$

If $\mathcal{S}_k \subset \mathcal{S}_0$, then only truly irrelevant coordinates are excluded when evaluating r_k , and thus ψ_k is equal to

$$\psi_0 = \min_{\boldsymbol{\beta} \in \mathbb{R}^p} \frac{1}{n} \|\mathbf{Y} - \mathbf{X}^\top \boldsymbol{\beta}\|^2 = \frac{1}{n} \|\mathbf{Y} - \mathbf{X}^\top \boldsymbol{\beta}_0\|^2,$$

the mean squared error under the true signal $\boldsymbol{\beta}_0$. However, if $\mathcal{S}_0 \subset \mathcal{S}_k$, then some relevant coordinates that are not in \mathcal{S}_0 may be excluded when computing r_k and thus $\psi_0 \leq \psi_k$. Observe further that since $\delta_1 > \dots > \delta_K$ is decreasing, we have $\mathcal{S}_1 \supset \dots \supset \mathcal{S}_K$, and as per the argument above, it holds that $\psi_1 \geq \dots \geq \psi_K \geq \psi_0$. The following assumption is taken to guarantee that $\mathcal{S}_K = \mathcal{S}_0$, and thus $\psi_K = \psi_0$.

Assumption 1. For each $j \in [p]$,

$$\min_{j \notin \mathcal{S}_0} |\beta_{0,j}| > \delta_K.$$

The assumption states that our choice of thresholds can distinguish between no signal and

the minimum signal. In the sparse regression literature, this corresponds to the common beta-min condition, which bounds the smallest signal away from zero (cf. Bühlmann and van de Geer, 2011).

We can propose the “variable selection” problem of determining

$$k_0 = \min \left\{ \arg \min_{k \in [K]} \psi_k \right\}.$$

That is, we wish to choose the largest threshold $\delta_{k_0} \in \{\delta_k : k \in [K]\}$ for which $\mathcal{S}_0 \subset \mathcal{S}_{k_0}$, thus yielding a model that excludes the largest possible number of irrelevant variables whilst achieving the minimum limiting risk value.

To estimate k_0 , we construct the information criterion-like estimator

$$\hat{K}_n = \min \left\{ \arg \min_{k \in [K]} \left(\hat{\psi}_{n,k} + P_{n,k} \right) \right\}.$$

Here, for each $k \in [K]$,

$$\hat{\psi}_{n,k} = \min_{\beta \in \mathbb{R}^p} R_{n,k}(\beta; \hat{\beta}_n)$$

is the minimum empirical risk under the k th thresholding transformation and estimator of ψ_k , and $P_{n,k} : \Omega \times \Theta \rightarrow \mathbb{R}$ is its corresponding penalty function, which can be chosen so that

$$\lim_{n \rightarrow \infty} \mathbb{P}(\hat{K}_n = k_0) = 1. \tag{4}$$

For each k and n , we can estimate \mathcal{S}_k by

$$\mathcal{S}_{k,n} = \left\{ j \in [p] : \left| \hat{\beta}_{n,j} \right| \leq \delta_k \right\}.$$

With this notation, our information criterion then yields the estimate $\hat{\mathcal{S}}_n = \mathcal{S}_{\hat{K}_n,n}$ of \mathcal{S}_0 which constitutes the output of our thresholding algorithm and our primary object of interest.

It follows from (4) that $\hat{\mathcal{S}}_n$ satisfies the variable screening property Bühlmann and van de Geer [2011, Sec. 2.7]:

$$\lim_{n \rightarrow \infty} \mathbb{P}(\hat{\mathcal{S}}_n \subset \mathcal{S}_0) = 1.$$

In the sequel, we will demonstrate that $\hat{\mathcal{S}}_n$ is moreover consistent in the sense of (1).

2.1 Thresholding functions

Naively, thresholding of irrelevant coordinates from the signal vector is conducted using the step function (2), as often applied in practice. However, to prove the necessary limit theorems, we require that the functions t_k are sufficiently smooth as per the following assumption.

Assumption 2. For each $k \in [K]$, the thresholding function $t_k : \mathbb{R} \rightarrow [0, 1]$ is twice continuously differentiable and has the form $t_k = \tau_k \circ |\cdot|$, where $\tau_k : \mathbb{R}_{\geq 0} \rightarrow [0, 1]$ is an increasing function, such that $\tau_k(0) = 0$ and for each $b > 0$, $\tau_k(b) > 0$.

For any $h > 0$, define the cubic splines

$$\tau_k(b) = \begin{cases} 0 & \text{if } b \leq \delta_k, \\ \frac{4}{h^3} (x - \delta_k)^3 & \text{if } \delta_k < b \leq \delta_k + \frac{h}{2}, \\ \frac{4}{h^3} (x - \delta_k - h)^3 + 1 & \text{if } \delta_k + \frac{h}{2} < b < \delta_k + h, \\ 1 & \text{if } b \geq \delta_k + h. \end{cases}$$

We can check that $t_k = \tau_k \circ |\cdot|$ satisfy Assumption 2, as required. For the remainder of the manuscript, we will use this spline construction as our choice for thresholding. However, as will be apparent in the sequel, the thresholding function serves as a purely technical device, because in practice, the thresholding procedure can be implemented directly with the step function. We shall elaborate on this point in Section 4.1.1.

2.2 Penalty functions

We will require that the penalty functions satisfy the following assumption.

Assumption 3. For each pair $k, l \in [K]$, $P_{k,n} > 0$ for all sufficiently large n , $P_{k,n} = o_{\mathbb{P}}(1)$, and if $l > k$, then $\sqrt{n}(P_{l,n} - P_{k,n}) \xrightarrow{\mathbb{P}} \infty$.

Let $\alpha : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ be a decreasing function. Then, we can satisfy Assumption 3 by taking

$$P_{n,k} = \alpha(\delta_k) \frac{\log n}{\sqrt{n}}. \quad (5)$$

Penalties $P_{n,k} = O(n^{-1/2} \log n)$ were proposed in Sin and White [1996] and adopted in Nguyen [2024] and Westerhout et al. [2024], where they are referred to as Sin–White information criteria (SWIC). Notice that SWIC differ from the more commonly used BIC-type (Bayesian information criteria; Schwarz, 1978) penalties, which have the form $P_{n,k} = O(n^{-1} \log n)$. Although BIC penalties provide milder penalization of the model complexity, they approach zero too quickly to distinguish between the cases where $\mathcal{S}_k \subset \mathcal{S}_0$, unless stronger assumptions are made. However, as long as $P_{k,n} = o_{\mathbb{P}}(1)$, which is satisfied by the BIC and AIC-type (Akaike information criterion; Akaike, 1974), we are guaranteed at least to have the variable screening property for $\hat{\mathcal{S}}_n$.

We note that there is some room to experiment with the functional form of α , which we will investigate in our numerical studies. However, a principled approach can be taken, such as via some analogy of the “slope heuristic” of Birgé and Massart [2007], as demonstrated in Baudry et al. [2012].

3 Theoretical results

The first step of our analysis is to establish that $\sqrt{n}(\hat{\psi}_{n,k} - \psi_k)$ converges weakly, for each $k \in [K]$. To facilitate our technical exposition, we require the following notation. Let $\Theta \subset \mathbb{R}^p$ be a compact subset, which we can choose to be arbitrarily large as to cause no practical restriction. For each $k \in [K]$, $i \in [n]$, and $\boldsymbol{\beta}, \boldsymbol{\beta}' \in \Theta$, we denote the individual level losses by

$$l_{\boldsymbol{\beta}, \boldsymbol{\beta}'}(\mathbf{X}_i, Y_i) = \{Y_i - \mathbf{X}_i^\top \mathbf{T}_k(\boldsymbol{\beta}; \boldsymbol{\beta}')\}^2,$$

where $l_{\boldsymbol{\beta}, \boldsymbol{\beta}'}(\mathbf{X}_i, Y_i)$ has the same law as $l_{\boldsymbol{\beta}, \boldsymbol{\beta}'}(\mathbf{X}, Y)$. We then define the empirical and expected thresholded risk functions by

$$G_{n,k}(\boldsymbol{\beta}; \boldsymbol{\beta}') = \frac{1}{n} \sum_{i=1}^n l_{\boldsymbol{\beta}, \boldsymbol{\beta}'}(\mathbf{X}_i, Y_i) \text{ and } g_k(\boldsymbol{\beta}; \boldsymbol{\beta}') = \mathbb{E}[l_{\boldsymbol{\beta}, \boldsymbol{\beta}'}(\mathbf{X}, Y)],$$

which allows us to write

$$R_{n,k} = G_{n,k}(\cdot; \hat{\boldsymbol{\beta}}_n) \text{ and } r_k = g_k(\cdot; \boldsymbol{\beta}_0).$$

Since our analysis is the same for each fixed k , we will also find it convenient to remove the index and write G_n and g in place of $G_{n,k}$ and g_k , and R_n and r in place of $R_{n,k}$ and r , respectively, for example.

Consider that the thresholding function $\mathbf{T}(\boldsymbol{\beta}; \boldsymbol{\beta}_0) = \mathbf{D}(\boldsymbol{\beta}_0)\boldsymbol{\beta}$ partitions the coordinates $j \in [p]$ into the set of irrelevant coordinates \mathcal{S} and its complement, the set of relevant coordinates \mathcal{S}^c . Let us write the parameter $\boldsymbol{\beta}$ and design vectors restricted to the relevant coordinates as $\boldsymbol{\beta}_R$ and \mathbf{X}_R , respectively, along with the relevant thresholded expected risk

$$r_R(\boldsymbol{\beta}_R) = \mathbb{E}[\{Y - \mathbf{X}_R \mathbf{D}_R(\boldsymbol{\beta}_0)\boldsymbol{\beta}_R\}^2],$$

where $\mathbf{D}_R(\boldsymbol{\beta}_0)$ is the same as $\mathbf{D}(\boldsymbol{\beta}_0)$ with empty rows and columns removed.

Observe that we can consider $R_n : \Theta \rightarrow \mathbb{R}$ and $r : \Theta \rightarrow \mathbb{R}$ to be elements of the set $\ell^\infty(\Theta)$ of bounded functionals supported on Θ (cf. van der Vaart and Wellner, 2023, Sec. 1.5). As such, we start by showing that $\sqrt{n}(R_n - r)$ converges weakly in $\ell^\infty(\Theta)$, as $n \rightarrow \infty$. Denoting weak convergence by \rightsquigarrow , we make the following assumptions.

Assumption 4. *The variables \mathbf{X} and ϵ have finite fourth moments: i.e., $\mathbb{E}[\|\mathbf{X}\|^4] < \infty$ and $\mathbb{E}[\epsilon^4] < \infty$.*

Assumption 5. *The estimator $\hat{\boldsymbol{\beta}}_n$ converges in distribution to $\boldsymbol{\beta}_0$: i.e.,*

$$\sqrt{n}(\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta}_0) \rightsquigarrow \mathbf{Z},$$

for some random vector $\mathbf{Z} : \Omega \rightarrow \mathbb{R}^p$.

Assumption 4 is a moment condition required for our empirical process weak convergence results, and Assumption 5 is satisfied, for example, whenever $\hat{\beta}_n$ is an asymptotically normal estimator, as in the case of the ordinary least squares (OLS) estimator:

$$\hat{\beta}_n = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{Y}. \quad (6)$$

Let $\beta \mapsto \nabla_2 g(\beta; \beta_0)$ be the gradient of $g(\beta; \beta')$ in β' , evaluated at $\beta' = \beta_0$, as a function of the β . The following theorem is technical but forms the foundation of our subsequent analysis.

Theorem 1. *Under Assumptions 2, 4 and 5,*

$$\sqrt{n}(R_n - r) \rightsquigarrow R + \mathbf{Z}^\top \nabla_2 g(\cdot, \beta_0),$$

where $R : \Omega \rightarrow \ell^\infty(\Theta)$ is a Gaussian process.

The proof of Theorem 1 follows from the theory of random empirical processes and is provided in Appendix A. Intuitively, the theorem states that for each k , when centered by the limiting risk and thresholded by the true signal β_0 , the empirical thresholded least squares converges to a random process in $\ell^\infty(\Theta)$.

Let $\iota : \ell^\infty(\Theta) \rightarrow \mathbb{R}$ denote the infimum function, whereby for each $h : \Theta \rightarrow \mathbb{R}$,

$$\iota(h) = \inf_{\beta \in \Theta} h(\beta).$$

With this notation, we can also define the ε -minimizers of h as the set

$$\mathcal{B}_\varepsilon(h) = \{\beta \in \Theta : h(\beta) \leq \iota(h) + \varepsilon\}.$$

Westerhout et al. [2024, Thm. 3.1] then states that the directional derivative of ι at h in the direction of $\eta \in \ell^\infty(\Theta)$ is

$$i_h(\eta) = \lim_{\varepsilon \searrow 0} \inf_{\beta \in \mathcal{B}_\varepsilon(h)} \eta(\beta).$$

We can apply the directional functional delta method of Römisch [2014] (see also Westerhout et al., 2024, Fact 3.2) to obtain the following result.

Lemma 1. *Under the assumptions of Theorem 1, we can choose a compact $\Theta \subset \mathbb{R}^p$ such that*

$$\sqrt{n}(\iota(R_n) - \psi) \rightsquigarrow F,$$

where $F = i_r(R + \mathbf{Z}^\top \nabla_2 g)$ is a random variable.

Proof. The result follows directly from the directional functional delta method to show that $\sqrt{n}(\iota(R_n) - \iota(r)) \rightsquigarrow F$. It then suffices to check that we can choose a large enough Θ so that

$\iota(r) = \psi$. Indeed, r is quadratic and is minimized on \mathbb{R}^p at any $\boldsymbol{\beta}^*$ that solves the Fermat condition

$$\mathbf{D}(\boldsymbol{\beta}_0) \mathbb{E}[\mathbf{X}Y] = \mathbf{D}(\boldsymbol{\beta}_0) \mathbb{E}[\mathbf{X}\mathbf{X}^\top] \mathbf{D}(\boldsymbol{\beta}_0) \boldsymbol{\beta}^*.$$

The system has potentially infinitely many solutions $\boldsymbol{\beta}^*$, although the relevant coordinates of $\boldsymbol{\beta}^*$ will be the same as the solution

$$\boldsymbol{\beta}_R^* = \{\mathbf{D}_R(\boldsymbol{\beta}_0) \mathbb{E}[\mathbf{X}_R \mathbf{X}_R^\top] \mathbf{D}_R(\boldsymbol{\beta}_0)\}^{-1} \mathbf{D}_R(\boldsymbol{\beta}_0) \mathbb{E}[\mathbf{X}_R Y],$$

the minimizer of r_R on $\mathbb{R}^{|\mathcal{S}^c|}$. In particular, r is minimized at $\boldsymbol{\beta}^{**}$ which equals $\boldsymbol{\beta}_R^*$ on the relevant coordinates and 0 elsewhere. We thus have $\boldsymbol{\beta}^{**} \in [-m, m]^p \subset \Theta$, where $m = \max_{j \in [p]} |\beta_j^{**}|$. \square

Note that since K is finite, a compact Θ large enough so that Lemma 1 holds for all $k \in [K]$ can be constructed as a union over the choice for each k . As a consequence of Lemma 1, the same proof as that of Nguyen [2024, Thm. 1] yields the consistency of \hat{K}_n and thus the variable screening property of $\hat{\mathcal{S}}_n$.

Theorem 2. *Under Assumptions 2–5, we can choose a compact $\Theta \subset \mathbb{R}^p$ such that*

$$\lim_{n \rightarrow \infty} \mathbb{P}(\hat{K}_n = k_0) = 1.$$

The following consistency theorem comprises our main technical result and is a direct consequence of the consistency of \hat{K}_n .

Theorem 3. *Under Assumptions 1–5, we can choose a compact $\Theta \subset \mathbb{R}^p$ such that*

$$\lim_{n \rightarrow \infty} \mathbb{P}(\hat{\mathcal{S}}_n = \mathcal{S}_0) = 1.$$

Proof. We first show that $\mathcal{S}_{k^0} = \mathcal{S}_0$. Under Assumption 1, $\min_{j \notin \mathcal{S}_0} |\beta_{0,j}| > \delta_K$, we have that the least squares loss

$$\tilde{r}(\boldsymbol{\beta}) = \mathbb{E}[\{Y - \mathbf{X}^\top \boldsymbol{\beta}\}^2]$$

has the same minimum as that of the relevant loss function

$$\tilde{r}_R(\boldsymbol{\beta}_R) = \mathbb{E}[\{Y - \mathbf{X}_R^\top \boldsymbol{\beta}_R\}^2],$$

where the relevant set is $\mathcal{S}_0^c = \mathcal{S}_{K^c}^c$. However, we note that that this minimum is also the same as that of the relevant thresholded loss $r_R = r_{K,R}$ since, for each $j \in \mathcal{S}_0^c$, $\beta_j \mapsto \beta_j t_K(\beta_{0,j})$ is surjective because $t_K(\beta_{0,j})$ is positive by Assumption 1 (cf. Hinderer et al., 2016, Sec. A.4.11). If we write the minimizer of \tilde{r}_R as $\tilde{\boldsymbol{\beta}}_R^*$, i.e., the least square solution

$$\tilde{\boldsymbol{\beta}}_R^* = \mathbb{E}[\mathbf{X}_R \mathbf{X}_R^\top]^{-1} \mathbb{E}[\mathbf{X}_R Y],$$

then the minimizer of $r_{\mathbf{R}}$ is simply a scaling of $\tilde{\boldsymbol{\beta}}_{\mathbf{R}}^*$, of the form

$$\boldsymbol{\beta}_{\mathbf{R}}^* = \left[\tilde{\boldsymbol{\beta}}_{\mathbf{R},j}^* / t_K(\beta_{0,j}) \right]_{j \in \mathcal{S}_0^c}.$$

In the same way, for each $k \in [K]$, we can argue that k th relevant thresholded loss $r_{k,\mathbf{R}}$ has the same minimum as that of k th relevant least squares loss

$$\tilde{r}_{k,\mathbf{R}}(\boldsymbol{\beta}_{k,\mathbf{R}}) = \mathbb{E} \left[\left\{ Y - \mathbf{X}_{k,\mathbf{R}}^\top \boldsymbol{\beta}_{k,\mathbf{R}} \right\}^2 \right],$$

defined by the relevant set $\mathcal{S}_k^c = \{j \in [p] : |\beta_{0,j}| > \delta_k\}$. Thus, with the minimizer of $\tilde{r}_{k,\mathbf{R}}$ written as $\tilde{\boldsymbol{\beta}}_{k,\mathbf{R}}^*$, we can obtain the minimizer $\boldsymbol{\beta}_{k,\mathbf{R}}^*$ of $r_{k,\mathbf{R}}$ via the same scaling argument. Finally, since r_k has the same minimum as $r_{k,\mathbf{R}}$, where we can take the minimizer $\boldsymbol{\beta}_k^*$ to have relevant elements equal to $\boldsymbol{\beta}_{k,\mathbf{R}}^*$ and irrelevant elements 0. But since $t_k(\beta_{0,j}) > 0$, for each $k \in [K]$ and $j \in \mathcal{S}_0^c$, we can construct a sufficiently large $\Theta \supset [-m, m]^p$ such that $\boldsymbol{\beta}_k^* \in \Theta$ for all k , by choosing

$$m = \max \left\{ |\beta_{k,j}^*| / t_k(\beta_{0,j}) : j \in [p], k \in [K] \right\},$$

with the convention that $0/0 = 0$. We have thus shown that $\iota(r_k) = \psi_k$ for each $k \in [K]$.

Due to Assumption 1, we have that $\psi_k \geq \psi_K = \psi_0$, with equality if and only if $\mathcal{S}_k \subset \mathcal{S}_K = \mathcal{S}_0$. But by definition of k_0 , it must hold that $\psi_{k_0} = \psi_K$, and thus $\mathcal{S}_{k_0} \subset \mathcal{S}_0$. The fact that $\mathcal{S}_0 \subset \mathcal{S}_{k_0}$ is trivial by observation that $t_k(0) = 0$ for each k , under Assumption 2.

Next, observe that

$$\mathbb{P}(\hat{\mathcal{S}}_n = \mathcal{S}_0) = \mathbb{P}(\mathcal{S}_{\hat{K}_n, n} = \mathcal{S}_{k_0}) \geq \mathbb{P}(\hat{K}_n = k_0, \mathcal{S}_{k_0, n} = \mathcal{S}_{k_0}),$$

and by Bonferonni's inequality,

$$\mathbb{P}(\hat{K}_n = k_0, \mathcal{S}_{k_0, n} = \mathcal{S}_{k_0}) \geq 1 - \mathbb{P}(\hat{K}_n \neq k_0) - \mathbb{P}(\mathcal{S}_{k_0, n} \neq \mathcal{S}_{k_0}).$$

Since $\mathbb{P}(\hat{K}_n = k_0) \rightarrow 1$ by Theorem 2, it suffices to show that $\mathbb{P}(\mathcal{S}_{k_0, n} = \mathcal{S}_{k_0}) \rightarrow 1$. Writing $\mathbf{1}(\mathbf{A})$ as the indicator function of the statement \mathbf{A} , we have that

$$\begin{aligned} \mathbb{P}(\mathcal{S}_{k_0, n} \neq \mathcal{S}_{k_0}) &= \mathbb{P} \left(\bigcup_{j=1}^p \left\{ \mathbf{1}(t_{k_0}(\hat{\beta}_{n,j}) = 0) \neq \mathbf{1}(t_{k_0}(\beta_{0,j}) = 0) \right\} \right) \\ &\leq \sum_{j=1}^p \mathbb{P} \left(\mathbf{1}(t_{k_0}(\hat{\beta}_{n,j}) = 0) \neq \mathbf{1}(t_{k_0}(\beta_{0,j}) = 0) \right) \\ &\leq \sum_{j=1}^p \mathbb{P} \left(\left| |\hat{\beta}_{n,j}| - |\beta_{0,j}| \right| > c \right), \end{aligned}$$

where $c = \min_{j \in [p]} |\delta_{k_0} - |\beta_{0,j}||$. But by Assumption 5, $\hat{\boldsymbol{\beta}}_n \xrightarrow{\mathbb{P}} \boldsymbol{\beta}_0$ and it follows from continuous mapping that

$$\lim_{n \rightarrow \infty} \sum_{j=1}^p \mathbb{P} \left(\left| |\hat{\beta}_{n,j}| - |\beta_{0,j}| \right| > c \right) = 0,$$

which completes the proof. \square

Finally, we return to one of our initial motivations: to justify the ad-hoc procedure of thresholding out “small” estimates of $\hat{\boldsymbol{\beta}}_n$. Using the estimate \hat{K}_n , we define the (hard) thresholded estimator $\bar{\boldsymbol{\beta}}_n$ component-wise by $\bar{\beta}_{n,j} = \hat{\beta}_{n,j} \mathbf{1} \left(t_{\hat{K}_n}(\hat{\beta}_{n,j}) > 0 \right)$, for each $j \in [p]$. This is equivalent to setting the j th coordinate of $\bar{\boldsymbol{\beta}}_n$ to zero if $j \in \hat{\mathcal{S}}_n$ and declaring the elements of all other coordinates equal to the corresponding elements of $\hat{\boldsymbol{\beta}}_n$. Below, we demonstrate a so-called oracle property for $\bar{\boldsymbol{\beta}}_n$, by combining the convergence in distribution of $\hat{\boldsymbol{\beta}}_n$ and the consistency of $\hat{\mathcal{S}}_n$. Using the true irrelevant set \mathcal{S}_0 , we can write $\boldsymbol{\beta}_0 = (\boldsymbol{\beta}_{0,R}, \boldsymbol{\beta}_{0,I}) = (\boldsymbol{\beta}_{0,R}, \mathbf{0})$, where $\boldsymbol{\beta}_I$ denotes the subvector of $\boldsymbol{\beta}$ that consists of only the irrelevant coordinates. The same notation extends to other vectors of length p ; e.g., \mathbf{Z}_R and \mathbf{Z}_I denote the relevant and irrelevant subvectors of \mathbf{Z} , respectively. The following result is proved in Appendix B.

Theorem 4. *Under Assumptions 1–5, $\sqrt{n}(\bar{\boldsymbol{\beta}}_{n,R} - \boldsymbol{\beta}_{0,R}) \rightsquigarrow \mathbf{Z}_R$ and $\lim_{n \rightarrow \infty} \mathbb{P}(\bar{\boldsymbol{\beta}}_{n,I} = \mathbf{0}) = 1$.*

The result of Theorem 4 therefore justifies that the thresholding process is a one-step improvement to any non-sparse estimator that converges in distribution, producing a sparse estimator that has the same asymptotic distribution in the relevant coordinates. Furthermore, if \mathbf{Z} is multivariate normal then so is \mathbf{Z}_R , and thus $\bar{\boldsymbol{\beta}}_{n,R}$ is an asymptotically normal estimator of $\boldsymbol{\beta}_{0,R}$. Observe that this result provides the same oracle guarantees as those for sparse penalization estimators under similar conditions, such as those considered in Fan and Li [2001].

4 Numerical studies

All numerical results were produced in the R programming environment and all codes are made available at https://github.com/karfho/Threshold_IC.

4.1 Simulation studies

4.1.1 Ordinary least squares

We begin by considering the performance of the thresholding procedure in the linear regression setting, with various sample sizes and covariate dimensions. Namely, we consider $n \in \{100, 1000, 10000\}$ and $p \in \{20, 50\}$, where we replicated each simulation scenario 100 times. For each n and p , we simulate covariates $(\mathbf{X}_i)_{i \in [n]}$ IID with the same distribution as $\mathbf{X} \sim N(0, \boldsymbol{\Sigma})$, and we simulate errors $(\epsilon_i)_{i \in [n]}$ IID with standard normal distribution. The covariate matrix is taken

to have 1s along the diagonal and $1/5$ in all off-diagonal elements. This setup is representative of mildly correlated covariate scenarios.

We generate $(Y_i)_{i \in [n]}$ as $Y_i = \mathbf{X}_i^\top \boldsymbol{\beta}_0 + \epsilon_i$, where we consider two scenarios for $\boldsymbol{\beta}_0$:

$$\text{S1} \quad \boldsymbol{\beta}_0 = (0.2, 0.4, \dots, 1.8, 2.0, 0, \dots, 0),$$

$$\text{S2} \quad \boldsymbol{\beta}_0 = (0.05, 0.1, \dots, 0.45, 0.5, 0, \dots, 0),$$

with S1 and S2 corresponding to high and low signal-to-noise ratio scenarios, respectively. Note that in both cases, the set of relevant coordinates is $\mathcal{S}_0^c = [10]$, which is independent of both p and n . For each simulation replication, we estimate $\boldsymbol{\beta}_0$ by the OLS (6): $\hat{\boldsymbol{\beta}}_n$, set $K = p$, and use threshold values

$$(\delta_k)_{k \in [K]} = \left(\left| \hat{\beta}_{n,(j)} \right| \right)_{j \in [p]}, \quad (7)$$

where $\left(\hat{\beta}_{n,(j)} \right)_{j \in [p]}$ are the elements of the estimator $\hat{\boldsymbol{\beta}}_n$, ordered from largest to smallest: i.e., $\left| \hat{\beta}_{n,(1)} \right| < \left| \hat{\beta}_{n,(2)} \right| \cdots < \left| \hat{\beta}_{n,(p)} \right|$. We note that although our theory characterizes the behavior under a deterministic sequence $(\delta_k)_{k \in [K]}$, the choice is sensible because the procedure will threshold the signals at exactly where the increase in maximum signal magnitude of the relevant set occurs, which allows us to consider all possible leveled sets with minimal computation. However, if we operate with a predetermined set of thresholding values, then it is possible to miss out on certain leveled sets if multiple components of $\left| \hat{\beta}_n \right|$ lie in between some $[\delta_k, \delta_{k+1})$, and if no component falls in between $[\delta_k, \delta_{k+1})$, recomputing the minimal empirical risk is simply redundant.

To satisfy Assumption 3, we noted that one can take penalties of SWIC form: (5). For our purposes, we will consider a further restriction to the case where

$$\alpha(\delta_k) = \frac{c}{\delta_k^r}, \quad (8)$$

for some choices of $c, r > 0$. In particular, to represent weak, medium, and strong penalties, respectively, we take $(c, r) = (0.5, 0.25)$, $(0.75, 0.4)$, and $(1, 0.5)$.

To measure the performance of our method, in the face of our theoretical result that \hat{K}_n exhibits model selection consistency, we report the average (over replications) estimated threshold value $\delta_{\hat{K}_n}$, average percentage of covariates that are relevant that are incorrectly estimated to be irrelevant, and the average percentage of irrelevant covariates that are correctly estimated to be irrelevant. The two latter indices can be characterized as the average false negative rate (FNR) and average true negative rate (TNR), respectively, as percentages where

$$\text{FNR} = 1 - \frac{|\hat{\mathcal{S}}_n^c \cap \mathcal{S}_0^c|}{|\mathcal{S}_0^c|}, \text{ and } \text{TNR} = \frac{|\hat{\mathcal{S}}_n \cap \mathcal{S}_0|}{|\mathcal{S}_0|}.$$

We report the outcomes in Tables 1 and 2.

Table 1: Results for S1 using OLS estimators. All averages are computed over 100 replications. Note that $\delta_{k_0} < 0.2$.

Measure	n	p	(c, r)		
			(0.5, 0.25)	(0.75, 0.4)	(1, 0.5)
$\delta_{\hat{K}_n}$	100	20	0.0995	0.1726	0.2181
	1000	20	0.06332	0.09462	0.1606
	10000	20	0.0207	0.0207	0.0207
	100	50	0.07655	0.1217	0.1729
	1000	50	0.06087	0.121	0.156
	10000	50	0.02739	0.02739	0.02739
$\text{FNR} \times 100\%$	100	20	2.8	6.3	10.2
	1000	20	0.3	3.3	7.8
	10000	20	0	0	0
	100	50	2.1	3.3	4.1
	1000	50	0.2	4.2	7
	10000	50	0	0	0
$\text{TNR} \times 100\%$	100	20	64.8	87	96.4
	1000	20	99.7	100	100
	10000	20	100	100	100
	100	50	38.075	55.85	70.425
	1000	50	84.75	100	100
	10000	50	100	100	100

Table 2: Results for S2 using OLS estimators. All averages are computed over 100 replications. Note that $\delta_{k_0} < 0.05$.

Measure	n	p	(c, r)		
			(0.5, 0.25)	(0.75, 0.4)	(1, 0.5)
$\delta_{\hat{K}_n}$	100	20	0.08472	0.1413	0.17
	1000	20	0.0624	0.09443	0.12
	10000	20	0.04549	0.0676	0.08767
	100	50	0.07323	0.1143	0.1474
	1000	50	0.04186	0.09816	0.1262
	10000	50	0.04349	0.06519	0.09206
$\text{FNR} \times 100\%$	100	20	16.9	27.7	34.2
	1000	20	11	20.1	25.3
	10000	20	8.8	14.3	18
	100	50	11.4	19	23.4
	1000	50	5.4	18.8	27.3
	10000	50	8.2	13.6	18.8
$\text{TNR} \times 100\%$	100	20	56.2	78.3	85.2
	1000	20	96.2	100	100
	10000	20	100	100	100
	100	50	36.85	53.5	64.7
	1000	50	74.325	99.675	100
	10000	50	100	100	100

The results of Tables 1 and 2 appear to indicate that the thresholding method usefully performs variable selection, especially for larger values of n . As expected, we observe that weaker penalization yields more conservative thresholding, and thus lowers the FNR, although at the expense of decreased TNR, and vice versa for stronger penalization. Supporting the asymptotic theory, we observe that both $(1 - \text{TNR})$ and FNR decrease as n increases, and similarly, the optimal selected thresholds $\delta_{\hat{K}_n}$ get closer to the upper bound of δ_{k_0} . Comparing Table 2 to Table 1, we observe, as expected that the overall performance of the thresholding method declines when the signal is weaker, and thus for comparable performance, one requires larger sample sizes. Interestingly, we note that the performance in TNR does not decline as severely as that for FNR. With better calibration of the penalty function α , one may anticipate better overall performance in each scenario, although we know of no general and theoretically supported approach for such a task in our setting.

4.1.2 Ridge and adaptive ridge estimators

As an alternative to OLS, we may consider instead taking $\hat{\beta}_n$ to be a ridge regression-type estimator, which naturally benefits from the use of a thresholding scheme. A noted shortcoming of l_2 regularization approaches, such as ridge regression and adaptive ridge regression, is that the shrinkage estimators are not sparse, like the LASSO, but instead merely shrink weak signals arbitrarily close to zero. As such, many practitioners have taken to simply conduct arbitrary thresholding of small signals as an ad hoc approach to variable selection, which our thresholding approach now makes rigorous.

In this section, we consider the same simulation as in Section 4.1.1, but instead using the classic ridge estimator [Hoerl and Kennard, 1970] and the adaptive ridge (AR) estimator of Dai et al. [2018] and Ho and Masuda [2024]. In particular, for tuning parameters $\lambda_n, \xi_n > 0$, we define the ridge estimator as

$$\hat{\beta}_{\text{Ridge},n} = (\mathbf{X}^\top \mathbf{X} + \lambda_n \mathbf{I})^{-1} \mathbf{X}^\top \mathbf{Y},$$

and for $s \in \mathbb{N}$, we define the AR estimator at the s th step by

$$\hat{\beta}_{\text{AR},n}^{(s)} = \left(\mathbf{X}^\top \mathbf{X} + \xi_n \mathfrak{D} \left(\hat{\beta}_{\text{AR},n}^{(s-1)} \right) \right)^{-1} \mathbf{X}^\top \mathbf{Y},$$

where $\mathfrak{D}(\beta) = \text{diag}(1/\beta_1^2, \dots, 1/\beta_p^2)$, and $\hat{\beta}_{\text{AR},n}^{(0)} = \hat{\beta}_{\text{Ridge},n}$. Using $\lambda_n = \sqrt{n}$ and $\xi_n = 1$, we compare the ridge and AR estimator, with $s = 5$, to the OLS for S1 when $n = 100$ and S2 when $n = 1000$, with results reported in Tables 3 and 4, respectively.

From Tables 3 and 4, we observe that the OLS and ridge-type estimators perform similarly, although there appear to be slight improvements when using the ridge-type estimators. We note that since ridge and AR estimators conduct shrinkage, more signals tend to be eliminated under each threshold, and thus the thresholding algorithm is better at eliminating irrelevant signals.

Table 3: Comparison of performance between OLS, ridge, and AR estimators for scenario S1 with $n = 100$. Note that $\delta_{k_0} < 0.2$ and $s = 5$.

Method	Measure	(c, r)		
		(0.5, 0.25)	(0.75, 0.4)	(1, 0.5)
OLS	$\delta_{\hat{K}_n}$	0.0995	0.1726	0.2181
	FNR $\times 100\%$	2.8	6.3	10.2
	TNR $\times 100\%$	64.8	87	96.4
Ridge	$\delta_{\hat{K}_n}$	0.1213	0.1962	0.2335
	FNR $\times 100\%$	2.7	5.9	9.1
	TNR $\times 100\%$	69.2	91	96.8
AR	$\delta_{\hat{K}_n}$	0.04553	0.09792	0.1506
	FNR $\times 100\%$	6	8.9	11.9
	TNR $\times 100\%$	94.2	97.2	98.7

Table 4: Comparison of performance between OLS, ridge, and AR estimators for scenario S2 with $n = 1000$. Note that $\delta_{k_0} < 0.05$ and $s = 5$.

Method	Measure	(c, r)		
		(0.5, 0.25)	(0.75, 0.4)	(1, 0.5)
OLS	$\delta_{\hat{K}_n}$	0.04186	0.09816	0.1262
	FNR $\times 100\%$	5.4	18.8	27.3
	TNR $\times 100\%$	74.325	99.675	100
Ridge	$\delta_{\hat{K}_n}$	0.04006	0.09563	0.1225
	FNR $\times 100\%$	5	18.9	27.3
	TNR $\times 100\%$	74.05	99.725	100
AR	$\delta_{\hat{K}_n}$	0.05081	0.09292	0.1174
	FNR $\times 100\%$	10.5	21.8	27.7
	TNR $\times 100\%$	96.825	99.975	100

Table 5: Sets of relevant coordinates $\hat{\mathcal{S}}_n$ obtained from thresholding of prostate cancer data using OLS, ridge, and AR estimators.

Method	(c, r)		
	(0.5, 0.25)	(0.75, 0.4)	(1, 0.5)
OLS	{1, 2, 3, 4, 5, 6, 8}	{1, 2, 5}	{1}
Ridge	{1, 2, 5}	{1, 2, 5}	{1}
AR	{1, 2, 5}	{1}	{1}

Table 6: Size of relevant covariate sets $\hat{\mathcal{S}}_n$ obtained from thresholding of prostate cancer data using OLS, ridge, and AR estimators, with pairwise interactions.

Method	(c, r)		
	(0.5, 0.25)	(0.75, 0.4)	(1, 0.5)
OLS	15	14	14
Ridge	30	14	2
AR	3	1	1

This can be seen in the improvement of TNR in both tables. In particular, we observe that the AR estimator conducts very aggressive shrinkage, causing much higher TNR than both OLS and ridge estimators, even when the penalty is small.

4.2 Prostate cancer data example

To illustrate the application of our method in practice, we consider the prostate data set of Stamey et al. [1989]. These data consist of $n = 97$ observations, each with covariates consisting of $p = 8$ clinical measurements from individual patients: the logarithm of cancer volume (X_1), logarithm of weight (X_2), age (X_3), logarithm of the amount of benign prostatic hyperplasia (X_4), seminal vesicle invasion (X_5), logarithm of capsular penetration (X_6), Gleason score (X_7), and percentage Gleason scores of four or five (X_8). The response variable is the logarithm of prostate-specific antigen (Y).

Following typical preprocessing protocols, we standardize the data and, with $K = 8$, we apply our thresholding method with $(\delta_k)_{k \in [K]}$ chosen as per (7), using SWIC penalties with α functions (8), taking $(c, r) = (0.5, 0.25)$, $(0.75, 0.4)$, and $(1, 0.5)$, as in the simulations. The OLS, ridge and AR estimators are used with $\lambda_n = \sqrt{n}$, $\xi_n = 1$, and $s = 5$, as per Section 4.1.2. We report the sets of relevant covariates obtained via the thresholding algorithm $\hat{\mathcal{S}}_n^c$ in Table 5. Further, expanding the set of covariates by also considering pairwise interactions, yielding a total of $p = 36$ predictors, we report the sizes of the sets of relevant covariates obtained from thresholding $|\hat{\mathcal{S}}_n^c|$ in Table 6.

From both Tables 5 and 6, we observe as expected, that stronger penalization implies less variables being declared relevant, regardless of initial estimator used, while in all but the first column of Table 6, we observe that the hierarchy of shrinkage from OLS, ridge, and AR estimators

yields decreasing numbers of relevant covariates, as anticipated. From Table 5, we conclude that the important variables are X_1, X_2, X_5 , with X_1 appearing to be the most relevant of all covariates. This corroborates with the outcomes from Zou and Hastie [2005], Lin [2010] and Sun et al. [2013], which identify the same variables as being most important.

5 Conclusion

Variable selection is a crucial step in many modern inference workflows, with shrinkage estimators forming a useful class of methods for identifying important covariates in the sparse signal setting. We have introduced a systematic approach that takes any initial regression estimator that exhibits convergence in distribution and outputs a consistent estimator of the set of relevant covariates, under practical regularity conditions. Our numerical studies demonstrate that our approach performs as theoretically anticipated, and functions well using OLS, ridge, and AR estimators used as inputs, although our theory suggests that the class of input estimators whose performance are guaranteed by theory is far larger.

We anticipate three immediate avenues for future research. First, to permit applications to infinite dimensional settings, we need to allow for K to be infinitely large, either countably or uncountably. It is known that in some nested regression settings, model selection procedures with uncountably large K are possible (see, e.g., Zhang and Nguyen, 2024), although the situations are quite different and we do not anticipate the proof techniques to transfer without additional technical development. Secondly, we have developed our theory in the linear regression setting for ease of exposition and to keep the scope manageable. However, with necessary modifications, there is little else that inhibits the application of our approach to the generalized linear model, nonlinear regression, linear functional regression, or other parametric regression modeling settings where the notion of variable selection applies. Thirdly, our current theory makes few restrictions on the class of allowable penalty functions, and in effect, makes no recommendations regarding the design of an efficient penalty function. Similar questions of optimal penalization appears in the setting of model selection oracle bounds, manifesting as the so called slope heuristic omnibus procedure (cf. Birgé and Massart, 2007, Baudry et al., 2012, and Arlot, 2019). Although the breadth of application of the slope heuristic does not apply in our asymptotic setting, it is possible that an analogous procedure for automatic penalty calibration exists.

Acknowledgments

KLKH is supported by the GPMI program at Kyushu University. HDN is funded by Australian Research Council grants: DP230100905 and DP250100860.

Appendix

A Proof of Theorem 1

Following the decomposition of van der Vaart and Wellner [2023, Sec. 3.13], we begin by writing:

$$\begin{aligned} \sqrt{n} \left\{ G_n(\boldsymbol{\beta}; \hat{\boldsymbol{\beta}}_n) - g(\boldsymbol{\beta}; \boldsymbol{\beta}_0) \right\} &= \sqrt{n} \left[\left\{ G_n(\boldsymbol{\beta}; \hat{\boldsymbol{\beta}}_n) - g(\boldsymbol{\beta}; \hat{\boldsymbol{\beta}}_n) \right\} - \left\{ G_n(\boldsymbol{\beta}; \boldsymbol{\beta}_0) - g(\boldsymbol{\beta}; \boldsymbol{\beta}_0) \right\} \right] \\ &\quad + \sqrt{n} \left\{ G_n(\boldsymbol{\beta}; \boldsymbol{\beta}_0) - g(\boldsymbol{\beta}; \boldsymbol{\beta}_0) \right\} + \sqrt{n} \left\{ g(\boldsymbol{\beta}; \hat{\boldsymbol{\beta}}_n) - g(\boldsymbol{\beta}; \boldsymbol{\beta}_0) \right\} \\ &= (1A) + (1B) + (1C). \end{aligned}$$

Our program is to prove the weak convergence of terms (1B) and (1C) to R and $\mathbf{Z}^\top \nabla_2 g(\cdot, \boldsymbol{\beta}_0)$, respectively, and to demonstrate that (1A) = $o_{\mathbb{P}}(1)$. The following result follows from van der Vaart and Wellner [2023, Thm. 3.13.4].

Proposition 1. *Under Assumptions 2, 4, and 5, it holds that*

$$\sup_{\boldsymbol{\beta} \in \Theta} \left| \sqrt{n} \left[\left\{ G_n(\boldsymbol{\beta}; \hat{\boldsymbol{\beta}}_n) - g(\boldsymbol{\beta}; \hat{\boldsymbol{\beta}}_n) \right\} - \left\{ G_n(\boldsymbol{\beta}; \boldsymbol{\beta}_0) - g(\boldsymbol{\beta}; \boldsymbol{\beta}_0) \right\} \right] \right| \xrightarrow{\mathbb{P}} 0.$$

By van der Vaart and Wellner [2023, Thm. 3.13.4], it suffices to check that the set

$$\mathcal{L} = \left\{ l_{\boldsymbol{\beta}, \boldsymbol{\beta}'}(\mathbf{X}, Y) = \{Y - \mathbf{X}^\top \mathbf{T}_k(\boldsymbol{\beta}; \boldsymbol{\beta}')\}^2 : (\boldsymbol{\beta}, \boldsymbol{\beta}') \in \Theta \times \Theta \right\}$$

is \mathbb{P} -Donsker and it holds that

$$\bar{\Delta}_n = \sup_{\boldsymbol{\beta} \in \Theta} P \left[\left\{ l_{\boldsymbol{\beta}, \hat{\boldsymbol{\beta}}_n}(\mathbf{X}, Y) - l_{\boldsymbol{\beta}, \boldsymbol{\beta}_0}(\mathbf{X}, Y) \right\}^2 \right] \xrightarrow{\mathbb{P}} 0,$$

where we use P to denote the expectation only over (\mathbf{X}, Y) and not $\hat{\boldsymbol{\beta}}_n$. To verify that \mathcal{L} is \mathbb{P} -Donsker, we apply Shapiro et al. [2021, Prop. 9.79] which requires that we check the conditions that (i) $\mathbb{E} [l_{\boldsymbol{\beta}_1, \boldsymbol{\beta}'_1}(\mathbf{X}, Y)^2] < \infty$ and (ii) there exists a function $\bar{L} : \mathbb{R}^{p+1} \rightarrow \mathbb{R}$ with $\mathbb{E} [\bar{L}(\mathbf{X}, Y)^2] < \infty$, where

$$|l_{\boldsymbol{\beta}_1, \boldsymbol{\beta}'_1}(\mathbf{X}, Y) - l_{\boldsymbol{\beta}_2, \boldsymbol{\beta}'_2}(\mathbf{X}, Y)| \leq \bar{L}(\mathbf{X}, Y) [\|\boldsymbol{\beta}_1 - \boldsymbol{\beta}_2\| + \|\boldsymbol{\beta}'_1 - \boldsymbol{\beta}'_2\|],$$

for every $\boldsymbol{\beta}_1, \boldsymbol{\beta}_2, \boldsymbol{\beta}'_1, \boldsymbol{\beta}'_2 \in \Theta$.

To check (i), we start by writing

$$\begin{aligned}
l_{\boldsymbol{\beta}, \boldsymbol{\beta}'}(\mathbf{X}, Y) &= \{Y - \mathbf{X}^\top \mathbf{T}_k(\boldsymbol{\beta}; \boldsymbol{\beta}')\}^2 \\
&= \{\mathbf{X}^\top [\boldsymbol{\beta}_0 - \mathbf{T}_k(\boldsymbol{\beta}; \boldsymbol{\beta}')] + \epsilon\}^2 \\
&= \left\{ \sum_{j=1}^p X_j \beta_{0j} - \sum_{j=1}^p X_j t_k(\beta'_j) \beta_j + \epsilon \right\}^2
\end{aligned}$$

and thus

$$\begin{aligned}
\mathbb{E} [l_{\boldsymbol{\beta}_1, \boldsymbol{\beta}'_1}(\mathbf{X}, Y)^2] &= \mathbb{E} \left[\{\mathbf{X}^\top [\boldsymbol{\beta}_0 - \mathbf{T}_k(\boldsymbol{\beta}_1; \boldsymbol{\beta}'_1)] + \epsilon\}^4 \right] \\
&\leq (1+p) \left\{ \mathbb{E} [|\epsilon|^4] + \sum_{j=1}^p (|\beta_{0j}| + t_k(\beta'_{1j}) |\beta_{1j}|)^4 \mathbb{E} [|X_j|^4] \right\}, \\
&\leq (1+p) \left\{ \mathbb{E} [|\epsilon|^4] + 16B^4 \sum_{j=1}^p \mathbb{E} [|X_j|^4] \right\},
\end{aligned}$$

where $B = \sup_{\boldsymbol{\beta} \in \Theta} \|\boldsymbol{\beta}\|_1$, since $0 \leq t_k \leq 1$. Thus, under the moment condition of Assumption 4, (i) is verified. Next, observe that the derivative of l in $(\boldsymbol{\beta}, \boldsymbol{\beta}')$ is

$$\nabla l_{\boldsymbol{\beta}, \boldsymbol{\beta}'}(\mathbf{X}, Y) = -2 \left(\sum_{j=1}^p X_j \beta_{0j} - \sum_{j=1}^p X_j t_k(\beta'_j) \beta_j + \epsilon \right) (\dots, X_j t_k(\beta'_j), \dots, X_j \beta_j \dot{t}_k(\beta'_j), \dots)^\top,$$

where \dot{t}_k is the derivative of t_k , which exists and is continuous due to Assumption 2. Thus, we can write

$$\|\nabla l_{\boldsymbol{\beta}, \boldsymbol{\beta}'}(\mathbf{X}, Y)\|_1 \leq C \sum_{j=1}^p \sum_{j'=1}^p a_{j,j'}(\boldsymbol{\beta}, \boldsymbol{\beta}') |X_j X_{j'}| + \sum_{j=1}^p b_j(\boldsymbol{\beta}, \boldsymbol{\beta}') |X_j \epsilon|, \quad (9)$$

for some large constant $C < \infty$, and continuous functions $a_{j,j'}, b_j : \Theta \times \Theta \rightarrow \mathbb{R}_{\geq 0}$. Since Θ is compact, we can therefore bound (9) by the supremum over $\Theta \times \Theta$ which exists and is finite by the Weierstrass extreme value theorem. Thus, there is a large constant $C < \infty$ such that

$$\|\nabla l_{\boldsymbol{\beta}, \boldsymbol{\beta}'}(\mathbf{X}, Y)\|_1 \leq C \left\{ \sum_{j=1}^p \sum_{j'=1}^p |X_j X_{j'}| + \sum_{j=1}^p |X_j \epsilon| \right\}.$$

But by the mean-value theorem, since $l_{\boldsymbol{\beta}, \boldsymbol{\beta}'}$ (Y, \mathbf{X}) is continuously differentiable and Θ is compact, we can write

$$|l_{\boldsymbol{\beta}_1, \boldsymbol{\beta}'_1}(\mathbf{X}, Y) - l_{\boldsymbol{\beta}_2, \boldsymbol{\beta}'_2}(\mathbf{X}, Y)| \leq \bar{L}(\mathbf{X}, Y) [\|\boldsymbol{\beta}_1 - \boldsymbol{\beta}_2\| + \|\boldsymbol{\beta}'_1 - \boldsymbol{\beta}'_2\|],$$

for each $\boldsymbol{\beta}_1, \boldsymbol{\beta}_2, \boldsymbol{\beta}'_1, \boldsymbol{\beta}'_2 \in \Theta$, where $\bar{L}(\mathbf{X}, Y) = C \|\nabla l_{\boldsymbol{\beta}, \boldsymbol{\beta}'}(\mathbf{X}, Y)\|_1$, for some large $C < \infty$, and

therefore we can take

$$\bar{L}(\mathbf{X}, Y) = C \left\{ \sum_{j=1}^p \sum_{j'=1}^p |X_j X_{j'}| + \sum_{j=1}^p |X_j \epsilon| \right\}.$$

Clearly, we have that for a constant $C < \infty$,

$$\bar{L}(\mathbf{X}, Y)^2 \leq C \left\{ \sum_{j=1}^p \sum_{j'=1}^p |X_j X_{j'}|^2 + \sum_{j=1}^p |X_j \epsilon|^2 \right\}.$$

Assumption 4 then implies that

$$\mathbb{E} [\bar{L}(\mathbf{X}, Y)^2] \leq C \left\{ \sum_{j=1}^p \sum_{j'=1}^p \mathbb{E} [|X_j X_{j'}|^2] + \sum_{j=1}^p [\mathbb{E} |X_j \epsilon|^2] \right\} < \infty,$$

and thus (ii) holds, implying that \mathcal{L} is \mathbb{P} -Donsker.

Next, to check that $\bar{\Delta}_n \xrightarrow{\mathbb{P}} 0$, we define

$$\Delta_n(\boldsymbol{\beta}) = P \left[\left\{ l_{\boldsymbol{\beta}, \hat{\boldsymbol{\beta}}_n}(\mathbf{X}, Y) - l_{\boldsymbol{\beta}, \boldsymbol{\beta}_0}(\mathbf{X}, Y) \right\}^2 \right].$$

We require pair of results from Davidson [2021, Thms. 22.9 and 22.10], which we specify in the context of our problem.

Lemma 2. *If $\Delta_n(\boldsymbol{\beta}) \xrightarrow{\mathbb{P}} 0$ for each $\boldsymbol{\beta} \in \Theta$ and (Δ_n) is stochastic equicontinuous, then $\bar{\Delta}_n \xrightarrow{\mathbb{P}} 0$.*

Lemma 3. *For sufficiently large $n \in \mathbb{N}$, if*

$$|\Delta_n(\boldsymbol{\beta}) - \Delta_n(\boldsymbol{\beta}')| \leq \bar{D}_n \|\boldsymbol{\beta} - \boldsymbol{\beta}'\|,$$

for each $\boldsymbol{\beta}, \boldsymbol{\beta}' \in \Theta$, where $\bar{D}_n = O_{\mathbb{P}}(1)$, then (Δ_n) is stochastic equicontinuous.

We begin by checking the first condition of Lemma 2. Let us write:

$$\Delta_n(\boldsymbol{\beta}) = P \left[l_{\boldsymbol{\beta}, \hat{\boldsymbol{\beta}}_n}(\mathbf{X}, Y)^2 \right] - 2P \left[l_{\boldsymbol{\beta}, \hat{\boldsymbol{\beta}}_n}(\mathbf{X}, Y) l_{\boldsymbol{\beta}, \boldsymbol{\beta}_0}(\mathbf{X}, Y) \right] + P \left[l_{\boldsymbol{\beta}, \boldsymbol{\beta}_0}(\mathbf{X}, Y)^2 \right].$$

But since $P \left[l_{\boldsymbol{\beta}, \boldsymbol{\beta}'}(\mathbf{X}, Y)^2 \right] = \mathbb{E} \left[l_{\boldsymbol{\beta}, \boldsymbol{\beta}'}(\mathbf{X}, Y)^2 \right] < \infty$ uniformly in $(\boldsymbol{\beta}, \boldsymbol{\beta}')$, as already checked, we have that $P \left[l_{\boldsymbol{\beta}, \boldsymbol{\beta}'}(\mathbf{X}, Y)^2 \right]$ is a continuous function in $(\boldsymbol{\beta}, \boldsymbol{\beta}')$, by Shapiro et al. [2021, Thm. 9.55]. We can similarly check that the middle term is continuous. Since $\hat{\boldsymbol{\beta}}_n$ is a consistent estimator by Assumption 5, we have that

$$\Delta_n(\boldsymbol{\beta}) \xrightarrow{\mathbb{P}} P \left[l_{\boldsymbol{\beta}, \boldsymbol{\beta}_0}(\mathbf{X}, Y)^2 \right] - 2P \left[l_{\boldsymbol{\beta}, \boldsymbol{\beta}_0}(\mathbf{X}, Y) l_{\boldsymbol{\beta}, \boldsymbol{\beta}_0}(\mathbf{X}, Y) \right] + P \left[l_{\boldsymbol{\beta}, \boldsymbol{\beta}_0}(\mathbf{X}, Y)^2 \right] = 0,$$

for each β , as required. To apply Lemma 3, we observe that we can write

$$|\Delta_n(\beta) - \Delta_n(\beta')| \leq (2A) + (2B) + (2C),$$

where

$$\begin{aligned} (2A) &= \left| P \left[l_{\beta, \hat{\beta}_n}(\mathbf{X}, Y)^2 \right] - P \left[l_{\beta', \hat{\beta}_n}(\mathbf{X}, Y)^2 \right] \right|, \\ (2B) &= 2 \left| P \left[l_{\beta, \hat{\beta}_n}(\mathbf{X}, Y) l_{\beta', \beta_0}(\mathbf{X}, Y) \right] - P \left[l_{\beta', \hat{\beta}_n}(\mathbf{X}, Y) l_{\beta', \beta_0}(\mathbf{X}, Y) \right] \right|, \\ (2C) &= \left| P \left[l_{\beta, \beta_0}(\mathbf{X}, Y)^2 \right] - P \left[l_{\beta', \beta_0}(\mathbf{X}, Y)^2 \right] \right|. \end{aligned}$$

We can thus verify the condition of Lemma 3 for each component. We will do so using the mean value theorem. In the case of (2A) and (2C), this requires that $P \left[l_{\beta, \beta'}(\mathbf{X}, Y)^2 \right]$ is differentiable in β , which holds if it satisfies the Lipschitz condition:

$$|l_{\beta_1, \beta'}(\mathbf{X}, Y)^2 - l_{\beta_2, \beta'}(\mathbf{X}, Y)^2| \leq C(\mathbf{X}, Y) \|\beta_1 - \beta_2\|,$$

for each $\beta_1, \beta_2, \beta' \in \Theta$, where $\mathbb{E}[C(\mathbf{X}, Y)] < \infty$ and if $l_{\beta, \beta'}(\mathbf{X}, Y)^2$ is differentiable for each (\mathbf{X}, Y) (cf. Shapiro et al. 2021, Thm. 9.56). Differentiability follows since $l_{\beta, \beta'}(\mathbf{X}, Y)^2$ is quadratic in β . To verify the Lipschitz condition, we follow the same approach as when checking the \mathbb{P} -Donsker property and write

$$l_{\beta, \beta'}(\mathbf{X}, Y)^2 = \left\{ \sum_{j=1}^p X_j \beta_{0j} - \sum_{j=1}^p X_j t_k(\beta'_j) \beta_j + \epsilon \right\}^4,$$

where the derivative with respect to β is

$$\nabla_1 [l_{\beta, \beta'}(\mathbf{X}, Y)^2] = \left[4X_j t_k(\beta'_j) \left(\sum_{j=1}^p X_j \beta_{0j} - \sum_{j=1}^p X_j t_k(\beta'_j) \beta_j + \epsilon \right)^3 \right]_{j \in [p]},$$

therefore, since Θ is compact, and the highest power of X_j and ϵ is 4, the extreme value theorem and Holder's inequality then yields a positive function $a : \mathbb{R}^{2+p} \rightarrow \mathbb{R}_{\geq 0}$ that is linear in its first two inputs, such that

$$\|\nabla_1 [l_{\beta, \beta'}(\mathbf{X}, Y)^2]\|_1 \leq a(\|\mathbf{X}\|^4, \epsilon^4, \beta').$$

The mean value theorem then suggests that we can take $C(\mathbf{X}, Y) = C \times a(\|\mathbf{X}\|^4, \epsilon^4, \beta')$, for some $C < \infty$, where $\mathbb{E}[a(\|\mathbf{X}\|^4, \epsilon^4, \beta')] < \infty$, as required, by Assumption 4. Shapiro et al.

[2021, Thm. 9.56] then implies that

$$\begin{aligned} \nabla_1 P [l_{\beta, \beta'} (\mathbf{X}, Y)^2] &= P [\nabla_1 \{l_{\beta, \beta'} (\mathbf{X}, Y)^2\}] \\ &= \left[4P \left\{ X_j t_k (\beta'_j) \left(\sum_{j=1}^p X_j \beta_{0j} - \sum_{j=1}^p X_j t_k (\beta'_j) \beta_j + \epsilon \right)^3 \right\} \right]_{j \in [p]}. \end{aligned} \quad (10)$$

The extreme value theorem and Holder's inequality again yield a positive continuous function $a : \mathbb{R}^{2+p} \rightarrow \mathbb{R}_{\geq 0}$, such that

$$\|\nabla_1 P [l_{\beta, \beta'} (\mathbf{X}, Y)^2]\|_1 \leq a (\mathbb{E} \|\mathbf{X}\|^4, \mathbb{E} [\epsilon^4], \beta'),$$

where the continuity is due to Assumption 2. By the mean value theorem and the compactness of Θ , we can choose

$$\bar{D} = C \times \sup_{\beta' \in \Theta} a (\mathbb{E} \|\mathbf{X}\|^4, \mathbb{E} [\epsilon^4], \beta') < \infty,$$

such that

$$(2A), (2C) \leq \bar{D} \|\beta - \beta'\|,$$

for each $\beta, \beta' \in \Theta$, as required. The Lipschitz condition for (2B) follows similarly, thus Lemma 3 yields the equicontinuity of (Δ_n) and therefore $\bar{\Delta}_n \xrightarrow{\mathbb{P}} 0$ follows from Lemma 2, thus proving Proposition 1.

Next, we verify the weak convergence of (1B). Indeed, since \mathcal{L} is \mathbb{P} -Donsker, and

$$\mathcal{L}_{\beta_0} = \left\{ l_{\beta, \beta_0} (\mathbf{X}, Y) = \{Y - \mathbf{X}^\top \mathbf{T}_k (\beta; \beta_0)\}^2 : \beta \in \Theta \right\} \subset \mathcal{L},$$

we have that \mathcal{L}_{β_0} is also \mathbb{P} -Donsker by van der Vaart and Wellner [2023, Thm. 2.10.1]. Thus, by definition of a \mathbb{P} -Donsker class (cf. van der Vaart and Wellner, 2023, Sec. 2.1), we have our required result.

Proposition 2. *Under Assumptions 2 and 4, (1B) $\rightsquigarrow R$, where $R : \Omega \rightarrow \ell^\infty (\Theta)$.*

It remains to show that (1C) $\rightsquigarrow \mathbf{Z}^\top \nabla_2 g(\cdot, \beta_0)$. To this end, we follow the suggestion from van der Vaart and Wellner [2023, Sec. 3.13] and apply the delta method, to the function $\beta' \mapsto g(\cdot; \beta') \in \ell^\infty (\Theta)$. We will again use the directional functional delta method from Westerhout et al. [2024, Fact 3.2].

We first obtain the Hadamard directional derivative of $g(\cdot; \beta')$ in β' and arbitrary direction $\eta \in \mathbb{R}^p$. We note that it is easy to check that for any compact set $\Theta' \subset \mathbb{R}^p$, the Lipschitz condition

$$\|g(\cdot; \beta) - g(\cdot; \beta')\|_\infty \leq C \|\beta - \beta'\|,$$

is satisfied, for some $C > 0$ and every $\beta, \beta' \in \Theta'$. Here the supremum norm is taken over the

compact domain Θ . As such, $g(\cdot; \beta')$ is locally Lipschitz and the Hadamard and Gateau derivatives coincide (cf. Penot, 2016, Prop. 5.9).

For brevity, we write

$$\mathbf{t}(\beta) = [t(\beta_j)]_{j \in [p]},$$

where $t = t_k$, for the same fixed k as in g . Under Assumptions 2 and 4, we have

$$\nabla_2 g(\beta; \beta') = 2 \left[\mathbb{E}Y - \{\mathbb{E}\mathbf{X}\}^\top \mathbf{D}(\beta') \beta \right] \nabla \mathbf{t}(\beta'),$$

for each $\beta \in \Theta$, where the derivative is taken with respect to β' . By definition, $\nabla_2 g(\cdot; \beta')^\top \boldsymbol{\eta}$ is the Gateau, and thus Hadamard derivative in the direction $\boldsymbol{\eta} \in \mathbb{R}^p$ if

$$\lim_{s \searrow 0} \left\| \frac{g(\cdot; \beta' + s\boldsymbol{\eta}) - g(\cdot; \beta')}{s} - \nabla g(\cdot; \beta')^\top \boldsymbol{\eta} \right\|_\infty = 0.$$

By Taylor's theorem, for each $\beta, \beta' \in \Theta$, it holds that

$$\frac{g(\beta; \beta' + s\boldsymbol{\eta}) - g(\beta; \beta')}{s} - \nabla g(\beta; \beta')^\top \boldsymbol{\eta} = s\boldsymbol{\eta}^\top \mathbf{H}(\beta; \bar{\beta}) \boldsymbol{\eta}, \quad (11)$$

for some $\bar{\beta} \in \Theta$, where $\mathbf{H}(\beta; \bar{\beta})$ is the Hessian of $g(\beta; \beta')$ in β' , where the righthand side of (11) is continuous in both $(\beta, \bar{\beta}) \in \Theta \times \Theta$, for each s and $\boldsymbol{\eta}$, by Assumptions 2. Thus,

$$\left\| \frac{g(\cdot; \beta' + s\boldsymbol{\eta}) - g(\cdot; \beta')}{s} - \nabla g(\cdot; \beta')^\top \boldsymbol{\eta} \right\|_\infty \leq s \sup_{\bar{\beta} \in \Theta} \left\| \boldsymbol{\eta}^\top \mathbf{H}(\cdot; \bar{\beta}) \boldsymbol{\eta} \right\|_\infty \rightarrow 0$$

as $s \searrow 0$, by the extreme value theorem as required. Since $\sqrt{n}(\hat{\beta}_n - \beta_0) \rightsquigarrow \mathbf{Z}$, from Assumption 5, the delta method of Westerhout et al. [2024, Fact 3.2] then yields the following result, which completes the proof.

Proposition 3. *Under Assumptions 2, 4, and 5, it holds that (1C) $\rightsquigarrow \mathbf{Z}^\top \nabla_2 g(\cdot, \beta_0)$.*

B Proof of Theorem 4

We first prove that $\mathbb{P}(\bar{\boldsymbol{\beta}}_{n,I} = \mathbf{0}) \rightarrow 1$, as $n \rightarrow \infty$. By the definition of $\bar{\boldsymbol{\beta}}_n$ and the union bound,

$$\begin{aligned} \mathbb{P}(\bar{\boldsymbol{\beta}}_{n,I} \neq \mathbf{0}) &= \mathbb{P}\left(\bigcup_{j \in \mathcal{S}_0} \{\mathbf{1}(t_{\hat{K}_n}(\hat{\beta}_{n,j}) = 0) \neq 0\}\right) \\ &= \mathbb{P}\left(\bigcup_{j \in \mathcal{S}_0} \{|\hat{\beta}_{n,j}| > \delta_{\hat{K}_n}\}\right) \\ &\leq \sum_{j \in \mathcal{S}_0} \mathbb{P}\left(|\hat{\beta}_{n,j}| > \delta_{\hat{K}_n}\right). \end{aligned}$$

Then for each $j \in \mathcal{S}_0$, by the consistency of \hat{K}_n and $\hat{\boldsymbol{\beta}}_n$, we have that

$$\begin{aligned} \mathbb{P}\left(|\hat{\beta}_{n,j}| > \delta_{\hat{K}_n}\right) &= \sum_{k \neq k_0} \left\{ \mathbb{P}\left(|\hat{\beta}_{n,j}| > \delta_k, \hat{K}_n = k\right) \right\} + \mathbb{P}\left(|\hat{\beta}_{n,j}| > \delta_{k_0}, \hat{K}_n = k_0\right) \\ &\leq \sum_{k \neq k_0} \left\{ \mathbb{P}\left(\hat{K}_n = k\right) \right\} + \mathbb{P}\left(|\hat{\beta}_{n,j}| > \delta_{k_0}\right) \rightarrow 0. \end{aligned}$$

As $|\mathcal{S}_0| < \infty$, this completes the proof regarding the consistency on the irrelevant set. For the relevant set, we write

$$\sqrt{n}(\bar{\boldsymbol{\beta}}_{n,R} - \boldsymbol{\beta}_{0,R}) = \sqrt{n}(\bar{\boldsymbol{\beta}}_{n,R} - \hat{\boldsymbol{\beta}}_{n,R}) + \sqrt{n}(\hat{\boldsymbol{\beta}}_{n,R} - \boldsymbol{\beta}_{0,R}).$$

The second term on the RHS converges weakly to \mathbf{Z}_R by Assumption 5, and so, by Slutsky's theorem, it suffices to show that

$$\lim_{n \rightarrow \infty} \mathbb{P}\left(\sqrt{n}(\bar{\boldsymbol{\beta}}_{n,R} - \hat{\boldsymbol{\beta}}_{n,R}) = \mathbf{0}\right) = 1.$$

Similarly to the irrelevant case, the definition of $\bar{\boldsymbol{\beta}}_n$ and the union bound yields

$$\begin{aligned} \mathbb{P}\left(\sqrt{n}(\bar{\boldsymbol{\beta}}_{n,R} - \hat{\boldsymbol{\beta}}_{n,R}) \neq \mathbf{0}\right) &= \mathbb{P}\left(\bigcup_{j \in \mathcal{S}_0^c} \{|\hat{\beta}_{n,j}| \leq \delta_{\hat{K}_n}\}\right) \\ &\leq \sum_{j \in \mathcal{S}_0^c} \mathbb{P}\left(|\hat{\beta}_{n,j}| \leq \delta_{\hat{K}_n}\right). \end{aligned}$$

For each $j \in \mathcal{S}_0^c$, an identical argument to the irrelevant case yields

$$\lim_{n \rightarrow \infty} \mathbb{P}\left(|\hat{\beta}_{n,j}| \leq \delta_{\hat{K}_n}\right) = 0,$$

which completes the proof.

References

- Rémy Abergel, Olivier Bouaziz, and Grégory Nuel. A review on the adaptive-ridge algorithm with several extensions. *Statistics and Computing*, 34:140, 2024.
- Hirotsugu Akaike. A new look at the statistical model identification. *IEEE Transactions on Automatic Control*, 19:716–723, 1974.
- Sylvain Arlot. Minimal penalties and the slope heuristics: a survey. *Journal de la Société Française de Statistique*, 160:1–106, 2019.
- Jean-Patrick Baudry, Cathy Maugis, and Bertrand Michel. Slope heuristics: overview and implementation. *Statistics and Computing*, 22:455–470, 2012.
- Lucien Birgé and Pascal Massart. Minimal penalties for Gaussian model selection. *Probability theory and related fields*, 138:33–73, 2007.
- Peter Bühlmann and Sara van de Geer. *Statistics for High-Dimensional Data: Methods, Theory and Applications*. Springer, 2011.
- Linlin Dai, Kani Chen, Zhihua Sun, Zhenqiu Liu, and Gang Li. Broken adaptive ridge regression and its asymptotic properties. *Journal of Multivariate Analysis*, 168:334–351, 2018.
- James Davidson. *Stochastic Limit Theory: An Introduction for Econometricians*. Oxford University Press, 2021.
- Jianqing Fan and Runze Li. Variable selection via nonconcave penalized likelihood and its oracle properties. *Journal of the American statistical Association*, 96:1348–1360, 2001.
- Jianqing Fan and Jinchi Lv. A selective overview of variable selection in high dimensional feature space. *Statistica Sinica*, 20(1):101, 2010.
- Florian Frommlet and Grégory Nuel. An adaptive ridge procedure for l0 regularization. *PloS one*, 11:e0148620, 2016.
- Wenjiang J Fu. Penalized regressions: the bridge versus the lasso. *Journal of computational and graphical statistics*, 7(3):397–416, 1998.
- Edward I George. The variable selection problem. *Journal of the American Statistical Association*, 95(452):1304–1308, 2000.
- Trevor Hastie. *The Elements of Statistical Learning: Data Mining, Inference, and Prediction*. Springer, 2009.

- Trevor Hastie, Robert Tibshirani, and Martin Wainwright. *Statistical Learning with Sparsity*. CRC Press, 2015.
- Georg Heinze, Christine Wallisch, and Daniela Dunkler. Variable selection—a review and recommendations for the practicing statistician. *Biometrical journal*, 60(3):431–449, 2018.
- Karl Hinderer, Ulrich Rieder, and Michael Stieglitz. *Dynamic Optimization*. Springer, 2016.
- K L K Ho and H Masuda. Adaptive ridge approach to heteroscedastic regression. *ArXiv*, (2402.13642), 2024.
- Arthur E Hoerl and Robert W Kennard. Ridge regression: Biased estimation for nonorthogonal problems. *Technometrics*, 12:55–67, 1970.
- David R Hunter and Runze Li. Variable selection using mm algorithms. *Annals of Statistics*, 33:1617, 2005.
- Q Lin. The bayesian elasticnet. *Bayesian Analysis*, 5:151–170, 2010.
- Luke R Lloyd-Jones, Hien D Nguyen, and Geoffrey J McLachlan. A globally convergent algorithm for lasso-penalized mixture of linear regression models. *Computational Statistics & Data Analysis*, 119:19–38, 2018.
- Hien Duy Nguyen. PanIC: Consistent information criteria for general model selection problems. *Australian & New Zealand Journal of Statistics*, 2024.
- Jean-Paul Penot. *Analysis: From Concepts to Applications*. Springer, 2016.
- Piotr Pokarowski and Jan Mielniczuk. Combined l1 and greedy l0 penalized least squares for linear model selection. *Journal of Machine Learning Research*, 16:961–992, 2015.
- Benedikt M Pötscher and Hannes Leeb. On the distribution of penalized maximum likelihood estimators: The lasso, scad, and thresholding. *Journal of Multivariate Analysis*, 100(9):2065–2082, 2009.
- Benedikt M Pötscher and Ulrike Schneider. Distributional results for thresholding estimators in high-dimensional gaussian regression models. *Electronic Journal of Statistics*, 5:1876–1934, 2011.
- Irina Rish and Genady Grabarnik. *Sparse Modeling: Theory, Algorithms, and Applications*. CRC Press, 2014.
- Werner Römisch. *Delta Method, Infinite Dimensional*. Wiley, 2014.
- Ulrike Schneider. Confidence sets based on thresholding estimators in high-dimensional gaussian regression models. *Econometric Reviews*, 35:1412–1455, 2016.

- Gideon Schwarz. Estimating the dimension of a model. *Annals of Statistics*, 6:461–464, 1978.
- Alexander Shapiro, Darinka Dentcheva, and Andrzej Ruszczyński. *Lectures on Stochastic Programming: Modeling and Theory*. SIAM, 2021.
- Chor-Yiu Sin and Halbert White. Information criteria for selecting possibly misspecified parametric models. *Journal of Econometrics*, 71:207–225, 1996.
- Martin Slawski and Matthias Hein. Non-negative least squares for high-dimensional linear models: Consistency and sparse recovery without regularization. *Electronic Journal of Statistics*, 7:3004, 2013.
- Thomas A Stamey, John N Kabalin, John E McNeal, Iain M Johnstone, Fuad Freiha, Elise A Redwine, and Norman Yang. Prostate specific antigen in the diagnosis and treatment of adenocarcinoma of the prostate. ii. radical prostatectomy treated patients. *The Journal of Urology*, 141:1076–1083, 1989.
- Qiang Sun, Bai Jiang, Hongtu Zhu, and Joseph G Ibrahim. Hard thresholding regression. *Scandinavian Journal of Statistics*, 46:314–328, 2019.
- Wei Sun, Junhui Wang, and Yixin Fang. Consistent selection of tuning parameters via variable selection stability. *J. Mach. Learn. Res.*, 14(1):3419–3440, January 2013. ISSN 1532-4435.
- Robert Tibshirani. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 58(1):267–288, 1996.
- Sara van de Geer, Peter Bühlmann, and Shuheng Zhou. The adaptive and the thresholded lasso for potentially misspecified models (and a lower bound for the lasso). *Electronic Journal of Statistics*, 5:688–749, 2011.
- Sara A van de Geer. *Estimation and testing under sparsity*. Springer, 2016.
- A W van der Vaart and Jon A Wellner. *Weak Convergence and Empirical Processes: With Applications to Statistics*. Springer, 2023.
- Aad W van der Vaart and Jon A Wellner. Empirical processes indexed by estimated functions. *Lecture Notes Monograph Series*, 55:234–252, 2007.
- Jacob Westerhout, TrungTin Nguyen, Xin Guo, and Hien Duy Nguyen. On the asymptotic distribution of the minimum empirical risk. In *Forty-first International Conference on Machine Learning*, 2024.
- Qingyuan Zhang and Hien Duy Nguyen. Consistent information criteria for regularized regression and loss-based learning problems. *ArXiv*, (2404.17181), 2024.

Yun-Bin Zhao. *Sparse Optimization Theory and Methods*. CRC Press, 2018.

Zemin Zheng, Yingying Fan, and Jinchi Lv. High dimensional thresholded regression and shrinkage effect. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 76:627–649, 2014.

Shuheng Zhou. Thresholding procedures for high dimensional variable selection and statistical estimation. *Advances in Neural Information Processing Systems*, 22, 2009.

Hui Zou and Trevor Hastie. Regularization and variable selection via the elastic net. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 67(2):301–320, 2005.