

OPERATORS ON INJECTIVE TENSOR PRODUCTS OF SEPARABLE BANACH SPACES AND SPACES WITH FEW OPERATORS

ANTONIO ACUAVIVA

ABSTRACT. We give a characterization of the operators on the injective tensor product $E \hat{\otimes}_\varepsilon X$ for E a separable Banach space and X a (non-separable) Banach space with few operators, in the sense that any operator $T : X \rightarrow X$ takes the form $T = \lambda I + S$ for a scalar $\lambda \in \mathbb{K}$ and an operator S with separable range. This is used to give a classification of the complemented subspaces and closed operator ideals of spaces of the form $C_0(\omega \times K_{\mathcal{A}})$, where $K_{\mathcal{A}}$ is a locally compact Hausdorff space induced by an almost disjoint family \mathcal{A} such that $C_0(K_{\mathcal{A}})$ has few operators.

1. INTRODUCTION.

A common characteristic of many Banach spaces is the rich structure of their spaces of operators. This has, in turn, led to the question of how simple the space of operators on a Banach space can be. Efforts to explore this question have spurred a rich vein of research, culminating in the construction of several exotic Banach spaces.

Among the most notable examples are the space constructed by Gowers and Maurey [9], that does not contain an unconditional basic sequence, and the space built by Argyros and Haydon, which solved the scalar-plus-compact problem [1].

In the non-separable setting, a similar question has been explored, focusing on spaces X where every operator is a separable perturbation of a scalar multiple of the identity; that is, $T = \lambda I + S$, where S is an operator with separable range. Such spaces, which we will refer to as *spaces with few operators*, were first constructed by Shelah [19] under the assumption of the diamond axiom, which was later removed

Date: March 28, 2025.

2020 Mathematics Subject Classification. 46B26, 47L10, 46B20, 46M05 (primary); 46E15, 54G12 54B10 (secondary).

Key words and phrases. Tensor products, Few Operators, Banach space, $C(K)$ -spaces, Scattered spaces, Complemented subspaces, Closed operator ideals.

by Shelah and Steprāns [20]. Another space of this form, realized as a dual space, was constructed by Argyros and Tolias [2], while a reflexive example was built by Wark [21] (note that this implies that its dual also has few operators).

This question has recently attracted significant interest in the context of spaces of continuous functions, leading to the construction of many exotic locally compact spaces. In this vein, Koszmider constructed, under the Continuum Hypothesis or Martin's Axiom, a scattered, locally compact Hausdorff space L such that $C_0(L)$ admits few operators [12]. These set-theoretic assumptions were later removed by Koszmider and Laustsen [13]. Additionally, under the assumption of Ostaszewski's \clubsuit -principle, Koszmider and Zieliński [14] constructed a different example of a $C_0(L)$ spaces with few operators. This construction was further developed in work of Candido [6] and [7].

Given the simple structure of the operators on these spaces, a natural question arises: can they be used to construct new spaces that still retain easily characterizable operators? We show that this is the case when considering the injective tensor product with a separable Banach space E . This result becomes particularly interesting when $X = C_0(L)$ and $E = C(M)$, where M is a compact metric space and L is a locally compact Hausdorff space since in this case $C(M) \hat{\otimes}_\varepsilon C_0(L) \cong C_0(M \times L)$.

We present our main result in slightly more general terms, as this does not complicate the proof. In the following theorem, $\hat{\otimes}_\alpha$ denotes any tensor completion for a *uniform cross-norm* $\alpha(\cdot)$, see [18, Chapter 6].

Theorem 1.1. *Let E, F be separable Banach spaces and X a non-separable Banach space with few operators. Then any operator $T : E \hat{\otimes}_\alpha X \rightarrow F \hat{\otimes}_\varepsilon X$ can be uniquely expressed as $T = U \hat{\otimes} I + S$, where $U : E \rightarrow F$ and S has separable range.*

The essence of the previous theorem is as follows: when we take the injective tensor product of a separable Banach space with a space with few operators, we obtain, up to separable elements, the original space we started with. In particular, taking the injective tensor product of separable Banach spaces with spaces with few operators is an injective operation, in the following sense.

Corollary 1.2. *Let E, F be separable Banach spaces and X be a non-separable Banach space with few operators. Then*

$$E \hat{\otimes}_\varepsilon X \sim F \hat{\otimes}_\varepsilon X \iff E \sim F.$$

As previously mentioned, we are particularly interested in the case of spaces of continuous functions. Theorem 1.1 becomes especially relevant when we take $E = C(M_1)$, $F = C(M_2)$, $\hat{\otimes}_\alpha = \hat{\otimes}_\varepsilon$ the injective tensor product and $X = C_0(L)$ for some locally compact Hausdorff space L . This gives information about operators on spaces of the form $C_0(M \times L)$.

Corollary 1.3. *Let L be a non-metrizable locally compact Hausdorff space such that $C_0(L)$ admits few operators, and M_1, M_2 be compact metric spaces. Then any operator $T : C_0(M_1 \times L) \rightarrow C_0(M_2 \times L)$ can be uniquely expressed as*

$$T = U \hat{\otimes} I + S,$$

where $U : C(M_1) \rightarrow C(M_2)$ and S has separable range.

The previous corollary generalizes Candido's result [6, Theorem 1.1], originally proved under \clubsuit . It extends the result to also include the case $M = [0, 1]$ and demonstrates that the key property is that $C_0(L)$ has few operators. For example, it applies to the space built by Koszmider and Laustsen, whose construction does not require additional set-theoretic assumptions.

As an application of Theorem 1.1, we expand the list of spaces of continuous functions with completely understood complemented subspaces, obtaining the following.

Theorem 1.4. *Let $K_{\mathcal{A}}$ be a locally compact Hausdorff space induced by an almost disjoint family $\mathcal{A} \subseteq [\mathbb{N}]^\omega$ of cardinality the continuum such that $C_0(K_{\mathcal{A}})$ admits few operators. Then any infinite-dimensional complemented subspace of $C_0(\omega \times K_{\mathcal{A}})$ is isomorphic to exactly one of the following: c_0 , $C_0(K_{\mathcal{A}})^n$ for some $n \in \mathbb{N}$, or $C_0(\omega \times K_{\mathcal{A}})$.*

Remark 1.5. Note that, if $C(L)$ is a Grothendieck space such that its complemented subspaces are classified, then we also obtain a classification of the complemented subspaces of $C_0((\omega \times K_{\mathcal{A}}) \sqcup L) \cong C_0((\omega \times K_{\mathcal{A}}) \oplus C(L)$, see [10]. For example, we get a classification of the complemented subspaces of $\ell_\infty^c(\Gamma) \oplus C(\omega \times K_{\mathcal{A}})$ for any set Γ .

We believe that it is possible to extend the previous classification of complemented subspaces to $C_0(\omega^\omega \times K_{\mathcal{A}})$, by using arguments similar to those of Benyamini's classification of the complemented subspaces of $C(\omega^\omega)$ [4]. However, the proof becomes significantly technical and involved, and we are not aware of any immediate applications. We conjecture that it may even be possible to extend the classification

to $C_0(M \times K_{\mathcal{A}})$ for M a compact metric space, in terms of the complemented subspaces of $C(M)$, as in [6, Theorem 1.3], though we are unable to determine a general approach to achieve this.

Finally, we can fully describe the lattice of closed operator ideals on $C_0(\omega \times K_{\mathcal{A}})$. For a Banach space Y , we denote by $\mathcal{K}(Y)$ the ideal of compact operators, $\mathcal{X}(Y)$ denotes the ideal of operators with separable range, and if Z is another Banach space, we define the ideal $\mathcal{G}_Z(Y)$ by

$$\mathcal{G}_Z(Y) = \text{span}\{ST : T \in \mathcal{B}(Y, Z), S \in \mathcal{B}(Z, Y)\}.$$

Theorem 1.6. *Let $K_{\mathcal{A}}$ be a locally compact Hausdorff space induced by an almost disjoint family $\mathcal{A} \subseteq [\mathbb{N}]^{\omega}$ of cardinality the continuum such that $C_0(K_{\mathcal{A}})$ admits few operators. Then the lattice of closed ideals of $\mathcal{B}(C_0(\omega \times K_{\mathcal{A}}))$ is given by*

$$\begin{aligned} \{0\} \subsetneq \mathcal{K}(C_0(\omega \times K_{\mathcal{A}})) \subsetneq \mathcal{G}_{c_0}(C_0(\omega \times K_{\mathcal{A}})) = \mathcal{X}(C_0(\omega \times K_{\mathcal{A}})) \\ \subsetneq \overline{\mathcal{T}}_{C_0(K_{\mathcal{A}})}(C_0(\omega \times K_{\mathcal{A}})) \subsetneq \mathcal{B}(C_0(\omega \times K_{\mathcal{A}})). \end{aligned}$$

Remark 1.7. It is worth noting that even more can be deduced from this result. Since the operators on c_0 are particularly well-behaved, the same is true for the operators on $C_0(\omega \times K_{\mathcal{A}})$. Specifically, note that $\mathcal{B}(C_0(\omega \times K_{\mathcal{A}}))/\mathcal{X}(C_0(\omega \times K_{\mathcal{A}}))$ and $\mathcal{B}(c_0)$ are isometrically isomorphic as Banach algebras. This allows us to conclude that many properties of operators on c_0 also hold for those on $C_0(\omega \times K_{\mathcal{A}})$. For example, one can easily deduce that the quotient $\mathcal{B}(C_0(\omega \times K_{\mathcal{A}}))/\mathcal{I}$ has a unique algebra norm for every closed ideal \mathcal{I} of $\mathcal{B}(C_0(\omega \times K_{\mathcal{A}}))$, in the spirit studied in [3].

Remark 1.8. Note that $C_0(\omega \times K_{\mathcal{A}}) \cong C_0(K_{\mathcal{B}})$ where $K_{\mathcal{B}} \sim \bigsqcup_{n=1}^{\infty} K_{\mathcal{A}}$ is a locally compact Hausdorff space induced by the almost disjoint family \mathcal{B} , obtained by breaking up \mathbb{N} into countably many copies of itself and using the order isomorphism to create a copy of \mathcal{A} in each of these copies of \mathbb{N} . Thus, the previous argument shows that the unital Banach algebra $\mathcal{B}(c_0)$ can be realized as a quotient $\mathcal{B}(C_0(K_{\mathcal{B}}))/\mathcal{X}(C_0(K_{\mathcal{B}}))$, addressing a question of Koszmider and Laustsen [13, Question 46 (iii)].

2. ORGANIZATION AND NOTATION

All normed spaces and algebras are over the scalar field \mathbb{K} , either the real or complex numbers, and we adhere to standard notational conventions. The term *operator* will refer to a bounded linear map between normed spaces.

From now on, X will denote a non-separable Banach space with few operators, meaning that every operator $T : X \rightarrow X$ can be expressed as $T = \lambda I + S$, where $\lambda \in \mathbb{K}$ and S is an operator with separable range. Meanwhile, E and F will always refer to separable Banach spaces, while M will be a metrizable compact Hausdorff space. We denote by B_Y the unit ball of a Banach space Y .

We will use some known facts about tensor products in general and injective tensor products in particular, see for example [18]. From now on, $\hat{\otimes}_\alpha$ will denote the tensor product taken for a uniform cross-norm $\alpha(\cdot)$ and $\hat{\otimes}_\varepsilon$ will denote the injective tensor product. In particular, given operators $U : E \rightarrow F$ and $R : X \rightarrow X$, we can define the tensor product operator $U \hat{\otimes} R : E \hat{\otimes}_\alpha X \rightarrow F \hat{\otimes}_\varepsilon X$ by extending the algebraic tensor product operator $U \otimes R : E \otimes X \rightarrow F \otimes X$. Further, recall that $C(M) \hat{\otimes}_\varepsilon X$ is isometrically isomorphic to $C(M, X)$ under the identification $(f \otimes x)(k) = f(k)x$ for $f \in C(M)$ and $x \in X$.

More specialized notation will be introduced as and when needed.

We give a brief layout of the structure of this paper. In Section 3, we present the proof of Theorem 1.1. First, we prove the result in the case $F = C(M)$. For this, the key is to construct the operator $U : E \rightarrow C(M)$ and then verify that $T - U \hat{\otimes} I$ has separable range. The general case when F is any separable Banach space will readily follow using the universality of $C[0, 1]$ for separable Banach spaces, together with the fact that the injective tensor product preserves subspaces. We also provide a proof of Corollary 1.2.

In Section 4, we explore some applications of our theorem in the context of Banach spaces induced by an almost disjoint family, including the proofs of Theorems 1.4 and 1.6.

3. PROOF OF THEOREM 1.1

The strategy for the proof is to exploit the simple structure of operators on X . We first assume that $F = C(M)$ and we need to transform an operator $T : E \hat{\otimes}_\alpha X \rightarrow C(M, X)$ into a collection of operators from X to X , which encapsulate all the information of the original operator. We will need the following definition.

Definition 3.1. For each $k \in M$, define the *evaluation operator*

$$\pi_k : C(M, X) \rightarrow X, \quad g \mapsto g(k).$$

For each $e \in E$, define the *inclusion operator*

$$\iota_e : X \rightarrow E \hat{\otimes}_\alpha X, \quad x \mapsto e \otimes x.$$

Note that $\|\iota_e\| \leq \|e\|$ and $\|\pi_k\| = 1$.

Using the evaluation and inclusion operators, it is easy to obtain operators from X to X using our original operator T . We will then use these operators to build $U : E \rightarrow C(M)$.

Definition 3.2. Let $e \in E$, $k \in M$ and $T : E \hat{\otimes}_\alpha X \rightarrow C(M, X)$ be an operator. Then the (e, k) -coordinate of T is given by

$$T_{e,k} = \pi_k T \iota_e : X \rightarrow X,$$

so that in particular there is a unique decomposition $T_{e,k} = \lambda_{e,k} I + S_{e,k}$ where $\lambda_{e,k} \in \mathbb{K}$ and $S_{e,k} : X \rightarrow X$ is an operator with separable range.

From now on fix a countable dense set $M_0 \subseteq M$. We have the following elementary observation.

Lemma 3.3. *Let Y_{E_0} be the closed linear span of $\cup\{S_{e,k}[X] : e \in E_0, k \in M_0\}$ for any countable $E_0 \subseteq E$. Then there exist $x \in B_X$ and $\varphi \in B_{X^*}$ such that $\varphi(x) \geq 1/2$ and $\varphi(Y_{E_0}) = 0$.*

In particular, for any finite collection of vectors $e_1, \dots, e_n \in E$, there exists $x \in B_X$ such that

$$\left| \sum_{i=1}^n c_i \lambda_{e_i, k_i} \right| \leq 2 \left\| \sum_{i=1}^n c_i \pi_{k_i} T(e_i \otimes x) \right\|$$

for any choice $k_1, \dots, k_n \in M_0$ and scalars $c_1, \dots, c_n \in \mathbb{K}$.

Proof. Note that $Y_{E_0} \subsetneq X$, since the latter is non-separable. Thus an application of the Hahn-Banach theorem gives the first part.

For the second part, take $E_0 = \{e_1, \dots, e_n\}$. There exists $x \in B_X$ and $\varphi \in B_{X^*}$ such that $\varphi(Y_{E_0}) = 0$ and $\varphi(x) \geq 1/2$. It follows that for any $k_1, \dots, k_n \in M_0$ and $c_1, \dots, c_n \in \mathbb{K}$, we have

$$\begin{aligned} \left| \sum_{i=1}^n c_i \lambda_{e_i, k_i} \right| &\leq 2 |\varphi(x)| \left| \sum_{i=1}^n c_i \lambda_{e_i, k_i} \right| = 2 \left| \varphi \left(\sum_{i=1}^n c_i \lambda_{e_i, k_i} x + c_i S_{e_i, k_i} x \right) \right| \\ &\leq 2 \left\| \sum_{i=1}^n c_i (\lambda_{e_i, k_i} x + S_{e_i, k_i} x) \right\| = 2 \left\| \sum_{i=1}^n c_i \pi_{k_i} T(e_i \otimes x) \right\|. \quad \square \end{aligned}$$

We can now build the operator $U : E \rightarrow C(M)$, using $(\lambda_{e,k})_{e \in E, k \in M_0}$ together with an extension via density.

Lemma 3.4. *For each $e \in E$ the function $U_0 e : M_0 \rightarrow \mathbb{K}$ defined by $U_0 e(k) = \lambda_{e,k}$ is uniformly continuous and thus admits a unique continuous extension to M , which we denote by $Ue \in C(M)$.*

Proof. Let $e \in E$ be fixed, applying the second part of Lemma 3.3 for any $k_1, k_2 \in M_0$ we have

$$\begin{aligned} |U_0e(k_1) - U_0e(k_2)| &= |\lambda_{e,k_1} - \lambda_{e,k_2}| \leq 2\|\pi_{k_1}T(e \otimes x) - \pi_{k_2}T(e \otimes x)\| \\ &= 2\|(T(e \otimes x))(k_1) - (T(e \otimes x))(k_2)\|, \end{aligned}$$

and since $T(e \otimes x) \in C(M, X)$ is uniformly continuous by the Heine-Cantor theorem, so is U_0e . \square

Lemma 3.5. *The map $U : E \rightarrow C(M)$, $e \mapsto Ue$, where Ue is given as in Lemma 3.4, defines an operator of norm at most $2\|T\|$.*

Proof. We start by showing that U is bounded, by arguing as in the proof of Lemma 3.4. For a fixed $e \in E$, an application of Lemma 3.3 gives

$$\|Ue\| = \sup_{k \in M_0} \|U_0e(k)\| = \sup_{k \in M_0} |\lambda_{e,k}| \leq \sup_{k \in M_0} \|\pi_k T(e \otimes x)\| \leq 2\|T\|\|e\|.$$

To show linearity, let $e_1, e_2 \in E$ and $c \in \mathbb{K}$. An application of Lemma 3.3, together with the linearity of T and the tensor product, gives $|\lambda_{e_1+k, c} + c\lambda_{e_2, k} - \lambda_{e_1+ce_2, k}| = 0$ for each $k \in M_0$. Therefore, the continuous function $(Ue_1 + cUe_2) - U(e_1 + ce_2)$ vanishes on the dense set M_0 , and thus it is the null function. This proves the linearity of U . \square

Once we have found the operator $U : E \rightarrow C(M)$, we simply need to check that it gives the desired representation, that is, we need to check that the operator $S = T - U \hat{\otimes} I$ has separable range. For that, we need the following lemma.

Lemma 3.6. *Let Y be a subspace of $C(M, X)$. Then Y is separable if and only if $\pi_k[Y]$ is separable for each $k \in M_0$.*

Proof. If Y is separable, clearly $\pi_k[Y]$ is separable, so we focus on the reverse implication. Let $Z = \overline{\text{span}} \bigcup_{k \in M_0} \pi_k[Y]$ which is separable, since by assumption each $\pi_k[Y]$ is. We naturally view $C(M, Z)$ as a closed subspace of $C(M, X)$. Observe that $C(M, Z)$ is also separable, since $C(M, Z) \cong C(M) \hat{\otimes}_\varepsilon Z$ and the tensor product of separable spaces is itself separable. Therefore, it is enough to show that $Y \subseteq C(M, Z)$,

For this, we need to show that every $y \in Y$ takes values in Z . For any $k \in M$, take $k_n \in M_0$, $k_n \rightarrow k$. By continuity of y , $\pi_{k_n}(y) \rightarrow \pi_k y$. Since $\pi_{k_n}(y) \in Z$ for every $n \in \mathbb{N}$ and Z is a closed subspace, then $\pi_k y \in Z$, finishing the proof. \square

We are ready to prove Theorem 1.1 in the case $F = C(M)$.

Proof of Theorem 1.1, for $F = C(M)$. Let $U : E \rightarrow C(M)$ be the operator given by Lemma 3.5. We claim that the operator $S = T - U \hat{\otimes} I$ has separable range. The uniqueness of the representation is clear, once its existence has been shown.

Thus, we only need to prove that S has separable range. Let E_0 be a countable dense set of E and recall that Y_{E_0} denotes the closed linear span of $\cup\{S_{e,k}[X] : e \in E_0, k \in M_0\}$. By Lemma 3.6, we need to show that $\pi_k S[E \hat{\otimes}_\alpha X]$ is separable for each $k \in M_0$, and for this, it is enough to show $\pi_k S[E \hat{\otimes}_\alpha X] \subseteq Y_{E_0}$.

Since E_0 is dense in E , $\text{span}\{e \otimes x : e \in E_0, x \in X\}$ is dense in $E \hat{\otimes}_\alpha X$. Therefore, by continuity and linearity, it is enough to show that $\pi_k S(e \otimes x) \in Y_{E_0}$ for each $e \in E_0, x \in X$. We have

$$\begin{aligned} \pi_k S(e \otimes x) &= \pi_k (T - U \hat{\otimes} I)(e \otimes x) = \pi_k T(e \otimes x) - \pi_k ((Ue) \otimes x) \\ &= (\lambda_{e,k}x + S_{e,k}x) - \lambda_{e,k}x = S_{e,k}x \in Y_{E_0}, \end{aligned}$$

as required. \square

Proof of Theorem 1.1, general case. Take $M = [0, 1]$ and let $j : F \hookrightarrow C(M)$ be an isometric embedding. Since the injective tensor product preserves subspaces, we can identify $F \hat{\otimes}_\varepsilon X$ with a closed subspace of $C(M, X) \cong C(M) \hat{\otimes}_\varepsilon X$, specifically $j \hat{\otimes} I : F \hat{\otimes}_\varepsilon X \hookrightarrow C(M) \hat{\otimes}_\varepsilon X$ is an isometric embedding.

Let $T : E \hat{\otimes}_\alpha X \rightarrow F \hat{\otimes}_\varepsilon X$ be an operator. By the previous case applied to $(j \hat{\otimes} I) \circ T$, there exists $U : E \rightarrow C(M)$ and an operator S with separable range such that

$$(j \hat{\otimes} I) \circ T = U \hat{\otimes} I + S.$$

Assume we have shown that $U[E] \subseteq j[F] \subseteq C(M)$, so that $U = j \circ V$ for some $V : E \rightarrow F$. Therefore

$$U \hat{\otimes} I = (j \circ V) \hat{\otimes} I = (j \hat{\otimes} I) \circ (V \hat{\otimes} I),$$

which combined with the previous equation gives

$$(j \hat{\otimes} I) \circ (T - V \hat{\otimes} I) = S.$$

Since $j \hat{\otimes} I$ is an isometry, we obtain that $T - V \hat{\otimes} I$ has separable range, and the result follows. To finish, we only need to show that $U[E] \subseteq j[F]$.

Since $j(F)$ is closed, it is enough to show that for any $\varepsilon > 0$ and $e \in E$ we have $d(Ue, j(F)) < \varepsilon$. From now on, fix $\varepsilon > 0$ and $e \in E$. Apply the first part of Lemma 3.3 to $E_0 = \{e\}$, to obtain $x \in B_X$ and $\varphi \in B_{X^*}$ such that $\varphi(x) \geq 1/2$ and $\varphi(Y_{E_0}) = 0$.

Since $\text{span}\{f \otimes x : f \in F, x \in X\}$ is dense in $F \hat{\otimes}_\varepsilon X$ we can choose $f_1, \dots, f_n \in F, x_1, \dots, x_n \in X$ such that

$$\|T(e \otimes x) - \sum_{i=1}^n f_i \otimes x_i\| < \varepsilon/2.$$

On the other hand, since $j \hat{\otimes} I$ is an isometry, we can compute this norm as

$$\begin{aligned} & \sup_{k \in M_0} \|\pi_k((j \hat{\otimes} I)T)(e \otimes x) - \sum_{i=1}^n j(f_i)(k)x_i\| \\ &= \sup_{k \in M_0} \|(\lambda_{e,k}x + S_{e,k}x) - \sum_{i=1}^n j(f_i)(k)x_i\| \\ &\geq \sup_{k \in M_0} |\varphi((\lambda_{e,k}x + S_{e,k}x) - \sum_{i=1}^n j(f_i)(k)x_i)| \\ &= \sup_{k \in M_0} |\varphi(x)\lambda_{e,k} - \sum_{i=1}^n \varphi(x_i)j(f_i)(k)| \\ &\geq \frac{1}{2} \sup_{k \in M_0} |\lambda_{e,k} - \sum_{i=1}^n \frac{\varphi(x_i)}{\varphi(x)}j(f_i)(k)| \\ &\geq \frac{1}{2}d(Ue, j(F)), \end{aligned}$$

so that $d(Ue, j(F)) < \varepsilon$, as desired. \square

Finally, we deduce Corollary 1.2.

Proof of Corollary 1.2. Clearly, $E \sim F$ implies $E \hat{\otimes}_\varepsilon X \sim F \hat{\otimes}_\varepsilon X$, so we focus on the reverse implication.

Let $T : E \hat{\otimes}_\varepsilon X \rightarrow F \hat{\otimes}_\varepsilon X$ be an isomorphism. Using Theorem 1.1 we can express $T = U \hat{\otimes} I + S$ where $U : E \rightarrow F$ and S has separable range. We claim that U is an isomorphism, which will finish the proof. For this, we will show that U is bounded below and surjective.

We start by showing that U is bounded below, so fix $e \in E$ with $\|e\| = 1$. Since X is non-separable, by transfinite induction we can find $(x_\alpha)_{\alpha < \omega_1} \subseteq B_X$ such that $\|x_\alpha - x_\beta\| > 1/2$ whenever $\alpha \neq \beta$. Let $y_\alpha = e \otimes x_\alpha$. Observe that $\|y_\alpha - y_\beta\| > 1/2$ whenever $\alpha \neq \beta$ and that

$$\|(U \hat{\otimes} I)(y_\alpha - y_\beta)\| = \|(Ue) \otimes (x_\alpha - x_\beta)\| = \|Ue\| \|x_\alpha - x_\beta\| \leq 2\|Ue\|.$$

Since S has separable range, we can find $\alpha < \beta < \omega_1$ such that $\|Sy_\alpha - Sy_\beta\| < 1/(4\|T^{-1}\|)$. We have that

$$\begin{aligned} 1/(2\|T^{-1}\|) &\leq \|T(y_\alpha - y_\beta)\| = \|(U \hat{\otimes} I)(y_\alpha - y_\beta) + S(y_\alpha - y_\beta)\| \\ &\leq \|(U \hat{\otimes} I)(y_\alpha - y_\beta)\| + \|S(y_\alpha - y_\beta)\| \leq 2\|Ue\| + 1/(4\|T^{-1}\|), \end{aligned}$$

in other words $\|Ue\| \geq 1/(8\|T^{-1}\|)$, so that U is bounded below.

We show now that U is surjective. Since U is bounded below, it has closed image, so it is enough to show that $U[E]$ is dense in F . Thus, fix $f \in F$ and $\varepsilon > 0$, we will show that $d(f, U[E]) < \varepsilon$.

As in the proof of Theorem 1.1 we can isometrically embed $F \hat{\otimes}_\varepsilon X$ into $C([0, 1], X)$. Let E_0 be a countable dense set in E and apply the first part of Lemma 3.3 to obtain $x \in B_X$ and $\varphi \in B_{X^*}$, where we identify S with $(j \hat{\otimes} I) \circ S$ whenever appropriate.

Since T is an isomorphism and E_0 is dense in E , we can find $e_1, \dots, e_n \in E_0$ and $x_1, \dots, x_n \in X$ such that

$$\|(f \otimes x) - T \left(\sum_{i=1}^n e_i \otimes x_i \right)\| < \varepsilon/2.$$

Since $j \hat{\otimes} I$ is an isometry, this norm can be computed as

$$\begin{aligned} & \sup_{k \in M_0} \|\pi_k \left((j \hat{\otimes} I)(f \otimes x) - (j \hat{\otimes} I) \circ T \left(\sum_{i=1}^n (e_i \otimes x_i) \right) \right)\| \\ &= \sup_{k \in M_0} \left\| j(f)(k)x - \sum_{i=1}^n j(Ue_i)(k)x_i + \pi_k \circ (j \hat{\otimes} I) \circ S((e_i \otimes x_i)) \right\| \\ &\geq \frac{1}{2} \sup_{k \in M_0} \left| j(f)(k) - \sum_{i=1}^n \frac{\varphi(x_i)}{\varphi(x)} j(Ue_i)(k) \right| \\ &\geq \frac{1}{2} d(j(f), j(U[E])) = \frac{1}{2} d(f, U[E]), \end{aligned}$$

which finishes the proof. \square

4. APPLICATIONS

In this section, we follow the notational conventions established in [13] and denote by $K_{\mathcal{A}}$ the locally compact Hausdorff space induced by the almost disjoint family $\mathcal{A} \subseteq [\mathbb{N}]^\omega$, as defined in [13, Definition 3], which we recall below.

Definition 4.1. Let $\mathcal{A} \subseteq [\mathbb{N}]^\omega$ be an almost disjoint family. Then $K_{\mathcal{A}}$ denotes the topological space consisting of distinct points $\{x_n : n \in \mathbb{N}\} \cup \{y_A : A \in \mathcal{A}\}$, where x_n is isolated for every $n \in \mathbb{N}$, and the sets

$$U(A, G) = \{x_n : n \in A \setminus G\} \cup \{y_A\}$$

for $G \in [\mathbb{N}]^{<\omega}$ form a neighbourhood basis at each point y_A for $A \in \mathcal{A}$. We write $U(A)$ for $U(A, \emptyset)$.

We begin by summarizing some fundamental properties of these spaces (see [13, Lemma 4]) and extending certain results from Section 3.2 of the same paper for $C_0(K_{\mathcal{A}})$ to their natural counterparts in the context of $C_0(M \times K_{\mathcal{A}})$, where we recall that M is a compact metric space.

Lemma 4.2. *Let $\mathcal{A} \subseteq [\mathbb{N}]^\omega$ be an almost disjoint family. Then:*

- $K_{\mathcal{A}}$ is a locally compact, scattered Hausdorff space.
- $K_{\mathcal{A}}$ is compact if and only if \mathcal{A} and $\mathbb{N} \setminus \bigcup \mathcal{A}$ are both finite.
- $\{x_n : n \in \mathbb{N}\}$ is the set of isolated points of $K_{\mathcal{A}}$; it is dense in $K_{\mathcal{A}}$, and so $K_{\mathcal{A}}$ is separable. Hence $K_{\mathcal{A}}$ is metrizable if and only if it is second countable, if and only if \mathcal{A} is countable.
- The subspace $K_{\mathcal{A}} \setminus \{x_n : n \in \mathbb{N}\} = \{y_A : A \in \mathcal{A}\}$ is closed and discrete.
- The sequence $(x_n)_{n \in \mathbb{N}}$ converges to y_A in $K_{\mathcal{A}}$ for every $A \in \mathcal{A}$.

Definition 4.3. Let $\mathcal{A} \subseteq [\mathbb{N}]^\omega$ be an almost disjoint family. For $f \in C_0(M \times K_{\mathcal{A}})$ and $\mathcal{X} \subseteq C_0(M \times K_{\mathcal{A}})$, we define

$$s(f) = \{A \in \mathcal{A} : \exists k \in M, f(k, y_A) \neq 0\}, \quad s(\mathcal{X}) = \bigcup \{s(f) : f \in \mathcal{X}\}.$$

Lemma 4.4. *The set $s(f)$ is countable for every $f \in C_0(M \times K_{\mathcal{A}})$.*

Proof. Let $M_0 \subseteq M$ be a countable dense set. We claim that

$$s(f) = \{A \in \mathcal{A} : \exists k \in M_0, f(k, y_A) \neq 0\} = \bigcup_{k \in M_0} \{A \in \mathcal{A} : f(k, y_A) \neq 0\}.$$

Indeed, if $A \in s(f)$, then there exists $k \in M$ such that $f(k, y_A) \neq 0$, and taking $(k_n) \subseteq M_0$, $k_n \rightarrow k$ we have $f(k_n, y_A) \rightarrow f(k, y_A)$. So $f(k_N, y_A) \neq 0$ for some $N \in \mathbb{N}$

Thus, it is enough to show that for each $k \in M_0$ the set $\{A \in \mathcal{A} : f(k, y_A) \neq 0\}$ is countable. Fix $k \in M_0$, and note that only finite subsets of the discrete closed set $\{(k, y_A) : A \in \mathcal{A}\}$ are compact, while for each $\varepsilon > 0$, the set $\{(k, y_A) \in M \times K_{\mathcal{A}} : |f(k, y_A)| \geq \varepsilon\}$ is compact. The result follows. \square

Lemma 4.5. *Let $\mathcal{A} \subseteq [\mathbb{N}]^\omega$ be an almost disjoint family. A closed subspace \mathcal{X} of $C_0(M \times K_{\mathcal{A}})$ is separable if and only if $s(\mathcal{X})$ is countable.*

Proof. Suppose that \mathcal{X} is separable and let \mathcal{X}_0 be a countable dense subset of \mathcal{X} , then by Lemma 4.4 $s(\mathcal{X}_0)$ is countable. It is easy to see that $s(\mathcal{X}) = s(\mathcal{X}_0)$, which proves the claim.

On the other hand, if $s(\mathcal{X})$ is countable, then \mathcal{X} is isomorphic to a subspace of

$$C_0(M \times K_{s(\mathcal{X})}) \cong C(M) \hat{\otimes}_\varepsilon C_0(K_{s(\mathcal{X})}),$$

and by Lemma 4.2, $C_0(K_{s(\mathcal{X})})$ is separable and thus so is $C_0(M \times K_{s(\mathcal{X})})$. \square

Lemma 4.6. *Suppose that $\mathcal{A} \subseteq [\mathbb{N}]^\omega$ is an almost disjoint family and $\mathcal{X} \subseteq C_0(M \times K_{\mathcal{A}})$ is separable. Then there is a closed subspace \mathcal{Y} of $C_0(M \times K_{\mathcal{A}})$ such that $\mathcal{X} \subseteq \mathcal{Y}$ and $\mathcal{Y} \sim C_0(\omega \times M)$. Specifically, when M is infinite, $\mathcal{Y} \sim C(M)$, while for finite M , $\mathcal{Y} \sim c_0$.*

Proof. This follows by looking at the second part of the proof of Lemma 4.5, and noticing that since $K_{s(\mathcal{X})}$ has finite Cantor–Bendixson index, then $C_0(K_{s(\mathcal{X})}) \sim C_0(\omega)$ by the classical classification of continuous functions on countable compact Hausdorff spaces [5]. \square

Our next lemma is a reformulation of [6, Proposition 3.11] in the context of spaces of the form $C_0(M \times K_{\mathcal{A}})$. Since the proof is essentially identical, we omit it here.

Lemma 4.7. *Let $P : C_0(M \times K_{\mathcal{A}}) \rightarrow C_0(M \times K_{\mathcal{A}})$ be a projection and suppose $P = Q \hat{\otimes} I + S$ for an operator $Q : C(M) \rightarrow C(M)$ and an operator S with separable range. Then Q is also a projection.*

Lastly, we will need the following well-known result, see for example [15, Lemma 3.6 (ii)].

Lemma 4.8. *Let \mathcal{X}, \mathcal{Y} be Banach spaces and $P : \mathcal{X} \rightarrow \mathcal{X}$ be a projection. Suppose P factors through \mathcal{Y} , then $P[\mathcal{X}]$ is isomorphic to a complemented subspace of \mathcal{Y} .*

Proposition 4.9. *Let $n \in \mathbb{N}$. Then every infinite-dimensional complemented subspace of $C_0(K_{\mathcal{A}})^n$ is isomorphic to either c_0 or $C_0(K_{\mathcal{A}})^m$ for some $m \leq n$.*

Proof. We proceed by induction, the case $n = 1$ follows by [12, Lemma 4]. Thus, assume the result is true for $m < n$, and we will show it holds for n .

Let $P : C_0(K_{\mathcal{A}})^n \rightarrow C_0(K_{\mathcal{A}})^n$ be a projection with infinite-dimensional image, and write $P = Q \hat{\otimes} I - S$ using Corollary 1.3, where S has separable range. By Lemma 4.7, the operator $Q : \ell_\infty^n \rightarrow \ell_\infty^n$ is also a projection and let $d = \dim Q[\ell_\infty^n]$. Since S factors through c_0 by Lemma 4.6, it follows that P factors through $C_0(K_{\mathcal{A}}, Q[\ell_\infty^n]) \oplus c_0 \sim C_0(K_{\mathcal{A}})^d \oplus c_0$. We distinguish three cases: $d = 0$, $0 < d < n$ or $d = n$.

In the first case, $Q = 0$, so $P = S$ which factors through c_0 , so that $P[C_0(K_{\mathcal{A}})^n]$ is isomorphic to a complemented subspace of c_0 and thus isomorphic to c_0 [17, Theorem 1].

If $0 < d < n$, P factors through $C_0(K_{\mathcal{A}})^d \oplus c_0 \sim C_0(K_{\mathcal{A}})^d$, so that an application of the induction hypothesis, together with Lemma 4.8, gives the result.

In the case $d = n$ then $Q = I$, so that $P = I - S$. Observe that $I - P = S$, implying that S is a projection that factors through c_0 . By Lemma 4.8, the space $S[C_0(K_{\mathcal{A}})^n]$ is isomorphic to a complemented subspace of c_0 and thus either finite-dimensional or isomorphic to c_0 .

Since $P[C_0(K_{\mathcal{A}})^n]$ is an infinite-dimensional complemented subspace of a space of continuous functions over a scattered locally compact space, it contains a complemented copy of c_0 [16]. Therefore

$$C_0(K_{\mathcal{A}})^n \sim P[C_0(K_{\mathcal{A}})^n] \oplus S[C_0(K_{\mathcal{A}})^n] \sim P[C_0(K_{\mathcal{A}})^n],$$

which completes the proof. \square

We are ready to provide the classification of the complemented subspaces of $C_0(\omega \times K_{\mathcal{A}})$.

Proof of Theorem 1.4. Let $P : C_0(\omega \times K_{\mathcal{A}}) \rightarrow C_0(\omega \times K_{\mathcal{A}})$ be a projection and express $P = Q \hat{\otimes} I - S$ according to Corollary 1.3, so that by Lemma 4.7 Q is also a projection. If $Q[C_0(\omega)]$ is finite dimensional, then P factors through $C_0(K_{\mathcal{A}})^{\dim Q[C_0(\omega)]+1}$, and the result follows from Lemma 4.8 and Proposition 4.9.

Otherwise, $Q[C_0(\omega)] \sim C_0(\omega)$, so after conjugation if necessary, we can assume $Q = I$, which gives $P = I - S$. Arguing as in the proof of Proposition 4.9 gives $P[C_0(\omega \times K_{\mathcal{A}})] \sim C_0(\omega \times K_{\mathcal{A}})$. \square

Finally, we can classify the closed ideals of $\mathcal{B}(C_0(\omega \times K_{\mathcal{A}}))$.

Proof of Theorem 1.6. Observe that $C_0(\omega \times K_{\mathcal{A}})$ has the bounded approximation property, being a \mathcal{L}_{∞} -space, and thus $\mathcal{H}(C_0(\omega \times K_{\mathcal{A}}))$ is the minimal non-zero closed ideal. By Lemma 4.6, operators with separable range factor through c_0 , so that $\mathcal{G}_{c_0}(C_0(\omega \times K_{\mathcal{A}})) = \mathcal{X}(C_0(\omega \times K_{\mathcal{A}}))$. In particular $\mathcal{G}_{c_0}(C_0(\omega \times K_{\mathcal{A}}))$ is a closed operator ideal.

Since $\omega \times K_{\mathcal{A}}$ is scattered, it is well-known that non-compact operators in $C_0(\omega \times K_{\mathcal{A}})$ factor the identity of c_0 [11, Proposition 5.4 (ii)], so that any ideal \mathcal{I} which is not contained in the ideal of compact operators, also contains the ideal of separable operators. Therefore for any closed ideal \mathcal{I} of $\mathcal{B}(C_0(\omega \times K_{\mathcal{A}}))$, $\mathcal{I} \neq \mathcal{H}(C_0(\omega \times K_{\mathcal{A}}))$, we have

$$\{0\} \subsetneq \mathcal{H}(C_0(\omega \times K_{\mathcal{A}})) \subsetneq \mathcal{X}(C_0(\omega \times K_{\mathcal{A}})) \subseteq \mathcal{I}.$$

Since $\varphi : \mathcal{B}(C_0(\omega \times K_{\mathcal{A}})) \rightarrow \mathcal{B}(c_0)$, defined by $\varphi(T) = \varphi(U \hat{\otimes} I + S) = U$, is a surjective algebra homomorphism with $\ker \varphi = \mathcal{X}((C_0(\omega \times K_{\mathcal{A}}))$, it follows that there is a one-to-one correspondence between the closed ideals of $\mathcal{B}(c_0)$ and the closed ideals of $\mathcal{B}(C_0(\omega \times K_{\mathcal{A}}))$ which contain the ideal $\mathcal{X}(C_0(\omega \times K_{\mathcal{A}}))$. The only closed operator ideals on c_0 are $\{0\} \subsetneq \mathcal{H}(c_0) \subsetneq \mathcal{B}(c_0)$ [8]; we obtain the rest of the ideals of

$\mathcal{B}(C_0(\omega \times K_{\mathcal{A}}))$ as the preimages of those. The only non-trivial case is

$$\begin{aligned} \varphi^{-1}(\mathcal{K}(c_0)) &= \{Q\hat{\otimes}I + S : Q \in \mathcal{K}(c_0), S \in \mathcal{X}(C_0(\omega \times K_{\mathcal{A}}))\} \\ &= \overline{\mathcal{G}}_{C_0(K_{\mathcal{A}})}(C_0(\omega \times K_{\mathcal{A}})), \end{aligned}$$

which follows since finite rank operators uniformly approximate every compact operator in c_0 , while separable range operators on $C_0(\omega \times K_{\mathcal{A}})$ factor through c_0 , and thus through $C_0(K_{\mathcal{A}})$. \square

Acknowledgements. This paper forms part of the author's PhD research at Lancaster University under the supervision of Professor N. J. Laustsen. The author extends sincere gratitude to Professor Laustsen for his insightful comments and valuable suggestions, which greatly improved the manuscript's presentation. Furthermore, his identification of Lemma 4.8 significantly simplified the proof of Theorem 1.4.

He also acknowledges with thanks the funding from the EPSRC (grant number EP/W524438/1) that has supported his studies.

REFERENCES

- [1] S. A. ARGYROS and R. G. HAYDON, A hereditarily indecomposable \mathcal{L}_∞ -space that solves the scalar-plus-compact problem, *Acta Math.* **206** (2011), 1–54.
- [2] S. A. ARGYROS and A. TOLIAS, Methods in the theory of hereditarily indecomposable Banach spaces, *Mem. Amer. Math. Soc.* **170** (2004).
- [3] M. ARNOTT and N. J. LAUSTSEN, Uniqueness of algebra norm on quotients of the algebra of bounded operators on a Banach space, *J. Funct. Anal.* **287** (2024), 110559.
- [4] Y. BENYAMINI, An extension theorem for separable Banach spaces, *Israel Journal of Mathematics* **29** (1978), 53–62.
- [5] C. BESSAGA and A. PELCZYŃSKI, Spaces of continuous functions (IV). (On isomorphical classification of spaces of continuous functions), *Studia Math.* **19** (1960), 53–62. Available at: <http://eudml.org/doc/216968>.
- [6] L. CANDIDO, On Banach spaces of the form $C_0(\alpha \times L)$ with few operators, *Banach J. Math. Anal.* **15** (2021), 1–23.
- [7] L. CANDIDO, Few operators on Banach spaces $C_0(L \times L)$, *arXiv preprint arXiv:2409.10477* (2024).
- [8] I. C. GOHBERG, A. S. MARKUS, and I. A. FELDMAN, Normally solvable operators and ideals associated with them, *Amer. Math. Soc. Transl. Ser. 2* **61** (1967), 63–84.
- [9] W. T. GOWERS and B. MAUREY, The unconditional basic sequence problem, *J. Amer. Math. Soc.* **6** (1993), 851–874.
- [10] W. B. JOHNSON, T. KANIA, and G. SCHECHTMAN, Closed ideals of operators on and complemented subspaces of Banach spaces of functions with countable support, *Proc. Amer. Math. Soc.* **144** (2016), 4471–4485.
- [11] T. KANIA and T. KOCHANEK, The ideal of weakly compactly generated operators acting on a Banach space, *J. Oper. Theory* **71** (2014), 455–477.

- [12] P. KOSZMIDER, On decompositions of Banach spaces of continuous functions on Mrówka's spaces, *Proc. Amer. Math. Soc.* **133** (2005), 2137–2146.
- [13] P. KOSZMIDER and N. J. LAUSTSEN, A Banach space induced by an almost disjoint family, admitting only few operators and decompositions, *Adv. Math.* **381** (2021), 107613.
- [14] P. KOSZMIDER and P. ZIELIŃSKI, Complementation and decompositions in some weakly Lindelöf Banach spaces, *J. Math. Anal. Appl.* **376** (2011), 329–341.
- [15] N. J. LAUSTSEN, Maximal ideals in the algebra of operators on certain Banach spaces, *Proc. Edinb. Math. Soc.* **45** (2002), 523–546.
- [16] H. P. LOTZ, N. T. PECK and H. PORTA, Semi-embeddings of Banach spaces, *Proc. Edinb. Math. Soc.* **22** (1979), 233–240.
- [17] A. PEŁCZYŃSKI, Projections in certain Banach spaces, *Studia Math.* **19** (1960), no. 2, 209–228. Available at: <http://eudml.org/doc/216957>.
- [18] R. A. RYAN, *Introduction to tensor products of Banach spaces*, Springer, 2002.
- [19] S. SHELAH, A Banach space with few operators, *Isr. J. Math.* **30** (1978), 181–191.
- [20] S. SHELAH and J. STEPRĀNS, A Banach space on which there are few operators, *Proc. Amer. Math. Soc.* **104** (1988), 101–105.
- [21] H. M. WARK, A non-separable reflexive Banach space on which there are few operators, *J. London Math. Soc.* **64** (2001), 675–689.

SCHOOL OF MATHEMATICAL SCIENCES, FYLDE COLLEGE, LANCASTER UNIVERSITY, LA1 4YF, UNITED KINGDOM
Email address: ahacua@gmail.com