# YOUNG STRUCTURES FOR PARTIALLY HYPERBOLIC SYSTEMS WITH MOSTLY CONTRACTING CENTRAL DIRECTION

JOSÉ F. ALVES AND JOÃO S. MATIAS

ABSTRACT. We establish the existence of Young structures for a broad class of partially hyperbolic diffeomorphisms with a splitting  $TM = E^{cs} \oplus E^{uu}$ , under exactly the same conditions that ensure the existence of SRB measures in a previous work by Bonatti and Viana. This extends the applicability of statistical techniques to systems where statistical properties remained largely unexplored. Our approach refines previous methods, introducing key adaptations to the partially hyperbolic setting. These results provide a foundation for obtaining decay of correlations, Central Limit Theorem, large deviations and a vector-valued almost sure invariance principle in this class of dynamical systems.

## 1. INTRODUCTION

The study of chaotic systems and their unpredictable behavior has shifted the focus of dynamical systems research toward understanding their statistical properties. A key tool in this pursuit is the concept of inducing, which involves simplifying a complex dynamical system by substituting it with one that is more manageable while retaining the ability to extract meaningful insights about the original system. This approach originated in the 1970s with the use of Markov partitions, enabling the analysis of uniformly hyperbolic systems through their conjugation with symbolic shifts. Since then, expanding this methodology to a broader range of systems has been a primary objective in the field.

A significant milestone in the study of dynamical systems was achieved with Young's works [32, 33] in the late 1990s, where the abstract framework of *Young towers* or *Young structures* has been introduced. This framework has been crucial in systematically analysing a diverse array of dynamical systems, including Axiom A attractors, piecewise hyperbolic maps, billiards with convex scatterers, logistic maps, intermittent maps, and Hénon-like attractors – the latter explored in detail by Benedicks and Young in [10]. Young's methodology facilitates the rigorous investigation of statistical properties such as the existence of SRB measures, decay of correlations, large deviations and the Central Limit Theorem. Central to this approach is the partitioning of the phase space into subsets with associated *recurrence times*, providing a robust tool for studying nonuniformly hyperbolic systems.

<sup>2020</sup> Mathematics Subject Classification. 37A05, 37A25, 37D25, 37D30, 60F05, 60F10.

*Key words and phrases.* Partial hyperbolicity, Young structures, Decay of correlations, Large deviations, Central Limit Theorem, Invariance Principle.

JFA and JSM are partially supported by CMUP (UID/MAT/00144/2019), PTDC/MAT-PUR/28177/2017 and PTDC/MAT-PUR/4048/2021, which are funded by FCT (Portugal) with national (MEC) and European structural funds through the program FEDER, under the partnership agreement PT2020. JSM is also supported by the FCT doctoral scholarship 2021.07090.BD.

The relationship between Young structures and the statistical properties of systems without contracting directions – noninvertible systems – was further developed in [4]. More recently, [7] expanded these results by demonstrating analogous conclusions for certain dynamical systems with contracting directions, thereby extending the scope and impact of this foundational framework.

Smooth dynamical systems  $f: M \to M$  on a Riemannian manifold M that do not belong to the class of uniformly hyperbolic systems, yet exhibit a combination of non-uniformly expanding or contracting central directions alongside other directions with uniform hyperbolic behavior, give rise to a broader category of *partially hyperbolic diffeomorphisms*. These systems naturally extend the classical theory of hyperbolic dynamics and encompass a rich variety of behaviors that bridge uniform hyperbolicity and more general nonuniformly hyperbolic phenomena. Partially hyperbolic diffeomorphisms can be categorised into two main types, depending on the structure of their invariant splitting: the tangent bundle TM admits a Df-invariant dominated splitting of the form

$$TM = E^{ss} \oplus E^{cu}$$
 or  $TM = E^{cs} \oplus E^{uu}$ ,

where  $E^{ss}$  is the strongly stable sub-bundle, exhibiting uniform contraction under Df;  $E^{uu}$  is the strongly unstable sub-bundle, displaying uniform expansion under Df;  $E^{cu}$  (center-unstable) and  $E^{cs}$  (center-stable) contain directions that may exhibit nonuniform expansion or contraction, respectively. The two types above encapsulate distinct dynamical behaviors, where the interplay between uniform and nonuniform hyperbolicity plays a crucial role in determining the ergodic and statistical properties of the system. Understanding the structure and dynamics of these sub-bundles is essential for analysing the long-term behavior of trajectories, particularly in the context of SRB measures and statistical stability.

In the case where the tangent bundle splits as  $TM = E^{ss} \oplus E^{cu}$ , the existence of SRB measures was established by Alves, Bonatti, and Viana [2] under the crucial assumption that Df exhibits nonuniform expansion along the  $E^{cu}$  direction. Their approach introduced the innovative concept of hyperbolic times, a key technique that allowed them to capture nonuniform expansion in a rigorous manner and construct SRB measures in this setting. Further developments in [3, 5, 6, 8, 9] demonstrated that hyperbolic times not only play a fundamental role in establishing the existence of SRB measures but also serve as a powerful tool for constructing Young structures, which provide a geometric framework for studying statistical properties of dynamical systems. In addition, these works uncovered a deep connection between the tail distribution of hyperbolic times and the tail behavior of recurrence times, further enriching the understanding of the interplay between nonuniform expansion and recurrence properties. By integrating these findings with earlier results from [24, 26, 32, 33], several fundamental statistical properties of SRB measures were derived, including exponential decay of correlations, the Central Limit Theorem, large deviations and a vector-valued almost sure invariance principle. Collectively, these contributions have led to a more comprehensive understanding of the statistical behavior of partially hyperbolic diffeomorphisms in the  $E^{ss} \oplus E^{cu}$  setting, under essentially the same assumptions that originally ensured the existence of SRB measures in [2]. This body of work has significantly expanded the scope of ergodic theory in nonuniformly hyperbolic dynamics, bridging the gap between theoretical existence results and concrete statistical properties.

Building upon the foundational work of Pesin and Sinai [28], Bonatti and Viana [11] established the existence of SRB measures for a class of partially hyperbolic diffeomorphisms with a splitting  $TM = E^{cs} \oplus E^{uu}$ , under the additional assumption that all Lyapunov exponents along the  $E^{cs}$  direction are negative. This result provided a significant advance in the understanding of the ergodic properties of such systems. However, despite the existence of SRB measures, a comprehensive statistical description of these dynamical systems remained largely unexplored. Specifically, general results concerning the statistical properties of SRB measures in a wide setting are still not available. To our knowledge, the most notable progress in this direction was achieved in the early 2000s, for specific subclasses of partially hyperbolic diffeomorphisms and, more recently, for fast-slow systems. The following cases represent the main instances where statistical properties were successfully derived in this context:

- Three-dimensional manifolds: Partially hyperbolic diffeomorphisms for which the central-stable bundle  $E^{cs}$  further decomposes as  $E^{cs} = E^{ss} \oplus E^{cs}$ , thereby introducing an additional level of hyperbolicity; see [20].
- **Derived from Axiom A systems**: Partially hyperbolic systems constructed from Axiom A dynamics with a *Markov partition*, where nonuniform contraction is restricted to a single rectangle; see [15].
- Systems with a pseudo-product structure: Partially hyperbolic diffeomorphisms exhibiting weak contraction in a region with a *pseudo-product structure* and a *one-dimensional* unstable sub-bundle  $E^{uu}$ ; see [16].
- Systems with an almost Markov family: Partially hyperbolic diffeomorphisms with an *almost Markov family* for which the images of all elements under iterations become uniformly distributed; see [21].
- **Two-dimensional fast-slow systems**: Partially hyperbolic endomorphisms of the two-torus with a strongly dominated direction; see [14, 18, 19].

The introduction of Young structures in [15, 16] provided a powerful framework for analysing statistical properties of SRB measures. This approach enabled the derivation of key probabilistic results, including the Central Limit Theorem, decay of correlations, large deviations and almost sure invariance principles, offering a deeper insight into the long-term statistical behavior of trajectories.

The primary objective of this work is to develop a comprehensive framework for establishing the existence of Young structures in a broader class of partially hyperbolic diffeomorphisms. Our approach builds upon and refines existing statistical techniques, incorporating new insights to extend their applicability beyond previously studied settings. To be specific, we construct Young structures under the same conditions that originally guaranteed the existence of SRB measures in [11]. While our methodology is inspired by ideas by Chernov [17] and Young [32], we introduce modifications to adapt them to the partially hyperbolic setting, ensuring a more general and flexible application.

In the following three subsections, we rigorously define the key mathematical objects involved and present precise statements of our main results. This provides a solid foundation for a systematic analysis of the statistical properties of these dynamical systems, paving the way for a deeper understanding of their ergodic and probabilistic behavior. 1.1. Young structures. Consider M a finite dimensional compact Riemannian manifold. Let d denote the distance on M and m the Lebesgue measure on the Borel sets of M, both induced by the Riemannian metric. Given a submanifold  $\gamma \subset M$ , we use  $m_{\gamma}$  to denote the Lebesgue measure on  $\gamma$ , induced by the restriction of the Riemannian metric to  $\gamma$ .

Consider a diffeomorphism  $f: M \to M$ . We introduce continuous families  $\Gamma^s$  and  $\Gamma^u$  of  $C^1$  disks of M. We will impose some mild forward contraction on the disks of  $\Gamma^s$  (stable disks) and some mild backward contraction on those of  $\Gamma^{u}$  (unstable disks). We say that  $\Gamma$  is a continuous family of  $C^1$  disks in M if there are a compact metric space K, a unit disk D in some  $\mathbb{R}^k$  and an injective continuous function  $\Phi: K \times D \to M$  such that

- $\Gamma = \{ \Phi(\{x\} \times D) : x \in K \};$
- $\Phi$  maps  $K \times D$  homeomorphically onto its image;
- $x \mapsto \Phi|_{\{x\} \times D}$  defines a continuous map from K into  $\operatorname{Emb}^1(D, M)$ , where  $\operatorname{Emb}^1(D, M)$ denotes the space of  $C^1$  embeddings of D into M.

All the disks in  $\Gamma$  have the same dimension as D, that we denote by dim  $\Gamma$ . We say that a compact set  $\Lambda \subset M$  has a product structure if there are continuous families of  $C^1$  disks  $\Gamma^s$ and  $\Gamma^u$  such that

- $\Lambda = \left(\bigcup_{\gamma \in \Gamma^s} \gamma\right) \bigcap \left(\bigcup_{\gamma \in \Gamma^u} \gamma\right);$   $\dim \Gamma^s + \dim \Gamma^u = \dim M;$
- each  $\gamma \in \Gamma^s$  meets each  $\gamma \in \Gamma^u$  in exactly one point.

We say that  $\Lambda_0 \subset \Lambda$  is an *s*-subset if  $\Lambda_0$  has a product structure with respect to families  $\Gamma_0^s$ and  $\Gamma_0^u$  such that  $\Gamma_0^s \subset \Gamma^s$  and  $\Gamma_0^u = \Gamma^u$ ; *u*-subsets are defined similarly. Let  $\gamma^*(x)$  denote the disk in  $\Gamma^*$  containing  $x \in \Lambda$ , for \* = s, u. Consider the holonomy map  $\Theta_{\gamma,\gamma'}: \gamma \cap \Lambda \to \gamma' \cap \Lambda$ , given for each  $x \in \gamma \cap \Lambda$  by

$$\Theta_{\gamma,\gamma'}(x) = \gamma^s(x) \cap \gamma'.$$

We say that a compact set  $\Lambda$  has a Young structure (with respect to f), if  $\Lambda$  has a product structure given by continuous families of  $C^1$  disks  $\Gamma^s$  and  $\Gamma^u$  such that

$$m_{\gamma}(\Lambda \cap \gamma) > 0$$
, for all  $\gamma \in \Gamma^{u}$ ,

and the conditions  $(\mathbf{Y}_1)$ - $(\mathbf{Y}_5)$  below are satisfied.

(Y<sub>1</sub>) Markov: there are pairwise disjoint s-subsets  $\Lambda_1, \Lambda_2, \dots \subset \Lambda$  such that

- $m_{\gamma}((\Lambda \setminus \bigcup_i \Lambda_i) \cap \gamma) = 0$  for all  $\gamma \in \Gamma^u$ ;
- for each  $i \ge 1$ , there is  $\tau_i \in \mathbb{N}$  such that  $f^{\tau_i}(\Lambda_i)$  is a u-subset and, moreover, for all  $x \in \Lambda_i$ ,

$$f^{\tau_i}(\gamma^s(x)) \subset \gamma^s(f^{\tau_i}(x)) \text{ and } f^{\tau_i}(\gamma^u(x)) \supset \gamma^u(f^{\tau_i}(x)).$$

This Markov property allows us to introduce a recurrence time  $\tau \colon \Lambda \to \mathbb{N}$  and a return map  $f^{\tau} \colon \Lambda \to \Lambda$ , defined for each  $i \ge 1$  by

$$\tau|_{\Lambda_i} = \tau_i$$
 and  $f^{\tau}|_{\Lambda_i} = f^{\tau_i}|_{\Lambda_i}$ .

We remark that  $\tau$  and  $f^{\tau}$  are defined on a full  $m_{\gamma}$  measure subset of  $\Lambda \cap \gamma$ , for each  $\gamma \in \Gamma^{u}$ . Thus, there is a set  $\Lambda' \subset \Lambda$  intersecting each  $\gamma \in \Gamma^u$  in a full  $m_{\gamma}$  measure subset, such that  $(f^{\tau})^n(x)$  belongs in some  $\Lambda_i$  for all  $n \ge 0$  and  $x \in \Lambda'$ . For points  $x, y \in \Lambda'$ , we define the separation time

$$s(x,y) = \min \{n \ge 0: (f^{\tau})^n(x) \text{ and } (f^{\tau})^n(y) \text{ lie in distinct } \Lambda_i\text{'s}\},\$$

with the convention that  $\min(\emptyset) = \infty$ . For definiteness, we set the separation time equal to zero for all other points. For the remaining conditions, we consider constants C > 0 and  $0 < \beta < 1$  only depending on f and  $\Lambda$ .

- (Y<sub>2</sub>) Contraction on stable disks: for all  $i \ge 1$ ,  $\gamma \in \Gamma^s$  and  $x, y \in \gamma \cap \Lambda_i$ ,
  - $d\left(\left(f^{\tau}\right)^{n}(x), \left(f^{\tau}\right)^{n}(y)\right) \leqslant C\beta^{n}$ , for all  $n \ge 0$ ;
  - $d(f^j(x), f^j(y)) \leq Cd(x, y)$ , for all  $1 \leq j \leq R_i$ .
- (Y<sub>3</sub>) Expansion on unstable disks: for all  $i \ge 1$ ,  $\gamma \in \Gamma^u$  and  $x, y \in \gamma \cap \Lambda_i$ , •  $d((f^{\tau})^n(x), (f^{\tau})^n(y)) \le C\beta^{s(x,y)-n}$ , for all  $n \ge 0$ ;
  - $d(f^j(x), f^j(y)) \leq Cd(f^\tau(x), f^\tau(y))$ , for all  $1 \leq j \leq R_i$ .
- (Y<sub>4</sub>) Gibbs: for all  $i \ge 1$ ,  $\gamma \in \Gamma^u$  and  $x, y \in \gamma \cap \Lambda_i$ ,

$$\log \frac{\det Df^{\tau}|_{T_x\gamma}}{\det Df^{\tau}|_{T_y\gamma}} \leqslant C\beta^{s(f^{\tau}(x),f^{\tau}(y))}.$$

(Y<sub>5</sub>) Regularity of the stable holonomy: for all  $\gamma, \gamma' \in \Gamma^u$ , the measure  $(\Theta_{\gamma,\gamma'})_* m_{\gamma}$  is absolutely continuous with respect to  $m_{\gamma'}$  and its density  $\rho_{\gamma,\gamma'}$  satisfies

$$\frac{1}{C} \leqslant \rho_{\gamma,\gamma'}(x) \leqslant C \quad \text{and} \quad \log \frac{\rho_{\gamma,\gamma'}(x)}{\rho_{\gamma,\gamma'}(y)} \leqslant C\beta^{s(x,y)}$$

for all  $x, y \in \gamma' \cap \Lambda$ .

We define the *tail of recurrence times* as the set

$$\{\tau > n\} := \{x \in \Lambda \colon \tau(x) > n\}.$$

In the case considered here, we will show that the Lebesgue measure of  $\{\tau > n\}$  on the unstable disks of  $\Gamma^u$  decays exponentially fast to zero as  $n \to \infty$ . This in particular gives the integrability of the return time  $\tau$  with respect to the Lebesgue measure on such disks.

1.2. Partially hyperbolic systems. Now we will clarify what we mean by partial hyperbolicity. Let  $f: M \to M$  be a  $C^2$  diffeomorphism on a finite-dimensional Riemannian manifold M, which admits a forward invariant compact set  $K \subset M$ , a Df-invariant splitting  $T_K M = E^{uu} \oplus E^{cs}$  and a constant  $0 < \lambda < 1$  such that, for some choice of a Riemannian metric on M,

- (1) **E<sup>uu</sup> is uniformly expanding**:  $||Df^{-1}|_{E^{uu}_{f(x)}}|| < \lambda$ , for all  $x \in K$ ;
- (2)  $\mathbf{E^{cs}}$  is dominated by  $\mathbf{E^{uu}}$ :  $\|Df|_{E_x^{cs}}\| \cdot \|Df^{-1}|_{E_{f(x)}^{uu}}\| < \lambda$ , for all  $x \in K$ .

We say that K is a partially hyperbolic set, and say that  $E^{uu}$  is the strong-unstable subbundle and  $E^{cs}$  centre-stable sub-bundle. This is the only type of partial hyperbolicity that we will consider and so when we write partial hyperbolic system or set, we are referring to systems or sets of this type. Throughout the text, we assume that  $f: M \to M$  is a  $C^2$ diffeomorphism satisfying the following properties:

 $(\mathbf{H}_1)$  f has an attractor: there exists a (maximal invariant) compact set

$$\mathcal{A} = \bigcap_{n \in \mathbb{N}} f^n\left(U\right),$$

for some open neighbourhood U of  $\mathcal{A}$  with  $\overline{f(U)} \subset U$ .

 $(\mathbf{H}_2)$  The attractor  $\mathcal{A}$  is partially hyperbolic.

In particular, there exists an unique foliation  $\mathcal{F}^{uu}$  of  $\mathcal{A}$  which is tangent to the strongunstable sub-bundle  $E_x^{uu}$ , at every  $x \in \mathcal{A}$ ; see [13, 23, 29]. Its leaves are  $C^2$  immersed submanifolds of M with uniformly bounded curvature. Given a point  $x \in \mathcal{A}$ , set

$$\lambda_+^{cs}(x) = \limsup_{n \to +\infty} \frac{1}{n} \log \|Df^n|_{E_x^{cs}}\|.$$

This is the largest Lyapunov exponent of f along the central-stable direction, wherever it is defined. By Oseledts Theorem, see [31, Theorem 3.3.10], Lyapunov exponents are defined almost everywhere, with respect to any invariant measure.

(H<sub>3</sub>) For every disk  $D^{uu}$  contained in a leaf of  $\mathcal{F}^{uu}$ , we have  $\lambda^{cs}_+(x) < 0$ , for a positive Lebesgue measure subset of points  $x \in D^{uu}$ .

These are the conditions presented by Bonatti and Viana in [11], under which they establish the existence of SRB measures; see Subsection 1.3 below. Our primary objective is to obtain Young structures with exponential recurrence tails and derive additional statistical properties for such systems.

**Theorem A.** Let  $f: M \to M$  be a  $C^2$  diffeomorphism with a partially hyperbolic attractor  $\mathcal{A} \subset M$  for which  $(\mathbf{H}_1)$ - $(\mathbf{H}_3)$  hold. Then f admits a Young structure for which there exist constants C > 0 and  $0 < \theta < 1$  such that  $m_{\gamma} \{\tau > n\} \leq C\theta^n$ , for all  $\gamma \in \Gamma^u$ .

The proof of Theorem A is structured into four key parts. The first one, detailed in Section 5, involves adapting the construction from [17, 32] to obtain an auxiliary partition. Although this partition does not give rise to a Young structure, it plays a crucial role in our approach. Section 6 establishes the corresponding exponential recurrence rates. In Section 7, we refine the auxiliary partition to construct a new one, also exhibiting exponential recurrence rates, which can be used to build a Young structure. Finally, in Section 8, we check that we have indeed a Young structure, by verifying that  $(\mathbf{Y}_1)$ - $(\mathbf{Y}_5)$ are valid. In Sections 2, 3 and 4 we present preliminary results that are used throughout the text.

1.3. Statistical properties. A useful approach to describing the dynamical behaviour of chaotic systems is through invariant probability measures. In this context, physical measures play a fundamental role, as they characterise the time averages of observables along a large set of orbits with respect to the volume measure. An invariant probability measure  $\mu$  is called a *physical measure* for  $f: M \to M$  if, for a set of positive Lebesgue measure, the following holds for any continuous observable  $\varphi: M \to \mathbb{R}$ :

$$\lim_{n \to +\infty} \frac{1}{n} \sum_{j=0}^{n-1} \varphi(f^j(x)) = \int \varphi \, d\mu.$$

A distinguished subclass of physical measures consists of the *Sinai-Ruelle-Bowen (SRB)* measures, which are ergodic invariant probability measures with at least one nonzero Lyapunov exponent almost everywhere and whose conditional measures on unstable manifolds are absolutely continuous with respect to the Lebesgue measure on these manifolds. SRB measures for partially hyperbolic diffeomorphisms with a mostly contracting central direction were established in [11].

While SRB measures establish laws of large numbers for dynamically defined processes, finer statistical properties necessitate mixing behaviour. The map f is said to be mixing with respect to the probability measure  $\mu$  if

$$\left|\mu\left(f^{-n}(A)\cap B\right)-\mu(A)\mu(B)\right|\to 0 \text{ as } n\to +\infty,$$

for any measurable sets A and B. In general, no specific rate of convergence can be guaranteed. The correlation function of observables  $\varphi, \psi \colon M \to \mathbb{R}$  is defined as

$$\operatorname{Cor}_{\mu}(\varphi,\psi\circ f^{n}) = \left| \int (\varphi\circ f^{n})\psi d\mu - \int \varphi d\mu \int \psi d\mu \right|$$

In some cases, it is possible to determine explicit rates at which  $\operatorname{Cor}_{\mu}(\varphi, \psi \circ f^n)$  decays to zero as n tends to infinity, provided that the observables possess sufficient regularity. It is worth noting that when the observables are taken as characteristic functions of Borel sets, this definition recovers the notion of mixing.

We are also interested in the validity of the *Central Limit Theorem*, which asserts that the deviation of the time-averaged values of an observable  $\varphi \colon M \to \mathbb{R}$  along an orbit from its asymptotic mean follows a normal distribution: there exists  $\sigma \ge 0$  such that

$$\frac{1}{\sqrt{n}}\sum_{j=0}^{n-1} \left(\varphi \circ f^j - \int \varphi d\mu\right) \xrightarrow{\text{distr}} N(0,\sigma), \quad \text{as } n \to \infty.$$

The standard deviation  $\sigma$  is obtained from the variance  $\sigma^2$ , frequently given by

$$\sigma^{2} = \lim_{n \to +\infty} \frac{1}{n} \int \left( \sum_{i=0}^{n-1} \varphi \circ f^{i} - n \int \varphi d\mu \right)^{2} d\mu$$

We say that the exponential large deviations principle holds for  $\varphi: M \to \mathbb{R}$  if, for every  $\epsilon > 0$ , there exists a rate function  $c(\epsilon)$  such that

$$\lim_{n \to +\infty} \frac{1}{n} \log \mu \left( \left| \frac{1}{n} \sum_{i=0}^{n-1} \varphi \circ f^i - \int \varphi d\mu \right| > \epsilon \right) = -c(\epsilon).$$

Finally, given a vector-valued observable  $\varphi \colon M \to \mathbb{R}^p$ , for some  $p \ge 1$ , set

$$\varphi_n = \sum_{j=0}^{n-1} \left( \varphi \circ f^j - \int \varphi \, d\mu \right), \quad \text{for each } n \ge 1.$$

We say that the sequence  $\{\varphi_n\}_n$  satisfies a *p*-dimensional almost sure invariance principle if there exist  $\lambda > 0$  and a probability space supporting a sequence of random variables  $\{\varphi_n^*\}_n$  along with a p-dimensional Brownian motion W(t) such that

- (1)  $\{\varphi_n\}_n$  and  $\{\varphi_n^*\}_n$  are equally distributed; (2)  $\varphi_n^* = W(n) + O(n^{1/2-\lambda})$  as  $n \to \infty$ , almost surely.

As a consequence of Theorem A and previously established results in the literature, we obtain Corollary B below, which addresses the statistical properties of the aforementioned partially hyperbolic diffeomorphisms. We emphasize that none of these results have been obtained in previous related works with the same level of generality as achieved here.

**Corollary B.** Let  $f: M \to M$  be a  $C^2$  diffeomorphism with a partially hyperbolic attractor  $\mathcal{A} \subset M$  satisfying  $(\mathbf{H}_1)$ - $(\mathbf{H}_3)$ . Then, f has some SRB measure  $\mu$ . Moreover,

- (a) there exist  $k \in \mathbb{N}$  and  $0 < \theta < 1$  such that, for each Hölder continuous  $\varphi, \psi \colon M \to \mathbb{R}$ , there exists C > 0 such that  $\operatorname{Cor}_{\mu}(\varphi, \psi \circ f^{kn}) \leq C\theta^{n}$ ;
- (b) the variance  $\sigma^2$  is well defined for all Hölder continuous  $\varphi \colon M \to \mathbb{R}$ , and  $\sigma = 0$  if and only if  $\varphi$  is a coboundary, i.e  $\varphi = \psi \circ f - \psi$ , for some  $\psi \in L^2(\mu)$ ;
- (c) the Central Limit Theorem holds for all Hölder continuous  $\varphi \colon M \to \mathbb{R}$ ;
- (d) if  $\sigma^2 > 0$ , then the exponential large deviations principle holds for all Hölder continuous  $\varphi \colon M \to \mathbb{R}$ ;
- (e) the p-dimensional almost sure invariance principle holds for all Hölder continuous  $\varphi \colon M \to \mathbb{R}^p$  such that  $\varphi \notin Z$ , where Z is a closed subspace of infinite codimension within the space of all Hölder functions.

The existence of the SRB measure and conditions guaranteeing either its finiteness or uniqueness were previously established in [11, Theorem A]. Here, its existence follows from Theorem A and [32, Theorem 1]. The conclusion regarding the exponential decay of correlations is a consequence of Theorem A and [32, Theorem 2]. The results concerning the variance follow from Theorem A and the Lemma in [32, Section 5.1.C]; see also [27, Proposition 4.12]. The conclusion on the Central Limit Theorem is derived from Theorem A and [32, Theorem 3]. The statement on exponential large deviations follows from Theorem A and [25, Theorem 4.1]. Finally, the vector-valued almost sure invariance principle is a consequence of Theorem A and [26, Theorem 1.6].

In general, we can only guarantee that the correlation decay in item (a) holds for some power  $k \ge 1$  of f. However, if the return times associated with the elements of the Young structure given by Theorem A are relatively prime, i.e.  $gcd\{R_i\} = 1$ , then the same result holds for f itself.

See [11, Section 6] for examples of open sets of partially diffeomorphisms of the type considered here, for which the conclusions of Theorem A and Corollary B can be derived; namely, DA attractors and transitive diffeomorphisms where the entire manifold has a partially hyperbolic structure.

# 2. Preliminaries

In this section we state some useful and well known results regarding dynamical systems that satisfy conditions  $(\mathbf{H}_1)$ - $(\mathbf{H}_3)$ . It was proved in [11] that the points x for which  $\lambda_+^{cs}(x) < 0$  admit a stable disk, transverse to the strong-unstable leaf passing through x. Before we address that specific result, we need some definitions.

Definition 2.1. Let  $D^{uu}$  be an unstable disk. Given  $\epsilon > 0$ , let  $D^{uu}(\epsilon)$  denote the tubular neighbourhood of radius  $\epsilon > 0$  around  $D^{uu}$ , defined as the image under the exponential map of M of all the vectors of norm less than  $\epsilon$  in the orthogonal complement of  $E_x^{uu}$ , for all  $x \in D^{uu}$ . If  $\epsilon > 0$  is small enough, then the tubular neighbourhood is diffeomorphic to a cylinder. Associated with it there is a canonical projection,  $\pi : D^{uu}(\epsilon) \to D^{uu}$ , that is a  $C^1$  map. We say that a  $C^1$  disk  $\gamma$  crosses  $D^{uu}(\epsilon)$  if it is contained in  $D^{uu}(\epsilon)$  and  $\pi$  induces a diffeomorphism from  $\gamma$  onto  $D^{uu}$ . Definition 2.2. Given a subset  $A_0$  of some unstable disk  $D^{uu}$ , we say that the foliation  $\{W_{loc}^s(x): x \in A_0\}$  is absolutely continuous if there exists a sequence  $(K_n)_{n \in \mathbb{N}}$  of compact subsets of  $A_0$  with  $m(A_0 \setminus K_n)$  converging to zero as  $n \to +\infty$ , and for each n, there exist maps

$$K_n \ni x \mapsto W^s_{loc}(x),$$

associating to every  $x \in K_n$  an embedded  $C^1$  disk  $W^s(x)$  and satisfying:

- (1)  $W_{loc}^s(x)$  depends continuously on the point  $x \in K_n$ . In particular, there is an uniform lower bound for the size of  $W_{loc}^s(x)$ , in the sense that there exists constants  $\delta(n) > 0$  such that the preimage of  $W_{loc}^s(x)$  under the exponential map of M at x contains the graph of a  $C^1$  map defined from the  $\delta(n)$  neighbourhood of 0 in  $E_x^{cs}$  to  $E_x^{uu}$ ;
- (2) given any  $0 < \epsilon < \delta(n)/2$  and any unstable disk  $\gamma$  crossing the tubular neighbourhood  $D^{uu}(\epsilon)$  the holonomy map

$$\Theta_{\gamma} \colon \bigcup_{x \in K_n} (\gamma \cap W^s_{loc}(x)) \to K_n,$$

defined by projection along the leaves of the foliation  $\{W_{loc}^s(x): x \in K_n\}$  is absolutely continuous with respect to the Lebesgue measure, i.e it preserves sets with zero Lebesgue measure.

The proof of the next result can be found in [11, Proposition 2.1].

**Proposition 2.1.** Let  $\lambda_+^{cs}(x) < 0$  for every point in a positive Lebesgue measure subset  $A_0$  of some unstable disk  $D^{uu}$ . Then,

- (a) for every  $x \in A_0$  there exists a  $C^1$  embedded disk  $W^s_{loc}(x)$  tangent to  $E^{cs}_x$  at x such that the diameter of  $f^n(W^s_{loc}(x))$  converges exponentially fast to zero as  $n \to +\infty$ ;
- (b) the  $C^1$  disk  $W^s_{loc}(x)$  depends in a measurable way on the point x and the foliation  $\{W^s_{loc}(x): x \in A_0\}$  is absolutely continuous.

Regarding the unstable direction, in [11] some distortion bounds of iterates of f restricted to strong-unstable leaves are proven. Before we precisely state that property, let us set up some context by introducing cone fields.

Definition 2.3. A continuous cone field  $C = (C_x)_x$  defined on a subset  $V \subset M$  is called *centre-unstable* if it is forward invariant

$$Df(x)C_x \subset C_{f(x)}$$
, for every  $x \in V \cap f^{-1}(V)$ .

We call it *strong-unstable* if it is strictly invariant,  $Df(x)C_x$  is contained in int  $(C_{f(x)} \cup \{0\})$ and every vector in it is uniformly expanding. A continuous cone field is *centre-stable*, respectively, *strong stable* for f if it is centre-unstable, respectively, strong-unstable for f.

From hypothesis (**H**<sub>1</sub>) and (**H**<sub>2</sub>) one deduce that there exists a strong-unstable cone field  $C^{uu}$  on  $\mathcal{A}$ ; see [11]. We say that a disk  $\gamma$  is tangent to  $C^{uu}$  if the tangent space to  $\gamma$  at every point  $x \in \gamma$  is contained in  $C^{uu}$ . Many properties of such tangent curves are studied in various works; see e.g. [11, 15, 16]. The next result is obtained in [11, Lemma 3.1].

**Lemma 2.2.** Given L > 0 there exists L' > 0 such that, given any unstable disk  $\gamma$  with curvature less than L, then every iteration  $f^{j}(\gamma)$  has curvature bounded by L', for all  $j \ge 1$ .

#### J. F. ALVES AND J. S. MATIAS

For the next result, see the proof of [11, Lemma 3.3].

**Proposition 2.3.** Given L > 0 there exists C > 0 such that given any  $C^2$  unstable disk  $\gamma \subset V$ , whose curvature is less than L, and given any n such that diam  $(f^n(\gamma)) \leq 2L$ , then

$$\log \frac{\det Df\left(f^{n}(x)\right) |T_{f^{n}(x)}f^{n}(\gamma)}{\det Df\left(f^{n}(y)\right) |T_{f^{n}(y)}f^{n}(\gamma)} \leqslant Cd\left(f^{n}(x), f^{n}(y)\right),$$

for every  $x, y \in \gamma$ .

### 3. UNSTABLE DISKS

In order to start constructing the Young structure we need to have some control on the growth of certain unstable disks. With that in mind, in this section we prove some results regarding the growth of unstable disks, in a way very similar to what is done in [17]. Heuristically speaking we aim to prove that the images of certain small unstable disks grow exponentially fast "on average" until they reach a certain size.

Definition 3.1. Given  $\delta_0 > 0$ , we say that a connected unstable disk  $\gamma$  is admissible if

- (1) its sectional curvature is at most L everywhere;
- (2) diam $(\gamma) \leq \delta_0$ ;
- (3) its boundary  $\partial \gamma$  is piecewise smooth.

The constant  $\delta_0 > 0$  will be made small along the text. This allows us in particular to assume that the tangent spaces  $T_x \gamma$  are "almost parallel", for every  $x \in \gamma$ . This way, and because of Proposition 2.3, we have that if  $\gamma' \subset f^n(\gamma)$  is another admissible unstable disk, then, for every  $n \ge 1$ ,  $f_*^n m_{\gamma}|_{\gamma'}$  has almost constant density with respect to  $m_{\gamma'}$ . In particular, not only we are able to approximate any admissible unstable disk by a  $d_u$ -dimensional flat domain in  $\mathbb{R}^d$  but we also are able to easily obtain estimates for the measure  $m_{\gamma'}$  using the estimates from the measures  $f_*^n m_{\gamma}|_{\gamma'}$ .

We must keep in mind that the iterates of an admissible unstable disk may not be admissible, since they grow exponentially in size. In order to keep some control, we will partition them into smaller admissible disks.

Definition 3.2. Let  $\gamma$  be an admissible unstable disk and  $\gamma' \subset \gamma$  be an open subset with piecewise smooth boundary. Given  $x \in \gamma'$  we denote by  $\gamma'(x)$  the connected component of  $\gamma'$  that contains x. We say that  $\gamma'$  is *n*-admissible, if the unstable disk  $f^n(\gamma'(x))$  is admissible, for every  $x \in \gamma'$ .

Notice that  $\gamma'$  does not need to be connected. Before we continue, let us fix some notation that will be very useful throughout the text.

Given an unstable disk  $\gamma$  and a *n*-admissible open subset  $\gamma' \subset \gamma$ , we denote by  $r_{\gamma',n}$  the distance from  $f^n(x)$  to the boundary of the connected component of  $f^n(\gamma')$  that contains that point, i.e.

$$r_{\gamma',n}(x) := d\left(f^n(x), \partial\left(f^n(\gamma')\right)\right). \tag{3.1}$$

We also set

$$Z[\gamma, \gamma', n] := \sup_{\epsilon > 0} \frac{m\left(\{x \in \gamma' \colon r_{\gamma', n}(x) < \epsilon\}\right)}{\epsilon m(\gamma)}.$$
(3.2)

Since  $\partial (f^n(\gamma'(x)))$  is piecewise smooth for every  $x \in \gamma'$  we have that the previous supremum is finite. Whenever  $m(\gamma \setminus \gamma') = 0$ , the value  $Z[\gamma, \gamma', n]$  characterizes the "average size" of the components of  $f^n(\gamma')$ , the larger they are, the smaller  $Z[\gamma, \gamma', n]$  is. In particular,  $Z[\gamma, \gamma, 0]$  characterizes the size of  $\gamma$ . Some examples can be found on [17]. While most of the time it will be clear which measure we are considering when we use or compute  $Z[\gamma, \gamma', n]$ , in some proofs we will need to use  $Z[\gamma, \gamma', n]$  in two different contexts. To avoid confusion we will write  $Z_m[\gamma, \gamma', n]$  for it to be clear which measure we are considering.

Definition 3.3. Let  $\gamma$  be an admissible unstable disk. A decreasing sequence of open subsets  $\gamma = \gamma_0 \supset \gamma_1 \supset \gamma_2 \supset \cdots$  is called a *u*-filtration of  $\gamma$  if

- (1) the set  $\gamma_n$  is *n*-admissible, for all  $n \ge 0$ ;
- (2) the set  $\gamma_n$  is dense in  $\gamma$ , for all  $n \ge 0$ .

Set also  $\gamma_{\infty} = \bigcap_{n \ge 0} \gamma_n$ . Notice that every  $\gamma_n$  and  $\gamma_{\infty}$  have full Lebesgue measure. Moreover,  $\gamma_{\infty}$  is totally disconnected. Next, adapting the construction made in [17, Theorem 2.1], we construct a u-filtration of an admissible unstable disk  $\gamma$ , that is going to be used on the construction of the Young structure. To simplify our notation, let us write

$$r_n = r_{\gamma_n, n}$$
 and  $Z_n = Z[\gamma, \gamma_n, n],$  (3.3)

for every  $n \ge 0$ . In this case, the value of  $Z_n$  characterizes the "average size" of the connected components of  $f^n(\gamma_n)$ .

**Proposition 3.1.** There exist constants  $0 < \alpha < 1$  and  $\beta > 0$ , depending only on the map f, such that for any admissible unstable disk  $\gamma$  there is an u-filtration  $(\gamma_n)_{n\geq 0}$  such that, for all  $n \geq 1$ ,

$$Z_n \leqslant \alpha^n Z_0 + \frac{\beta}{\delta_0} \sum_{j=0}^{n-1} \alpha^j.$$

*Proof.* We begin by constructing the dense and open 1-admissible subset  $\gamma_1 \subset \gamma$  satisfying the conclusion. To do so, we consider two separate cases, where the first one is significantly easier to deal with than the last one.

Case 1: Consider first that  $\operatorname{diam}(\gamma) \leq \delta_0 \lambda$ , where  $\lambda$  is the same as in the definition of partial hyperbolicity. In this case  $f(\gamma)$  is admissible and we set  $\gamma_1 = \gamma$ . Thus, for every  $x \in \gamma$ 

$$r_1(x) = d\left(f(x), \partial\left(f(\gamma)\right)\right) \geqslant \lambda^{-1}d\left(x, \partial\gamma\right) = \lambda^{-1}r_0 \tag{3.4}$$

and so

$$m\left(\{x \in \gamma \colon r_1(x) < \epsilon\}\right) \leqslant m\left(\{x \in \gamma \colon r_0(x) < \lambda\epsilon\}\right) \leqslant Z_0 m(\gamma) \lambda\epsilon, \tag{3.5}$$

which proves the conclusion by setting  $\alpha = \lambda$  and any  $\beta > 0$ .

Case 2: Now we assume that  $\operatorname{diam}(\gamma) > \delta_0 \lambda$ . We aim to define an open and dense subset  $\gamma'_1 \subset \gamma$  whose connected components have diameter less or equal to  $\delta_0 \lambda$ . Since  $\delta_0$ is chosen to be sufficiently small, among other properties,  $\gamma$  being admissible implies that  $\gamma$  must be almost flat. Denote by  $d_u$  the dimension of the strong-unstable sub-bundle  $E^{uu}$ . Without loss of generality, we assume that  $\gamma$  is a flat  $d_u$ -dimensional surface in  $\mathbb{R}^d$ with smooth boundary. Fix a coordinate system in  $\mathbb{R}^d$  so that  $\gamma$  is parallel to the first  $d_u$ coordinate axes. This means that, on  $\gamma$ ,  $x_{d_u+1} = \cdots = x_d = 0$ . Assume also that m is the  $d_u$ -dimensional volume on  $\gamma$ . For each  $i = 1, ..., d_u$  we consider the family of parallel hyperplanes  $\{x_i = a_i + m\delta'\}$ , with  $m \in \mathbb{Z}$  and  $\delta' = \delta_0 \lambda / \sqrt{2d_u}$ . For now  $a_i \in [0, \delta')$  are fixed constants which will be determined during this proof. All these hyperplanes together "slice"  $\gamma$  into cubic pieces of diameter  $\delta_0 \lambda / \sqrt{2} < \delta_0 \lambda$ . Thus, the set

$$\gamma_1 := \gamma \setminus \bigcup_{\substack{m \in \mathbb{Z}\\i=1,\dots,d_u}} \{x_i = a_i + k\delta'\}$$

is open and dense in  $\gamma$  and is completely determined by the vector  $(a_1, \ldots, a_{d_u})$ , which is fixed. For each  $i = 1, \ldots, d_u$  and  $k \in \mathbb{Z}$  set

$$D_{k,a_i} = \gamma \cap \{x_i = a_i + k\delta'\}$$

Set also, for  $\epsilon > 0$ 

$$\gamma_{\epsilon}^{0} = \{x \in \gamma : d(x, \partial \gamma) < \epsilon\}$$
 and  $\gamma_{\epsilon}^{1} = \{x \in \gamma : d(x, \partial \gamma_{1}) < \epsilon\}$ .

Notice that  $\partial \gamma_1 = \partial \gamma \cup (\bigcup_{k,i} D_{k,a_i})$ . Now we determine the vector  $(a_1, \ldots, a_{d_u})$  in such a way that  $m(\gamma_{\epsilon}^1)$  is sufficiently small,  $\forall \epsilon > 0$ . With that in mind, let

$$\mathcal{C}_{k,a_i}(\epsilon) = D_{k,a_i} \times [a_i + k\delta' - \epsilon \leqslant x_i \leqslant a_i + k\delta' + \epsilon]$$

be the solid cylinder in  $\mathbb{R}^{d_u}$  of height  $2\epsilon$  whose middle cross-section is  $D_{k,a_i}$ . Since for any  $x \in \gamma_{\epsilon}^1 \setminus \gamma_{\epsilon}^0$  the  $d_u$ -dimensional ball in  $\gamma$  of radius  $d(x, \partial U_1)$ , centred at x, intersects one of the domains  $D_{k,a_i}$ , we deduce that  $\gamma_{\epsilon}^1 \setminus \gamma_{\epsilon}^0$  is covered by the union of the cylinders  $\mathcal{C}_{k,a_i}(\epsilon)$ . Hence,

$$m\left(\gamma_{\epsilon}^{1}\setminus\gamma_{\epsilon}^{0}\right)\leqslant 2\epsilon\sum_{i=1}^{d_{u}}V_{a_{i}},$$

where  $V_{a_i}$  denotes the total  $(d_u - 1)$ -dimensional volume of the domains  $D_{k,a_i}$ , for  $k \in \mathbb{Z}$ . We fix  $a_i \in [0, \delta']$  so that  $V_{a_i}$  takes its minimum value. This way, the set  $\gamma_1$  is finally fixed. Now we estimate  $Z_i$ . To do so, we start to notice that

Now we estimate  $Z_1$ . To do so, we start to notice that

$$m\left(x \in \gamma_1: d(x, \partial \gamma_1) < \epsilon\right) \leqslant m\left(x \in \gamma: d(x, \partial \gamma_1) < \epsilon\right) \leqslant m(\gamma_{\epsilon}^0) + m\left(\gamma_{\epsilon}^1 \setminus \gamma_{\epsilon}^0\right).$$
(3.6)

To estimate  $m(\gamma_{\epsilon}^1 \setminus \gamma_{\epsilon}^0)$ , first observe also that since

$$m(\gamma) = \int_0^{\delta'} V_{a_i} da_i$$

we have that

$$\min_{a_i} V_{a_i} \leqslant \frac{m(\gamma)}{\delta'}$$

Therefore,

$$m\left(\gamma_{\epsilon}^{1} \setminus \gamma_{\epsilon}^{0}\right) \leqslant 2\epsilon d_{u} \frac{m(\gamma)}{\delta'} \leqslant 4d_{u}^{\frac{3}{2}} \delta_{0}^{-1} \lambda^{-1} m(\gamma) \epsilon.$$

$$(3.7)$$

For estimating  $m(\gamma_{\epsilon}^0)$ , notice that, by the definition of  $Z_0$ ,

$$n\left(\gamma_{\epsilon}^{0}\right) \leqslant Z_{0}m(\gamma)\epsilon.$$

Substitution these two last estimates on (3.6) we deduce that

$$m\left(\left\{x \in \gamma_1 \colon d(x, \partial \gamma_1) < \epsilon\right\}\right) \leqslant m(\gamma)\epsilon\left(Z_0 + 4d_u^{\frac{3}{2}}\delta_0^{-1}\lambda^{-1}\right).$$

Therefore,

$$Z_1 = \sup_{\epsilon > 0} \frac{m\left(\left\{x \in \gamma_1 \colon d\left(x, \partial\left(f(\gamma_1)\right)\right) < \epsilon\right\}\right)}{\epsilon m(\gamma)}$$
$$\leqslant \left(Z_0 + 4d_u^{\frac{3}{2}} \delta_0^{-1} \lambda^{-1}\right) \lambda = \alpha Z_0 + \frac{\beta}{\delta_0},$$

where  $\alpha = \lambda$  and  $\beta = 4d_u^{\frac{3}{2}}$ .

Now, to obtain the general case, we proceed by induction. Assume that  $\gamma_{n-1}$  is defined. Thus, every connected component  $\gamma' \subset f^{n-1}(\gamma_{n-1})$  is an admissible unstable disk and so, by the previous argument, it admits a open dense subset  $\gamma'_1 \subset \gamma'$ . Define  $\gamma_n$  as the union of all of  $f^{-n+1}(\gamma'_1)$  over all  $\gamma' \subset f^{n-1}(\gamma_{n-1})$ , i.e.,

$$\gamma_n = \bigcup_{\gamma' \subset f^{n-1}(\gamma_{n-1})} f^{-n+1}(\gamma_1').$$

It is clear that  $\gamma_n$  is *n*-admissible. Moreover for each  $\gamma_n$  we have

$$Z_{f_*^{n-1}m} \left[ \gamma_{n-1}, \gamma_n, 1 \right] \leqslant \alpha Z_{f_*^{n-1}m} \left[ \gamma_{n-1}, \gamma_{n-1}, 0 \right] + \frac{\beta}{\delta_0}$$

Because of the comment after Definition 3.2, and slightly increasing  $\alpha$  and  $\beta$  if needed, we may write, for every  $n \ge 1$ ,

$$Z_n \leqslant \alpha Z_{n-1} + \beta \delta_0^{-1}$$
  
$$\leqslant \alpha \left( \alpha^{n-1} Z_0 + \beta \delta_0^{-1} \left( 1 + \alpha + \dots + \alpha^{n-2} \right) \right) + \beta \delta_0^{-1}$$
  
$$= \alpha^n Z_n + \beta \delta_0^{-1} \left( 1 + \alpha + \dots + \alpha^{n-1} \right),$$

completing the proof.

When n is sufficiently large we can obtain a simpler estimate for  $Z_n$ .

**Corollary 3.2.** Let  $a = -(\log(\alpha))^{-1}$  and  $b = \max\{0, -\log(\delta_0(1-\alpha)/\beta) / \log \alpha\}$ . If  $n \ge a \log(Z_0) + b$ , then

$$Z_n \leqslant \frac{\beta}{\delta_0},$$

where  $\bar{\beta} = 2\beta/(1-\alpha)$ .

*Proof.* Using the sum of a geometric series, we deduce that

$$\sum_{j=0}^{n-1} \alpha^j \leqslant \frac{1-\alpha^n}{1-\alpha} \leqslant \frac{1}{1-\alpha}.$$

Thus, from Theorem 3.1, we have that, for every n,

$$Z_n \leqslant \alpha^n Z_0 + \frac{\beta}{\delta_0(1-\alpha)}$$

Therefore, we are left to prove that if  $n \ge a \log(Z_0) + b$ , then

$$\alpha^n Z_0 \leqslant \frac{\beta}{\delta_0(1-\alpha)}.$$

Indeed, since  $\log(\alpha) < 0$ , we have that

$$\alpha^{n} Z_{0} \leqslant \frac{\beta}{\delta_{0}(1-\alpha)} \iff n \log(\alpha) \leqslant \log\left(\frac{\beta}{\delta_{0}(1-\alpha)Z_{0}}\right)$$
$$\iff n \geqslant -\frac{\log\left(\frac{\delta_{0}(1-\alpha)}{\beta}\right)}{\log(\alpha)} - \frac{\log(Z_{0})}{\log(\alpha)}$$
$$\iff n \geqslant a \log(Z_{0}) + b.$$

From now on, we denote the quantity

$$a\log(Z_0) + b \tag{3.8}$$

by  $n_0$ . For most of the results, the exact value of  $n_0$  will not be important, however, when proving the decay estimates for the recurrence times, this quantity will be useful. Now, take

$$\delta_1 = \frac{\delta_0}{2\bar{\beta}} > 0. \tag{3.9}$$

With this  $\delta_1$ , we can obtain some more information from Corollary 3.2, regarding the fraction of points of  $f^n(\gamma_n)$  that are far away from its boundary.

**Corollary 3.3.** For all  $n \ge n_0$ ,

$$m_{\gamma}\left(\left\{x \in \gamma_n \colon d\left(f^n(x), \partial\left(f^n(\gamma_n)\right)\right) > \delta_1\right\}\right) \geqslant \frac{1}{2}m_{\gamma}(\gamma_n),$$

*Proof.* It follows from Corollary 3.2 that, for  $n \ge n_0$ ,

$$Z_n \leqslant \frac{\beta}{\delta_0} \iff Z[\gamma, \gamma_n, n] \leqslant \frac{\beta}{\delta_0}$$
  
$$\implies \frac{m_{\gamma} \left( \left\{ x \in \gamma_n : d \left( f^n(x), \partial \left( f^n(\gamma_n) \right) \right) < \delta_1 \right\} \right)}{\delta_1 m_{\gamma}(\gamma)} \leqslant \frac{\overline{\beta}}{\delta_0}$$
  
$$\iff m_{\gamma} \left( \left\{ x \in \gamma_n : d \left( f^n(x), \partial \left( f^n(\gamma_n) \right) \right) < \delta_1 \right\} \right) \leqslant \frac{\overline{\beta}}{\delta_0} m_{\gamma}(\gamma) \frac{\delta_0}{2\overline{\beta}}$$
  
$$\iff m_{\gamma} \left( \left\{ x \in \gamma_n : d \left( f^n(x), \partial \left( f^n(\gamma_n) \right) \right) < \delta_1 \right\} \right) \leqslant \frac{1}{2} m_{\gamma}(\gamma).$$

Since  $m_{\gamma}(\gamma) \ge m_{\gamma}(\gamma_n)$ , we get the desired conclusion.

This means that at least one half of the points of  $f^n(\gamma_n)$  are at least at a distance of  $\delta_1$  of the boundary  $\partial f^n(\gamma_n)$ .

### 4. Rectangular Structures

We now focus on sets with a product structure. In our framework, these sets can be seen as rectangles in the Sinai-Bowen sense; see [12, 30]. Here we give some important definitions regarding such rectangles, as well as establish some of its important properties.

Definition 4.1. Fix  $0 < \epsilon < \delta_0$ , with  $\delta_0$  as in Definition 3.1. We say that  $R \subset M$  is a rectangle if, for any  $x, y \in R$ , there exists a stable disk  $W^s_{\epsilon}(x)$  and an unstable disk  $W^u_{\epsilon}(y)$  both with diameter less than or equal to  $\epsilon$ , such that  $W^s_{\epsilon}(x)$  and  $W^u_{\epsilon}(y)$  intersect in exactly one point, that also belongs in R. This point is denoted by [x, y].

In the context of uniform hyperbolicity, the existence of local stable manifolds is standard. As we noticed in Proposition 2.1, the issue is not so simple in our context of partial hyperbolicity. From now on, and in order to guarantee the existence of local stable manifolds with an adequate length, instead of considering the phase space M we work directly in a subspace  $K_n \subset A_0$  (as in Proposition 2.1) whose lower bound on the size of the stable manifolds is at least  $\delta_0$ .

Let us recall some important definitions regarding the rectangles that we will consider.

Definition 4.2. A subrectangle  $R' \subset R$  is called a *u*-subrectangle if  $W^u(x) \cap R = W^u(x) \cap R'$ for all  $x \in R'$ . An *s*-subrectangle can be defined in a very similar way, substituting the unsatable disk  $W^u(x)$  by the stable disk  $W^s(x)$ . We say that a rectangle R' *u*-crosses another rectangle R if  $R \cap R'$  is an *u*-subrectangle in R and an *s*-subrectangle in R'.

Now we introduce some more notation that will be useful on Section 5 to define the return time function of our inducing scheme. We start by remarking that, given an unstable disk  $\gamma$ of diameter smaller than  $\delta_0 > 0$  and a point  $x \in M$ , we can see that any stable disk  $\gamma_{\delta_0}^s(x)$ will intersect  $\gamma$  in at most one point, as long as we assume  $\delta_0$  small enough. Given an unstable disk  $\gamma$  and a point  $x \in M$ , the set

$$\Theta_x(\gamma) = \left\{ y \in \gamma \colon y = \gamma^s_{\delta_0}(x) \cap \gamma \right\}$$

is either empty or contains only one point, obtained via the holonomy map  $\Theta_{\gamma_{\delta_0}(x),\gamma}(x)$ . Whenever  $\Theta_x(\gamma) \neq \emptyset$ , we can define the *s*-distance from x to  $\gamma$  in the following way

$$d^{s}(x,\gamma) = d_{\gamma^{s}_{\delta_{0}}(x)}\left(x,\gamma^{s}_{\delta_{0}}(x)\cap\gamma\right).$$

Notice that, when defined, this distance is always at most  $\delta_0$ .

Definition 4.3. Let  $\gamma$  and  $\gamma'$  be distinct unstable disks with diameter smaller than  $\delta_0$ . We say that  $\gamma'$  overshadows  $\gamma$  if  $\Theta_x(\gamma') \neq \emptyset$  for every point  $x \in \gamma$ . In this case we define

$$d^{s}(\gamma, \gamma') = \sup_{x \in \gamma} d^{s}(x, \gamma')$$

as the *s*-distance between  $\gamma$  and  $\gamma'$ .

Now that we have stated the main definitions for this section, we apply them to the unstable manifolds that play the major role in our construction. Given  $\delta_1 > 0$  as in (3.9), consider

$$A_{\delta_1} := \left\{ x \in M \colon W^u_{\delta_1}(x) \text{ exists } \right\}.$$

Notice that  $\delta_1 > 0$  can be made arbitrarily small, by taking  $\delta_0$  small enough. In particular, we may have  $A_{\delta_1} \neq \emptyset$  and, for  $z \in A_{\delta_1}$ , an admissible local unstable disk

$$W(z) := W^u_{\delta_1/3}(z).$$

It follows from Proposition 3.1 that there exists a u-filtration  $(W_n(z))_{n\in\mathbb{N}}$  of the admissible disk W(z). Set

$$W_{\infty}(z) = \bigcap_{n \ge 1} W_n(z).$$

Let also  $Z_n$  be as in (3.3), with  $\gamma = W(z)$  and  $\gamma_n = W_n(z)$ . Applying Corollary 3.2 and Corollary 3.3, we obtain:

**Corollary 4.1.** For every  $n \ge n_0$ , we have

(a)  $Z_n \leq 1/(2\delta_1);$ (b)  $m_{W(z)}(\{x \in W_n(z) : d(f^n(x), (\partial((f^n(W))(x))) > \delta_1\}) > m_{W(z)}(W_\infty(z))/2.$ 

Consider now a small constant  $\delta_2 \in (0, \delta_1)$  to be specified later.

Definition 4.4. Given  $z \in A_{\delta_1}$ , we define the canonical rectangle R(z) in the following way: we say that  $y \in R(z)$  if and only if  $y = W^s_{\delta_1}(x) \cap \gamma$  for some  $x \in W_{\infty}(z)$  and some unstable disk  $\gamma$  that overshadows W(z) with  $d^s(W(z), \gamma) < \delta_2$ .

Assuming  $\delta_2/\delta_1 < c'$ , where c' > 0 is a constant determined by the minimum angle between the centre-stable and unstable directions, then every unstable disk  $W^u$  that overshadows W(z) and is  $\delta_2$ -close to it will meet all stable manifolds  $W^s_{\delta_1}(x)$ , for every  $x \in W_{\infty}(z)$ . In this case R(z) will be, indeed, a rectangle as we defined in the beginning of this section. For any connected subdomain  $V \subset W(z)$  the set

$$R_V(z) = \{ y \in R(z) \colon W^s(y) \cap V \neq \emptyset \}$$

is an s-subrectangle in R(z). We say that  $R_V(z)$  is the *s*-subrectangle in R(z) with base V. For each element  $W_n(z)$  of the u-filtration, its partition into connected components  $\{V_n\}$ induces a partition of R(z) into s-subrectangles  $\{R_{V_n}\}$  with those connected components as bases. Notice that  $f^n(R_{V_n})$  is still a rectangle that is a component of  $f^n(R(z))$ . On the other hand, it may happen that  $f^n(R(z))$  is not a rectangle, since its size grows exponentially fast.

Next, we establish some results regarding the intersection of the sets  $f^n(R_{V_n})$  and some other canonical rectangle R(z'), for some  $z' \in A_{\delta_1}$ . The next lemma is a straightforward consequence of the continuity of the stable and unstable local disks.

**Lemma 4.2.** There exists  $c_1 > 0$  such that, for every  $z, z' \in A_{\delta_1}$  with  $d(z, z') < c_1 \delta_2$ , the unstable disk  $W^u_{\delta_1}(z)$  overshadows the unstable disk  $W^u_{\delta_1/2}(z')$  and

$$d^{s}\left(W^{u}_{\delta_{1}/2}(z'), W^{u}_{\delta_{1}}(z)\right) \leqslant \delta_{2}.$$

Heuristically, it states that if two unstable manifolds come close to each other at their central points, then they must be close enough to each other all along. Now, we are ready to study the intersection of  $f^n(R_{V_n})$  with R(z'), for some  $z' \in A_{\delta_1}$ . For that, fix  $\delta_3 = c_1 \delta_2$ .

**Proposition 4.3.** Given  $z \in A_{\delta_1}$  and  $n \ge 1$ , let  $V_n$  be a connected component of  $W_n(z)$ and  $x \in V_n$  be such that  $r_{V_n,n}(x) > \delta_1$  and  $d(f^n(x), z') < \delta_3$ , for some  $z' \in A_{\delta_1}$ . Then,  $f^n(R_{V_n})$  u-crosses the rectangle R(z').

*Proof.* There are two things that need to be proved: i)  $f^n(R_{V_n}) \cap R(z')$  is an s-subrectangle of  $f^n(R_{V_n})$ , and ii)  $f^n(R_{V_n}) \cap R(z')$  is a u-subrectangle of R(z').

We begin by proving i). By Lemma 4.2,  $f^n(V_n)$  overshadows W(z') and

$$d^s(W(z'), f^n(V_n)) \leq \delta_2.$$

Thus, for every  $y \in f^n(R_{V_n})$ , we have that  $W^u(y)$  overshadows W(z') and

$$d^{s}(W(z'), W^{u}(y)) \leq \delta_{2}\lambda^{n} \leq \delta_{2}.$$

Since  $2\delta_2$  is the length (along the center-stable direction) of the rectangle R(z'), whose central strip is  $W^u(z')$ , we have that  $f^n(R_{V_n}) \cap R(z')$  is an s-subrectangle of  $f^n(R_{V_n})$ .

Finally, for item *ii*), since  $d^s(W(z'), f^n(V)) \leq \delta_2$  and  $f^n(V_n)$  overshadows W(z'), we have that the  $f^n(R_{V_n})$  extends a little further beyond the rectangle R(z') (along the unstable direction). So, by the definition of u-subrectangle, we deduce that  $f^n(R_{V_n}) \cap R(z')$  is an u-subrectangle of R(z').

### 5. AUXILIARY PARTITION

We are now ready to present the algorithm that will originate a partition on certain unstable disks, which will then give rise to a partition on certain canonical rectangles. The various steps presented here are an adaptation to the partially hyperbolic context of the technique used in [17].

Let us consider a finite  $\delta_3$ -dense subset of  $A_{\delta_1}$ ,

$$\mathcal{Z} = \{z_1, z_2, \dots z_p\}.$$

Using these points, we define  $\mathcal{R} = \bigcup_i R(z_i)$ . Note that  $\mathcal{R}$  is a finite union of rectangles that most likely overlap and do not cover M, or even the support of m. We will partition the sets  $W_{\infty}(z_i)$  into a countable collection of subsets  $W_{\infty}^k(z_i)$ , for  $k \ge 0$ , such that for every  $k \ge 1$  there is an integer  $\tau_k \ge 1$  such that for the s-subrectangle  $R_k \subset R(z_i)$  with base  $W_{\infty}^k(z_i)$  the set  $f^{\tau_k}(R_k)$  will be a u-subrectangle in some  $R(z_i)$ . When this happens we say that we have a proper return of  $R_k$  into  $\mathcal{R}$  and we define a function  $\tau(x)$  on  $W_{\infty}(z_i)$ by  $\tau(x) = \tau_k$  for  $x \in W_{\infty}^k(z_i)$  and  $\tau(x) = +\infty$  for  $x \in W_{\infty}^0(z_i)$ . This map is called the return time. It is important to notice that the sets  $W_{\infty}^k(z_i)$  are  $d_u$ -dimensional Cantor sets. At the end of the process the rectangles  $R_k$  will form a partition of  $\mathcal{R}$ , that we call  $\mathcal{P}$ . Naively, one may think that  $\mathcal{P}$  will give rise to a Young structure, however, as we will see on Section 7, that will not be the case.

The construction of the partition, and the definition of the return time function  $\tau(x)$  will be done in several steps. Moreover, as we will see, this construction is independent of the point  $z_i \in \mathbb{Z}$  we consider. In order to simplify the algorithm we only present it for  $z_1$ . Given this, write

$$R = R(z_1), \quad W = W(z_1), \quad W_{\infty} = W_{\infty}(z_1) \text{ and } W_{\infty}^k = W_{\infty}^k(z_1).$$

During this construction we will also prove some decay estimates, that ultimately will lead us to an exponential return tail bound

$$m_W\left(\{\tau(x) > n\}\right) \leqslant C\theta^n. \tag{5.1}$$

5.1. Initial Growth. First of all, consider  $n_0$  as in (3.8). By Corollaries 3.2 and 3.3 we have  $Z_{n_0} < (2\delta_1)^{-1}$  and  $m_W \left( \left\{ x \in W_{n_0} : r_{W_{n_0},n}(x) \ge \delta_1 \right\} \right) \ge 1/2m_W (W_{n_0})$ . In other words, we take  $n_0$  iterates in order to guarantee that the components of  $f^{n_0}(W_{n_0})$  are large enough and at least half of the points in  $f^{n_0}(W_{n_0})$  are in a distance of at least  $\delta_1$  from the boundary. To simplify our notation, let us write  $W^g = f^{n_0}(W_{n_0})$  and  $\widetilde{m}_{W^g} = f_*^{n_0}m_W|_{W^g}$ , where  $\widetilde{m}_{W^g}$  is the induced Lebesgue measure on  $W^g$ . Now, wherever possible, for every connected component  $V \subset W^g$  we arbitrarily fix a point  $x_V$  satisfying  $d(x_V, \partial V) \ge \delta_1$ . This way  $x_V \in A_{\delta_1}$ , and so there exists  $z_V \in \mathcal{Z}$  such that  $d(x_V, z_V) < \delta_3$ . We fix  $z_V$  as well. At this point we can define the first elements of our partition. For each  $z_V$  fixed we define the set

$$f^{-n_0}\left(V \cap R(z_V)\right)$$

as one of the  $W_{\infty}^k$ 's, defining also  $\tau_k = n_0$ . By Proposition 4.3, we have that  $f^{\tau_k}(R_k)$  is an u-subrectangle in  $R(z_V)$ . As a final remark, notice that we define at most one element of the partition in each component V of  $W^g$ .

Before we address the next step in our construction, let us estimate the proportion of points that make a successful return after  $n_0$  iterations. Every time we define a new element of the partition, we require by construction that the rectangle  $R_k$  that originates it u-crosses some canonical rectangle R(z). Also by construction, each of the rectangles R(z) have the same size. This means that, independently of the choice of the points  $x_V$ and  $z_V$  in the components  $V \subset W^g$ , the elements of the partition,  $W^k_{\infty}$ , defined at time  $n_0$ satisfy

$$\frac{m_W\left(\cup W_\infty^k\right)}{m_W\left(W_{n_0}\right)} > \epsilon_1,\tag{5.2}$$

for some constant  $0 < \epsilon_1 < 1$ .

5.2. Capture. Now that we have defined the earliest return in our construction we proceed to the next step. In a first approach one may think that a good option is to keep iterating the remaining points, until the components of  $f^n(W_n)$  are sufficiently large. However, if we pick arbitrary points  $x_V$  and  $z_V$  the sets  $f^{-n}(V \cap R(z_V))$  may overlap with elements defined at previous iterations. To avoid those possible overlaps we proceed as follows. For every connected component  $V \subset W^g$ , where a point  $x_V$  was picked, we divide it into two sets:

$$V^c = W^u_{\delta_1/2}(x_V)$$
 and  $V^f = W^g \setminus V^c$ .

By definition,  $V^c$  overshadows  $W(z_V)$  and so the element  $f^{-n_0}(V \cap R(z_V))$  defined in the previous step is completely contained inside  $V^c$ . Because of that we say that  $V^c$  is *captured* at the  $n_0$ -th iteration. The component  $V^f$  is said to be *free to move*. For the moment we will ignore the captured components and focus only on the free to move parts of  $W^g$ . We call them  $W^f$ , which we keep iterating by f. Notice that  $W^f$  does not contain any point of the previously defined elements of the partition. Define

$$W_n^f = W^f \cap f^{n_0}(W_{n_0+n}).$$

Notice that  $(W_n^f)_{n\geq 0}$  corresponds to a u-filtration of  $W^f$ , in the sense of Proposition 3.1.

Back to the algorithm, first we need to take a certain number of iterations in order to be sure that the components of  $f^n(W_n^f)$  are sufficiently large. For that we make use, once more, of Corollary 3.2, this time applied to the u-filtration  $(W_n^f)_{n\geq 0}$  and the measure  $\widetilde{m}_{W^g}$ . Thus, there exists some iterate  $n_2 \geq 0$  such that, for all  $n \geq n_2$ 

$$Z\left[W^f, W_n^f, n\right] \leqslant \frac{1}{2\delta_1}$$

In other words, it takes  $n_2$  iterations to recover the average size of the manifolds  $f^n(W_n^f)$ , after we remove the components  $V^c$  of  $W^g$ . Moreover, in the light of Corollary 3.3 we also have that, for  $n \ge n_2$ 

$$\widetilde{m}_{W^g}\left(\left\{x\in W_n^f\colon r_{W_n^f,n}(x)\geqslant \delta_1\right\}\right)\geqslant 1/2\widetilde{m}_{W^g}\left(W_n^f\right),$$

which means that after  $n_2$  iterates at least half of the points in  $f^n(W_n^f)$  are in a distance of at least  $\delta_1$  from its boundary.

The next step is repeating inductively this process of picking points  $x_V$  and  $z_V$  in the large components of the free to move manifold, defining new elements  $f^{-n}(V \cap R(z_V))$ , capturing the disks covering these new elements, moving the remaining part  $n_2$  iterations by f, until the components are large enough for this process to be repeated, and so on.

We remark that the argument used to deduce (5.2) can also be used in this context, allowing us to deduce that the freely moving points are being captured at an exponential rate, i.e., every  $n_2$  iterations of f at least a fraction  $\epsilon_1$  of points is being captured. For each  $x \in W_{\infty}$ , let  $t_0(x)$  be the number of iterations that it takes to capture the image of x.

**Lemma 5.1.** There exist some constants  $0 < \theta_0 < 1$  and  $C_0 > 0$  such that

$$m_W\left(\{t_0 > n\}\right) \leqslant C_0 \theta_0^n m_W\left(W_\infty\right),\tag{5.3}$$

for all  $n \ge 1$ . In particular,  $t_0(x) < +\infty$  for almost every  $x \in W_{\infty}$ .

*Proof.* First, notice that  $t_0(x) = n_0 + kn_2$ , for some  $k \in \mathbb{N}$ . Moreover, from (5.2) we deduce that, after each  $n_2$  iterations, the proportion of points that is not captured is at most  $1 - \epsilon_1$ . Thus,

$$m_W(\{t_0 > n\}) \leqslant C_0(1 - \epsilon_1)^{n/n_2} m_W(W_\infty),$$

for every  $n \ge 0$ .

5.3. **Release.** At this point we deal with the captured parts of the manifolds  $f^n(W_n)$ . With that in mind, suppose that you have a captured component  $V^c$  at time  $n_c \ge n_0$ . By definition  $V^c$  contains an element of the partition built in the previous steps. We denote such set by  $V_R^c$ . Thus, there exists some point  $x_c \in V^c$  that also belongs in  $A_{\delta_1}$  and some point  $z_c \in \mathcal{Z}$  such that  $d(x_c, z_c) < \delta_3$ . Therefore, we may write  $V_R^c = f^{-n_c} (V^c \cap R(z_c))$ . We also define  $V_{\infty}^c = V^c \cap f^{n_c}(W_{\infty})$ . Now we take care of  $V_{\infty}^c \setminus V_R^c$ . To do so, first we define a new u-filtration

$$V_n^c = V^c \cap f^{n_c}(W_{n_c+n})$$

associated with the measure  $\widetilde{m}_{V^c} = f_*^{n_c} m|_{V_c}$ . Next, and similarly to what we have done before, we take  $n_0$  more iterations of f, in order to deduce that, in the light of Corollary 3.2,

$$Z[V^c, V_n^c, n] < \frac{1}{2\delta_1}, \quad \forall n \ge n_0,$$

meaning that it takes more  $n_0$  iterations of f for the components of  $f^n(V_n^c)$  to be sufficiently large. The main difference is that now we need to be sure that when we build new elements

they do not overlap with the ones built before. To solve this possible problem, we will define the point release time  $l(x) \in \mathbb{R}$ , for points  $x \in V_{\infty}^c \setminus V_R^c$ . We say that a point is released at l(x) if the component of  $f^n(V_n^c)$  containing  $f^n(x)$  does not intersect  $f^n(V_n^c)$ , for all  $n \ge l(x)$ . Considering a point  $x \in V_{\infty}^c \setminus V_R^c$ , we have two possibilities: either  $W^s(x)$  exists and intersects  $W_{\delta_1}^u(z_c)$ , or x does not have a local stable disk intersecting  $W_{\delta_1}^u(z_c)$ .

We begin with the first possibility. For those points, set h(x) as the intersection of  $W^s(x)$ with  $W^u_{\delta_1}(z_c)$ , i.e.  $h(x) = [z_c, x]$ . Notice that  $h(x) \notin W_{\infty}(z_c)$ , otherwise h(x) would be a point in  $V^c_R$ . Thus,  $h(x) \in W^u_{\delta_1}(z_c) \setminus W^u_{\delta_1/3}(z_c)$  and so  $\epsilon(x) := d(h(x), W^u_{\delta_1/3}(z_c)) > 0$ . Set

$$l(x) = \log_{\lambda^{-1}} \left( \frac{\delta_0}{\epsilon(x)} \right).$$

Thus,  $f^n(x)$  is at least at a distance of  $\delta_0$  from  $f^n(V_R^c)$ , for each  $n \ge l(x)$ . Hence, the connected component that contains  $f^n(x)$  does not intersect  $f^n(V_R^c)$ .

For the second type of points, the point h(x) does not exist. However, we have two possible scenarios (we are excluding the case of x not having a local stable disk, since those points have measure zero in our domain  $K_n$ ): either  $W^s(x)$  does not intersect such unstable disk, or  $W^s(x)$  intersects some  $W^u(y)$  outside of the rectangle  $R(z_c)$ , with  $y \in R(z_c)$ . In the first case, there is nothing to be done. In the second one, notice that, by the fixed size of the stable manifolds in  $R(z_c)$ , the distance along the center-stable direction between  $W(z_c)$  and  $W^u(x)$  is at least  $\delta_2 > \delta_2/2$ . Moreover, the distance along the unstable direction between yand the intersected unstable disk is at least  $(\delta_1/3)/2 > \delta_2/2$ . Thus,  $d(W^u_{\delta_1/3}(z_c), x) > c'\delta_2/2$ , where c' > 0 is a constant that depends only on the minimum angle between the centrestable and the unstable directions. Hence, taking at least  $\log_{\lambda^{-1}}(2\delta_0/c'\delta_2)$  iterates we guarantee that the image of x will be  $\delta_0$  apart from the image of  $V^c_R$ . Hence, for the second type of points, we may set

$$l(x) = \log_{\lambda^{-1}} \left( \frac{2\delta_0}{c'\delta_2} \right),$$

which does not depend on the point considered, contrasting with the first case. This observation will be relevant later on, when we study the recurrence tails.

5.4. Growth. Now that we have components far apart from the captured components, we need to have control on the size of  $f^n(V_n^c)$ , for each  $n \ge l(x)$ . For that we need to define some new functions.

We say that a connected component V is released at time  $n \ge 0$  if at least one point of V is released at such iterate and none of the points of the component of  $f^i(V_i^c)$  that contains  $f^{-(n-i)}(V)$  is released on a previous iteration. When this happens, we set s(x) = n, for each  $x \in V_{\infty}^c \cap f^{-n}(V)$ . This new function s is called the *component release time*. Notice that  $s(x) \le l(x)$ .

From now on, fix a component release time  $s \ge 0$  and set

$$\widetilde{W} = \widetilde{W}(s) = \bigcup \left\{ V \subset f^s(V_s^c) \colon s(x) = s, \text{ for } x \in V_\infty^c \cap f^{-s}(V) \right\}.$$

Note that W is the collection of all the components of  $f^s(V_s^c)$  that are released on the s-th iterate. Consider the associated measure  $\widetilde{m}_{\widetilde{W}} = f_*^s \widetilde{m}_{V^c}|_{\widetilde{W}}$ . Set also for  $n \ge 0$ 

$$\overline{W}_n = \overline{W} \cap f^s(V_{s+n}^c).$$

Notice that  $(\widetilde{W}_n)_{n \ge 0}$  is a u-filtration of  $\widetilde{W}$ . By Corollary 3.2, there exists some n' > 0 such that, for every  $n \ge n'$ ,

$$Z\left[\widetilde{W},\widetilde{W}_n,n\right] \leqslant \frac{1}{2\delta_1}$$

Set the growth time g(x) as the smallest natural number n' with this property. We remark that g(x) is constant on  $V^c_{\infty} \cap f^{-s}(\widetilde{W})$ , depending only on s. Considering the unstable disk  $\widehat{W} = f^g(\widetilde{W}_g)$ , let  $\widetilde{m}_{\widehat{W}} = f^g_* \widetilde{m}_{\widetilde{W}}|_{\widehat{W}}$ . Corollary 3.2 applied to the measure  $\widetilde{m}_{\widehat{W}}$  gives

$$Z\left[\widehat{W},\widehat{W},0\right] \leqslant \frac{1}{2\delta_1},$$

which means that the components of  $\widehat{W}$  are large enough in order to define new elements of our partition. Afterwards we just repeat all this steps of defining new elements, capturing disks and letting them grow. This iterative process leads us to another function t(x), that represents the *capture time* for  $x \in \widehat{W}$ . Note that t(x) gives the smallest  $t \ge 0$  such that  $f^t(x)$  belongs in a captured disk. It may happen that  $f^t(x)$  is already on a defined element, returning to  $\mathcal{R}$ . Otherwise we need to further iterate.

The idea of the proof of Lemma 5.1 can also be used for  $\widehat{W}$  instead of  $W^g$ , yielding exponential decay for  $\{t(x) > n\}$ .

**Lemma 5.2.** For  $n \in \mathbb{N}$ , we have

$$\widetilde{m}_{\widehat{W}}\left(\{t(x) > n\}\right) \leqslant C_0 \theta_0^n,$$

where  $C_0 > 0$  and  $0 < \theta_0 < 1$  are the constants in Lemma 5.1.

5.5. Growth-Capture-Release Cycle. In this final part we summarize the growthcapture-release cycle that we have defined in the previous steps. Fixing a release time s, we take the union of all components of  $f^s(V_s^c)$  that are released exactly at time s. Those components are iterated g = g(s) times without defining new elements nor capturing disks, until they become sufficiently large. This process is then repeated inductively. On  $f^t(\widehat{W})$ , with  $t \ge g$ , we define new elements that make a successful return to  $\mathcal{R}$  at the time they are defined and the captured points are further iterated, until they are released, starting a new cycle. For  $m_W$ -almost every  $x \in W_\infty$ , this cycle is repeated until the point returns to  $\mathcal{R}$  at the moment of a capture. The points that never return are put on the set  $W_\infty^0$ and we set  $\tau(x) = +\infty$  for those points. This concludes the construction of the partition  $W_\infty = \bigcup_k W_\infty^k$  and the respective return time  $\tau(x)$ .

## 6. RECURRENCE TAILS

The goal of this section is to prove the exponential decay in (5.1) above for  $m_W \{\tau(x) > n\}$ . First, we prove that the points of any captured disk  $V^c$  are released at exponential rate.

**Lemma 6.1.** There exist  $C_1 > 0$  and  $0 < \theta_1 < 1$  such that, for any capture disk  $V^c$ ,

$$\widetilde{m}_{V^c}\left(\{l(x) > n\}\right) \leqslant C_1 \theta_1^n \widetilde{m}_{V^c}(V^c),$$

for  $n \in \mathbb{N}$  sufficiently large.

*Proof.* First, recall that for defining l(x) we considered two families of points. For the points in  $V^c$  such that  $W^s(x)$  exists and intersects  $W^u_{\delta_1}(z_c)$ , we have taken

$$l(x) = \log_{\lambda^{-1}} \left( \frac{\delta_0}{\epsilon(x)} \right).$$

If, on the other hand, x does not have a local stable disk intersecting  $W^u_{\delta_1}(z_c)$ ,

$$l(x) = \log_{\lambda^{-1}} \left( \frac{2\delta_0}{c'\delta_2} \right).$$

Since the right hand side is constant, it will not interfere in the desired estimate, so we only focus on the first type of points. For such a point x, we have

$$l(x) > n \iff \log_{\lambda^{-1}} \left( \frac{\delta_0}{\epsilon(x)} \right) > n \iff \frac{\delta_0}{\epsilon(x)} > \lambda^{-n} \iff \epsilon(x) < \delta_0 \lambda^n$$
$$\iff x \in \left\{ x \in V^c \setminus V_R^c \colon d\left(h(x), W_{\delta_1/3(z_c)}^u\right) < \delta_0 \lambda^n \right\}.$$

To simplify our notation, set

$$A = \left\{ x \in V^c \setminus V_R^c \colon d\left(h(x), W_{\delta_1/3}^u(z_c)\right) < \delta_0 \lambda^n \right\}.$$

Noting that the boundary of  $W^u_{\delta_1/3}(z_c)$  is included in the boundary of  $W^u_{\delta_1}(z_c) \setminus W^u_{\delta_1/3}(z_c)$ , we have

$$A \subset \left\{ x \in W^u_{\delta_1}(z_c) \setminus W^u_{\delta_1/3}(z_c) \colon d\left(x, \partial\left(W^u_{\delta_1}(z_c) \setminus W^u_{\delta_1/3}(z_c)\right)\right) < \delta_0 \lambda^n \right\} =: B.$$

Since the quantity  $Z\left[V^c, W^u_{\delta_1}(z_c) \setminus W^u_{\delta_1/3}(z_c), 0\right]$  is finite, we deduce that

$$\widetilde{m}_{V^c}(A) \leqslant \widetilde{m}_{V^c}(B) \leqslant Z\left[V^c, W^u_{\delta_1}(z_c) \setminus W^u_{\delta_1/3}(z_c), 0\right] \delta_0 \lambda^n \widetilde{m}_{V^c}(V^c),$$

thus obtaining the desired estimate.

The next lemma shows that the released components in the images of a captured disk  $V^c$  grow at an exponential rate. To simplify, we assume  $\widetilde{m}_{V^c}(V^c) = 1$ .

**Lemma 6.2.** There exist  $C_2 > 0$  and  $0 < \theta_2 < 0$  such that, for any captured disk  $V^c$  and any  $n \ge 0$ , we have

$$\widetilde{m}_{V^c}\left(\{s(x) + g(x) > n\}\right) < C_2 \theta_2^n.$$

*Proof.* Fix the value  $s(x) = s \ge 0$  and set

$$p(s) = \frac{\widetilde{m}_{\widetilde{W}}\left(\widetilde{W} \cap f^s V_{\infty}^c\right)}{\widetilde{m}_{\widetilde{W}}\left(\widetilde{W}\right)}$$

Recall that on  $V_{\infty}^c \cap f^{-s}\widetilde{W}(s)$  the functions s(x) = s and g(x) = g(s) = g are constant. By Corollary 3.2, we have

$$g \leqslant a \log \left( Z_{\widetilde{m}_{\widetilde{W}}} \left[ \widetilde{W}, \widetilde{W}, 0 \right] \right) + b.$$

Slightly changing the constants a and b, we can also write

$$g \leq a_1 \log \left( Z_{\widetilde{m}_{\widetilde{W}}} \left[ \widetilde{W}, \widetilde{W}, 0 \right] - \log(p(s)) \right) + b_1.$$

Set also  $q(s) = \widetilde{m}_{V^c}(f^{-s}\widetilde{W}(s)) = \widetilde{m}_{\widetilde{W}}(\widetilde{W})$ . By Corollary 3.2 applied to the measure  $\widetilde{m}_{V^c}$ , we have

$$Z_{\widetilde{m}_{V^c}}\left[V^c, \widetilde{W}, s\right] \leqslant C',$$

for some constant C' > 0. Thus,

$$Z_{\widetilde{m}_{\widetilde{W}}}\left[\widetilde{W},\widetilde{W},0\right] = Z_{\widetilde{m}_{V^c}}\left[V^c,\widetilde{W},s\right]q(s)^{-1} \leqslant C'q(s)^{-1},$$

where the equality is true since  $\widetilde{m}_{V^c}(V^c) = 1$ . Therefore,  $g \leq -a_1 \log(p(s)q(s)) + k$ , for some k > 0. Thus,

$$p(s)q(s) \leqslant ke^{-\frac{g}{a_1}}.$$

On the other hand, since  $\widetilde{m}_{V^c}(f^{-s}\widetilde{W}(s)) = \widetilde{m}_{\widetilde{W}}(\widetilde{W})$  and  $\widetilde{m}_{\widetilde{W}} = f_*^s \widetilde{m}_{V^c}$ , we get

$$p(s)q(s) = \frac{\widetilde{m}_{\widetilde{W}}\left(W \cap f^{s}(V_{\infty}^{c})\right)}{\widetilde{m}_{\widetilde{W}}\left(\widetilde{W}\right)} \widetilde{m}_{V^{c}}\left(f^{-s}\left(\widetilde{W}(s)\right)\right) = \widetilde{m}_{\widetilde{W}}\left(\widetilde{W} \cap f^{s}(V_{\infty}^{c})\right)$$
$$= \widetilde{m}_{V^{c}}\left(f^{-s}\left(\widetilde{W} \cap f^{s}(V_{\infty}^{c})\right)\right) = \widetilde{m}_{V^{c}}\left(f^{-s}\left(\widetilde{W}\right) \cap V_{\infty}^{c}\right).$$

Given that  $l(x) \ge s(x)$ , from Lemma 6.1, we deduce that

$$p(s)q(s) = \widetilde{m}_{V^c} \left( f^{-s} \left( \widetilde{W} \right) \cap V_{\infty}^c \right) \leqslant C_1 \theta_1^s.$$

Setting  $\theta_2 = \max\left\{\theta_1, e^{-a_1^{-1}}\right\}$ , we obtain

$$p(s)q(s) \leqslant K'\theta_2^{s+g}$$

hence yielding the desired estimate.

Now we are ready to establish the decay rate of  $m_W(\{\tau(x) > n\})$ . Take  $x \in W_\infty$ . First, we iterate this point  $t_0(x)$  times, going through the initial growth period. Being captured at the  $t_0(x)$ -th iterate either the point returns to  $\mathcal{R}$  or it goes through at least one cycle of release-growth-capture. Let N(x) be the number of cycles the point x goes through before it makes a return. Set also  $s_i(x), g_i(x)$  and  $t_i(x)$  as the number of iterations that are necessary on the *i*-th cycle so that x goes through the release, growth and capture phases, respectively. We may write

$$\tau(x) = t_0(x) + \sum_{i=1}^{N(x)} s_i(x) + g_i(x) + t_i(x).$$

To prove the desired decay rate we make use of the next general result for any probability measure  $\mathbb{P}$ , whose proof can be found in [17, Proposition 7.3]

**Proposition 6.3.** Let  $(\xi_n)_{n\geq 0}$  be a sequence of independent identically distributed random variables taking values in  $\mathbb{N}$  and satisfying an exponential tail bound

$$\mathbb{P}\left(\{\xi_i=n\}\right)\leqslant C_1\theta_1^n,$$

for some  $C_1 > 0$  and  $0 < \theta_1 < 1$ . Let also K be a random variable independent form all  $\xi_i$ 's, taking values in  $\mathbb{N}$  and satisfying an exponential tail bound

$$\mathbb{P}\left(\{K=n\}\right) \leqslant C_2 \theta_2^n$$

for some  $C_2 > 0$  and  $0 < \theta_2 < 1$ . Then,

$$\mathbb{P}\left(\left\{\sum_{i=0}^{K}\xi_{i}=n\right\}\right)\leqslant C\theta^{n},$$

for some C > 0 and  $0 < \theta < 1$ .

We are going to apply this result with  $\mathbb{P} = m_W$ ,  $\xi_0 = t_0$  and  $\xi_i = s_i + g_i + t_i$ , for  $i \ge 1$ , and K = N. In Lemma 5.1 we obtained the exponential decay rate for  $t_0$ . Lemma 5.2 gives the exponential decay rate for each  $t_i$  and Lemma 6.2 gives the decay rate for each  $s_i + g_i$ . We are left to obtain the exponential decay rate for  $m_W(\{N(x) > n\})$ . For that, we use ideas similar to those in the proof of Lemma 5.1. As remarked to deduce (5.2), a proportion ( $\ge \epsilon_1$ ) of the points in every captured disk makes a return. Since the cycle release, growth, capture ends in the capture phase, at the end of each cycle we know that the fraction of points that do not return, and hence need to go through more cycles, is at most  $1 - \epsilon_1$ . Therefore, for every  $n \ge 0$ , we have

$$m_W(\{N(x) \ge n\}) \le (1 - \epsilon_1)^n.$$

Using Proposition 6.3, we deduce that there exist constants C > 0 and  $0 < \theta < 1$  such that

$$m_W(\{\tau(x)=n\}) \leqslant C\theta^n$$

Thus,

$$m_W\left(\{\tau(x) > n\}\right) \leqslant \sum_{i=n}^{+\infty} m_W\left(\{\tau(x) = i\}\right) \leqslant \sum_{i=n}^{+\infty} C\theta^i \approx C\theta^n,$$

yielding the desired estimate (5.1) for the tail of  $\tau$ .

## 7. The Inducing Scheme

Even though the partition constructed in Section 5 is sufficient to establish the existence of SRB measures (see [17, 32]), it does not satisfy property  $(\mathbf{Y}_1)$  in the definition of a Young structure, since  $f^{\tau}(R_k)$  does not u-cross all the rectangles  $R(z_i)$ . To construct an inducing scheme that satisfies all the required properties, we refine the previously built partition  $\mathcal{P}$ . For this, we employ some ideas from [22].

As we consider a finite collection of canonical rectangles  $R(z_i)$ , there exists a subfamily of rectangles, say  $R_1, \ldots, R_q$ , such that  $\bigcup_{1 \leq i \leq q} R_i$  is  $f^{\tau}$ -invariant and, for every  $1 \leq i, j \leq q$ , there exists some n > 0 such that

$$m\left(\left(f^{\tau}\right)^{n}(R_{i})\cap R_{j}\right)>0.$$
 (7.1)

Fix one of those rectangles satisfying  $m(R_i) > 0$ . Without loss of generality, we may assume that the selected rectangle is  $R_1$  and denote it by  $R^*$ . This rectangle  $R^*$  will be

the base set for the Young structure. For the sake of notational simplicity we still denote by  $\mathcal{P}$  the partition obtained before restricted to  $\bigcup_{1 \le i \le q} R_i$ . Set also

$$\mathcal{P}^n = \bigvee_{i=0}^{n-1} \left( f^{\tau} \right)^{-i} \left( \mathcal{P} \right).$$

Notice that the iterations of an element of  $\mathcal{P}^n$  by  $f^{\tau}$  always u-cross one of the canonical rectangles  $R_i$ , with  $1 \leq i \leq q$ . For each  $x \in R^*$ , we are going to define a sequence of stopping times

$$S_0(x) = 0 < S_1(x) < S_2(x) < \cdots$$

that will allow us to define a new return time  $\tau^*$  and also help us to estimate its tails. Moreover, we will also define a positive integer k(x) that, in some sense, tell us the number of iterations needed for x to make a return to  $R^*$ ; the meaning of k(x) will be made clear through out the construction.

The partition of the rectangle  $R^*$ , that we denote by  $\mathcal{P}^*$ , will be obtained inductively in the following way: given  $x \in \omega \in \mathcal{P}$  and  $\tau(x)$  the return time of x (recall that  $\tau$  is constant on  $\omega$ ), set  $S_1(x) = \tau(x)$ . If  $f^{S_1(x)}(\omega)$  is a u-subset of  $R^*$ , then we define k(x) = 1 and stop the process for x and, consequently, for every point in  $\omega$ . In this case  $\omega$  is selected as an element for  $\mathcal{P}^*$  with return time  $S_1(x)$ . If, on the other hand,  $f^{S_1(x)}(\omega)$  is not a u-subset of  $R^*$ , then, by the construction of  $\mathcal{P}$ ,  $f^{S_1(x)}(\omega) = f^{\tau(x)}(\omega)$  is a u-subset of some other rectangle  $R_i$ , with  $2 \leq i \leq q$ . Let  $\omega'$  be the element of  $\mathcal{P}$  where  $f^{\tau}(x)$  belongs and set  $\omega^1 = \omega \cap f^{-\tau}(\omega')$ . Notice that  $x \in \omega^1 \in \mathcal{P}^1$ . In this case, we define

$$S_2(x) = S_1(x) + \tau \left( f^{\tau}(x) \right) = S_1(x) + S_1 \left( f^{\tau}(x) \right).$$

Now, if  $f^{S_2(x)}(\omega^1)$  is a u-subset of  $R^*$ , then we fix k(x) = 2 and stop the process for x and, consequently, for every point in  $\omega^1$ . The set  $w^1 \in \mathcal{P}^1$  is then selected as an element of the new partition  $\mathcal{P}^*$  with return time equal to  $S_2(x)$ . In case  $f^{S_2(x)}(\omega^1)$  is not a u-subset of  $R^*$ , then we keep iterating by  $f^{\tau}$  until it u-crosses  $R^*$ ; note that by (7.1) this will eventually happen. Proceeding inductively, we obtain the subsequent  $S_i$ 's and k so that

$$k(x) = k\left(f^{\tau}(x)\right) + 1$$

and, for  $j \leq k(x)$ ,

$$S_{i}(x) = S_{i-1} \left( f^{\tau}(x) \right) + S_{1}(x).$$

By definition, k(x) is the smallest positive integer  $n \ge 1$  such that the element of  $\mathcal{P}^n$  that contains x is a u-subset of  $R^*$ . That element  $\omega \in \mathcal{P}^n$  with  $x \in \omega$  is selected to be part of the new partition of  $R^*$  with return time  $\tau^*(x) = S_{k(x)}(x)$ . Notice that  $S_{k(x)}(x)$  is constant on the element of the partition  $\mathcal{P}^n$  to which x belongs. We remark that  $\tau^*$  can be seen as the first return to  $R^*$  under the map  $f^{\tau}$ . Thus, since (7.1) holds, we have  $\tau^*(x) < +\infty$  for  $m_W$  almost every point.

In contrast with the partition  $\mathcal{P}$ , this new partition  $\mathcal{P}^*$  satisfies property  $(\mathbf{Y}_1)$ . Before we address the study of the properties  $(\mathbf{Y}_1)$ - $(\mathbf{Y}_5)$ , we state the following result on the tail of  $\tau^*$ , whose proof can be found in [22, Lemma 4.2] in a more general version.

**Lemma 7.1.** Let  $k: \mathbb{R}^* \to \mathbb{N}$  and  $S_0, S_1, \ldots : \mathbb{R}^* \to \mathbb{N}$  be measurable maps such that  $0 = S_0 < S_1 < \cdots$  almost everywhere and set  $\tau^*(x) = S_{k(x)}(x)$ . Suppose that there exists

 $\epsilon > 0$  such that, for any  $\omega \in \mathcal{P}^{j-1}$ ,

$$\frac{m_W\left(\omega \cap \{\tau^* = S_j\}\right)}{m_W\left(\omega\right)} > \epsilon,\tag{7.2}$$

and there exist C > 0 and  $0 < \theta < 1$  such that, for any  $\omega \in \mathcal{P}^{j-1}$ ,

$$\frac{m_W\left(\omega \cap \{S_j - S_{j-1} > n\}\right)}{m_W\left(\omega\right)} \leqslant C\theta^n.$$
(7.3)

Then,

$$m_W(\{\tau^* > n\}) \leqslant C'\theta^n,$$

for some constant C' > 0.

We are now ready to establish the desired tail estimates.

**Proposition 7.2.** There exist constants C > 0 and  $0 < \theta < 1$  such that for every  $n \ge 1$ 

 $m_W\left(\{\tau^* > n\}\right) \leqslant C\theta^n.$ 

*Proof.* By the previous lemma, it is enough to verify conditions (7.2) and (7.3). First of all, notice that since  $\mathcal{P}^*$  is a refinement of the previous partition  $\mathcal{P}$ , just as in the construction of  $\mathcal{P}$ , the proportion of points that is selected at each iteration has a lower bound, say  $0 < \epsilon_2 < 1$ ; see (5.2). In other words, given  $\omega \in \mathcal{P}^{n-1}$ , we have

$$\frac{m_{W}\left(\omega \cap \{\tau^{*} = S_{n}\}\right)}{m_{W}\left(\omega\right)} > \epsilon_{2}$$

This yields condition (7.2) in Lemma 7.1.

Now, if  $k(x) \ge j$  and  $S_j(x) - S_{j-1}(x) > n$ , then we must take at least *n* iterations before  $f^{S_{j-1}(x)}(x)$  makes a return, meaning that  $\tau(f^{S_{j-1}(x)}(x)) > n$ . Therefore,

$$\frac{m_{\gamma}\left(\omega \cap \{S_j - S_{j-1} > n\}\right)}{m_{\gamma}\left(\omega\right)} \leqslant C\theta_3^n,$$

which gives condition (7.3).

## 8. Young Structure

In this section, we construct a Young structure  $\Lambda$  making use of the partition  $\mathcal{P}^*$  introduced in Section 7.

8.1. **Product structure.** We begin by defining the families  $\Gamma^s$  and  $\Gamma^u$  of stable and unstable disks, respectively, that originate our Young structure  $\Lambda$ . Let  $z \in \mathcal{Z}$  be such that  $R(z) = R^*$ . Set

$$\Gamma^s = \left\{ W^s_{\delta_1}(x) \colon x \in W_\infty(z) \text{ and } W^s_{\delta_1} \neq \emptyset \right\}$$

and

$$\Gamma^{u} = \left\{ W^{u}_{loc} : \begin{array}{l} W^{u}_{loc} \text{ intersects every } \gamma^{s} \in \Gamma^{s}(z) \text{ and} \\ \text{overshadows } W^{u}_{\delta_{1}/3}(z) \text{ with } d^{s} \left( W^{u}_{\delta_{1}/3}(z), W^{u}_{loc} \right) \leqslant \delta_{2} \end{array} \right\}.$$

26

The set with a Young structure is then

$$\Lambda = \left(\bigcup_{\gamma \in \Gamma^s} \gamma\right) \bigcap \left(\bigcup_{\gamma \in \Gamma^u} \gamma\right).$$

Property  $(\mathbf{Y}_1)$  holds by the construction of  $\mathcal{P}^*$ .

8.2. Contraction, expansion and bounded distortion. Regarding  $(\mathbf{Y}_2)$ - $(\mathbf{Y}_4)$ , note that since  $\Gamma^s$  and  $\Gamma^u$  are collections of local stable and unstable disks, respectively, there is nothing to verify in  $(\mathbf{Y}_2)$  and  $(\mathbf{Y}_3)$ . Finally,  $(\mathbf{Y}_4)$  is a consequence of the backward contraction  $(\mathbf{Y}_3)$  together with the next result.

**Proposition 8.1.** For every  $\gamma \in \Gamma^u$ ,  $i \ge 1$  and  $x, y \in \gamma \cap \Lambda_i$  we have that

$$\log \frac{\det Df^{\tau^*}(x)|T_x\gamma}{\det Df^{\tau^*}(y)|T_y\gamma} \leqslant Cd\left(f^{\tau^*}(x), f^{\tau^*}(y)\right).$$

*Proof.* By the Lipschitz continuity of  $\log |\det Df(f^k(x))|T_{f^k(x)}f^k(\gamma)|$  given by Proposition 2.3, we may write

$$\begin{split} \log \frac{\det Df^{\tau^*}(x)|T_x\gamma}{\det Df^{\tau^*}(y)|T_y\gamma} &= \sum_{k=0}^{\tau^*-1} \log \frac{\det Df\left(f^k(x)\right)|T_{f^k(x)}f^k(\gamma)}{\det Df\left(f^k(y)\right)|T_{f^k(y)}f^k(\gamma)} \\ &\leqslant \sum_{k=0}^{\tau^*-1} Ld\left(f^k(x), f^k(y)\right). \end{split}$$

Since x, y are in the same unstable disk  $\gamma$ , it follows that, for every  $k \ge 1$ ,

$$d(f^k(x), f^k(y)) \leqslant \lambda^{\tau^* - k} d\left(f^{\tau^*}(x), f^{\tau^*}(y)\right).$$

Therefore,

$$\log \frac{\det Df^{\tau^*}(x)|T_x\gamma}{\det Df^{\tau^*}(y)|T_y\gamma} \leqslant \sum_{k=0}^{\tau^*-1} Ld\left(f^k(x), f^k(y)\right)$$
$$\leqslant \sum_{k=0}^{\tau^*-1} L(\lambda^{(\tau^*-k)}d\left(f^{\tau^*}(x), f^{\tau^*}(y)\right)$$
$$= \frac{L}{1-\lambda}d\left(f^{\tau^*}(x), f^{\tau^*}(y)\right).$$

Take  $C = L/(1 - \lambda)$ .

8.3. Regularity of the stable holonomy. Here we prove that  $(Y_5)$  holds. As we have already proved Proposition 8.1, by [1, Theorem 4.20], we only have to verify that

**Proposition 8.2.** Let  $\gamma, \gamma' \in \Gamma^u$ ,  $x \in \gamma \cap \Lambda_i$ , for some  $i \ge 1$  and  $x' = \Theta_{\gamma,\gamma'}(x)$ , then, for  $n \in \mathbb{N}$ ,

$$\sum_{i=n}^{+\infty} \log \frac{\det Df\left(f^{i}(x)\right) |T_{f^{i}(x)}f^{i}(\gamma)|}{\det Df\left(f^{i}(x')\right) |T_{f^{i}(x')}f^{i}(\gamma)|} \leqslant C'\beta^{n}.$$

#### J. F. ALVES AND J. S. MATIAS

*Proof.* By the Lipschitz continuity of  $\log |\det Df(f^i(x))|T_{f^i(x)}f^i(\gamma)|$  given by Proposition 2.3, we have

$$\sum_{i=n}^{+\infty} \log \frac{\det Df(f^{i}(x)) | T_{f^{i}(x)} f^{i}(\gamma)}{\det Df(f^{i}(x')) | T_{f^{i}(x')} f^{i}(\gamma)} \leqslant \sum_{i=n}^{+\infty} Ld(f^{i}(x), f^{i}(x'))$$

Since x' is obtained from x by the holonomy transformation, it means that  $x' \in \gamma^s(x)$ . In particular (Y<sub>2</sub>) holds and so, for every  $n \in \mathbb{N}$ ,

$$d\left(f^n(x), f^n(x')\right) \leqslant C\beta^n$$

Hence,

$$\sum_{i=n}^{+\infty} d\left(f^{i}(x), f^{i}(x')\right) \leqslant C \sum_{i=n}^{+\infty} \beta^{n} = \frac{C}{1-\beta} \beta^{n}$$

Setting  $C' = LC/(1 - \beta)$ , we get the conclusion.

At this point we have constructed a Young structure whose recurrence tails are of exponential type, thus having proved Theorem A.

### References

- [1] ALVES, J. F. Nonuniformly hyperbolic attractors. Geometric and probabilistic aspects. Springer Monographs in Mathematics. Springer International Publishing, 2020.
- [2] ALVES, J. F., BONATTI, C., AND VIANA, M. SRB measures for partially hyperbolic systems whose central direction is mostly expanding. *Invent. Math.* 140, 2 (2000), 351–398.
- [3] ALVES J. F., DIAS C. L., LUZZATTO S., AND PINHEIRO V. SRB measures for partially hyperbolic systems whose central direction is weakly expanding. J. Eur. Math. Soc. (JEMS) 19, 10 (2017), 2911–2946.
- [4] ALVES, J. F., FREITAS, J. M., LUZZATTO, S., AND VAIENTI, S. From rates of mixing to recurrence times via large deviations. Adv. Math. 228, 2 (2011), 1203–1236.
- [5] ALVES, J. F., AND LI, X. Gibbs-Markov-Young structures with (stretched) exponential tail for partially hyperbolic attractors. Adv. Math. 279, 0 (2015), 405 – 437.
- [6] ALVES, J. F., LUZZATTO, S., AND PINHEIRO, V. Markov structures and decay of correlations for non-uniformly expanding dynamical systems. Ann. Inst. H. Poincaré Anal. Non Linéaire 22, 6 (2005), 817–839.
- [7] ALVES, J. F., AND MATIAS, J. S. From decay of correlations to recurrence times in invertible dynamical systems. *preprint arXiv:2401.06024* (2024).
- [8] ALVES J. F., AND PINHEIRO, V. Slow rates of mixing for dynamical systems with hyperbolic structures. J. Stat. Phys. 131, 3 (2008), 505–534.
- [9] ALVES, J. F., AND PINHEIRO, V. Gibbs-Markov structures and limit laws for partially hyperbolic attractors with mostly expanding central direction. *Adv. Math.* 223, 5 (2010), 1706–1730.
- [10] BENEDICKS, M., AND YOUNG, L.-S. Markov extensions and decay of correlations for certain Hénon maps. Astérisque 261 (2000), 13–56. Géométrie complexe et systèmes dynamiques (Orsay, 1995).
- [11] BONATTI, C., AND VIANA, M. SRB measures for partially hyperbolic systems whose central direction is mostly contracting. *Israel J. Math.* 115 (2000), 157–193.
- [12] BOWEN, R. Equilibrium states and the ergodic theory of Anosov diffeomorphisms. Lecture Notes in Mathematics, Vol. 470. Springer-Verlag, Berlin, 1975.
- [13] BRIN, M. I., AND PESIN, J. B. Partially hyperbolic dynamical systems. Izv. Akad. Nauk SSSR Ser. Mat. 38 (1974), 170–212.
- [14] CASTORRINI, R., AND LIVERANI, C. Quantitative statistical properties of two-dimensional partially hyperbolic systems. Adv. Math. 409 (2022), Paper No. 108625, 122 pp.

- [15] CASTRO, A. Backward inducing and exponential decay of correlations for partially hyperbolic attractors. Israel J. Math. 130 (2002), 29–75.
- [16] CASTRO, A. Fast mixing for attractors with a mostly contracting central direction. Ergodic Theory Dynam. Systems 24, 1 (2004), 17–44.
- [17] CHERNOV, N. Statistical properties of piecewise smooth hyperbolic systems in high dimensions. Discrete and Continuous Dynamical Systems 5 (1999), 425–448.
- [18] DE SIMOI, J., AND LIVERANI, C. Statistical properties of mostly contracting fast-slow partially hyperbolic systems. *Invent. Math. 206*, 1 (2016), 147–227.
- [19] DE SIMOI, J., AND LIVERANI, C. Limit theorems for fast-slow partially hyperbolic systems. Invent. Math. 213, 3 (2018), 811–1016.
- [20] DOLGOPYAT, D. On dynamics of mostly contracting diffeomorphisms. Comm. Math. Phys. 213, 1 2000, 181–201.
- [21] DOLGOPYAT, D. Limit theorems for partially hyperbolic systems. Trans. Amer. Math. Soc. 356, 4 (2004), 1637–1689.
- [22] GOUËZEL, S. Decay of correlations for nonuniformly expanding systems. Bull. Soc. Math. France 134, 1 (2006), 1–31.
- [23] HIRSCH, M. W., PUGH, C. C., AND SHUB, M. Invariant manifolds. Lecture Notes in Mathematics, Vol. 583. Springer-Verlag, Berlin, 1977.
- [24] MELBOURNE, I. Large and moderate deviations for slowly mixing dynamical systems. Proc. Amer. Math. Soc. 137, 5 (2009), 1735–1741.
- [25] MELBOURNE, I., AND NICOL, M. Large deviations for nonuniformly hyperbolic systems. Trans. Amer. Math. Soc. 360, 12 (2008), 6661–6676.
- [26] MELBOURNE, I., AND NICOL, M. A vector-valued almost sure invariance principle for hyperbolic dynamical systems. Ann. Probab. 37, 2 (2009), 478–505.
- [27] PARRY, W. AND POLLICOTT, M. Zeta functions and the periodic orbit structure of hyperbolic dynamics. Astérisque, 187-188 (1990), 268 pp.
- [28] PESIN, Y. B. AND SINAI, Y. G. Gibbs measures for partially hyperbolic attractors. Ergodic Theory Dynam. Systems 2, 3-4 (1982), 417–438.
- [29] SHUB, M. Global stability of dynamical systems. Springer Science & Business Media, 2013.
- [30] SINAI, J. G. Markov partitions and U-diffeomorphisms. Funkcional. Anal. i Priložen 2, 1 (1968), 64–89.
- [31] VIANA, M., AND OLIVEIRA, K. Foundations of ergodic theory. Cambridge Stud. Adv. Math., 151. Cambridge University Press, Cambridge, 2016, xvi+530 pp.
- [32] YOUNG, L.-S. Statistical properties of dynamical systems with some hyperbolicity. Ann. of Math. (2) 147, 3 (1998), 585–650.
- [33] YOUNG, L.-S. Recurrence times and rates of mixing. Israel J. Math. 110 (1999), 153–188.

JOSÉ F. ALVES, CENTRO DE MATEMÁTICA DA UNIVERSIDADE DO PORTO, RUA DO CAMPO ALEGRE 687, 4169-007 PORTO, PORTUGAL.

Email address: jfalves@fc.up.pt URL: http://www.fc.up.pt/cmup/jfalves

JOÃO S. MATIAS, CENTRO DE MATEMÁTICA DA UNIVERSIDADE DO PORTO, RUA DO CAMPO ALEGRE 687, 4169-007 PORTO, PORTUGAL.

Email address: up201504959@fc.up.pt