

# Reducing of system of partial differential equations and generalized symmetry of ordinary differential equations

I. M. Tsyfra<sup>1</sup>, P. Sitko<sup>1</sup>

March 28, 2025

## Abstract.

Symmetry reductions of systems of two nonlinear partial differential equations are studied. We find ansatzes reducing system of partial differential equations to system of ordinary differential equations. The method is applied to system related to Korteweg – de Vries (KdV) equation, and reaction–diffusion equations. We have also shown the possibility of constructing solution to system of non-evolutionary equations (9), and (13), which contains one or two arbitrary functions.

A well-known method for constructing solutions to nonlinear differential equations, based on the generalized conditional symmetry of differential equations, was proposed in the works [3, 10]. It allows reducing a partial differential equation to a system of ordinary differential equations and thus constructing its solutions. We use the approach of reduction proposed in [4, 7] and their generalization [8, 9]. In this context, it is necessary to mention the important works [5, 6]. Nonlinear evolutionary equation systems related to the KdV equation, non-evolutionary equation systems, and reaction-diffusion equation systems are studied in the article.

Consider the system of ordinary differential equations.

$$\begin{cases} u_{xx} - a(t, x)u = 0, \\ v_{xx} - a(t, x)v = 0, \end{cases} \quad (1)$$

where  $a(t, x)$  is a function on  $t, x$ . Following reference [9], we study the generalized symmetry of system (1). We prove that system (1) admits the following

generalized vector field

$$Q = \left( u_t - \frac{1}{2} \left( 3 \frac{u_x u_{xx}}{u} - u_{xxx} - \frac{u v_{xxx}}{v} + 2 \frac{u_x v_{xx}}{v} + \frac{u v_x v_{xx}}{v^2} \right) \right) \partial_u + \left( v_t - \frac{1}{2} \left( 3 \frac{v_x v_{xx}}{v} - v_{xxx} - \frac{v u_{xxx}}{u} + 2 \frac{v_x u_{xx}}{u} + \frac{v u_x u_{xx}}{u^2} \right) \right) \partial_v, \quad (2)$$

if and only if  $a(t, x)$  satisfies the KdV equation

$$a_t = 6a a_x - a_{xxx}. \quad (3)$$

It means that the infinitesimal criterion of invariance [2]

$$Q^{(2)} \left( u_{xx} - a(t, x)u \right) \Big|_{[K]=0} \equiv 0, \\ Q^{(2)} \left( v_{xx} - a(t, x)v \right) \Big|_{[K]=0} \equiv 0,$$

where  $Q^{(2)}$  is the second prolongation of the generalized vector field  $Q$ , and  $K$  denotes the differential consequences of the system of equations (1) with respect to the variables  $t, x$ , is satisfied in this case. It should be noted that the function included in the system (1) is an arbitrary particular solution of the KdV equation (3). If we succeed in constructing the general solution of system (1), we will derive an ansatz that reduces the system of evolutionary equations

$$\begin{cases} u_t = \frac{1}{2} \left( 3 \frac{u_x u_{xx}}{u} - u_{xxx} - \frac{u v_{xxx}}{v} + 2 \frac{u_x v_{xx}}{v} + \frac{u v_x v_{xx}}{v^2} \right), \\ v_t = \frac{1}{2} \left( 3 \frac{v_x v_{xx}}{v} - v_{xxx} - \frac{v u_{xxx}}{u} + 2 \frac{v_x u_{xx}}{u} + \frac{v u_x u_{xx}}{u^2} \right), \end{cases} \quad (4)$$

to a system of four ordinary differential equations. Note that system (4) is invariant with respect to the Lie transformation groups with infinitesimal generators  $Q_1 = u\partial_u + u\partial_v$ , and  $Q_2 = v\partial_u + v\partial_v$ . Then the system of equations (4) is modified and takes the form

$$\begin{cases} u_t = \frac{1}{2} \left( 3 \frac{u_x u_{xx}}{u} - u_{xxx} - \frac{u v_{xxx}}{v} + 2 \frac{u_x v_{xx}}{v} + \frac{u v_x v_{xx}}{v^2} \right) + \lambda_1 u + \lambda_2 v, \\ v_t = \frac{1}{2} \left( 3 \frac{v_x v_{xx}}{v} - v_{xxx} - \frac{v u_{xxx}}{u} + 2 \frac{v_x u_{xx}}{u} + \frac{v u_x u_{xx}}{u^2} \right) + \lambda_1 u + \lambda_2 v. \end{cases} \quad (5)$$

Given the specific form of the function  $a(t, x)$ , one can find an ansatz that is a solution to the system (1). The variable  $t$  in (1), is then treated as a parameter. Let us take the stationary solution  $a(t, x) = \frac{2}{x^2}$  of equation (3). Then system (1) is integrable by quadratures and the general solution has the form

$$\begin{cases} u(t, x) = c_1(t)x^2 + \frac{c_2(t)}{x}, \\ v(t, x) = c_3(t)x^2 + \frac{c_4(t)}{x}. \end{cases} \quad (6)$$

Using the obtained ansatz, one can then find the system of reduced equations of system (5). This is a system of ordinary differential equations

$$\begin{cases} c_1'(t) = \frac{1}{2}(\lambda_1 c_1(t) + \lambda_2 c_3(t)), \\ c_2'(t) = \frac{1}{2}(24c_1(t) + \lambda_1 c_2(t) + \lambda_2 c_4(t)), \\ c_3'(t) = \frac{1}{2}(\lambda_1 c_1(t) + \lambda_2 c_3(t)), \\ c_4'(t) = \frac{1}{2}(24c_1(t) + \lambda_1 c_2(t) + \lambda_2 c_4(t)). \end{cases} \quad (7)$$

By integrating the reduced system (7) and using (6), we obtain the solution of system of equations (5)

$$\begin{aligned} u(t, x) = & (s_1 + s_2 e^{\frac{1}{2}(\lambda_1 + \lambda_2)t})x^2 + \\ & + \frac{1}{x(\lambda_1 + \lambda_2)}(12s_2\lambda_1 t e^{\frac{1}{2}(\lambda_1 + \lambda_2)t} + 12s_2\lambda_2 t e^{\frac{1}{2}(\lambda_1 + \lambda_2)t} + 12s_1\lambda_1 t + 12s_1\lambda_2 t + \\ & + 2s_3 e^{\frac{1}{2}(\lambda_1 + \lambda_2)t} - 24s_2 e^{\frac{1}{2}(\lambda_1 + \lambda_2)t} + s_4\lambda_1 + s_4\lambda_2), \\ v(t, x) = & \frac{1}{\lambda_2}(s_2\lambda_2 e^{\frac{1}{2}(\lambda_1 + \lambda_2)t} - s_1\lambda_1)x^2 + \\ & + \frac{1}{\lambda_2 x(\lambda_1 + \lambda_2)}(12s_2\lambda_1\lambda_2 t e^{\frac{1}{2}(\lambda_1 + \lambda_2)t} + 12s_2\lambda_2^2 t e^{\frac{1}{2}(\lambda_1 + \lambda_2)t} - 12s_1\lambda_1^2 t \\ & - 12s_1\lambda_1\lambda_2 t + 2s_3\lambda_2 e^{\frac{1}{2}(\lambda_1 + \lambda_2)t} - 24s_2\lambda_2 e^{\frac{1}{2}(\lambda_1 + \lambda_2)t} - s_4\lambda_1^2 - s_4\lambda_1\lambda_2), \end{aligned}$$

where  $s_1, s_2, s_3, s_4$  are real constants.

In what follows, we define the system of ordinary differential equations [9]

$$\begin{cases} u_{xx} + u_x^2 = 0, \\ v_x = vu_x. \end{cases} \quad (8)$$

The second equation in (8) was selected to demonstrate the method's effectiveness. We show that system (8) possesses generalized symmetry given by vector field

$$Q_{ne1} = \left( v^2 u_{xt} - F(u - \ln v) \right) \partial_u + \left( v v_{xt} - e^u H(u - \ln v) \right) \partial_v,$$

so the conditions hold

$$Q_{ne1}^{(2)}(u_{xx} + u_x^2) \Big|_{[K]=0} \equiv 0$$

and

$$Q_{ne1}^{(2)}(v_x - vu_x) \Big|_{[K]=0} \equiv 0,$$

where  $K$  denotes the differential consequences of the system of equations (8) with respect to the variables  $t, x$ . Then, the example of a non–evolutionary system to which the reduction method can be applied is the system

$$\begin{cases} u_{xt} = \frac{1}{v^2} F(u - \ln v), \\ v_{xt} = \frac{e^u}{v} H(u - \ln v). \end{cases} \quad (9)$$

The ansatz reducing system (9) is obtained by solving the system of ordinary equations (8), and it has the form

$$\begin{cases} u(t, x) = \ln(x + c_1(t)) + c_2(t), \\ v(t, x) = c_3(t)(x + c_1(t)). \end{cases} \quad (10)$$

System (9) is reduced to a system of ordinary differential equations

$$\begin{cases} -c_3^2(t)c_1'(t) = F(c_2(t) - \ln c_3(t)), \\ c_3(t)c_3'(t) = e^{c_2(t)} H(c_2(t) - \ln c_3(t)) \end{cases}$$

by virtue of the ansatz (10). Thus, we obtained a system of two equations with three unknown functions  $c_1, c_2, c_3$ . Obviously, the solution will then depend on one arbitrary function on variable  $t$ .

We show also that the generalized vector field is admitted by system of ordinary differential equations (8)

$$Q_{ne2} = \left( v u_{xt} + v_x v_t - F(u - \ln v) \right) \partial_u + \left( v v_{xt} - v H(u - \ln v) \right) \partial_v.$$

From this it follows that ansatz (10) reduces non–evolutionary system

$$\begin{cases} u_{xt} = \frac{1}{v} F(u - \ln v) - \frac{u_x u_t}{v}, \\ v_{xt} = H(u - \ln v), \end{cases} \quad (11)$$

to the system of ordinary differential equations. Indeed, substituting (10) into (11) gives

$$\begin{cases} c_3'(t) = F(c_2(t) - \ln c_3(t)), \\ c_3'(t) = H(c_2(t) - \ln c_3(t)). \end{cases} \quad (12)$$

Thus, we obtained an inconsistent system of two ordinary differential equations for three unknown functions  $c_1, c_2, c_3$  in the case when  $F \neq H$ . We see that the generalized symmetry ensures a reduction to a system of ordinary differential equations, but does not guarantee the consistency of the reduced

system. System (10) is consistent if and only if  $F = H$ . In this case, the system (11) takes the form.

$$\begin{cases} u_{xt} = \frac{1}{v}F(u - \ln v) - \frac{u_x v_t}{v}, \\ v_{xt} = F(u - \ln v). \end{cases} \quad (13)$$

Ansatz (10) reduces this system to the single ordinary differential equation

$$c_3'(t) = F(c_2(t) - \ln c_3(t)). \quad (14)$$

Then we conclude that the solution of (13) will depend on two arbitrary functions on variable  $t$ .

Next, the system of partial differential equations is considered [1]

$$\begin{cases} u_t = (F_1(u, v)u_x)_x + G_1(u, v)u_x + H_1(u, v), \\ v_t = (F_2(u, v)v_x)_x + G_2(u, v)v_x + H_2(u, v). \end{cases} \quad (15)$$

The desired vector field is defined in the form of

$$\begin{aligned} Q = & [(F_1(u, v)u_x)_x + G_1(u, v)u_x + H_1(u, v)]\partial_u + \\ & + [(F_2(u, v)v_x)_x + G_2(u, v)v_x + H_2(u, v)]\partial_v. \end{aligned} \quad (16)$$

Let us also consider a system of ordinary differential equations

$$\begin{cases} u_{xx} = \frac{1}{u}u_x^2, \\ v_{xx} = \frac{1}{v}v_x^2. \end{cases} \quad (17)$$

We prove that system (17) admits the generalized vector field (16) if and only if

$$\begin{cases} F_1(u, v) = (\lambda_{11} \ln u + \delta_{11} \ln v + \mu_{11} + \frac{s_{11}}{u})u_x, \\ G_1(u, v) = (\lambda_{21} \ln u + \delta_{21} \ln v + \mu_{21}), \\ H_1(u, v) = u(\delta_{31} \ln v + \lambda_{31} \ln u + \mu_{31}), \\ F_2(u, v) = (\lambda_{12} \ln u + \delta_{12} \ln v + \mu_{12} + \frac{s_{12}}{v})v_x, \\ G_2(u, v) = (\lambda_{22} \ln u + \delta_{22} \ln v + \mu_{22}), \\ H_2(u, v) = v(\delta_{32} \ln v + \lambda_{32} \ln u + \mu_{32}). \end{cases} \quad (18)$$

Then we obtain the following system of evolution equations

$$\begin{cases} u_t = [(\lambda_{11} \ln u + \delta_{11} \ln v + \mu_{11} + \frac{s_{11}}{u})u_x]_x + (\lambda_{21} \ln u + \delta_{21} \ln v + \mu_{21})u_x + \\ + u(\delta_{31} \ln v + \lambda_{31} \ln u + \mu_{31}), \\ v_t = [(\lambda_{12} \ln u + \delta_{12} \ln v + \mu_{12} + \frac{s_{12}}{v})v_x]_x + (\lambda_{22} \ln u + \delta_{22} \ln v + \mu_{22})v_x + \\ + v(\delta_{32} \ln v + \lambda_{32} \ln u + \mu_{32}). \end{cases} \quad (19)$$

Solving system of ordinary differential equations (17) we obtain the corresponding ansatz

$$u(t, x) = c_2(t)e^{c_1(t)x}, \quad v(t, x) = c_4(t)e^{c_3(t)x}. \quad (20)$$

Substituting (20) into (19) we obtain reduced system of ordinary differential equations

$$\begin{cases} c_1' = \lambda_{11}c_1^3 + \delta_{11}c_1^2c_3 + \lambda_{21}c_1^2 + \delta_{21}c_1c_3 + \lambda_{31}c_1 + \delta_{32}c_3, \\ c_2' = c_2(\lambda_{11}c_1^2 \ln c_2 + \delta_{11}c_1^2 \ln c_4 + \lambda_{21}c_1 \ln c_2 + \delta_{21}c_1 \ln c_4 + \delta_{11}c_1c_3 + \lambda_{11}c_1^2 + \\ + \mu_{11}c_1^2 + \lambda_{31} \ln c_2 + \delta_{32} \ln c_4 + \mu_{21}c_1 + \mu_{31}), \\ c_3' = \lambda_{12}c_1c_3^2 + \delta_{12}c_3^3 + \lambda_{22}c_1c_3 + \delta_{22}c_3^2 + \lambda_{32}c_1 + \delta_{32}c_3, \\ c_4' = c_4(\lambda_{12}c_3^2 \ln c_2 + \delta_{12}c_3^2 \ln c_4 + \lambda_{22}c_3 \ln c_2 + \delta_{22}c_3 \ln c_4 + \lambda_{12}c_1c_3 + \delta_{12}c_3^2 + \\ + \mu_{12}c_3^2 + \lambda_{32} \ln c_2 + \delta_{32} \ln c_4 + \mu_{22}c_3 + \mu_{32}). \end{cases}$$

We show the application of the inverse symmetry reuction method for reducing systems of two partial differential equations to the systems of ordinary differential equations. We show that the system of evolutionary equations (5) can be reduced to a system of ordinary equations, parameterized by an arbitrary solution of the KdV equation. We have also demonstrate the possibility of constructing the solution to the system of non-evolutionary equations (9), and (13) which contains one or two arbitrary functions.

## References

- [1] A. F. Barannyk, T. A. Barannyk, and I. I. Yuryk, *A method for the construction of exact solutions to the nonlinear heat equation*  $U_T = (F(U)U_X)_X + G(U)U_X + H(U)$ , Ukrainian Mathematical Journal, Vol. 71, No. 11, April, 2020 (Ukrainian Original Vol. 71, No. 11, November, 2019).
- [2] G. W. Bluman, S. Kumei, *Symmetries and differential equations*, Springer-Verlag, New York, 1981.
- [3] A. S. Fokas, Q. M. Liu, *Nonlinear interaction of traveling waves of non-integrable equations*, Phys. Rev. Lett., **72** (1994), 3293–3296. <https://doi.org/10.1103/PhysRevLett.72.3293>
- [4] N. Kamran, R. Milson, P. J. Olver, *Invariant Moduler and the Reduction of Nonlinear Partial Differential Equations to Dynamical Systems*, Advances in Mathematics, **156**, 286-319 (2000).

- [5] A. Samokhin, *Full Symmetry Algebra for ODEs and Control Systems*, Acta Applicandae Mathematicae 72: 87-99, 2002.
- [6] A. Sergyeyev, *Constructing conditionally integrable evolution systems in (1+1) dimensions: a generalization of invariant modules approach*, J. Phys. A: Math. Gen. **35** (2002) 7653-7660.
- [7] S. R. Svirshchevskii, *Lie-Bäcklund symmetries of linear ODEs and generalized separation of variables in nonlinear equations*, Physics Letters A 199 (1995) 344-348.
- [8] I. Tsyfra, *On the Symmetry Approach to Reduction of Partial Differential Equations*, Proceeding of Institute of Mathematics of NAS of Ukraine, 2004.
- [9] Ivan Tsyfra, Wojciech Rzeszut *On reducing and finding solutions of non-linear evolutionary equations via generalized symmetry of ordinary differential equation*, MBE, 19(7): 6962–6984, DOI: 10.3934/mbe.2022328
- [10] R. Z. Zhdanov, *Conditional Lie-Bäcklund symmetry and reduction of evolution equations*, J. Phys. A: Math. Gen., **28** (1995), 3841–3850. <https://doi.org/10.1088/0305-4470/28/13/027>.

<sup>1</sup>Ivan Tsyfra  
 tsyfra@agh.edu.pl  
 ORCID: <https://orcid.org/0000-0001-6665-3934>

AGH University of Krakow  
 Faculty of Applied Mathematics  
 30 Mickiewicza Avenue, 30-059, Krakow, Poland.

<sup>1</sup>Patryk Sitko  
 psitko@agh.edu.pl  
 ORCID: <https://orcid.org/0000-0002-2510-6528>

AGH University of Krakow  
 Faculty of Applied Mathematics  
 30 Mickiewicza Avenue, 30-059, Krakow, Poland.