arXiv:2503.22043v1 [math.CO] 27 Mar 2025

Shuffle squares and nest-free graphs

Jarosław Grytczuk^{1*}, Bartłomiej Pawlik², and Andrzej Ruciński^{3**}

¹ Faculty of Mathematics and Information Science, Warsaw University of Technology, 00-662 Warsaw, Poland

jaroslaw.grytczuk@pw.edu.pl

² Institute of Mathematics, Silesian University of Technology, 44-100 Gliwice, Poland

bpawlik@polsl.pl

³ Faculty of Mathematics and Computer Science, Adam Mickiewicz University, 61-614 Poznań, Poland rucinski@amu.edu.pl

ucinski@amu.edu.pi

Abstract. A *shuffle square* is a word consisting of two shuffled copies of the same word. For instance, the French word **tuteurer** is a shuffle square, as it can be split into two copies of the word **tuer**. An *ordered graph* is a graph with a fixed linear order of vertices.

We propose a representation of shuffle squares in terms of special nest-free ordered graphs and demonstrate the usefulness of this approach by applying it to several problems. Among others, we prove that binary words of the type $(1001)^n$, n odd, are not shuffle squares and, moreover, they are the only such words among all binary words whose every 1-run has length one or two, while every 0-run has length two. We also provide a counterexample to a believable stipulation that binary words of the form $1^n 0^{n-2} 1^{n-4} \cdots$, n odd, are far from being shuffle squares (the distance measured by the minimum number of letters one has to delete in order to turn a word into a shuffle square).

1 Introduction

1.1 Definitions

Let $k \ge 1$ be an integer and let A be an alphabet of letters, with |A| = k. By a k-ary word of length n we mean any sequence $W = w_1 \cdots w_n$ with $w_i \in A$ for all $1 \le i \le n$. For binary words we typically take the alphabet $\{0, 1\}$ or $\{A, B\}$, while for bigger alphabets we rather use the set $[k] := \{1, 2, \dots, k\}$. The length of a word W will be denoted by |W|.

Any subsequence $W' = w_{i_1} \cdots w_{i_t}$ of a word $W = w_1 \cdots w_n$ is called a *subword* of W and its set of indices, supp $(W') = \{i_1, \ldots, i_t\}$, is called the *support* of W'. If the support of a subword W' consists of consecutive integers, then we call such W' a *block* of W.

A pair of subwords X, Y of W with X = Y and disjoint supports will be called *twins*. We will be using the shorthand $\operatorname{supp}(X, Y) = \operatorname{supp}(X) \cup \operatorname{supp}(Y)$ wherever convenient. The length of twins is defined as the length of either one of them, |X| = |Y|. We will sometime refer also to the *double length* of twins which equals twice the length and measures how much of the word is taken by the twins. The elements of W which do not belong to either of the twins are called *gaps* (with respect to $\{X, Y\}$).

Given two words $U = u_1 \cdots u_n$ and $W = w_1 \cdots w_m$ their concatenation is defined as the word $UW = u_1 \cdots u_n w_1 \cdots w_m$. This definition extends naturally to any finite number of words. In particular, W^r denotes the concatenation of r copies of the word W. For $w \in A$, we may write w^r instead of $ww \cdots w$. For instance, $W = 111001000110 = 1^30^2 10^3 1^2 0$. This notation comes in handy when there are long blocks of identical letters.

A *w*-run in a word W is a maximal block F of W of the form $F = w^r$ for some $r \ge 1$. Here "maximal" means that no other block F' of W which contains F is of the form $F' = w^{r'}$, with r' > r. By a run we mean

^{*} Supported by Narodowe Centrum Nauki, grant 2020/37/B/ST1/03298.

^{**} Supported by Narodowe Centrum Nauki, grant 2024/53/B/ST1/00164.

a *w*-run for any $w \in A$. Every word can be uniquely decomposed into a concatenation of its runs. For instance, in 011011100 there are two 1-runs (11, 111) and three 0-runs (0, 0, 00), and the 5-block decomposition is $0|11|0|111|00 = 01^201^30^2$.

For every non-empty word X we call the word $X^2 = XX$ a square. For example couscous is a square for X = cous, as is 01100110 for X = 0110. In this note, however, we are interested in a (much) broader notion. A shuffle square is any word W which contains twins of double length |W|; such twins will be called *perfect*. For instance, 00001001 is a shuffle square with the twins equal to 0001.

Obviously, a necessary condition for a word W to be a shuffle square is that each letter appears in W an even number of times. We will sometimes refer to such a word as *even*. For example, words **reappear** and **tattletale** are even. (Elsewhere, cf. [4,8], such words are call *tangrams*.)

Note that if there are twins of length t in a word W, then there are also monotone twins X, Y of length t in W, that is, $i_h < j_h$ for all $1 \le h \le t$, where $\operatorname{supp}(X) = \{i_1, \ldots, i_t\}$ and $\operatorname{supp}(Y) = \{j_1, \ldots, j_t\}$. Moreover, there are also canonical twins which, in addition to being monotone, partition each run of W into three consecutive blocks (some might be empty) with the elements of X to the left, followed by the elements of Y, followed by gaps. It is crucial for us that among the longest twins in a word there are always canonical ones. For a formal explanation see Appendix.

Example 2. In Example 1 the twins are not monotone, but after swapping four pairs of elements between X and Y (all but the first one), the pair of twins indicated here: $W = \underline{1} \ \overline{1} \ \underline{0} \ \underline{0} \ \overline{1} \ \underline{0} \ \overline{0} \ \overline{0} \ 1 \ 1 \ \overline{0}$, is monotone, though not yet canonical due to the (dis)order in the first 1-run. A single "rewiring" operation makes it canonical: $W = \underline{1} \ \overline{1} \ \underline{0} \ \underline{0} \ \overline{1} \ \underline{0} \ \overline{0} \ \overline{0} \ 1 \ 1 \ \overline{0}$ (see Fig. 1).



Fig. 1. Non-canonical and canonical (red dashed) twins in the word W = 111001000110.

1.2 Results and method

In this paper we establish conditions under which binary words belonging to a specified class are shuffle squares. Among others, in Section 3 we consider words with four runs and words with all 1-runs of length 1 (see Propositions 2 and 3). In Section 4, we give our main result. It states (see Proposition 4 and Theorem 1) that among the binary words with all 0-runs of length two and all 1-runs of length at most two, only the words $(1001)^n$ for n odd, are *not* shuffle squares. That any odd power of the word 1001 is not a shuffle square was conjectured by Komisarski [7], who used to call it the *odd ABBA problem*. Let us also mention that an alleged proof was presented by Fortuna [5], though, up to our knowledge, it has not yet been written down. Actually, Theorem 1 is much more general as it covers not only "odd abbas", but many other binary words with an odd number of 0-runs of the same length.

In the last section, we tackle the question of how far a binary word may be from a shuffle square. Given a word W of length n, let f(W) be the length of the longest twins in W and set g(W) = n - 2f(W). Further, let $g(n) = \max g(W)$ over all W, with |W| = n. By a sophisticated approach via a regularity lemma for words, Axenovich, Person, and Puzynina [1] showed that g(n) = o(n) and, by providing a construction, that $g(n) = \Omega(\log n)$. The lower bound was improved by Bukh [3] to $g(n) = \Omega(n^{1/3})$ by refining the construction from [1] (see [2] for a proof). In a personal communication Bukh has expressed his belief that, in fact, $g(n) = \Omega(\sqrt{n})$ and that the real challenge would be to improve the upper bound, where one would need to refine the regularity approach from [1]. Focusing on the lower bound, in [2] we shared that belief and implicitly stated the following conjecture.

Conjecture 1. There is a constant c such that for every n there is a binary word of length n with largest twins leaving out at least $c\sqrt{n}$ elements. In short, $g(n) = \Omega(\sqrt{n})$.

In view of the constructions from [1] and [3], we had believed for long that a natural candidate to facilitate Conjecture 1 is the binary word $O_{m,r} = 1^m 0^{m-2} 1^{m-4} \cdots$ consisting of r runs whose lengths form a decreasing sequence of *consecutive* odd numbers, for some odd m and $r \leq (m+1)/2$. Here we rebuke this belief by finding twins in $O_{m,r}$ of the double length at least $|O_{m,r}| - 23$, for all m and r. This result does not mean that Conjecture 1 is false. In search for better constructions, one should perhaps allow varying differences between consecutive, still odd, terms.

But at least as important as the obtained results, we believe, is the method we propose to prove them with. To each pair of canonical twins we uniquely assign an *ordered graph* (see the formal definition in he next section). Thus, instead of looking for twins in a word or deciding if it is a shuffle square, we construct a corresponding graph or show that such a graph does not exist. We believe that this approach is more transparent, insightful, and ultimately has some potential in this area of research.

2 Graphic representations

Here we introduce our main tool – a representation of shuffle squares and, more generally, canonical twins in terms of ordered graphs with no nests. An *ordered graph* is just a graph with its vertex set linearly ordered. We allow loops and parallel edges. A *nest* in an ordered graph consists of a pair of disjoint edges (including loops) e and f with min $e < \min f$ and max $e > \max f$. An ordered graph is *nest-free* if no pair of its edges forms a nets.

Let a k-ary word $W = w_1 \cdots w_n$ have canonical twins $X = w_{i_1} \cdots w_{i_t}$ and $Y = w_{j_1} \cdots w_{j_t}$. In particular, $\{i_1, \ldots, i_t\} \cap \{j_1, \ldots, j_t\} = \emptyset$, and for all $h = 1, \ldots, t$, we have $i_h < j_h$ and $w_{i_h} = w_{j_h}$.

As an intermediate step, define an ordered graph $M = M_{X,Y}$ on the set of vertices $\{1, \ldots, n\}$ with edges $\{i_1, j_1\}, \ldots, \{i_t, j_t\}$. So, M consists of a *matching* of size t and n - 2t isolated vertices. Clearly, there are no nests in M.

Let $W = U_1 \cdots U_m$, where each block U_h is a run in W. Now, contract each set $\sup(U_h)$ into a single vertex u_h , keeping all the edges of M in place and keeping the order $u_1 < \cdots < u_m$. We denote the obtained ordered graph by $G = G_{X,Y}$. Observe that G is still nest-free and, moreover, consists of k separated graphs on vertex sets $N_w = \{h : U_h \text{ is a } w\text{-run}\}$, for each letter $w \in [k]$. The degrees of vertices in G are $\deg_G(u_h) = |\operatorname{supp}(U_h) \cap \operatorname{supp}(X,Y)|, h = 1, \ldots, m$.

From G it is straightforward to uniquely reconstruct (X, Y). The uniqueness is a consequence of the fact that we restrict ourselves to canonical twins. We simply scan the vertices u of G from left to right and record their right degrees $\deg_{G}^{(right)}(u)$ defined as the number of edges going from u to the right and left degrees $\deg_{G}^{(left)}(u)$ defined analogously. Then from each run U_h , assign the first $\deg_{G}^{(right)}(u)$ elements to X and the next $\deg_{G}^{(left)}(u)$ to Y, leaving the remaining vertices of U_h , if there are any, unassigned (gaps). But if we are only interested in the length of the twins, then just by looking at G we learn that there is a pair of twins in W of double length equal to $\sum \deg_{G}(u_h)$, where loops count as two.

This allows us to apply this "graphic" approach also in situations when we do not know any twins in Wand our goal is to find them as large as possible. Then we simply attempt to construct a nest-free ordered graph G on vertices u_h , $h = 1, \ldots, m$, with no edges joining vertices from different sets N_w , with degrees as close as possible to the lengths of the corresponding runs U_h . In particular, if we succeed to create such a graph G with all $\deg_G(u_h) = |U_h|$, then we know that W is a shuffle square and we can produce a pair of perfect twins witnessing this property.

Let us summarize the above discussion in the following statement characterizing shuffle squares in terms of their ordered graph representations.

Proposition 1. Let W be a k-ary word and let $W = U_1 \cdots U_m$ be a decomposition of W into runs U_i . Then W is a shuffle square if and only if there exists an ordered graph G on the set $\{u_1, \ldots, u_m\}$ satisfying the following conditions.

- 1. If u_i is joined to u_j by an edge, then the corresponding runs U_i and U_j are w-runs for the same letter w.
- 2. The degree of each vertex u_i satisfies $\deg_G(u_i) = |U_i|$.
- 3. Graph G is nest-free.

Let us look at a couple of examples.

Example 3. Let $W = \underline{1} \underline{1} \overline{1} \underline{2} \underline{2} \underline{3} 3 3 \overline{1} \overline{2} \overline{2} \overline{3}$ with canonical twins as indicated. The corresponding matching M and graph G are given in Fig. 2. Note that there is no way to change G so that $\deg_G(u_3) = 3$ while keeping all other degrees intact. Indeed, u_6 has to be adjacent to u_3 , so there cannot be a loop at u_5 which, in turn, implies that there is no loop at u_3 either. If we swapped the 3-runs around and instead consider the word W' = 111223122333, then the graph in Fig. 3 shows that W' is a shuffle square: $W' = \underline{1} \underline{1} \overline{1} \underline{2} \underline{2} \underline{3} \overline{1} \overline{2} \overline{2} \underline{3} \overline{3} \overline{3}$.



Fig. 2. Matching *M* and graph *G* for $W = \underline{1} \underline{1} \overline{1} \underline{2} \underline{2} \underline{3} 3 \overline{3} \overline{1} \overline{2} \overline{2} \overline{3}$.



Fig. 3. The word W' = 1 1 1 2 2 3 1 2 2 3 3 3 is a shuffle square.

From now on we will be exclusively focusing on binary words.

Example 4. Let $W = \underline{1} \ 1 \ \underline{0} \ \underline{0} \ \overline{1} \ \overline{0} \ \overline{0} \ \underline{0} \ \underline{1} \ \overline{1} \ 0$ and (X, Y) be monotone twins in W of length 4 given by $\sup (X) = \{1, 4, 5, 10\}$ and $\sup (Y) = \{6, 7, 8, 11\}$ (so, both equal 1001) as indicated. The corresponding matching M and graph G are given in Fig. 4. By adding a single loop at vertex u_1 , we immediately obtain a pair of twins of length 5 in W, X' = Y' = 11001, a bit different from those in Example 2.

As W is even, we might wonder if it is a shuffle square, that is, if it possesses a pair of twins of length 6. This can be quickly answered in negative, using the graph representation as a tool. If W were a shuffle square, there would exist an ordered nest-free graph G on vertex set $\{u_1, \ldots, u_6\}$ with no edges between $\{u_1, u_3, u_5\}$ and $\{u_2, u_4, u_6\}$ and degrees deg_G $(u_h) = |U_h|, h = 1, \ldots, 6$. However, in Fig. 4 we see that such a graph does not exist. Indeed, vertex u_1 cannot be connected to u_5 as then all edges at the bottom part would need to be adjacent to u_6 which is impossible. So, there must be a loop at u_5 which cuts off vertex u_6 at the bottom.



Fig. 4. Matching M and graph G for $W = \underline{1} \ 1 \ 1 \ \underline{0} \ \underline{0} \ \overline{1} \ \overline{0} \ 0 \ \underline{1} \ \overline{1} \ 0.$

On a positive side, consider the next example.

Example 5. Is $W = 110^5 10^6 1110^4 110^7$ a shuffle square? One may try a tedious trial-and-error approach, but instead we draw a nest-free graph G, see Fig. 5, with degrees equal to the sizes of the blocks in W. This yields the answer "yes" to the above question, and we may still construct a pair of perfect twins in W based on G. It is $W = 1 \overline{1} \ 0^5 \ 1 \ 0 \ \overline{0^5} \ 1^2 \ \overline{1} \ 0^2 \ \overline{0^5}$.



Fig. 5. The word $W = 1 \ 1 \ 0^5 \ 1 \ 0 \ 0^5 \ 1^2 \ 1 \ 0^3 \ 0 \ 1^2 \ 0^2 \ 0^5$ is a shuffle square.

3 Two easy pieces

In this section we prove two relatively simple results on shuffle squares which demonstrate the elegance and simplicity of the graphic representation. An even word is *dull* if all its runs have even length. It is called dull, because it is trivially a shuffle square with the perfect twins just alternating.

Note that no even binary word with the total of *three* runs is a shuffle square unless it is dull. Indeed, the only way to construct a graph G with vertices $u_1 < u_2 < u_3$ of degrees $|U_1|, |U_2|, |U_3|$, respectively, and with no edges between $\{u_1, u_3\}$ and u_2 , is to put $|U_i|/2$ loops at each vertex u_i , i = 1, 2, 3, as otherwise nests would be created.

We now settle the case of *four* runs. The result appeared already in [6].

Proposition 2. Let a, b, c, d be positive integers, not all even, but with a + c and b + d both even. Then the (even but not dull) word $W = 1^a 0^b 1^c 0^d$ is a shuffle square if and only if $a \ge c$ and $b \le d$.

Proof. Consider a nest-free ordered graph on vertex set $u_1 < u_2 < u_3 < u_4$ with capacities a, b, c, d, respectively, and edges permissible only between pairs u_1, u_3 and u_2, u_4 (in addition to loops). If there is a shuffle square in W, then both edges, u_1, u_3 and u_2, u_4 , must be present (with some multiplicities), as loops can only



Fig. 6. Graph G for the shuffle square $1^a 0^b 1^c 0^d$.

accommodate even degrees. But then there cannot be any loops either at u_2 or at u_3 , implying that $a \ge c$ and $b \le d$ (see Fig. 6).

On the other hand, if $a \ge c$ and $b \le d$, then we draw edge u_1u_3 with multiplicity c, edge u_2u_4 with multiplicity b and add (a - c)/2 loops at u_1 and (d - b)/2 loops at u_4 (see Fig. 6). The obtained ordered graph is nest-free and thus corresponds to a pair of perfect twins in W of the form

$$1^{(a-c)/2} \overline{1^{(a-c)/2}} 1^c 0^b \overline{1^c} \overline{0^b} 0^{(d-b)/2} \overline{0^{(d-b)/2}}.$$

Note that both twins form the word $1^{(a+c)/2}0^{(b+d)/2}$.

As a corollary of Proposition 2 we infer that every word $w = 1^a 0^b 1^c 0^d$ with all a, b, c, d positive, can be rotated (cyclically permuted) so that the outcome is a shuffle square. Indeed, the orderings *abcd*, *bcda*, *cdab*, *dabc* represent all four possible pairwise relations between a, c and between b, d, so at least one of them will satisfy the conditions in Proposition 2 (with a, b, c, d renamed if needed).

Finally in this section, we consider even binary words with all 1's separated from each other, that is with all 1-runs of length 1, but with no restrictions on the 0's. That is, we are looking at words of the form

$$W = 1 \ 0^{a_1} 1 \ 0^{a_2} \cdots 1 \ 0^{a_{2m}},\tag{1}$$

for positive integers m, a_1, \ldots, a_{2m} . Our goal is to define conditions for the values of a_i that make W a shuffle square with the 1's belonging alternately to perfect twins.

This could be solved without referring to the ordered graphs, but we prefer the latter approach as it seems to be more insightful. Any graph representing perfect twins in word W must attain a structure presented in Fig. 7. with the additional constraint that for each $i = 1, 3, \ldots, 2m - 3$, either y_i or z_{i+1} must be 0. The variables x_i, y_i, z_i represent multiplicities of the respective edges, so we have altogether 4m - 2 unknowns to find. Let us denote the top vertices by u_1, \ldots, u_{2m} and the bottom vertices by v_1, \ldots, v_{2m} .



 $r_i = \deg^{(right)}(v_i), \quad \ell_i = \deg^{(left)}(v_i), \quad r_i + r_{i+1} = \ell_{i+1} + \ell_{i+2}$

Fig. 7. The structure of the graph representing perfect twins in the shuffle square $1 \ 0^{a_1} 1 \ 0^{a_2} \cdots 1 \ 0^{a_{2m}}$.

One way to proceed from this would be to set up a system of 2m equations guaranteeing that for each i = 1, ..., 2m, the degree of vertex v_i must be equal to a_i , supplemented by m - 1 equations of the form

 $y_i z_{i+1} = 0, i = 1, 3, ..., 2m - 3$. However, it seems simpler to apply more coarse setting in terms of the right degrees $r_i = \deg_G^{(right)}(v_i)$ and left degrees $\ell_i = \deg_G^{(left)}(v_i)$ and utilize the observation that $r_1 = a_1$ and, for each $i = 1, 3, ..., 2m - 1, r_i + r_{i+1} = \ell_{i+1} + \ell_{i+2}$. Thus, after getting rid of the ℓ_i 's (as $r_i + \ell_i = a_i$ for all i), we have just 2m - 1 variables and m equations. Moreover, we avoid the annoying system of m - 1 alternatives: $y_i = 0$ or $z_{i+1} = 0$. But the biggest benefit we earn from this substitution is that the m equations each involve a separate set of variables.

So, it remains to solve this system of m equations. We arbitrarily choose the r_i 's with odd i as free variables which determine the remaining ones. Then, the system becomes

$$\begin{cases} r_2 = \frac{1}{2}(-a_1 + a_2 + a_3 - r_3) \\ r_{2i+2} = \frac{1}{2}(a_{2i+2} + a_{2i+3} - r_{2i+1} - r_{2i+3}), 1 \le i \le m - 2 \\ r_{2m} = \frac{1}{2}(a_{2m} - r_{2m-1}) \end{cases}$$
(2)

As for all i we have the natural constraints $0 \leq r_i \leq a_i$, we need to show that the system of inequalities

$$\begin{cases} (a_3 - a_1) - u_2 \leqslant r_3 \leqslant (a_3 - a_1) + a_2\\ a_{2i+3} - a_{2i+2} \leqslant r_{2i+1} + r_{2i+3} \leqslant a_{2i+3} + a_{2i+2}, \ 1 \leqslant i \leqslant m - 2\\ r_{2m-1} \leqslant a_{2m} \end{cases}$$
(3)

has integer-valued solutions r_3, \ldots, r_{2m-1} such that for all $i = 3, \ldots, 2m-1$ we have $0 \leq r_i \leq a_i$ and the quantities in the parenthesis in (2) are all even (so that the values of r_2, r_4, \ldots, r_{2m} are integers). If for a given sequence of a_i 's there exist r_i 's satisfying the above conditions, then W has perfect twins of the form

$$1 0^{a_1+r_2} 1 0^{r_3+r_4} \dots 1 0^{r_{2m-1}+r_{2m}} = 1 0^{\ell_2+\ell_3} 1 0^{\ell_4+\ell_5} \dots 1 0^{\ell_{2m}}.$$

Thus we proved the following result.

Proposition 3. A binary word given by (1) is a shuffle square with alternating 1's if and only if the combined systems of inequalities (3) and equations (2) have integer solutions with $0 \le r_i \le a_i$ for all i.

In particular, for m = 1, the sole condition for $W = 1 \ 0^{a_1} \ 1 \ 0^{a_2}$ to be a shuffle square is, trivially, that $a_1 \leq a_2$. For m = 2, the above system of inequalities is reduced to just

$$\begin{cases} (a_3 - a_1) - a_2 \leqslant r_3 \leqslant (a_3 - a_1) + a_2 \\ r_3 \leqslant a_4 \end{cases}$$

So, whether W is a shuffle square is determined by the validity of the single inequality $a_3 + a_2 - a_1 \ge 0$. Then any choice of

$$r_3 \in \{\max(0, a_3 - a_2 - a_1), \min(a_3 + a_2 - a_1, a_3, a_4)\}\$$

which has the same parity as $a_3 + a_2 - a_1$ (and so, the same parity as a_4) yields a feasible solution (r_2, r_3, r_4) , where

$$\begin{cases} r_2 = \frac{1}{2}(-a_1 + a_2 + a_3 - r_3) \\ r_4 = \frac{1}{2}(a_4 - r_3) \end{cases}$$

Interestingly, if a word $W = 1 \ 0^{a_1} 1 \ 0^{a_2} 1 \ 0^{a_3} 1 \ 0^{a_3}$ does not the satisfy the condition $a_3 + a_2 - a_1 \ge 0$, then there is always a rotation so that the new word still begins with a 1, but now is a shuffle square with the 1's alternating. Indeed, it suffices to redefine a_1 as the minimum of all four parameters and start the word there. For instance, let $W = 1 \ 0^{16} 1 \ 0^9 1 \ 0^5 1 \ 0^4$. Here $a_3 + a_2 - a_1 = 5 + 9 - 16 < 0$, so this word is not a shuffle square. But beginning at, say, $1 \ 0^4$ we get a new word $W' = 1 \ 0^4 1 \ 0^{16} 1 \ 0^9 1 \ 0^5$, a rotation of W, for which the condition $a_3 + a_2 - a_1 \ge 0$ holds. Thus, W' is a desired shuffle square.

This is no longer true for $m \ge 3$. For instance, $W = 1 \ 0 \ 1 \ 0 \ 1 \ 0^5 \ 1 \ 0 \ 1 \ 0^5$ does not have a shuffle square consisting of twins which get the 1's alternately, no matter at which 1 we begin. (On the other hand, W is a perfect square.)

4 The odd ABBA problem

Recall that if every run of a binary word W has an even length, in particular, length two, then, trivially, W is a shuffle square. It is perhaps a bit surprising that a modest relaxation allowing runs of length one, but only for one of the two letters, say letter 1, becomes quite non-trivial.

One can quickly come up with such words which are not shuffle squares, like 1001 or 100110011001. Our main result in this section states that words of that form, more generally, of the form $(1001)^n$, for n odd, are indeed *not* shuffle squares.

But first, as a sort of complementary result, we show that all other even binary words with all 0-runs of length exactly two and all 1-runs of length at most two, are shuffle squares. So, in fact, the odd repeats of the word 1001 are the sole exception in this class.

Proposition 4. Every even binary word in which every 0-run has length two, while every 1-run has length at most two is a shuffle square, unless it is of the form $(1001)^n$ for some odd n.

Proof. Let W be an even binary word with every 0-run of length two and every 1-run of length at most 2. Assume that $W \neq (1001)^n$ for any odd n.

We are going to construct a corresponding nest-free graph G on vertices u_1, \ldots, u_m , where m is the total number of runs in W. As always G consists of two separated subgraphs, G_1 and G_0 whose vertices interlace and correspond to the 1-runs and 0-runs of W, respectively. Within G_1 we distinguish the vertices with capacity 1, call them *singles*, and with capacity 2, call them *doubles*. As G_1 (the top graph in all figures) we take a union of paths connecting consecutive singles and going through all the doubles along the way, plus loops at all other doubles.

Through the rest of the proof we are going to convince the reader that such a G_1 can be always complemented by a suitable 2-regular graph G_0 , so that the resulting graph $G = G_1 \cup G_2$ is nest-free.

We begin with two observations that can be verified by just looking at the corresponding illustrations.

Observation 1. If $P = (u_h, u_{h+2}, \ldots, u_{h+2s})$ is a path of an odd length s in G_1 and $h \ge 2$, then there is a nest-free graph on $\{u_{h-1}, u_h, u_{h+1}, \ldots, u_{h+2s}\}$ which, in addition to P, consists of a 2-regular subgraph on $\{u_{h-1}, u_{h+1}, \ldots, u_{h+2s-1}\}$. An analogous statement is also true in the symmetric case when h + 2s < m with u_{h-1} replaced by u_{h+2s+1} .

For the proof see Fig. 8.



Fig. 8. An illustration to the proof of Observation 1.

Observation 2. If $P = (u_h, u_{h+2}, \ldots, u_{h+2s})$ and $Q = (u_{h+2s+2}, \ldots, u_{h+2s+2t})$ are two consecutive odd paths P and Q in G_1 , then there is a nest-free graph on $\{u_h, u_{h+1}, \ldots, u_{h+2s+2t}\}$ which, in addition to P and Q, consists of a 2-regular subgraph on $\{u_{h+1}, u_{h+3}, \ldots, u_{h+2s-1}, u_{h+2s+1}, \ldots, u_{h+2s+2t-1}\}$.

For the proof see Fig. 9.

Now we may finish quickly the proof of Proposition 4 by induction on N—the number of paths in G_1 . The case N = 0, that is when W is a dull word, is trivial. For N = 1, if the sole path P in G_1 has even length, then we draw double edges between consecutive pairs of G_0 of vertices lying "under P" and loops elsewhere (see Fig. 10). Otherwise, as W is not of the form $(1001)^n$, n odd, the conclusion follows by Observation 1.

Assume now that $N \ge 2$ and the statement is true for fewer than N paths in G_1 . Let P and Q be the first two (from left) paths in G_1 . Let p be the leftmost vertex of P and let q be the rightmost vertex of Q. If there



Fig. 10. An illustration to the proof of Proposition 4, case m = 1.

are double vertices between P and Q, then we may apply Observations 1 separately to each of the paths if needed, that is, if it has odd length (see Fig. 11). Otherwise, we apply Observation 2. In either case, one can build a nest-free subgraph G[P,Q] on the vertex set $\{p, p+1, \ldots, q-1, q\}$ with all vertices of degrees equal to their capacities. After removing all the vertices of G[P,Q], if there is still a nonempty remaining part, we apply the inductive assumption.



Fig. 11. An illustration to the proof of Proposition 4, case $m \ge 2$.

9

The proof of our main result relies on the following lemma.

Lemma 1. Let H be an ordered graph on the set of vertices $\{1, 2, ..., n\}$ consisting of two vertex disjoint subgraphs: a path P starting at vertex 1 and ending at vertex n, and a cycle C. If H is nest-free, then C must be an even cycle.

Proof. Let us assume that the graph G satisfying the assumptions of the lemma is nest-free. Let E(P) denote the set of the edges of the path P. Consider an order relation \prec on E(P) defined by $e \prec f$ if min $e \leq \min f$ and max $e \leq \max f$, for each pair of distinct edges $e, f \in E(P)$. Since P is nest-free, this relation is a linear order and we may enumerate the edges of P accordingly to that order as $E(P) = \{e_1 \prec e_2 \prec \cdots \prec e_p\}$. Next, we assign signs $\{-1, +1\}$ to these edges in an alternating manner:

$$s(e_1) = -1, s(e_2) = +1, s(e_3) = -1, \dots, s(e_p) = (-1)^p.$$

In particular, $S(P) = s(e_1)s(e_2)\cdots s(e_p) = (-1)(+1)(-1)\cdots (-1)^p$ is an alternating string of signs.

Let $V(C) = \{x_1, \ldots, x_q\}$ be the vertex set of the cycle C, where the vertices are numbered in the order they are traversed by C. We say that a vertex x of C is *hugged* by an edge e of P if $\min e < x < \max e$. For each $i = 1, \ldots, q$, let $X_i = \{e \in P : x_i \text{ is hugged by } e\}$.

A crucial observation stemming from the nest-freeness of G is that each X_i is an *interval* consisting of consecutive edges of P in the linear order \prec . Moreover, each of these intervals has odd cardinality, $X_i \cap X_{i+1} =$

 \emptyset and the edges in X_{i+1} either directly follow or directly precede the edges of X_i . Thus, denoting $r_i = |X_i|$, we may write $X_i = \{e_{j_i} \prec e_{j_i+1} \prec \cdots \prec e_{j_i+r_i-1}\}$ and

$$S(X_i) = s(e_{j_i})s(e_{j_i+1})\cdots s(e_{j_i+r_i-1}).$$

Notice that each $S(X_i)$ is a *block* of *odd* length in the string S(P) and each pair of consecutive strings, $S(X_i)$ and $S(X_{i+1})$ (in the cyclic order of C), must be *adjacent* in the string S(P). Therefore $S(X_1) \cdots S(X_q)$ is also an alternating string of signs whose first and last terms must be different, since the blocks $S(X_1)$ and $S(X_q)$ are adjacent in S(P) as well. This forces q to be even.

Using this lemma we may easily get our main result.

Theorem 1. Let $Let r \ge 2$ and m be arbitrary positive integers and let a_1, \ldots, a_{2m} be arbitrary positive even integers. Then the word $W = 10^r 1^{a_1} 0^r 1^{a_2} \cdots 1^{a_{2m}} 0^r 1$, is not a shuffle square. In particular, for every odd $n \ge 1$, the word $(1001)^n$ is not a shuffle square.

Proof. Suppose on the contrary that W is a shuffle square and let G be an ordered nest-free graph representing W as in Proposition 1. Next let G_0 and G_1 be the subgraphs of G corresponding to 0-runs and 1-runs respectively. Clearly the degree sequences of these graphs are (r, r, \ldots, r) and $(1, a_1, a_2, \ldots, a_{2m}, 1)$, respectively. Since, G_0 is a regular graph of degree at least 2 with an odd number of vertices, it must contain and odd cycle C (as it cannot be *bipartite*). The second graph G_1 must have a path P whose ends are the only vertices of G_1 with odd degrees, namely the first and the last vertex. It follows that $H = C \cup P \subseteq G$ is a nest-free ordered graph, which contradicts Lemma 1.

In particular, with all $a_i = 2$, Theorem 1 covers words of the form $(10^r 1)^{2m+1}$ which with r = 2 becomes the family of "odd abbas".

5 Odds at the end

We first describe a general class of words whose two particular instances yielded lower bounds on g(n) in [1] and [3]. For positive integers n_1, \ldots, n_r , let $W(n_1, \ldots, n_r) = U_1 \cdots U_r$ be the binary word whose consecutive runs U_i have lengths $|U_i| = n_i$, $i = 1, \ldots, r$. We adopt the convention that U_1 is a 1-run. The main features of the words used in [1] and [3] were that the runs rapidly decrease in length and that their lengths are odd.

An easy generalization of these two special words led us in [2] to an observation that if $n_1 > n_2 > \cdots > n_r$ are all odd, then $g(W(n_1, \ldots, n_r)) \ge \min\{r, \delta\}$, where δ is the smallest difference between two consecutive terms (see [2, Lemma 3.1]). The proof idea is that if twins take equally from each run, then they "lose" at least one element per run. Otherwise, the leading twin is ahead at some point and the follower has to "jump" over the second next run, a loss that can never be made up, due to the run lengths getting smaller and smaller.

The particular words used in [1] and [3] were $A_r := W(3^{r-1}, 3^{r-2}, \ldots, 3^2, 3, 1)$ and $B_r := W(r^2 + 1, (r - 1)r + 1, \ldots, 2r + 1, r + 1)$, respectively. Simple calculations reveal that $r = \Theta(\log |A_r|)$ and $r = \Theta(|B_r|^{1/3})$, leading to the desired bounds. (For *n* not expressible as $|A_r|$ or $|B_r|$ for any *r* an obvious extrapolation has been applied.)

In view of the constructions from [1] and [3] a natural candidate to facilitate Conjecture 1 seemed to be, for an odd integer m and $r \leq (m+1)/2$, a binary word $O_{m,r} := W(m, m-2, m-4, \ldots, m-2r+2)$ consisting of r consecutive odd numbers. Indeed, $n := |O_{m,r}| = r(m-r+1) = \Theta(m^2)$ for $r = \Theta(m)$ and thus, showing that $g(O_{m,r}) = \Omega(m)$ would do the job. Below we show that, although indeed $O_{m,r}$ is not a shuffle square, we have $g(O_{m,r}) \leq 23$ independently of m and r, so it is really close to a shuffle square even for large m.

We begin with an easy and more general fact which might be of independent interest.

Claim 1. Let $W = W(n_1, n_2, ..., n_r)$. If n_1 is odd and $n_2 > \cdots > n_r$ form a decreasing sequence (but are not necessarily odd), then W is not a shuffle square.

Proof. We will show that a graph corresponding to canonical perfect twins does not exists. Such a graph if exists has vertices u_1, \ldots, u_r with degrees $\deg_G(u_i) = n_i$ and is nest-free. Since n_1 is odd, there must be an edge from u_1 to the right (and not just loops). This implies that there are no loops at u_2 and, thus, $\deg_G^{(left)}(u_2) = 0$ (see Fig. 12 - left side). Let u_h be the rightmost vertex with $\deg_G^{(left)}(u_h) = 0$. To have



Fig. 12. An illustration to the proof of Claim 1.

 $\deg_G(u_h) = n_h$, vertex u_h must send edges to at least two vertices to the right, say, u_i and u_j , h < i < j, all indices with the same parity. But then, owing to the nestlessness of G, we have $\deg_G^{(left)}(u_{i+1}) = 0$, a contradiction with the definition of u_h (see Fig. 12 - right side).

This, in particular, implies that the word $O_{m,r}$ defined above is not a shuffle square. Our final result shows that $O_{m,r}$, quite surprisingly, is very close to being a shuffle square.

Proposition 5. For every $m \ge 2r-1$, m odd, the word $O_{m,r}$ contains twins of total length at least $|O_{m,r}|-23$, that is, $g(O_{m,r}) \le 23$.

Proof. Our aim is to find canonical twins in $O_{m,r}$ of double length at least $|O_{m,r}| - 23$. To this end, we are going to construct a nest-free graph on r vertices whose *deficit* defined as $|O_{m,r}| - \sum_{u \in V(G)} \deg_G(v)$ is at most 23. We may and do assume that $r \ge 24$, as otherwise we could draw as many as possible loops at every vertex, getting the deficit of $r \le 23$. For sheer convenience, we also assume that r is divisible by 8. If not, we add loops at up to the first seven vertices on, say, the left end of $O_{m,r}$, procuring a deficit of at most 7, and build G on the remaining vertices. So, let r = 8k for some integer $k \ge 3$.

As usual, there are two kinds of vertices, say, red and blue, alternating, and there are no red-blue edges whatsoever. Let us identify the vertices with their capacities, so we have the red set $V_1 = \{m, m-4, m-8, \ldots, m-2r+4\}$ and the blue set $V_2 = \{m-2, m-6, \ldots, m-2r+2\}$.

We begin by describing the "red" subgraph $G[V_1]$. We split V_1 into k consecutive 4-tuples $U_i = \{i, i - 4, i - 8, i - 12\}$, $i = m, m - 16, \ldots, m - 2r + 16$, with edge sets $E_i = \{\{i, i - 4\}, \{i, i - 8\}, \{i - 4, i - 12\}\}$ and the edges bearing multiplicities $\mu(i, i - 4) = 8$, $\mu(i, i - 8) = i - 8$, and $\mu(i - 4, i - 12) = i - 12$ (see Fig. 13 - the top part). Note that this way the deficit of the red subgraph $G[V_1]$ is zero. Indeed, for each i, $\deg_G(i) = 8 + (i - 8) = i$, $\deg(i - 4) = 8 + (i - 12) = i - 4$, d(i - 8) = i - 8, and d(i - 12) = i - 12.



Fig. 13. Construction from the proof of Proposition 5.

The "blue" subgraph is a bit more complicated, as we have to avoid traps set up by the red graph. Nevertheless, its structure is almost the same: it consists of disjoint paths of lengths three, except that now these paths are slightly more stretched out, and two isolated edges, one at the beginning and one at the end. It is these two edges what raises the deficit by 16. Precisely, we have $\mu(m-2, m-10) = m-10$ and $\mu(m-2r+10, m-2r+1) = m-2r+1$, while the remaining vertices are broken up into k-1 4-tuples $W_i = \{i, i-8, i-12, i-20\}, i = m-6, m-22, \ldots, m-2r+26$, with edges sets $E_i = \{\{i, i-8\}, \{i, i-12\}, \{i-12, i-20\}\}$ and the edges of multiplicities $\mu(i, i-8) = i-8, \mu(i, i-12) = 8$, and $\mu(i-12, i-18) = i-18$ (see Fig. 13 – the bottom part). The subgraphs $G[W_i]$ contribute no deficit, so that the total deficit of G is at most 7+8+8=23.

To illustrate the above construction we present in Fig. 14 the graph G for the word

$$O_{47,24} = 1^{47} 0^{45} 1^{43} 0^{41} 1^{39} 0^{37} 1^{35} 0^{33} 1^{31} 0^{29} 1^{27} 0^{25} 1^{23} 0^{21} 1^{19} 0^{17} 1^{15} 0^{13} 1^{11} 0^{9} 1^{7} 0^{5} 1^{3} 0^{1}.$$



Fig. 14. The graphic representation of the word $O_{47,24}$.

The corresponding pair of canonical twins of double length $24^2 - 16$ looks like this:

 $1^{47}0^{37}0^{8}1^{35}\overline{1^{8}}0^{41}\overline{1^{39}0^{37}1^{35}0^{33}}1^{31}0^{21}\overline{0^{8}}1^{19}1^{8}0^{25}\overline{1^{23}0^{21}1^{19}0^{17}}1^{15}0^{5}0^{8}1^{3}\overline{1^{8}}0^{1}0^{8}\overline{1^{7}0^{5}1^{3}0^{1}},$

with each single twin equal to $1^{47}0^{37}1^{35}0^{41}1^{31}0^{21}1^{19}0^{25}1^{15}0^{5}1^{3}0^{1}$. As you may see, the only gaps, each of size 8, appear in the second run and the fifth from the end. It would not be so easy to get these twins if not for the use of the graphs.

Acknowledgment

We would like to thank Andrzej Komisarski for sharing with us his lovely ABBA problem, as well as some results of his impressive computer experiments.

References

- Axenovich, M., Person, Y., Puzynina, S.: A regularity lemma and twins in words. J. Combin. Theory Ser. A 120 (2012) 733–743. 1.2, 1.2, 5
- Basu, A., Ruciński, A.: How far are ternary words from shuffle squares? Ars Math. Contemp. (2024), https://doi.org/10.26493/1855-3974.3211.86d. 1.2, 5
- 3. Bukh, B.: unpublished note (2013) (1 page). 1.2, 1.2, 5
- Dębski, M., Grytczuk, J., Pawlik, B., Przybyło, J., Śleszyńska-Nowak, M.: Words avoiding tangrams. Ann. Comb. (2024). 1.1
- 5. Fortuna, W.: presentation at The VIII Students Conference "Zmirłacz", AGH Cracow, December 2024. 1.2
- 6. Grytczuk, J., Pawlik, B., Pleszczyński, M.: Variations on Shuffle Squares. arXiv preprint arxiv:2308.13882, (2023). 3
- 7. Komisarski, A.: personal communication. 1.2
- 8. Ochem, P., Pierron, T.: 4-tangrams are 4-avoidable. arxiv preprint arxiv:2502.20774, (2025). 1.1

Appendix

Operations on twins

Note that if there are twins of length t in a word W, then there are 2^t twins of length t whose union occupies the very same set of positions and the words they represent are the same. We obtain them by swapping (or not) every pair of corresponding elements. More precisely, let X, Y be a pair of twins in W with supports $I = \text{supp}(X) = \{i_1, \ldots, i_t\}$ and $J = \text{supp}(Y) = \{j_1, \ldots, j_t\}$. Then, for every subset $S \subseteq [t]$, the sets $I_S = \{i_h : h \notin S\} \cup \{j_h : h \in S\}$ and $J_S = \{j_h : h \notin S\} \cup \{i_h : h \in S\}$ support another pair of twins X_S, Y_S with $I_S \cup J_S = I \cup J$. Among all 2^t possibilities there is one where $i_h < j_h$ for all $h \in [t]$. Such twins are called *monotone*.

Besides swapping there are other operations which do not change the twins except for their location and mutual entanglement. Given $i_g, i_h \in \operatorname{supp}(X)$ and $j_g, j_h \in \operatorname{supp}(Y)$, g < h, with $j_g < i_h$, $w_{i_h} = w_{j_g}$, and $\{j_g + 1, \ldots, i_h - 1\} \cap \operatorname{supp}(X, Y) = \emptyset$, the (g, h)-rewiring of the monotone twins (X, Y) in W is a new pair of twins, (X', Y') with $\operatorname{supp}(X') = (\operatorname{supp}(X) \setminus \{i_h\}) \cup \{j_g\}$ and $\operatorname{supp}(Y') = (\operatorname{supp}(Y) \setminus \{j_g\}) \cup \{i_h\}$. Note that (X', Y') are still monotone. Monotone twins (X, Y) for which a rewiring operation is no longer possible, enjoy the property that within each run all elements of X precede all elements of Y.

Finally, a trivial operation of *shifting* moves all the gaps to the right end of each run. It applies whenever there is a pair of consecutive positions, i, i + 1 such that i is a gap, while, say $i + 1 \in \text{supp}(X)$. Then we switch them around, that is, for the new X', $\text{supp}(X') = (\text{supp}(X) \setminus \{i + 1\}) \cup \{i\}$. When $i + 1 \in \text{supp}(Y)$, the operation is analogous. If this operation is also no longer possible, then the resulted twins partition each run of W into three consecutive blocks (some might be empty) with the elements of X to the left, followed by the elements of Y, followed by gaps. Such twins will be called *canonical*.

Example 6. In the word $\underline{0}$ 1 $\underline{1}$ $\overline{0}$ $\overline{1}$ $\underline{1}$ $\overline{1}$ $\underline{0}$ $\overline{0}$, the indicated twins are not canonical for two reasons: a gap in front of the first 1-run and an element of Y preceding an element of X in the second 1-run. Two operations, one shifting and one rewiring brings it to the canonical form $\underline{0}$ $\underline{1}$ $\overline{1}$ $\overline{0}$ $\overline{1}$ $\overline{1}$ $\overline{0}$ (see Fig. 15).



Fig. 15. Non-canonical and canonical (red dashed) twins in the word W = 011011100.

Shuffle squares with five runs

In Proposition 2 we characterized all shuffle squares with four runs. A similar, though a bit more involved analysis can be preformed for words with *five* runs, leading to the following characterization (proof omitted).

Proposition 6. Let a, b, c, d, e be positive integers, not all even, but with a + c + e and b + d both even. Then the word $W = 1^a 0^b 1^c 0^d 1^e$ is a shuffle square if and only if $(b = d \text{ and } c \ge a + e)$ OR $(b \le d \text{ and } a \ge c \text{ and } e$ is even) OR $(b \ge d \text{ and } c \le e \text{ and } a \text{ is even})$ (see Fig. 16).

Example 7. Let $W = 1 \ 0^3 \ 1 \ 0^9 \ 1 \ 0^{11} \ 1 \ 0^7$. We have $a_3 + a_2 - a_1 = 11 + 9 - 3 = 17$ and $a_3 - a_2 - a_1 = -1$ and, thus, four solutions determined by the odd positive integers in $\{0, \ldots, 7\}$. They are, remembering that $r_1 = a_1 = 3$,

$$r_{3} = 1, r_{2} = 8, r_{4} = 3$$

$$r_{3} = 3, r_{2} = 7, r_{4} = 2,$$

$$r_{3} = 5, r_{2} = 6, r_{4} = 1,$$

$$r_{3} = 7, r_{2} = 5, r_{4} = 0$$



Fig. 16. Variants of the graph G of the shuffle square $1^a 0^b 1^c 0^d 1^e$.

and correspond, respectively, to the following four pairs of perfect twins:

The graphs corresponding to all four solutions are presented in Fig. 17.



Fig. 17. The possible graphs representing perfect twins in the shuffle square $10^{a_1}10^{a_2}10^{a_3}10^{a_4}$.