EQUIVALENCES AND DISTINCTIONS IN LEXICOGRAPHIC SHELLABILITY OF POSETS

STEPHEN LACINA AND GRACE STADNYK*

ABSTRACT. We present two perhaps surprisingly small posets, one graded and one non-graded, that are CC-shellable in the sense of Kozlov and TCL-shellable in the sense of Hersh, but not CL-shellable in the sense of Björner and Wachs. In the spirit of Björner and Wachs' recursive atom orderings (RAO) and Hersh and Stadnyk's generalized recursive atom orderings (GRAO), we also introduce a notion called recursive first atom sets (RFAS). An RFAS is a set of conditions on the atoms of each interval in a finite bounded poset P that are necessary for CC-shellability of P and sufficient for shellability of P. We also prove that under an extra condition, P is CC-shellable if and only if it admits an RFAS, in the same way that RAOs provide a reformulation of CL-shellability.

1. INTRODUCTION

Since the 1980s, lexicographic shellability has been a popular and powerful tool to establish poset shellability, a property with important topological, combinatorial, and algebraic implications. These implications stem largely from the relationship between a poset P and an associated simplicial complex, $\Delta(P)$, called the order complex of P. For instance, $\Delta(P)$ is naturally associated with a commutative ring called the Stanley-Reisner ring. If P is shellable, then $\Delta(P)$ has the homotopy type of a wedge of spheres and the Stanley-Reisner ring is Cohen-Macaulay. Poset shellability, particularly lexicographic shellability, also has applications in other mathematical contexts; many of these arise when considering the shellability of specific posets. For instance, the order complex of the subgroup lattice of a finite group is shellable if and only if the group is solvable ([13]), and a lexicographic shelling of a poset associated to a specific subspace arrangement was used to prove a complexity theory lower bound in [3].

Lexicographic shellability is a family of methods that involves labeling the cover relations of a poset P in a way that yields a shelling order of $\Delta(P)$. In particular, these labelings induce a sequence of labels for each maximal chain of P. As the faces of $\Delta(P)$ are the chains of P, when these label sequences are lexicographically ordered, the corresponding order on maximal chains gives a shelling order of $\Delta(P)$. Björner introduced the first of these methods, called EL-labeling, in [1]. At the heart of how EL-labelings produce shellings are pairs of adjacent cover relations where the corresponding pair of labels increase from bottom to top (ascents) or do not increase from bottom to top (descents).

EL-labelings have been generalized in several ways in the past few decades giving other types of lexicographic shellings. Björner and Wachs introduced CL-labelings,

^{*}Support for this research was provided by an AMS-Simons Research Enhancement Grant for Primarily Undergraduate Institution Faculty.



FIGURE 1. Relationships between different notions of lexicographic shellability and where the relationships are proved in this paper and elsewhere in the literature.

which allows for edge labels to depend on the path taken from the bottom of the poset (the root) to the edge being labeled ([4]). These types of labelings are called chain-edge labelings. As with EL-labelings, the heart of the matter are ascents and descents in the label sequences of maximal chains.

Another of Björner and Wachs' significant contributions to the theory of lexicographic shellability was a recursive perspective first introduced in [5]. They gave a formulation of CL-shellability called recursive atom ordering (RAO), which consists of the specification of total orderings of atoms of upper intervals for a bounded poset. These atom orderings are recursive in that for nested intervals, the respective atom orderings agree with each other in a particular sense. Hersh and Stadnyk introduced generalized recursive atom orderings (GRAOs) in [8] as a more flexible version of RAOs that is nonetheless sufficient for CL-shellability.

Kozlov introduced a notion of lexicographic shellability called CC-shellability in [9]. His method amounts to showing that actual ascents and descents are not strictly necessary to produce a shelling order from an edge or chain-edge labeling. Hersh then introduced in [7] a slightly different version of lexicographic shellability, what has come to be known as TCL-shellability, which similarly relies on a broader notion of ascents and descents. More specifically, for a particular labeling, it is enough to be able to identify pairs of adjacent cover relation that behave like ascents (resp. descents) though the label sequences on these chains may not actually be increasing (resp. not increasing). Hersh calls these pairs of adjacent cover relations topological ascents (resp. topological descents).

There have been several results exploring the relationships between these different formulations of lexicographic shellability. For example, Vince and Wachs in [14] and Walker in [16] constructed posets that are shellable but not lexicographically shellable. Fig. 1 illustrates the known relationships among these different versions of lexicographic shellability and their different formulations, including the relationships known from previous work and those proven in this paper.

In Theorem 4.10, we show that Kozlov's CC-shellability is equivalent to Hersh's TCL-shellability. While it is clear that any CC-labeling is also a TCL-labeling, we construct a CC-labeling for a TCL-shellable poset by introducing a relabeling technique that is intuitive in spirit but quite technical in detail. This relabeling technique further plays a central role in the proof of our recursive formulation of CC-shellability.

Hersh and Stadnyk in [8] showed that certain CC-shellable posets, those that they call "self-consistent" CC-shellable posets, are also CL-shellable. They additionally asked the question are all CC-shellable posets CL-shellable? In Theorem 3.1, we show that CC-shellability (and thus TCL-shellability) and CL-shellability, are actually distinct. In particular, we construct quite small graded and non-graded posets (see Figures 3, 4) that are CC-shellable but not CL-shellable. Our examples, together with Li's examples of posets that are CL-shellable but not EL-shellable ([12]), help to establish non-equivalence of (and in fact, strict containment among) various notions of lexicographic shellability.

Building on Björner and Wachs' recursive perspective on lexicographic shellability, we introduce in Definition 5.1 a new recursive method for shelling a poset called recursive first atom sets (RFAS). Notably, there exist posets (see Example 6.7) that admit an RFAS but for which there is no labeling that could yield any of the shellings that arise from the RFAS. Thus we specify a compatibility condition in Definition 6.1—when an RFAS satisfies this condition we call it an LCRFAS—and show in Theorem 6.5 that a bounded poset admits an LCRFAS if and only if it is CC-shellable.

As the names suggest, the essence of our recursive formulations is that we only need specify the first atom in every rooted interval rather than specifying a total order on all atoms of upper rooted intervals as in RAOs and GRAOs. In this way, for a particular interval, RFASs and LCRFASs require much less data than RAOs and GRAOs. There is another important contrast between RAOs, GRAOs and LCRFASs that illustrates that there is still substantial subtlety to LCRFASs. In RAOs and GRAOs, the atom ordering depends on the root to the bottom of an interval but it does not depend on the top element of the interval. On the other hand, the first atoms in an LCRFAS can depend on both the root to the bottom of an interval and the top element of the interval.

The structure of this paper is as follows: Section 2 provides the necessary background on posets and lexicographic shellability. In Section 3, we give two examples of posets that are CC-shellable and prove in Theorem 3.1 why these examples cannot be CL-shellable by showing neither can admit an RAO. In Section 4, we show that CC-shellability and TCL-shellability are equivalent in Theorem 4.10 by providing a method for taking a TCL-shellable poset and relabeling it with a CC-labeling. In Section 5 we define RFASs and prove that an RFAS implies shellability in Theorem 5.13. In Section 6, we introduce a LCRFASs and show that an LCRFAS is equivalent to CC-shellability in Theorem 6.5; the proof relies on the relabeling method used in the proof of Theorem 4.10. Lastly, Section 7 discusses some open questions.

2. Background

A poset (P, <) is a set, together with an order relation < that is reflexive, transitive, and antisymmetric. We will often refer to a poset (P, \leq) as P when the order relation \leq is understood. If x < y in P and there exists no z such that x < z < y, then we call x < y a cover relation. We will usually denote a cover relation by $x \leq y$. If $x \leq y$ in P, then the **interval** [x, y] is the poset $[x,y] = \{z : x \le z \le y\}$ with order relations inherited from P. The poset P is **bounded** if there exist in P a unique maximal element, denoted $\hat{1}$, and a unique minimal element, denoted $\hat{0}$. In this paper, we consider only bounded posets. The elements covering $\hat{0}$ are called the **atoms** of P while the elements covered by $\hat{1}$ are call the **coatoms** of P. A **chain** of P is a totally ordered subset of P. We will denote a chain both by $x_1 < x_2 < \ldots < x_k$ and by $\{x_1, x_2, \ldots, x_k\}$. The length of a chain c is one less than the number of elements in c. A chain is **maximal** if it is not properly contained in any other chain of P. A chain is **saturated** if it is a maximal chain of an interval of P. If all maximal chains in P have the same length, then P is said to be graded. Otherwise, P is nongraded. A rooted interval $[x,y]_r$ is the interval [x,y] together with a maximal chain r in [0,x]; r is called the root. A rooted cover relation is a rooted interval $[x, y]_r$ where x < y. If m is a chain containing x in P, we will denote by m^x the subchain of m given by $\{y: y \in m, y \le x\}.$

We will assume the reader has a basic understanding of simplicial complexes, though they may choose to refer to [15] for more details that are helpful in the context of poset shellability. We simply remind the reader that every poset P can be associated with a simplicial complex $\Delta(P)$, called the **order complex of** P, whose k-faces are the chains of length k in P. A simplicial complex Δ is said to be **shellable** if its facets can be ordered F_1, F_2, \ldots, F_s such that for $1 < j \leq s$, $\overline{F_j} \cap (\bigcup_{1 \leq i \leq j-1} \overline{F_i})$ is pure and $(\dim(F_j) - 1)$ -dimensional. A poset is shellable if its order complex is shellable.

One common method for proving a poset is shellable is by labeling the cover relations of the poset in a particular way, which then induces a total order on the maximal chains of the poset and hence the facets of the order complex; this family of methods is called **lexicographic shellability**. We now provide a brief overview of these various methods. For more complete but concise references on different types lexicographic shellability see [8] and [15].

The simplest version of lexicographic shellability arises from particular kinds of edge labelings called EL-labelings, which were introduced by Björner in [1]; a version for nongraded posets was introduced in [6]. An **edge labeling** of a poset Pis a map λ from the set of cover relations of P to the elements of another poset Λ . For any edge labeling λ of P and any saturated chain c in P, the **label sequence** $\lambda(c)$ associated to c by λ is the sequence obtained by reading the labels assigned by λ to the cover relations in *c* from the bottom of the chain to the top. An edge labeling induces a partial order on the maximal chains of the poset by taking the lexicographic order on label sequences on maximal chains. (Note when we say **lexicographic order** we mean dictionary order; in particular *aa* comes before *aaa* in lexicographic order.) An **EL-labeling** of the bounded poset *P* is an edge labeling of *P* satisfying the requirement that in any interval, exactly one maximal chain has a strictly increasing label sequence, and this label sequence comes lexicographically earlier than the label sequence on all other maximal chains in the interval. Any poset that admits an EL-labeling is said to be **EL-shellable** and is shellable [1].

More flexibility in labeling is afforded by instead labeling cover relations in chains, which allows one to label the same cover relation differently depending on which chain is being labeled. This more flexible labeling scheme is called a CElabeling. A **CE-labeling** (chain-edge labeling) of a bounded poset P is a map λ from the set of rooted cover relations of P to a poset Λ . For a CE-labeling λ , we will let $\lambda(r, x, u)$ be the label on the rooted cover relation $[x, u]_r$. For the rooted interval $[x, y]_r$ and a maximal chain c given by $x \leq x_1 \leq \ldots \leq x_t = y$ in $[x, y]_r$, the label sequence associated to c is $\lambda(r, x, x_1), \lambda(r \cup \{x_1\}, x_1, x_2), \ldots, \lambda(r \cup \{x_1\}, x_2), \ldots, \lambda(r \cup \{x_2\}, x_2), \ldots, \lambda(r \cup \{$ $\{x_1,\ldots,x_{t-1}\}, x_{t-1}, x_t\}$. We will let $\lambda(r)$ denote this label sequence for a root r. A CE-labeling λ is a **CL-labeling** if for every rooted interval $[x, y]_r$ in P, there is a unique maximal chain c with a strictly increasing label sequence in $[x, y]_r$, and the label sequence associated to c lexicographically precedes the label sequence associated to all other maximal chains in $[x, y]_r$. If the poset P admits such a labeling, it is said to be **CL-shellable** and it is shellable [4]. In [5], Björner and Wachs introduced the notion of recursive atom ordering, and showed that a poset admits a recursive atom ordering if and only if it is CL-shellable.

Definition 2.1 ([5]). A bounded poset P admits a **recursive atom ordering** (RAO) if the length of the longest chain in P is 1, or if the length of the longest chain in P is greater than 1 and there is an ordering a_1, a_2, \ldots, a_t of the atoms of P satisfying the following:

- (1) For j = 1, 2, ..., t, $[a_j, \hat{1}]$ admits a recursive atom ordering such that the atoms of $[a_j, \hat{1}]$ covering some a_i for i < j come first.
- (2) For i < j, if $a_i, a_j < y$ for some $y \in P$, then there exists some k and some $z \in P$ where $k < j, a_j < z \le y$, and $a_k < z$.

In [7], Hersh introduced the language of topological ascents and descents to describe a more general version of lexicographic shellability. We will define both CC-labelings and TCL-labelings using this language, though CC-labelings were introduced first by Kozlov in [9] without explicitly using the language of ascents and descents. Suppose that λ is a CE-labeling. Then a **topological ascent** is a pair of rooted cover relations $[u, v]_r$, $[v, w]_{r\cup v}$ in P for which the pair of labels $\lambda(r, u, v), \lambda(r \cup \{v\}, v, w)$ lexicographically precedes all other label sequences on maximal chains in $[u, w]_r$. If this pair of rooted cover relations is not a topological ascent, then it is called a **topological descent**. If a chain c in $[x, y]_r$ consists entirely of topological ascents, we say that c is **topologically ascending**.

Definition 2.2 ([9]). A CE-labeling is a **CC-labeling** if every rooted interval $[x, y]_r$ has a unique topologically ascending maximal chain, the label sequences for the maximal chains in $[x, y]_r$ are all distinct, and no label sequence is the prefix for any other. If P admits a CC-labeling, then P is called **CC-shellable**.



FIGURE 2. A poset P with an EL-labeling (left), a CL-labeling that is not an EL-labeling (middle), and a CC-labeling that is neither an EL-labeling nor a CL-labeling (right), as described in Example 2.6.

Kozlov showed that any poset that is CC-shellable is in fact shellable. He also defined EC-labelings in [9], which are CC-labelings whose labels on cover relations do not vary depending on the root, in the same way that EL-labelings can be viewed as CL-labelings whose labels do not depend on roots.

Definition 2.3 ([7]). A CE-labeling is a **TCL-labeling** if every rooted interval $[x, y]_r$ has a unique, topologically ascending maximal chain. If P admits a TCL-labeling, P is said to be **TCL-shellable**.

Hersh showed that TCL-shellable posets are shellable in [7]. Note that there exist TCL-labelings that are not CC-labelings because of the requirement that CC-labelings have distinct label sequences. There are some important consequences of the added flexibility afforded by TCL-labelings which we highlight in the following two remarks.

Remark 2.4. Notice that if λ is a TCL-labeling of the poset P and λ' is another CE-labeling of P with a topological descent at $u \leq v \leq w$ exactly when $u \leq v \leq w$ is a topological descent with respect to λ , then λ' must also be a TCL-labeling of P.

Remark 2.5. Let P be a finite bounded poset with TCL-labeling λ taking labels in a poset Λ . Observe that ordering the labels by any linear extension of Λ preserves the property of λ being a TCL-labeling. This is because the topologically ascending maximal chain in any rooted interval must strictly lexicographically precede any other maximal chain in the interval and this is preserved after taking a linear extension of Λ . Thus, we may assume that the labels of all TCL-labelings are integers. Kozlov observes this for CC-labelings in [9]. Note that this does not hold for either EL- or CL-labelings.

Example 2.6. Figure 2 gives three different labelings on a poset P. The poset on the left of the figure is labeled with an EL-labeling. The poset in the middle of the figure is labeled with a CL-labeling that is not an EL-labeling, since the label on the top left cover relation in the poset depends on the root leading from the bottom of the poset to the cover relation. Specifically, the circled labels correspond to cover relations within chains that include the leftmost atom of the poset, whereas the boxed labels give the labels on cover relations within chains that include the right of the figure is labeled by

a CC-labeling that is neither an EL-labeling nor a CL-labeling, since there exist intervals in the poset where several maximal chains in the interval have increasing label sequences. Note that the lexicographic ordering on maximal chains induced by each of the labelings in the figure is different.

In [8], Hersh and Stadnyk introduced the notion of generalized recursive atom ordering:

Definition 2.7 ([8]). A poset P admits a generalized recursive atom ordering (or GRAO) if the length of P is 1 or if the length of P is greater than 1 and there exists an ordering $a_1, a_2, \ldots a_t$ of the atoms of P satisfying the following:

- (1) (a) For $1 \le j \le t$, $[a_j, \hat{1}]$ admits a GRAO
 - (b) For atom a_j and x, w such that $a_j < x < w$, the following holds when the ordering from (i)(a) is restricted to $[a_j, w]$: either the first atom of $[a_j, w]$ is above some a_i with i < j, or no atom of $[a_j, w]$ is above any a_i with i < j.
- (2) For any $y \in P$ and $a_i, a_j < y$ with i < j, there exists some $z \in P$ and some atom $a_k \in P$ where $a_j < z \le y$, $a_k < z$, and k < j.

Furthermore, they showed that a poset admits a GRAO if and only if it is TCLshellable using a TCL-labeling that is self-consistent. A **self-consistent** TCLlabeling is a TCL-labeling with the property that whenever a and b are atoms in a rooted interval $[x, y]_r$ and a is in the lexicographically first saturated chain of $[x, y]_r$, any saturated chain containing the atom a in a rooted interval $[x, y']_r$ comes lexicographically earlier than any saturated chain containing b in $[x, y']_r$. In [8], Hersh and Stadnyk also showed that any GRAO can be transformed into an RAO, thereby showing that any poset that admits a self-consistent TCL-labeling is also CL-shellable.

3. Posets that are CC-shellable but not CL-shellable

Here we provide two examples of posets—one graded and one nongraded—that are TCL-shellable but not CL-shellable as neither admits a recursive atom ordering.

Theorem 3.1. There exist graded and non-graded posets that are CC-shellable but not CL-shellable.

Proof. Fig. 3 and Fig. 4 show examples of a graded poset P and a nongraded poset Q, respectively, that are each TCL-shellable but not CL-shellable. A TCL-labeling is given for each poset (for P, this labeling is given by bold labels in Fig. 3; for Q, the labeling is given on the left of Fig. 4). In fact, each of these labelings is a CC-labeling, and even more specifically, an EC-labeling. Both examples fail to be CL-shellable because no atom order satisfies condition 2 of a GRAO. For P, observe that for any pair of atoms a and b, there is some rank 3 element y such that a, b < y while no z in $[\hat{0}, y]$ covers both a and b. Thus, no pair of atoms can begin a recursive atom ordering of P.

Similarly, for the two atoms c and d of the nongraded poset Q, there are elements y and y' such that c < y and d < y', but y does not cover d and y' does not cover c. Thus, neither c nor d can be the first atom in a recursive atom ordering of Q. \Box

While neither poset P given in Fig. 3 nor poset Q given in Fig. 4 is CL-shellable, these posets have TCL-labelings as illustrated in their respective figures. In light of



FIGURE 3. A graded TCL-shellable poset that is not CL-shellable. A TCL-labeling of the poset is given by the bold labels. A dual EL-labeling of the poset is given by the labels in parentheses.

this distinction between TCL- and CL-shellable posets and the examples of graded and non-graded posets that are CL-shellable but not EL-shellable provided by Li in [12], one might wonder whether there exist posets that are CC-shellable but not EC-shellable. We immediately see that this is in fact the case, at least for nongraded posets, by considering Li's non-graded example:

Remark 3.2. Replacing "EL" with "EC" in the proof of Theorem 2.1 in [12] shows that the poset is CC-shellable but not EC-shellable. However, the proof technique used in the proof of Theorem 3.1 in [12] does not clearly go through for Li's graded example because an EC-labeling does not induce an RAO as shown by our Theorem 3.1.

Observe that both P and Q are dual CL-shellable as witnessed by the dual EL-labelings given in Fig. 3 and the labeling on the right of Fig. 4, respectively. However, we can easily construct posets that are TCL-shellable but neither CL nor dual CL-shellable. Take the ordinal sum of P with its dual P^* . Label this ordinal sum with the given TCL-labeling of P, the given EL-labeling of P^* , and any integer labeling the cover relation between the $\hat{1}$ of P and the $\hat{0}$ of P^* . The resulting labeling is clearly a TCL-labeling (an EC-labeling more specifically) while

9



FIGURE 4. An nongraded TCL-shellable poset that is not CL-shellable. A TCL-labeling of the poset is shown on the left. A dual EL-labeling of the poset is shown on the right.

the resulting poset is graded and neither CL nor dual CL-shellable. The same construction using Q, Q^* , and their labelings produces a nongraded example.

Of further note, neither P nor its dual is the face lattice of a regular CW-complex. To see this, recall the following consequence of Björner's Proposition 3.1 in [2]. In the face poset of a regular CW-complex, every non-trivial open lower interval (\emptyset, x) is homeomorphic to a sphere. Observe that for the element $x \in P$ with rank 3 and degree 7, $[\hat{0}, x]$ has three topologically descending chains, namely the saturated chains labeled (1, 6, 1), (1, 7, 1), and (1, 9, 1). Thus, $(\hat{0}, x)$ is homotopy equivalent to a wedge of three 1-spheres, so P is not the face lattice of a regular CW-complex. On the other hand, for any atom x of P the order complex of $(x, \hat{1})$ is shown in Fig. 5. This is clearly not homeomorphic to a sphere despite being homotopy equivalent to a 1-sphere. Thus, the dual of P is also not the face lattice of a regular CWcomplex. This shows that P is not CL-shellable for a somewhat different reason than the examples due to Vince and Wachs [14] and Walker [16].



FIGURE 5. The order complex of $(x, \hat{1})$.

4. TCL-shellable is Equivalent to CC-shellable

We begin by introducing a labeling technique based on any total order of the maximal chains of a poset. We use this technique later in this section to show that TCL-shellability is equivalent to CC-shellability. Then, in Section 6, we use the same technique to build CE-labelings from certain sets that we will call recursive first atom sets.

While this labeling is somewhat technical, the essence of the labeling is more straightforward. First, we fix a total order on the maximal chains of P. We wish to label a rooted cover relation $(r, x \leq y)$ by the index of the first maximal chain m_i in this total order that contains both r and y. We do this except when a different rooted cover relation $(r, x \leq y')$ appears in maximal chains before m_i and again after m_i in the total order, that is, when the chain m_i is "sandwiched" by two other chains containing the rooted cover relation $(r, x \leq y')$ not $(r, x \leq y')$. In this case, the distinct rooted cover relations $(r, x \leq y)$ and $(r, x \leq y')$ receive the same label. In any case, a rooted cover relation is always labeled by the position in the total order of some maximal chain which contains the root.

Definition 4.1. Let P be a finite bounded poset. Fix a total order $\Gamma : m_1, \ldots, m_t$ on the maximal chains of P. Define a CE-labeling λ of P in the following manner. For a root r from $\hat{0}$ to x, let a_1, \ldots, a_s be the atoms of $[x, \hat{1}]_r$ in the order in which they first appear with r in a maximal chain in Γ . Label the rooted edge $x < a_j$ with label $\lambda(r, x, a_j)$ defined as follows: Let m_{i_1} be the earliest chain of Γ containing rand a_1 . Set $\lambda(r, x, a_1) = i_1$. Say the first j - 1 rooted edges have been labeled. Let m_{i_j} be the first chain of Γ containing r and a_j . If there exists an atom a_h with h < j such that a_h and r are contained in maximal chains m_k and m_l with $k < i_j < l$, set $\lambda(r, x, a_j) = \lambda(r, x, a_h)$. Otherwise, set $\lambda(r, x, a_j) = i_j$.

Throughout the remainder of this section, given a root r, we routinely use the notation i_j to denote the index of the earliest maximal chain (in the total order on maximal chains) containing the atom a_j and the root r. Two important consequences of Definition 4.1, both of which are used in proofs later in this section, are provided in the following remarks.

Remark 4.2. Notice that using Definition 4.1, a rooted cover relation $x \leq y$ with root r can never have a label larger than the index of the first chain in Γ that contains $r \cup \{x, y\}$.

Remark 4.3. Observe that by Definition 4.1 we have the following: if a rooted cover relation $x \leq y$ with root r is not labeled by the position of the first chain m_j in Γ containing $r \cup \{x, y\}$, then there is an element y' covering x such that the first chain m_i in Γ containing $r \cup \{x, y'\}$ and some chain m_k containing $r \cup \{x, y'\}$ have i < j < k.

The following proposition is used as a key component in many of the proofs that follow, both in this section and in Section 6. In essence, it describes characteristics of the total order Γ that force a set of rooted cover relations to share the same label when labeling the poset using Definition 4.1. It also shows that Definition 4.1 yields a CE-labeling with the property that no two maximal chains in any rooted have the same label sequence. Here and throughout the rest of the paper, we use the notation m^x to refer to the the elements not above x in a chain m when m contains x. **Proposition 4.4.** Let P be a finite bounded poset. Fix a total order $\Gamma : m_1, \ldots, m_t$ on the maximal chains of P. Let λ be the CE-labeling of P determined by Γ as in Definition 4.1. The following then hold:

- (i) Let a_1, \ldots, a_s be the atoms of the rooted interval $[x, \hat{1}]_r$ in the order induced by Γ as in Definition 4.1. If $\lambda(r, x, a_k) = \lambda(r, x, a_l)$ with $k \leq l$, then we have $\lambda(r, x, a_k) = \lambda(r, x, a_j)$ for all j such that $k \leq j \leq l$.
- (ii) If maximal chains m and m' of the rooted interval [x, 1]_r contain distinct atoms of [x, 1], then the label sequence associated to m is distinct from the label sequence associated to m'. Further, if P is not graded, then for any distinct maximal chains c and c' of [x, y]_r, the label sequence associated to c is not a prefix of the label sequence associated to c'.

Proof. We show (i) by induction on l - k. If l - k = 0, there is nothing to check. Suppose that for some $n \ge 1$ whenever l - k < n, we have that for all j such that $k \le j \le l$, $\lambda(r, x, a_k) = \lambda(r, x, a_j)$. Now suppose l - k = n. Since $n \ge 2$, there is a smallest j' such that j' < l and $r \cup a_{j'}$ is contained in some m_p , where $i_{j'} < i_l < p$. Thus, we have $\lambda(r, x, a_{j'}) = \lambda(r, x, a_l)$ by Definition 4.1. Then since $\lambda(r, x, a_l) = \lambda(r, x, a_k)$, we have $\lambda(r, x, a_{j'}) = \lambda(r, x, a_k)$ as well.

If $j' \leq k$, then for all j such that k < j < l we have $i_{j'} \leq i_k < i_j < i_l < p$. We therefore have $\lambda(r, x, a_k) = \lambda(r, x, a_j)$ by Definition 4.1 for all j such that $k \leq j \leq l$.

Otherwise, k < j' < l. Since j' - k < l - k = n and $\lambda(r, x, a_{j'}) = \lambda(r, x, a_k)$, the induction hypothesis says that $\lambda(r, x, a_k) = \lambda(r, x, a_j)$ for all j such that $k \leq j \leq j'$. And since l - j' < l - k = n and $\lambda(r, x, a_{j'}) = \lambda(r, x, a_l)$, the induction hypothesis says that $\lambda(r, x, a_j) = \lambda(r, x, a_l)$ for all j such that $j' \leq j \leq l$. Therefore, $\lambda(r, x, a_k) = \lambda'(r, x, a_j)$ for all $k \leq j \leq l$.

For (ii), observe the following. First, by Definition 4.1, a cover relation with root r is labeled by the position in Γ of some maximal chain that contains r. Let $x \leq a_1 \leq b_1$ be contained in m and $x \leq a_2 \leq b_2$ be contained in m' with $a_1 \neq a_2$. The label $\lambda(r \cup a_1, a_1, b_1)$ is the position in Γ of some maximal chain which contains $r \cup a_1$ while the label $\lambda(r \cup a_2, a_2, b_2)$ is the position in Γ of some maximal chain which contains $r \cup a_2$. No maximal chain of P contains both $r \cup a_1$ and $r \cup a_2$ because a_1 and a_2 are distinct. Therefore, $\lambda(r \cup a_1, a_1, b_1) \neq \lambda(r \cup a_2, a_2, b_2)$. This in turn implies that the label sequence associated to m is distinct from the label sequence associated to m'.

Now for any distinct maximal chains c and c' of a rooted interval $[x, y]_r$, there is some minimal z < y such that $c^z = c'^z$ but c and c' contain distinct atoms of [z, y]. Then by the previous argument the label sequence associated to c is not a prefix of the label sequence associated to c'.

We now turn to showing that any poset that admits a TCL-labeling also admits a CC-labeling in Lemma 4.8. We first make an observation about initial sections of label sequences that arise from TCL-labelings. It is used in the two upcoming technical propositions concerning topological descents and their preservation under a relabeling given by Definition 4.1. Both propositions are necessary for the proof of Lemma 4.8.

Remark 4.5. Let P be a finite bounded poset with TCL-labeling λ taking values in the integers. Fix a total order $\Gamma: m_1, \ldots, m_t$ on the maximal chains of P which is consistent with the lexicographic order on maximal chains induced by λ . If r is a saturated chain containing $\hat{0}$ which is contained in m_i and m_k with i < k, then for each j with $i \leq j \leq k$, $\lambda(r)$ is an initial section of the label sequence $\lambda(m_j)$.

Proposition 4.6. Let *P* be a finite bounded poset with TCL-labeling λ taking values in the integers. Suppose that maximal chain *m* has a topological descent at x < y < z and that *c* given by $x < x_1 < x_2 < \cdots < z$ is the topologically ascending maximal chain of $[x, z]_{m^x}$. Then either $\lambda(m^x, x, x_1) < \lambda(m^x, x, y)$ or $\lambda(m^x, x, x_1) = \lambda(m^x, x, y)$ and $\lambda(m^x \cup x_1, x_1, x_2) < \lambda(m^x \cup y, y, z)$.

Proof. This follows because the label sequence associated to c with respect to m^x strictly precedes lexicographically the label sequence associated to x < y < z with respect to m^x

Proposition 4.7. Let *P* be a finite, bounded poset with a TCL-labeling λ . By Remark 2.5, we may take λ to be a labeling by integers. Fix a total order Γ : m_1, \ldots, m_t on the maximal chains of *P* that agrees with the lexicographic order on maximal chains induced by λ . Let λ' be the CE-labeling of *P* determined by Γ as in Definition 4.1. Suppose $x \le u \le b$ and $x \le y \le d$ with $u \ne y$ and let *r* be a root of *x*. If $\lambda(r, x, u) < \lambda(r, x, y)$, then $\lambda'(r, x, u) < \lambda'(r, x, y)$. Also, if $\lambda(r, x, u) = \lambda(r, x, y)$ and $\lambda(r \cup u, u, b) < \lambda(r \cup y, y, d)$, then $\lambda'(r \cup u, u, b) < \lambda'(r \cup y, y, d)$.

Proof. Let a_1, \ldots, a_s be the atoms of $[x, \hat{1}]_r$ in the order in which they first appear with r in a maximal chain in Γ .

Suppose that $\lambda(r, x, u) < \lambda(r, x, y)$. Let a_j be the earliest atom such that $\lambda'(r, x, a_j) = \lambda'(r, x, u)$. Notice that by Definition 4.1, $\lambda'(r, x, a_j) = i_j$ where m_{i_j} is the first maximal chain in Γ containing $r \cup a_j$. Otherwise, there would be some earlier atom than a_j , say a, such that $\lambda'(r, x, a) = \lambda'(r, x, u)$. Similarly, let a_k be the earliest atom such that $\lambda'(r, x, a_k) = \lambda'(r, x, y)$. So, $\lambda'(r, x, a_k) = i_k$ where m_{i_k} is the first maximal chain in Γ containing $r \cup a_k$. By Proposition 4.4 (i) and Remark 4.5, $\lambda(r, x, a_j) = \lambda(r, x, u)$ and $\lambda(r, x, a_k) = \lambda(r, x, y)$. Since $\lambda(r, x, u) < \lambda(r, x, y)$, we have $\lambda(r, x, a_j) < \lambda(r, x, a_k)$ and thus $i_j < i_k$. Thus, $\lambda'(r, x, u) < \lambda'(r, x, y)$.

Now suppose that $\lambda(r, x, u) = \lambda(r, x, y)$ and $\lambda(r \cup u, u, b) < \lambda(r \cup y, y, d)$. Notice that $r \cup u \neq r \cup y$, but $\lambda(r \cup u) = \lambda(r \cup y)$, so $\lambda(r \cup u \cup b)$ strictly precedes $\lambda(r \cup y \cup d)$ lexicographically. Let v_l be the earliest atom of $[u, \hat{1}]_{r \cup u}$ such that $\lambda'(r \cup u, u, v_l) = \lambda'(r \cup u, u, b)$. Notice that by Definition 4.1, $\lambda'(r \cup u, u, v_l) = i_l$ where m_{i_l} is the first maximal chain in Γ containing $r \cup u \cup v_l$. Similarly, let w_n be the earliest atom of $[y, \hat{1}]_{r \cup y}$ such that $\lambda'(r \cup y, y, w_n) = \lambda'(r \cup y, y, d) = i_n$ where m_{i_n} is the first maximal chain in Γ containing $r \cup y \cup w_n$. Again by Proposition 4.4 (i) and Remark 4.5, we have $\lambda(r \cup u, u, v_l) = \lambda(r \cup u, u, b)$ and $\lambda(r \cup y, y, w_n) = \lambda(r \cup y, y, d)$. Since $\lambda(r \cup u, u, b) < \lambda(r \cup y, y, d)$, we have $\lambda(r \cup u, u, v_l) < \lambda(r \cup y, y, w_n)$ and thus $i_l < i_n$. Thus, $\lambda'(r \cup u, u, b) < \lambda'(r \cup y, y, d)$.

We are now ready to prove our main result of this section, Lemma 4.8, which yields Theorem 4.10 when combined with Theorem 5.8 in [8].

Lemma 4.8. Let P be a finite bounded poset with a TCL-labeling λ . By Remark 2.5, we may take λ to be a labeling by integers. Fix a total order $\Gamma : m_1, \ldots, m_t$ on the maximal chains of P that agrees with the lexicographic order on maximal chains induced by λ . Let λ' be the CE-labeling of P determined by Γ as in Definition 4.1. Then λ' is a CC-labeling.

Proof. By Proposition 4.4 (ii), in each rooted interval, λ' gives distinct label sequences to distinct maximal chains and no label sequence of a maximal chain is a prefix of the label sequence of any other maximal chain. Thus, it suffices to show that λ' preserves the topological descents of λ (see Remark 2.4). Suppose that the maximal chain m has a topological descent at $x \leq y \leq z$ with respect to λ . Let $r = m^x$ be the chain that agrees with m up to x. As in Definition 4.1, take the atoms a_1, \ldots, a_s of $[x, \hat{1}]_r$ in the order in which they first appear with r in a maximal chain in Γ . So, $y = a_l$ for some l. Let $c : x \leq a_k \leq w \leq \cdots \leq z$ be the topologically ascending maximal chain of $[x, z]_r$ with respect to λ . By Proposition 4.6, either

- (a) $\lambda(r, x, a_k) < \lambda(r, x, a_l)$ or
- (b) $\lambda(r, x, a_k) = \lambda(r, x, a_l)$ and $\lambda(r \cup a_k, a_k, w) < \lambda(r \cup a_l, a_l, z)$.

If (a) holds, Proposition 4.7 implies $\lambda'(r, x, a_k) < \lambda'(r, x, a_l)$, so *m* has a topological descent at $x < a_l < z$ with respect to λ' .

If (b) holds, we will show that either we have $\lambda'(r, x, a_k) < \lambda'(r, x, a_l)$ or both $\lambda'(r, x, a_k) = \lambda'(r, x, a_l)$ and $\lambda'(r \cup a_k, a_k, w) < \lambda'(r \cup a_l, a_l, z)$. In either case, $x < a_l < z$ is a topological descent with respect to λ' .

Suppose that (b) holds. We first establish some notation. Let $a_{l'}$ be the earliest atom of $[x, \hat{1}]_r$ such that $\lambda'(r, x, a_{l'}) = \lambda'(r, x, a_l)$ and let $a_{k'}$ be the earliest atom of $[x, \hat{1}]_r$ such that $\lambda'(r, x, a_{k'}) = \lambda'(r, x, a_k)$. Let $m_{i_{l'}}$ be the first chain in Γ containing $r \cup a_{l'}$ and let $m_{i_{k'}}$ be the first chain in Γ containing $r \cup a_{k'}$. Thus, we have $\lambda'(r, x, a_{k'}) = \lambda'(r, x, a_k) = i_{k'}$ and $\lambda'(r, x, a_{l'}) = \lambda'(r, x, a_l) = i_{l'}$. We also necessarily have $k' \leq k$ and $l' \leq l$. We are now ready to show that $\lambda'(r, x, a_k) \leq \lambda'(r, x, a_l)$.

If $k' \leq l'$, then $i_{k'} \leq i_{l'}$ so $\lambda'(r, x, a_k) \leq \lambda'(r, x, a_l)$. If l' < k', we must consider the two cases that either l < k or k < l. Suppose k < l. Observe that $r \cup c$ followed by the portion of m above z is some maximal chain containing $r \cup a_k$ which precedes m, so m_{i_k} , the first maximal chain containing $r \cup a_k$, precedes m. Since l < k, m_{i_l} precedes m_{i_k} and m_{i_k} precedes m. Since both m_{i_l} and m contain $r \cup a_l$ while m_{i_k} is the first maximal chain containing $r \cup a_k$, Definition 4.1 gives $\lambda'(r, x, a_k) =$ $\lambda'(r, x, a_l)$. If k < l, then knowing that $\lambda'(r, x, a_{l'}) = \lambda'(r, x, a_l)$ and $l' < k' \leq k < l$ allows us to use Proposition 4.4 (i) to conclude that $\lambda'(r, x, a_k) = \lambda'(r, x, a_l)$.

If $\lambda'(r, x, a_k) < \lambda'(r, x, a_l)$, then $x < a_l < z$ is a topological descent in m with respect to λ' and we are done. It remains to consider the case where $\lambda'(r, x, a_k) = \lambda'(r, x, a_l)$. Recall that we are in case (b), that is, $\lambda(r, x, a_k) = \lambda(r, x, a_l)$ and $\lambda(r \cup a_k, a_k, w) < \lambda(r \cup a_l, a_l, z)$. So, by Proposition 4.7, $\lambda'(r \cup a_k, a_k, w) < \lambda'(r \cup a_l, a_l, z)$. Thus, $x < a_l < z$ is a topological decent in m with respect to λ' .

Remark 4.9. Notice that the proof of Lemma 4.8 shows that for a given TCL-labeling of a bounded poset P, the corresponding CC-labeling defined by Definition 4.1 has topological descents in exactly the same locations as the TCL-labeling.

Theorem 4.10. A finite bounded poset P is TCL-shellable if and only if P is CC-shellable.

Proof. The forward direction is Lemma 4.8. The backward direction is Theorem 5.8 in [8]. \Box



FIGURE 6. The posets P (left) and Q (right) discussed in Example 5.2 and Example 5.3, respectively.

5. Recursive First Atom Sets

In the spirit of recursive atom orderings and generalized recursive atom orderings, we now introduce recursive first atom sets (RFAS) as a new method for producing shelling orders of finite bounded posets. Posets exist that admit an RFAS but that do not admit an RAO or GRAO (see Figure 3 and Theorem 6.5). These RFASs have the benefit of only requiring us to specify a first atom for each rooted interval, as opposed to requiring us to specify total orderings on atoms of upper rooted intervals.

As the following definition is somewhat technical, the reader might find it useful to refer to Figures 6 and 7 and Examples 5.2 and 5.3 as they read.

Definition 5.1. The bounded poset P admits a **recursive first atom set** (**RFAS**) if the length of P is 1 or if for each x < y in P and each root r for [x, y], there exists an atom of [x, y], denoted $\Omega(r, x, y)$ and called the **first atom** of $[x, y]_r$, satisfying the following whenever a is an atom of $[x, y]_r$ and b is the first atom of $[a, y]_{r\cup a}$:

- (i) a is the first atom in $[x, y]_r$ if and only if a is the first atom in $[x, b]_r$
- (ii) if a is not the first atom in $[x, y]_r$ and a' is the first atom in $[x, y]_r$, then there exist atoms a_1, \ldots, a_p of [x, y] and elements b_1, \ldots, b_{p-1} such that:
 - (a) a_p is the first atom of $[x, b]_r$,
 - (b) $a_1 = a'$, and
 - (c) b_i is the first atom of $[a_{i+1}, y]_{r \cup a_i}$ and a_i is the first atom of $[x, b_i]_r$ for all i with $1 \le i \le p-1$

The goal of Definition 5.1 is to specify an atom in every rooted interval of the bounded poset P in such a way that (1) a total order on the maximal chains of Pis completely determined by these specified atoms, and (2) this total order on the maximal chains is a shelling order. The sense in which we determine a total order on maximal chains from a specification of atoms is given by the following notion of consistency. Let C be such a specification of atoms. Say that a total order on maximal chains of P is consistent with C if, for any rooted interval $[u, v]_r$ in P, the earliest maximal chain in the total order that contains u, v, and r also contains the specified atom in the interval $[u, v]_r$. Condition (i) is necessary to guarantee that there exists a total order that is consistent with a given specification of atoms.

15



FIGURE 7. A diagram illustrating condition (ii) of Definition 5.1

Condition (ii), which is illustrated in Fig. 7, is necessary to ensure that the resulting total order is a shelling order. The following two examples highlight these points.

Example 5.2. Consider the poset P shown in Figure 6 and the collection C of first atoms for intervals in P where a is first in [x, b], a' is first in [x, b'] and [x, y], and b is first in [a', y] and [a, y]. Note there is only one root for each of these intervals, so we are not specifying the root here, and we are not specifying first atoms in intervals containing only one atom. Notice that C does not satisfy the backwards direction of Definition 5.1 (i). Since a' is first in [x, y] and b is first in [a', y], any total order on maximal chains that is consistent with C must begin with the chain $m = \{x, a', b, y\}$. Notice that m is then the earliest chain containing x and b. However, m does not contain a, the first atom in [x, b]. So, any such total order is not consistent with C.

Likewise, consider the collection \mathcal{C}' of first atoms where a' is first in [x, b], a is first in [x, b'] and [x, y], and b is first in [a', y] and [a, y]. Notice \mathcal{C}' does not satisfy the forward direction of Definition 5.1 (i). Since a is first in [x, y] and b is first in [a, y], any total order on maximal chains that is consistent with \mathcal{C}' must begin with the chain $m' = \{x, a, b, y\}$. Notice that m' is then the earliest chain containing x and b. However, m' does not contain a', the first atom in [x, b]. So, any such total order is not consistent with \mathcal{C}' .

Example 5.3. For the poset Q in Fig. 6, define a first atom set as follows: a'' is first in [x, b''] and [x, y], a is first in [x, b], a' is first in [x, b'], b'' is first in [a'', y], b' is first in [a, y], and b is first in [a', y]. Just as in Example 5.2, we need not specify roots for any of these intervals. It can easily be checked that this first atom set satisfies condition (i) but not condition (ii) of Definition 5.1. Further, Q is clearly not shellable.

Remark 5.4. Observe that if P admits an RFAS Ω , then restricting Ω to any closed rooted interval $[x, y]_r$ of P clearly gives an RFAS of [x, y].

The following several definitions and propositions are used to prove Theorem 5.13 which says that an RFAS produces a shelling order of a finite bounded poset. While these intermediate steps are quite technical, in essence they are identifying structures from an RFAS that behave like ascents and descents (or topological ascents and topological descents) in a lexicographic shelling. First we note that although we have shown through Theorem 3.1 that topological ascents and topological descents are structurally different than ascents and descents, these two different notions of ascents and descents function in exactly the same way in terms of proving that the lexicographic orders from the different labeling types are shelling orders. It is for this purpose of producing shellings that we want to draw parallels between structures in an RFAS and ascents and descents, so we only use the words "ascent" and "descent" throughout the rest of this section, even though we could correctly say "topological ascent" as well.

For a finite bounded poset with an RFAS, the following definition identifies a chain in each rooted interval that plays a role similar to that of an ascending chain in a lexicographic shelling.

Definition 5.5. Let *P* be a finite, bounded poset and let Ω be an RFAS of *P*. For any rooted interval $[x, y]_r$, define the **first atom chain of** $[x, y]_r$, denoted c(r, x, y), to be the unique maximal chain c(r, x, y) of $[x, y]_r$ given by $x = z_0 < z_1 < \cdots < z_q = y$ satisfying z_{i+1} is the first atom of $[z_i, y]_{r \cup \{z_0, \dots, z_i\}}$ according to Ω for all *i* such that $0 \le i \le q-2$.

Now we use an RFAS to identify chains that function the same way that descents do in a lexicographic shelling.

Definition 5.6. Let *P* be a finite, bounded poset and let Ω be an RFAS of *P*. For a maximal chain *m* of *P* given by $\hat{0} = x_0 < x_1 < \cdots < x_n = \hat{1}$, say that $x_i < x_{i+1} < x_{i+2}$ is a **pseudo descent** of *m* if x_{i+1} is not the first atom of $[x_i, x_{i+2}]_{m^{x_i}}$ according to Ω .

The following proposition shows that for an RFAS, any chain that does not behave like an ascending chain must contain a chain that behaves like a descent.

Proposition 5.7. Let *P* be a finite, bounded poset and let Ω be an RFAS of *P*. Let $[x, y]_r$ be a rooted interval of *P* and let *m* be a maximal chain of $[x, y]_r$. If $m \neq c(r, x, y)$, then *m* contains a pseudo descent.

Proof. Say that m is given by $x = x_0 < x_1 < \cdots < x_p = y$ and $m \neq c(r, x, y)$. We prove that m contains a pseudo descent by induction on the length of the longest chain in [x, y]. It holds vacuously if the longest chain in P is length 1. Suppose [x, y] has some chain longer than length 1. Since $m \neq c(r, x, y)$, there exists a unique k with $0 \leq k \leq p-2$ such that $x_k \in c(r, x, y)$ and $c(r, x, y)^{x_k} = m^{x_k}$ while c(r, x, y) and m contain different atoms of $[x_k, \hat{1}]$. By definition, c(r, x, y) contains $\Omega(m^{x_k}, x_k, \hat{1})$ the first atom of $[x_k, \hat{1}]_{m^{x_k}}$, while m contains $x_{k+1} \neq \Omega(m^{x_k}, x_k, \hat{1})$. So, x_{k+1} is not the first atom of $[x_k, \hat{1}]_{m^{x_k}}$ according to Ω . If x_{k+2} is not the first atom of $[x_{k+1}, \hat{1}]_{m^{x_{k+1}}}$ according to Ω , then above x_{k+1} , m is not $c(r \cup$ $\{x_1, \ldots, x_{k+1}\}, x_{k+1}, \hat{1})$. Thus by induction, m has a pseudo descent above x_{k+1} . If x_{k+2} is the first atom of $[x_{k+1}, \hat{1}]_{m^{x_{k+1}}}$ according to Ω , then by Definition 5.1 (i) the first atom $\Omega(m^{x_k}, x_k, x_{k+2})$ of $[x_k, x_{k+2}]_{m^{x_k}}$ is not x_{k+1} . In this case, $x_k < x_{k+1} < x_{k+2}$ is a pseudo descent of m.

On the other hand, the next proposition shows that in an RFAS, the chains that function as ascending chains do not contain any of the chains that behave like descents.

Proposition 5.8. Let P be a finite, bounded poset and let Ω be an RFAS of P. For any rooted interval $[x, y]_r$, the first atom chain c(r, x, y) does not contain a pseudo descent.

Proof. This follows directly from the forward direction of condition (i) in Definition 5.1.

Now we use these ascent-like (first atom chains) and descent-like (pseudo descents) structures to construct a total order on the maximal chains of a finite bounded poset P with an RFAS Ω . In particular, in the following definition and the next three propositions, we use the first atom chains and pseudo descents to define a partial order on the maximal chains of the poset, denoted \mathcal{M}_{Ω} . Any linear extension of \mathcal{M}_{Ω} will be shown to give a shelling order in Theorem 5.13. Note that this partial order is in essence a maximal chain descent order (MCDO) as introduced in [10]. Theorem 5.13 shows that \mathcal{M}_{Ω} encodes shellings of P the same way that an MCDO does.

Definition 5.9. Let P be a finite, bounded poset with an RFAS Ω . For two maximal chains m and m' of P, define a relation $m \to m'$ exactly when there exist $x \leq y \leq z$ such that x and z are both in m and m', m and m' agree below x and above z, $x \leq y \leq z$ is a pseudo descent of m', and m between x and z is the first atom chain $c(m^x, x, z)$. Denote by \preceq the reflexive and transitive closure of all relations of the form $m \to m'$. This relation together with the set of maximal chains of P will be denoted \mathcal{M}_{Ω} .

We will show M_{Ω} is a partial order in Proposition 5.12. The next two propositions are technical results necessary for the proof of Proposition 5.12.

Proposition 5.10. Let *P* be a finite, bounded poset with an RFAS Ω . Suppose *m* is a maximal chain of *P* that is not the first atom chain $c(\{\hat{0}\}, \hat{0}, \hat{1})$. Then there exists a sequence of maximal chains $c(\{\hat{0}\}, \hat{0}, \hat{1}) \to m_1 \to m_2 \to \cdots \to m$.

Proof. Let m be a maximal chain of P given by $\hat{0} = x_0 < x_1 < \cdots < x_s = \hat{1}$ such that m is not the first atom chain $c(\{\hat{0}\}, \hat{0}, \hat{1})$. We show there is a sequence of maximal chains $c(\{\hat{0}\}, \hat{0}, \hat{1}) \to m_1 \to m_2 \to \cdots \to m$ by induction on the length of the longest chain in P. If the length of the longest chain in P is 1, this is vacuously true. If the length of the longest chain in P is 2, this is clear by Proposition 5.7. Suppose that for some $n \geq 3$, whenever the length of the longest chain of P is at most n-1, there is a sequence of maximal chains $c(\{\hat{0}\}, \hat{0}, \hat{1}) \to m_1 \to m_2 \to \cdots \to m$.

Say the length of the longest maximal chain in P is n. Since the longest maximal chain of $[x_1, \hat{1}]$ has length less than n, by induction there is a sequence of maximal chains $\hat{0} \cup c(\{\hat{0}, x_1\}, x_1, \hat{1}) \rightarrow m'_1 \rightarrow m'_2 \rightarrow \cdots \rightarrow m$. Now there are two cases: either x_1 is the first atom in $[\hat{0}, \hat{1}]$ according to Ω or not. Suppose x_1 is the first atom in $[\hat{0}, \hat{1}]$ according to Ω . Then $\hat{0} \cup c(\{\hat{0}, x_1\}, x_1, \hat{1}) = c(\{\hat{0}\}, \hat{0}, \hat{1})$, so we have the desired sequence of maximal chains.

Suppose that x_1 is not the first atom in $[\hat{0}, \hat{1}]$ according to Ω . Let b be the first atom of $[x_1, \hat{1}]_{\{\hat{0}, x_1\}}$, that is, b is the element of $c(\{\hat{0}, x_1\}, x_1, \hat{1})$ covering x_1 . Then by Definition 5.1 (ii), there exist atoms a_1, \ldots, a_p of P and elements b_1, \ldots, b_{p-1} such that a_p is the first atom of $[x, b]_r$, a_1 is the first atom of $[\hat{0}, \hat{1}]$ according to Ω , b_i is the first atom of $[a_{i+1}, y]_{r\cup a_i}$, and a_i is the first atom of $[x, b_i]_r$ for all i with $1 \leq i \leq p-1$. Then we have $c(\{\hat{0}\}, \hat{0}, b) \cup c(\{\hat{0}, x_1\}, x_1, \hat{1})_b \to \hat{0} \cup c(\{\hat{0}, x_1\}, x_1, \hat{1})$ and $c(\{\hat{0}\}, \hat{0}, b)$ contains a_p . Then again by induction we have a sequence of maximal chains $\hat{0} \cup c(\{\hat{0}, a_p\}, a_p, \hat{1}) \to \tilde{m_1} \to \tilde{m_2} \to \cdots \to c(\{\hat{0}\}, \hat{0}, b) \cup c(\{\hat{0}, x_1\}, x_1, \hat{1})_b$. Note that $\hat{0} \cup c(\{\hat{0}, a_p\}, a_p, \hat{1})$ contains b_{p-1} , so we can repeat this process with b_{p-1} and a_{p-1} and so on until reaching $\hat{0} \cup c(\{\hat{0}, a_1\}, a_1, \hat{1})$ which is the first atom chain $c(\{\hat{0}\}, \hat{0}, \hat{1})$. Together, this gives a sequence of maximal chains $c(\{\hat{0}\}, \hat{0}, \hat{1}) \to m_1 \to m_2 \to \cdots \to m$. This shows that there exists such a sequence which begins with $c(\{\hat{0}\}, \hat{0}, \hat{1})$.

Proposition 5.11. Let P be a finite, bounded poset with an RFAS Ω . Then any sequence of maximal chains obtained by replacing pseudo descents in maximal chains of P with the corresponding first atom chains must eventually reach $c(\{\hat{0}\}, \hat{0}, \hat{1})$.

Proof. First, note that every maximal chain m has some sequence of maximal chains $c(\{\hat{0}\}, \hat{0}, \hat{1}) \to m_1 \to m_2 \to \cdots \to m_k = m$ by Proposition 5.10. Let k be maximal among such sequences. We will show the desired result by induction on k, that is, on the length of the longest such sequence. We first show the base case of k = 1. We do this by showing that if $c(\{\hat{0}\}, \hat{0}, \hat{1}) \to m$ is the longest such sequence, then m has exactly one pseudo descent. Suppose x < y < z is the pseudo descent of m which gives $c(\{\hat{0}\}, \hat{0}, \hat{1}) \to m$, so y is not in $c(\{\hat{0}\}, \hat{0}, \hat{1})$. Thus, we have $m^x = c(\{\hat{0}\}, \hat{0}, x)$. Then by Proposition 5.8, m^x has no pseudo descents below x. Assume seeking contradiction that m has a pseudo descent x' < y' < z'with $y \leq x'$. Let m' be the maximal chain obtained from m by replacing the pseudo descent above y with the corresponding first atom chain. Then we have $m' \to m$ while $m' \neq c(\{\hat{0}\}, \hat{0}, \hat{1})$ since m' contains y. By Proposition 5.10, there is a sequence of maximal chains $c(\{\hat{0}\}, \hat{0}, \hat{1}) \to m'_1 \to m'_2 \to \cdots \to m'$. But this contradicts that the longest sequence from $c(\{\hat{0}\}, \hat{0}, \hat{1})$ to m has length one, since $c(\{0\}, 0, 1) \to m'_1 \to m'_2 \to \cdots \to m' \to m$ has length at least two. Similarly, assuming that m contains a pseudo descent x' < x < y, contradicts the fact that the longest sequence from $c(\{0\}, 0, 1)$ to m has length one. Hence, m has a single pseudo descent which means that $c({\hat{0}}, \hat{0}, \hat{1}) \to m$ is the only such sequence for m proving the base case.

Now suppose that for some $n \ge 2$, whenever a maximal chain m has k < n, any sequence of maximal chains obtained by beginning with m and replacing pseudo descents with the corresponding first atom chains must eventually reach $c(\{\hat{0}\}, \hat{0}, \hat{1})$. Say m is a maximal chain with k = n. Since $m \ne c(\{\hat{0}\}, \hat{0}, \hat{1})$, m has at least one pseudo descent by Proposition 5.7. For any maximal chain m' such that $m' \rightarrow m$, that is, for any maximal chain obtained from m by replacing a pseudo descent with the corresponding first atom chain, the longest sequence $c(\{\hat{0}\}, \hat{0}, \hat{1}) \rightarrow \tilde{m}_1 \rightarrow \tilde{m}_2 \rightarrow$ $\dots \rightarrow m'$ is at most length n - 1. Otherwise, there would be such a sequence from $c(\{\hat{0}\}, \hat{0}, \hat{1})$ to m of length at least n + 1. Then by the induction hypothesis, any sequence of maximal chains obtained by beginning with m' and replacing pseudo descents with the corresponding first atom chains must eventually reach $c(\{\hat{0}\}, \hat{0}, \hat{1})$. Then since m' was obtained from an arbitrary pseudo descent of m, any sequence of maximal chains obtained by beginning with m and replacing pseudo descents with the corresponding first atom chains must eventually reach $c(\{\hat{0}\}, \hat{0}, \hat{1})$.

Proposition 5.12. Let *P* be a finite, bounded poset with an RFAS Ω . Let \mathcal{M}_{Ω} be as defined in Definition 5.9. Then \mathcal{M}_{Ω} is a partial order and has a unique minimal element given by the first atom chain $c(\{\hat{0}\}, \hat{0}, \hat{1})$.

Proof. To show that \leq is a partial order, it suffices to show that \leq is antisymmetric. We prove this by showing that no sequence of maximal chains $m_1 \to m_2 \to \cdots \to m$ of at least length one has $m_1 = m$. Recall that $c(\{\hat{0}\}, \hat{0}, \hat{1})$ contains no pseudo descents by Proposition 5.8. Thus, $m_i \neq c(\{\hat{0}\}, \hat{0}, \hat{1})$ for any *i* such that 1 < i. Assume that there is some sequence $m_1 \to m_2 \to \cdots \to m$ of at least length one that has $m_1 = m$. Then $c(\{\hat{0}\}, \hat{0}, \hat{1})$ is not one of the chains in this sequence and the sequence can be repeated forever without reaching $c(\{\hat{0}\}, \hat{0}, \hat{1})$. This contradicts Proposition 5.11. Hence, \leq is antisymmetric and thus is a partial order. Moreover, Proposition 5.10 implies that $c(\{\hat{0}\}, \hat{0}, \hat{1}) \preceq m$ for all maximal chains *m* of *P*, so $c(\{\hat{0}\}, \hat{0}, \hat{1})$ is the unique minimal element with respect to \leq .

Now we come to the main result of this section, namely that a finite bounded poset that admits an RFAS is shellable. The proof of this theorem essentially follows the structure of Björner and Wachs' proof of lexicographic shellability in [6].

Theorem 5.13. Suppose finite, bounded poset P admits an RFAS Ω . Let M_{Ω} be as defined in Definition 5.9. Then any linear extension of M_{Ω} gives a shelling of the order complex $\Delta(P)$.

Proof. Let the total order on maximal chains $\Gamma : m_1, \ldots, m_t$ be a linear extension of \preceq . Consider maximal chains m_i and m_j such that i < j. We will find a maximal chain $m_{i'}$ such that i' < j and $m_i \cap m_j \subseteq m_{i'} \cap m_j$ with $m_{i'} \cap m_j = m_j \setminus \{y\}$ for some $y \in m_j$. This then shows that Γ gives a shelling order of $\Delta(P)$.

We first identify all of the intervals in which m_i and m_j disagree. Say that [u, v] is an interval where m_i and m_j differ when u and v are contained in both m_i and m_j but m_i and m_j share no common elements in (u, v). Let s be the number of intervals where m_i and m_j differ. We show by induction on s that m_j contains a pseudo descent in at least one of the intervals where m_i and m_i differ. Suppose s = 1 and the one interval where m_i and m_j disagree is [u, v]. Assume seeking contradiction that m_i does not contain a pseudo descent in the interval [u, v]. Then m_j restricted to [u, v] is the first atom chain of $[u, v]_{m_i^u}$. Then by Remark 5.4 and Proposition 5.12, we have $m_j \prec m_i$. But this contradicts that Γ is a linear extension of \leq since m_i precedes m_j in Γ . Thus, m_j contains a pseudo descent in the interval [u, v]. Now suppose s > 1. Let [u, v] be the first interval on which m_i and m_j differ in the sense that m_i and m_j agree everywhere below u. If m_j contains a pseudo descent in [u, v], then we are done. Otherwise m_j restricted to [u, v] is the first atom chain of $[u, v]_{m_i^u}$. Let d' be the portion of m_i above v. Then again by Remark 5.4 and Proposition 5.12, we have $m_i^v \cup d' \prec m_i$. This implies that $m_i^v \cup d'$ precedes m_i in Γ , so $m_i^v \cup d'$ precedes m_j in Γ . Further, $m_i^v \cup d'$ and m_j disagree in only s-1 intervals and those intervals are among the intervals in which m_i and m_j differ. Thus, by induction m_j has a pseudo descent in an interval where m_i and m_j disagree.

Now let [u, v] be an interval where m_i and m_j differ such that m_j has a pseudo descent in [u, v]. Let x < y < z be this pseudo descent of m_j in [u, v]. Let $c(m_j^x, x, z)$ be the first atom chain of $[x, z]_{m_j^x}$ and let d be the portion of m_j above z. Then again by Remark 5.4 and Proposition 5.12, we have $m_j^x \cup c(m_j^x, x, z) \cup d \prec m_j$. Thus, $m_j^x \cup c(m_j^x, x, z) \cup d = m_{i'}$ for some i' < j since Γ is a linear extension of \preceq . Finally, observe that $m_i \cap m_j \subseteq m_{i'} \cap m_j$ with $m_{i'} \cap m_j = m_j \setminus \{y\}$ by construction. Therefore, Γ gives a shelling of $\Delta(P)$.

Remark 5.14. The proof of Theorem 5.13 also shows that the restriction map of any of these linear extension shellings from an RFAS is given by the pseudo descents of the RFAS. This is the analog of the fact that in a lexicographic shelling, the restriction map is given by the (topological) descents.

6. Labeling-Compatible Recursive First Atom Sets (LCRFAS) and Equivalence to TCL-Shellability

In this section, we add a condition to the definition of an RFAS, producing what we call labeling-compatible recursive first atom sets (LCRFAS) in Definition 6.1. In Theorem 6.5, we show that posets have sets of atoms satisfying this definition if and only if they are TCL-shellable. We also provide in Example 6.7 a poset with an RFAS for which there is no labeling "compatible" with the given RFAS. This shows that in Theorem 6.5, an LCRFAS cannot be replaced with an RFAS.

Definition 6.1. The finite bounded poset P admits a **labeling compatible re**cursive first atom set (LCRFAS) if P admits an RFAS Ω (as in Definition 5.1) and Ω satisfies the following condition on the partial order \leq on the maximal chains of P induced by Ω (as in Definition 5.9).

(LC) There exists a linear extension $\Gamma : m_1, \ldots, m_t$ of \leq such that for maximal chains m_i, m_j , and m_k with i < j < k, whenever m_i and m_k contain $r \cup \{x \leq y \leq z\}$, and m_j contains $r \cup \{x \leq y'\}$, then y' = y where x, y, y', z are in P and r is a root of x in P.

The following lemma is one direction of Theorem 6.5, that is, a poset that is TCL-shellable admits an LCRFAS. The main idea is to let the first atom in a rooted interval be the atom contained in the topologically ascending chain of the rooted interval. This is similar in spirit to the main idea of Björner and Wachs' proof that a poset is CL-shellable if and only if it admits an RAO.

Lemma 6.2. If a finite, bounded poset P admits a TCL-labeling, then P admits an LCRFAS.

Proof. Suppose that the finite, bounded poset P admits a TCL-labeling λ . By Theorem 4.10, P admits a CC-labeling λ' as in Definition 4.1. Note that λ' takes values in the integers by definition. Define Ω as follows: for each rooted interval $[x, y]_r$, set $\Omega(r, x, y) = u$ where u is the atom of [x, y] contained in the unique topologically ascending chain of $[x, y]_r$ with respect to λ' . We show that Ω is an LCRFAS (satisfies Definition 6.1).

Consider any atom a of $[x, y]_r$. Suppose $\Omega(r \cup a, a, y) = b$, that is, b is the atom of $[a, y]_{r \cup a}$ contained in the topologically ascending chain c of $[a, y]_{r \cup a}$ with respect to λ' . First we show that Definition 5.1 (i) holds by considering the following two cases: either a is the first atom of $[x, y]_r$ or a is not the first atom of $[x, y]_r$.

20

21

First suppose that a is the first atom of $[x, y]_r$ according to Ω . Since b is the atom of $[a, y]_{r \cup u}$ contained in the topologically ascending chain c of $[a, y]_{r \cup u}$ with respect to λ' and a is the atom contained in the topologically ascending chain of $[x, y]_r$ with respect to λ' , $a \cup c$ is the topologically ascending chain of $[x, y]_r$. Thus, $x \leq a \leq b$ is a topological ascent with respect to r and λ' , so $a = \Omega(r, x, b)$.

Next suppose that a is not the first atom of $[x, y]_r$ with respect Ω . Observe that $x \cup c$ is not the topologically ascending chain of $[x, y]_r$ because a is in c. Thus, $x \cup c$ must have a topological descent with respect to λ' . Since c is topologically ascending in $[a, y]_{r \cup a}$, x < a < b must be the topological descent in $x \cup c$ with respect to r and λ' . Let d be the topologically ascending chain of $[x, b]_r$ with respect to λ' and let u' be the atom of [x, b] in d. Then $\Omega(r, x, b) = u'$ by definition of Ω and $u' \neq a$.

Next we show that Definition 5.1 (ii) holds. Let us suppose that a is not the first atom of $[x, y]_r$ and a' is the first atom of $[x, y]_r$. Then, as previously observed, $x \le a \le b$ must be a topological descent with respect to r and λ' . Let a_p be the first atom of $[x, b]_r$ and b_{p-1} be the first atom $[a_p, y]_{r \cup a_p}$. If $a_p \ne a'$, then we can repeat the same argument with a_p and b_{p-1} to produce a_{p-1} and b_{p-2} where a_{p-1} is the first atom in $[x, b_{p-1}]_r$ and b_{p-2} is the first atom in $[a_{p-1}, y]_{r \cup a_{p-1}}$. If $a_{p-1} \ne a'$, then we repeat this process to produce a_{p-2} and b_{p-3} , and so on. Notice that for any $i \le p$, the lexicographically first chain in $[x, y]_r$ with respect to λ' containing a_{i-1} and b_{i-2} comes strictly earlier lexicographically than the lexicographically first chain in $[x, y]_r$. Thus, Definition 5.1 (ii) holds and Ω is an RFAS of P.

Lastly we show that condition (LC) of Definition 6.1 holds so that Ω is an LCRFAS. Let \preceq be the partial order on the maximal chains of P induced by Ω as in Definition 5.9. Observe that any total order $\Gamma : m_1, \ldots, m_t$ on the maximal chains of P that is compatible with the lexicographic order induced by λ' is a linear extension of \preceq . Assume seeking contradiction that condition (LC) of Definition 6.1 does not hold. Then there are maximal chains m_i, m_j, m_k in Γ with i < j < k where m_i and m_k contain $r \cup \{x < y < z\}$, while m_j contains $r \cup \{x < y' < z'\}$ where $y \neq y'$. Then by Remark 4.5, $\lambda'(r, x, y) = \lambda'(r, x, y')$ and $\lambda'(r \cup y, y, z) = \lambda'(r \cup y', y', z')$. However, this contradicts Proposition 4.4 (ii) since $y \neq y'$. Therefore, Ω is an LCRFAS.

Now we precisely define what we mean when we say that a CE-labeling is "compatible" with an RFAS. This definition and the following lemma are necessary for the opposite direction of Theorem 6.5. We show in the subsequent, quite technical lemma that a poset that admits an LCRFAS also admits a CE-labeling that is compatible with the LCRFAS. It is then somewhat straightforward to prove that a CE-labeling that is compatible with an RFAS is a TCL-labeling, which we do in the proof of Theorem 6.5.

Definition 6.3. Let P be a finite, bounded poset. Let λ be a CE-labeling on P and let Ω be an RFAS for P. We say λ is **compatible** with Ω if the following holds for all rooted intervals $[x, y]_r$: whenever Ω gives a as the first atom of $[x, y]_r$, a is contained in a maximal chain in $[x, y]_r$ that has the lexicographically smallest label sequence with respect to λ .



FIGURE 8. A schematic illustrating the argument in the proof of Lemma 6.4 in which $m_h, m_i, m_j, m', m_k, m$ is the relative order of the exhibited maximal chains. Chains are denoted with bold text, while elements are denoted with italic text.

Lemma 6.4. Suppose finite, bounded poset P admits an LCRFAS Ω . Then P admits a CE-labeling λ such that λ and Ω are compatible.

Proof. Let \leq be the partial order on the maximal chains induced by Ω as in Definition 5.9 and let Γ be a linear extension of \leq satisfying (LC) of Definition 6.1. Let λ be the CE-labeling of P induced by Γ as in Definition 4.1. We will show that λ is compatible with Ω .

Say that Γ is given by m_1, \ldots, m_t . We show that λ is compatible with Ω by first showing that if, according to Ω , u < v < w is a pseudo descent with respect to the root r, then the label sequence of the first atom chain c(r, u, w) with respect to r and λ strictly lexicographically precedes the label sequence of u < v < w with respect to r and λ . We will then show that λ and Ω are compatible for an arbitrary rooted interval $[x, y]_r$ using Proposition 5.10.

Let $u \leq v \leq w$ be a pseudo descent with respect to the root r and Ω . Let v' be the element of c(r, u, w) covering u, that is, v' is the first atom of $[u, w]_r$. Note that $v' \neq v$. Just as in the proof of Lemma 4.8, Proposition 4.4 (i) implies that $\lambda(r, u, v') \leq \lambda(r, u, v)$. Also, let w' be the element of c(r, u, w) covering v'.

If $\lambda(r, u, v') < \lambda(r, u, v)$, then the label sequence of the first atom chain c(r, u, w)with respect to r and λ strictly lexicographically precedes the label sequence of u < v < w with respect to r and λ . Suppose then that $\lambda(r, u, v') = \lambda(r, u, v)$. It suffices to show that $\lambda(r \cup v', v', w') < \lambda(r \cup v, v, w)$. By the proof of Proposition 4.4 (ii), $\lambda(r \cup v', v', w') \neq \lambda(r \cup v, v, w)$ since $v' \neq v$. Assume seeking contradiction that $\lambda(r \cup v, v, w) < \lambda(r \cup v', v', w')$. We will produce three chains that violate (LC) of Definition 6.1, which contradicts the fact that Ω is an LCRFAS. Fig. 8 shows a schematic of these chains that can be followed for the remainder of the argument. Let m_j be the earliest chain in Γ that contains $r \cup \{v', w'\}$. Observe that since Γ is a linear extension of \preceq and $c(r \cup \{v', w'\}, w', \hat{1})$ is the unique minimal element of \preceq restricted to $[w', \hat{1}]_{r \cup \{v', w'\}}$ (by Proposition 5.12), we have that the portion of m_j above w' is $c(r \cup \{v', w'\}, w', \hat{1})$. Further observe that for the same reasons, any maximal chain containing $r \cup \{v', w'\}$ follows m_j in Γ . We will produce two maximal chains m_i and m_k that both contain $u \lessdot v \lt \tilde{w}$ for some \tilde{w} such that $i \lt j \lt k$.

Recall that by Definition 4.1, $\lambda(r \cup v, v, w) = h$ where h is the position of some maximal chain m_h in Γ that contains $r \cup v$. Notice that the label of the cover relation above v in m_h is h, otherwise $\lambda(r \cup v, v, w)$ would be the position of some earlier chain in Γ than m_h . Observe that since m_j is the first chain in Γ containing $r \cup \{v', w'\}$, Remark 4.2 implies that $\lambda(r \cup v', v', w') \leq j$. Since we have assumed $\lambda(r \cup v, v, w) < \lambda(r \cup v', v', w')$, we have $h = \lambda(r \cup v, v, w) < \lambda(r \cup v', v', w') \leq j$. Thus m_h precedes m_j in Γ . Now let m be the earliest chain in Γ that contains $r \cup \{v, w\}$ and let d be the portion of m above w. Observe that since $r \cup c(r, u, w) \cup d$ contains $r \cup \{v', w'\}$, we have $m_j \leq r \cup c(r, u, w) \cup d$. Then since $r \cup c(r, u, w) \cup d \to m$, we have that m_j precedes m in Γ . Notice that in Γ , m_h is followed by m_j which is followed by m. Furthermore, $r \cup v$ is contained in both m_h and m and the labels of the cover relations above v in both m_h and m are h.

Now let m' be the first maximal chain after m_j in Γ that contains both $r \cup v$ and an element \hat{w} covering v such that $\lambda(r \cup v, v, \hat{w}) = h$. We know m' exists because m satisfies these conditions. We consider two cases: either m' is not the first chain in Γ containing $r \cup \{v, \hat{w}\}$ or m' is the first chain containing $r \cup \{v, \hat{w}\}$. If m' is not the first chain in Γ containing $r \cup \{v, \hat{w}\}$, then the first chain m_i containing $r \cup \{v, \hat{w}\}$ precedes m_j in Γ by choice of m'. Consider m_i, m_j , and $m_k = m'$. We have i < j < k and these three chains violate (LC) of Definition 6.1 since m_i and $m_k = m'$ both contain $r \cup \{v, \hat{w}\}$ while m_j contains $r \cup \{v', w'\}$ and $v \neq v'$.

On the other hand, suppose m' is the first chain in Γ containing $r \cup \{v, \hat{w}\}$. We will find new maximal chains m_i and m_k that together with m_j violate (LC). Since $\lambda(r \cup v, v, \hat{w}) = \lambda(r \cup v, v, w) = h$ and $m' \neq m_h$, Remark 4.3 implies that there is an element \tilde{w} that covers v and satisfies the following properties: there exist maximal chains m_i and m_k , both containing $r \cup \{v, \tilde{w}\}$, such that m_i precedes m' and m' precedes m_k in Γ . This means that $\lambda(r \cup v, v, \tilde{w}) = \lambda(r \cup v, v, w) = h$. By choice of m', we have that m_i precedes m_j in Γ . We also have that m_j precedes m' which precedes m_k . So, we have i < j < k and m_i, m_j, m_k violate (LC) of Definition 6.1 since m_i and m_k both contain $r \cup \{v, \tilde{w}\}$ while m_j contains $r \cup \{v', w'\}$ and $v \neq v'$. Therefore, $\lambda(r \cup v', v', w') < \lambda(r \cup v, v, w)$.

Now we show that λ is compatible with Ω . Suppose m' is a maximal chain in the rooted interval $[x, y]_r$ that is not the first atom chain. By Remark 5.4 and Proposition 5.10, there is a sequence d_1, d_2, \ldots, d_s of maximal chains in $[x, y]_r$ such that $r \cup c(r, x, y) \rightarrow r \cup d_1 \rightarrow r \cup d_2 \rightarrow \cdots \rightarrow r \cup d_s \rightarrow r \cup m'$. Recall from Definition 5.9 that each step in the sequence corresponds to replacing a first atom chain with a single pseudo descent. By the argument above the label sequences strictly increase lexicographically at each step.

We are now ready to prove the main result of the section.

Theorem 6.5. A bounded poset P admits an LCRFAS if and only if P is TCLshellable.

Proof. We begin by proving that if P admits an LCRFAS Ω , then P is TCL-shellable. Let λ be a CE-labeling of P that is compatible (see Definition 6.3) with Ω ; such a labeling exists by Lemma 6.4.

We have that λ is a CE-labeling of P by integers, and no two maximal chains have the same label sequence by Proposition 4.4 (ii). We must show that for each interval $[x, y]_r$, there is a unique, topologically ascending chain and it is lexicographically first.

By the final paragraph of the proof of Lemma 6.4, the first atom chain c(r, x, y) has the lexicographically smallest label sequence with respect to r and λ among all maximal chains of $[x, y]_r$. Thus, c(r, x, y) is topologically ascending with respect to λ . Further, this implies that every pseudo descent with respect to Ω (see Definition 5.6) is a topological descent with respect to λ . Now by Proposition 5.7, any maximal chain m of $[x, y]_r$ that is distinct from c(r, x, y) contains a pseudo descent, and so contains a topological descent with respect to λ . Hence, c(r, x, y) is the unique topologically ascending chain in the rooted interval and has the lexicographically smallest label sequence. Therefore, λ is a TCL-labeling.

Lemma 6.2 is the reverse direction.

Remark 6.6. Observe that in the proof of Theorem 6.5 the only place where condition (LC) of an LCRFAS (Definition 6.1) is used is to produce a labeling that is compatible with the LCRFAS. Once we know there exists a labeling λ that is compatible with the LCRFAS, only the RFAS conditions (Definition 5.1) are necessary to show that λ is a TCL-labeling. In particular, only the RFAS conditions are necessary to show that every pseudo descent is a topological descent with respect to λ . Thus, any labeling that is compatible with an RFAS, in the sense of Definition 6.3, is a TCL-labeling. However, without condition (LC), there may not exist a labeling that is compatible with an RFAS as the following Example 6.7 shows.

Example 6.7. Consider the poset P shown in Fig. 9. There is an RFAS Ω of P for which there does not exist a labeling compatible with Ω . This shows an LCRFAS rather than an RFAS is necessary in Theorem 6.5. Define Ω as follows. In each rooted interval $[x, y]_r$, the first atom $\Omega(r, x, y)$ is the leftmost atom in the interval when P is drawn as in Fig. 9 except for the following: $\Omega(\{\hat{0}\}, \hat{0}, e) = b, \Omega(\{\hat{0}\}, \hat{0}, i) = c, \Omega(\{\hat{0}\}, \hat{0}, n) = b, \Omega(\{\hat{0}, b\}, b, j) = i, \Omega(\{\hat{0}, c\}, c, k) = g, \Omega(\{\hat{0}, c\}, c, \hat{1}) = g, \text{ and } \Omega(r, f, \hat{1}) = k$ for both roots r of f.

While it is somewhat tedious to verify that this is an RFAS, it is fairly straightforward to verify that the partial order \leq on the maximal chains of P induced by Ω as in Definition 5.9 contains the chain $\{\hat{0}, b, d, r, \hat{1}\} \rightarrow \cdots \rightarrow \{\hat{0}, c, i, k, \hat{1}\} \rightarrow \{\hat{0}, c, i, j, \hat{1}\} \rightarrow \{\hat{0}, b, i, j, \hat{1}\} \rightarrow \{\hat{0}, b, d, j, \hat{1}\}$. Observe that Ω violates (LC) of Definition 6.1 since any linear extension of \leq has some chain containing $\{\hat{0}, b, d\}$ coming before a chain containing $\{\hat{0}, c, i\}$ which comes before another chain containing $\{\hat{0}, b, d\}$. Further, by Remark 6.6, any CE-labeling λ that is consistent with Ω is a TCL-labeling. Also, as observed in the proof of Theorem 6.5, every pseudo descent with respect to Ω is a topological descent with respect to λ . Thus, the lexicographic order on maximal chains of P induced by λ is at least as fine as \leq . Then by the proof of Theorem 4.10, there is a CC-labeling λ' of P by integers obtained from λ via Definition 4.1 such that the topological descents with respect to λ' are the



FIGURE 9. A poset that admits an RFAS Ω such that there is no CE-labeling compatible with Ω .

same as the topological descents with respect to λ . Thus, the lexicographic order induced by λ' is also at least as fine as \leq . Now by Remark 4.5, the label sequences of $\{\hat{0}, b, d\}$, $\{\hat{0}, b, i\}$ and $\{\hat{0}, c, i\}$ assigned by λ' are the same. This contradicts that λ' is a CC-labeling since $\{\hat{0}, b, i\}$ and $\{\hat{0}, c, i\}$ are the only maximal chains in the interval $[\hat{0}, i]$, so $[\hat{0}, i]$ does not have a unique topologically ascending maximal chain. (Notice that this also contradicts Proposition 4.4 (ii).) Therefore, there is no CE-labeling that is consistent with Ω . This implies that (LC) of Definition 6.1 is necessary for the existence of a CE-labeling that is compatible with an RFAS.

7. Open Questions

A natural question that arises in light of this work is whether there exist posets that admit RFASs but that do not admit LCRFASs. To this end, one may consider looking at the examples presented in [14] and [16], which are shellable but not CL-shellable and may or may not be CC-shellable. Another question that remains is whether there exists a graded poset that is CC-shellable but not EC-shellable. It is possible that the graded example presented in [12] is one such example, but the method used there for proving this poset is not EL-shellable cannot be easily adapted, at least as far as we can see, to show that the poset is not EC-shellable.

References

 Anders Björner, <u>Shellable and Cohen-Macaulay partially ordered sets</u>, Trans. Amer. Math. Soc. **260** (1980), no. 1, 159–183.

^{2.} _____, <u>Posets, regular CW complexes and Bruhat order</u>, European J. Combin. 5 (1984), no. 1, 7–16.

Anders Björner, László Lovász, and Andrew Yao, <u>Linear decision trees: volume estimates and topological bounds</u>, Proc. 24th ACM Symp. on Theory of Computing (May 1992), ACM Press, New York (1992), 170–177.

Anders Björner and Michelle Wachs, <u>Bruhat order of Coxeter groups and shellability</u>, Adv. in Math. 43 (1982), no. 1, 87–100.

- 5. <u>On lexicographically shellable posets</u>, Trans. Amer. Math. Soc. **277** (1983), no. 1, 323–341.
- <u>_____</u>, <u>Shellable nonpure complexes and posets. I</u>, Trans. Amer. Math. Soc. **348** (1996), no. 4, 1299–1327.
- Patricia Hersh, <u>Lexicographic shellability for balanced complexes</u>, J. Algebraic Combin. 17 (2003), no. 3, 225–254.
- Patricia Hersh and Grace Stadnyk, <u>Generalized recursive atom ordering and equivalence to</u> CL-shellability, Comb. Theory 4 (2024), no. 1, Paper No. 18, 44. MR 4770597
- Dmitry N. Kozlov, <u>General lexicographic shellability and orbit arrangements</u>, Ann. Comb. 1 (1997), no. 1, 67–90.
- 10. Stephen Lacina, Maximal chain descent orders, arXiv: 2209.15142 (2022).
- Tiansi Li, <u>EL-shelling on comodernistic lattices</u>, J. Combin. Theory Ser. A **177** (2021), Paper No. 105334, 8. MR 4151554
- 12. _____, <u>CL-shellable posets with no EL-shellings</u>, Proc. Amer. Math. Soc. **152** (2024), no. 5, 1821–1830. MR 4728454
- 13. John Shareshian, On the shellability of the order complex of the subgroup lattice of a finite group, Transactions of the American Mathematical Society **353** (2001), no. 7, 2689–2703.
- Andrew Vince and Michelle Wachs, <u>A shellable poset that is not lexicographically shellable</u>, Combinatorica 5 (1985), no. 3, 257–260. MR 837069
- Michelle L. Wachs, <u>Poset topology: tools and applications</u>, Geometric combinatorics, IAS/Park City Math. Ser., vol. 13, Amer. Math. Soc., Providence, RI, 2007, pp. 497–615.
- James W. Walker, <u>A poset which is shellable but not lexicographically shellable</u>, European J. Combin. 6 (1985), no. 3, 287–288. MR 818604

Email address: slacina@truman.edu

Email address: grace.stadnyk@furman.edu