Persistent homology of Morse decomposition in Markov chains based on combinatorial multivector fields

Donald Woukeng

Division of Computational Mathematics, Faculty of Mathematics and Computer Science, Jagiellonian University, ul. St. Lojasiewicza 6, Krakow, 30-348, Poland, donald.woukeng@aims.ac.rw

Abstract

In this paper, we introduce a novel persistence framework for Morse decompositions in Markov chains using combinatorial multivector fields. Our approach provides a structured method to analyze recurrence and stability in finite-state stochastic processes. In our setting filtrations are governed by transition probabilities rather than spatial distances. We construct multivector fields directly from Markov transition matrices, treating states and transitions as elements of a directed graph. By applying Morse decomposition to the induced multivector field, we obtain a hierarchical structure of invariant sets that evolve under changes in transition probabilities. This structure naturally defines a persistence diagram, where each Morse set is indexed by its topological and dynamical complexity via homology and Conley index dimensions.

1 Introduction

1.1 Motivation

The study of dynamics has seen a growing interaction between topological methods and discrete combinatorial approaches, especially in understanding recurrence and stability in finitestate models. One of the key frameworks in this domain is Morse decomposition, which provides a hierarchical structure for understanding long-term system behavior. Persistent homology has been instrumental in tracking topological features across scales, but its direct application to Morse decompositions has remained a challenge.

Several works have addressed the persistence of topological invariant for dynamics data or dynamical systems, it is the case for example of [5, 7, 8] where the study of persistence is done for Morse decomposition, Conley index and Conley-Morse graph. However, a key missing component in these works is a stability theorem that ensures that Morse decompositions remain robust under small perturbations in the system's transition dynamics. In particular, no prior work has established a formal stability result for Morse set persistence in Markov chains. Our work addresses this gap by providing a rigorous foundation for the stability of Morse decompositions in combinatorial multivector fields constructed from Markov chains. Markov chains are widely used to model stochastic processes in various real-world domains, including biological networks, financial markets, climate systems, and reinforcement learning. In these applications, understanding long-term behavior and recurrent structures is crucial for predicting trends, identifying stable states, and classifying different system dynamics. Some traditional methods for analyzing Markov chains focus on stationary distributions, spectral properties, and absorbing states [11, 2, 12], but they often lack a topological and hierarchical perspective on the structure of state transitions. Some recent work have been done for the analysis of Markov chains using classical persistence [13, 17, 9], but they do not take into account the potential dynamics hidden in the Markov chains.

Our method introduces a new classification approach for Markov chains by applying persistent homology to Morse decompositions, allowing us to track how recurrent structures change as transition probabilities vary. By constructing persistence diagrams indexed by homology and Conley indices, we provide a robust way to classify Markov chains based on their intrinsic recurrence structures rather than just numerical transition probabilities.

1.2 Overview of the result

In this paper, we develop a persistence framework for Morse decompositions in Markov chains using combinatorial multivector fields. Our main contributions is as follows:

Theorem 1 (Stability of Morse Set Persistence Diagrams). Let P and P' be two transition matrices of a Markov process with n states such that:

$$\|P - P'\|_{\infty} < \delta.$$

Then, the bottleneck distance between their Morse set persistence diagrams satisfies:

$$d_B(D(P), D(P')) < C\delta,\tag{1}$$

where C depends only on the structure of the Markov process.

2 **Preliminaries**

2.1 Relations and Posets

Let X be a set. A *binary relation* on X is defined as a subset $R \subset X \times X$. We use the standard notation xRy to indicate that $(x, y) \in R$.

A relation \leq that is reflexive, antisymmetric, and transitive is called a *partial order*, and the pair (X, \leq) is known as a *poset*. A subset $A \subset X$ of a poset (X, \leq) is called an *upper set* if $\{z \in X \mid \exists_{x \in A} x \leq z\} \subset A$. Similarly, $A \subset X$ is a *lower set* if $\{z \in X \mid \exists_{x \in A} z \leq x\} \subset A$.

A subset $A \subset X$ is said to be *convex* in a poset (X, \leq) if for every $x, z \in A$ and $y \in X$ satisfying $x \leq y \leq z$, it follows that $y \in A$. Equivalently, a convex set is the intersection of a lower set and an upper set.

A relation R is called an *equivalence relation* if it satisfies reflexivity, symmetry, and transitivity. The *equivalence class* of an element x under R is given by $[x]_R := \{y \in X \mid xRy\}$.

A partition of X is a collection \mathcal{V} of non-empty subsets of X such that for every distinct $A, B \in \mathcal{V}$, we have $A \cap B = \emptyset$ and $\bigcup \mathcal{V} = X$. A partition \mathcal{V} induces an equivalence relation R defined by xRy if there exists some $A \in \mathcal{V}$ such that $x, y \in A$. We denote the equivalence class of a point $x \in X$ under partition \mathcal{V} as $[x]_{\mathcal{V}}$.

2.2 **Topological Spaces**

Given a topology \mathcal{T} on X, the pair (X, \mathcal{T}) is referred to as a *topological space*. When the topology \mathcal{T} is understood, we may simply refer to X as a topological space. The interior, closure, and boundary of a subset $A \subset X$ with respect to \mathcal{T} are denoted by $\operatorname{int}_{\mathcal{T}} A$, $\operatorname{cl}_{\mathcal{T}} A$, and $\operatorname{bd}_{\mathcal{T}} A$, respectively. The *mouth* of A is defined as $\operatorname{mo}_{\mathcal{T}} A := \operatorname{cl}_{\mathcal{T}} A \setminus A$. When the topology is clear from context, we may use the shorthand $\operatorname{cl}_X A$ or omit the subscript altogether.

For a subset $Y \subset X$, the induced topology on Y is given by $\mathcal{T}_Y := \{U \cap Y \mid U \in \mathcal{T}\}$. The closure of $A \subset Y$ with respect to \mathcal{T}_Y is denoted as $\operatorname{cl}_{\mathcal{T}_Y} A$ or simply $\operatorname{cl}_Y A$. The same applies to the interior, boundary, and mouth.

A subset $A \subset X$ is called *locally closed* if every point $x \in A$ has a neighborhood U in X such that $A \cap U$ is closed in U.

Proposition 2.1. [10, Problem 2.7.1]. A subset A of a topological space X satisfies the following equivalent conditions:

- (i) A is locally closed,
- (ii) The mouth $\operatorname{mo} A = \operatorname{cl} A \setminus A$ is closed,
- (iii) A can be expressed as the difference of two closed sets in X,
- (iv) A is the intersection of an open set and a closed set in X.

Moreover, the finite intersection of locally closed sets remains locally closed.

A topological space is called a T_0 space if for every pair of distinct points in X, at least one of them has a neighborhood that does not contain the other. We are particularly interested in finite T_0 spaces, which, by Alexandroff's theorem [1], can be associated with posets.

Theorem 2.2. [1] For a finite poset (P, \leq) , the collection of all upper sets of \leq forms a T_0 topology \mathcal{T}_{\leq} on P. Conversely, for any finite T_0 space (X, \mathcal{T}) , defining $x \leq_{\mathcal{T}} y$ by $x \in cl_{\mathcal{T}}\{y\}$ results in a partial order on X. These two correspondences are mutually inverse.

This equivalence allows all topological properties of a finite T_0 space to be expressed in terms of posets and vice versa. In particular, given a finite T_0 space (X, \mathcal{T}) with an associated poset (X, \leq) , the closure and boundary of $A \subset X$ can be described as:

$$\operatorname{cl}_{\mathcal{T}} A = \{ x \in X \mid \exists_{a \in A} x \le a \},$$
(2)

$$\operatorname{bd}_{\mathcal{T}} A = \{ x \in X \mid \exists_{a \in A, b \in X \setminus A} x \le a \text{ and } x \le b \}.$$
(3)

In the context of finite topological spaces, the notions of local closedness and convexity coincide [14, Proposition 1.4.10]. Thus, a subset is locally closed in a T_0 topology if and only if it is convex in the corresponding poset structure. Throughout this paper, we use these terms interchangeably, favoring "locally closed" in topological discussions and "convex" in combinatorial or algorithmic contexts.

2.3 Combinatorial multivector fields

All of the definitions in this subsection can be found in [15].

Let X be a finite topological space. A combinatorial multivector or briefly a multivector is a locally closed subset $V \subset X$. A combinatorial multivector field on X, or briefly a multivector field, is a partition \mathcal{V} of X into multivectors.

Since \mathcal{V} is a partition, we can denote by $[x]_{\mathcal{V}}$ the unique multivector in \mathcal{V} that contains $x \in X$. If the multivector field \mathcal{V} is clear from the context, we write briefly [x]. We say that a multivector $V \in \mathcal{V}$ is *critical* if the relative singular homology $H(\operatorname{cl} V, \operatorname{mo} V)$ is non-trivial. A multivector V which is not critical is called *regular*. We say that a set $A \subset X$ is \mathcal{V} -compatible if for every $x \in X$ either $[x] \cap A = \emptyset$ or $[x] \subset A$.

Multivector field \mathcal{V} on X induces a multivalued map $\Pi_{\mathcal{V}}: X \multimap X$ given by

$$\Pi_{\mathcal{V}}(x) = [x]_{\mathcal{V}} \cup \operatorname{cl} x . \tag{4}$$

We consider a combinatorial dynamical system given by the iterates of $\Pi_{\mathcal{V}}$.

A solution of a combinatorial dynamical system $\Pi_{\mathcal{V}} : X \multimap X$ in $A \subset X$ is a partial map $\varphi : \mathbb{Z} \nrightarrow A$ whose domain, denoted dom φ , is a \mathbb{Z} -interval and for any $i, i + 1 \in \operatorname{dom} \varphi$ the inclusion $\varphi(i + 1) \in \Pi_{\mathcal{V}}(\varphi(i))$ holds. Let us denote by $\operatorname{Sol}(A)$ the set of all solutions φ such that im $\varphi \subset A$. $\operatorname{Sol}(X)$ is the set of all solution of $\Pi_{\mathcal{V}}$. If dom φ is a bounded interval then we say that φ is a *path*. If dom $\varphi = \mathbb{Z}$ then φ is a *full solution*.

A full solution $\varphi : \mathbb{Z} \to X$ is *left-essential* (respectively *right-essential*) if for every regular $x \in \operatorname{im} \varphi$ the set $\{t \in \mathbb{Z} \mid \varphi(t) \notin [x]_{\mathcal{V}}\}$ is left-infinite (respectively right-infinite). We say that φ is *essential* if it is both left- and right-essential. The collection of all essential solutions φ such that $\operatorname{im} \varphi \subset A$ is denoted by $\operatorname{eSol}(A)$.

The *invariant* part of a set $A \subset X$ is $\operatorname{Inv} A := \bigcup \{ \operatorname{im} \varphi \mid \varphi \in \operatorname{eSol}(A) \}$. In particular, if $\operatorname{Inv} A = A$ we say that A is an *invariant set* for a multivector field \mathcal{V} .

A closed set $N \subset X$ isolates invariant set $S \subset N$ if the following conditions are satisfied:

- (i) every path in N with endpoints in S is a path in S,
- (ii) $\Pi_{\mathcal{V}}(S) \subset N$.

In this case, N is an *isolating set* for S. If an invariant set S admits an isolating set then we say that S is an *isolated invariant set*. The *homological Conley index* of an isolated invariant set S is defined as Con(S) := H(cl S, mo S).

Let $A \subset X$. By $\langle A \rangle_{\mathcal{V}}$ we denote the intersection of all locally closed and \mathcal{V} -compatible sets in X containing A. We call this set the \mathcal{V} -hull of A. The combinatorial α -limit set and ω -limit set for a full solution φ are defined as

$$\begin{aligned} \alpha(\varphi) &:= \left\langle \bigcap_{t \in \mathbb{Z}^-} \varphi((-\infty, t]) \right\rangle_{\mathcal{V}}, \\ \omega(\varphi) &:= \left\langle \bigcap_{t \in \mathbb{Z}^+} \varphi([t, \infty)) \right\rangle_{\mathcal{V}}. \end{aligned}$$

Let $S \subset X$ be a \mathcal{V} -compatible, invariant set. Then, a finite collection $\mathcal{M} = \{M_p \subset S \mid p \in \mathbb{P}\}$ is called a *Morse decomposition* of S if there exists a finite poset (\mathbb{P}, \leq) such that the following conditions are satisfied:

- (i) \mathcal{M} is a family of mutually disjoint, isolated invariant subsets of S,
- (ii) for every $\varphi \in \operatorname{eSol}(S)$ either $\operatorname{im} \varphi \subset M_r$ for an $r \in \mathbb{P}$ or there exist $p, q \in \mathbb{P}$ such that $q > p, \alpha(\varphi) \subset M_q$, and $\omega(\varphi) \subset M_p$.

We refer to the elements of \mathcal{M} as *Morse sets*.

Let \mathcal{V} and \mathcal{V}' be two multivector fields on the same underlying space. We say that \mathcal{V} is a **coarsening** of \mathcal{V}' (see [6], denoted $\mathcal{V}' \leq \mathcal{V}$, if every multivector in \mathcal{V} is a union of one or more multivectors in \mathcal{V}' . Formally,

$$\forall V \in \mathcal{V}, \quad \exists \{W_i\} \subseteq \mathcal{V}' \text{ such that } V = \bigcup_i W_i.$$

Coarsening corresponds to merging finer multivectors into larger ones, effectively reducing the resolution of the multivector field.

2.4 Markov Chains

The definitons on this subsection can be found in [16, Chapter 2]

2.4.1 Definition of a Markov Chain

A discrete-time Markov chain is a stochastic process $\{X(n)\}_{n \in \mathbb{N}_0}$ taking values in a finite or countable state space S. The process satisfies the Markov property, meaning that the probability of transitioning to the next state depends only on the present state and not on the past history:

$$P(X(n+1) = j \mid X(n) = i, X(n-1) = i_{n-1}, \dots, X(0) = i_0) = P_{i,j},$$
(5)

for all $i, j, i_{n-1}, \ldots, i_0 \in S$ and $n \in \mathbb{N}_0$.

A Markov chain is called *time-homogeneous* if the transition probabilities do not depend on n, i.e.,

$$P(X(n+1) = j \mid X(n) = i) = P_{i,j}.$$
(6)

The dynamics of a discrete-time Markov chain on a finite state space is fully characterized by its *transition matrix* $P = (P_{i,j})$, where each entry $P_{i,j}$ represents the probability of transitioning from state *i* to state *j*:

$$P = \begin{bmatrix} P_{1,1} & P_{1,2} & \dots & P_{1,N} \\ P_{2,1} & P_{2,2} & \dots & P_{2,N} \\ \vdots & \vdots & \ddots & \vdots \\ P_{N,1} & P_{N,2} & \dots & P_{N,N} \end{bmatrix}.$$
(7)

Each row of P sums to 1:

$$\sum_{j \in S} P_{i,j} = 1, \quad \forall i \in S.$$
(8)

The probability of the system being in a specific state at step n is given by the vector $\mathbf{p}(n)$, which evolves according to the equation:

$$\mathbf{p}(n) = \mathbf{p}(0)P^n. \tag{9}$$

where $\mathbf{p}(0)$ is the initial probability distribution over states.

2.4.2 Graph Representation of a Markov Chain

A Markov chain can be represented as a *directed graph*, called a *transition diagram*. In this representation:

- The nodes (vertices) correspond to the states of the Markov chain. - A directed edge $i \rightarrow j$ exists if $P_{i,j} > 0$, meaning there is a positive probability of transitioning from state i to state j. - The edges are labeled by the corresponding transition probabilities $P_{i,j}$.

For example, the transition diagram of a three-state Markov chain with the transition matrix:

$$P = \begin{bmatrix} 1/3 & 1/3 & 1/3 \\ 1/2 & 1/2 & 0 \\ 1 & 0 & 0 \end{bmatrix}$$
(10)

is represented as:



where the directed edges indicate possible transitions and their associated probabilities.

3 Morse decomposition for Markov chains

We introduce the construction of *multivector fields* from Markov chains, explore the notion of *coarsening*, and establish the *Morse decomposition* of a multivector field. These concepts provide a structured way to analyze the dynamics of state transitions within a Markov process and extract meaningful topological features from the induced flow structure.

3.1 Multivector field construction

A *multivector field* is constructed by merging *vertices and edges* based on transition probabilities:

- Each state in the Markov chain is represented as a vertex.
- Each *directed edge* represents a transition between states.
- Edges and vertices are merged into multivectors if the transition probability is less than or equal a chosen threshold γ .
- Any element (vertex or edge) that is not merged remains a singleton multivector.

This construction ensures that *all vertices and edges are included* in the multivector field, forming a *directed partition of the state space and the edges that have non-zero transition probability*. The process is *iterative*, merging multivectors when they *overlap*, ensuring a well-defined decomposition.

A formal *algorithm* is presented here, defining the precise steps to construct the multivector field. We will follow a method similar to the one use in [18, 4].

Algorithm 1 Construction of the Multivector Field from a Markov Chain

Require: Transition matrix P, threshold γ **Ensure:** Multivector field V1: Initialize $V \leftarrow \emptyset$ ▷ Start with an empty set of multivectors 2: Define S as the set of all states (nodes) in the Markov chain 3: Define E as the set of all directed edges (N_i, N_j) where $P(N_i \rightarrow N_j) > 0$ ▷ Step 1: Initialize all vertices and edges as separate multivectors 4: for all $N_i \in S$ do $V \leftarrow V \cup \{\{N_i\}\}$ 5: 6: **end for** 7: for all $(N_i, N_j) \in E$ with i < j do $V \leftarrow V \cup \{\{(N_i, N_j)\}\}$ > Avoid duplicate bidirectional edges 8: 9: end for \triangleright Step 2: Merge vertices with edges if transition probability is below γ 10: for all $(N_i, N_j) \in E$ with i < j do if $P(N_i \to N_j) \leq \gamma$ then 11: Merge N_i and the edge (N_i, N_j) into one multivector 12: $V \leftarrow V \setminus \{\{N_i\}, \{(N_i, N_j)\}\} \cup \{\{N_i, (N_i, N_j)\}\}$ 13: end if 14: if $P(N_i \to N_i) \leq \gamma$ then 15: Merge N_i and the edge (N_i, N_j) into one multivector 16: $V \leftarrow V \setminus \{\{N_i\}, \{(N_i, N_j)\}\} \cup \{\{N_i, (N_i, N_j)\}\}$ 17: end if 18: 19: end for ▷ Step 3: Merge overlapping multivectors to maintain the partition 20: for all pairs $V_k, V_m \in V$ do if $V_k \cap V_m \neq \emptyset$ then 21: Merge V_k and V_m into a single multivector 22: $V \leftarrow V \setminus \{V_k, V_m\} \cup \{V_k \cup V_m\}$ 23: end if 24: 25: end for 26: return V ▷ Return the Multivector field V

Theorem 2. Let P be the transition matrix of a finite Markov process with state space S, and let $\mathcal{V}\gamma$ be the multivector field constructed using the threshold γ . Then:

- The algorithm always produces a valid multivector field $V\gamma$, where each state and transition appears exactly once.
- The sequence of multivector fields $\mathcal{V}\gamma_1, \mathcal{V}\gamma_2, \dots$ forms a decreasing filtration:

$$\mathcal{V}\gamma_1 \preceq \mathcal{V}\gamma_2 \preceq \dots$$

for $\gamma_1 < \gamma_2$, meaning that as γ increases.

Proof:

• We consider our finite space being the state space of our Markov process plus edges between states, if the probability of going from one state to another is non-zero. We will denote that space X.

$$X = \bigcup \{N_i\} \cup \{(N_i, N_j)\} \text{ if } P(N_i \to N_j) > 0 \text{ or } P(N_j \to N_i) > 0, \ i < j$$

In Algorithm 1, from line 4 to 9 by adding all the elements of our space X, vertices representing states, and edges representing connection between them with non-zero transition probability. From line 10 to 19, we merge edges and vertices of our space according to the fix given threshold γ , and the last part from line 20 to 24, we ensure that what we created is partition, by merging together elements that have non empty intersection with other elements. We then have a partition of our space X, and since we only deal with edges and vertices, we are sure to always have the local closedness property. Since V returned at the line 26 is a partition of our space X, into locally closed subset, V is then a multivector field.

• The second part of the theorem follows naturally from the construction of the multivector field, since it is done by merging edge to vertices for a certain threshold. If we have a bigger threshold, it is only natural that the multivectors will contain at least all the element contains, for a lower threshold, and maybe we will have multivector that are just union of multivectors in a for a lower threshold.

Example 1. Consider a Markov process with three states $S = \{N_1, N_2, N_3\}$ and the transition *matrix:*

$$P = \begin{bmatrix} 0.5 & 0.17 & 0.33\\ 0.17 & 0.6 & 0.23\\ 0.15 & 0.15 & 0.7 \end{bmatrix}$$

We choose a threshold $\gamma = 0.2$ for merging.

1. Step 1: Initialize Multivectors

Each state starts as its own singleton:

$$V1 = \{N_1\}, \quad V2 = \{N_2\}, \quad V3 = \{N_3\},$$
$$V_4 = \{(N_1, N_2)\}, \quad V_5 = \{(N_1, N_3)\}, \quad V_6 = \{(N_2, N_3)\}.$$

2. Step 2: Merge Based on Transition Probabilities

- Since $P_{12} = 0.17 \leq \gamma$, we merge V_4 into V_1 . Since $P_{31} = 0.15 \leq \gamma$, we merge V_5 into V_3 . Since $P_{32} = 0.15 \leq \gamma$, we merge V_6 into V_3 . - Since $P_{21} = 0.17 \leq \gamma$, we merge V_4 into V_2 .

The new candidates are:

$$V_1^* = \{N_1, (N_1, N_2)\}, \quad V_2^* = \{N_2, (N_1, N_2)\}, \quad V_3^* = \{N_3, (N_2, N_3), (N_1, N_3)\}.$$

3. Step 3: Partition and Merge Overlapping Multivectors Since (N_1, N_2) appears in V_1^* and V_2^* , we merge them:

$$V_3^* = \{N_3, (N_1, N_3), (N_2, N_3)\}, \quad V_1^* = \{N_1, N_2, (N_1, N_2)\}.$$

4. Step 4: Final Multivector Field The final set of multivectors is:

 $\mathcal{V} = \{V_1^*, V_3^*\}.$

3.2 Morse Decomposition in Multivector Fields

Once the multivector field is constructed, we analyze its *global structure* through Morse decompositions. A *Morse set* is defined as a *strongly connected component* (see[15]) in the *graph induced by the multivector field (M-graph)*. These Morse sets represent *regions of recurrence and stability* in the system.

- The Morse sets are computed from the *M*-graph, where each multivector is a node, and edges exist based on the mouth condition (a multivector *M* transitions to another *M'* if *M'* contains an element in *M*'s mouth).
- The resulting Morse decomposition provides a *hierarchical structure* of stability regions.
- Morse sets can be *persistent* under coarsening, allowing us to track their evolution through different scales.

A key property of this approach is that Morse decompositions naturally arise from the multivector field and do not require predefined distance metrics, making them applicable to discrete-state systems such as Markov chains.

A Morse set M persists as long as it remains an SCC in G_{γ} .

Remark 1 (Persistence of Morse Sets). A Morse set M has a **birth threshold** γ_b where it first appears and a **death threshold** γ_d , where:

- 1. *M* merges into another Morse set at γ_d .
- 2. The index of M (given by the dimension of its Conley index or its homology) changes at γ_d .

Thus, the lifespan of a Morse set is defined as:

Persistence $(M) = \gamma_d - \gamma_b$.

Remark 2. The change in Conley index and homology signals a **topological transition** in the Morse set structure. This can indicate a loss or gain of significant recurrent dynamics in the Markov process.

Theorem 3. Let $\mathcal{V}\gamma_1$ and $\mathcal{V}\gamma_2$ be two multivector fields such that $\gamma_1 < \gamma_2$. Then for any Morse set $M_1 \in \mathcal{M}(\mathcal{V}\gamma_1)$, there exists a Morse set $M_2 \in \mathcal{M}(\mathcal{V}\gamma_2)$ such that:

$$M_1 \subset M_2$$

Proof: Since $\gamma_1 < \gamma_2$, the construction of $\mathcal{V}\gamma_2$ involves additional merging compared to $\mathcal{V}\gamma_1$. From that we are sure that G_{γ_1} is actually a subgraph of G_{γ_2} and then Any strongly connected component (SCC) in G_{γ_1} remains connected in G_{γ_2} or merges into a larger SCC.

Thus, the Morse sets at γ_2 is a union of Morse sets at γ_1 , establishing the filtration property.

Remark 3. This result shows that Morse sets **never split** as γ increases. Instead, they follow a strict merging hierarchy, revealing the most **persistent structures** in the system.

4 Stability of Persistence diagrams for Morse decomposition

A fundamental property of any meaningful topological or combinatorial structure in applied mathematics is *stability*: small perturbations in the input data should not lead to large variations in the extracted features. In the context of multivector fields constructed from Markov chains, stability ensures that Morse decompositions remain robust under perturbations in the transition probabilities.

The goal here is to establish a formal stability result for Morse set persistence in multivector fields. Specifically, we consider how small perturbations in the transition matrix of a Markov process affect the birth and death times of Morse sets in the associated persistence diagram. To quantify this effect, we introduce an appropriate bottleneck distance that measures the similarity between Morse set persistence diagrams obtained from different Markov chains. But before that we need to define the persistence diagram in our setting since it is a bit different from the natural one (see [3]).

4.1 Persistence Diagram for Morse Sets

Definition 1. Let $\mathcal{M} = \{M_i\}$ be the set of Morse sets obtained from the **M-graph** of a multivector field at different values of γ . Each Morse set M_i is characterized by:

- A birth threshold γ_b , where it first appears.
- A death threshold γ_d, where it either merges with another Morse set or undergoes an index change.
- A dimensional topological index k_i , defined as the pair:

$$k_i = \left(\dim H_1(\overline{M_i}), \dim \operatorname{Ind}_1(M_i)\right), \tag{11}$$

where:

- dim $H_1(\overline{M_i})$ is the dimension of the first homology group of the closure $\overline{M_i}$ of the Morse set.
- dim $Ind_1(M_i)$ is the dimension of the first relative homology group of the closure of M_i to its mouth.

The **persistence diagram** consists of a set of points $(\gamma_b, \gamma_d, k_i)$ in the **birth-death plane**, where the index k_i encodes the **topological complexity** of the Morse set across different thresholds.

Remark 4. k_i is not really a part of the coordinate for the persistence of our Morse set M_i in our persistence diagram. It just give you informations about the nature if the Morse set M_i .

4.2 Bottleneck Distance for Morse Set Persistence Diagrams

To compare the persistence diagrams of Morse sets obtained from different Markov chains, we define a bottleneck distance that captures differences in their birth and death times, ensuring that the measure remains independent of any fixed threshold γ .

4.2.1 Preliminaries and Notation

Let P and P' be two transition matrices representing two Markov chains on the same state space:

$$P = (p_{ij})_{1 \le i,j \le n}, \quad P' = (p'_{ij})_{1 \le i,j \le n}.$$

Each Markov chain induces a multivector field $\mathcal{V}\gamma(P)$ and $\mathcal{V}\gamma(P')$ at each threshold γ , leading to the construction of an M-graph and the computation of Morse sets. The persistence diagram tracks the birth and death of each Morse set as γ varies.

Let:

- $D(P) = \{(\gamma_b^i, \gamma_d^i, k_i)\}$ be the persistence diagram of Morse sets obtained from P.
- $D(P') = \{(\gamma_b'^{j}, \gamma_d'^{j}, k_j')\}$ be the persistence diagram obtained from P'.

Each point in D(P) and D(P') represents a Morse set M_i with:

- Birth time γ_b^i : the smallest threshold γ at which M_i appears.
- **Death time** γ_d^i : the largest threshold γ at which M_i exists before merging or disappearing.
- Topological index $k_i = (\dim H_1(\overline{M_i}), \dim \operatorname{Ind}_1(M_i))$, encoding the first homology and Conley index dimensions.

4.2.2 Definition of the Bottleneck Distance

The **bottleneck distance** between D(P) and D(P') is given by:

$$d_B(D(P), D(P')) = \inf_{\varphi} \sup_i \|\mathbf{p}_i - \mathbf{p}'_{\varphi(i)}\|_{\infty},$$
(12)

where:

• φ is a bijection matching points only if $k_i = k'_{\varphi(i)}$.

- $\mathbf{p}_i = (\gamma_b^i, \gamma_d^i)$ and $\mathbf{p}'_{\varphi(i)} = (\gamma_b'^{\varphi(i)}, \gamma_d'^{\varphi(i)})$ are the birth-death coordinates of matched Morse sets.
- The distance between matched pairs is computed in the ℓ_{∞} norm:

$$\|\mathbf{p}_i - \mathbf{p}'_{\varphi(i)}\|_{\infty} = \max\left(|\gamma_b^i - \gamma_b'^{\varphi(i)}|, |\gamma_d^i - \gamma_d'^{\varphi(i)}|\right).$$
(13)

If a Morse set has no valid match in the other persistence diagram, it is assigned to the diagonal (γ_b = γ_d), representing a zero-persistence feature.

Unlike classical persistence in topological data analysis, where stability is typically analyzed with respect to an underlying metric space, our setting is purely combinatorial: the Markov chain defines a weighted graph, and persistence is extracted through a filtration on the multivector field induced by transition probabilities.

4.2.3 The Stability Theorem

Theorem 4 (Stability of Morse Set Persistence Diagrams). Let P and P' be two transition matrices of a Markov process with n states such that:

$$\|P - P'\|_{\infty} < \delta.$$

Then, the bottleneck distance between their Morse set persistence diagrams satisfies:

$$d_B(D(P), D(P')) < C\delta, \tag{14}$$

where C depends only on the structure of the Markov process.

Before proving the theorem, we establish a key intermediate result. Since a global perturbation δ in P may affect multiple transitions simultaneously, we first consider the effect of a single-entry perturbation in P, which will allow us to extend the result to the full transition matrix.

Lemma 1 (Local Stability Under Single Transition Change). Let P and P' be two transition matrices that differ at only one entry (i, j):

 $p'_{ii} = p_{ij} + \delta$, where $|\delta| < \epsilon$, and $p_{kl} = p'_{kl}$ for all $(k, l) \neq (i, j)$.

Then, the bottleneck distance between their Morse set persistence diagrams satisfies:

$$d_B(D(P), D(P')) < \epsilon.$$
(15)

Proof: Since only one transition probability p_{ij} is modified, the global structure of the multivector field remains unchanged except in one location:

- The modification of p_{ij} affects only the relationship between states N_i and N_j .
- The change in the multivector field may result in:
 - A Morse set being born earlier or later.
 - A Morse set disappearing earlier or later.
 - A merging of multiple Morse sets into a single one.

• However, in all cases, each affected Morse set undergoes a shift in birth or death time by at most $|\delta|$.

Thus, for any affected Morse set M, we have:

$$\gamma_b(M) - \gamma_b(M')| \le |\delta|, \quad |\gamma_d(M) - \gamma_d(M')| \le |\delta|.$$

Now, we define a bijection φ between Morse sets in D(P) and D(P'):

- For all unchanged Morse sets, we match them with their identical counterparts.
- For affected Morse sets, we match each of them to the corresponding Morse set in D(P'), whether it remains the same or has merged into a new Morse set.

Since the maximum shift in birth or death time is at most $|\delta|$, we obtain:

$$\|\mathbf{p}_i - \mathbf{p}'_{\varphi(i)}\|_{\infty} = \max\left(|\gamma_b(M) - \gamma_b(M')|, |\gamma_d(M) - \gamma_d(M')|\right) \le |\delta|$$

By the definition of the bottleneck distance, which takes the infimum over all bijections, we conclude:

$$d_B(D(P), D(P')) \le \sup_M \max\left(|\gamma_b(M) - \gamma_b(M')|, |\gamma_d(M) - \gamma_d(M')|\right) \le |\delta|.$$

and

$$d_B(D(P), D(P')) < \epsilon$$

4.2.4 **Proof of the sability theorem**

The proof follows by decomposing the perturbation into single-entry modifications in the transition matrix and applying the local stability lemma iteratively:

• Step 1: Decomposition of Perturbation We write the transition matrix perturbation as a sequence of stepwise changes:

$$P = P_0 \to P_1 \to P_2 \to \cdots \to P_l = P',$$

where each P_k differs from P_{k-1} in at most one transition probability.

Step 2: Application of the Local Stability Lemma From the local stability lemma, we know that if *P* and *P'* differ by a single entry change δ, then:

$$d_B(D(P_k), D(P_{k+1})) \le |\delta_k|.$$

Summing over all steps, we get:

$$d_B(D(P), D(P')) \le \sum_{k=0}^{l-1} d_B(D(P_k), D(P_{k+1})) \le l \times \max_k(\delta_k).$$

Since $||P - P'||_{\infty} < \delta$, we have:

$$d_B(D(P), D(P')) < l\delta$$

l being the number of non-zero entries of the matrix P - P', with $l \le n^2$

5 Example of computation

Here we will show how the compution of the persistence diagram for this setting looks like. We will continue with the Example we used before(Example 1).

We consider a Markov process with three states $S = \{N_1, N_2, N_3\}$ and the transition matrix:

$$P = \begin{bmatrix} 0.5 & 0.17 & 0.33\\ 0.17 & 0.6 & 0.23\\ 0.15 & 0.15 & 0.7 \end{bmatrix}.$$

Here the different values of gamma we are interested in are: $\{0.15, 0.17, 0.23, 0.33\}$.

Note that we are not interested by value at the diagonal since while creating the M-graph, we always have a self loop for each multivector.

5.1 Multivector field construction

As we have already shown how we construct for $\gamma = 0.2$ before, I am only going to give the final construction for each γ .

5.1.1 For $\gamma < 0.15$, γ_0

We have:

$$\mathcal{V} = \{\{N_1\}, \{N_2\}, \{N_3\}, \{(N_1, N_2)\}, \{(N_1, N_3)\}, \{(N_2, N_3)\}\}$$

Here we only have singletons since we did not cross any of the transition probabilities in the transition matrix yet

5.1.2 For $\gamma = 0.15$, γ_1

We have:

 $\mathcal{V} = \{\{N_1\}, \{N_2\}, \{N_3, (N_1, N_3), (N_2, N_3)\}, \{(N_1, N_2)\}$

5.1.3 For $\gamma = 0.17, \gamma_2$

We have:

$$\mathcal{V} = \{\{N_1, N_2, (N_1, N_2)\}, \{N_3, (N_1, N_3), (N_2, N_3)\}\}$$

5.1.4 For $\gamma = 0.23$, γ_3

We have:

$$\mathcal{V} = \{\{N_1, N_2, (N_1, N_2), N_3, (N_1, N_3), (N_2, N_3)\}\}$$

For $\gamma \ge 0.23$ which is the greastest element not in the diagonal, our multivector field will only always consist in one big multivector containing the whole space.

5.2 Morse set computations

Here we will also compute Morse set from the multivector field for each value of γ and their index, which is just computing each time strongly connected component of the M-graph.

5.2.1 For $\gamma < 0.15$

•

•

Since we said that in the M-graph each multivector have self loop, they are then also strongly connected and then we have as Morse set for this value of γ :

$$M_1^0, M_2^0, M_3^0 = \{N_1\}, \{N_2\}, \{N_3\}$$

all of them have the same index so:

$$k_1^0 = k_2^0 = k_3^0 = (0,0)$$

, since the dimensions of the first homology and relative homology of a point are 0.

 $M_4^0, M_5^0, M_6^0 = \{(N_1, N_2)\}, \{(N_1, N_3)\}, \{(N_2, N_3)\}$

all of them have the same index so:

$$k_4^0 = k_5^0 = k_6^0 = (0, 1)$$

, since the dimensions of the first homology of an edge is zero, but the first relative homology of an edge to its two vertices is actually the same as the homology of a circle.

- **5.2.2** For $\gamma = 0.15$
 - M_1^1 , $M_2^1 = \{N_1\}$, $\{N_2\}$ and $k_1^1 = k_2^1 = (0,0)$
 - $M_3^1, M_4^1 = \{(N_1, N_2)\}, \ \{N_3, (N_1, N_3), (N_2, N_3)\} \text{ and } k_3^1 = k_4^1 = (0, 1)$

5.2.3 For $\gamma = 0.17$

- $M_1^2 = \{N_1, N_2, (N_1, N_2)\}$ and $k_1^2 = (0, 0)$
- $M_2^2 = \{N_3, (N_1, N_3), (N_2, N_3)\}$ and $k_2^2 = (0, 1)$

5.2.4 For $\gamma = 0.23$

• $M_1^3 = \{N_1, N_2, (N_1, N_2), N_3, (N_1, N_3), (N_2, N_3)\}$ and $k_1^3 = (1, 1)$

Note that the Morse set remains unchanged for greater values of γ . In general Morse sets are union of multivectors, here we just have a special case where Morse sets correspond to multivectors for the sake of simplicity.

5.3 Persistence diagram computation

From the computations of Morse sets we have:

- M₁⁰ ⊂ M₁¹ ⊂ M₁² ⊂ M₁³ M₁⁰ dies at γ₃ because of the change in index.
- $M_2^0 \subset M_2^1 \subset M_1^2 \subset M_1^3 \ M_2^0$ dies at γ_2 because it merges with M_1^0



Figure 1: Persistence diagram of the Morse sets for P

- $M_3^0 \subset M_4^1 \subset M_2^2 \subset M_1^3$ M_3^0 dies at γ_1 because there is a change of index.
- $M_4^0 \subset M_2^1 \subset M_1^2 \subset M_1^3$; M_4^0 dies at γ_2 because it merges with M_1^0
- $M_5^0 \subset M_4^1 \subset M_2^2 \subset M_1^3$ M_5^0 dies at γ_3 because there is a change of index.
- $M_6^0 \subset M_4^1 \subset M_2^2 \subset M_1^3$ M_6^0 dies at γ_1 because it merges with M_5^0 .
- M_1^3 is born at γ_3 and never dies.

The persistence diagram for the transition Matrix P will have the points: $p_1 = (0, 0.23, (0, 0)), p_2 = (0, 0.17, (0, 0)), p_3 = (0, 0.15, (0, 0)), p_4 = (0, 0.17, (0, 1)), p_5 = (0, 0.23, (0, 1)), p_6 = (0, 0.15, (0, 1)), p_7 = (0.23, \infty, (1, 1))$

Competing interests

Research of D.W. is partially supported by the Polish National Science Center under Opus Grant No. 2019/35/B/ST1/00874.

References

- [1] P. Alexandroff. Diskrete Räume. Rec. Math. Moscou, n. Ser., 2:501–519, 1937.
- [2] Iddo Ben-Ari and Ningwei Jiang. Representation and characterization of quasistationary distributions for markov chains. *arXiv preprint arXiv:2402.11154*, 2024.
- [3] David Cohen-Steiner, Herbert Edelsbrunner, and John Harer. Stability of persistence diagrams. In *Proceedings of the twenty-first annual symposium on Computational geometry*, pages 263–271, 2005.
- [4] Dominic Desjardins Côté and Donald Woukeng. From data to combinatorial multivector field through an optimization-based framework. *arXiv preprint arXiv:2501.02023*, 2025.
- [5] Tamal Dey, Mateusz Juda, Tomasz Kapela, Jacek Kubica, Michał Lipiński, and Marian Mrozek. Persistent homology of Morse decompositions in combinatorial dynamics. *SIAM Journal on Applied Dynamical Systems*, 18:510–530, January 2018.
- [6] Tamal K Dey, Michał Lipiński, Marian Mrozek, and Ryan Slechta. Tracking dynamical features via continuation and persistence. *arXiv preprint arXiv:2203.05727*, 2022.
- [7] Tamal K Dey, Marian Mrozek, and Ryan Slechta. Persistence of the conley index in combinatorial dynamical systems. *arXiv preprint arXiv:2003.05579*, 2020.
- [8] Tamal K Dey, Marian Mrozek, and Ryan Slechta. Persistence of conley-morse graphs in combinatorial dynamical systems. SIAM Journal on Applied Dynamical Systems, 21(2):817–839, 2022.
- [9] Irene Donato, Giovanni Petri, Martina Scolamiero, Lamberto Rondoni, and Francesco Vaccarino. Decimation of fast states and weak nodes: topological variation via persistent homology. In *Proceedings of the European Conference on Complex Systems 2012*, pages 295–301. Springer, 2013.
- [10] R. Engelking. General Topology. Heldermann Verlag, Berlin, 1989.
- [11] Roberto Fernandez, Francesco Manzo, Matteo Quattropani, and Elisabetta Scoppola. Quasi-stationary distributions of non-absorbing markov chains. *Journal of Statistical Physics*, 192(3):43, 2025.
- [12] Lixing Han and Jianhong Xu. On classification of states in higher order markov chains. *Linear Algebra and its Applications*, 685:24–45, 2024.
- [13] Minh Quang Le and Dane Taylor. Persistent homology of convection cycles in network flows. *Physical Review E*, 105(4):044311, 2022.
- [14] Michał Lipiński. Morse-Conley-Forman theory for generalized combinatorial multivector fields on finite topological spaces. PhD thesis, Jagiellonian University, 2021.
- [15] Michał Lipiński, Jacek Kubica, Marian Mrozek, and Thomas Wanner. Conley-Morse-Forman theory for generalized combinatorial multivector fields on finite topological spaces. *Journal of Applied and Computational Topology*, 2022.
- [16] Anders Tolver. An introduction to markov chains. *Department of Mathematical Sciences, University of Copenhagen*, 2016.

- [17] Sarah Tymochko, Kritika Singhal, and Giseon Heo. Classifying sleep states using persistent homology and markov chains: a pilot study. *Advances in Data Science*, pages 253–289, 2021.
- [18] Donald Woukeng, Damian Sadowski, Jakub Leśkiewicz, Michał Lipiński, and Tomasz Kapela. Rigorous computation in dynamics based on topological methods for multivector fields. *Journal of Applied and Computational Topology*, 8(4):875–908, 2024.