

CLT FOR LES OF CORRELATED NON-HERMITIAN RANDOM MATRICES

INDRAJIT JANA AND SUNITA RANI

ABSTRACT. We consider two $n \times n$ non-Hermitian random matrices such that the ij th entry of one matrix is correlated with the ij th entry of the other matrix. However, the entries of any particular matrix are i.i.d. random variables. We study the asymptotic behavior of the combined spectrum, and the limit of the linear eigenvalue statistic defined on the combined spectrum. We show that if the random variables are centered with variance $1/n$ and having finite moments, then the centered *Linear Eigenvalue Statistics* (LESs) converge jointly to a bivariate Gaussian distribution. We assumed that the test function used in the LES belongs to Sobolev $H^{2+\delta}$ space. The variance of the limiting Gaussian distribution depends on correlation structure of the matrix entries and the fourth order mixed cumulants of the matrix entries. This generalizes the previous results by Rider, Silverstein[22], Cipolloni, Erdős, Schröder [14, 15]. In particular, we obtain the limiting LES of random centrosymmetric matrices.

Keywords: *Random matrix, Centrosymmetric, Central limit theorem, Linear eigenvalue statistic*

Mathematics Subject Classification 2020: 15B52, 60BXX, 60FXX

CONTENTS

1. Introduction	1
2. Notations	2
3. Main result	3
4. Primary reductions	8
5. Central limit theorem for linear eigenvalue statistic	12
6. Central limit theorem for resolvents	17
7. Independence of resolvents	49
References	52

1. INTRODUCTION

In this paper, we focus on analyzing the combined spectrum of two correlated non-Hermitian random matrices. While numerous studies analyzed random matrices with correlated entries [12, 1, 17, 5, 7, 13, 9, 11], to the best of our knowledge, there are relatively few works specifically on the joint convergence of correlated random matrices. Joint convergence of sample cross-covariance matrices [10], independent random elliptic matrices [2],

Indrajit Jana's research is partially supported by INSPIRE Fellowship DST/INSPIRE/04/2019/000015, Dept. of Science and Technology, Govt. of India.

Sunita Rani's research is fully supported by the University Grant Commission (UGC), New Delhi.

independent random Toeplitz matrices [3] were studied in the cited articles. In the last three articles, a correlation structure was assumed for intra-matrix elements; however, no correlation structure was considered for inter-matrix elements. Before continuing any further, let us introduce the following two notions related to the spectrum of a random matrix.

Let $\lambda_1, \lambda_2, \dots, \lambda_n$ be the eigenvalues of an $n \times n$ random matrix M . We define the *Empirical Spectral Measure* (ESM) on the Borel sigma algebra of \mathbb{C} as

$$\mu_n(\cdot) = \frac{1}{n} \sum_{i=1}^n \delta_{\lambda_i}(\cdot),$$

where $\delta_x(A) = \mathbf{1}_{\{x \in A\}}$. This is a random probability measure. It is known that when M is a random non-Hermitian matrix, μ_n converges almost surely in weak topology of measures to the uniform distribution on the unit disc $\{z \in \mathbb{C} : |z| \leq 1\}$. In other words, if $f : \mathbb{C} \rightarrow \mathbb{C}$ is a bounded continuous function, then

$$\frac{1}{n} \sum_{i=1}^n f(\lambda_i) \xrightarrow{\text{a.s.}} \frac{1}{\pi} \int_{\mathbb{C}} f(z) d^2z \text{ as } n \rightarrow \infty,$$

where $d^2z = d\Re z d\Im z$. This is known as *Circular law* [18, 8, 24, 21, 19].

In the above, $\sum_{i=1}^n f(\lambda_i) =: L_n(f)$ is referred as the *Linear Eigenvalue Statistic* (LES) of the matrix M . The fluctuations of the above convergence can be analyzed by studying the asymptotic behavior of $a_n(L_n(f) - b_n)$ for some sequences $\{a_n\}_n, \{b_n\}_n$ and some test function f . Typically, $b_n = \mathbb{E}[L_n(f)]$, but unlike the classical Central Limit Theorem (CLT), a_n is not $\mathcal{O}(1/\sqrt{n})$, primarily because the eigenvalues $\{\lambda_i\}_{1 \leq i \leq n}$ are not independent.

In our case, we have two $n \times n$ random non-Hermitian matrices whose entries are correlated with each other. We study the LES of the joint spectrum via a linear combination of the individual LESs as mentioned in (3.3). The main contribution of this paper is the Theorem 3.2, which says that the limiting LES follows a Gaussian distribution. The variance of the limiting Gaussian distribution depends on the fourth mixed cumulant and the correlation structure of the matrix entries. Our proof relies on the methods developed by Erdős et. al [14, 15]. Broadly, the LES is converted into an integral of trace of resolvent by using Girko's representation formula (3.7). Then it is shown that asymptotically the product of the traces of the resolvents at different points decomposes as per the Isserlis' theorem, which in turn implies that the limiting trace of the resolvent is a Gaussian process.

As a consequence of Theorem 3.2, we also obtain the limiting LES for random Centrosymmetric matrices, which was discussed in our previous work [20]. This article is organized as follows. The main result and the consequences are stated in Section 3. The proof of the CLT for LES is provided in Section 5. However, the key part of the proof relies on establishing the CLT for resolvents, which is presented in Section 6. Let us begin with introducing the notations, which will be frequently used in this article.

2. NOTATIONS

Throughout the paper, we adopt the following notations. Let $\mathbf{H} := \{z \in \mathbb{C} \mid \Im z > 0\}$ denote the upper half-plane, and let $\mathbf{D} \subset \mathbb{C}$ be the open unit disk. The integration with respect to d^2z is equivalent to the integration with respect to $d\Re z d\Im z$. For positive functions f and g , we use the notations $f \lesssim g$ and $f \sim g$ to indicate that $f \leq Cg$ and $df \leq g \leq Df$

for some constants $C, d, D > 0$. For $\nu_1, \nu_2 > 0$, the notation $\nu_1 \ll \nu_2$ means that $\nu_1 \leq c\nu_2$ for sufficiently small constant $c > 0$.

In this paper, we are considering two correlated random matrices $X^{(1)}, X^{(2)}$. We shall use the notation $\mathcal{M}^{(t)}$ to represent a matrix which is a function of the matrix $X^{(t)}$ for $t = 1, 2$. However, \mathcal{M}^t denotes the transpose of the matrix \mathcal{M} . The symbol \mathbb{I} denotes the identity matrix of an appropriate size, while $\mathbf{1} = (1, 1, \dots, 1)$ denotes a column vector of an appropriate size. For an $n \times n$ matrix \mathcal{M} , we define $\langle \mathcal{M} \rangle := \frac{1}{n} \text{Tr } \mathcal{M}$. For vectors $x, y \in \mathbb{C}$, the inner product is defined by $\langle x, y \rangle := \sum \bar{x}_i y_i$, and for matrices $\mathcal{M}_1, \mathcal{M}_2 \in \mathbb{C}^{2n \times 2n}$, we define $\langle \mathcal{M}_1, \mathcal{M}_2 \rangle := \langle \mathcal{M}_1^* \mathcal{M}_2 \rangle$, where \mathcal{M}_1^* is the conjugate transpose of \mathcal{M}_1 .

We use the notation $\kappa(\zeta_{i_1}, \zeta_{i_2}, \dots, \zeta_{i_l})$ to denote a l -th order joint cumulant of the random variables $\zeta_1, \zeta_2, \dots, \zeta_p$. It is defined as follows;

$$\log \left(\mathbb{E} \left[\prod_{k=1}^p s_k \zeta_k \right] \right) = \sum_{l \geq 0} l! \sum_{i_1, \dots, i_l=1}^p \frac{\kappa(\zeta_{i_1}, \zeta_{i_2}, \dots, \zeta_{i_l})}{g_1! g_2! \dots g_p!} s_1^{g_1} s_2^{g_2} \dots s_p^{g_p},$$

where $g_j = \#\{i_q, 1 \leq q \leq l \mid i_q = j\}$. We note down a fact that if the random variables $\zeta_1, \zeta_2, \dots, \zeta_p$ are independent, then $\kappa(\zeta_{i_1}, \zeta_{i_2}, \dots, \zeta_{i_l}) = 0$ whenever $g_j < l$ for all $1 \leq j \leq p$ [23, Theorem 7.12].

We say that a sequence of events $\{F_n\}_n$ occur with high probability if for any arbitrary $K > 0$, there exist $n_0 = n_0(K) \in \mathbb{N}$ such that $\mathbb{P}(F_n^c) < n^{-K}$ for all $n \geq n_0$. We also use the notion of stochastic domination, which is defined in Definition 2.1.

Definition 2.1 (Stochastic domination). Consider families of non-negative random variables,

$$X = \{X_n(\vartheta) \mid n \in \mathbb{N}, \vartheta \in \Theta^{(n)}\}, \quad Y = \{Y_n(\vartheta) \mid n \in \mathbb{N}, \vartheta \in \Theta^{(n)}\},$$

where $\Theta^{(n)}$ is a parameter set. If for any $\delta, K > 0$, there exists $n_0 = n_0(\delta, K) \in \mathbb{N}$ such that

$$\sup_{\vartheta \in \Theta^{(n)}} \mathbb{P} [X_n(\vartheta) > n^\delta Y_n(\vartheta)] \leq n^{-K}, \quad \forall n \geq n_0,$$

then X is said to be stochastically dominated by Y , denoted by $X \prec Y$.

3. MAIN RESULT

Let us consider a $2n \times 2n$ ordered block matrix X defined as follows;

$$(3.1) \quad X = \begin{pmatrix} (X^{(1)})_{n \times n} & 0 \\ 0 & (X^{(2)})_{n \times n} \end{pmatrix},$$

where $X^{(1)}$ and $X^{(2)}$ are i.i.d. random matrices with mean 0 and variance $1/n$ for $i = 1, 2$ respectively. The $(i, j)^{th}$ entry of $X^{(1)}$ is correlated to the $(i, j)^{th}$ entry of $X^{(2)}$ as follows;

$$(3.2) \quad \gamma := n \text{Cov} \left((X^{(1)})_{(i,j)}, (X^{(2)})_{(i,j)} \right) = n \text{Cov} \left(X_{(i,j)}, X_{(n+i, n+j)} \right),$$

and assume that $\{\sigma_i\}_{i=1}^{2n}$ are eigenvalues of X . Without loss of generality, let us assume that $\{\sigma_i\}_{i=1}^n$ are eigenvalues of $X^{(1)}$ and $\{\sigma_i\}_{i=n+1}^{2n}$ are eigenvalues of $X^{(2)}$. The limiting ESM of $X^{(1)}$ and $X^{(2)}$ follow the Circular law. Consequently, the limiting ESM of X follows the Circular law as well.

For test functions f defined on the spectrum of X , let us define the linear eigenvalue statistics as

$$(3.3) \quad \mathcal{L}_n(f) := c\mathcal{L}_n^{(1)}(f) + d\mathcal{L}_n^{(2)}(f),$$

where $c, d \in \mathbb{C}$ and

$$\begin{aligned} \mathcal{L}_n^{(1)}(f) &:= \sum_{i=1}^n f(\sigma_i) - \mathbb{E} \sum_{i=1}^n f(\sigma_i), \\ \mathcal{L}_n^{(2)}(f) &:= \sum_{i=n+1}^{2n} f(\sigma_i) - \mathbb{E} \sum_{i=n+1}^{2n} f(\sigma_i). \end{aligned}$$

From [14, Theorem 2.2] and [15, Theorem 2.2], we have

$$(3.4) \quad \mathcal{L}_n^{(1)}(f) \xrightarrow{d} \mathcal{N}(0, V_f^{(1)}),$$

$$(3.5) \quad \mathcal{L}_n^{(2)}(f) \xrightarrow{d} \mathcal{N}(0, V_f^{(2)}),$$

as $n \rightarrow \infty$, where \xrightarrow{d} denotes convergence in distribution, and $V_f^{(1)}, V_f^{(2)}$ are the variances of the limiting normal distribution, which depend on the test function f and cumulants of the matrix entries. However, $\mathcal{L}_n^{(1)}(f)$ and $\mathcal{L}_n^{(2)}(f)$ are not independent due to (3.2). Therefore, the limiting behavior of $\mathcal{L}_n(f)$ can not be derived from the above limits. The main result of the current paper is the proof of

$$(3.6) \quad \mathcal{L}_n(f) \xrightarrow{d} \mathcal{N}(0, V_f),$$

for non-Hermitian random matrices with correlated entries (3.2) and for general test functions f . This also shows that $(\mathcal{L}_n^{(1)}(f), \mathcal{L}_n^{(2)}(f))$ converges to a bivariate normal distribution. The main idea of this article makes use of the following Girko's formula [25]

$$(3.7) \quad \sum_{\sigma \in \text{Spec}(X^{(t)})} f(\sigma) = -\frac{1}{4\pi} \int_{\mathbb{C}} \Delta f(z) \int_0^\infty \Im \text{Tr}(G^{(t)})^z(\iota\eta) d\eta d^2z,$$

which expresses the linear eigenvalue statistics for any smooth compactly supported test functions in terms of the integral of the trace of the resolvent of

$$(3.8) \quad (H^{(t)})^z := \begin{pmatrix} 0 & X^{(t)} - z \\ (X^{(t)})^* - \bar{z} & 0 \end{pmatrix}, t = 1, 2,$$

where $z \in \mathbb{C}$. The resolvent of $(H^{(t)})^z$ is denoted by $(G^{(t)})^z(\omega) := ((H^{(t)})^z - \omega)^{-1}$.

Before proceeding any further, we list down our assumptions on the matrix X , which is defined in (3.1).

Condition 3.1. Let $X^{(t)} = [x_{ab}^{(t)}]_{1 \leq a, b, \leq n}$, $t = 1, 2$, be two random matrices such that $x_{ab}^{(t)} \stackrel{\text{i.i.d.}}{\sim} n^{-1/2} \chi^{(t)}$, where $\chi^{(t)}$ satisfy the following conditions;

(i) $\mathbb{E}[\chi^{(t)}] = 0$, $\mathbb{E}[|\chi^{(t)}|^2] = 1$, and $\mathbb{E}[(\chi^{(t)})^2] = \gamma_1$, where

$$\gamma_1 = \begin{cases} 1, & \text{if } \chi^{(t)} \text{ is a real valued,} \\ 0, & \text{if } \chi^{(t)} \text{ is a complex valued.} \end{cases}$$

(ii) $\mathbb{E}[\chi^{(1)}\chi^{(2)}] = \rho$, $\mathbb{E}[\chi^{(1)}\overline{\chi^{(2)}}] = \gamma$.

(iii) There exist constants $C_p > 0$, for any $p \in \mathbb{N}$, such that

$$\mathbb{E}[|\chi^{(t)}|^p] \leq C_p.$$

Essentially, we start with an i.i.d. set of bivariate random variables $(\chi^{(1)}, \chi^{(2)})$. Then we create the matrix $X^{(1)}$ out of $n^{-1/2}\chi^{(1)}$ s and $X^{(2)}$ out of $n^{-1/2}\chi^{(2)}$ s. Individually, each matrix $X^{(1)}$ and $X^{(2)}$ are matrices with i.i.d. entries. Therefore, individually the limiting empirical distribution of the eigenvalues of $X^{(1)}$ and $X^{(2)}$ will follow the circular law. To analyze the fluctuations around the circular law, we consider the linear eigenvalue statistic defined in (3.3). We take the test functions from the Sobolev space $H_0^{2+\delta}(\Omega)$, where $\Omega \supset \overline{\mathbf{D}} := \{z \in \mathbb{C} : |z| \leq 1\}$. Here the space $H_0^{2+\delta}(\Omega)$ is the closure of the space of compactly supported smooth functions having the following finite norm

$$\|f\|_{H^{2+\delta}(\Omega)} := \left\| (1 + |\xi|)^{2+\delta} \hat{f}(\xi) \right\|_{L^2(\Omega)},$$

where \hat{f} is the Fourier transform of f .

Theorem 3.2 (Central Limit Theorem for linear eigenvalue statistics). *Let $X^{(1)}$ and $X^{(2)}$ be two $n \times n$ random matrices satisfying Condition 3.1. Let $f \in H^{2+\delta}(\Omega)$ be a test function, where $\delta > 0$ and $\overline{\mathbf{D}} \subset \Omega \subset \mathbb{C}$. Then the centered linear statistics $\mathcal{L}_n(f)$, defined in 3.3, converges in distribution to a complex Gaussian random variable $\mathcal{L}(f)$, where $\mathbb{E}[\mathcal{L}(f)] = 0$, $\mathbb{E}[\mathcal{L}(f)^2] = C(\bar{f}, f)$, and $\mathbb{E}[|\mathcal{L}(f)|^2] = C(f, f) := V_f$. The covariance kernel $C(g, f)$ is given by*

$$(3.9) \quad C(g, f) := -\frac{1}{8\pi^2} \int_{\mathbb{C}} d^2 z_1 \int_{\mathbb{C}} d^2 z_2 \Delta f(z_1) \Delta \overline{g(z_2)} \int_0^\infty d\eta_1 \int_0^\infty d\eta_2 L.$$

The above notations are defined as follows;

$$(3.10) \quad \begin{aligned} L &= c^2 \widehat{V}_{ij}^{(1)} + d^2 \widehat{V}_{ij}^{(2)} + 8cd\gamma(L_{ji}^{12} + L_{ji}^{21}) + 2cd\rho N_{ij} + c^2(\kappa_4)_{11} U_i^{(1)} U_j^{(1)} \\ &\quad + d^2(\kappa_4)_{22} U_i^{(2)} U_j^{(2)} + cd(\kappa_4)_{12} (U_i^{(1)} U_j^{(2)} + U_j^{(1)} U_i^{(2)}), \\ (\kappa_4)_{ts} &= \kappa(\chi^{(t)}, \bar{\chi}^{(t)}, \chi^{(s)}, \bar{\chi}^{(s)}), \\ \widehat{V}_{ij}^{(t)} &= \widehat{V}_{ij}^{(t)}(z_i, z_j, \eta_i, \eta_j) \\ &= \begin{cases} V_{ij}^{(t)}(z_i, z_j, \eta_i, \eta_j), & \text{if } X^{(t)} \text{ is } \mathbb{C}\text{-valued,} \\ V_{ij}^{(t)}(z_i, z_j, \eta_i, \eta_j) + V_{ij}^{(t)}(z_i, \bar{z}_j, \eta_i, \eta_j), & \text{if } X^{(t)} \text{ is } \mathbb{R}\text{-valued,} \end{cases} \\ V_{ij}^{(t)} &= V_{ij}^{(t)}(z_i, z_j, \eta_i, \eta_j) \\ &= \frac{1}{2} \partial_{\eta_i} \partial_{\eta_j} \log \left[1 + (u_i^{(t)} u_j^{(t)} |z_i| |z_j|)^2 - (m_i^{(t)})^2 (m_j^{(t)})^2 - 2u_i^{(t)} u_j^{(t)} \Re z_i \bar{z}_j \right], \\ L_{ji}^{12} &= \frac{m_j^{(1)} m_i^{(2)} u_j^{(1)} u_i^{(2)} \Re(\bar{z}_i z_j)}{\beta_j^{(1)} \beta_i^{(2)}}, \quad L_{ji}^{21} = \frac{m_j^{(2)} m_i^{(1)} u_j^{(2)} u_i^{(1)} \Re(\bar{z}_i z_j)}{\beta_j^{(2)} \beta_i^{(1)}}, \\ N_{ij} &= (\partial_{\eta_i} (m_i^{(1)})) (\partial_{\eta_j} (m_j^{(2)})) + (\partial_{\eta_i} (m_i^{(2)})) (\partial_{\eta_j} (m_j^{(1)})), \\ U_i^{(t)} &= \frac{i}{\sqrt{2}} \partial_{\eta_i} (m_i^{(t)})^2, \end{aligned}$$

$m_i^{(t)} = (m^{z_i})^{(t)}(\iota\eta_i)$ is the solution of the equation (4.2), $u_i = u^{z_i}(\iota\eta_i) = -\frac{\iota\Im m_i}{\eta_i + \Im m_i}$, and $\beta_i^{(t)} = 1 - (m_i^{(t)})^2 - (u_i^{(t)})^2 |z_i|^2$.

The above theorem is proved using the resolvents of the matrix $(H^{(t)})^z$, which is $(G^{(t)})^z$. Using (3.7), the linear eigenvalue statistics $\mathcal{L}_n(f)$ can be expressed in terms of $(G^{(t)})^z$ as follows;

$$\begin{aligned}
(3.11) \quad & \mathcal{L}_n^{(t)}(f) \\
&= \frac{1}{4\pi} \int_{\mathbb{C}} \Delta f(z) [\log |\det((H^{(t)})^z - iT)| - \mathbb{E} \log |\det((H^{(t)})^z - iT)|] d^2z \\
&\quad - \frac{n}{2\pi i} \int_{\mathbb{C}} \Delta f \left[\left(\int_0^{\eta_0} + \int_{\eta_0}^{\eta_c} + \int_{\eta_c}^T \right) [\langle (G^{(t)})^z(\iota\eta) - \mathbb{E}(G^{(t)})^z(\iota\eta) \rangle] d\eta \right] d^2z \\
&=: J_T^{(t)} + (I_0^{\eta_0})^{(t)} + (I_{\eta_0}^{\eta_c})^{(t)} + (I_{\eta_c}^T)^{(t)}.
\end{aligned}$$

Here, we have decomposed the whole integral into four parts, namely $(0, \eta_0)$, $[\eta_0, \eta_c)$, $[\eta_c, T)$, and $[T, \infty)$, where $\eta_0 = n^{-1-\delta_0}$, $\eta_c = n^{-1+\delta_1}$, and $T = n^C$, for some small constants $\delta_0, \delta_1 > 0$ and a large positive constant C . In our context, the linear eigenvalue statistic $\mathcal{L}_n(f) = c\mathcal{L}_n^{(1)}(f) + d\mathcal{L}_n^{(2)}(f)$, which can similarly be expressed as

$$\begin{aligned}
(3.12) \quad & \mathcal{L}_n(f) \\
&= c\mathcal{L}_n^{(1)}(f) + d\mathcal{L}_n^{(2)}(f) \\
&=: J_T + I_0^{\eta_0} + I_{\eta_0}^{\eta_c} + I_{\eta_c}^T,
\end{aligned}$$

where

$$\begin{aligned}
J_T &= cJ_T^{(1)} + dJ_T^{(2)}, & I_0^{\eta_0} &= c(I_0^{\eta_0})^{(1)} + d(I_0^{\eta_0})^{(2)}, \\
I_{\eta_0}^{\eta_c} &= c(I_{\eta_0}^{\eta_c})^{(1)} + d(I_{\eta_0}^{\eta_c})^{(2)}, & I_{\eta_c}^T &= c(I_{\eta_c}^T)^{(1)} + d(I_{\eta_c}^T)^{(2)}.
\end{aligned}$$

Now, we handle J_T , $I_0^{\eta_0}$, $I_{\eta_0}^{\eta_c}$, and $I_{\eta_c}^T$ using different techniques. In particular, from (5.1), we observe that the contribution of J_T is negligible. To estimate $I_0^{\eta_0}$, we use the fact that, with high probability, there are no eigenvalues in the interval $[0, \eta_0]$, as established in [26, Theorem 3.2]. The contribution of $I_{\eta_0}^{\eta_c}$ is analyzed in Proposition 7.3. The main contributing factor in (3.12) is $I_{\eta_c}^T$, which constitutes the majority of this paper. Essentially, we need to analyze the limiting behaviors of $\langle (G^{(1)})^z(\iota\eta) - \mathbb{E}(G^{(1)})^z(\iota\eta) \rangle$, and $\langle (G^{(2)})^z(\iota\eta) - \mathbb{E}(G^{(2)})^z(\iota\eta) \rangle$ in the regime $[\eta_c, T]$. This is formulated in the following Theorem 3.3.

Theorem 3.3 (CLT for resolvents). *Let $G_i^{(t)} := (G^{(t)})^{z_i}(\iota\eta_i)$, where $\eta_i > 0$ for $1 \leq i \leq p$. Let $z_1, z_2, \dots, z_p \in \mathbb{C}$ be distinct complex numbers such that whenever $i \neq j$, $\min\{\eta_i, \eta_j\} \geq$*

$n^{\epsilon-1}|z_i - z_j|^{-2}$ for some $\epsilon > 0$. Then we have

$$\begin{aligned}
& \mathbb{E} \left[\prod_{i=1}^p c \langle G_i^{(1)} - \mathbb{E}(G_i^{(1)}) \rangle + d \langle G_i^{(2)} - \mathbb{E}(G_i^{(2)}) \rangle \right] \\
&= \sum_{\mathcal{P} \in \text{Parings}[p]} \prod_{\{i,j\} \in \mathcal{P}} \mathbb{E} \left[\{ c \langle G_i^{(1)} - \mathbb{E}(G_i^{(1)}) \rangle + d \langle G_i^{(2)} - \mathbb{E}(G_i^{(2)}) \rangle \} \right. \\
&\quad \left. \times \{ c \langle G_j^{(1)} - \mathbb{E}(G_j^{(1)}) \rangle + d \langle G_j^{(2)} - \mathbb{E}(G_j^{(2)}) \rangle \} \right] + \mathcal{O}_{\prec}(\Psi) \\
(3.13) \quad &= \frac{1}{n^p} \sum_{\mathcal{P} \in \text{Parings}[p]} \prod_{\{i,j\} \in \mathcal{P}} \left\{ \frac{c^2 \widehat{V}_{ij}^{(1)} + d^2 \widehat{V}_{ij}^{(2)} + 8cd\gamma(L_{ji}^{12} + L_{ji}^{21}) + 2cd\rho N_{ij}}{2} \right. \\
&\quad \left. + \frac{c^2(\kappa_4)_{11} U_i^{(1)} U_j^{(1)} + d^2(\kappa_4)_{22} U_i^{(2)} U_j^{(2)} + cd(\kappa_4)_{12} (U_i^{(1)} U_j^{(2)} + U_j^{(1)} U_i^{(2)})}{2} \right\} \\
&\quad + \mathcal{O}_{\prec}(\Psi),
\end{aligned}$$

where

$$\begin{aligned}
(3.14) \quad & \widehat{V}_{ij}^{(t)}(z_i, z_j, \eta_i, \eta_j) \\
&= \begin{cases} V_{ij}^{(t)}(z_i, z_j, \eta_i, \eta_j), & \text{if } X^{(t)} \text{ is } \mathbb{C}\text{-valued,} \\ V_{ij}^{(t)}(z_i, z_j, \eta_i, \eta_j) + V_{ij}^{(t)}(z_i, \bar{z}_j, \eta_i, \eta_j), & \text{if } X^{(t)} \text{ is } \mathbb{R}\text{-valued,} \end{cases} \\
(3.15) \quad & \text{and } V_{ij}^{(t)}(z_i, z_j, \eta_i, \eta_j) \\
&= \frac{1}{2} \partial_{\eta_i} \partial_{\eta_j} \log \left[1 + (u_i^{(t)} u_j^{(t)} |z_i| |z_j|)^2 - (m_i^{(t)})^2 (m_j^{(t)})^2 - 2u_i^{(t)} u_j^{(t)} \Re z_i \bar{z}_j \right],
\end{aligned}$$

$$\begin{aligned}
(3.16) \quad & \Psi \\
&= \left(\frac{1}{(n\eta_*)^2} + \frac{1}{n^2 \eta_* |z_i - z_j|^4} + \frac{1}{(n\eta_*)^3 |z_i - z_j|^4} + \frac{1}{\Im z_i (n\eta_*)^{3/2}} \right) \prod_{i \in [p]} \frac{(n\eta_i)^{-1/2}}{|1 - |z_i||},
\end{aligned}$$

where $\eta_* := \min_i \{\eta_i\}$. The other notations are same as defined in (3.10).

We would like to conclude this section by noting down the following consequences.

- (A) Recall from (3.3) that $\mathcal{L}_n(f) = c\mathcal{L}_n^{(1)}(f) + d\mathcal{L}_n^{(2)}(f)$. Since Theorem 3.2 is true for all c, d , it implies that $(\mathcal{L}_n^{(1)}(f), \mathcal{L}_n^{(2)}(f))$ converges to a bivariate Gaussian distribution.
- (B) If the random matrices defined in the Condition 3.1 are independent of each other, then $\gamma = 0 = \rho$. Moreover, the joint cumulants $(\kappa_4)_{12} = 0$. As a result the Theorem 3.3 becomes equivalent to [14, Proposition 3.3] when the matrices are complex valued, and [15, Proposition 3.3] when the matrices are real valued.
- (C) If we begin with a $2n \times 2n$ random centrosymmetric matrix X , then X can be represented as (3.1), where the matrices $X^{(1)}$ and $X^{(2)}$ are $n \times n$ random matrices with i.i.d. entries. Moreover, there exist independent random variables ζ and θ such that the entries of $X^{(1)}$ have the same distribution as $\zeta + \theta$, and the entries of $X^{(2)}$ have the same distribution as $\zeta - \theta$ (see [27, Theorem 9]). This impels that the (ab) th entry of $X^{(1)}$ and (ab) th

entry of $X^{(2)}$ are uncorrelated but may not be independent. In other words, according to the notations used in Condition 3.1, $\gamma = 0 = \rho$. But $(\kappa_4)_{12}$ is not necessarily zero.

However, if in addition, we assume that the test function f is analytic over $\overline{\mathbf{D}}$, then in the expression of the covariance kernel $C(f, g)$ (see (3.9)), only the integration against $c^2\widehat{V_{ij}^{(1)}} + d^2\widehat{V_{ij}^{(2)}}$ will survive. Following the calculations outlined in [14, Lemma 4.8], we obtain the expression of the limiting variance as

$$C(f, f) = \frac{2}{\pi} \int_{\mathbf{D}} |f'(z)|^2 d^2z.$$

The above expression was proved independently in a previous work [20, Theorem 3.4].

4. PRIMARY REDUCTIONS

Proof of the main theorem requires a careful analysis of the resolvent of the matrix $(H^{(t)})^z$, which is defined in (3.8). The asymptotic behavior of the resolvent $(G^{(t)})^z(\omega)$ is given by the following matrix

$$(4.1) \quad \begin{aligned} (M^{(t)})^z(\omega) &:= \begin{pmatrix} (m^{(t)})^z(\omega)\mathbb{I} & -z(u^{(t)})^z(\omega)\mathbb{I} \\ -\bar{z}(u^{(t)})^z(\omega)\mathbb{I} & (m^{(t)})^z(\omega)\mathbb{I} \end{pmatrix}, \\ (u^{(t)})^z(\omega) &:= \frac{(m^{(t)})^z(\omega)}{\omega + (m^{(t)})^z(\omega)}, \end{aligned}$$

where \mathbb{I} is the $n \times n$ identity matrix, and $(m^{(t)})^z(\omega)$ is the unique solution of the following equation

$$(4.2) \quad -\frac{1}{(m^{(t)})^z} = \omega + (m^{(t)})^z - \frac{|z|^2}{\omega + (m^{(t)})^z}, \quad \eta \Im(m^{(t)})^z(\omega) > 0, \quad \eta = \Im\omega \neq 0.$$

The above follows from the analysis of *matrix Dyson equation* (MDE) (see [4]). The exact quantification of the error $(G^{(t)})^z(\omega) - (M^{(t)})^z(\omega)$ is given by the local law, which is mentioned in Theorem 4.1.

As it is mentioned in the statement of Theorem 3.2, the derivative of $(m^{(t)})^z(\omega)$ with respect to ω and $\Im\omega = \eta$ appears frequently in the rest of the article. We note down the expression below;

$$(4.3) \quad \partial_\omega(m^{(t)}) = \frac{1 - \beta^{(t)}}{\beta^{(t)}} = -\iota\partial_\eta(m^{(t)}), \quad \beta^{(t)} := 1 - (m^{(t)})^2 - (u^{(t)})^2|z|^2.$$

Along the same line, we define yet another quantity

$$\beta_*^{(t)} = 1 - |m^{(t)}|^2 - |u^{(t)}|^2|z|^2,$$

which satisfies

$$(4.4) \quad \beta_*^{(t)} \Im m^{(t)} = (1 - \beta_*^{(t)}) \Im \omega,$$

$$(4.5) \quad (|m^{(t)}|^2 - |u^{(t)}|^2|z|^2 + 1) \Re m^{(t)}(\iota\eta) = 0.$$

$$(4.6) \quad (\beta^{(t)})_* = \frac{\eta}{\Im m^{(t)} + \eta}, \quad \beta^{(t)} = (\beta^{(t)})_* + 2(\Im m^{(t)})^2.$$

The above identities are obtained by taking the imaginary and real part of (4.2) on the both sides. Now, since $\Im\omega$ and $\Im m^{(t)}$ have the same sign (from (4.2)), $\beta_*^{(t)}$ and $(1 - \beta_*^{(t)})$ must

also have the same sign. However, as per the definition, $1 - \beta_*^{(t)} > 0$. Consequently, $\beta_*^{(t)} > 0$, which is equivalent to

$$(4.7) \quad |m^{(t)}|^2 + |(u^{(t)})^z|^2 |z|^2 < 1,$$

which implies $|(u^{(t)})^z|^2 |z|^2 < 1$ as well. Using this fact in (4.5), we obtain

$$\Re m^{(t)}(\iota\eta) = 0.$$

Moreover, from (4.2), we have $u^{(t)} = -(m^{(t)})^2 + (u^{(t)})^2 |z|^2$, which in conjunction with (4.7) implies that $|u^{(t)}| < 1$. Therefore, we obtain the following uniform bound in z and ω

$$(4.8) \quad \|(M^{(t)})^z(\omega)\| + |(m^{(t)})^z(\omega)| + |(u^{(t)})^z(\omega)| \lesssim 1.$$

Furthermore, from [6, eq. (3.13)] we have

$$(4.9) \quad \Im m^{(t)}(\iota\eta) \sim \begin{cases} \eta^{1/3} + |1 - |z|^2|^{1/2} & \text{if } |z| \leq 1 \\ \frac{\eta}{|z|^2 - 1 + \eta^{2/3}} & \text{if } |z| > 1 \end{cases} \quad \eta \lesssim 1.$$

Now, using (4.7), (4.9), we get a lower bound on $|\beta^{(t)}|$ as follows;

$$(4.10) \quad \begin{aligned} |\beta^{(t)}| &\geq 1 - \Re((m^{(t)})^2) - \Re((u^{(t)})^2) |z|^2 \\ &= 1 - (\Re(m^{(t)}))^2 + (\Im(m^{(t)}))^2 - \Re((u^{(t)})^2) |z|^2 \\ &\geq (1 - |m^{(t)}|^2 - |u^{(t)}|^2 |z|^2) + (\Im(m^{(t)}))^2 \\ &\geq (\Im(m^{(t)}))^2 \\ &\gtrsim \eta^{2/3} + |1 - |z|^2| \\ &\geq \eta^{2/3} + |1 - |z||. \end{aligned}$$

We conclude this subsection by stating the local law of the resolvent.

Theorem 4.1 (Optimal local law for $G^{(t)}$, [16], Theorem 3.1). *For any $\epsilon > 0$ and $z \in \mathbb{C}$ with $|1 - |z|| \geq \epsilon$ the resolvent $(G^{(t)})^z$ at $\omega \in \mathbf{H}$ with $\eta = \Im \omega$ is very well approximated by the deterministic matrix $(M^{(t)})^z$ in the sense*

$$\begin{aligned} |\langle ((G^{(t)})^z(\omega) - (M^{(t)})^z(\omega))A \rangle| &\leq \frac{C_\epsilon \|A\| n^\xi}{n\eta} \\ |\langle \mathbf{x}, ((G^{(t)})^z(\omega) - (M^{(t)})^z(\omega)) \mathbf{y} \rangle| &\leq C_\epsilon \|\mathbf{x}\| \|\mathbf{y}\| n^\xi \left(\frac{1}{\sqrt{n\eta}} + \frac{1}{n\eta} \right), \end{aligned}$$

with very high probability for some $C_\epsilon \leq \epsilon^{-100}$, uniformly for $\eta > n^{-100}$, and for any deterministic matrices A and any vectors \mathbf{x}, \mathbf{y} and $\xi > 0$.

This local law will be used in various estimates. The main idea is to write $G^{(t)} = M^{(t)} + G^{(t)} - M^{(t)}$. Then, any particular entry of $G^{(t)}$ can be written as $G_{ab}^{(t)} = M_{ab}^{(t)} + (G^{(t)} - M^{(t)})_{ab}$. Now, the estimate of $(G^{(t)} - M^{(t)})_{ab}$ follows from the second part of the local law by taking $\mathbf{x} = \mathbf{e}_a$ and $\mathbf{y} = \mathbf{e}_b$. Moreover, since the matrix $M^{(t)}$ is mostly a diagonal matrix, $M_{ab}^{(t)} = 0$ for most of the off-diagonal $M_{ab}^{(t)}$. A combination of these two facts yield the required estimates. Additionally, when we have $(G^{(t)}A)_{ab}$, we decompose $G^{(t)}$ as before $G^{(t)} = M^{(t)} + G^{(t)} - M^{(t)}$. Then, we apply the first part of the local law to estimate $((G^{(t)} - M^{(t)})A)_{ab}$. This idea is being repeatedly used in the estimates such as (6.9), (6.52), (6.53) etc..

4.1. **Local law for product of resolvents.** We begin with the shorthand notations

$$(4.11) \quad G_i^{(t)} := (G^{(t)})^{z_i}(\omega_i), \quad M_i^{(t)} := (M^{(t)})^{z_i}(\omega_i).$$

From [6], we note down the deviation of $G^{(t)}$ from $M^{(t)}$ in the equation (4.14), which requires the following notions

$$W^{(t)} := \begin{pmatrix} 0 & X^{(t)} \\ (X^{(t)})^* & 0 \end{pmatrix}.$$

The linear covariance operator $\mathcal{S} : \mathbb{C}^{2n \times 2n} \rightarrow \mathbb{C}^{2n \times 2n}$ is defined by

$$(4.12) \quad \mathcal{S} \left[\begin{pmatrix} A & B \\ C & D \end{pmatrix} \right] := \tilde{\mathbb{E}}^{(t)} \tilde{W}^{(t)} \begin{pmatrix} A & B \\ C & D \end{pmatrix} \tilde{W}^{(t)} = \begin{pmatrix} \langle D \rangle & 0 \\ 0 & \langle A \rangle \end{pmatrix},$$

where $\tilde{W}^{(t)} = \begin{pmatrix} 0 & \tilde{X}^{(t)} \\ (\tilde{X}^*)^{(t)} & 0 \end{pmatrix},$

and $\tilde{X}^{(t)} \sim \text{Gin}_{\mathbb{C}}$. Here $\text{Gin}_{\mathbb{C}}$ stands for the standard complex Ginibre ensemble, $\tilde{\mathbb{E}}^{(t)}$ denotes the expectation with respect to $\tilde{X}^{(t)}$.

For any given function $f : \mathbb{C}^{2n \times 2n} \rightarrow \mathbb{C}^{2n \times 2n}$, we define the self renormalization as follows;

$$(4.13) \quad \underline{W^{(t)} f(W^{(t)})} := W^{(t)} f(W^{(t)}) - \tilde{\mathbb{E}}^{(t)} \tilde{W}^{(t)} (\partial_{\tilde{W}^{(t)}} f) (W^{(t)}),$$

where $\partial_{\tilde{W}^{(t)}}$ indicates a directional derivative in the direction $\tilde{W}^{(t)}$ and $\tilde{W}^{(t)}$ denotes an independent random matrix as in (4.12) with $\tilde{X}^{(t)}$ a complex Ginibre matrix with expectation $\tilde{\mathbb{E}}^{(t)}$.

4.1.1. *Approximation of $G_1^{(t)} B G_2^{(t)}$ via stability operators.* According to the definition above, we note that for any deterministic matrix B with bounded norm, we have

$$\begin{aligned} \underline{W^{(t)} G^{(t)}} &= W^{(t)} G^{(t)} + \mathcal{S}[G^{(t)}] G^{(t)} \\ \underline{W^{(t)} G_1^{(t)} B G_2^{(t)}} &= W^{(t)} G_1^{(t)} B G_2^{(t)} + \mathcal{S} \left[G_1^{(t)} \right] G_1^{(t)} B G_2^{(t)} + \mathcal{S} \left[G_1^{(t)} B G_2^{(t)} \right] G_2^{(t)} \\ &= \underline{W^{(t)} G_1^{(t)} B G_2^{(t)}} + \mathcal{S} \left[G_1^{(t)} B G_2^{(t)} \right] G_2^{(t)}. \end{aligned}$$

Now, using the above equations, (4.2), (4.1), along with the observation $\mathcal{S}[M_i^{(t)}] = m_i^{(t)} \mathbb{I}$, we get the following identity

$$(4.14) \quad G_i^{(t)} = M_i^{(t)} - M_i^{(t)} \underline{W^{(t)} G_i^{(t)}} + M_i \mathcal{S}[G_i^{(t)} - M_i^{(t)}] G_i^{(t)}.$$

Similarly, we can calculate the deviation of $G_1^{(t)} B G_2^{(t)}$ from $M_1^{(t)} B M_2^{(t)}$ as follows;

$$(4.15) \quad \begin{aligned} G_1^{(t)} B G_2^{(t)} &= M_1^{(t)} B G_2^{(t)} - M_1^{(t)} \underline{W^{(t)} G_1^{(t)} B G_2^{(t)}} + M_1^{(t)} \mathcal{S}[G_1^{(t)} - M_1^{(t)}] G_1^{(t)} B G_2^{(t)} \\ &= M_1^{(t)} B M_2^{(t)} + M_1^{(t)} B (G_2^{(t)} - M_2^{(t)}) - M_1^{(t)} \underline{W^{(t)} G_1^{(t)} B G_2^{(t)}} \\ &\quad + M_1^{(t)} \mathcal{S}[G_1^{(t)} B G_2^{(t)}] M_2^{(t)} + M_1^{(t)} \mathcal{S}[G_1^{(t)} B G_2^{(t)}] (G_2^{(t)} - M_2^{(t)}) \\ &\quad + M_1^{(t)} \mathcal{S}[G_1^{(t)} - M_1^{(t)}] G_1^{(t)} B G_2^{(t)}. \end{aligned}$$

It can be shown that the terms $G_i^{(t)} - M_i^{(t)}$ and the self renormalized terms are small, which is formalized in Theorem 4.2. As a result, we obtain the following approximation

$$(4.16) \quad G_1^{(t)} B G_2^{(t)} \approx (M_B^{(t)})^{z_1, z_2},$$

where

$$(4.17) \quad (M_B^{(t)})^{z_1, z_2}(\omega_1, \omega_2) := (\mathbb{I} - (M^{(t)})^{z_1}(\omega_1) \mathcal{S}[\cdot] (M^{(t)})^{z_2}(\omega_2))^{-1} [(M^{(t)})^{z_1}(\omega_1) B (M^{(t)})^{z_2}(\omega_2)].$$

In this context, we define the 2-body stability operator $\hat{\mathcal{B}}_{12}^{(t)}(z_1, z_2, \omega_1, \omega_2) : GL_{2n}(\mathbb{C}) \rightarrow GL_{2n}(\mathbb{C})$ as follows;

$$(4.18) \quad \hat{\mathcal{B}}_{12}^{(t)}(z_1, z_2, \omega_1, \omega_2) := \mathbb{I} - M_1^{(t)} \mathcal{S}[\cdot] M_2^{(t)}.$$

Here, $GL_{2n}(\mathbb{C})$ denotes the space of $2n \times 2n$ matrices with the usual matrix norm. In the rest of the article, for convenience, we denote $\hat{\mathcal{B}}_{12}^{(t)}(z_1, z_2, \omega_1, \omega_2)$ as $\hat{\mathcal{B}}_{12}^{(t)}$ or $\hat{\mathcal{B}}^{(t)}$. Similarly, for $t \neq s$, we can have the following approximation

$$(4.19) \quad G_1^{(t)} B G_2^{(s)} \approx M_1^{(t)} B M_2^{(s)}.$$

Theorem 4.2. [16, Theorem 3.5] *Fix $z_1, z_2 \in \mathbb{C}$ with $|1 - |z_i|| \geq \epsilon$, for some $\epsilon > 0$ and $\omega_1, \omega_2 \in \mathbb{C}$ with $|\eta_i| := |\Im w_i| \geq n^{-1}$ such that*

$$\eta_* := \min\{|\eta_1|, |\eta_2|\} \geq n^{-1+\epsilon_*} |(\hat{\beta}_*^{(t)})^{-1}|$$

for some $\epsilon_* > 0$, where $|\hat{\beta}_*^{(t)}|$ is the smallest eigen value of $\hat{\mathcal{B}}_{12}^{(t)}$. Then, for any bounded deterministic matrix B , with $\|B\| \lesssim 1$, the product of resolvents

$$(G^{(t)})^{z_1} B (G^{(t)})^{z_2} = (G^{(t)})^{z_1}(\omega_1) B (G^{(t)})^{z_2}(\omega_2)$$

is approximated by $(M^{(t)})_B^{z_1, z_2} = (M^{(t)})_B^{z_1, z_2}(\omega_1, \omega_2)$ defined in (4.17) in the sense that

$$(4.20) \quad |\langle A((G^{(t)})^{z_1} B (G^{(t)})^{z_2} - (M^{(t)})_B^{z_1, z_2}) \rangle| \lesssim \frac{C_\epsilon \|A\| n^\xi}{n \eta_* (\eta_1 \eta_2)^{1/2} |\hat{\beta}_*^{(t)}|} \left((\eta_*)^{1/12} + \frac{\eta_*^{1/4}}{|\hat{\beta}_*^{(t)}|} + \frac{1}{\sqrt{n \eta_*}} + \frac{1}{(|\hat{\beta}_*^{(t)}| n \eta_*)^{1/4}} \right),$$

$$(4.21) \quad |\langle \mathbf{x}, ((G^{(t)})^{z_1} B (G^{(t)})^{z_2} - (M^{(t)})_B^{z_1, z_2}) \mathbf{y} \rangle| \lesssim \frac{C_\epsilon \|\mathbf{x}\| \|\mathbf{y}\| n^\xi}{(n \eta_*)^{1/2} (\eta_1 \eta_2)^{1/2} |\hat{\beta}_*^{(t)}|}$$

for some C_ϵ with very high probability for any deterministic $A, \mathbf{x}, \mathbf{y}$, and $\epsilon \geq 0$. If $\omega_1, \omega_2 \in i\mathbb{R}$ we may choose $C_\epsilon = 1$, otherwise we can choose $C_\epsilon \leq \epsilon^{-100}$.

In continuation with the two-body stability operator, we also note down the one-body stability operator as follows;

$$(4.22) \quad \mathcal{B}^{(t)} := \mathbb{I} - M^{(t)} \mathcal{S}[\cdot] M^{(t)}.$$

Recalling the action of \mathcal{S} from (4.12), we see that \mathcal{B} acting on a block matrix $\begin{pmatrix} a\mathbb{I} & b\mathbb{I} \\ c\mathbb{I} & d\mathbb{I} \end{pmatrix}$ is given by

$$\mathcal{B} \begin{pmatrix} a\mathbb{I} & b\mathbb{I} \\ c\mathbb{I} & d\mathbb{I} \end{pmatrix} = \begin{pmatrix} (1 - d(m^{(t)})^2 - a|z|^2(u^{(t)})^2)\mathbb{I} & (d+a)zu^{(t)}m^{(t)}\mathbb{I} \\ (d+a)\bar{z}u^{(t)}m^{(t)}\mathbb{I} & (1 - d|z|^2u^{(t)} - a(m^{(t)})^2)\mathbb{I} \end{pmatrix}.$$

From the above expression, we notice that any eigenvector of \mathcal{B} will be of the form $\begin{pmatrix} a\mathbb{I} & 0 \\ 0 & d\mathbb{I} \end{pmatrix}$.

We now enlist the following lemmas, which estimate the bounds on the eigenvalues of the stability operators.

Lemma 4.3. [16, Lemma 5.2] *Let $\omega \in \mathbf{H}, z \in \mathbb{C}$ be bounded spectral parameters, $|\omega| + |z| \lesssim 1$. Then the operator $\mathcal{B}^{(t)}$ has the trivial eigenvalue 1 and two non-trivial eigenvalues $1 + (m^{(t)})^2 - |z|^2(u^{(t)})^2$ and $1 - (m^{(t)})^2 - (u^{(t)})^2|z|^2$, which is same as $\beta^{(t)}$ defined in (4.3).*

Moreover, $\bar{\beta}^{(t)}$ is an eigenvalue of \mathcal{B}^ corresponding to the eigenvector $\begin{pmatrix} \mathbb{I} & 0 \\ 0 & \mathbb{I} \end{pmatrix}$. In addition, we have the bound*

$$|\beta^{(t)}| \gtrsim |1 - |z|| + \eta^{2/3}.$$

Lemma 4.4. [16, Lemma 6.1] *For $z_1, z_2 \in \mathbb{C}, \omega_1, \omega_2 \in \mathbb{C} \setminus \mathbb{R}$ such that $|z_i|, |\omega_i| \lesssim 1$, the two non-trivial eigenvalues $\hat{\beta}^{(t)}, \hat{\beta}_*^{(t)}$ of $\hat{\mathcal{B}}^{(t)}$ satisfy*

$$\min \{ \Re \hat{\beta}^{(t)}, \Re \hat{\beta}_*^{(t)} \} \gtrsim |z_1 - z_2|^2 + \min \{ |\omega_1 + \bar{\omega}_2|, |\omega_1 - \bar{\omega}_2| \}^2 + |\Im \omega_1| + |\Im \omega_2|.$$

5. CENTRAL LIMIT THEOREM FOR LINEAR EIGENVALUE STATISTIC

In this section, we prove our main result, Theorem 3.2. Before proceeding, let us state the following result, which asserts that the main contribution comes from the regime $I_{\eta_c}^T$.

Lemma 5.1 ([14], Lemma 4.3, 4.4). *For some bounded open $\bar{\mathbf{D}} \subset \Omega \subset \mathbb{C}$, let $f \in H_0^{2+\delta}(\Omega)$. Then, for any $\xi > 0$ the following bound holds with high probability;*

$$\begin{aligned} |J_T^{(t)}| &\lesssim \frac{n^{1+\xi} \|\Delta f\|_{L^1(\Omega)}}{T^2}, \\ |(I_0^{\eta_0})^{(t)}| + |(I_{\eta_0}^{\eta_c})^{(t)}| + |(I_{\eta_c}^T)^{(t)}| &\lesssim n^\xi \|\Delta f\|_{L^2(\Omega)} |\Omega|^{1/2}, \end{aligned}$$

where $J_T^{(t)}, (I_0^{\eta_0})^{(t)}, (I_{\eta_0}^{\eta_c})^{(t)}, (I_{\eta_c}^T)^{(t)}$ are same as defined in (3.11). Moreover, there exists $\delta' > 0$ such that

$$\mathbb{E}[|(I_0^{\eta_0})^{(t)}|] + \mathbb{E}[|(I_{\eta_0}^{\eta_c})^{(t)}|] \lesssim n^{-\delta'} \|\Delta f\|_{L^2(\Omega)}.$$

Applying the above lemma for $t = 1, 2$, we have the following estimates

$$(5.1) \quad |J_T| \lesssim \frac{n^{1+\xi} \|\Delta f\|_{L^1(\Omega)}}{T^2}, \quad |I_0^{\eta_0}| + |I_{\eta_0}^{\eta_c}| + |I_{\eta_c}^T| \lesssim n^\xi \|\Delta f\|_{L^2(\Omega)} |\Omega|^{1/2},$$

with high probability, and

$$(5.2) \quad \mathbb{E}[|I_0^{\eta_0}|] + \mathbb{E}[|I_{\eta_0}^{\eta_c}|] \lesssim n^{-\delta'} \|\Delta f\|_{L^2(\Omega)},$$

where $J_T, I_0^{\eta_0}, I_{\eta_0}^{\eta_c}, I_{\eta_c}^T$ are defined in (3.12).

The above estimates (5.1), and (5.2) imply that the main contribution in $\mathcal{L}_n(f)$ comes from $I_{\eta_c}^T$, which leads to the following corollary.

Corollary 5.2. *Let $f_i \in H^{2+\delta}(\Omega)$, $1 \leq i \leq p$ for some bounded open $\bar{\mathbf{D}} \subset \Omega \subset \mathbb{C}$. Then*

$$\mathbb{E} \left[\prod_{i \in [p]} \mathcal{L}_n(f_i) \right] = \mathbb{E} \left[\prod_{i \in [p]} I_{\eta_c}^T(f_i) \right] + \mathcal{O}_{\prec}(n^{-c(p)}),$$

where $\mathcal{L}_n(f_i)$ and $I_{\eta_c}^T(f_i)$ defined in (3.11) and $c(p) > 0$.

The above corollary leads to the following;

$$\begin{aligned}
& \mathbb{E} \left[\prod_{i \in [p]} \mathcal{L}_n(f_i) \right] \\
&= \mathbb{E} \left[\prod_{i \in [p]} I_{\eta_c}^T(f_i) \right] + \mathcal{O}_{\prec}(n^{-c(p)}) \\
(5.3) \quad &= \mathbb{E} \left[\prod_{i \in [p]} \left(-\frac{n}{2\pi\iota} \int_{\mathbb{C}} \Delta f_i(z) \int_{\eta_c}^T (c \langle (G^{(1)})^z(\iota\eta) - \mathbb{E}(G^{(1)})^z(i\eta) \rangle \right. \right. \\
&\quad \left. \left. + d \langle (G^{(2)})^z(\iota\eta) - \mathbb{E}(G^{(2)})^z(\iota\eta) \rangle) d\eta d^2 z \right) \right] + \mathcal{O}_{\prec}(n^{-c(p)}).
\end{aligned}$$

Lemma 5.3. *Let f_i be as in Corollary 5.2. Then*

$$\begin{aligned}
(5.4) \quad & \mathbb{E} \left[\prod_{i \in [p]} \mathcal{L}_n(f_i) \right] \\
&= \mathbb{E} \left[\prod_{i \in [p]} \left(-\frac{n}{2\pi\iota} \int_{\mathbb{C}} \Delta f_i(z) \int_{\eta_c}^T (c \langle (G^{(1)})^z(\iota\eta) - \mathbb{E}(G^{(1)})^z(\iota\eta) \rangle \right. \right. \\
&\quad \left. \left. + d \langle (G^{(2)})^z(\iota\eta) - \mathbb{E}(G^{(2)})^z(\iota\eta) \rangle) d\eta d^2 z \right) \right] \\
&= \frac{1}{8\pi^2} \sum_{p \in \prod_p} \prod_{\{i,j\} \in P} \left[\int_{\mathbb{C}} -d^2 z_i \Delta f_i \int_{\mathbb{C}} d^2 z_j \Delta f^{(j)} \right. \\
&\quad \times \int_0^\infty d\eta_i \int_0^\infty d\eta_j \left(c^2 \widehat{V}_{ij}^{(1)} + d^2 \widehat{V}_{ij}^{(2)} + 8cd\gamma(L_{ji}^{12} + L_{ji}^{21}) + 2cd\rho N_{ij} \right. \\
&\quad \left. \left. + c^2(\kappa_4)_{11} U_i^{(1)} U_j^{(1)} + d^2(\kappa_4)_{22} U_i^{(2)} U_j^{(2)} + cd(\kappa_4)_{12} (U_i^{(1)} U_j^{(2)} + U_j^{(1)} U_i^{(2)}) \right) \right] \\
&\quad + \mathcal{O}_{\prec}(n^{-c(p)}),
\end{aligned}$$

for some small $c(p) > 0$, where $\widehat{V}_{ij}^{(1)}, \widehat{V}_{ij}^{(2)}, L_{ji}^{12}, L_{ji}^{21}, N_{ij}$, and $U_i^{(t)}$ are defined in (3.14) and (3.10).

Proof. This is proved by applying the CLT for resolvents (Theorem 3.3) on (5.3). However, it should be noted that while applying (3.13), we need to estimate the error term Ψ , which is given in (3.16). In addition, it should be noted that the domain of the integration in the last equality of (5.4) is changed from $[\eta_c, T]$ to $[0, \infty)$. Therefore, we need to argue that the value of the integration is negligible on the domains $[0, \eta_c)$ and (T, ∞) . Note that it is only sufficient to consider even p . Because the terms corresponding to odd p , are of lower order by (3.13). We now proceed to estimate the terms $V_{ij}^{(t)}, U_i^{(t)}, N_{ij}, L_{ij}$.

Borrowing the shorthand notations from (4.11), let us denote $m_i^{(t)} := (m^{(t)})^{z_i}(\iota\eta_i)$. Since $\Re(m^{(t)})^z(\omega) = 0$ on $\omega = \iota\eta$, we may write $m_i = \iota \Im m_i$. As a result, from (4.1), we may denote

$$(5.5) \quad u_i = -\frac{\iota \Im m_i}{\eta_i + \Im m_i}.$$

We now list down the estimates as follows;

$$\begin{aligned}
|\partial_{\eta_i}(m_i^{(t)})| &\lesssim \frac{1}{(\eta_i + (\Im m_i^{(t)})^2)^2}, \\
|\partial_{\eta_i} u_i^{(t)}| &\lesssim [\Im m_i^{(t)} + \eta_i]^{-2}, \\
|V_{ij}^{(t)}| &\lesssim \frac{[(\Im(m^{(t)z_i}(\nu\eta_i) + \eta_i) (\Im(m^{(t)z_j}(\nu\eta_j) + \eta_j))]^{-2}}{\left[|z_i - z_j|^2 + (\eta_i^{(t)} + \eta_j^{(t)}) \left(\min\{\Im m_i^{(t)}, \Im m_j^{(t)}\}\right)^2\right]^2}, \\
(5.6) \quad |U_i^{(t)}| &\lesssim \frac{1}{(\eta_i + (\Im m_i^{(t)})^2)^2}, \\
|N_{1i}| &\lesssim \frac{1}{(\eta_i + (\Im m_i^{(1)})^2)^2 (\eta_j + (\Im m_j^{(2)})^2)^2}, \\
|L_{ji}^{12}| &\lesssim \frac{1}{(\eta_i + (\Im m_i^{(2)})^2)^2 (\eta_j + (\Im m_j^{(1)})^2)^2} |\Re(\bar{z}_i z_j)|, \\
|L_{ji}^{21}| &\lesssim \frac{1}{(\eta_i + (\Im m_i^{(1)})^2)^2 (\eta_j + (\Im m_j^{(2)})^2)^2} |\Re(\bar{z}_i z_j)|.
\end{aligned}$$

Estimate of $\partial_{\eta_i} m_i^{(t)}$: By using the estimate (4.7) and asymptotes from (4.6), we obtain the following estimate;

$$\begin{aligned}
|\partial_{\eta_i}(m_i^{(t)})| &= \left| \frac{1 - \beta_i^{(t)}}{\beta_i^{(t)}} \right| \leq \frac{1}{|\beta_i^{(t)}|} = \frac{1}{|(\beta_*^{(t)})_i + 2(\Im m_i^{(t)})^2|} \\
&= \frac{1}{\left| \frac{\eta_i}{\Im m_i^{(t)} + \eta_i} + 2(\Im m_i^{(t)})^2 \right|} \lesssim \frac{1}{(\eta_i + (\Im m_i^{(t)})^2)^2},
\end{aligned}$$

where the last inequality follows from the fact that $\eta_i/(\eta_i + \Im m_i^{(t)}) \geq \eta_i/(1 + \eta_i) \gtrsim \eta_i$.

Estimate of $\partial_{\eta_i} u_i$: Differentiating both sides of (5.5) with respect to η_i , we obtain

$$\partial_{\eta_i}(u_i^{(t)}) = -\nu \frac{\eta_i \partial_{\eta_i}(\Im m_i^{(t)}) - \Im m_i^{(t)}}{(\eta_i + \Im m_i^{(t)})^2}.$$

Using (4.7), we have $|\Im m_i^{(t)}| < 1$. Moreover, using (4.3) in conjunction with the estimates (4.9) and (4.6) we obtain $|\eta_i \partial_{\eta_i} \Im m_i^{(t)}| < 1$, which gives us

$$|\partial_{\eta_i} u_i^{(t)}| \lesssim [\Im m_i^{(t)} + \eta_i]^{-2}.$$

Estimate of $V_{ij}^{(t)}$: Using the definition of $u_i^{(t)}$ from (4.1) and the equation (4.2), we notice that $-(m_i^{(t)})^2 + |z_i|^2 (u_i^{(t)})^2 = u_i^{(t)}$. Consequently, we may rewrite $V_{ij}^{(t)}$, whose expression is given in (3.14), as follows;

$$\begin{aligned}
&V_{ij}^{(t)} \\
&= \frac{1}{2} \partial_{\eta_i} \partial_{\eta_j} \log \left(1 - u_i^{(t)} u_j^{(t)} (1 - |z_i - z_j|^2 + (1 - u_i^{(t)})|z_i|^2 + (1 - u_j^{(t)})|z_j|^2) \right)
\end{aligned}$$

$$= \frac{1}{2} \frac{(\partial_{\eta_i} u_i^{(t)})(\partial_{\eta_j} u_j^{(t)})C(u_i^{(t)}, u_j^{(t)}, z_i, z_j)}{\left(1 - u_i^{(t)} u_j^{(t)} (1 - |z_i - z_j|^2 + (1 - u_i^{(t)})|z_i|^2 + (1 - u_j^{(t)})|z_j|^2)\right)^2},$$

where $C(u_i^{(t)}, u_j^{(t)}, z_i, z_j)$ is a function of $u_i^{(t)}, u_j^{(t)}, z_i, z_j$, which is uniformly bounded by a constant for all $u_i^{(t)}, u_j^{(t)}, z_i, z_j$.

We now just need to calculate a lower bound of the denominator. Using $-(m_i^{(t)})^2 + |z_i|^2(u_i^{(t)})^2 = u_i^{(t)}$ once again, we shall obtain (5.7), where the main factor to be analyzed is $1 - u_i^{(t)}$. Now, on the imaginary axis i.e., $\omega_i = \eta_i$, we have $m^{(t)}(\eta_i) = \iota \Im m^{(t)}(\eta_i)$. It should be noted that we are working on the regime where $\eta_i \in [\eta_c, T]$ and $\eta_c = n^{-1+\delta_1} > 0$. Since $\eta_i \Im m_i^{(t)} > 0$, we have $\Im m_i^{(t)} > 0$. Moreover, we also have $\Im m_i^{(t)} < 1$, which was noted down in the estimate of $\partial_{\eta_i} u_i^{(t)}$ as well. Consequently, $1 - u_i^{(t)} = \eta_i / (\eta_i + \Im m_i^{(t)}) \geq \eta_i / (1 + \eta_i)$, which implies that $\eta_i \lesssim 1 - u_i^{(t)} < 1$. Combining all, we obtain

$$(5.7) \quad \begin{aligned} & \left| 1 - u_i^{(t)} u_j^{(t)} (1 - |z_i - z_j|^2 + (1 - u_i^{(t)})|z_i|^2 + (1 - u_j^{(t)})|z_j|^2) \right| \\ &= \left| u_i^{(t)} u_j^{(t)} |z_i - z_j|^2 + (1 - u_i^{(t)})(1 - u_j^{(t)}) \right. \\ & \quad \left. - (m_i^{(t)})^2 u_j^{(t)} \left(\frac{1}{u_i^{(t)}} - 1 \right) - (m_j^{(t)})^2 u_i^{(t)} \left(\frac{1}{u_j^{(t)}} - 1 \right) \right| \\ & \geq u_i^{(t)} u_j^{(t)} |z_i - z_j|^2 + \min\{(\Im m_i^{(t)})^2, (\Im m_j^{(t)})^2\} (2 - u_i^{(t)} - u_j^{(t)}) \\ & \gtrsim u_i^{(t)} u_j^{(t)} |z_i - z_j|^2 + \min\{(\Im m_i^{(t)})^2, (\Im m_j^{(t)})^2\} (\eta_i + \eta_j). \end{aligned}$$

If both $u_i^{(t)}, u_j^{(t)}$ are bounded away from zero i.e., there exists a δ such that $u_i^{(t)}, u_j^{(t)} > \delta$, then $u_i^{(t)} u_j^{(t)} |z_i - z_j|^2 \gtrsim |z_i - z_j|^2$. Otherwise, $u_i^{(t)} u_j^{(t)} |z_i - z_j|^2 \geq 0$, but $2 - u_i^{(t)} - u_j^{(t)} \geq 1 - \delta$. So, in the worst case, we have the following bound

$$\begin{aligned} & \left| 1 - u_i^{(t)} u_j^{(t)} (1 - |z_i - z_j|^2 + (1 - u_i^{(t)})|z_i|^2 + (1 - u_j^{(t)})|z_j|^2) \right| \\ & \gtrsim |z_i - z_j|^2 + \min\{(\Im m_i^{(t)})^2, (\Im m_j^{(t)})^2\} (\eta_i + \eta_j). \end{aligned}$$

Estimate of $U_i^{(t)}$: From (5.6) using the estimate of $|\partial_{\eta_i}(m_i^{(t)})|$, we obtain the following estimate for $U_i^{(t)}$;

$$|U_i^{(t)}| = \left| \frac{i}{\sqrt{2}} \partial_{\eta_i} (m_i^{(t)})^2 \right| \lesssim |m_i^{(t)} (m_i^{(t)})'| \lesssim \frac{1}{(\eta_i + (\Im m_i^{(t)})^2)^2}.$$

Estimate of N_{ij} :

$$\begin{aligned} |N_{ij}| &= |\partial_{\eta_i} (m_i^{(1)}) \partial_{\eta_j} (m_j^{(2)}) + \partial_{\eta_i} (m_i^{(2)}) \partial_{\eta_j} (m_j^{(1)})| \\ &\lesssim \frac{1}{(\eta_i + (\Im m_i^{(1)})^2)^2 (\eta_j + (\Im m_j^{(2)})^2)^2}. \end{aligned}$$

Estimates of L_{ji}^{12} and L_{ji}^{21} :

$$|L_{ji}^{12}| = \left| \frac{m_j^{(1)} m_i^{(2)} (1 - 2(m_j^{(1)})^2 - \beta_j^{(1)}) (1 - 2(m_i^{(2)})^2 - \beta_i^{(2)}) \Re(\bar{z}_i z_j)}{\beta_j^{(1)} \beta_i^{(2)}} \right|$$

$$\begin{aligned}
&\lesssim \left| \frac{m_j^{(1)} m_i^{(2)} (1 - \beta_j^{(1)}) (1 - \beta_i^{(2)}) \Re(\bar{z}_i z_j)}{\beta_j^{(1)} \beta_i^{(2)}} \right| \\
&\lesssim \left| m_j^{(1)} m_i^{(2)} (m_j^{(1)})' (m_i^{(2)})' \Re(\bar{z}_i z_j) \right| \\
&\lesssim \frac{1}{(\eta_i + (\Im m_i^{(2)})^2)^2 (\eta_j + (\Im m_j^{(1)})^2)^2} \left| \Re(\bar{z}_i z_j) \right|.
\end{aligned}$$

Similarly,

$$|L_{ji}^{21}| \lesssim \frac{1}{(\eta_i + (\Im m_i^{(1)})^2)^2 (\eta_j + (\Im m_j^{(2)})^2)^2} \left| \Re(\bar{z}_i z_j) \right|.$$

We are now ready to prove the lemma. We first remove the region of singularity

$$Z_i := \{|1 - |z_i|^2| \leq n^{-2\nu}\} \bigcup_{j < i} \{z_j : |z_i - z_j| \leq n^{-2\nu}\}$$

from \mathbb{C} , for some small $\nu > 0$. We may now rewrite the integral of (5.4) as follows;

$$\begin{aligned}
(5.8) \quad &\frac{(-n)^p}{(2\pi\nu)^p} \prod_{i \in [p]} \int_{Z_i^c} d^2 z_i \Delta f_i(z_i) \mathbb{E} \prod_{i \in [p]} \int_{\eta_c}^T (c \langle (G^{(1)})^z(\nu\eta_i) \rangle - \mathbb{E} \langle (G^{(1)})^z(\nu\eta_i) \rangle) \\
&\quad + d \langle (G^{(2)})^z(\nu\eta_i) \rangle - \mathbb{E} \langle (G^{(2)})^z(\nu\eta_i) \rangle) d\eta_i + \mathcal{O}_{\prec} \left(\frac{n^{p\xi}}{n^\nu} \right).
\end{aligned}$$

Since the Lebesgue measure of Z_i is $\mathcal{O}(n^{-2\nu})$, the removal of the region Z_i introduced an error of $\mathcal{O}_{\prec}(n^{p\xi}/n^\nu)$, for any very small $\xi > 0$, which is a consequence of (5.1).

Now, by using (5.6) we have the following estimates over Z_i^c in the regime $\eta_i \in [0, \eta_c]$, where $\eta_c = n^{-1+\delta_1}$;

$$\begin{aligned}
(5.9) \quad &|V_{ij}^{(t)}| \lesssim \frac{n^{12\nu}}{(1 + \eta_i^2)(1 + \eta_j^2)}, \\
&|U_i^{(t)}| \lesssim \frac{n^{4\nu}}{1 + \eta_i^2}, \\
&|N_{ij}| \lesssim \frac{n^{8\nu}}{(1 + \eta_i^2)(1 + \eta_j^2)}, \\
&|L_{ji}^{12}| \lesssim \frac{n^{8\nu}}{(1 + \eta_i^2)(1 + \eta_j^2)}, \\
&|L_{ji}^{21}| \lesssim \frac{n^{8\nu}}{(1 + \eta_i^2)(1 + \eta_j^2)}.
\end{aligned}$$

Finally, using (3.13) in (5.8), we obtain the integrand displayed in the last equality of (5.4). However, the domain of the integration in (5.8) is not same as (5.4). We can add back the domains $(0, \eta_c)$ and (T, ∞) using the estimates (5.9) and the first estimate in (5.1). Lastly, the error Ψ from (3.16) is bounded by $\mathcal{O}(n^{\xi p + 2p\nu - \delta_1/2})$ on $\cap_i Z_i^c$. This concludes the lemma and yields the following equation

$$\begin{aligned}
& \frac{1}{(2\pi\iota)^p} \prod_{i \in [p]} \int_{Z_i^c \cap \widehat{Z}_i^c} d^2 z_i \Delta f_i \\
(5.10) \quad & \times \sum_{P \in \Pi_p\{i,j\} \in P} \int_0^\infty \int_0^\infty - \left(\frac{c^2 \widehat{V}_{ij}^{(1)} + d^2 \widehat{V}_{ij}^{(2)} + 8cd\gamma(L_{ji}^{12} + L_{ji}^{21}) + 2cd\rho N_{ij}}{8\pi^2} \right. \\
& \left. + \frac{c^2(\kappa_4)_{11} U_i^{(1)} U_j^{(1)} + d^2(\kappa_4)_{22} U_i^{(2)} U_j^{(2)} + cd(\kappa_4)_{12} (U_i^{(1)} U_j^{(2)} + U_j^{(1)} U_i^{(2)})}{8\pi^2} \right) d\eta_i d\eta_j \\
& + \mathcal{O}_{\prec} \left(\frac{n^{p\xi}}{n^\nu} + \frac{n^{12\nu p + \delta_1}}{n} + \frac{n^{\xi p + 2p\nu}}{n^{\delta_1/2}} \right).
\end{aligned}$$

Based on δ_1 , we can choose ν in such a way that the above error vanishes asymptotically. \square

6. CENTRAL LIMIT THEOREM FOR RESOLVENTS

In this section, we shall present the the proof of the CLT for the resolvents, which is stated in the Theorem 3.3. Recall that we are considering the resolvents $G_i^{(t)} = (G^{(t)})^{z_i}(\iota\eta_i)$, $1 \leq i \leq p$ at p different points on the imaginary axis. In addition, recall that $\Upsilon_i := c\langle G_i^{(1)} - \mathbb{E}(G_i^{(1)}) \rangle + d\langle G_i^{(2)} - \mathbb{E}(G_i^{(2)}) \rangle$. Before proceeding to the computation of $\mathbb{E}[\prod_{i=1}^p \Upsilon_i]$, we give an estimate of $\mathbb{E}[(c(G^{(1)})^z + d(G^{(2)})^z)(\iota\eta)]$. From the local law, we get the first order approximation $\mathbb{E}[\langle cG^{(1)} + dG^{(2)} \rangle] \sim \langle cM^{(1)} + dM^{(2)} \rangle$. The second order correction of $\mathbb{E}[\langle cG^{(1)} + dG^{(2)} \rangle]$ is of the order of $1/n$. This is formalized in the following lemma.

Lemma 6.1. *For $(\kappa_4)_{tt} = \kappa(\chi^{(t)}, \bar{\chi}^{(t)}, \chi^{(t)}, \bar{\chi}^{(t)}) \neq 0$, where $\chi^{(t)}$ is same as defined in Condition 3.1, we have*

$$(6.1) \quad \mathbb{E}\langle cG^{(1)} + dG^{(2)} \rangle = \langle cM^{(1)} + dM^{(2)} \rangle + c\xi^{(1)} + d\xi^{(2)} + Error_{exp},$$

where

$$\begin{aligned}
& \xi^{(t)} \\
& = \begin{cases} -\frac{i(\kappa_4)_{tt}}{4n} \partial_\eta((m^{(t)})^4), & \text{when } X^{(t)} \text{ in Condition 3.1 is } \mathbb{C}\text{-valued,} \\ \frac{i}{4n} \partial_\eta \left(1 - (u^{(t)})^2 + 2(u^{(t)})^3 |z|^2 - (u^{(t)})^2 (z^2 + \bar{z}^2) \right) \\ -\frac{i(\kappa_4)_{tt}}{4n} \partial_\eta((m^{(t)})^4), & \text{when } X^{(t)} \text{ in Condition 3.1 is } \mathbb{R}\text{-valued,} \end{cases}
\end{aligned}$$

and

$$\begin{aligned}
& Error_{exp} \\
& = \begin{cases} \mathcal{O}_{\prec} \left(\frac{1}{|1-|z||} \left(\frac{1}{n^{3/2}\eta} + \frac{1}{(n\eta)^2} \right) \right), & \text{when } X^{(t)} \text{ is } \mathbb{C}\text{-valued,} \\ \mathcal{O}_{\prec} \left(\left(\frac{1}{|1-|z||} + \frac{1}{\Im z^2} \right) \left(\frac{1}{n^{3/2}\eta} + \frac{1}{(n\eta)^2} \right) \right), & \text{when } X^{(t)} \text{ is } \mathbb{R}\text{-valued.} \end{cases}
\end{aligned}$$

Before proving the lemma, we note down the cumulant expansion as follows [23, Proposition 7.5].

Result 6.2 (Cumulant Expansion). (1) For a general scalar random variable X , the cumulant expansion is given as follows;

$$\mathbb{E}[Xf(X)] = \sum_l \frac{\kappa_{l+1}(X)}{l!} \mathbb{E}[f^{(l)}(X)],$$

where $\kappa_{l+1}(X)$ represents the $(l+1)$ -th cumulant of X , and $f^{(l)}$ denotes the l -th derivative of a smooth function f .

(2) In the multidimensional case, the cumulant expansion is defined as follows. Consider a collection of random variables X_1, X_2, \dots, X_p , and let κ denote their joint cumulants. For a smooth function $f : \mathbb{R}^p \rightarrow \mathbb{R}$, the cumulant expansions are given by

$$\begin{aligned} & \mathbb{E}[X_i f(X_1, X_2, \dots, X_p)] \\ &= \sum_{l \geq 0} \sum_{i_1, \dots, i_l=1}^p \frac{\kappa(X_i, X_{i_1}, \dots, X_{i_l})}{l!} \mathbb{E}[\partial_{i_1} \partial_{i_2} \dots \partial_{i_l} f(X_1, X_2, \dots, X_p)], \end{aligned}$$

where $\partial_{i_1} \partial_{i_2} \dots \partial_{i_l}$ represents the partial derivatives with respect to $X_{i_1}, X_{i_2}, \dots, X_{i_l}$.

Proof of Lemma 6.1. Using the stability operator $\mathcal{B}^{(t)}$ from (4.22), let us define $A^{(t)} := (((\mathcal{B}^{(t)})^*)^{-1}[\mathbb{I}])^* M^{(t)}$. Now using (4.14), we have

$$\begin{aligned} (6.2) \quad & \langle cG^{(1)} + dG^{(2)} - cM^{(1)} - dM^{(2)} \rangle \\ &= c\langle G^{(1)} - M^{(1)} \rangle + d\langle G^{(2)} - M^{(2)} \rangle \\ &= c\langle \mathbb{I}, (\mathcal{B}^{(1)})^{-1} \mathcal{B}^{(1)} [G^{(1)} - M^{(1)}] \rangle + d\langle \mathbb{I}, (\mathcal{B}^{(2)})^{-1} \mathcal{B}^{(2)} [G^{(2)} - M^{(2)}] \rangle \\ &= -c\langle (M^{(1)})^* ((\mathcal{B}^{(1)})^*)^{-1} [\mathbb{I}], \underline{W^{(1)}G^{(1)}} \rangle \\ &\quad + c\langle (M^{(1)})^* ((\mathcal{B}^{(1)})^*)^{-1} [\mathbb{I}], \mathcal{S}G^{(1)} - M^{(1)} \rangle \\ &\quad - d\langle (M^{(2)})^* ((\mathcal{B}^{(2)})^*)^{-1} [\mathbb{I}], \underline{W^{(2)}G^{(2)}} \rangle \\ &\quad + d\langle (M^{(2)})^* ((\mathcal{B}^{(2)})^*)^{-1} [\mathbb{I}], \mathcal{S}G^{(2)} - M^{(2)} \rangle \\ &= -c\langle (M^{(1)})^* ((\mathcal{B}^{(1)})^*)^{-1} [\mathbb{I}], \underline{W^{(1)}G^{(1)}} \rangle + \mathcal{O}_{\prec} \left(\frac{\|((\mathcal{B}^{(1)})^*)^{-1} [\mathbb{I}]\|}{(n\eta)^2} \right) \\ &\quad - d\langle (M^{(2)})^* ((\mathcal{B}^{(2)})^*)^{-1} [\mathbb{I}], \underline{W^{(2)}G^{(2)}} \rangle + \mathcal{O}_{\prec} \left(\frac{\|((\mathcal{B}^{(2)})^*)^{-1} [\mathbb{I}]\|}{(n\eta)^2} \right) \\ (6.3) \quad &= -\langle c\underline{W^{(1)}G^{(1)}}A^{(1)} + d\underline{W^{(2)}G^{(2)}}A^{(2)} \rangle + \mathcal{O}_{\prec} \left(\frac{\|((\mathcal{B}^{(1)})^*)^{-1} [\mathbb{I}]\|}{(n\eta)^2} \right) \\ &\quad + \mathcal{O}_{\prec} \left(\frac{\|((\mathcal{B}^{(2)})^*)^{-1} [\mathbb{I}]\|}{(n\eta)^2} \right). \end{aligned}$$

Now, we use the cumulant expansion from the Result 6.2 to evaluate $\mathbb{E}[\langle \underline{W^{(t)}G^{(t)}}A^{(t)} \rangle]$. This will lead us to calculate the derivatives of $G^{(t)}A^{(t)}$ with respect to the entries of $W^{(t)}$. This calculation is demonstrated below, where for brevity we use the notation $\Delta^{ba} := e_b^t e_a$, which is essentially a $2n \times 2n$ matrix with all zero entries except the (ba) th entry being equal to 1. Here, we introduce a shorthand notation for joint cumulants of the entries of the matrices

$W^{(1)}$ and $W^{(2)}$. We shall denote the joint cumulant $\kappa(w_{ab}^{(t)}, w_{ij}^{(s)}, \dots)$ as $\kappa((ab)_t, (ij)_s, \dots)$, where $t, s \in \{1, 2\}$.

For $\alpha = (\alpha_1, \dots, \alpha_k)$, we note that $\kappa((ba)_t, \alpha)$ is non-zero only for $\alpha \in \{(ab)_t, (ba)_t\}^k$. The derivative ∂_α denotes the derivative with respect to $w_{\alpha_1}^{(t)}, w_{\alpha_2}^{(t)}, \dots, w_{\alpha_k}^{(t)}$.

$$\begin{aligned}
(6.4) \quad & \mathbb{E} [\langle \underline{W^{(t)} G^{(t)} A^{(t)}} \rangle] \\
&= \mathbb{E} [\langle W^{(t)} G^{(t)} A^{(t)} \rangle] + \mathbb{E} [\langle \mathcal{S}[G^{(t)}] G^{(t)} A^{(t)} \rangle] \\
&= \mathbb{E} \left[\sum_{ab} w_{ba}^{(t)} \langle \Delta^{ba} G^{(t)} A^{(t)} \rangle \right] + \mathbb{E} [\langle \mathcal{S}[G^{(t)}] G^{(t)} A^{(t)} \rangle] \\
&= \sum_{k \geq 0} \sum_{ab} \sum_{\alpha \in \{(ab)_t, (ba)_t\}^k} \frac{\kappa((ba)_t, \alpha)}{k!} \mathbb{E} [\partial_\alpha \langle \Delta^{ba} G^{(t)} A^{(t)} \rangle] + \mathbb{E} [\langle \mathcal{S}[G^{(t)}] G^{(t)} A^{(t)} \rangle] \\
&= \sum_{ab} \frac{\kappa((ba)_t, (ab)_t)}{1!} \mathbb{E} [\partial_{(ab)_t} \langle \Delta^{ba} G^{(t)} A^{(t)} \rangle] \\
&\quad + \sum_{ab} \frac{\kappa((ba)_t, (ba)_t)}{1!} \mathbb{E} [\partial_{(ba)_t} \langle \Delta^{ba} G^{(t)} A^{(t)} \rangle] + \mathbb{E} [\langle \mathcal{S}[G^{(t)}] G^{(t)} A^{(t)} \rangle] \\
&\quad + \sum_{k \geq 2} \sum_{ab} \sum_{\alpha \in \{(ab)_t, (ba)_t\}^k} \frac{\kappa((ba)_t, \alpha)}{k!} \mathbb{E} [\partial_\alpha \langle \Delta^{ba} G^{(t)} A^{(t)} \rangle] \\
&= -\frac{1}{2n^2} \sum'_{ab} \mathbb{E} [G_{aa}^{(t)} (G^{(t)} A^{(t)})_{bb}] + \sum_{ab} \frac{\kappa((ba)_t, (ba)_t)}{1!} \mathbb{E} [\partial_{(ba)_t} \langle \Delta^{ba} G^{(t)} A^{(t)} \rangle] \\
&\quad + \sum_{k \geq 2} \sum_{ab} \sum_{\alpha \in \{(ab)_t, (ba)_t\}^k} \frac{\kappa((ba)_t, \alpha)}{k!} \mathbb{E} [\partial_\alpha \langle \Delta^{ba} G^{(t)} A^{(t)} \rangle] \\
&\quad + \frac{1}{2n^2} \sum'_{ab} \mathbb{E} [G_{aa}^{(t)} (G^{(t)} A^{(t)})_{bb}] \\
&= \sum_{ab} \frac{\kappa((ba)_t, (ba)_t)}{1!} \mathbb{E} [\partial_{(ba)_t} \langle \Delta^{ba} G^{(t)} A^{(t)} \rangle] \\
&\quad + \sum_{k \geq 2} \sum_{ab} \sum_{\alpha \in \{(ab)_t, (ba)_t\}^k} \frac{\kappa((ba)_t, \alpha)}{k!} \mathbb{E} [\partial_\alpha \langle \Delta^{ba} G^{(t)} A^{(t)} \rangle] \\
&= -\frac{1}{n} \sum'_{ab} \mathbb{E} [\langle \Delta^{ba} G^{(t)} \Delta^{ba} G^{(t)} A^{(t)} \rangle] \\
&\quad + \sum_{k \geq 2} \sum_{ab} \sum_{\alpha \in \{(ab)_t, (ba)_t\}^k} \frac{\kappa((ba)_t, \alpha)}{k!} \mathbb{E} [\partial_\alpha \langle \Delta^{ba} G^{(t)} A^{(t)} \rangle] \\
&=: -\frac{1}{n} F_1^{(t)} + F_{k \geq 2}^{(t)},
\end{aligned}$$

where, the summation notation \sum'_{ab} is defined as

$$\sum'_{ab} = \sum_{a \leq n} \sum_{b > n} + \sum_{a > n} \sum_{b \leq n},$$

which is not leaving out any terms of the double sum \sum_{ab} , though it seems so. This is a simple consequence of the structure of $W^{(t)} = \begin{bmatrix} 0 & X^{(t)} \\ X^{(t)*} & 0 \end{bmatrix}_{2n \times 2n}$.

Moreover, according to the scaling and the definition of the cumulants, we notice that

$$\kappa((ba)_t, (ba)_t) = \begin{cases} 0 & \text{if } X^{(t)} \text{ is } \mathbb{C}\text{-valued} \\ \frac{1}{n} & \text{if } X^{(t)} \text{ is } \mathbb{R}\text{-valued,} \end{cases}$$

and

$$(6.5) \quad |\kappa(\alpha_1, \alpha_2, \dots, \alpha_k)| \lesssim \frac{1}{n^{k/2}}.$$

Before evaluating the terms $F_1^{(t)}, F_{k \geq 2}^{(t)}$, let us define $2n \times 2n$ ordered block matrices E_1 and E_2 as follows;

$$(6.6) \quad E_1 = \begin{pmatrix} \mathbb{I} & 0 \\ 0 & 0 \end{pmatrix}, \text{ and } E_2 = \begin{pmatrix} 0 & 0 \\ 0 & \mathbb{I} \end{pmatrix},$$

where \mathbb{I} is $n \times n$ identity matrix. We shall follow the convention

$$\langle AE_1BE_2 + AE_2BE_1 \rangle =: \langle AEBE^\dagger \rangle,$$

for any $2n \times 2n$ matrices A and B . Now, let us move to evaluate the terms $F_1^{(t)}, F_{k \geq 2}^{(t)}$.

We shall use the convention $((G^{(t)})^z)^t = (G^{(t)})^{\bar{z}}$, along with Theorem 4.2 (the local law for the product of resolvents), and the bound on $|(\hat{\beta}^{(t)})_*|$ from Lemma 4.4 to obtain the following estimate;

$$(6.7) \quad \begin{aligned} & \left| \langle (G^{(t)})^z A^{(t)} E (G^{(t)})^{\bar{z}} E^\dagger - (M^{(t)})_{A^{(t)}E}^{z, \bar{z}} E^\dagger \rangle \right| \\ & \lesssim \frac{C_\epsilon \|A^{(t)} E\| \|E^\dagger\| n^\xi}{n\eta^2 |z - \bar{z}|^2} \left(\eta^{1/12} + \frac{\eta^{1/4}}{|z - \bar{z}|^2} + \frac{1}{\sqrt{n\eta}} + \frac{1}{(|z - \bar{z}|^2 |n\eta|)^{1/4}} \right) \\ & = \mathcal{O}_\prec \left(\frac{1}{|z - \bar{z}|^2 |\beta^{(t)}| (n\eta)^2} \right). \end{aligned}$$

Estimate of $F_1^{(t)}$: Using (6.7) in the fourth equality and (4.17) in the the fifth equality in the following calculation, we conclude

$$(6.8) \quad \begin{aligned} & F_1^{(t)} \\ & = \sum'_{ab} \mathbb{E} [\langle \Delta^{ba} G^{(t)} \Delta^{ba} G^{(t)} A^{(t)} \rangle] \\ & = \mathbb{E} [\langle G^{(t)} A^{(t)} E (G^{(t)})^t E^\dagger \rangle] \\ & = \mathbb{E} [\langle (G^{(t)})^z A^{(t)} E (G^{(t)})^{\bar{z}} E^\dagger \rangle] \\ & = \langle (M^{(t)})_{A^{(t)}E}^{z, \bar{z}} E^\dagger \rangle + \mathcal{O}_\prec \left(\frac{1}{|z - \bar{z}|^2 |\beta^{(t)}| (n\eta)^2} \right) \\ & = \langle (\mathbb{I} - (M^{(t)})^z \mathcal{S}[\cdot] (M^{(t)})^{\bar{z}})^{-1} [(M^{(t)})^z A^{(t)} E (M^{(t)})^{\bar{z}}] E^\dagger \rangle \\ & \quad + \mathcal{O}_\prec \left(\frac{1}{|z - \bar{z}|^2 |\beta^{(t)}| (n\eta)^2} \right) \end{aligned}$$

$$\begin{aligned}
&= m^{(t)} \frac{(m^{(t)})^4 + (m^{(t)})^2(u^{(t)})^2|z|^2 - 2(u^{(t)})^4|z|^4 + 2(u^{(t)})^2(x^2 - y^2)}{(1 - (m^{(t)})^2 - (u^{(t)})^2|z|^2)(1 + (u^{(t)})^4|z|^4 - (m^{(t)})^4 - 2(u^{(t)})^2(x^2 - y^2))} \\
&\quad + \mathcal{O}_{\prec} \left(\frac{1}{|z - \bar{z}|^2 |\beta^{(t)}| (n\eta)^2} \right) \\
&= -\frac{i(u^{(t)})'}{2} \frac{u^{(t)} - 3|z|^2(u^{(t)})^2 + 2u^{(t)}(x^2 - y^2)}{1 - (u^{(t)})^2 + 2(u^{(t)})^3|z|^2 - 2(u^{(t)})^2(x^2 - y^2)} \\
&\quad + \mathcal{O}_{\prec} \left(\frac{1}{|z - \bar{z}|^2 |\beta^{(t)}| (n\eta)^2} \right) \\
&= \frac{i}{4} \partial_{\eta} \log \left(1 - (u^{(t)})^2 + (u^{(t)})^3|z|^2 - (u^{(t)})^2(z^2 + \bar{z}^2) \right) \\
&\quad + \mathcal{O}_{\prec} \left(\frac{1}{|z - \bar{z}|^2 |\beta^{(t)}| (n\eta)^2} \right),
\end{aligned}$$

where $z = x + iy$. The sixth equality follows from the direct calculation of the inverse operator from the previous step. The second last step follows from the facts $(m^{(t)})^z = (m^{(t)})^{\bar{z}}$, $(u^{(t)})^z = (u^{(t)})^{\bar{z}}$, and the following identities,

$$\begin{aligned}
|z|^2(u^{(t)})^2 - (m^{(t)})^2 &= u^{(t)}, \\
(u^{(t)})' &= \partial_{\eta} u^{(t)} = \frac{2iu^{(t)}m^{(t)}}{1 + u^{(t)} - 2|z|^2(u^{(t)})^2}.
\end{aligned}$$

Estimate of $F_{k \geq 2}^{(t)}$: Let us first consider the case when $k = 2$, and let $\alpha = (\alpha_1, \alpha_2)$. Then

$$\begin{aligned}
&\sum_{ab} \sum_{\alpha} \frac{\kappa((ba)_t, \alpha_1, \alpha_2)}{2!} \mathbb{E} [\partial_{\alpha_1, \alpha_2} \langle \Delta^{ba} G^{(t)} A^{(t)} \rangle] \\
&= \sum_{\alpha} \frac{\kappa((ba)_t, \alpha_1, \alpha_2)}{2!} \sum'_{ab} \mathbb{E} [\langle \Delta^{ba} G^{(t)} \Delta^{\alpha_1} G^{(t)} \Delta^{\alpha_2} G^{(t)} A^{(t)} \rangle].
\end{aligned}$$

To obtain a significant contribution, we need more diagonal terms than off-diagonal ones (as explained in Remark 6.4). Here, by parity, at least one $G^{(t)}$ factor in $\Delta^{ba} G^{(t)} \Delta^{\alpha_1} G^{(t)} \Delta^{\alpha_2} G^{(t)} A^{(t)}$ is off-diagonal. For example, taking $(\alpha_1, \alpha_2) = ((ba)_t, (ab)_t)$ and using $G^{(t)} = M^{(t)} + G^{(t)} - M^{(t)}$ along with the local law bound of $G^{(t)} - M^{(t)}$ from Theorem 4.1, we obtain

$$\begin{aligned}
(6.9) \quad &\frac{1}{n^{1+3/2}} \sum'_{ab} \mathbb{E} [(G^{(t)})_{ab} (G^{(t)})_{aa} (G^{(t)} A^{(t)})_{bb}] \\
&= \frac{1}{n^{5/2}} \sum'_{ab} \mathbb{E} \left[(G^{(t)})_{ab} \left(m^{(t)} (M^{(t)} A^{(t)})_{bb} + m^{(t)} ((G^{(t)} - M^{(t)}) A^{(t)})_{bb} \right. \right. \\
&\quad \left. \left. + (M^{(t)} A^{(t)})_{bb} (G^{(t)} - M^{(t)})_{aa} + (G^{(t)} - M^{(t)})_{aa} ((G^{(t)} - M^{(t)}) A^{(t)})_{bb} \right) \right] \\
&= \frac{1}{n^{5/2}} \sum'_{ab} \mathbb{E} \left[m^{(t)} (M^{(t)})_{ab} (M^{(t)} A^{(t)})_{bb} + m^{(t)} (G^{(t)} - M^{(t)})_{ab} (M^{(t)} A^{(t)})_{bb} \right. \\
&\quad \left. + m^{(t)} (M^{(t)})_{ab} ((G^{(t)} - M^{(t)}) A^{(t)})_{bb} + m^{(t)} (G^{(t)} - M^{(t)})_{ab} ((G^{(t)} - M^{(t)}) A^{(t)})_{bb} \right. \\
&\quad \left. + (M^{(t)})_{ab} (M^{(t)} A^{(t)})_{bb} (G^{(t)} - M^{(t)})_{aa} + (G^{(t)} - M^{(t)})_{ab} (M^{(t)} A^{(t)})_{bb} (G^{(t)} - M^{(t)})_{aa} \right]
\end{aligned}$$

$$\begin{aligned}
& + (M^{(t)})_{ab}(G^{(t)} - M^{(t)})_{aa}((G^{(t)} - M^{(t)})A^{(t)})_{bb} \\
& + (G^{(t)} - M^{(t)})_{ab}(G^{(t)} - M^{(t)})_{aa}((G^{(t)} - M^{(t)})A^{(t)})_{bb} \Big] \\
& = \frac{1}{n^{5/2}} \mathbb{E} \left[m^{(t)}(M^{(t)}A^{(t)})_{(n+1, n+1)} \langle E\mathbf{1}, M^{(t)}E^\dagger\mathbf{1} \rangle + m^{(t)}(M^{(t)}A^{(t)})_{bb} \langle \mathbf{e}_a, (G^{(t)} - M^{(t)})\mathbf{e}_b \rangle \right. \\
& \quad + m^{(t)}(M^{(t)})_{ab}((G^{(t)} - M^{(t)})A^{(t)})_{bb} + m^{(t)}(G^{(t)} - M^{(t)})_{ab}((G^{(t)} - M^{(t)})A^{(t)})_{bb} \\
& \quad + (M^{(t)})_{ab}(M^{(t)}A^{(t)})_{bb}(G^{(t)} - M^{(t)})_{aa} + (G^{(t)} - M^{(t)})_{ab}(M^{(t)}A^{(t)})_{bb}(G^{(t)} - M^{(t)})_{aa} \\
& \quad + (M^{(t)})_{ab}(G^{(t)} - M^{(t)})_{aa}((G^{(t)} - M^{(t)})A^{(t)})_{bb} \\
& \quad \left. + (G^{(t)} - M^{(t)})_{ab}(G^{(t)} - M^{(t)})_{aa}((G^{(t)} - M^{(t)})A^{(t)})_{bb} \right] \\
& = \mathcal{O}_{\prec} \left(\frac{1}{n^{5/2}} \left(\frac{\sqrt{2n}\sqrt{2n}}{|\beta^{(t)}|(1+\eta)} + \frac{1}{|\beta^{(t)}|} \frac{n^2}{\sqrt{n\eta}} + \frac{n}{\eta|\beta^{(t)}|} + \frac{1}{\sqrt{n\eta}} \frac{n}{\eta|\beta^{(t)}|} \right. \right. \\
& \quad \left. \left. + \frac{n}{(1+\eta)} \frac{1}{|\beta^{(t)}|} \frac{1}{\sqrt{n\eta}} + \frac{1}{\sqrt{n\eta}} \frac{1}{|\beta^{(t)}|} \frac{n}{\eta} + \frac{1}{\sqrt{n\eta}} \frac{n}{\eta|\beta^{(t)}|} + \frac{1}{\sqrt{n\eta}} \frac{1}{\sqrt{n\eta}} \frac{n}{\eta|\beta^{(t)}|} \right) \right) \\
& = \mathcal{O}_{\prec} \left(\frac{1}{|\beta^{(t)}|} \left(\frac{1}{n^{3/2}(1+\eta)} + \frac{1}{n^2\eta^{3/2}} \right) \right).
\end{aligned}$$

Thus, we can bound all $k = 2$ terms in $F_{k \geq 2}^{(t)}$ by

$$|\beta^{(t)}|^{-1} \times (n^{-3/2}(1+\eta)^{-1} + n^{-2}\eta^{-3/2}).$$

In the following paragraph, we argue that the terms corresponding to $k \geq 4$ have insignificant contribution to $F_{k \geq 2}^{(t)}$, and the only significant contribution comes from the terms corresponding to $k = 3$. When $k = 4$, proceeding as (6.9), we shall get terms of the form $n^{-7/2} \sum'_{ab} \mathbb{E}[(G^{(t)})_{ab}(G^{(t)})_{aa}(G^{(t)})_{bb}(G^{(t)})_{aa}(G^{(t)}A^{(t)})_{bb}]$. Now, we can apply individual naive bounds on the each $G^{(t)}$ term, which will yield the bound of $\mathcal{O}_{\prec}(|\beta^{(t)}|^{-1}n^{-7/2} \cdot n^2) = \mathcal{O}_{\prec}(|\beta^{(t)}|^{-1}n^{-3/2})$. For $k \geq 4$, the pre-factor will be even smaller than $n^{-7/2}$. Thus, the terms do not contribute to $F_{k \geq 2}^{(t)}$.

Now, we estimate the contribution of the terms in $F_{k \geq 2}^{(t)}$ when $k = 3$. We begin with the observation from the Condition 3.1 that $x_{ab}^{(t)} \stackrel{\text{i.i.d.}}{\sim} n^{-1/2}\chi^{(t)}$, implying $w_{(ab)}^{(t)} \sim n^{-1/2}\chi^{(t)}$ for $a \leq n, b > n$ or $a > n, b \leq n$. Therefore

$$\kappa((ba)_t, (ab)_t, (ab)_s, (ba)_s) = \frac{1}{(n^{1/2})^4} \kappa(\chi^{(t)}, \bar{\chi}^{(t)}, \chi^{(s)}, \bar{\chi}^{(s)}) = \frac{1}{n^2} (\kappa_4)_{ts}.$$

When $k = 3$, we get a product of three $G^{(t)}$ terms and one $G^{(t)}A^{(t)}$ term in the expansion. In that scenario, all the choices of α s except $\alpha = ((ab)_t, (ba)_t, (ba)_t)$ will have a off-diagonal entry of $G^{(t)}$ in the product. When $\alpha = ((ab)_t, (ba)_t, (ba)_t)$, we get the term

$$\mathbb{E}[(G^{(t)})_{aa}(G^{(t)})_{bb}(G^{(t)})_{aa}(G^{(t)}A^{(t)})_{bb}],$$

which results in the following contribution

$$\begin{aligned}
(6.10) \quad & -\frac{(\kappa_4)_{tt}}{2n^3} \sum'_{ab} (M^{(t)})_{aa} (M^{(t)})_{bb} (M^{(t)})_{aa} (M^{(t)} A^{(t)})_{bb} \\
& = -\frac{(\kappa_4)_{tt}}{n} \langle M^{(t)} \rangle^3 \langle M^{(t)} A^{(t)} \rangle \\
& = \frac{i(\kappa_4)_{tt}}{4n} \partial_\eta ((m^{(t)})^4),
\end{aligned}$$

where we have used

$$\langle M^{(t)} A^{(t)} \rangle = \frac{1 - \beta^{(t)}}{\beta^{(t)}} = -i \partial_\eta m^{(t)}.$$

Thus, combining (6.4) (6.8), and (6.10) we have

$$\begin{aligned}
& \mathbb{E} [\langle \underline{W^{(t)} G^{(t)} A^{(t)}} \rangle] \\
& = -\frac{i}{4n} \partial_\eta \log (1 - (u^{(t)})^2 + 2(u^{(t)})^3 |z|^2 - (u^{(t)})^2 (z^2 + \bar{z}^2)) \\
& \quad + \mathcal{O}_\prec \left(\frac{1}{|z - \bar{z}|^2 |\beta^{(t)}| (n\eta)^2} \right) + \frac{i(\kappa_4)_{tt}}{4n} \partial_\eta ((m^{(t)})^4) \\
& \quad + \mathcal{O}_\prec \left(\frac{1}{|\beta^{(t)}| n^{3/2} (1 + \eta)} + \frac{1}{|\beta^{(t)}| n^2 \eta^{3/2}} \right).
\end{aligned}$$

Finally,

$$\begin{aligned}
& \mathbb{E} [\langle G^{(t)} - M^{(t)} \rangle] \\
& = \mathbb{E} [\langle \underline{-W^{(t)} G^{(t)} A^{(t)}} \rangle] + \mathcal{O}_\prec \left(\frac{\|((\mathcal{B}^{(t)})^*)^{-1}\| [\mathbb{I}]}{(n\eta)^2} \right) \\
& = -\frac{i(\kappa_4)_{tt}}{4n} \partial_\eta ((m^{(t)})^4) + \frac{i}{4n} \partial_\eta \log (1 - (u^{(t)})^2 + 2(u^{(t)})^3 |z|^2 - (u^{(t)})^2 (z^2 + \bar{z}^2)) \\
& \quad + \mathcal{O}_\prec \left(\frac{1}{|\beta^{(t)}| n^{3/2} (1 + \eta)} + \frac{1}{|\beta^{(t)}| n^2 \eta^{3/2}} \right) \\
& \quad + \mathcal{O}_\prec \left(\frac{1}{|z - \bar{z}|^2 |\beta^{(t)}| (n\eta)^2} \right) + \mathcal{O}_\prec \left(\frac{1}{|\beta^{(t)}| (n\eta)^2} \right) \\
& =: \xi^{(t)} + \mathcal{O}_\prec \left(\left(\frac{1}{|1 - |z||} + \frac{1}{|\Im z|^2} \right) \left(\frac{1}{n^{3/2} (1 + \eta)} + \frac{1}{(n\eta)^2} \right) \right) \\
& =: \xi^{(t)} + \text{Error}_{exp}.
\end{aligned}$$

Therefore,

$$(6.11) \quad \mathbb{E} [\langle cG^{(1)} + dG^{(2)} - cM^{(1)} - dM^{(2)} \rangle] = c\xi^{(1)} + d\xi^{(2)} + \text{Error}_{exp},$$

which completes the proof of Lemma 6.1. \square

Now, we are ready to present the proof of the Theorem 3.3. Using Lemma 6.1 along with the equation (6.2), $\mathbb{E} \left[\prod_{i=1}^p \{c \langle G_i^{(1)} - \mathbb{E}(G_i^{(1)}) \rangle + d \langle G_i^{(2)} - \mathbb{E}(G_i^{(2)}) \rangle\} \right]$ is equivalent to

computing

$$\begin{aligned} & \mathbb{E} \left[\prod_{i \in [p]} \langle -c \underline{W^{(1)} G_i^{(1)} A_i^{(1)}} - c \xi_i^{(1)} - d \underline{W^{(2)} G_i^{(2)} A_i^{(2)}} - d \xi_i^{(2)} \rangle \right] \\ & + \mathcal{O}_{\prec} \left(\frac{\psi^{(1)}}{n \eta_i} \right) + \mathcal{O}_{\prec} \left(\frac{\psi^{(2)}}{n \eta_i} \right), \end{aligned}$$

where

$$(6.12) \quad \psi^{(t)} = \prod_{i \in [p]} \left(\frac{1}{|\beta_i^{(t)}|} + \frac{1}{(\Im z_i)^2} \right) \frac{1}{n \eta_i} \leq \prod_{i \in [p]} \left(\frac{1}{|1 - |z_i||} + \frac{1}{(\Im z_i)^2} \right) \frac{1}{n \eta_i},$$

and $\psi^{(1)} + \psi^{(2)} \leq \psi$, where ψ is defined in (6.30).

Recall that

$$\Upsilon_i = c \langle -\underline{W^{(1)} G_i^{(1)} A_i^{(1)}} - \xi_i^{(1)} \rangle + d \langle -\underline{W^{(2)} G_i^{(2)} A_i^{(2)}} - \xi_i^{(2)} \rangle.$$

We now begin with the expansion of $\mathbb{E}[\prod_{i \in [p]} \Upsilon_i]$ as follows;

$$\begin{aligned} (6.13) \quad & \mathbb{E} \left[\prod_{i \in [p]} \Upsilon_i \right] \\ & = \mathbb{E} \left[\Upsilon_1 \prod_{i \neq 1} \Upsilon_i \right] \\ & = \mathbb{E} \left[c \langle -\underline{W^{(1)} G_1^{(1)} A_1^{(1)}} - \xi_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i \right] + \mathbb{E} \left[d \langle -\underline{W^{(2)} G_1^{(2)} A_1^{(2)}} - \xi_1^{(2)} \rangle \prod_{i \neq 1} \Upsilon_i \right] \\ & =: I^{(1)} + I^{(2)}. \end{aligned}$$

Now, we shall evaluate $I^{(1)}$.

$$\begin{aligned} (6.14) \quad & I^{(1)} \\ & = \mathbb{E} \left[c \langle -\underline{W^{(1)} G_1^{(1)} A_1^{(1)}} - \xi_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i \right] \\ & = \mathbb{E} \left[c \langle -\underline{W^{(1)} G_1^{(1)} A_1^{(1)}} - \mathcal{S}[G_1^{(1)}] G_1^{(1)} A_1^{(1)} - \xi_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i \right] \\ & = -c \langle \xi_1^{(1)} \rangle \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] + c \mathbb{E} \left[\langle -\mathcal{S}[G_1^{(1)}] G_1^{(1)} A_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i \right] \\ & \quad + c \mathbb{E} \left[\langle -\underline{W^{(1)} G_1^{(1)} A_1^{(1)}} \rangle \prod_{i \neq 1} \Upsilon_i \right] \\ & = -c \langle \xi_1^{(1)} \rangle \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] - c \frac{1}{2n^2} \mathbb{E} \left[\sum_{ab} (G_1^{(1)})_{aa} (G_1^{(1)} A_1^{(1)})_{bb} \prod_{i \neq 1} \Upsilon_i \right] \\ & \quad + c \mathbb{E} \left[\langle -\underline{W^{(1)} G_1^{(1)} A_1^{(1)}} \rangle \prod_{i \neq 1} \Upsilon_i \right] \\ & =: c(I_1^{(1)} + I_2^{(1)} + I_3^{(1)}). \end{aligned}$$

We shall now estimate $I_1^{(1)}$, $I_2^{(1)}$, and $I_3^{(1)}$ individually. From a glimpse of the upcoming calculations, we shall see from (6.17) that $I_2^{(1)}$ actually gets canceled by a part of $I_3^{(1)}$. The primary ingredient in these estimations is the cumulant expansion as mentioned in the Result 6.2.

6.1. Evaluation of $I_3^{(1)}$.

$$\begin{aligned}
(6.15) \quad I_3^{(1)} &= \mathbb{E} \left[\langle -W^{(1)} G_1^{(1)} A_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i \right] \\
&= \mathbb{E} \left[\sum_{ab} w_{ba}^{(1)} \langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i \right] \\
&= \sum_{ab} \sum_{k \geq 0} \sum_{\alpha \in \{(ab)_1, (ba)_1, (ab)_2, (ba)_2\}^k} \frac{\kappa((ba)_1, \alpha)}{k!} \mathbb{E} \left[\partial_\alpha (\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i) \right].
\end{aligned}$$

Notice that in (6.15), the term with $k = 0$ shall not contribute since according to the Condition 3.1, $\mathbb{E}[w_{ba}^{(1)}] = 0$. Thus, from (6.15) we have

$$\begin{aligned}
(6.16) \quad I_3^{(1)} &= \sum_{ab} \kappa((ba)_1, (ab)_1) \mathbb{E} \left[\partial_{(ab)_1} [\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i] \right] + \sum_{ab} \kappa((ba)_1, (ba)_1) \\
&\quad \mathbb{E} \left[\partial_{(ba)_1} [\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i] \right] + \sum_{ab} \kappa((ba)_1, (ab)_2) \mathbb{E} \left[\partial_{(ab)_2} [\langle -\Delta^{ba} \right. \\
&\quad \left. G_1^{(1)} A_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i] \right] + \sum_{ab} \kappa((ba)_2, (ba)_2) \mathbb{E} \left[\partial_{(ba)_2} [\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \prod_{i \neq 1} \right. \\
&\quad \left. \Upsilon_i] \right] + \sum_{ab} \sum_{k \geq 2} \sum_{\alpha \in \{(ab)_1, (ba)_1, (ab)_2, (ba)_2\}^k} \frac{\kappa((ba)_1, \alpha)}{k!} \mathbb{E} \left[\partial_\alpha (\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i) \right] \\
&= \frac{1}{n} \sum_{ab} \mathbb{E} \left[\partial_{(ab)_1} [\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i] \right] + \frac{\gamma_1}{n} \sum_{ab} \mathbb{E} \left[\partial_{(ba)_1} [\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \right. \\
&\quad \left. \prod_{i \neq 1} \Upsilon_i] \right] + \frac{\gamma}{n} \sum_{ab} \mathbb{E} \left[\partial_{(ab)_2} [\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i] \right] \\
&\quad + \frac{\rho}{n} \sum_{ab} \mathbb{E} \left[\partial_{(ba)_2} [\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i] \right] \\
&\quad + \sum_{ab} \sum_{k \geq 2} \sum_{\alpha \in \{(ab)_1, (ba)_1, (ab)_2, (ba)_2\}^k} \frac{\kappa((ba)_1, \alpha)}{k!} \mathbb{E} \left[\partial_\alpha (\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i) \right] \\
&=: \frac{1}{n} T_1^{(1)} + \frac{1}{n} \gamma_1 T_2^{(1)} + \frac{1}{n} \gamma T_3^{(1)} + \frac{1}{n} \rho T_4^{(1)} + T_5^{(1)}.
\end{aligned}$$

The symbols γ_1, γ, ρ are same as defined in the Condition 3.1. For convenience, we list those down below,

$$\begin{aligned}\kappa((ba)_1, (ab)_1) &= \mathbb{E}[(ba)_1, (b\bar{a})_1] = \mathbb{E}[|(ba)_1|^2] = \frac{1}{n}, \\ \kappa((ba)_1, (ba)_2) &= \mathbb{E}[(ba)_1, (ba)_2] =: \frac{1}{n}\rho, \\ \kappa((ba)_1, (ab)_2) &= \mathbb{E}[(ba)_1, (ab)_2] =: \frac{1}{n}\gamma,\end{aligned}$$

and

$$\kappa((ba)_1, (ba)_1) = \frac{1}{n}\gamma_1 = \begin{cases} 0 & \text{when } X^{(t)} \text{ in condition 3.1 is } \mathbb{C}\text{-valued,} \\ 1/n & \text{when } X^{(t)} \text{ in condition 3.1 is } \mathbb{R}\text{-valued.} \end{cases}$$

We delegate the calculations of the factors $T_1^{(1)}, T_2^{(1)}, T_3^{(1)}, T_4^{(1)}, T_5^{(1)}$ in the following subsections.

6.1.1. *Evaluation of $T_1^{(1)}, T_2^{(1)}, T_3^{(1)}, T_4^{(1)}$.* Let us first evaluate $T_1^{(1)}$ as follows;

$$\begin{aligned}(6.17) \quad T_1^{(1)} &= \sum_{ab} \mathbb{E} \left[\partial_{(ab)_1} \left(\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i \right) \right] \\ &= \frac{1}{2n} \sum'_{ab} \mathbb{E} \left[(G_1^{(1)})_{aa} (G_1^{(1)} A_1^{(1)})_{bb} \prod_{i \neq 1} \Upsilon_i \right] \\ &\quad + \sum_{ab} \mathbb{E} \left[\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \sum_{i \neq 1} \partial_{(ab)_1} \left(c \langle -W^{(1)} G_i^{(1)} A_i^{(1)} \rangle \right) \prod_{j \neq 1, i} \Upsilon_j \right] \\ &= -nI_2^{(1)} + \sum'_{ab} \mathbb{E} \left[\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \sum_{i \neq 1} c \langle -\Delta^{ab} G_i^{(1)} A_i^{(1)} \rangle \right. \\ &\quad \left. + \underline{W^{(1)} G_i^{(1)} \Delta^{ab} G_i^{(1)} A_i^{(1)}} \prod_{j \neq 1, i} \Upsilon_j \right] \\ &= -nI_2^{(1)} + c \sum_{i \neq 1} \mathbb{E} \left[\sum'_{ab} \left(\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \langle -\Delta^{ab} G_i^{(1)} A_i^{(1)} \rangle \right. \right. \\ &\quad \left. \left. + \langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \langle \underline{W^{(1)} G_i^{(1)} \Delta^{ab} G_i^{(1)} A_i^{(1)}} \rangle \right) \prod_{j \neq 1, i} \Upsilon_j \right] \\ &= -nI_2^{(1)} + \frac{c}{2n} \sum_{i \neq 1} \mathbb{E} \left[\langle G_1^{(1)} A_1^{(1)} E_1 G_i^{(1)} A_i^{(1)} E_2 + G_1^{(1)} A_1^{(1)} E_2 G_i^{(1)} A_i^{(1)} E_1 \rangle \right. \\ &\quad \left. \prod_{j \neq 1, i} \Upsilon_j \right] + \frac{c}{2n} \sum_{i \neq 1} \mathbb{E} \left[\langle -G_1^{(1)} A_1^{(1)} E_1 \underline{G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)} E_2} \right. \\ &\quad \left. - G_1^{(1)} A_1^{(1)} E_2 \underline{G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)} E_1} \rangle \prod_{j \neq 1, i} \Upsilon_j \right] \\ &= -nI_2^{(1)} + \frac{c}{2n} \sum_{i \neq 1} \mathbb{E} \left[\langle G_1^{(1)} A_1^{(1)} E G_i^{(1)} A_i^{(1)} E^\dagger \right.\end{aligned}$$

$$- G_1^{(1)} A_1^{(1)} \underline{E G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)} E^\dagger} \prod_{j \neq 1, i} \Upsilon_j \Big].$$

Here, $I_2^{(1)}$ is same as defined in (6.14), and E_1, E_2, E, E^\dagger are same as defined in (6.6). Similarly, we can calculate

$$(6.18) \quad \begin{aligned} & T_2^{(1)} \\ &= \frac{1}{2n} \sum'_{ab} \mathbb{E} \left[(G_1^{(1)})_{ab} (G_1^{(1)} A_1^{(1)})_{ab} \prod_{i \neq 1} \Upsilon_i \right] + \frac{c}{2n} \sum'_{i \neq 1} \mathbb{E} \left[\langle G_1^{(1)} A_1^{(1)} E (A_i^{(1)})^t (G_i^{(1)})^t E^\dagger \right. \\ & \quad \left. - G_1^{(1)} A_1^{(1)} \underline{E (G_i^{(1)})^t W^{(1)} (A_i^{(1)})^t (G_i^{(1)})^t E^\dagger} \prod_{j \neq 1, i} \Upsilon_j \right], \end{aligned}$$

$$(6.19) \quad \begin{aligned} & T_3^{(1)} \\ &= \frac{d}{2n} \sum'_{i \neq 1} \mathbb{E} \left[\langle G_1^{(1)} A_1^{(1)} E G_i^{(2)} A_i^{(2)} E^\dagger + \underline{G_1^{(1)} A_1^{(1)} E G_i^{(2)} A_i^{(2)} W^{(2)} G_i^{(2)} E^\dagger} \prod_{j \neq 1, i} \Upsilon_j \right], \end{aligned}$$

$$(6.20) \quad \begin{aligned} & T_4^{(1)} \\ &= \frac{d}{2n} \sum'_{i \neq 1} \mathbb{E} \left[\left(\langle G_1^{(1)} A_1^{(1)} E_1 G_i^{(2)} A_i^{(2)} E_1 \rangle + \langle G_1^{(1)} A_1^{(1)} E_2 G_i^{(2)} A_i^{(2)} E_2 \rangle \right. \right. \\ & \quad \left. \left. + \langle \underline{G_1^{(1)} A_1^{(1)} E_1 G_i^{(2)} A_i^{(2)} W^{(2)} G_i^{(2)} E_1} \rangle + \langle \underline{G_1^{(1)} A_1^{(1)} E_2 G_i^{(2)} A_i^{(2)} W^{(2)} G_i^{(2)} E_2} \rangle \right) \right. \\ & \quad \left. \prod_{j \neq 1, i} \Upsilon_j \right], \end{aligned}$$

where A^t denotes the transpose of matrix A .

6.2. Reduced form of $I^{(1)} + I^{(2)}$. Now, we use the above expressions of $T_1^{(1)}, T_2^{(1)}, T_3^{(1)}, T_4^{(1)}$ and $T_5^{(1)}$ in conjunction with (6.16) to evaluate $I^{(1)} + I^{(2)}$. First, we notice that the expression of $I_3^{(1)}$ in equation (6.16) contains the factor $T_1^{(1)} + \gamma_1 T_2^{(1)}$. But, as per the definition of γ_1 , it can take two possible values; $\gamma_1 = 0$ (complex value case), and $\gamma_1 = 1$ (real value case). In the first case, as per the equation (6.16), we observe that $T_2^{(1)}$ will not have any contribution to the $I_3^{(1)}$. In the rest of the section, we present the calculation for the second case i.e., $\gamma_1 = 1$. The other case when $\gamma_1 = 0$ is discussed in Remark 6.3.

$$(6.21) \quad \begin{aligned} & T_1^{(1)} + \gamma_1 T_2^{(1)} \\ &= T_1^{(1)} + T_2^{(1)} \\ &= -n I_2^{(1)} + \frac{1}{2n} \sum'_{ab} \mathbb{E} \left[(G_1^{(1)})_{ab} (G_1^{(1)} A_1^{(1)})_{ab} \prod_{i \neq 1} \Upsilon_i \right] \\ & \quad + \frac{c}{2n} \sum'_{i \neq 1} \mathbb{E} \left[\langle G_1^{(1)} A_1^{(1)} E (G_i^{(1)} A_i^{(1)} + (A_i^{(1)})^t (G_i^{(1)})^t) E^\dagger \right. \\ & \quad \left. - G_1^{(1)} A_1^{(1)} \underline{E (G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)} + (G_i^{(1)})^t W^{(1)} (A_i^{(1)})^t (G_i^{(1)})^t) E^\dagger} \prod_{j \neq 1, i} \Upsilon_j \right]. \end{aligned}$$

Thus, from (6.14), (6.16), (6.19), (6.20), and (6.21) we have

$$\begin{aligned}
(6.22) \quad I^{(1)} &= c(I_1^{(1)} + I_2^{(1)} + I_3^{(1)}) \\
&= c\left(I_1^{(1)} + I_2^{(1)} + \frac{1}{n}T_1^{(1)} + \frac{1}{n}T_2^{(1)} + \frac{\gamma}{n}T_3^{(1)} + \frac{\rho}{n}T_4^{(1)} + T_5^{(1)}\right) \\
&= cT_5^{(1)} - c\langle \xi_1^{(1)} \rangle \mathbb{E}\left[\prod_{i \neq 1} \Upsilon_i\right] + \frac{c}{2n^2} \sum'_{ab} \mathbb{E}\left[(G_1^{(1)})_{ab}(G_1^{(1)} A_1^{(1)})_{ab} \prod_{i \neq 1} \Upsilon_i\right] \\
&\quad + \frac{c^2}{2n^2} \sum_{i \neq 1} \mathbb{E}\left[\langle G_1^{(1)} A_1^{(1)} E(G_i^{(1)} A_i^{(1)} + (A_i^{(1)})^t(G_i^{(1)})^t) E^\dagger \right. \\
&\quad \left. - G_1^{(1)} A_1^{(1)} E(\underline{G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)}} + \underline{(G_i^{(1)})^t W^{(1)} (A_i^{(1)})^t (G_i^{(1)})^t}) E^\dagger) \prod_{j \neq 1, i} \Upsilon_j\right] \\
&\quad + \frac{cd\gamma}{2n^2} \sum_{i \neq 1} \mathbb{E}\left[\langle \underline{G_1^{(1)} A_1^{(1)} E G_i^{(2)} A_i^{(2)} E^\dagger} \rangle + \langle \underline{G_1^{(1)} A_1^{(1)} E G_i^{(2)} A_i^{(2)} W^{(2)} G_i^{(2)} E^\dagger} \rangle\right) \\
&\quad \prod_{j \neq 1, i} \Upsilon_j \left. + \frac{cd\rho}{2n^2} \sum_{i \neq 1} \mathbb{E}\left[\langle \underline{G_1^{(1)} A_1^{(1)} E_1 G_i^{(2)} A_i^{(2)} E_1} \rangle + \langle \underline{G_1^{(1)} A_1^{(1)} E_2 G_i^{(2)} A_i^{(2)} E_2} \rangle\right) \right. \\
&\quad \prod_{j \neq 1, i} \Upsilon_j \left. + \frac{cd\rho}{2n^2} \sum_{i \neq 1} \mathbb{E}\left[\langle \underline{G_1^{(1)} A_1^{(1)} E_1 G_i^{(2)} A_i^{(2)} W^{(2)} G_i^{(2)} E_1} \rangle \right. \right. \\
&\quad \left. \left. + \langle \underline{G_1^{(1)} A_1^{(1)} E_2 G_i^{(2)} A_i^{(2)} W^{(2)} G_i^{(2)} E_2} \rangle\right) \prod_{j \neq 1, i} \Upsilon_j\right].
\end{aligned}$$

Similarly, an equivalent expression of $I^{(2)}$ can be obtained as well. Now, by using (6.13), the expressions of $I^{(1)}$ from (6.22), and an equivalent expression for $I^{(2)}$, we have

$$\begin{aligned}
&\mathbb{E}\left[\prod_{i \in [p]} \Upsilon_i\right] \\
&= cT_5^{(1)} + dT_5^{(2)} - c\langle \xi_1^{(1)} \rangle \mathbb{E}\left[\prod_{i \neq 1} \Upsilon_i\right] - d\langle \xi_1^{(2)} \rangle \mathbb{E}\left[\prod_{i \neq 1} \Upsilon_i\right] \\
&\quad + \frac{c}{2n^2} \sum'_{ab} \mathbb{E}\left[(G_1^{(1)})_{ab}(G_1^{(1)} A_1^{(1)})_{ab} \prod_{i \neq 1} \Upsilon_i\right] + \frac{d}{2n^2} \sum'_{ab} \mathbb{E}\left[(G_1^{(2)})_{ab}(G_1^{(2)} A_1^{(2)})_{ab} \prod_{i \neq 1} \Upsilon_i\right] \\
&\quad + \frac{c^2}{2n^2} \sum_{i \neq 1} \mathbb{E}\left[\langle G_1^{(1)} A_1^{(1)} E(G_i^{(1)} A_i^{(1)} + (A_i^{(1)})^t(G_i^{(1)})^t) E^\dagger - G_1^{(1)} A_1^{(1)} E(\underline{G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)}} \right. \\
&\quad \left. + \underline{(G_i^{(1)})^t W^{(1)} (A_i^{(1)})^t (G_i^{(1)})^t}) E^\dagger) \prod_{j \neq 1, i} \Upsilon_j\right] + \frac{d^2}{2n^2} \sum_{i \neq 1} \mathbb{E}\left[\langle G_1^{(2)} A_1^{(2)} E(G_i^{(2)} A_i^{(2)} \right. \\
&\quad \left. + (A_i^{(2)})^t(G_i^{(2)})^t) E^\dagger - G_1^{(2)} A_1^{(2)} E(\underline{G_i^{(2)} A_i^{(2)} W^{(2)} G_i^{(2)}} + \underline{(G_i^{(2)})^t W^{(2)} (A_i^{(2)})^t (G_i^{(2)})^t}) E^\dagger) \right. \\
&\quad \left. \prod_{j \neq 1, i} \Upsilon_j\right] + \frac{cd\gamma}{2n^2} \sum_{i \neq 1} \mathbb{E}\left[\langle \underline{G_1^{(1)} A_1^{(1)} E G_i^{(2)} A_i^{(2)} E^\dagger} \rangle \prod_{j \neq 1, i} \Upsilon_j\right]
\end{aligned}$$

$$\begin{aligned}
& + \frac{cd\gamma}{2n^2} \sum_{i \neq 1} \mathbb{E} \left[\langle G_1^{(2)} A_1^{(2)} E G_i^{(1)} A_i^{(1)} E^\dagger \rangle \prod_{j \neq 1, i} \Upsilon_j \right] + \frac{cd\rho}{2n^2} \sum_{i \neq 1} \mathbb{E} \left[\langle (G_1^{(1)} A_1^{(1)} E_1 G_i^{(2)} A_i^{(2)} E_1) \right. \\
& + \langle G_1^{(1)} A_1^{(1)} E_2 G_i^{(2)} A_i^{(2)} E_2 \rangle) \prod_{j \neq 1, i} \Upsilon_j \left. \right] + \frac{cd\rho}{2n^2} \sum_{i \neq 1} \mathbb{E} \left[\langle (G_1^{(2)} A_1^{(2)} E_1 G_i^{(1)} A_i^{(1)} E_1) \right. \\
& + \langle G_1^{(2)} A_1^{(2)} E_2 G_i^{(1)} A_i^{(1)} E_2 \rangle) \prod_{j \neq 1, i} \Upsilon_j \left. \right] + \frac{cd\gamma}{2n^2} \left(\sum_{i \neq 1} \mathbb{E} \left[\langle \underline{G_1^{(1)} A_1^{(1)} E G_i^{(2)} A_i^{(2)} W^{(2)} G_i^{(2)} E^\dagger} \rangle \right. \right. \\
& \left. \left. \prod_{j \neq 1, i} \Upsilon_j \right] + \sum_{i \neq 1} \mathbb{E} \left[\langle \underline{G_1^{(2)} A_1^{(2)} E G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)} E^\dagger} \rangle \prod_{j \neq 1, i} \Upsilon_j \right] \right) \\
& + \frac{cd\rho}{2n^2} \sum_{i \neq 1} \mathbb{E} \left[\langle \underline{(G_1^{(1)} A_1^{(1)} E_1 G_i^{(2)} A_i^{(2)} W^{(2)} G_i^{(2)} E_1) + (G_1^{(1)} A_1^{(1)} E_2 G_i^{(2)} A_i^{(2)} W^{(2)} G_i^{(2)} E_2)} \rangle \right. \\
& \left. \prod_{j \neq 1, i} \Upsilon_j \right] + \frac{cd\rho}{2n^2} \sum_{i \neq 1} \mathbb{E} \left[\langle \underline{(G_1^{(2)} A_1^{(2)} E_1 G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)} E_1) \right. \\
& \left. + \underline{(G_1^{(2)} A_1^{(2)} E_2 G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)} E_2)} \rangle \prod_{j \neq 1, i} \Upsilon_j \right]
\end{aligned}$$

(6.23)

$$\begin{aligned}
& = : cT_5^{(1)} + dT_5^{(2)} - c\langle \xi_1^{(1)} \rangle \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] - d\langle \xi_1^{(2)} \rangle \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] + \frac{c}{2n^2} S_1^{(1)} \\
& + \frac{d}{2n^2} S_1^{(2)} + \frac{c^2}{2n^2} S_2^{(1)} + \frac{d^2}{2n^2} S_2^{(2)} + \frac{cd\gamma}{2n^2} S_3^{(1)} + \frac{cd\gamma}{2n^2} S_3^{(2)} + \frac{cd\rho}{2n^2} S_4^{(1)} \\
& + \frac{cd\rho}{2n^2} S_4^{(2)} + \frac{cd\gamma}{2n^2} (S_5^{(1)} + S_5^{(2)}) + \frac{cd\rho}{2n^2} S_6^{(1)} + \frac{cd\rho}{2n^2} S_6^{(2)}.
\end{aligned}$$

We note that the $S_i^{(t)}$ terms actually originated from the cumulant expansion (6.16), when $k = 1$, and the $T_i^{(t)}$ terms originated from the terms corresponding to $k \geq 2$.

6.3. Evaluation of $S_i^{(t)}$ terms in (6.23). Now, we proceed to the evaluations of $S_i^{(t)}$. The evaluation of $S_2^{(t)}$ s are explained in the Subsection 6.3.1, and rest are calculated in the Subsection 6.3.2.

6.3.1. Evaluation of $S_2^{(t)}$ s. We shall use the notation $G_i^{(t)} = (G^{(t)})^{z_i}$. We first identify the main contributing terms in $S_2^{(1)}$ as follows;

$$\begin{aligned}
(6.24) \quad & S_2^{(1)} \\
& = \sum_{i \neq 1} \mathbb{E} \left[\langle G_1^{(1)} A_1^{(1)} E (G_i^{(1)} A_i^{(1)} + (A_i^{(1)})^t (G_i^{(1)})^t) E^\dagger \right. \\
& \quad \left. - G_1^{(1)} A_1^{(1)} E (\underline{G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)}} + \underline{(G_i^{(1)})^t W^{(1)} (A_i^{(1)})^t (G_i^{(1)})^t}) E^\dagger \rangle \prod_{j \neq 1, i} \Upsilon_j \right] \\
& = \sum_{i \neq 1} \mathbb{E} \left[\left\{ \langle G_1^{(1)} A_1^{(1)} E (G_i^{(1)} A_i^{(1)} + (A_i^{(1)})^t (G_i^{(1)})^t) E^\dagger \rangle E \right. \right.
\end{aligned}$$

$$\begin{aligned}
& + \langle G_1^{(1)} \mathcal{S}[G_1^{(1)} A_1^{(1)} G_i^{(1)} A_i^{(1)}] G_i^{(1)} E^\dagger \rangle + G_1^{(1)} \mathcal{S}[G_1^{(1)} A_1^{(1)} E G_i^{(1)}] (A_i^{(1)})^t G_i^{(1)} E^\dagger \\
& - \langle \underline{G_1^{(1)} A_1^{(1)} E G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)} E^\dagger} - \underline{G_1^{(1)} A_1^{(1)} E G_i^{(1)} W^{(1)} (A_i^{(1)})^t G_i^{(1)} E^\dagger} \rangle \\
& \quad \left. \prod_{j \neq 1, i} \Upsilon_j \right] \\
= & \sum_{i \neq 1} \mathbb{E} \left[\left\{ \langle G_1^{(1)} A_1^{(1)} E (G_i^{(1)} A_i^{(1)} + (A_i^{(1)})^t (G_i^{(1)})^t) E^\dagger \rangle + \langle G_1^{(1)} \mathcal{S}[G_1^{(1)} A_1^{(1)} E \right. \right. \\
& \quad \left. \left. G_i^{(1)} A_i^{(1)}] G_i^{(1)} E^\dagger \rangle + G_1^{(1)} \mathcal{S}[G_1^{(1)} A_1^{(1)} E G_i^{(1)}] (A_i^{(1)})^t G_i^{(1)} E^\dagger \rangle \right\} \prod_{j \neq 1, i} \Upsilon_j \right] \\
& - \sum_{i \neq 1} \mathbb{E} \left[\left\{ \langle \underline{G_1^{(1)} A_1^{(1)} E G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)} E^\dagger} \right. \right. \\
& \quad \left. \left. + \underline{G_1^{(1)} A_1^{(1)} E G_i^{(1)} W^{(1)} (A_i^{(1)})^t G_i^{(1)} E^\dagger} \rangle \right\} \prod_{j \neq 1, i} \Upsilon_j \right].
\end{aligned}$$

In the following discussion, we show that the main contribution comes only from the terms of the form $G_1^{(1)} A_1^{(1)} E G_i^{(1)} A_i^{(1)} E^\dagger$ and $G_1^{(1)} \mathcal{S}[G_1^{(1)} A_1^{(1)} E G_i^{(1)}] (A_i^{(1)})^t G_i^{(1)} E^\dagger$. The main ingredient here is the local law as stated in the Theorem 4.2, which gives us the following estimates;

$$\begin{aligned}
(6.25) \quad & \left| \langle G_1^{(1)} A_1^{(1)} E G_i^{(1)} A_i^{(1)} E^\dagger - (M^{(1)})_{A_1^{(1)} E}^{z_1, z_i} A_i^{(1)} E^\dagger \rangle \right| \\
& \prec \frac{C_\epsilon \|A_i^{(1)} E^\dagger\| \|A_1^{(1)} E\|}{n \eta_*^{1i} (\eta_1 \eta_i)^{1/2} |\hat{\beta}_{1i}^*(1)|} \left((\eta_*^{1i})^{1/12} + \frac{(\eta_*^{1i})^{1/4}}{|\hat{\beta}_{1i}^*(1)|} + \frac{1}{\sqrt{n \eta_*^{1i}}} + \frac{1}{(|\hat{\beta}_*^{(1)}| n \eta_*^{1i})^{1/4}} \right) \\
& = \mathcal{O}_\prec \left(\frac{1}{n |z_1 - z_i|^4 \eta_*^{1i} (\eta_1 \eta_i)^{1/2} |\beta_i^{(1)}| |\beta_1^{(1)}|} \right),
\end{aligned}$$

where $|\hat{\beta}_{1i}^*(1)| \gtrsim |z_1 - z_i|^2$ is from Lemma 4.4 and $\eta_*^{1i} = \min\{\eta_1, \eta_i\}$.

$$\begin{aligned}
& \left| \langle G_1^{(1)} \mathcal{S}[G_1^{(1)} A_1^{(1)} E G_i^{(1)}] (A_i^{(1)})^t G_i^{(1)} E^\dagger - \mathcal{S}[(M^{(1)})_{A_1^{(1)} E}^{z_1, \bar{z}_i}] (A_i^{(1)})^t (M^{(1)})_{E^\dagger}^{\bar{z}_i, z_1} \rangle \right| \\
= & \left| \langle \mathcal{S}[G_1^{(1)} A_1^{(1)} E G_i^{(1)}] (A_i^{(1)})^t G_i^{(1)} E^\dagger G_1^{(1)} - \mathcal{S}[(M^{(1)})_{A_1^{(1)} E}^{z_1, \bar{z}_i}] (A_i^{(1)})^t (M^{(1)})_{E^\dagger}^{\bar{z}_i, z_1} \rangle \right| \\
& \prec \left(\frac{C_\epsilon \|A_1^{(1)} E\| \|A_i^{(1)} E^\dagger\|}{n \eta_*^{1i} (\eta_1 \eta_i)^{1/2} |\hat{\beta}_{1i}^*(1)|} \left((\eta_*^{1i})^{1/2} + \frac{(\eta_*^{1i})^{1/4}}{|\hat{\beta}_{1i}^*(1)|} + \frac{1}{\sqrt{n \eta_*^{1i}}} + \frac{1}{(|\hat{\beta}_{1i}^*(1)| n \eta_*^{1i})^{1/4}} \right) \right)^2 \\
(6.26) \quad & = \mathcal{O}_\prec \left(\frac{1}{n^2 |z_1 - z_i|^4 (\eta_*^{1i})^2 \eta_1 \eta_i |\beta_i^{(1)}| |\beta_1^{(1)}|} \right).
\end{aligned}$$

Now, for the error terms in (6.31), we shall use the bound

$$\begin{aligned}
(6.27) \quad & \mathbb{E} \left| \langle \underline{G_1^{(1)} A_1^{(1)} E G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)} E^\dagger} \rangle \right|^2 + \mathbb{E} \left| \langle \underline{G_1^{(1)} A_1^{(1)} E G_i^{(1)} W^{(1)} (A_i^{(1)})^t G_i^{(1)} E^\dagger} \rangle \right|^2 \\
& \lesssim \left(\frac{1}{n \eta_1 \eta_i \eta_*^{1i} |\beta_1^{(1)}| |\beta_i^{(1)}|} \right)^2,
\end{aligned}$$

which we shall prove at the end of this section in Corollary 6.7. In addition, using the (6.2), $|\xi_i^{(t)}| \lesssim \frac{1}{n}$, and the local law Theorem 4.1, we obtain that

$$\begin{aligned}
& |\Upsilon_i| \\
&= |c\langle -\underline{W}^{(1)}G_i^{(1)}A_i^{(1)} - \xi_i^{(1)} \rangle + d\langle -\underline{W}^{(2)}G_i^{(2)}A_i^{(2)} - \xi_i^{(2)} \rangle| \\
&= |c\langle G_i^{(1)} - M_i^{(1)} - \xi_i^{(1)} \rangle + d\langle G_i^{(2)} - M_i^{(2)} - \xi_i^{(2)} \rangle| + \mathcal{O}_{\prec}\left(\frac{1}{|\beta_i^*|(n\eta_i)^2}\right) \\
(6.28) \quad &= \mathcal{O}_{\prec}\left(\frac{1}{\sqrt{n\eta_i}}\right) + \mathcal{O}_{\prec}\left(\frac{1}{|\beta_i^*|(n\eta_i)^2}\right) = \mathcal{O}_{\prec}\left(\frac{1}{|\beta_i^*|\sqrt{n\eta_i}}\right),
\end{aligned}$$

where $\beta_i^* := \min\{\beta_i^{(1)}, \beta_i^{(2)}\}$.

Thus, by using (6.27) and (6.28), we have

$$\begin{aligned}
(6.29) \quad & \left| n^{-2}\mathbb{E}\left[\langle \underline{G}_1^{(1)}A_1^{(1)}E \underline{G}_i^{(1)}A_i^{(1)}W^{(1)}G_i^{(1)}E^\dagger \rangle + \langle \underline{G}_1^{(1)}A_1^{(1)}E \underline{G}_i^{(1)}W^{(1)}(A_i^{(1)})^t \underline{G}_i^{(1)}E^\dagger \rangle\right] \prod_{j \neq 1, i} \Upsilon_j \right| \\
& \prec \frac{1}{n^2} \left[\prod_{j \neq 1, i} \frac{1}{|\beta_j^*|\sqrt{n\eta_j}} \right] \left((\mathbb{E}|\langle \underline{G}_1^{(1)}A_1^{(1)}E \underline{G}_i^{(1)}A_i^{(1)}W^{(1)}G_i^{(1)}E^\dagger \rangle|^2)^{1/2} \right. \\
& \quad \left. + (\mathbb{E}|\langle \underline{G}_1^{(1)}A_1^{(1)}E \underline{G}_i^{(1)}W^{(1)}(A_i^{(1)})^t \underline{G}_i^{(1)}E^\dagger \rangle|^2)^{1/2} \right) \\
& \prec \frac{1}{n^2} \left[\prod_{j \neq 1, i} \frac{1}{|\beta_j^*|\sqrt{n\eta_j}} \right] \left(\frac{1}{n\eta_1\eta_i\eta_*^{1i}|\beta_1^*||\beta_i^*|} \right) \\
&= \frac{1}{n^2} \prod_j \frac{|\beta_1^*|\sqrt{n\eta_1}|\beta_i^*|\sqrt{n\eta_i}}{|\beta_j^*|\sqrt{n\eta_j}} \left(\frac{1}{n\eta_1\eta_i\eta_*^{1i}|\beta_1^*||\beta_i^*|} \right) \\
&= \frac{1}{n^2\eta_*^{1i}\sqrt{\eta_1\eta_i}} \prod_j \frac{1}{|\beta_j^*|\sqrt{n\eta_j}} \\
& \prec \frac{1}{(n\eta_*^{1i})^2} \prod_j \frac{1}{|\beta_j^*|\sqrt{n\eta_j}} \\
&= \frac{\psi}{(n\eta_*^{1i})^2},
\end{aligned}$$

where ψ is defined as follows;

$$(6.30) \quad \psi := \prod_j \frac{1}{|\beta_j^*|\sqrt{n\eta_j}}.$$

Now, in (6.24), using the estimates (6.25), (6.26), (6.29) along with the notations defined in (6.33), we obtain the expression of $n^{-2}S_2^{(1)}$ as follows;

$$\begin{aligned}
(6.31) \quad & \frac{1}{n^2}S_2^{(1)} \\
&= \frac{1}{n^2} \sum_{i \neq 1} \mathbb{E} \left[\left\{ V_{1i}^{(1)}(z_i, z_j, \eta_i, \eta_j) + V_{1i}^{(1)}(z_i, \bar{z}_j, \eta_i, \eta_j) \right\} \right]
\end{aligned}$$

$$\begin{aligned}
& + \mathcal{O}_{\prec} \left(\frac{1}{n|z_1 - z_i|^4 \eta_*^{1i} (\eta_1 \eta_i)^{1/2} |\beta_i^{(1)}| |\beta_1^{(1)}|} + \frac{1}{n^2 |z_1 - z_i|^4 (\eta_*^{1i})^2 \eta_1 \eta_i |\beta_i^{(1)}| |\beta_1^{(1)}|} \right) \Big\} \\
& \prod_{j \neq 1, i} \Upsilon_j \Big] - \frac{1}{n^2} \sum_{i \neq 1} \mathbb{E} \left[\left\{ \langle \underline{G_1^{(1)} A_1^{(1)} E G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)} E^\dagger} \right. \right. \\
& \left. \left. + \underline{G_1^{(1)} A_1^{(1)} E G_i^{(1)} W^{(1)} (A_i^{(1)})^t G_i^{(1)} E^\dagger} \rangle \right\} \prod_{j \neq 1, i} \Upsilon_j \right] \\
& = \frac{1}{n^2} \sum_{i \neq 1} \left\{ \widehat{V_{1i}^{(1)}} + \mathcal{O}_{\prec} \left(\frac{1}{n|z_1 - z_i|^4 \eta_*^{1i} (\eta_1 \eta_i)^{1/2} |\beta_i^{(1)}| |\beta_1^{(1)}|} \right. \right. \\
& \left. \left. + \frac{1}{n^2 |z_1 - z_i|^4 (\eta_*^{1i})^2 \eta_1 \eta_i |\beta_i^{(1)}| |\beta_1^{(1)}|} \right) \right\} \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] + \mathcal{O}_{\prec} \left(\frac{\psi}{(n\eta_*)^2} \right) \\
& = \sum_{i \neq 1} \left\{ \frac{\widehat{V_{1i}^{(1)}}}{n^2} \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] + \mathcal{O}_{\prec} \left(\psi \left(\frac{1}{(n\eta_*)^2} + \frac{1}{n^2 \eta_*^{1i} |z_1 - z_i|^4} + \frac{1}{(n\eta_*^{1i})^3 |z_1 - z_i|^4} \right) \right) \right\} \\
& = : \frac{1}{n^2} \sum_{i \neq 1} \widehat{V_{1i}^{(1)}} \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] + error_1,
\end{aligned}$$

where $\widehat{V_{1i}^{(1)}}$ is defined below. Similarly, we can evaluate $n^{-2} S_2^{(2)}$ as follows;

$$\begin{aligned}
(6.32) \quad & \frac{1}{n^2} \sum_{i \neq 1} \mathbb{E} \left[\langle G_1^{(2)} A_1^{(2)} E (G_i^{(2)} A_i^{(2)} + (A_i^{(2)})^t (G_i^{(2)})^t) E^\dagger \right. \\
& \left. - G_1^{(2)} A_1^{(2)} E \underline{(G_i^{(2)} A_i^{(2)} W^{(2)} G_i^{(2)})} + \underline{(G_i^{(2)})^t W^{(2)} (A_i^{(2)})^t (G_i^{(2)})^t} E^\dagger \rangle \prod_{j \neq 1, i} \Upsilon_j \right] \\
& = \sum_{i \neq 1} \left\{ \frac{\widehat{V_{1i}^{(2)}}}{n^2} \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] + \mathcal{O}_{\prec} \left(\psi \left(\frac{1}{(n\eta_*)^2} + \frac{1}{n^2 \eta_*^{1i} |z_1 - z_i|^4} + \frac{1}{(n\eta_*^{1i})^3 |z_1 - z_i|^4} \right) \right) \right\} \\
& = \frac{1}{n^2} \sum_{i \neq 1} \widehat{V_{1i}^{(2)}} \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] + error_1,
\end{aligned}$$

where $\widehat{V_{1i}^{(t)}}(z_i, z_j, \eta_i, \eta_j) = V_{1i}^{(t)}(z_i, z_j, \eta_i, \eta_j) + V_{1i}^{(t)}(z_i, \bar{z}_j, \eta_i, \eta_j)$ with

$$\begin{aligned}
(6.33) \quad & V_{1i}^{(t)} := \langle (M^{(t)})_{A_1^{(t)} E}^{z_1, z_i} A_i^{(t)} E^\dagger \rangle + \langle (M^{(t)})_{A_1^{(t)} E}^{z_1, \bar{z}_i} (A_i^{(t)})^t E^\dagger \rangle \\
& + \langle \mathcal{S}[(M^{(t)})_{A_1^{(t)} E}^{z_1, z_i} A_i^{(t)}] (M^{(t)})_{E^\dagger}^{z_i, z_1} \rangle + \langle \mathcal{S}[(M^{(t)})_{A_1^{(t)} E}^{z_1, \bar{z}_i} (A_i^{(t)})^t] (M^{(t)})_{E^\dagger}^{\bar{z}_i, z_1} \rangle \\
& = \frac{2m_1^{(t)} m_i^{(t)} (2u_1^{(t)} u_i^{(t)} \Re z_1 \bar{z}_i + (u_1^{(t)} u_i^{(t)} |z_1| |z_i|)^2 (r_1^{(t)} r_i^{(t)} - 4))}{\beta_1^{(t)} \beta_i^{(t)} [1 + (u_1^{(t)} u_i^{(t)} |z_1| |z_i|)^2 - (m_1^{(t)} m_i^{(t)})^2 - 2u_1^{(t)} u_i^{(t)} \Re z_1 \bar{z}_i]^2} \\
& + \frac{2m_1^{(t)} m_i^{(t)} ((m_1^{(t)})^2 + (u_1^{(t)})^2 |z_1|^2) ((m_i^{(t)})^2 + (u_i^{(t)})^2 |z_i|^2)}{\beta_1^{(t)} \beta_i^{(t)} [1 + (u_1^{(t)} u_i^{(t)} |z_1| |z_i|)^2 - (m_1^{(t)} m_i^{(t)})^2 - 2u_1^{(t)} u_i^{(t)} \Re z_1 \bar{z}_i]^2},
\end{aligned}$$

where $r_i^{(t)} = (m_i^{(t)})^2 - (u_i^{(t)})^2 |z_i|^2$, $\beta_i^{(t)} = 1 - (m_i^{(t)})^2 - (u_i^{(t)})^2 |z_i|^2$.

Remark 6.3. If the matrix $X^{(t)}$ from Condition 3.1 is considered to be complex-valued, then $\gamma_1 = 0$. In this case, the term $T_2^{(1)}$ does not contribute to $I_3^{(1)}$ in (6.16). Consequently, the terms $(A_i^{(1)})^t (G_i^{(1)})^t$ and $\overline{(G_i^{(1)})^t W^{(1)} (A_i^{(1)})^t (G_i^{(1)})^t}$ won't appear in the expression of $S_2^{(1)}$ and $S_2^{(2)}$ in (6.24). As a result, $V_{1i}^{(1)}(z_i, \bar{z}_j, \eta_i, \eta_j)$ and $V_{1i}^{(2)}(z_i, \bar{z}_j, \eta_i, \eta_j)$ do not appear in the expressions for $n^{-2} S_2^{(1)}$ and $n^{-2} S_2^{(2)}$ in (6.31) and (6.32), respectively. Therefore, in this case, we have $\widehat{V_{1i}^{(t)}}(z_i, z_j, \eta_i, \eta_j) = V_{1i}^{(t)}(z_i, z_j, \eta_i, \eta_j)$.

6.3.2. *Evaluation of $S_i^{(t)}$ s, when $i = 1, 3, 4, 5, 6$.* Let us first start with $S_3^{(1)}$ and $S_3^{(2)}$. We shall follow the local law stated in Theorem 4.1

$$(6.34) \quad |\langle (G_1^{(t)} - M_1^{(t)}) A \rangle| \prec \frac{C_\epsilon \|A\|}{n\eta_1},$$

which gives us the following estimate;

$$(6.35) \quad \begin{aligned} & \langle G_1^{(t)} A_1^{(t)} E G_i^{(s)} A_i^{(s)} E^\dagger \rangle \\ &= M_1^{(t)} A_1^{(t)} E M_i^{(s)} A_i^{(s)} E^\dagger + M_1^{(t)} A_1^{(t)} E (G_i^{(s)} - M_i^{(s)}) A_i^{(s)} E^\dagger \\ & \quad + (G_1^{(t)} - M_1^{(t)}) A_1^{(t)} E M_i^{(s)} A_i^{(s)} E^\dagger + (G_1^{(t)} - M_1^{(t)}) A_1^{(t)} E (G_i^{(s)} - M_i^{(s)}) A_i^{(s)} E^\dagger \\ &= M_1^{(t)} A_1^{(t)} E M_i^{(s)} A_i^{(s)} E^\dagger + \mathcal{O}_\prec \left(\frac{1}{|\beta_1^*| |\beta_i^*| n \eta_*^{1i}} \right). \end{aligned}$$

Now, we shall use this estimate to evaluate $S_3^{(1)}$ and $S_3^{(2)}$ as follows;

$$(6.36) \quad \begin{aligned} & \frac{1}{n^2} (S_3^{(1)} + S_3^{(2)}) \\ &= \frac{1}{n^2} \sum_{i \neq 1} \mathbb{E} \left[\left(\langle G_1^{(1)} A_1^{(1)} E G_i^{(2)} A_i^{(2)} E^\dagger \rangle + \langle G_1^{(2)} A_1^{(2)} E G_i^{(1)} A_i^{(1)} E^\dagger \rangle \right) \prod_{j \neq 1, i} \Upsilon_j \right] \\ &= \frac{1}{n^2} \sum_{i \neq 1} \mathbb{E} \left[\left(\langle M_1^{(1)} A_1^{(1)} E M_i^{(2)} A_i^{(2)} E^\dagger \rangle + \langle M_1^{(2)} A_1^{(2)} E M_i^{(1)} A_i^{(1)} E^\dagger \rangle \right) \right. \\ & \quad \left. + \mathcal{O}_\prec \left(\frac{1}{|\beta_1^*| |\beta_i^*| n \eta_*^{1i}} \right) \right] \prod_{j \neq 1, i} \Upsilon_j \\ &= \sum_{i \neq 1} \left\{ \frac{1}{n^2} \mathbb{E} \left[\left(\langle M_1^{(1)} A_1^{(1)} E_1 M_i^{(2)} A_i^{(2)} E_2 \rangle + \langle M_1^{(1)} A_1^{(1)} E_2 M_i^{(2)} A_i^{(2)} E_1 \rangle \right) \right. \right. \\ & \quad \left. \left. + \langle M_1^{(2)} A_1^{(2)} E_1 M_i^{(1)} A_i^{(1)} E_2 \rangle + \langle M_1^{(2)} A_1^{(2)} E_2 M_i^{(1)} A_i^{(1)} E_1 \rangle \right) \prod_{j \neq 1, i} \Upsilon_j \right] \\ & \quad \left. + \mathcal{O}_\prec \left(\frac{\psi |\beta_1^*| |\beta_i^*| \sqrt{\eta_1 \eta_i}}{n^2 \eta_*^{1i} |\beta_1^*| |\beta_i^*|} \right) \right\} \end{aligned}$$

$$\begin{aligned}
&= \sum_{i \neq 1} \left\{ \frac{1}{n^2} \left(\frac{1}{\beta_1^{(1)} \beta_i^{(2)}} (4m_i^{(2)} m_1^{(1)} u_i^{(2)} u_1^{(1)} z_i \bar{z}_1) + \frac{1}{\beta_1^{(1)} \beta_i^{(2)}} (4m_i^{(2)} m_1^{(1)} u_i^{(2)} u_1^{(1)} \bar{z}_i z_1) \right. \right. \\
&\quad \left. \left. + \frac{1}{\beta_1^{(2)} \beta_i^{(1)}} (4m_i^{(1)} m_1^{(2)} u_i^{(1)} u_1^{(2)} z_i \bar{z}_1) + \frac{1}{\beta_1^{(2)} \beta_i^{(1)}} (4m_i^{(1)} m_1^{(2)} u_i^{(1)} u_1^{(2)} \bar{z}_i z_1) \right) \right. \\
&\quad \left. \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] + \mathcal{O}_{\prec} \left(\frac{\psi \sqrt{\eta_1 \eta_i}}{n^2 \eta_*^{1i}} \right) \right\} \\
&= : \frac{1}{n^2} \sum_{i \neq 1} \left\{ \left(\frac{8m_i^{(2)} m_1^{(1)} u_1^{(1)} u_i^{(2)} \Re(\bar{z}_1 z_i)}{\beta_i^{(2)} \beta_1^{(1)}} + \frac{8m_i^{(1)} m_1^{(2)} u_i^{(1)} u_1^{(2)} \Re(\bar{z}_1 z_i)}{\beta_i^{(1)} \beta_1^{(2)}} \right) \right. \\
&\quad \left. \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] \right\} + error_2.
\end{aligned}$$

Similarly, we can evaluate $S_4^{(1)}$ and $S_4^{(2)}$ as follows;

$$\begin{aligned}
(6.37) \quad &\frac{1}{n^2} (S_4^{(1)} + S_4^{(2)}) \\
&= \sum_{i \neq 1} \left\{ \frac{1}{n^2} \left(\frac{(1 - \beta_1^{(1)})}{\beta_1^{(1)} \beta_i^{(2)}} ((m_i^{(2)})^2 + (u_i^{(2)})^2 |z_i|^2) \right. \right. \\
&\quad \left. \left. + \frac{(1 - \beta_1^{(1)})}{\beta_i^{(2)} \beta_1^{(1)}} ((m_i^{(2)})^2 + (u_i^{(2)})^2 |z_i|^2) + \frac{(1 - \beta_1^{(2)})}{\beta_1^{(2)} \beta_i^{(1)}} ((m_i^{(1)})^2 + (u_i^{(1)})^2 |z_i|^2) \right. \right. \\
&\quad \left. \left. + \frac{(1 - \beta_1^{(2)})}{\beta_i^{(1)} \beta_1^{(2)}} ((m_i^{(1)})^2 + (u_i^{(1)})^2 |z_i|^2) \right) \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] + \mathcal{O}_{\prec} \left(\frac{\psi \sqrt{\eta_1 \eta_i}}{n^2 \eta_*^{1i}} \right) \right\} \\
&= \sum_{i \neq 1} \left\{ \frac{1}{n^2} \left(\frac{2(1 - \beta_1^{(1)})(1 - \beta_i^{(2)})}{\beta_1^{(1)} \beta_i^{(2)}} + \frac{2(1 - \beta_1^{(2)})(1 - \beta_i^{(1)})}{\beta_1^{(2)} \beta_i^{(1)}} \right) \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] \right. \\
&\quad \left. + \mathcal{O}_{\prec} \left(\frac{\psi \sqrt{\eta_1 \eta_i}}{n^2 \eta_*^{1i}} \right) \right\} \\
&= \frac{1}{n^2} \sum_{i \neq 1} (-2) \{ \partial_{\eta_1} (m_1^{(1)}) \partial_{\eta_i} (m_i^{(2)}) + \partial_{\eta_1} (m_1^{(2)}) \partial_{\eta_i} (m_i^{(1)}) \} \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] + error_2,
\end{aligned}$$

where in the last equality, we have used $(1 - \beta^{(t)})/\beta^{(t)} = -\iota \partial_{\eta_1} (m^{(t)})$ from (4.3). Now, let us move to evaluate the terms $S_1^{(1)}$, $S_1^{(2)}$. As per the definition,

$$\begin{aligned}
S_1^{(1)} &= \sum'_{ab} \mathbb{E} \left[(G_1^{(1)})_{ab} (G_1^{(1)} A_1^{(1)})_{ab} \prod_{i \neq 1} \Upsilon_i \right] \\
&= 2n \mathbb{E} \left[\langle G_1^{(1)} A_1^{(1)} E (G_1^{(1)})^t E^\dagger \rangle \prod_{i \neq 1} \Upsilon_i \right].
\end{aligned}$$

We observe that $\langle G_1^{(1)} A_1^{(1)} E (G_1^{(1)})^t E^\dagger \rangle$ is same as the expression in the second equality in (6.8). Therefore, using (6.8), we obtain

$$\begin{aligned}
& S_1^{(1)} \\
&= \sum'_{ab} \mathbb{E} \left[(G_1^{(1)})_{ab} (G_1^{(1)} A_1^{(1)})_{ab} \prod_{i \neq 1} \Upsilon_i \right] \\
&= \left\{ \frac{in}{2} \partial_{\eta_1} \log (1 - (u_1^{(1)})^2 + 2(u_1^{(1)})^3 |z_1|^2 - (u_1^{(1)})^2 (z_1^2 + \bar{z}_1^2)) \right. \\
&\quad \left. + \mathcal{O}_{\prec} \left(\frac{1}{|z_1 - \bar{z}_1|^2} \frac{1}{|\beta_1^{(1)}| (n\eta_1)^2} \right) \right\} \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] \\
&= \left\{ \frac{in}{2} \partial_{\eta_1} \log (1 - (u_1^{(1)})^2 + 2(u_1^{(1)})^3 |z_1|^2 - (u_1^{(1)})^2 (z_1^2 + \bar{z}_1^2)) \right\} \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] \\
&\quad + \mathcal{O}_{\prec} \left(\frac{\psi}{|\Im z_1| (n\eta_1)^{3/2}} \right) \\
(6.38) \quad &= : \frac{in}{2} \partial_{\eta_1} \log (1 - (u_1^{(1)})^2 + 2(u_1^{(1)})^3 |z_1|^2 - (u_1^{(1)})^2 (z_1^2 + \bar{z}_1^2)) \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] \\
&\quad + error_3,
\end{aligned}$$

Similarly, we can evaluate $S_1^{(2)}$ as well. Now, for all the other terms in (6.23), namely $S_5^{(1)}, S_5^{(2)}, S_6^{(1)}$, and $S_6^{(2)}$, we shall use the following bound from Lemma 6.6;

$$\begin{aligned}
(6.39) \quad & \mathbb{E} |\langle \underline{G_1^{(t)} A_1^{(t)} E G_i^{(s)} A_i^{(s)} W^{(s)} G_i^{(s)} E^\dagger} \rangle|^2 \\
&= \mathcal{O}_{\prec} \left(\frac{1}{|\beta_1^{(t)}|^2 |\beta_i^{(s)}|^2 n^2 \eta_i^4} \right), \text{ for } 1 \leq t \neq s \leq 2.
\end{aligned}$$

Now, using (6.39), similar to the (6.29), we have

$$\begin{aligned}
(6.40) \quad & \frac{1}{n^2} (S_5^{(1)} + S_5^{(2)} + S_6^{(1)} + S_6^{(2)}) \\
&= \mathcal{O}_{\prec} \left(\frac{\psi \sqrt{\eta_1 \eta_i}}{(n\eta_i)^2} + \frac{\psi \sqrt{\eta_1 \eta_i}}{(n\eta_1)^2} \right) \\
&= \mathcal{O}_{\prec} \left(\frac{\psi \sqrt{\eta_1 \eta_i}}{(n\eta_*^{1i})^2} \right) =: error_4.
\end{aligned}$$

Thus, from (6.23), (6.31), (6.32), (6.36), (6.37), (6.38), (6.40), and using the expressions of $\xi_1^{(1)}, \xi_1^{(2)}$ from Lemma 6.1 we have

$$\begin{aligned}
& \mathbb{E} \left[\prod_{i \in [p]} c \langle -\underline{W^{(1)} G_i^{(1)} A_i^{(1)}} - \xi_i^{(1)} \rangle + d \langle -\underline{W^{(2)} G_i^{(2)} A_i^{(2)}} - \xi_i^{(2)} \rangle \right] \\
&= cT_5^{(1)} + dT_5^{(2)} - c \langle \xi_1^{(1)} \rangle \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] - d \langle \xi_1^{(2)} \rangle \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] + \frac{c}{2n^2} S_1^{(1)}
\end{aligned}$$

$$\begin{aligned}
& + \frac{d}{2n^2} S_1^{(2)} + \frac{c^2}{2n^2} S_2^{(1)} + \frac{d^2}{2n^2} S_2^{(2)} + \frac{cd\gamma}{2n^2} (S_3^{(1)} + S_3^{(2)} + S_5^{(1)} + S_5^{(2)}) \\
& + \frac{cd\rho}{2n^2} (S_4^{(1)} + S_4^{(2)} + S_6^{(1)} + S_6^{(2)}) \\
= & cT_5^{(1)} + dT_5^{(2)} - c \langle \frac{i}{4n} \partial_{\eta_1} (1 - (u_1^{(1)})^2 + 2(u_1^{(1)})^3 |z_1|^2 \\
& - (u_1^{(1)})^2 (z_1^2 + \bar{z}_1^2)) \rangle \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] + c \langle \frac{i(\kappa_4)_{11}}{4n} \partial_{\eta_1} (m_1^{(1)})^4 \rangle \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] \\
& - d \langle \frac{i}{4n} \partial_{\eta_1} (1 - (u_1^{(2)})^2 + 2(u_1^{(2)})^3 |z_1|^2 - (u_1^{(2)})^2 (z_1^2 + \bar{z}_1^2)) \rangle \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] \\
& + d \langle \frac{i(\kappa_4)_{22}}{4n} \partial_{\eta_1} (m_1^{(2)})^4 \rangle \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] \\
& + \frac{c}{2n^2} \frac{in}{2} \partial_{\eta} \log (1 - (u_1^{(1)})^2 + 2(u_1^{(1)})^3 |z_1|^2 - (u_1^{(1)})^2 (z_1^2 + \bar{z}_1^2)) \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] \\
& + error_3 + \frac{d}{2n^2} \frac{in}{2} \partial_{\eta} \log (1 - (u_1^{(2)})^2 + 2(u_1^{(2)})^3 |z_1|^2 - (u_1^{(2)})^2 (z_1^2 + \bar{z}_1^2)) \\
& \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] + error_3 + \sum_{i \neq 1} \frac{c^2 \widehat{V}_{1i}^{(1)}}{2n^2} \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] \\
& + \sum_{i \neq 1} \frac{d^2 \widehat{V}_{1i}^{(2)}}{2n^2} \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] + error_1 + \sum_{i \neq 1} \left\{ \frac{8cd\gamma m_i^{(2)} m_1^{(1)} u_1^{(1)} u_i^{(2)} \Re(\bar{z}_1 z_i)}{2n^2 \beta_i^{(2)} \beta_1^{(1)}} \right. \\
& \left. + \frac{8cd\gamma m_i^{(1)} m_1^{(2)} u_i^{(1)} u_1^{(2)} \Re(\bar{z}_1 z_i)}{2n^2 \beta_i^{(1)} \beta_1^{(2)}} \right\} \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] + error_2 \\
& + \frac{cd\rho}{2n^2} \sum_{i \neq 1} (-2) \{ \partial_{\eta_1} (m_1^{(1)}) \partial_{\eta_i} (m_i^{(2)}) + \partial_{\eta_1} (m_1^{(2)}) \partial_{\eta_i} (m_i^{(1)}) \} \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] \\
& + error_2 + error_4 \\
(6.41) \quad = & cT_5^{(1)} + dT_5^{(2)} + c \frac{\iota(\kappa_4)_{11}}{4n} \partial_{\eta_1} (m_1^{(1)})^4 \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] + d \frac{\iota(\kappa_4)_{22}}{4n} \partial_{\eta_1} (m_1^{(2)})^4 \\
& \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] + \sum_{i \neq 1} \frac{c^2 \widehat{V}_{1i}^{(1)}}{2n^2} \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] + \sum_{i \neq 1} \frac{d^2 \widehat{V}_{1i}^{(2)}}{2n^2} \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] \\
& + \sum_{i \neq 1} \left\{ \frac{8cd\gamma m_i^{(2)} m_1^{(1)} u_1^{(1)} u_i^{(2)} \Re(\bar{z}_1 z_i)}{2n^2 \beta_i^{(2)} \beta_1^{(1)}} + \frac{8cd\gamma m_i^{(1)} m_1^{(2)} u_i^{(1)} u_1^{(2)} \Re(\bar{z}_1 z_i)}{2n^2 \beta_i^{(1)} \beta_1^{(2)}} \right\} \\
& \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] + \frac{cd\rho}{n^2} \sum_{i \neq 1} \left(- \partial_{\eta_1} (m_1^{(1)}) \partial_{\eta_i} (m_i^{(2)}) - \partial_{\eta_1} (m_1^{(2)}) \partial_{\eta_i} (m_i^{(1)}) \right)
\end{aligned}$$

$$\mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] + error_1 + error_2 + error_3 + error_4.$$

6.4. Evaluation of $T_5^{(t)}$ terms in (6.23). We evaluate $T_5^{(1)}$ on the RHS of (6.16) for $k = 2$ and $k \geq 3$ separately. Recall $T_5^{(1)}$ from (6.16)

$$(6.42) \quad \sum_{ab} \sum_{k \geq 2} \sum_{\alpha \in \{(ab)_1, (ba)_1, (ab)_2, (ba)_2\}^k} \frac{\kappa((ba)_1, \alpha)}{k!} \mathbb{E} \left[\partial_\alpha \left(\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i \right) \right].$$

Now, let us consider the case when $k = 2$. In that case, letting $\alpha = (\alpha_1, \alpha_2)$, we have the following;

$$(6.43) \quad \begin{aligned} & \partial_{\alpha_1, \alpha_2} \left(\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i \right) \\ &= \left(\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \right)'' \left(\prod_{i \neq 1} \Upsilon_i \right) \\ & \quad + \left(\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \right)' \left(\prod_{i \neq 1} \Upsilon_i \right)' \\ & \quad + \langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \left(\prod_{i \neq 1} \Upsilon_i \right)'' \\ &=: H_1 + H_2 + H_3. \end{aligned}$$

Recall from page 19 that ∂_{α_i} denotes the derivative with respect to $w_{\alpha_i}^{(t)}$. In general, we use the shorthand notation α_i to denote $w_{\alpha_i}^{(t)}$. The same convention is used in the above, where $\partial_{\alpha_i} f$ is denoted by f' , $\partial_{\alpha_i} \partial_{\alpha_j} f$ is denoted by f'' and so on.

Notice that H_1 is non-zero only if both α_1, α_2 , belong to $\{(ab)_1, (ba)_1\}$. In that case, we have the following expression;

$$(6.44) \quad \begin{aligned} H_1 &= \left(\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \right)'' \prod_{i \neq 1} \Upsilon_i \\ &= \left(\partial_{\alpha_1, \alpha_2} \langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \right) \prod_{i \neq 1} \Upsilon_i \\ &= \langle -\Delta^{ba} G_1^{(1)} (\Delta G_1^{(1)})^2 A_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i, \end{aligned}$$

where $(\Delta G_1^{(1)})^k = \Delta G_1^{(1)} \Delta G_1^{(1)} \cdots \Delta G_1^{(1)}$, and each Δ is either Δ^{ab} or Δ^{ba} depending on the context.

Similarly, H_2 is non-zero when at least one of α_1 and α_2 does not belong to $\{(ab)_2, (ba)_2\}$. Thus, taking derivatives with respect appropriate variables, we have

$$H_2 = \left(\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \right)' \left(\prod_{i \neq 1} \Upsilon_i \right)'$$

$$\begin{aligned}
(6.45) \quad &= \langle \Delta^{ba} G_1^{(1)} \Delta G_1^{(1)} A_1^{(1)} \rangle \sum_{j=2}^p \left(c \langle \underline{-W^{(1)} G_j^{(1)} \Delta G_j^{(1)} A_j^{(1)}} + \Delta G_j^{(1)} A_j^{(1)} \rangle \right) \prod_{i \neq 1, j} \Upsilon_i \\
&+ \langle \Delta^{ba} G_1^{(1)} \Delta G_1^{(1)} A_1^{(1)} \rangle \sum_{j=2}^p \left(d \langle \underline{-W^{(2)} G_j^{(2)} \Delta G_j^{(2)} A_j^{(2)}} + \Delta G_j^{(2)} A_j^{(2)} \rangle \right) \prod_{i \neq 1, j} \Upsilon_i.
\end{aligned}$$

Finally, no particular choice of α_1, α_2 completely nullifies H_3 . Therefore, taking all possible derivatives,

$$\begin{aligned}
(6.46) \quad &H_3 = \langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \left(\prod_{i \neq 1} \Upsilon_i \right)'' \\
&= \langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \sum_{j=2}^p \left(2c \langle \underline{W^{(1)} (G_j^{(1)} \Delta)^2 G_j^{(1)} A_j^{(1)}} - \Delta (G_j^{(1)})^2 A_j^{(1)} \rangle \right) \prod_{i \neq 1, j} \Upsilon_i \\
&+ \langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \sum_{j=2}^p \left(2d \langle \underline{W^{(2)} (G_j^{(2)} \Delta)^2 G_j^{(2)} A_j^{(2)}} - (\Delta G_j^{(2)})^2 A_j^{(2)} \rangle \right) \prod_{i \neq 1, j} \Upsilon_i \\
&+ \langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \sum_{\substack{j, l=2 \\ j \neq l}}^p \left(c \langle \underline{-W^{(1)} G_j^{(1)} \Delta G_l^{(1)} A_j^{(1)}} + \Delta G_j^{(1)} A_j^{(1)} \rangle \right) \\
&\quad \left(d \langle \underline{-W^{(2)} G_l^{(2)} \Delta G_l^{(2)} A_l^{(2)}} + \Delta G_l^{(2)} A_l^{(2)} \rangle \right) \prod_{i \neq 1, j, l} \Upsilon_i.
\end{aligned}$$

When we perform the α -derivatives in (6.42) using the Leibniz rule, we get sum of product of normalized traces. Each product in the sum is a product of p terms. Among those p terms, at least one or more (say, r many) are of the form $\langle (\Delta G_i^{(1)})^{k_i} A_i^{(1)} \rangle$, $\langle \underline{W^{(1)} (G_i^{(1)} \Delta)^{k_i} G_i^{(1)} A_i^{(1)}} \rangle$, or $\langle \underline{W^{(2)} (G_i^{(2)} \Delta)^{k_i} G_i^{(2)} A_i^{(2)}} \rangle$ with $k_i \geq 0$, $\sum k_i = k + 1$. Rest (i.e., $p - r$ many) are of the form $\langle \underline{W^{(1)} G_i^{(1)} A_i^{(1)} + \xi_i^{(1)}} \rangle + \langle \underline{W^{(2)} G_i^{(2)} A_i^{(2)} + \xi_i^{(2)}} \rangle$. For example, in the expression of H_3 , we have a term of the form $\langle \Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \langle (\Delta G_j^{(1)})^2 A_j^{(1)} \rangle \prod_{i \neq 1, j} \Upsilon_i$. For this particular term, $k = 2$ and $r = 2$.

In the rest of this section, we shall analyze the cases for different values of k and r . We shall see that the significant contribution comes only from the case when $k = 3$. Prior to that, we rewrite the self-renormalized terms as

$$\begin{aligned}
&\langle \underline{W^{(t)} (G^{(t)} \Delta)^k G^{(t)} A^{(t)}} \rangle \\
&= \langle \underline{G^{(t)} A^{(t)} W^{(t)} (G^{(t)} \Delta)^k} \rangle \\
&= \langle \underline{G^{(t)} A^{(t)} W^{(t)} G^{(t)} \Delta (G^{(t)} \Delta)^{k-1}} \rangle + \sum_{j=1}^{k-1} \langle G^{(t)} A^{(t)} S [(G^{(t)} \Delta)^j G^{(t)}] (G^{(t)} \Delta)^{k-j} \rangle \\
(6.47) \quad &= \langle \underline{G^{(t)} A^{(t)} W^{(t)} G^{(t)} \Delta (G^{(t)} \Delta)^{k-1}} \rangle + \sum_{j=1}^{k-1} \langle G^{(t)} A^{(t)} E (G^{(t)} \Delta)^{k-j} \rangle \\
&\quad \langle G^{(t)} E^\dagger (G^{(t)} \Delta)^j \rangle.
\end{aligned}$$

Now, we analyze the cases for different values of k and r .

6.4.1. *Contribution of (6.42) to $T_5^{(1)}$ when $k = 2, r = 1$.* This is essentially H_1 from (6.44). We argue that the term $\langle \Delta G_1^{(1)} \Delta G_1^{(1)} \Delta G_1^{(1)} A_1^{(1)} \rangle$ will have insignificant contribution. Since it is a product of three $\Delta^{\alpha_i} G_1^{(1)}$ s, and $\alpha_i \in \{(ab)_1, (ba)_1\}$, at least one of the term in this product will have an off-diagonal entry of $G_1^{(1)}$. For example, the term $\langle \Delta^{ba} G_1^{(1)} \Delta^{ab} G_1^{(1)} \Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle$ simplifies to $(2n)^{-1} \sum' (G_1^{(1)})_{aa} (G_1^{(1)})_{bb} (G_1^{(1)} A_1^{(1)})_{ab}$, which contains an off-diagonal entry. Such a term (along with the prefactor $n^{-3/2}$) can further be estimated by proceeding in the same way as in (6.9), as follows;

$$\begin{aligned}
(6.48) \quad & n^{-1-3/2} \sum_{a \leq n} \sum_{b > n} (G_1^{(1)})_{aa} (G_1^{(1)})_{bb} (G_1^{(1)} A_1^{(1)})_{ab} \\
&= \frac{m^2}{n^{5/2}} \langle E_1 \mathbf{1}, G_1^{(1)} A_1^{(1)} E_2 \mathbf{1} \rangle + \mathcal{O}_{\prec} \left(\frac{1}{n^{1/2}} \frac{1}{|\beta_1^{(1)}| (n\eta_1)^{3/2}} \right) \\
&= \mathcal{O}_{\prec} \left(\frac{1}{|\beta_1^{(1)}| n^{3/2}} + \frac{1}{|\beta_1^{(1)}| n^2 \eta_1^{3/2}} \right).
\end{aligned}$$

6.4.2. *Contribution of (6.42) to $T_5^{(1)}$ when $k = 2, r = 2$.* From (6.45) and (6.46), we see that this case consists of the expression of H_2 and the first two terms of H_3 , which is written as follows;

$$\begin{aligned}
(6.49) \quad & \langle \Delta^{ba} G_1^{(1)} \Delta G_1^{(1)} A_1^{(1)} \rangle \langle \Delta G_i^{(1)} A_i^{(1)} - \underline{W^{(1)} G_i^{(1)} \Delta G_i^{(1)} A_i^{(1)}} \rangle \\
&+ \langle \Delta^{ba} G_1^{(1)} \Delta G_1^{(1)} A_1^{(1)} \rangle \langle \Delta G_i^{(2)} A_i^{(2)} - \underline{W^{(2)} G_i^{(2)} \Delta G_i^{(2)} A_i^{(2)}} \rangle \\
&+ \langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle (2 \langle -\Delta G_i^{(1)} \Delta G_i^{(1)} A_i^{(1)} + \underline{W^{(1)} G_i^{(1)} \Delta G_i^{(1)} \Delta G_i^{(1)} A_i^{(1)}} \rangle) \\
&+ \langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle (2 \langle -\Delta G_i^{(2)} \Delta G_i^{(2)} A_i^{(2)} + \underline{W^{(2)} G_i^{(2)} \Delta G_i^{(2)} \Delta G_i^{(2)} A_i^{(2)}} \rangle).
\end{aligned}$$

In the rest of the subsection, we find the most contributing factors of the above expression. The essential tools are the isotropic local law, and accumulation of the diagonal terms from $G_1^{(1)}$ or $G_1^{(1)} A_1^{(1)}$ in the product, and the following estimate (6.50)

$$(6.50) \quad |\langle \mathbf{x}, \underline{G^{(t)} B W^{(t)} G^{(t)} \mathbf{y}} \rangle| \prec \frac{\|\mathbf{x}\| \|\mathbf{y}\| \|B\|}{n^{1/2} \eta^{3/2}},$$

which can be obtained by following the techniques in Lemma 6.6. Let's see how these principles are applied to obtain the estimates. For example, the highest contribution from the first four factors may arise from one of the following scenarios;

$$\begin{aligned}
(6.51) \quad & \frac{1}{n^{7/2}} \sum'_{ab} \left((G_1^{(1)})_{aa} (G_1^{(1)} A_1^{(1)})_{bb} \left((G_i^{(1)} A_i^{(1)})_{ab} - \underline{(G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)})_{ab}} \right) \right. \\
&\quad \left. + (G_1^{(1)})_{aa} (G_1^{(1)} A_1^{(1)})_{bb} \left((G_i^{(2)} A_i^{(2)})_{ab} - \underline{(G_i^{(2)} A_i^{(2)} W^{(2)} G_i^{(2)})_{ab}} \right) \right) \\
&= \mathcal{O}_{\prec} \left(\frac{1}{n^3 \eta_1 \eta_i^{3/2} |\beta_1^{(1)}| |\beta_i^{(1)}|} \right).
\end{aligned}$$

The above is obtained from the following estimates (6.52) and (6.53). The following calculations are similar to (6.9).

$$\begin{aligned}
(6.52) \quad & \frac{1}{n^{7/2}} \sum'_{ab} (G_1^{(1)})_{aa} (G_1^{(1)} A_1^{(1)})_{bb} (G_i^{(1)} A_i^{(1)})_{ab} \\
&= \frac{1}{n^{7/2}} \sum'_{ab} \left\{ (m_1^{(1)} + (G_1^{(1)} - M_1^{(1)})_{aa}) \left((M_1^{(1)} A_1^{(1)})_{bb} \right. \right. \\
&\quad \left. \left. + ((G_1^{(1)} - M_1^{(1)}) A_1^{(1)})_{bb} \right) \left((M_i^{(1)} A_i^{(1)})_{ab} + ((G_i^{(1)} - M_i^{(1)}) A_i^{(1)})_{ab} \right) \right\} \\
&= \frac{1}{n^{7/2}} \sum'_{ab} \left\{ m_1^{(1)} (M_1^{(1)} A_1^{(1)})_{bb} ((M_i^{(1)} A_i^{(1)})_{ab} + ((G_i^{(1)} - M_i^{(1)}) A_i^{(1)})_{ab}) \right. \\
&\quad + (G_1^{(1)} - M_1^{(1)})_{aa} (M_1^{(1)} A_1^{(1)})_{bb} ((M_i^{(1)} A_i^{(1)})_{ab} + ((G_i^{(1)} - M_i^{(1)}) A_i^{(1)})_{ab}) \\
&\quad + m_1^{(1)} ((G_1^{(1)} - M_1^{(1)}) A_1^{(1)})_{bb} ((M_i^{(1)} A_i^{(1)})_{ab} + ((G_i^{(1)} - M_i^{(1)}) A_i^{(1)})_{ab}) \\
&\quad \left. + ((G_1^{(1)} - M_1^{(1)}) A_1^{(1)})_{bb} (G_1^{(1)} - M_1^{(1)})_{aa} ((M_i^{(1)} A_i^{(1)})_{ab} \right. \\
&\quad \left. + ((G_i^{(1)} - M_i^{(1)}) A_i^{(1)})_{ab}) \right\} \\
&= \mathcal{O}_{\prec} \left(\frac{1}{n^{7/2}} \left(\frac{1}{|\beta_1^{(1)}|} \frac{n}{(1+\eta_i) |\beta_i^{(1)}|} + \frac{1}{|\beta_1^{(1)}|} \frac{n^2}{\sqrt{n\eta_i} |\beta_i^{(1)}|} + \frac{1}{|\beta_1^{(1)}| |\beta_i^{(1)}|} \frac{n}{\eta_1} \right. \right. \\
&\quad \left. \left. + \frac{1}{|\beta_1^{(1)}| |\beta_i^{(1)}|} \frac{n}{\eta_1} \frac{1}{\sqrt{n\eta_i}} + \frac{1}{|\beta_1^{(1)}| |\beta_i^{(1)}|} \frac{n}{\eta_1} + \frac{1}{|\beta_1^{(1)}| |\beta_i^{(1)}|} \frac{n}{\eta_1} \frac{1}{\sqrt{n\eta_i}} \right. \right. \\
&\quad \left. \left. + \frac{1}{|\beta_1^{(1)}| |\beta_i^{(1)}|} \frac{n}{\eta_1} \frac{1}{\sqrt{n\eta_1}} + \frac{1}{|\beta_1^{(1)}| |\beta_i^{(1)}|} \frac{n}{\eta_1} \frac{1}{\sqrt{n\eta_1} \sqrt{n\eta_i}} \right) \right) \\
&= \mathcal{O}_{\prec} \left(\frac{1}{n^3 \eta_1 \eta_i^{1/2} |\beta_1^{(1)}| |\beta_i^{(1)}|} \right),
\end{aligned}$$

and

$$\begin{aligned}
(6.53) \quad & \frac{1}{n^{7/2}} \sum'_{ab} (G_1^{(1)})_{aa} (G_1^{(1)} A_1^{(1)})_{bb} \underline{(G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)})_{ab}} \\
&= \frac{1}{n^{7/2}} \sum'_{ab} \left\{ m_1^{(1)} (M_1^{(1)} A_1^{(1)})_{bb} \underline{(G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)})_{ab}} \right. \\
&\quad + (G_1^{(1)} - M_1^{(1)})_{aa} (M_1^{(1)} A_1^{(1)})_{bb} \underline{(G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)})_{ab}} \\
&\quad + m_1^{(1)} (G_1^{(1)} - M_1^{(1)}) A_1^{(1)}_{bb} \underline{(G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)})_{ab}} \\
&\quad \left. + (G_1^{(1)} - M_1^{(1)}) A_1^{(1)}_{bb} (G_1^{(1)} - M_1^{(1)})_{aa} \underline{(G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)})_{ab}} \right\} \\
&= \mathcal{O}_{\prec} \left(\frac{1}{n^{7/2}} \frac{1}{|\beta_1^{(1)}|} \frac{n}{n^{1/2} \eta_i^{3/2} |\beta_i^{(1)}|} + \frac{1}{n^{7/2}} \left(\frac{1}{\sqrt{n\eta_1}} \frac{1}{|\beta_1^{(1)}|} \frac{n}{n^{1/2} \eta_i^{3/2} |\beta_i^{(1)}|} \right. \right.
\end{aligned}$$

$$\begin{aligned}
& + \frac{n^2}{n\eta_1|\beta_1^{(1)}|} \frac{1}{n^{1/2}\eta_i^{3/2}|\beta_i^{(1)}|} + \frac{n^2}{n\eta_1|\beta_1^{(1)}|} \frac{1}{\sqrt{n\eta_1}} \frac{1}{n^{1/2}\eta_i^{3/2}|\beta_i^{(1)}|} \Big) \\
& = \mathcal{O}_{\prec} \left(\frac{1}{n^3\eta_1\eta_i^{3/2}|\beta_1^{(1)}||\beta_i^{(1)}|} \right).
\end{aligned}$$

Similarly, by using (6.47), (6.50), and the estimates from Lemma 4.1, we obtain the following estimate for the remaining terms of (6.49);

$$\begin{aligned}
(6.54) \quad & \frac{1}{n^{7/2}} \mathbb{E} \sum'_{ab} (G_1^{(1)} A_1^{(1)})_{ab} \left(\underline{(G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)})_{ab}} (G_i^{(1)})_{bb} \right. \\
& \quad \left. + \frac{1}{n} (G_i^{(1)} A_i^{(1)} E G_i^{(1)})_{ab} (G_i^{(1)} E^\dagger G_i^{(1)})_{ab} \right) + \frac{1}{n^{7/2}} \mathbb{E} \sum'_{ab} (G_1^{(1)} A_1^{(1)})_{ab} \\
& \quad \left(\underline{(G_i^{(2)} A_i^{(2)} W^{(2)} G_i^{(2)})_{ab}} (G_i^{(2)})_{bb} + \frac{1}{n} (G_i^{(2)} A_i^{(2)} E G_i^{(2)})_{ab} (G_i^{(2)} E^\dagger G_i^{(2)})_{ab} \right) \\
& = \mathcal{O}_{\prec} \left(\frac{1}{n^3\eta_1^{1/2}\eta_i^2|\beta_i^{(1)}||\beta_1^{(1)}|} \right).
\end{aligned}$$

6.4.3. *Contribution of (6.42) to $T_5^{(1)}$ when $k = 2, r = 3$.* This case consists of the remaining terms of H_3 (see (6.46)), which were not considered in the previous cases (i.e., $k = 2, r \leq 2$). The expression becomes

$$\begin{aligned}
& \langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \langle \Delta G_i^{(1)} A_i^{(1)} - \underline{W^{(1)} G_i^{(1)} \Delta G_i^{(1)} A_i^{(1)}} \rangle \langle \Delta G_j^{(1)} A_j^{(1)} - \underline{W^{(1)} G_j^{(1)} \Delta G_j^{(1)} A_j^{(1)}} \rangle \\
& + \langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \langle \Delta G_i^{(2)} A_i^{(2)} - \underline{W^{(2)} G_i^{(2)} \Delta G_i^{(2)} A_i^{(2)}} \rangle \langle \Delta G_j^{(2)} A_j^{(2)} - \underline{W^{(2)} G_j^{(2)} \Delta G_j^{(2)} A_j^{(2)}} \rangle \\
& + \langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \langle \Delta G_i^{(1)} A_i^{(1)} - \underline{W^{(1)} G_i^{(1)} \Delta G_i^{(1)} A_i^{(1)}} \rangle \langle \Delta G_j^{(2)} A_j^{(2)} - \underline{W^{(2)} G_j^{(2)} \Delta G_j^{(2)} A_j^{(2)}} \rangle
\end{aligned}$$

which, using (6.50), we estimate by

$$\begin{aligned}
(6.55) \quad & \frac{1}{n^{9/2}} \sum'_{ab} \left\{ - (G_1^{(1)} A_1^{(1)})_{ab} \left((G_i^{(1)} A_i^{(1)})_{ab} - \underline{(G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)})_{ab}} \right) \right. \\
& \quad \left((G_j^{(1)} A_j^{(1)})_{ab} - \underline{(G_j^{(1)} A_j^{(1)} W^{(1)} G_j^{(1)})_{ab}} \right) - (G_1^{(1)} A_1^{(1)})_{ab} \left((G_i^{(2)} A_i^{(2)})_{ab} \right. \\
& \quad \left. - \underline{(G_i^{(2)} A_i^{(2)} W^{(2)} G_i^{(2)})_{ab}} \right) \left((G_j^{(2)} A_j^{(2)})_{ab} - \underline{(G_j^{(2)} A_j^{(2)} W^{(2)} G_j^{(2)})_{ab}} \right) \\
& \quad - (G_1^{(1)} A_1^{(1)})_{ab} \left((G_i^{(1)} A_i^{(1)})_{ab} - \underline{(G_i^{(1)} A_i^{(1)} W^{(1)} G_i^{(1)})_{ab}} \right) \left((G_j^{(2)} A_j^{(2)})_{ab} \right. \\
& \quad \left. \left. - \underline{(G_j^{(2)} A_j^{(2)} W^{(2)} G_j^{(2)})_{ab}} \right) \right\} \\
& = \mathcal{O}_{\prec} \left(\frac{1}{n^4\eta_1^{1/2}\eta_i^{3/2}\eta_j^{3/2}|\beta_1^{(1)}||\beta_i^{(1)}||\beta_j^{(2)}|} \right).
\end{aligned}$$

In summary, from the (6.48), (6.51), (6.54), and (6.55), we have the essential estimates for the terms of (6.42) when $k = 2$. However, none of estimates consider the attached partial product $\prod \Upsilon_i$, where some of Υ_i s are omitted. But this can be bounded by some form of the estimate given in (6.30). Consequently, the updated estimates of (6.48), (6.51), (6.54),

and (6.55) will become as

$$\mathcal{O}_{\prec}\left(\frac{\psi}{n^{3/2}\eta_1}\right), \mathcal{O}_{\prec}\left(\frac{\psi}{n^2\eta_i\sqrt{\eta_1}}\right), \mathcal{O}_{\prec}\left(\frac{\psi}{n^2\eta_i^{3/2}}\right), \text{ and } \mathcal{O}_{\prec}\left(\frac{\psi}{n^{5/2}\eta_i\eta_j}\right)$$

respectively. Hence, the final contribution of (6.42) in the case of $k = 2$ is $\mathcal{O}_{\prec}\left(\frac{\psi}{n^{3/2}\eta_*}\right)$.

6.4.4. *Contribution of (6.42) to $T_5^{(1)}$ when $k \geq 3$.* Now, for $k = 3$, the RHS term of (6.42) is

$$(6.56) \quad \sum_{ab} \sum_{\alpha \in \{(ab)_1, (ba)_1, (ab)_2, (ba)_2\}^3} \frac{\kappa((ba)_1, \alpha)}{3!} \mathbb{E} \left[\partial_\alpha \left(\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i \right) \right].$$

Before considering this case, let us look at the following remark which summarizes the estimate types from previous three subsections.

Remark 6.4. (1) *If we have the following sum;*

$$\sum_{ab} \prod_{i=1}^l \Delta^{\alpha_i} G_i^{(t)},$$

where $\alpha_i \in \{ab, ba\}$, then we have the following two cases.

- If l is odd, no product term in the sum can consist entirely of diagonal entries; there must be at least one off-diagonal entry in each term.
 - If l is even, it is possible to have a product term where all the entries are diagonal.
- (2) *If we have entries such as $(G_i^{(t)})_{ab}$ or $(G_i^{(t)} A_i^{(t)})_{ab}$, then these are decomposed as $(M_i^{(t)})_{ab} + (G_i^{(t)} - M_i^{(t)})_{ab}$ or $(M_i^{(t)} A_i^{(t)})_{ab} + ((G_i^{(t)} - M_i^{(t)}) A_i^{(t)})_{ab}$ respectively. Now, the terms $(M_i^{(t)})_{ab}$ or $(M_i^{(t)} A_i^{(t)})_{ab}$ yield a significant contribution if these are diagonal terms. Therefore if a product term contains a off-diagonal term, it does not give any significant contribution, which is further explained in the following point.*
- (3) *If we have off-diagonal factors such as $(G_i^{(t)} - M_i^{(t)})_{ab}$, $((G_i^{(t)} - M_i^{(t)}) A_i^{(t)})_{ab}$, $(G_i^{(t)} A_i^{(t)} W^{(t)} G_i^{(t)})_{ab}$ etc. in the product terms, then using the local law (Theorem 4.1) or (6.50), we argue that such terms do not give any contribution. This was demonstrated in the derivations (6.48), (6.51), (6.54), and (6.55).*
- (4) *In (6.42), after performing the α -derivatives, we obtain a sum of product of terms, where each of the product is a product of p -terms. Essentially, the terms in the product can be categorized into two categories (a) the terms in $\prod_i \Upsilon_i$ which are affected by the derivatives (b) terms in $\prod_i \Upsilon_i$, which are unaffected by the derivatives. The terms, which are affected by the differentiation, will be a normalized trace of the form $\langle (\Delta G_i^{(t)})^{k_i} A_i \rangle$ or $\langle W^{(t)} (G_i^{(t)} \Delta)^{k_i} G_i^{(t)} A_i^{(t)} \rangle$, where $k_i \in \mathbb{N}$. Moreover, the first term $\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle$ is also a normalized trace. Overall if there are r many normalized traces, this will give a pre-factor of $n^{-r-(k+1)/2}$ (including the cumulant estimate (6.5)). This is also mentioned on page 38. Additionally, $\sum_{i=1}^r k_i = k + 1$.*
- (5) *Now, if the product contains all the diagonal entries (similar to (6.10)), we may get some contribution. In such cases, we still have the pre-factor $n^{-r-(k+1)/2}$. Giving a naive bound (of constant order) to each of those diagonal entries, the prefactor will get reduced to $n^{-r-(k-3)/2}$. We loose a factor of n^2 because the sum \sum'_{ab} contains $2n^2$*

many terms. From here, it is clear that the leading order contribution comes from the case $k = 3$.

Now, for $k = 3$ proceeding as (6.43), we can identify the terms which contain all diagonal entries. The terms which can possibly contain all diagonal entries are of the following forms;

$$\langle \Delta^{ba} G_1^{(1)} \Delta G_1^{(1)} \Delta G_1^{(1)} \Delta G_1^{(1)} A_1^{(1)} \rangle, \quad \langle \Delta^{ba} G_1^{(1)} \Delta G_1^{(1)} A_1^{(1)} \rangle \langle \Delta G_i^{(t)} \Delta G_i^{(t)} A_i^{(t)} \rangle.$$

For the second types of terms, we notice that $\alpha = ((ab)_1, (ab)_1, (ba)_1)$ yields a terms having product of diagonal entries, which can be evaluated as follows;

$$\begin{aligned} & \kappa((ba)_1, (ab)_1, (ab)_1, (ba)_1) \sum'_{ab} \langle \Delta^{ba} G_1^{(1)} \Delta^{ab} G_1^{(1)} A_1^{(1)} \rangle \langle \Delta^{ab} G_i^{(1)} \Delta^{ba} G_i^{(1)} A_i^{(1)} \rangle \\ &= \frac{(\kappa_4)_{11}}{4n^4} \sum'_{ab} (G_1^{(1)})_{aa} (G_1^{(1)} A_1^{(1)})_{bb} (G_i^{(1)})_{bb} (G_i^{(1)} A_i^{(1)})_{aa} \\ &= \frac{(\kappa_4)_{11}}{4n^4} \sum'_{ab} m_1^{(1)} m_i^{(1)} (M_1^{(1)} A_1^{(1)})_{bb} (M_i^{(1)} A_i^{(1)})_{aa} + \mathcal{O}_{\prec} \left(\frac{1}{n^{5/2} \eta_*^{1/2} |\beta_{1i}^*|^2} \right) \\ &= \frac{(\kappa_4)_{11}}{2n^2} \langle M_1^{(1)} \rangle \langle M_i^{(1)} \rangle \langle M_1^{(1)} A_1^{(1)} \rangle \langle M_i^{(1)} A_i^{(1)} \rangle + \mathcal{O}_{\prec} \left(\frac{1}{n^{5/2} \eta_*^{1/2} |\beta_{1i}^*|^2} \right). \end{aligned}$$

The other choices of α which give a significant contribution are $\alpha = ((ab)_1, (ba)_1, (ab)_1)$, $((ab)_1, (ab)_2, (ba)_2)$, $((ab)_1, (ba)_2, (ab)_2)$. Combining all the contributions, we obtain

$$\begin{aligned} & \sum_{ab} \sum_{\alpha} \kappa((ba)_1, \alpha) \langle \Delta^{ba} G_1^{(1)} \Delta^{\alpha_1} G_1^{(1)} \rangle \langle \Delta^{\alpha_2} G_i^{(t)} \Delta^{\alpha_3} G_i^{(t)} A_i^{(t)} \rangle \\ &= \frac{(\kappa_4)_{11}}{n^2} \langle M_1^{(1)} \rangle \langle M_i^{(1)} \rangle \langle M_1^{(1)} A_1^{(1)} \rangle \langle M_i^{(1)} A_i^{(1)} \rangle \\ & \quad + \frac{(\kappa_4)_{12}}{n^2} \langle M_1^{(1)} \rangle \langle M_i^{(2)} \rangle \langle M_1^{(1)} A_1^{(1)} \rangle \langle M_i^{(2)} A_i^{(2)} \rangle \\ & \quad + \mathcal{O}_{\prec} \left(\frac{1}{n^{5/2} \eta_*^{1/2} |\beta_{1i}^*|^2} \right) + \mathcal{O}_{\prec} \left(\frac{1}{n^3 \eta_* |\beta_{1i}^*|^2} \right) \\ &=: \frac{(\kappa_4)_{11}}{2n^2} U_1^{(1)} U_i^{(1)} + \frac{(\kappa_4)_{12}}{2n^2} U_1^{(1)} U_i^{(2)} + \mathcal{O}_{\prec} \left(\frac{1}{n^{5/2} \eta_*^{1/2} |\beta_{1i}^*|^2} \right), \end{aligned}$$

where,

$$(6.57) \quad U_i^{(t)} := -\sqrt{2} \langle M_i^{(t)} \rangle \langle M_i^{(t)} A_i^{(t)} \rangle = \frac{\iota}{\sqrt{2}} \partial_{\eta_i} (m_i^{(t)})^2.$$

Similarly, for the first one, only $\alpha = ((ab)_1, (ba)_1, (ab)_1)$ gives significant contribution as follows;

$$\begin{aligned} & \sum_{ab} \sum_{\alpha} \kappa((ba)_1, \alpha) \langle \Delta^{ba} G_1^{(1)} \Delta^{ab} G_1^{(1)} \Delta^{ba} G_1^{(1)} \Delta^{ab} G_1^{(1)} A_1^{(1)} \rangle \\ &= \frac{(\kappa_4)_{11}}{n} \langle M_1^{(1)} \rangle^3 \langle M_1^{(1)} A_1^{(1)} \rangle + \mathcal{O}_{\prec} \left(\frac{1}{n^{3/2} \eta_1^{1/2} |\beta_1^{(1)}|} \right) + \mathcal{O}_{\prec} \left(\frac{1}{n^2 \eta_1 |\beta_1^{(1)}|} \right) \\ (6.58) \quad &= -\frac{\iota(\kappa_4)_{11}}{4n} \partial_{\eta_1} (m_1^{(1)})^4 + \mathcal{O}_{\prec} \left(\frac{1}{n^2 \eta_1 |\beta_1^{(1)}|} \right). \end{aligned}$$

The above calculation is same as (6.10). Thus, we can conclude for the $k \geq 3$ terms in (6.15) that

$$\begin{aligned}
(6.59) \quad & \sum_{ab} \sum_{k \geq 3} \sum_{\alpha \in \{(ab)_1, (ba)_1, (ab)_2, (ba)_2\}^k} \frac{\kappa((ba)_1, \alpha)}{k!} \mathbb{E} \left[\partial_\alpha \left(\langle -\Delta^{ba} G_1^{(1)} A_1^{(1)} \rangle \prod_{i \neq 1} \Upsilon_i \right) \right] \\
&= -\frac{\iota(\kappa_4)_{11}}{4n} \partial_{\eta_1} (m_1^{(1)})^4 \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] + \sum_{i \neq 1} \frac{1}{2n^2} (c(\kappa_4)_{11} U_1^{(1)} U_i^{(1)} + d(\kappa_4)_{12} U_1^{(1)} U_i^{(2)}) \\
&\quad \mathbb{E} \left[\prod_{j \neq i, 1} \Upsilon_j \right] + \mathcal{O}_{\prec} \left(\frac{1}{n^2 \eta_1 |\beta_1^{(1)}|} \right) \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] + \mathcal{O}_{\prec} \left(\frac{1}{n^{5/2} \eta_*^{1/2} |\beta_{1i}^*|^2} \right) \mathbb{E} \left[\prod_{j \neq i, 1} \Upsilon_j \right] \\
&= -\frac{\iota(\kappa_4)_{11}}{4n} \partial_{\eta_1} (m_1^{(1)})^4 \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] + \sum_{i \neq 1} \frac{1}{2n^2} (c(\kappa_4)_{11} U_1^{(1)} U_i^{(1)} + d(\kappa_4)_{12} U_1^{(1)} U_i^{(2)}) \\
&\quad \mathbb{E} \left[\prod_{j \neq i, 1} \Upsilon_j \right] + \mathcal{O}_{\prec} \left(\frac{\psi}{n^{3/2} \eta_1^{1/2}} + \frac{\psi(\eta_*)^{1/2}}{n^2} \right).
\end{aligned}$$

Thus, from page 41 and (6.59) we have

$$\begin{aligned}
(6.60) \quad & T_5^{(1)} \\
&= -\frac{\iota(\kappa_4)_{11}}{4n} \partial_{\eta_1} (m_1^{(1)})^4 \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] + \sum_{i \neq 1} \frac{1}{2n^2} (c(\kappa_4)_{11} U_1^{(1)} U_i^{(1)} \\
&\quad + d(\kappa_4)_{12} U_1^{(1)} U_i^{(2)}) \mathbb{E} \left[\prod_{j \neq i, 1} \Upsilon_j \right] + \mathcal{O}_{\prec} \left(\frac{\psi}{n^{3/2} \eta_*^{1/2}} \right).
\end{aligned}$$

Similarly, we can evaluate $T_5^{(2)}$ and we have the final expression for $T_5^{(1)} + T_5^{(2)}$ as follows;

$$\begin{aligned}
(6.61) \quad & T_5^{(1)} + T_5^{(2)} \\
&= -\frac{\iota(\kappa_4)_{11}}{4n} \partial_{\eta_1} (m_1^{(1)})^4 \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] + \sum_{i \neq 1} \frac{1}{2n^2} (c(\kappa_4)_{11} U_1^{(1)} U_i^{(1)} \\
&\quad + d(\kappa_4)_{12} U_1^{(1)} U_i^{(2)}) \mathbb{E} \left[\prod_{j \neq i, 1} \Upsilon_j \right] - \frac{\iota(\kappa_4)_{22}}{4n} \partial_{\eta_1} (m_1^{(2)})^4 \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] \\
&\quad + \sum_{i \neq 1} \frac{1}{2n^2} (c(\kappa_4)_{22} U_1^{(2)} U_i^{(2)} + d(\kappa_4)_{21} U_1^{(2)} U_i^{(1)}) \mathbb{E} \left[\prod_{j \neq i, 1} \Upsilon_j \right] + \mathcal{O}_{\prec} \left(\frac{\psi}{n^{3/2} \eta_*^{1/2}} \right) \\
&= -\frac{\iota(\kappa_4)_{11}}{4n} \partial_{\eta_1} (m_1^{(1)})^4 \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] + \sum_{i \neq 1} \frac{1}{2n^2} (c(\kappa_4)_{11} U_1^{(1)} U_i^{(1)} \\
&\quad + d(\kappa_4)_{12} U_1^{(1)} U_i^{(2)}) \mathbb{E} \left[\prod_{j \neq i, 1} \Upsilon_j \right] - \frac{\iota(\kappa_4)_{22}}{4n} \partial_{\eta_1} (m_1^{(2)})^4 \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] \\
&\quad + \sum_{i \neq 1} \frac{1}{2n^2} (c(\kappa_4)_{22} U_1^{(2)} U_i^{(2)} + d(\kappa_4)_{21} U_1^{(2)} U_i^{(1)}) \mathbb{E} \left[\prod_{j \neq i, 1} \Upsilon_j \right] + error_5.
\end{aligned}$$

6.5. **Concluding the proof of the Theorem 3.3.** From (6.41) and (6.61) we have

$$\begin{aligned}
& \mathbb{E} \left[\prod_{i \in [p]} \Upsilon_i \right] \\
&= c \left\{ -\frac{\iota(\kappa_4)_{11}}{4n} \partial_{\eta_1} (m_1^{(1)})^4 \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] + \sum_{i \neq 1} \frac{1}{2n^2} (c(\kappa_4)_{11} U_1^{(1)} U_i^{(1)} \right. \\
&\quad \left. + d(\kappa_4)_{12} U_1^{(1)} U_i^{(2)}) \mathbb{E} \left[\prod_{j \neq i, 1} \Upsilon_j \right] \right\} + d \left\{ -\frac{\iota(\kappa_4)_{22}}{4n} \partial_{\eta_1} (m_1^{(2)})^4 \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] \right. \\
&\quad \left. + \sum_{i \neq 1} \frac{1}{2n^2} (c(\kappa_4)_{22} U_1^{(2)} U_i^{(2)} + d(\kappa_4)_{21} U_1^{(2)} U_i^{(1)}) \mathbb{E} \left[\prod_{j \neq i, 1} \Upsilon_j \right] \right\} \\
&\quad + c \frac{\iota(\kappa_4)_{11}}{4n} \partial_{\eta_1} (m_1^{(1)})^4 \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] + d \frac{\iota(\kappa_4)_{22}}{4n} \partial_{\eta_1} (m_1^{(2)})^4 \mathbb{E} \left[\prod_{i \neq 1} \Upsilon_i \right] \\
&\quad + \sum_{i \neq 1} \frac{c^2 \widehat{V}_{1i}^{(1)}}{2n^2} \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] + \sum_{i \neq 1} \frac{d^2 \widehat{V}_{1i}^{(2)}}{2n^2} \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] \\
&\quad + \sum_{i \neq 1} \left\{ \frac{8cd\gamma m_i^{(2)} m_1^{(1)} u_1^{(1)} u_i^{(2)} \Re(\bar{z}_1 z_i)}{2n^2 \beta_i^{(2)} \beta_1^{(1)}} + \frac{8cd\gamma m_i^{(1)} m_1^{(2)} u_1^{(1)} u_i^{(2)} \Re(\bar{z}_1 z_i)}{2n^2 \beta_i^{(1)} \beta_1^{(2)}} \right\} \\
&\quad \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] + \frac{cd\rho}{n^2} \sum_{i \neq 1} \left(-\partial_{\eta_1} (m_1^{(1)}) \partial_{\eta_i} (m_i^{(2)}) - \partial_{\eta_1} (m_1^{(2)}) \partial_{\eta_i} (m_i^{(1)}) \right) \\
&\quad \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] + error_1 + error_2 + error_3 + error_4 + error_5 \\
&= \sum_{i \neq 1} \frac{1}{2n^2} (c^2(\kappa_4)_{11} U_1^{(1)} U_i^{(1)} + d^2(\kappa_4)_{22} U_1^{(2)} U_i^{(2)}) \mathbb{E} \left[\prod_{j \neq i, 1} \Upsilon_j \right] \\
&\quad + \sum_{i \neq 1} \frac{cd(\kappa_4)_{12}}{2n^2} (U_1^{(2)} U_i^{(1)} + U_1^{(1)} U_i^{(2)}) \mathbb{E} \left[\prod_{j \neq i, 1} \Upsilon_j \right] + \sum_{i \neq 1} \frac{c^2 \widehat{V}_{1i}^{(1)}}{2n^2} \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] \\
&\quad + \sum_{i \neq 1} \frac{d^2 \widehat{V}_{1i}^{(2)}}{2n^2} \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] + \sum_{i \neq 1} \left\{ \frac{8cd\gamma m_i^{(2)} m_1^{(1)} u_1^{(1)} u_i^{(2)} \Re(\bar{z}_1 z_i)}{2n^2 \beta_i^{(2)} \beta_1^{(1)}} \right. \\
&\quad \left. + \frac{8cd\gamma m_i^{(1)} m_1^{(2)} u_1^{(1)} u_i^{(2)} \Re(\bar{z}_1 z_i)}{2n^2 \beta_i^{(1)} \beta_1^{(2)}} \right\} \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] \\
&\quad + \frac{cd\rho}{n^2} \sum_{i \neq 1} \left(-\partial_{\eta_1} (m_1^{(1)}) \partial_{\eta_i} (m_i^{(2)}) - \partial_{\eta_1} (m_1^{(2)}) \partial_{\eta_i} (m_i^{(1)}) \right) \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] \\
&\quad + error_1 + error_2 + error_3 + error_4 + error_5 \\
&=: \sum_{i \neq 1} \frac{c^2 \widehat{V}_{1i}^{(1)} + d^2 \widehat{V}_{1i}^{(2)} + 8cd\gamma(L_{i1}^{12} + L_{i1}^{21}) + 2cd\rho N_{1i}}{2n^2} \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right]
\end{aligned}$$

$$\begin{aligned}
& + \sum_{i \neq 1} \frac{c^2(\kappa_4)_{11} U_1^{(1)} U_i^{(1)} + d^2(\kappa_4)_{22} U_1^{(2)} U_i^{(2)} + cd(\kappa_4)_{12} (U_1^{(1)} U_i^{(2)} + U_i^{(1)} U_1^{(2)})}{2n^2} \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] \\
& + \text{error}_1 + \text{error}_2 + \text{error}_3 + \text{error}_4 + \text{error}_5 \\
& =: \frac{1}{2n^2} \sum_{i \neq 1} \left\{ \left(c^2 \widehat{V}_{1i}^{(1)} + d^2 \widehat{V}_{1i}^{(2)} + 8cd\gamma(L_{i1}^{12} + L_{i1}^{21}) + 2cd\rho N_{1i} + c^2(\kappa_4)_{11} U_1^{(1)} U_i^{(1)} \right. \right. \\
& \left. \left. + d^2(\kappa_4)_{22} U_1^{(2)} U_i^{(2)} + cd(\kappa_4)_{12} (U_1^{(1)} U_i^{(2)} + U_i^{(1)} U_1^{(2)}) \right) \mathbb{E} \left[\prod_{j \neq 1, i} \Upsilon_j \right] \right\} \\
& + \mathcal{O}_{\prec} \left(\psi \left(\frac{1}{(n\eta_*)^2} + \frac{1}{n^2 \eta_*^{1i} |z_1 - z_i|^4} + \frac{1}{(n\eta_*^{1i})^3 |z_1 - z_i|^4} + \frac{1}{\Im z_1 (n\eta_1)^{3/2}} \right) \right),
\end{aligned}$$

where

$$(6.62) \quad L_{i1}^{12} = \frac{m_i^{(1)} m_1^{(2)} u_i^{(1)} u_1^{(2)} \Re(\overline{z_1} z_i)}{\beta_i^{(1)} \beta_1^{(2)}}, \quad L_{i1}^{21} = \frac{m_i^{(2)} m_1^{(1)} u_i^{(2)} u_1^{(1)} \Re(\overline{z_1} z_i)}{\beta_i^{(2)} \beta_1^{(1)}},$$

and

$$(6.63) \quad N_{1i} = \partial_{\eta_1} (m_1^{(1)}) \partial_{\eta_i} (m_i^{(2)}) + \partial_{\eta_1} (m_1^{(2)}) \partial_{\eta_i} (m_i^{(1)}).$$

Thus, by induction we have

$$\begin{aligned}
(6.64) \quad & \mathbb{E} \left[\prod_{i \in [p]} \Upsilon_i \right] \\
& = \frac{1}{2n^p} \sum_{P \in \Pi_p} \prod_{\{i, j\} \in P} \left\{ c^2 \widehat{V}_{ij}^{(1)} + d^2 \widehat{V}_{ij}^{(2)} + 8cd\gamma(L_{ji}^{12} + L_{ji}^{21}) + 2cd\rho N_{ij} \right. \\
(6.65) \quad & \left. + c^2(\kappa_4)_{11} U_i^{(1)} U_j^{(1)} + d^2(\kappa_4)_{22} U_i^{(2)} U_j^{(2)} + cd(\kappa_4)_{12} (U_i^{(1)} U_j^{(2)} + U_j^{(1)} U_i^{(2)}) \right\} \\
& + \mathcal{O}_{\prec} \left(\psi \left(\frac{1}{(n\eta_*)^2} + \frac{1}{n^2 \eta_* |z_i - z_j|^4} + \frac{1}{(n\eta_*)^3 |z_i - z_j|^4} + \frac{1}{\Im z_i (n\eta_*)^{3/2}} \right) \right),
\end{aligned}$$

where $\eta_* = \min_i \eta_i$.

Now, we will estimate the bound (6.27) in Lemma 6.6. However, before proceeding, we will state the following lemma;

Lemma 6.5. [14, Lemma 5.8] *Let ω_i and z_i be arbitrary spectral parameters with $\eta_i = \Im(\omega_i) > 0$. Define $G_j^{(t)} = (G^{(t)})^{z_j}(\omega_j)$, and let $G_{j_1, \dots, j_k}^{(t)}$ denote products of resolvents (or their adjoints/transposes) interspersed with bounded deterministic matrices, e.g., $G_{1i_1}^{(t)} = A_1^{(t)} G_1^{(t)} A_2^{(t)} G_i^{(t)} A_3^{(t)} G_1^{(t)} A_4^{(t)}$. Then for j_1, j_2, \dots, j_k and any $1 \leq s < t \leq k$, we obtain the following bounds;*

(1) *Isotropic Bound*

$$|\langle \mathbf{x}, G_{j_1, \dots, j_k}^{(t)} \mathbf{y} \rangle| \prec \|\mathbf{x}\| \|\mathbf{y}\| \sqrt{\eta_{j_1} \eta_{j_k}} \left(\prod_{i=1}^k \eta_{j_i}^{-1} \right),$$

(2) Averaged Bound

$$|\langle G_{j_1, \dots, j_k}^{(t)} \rangle| \prec \sqrt{\eta_{j_s} \eta_{j_t}} \left(\prod_{i=1}^k \eta_{j_i} \right)^{-1}.$$

Lemma 6.6. *Let $G_i^{(t)}$, $G_{j_1, \dots, j_k}^{(t)}$ be same as defined in Lemma 6.5, and $A_i^{(t)}$ be deterministic matrices with $\|A_i^{(t)}\| \leq |\beta_i^{(t)}|^{-1}$. Then we have the following estimate;*

$$\begin{aligned} & \mathbb{E}[|\langle \underline{G_1^{(t)} A_1^{(t)} E G_i^{(s)} A_i^{(s)} W^{(s)} G_i^{(s)} E^\dagger} \rangle|^2] \\ &= \mathcal{O}_{\prec} \left(\frac{1}{n^2 \eta_i^4 |\beta_1^{(t)}|^2 |\beta_i^{(s)}|^2} \right), \text{ for } 1 \leq t \neq s \leq 2. \end{aligned}$$

Proof. Consider

$$\begin{aligned} & \mathbb{E}[|\langle \underline{G_1^{(1)} A_1^{(1)} E G_i^{(2)} A_i^{(2)} W^{(2)} G_i^{(2)} E^\dagger} \rangle|^2] \\ &= \mathbb{E}[|\langle \underline{W^{(2)} G_i^{(2)} E^\dagger G_1^{(1)} A_1^{(1)} E G_i^{(2)} A_i^{(2)}} \rangle|^2] \\ &= \mathbb{E}[|\langle \underline{W^{(2)} G_{ii}^{(2)}} \rangle|^2]. \end{aligned}$$

Now, we compute $\mathbb{E}[|\langle \underline{W^{(2)} G_{ii}^{(2)}} \rangle|^2]$ by using a cumulant expansion 6.2. The following cumulant expansion is similar to (6.14), and the calculations thereafter.

$$\begin{aligned} & \mathbb{E}[|\langle \underline{W^{(2)} G_{ii}^{(2)}} \rangle|^2] \\ &= \frac{1}{n} \sum'_{ab} \mathbb{E}[\langle \Delta^{ba} G_{ii}^{(2)} \rangle \partial_{ab} (\langle \underline{W^{(2)} G_{ii}^{(2)}} \rangle)] + \frac{\gamma_1}{n} \sum'_{ab} \mathbb{E}[\partial_{ba} (\langle \Delta^{ba} G_{ii}^{(2)} \rangle) \langle \underline{W^{(2)} G_{ii}^{(2)}} \rangle] \\ & \quad + \frac{\gamma_1}{n} \sum'_{ab} \mathbb{E}[\langle \Delta^{ba} G_{ii}^{(2)} \rangle \partial_{ba} (\langle \underline{W^{(2)} G_{ii}^{(2)}} \rangle)] \\ & \quad + \sum'_{ab} \sum_{k \geq 2} \sum_{\alpha \in \{(ab)_2, (ba)_2\}^k} \frac{\kappa((ba)_2, \alpha)}{k!} \mathbb{E}[\partial_\alpha (\langle \Delta^{ba} G_{ii}^{(2)} \rangle) \langle \underline{W^{(2)} G_{ii}^{(2)}} \rangle] \\ (6.66) \quad &= \mathbb{E} \left[\widetilde{\mathbb{E}} \left[\langle \widetilde{W}^{(2)} G_{ii}^{(2)} \rangle (\langle \widetilde{W}^{(2)} G_{ii}^{(2)} \rangle + \langle \underline{W^{(2)} G_i^{(2)} \widetilde{W}^{(2)} G_{ii}^{(2)}} \rangle + \langle \underline{W^{(2)} G_{ii}^{(2)} \widetilde{W}^{(2)} G_i^{(2)}} \rangle) \right] \right] \\ & \quad + \frac{\gamma_1}{n} \sum'_{ab} \mathbb{E}[\langle \Delta^{ba} \partial_{ba} G_{ii}^{(2)} \rangle \langle \underline{W^{(2)} G_{ii}^{(2)}} \rangle] + \frac{\gamma_1}{n} \sum'_{ab} \mathbb{E}[\langle \Delta^{ba} G_{ii}^{(2)} \rangle \partial_{ba} (\langle \underline{W^{(2)} G_{ii}^{(2)}} \rangle)] \\ & \quad + \sum_{k \geq 2} \sum'_{ab} \sum_{k_1+k_2=k-1} \sum_{\alpha_1, \alpha_2} \frac{\kappa((ba)_2, \alpha_1, \alpha_2)}{k!} \mathbb{E}[\langle \Delta^{ba} \partial_{\alpha_1} G_{ii}^{(2)} \rangle \langle \Delta^{ba} \partial_{\alpha_2} G_{ii}^{(2)} \rangle] \\ & \quad + \sum_{k \geq 2} \sum'_{ab} \sum_{k_1+k_2=k} \sum_{\alpha_1, \alpha_2} \frac{\kappa((ba)_2, \alpha_1, \alpha_2)}{k!} \mathbb{E}[\langle \Delta^{ba} \partial_{\alpha_1} G_{ii}^{(2)} \rangle \langle \underline{W^{(2)} \partial_{\alpha_2} G_{ii}^{(2)}} \rangle], \end{aligned}$$

where $\alpha_i \in \{(ba)_2, (ba)_2\}^{k_i}$.

First, we bound the terms that do not contain self-renormalization terms, as we do not need to expand them further. So, first consider the term

$$(6.67) \quad \widetilde{\mathbb{E}} \left[\langle \widetilde{W}^{(2)} G_{ii}^{(2)} \rangle \langle \widetilde{W}^{(2)} G_{ii}^{(2)} \rangle \right] = \frac{1}{2n^2} \langle G_{ii}^{(2)} E_1 G_{ii}^{(2)} E_2 + G_{ii}^{(2)} E_2 G_{ii}^{(2)} E_1 \rangle$$

$$= \frac{\langle G_{iii}^{(2)} \rangle}{2n^2} = \mathcal{O}_{\prec} \left(\frac{1}{n^2 \eta_i^3 |\beta_1^{(1)}|^2 |\beta_i^{(2)}|^2} \right),$$

where we use the Lemma 6.5 in the last step.

Now, we consider the next non self-renormalization term

$$(6.68) \quad \sum_{k \geq 2} \frac{\kappa((ba)_2, \alpha_1, \alpha_2)}{k!} \sum'_{ab} \sum_{k_1+k_2=k-1} \sum_{\alpha_1, \alpha_2} \mathbb{E}[\langle \Delta^{ba} \partial_{\alpha_1} G_{ii}^{(2)} \rangle \langle \Delta^{ba} \partial_{\alpha_2} G_{ii}^{(2)} \rangle]$$

$$= \mathcal{O}_{\prec} \left(\frac{1}{n^{3/2} \eta_i^2 |\beta_1^{(1)}|^2 |\beta_i^{(2)}|^2} \right),$$

where we use the estimate $|\kappa((ba)_2, \alpha_1, \alpha_2)|/k! \lesssim n^{-(k+1)/2}$ of the cumulants from (6.5). Note that from Lemma 6.5, we have the bound $|(G_{ii}^{(2)})_{ab}| \prec 1/\eta_i$. After applying the α derivatives on these terms and applying Lemma 6.5, we get back the same bound. This justifies the last inequality in equation (6.68).

So far, we have examined the non self renormalized terms in the (6.66). To analyze the self renormalized terms, which are listed at the end of the first line, in the second line, and in the last line of (6.66), we do a further cumulant expansion as follows;

$$(6.69) \quad \mathbb{E}[\langle \Delta^{ba} \partial_{\alpha_1} G_{ii}^{(2)} \rangle \langle \widetilde{W}^{(2)} \partial_{\alpha_2} G_{ii}^{(2)} \rangle]$$

$$= \mathbb{E}[\widetilde{\mathbb{E}}[\langle \Delta^{ba} \partial_{\alpha_1} (G_i^{(2)} \widetilde{W}^{(2)} G_{ii}^{(2)} + G_{ii}^{(2)} \widetilde{W}^{(2)} G_i^{(2)}) \rangle \langle \widetilde{W}^{(2)} \partial_{\alpha_2} G_{ii}^{(2)} \rangle]]$$

$$+ \frac{\gamma_1}{n} \sum'_{cd} \mathbb{E}[\partial_{dc} (\langle \Delta^{dc} \partial_{\alpha_2} G_{ii}^{(2)} \rangle \langle \Delta^{ba} \partial_{\alpha_1} G_{ii}^{(2)} \rangle)]$$

$$+ \sum_{l \geq 2} \sum'_{cd} \sum_{l_1+l_2=l} \sum_{\vartheta_1, \vartheta_2} \mathbb{E}[\langle \Delta^{ba} \partial_{\alpha_1} \partial_{\vartheta_1} G_{ii}^{(2)} \rangle \langle \Delta^{cd} \partial_{\alpha_2} \partial_{\vartheta_2} G_{ii}^{(2)} \rangle]$$

$$= \frac{1}{n^2} \mathbb{E}[\langle \partial_{\alpha_1} (G_{ii}^{(2)} \Delta^{ba} G_i^{(2)} + G_i^{(2)} \Delta^{ba} G_{ii}^{(2)}) \partial_{\alpha_2} (G_{ii}^{(2)}) \rangle] + \frac{\gamma_1}{n} \sum'_{cd} \mathbb{E}[\partial_{dc} (\langle \Delta^{dc}$$

$$\partial_{\alpha_2} G_{ii}^{(2)} \rangle \langle \Delta^{ba} \partial_{\alpha_1} G_{ii}^{(2)} \rangle) + \frac{\gamma_1}{n} \sum'_{cd} \mathbb{E}[\langle \Delta^{dc} \partial_{\alpha_2} G_{ii}^{(2)} \rangle \partial_{dc} (\langle \Delta^{ba} \partial_{\alpha_1} G_{ii}^{(2)} \rangle)]$$

$$+ \sum_{l \geq 2} \sum'_{cd} \sum_{l_1+l_2=l} \sum_{\vartheta_1, \vartheta_2} \frac{\kappa(dc, \vartheta_1, \vartheta_2)}{l!} \mathbb{E}[\langle \Delta^{ba} \partial_{\alpha_1} \partial_{\vartheta_1} G_{ii}^{(2)} \rangle \langle \Delta^{cd} \partial_{\alpha_2} \partial_{\vartheta_2} G_{ii}^{(2)} \rangle],$$

where $\vartheta_i \in \{(cd)_2, (dc)_2\}^{l_i}$. After inserting the first line of (6.69) back into (6.66), we obtain an overall factor of $n^{-3-(k+1)/2}$ as well as the \sum'_{ab} summation over some $\partial_{\alpha}(\mathcal{G})_{ab}$, where \mathcal{G} is a product of five $G_i^{(2)}$'s. The sum \sum'_{ab} has $\mathcal{O}(n^2)$ many terms. Therefore, using the bound $|\partial_{\alpha}(\mathcal{G})_{ab}| \prec \eta_i^{-4}$, we can estimate the sum by $n^2 n^{-9/2} \eta_i^{-4} = n^{-5/2} \eta_i^{-4}$ since $k \geq 2$.

Now, we turn to the last line of (6.69). When this is inserted back into (6.66), we obtain a total prefactor of $n^{-(k+l)/2-3}$, and a summation $\sum'_{ab} \sum'_{cd}$ over

$$(\partial_{\alpha_1} \partial_{\vartheta_1} G_{ii}^{(2)})_{ab} (\partial_{\alpha_2} \partial_{\vartheta_2} G_{ii}^{(2)})_{cd}.$$

Using the fact that $\partial_{ab} G^{(t)} = -G^{(t)} \Delta^{ab} G^{(t)}$, we observe that a single derivative of $G^{(t)}$ yields one Δ . Therefore, the term $\Delta^{ba} \partial_{\alpha_1} \partial_{\vartheta_1} G_{ii}^{(2)}$ will have $|\alpha_1| + |\vartheta_1| + 1$ many Δ s. Similarly,

$\Delta^{cd}\partial_{\alpha_2}\partial_{\vartheta_2}G_{ii}^{(2)}$ will have $|\alpha_2| + |\vartheta_2| + 1$ many Δ s. The indices of Δ s, such as Δ^{ab}, Δ^{cd} etc, determine the number of diagonal and off diagonal factors of G in the product. When $k = l = 2$, for $|\alpha_1| + |\vartheta_1| = 3$ or $|\alpha_1| + |\vartheta_1| = 1$, all the entries are diagonal, which gives a bound of $n^{-1}\eta_i^{-2}$. In all other cases, at least two factors are off-diagonal, so the bound is $n^{-1}\eta_i^{-2}$ multiplied by a Ward improvement factor of $(n\eta_i)^{-1}$, which is $\mathcal{O}(n^{-2}\eta_i^{-3})$. Now, when $k + l \geq 5$, the bound are of smaller order.

Therefore, finally we obtain

$$(6.70) \quad \frac{1}{n^{(k+1)/2}} \sum_{k_1+k_2=k, k \geq 2} \sum_{ab} \sum_{\alpha_1, \alpha_2} \mathbb{E}[\langle \Delta^{ba} \partial_{\alpha_1} G_{ii}^{(2)} \rangle \langle \underline{W^{(2)}} \partial_{\alpha_2} G_{ii}^{(2)} \rangle] \\ = \mathcal{O}_{\prec} \left(\frac{1}{n^2 \eta_i^4 |\beta_1^{(1)}|^2 |\beta_i^{(2)}|^2} \right).$$

Finally, we consider the remaining terms in the first line of (6.66). After applying the cumulant expansion, we obtain the estimate

$$(6.71) \quad \frac{1}{n^2} \widetilde{\mathbb{E}}[\langle (G_{ii}^{(2)} \widetilde{W}^{(2)} G_i^{(2)} + G_i^{(2)} \widetilde{W}^{(2)} G_{ii}^{(2)})^2 \rangle] = \mathcal{O}_{\prec} \left(\frac{1}{n^2 \eta_i^4 |\beta_1^{(1)}|^2 |\beta_i^{(2)}|^2} \right).$$

The above bound is obtained by piecing the following bounds together;

$$|\langle G_{ii}^{(2)} G_i^{(2)} \rangle| \prec \frac{1}{\eta_i^2 |\beta_1^{(1)}| |\beta_i^{(2)}|},$$

which are from Lemma 6.5.

Similarly, we can estimate the remaining terms. For example, the second line of (6.69) can be estimated in the same way as we estimated the third line of (6.69), with $|\vartheta_1| \leq 1$ and $|\vartheta_2| = 0$ for the first term, and $|\vartheta_2| \leq 1$ and $|\vartheta_1| = 0$ for the second term. Likewise, the second line of (6.66) can be estimated as the last line of (6.66) in the cases $|\alpha_1| = 1, |\alpha_2| = 0$ or $|\alpha_1| = 0, |\alpha_2| = 1$. However, the estimates obtained from these will be less than or equal to the previously obtained estimates. Thus, by combining (6.67)-(6.71) we conclude the proof. \square

Similarly we can prove the following estimate.

Corollary 6.7. [15] *Let $G_i^{(t)}, G_{i1i}^{(t)}, A_i^{(t)}$ be same as defined in Lemma 6.5. Then we have*

$$\mathbb{E}|\langle \underline{G_1^{(t)} A_1^{(t)} E G_i^{(t)} A_i^{(t)} W^{(t)} G_i^{(t)} E^\dagger} \rangle|^2 = \mathbb{E}|\langle \underline{G_{i1i}^{(t)}} \rangle|^2 \\ \lesssim \left(\frac{1}{n \eta_1 \eta_i \eta_*^{1i} |\beta_1^{(t)}| |\beta_i^{(t)}|} \right)^2,$$

where $\eta_*^{1i} = \min\{\eta_1, \eta_i\}$.

7. INDEPENDENCE OF RESOLVENTS

Recall the definition of $I_{\eta_0}^{\eta_c}$ from (3.12), where $\eta_0 = n^{-1-\delta_0}$ and $\eta_c = n^{-1+\delta_1}$ for some small enough $\delta_0, \delta_1 > 0$. The contribution of $I_{\eta_0}^{\eta_c}$ to $\mathcal{L}_n(f)$ is negligible by the virtue of the fact that $(G^{(t)})^{z_1}(\iota\eta_1)$ and $(G^{(t)})^{z_2}(\iota\eta_2)$ are asymptotically independent if z_i, η_i are in appropriate regime. The exact technical conditions on z_i, η_i and the results are listed below.

Proposition 7.1. [14, Proposition 3.5] Fix $p \in \mathbb{N}$, and let $X^{(1)}$ is complex valued matrix. For any sufficiently small $\omega_d, \omega_h, \omega_f > 0$ such that $\omega_h \ll \omega_f$, there exist $\omega, \hat{\omega}, \delta_0, \delta_1 > 0$ such that $\omega_h \ll \delta_m \ll \hat{\omega} \ll \omega \ll \omega_f$, for $m = 0, 1$, such that for any $|z_l| \leq 1 - n^{-\omega_h}$, $|z_l - z_m| \geq n^{-\omega_d}$, with $l, m \in [p], l \neq m$, it holds

$$(7.1) \quad \begin{aligned} & \mathbb{E} \left[\prod_{l=1}^p \langle (G^{(1)})^{z_l}(i\eta_l) \rangle \right] \\ &= \prod_{l=1}^p \mathbb{E} [\langle (G^{(1)})^{z_l}(i\eta_l) \rangle] + \mathcal{O}_{\prec} \left(\frac{n^{p(\omega_l + \delta_0) + \delta_1}}{n^\omega} + \frac{n^{\omega_l + 3\delta_0}}{\sqrt{n}} \right), \end{aligned}$$

for any $\eta_1, \dots, \eta_p \in [n^{-1-\delta_0}, n^{-1+\delta_1}]$.

We also have following Proposition 7.2 which is real analogue of the Proposition 7.1.

Proposition 7.2. [15, Proposition 3.4] Fix $p \in \mathbb{N}$, and let $X^{(1)}$ is real valued matrix. For any sufficiently small $\omega_h, \omega_d > 0$, there exist $\omega_*, \delta_0, \delta_1$ with $\omega_h \ll \delta_m \ll \omega_* \ll 1$, for $m = 0, 1$, such that for any choice of z_1, z_2, \dots, z_p with

$$|z_l| \leq 1 - n^{-\omega_h}, |z_l - z_m| \geq n^{-\omega_d}, |z_l - \bar{z}_m| \geq n^{-\omega_d}, |z_l - \bar{z}_l| \geq n^{-\omega_d},$$

with $l, m \in [p], l \neq m$, it follows that

$$(7.2) \quad \mathbb{E} \left[\prod_{l=1}^p \langle (G^{(1)})^{z_l}(i\eta_l) \rangle \right] = \prod_{l=1}^p \mathbb{E} [\langle (G^{(1)})^{z_l}(i\eta_l) \rangle] + \mathcal{O}_{\prec} \left(\frac{n^{p(\omega_l + \delta_0) + \delta_1}}{n^{\omega_*}} \right),$$

for any $\eta_1, \dots, \eta_p \in [n^{-1-\delta_0}, n^{-1+\delta_1}]$.

Combining the above two propositions, we state the proposition for $(c(G^{(1)})^{z_i}(\nu\eta_i) + d(G^{(2)})^{z_i}(\nu\eta_i))$ as follows.

Proposition 7.3. Fix $p \in \mathbb{N}$, and let $X^{(t)}$ be either complex-valued or real-valued, and $\eta_1, \dots, \eta_p \in [n^{-1-\delta_0}, n^{-1+\delta_1}]$. Then, the following holds.

- (1) **Complex-valued case:** For any sufficiently small $\omega_d, \omega_h, \omega_f > 0$ such that $\omega_h \ll \omega_f$, there exist $\omega, \hat{\omega}, \delta_0, \delta_1 > 0$ such that $\omega_h \ll \delta_m \ll \hat{\omega} \ll \omega \ll \omega_f$, for $m = 0, 1$, such that for any $|z_l| \leq 1 - n^{-\omega_h}$, $|z_l - z_m| \geq n^{-\omega_d}$, with $l, m \in [p], l \neq m$, it holds

$$(7.3) \quad \begin{aligned} & \mathbb{E} \left[\prod_{l=1}^p \langle c(G^{(1)})^{z_l}(\nu\eta_l) + d(G^{(2)})^{z_l}(\nu\eta_l) \rangle \right] \\ &= \prod_{l=1}^p \mathbb{E} [\langle c(G^{(1)})^{z_l}(\nu\eta_l) + d(G^{(2)})^{z_l}(\nu\eta_l) \rangle] + \mathcal{O}_{\prec} \left(2^p \left(\frac{n^{p(\omega_l + \delta_0) + \delta_1}}{n^\omega} + \frac{n^{\omega_l + 3\delta_0}}{\sqrt{n}} \right) \right). \end{aligned}$$

- (2) **Real-valued case:** For any sufficiently small $\omega_h, \omega_d > 0$, there exist $\omega_*, \delta_0, \delta_1$ with $\omega_h \ll \delta_m \ll \omega_* \ll 1$, for $m = 0, 1$, such that for any choice of z_1, z_2, \dots, z_p with

$$|z_l| \leq 1 - n^{-\omega_h}, |z_l - z_m| \geq n^{-\omega_d}, |z_l - \bar{z}_m| \geq n^{-\omega_d}, |z_l - \bar{z}_l| \geq n^{-\omega_d},$$

with $l, m \in [p], l \neq m$, it holds

$$(7.4) \quad \mathbb{E} \left[\prod_{l=1}^p \langle c(G^{(1)})^{z_l}(\nu\eta_l) + d(G^{(2)})^{z_l}(\nu\eta_l) \rangle \right]$$

$$= \prod_{l=1}^p \mathbb{E}[\langle c(G^{(1)})^{z_l}(i\eta_l) + d(G^{(2)})^{z_l}(i\eta_l) \rangle] + \mathcal{O}_{\prec} \left(2^p \left(\frac{n^{p(\omega_l + \delta_0) + \delta_1}}{n^{\omega_*}} \right) \right).$$

Proof. Consider

$$(7.5) \quad \begin{aligned} & \mathbb{E} \left[\prod_{l=1}^p \langle c(G^{(1)})^{z_l}(i\eta_l) + d(G^{(2)})^{z_l}(i\eta_l) \rangle \right] \\ &= \sum_{\substack{i_1, i_2, \dots, i_p \in \{1, 2\} \\ s = |\{l: i_l = 1\}|}} \mathbb{E} \left[\prod_{l=1}^p \langle c^s d^{p-s} (G^{i_l})^{z_l}(i\eta_l) \rangle \right]. \end{aligned}$$

On the RHS of equation (7.5), we have terms like

$$\mathbb{E} \left[\langle t_1(G^{(i_1)})^{z_1}(i\eta_1) \rangle \langle t_2(G^{(i_2)})^{z_2}(i\eta_2) \rangle \langle t_3(G^{(i_3)})^{z_3}(i\eta_3) \rangle \dots \langle t_p(G^{(i_p)})^{z_p}(i\eta_p) \rangle \right]$$

where $i_1, i_2, \dots, i_p \in \{1, 2\}$, and $t_1, t_2, \dots, t_p \in \{c, d\}$ with $t_l = c$ if $i_l = 1$ and $t_l = d$ if $i_l = 2$. So, we have the following cases.

Case 1. When $i_1 = i_2 = \dots = i_p$. Then, from Proposition 7.1, we have

$$\begin{aligned} \mathbb{E} \left[\prod_{l=1}^p \langle c(G^{(1)})^{z_l}(i\eta_l) \rangle \right] &= \prod_{l=1}^p \mathbb{E}[\langle c(G^{(1)})^{z_l}(i\eta_l) \rangle] \\ &+ \mathcal{O}_{\prec} \left(\frac{n^{p(\omega_l + \delta_0) + \delta_1}}{n^{\omega}} + \frac{n^{\omega_l + 3\delta_0}}{\sqrt{n}} \right), \end{aligned}$$

Case 2. When not all i_1, i_2, \dots, i_p are equal. Then, for fix i_1, i_2, \dots, i_p , from Proposition 7.1, we have

$$\begin{aligned} \mathbb{E} \left[\prod_{l=1}^p \langle t_l(G^{(i_l)})^{z_l}(i\eta_l) \rangle \right] &= \prod_{l=1}^p \mathbb{E}[\langle t_l(G^{(i_l)})^{z_l}(i\eta_l) \rangle] \\ &+ \mathcal{O}_{\prec} \left(\frac{n^{p(\omega_l + \delta_0) + \delta_1}}{n^{\omega}} + \frac{n^{\omega_l + 3\delta_0}}{\sqrt{n}} \right). \end{aligned}$$

Notice that in Case 2, the matrices are different, i.e., $G^{(1)}$ and $G^{(2)}$, so the correlation between them is smaller in comparison with the Case 1, which has the same matrix $G^{(1)}$.

Thus, from Case 1 and Case 2 we have

$$\begin{aligned} & \mathbb{E} \left[\prod_{l=1}^p \langle c(G^{(1)})^{z_l}(i\eta_l) + d(G^{(2)})^{z_l}(i\eta_l) \rangle \right] \\ &= \sum_{\substack{i_1, i_2, \dots, i_p \in \{1, 2\} \\ s = |\{l: i_l = 1\}|}} \mathbb{E} \left[\prod_{l=1}^p \langle c^s d^{p-s} (G^{i_l})^{z_l}(i\eta_l) \rangle \right] \\ &= \sum_{\substack{i_1, i_2, \dots, i_p \in \{1, 2\} \\ s = |\{l: i_l = 1\}|}} \left\{ \prod_{l=1}^p \mathbb{E}[\langle c^s d^{p-s} (G^{i_l})^{z_l}(i\eta_l) \rangle] + \mathcal{O}_{\prec} \left(\frac{n^{p(\omega_l + \delta_0) + \delta_1}}{n^{\omega}} + \frac{n^{\omega_l + 3\delta_0}}{\sqrt{n}} \right) \right\} \end{aligned}$$

$$\begin{aligned}
&= \sum_{\substack{i_1, i_2, \dots, i_p \in \{1, 2\} \\ s = |\{l: i_l = 1\}|}} \left\{ \prod_{l=1}^p \mathbb{E}[\langle c^s d^{p-s} (G^{i_l})^{z_l}(i\eta_l) \rangle] \right\} \\
&\quad + \mathcal{O}_{\prec} \left(2^p \left(\frac{n^{p(\omega_l + \delta_0) + \delta_1}}{n^\omega} + \frac{n^{\omega_l + 3\delta_0}}{\sqrt{n}} \right) \right) \\
&= \prod_{l=1}^p \mathbb{E}[\langle c(G^1)^{z_l}(i\eta_l) + d(G^2)^{z_l}(i\eta_l) \rangle] + \mathcal{O}_{\prec} \left(2^p \left(\frac{n^{p(\omega_l + \delta_0) + \delta_1}}{n^\omega} + \frac{n^{\omega_l + 3\delta_0}}{\sqrt{n}} \right) \right).
\end{aligned}$$

Similarly, we can prove the independence of product of resolvents for the real-valued case (7.4). \square

REFERENCES

- [1] A. Adhikari and Z. Che. Edge universality of correlated gaussians. *Electron. J. Probab.*, 24(44):1–25, 2019.
- [2] K. Adhikari and A. Bose. Brown measure and asymptotic freeness of elliptic and related matrices. *Random Matrices: Theory and Applications*, 8(02):1950007, 2019.
- [3] K. Adhikari, A. Bose, and S. N. Maurya. Convergence of high dimensional toeplitz and related matrices with correlated inputs. *Latin American Journal of Probability & Mathematical Statistics*, 21:1867–1901, 2024.
- [4] O. H. Ajanki, L. Erdős, and T. Krüger. Stability of the matrix dyson equation and random matrices with correlations. *Probability Theory and Related Fields*, 173:293–373, 2019.
- [5] J. Alt, L. Erdős, T. Krüger, and D. Schröder. Correlated random matrices: band rigidity and edge universality. *The Annals of Probability*, 48(2):963–1001, 2020.
- [6] J. Alt, L. Erdős, and T. Krüger. Spectral radius of random matrices with independent entries. *Probability and Mathematical Physics*, 2:221–280, 05 2021.
- [7] J. Alt and T. Krüger. Inhomogeneous circular law for correlated matrices. *Journal of Functional Analysis*, 281(7):109120, 2021.
- [8] Z. D. Bai. Circular law. *The Annals of Probability*, 25(1):494–529, 1997.
- [9] D. Banerjee, S. S. Mukherjee, and D. Pal. Edge spectra of gaussian random symmetric matrices with correlated entries. *arXiv preprint arXiv:2409.11381*, 2024.
- [10] M. Bhattacharjee, A. Bose, and A. Dey. Joint convergence of sample cross-covariance matrices. *Latin American Journal of Probability & Mathematical Statistics*, 20:395–423, 2023.
- [11] R. Catalano, M. Fleermann, and W. Kirsch. Random band and block matrices with correlated entries. *Electronic Journal of Probability*, 29:1–32, 2024.
- [12] Z. Che. Universality of random matrices with correlated entries. *Electron. J. Probab.*, 22(30):1–38, 2017.
- [13] G. Cipolloni, L. Erdős, and S. J. Henheik. Out-of-time-ordered correlators for wigner matrices. *Advances in Theoretical and Mathematical Physics*, 28(6), 2024.
- [14] G. Cipolloni, L. Erdős, and D. Schröder. Central limit theorem for linear eigenvalue statistics of non-hermitian random matrices. *Communications on Pure and Applied Mathematics*, 76(5):946–1034, 2023.
- [15] G. Cipolloni, L. Erdős, and D. Schröder. Fluctuation around the circular law for random matrices with real entries. *Electronic Journal of Probability*, 26(none):1–61, 2021.
- [16] G. Cipolloni, L. Erdős, and D. Schröder. Fluctuation around the circular law for random matrices with real entries. *Electronic Journal of Probability*, 26(none):1–61, 2021.
- [17] L. Erdős, T. Krüger, and D. Schröder. Random matrices with slow correlation decay. In *Forum of Mathematics, Sigma*, volume 7, page e8. Cambridge University Press, 2019.
- [18] V. L. Girko. Circular law. *Theory of Probability & Its Applications*, 29(4):694–706, 1985.
- [19] F. Götze and A. Tikhomirov. The circular law for random matrices. *The Annals of Probability*, 38(4):1444–1491, 2010.
- [20] I. Jana and S. Rani. Spectrum of random centrosymmetric matrices; clt and circular law. *Random Matrices: Theory and Applications*, 14(01):2450026, 2025.

- [21] G. Pan and W. Zhou. Circular law, extreme singular values and potential theory. *Journal of Multivariate Analysis*, 101(3):645–656, 2010.
- [22] B. Rider and J. W. Silverstein. Gaussian fluctuations for non-Hermitian random matrix ensembles. *The Annals of Probability*, 34(6):2118 – 2143, 2006.
- [23] R. Speicher. High-dimensional analysis: Random matrices and machine learning. *Lecture Notes, Department of Mathematics, Saarland University*, 2023.
- [24] T. Tao and V. Vu. Random matrices: the circular law. *Communications in Contemporary Mathematics*, 10(02):261–307, 2008.
- [25] T. Tao and V. Vu. Random matrices: Universality of local spectral statistics of non-Hermitian matrices. *The Annals of Probability*, 43(2):782 – 874, 2015.
- [26] T. Tao and V. H. Vu. Smooth analysis of the condition number and the least singular value. In *Mathematics of Computation*, 2008.
- [27] J. R. Weaver. Centrosymmetric (cross-symmetric) matrices, their basic properties, eigenvalues, and eigenvectors. *The American Mathematical Monthly*, 92(10):711–717, 1985.

INDIAN INSTITUTE OF TECHNOLOGY, BHUBANESWAR
Email address: ijana@iitbbs.ac.in

INDIAN INSTITUTE OF TECHNOLOGY, BHUBANESWAR
Email address: s21ma09007@iitbbs.ac.in