

A Unified Approach for Estimating Various Treatment Effects in Causal Inference

Kuan-Hsun Wu and Li-Pang Chen¹

Department of Statistics, National Chengchi University

Abstract

In this paper, we introduce a unified estimator to analyze various treatment effects in causal inference, including but not limited to the average treatment effect (ATE) and the quantile treatment effect (QTE). The proposed estimator is developed under the statistical functional and cumulative distribution function structure, which leads to a flexible and robust estimator and covers some frequent treatment effects. In addition, our approach also takes variable selection into account, so that informative and network structure in confounders can be identified and be implemented in our estimation procedure. The theoretical properties, including variable selection consistency and asymptotic normality of the statistical functional estimator, are established. Various treatment effects estimations are also conducted in numerical studies, and the results reveal that the proposed estimator generally outperforms the existing methods and is more efficient than its competitors.

Keywords: Causal inference; counterfactual distributions; cumulative distribution function; network structures; statistical functional; propensity score; variable selection.

Short title: Estimating Various Causal Effects Using IPW CDF

¹Corresponding Author. Email: lchen723@nccu.edu.tw

1 Introduction

Causal inference aims to explore the causality of two variables, and it attract people’s attention in various fields recently, such as social science or biological studies. One of the important questions in causal inference us the estimations of treatment effects, which reflect how different treatments affect the outcomes. In the literature, some typical causal effects include but are not limited to the average treatment effect (ATE, Chen, 2020; Yi and Chen, 2023) and the quantile treatment effect (QTE, Firpo 2007; Donald and Hsu, 2014; Hsu et al., 2022). To estimate treatment effects, the inverse probability weight (IPW) method (e.g., Rosenbaum and Rubin, 1983) is one of the popular approaches, whose key idea is to model the propensity score and treat it as the weight to recover the ”missingness” in the outcome. This approach has also been widely used to the estimation procedure, such as ATE (e.g., Chen, 2020; Yi and Chen, 2023), QTE (e.g., Firpo 2007; Donald and Hsu, 2014; Hsu et al., 2022) and treatment effect for survival data (e.g., Chapfuwa et al., 2021).

In applications, one may encounter a scenario that the dataset contains multivariate or high-dimensional confounders. Among those confounders, few of them are informative to the outcome, so it is crucial to do variable selection. There are some methods available to address this challenge. To name a few, Ertefaie et al. (2018) proposed the weighted LASSO penalization with simultaneous consideration on outcome and treatment models. Yi and Chen (2023) proposed the penalized likelihood function with error-prone confounders. Following the spirit of adaptive LASSO from Zou (2006), the outcome-adaptive LASSO proposed by Shortreed and Ertefaie (2017) focused on the estimation of propensity score with the logit model adopted. Moreover, Bayesian approaches were also employed to causal inference, including Koch et al. (2020) who implemented variable selection under the similar framework of estimating causal effect with spike and slab priors applied.

In addition to selecting informative covariates, the other challenging feature in multivariate or high-dimensional variables is the complex network structure. To offer an intelligible interpretation of the dependencies among high-dimensional confounders, *graphical model* is a powerful tool and has been adopted for numerous applications in regression models or supervised learning; see details in Chen (2024, Section 4). However, the impact of network

structure in confounders and the resulting estimation for treatment effects have not been fully explored.

To address the challenges in the multivariate or high-dimensional settings and unify various treatment effect estimation into a general estimator, we proposed the IPW method under the cumulative distribution function (CDF) and statistical functional form. The proposed estimator is robust in dealing with outlier and is reduced to some well known treatment effects if the statistical functional is properly specified. Moreover, we also implement variable selection technique to detect informative confounders and network structure when estimating the propensity score. Theoretically, we establish the variable selection consistency and asymptotic properties for the proposed estimator.

The remainder is organized as follows. In Section 2, we introduce the fundamentals of causal inference including necessary assumptions and estimands of interest, along with the model formulation of network structure. Subsequently, we introduce the proposed approach and the theoretical results in Section 3. To verify the effectiveness of our method, we conduct the simulation studies and real data analysis in Sections 4 and 5, respectively. Finally, the highlights of this article are discussed and summarized in Section 6.

2 Preliminaries and Model Setup

2.1 Causal Effects and Propensity Score

Let $\mathbf{X} = (X_1, \dots, X_p)^\top$ denote the p -dimensional vector of covariates, or confounders with little case letter \mathbf{x} being a realized value, and let $A \in \{0, 1\}$ denote the binary treatment that indicates treatment ($A = 1$) or control ($A = 0$). Given the treatment effect $A = a$, let $Y^{(a)}$ denote the potential outcome. Specifically, when $a = 1$, $Y^{(1)}$ stands for the potential outcome if a subject is assigned to the treatment; when $a = 0$, then $Y^{(0)}$ is the potential outcome if a subject is assigned to the control. Moreover, let $F^{(a)}(y)$ denote the CDF of $Y^{(a)}$ for $a = 0, 1$.

Let $T(F)$ denote the statistical functional, which is a function of the CDF $F(y)$. Hence,

we consider a general class of the treatment effect

$$T(F^{(1)}) - T(F^{(0)}). \quad (1)$$

Then (1) can be reduced to some commonly used treatment effects in causal inference. For example, when $T(F) = \int y dF(y)$, then (1) is reduced to the *average treatment effect* (ATE), which is given by

$$\begin{aligned} \tau &\triangleq \int_{\mathbb{R}} y dF^{(1)}(y) - \int_{\mathbb{R}} y dF^{(0)}(y) \\ &= E(Y^{(1)}) - E(Y^{(0)}). \end{aligned} \quad (2)$$

When $T(F^{(1)})$ and $T(F^{(0)})$ are specified as the q th quantiles of $Y^{(1)}$ and $Y^{(0)}$, i.e.,

$$\xi^{(1)}(q) \triangleq \inf \left\{ y \in \mathbb{R} \mid F^{(1)}(y) \geq q \right\} \text{ and } \xi^{(0)}(q) \triangleq \inf \left\{ y \in \mathbb{R} \mid F^{(0)}(y) \geq q \right\} \quad (3)$$

for $q \in (0, 1)$, respectively, the (1) becomes the *q th quantile treatment effect* (QTE):

$$\Xi(q) \triangleq \xi^{(1)}(q) - \xi^{(0)}(q). \quad (4)$$

If $T(F)$ is the probability functional, say $T(F) = \int_{-\infty}^y dF(x)$ for a fixed y , then (1) gives the *distributional treatment effect* (DTE), which is defined as

$$\Delta(y) \triangleq F^{(1)}(y) - F^{(0)}(y) \quad (5)$$

for a given $y \in \mathbb{R}$.

To estimate (2), (4) and (5), a crucial issue is the estimation of $F^{(a)}(y)$. In this study, we employ the inverse probability weight (IPW) method. Specifically, the propensity score (PS) is defined as the conditional probability of treatment, given confounders, which is given by

$$\pi(\mathbf{x}) \triangleq P(A = 1 \mid \mathbf{X} = \mathbf{x}). \quad (6)$$

To further discuss the estimation as well as the relevant development, we impose the following conditions that are also commonly assumed in the framework of causal inference (e.g. Yi and Chen 2023):

(C1) *Stable Unit Treatment Value Assumption (SUTVA)*:

Response of a subject is not affected by responses of other subjects (noninterference). The treatment A could be assigned by different ways, but they all lead to the same outcome (consistency). In this assumption, we have $Y = AY^{(1)} + (1 - A)Y^{(0)}$.

(C2) *Strong Ignorable Treatment Assumption (SITA)*:

The treatment assignment A is independent of potential outcomes $(Y^{(1)}, Y^{(0)})$, given the covariates \mathbf{X} .

(C3) $0 < \pi(\mathbf{x}) < 1$.

(C4) (A, \mathbf{X}) is independent and identically distributed.

(C5) \mathbf{X} is bounded.

Conditions (C1)-(C5) allow us to characterize the distribution function $F^{(a)}$ by the variables Y, A and \mathbf{X} for $a \in \{0, 1\}$. Specifically, let $I(\cdot)$ denote the indicator function. Then for

$a = 1$, we have that

$$\begin{aligned}
E\left\{\frac{AI(Y \leq y)}{\pi(\mathbf{X})}\right\} &= E\left[\frac{1}{\pi(\mathbf{X})}E\left\{AI(Y \leq y) \mid \mathbf{X}\right\}\right] \\
&= E\left[\frac{E\left\{1 \times I(Y \leq y) \mid \mathbf{X}, A = 1\right\}P(A = 1 \mid \mathbf{X})}{\pi(\mathbf{X})}\right] \\
&\quad + E\left[\frac{E\left\{0 \times I(Y \leq y) \mid \mathbf{X}, A = 0\right\}P(A = 0 \mid \mathbf{X})}{\pi(\mathbf{X})}\right] \\
&= E\left[\frac{E\left\{I(AY^{(1)} + (1 - A)Y^{(0)} \leq y) \mid \mathbf{X}, A = 1\right\}\pi(\mathbf{X})}{\pi(\mathbf{X})}\right] \tag{7} \\
&= E\left[E\left\{I(Y^{(1)} \leq y) \mid \mathbf{X}, A = 1\right\}\right] \\
&= E\left[E\left\{I(Y^{(1)} \leq y) \mid \mathbf{X}\right\}\right] \\
&= E\left\{I(Y^{(1)} \leq y)\right\} \\
&= F^{(1)}(y),
\end{aligned}$$

where the first equality is established from the law of iterated expectations, the second equality comes from partitioning the conditional expectation $E\left\{AI(Y \leq y) \mid \mathbf{X}\right\}$ with the event $A = 1$ and $A = 0$, the third equality is a direct result of (C2) and $P(A = 1 \mid \mathbf{X}) = \pi(\mathbf{X})$, the fourth equality is obtained with basic operation, the fifth step utilizes (C2) again, and the sixth equality is derived from the law of iterated expectations with the inverse direction. The identification procedure of $F^{(0)}(y)$ can be obtained by (7) with A and $\pi(\mathbf{X})$ replaced by $1 - A$ and $1 - \pi(\mathbf{X})$, respectively. Consequently, under the independent and identically distributed (IID) samples with size n , denoted as $\mathcal{O} \triangleq \{\{Y_i, A_i, \mathbf{X}_i\} : i = 1, \dots, n\}$, with $\pi(\mathbf{X})$ being estimated consistently, the estimators of $F^{(1)}(y)$ and $F^{(0)}(y)$ are respectively given by

$$\widehat{F}^{(1)}(y) = \left(\sum_{i=1}^n \frac{A_i}{\widehat{\pi}(\mathbf{X}_i)}\right)^{-1} \sum_{i=1}^n \frac{A_i I(Y_i \leq y)}{\widehat{\pi}(\mathbf{X}_i)} \tag{8}$$

and

$$\widehat{F}^{(0)}(y) = \left(\sum_{i=1}^n \frac{1 - A_i}{1 - \widehat{\pi}(\mathbf{X}_i)}\right)^{-1} \sum_{i=1}^n \frac{(1 - A_i)I(Y_i \leq y)}{\widehat{\pi}(1 - \mathbf{X}_i)}. \tag{9}$$

The first notable remark for (8) and (9) is that we implement $\left(\sum_{i=1}^n \frac{A_i}{\hat{\pi}(\mathbf{X}_i)}\right)$ and $\left(\sum_{i=1}^n \frac{1 - A_i}{1 - \hat{\pi}(\mathbf{X}_i)}\right)$ to replace n when estimating the expectation (7) empirically because (8) and (9) form the estimated CDF; detailed justification can be found in Appendix C of the supporting information. While Donald and Hsu (2014) also adopted the similar rationales in (8) and (9), the difference is that our approach is for the estimation of CDF.

The other notable remark is that we focus on a scenario that $Y^{(a)}$ is continuous for $a \in \{0, 1\}$ in the current study since our development is based on CDF. When $Y^{(a)}$ is discrete, the estimation problem naturally reduces to finding the location of largest jump in counterfactual CDFs.

2.2 Network Structure in Covariates

According to the structure in Figure 1, the covariates \mathbf{X} are possibly formulated by the network structure. To characterize the dependence structure for \mathbf{X} , we use an undirected *graph*, denoted $\tilde{G} \triangleq (\tilde{V}, \tilde{E})$, where $\tilde{V} = \{1, \dots, p\}$ records the indexes of all components of \mathbf{X} , and $\tilde{E} \subset \tilde{V} \times \tilde{V}$ contains dependent pairs of covariates. Following Chen and Yi (2021), we consider the exponential family graphical model that is formulated as follows:

$$P(\mathbf{X}; \boldsymbol{\beta}, \boldsymbol{\Theta}) = \exp \left\{ \sum_{r \in \tilde{V}} \beta_r B(X_r) + \sum_{(s, \nu) \in \tilde{E}} \theta_{s\nu} B(X_s) B(X_\nu) + \sum_{r \in \tilde{V}} C(X_r) - \varphi(\boldsymbol{\beta}, \boldsymbol{\Theta}) \right\}, \quad (10)$$

where $\boldsymbol{\beta} = (\beta_1, \dots, \beta_p)^\top$ is a p -dimensional parameter vector, $\boldsymbol{\Theta} = [\theta_{s\nu}]$ is a $p \times p$ symmetric matrix, and $B(\cdot)$ and $C(\cdot)$ are given functions. The function $\varphi(\boldsymbol{\beta}, \boldsymbol{\Theta})$ is the normalizing constant that makes (10) be integrated as 1. For $r \in \tilde{V}$, the parameter β_r reflects the main effect associated with the covariate X_r ; for $(s, \nu) \in \tilde{E}$, the parameter $\theta_{s\nu}$ facilitates the association of X_s and X_ν in the sense that $\theta_{s\nu} \neq 0$ shows the *conditional dependence* of X_s and X_ν given other covariates.

3 Methodology

3.1 Variable Selection and Network Construction for Outcome Responses

Since the dataset contains multivariate confounders and some of them are not necessarily informative to estimate $\pi(\cdot)$ in (8) and (9) and the treatment effects in (1), it is crucial to determine the important variables and detect the network structures that are dependent on the outcome response Y (e.g., Shortreed and Ertefaie 2017). Let $V = \{r : \beta_r \neq 0, r = 1, \dots, p\}$ denote the indices set containing informative covariates, and let $E = \{(s, \nu) : \theta_{s\nu} \neq 0\}$ be the set containing the pairwise dependence, so that $G \triangleq (V, E)$. For the convenience of the presentation, we define \mathbb{X} as the $n \times p$ matrix of confounders with \mathbb{X}_j being the j th column vector in \mathbb{X} for $j = 1, \dots, p$, and let $\mathbb{X}^{\circ 2} \triangleq [\mathbb{X}_1 * \mathbb{X}_2 \ \mathbb{X}_1 * \mathbb{X}_3 \ \dots \ \mathbb{X}_{p-1} * \mathbb{X}_p]$, where $*$ is the entry-wise product of two column vectors. In addition, define two vectors $\mathbf{Y} = (Y_1 \ Y_2 \ \dots \ Y_n)^\top$ and $\boldsymbol{\varepsilon} = (\varepsilon_1 \ \varepsilon_2 \ \dots \ \varepsilon_n)^\top$ with $\varepsilon_i \sim N(0, \sigma^2)$, and denote $\text{vec}(\boldsymbol{\Theta})$ as the column vectorization of $\boldsymbol{\Theta}$. Then the matrix form can be expressed as

$$\mathbf{Y} = \mathbb{X}\boldsymbol{\beta} + \mathbb{X}^{\circ 2}\text{vec}(\boldsymbol{\Theta}) + \boldsymbol{\varepsilon}. \quad (11)$$

To select informative confounders and detect the corresponding network structure, we adopt the penalized likelihood method, and the estimators of $\boldsymbol{\beta}$ and $\boldsymbol{\Theta}$ are determined by

$$(\hat{\boldsymbol{\beta}}, \hat{\boldsymbol{\Theta}}) = \underset{\boldsymbol{\beta}, \boldsymbol{\Theta}}{\text{argmin}} \left\{ \|\mathbf{Y} - \mathbb{X}\boldsymbol{\beta} - \mathbb{X}^{\circ 2}\text{vec}(\boldsymbol{\Theta})\|_2^2 + \lambda_1 \|\boldsymbol{\beta}\|_1 + \lambda_2 \|\text{vec}(\boldsymbol{\Theta})\|_1 \right\}, \quad (12)$$

where $\|\cdot\|_2$ is the L_2 -norm and λ_1 and λ_2 are the tuning parameters associated with of the penalty terms $\|\cdot\|_1$ in the L_1 -norm. Moreover, when $\hat{\boldsymbol{\beta}}$ and $\hat{\boldsymbol{\Theta}}$ are obtained, we define $\hat{V} = \{r : \hat{\beta}_r \neq 0\}$ and $\hat{E} = \{(s, \nu) : \hat{\theta}_{s\nu} \neq 0\}$ as the corresponding estimated sets of V and E , respectively, which reflect informative main effects and pairwise interaction with respect to \mathbf{Y} . Accordingly, the estimated undirected graph is defined as $\hat{G} = (\hat{V}, \hat{E})$. Since $\|\mathbf{Y} - \mathbb{X}\boldsymbol{\beta} - \mathbb{X}^{\circ 2}\text{vec}(\boldsymbol{\Theta})\|_2^2$ and two penalty terms $\|\boldsymbol{\beta}\|_1$ and $\|\text{vec}(\boldsymbol{\Theta})\|_1$ are convex, then the objective function in (12) is convex as well, ensuring the existence of the minimizer. To compute (12), we adopt the coordinate-descent method to iteratively calculate the numerical values (e.g., Chen and Yi 2021).

3.2 Estimation of Propensity Scores

In this section, we estimate the propensity score with selected confounders and the network structure taken into account.

To model the propensity score $\pi(\mathbf{x})$ in (6), we consider the following *network-based logistic regression model*:

$$\text{logit} \{ \pi(\mathbf{x}) \} = \eta_0 + \sum_{r \in \widehat{V}} \eta_r x_r + \sum_{(s, \nu) \in \widehat{E}} \eta_{s\nu} x_s x_\nu, \quad (13)$$

where η_0 is an intercept, $\sum_{r \in \widehat{V}} \eta_r x_r$ reflects the informative confounders that are highly correlated with outcomes, and $\sum_{(s, \nu) \in \widehat{E}} \eta_{s\nu} x_s x_\nu$ reflects the dependence structures among confounders. Let $\boldsymbol{\eta} \triangleq (\eta_0, \boldsymbol{\eta}_V^\top, \boldsymbol{\eta}_E^\top)^\top$, where $\boldsymbol{\eta}_V \triangleq (\eta_r : r \in \widehat{V})^\top$ and $\boldsymbol{\eta}_E \triangleq (\eta_{s\nu} : (s, \nu) \in \widehat{E})^\top$ are two vectors of parameters. Based on the sample \mathcal{O} , we apply the maximum likelihood method based on logistic regression model to estimate $\boldsymbol{\eta}$, and the estimator is given by

$$\widehat{\boldsymbol{\eta}} = \underset{\boldsymbol{\eta}}{\text{argmax}} \left[\prod_{i=1}^n \{ \pi(\mathbf{x}_i) \}^{a_i} \{ 1 - \pi(\mathbf{x}_i) \}^{1-a_i} \right]. \quad (14)$$

Therefore, the proposed estimator of the propensity score is given by

$$\widehat{\pi}(\mathbf{x}) = \frac{\exp \left(\widehat{\eta}_0 + \sum_{j \in \widehat{V}} \widehat{\eta}_j x_j + \sum_{(s, \nu) \in \widehat{E}} \widehat{\eta}_{s\nu} x_s x_\nu \right)}{1 + \exp \left(\widehat{\eta}_0 + \sum_{r \in \widehat{V}} \widehat{\eta}_r x_r + \sum_{(s, \nu) \in \widehat{E}} \widehat{\eta}_{s\nu} x_s x_\nu \right)}. \quad (15)$$

(15) is an extended formulation from the conventional logistic regression models by incorporating network structures. This expression is also similar to Chen et al. (2019), which is numerically justified that incorporating network structures would improve the accuracy of the classification.

3.3 Estimation of IPW-CDFs

When propensity scores $\pi(\cdot)$ are estimated by (15), $F^{(a)}$ is estimated by (8) and (9) for $a \in \{0, 1\}$. Consequently, the estimator of (1) is given by

$$T(\widehat{F}^{(1)}) - T(\widehat{F}^{(0)}), \quad (16)$$

which reduces to the estimators of various treatment effects in Section 2.1. For example, the ATE (2) is estimated by

$$\hat{\tau} = \int_{\mathbb{R}} y d\hat{F}^{(1)}(y) - \int_{\mathbb{R}} y d\hat{F}^{(0)}(y). \quad (17)$$

The estimators of the q th quantiles in (3) are given by

$$\hat{\xi}^{(1)}(q) = \inf \left\{ y \in \mathbb{R} \mid \hat{F}^{(1)}(y) \geq q \right\} \text{ and } \hat{\xi}^{(0)}(q) = \inf \left\{ y \in \mathbb{R} \mid \hat{F}^{(0)}(y) \geq q \right\}, \quad (18)$$

yielding the estimator of (4)

$$\hat{\Xi}(q) = \hat{\xi}^{(1)}(q) - \hat{\xi}^{(0)}(q) \quad (19)$$

for some $q \in (0, 1)$. Finally, the estimator of DTE (5) is given by the difference of (8) and (9), i.e,

$$\hat{\Delta}(y) = \hat{F}^{(1)}(y) - \hat{F}^{(0)}(y) \quad (20)$$

for a given $y \in \mathbb{R}$.

Here we comment the proposed estimator (16). The first advantage of (16) is robustness. Specifically, as pointed out by Khan and Ugander (2023), $\left(\sum_{i=1}^n \frac{A_i}{\hat{\pi}(\mathbf{X}_i)} \right)$ and $\left(\sum_{i=1}^n \frac{1 - A_i}{1 - \hat{\pi}(\mathbf{X}_i)} \right)$ in the denominator of the estimators (8) and (9) are positively correlated with the numerators, the effect of the extreme values can be mitigated by such correlation. In addition, (16) is formulated by the CDF that lie in an interval $[0, 1]$, which can also avoid potential impact of outliers in the outcome Y . The second advantage of the proposed estimator (16) is to provide a general formulation that unifies a class of various causal effects and incorporates variable selection to avoid unnecessary bias induced by redundant covariates and pairwise dependence structure. When variable selection is not taken into account, (17) reduces to the estimator described in Lunceford and Davidan (2004). Furthermore, if the treatment model is assigned as series logit model (Hirano, Imbens and Ridder, 2003) with variable selection ignored, (19) reduces to the estimator proposed by Donald and Hsu (2014). Finally, the estimator (20) can be referred to the generalization of caused effects of survivor functions.

3.4 Theoretical Results

To establish the asymptotic properties, we first present the consistency of variable selection in the following theorem, which ensures that the truly informative confounders and the network structure can be identified.

Theorem 3.1 *Under regularity conditions stated in the Supporting Information, the following property holds:*

$$P(\widehat{G} = G) \rightarrow 1 \text{ as } n \rightarrow \infty.$$

Theorem 3.1 states that there is a high probability that our estimation procedure can successfully capture the underlying true confounders and dependence structure among covariates for the potential outcome. Consequently, our estimators are free of the potential bias incurred by non-informative confounders.

Before introducing the asymptotic properties for the proposed estimators, we first define some notations. We define the density functions of $Y^{(1)}$ and $Y^{(0)}$ as $f^{(1)}$ and $f^{(0)}$, respectively. With Theorem 3.1, we are capable of using the selected covariates. Let $\mathbf{X}_{\widehat{V}}$ and $\mathbf{X}_{\widehat{E}}$ be two matrices with dimensions $n \times k$ and $n \times m$ consisting of the chosen main and interaction effects, respectively. Consequently, we define $\mathbf{X}^* = \begin{bmatrix} \mathbf{1} & \mathbf{X}_{\widehat{V}} & \mathbf{X}_{\widehat{E}} \end{bmatrix}$ where $\mathbf{1}$ is the column vector with all components being 1.

To derive the theoretical properties, we further impose some assumptions to the statistical functional. Suppose that $T : \mathcal{F} \rightarrow \mathbb{R}$ satisfies Hadamard differentiability at $F \in \mathcal{F}$, where \mathcal{F} is a normed space consisting of probability distribution functions, i.e., there exists a continuous linear function T'_F , such that, as $n \rightarrow \infty$,

$$\frac{T(F + t_n h_n) - T(F)}{t_n} \rightarrow T'_F(h)$$

for all sequences $\{t_n\} \subset \mathbb{R}$ and $\{h_n\} \subset \mathcal{F}$ satisfying $t_n \rightarrow 0$, $h_n \rightarrow h \in \mathcal{F}$ and $F + t_n h_n \in \mathcal{F}$. In addition, when the statistical functional T is Hadamard differentiable, the Gateaux derivative, which is also called the von Mises derivative, of T exists and the values of these two types of derivative are the same (van der Vaart, 2000). Furthermore, the influence curve, or the influence function, of a statistical functional T is defined as

$$\phi_F(y) \triangleq \lim_{t \rightarrow 0} \frac{T((1-t)F + t\delta_y) - T(F)}{t}, \quad (21)$$

where δ_y is the distribution function with point mass 1 on y (Fernholz, 1983). In particular, if a statistical functional can be written as the form $\int \varphi(y)dF(y)$ for some function $\varphi(y)$, it is called a linear functional and its influence curve is $\phi_F(x) = \varphi(x) - T(F)$ (Wasserman, 2006).

We now present the asymptotic properties of the proposed estimator.

Theorem 3.2 *Under regularity conditions stated in the Supporting Information, the following properties hold as $n \rightarrow \infty$:*

$$(i) \sqrt{n}(\hat{\boldsymbol{\eta}} - \boldsymbol{\eta}) \xrightarrow{d} N(\mathbf{0}, \mathcal{A}^{-1});$$

$$(ii) \sqrt{n} \left[\left(T(\hat{F}^{(1)}) - T(\hat{F}^{(0)}) \right) - \left(T(F^{(1)}) - T(F^{(0)}) \right) \right] \xrightarrow{d} N\left(0, -\mathcal{B}\mathcal{A}^{-1}\mathcal{B}^\top + \mathcal{C}\right),$$

where

$$\mathcal{A} = E \left\{ \pi(\mathbf{X}^* | \boldsymbol{\eta}) \left(1 - \pi(\mathbf{X}^* | \boldsymbol{\eta}) \right) \mathbf{X}^{*\top} \mathbf{X}^* \right\},$$

$$\mathcal{B} = E \left[\left\{ \frac{\phi_{F^{(1)}}(Y^{(1)})}{\pi(\mathbf{X}^* | \boldsymbol{\eta})} + \frac{\phi_{F^{(0)}}(Y^{(0)})}{1 - \pi(\mathbf{X}^* | \boldsymbol{\eta})} \right\} \frac{\partial}{\partial \boldsymbol{\eta}} \pi(\mathbf{X}^* | \boldsymbol{\eta}) \right]$$

and

$$\mathcal{C} = E \left\{ \frac{\phi_{F^{(1)}}^2(Y^{(1)})}{\pi(\mathbf{X}^* | \boldsymbol{\eta})} + \frac{\phi_{F^{(0)}}^2(Y^{(0)})}{1 - \pi(\mathbf{X}^* | \boldsymbol{\eta})} \right\}.$$

Theorem 3.2 (i) shows the asymptotic normality of the logistic estimates derived in Section 3.2, which ensures the validity of the treatment model throughout the estimation procedure. Theorem 3.2 (ii) states that, when $n \rightarrow \infty$, the estimated treatment effect under a given statistical functional T follows a normal distribution with sandwich type variance. With the propensity score being estimated, the estimated treatment effect $T(\hat{F}^{(1)}) - T(\hat{F}^{(0)})$ is more efficient than that under the given propensity score, which is due to the additional term $-\mathcal{B}\mathcal{A}^{-1}\mathcal{B}^\top$ from the result (i). The novel contribution in Theorem 3.2 (ii) is that the treatment effect holds for any statistical functional $T(\cdot)$. To see the generality of Theorem 3.2, we revisit (17), (19) and (20), and connect these theoretical results with Theorem 3.2 in the following discussion.

Example 1 (Average Treatment Effect) Let $T(F) = \int x dF(x)$ with $\varphi(y) = y$, then (16) reduces to (17). Thus, the influence curve is $\phi_F(y) = y - T(F)$. By Theorem 3.2 (ii), we have that, as $n \rightarrow \infty$,

$$\sqrt{n}(\hat{\tau} - \tau) \xrightarrow{d} N\left(0, -\mathcal{B}_{\text{ATE}}\mathcal{A}^{-1}\mathcal{B}_{\text{ATE}}^\top + \mathcal{C}_{\text{ATE}}\right),$$

where the components of the asymptotic variance are given by

$$\mathcal{B}_{\text{ATE}} = E\left[\left\{\frac{Y^{(1)} - E(Y^{(1)})}{\pi(\mathbf{X}^* | \boldsymbol{\eta})} + \frac{Y^{(0)} - E(Y^{(0)})}{1 - \pi(\mathbf{X}^* | \boldsymbol{\eta})}\right\} \frac{\partial}{\partial \boldsymbol{\eta}} \pi(\mathbf{X}^* | \boldsymbol{\eta})\right]$$

and

$$\mathcal{C}_{\text{ATE}} = E\left[\frac{\{Y^{(1)} - E(Y^{(1)})\}^2}{\pi(\mathbf{X}^* | \boldsymbol{\eta})} + \frac{\{Y^{(0)} - E(Y^{(0)})\}^2}{1 - \pi(\mathbf{X}^* | \boldsymbol{\eta})}\right].$$

This is the same as that in Lunceford and Davidian (2004).

Example 2 (Quantile Treatment Effect) Let $T(F) = \inf\{y : F(y) \geq q\} \triangleq \xi(q)$ for some $0 < q < 1$, then (16) reduces to the estimator (19). By Wasserman (2006, p.21), the influence curve is given by $\phi_F(y) = \frac{q - I_{(-\infty, \xi(q)]}(y)}{f(\xi(q))}$, where f is the density function of F . Theorem 3.2 (ii) is reduced to

$$\sqrt{n}\left(\hat{\Xi}(q) - \Xi(q)\right) \xrightarrow{d} N\left(0, -\mathcal{B}_{\text{QTE}}\mathcal{A}^{-1}\mathcal{B}_{\text{QTE}}^\top + \mathcal{C}_{\text{QTE}}\right)$$

as $n \rightarrow \infty$, where the components of the asymptotic variance are given by

$$\mathcal{B}_{\text{QTE}} = E\left[\left\{\frac{q - I_{(-\infty, \xi^{(1)}(q)]}(Y^{(1)})}{f^{(1)}(\xi^{(1)}(q))\pi(\mathbf{X}^* | \boldsymbol{\eta})} + \frac{q - I_{(-\infty, \xi^{(0)}(q)]}(Y^{(0)})}{f^{(0)}(\xi^{(0)}(q))[1 - \pi(\mathbf{X}^* | \boldsymbol{\eta})]}\right\} \frac{\partial}{\partial \boldsymbol{\eta}} \pi(\mathbf{X}^* | \boldsymbol{\eta})\right]$$

and

$$\mathcal{C}_{\text{QTE}} = \frac{1}{f^{(1)}(\xi^{(1)}(q))^2} E\left[\frac{(q - I_{(-\infty, \xi^{(1)}(q)]}(Y^{(1)}))^2}{\pi(\mathbf{X}^* | \boldsymbol{\eta})}\right] + \frac{1}{f^{(0)}(\xi^{(0)}(q))^2} E\left[\frac{(q - I_{(-\infty, \xi^{(0)}(q)]}(Y^{(0)}))^2}{1 - \pi(\mathbf{X}^* | \boldsymbol{\eta})}\right].$$

Note that $\frac{1}{f^{(1)}(\xi^{(1)}(q))^2} E\left[\frac{(q - I_{(-\infty, \xi^{(1)}(q)]}(Y^{(1)}))^2}{\pi(\mathbf{X}^* | \boldsymbol{\eta})}\right]$ and $\frac{1}{f^{(0)}(\xi^{(0)}(q))^2} E\left[\frac{(q - I_{(-\infty, \xi^{(0)}(q)]}(Y^{(0)}))^2}{1 - \pi(\mathbf{X}^* | \boldsymbol{\eta})}\right]$ coincide with the similar form of the semiparametric efficiency bound derived by Firpo (2007).

The difference in the forms of asymptotic variance results from the different approaches to the estimation of propensity score. To model the relationship between treatment and confounders, we employ logistic regression whereas Firpo (2007) adopt the logit series approximation from Hirano, Imbens and Ridder (2003).

Example 3 (Distributional Treatment Effect) Let $T(F) = \int I_{(-\infty, y]}(x) dF(x) = F(y)$ with $\varphi(x) = I_{(-\infty, y]}(x)$, then (16) reduces to (20). The resulting influence curve is $\phi_F(x) = I_{(-\infty, y]}(x) - T(F)$. According to Theorem 3.2 (ii), we have that, as $n \rightarrow \infty$,

$$\sqrt{n} \left(\widehat{\Delta}(y) - \Delta(y) \right) \xrightarrow{d} N \left(0, -\mathcal{B}_{\text{DTE}} \mathcal{A}^{-1} \mathcal{B}_{\text{DTE}}^\top + \mathcal{C}_{\text{DTE}} \right),$$

where the components of the asymptotic variance are given by

$$\mathcal{B}_{\text{DTE}} = E \left[\left\{ \frac{I_{(-\infty, y]}(Y^{(1)}) - F^{(1)}(y)}{\pi(\mathbf{X}^* | \boldsymbol{\eta})} + \frac{I_{(-\infty, y]}(Y^{(0)}) - F^{(0)}(y)}{1 - \pi(\mathbf{X}^* | \boldsymbol{\eta})} \right\} \frac{\partial}{\partial \boldsymbol{\eta}} \pi(\mathbf{X}^* | \boldsymbol{\eta}) \right]$$

and

$$\mathcal{C}_{\text{DTE}} = E \left[\frac{\{I_{(-\infty, y]}(Y^{(1)}) - F^{(1)}(y)\}^2}{\pi(\mathbf{X}^* | \boldsymbol{\eta})} + \frac{\{I_{(-\infty, y]}(Y^{(0)}) - F^{(0)}(y)\}^2}{1 - \pi(\mathbf{X}^* | \boldsymbol{\eta})} \right].$$

4 Numerical Studies

In this section, we design a series of synthetic data to assess the performance of the proposed method. To show the advantage of the proposed method, we also examine several existing methods.

4.1 Simulation Design

Let the dimension of covariates be $p = 12$ and let \boldsymbol{x} denote the p -dimensional confounders. We consider four different sample sizes $n = 500, 1000, 2000$ or 10000 . In the numerical studies, we design two different scenarios to generate the covariates. In Scenario 1, we consider the independent covariates and independently generate \boldsymbol{x} by the multivariate normal distribution with mean zero and identity covariance matrix, which implies that there is no network structure among confounders.

In Scenario 2, we consider the dependent covariates, which are generated by the multivariate normal distribution with mean zero and the covariance matrix reflecting two network structures in Figure 2.

When \mathbf{X} is generated, we respectively generate A and Y by the following two models

$$\text{logit} \{P(A = 1 | \mathbf{X})\} = 1 + X_1 + X_3 + \sum_{(s, \nu) \in E} X_s X_\nu \quad (22)$$

and

$$Y = \gamma_0 A + 1 + X_1 + X_3 + \sum_{(s,\nu) \in E} X_S X_\nu + \varepsilon, \quad (23)$$

where $\varepsilon \sim N(0, 1)$, $\gamma_0 = 1$ and E is the set that is either empty under Scenario 1 or contains pairs in Scenario 2. From (22) and (23), variables X_1 and X_3 and pairs in E are confounders. Consequently, we obtain the IID sample $\mathcal{O} = \left\{ \{Y_i, A_i, \mathbf{X}_i : i = 1, \dots, n\} \right\}$ with size n .

In this study, we primarily examine the estimators (17), (19) with $q \in \{0.2, 0.25, 0.5, 0.75, 0.8\}$, and (20) with $y \in \{-3, 0, 3\}$. We compare the proposed method with existing approaches. For the estimation of the ATE, we primarily examine the inverse probability weighted estimator under the outcome Y_i

$$\tau_{\text{IPW}} = \frac{1}{n} \left(\sum_{i=1}^n \frac{A_i Y_i}{\hat{\pi}(\mathbf{X}_i)} \right) - \frac{1}{n} \left(\sum_{i=1}^n \frac{(1 - A_i) Y_i}{1 - \hat{\pi}(\mathbf{X}_i)} \right) \quad (24)$$

and the modified version proposed by Lunceford and Davidian (2004)

$$\tau_{\text{LD}} = \left(\sum_{i=1}^n \frac{A_i}{\hat{\pi}(\mathbf{X}_i)} \right)^{-1} \left(\sum_{i=1}^n \frac{A_i Y_i}{\hat{\pi}(\mathbf{X}_i)} \right) - \left(\sum_{i=1}^n \frac{1 - A_i}{1 - \hat{\pi}(\mathbf{X}_i)} \right)^{-1} \left(\sum_{i=1}^n \frac{(1 - A_i) Y_i}{1 - \hat{\pi}(\mathbf{X}_i)} \right), \quad (25)$$

where (24) and (25) do not take variable selection into account.

For the estimation of the QTE, we examine the approach proposed in Firpo (2007), which is given by

$$\hat{\Xi}_{\text{Firpo}}(q) = \hat{\xi}_{\text{Firpo}}^{(1)}(q) - \hat{\xi}_{\text{Firpo}}^{(0)}(q), \quad (26)$$

where

$$\hat{\xi}_{\text{Firpo}}^{(0)}(q) = \underset{q}{\operatorname{argmin}} \frac{1}{n} \sum_{i=1}^n \frac{1 - A_i}{1 - \hat{\pi}(\mathbf{X}_i)} \cdot \rho_q(Y_i - q) \quad \text{and} \quad \hat{\xi}_{\text{Firpo}}^{(1)}(q) = \underset{q}{\operatorname{argmin}} \frac{1}{n} \sum_{i=1}^n \frac{A_i}{\hat{\pi}(\mathbf{X}_i)} \cdot \rho_q(Y_i - q)$$

with ρ_q being the check function stated in Koenker (2005). Noting that the estimators (19) and (26) form a group of comparison, where the former incorporates variable selection in the estimation procedure while the latter does not.

To assess the performance of variable selection, we compute the specificity (SPE) and sensitivity (SEN), which are respectively given by

$$\text{SEN} = \frac{\#\left\{ j, (s, \nu) : \theta_{s\nu} = 0, \hat{\theta}_{s\nu} = 0, \beta_j = 0, \hat{\beta}_j = 0 \right\}}{\#\left\{ j, (s, \nu) : \theta_{s\nu} = 0, \beta_j = 0 \right\}}$$

and

$$\text{SPE} = \frac{\#\left\{j, (s, \nu) : \theta_{s\nu} \neq 0, \widehat{\theta}_{s\nu} \neq 0, \beta_j \neq 0, \widehat{\beta}_j \neq 0\right\}}{\#\left\{j, (s, \nu) : \theta_{s\nu} \neq 0, \beta_j \neq 0\right\}}.$$

In addition, to examine the performance of the estimators derived by the proposed or existing methods, we compute the biases (BIAS), standard errors (S.E.), mean squared errors (MSE) and coverage rate (CR). For each setting, we repeat the simulation 1000 times.

4.2 Simulation Results

The simulation results of ATE, QTE and DTE under Scenario 1 are summarized in Tables 1, 2 and 3, respectively; the results under Scenario 2 are recorded in Tables 4, 5 and 6, respectively. For variable selection, we find that both sensitivity and specificity are close or equal to one regardless of network structures, which indicate that the estimated graphical structure and selected informative confounders can be recovered to the true ones and are consistent with the finding in Theorem 3.1.

From the estimation of various treatments, our CDF approach generally outperforms the existing methods and is more efficient with smaller MSE. The CR of the proposed method is also close to 95%. To see the performance of each treatment effects, we first observe from ATE that our method shows numerical equivalence to the method from Lunceford and Davidian (2004), which matches the finding in Example 1 in Section 3.4. In addition, the proposed and Lunceford and Davidian’s methods are more efficient than the traditional IPW method with smaller S.E. and MSE. When we look at the QTE, we find that our method outperforms Firpo’s (2007) method with smaller bias and MSE even in Scenario 1. It indicates that the proposed CDF approach is robust. Finally, the performance for DTE shows that our method is capable of offering precise estimates due to the lower bias, S.E. and MSE and approximate 95% CR.

5 Analysis of NHEFS data

In this section, we implement the proposed method in Section 3 to analyze the data arising from NHANES I Epidemiologic Follow-up Study (NHEFS), which is a national longitudinal study collaboratively launched by the National Center for Health Statistics and the National Institute on Aging. The NHEFS aimed to analyze the links between clinical, nutritional, and behavioral variables. For the detailed description, one can refer to <https://www.cdc.gov/nchs/nhanes/nhefs/default.aspx> for the official information.

The full data set, which is available at <https://miguelhernan.org/whatifbook>, contains 1430 subjects and several variables including weight measured in kilograms (`wt`), the status of smoking behavior (`qsmk`), systolic blood pressure (`sbp`, in millimeter of mercury (mmHg)), serum cholesterol (`cholesterol`, in mg/100), number of cigarettes smoked per day in 1971 (`smokeintensity`), diastolic blood pressure (`dbp`, in mmHg), height in centimeters (`ht`), average tobacco price in the state of residence in 1982 (`price82`, in US dollars), years of smoking (`smokeyrs`), age in 1971 (`age`, in years), and total family income in 1971 (`income`). We aim to explore the causal effects of smoking habits on the change of weight. In this study, we aim to explore various types of causal effects of weights from different status of smoking behavior. Following the notation from Section 2, we take Y and A as `wt` and `qsmk`, respectively. Moreover, we let the remaining nine variables be confounders. Since those nine confounders may form pairwise dependence structure and part of them is informative to Y , it is crucial to implement variable selection technique in Section 3 and then estimate the treatment effects. In addition, to see the impact of variable selection, we apply the existing methods in Section 4 for the comparisons.

The proposed method suggests that the selected confounders are `ht` and `age`, and the network structure of nine confounders is displayed in Figure 3. The revealed dependence structure is centered around `race` and `sex` while `age` does not contribute to the dependence structure with other variables. This indicates that when it comes to smoking behavior and weight gain, `race` and `sex` are two critical variables that possess further underlying connections with other covariates and should be taken into account while analyzing causality in the data.

Our interest lies in estimating (2), (4) with $q = 0.2, 0.25, 0.5, 0.75$ and 0.8 , and (5) evaluated at $y = \bar{Y}$, the sample average of the outcome. The estimation results of the proposed and existing methods are recorded in Table 7, where EST represents the estimator, S.E. is the standard error obtained by the bootstrap method with 10000 repetitions, and p-value is used to examine two-tailed hypotheses $H_0 : \tau = 0$, $H_0 : \Xi(q) = 0$ or $H_0 : \Delta(y) = 0$ for some pre-specified q and y . We find all estimates are greater than 2 except for DTE. Consistent with the finding in simulation, the proposed CDF method is more efficient with smaller S.E. Under level of significance 0.05, all estimates are significant except for $\Xi(0.75)$, which indicate that smoking generally causes an increase in weight. Regarding the results of $\Xi(0.75)$, we comment that Firpo’s method might falsely reject the null hypothesis, i.e. $H_0 : \Xi(0.75) = 0$, as the simulation results suggested that Firpo’s method rarely captures the true parameter from the perspective of CR.

6 Summary

Treatment effect estimations have been an important issue in causal inference, which include but not are not limited to ATE, QTE and DTE. The other challenge for the estimation is the multivariate/high-dimensional data with potential network structure. To tackle these problems, we propose a flexible and unified methodology that is based on statistical functional structure and accommodates the estimation of various causal effects, including ATE, QTE, DTE. Our approach is under the CDF approach and takes the detection of network structure among covariates into account. Theoretically, the variable selection consistency and asymptotic normality properties of the statistical functional estimator are rigorously established, which ensure the validity of the proposed estimator. Numerical studies show that the finite sample performance of the proposed method is satisfactory and even outperform other existing methods.

The findings illustrate the framework’s flexibility in estimating causal effects. The proposed method offers a versatile and effective solution for exploring diverse causal measures in complex datasets. This unified approach represents a significant advance in the field of causal inference, providing researchers with a powerful tool for uncovering insights in

high-dimensional data.

References

- Chapfuwa, P., Assaad, S., Zeng, S., Pencina, M. J., Carin, L. and Henao, R. (2021, April). Enabling counterfactual survival analysis with balanced representations. In *Proceedings of the Conference on Health, Inference, and Learning* (pp. 133-145).
- Chen, L. P. (2020). Causal inference for left-truncated and right-censored data with covariate measurement error. *Computational and Applied Mathematics*, 39, 1-27.
- Chen, L.-P. (2024). Estimation of graphical models: an overview of selected topics. *International Statistical Review*, 92: 194-245.
- Chen, L.-P. and Yi, G. Y. (2021). Analysis of noisy survival data with graphical proportional hazards measurement error model. *Biometrics*, 77, 956-969.
- Chen, L.-P., Yi, G. Y., Zhang, Q. and He, W. (2019) Multiclass analysis and prediction with network structured covariates. *Journal of Statistical Distributions and Applications*, 6:6.
- Chernozhukov, V., Fernández-Val, I. and Melly, B. (2013). Inference on counterfactual distributions. *Econometrica*, 81, 2205-2268.
- Donald, S. G. and Hsu, Y. C. (2014). Estimation and inference for distribution functions and quantile functions in treatment effect models. *Journal of Econometrics*, 178, 383-397.
- Ertefaie, A., Asgharian, M. and Stephens, D. A. (2018). Variable selection in causal inference using a simultaneous penalization method. *Journal of Causal Inference*, 6, 20170010.
- Firpo, S. (2007). Efficient semiparametric estimation of quantile treatment effects. *Econometrica*, 75, 259-276.
- Hirano, K., Imbens, G. W. and Ridder, G. (2003). Efficient estimation of average treatment effects using the estimated propensity score. *Econometrica*, 71, 1161-1189.
- Hsu, Y. C., Lai, T. C. and Lieli, R. P. (2022). Counterfactual treatment effects: Estimation and inference. *Journal of Business & Economic Statistics*, 40, 240-255.

- Khan, S. and Ugander, J. (2023). Adaptive normalization for IPW estimation. *Journal of Causal Inference*, 11, 20220019.
- Koch, B., Vock, D. M., Wolfson, J. and Vock, L. B. (2020). Variable selection and estimation in causal inference using Bayesian spike and slab priors. *Statistical Methods in Medical Research*, 29, 2445-2469.
- Koenker, R. (2005). *Quantile Regression*. Cambridge University Press.
- Lunceford, J. K. and Davidian, M. (2004). Stratification and weighting via the propensity score in estimation of causal treatment effects: a comparative study. *Statistics in Medicine*, 23, 2937-2960
- Rosenbaum, P. R. and Rubin, D. B. (1983) The central role of the propensity score in observational studies for causal effects. *Biometrika*, 70, 41-55.
- Shortreed, S. M. and Ertefaie, A. (2017). Outcome-adaptive lasso: variable selection for causal inference. *Biometrics*, 73, 1111-1122.
- van der Vaart, A. W. (2000). *Asymptotic Statistics*. Cambridge university press.
- von Mises, R. (1947). On the asymptotic distribution of differentiable statistical functions. *The Annals of Mathematical Statistics*, 18, 309-348.
- Wasserman, L. (2006). *All of nonparametric statistics*. Springer Science & Business Media.
- Yi, G. Y. and Chen, L. P. (2023). Estimation of the average treatment effect with variable selection and measurement error simultaneously addressed for potential confounders. *Statistical Methods in Medical Research*, 32, 691-711.
- Zou, H. (2006). The adaptive lasso and its oracle properties. *Journal of the American Statistical Association*, 101, 1418-1429.

Table 1: Simulation results for ATE under Scenario 1

n	Method	SEN	SPE	BIAS	S.E.	MSE	CR
500	IPW	-	-	0.010	0.344	0.118	0.970
	LD	-	-	0.026	0.251	0.063	0.959
	CDF	0.998	1.000	0.026	0.251	0.063	0.959
1000	IPW	-	-	0.006	0.261	0.068	0.962
	LD	-	-	0.017	0.195	0.038	0.957
	CDF	0.999	1.000	0.017	0.195	0.038	0.957
2000	IPW	-	-	0.002	0.175	0.030	0.963
	LD	-	-	0.007	0.138	0.019	0.961
	CDF	0.999	1.000	0.007	0.138	0.019	0.961
10000	IPW	-	-	-0.000	0.088	0.007	0.969
	LD	-	-	0.000	0.070	0.005	0.968
	CDF	1.000	1.000	0.000	0.070	0.005	0.968

Table 2: Simulation results for QTE under Scenario 1

n	Estimand	Method	SEN	SPE	BIAS	S.E.	MSE	CR	
500	$\Xi(0.2)$	Firpo	-	-	1.457	0.209	2.124	0.000	
		CDF	0.998	1.000	0.004	0.222	0.000	0.953	
	$\Xi(0.25)$	Firpo	-	-	1.454	0.198	2.115	0.000	
		CDF	0.998	1.000	0.007	0.204	0.000	0.958	
	$\Xi(0.5)$	Firpo	-	-	1.475	0.179	2.177	0.000	
		CDF	0.998	1.000	-0.008	0.282	0.000	0.981	
	$\Xi(0.75)$	Firpo	-	-	1.509	0.196	2.279	0.000	
		CDF	0.998	1.000	-0.023	0.484	0.000	0.967	
	$\Xi(0.8)$	Firpo	-	-	1.525	0.213	2.325	0.000	
		CDF	0.998	1.000	-0.017	0.533	0.000	0.959	
	1000	$\Xi(0.2)$	Firpo	-	-	1.447	0.153	2.096	0.000
			CDF	0.948	1.000	0.005	0.150	0.000	0.946
$\Xi(0.25)$		Firpo	-	-	1.443	0.144	2.082	0.000	
		CDF	0.999	1.000	0.001	0.140	0.000	0.950	
$\Xi(0.5)$		Firpo	-	-	1.466	0.133	2.151	0.000	
		CDF	0.999	1.000	-0.003	0.163	0.000	0.953	
$\Xi(0.75)$		Firpo	-	-	1.495	0.145	2.236	0.000	
		CDF	0.999	1.000	-0.007	0.298	0.000	0.955	
$\Xi(0.8)$		Firpo	-	-	1.502	0.147	2.257	0.000	
		CDF	0.999	1.000	-0.007	0.350	0.000	0.956	

2000	$\Xi(0.2)$	Firpo	-	-	1.445	0.107	2.090	0.000	
		CDF	0.999	1.000	-0.006	0.108	0.006	0.951	
	$\Xi(0.25)$	Firpo	-	-	1.448	0.103	2.097	0.000	
		CDF	0.999	1.000	-0.006	0.100	0.003	0.944	
	$\Xi(0.5)$	Firpo	-	-	1.467	0.090	2.153	0.000	
		CDF	0.999	1.000	-0.007	0.125	0.004	0.954	
	$\Xi(0.75)$	Firpo	-	-	1.502	0.099	2.258	0.000	
		CDF	0.999	1.000	-0.010	0.237	0.037	0.967	
	$\Xi(0.8)$	Firpo	-	-	1.514	0.106	2.295	0.000	
		CDF	0.999	1.000	-0.017	0.291	0.100	0.962	
	10000	$\Xi(0.2)$	Firpo	-	-	1.443	0.047	2.084	0.000
			CDF	0.971	1.000	-0.002	0.047	0.000	0.949
		$\Xi(0.25)$	Firpo	-	-	1.445	0.045	2.089	0.000
			CDF	0.971	1.000	-0.001	0.044	0.000	0.948
$\Xi(0.5)$		Firpo	-	-	1.465	0.042	2.147	0.000	
		CDF	0.971	1.000	-0.001	0.050	0.001	0.956	
$\Xi(0.75)$		Firpo	-	-	1.499	0.045	2.247	0.000	
		CDF	0.971	1.000	-0.001	0.087	0.000	0.950	
$\Xi(0.8)$		Firpo	-	-	1.509	0.047	2.279	0.000	
		CDF	0.971	1.000	-0.000	0.103	0.000	0.947	

Table 3: Simulation results for DTE under Scenario 1

n	Estimand	SEN	SPE	BIAS	S.E.	MSE	CR
500	$\Delta(0)$	0.998	1.000	-0.000	0.037	0.000	0.954
	$\Delta(3)$	0.998	1.000	-0.004	0.070	0.000	0.962
	$\Delta(-3)$	0.998	1.000	0.149	0.008	0.022	0.961
1000	$\Delta(0)$	0.999	1.000	-0.000	0.028	0.000	0.958
	$\Delta(3)$	0.999	1.000	0.000	0.054	0.000	0.959
	$\Delta(-3)$	0.999	1.000	0.148	0.006	0.022	0.957
2000	$\Delta(0)$	0.999	1.000	-0.000	0.019	0.000	0.948
	$\Delta(3)$	0.999	1.000	-0.000	0.037	0.000	0.959
	$\Delta(-3)$	0.999	1.000	0.149	0.005	0.022	0.955
10000	$\Delta(0)$	1.000	1.000	-0.000	0.008	0.000	0.952
	$\Delta(3)$	1.000	1.000	-0.000	0.016	0.000	0.953
	$\Delta(-3)$	1.000	1.000	0.149	0.002	0.022	0.959

Table 4: Simulation results for ATE under Scenario 2

n	Method	Hub						Lattice					
		SEN	SPE	BIAS	S.E.	MSE	CR	SEN	SPE	BIAS	S.E.	MSE	CR
500	IPW	-	-	0.446	3.181	10.312	0.981	-	-	0.227	3.206	10.320	0.992
	LD	-	-	0.929	1.047	1.960	0.924	-	-	0.530	0.869	1.036	0.969
	CDF	0.911	1.000	0.929	1.047	1.960	0.924	0.961	1.000	0.530	0.869	1.036	0.969
1000	IPW	-	-	-0.018	6.771	45.808	0.992	-	-	-0.045	5.035	25.332	0.990
	LD	-	-	0.678	1.130	1.736	0.969	-	-	0.391	0.899	0.961	0.971
	CDF	0.945	1.000	0.678	1.130	1.736	0.969	0.982	1.000	0.391	0.899	0.961	0.971
2000	IPW	-	-	0.130	5.539	30.668	0.992	-	-	0.175	1.719	2.983	0.983
	LD	-	-	0.576	1.030	1.392	0.980	-	-	0.321	0.714	0.612	0.970
	CDF	0.970	1.000	0.576	1.030	1.392	0.980	0.992	1.000	0.321	0.714	0.612	0.970
10000	IPW	-	-	0.147	2.212	4.913	0.987	-	-	0.019	2.834	8.026	0.993
	LD	-	-	0.335	0.855	0.843	0.965	-	-	0.181	0.697	0.518	0.972
	CDF	0.996	1.000	0.335	0.855	0.843	0.965	0.999	1.000	0.181	0.697	0.518	0.972

Table 5: Simulation results for QTE under Scenario 2

n	Estimand	Method	Hub						Lattice						
			SEN	SPE	BIAS	S.E.	MSE	CR	SEN	SPE	BIAS	S.E.	MSE	CR	
500	$\Xi(0.2)$	Firpo	-	-	4.121	0.375	16.987	0.000	-	-	2.813	0.285	7.913	0.000	
		CDF	0.970	1.000	0.353	1.271	0.124	0.959	0.961	1.000	0.053	1.090	0.002	0.964	
	$\Xi(0.25)$	Firpo	-	-	3.934	0.338	15.482	0.000	-	-	2.751	0.261	7.572	0.000	
		CDF	0.970	1.000	0.263	1.276	0.069	0.961	0.961	1.000	0.035	1.040	0.001	0.966	
	$\Xi(0.5)$	Firpo	-	-	3.602	0.288	12.979	0.000	-	-	2.749	0.241	7.558	0.000	
		CDF	0.970	1.000	0.156	1.284	0.024	0.951	0.961	1.000	0.009	1.066	0.000	0.959	
	$\Xi(0.75)$	Firpo	-	-	3.973	0.321	15.786	0.000	-	-	3.277	0.295	10.742	0.000	
		CDF	0.970	1.000	0.460	1.343	0.211	0.951	0.961	1.000	0.215	1.275	0.046	0.947	
	$\Xi(0.8)$	Firpo	-	-	4.189	0.347	17.554	0.000	-	-	3.513	0.325	12.343	0.000	
		CDF	0.970	1.000	0.693	1.337	0.480	0.954	0.961	1.000	0.387	1.305	0.150	0.953	
	1000	$\Xi(0.2)$	Firpo	-	-	4.107	0.276	16.870	0.000	-	-	2.810	0.198	7.901	0.000
			CDF	0.948	1.000	0.243	1.177	0.059	0.957	0.992	1.000	-0.036	0.993	0.001	0.962
$\Xi(0.25)$		Firpo	-	-	3.918	0.246	15.356	0.000	-	-	2.751	0.189	7.568	0.000	
		CDF	0.948	1.000	0.197	1.114	0.039	0.961	0.992	1.000	-0.040	0.971	0.001	0.967	
$\Xi(0.5)$		Firpo	-	-	3.612	0.195	13.046	0.000	-	-	2.751	0.174	7.568	0.000	
		CDF	0.948	1.000	0.178	1.060	0.032	0.971	0.992	1.000	-0.005	1.002	0.000	0.967	
$\Xi(0.75)$		Firpo	-	-	4.001	0.232	16.008	0.000	-	-	3.279	0.210	10.751	0.000	
		CDF	0.948	1.000	1.397	1.237	0.158	0.956	0.992	1.000	0.154	1.176	0.023	0.945	
$\Xi(0.8)$		Firpo	-	-	4.214	0.255	17.759	0.000	-	-	3.512	0.228	12.335	0.000	
		CDF	0.948	1.000	0.541	1.289	0.293	0.967	0.992	1.000	0.292	1.184	0.085	0.951	

2000	$\Xi(0.2)$	Firpo	-	-	4.113	0.186	16.924	0.000	-	-	2.823	0.140	7.971	0.000	
		CDF	0.971	1.000	0.082	1.362	0.006	0.959	0.981	1.000	0.013	0.769	0.000	0.980	
	$\Xi(0.25)$	Firpo	-	-	3.928	0.170	15.431	0.000	-	-	2.762	0.134	7.628	0.000	
		CDF	0.971	1.000	0.056	1.312	0.003	0.965	0.981	1.000	0.021	0.971	0.000	0.983	
	$\Xi(0.5)$	Firpo	-	-	3.607	0.135	13.016	0.000	-	-	2.760	0.115	7.621	0.000	
		CDF	0.971	1.000	0.063	1.130	0.004	0.967	0.981	1.000	0.047	1.002	0.002	0.982	
	$\Xi(0.75)$	Firpo	-	-	3.997	0.155	15.981	0.000	-	-	3.277	0.140	10.742	0.000	
		CDF	0.971	1.000	0.193	1.265	0.037	0.945	0.981	1.000	0.085	1.176	0.007	0.953	
	$\Xi(0.8)$	Firpo	-	-	4.219	0.178	17.806	0.000	-	-	3.509	0.156	12.314	0.000	
		CDF	0.971	1.000	0.317	1.292	0.100	0.946	0.981	1.000	0.160	1.184	0.025	0.943	
	10000	$\Xi(0.2)$	Firpo	-	-	4.110	0.085	16.893	0.000	-	-	2.817	0.065	7.935	0.000
			CDF	0.971	1.000	-0.015	1.013	0.000	0.972	0.981	1.000	0.010	0.484	0.000	0.985
		$\Xi(0.25)$	Firpo	-	-	3.921	0.076	15.377	0.000	-	-	2.755	0.060	7.593	0.000
			CDF	0.971	1.000	0.003	0.896	0.000	0.977	0.981	1.000	0.056	1.312	0.000	0.985
$\Xi(0.5)$		Firpo	-	-	3.604	0.062	12.990	0.000	-	-	2.757	0.055	7.604	0.000	
		CDF	0.971	1.000	0.044	0.719	0.001	0.979	0.981	1.000	0.063	1.130	0.000	0.991	
$\Xi(0.75)$		Firpo	-	-	3.989	0.071	15.919	0.000	-	-	3.276	0.064	10.736	0.000	
		CDF	0.971	1.000	0.020	1.205	0.000	0.961	0.981	1.000	0.193	1.265	0.000	0.967	
$\Xi(0.8)$		Firpo	-	-	4.210	0.079	17.729	0.000	-	-	3.509	0.070	12.316	0.000	
		CDF	0.971	1.000	0.056	1.318	0.000	0.953	0.981	1.000	0.317	1.292	0.000	0.958	

Table 6: Simulation results for DTE under Scenario 2

n	Estimand	Hub						Lattice					
		SEN	SPE	BIAS	S.E.	MSE	CR	SEN	SPE	BIAS	S.E.	MSE	CR
500	$\Delta(0)$	0.913	1.000	-0.066	0.103	0.004	0.928	0.961	1.000	-0.039	0.079	0.001	0.939
	$\Delta(3)$	0.913	1.000	-0.088	0.150	0.007	0.957	0.961	1.000	-0.058	0.134	0.003	0.962
	$\Delta(-3)$	0.913	1.000	-0.044	0.084	0.002	0.975	0.961	1.000	-0.007	0.056	0.000	0.971
1000	$\Delta(0)$	0.945	1.000	-0.045	0.091	0.002	0.964	0.982	1.000	-0.024	0.071	0.000	0.959
	$\Delta(3)$	0.945	1.000	-0.052	0.137	0.002	0.960	0.982	1.000	-0.041	0.117	0.001	0.963
	$\Delta(-3)$	0.945	1.000	-0.026	0.096	0.000	0.959	0.982	1.000	0.001	0.050	0.000	0.950
2000	$\Delta(0)$	0.970	1.000	-0.031	0.079	0.000	0.966	0.992	1.000	-0.016	0.058	0.000	0.956
	$\Delta(3)$	0.970	1.000	-0.044	0.109	0.001	0.964	0.992	1.000	-0.029	0.103	0.000	0.961
	$\Delta(-3)$	0.970	1.000	-0.022	0.078	0.000	0.967	0.992	1.000	0.004	0.041	0.000	0.961
10000	$\Delta(0)$	0.996	1.000	-0.014	0.074	0.000	0.964	0.999	1.000	-0.008	0.036	0.000	0.963
	$\Delta(3)$	0.996	1.000	-0.023	0.074	0.000	0.973	0.999	1.000	-0.010	0.071	0.000	0.962
	$\Delta(-3)$	0.996	1.000	-0.009	0.061	0.000	0.972	0.999	1.000	0.009	0.028	0.000	0.946

Table 7: Real data analysis: estimation of various treatment effects by the proposed and existing methods.

Estimand	Method	EST	S.E.	p-value
ATE	IPW	3.841	1.128	0.000
	LD	3.568	0.894	0.000
	CDF	3.568	0.894	0.000
$\Xi(0.2)$	Firpo	3.430	1.107	0.000
	CDF	2.686	0.864	0.000
$\Xi(0.25)$	Firpo	3.933	1.344	0.001
	CDF	2.803	1.302	0.015
$\Xi(0.5)$	Firpo	4.618	1.500	0.001
	CDF	3.284	1.342	0.007
$\Xi(0.75)$	Firpo	3.263	1.521	0.015
	CDF	2.403	1.514	0.056
$\Xi(0.8)$	Firpo	3.270	1.409	0.010
	CDF	2.476	1.259	0.024
$\Delta(\bar{Y})$	CDF	-0.072	0.027	0.004




Figure 1: Diagram of causal inference with network covariates




Figure 2: Covariate network structures for simulation studies. The left panel displays the lattice structure, where X_1 , X_2 and X_3 form a triangle, X_4 , X_5 and X_6 form the other, and X_7 to X_{12} are not connected; the right panel displays the hub structure, which X_1 to X_6 and X_7 to X_{12} are interwoven, respectively.




Figure 3: Real data analysis result: estimated network structure for nine confounders in NHEFS data.

March 27, 2025

Supporting Information for “A Unified Approach for Estimating Various Treatment Effect in Causal Inference”

Kuan-Hsun Wu and Li-Pang Chen¹

Department of Statistics, National Chengchi University

Abstract

The supporting information contains a list of regularity conditions and the detailed proofs of theorems in the manuscript, entitled “A Unified Approach for Estimating Various Treatment Effect in Causal Inference”.

Keywords: Causal inference; counterfactual distributions; plug-in estimators; high-dimensional; network structures; graphical model; propensity score; variable selection.

Short title: Estimating Various Causal Effects Using IPW CDF

¹Corresponding Author. Email: lchen723@nccu.edu.tw

A Regularity Conditions

We list the following conditions that are imposed to the derivation:

(A1) The matrices $\mathbf{X}_V^\top \mathbf{X}_V$ and $(\mathbf{X}_E^{\circ 2})^\top \mathbf{X}_E^{\circ 2}$ are nonsingular, where \mathbf{X}_V and $\mathbf{X}_E^{\circ 2}$ are the matrices consisting of only the variables whose indices are recorded by sets V and E , respectively. That is, the matrices $(\mathbf{X}_V^\top \mathbf{X}_V)^{-1}$ and $[(\mathbf{X}_E^{\circ 2})^\top \mathbf{X}_E^{\circ 2}]^{-1}$ exist.

(A2) There exist two constants γ_1 and $\gamma_2 \in (0, 1)$, such that

$$\max_{i \in V^C} \|(\mathbf{X}_V^\top \mathbf{X}_V)^{-1} \mathbf{X}_V^\top \mathbf{X}_i\|_1 \leq 1 - \gamma_1 \text{ and } \max_{i \in E^C} \|[(\mathbf{X}_E^{\circ 2})^\top (\mathbf{X}_E^{\circ 2})]^{-1} (\mathbf{X}_E^{\circ 2})^\top \mathbf{X}_i^{\circ 2}\|_1 \leq 1 - \gamma_2,$$

where \mathbf{X}_i is the i th column of matrix \mathbf{X} and $\mathbf{X}_i^{\circ 2}$ is the i th column of matrix $\mathbf{X}^{\circ 2}$.

(A3) There exist four positive constants a_1, a_2, b_1 and b_2 , such that

$$E(\mathbb{X}_r) < a_1, E(\mathbb{X}_r^\top \mathbb{X}_r) < a_2, E(\mathbb{X}_r^{\circ 2}) < b_1 \text{ and } E\left\{(\mathbb{X}^{\circ 2})^\top (\mathbb{X}^{\circ 2})\right\} < b_2$$

where \mathbb{X}_r is the r th column of matrix \mathbb{X} .

(A4) There exist four positive constants c_1, c_2, d_1 and d_2 , such that

$$\Lambda_{\min}(\mathbb{X}) > c_1, \Lambda_{\max}(\mathbb{X}) < c_2, \Lambda_{\min}(\mathbb{X}^{\circ 2}) > d_1 \text{ and } \Lambda_{\max}(\mathbb{X}^{\circ 2}) < d_2.$$

where $\Lambda_{\min}(\mathbf{A})$ and $\Lambda_{\max}(\mathbf{A})$ represent the minimum and maximum eigenvalues of the matrix \mathbf{A} , respectively.

(A5) There exists a positive constant K_{clm} , such that

$$\max_{j=1,2,\dots,p} \frac{\|\mathbf{X}_j\|_2}{\sqrt{n}} \leq K_{\text{clm}}.$$

(A6) The true CDFs of $Y^{(1)}$ and $Y^{(0)}$ are continuous. Moreover, the derivatives of $F^{(1)}$ and $F^{(0)}$ exist, and the corresponding density functions are denoted by $f^{(1)}(y)$ and $f^{(0)}(y)$ of $Y^{(1)}$ and $Y^{(0)}$, respectively.

Assumptions (A1)-(A5) state the conditions on the behavior of design matrices $\mathbf{X}, \mathbf{X}^{\circ 2}, \mathbb{X}$ and $\mathbb{X}^{\circ 2}$, ensuring the sparsity recovery for main and interaction effects holds. Specifically, (A1) is necessary for computing the regression coefficients using \mathbf{X}_V and $\mathbf{X}_E^{\circ 2}$. (A2) is often

called *mutual incoherence*, which restricts the dependence of variables between in V (or E) and V^C (or E^C). (A3) is used to guarantee the existence of variance of confounders. The eigenvalues in (A4) are required to ensure that the covariates are not excessively linear dependent (e.g., Yang et al., 2015). In addition, (A5) states that the columns of the design matrix are normalized. Following the last remark of Section 2.1, (A6) concentrates our discussion on the continuous outcome.

B Some Lemmas

In this section, we introduce some useful lemmas for the derivations of the main theorems.

Lemma B.1 (Gaussian Tail Bound.) *Let $X \sim N(\mu, \sigma^2)$. The Gaussian tail bound is, for any $\epsilon > 0$,*

$$P(|X| > \epsilon) \leq 2 \exp\left(-\frac{(\epsilon - \mu)^2}{2\sigma^2}\right).$$

We demonstrate the derivation of Lemma B.1 by using the Chernoff's bound, which states that, for any $\epsilon > 0$,

$$P(X - \mu > \epsilon) \leq \exp\left(-\frac{\epsilon^2}{2\sigma^2}\right). \tag{B.1}$$

Let $\epsilon^* = \epsilon - \mu$. Directly applying (B.1) yields that

$$\begin{aligned} P(X > \epsilon) &= P(X - \mu > \epsilon - \mu) \\ &= P(X - \mu > \epsilon^*) \\ &\leq \exp\left(-\frac{\epsilon^{*2}}{2\sigma^2}\right) \\ &= \exp\left(-\frac{(\epsilon - \mu)^2}{2\sigma^2}\right). \end{aligned}$$

Due to the symmetry of normally distributed random variables and the union bound, the Gaussian tail bound is obtained.

Lemma B.2 (Expansion of Statistical Functionals.) *Assume that \widehat{F} , an estimated CDF derived by the IID sample, converges to F in probability uniformly over $y \in \mathbb{R}$, where F is a CDF of a random variable Y . Consider a statistical functional T with influence curve ϕ_F*

defined in Section 3.4. Therefore, the asymptotic variance of $\sqrt{n} \left(T(\widehat{F}) - T(F) \right)$ is determined only by the dominant term $\frac{1}{\sqrt{n}} \sum_{i=1}^n \phi_F(Y_i)$.

Proof:

Let $D[0, 1]$ be the space of right-continuous real-valued functions on $[0, 1]$ with left-hand limits. That is, for any $R \in D[0, 1]$, we have that

$$\lim_{t \rightarrow t^+} R(t) = R(t)$$

for any $t \in [0, 1)$ and

$$\lim_{t \rightarrow t^-} R(t)$$

exists for any $t \in (0, 1]$. We define $\rho : D[0, 1] \rightarrow \mathbb{R}$, the functional induced by a statistical functional T , as

$$\rho(R) = T(R \circ F)$$

Let U denote the distribution function from the uniform distribution in an interval $[0, 1]$, say

$$U(t) = \begin{cases} 0 & \text{if } t < 0; \\ t & \text{if } 0 \leq t < 1; \\ 1 & \text{if } t \geq 1. \end{cases} \quad (\text{B.2})$$

We have that $U = F \circ F^{-1}$ and $\widehat{U} \triangleq \widehat{F} \circ F^{-1}$, where F^{-1} is the inverse distribution function of F . By Lemma 4.4.1 of Fernholz (1983), if the induced functional ρ is Hadamard differentiable at the uniform distribution on $[0, 1]$, that is, there exists a continuous linear function ρ'_U such that

$$\frac{\rho(U + t_n h_n) - \rho(U)}{t_n} \rightarrow \rho'_U(h) \text{ as } n \rightarrow \infty,$$

for all sequences $\{t_n\} \subset \mathbb{R}$ and $\{h_n\} \in D[0, 1]$ satisfying $t_n \rightarrow 0$, $h_n \rightarrow h \in D[0, 1]$ and $U + t_n h_n \in D[0, 1]$, then combining with (21) gives that

$$\phi_F(y) = \rho'_U((\delta_y - U) \circ F^{-1}). \quad (\text{B.3})$$

An application of Hadamard derivative yields that

$$\begin{aligned}
\sqrt{n}\left(T(\widehat{F}) - T(F)\right) &= \sqrt{n}\left(\rho'_U(\widehat{U} - U)\right) + \sqrt{n}\text{Rem}(\widehat{U} - U) \\
&= \sqrt{n}\left[\rho'_U\left((\widehat{F} - F) \circ F^{-1}\right)\right] + \sqrt{n}\text{Rem}(\widehat{U} - U) \\
&= \sqrt{n}\left[\rho'_U\left(\frac{1}{n}\sum_{i=1}^n(\delta_{Y_i} - F) \circ F^{-1}\right)\right] + \sqrt{n}\text{Rem}(\widehat{U} - U) \\
&= \frac{1}{\sqrt{n}}\sum_{i=1}^n\phi_F(Y_i) + \sqrt{n}\text{Rem}(\widehat{U} - U),
\end{aligned} \tag{B.4}$$

where the first equality is shown in Fernholz (1983, p.40), last equality is obtained with (B.3) and $\text{Rem}(\cdot)$ is the remainder term defined in Fernholz (1983). By Proposition 4.3.3 and Theorem 4.4.2 in Fernholz (1983), we have that $\text{Rem}(\widehat{U} - U) \xrightarrow{P} 0$ as $n \rightarrow \infty$. Therefore, the claim is justified.

C Justification of CDF of (8) and (9)

To verify that (8) and (9) form CDF, it suffices to check the properties of CDF. Specifically,

$$\begin{aligned}
\lim_{y \rightarrow \infty} \widehat{F}^{(1)}(y) &= \left(\sum_{i=1}^n \frac{A_i}{\widehat{\pi}(\mathbf{X}_i)}\right)^{-1} \lim_{y \rightarrow \infty} \sum_{i=1}^n \frac{A_i I(Y_i \leq y)}{\widehat{\pi}(\mathbf{X}_i)} \\
&= \left(\sum_{i=1}^n \frac{A_i}{\widehat{\pi}(\mathbf{X}_i)}\right)^{-1} \left(\sum_{i=1}^n \frac{A_i}{\widehat{\pi}(\mathbf{X}_i)}\right) \\
&= 1
\end{aligned} \tag{C.1}$$

and

$$\begin{aligned}
\lim_{y \rightarrow \infty} \widehat{F}^{(0)}(y) &= \left(\sum_{i=1}^n \frac{1 - A_i}{1 - \widehat{\pi}(\mathbf{X}_i)}\right)^{-1} \lim_{y \rightarrow \infty} \sum_{i=1}^n \frac{(1 - A_i) I(Y_i \leq y)}{1 - \widehat{\pi}(\mathbf{X}_i)} \\
&= \left(\sum_{i=1}^n \frac{1 - A_i}{1 - \widehat{\pi}(\mathbf{X}_i)}\right)^{-1} \left(\sum_{i=1}^n \frac{(1 - A_i)}{1 - \widehat{\pi}(\mathbf{X}_i)}\right) \\
&= 1.
\end{aligned} \tag{C.2}$$

Similarly, taking the limit $y \rightarrow -\infty$ to (8) and (9) gives that

$$\begin{aligned} \lim_{y \rightarrow -\infty} \widehat{F}^{(1)}(y) &= \left(\sum_{i=1}^n \frac{A_i}{\widehat{\pi}(\mathbf{X}_i)} \right)^{-1} \lim_{y \rightarrow -\infty} \sum_{i=1}^n \frac{A_i I(Y_i \leq y)}{\widehat{\pi}(\mathbf{X}_i)} \\ &= \left(\sum_{i=1}^n \frac{A_i}{\widehat{\pi}(\mathbf{X}_i)} \right)^{-1} \times 0 \\ &= 0 \end{aligned} \tag{C.3}$$

and

$$\begin{aligned} \lim_{y \rightarrow -\infty} \widehat{F}^{(0)}(y) &= \left(\sum_{i=1}^n \frac{1 - A_i}{1 - \widehat{\pi}(\mathbf{X}_i)} \right)^{-1} \lim_{y \rightarrow -\infty} \sum_{i=1}^n \frac{(1 - A_i) I(Y_i \leq y)}{1 - \widehat{\pi}(\mathbf{X}_i)} \\ &= \left(\sum_{i=1}^n \frac{1 - A_i}{1 - \widehat{\pi}(\mathbf{X}_i)} \right)^{-1} \times 0 \\ &= 0. \end{aligned} \tag{C.4}$$

Finally, we check the non-decreasing property of (8) and (9). We denote the ordered sample of $\{Y_1, Y_2, \dots, Y_n\}$ as $\{Y_{(1)}, Y_{(2)}, \dots, Y_{(n)}\}$ with $Y_{(i)} \leq Y_{(j)}$ for any $i < j$. For $i = 1, 2, \dots, n-1$, define interval $\mathbf{I}_i \triangleq (Y_{(i)}, Y_{(i+1)}]$ with $\mathbf{I}_0 \triangleq (-\infty, Y_{(1)}]$ and $\mathbf{I}_n \triangleq (Y_{(n)}, \infty)$. For any $a < b$, let

$$a^* \triangleq \sup_m \left\{ m : a \in \bigcup_{i=0}^m \mathbf{I}_i \right\} \text{ and } b^* \triangleq \sup_m \left\{ m : b \in \bigcup_{i=0}^m \mathbf{I}_i \right\}.$$

As $\left\{ m : a \in \bigcup_{i=0}^m \mathbf{I}_i \right\} \subseteq \left\{ m : b \in \bigcup_{i=0}^m \mathbf{I}_i \right\}$, we have $a^* \leq b^*$. Therefore,

$$\begin{aligned} \widehat{F}^{(1)}(b) &= \left(\sum_{i=1}^n \frac{A_i}{\widehat{\pi}(\mathbf{X}_i)} \right)^{-1} \sum_{i=1}^n \frac{A_i I(Y_i \leq b)}{\widehat{\pi}(\mathbf{X}_i)} \\ &= \left(\sum_{i=1}^n \frac{A_i}{\widehat{\pi}(\mathbf{X}_i)} \right)^{-1} \sum_{i=1}^n \frac{A_i I(Y_{(i)} \leq b)}{\widehat{\pi}(\mathbf{X}_i)} \\ &= \left(\sum_{i=1}^n \frac{A_i}{\widehat{\pi}(\mathbf{X}_i)} \right)^{-1} \sum_{i=1}^{b^*} \frac{A_i I(Y_{(i)} \leq b)}{\widehat{\pi}(\mathbf{X}_i)} \\ &\geq \left(\sum_{i=1}^n \frac{A_i}{\widehat{\pi}(\mathbf{X}_i)} \right)^{-1} \sum_{i=1}^{a^*} \frac{A_i I(Y_{(i)} \leq b)}{\widehat{\pi}(\mathbf{X}_i)} \\ &= \left(\sum_{i=1}^n \frac{A_i}{\widehat{\pi}(\mathbf{X}_i)} \right)^{-1} \sum_{i=1}^{a^*} \frac{A_i I(Y_{(i)} \leq a)}{\widehat{\pi}(\mathbf{X}_i)} \\ &= \widehat{F}^{(1)}(a) \end{aligned} \tag{C.5}$$

and

$$\begin{aligned}
\widehat{F}^{(0)}(b) &= \left(\sum_{i=1}^n \frac{1 - A_i}{1 - \widehat{\pi}(\mathbf{X}_i)} \right)^{-1} \sum_{i=1}^n \frac{(1 - A_i)I(Y_i \leq b)}{1 - \widehat{\pi}(\mathbf{X}_i)} \\
&= \left(\sum_{i=1}^n \frac{1 - A_i}{1 - \widehat{\pi}(\mathbf{X}_i)} \right)^{-1} \sum_{i=1}^n \frac{(1 - A_i)I(Y_{(i)} \leq b)}{1 - \widehat{\pi}(\mathbf{X}_i)} \\
&= \left(\sum_{i=1}^n \frac{1 - A_i}{1 - \widehat{\pi}(\mathbf{X}_i)} \right)^{-1} \sum_{i=1}^{b^*} \frac{(1 - A_i)I(Y_{(i)} \leq b)}{\widehat{\pi}(\mathbf{X}_i)} \\
&\geq \left(\sum_{i=1}^n \frac{1 - A_i}{1 - \widehat{\pi}(\mathbf{X}_i)} \right)^{-1} \sum_{i=1}^{a^*} \frac{(1 - A_i)I(Y_{(i)} \leq b)}{\widehat{\pi}(\mathbf{X}_i)} \\
&= \left(\sum_{i=1}^n \frac{1 - A_i}{1 - \widehat{\pi}(\mathbf{X}_i)} \right)^{-1} \sum_{i=1}^{a^*} \frac{(1 - A_i)I(Y_{(i)} \leq a)}{\widehat{\pi}(\mathbf{X}_i)} \\
&= \widehat{F}^{(0)}(a),
\end{aligned} \tag{C.6}$$

where the third step in (C.5) and (C.6) is due to that $I(Y_{(i)} \leq b) = 0$ for $i = b^* + 1, b^* + 2, \dots, n$ and the fifth step in (C.5) and (C.6) is due to that $I(Y_{(i)} \leq a) = I(Y_{(i)} \leq b) = 1$ for $i = 1, 2, \dots, a^*$. Then, (C.5) and (C.6) imply that the CDFs are non-decreasing. Consequently, by (C.1) to (C.6), we verify that the proposed estimators (8) and (9) are proper CDFs.

D Proof of Theorem 3.1

We prove the variable selection consistency here. As Theorem 3.1 is a statement with respect to both \widehat{V} and \widehat{E} , we tackle the sparsity recoveries of \widehat{V} and \widehat{E} separately and eventually integrate the results to arrive at the desired theorem.

Part 1: Inference for \widehat{V}

We aim to show

$$P(\widehat{V} = V) \rightarrow 1 \text{ as } n \rightarrow \infty. \tag{D.1}$$

Let the true coefficient be $\boldsymbol{\beta} = \left(\boldsymbol{\beta}_V^\top, \boldsymbol{\beta}_{V^c}^\top \right)^\top$ where $\boldsymbol{\beta}_V$ is the vector comprising coefficients of informative confounders with cardinality $|\boldsymbol{\beta}_V| = k$ and $\boldsymbol{\beta}_{V^c}$ contains coefficients of non-informative confounders and hence is $\mathbf{0}_{(p-k)}$. Moreover, $\widehat{\boldsymbol{\beta}} = \left(\widehat{\boldsymbol{\beta}}_V^\top, \widehat{\boldsymbol{\beta}}_{V^c}^\top \right)^\top$ is the corresponding estimated version of $\boldsymbol{\beta}$. Let $\widehat{\mathbf{z}} = \left(\widehat{\mathbf{z}}_V^\top, \widehat{\mathbf{z}}_{V^c}^\top \right)^\top$ be the dual vector of $\widehat{\boldsymbol{\beta}}$, where the sub-vectors $\widehat{\mathbf{z}}_V^\top$ and $\widehat{\mathbf{z}}_{V^c}^\top$ are the dual vectors of $\widehat{\boldsymbol{\beta}}_V$ and $\widehat{\boldsymbol{\beta}}_{V^c}$, respectively, and the j th component is $\widehat{z}_j = \text{sign}(\widehat{\beta}_j)$ if $\widehat{\beta}_j \neq 0$ and $|z_j| \leq 1$ otherwise. Also, any pair of $(\widehat{\boldsymbol{\beta}}, \widehat{\mathbf{z}})$ satisfies

the zero sub-gradient condition:

$$-\frac{1}{n}\mathbf{X}^\top(\mathbf{Y} - \mathbf{X}\hat{\boldsymbol{\beta}}) + \lambda_1\hat{\mathbf{z}} = \mathbf{0}. \quad (\text{D.2})$$

According to Hastie et al. (2015), condition (A1) together with primal dual witness (PDW) procedure can be used to examine the sparsity of $\hat{\boldsymbol{\beta}}$. The PDW procedure is stated as follows:

1. Set $\hat{\boldsymbol{\beta}}_{V^c} = \mathbf{0}$.
2. Determine $\hat{\mathbf{z}}_{V^c}$ according to (D.2).
3. Verify whether or not

$$\|\hat{\mathbf{z}}_{V^c}\|_\infty < 1. \quad (\text{D.3})$$

As discussed in Lemma 11.2. of Hastie et al. (2015, p.307), if (D.3) is true, then the vector $(\hat{\boldsymbol{\beta}}_V, \mathbf{0})$ is the unique optimal solution to the lasso program, which guarantees (D.1). Therefore, the main focus of remaining proof is to show (D.3). Following the relationship in (D.2) with Θ fixed, we have that

$$\begin{aligned} \hat{\mathbf{z}} &= \frac{1}{n\lambda_1}\mathbf{X}(\mathbf{Y} - \mathbf{X}\hat{\boldsymbol{\beta}}) \\ &= \frac{1}{n\lambda_1}(\mathbf{X}^\top\mathbf{Y} - \mathbf{X}^\top\mathbf{X}\hat{\boldsymbol{\beta}}) \\ &= \frac{1}{n\lambda_1}\left\{\mathbf{X}^\top(\mathbf{X}\boldsymbol{\beta} + \mathbf{X}^{\circ 2}\text{vec}(\Theta) + \boldsymbol{\varepsilon}) - \mathbf{X}^\top\mathbf{X}\hat{\boldsymbol{\beta}}\right\} \\ &= \frac{1}{n\lambda_1}(\mathbf{X}^\top\mathbf{X}\boldsymbol{\beta} + \mathbf{X}^\top\mathbf{X}^{\circ 2}\text{vec}(\Theta) + \mathbf{X}^\top\boldsymbol{\varepsilon} - \mathbf{X}^\top\mathbf{X}\hat{\boldsymbol{\beta}}) \\ &= \frac{1}{n\lambda_1}\left\{(\mathbf{X}^\top\mathbf{X})(\boldsymbol{\beta} - \hat{\boldsymbol{\beta}}) + \mathbf{X}^\top\mathbf{X}^{\circ 2}\text{vec}(\Theta) + \mathbf{X}^\top\boldsymbol{\varepsilon}\right\}. \end{aligned} \quad (\text{D.4})$$

Since $\hat{\mathbf{z}}_{V^c}$ is the sub-vector of $\hat{\mathbf{z}}$, we write (D.4) in the block matrix form, so that $\hat{\mathbf{z}}_V$ and $\hat{\mathbf{z}}_{V^c}$ can be expressed as

$$\begin{bmatrix} \hat{\mathbf{z}}_V \\ \hat{\mathbf{z}}_{V^c} \end{bmatrix} = \frac{1}{n\lambda_1}\left\{\begin{bmatrix} \mathbf{X}_V^\top\mathbf{X}_V & \mathbf{X}_V^\top\mathbf{X}_{V^c} \\ \mathbf{X}_{V^c}^\top\mathbf{X}_V & \mathbf{X}_{V^c}^\top\mathbf{X}_{V^c} \end{bmatrix}\begin{bmatrix} \boldsymbol{\beta}_V - \hat{\boldsymbol{\beta}}_V \\ \mathbf{0}_{(p-k)} \end{bmatrix} + \begin{bmatrix} \mathbf{X}_V^\top\mathbf{X}^{\circ 2}\text{vec}(\Theta) \\ \mathbf{X}_{V^c}^\top\mathbf{X}^{\circ 2}\text{vec}(\Theta) \end{bmatrix} + \begin{bmatrix} \mathbf{X}_V^\top\boldsymbol{\varepsilon} \\ \mathbf{X}_{V^c}^\top\boldsymbol{\varepsilon} \end{bmatrix}\right\}.$$

It also implies that

$$\hat{\mathbf{z}}_V = \frac{1}{n\lambda_1}\left(\mathbf{X}_V^\top\mathbf{X}_V(\boldsymbol{\beta}_V - \hat{\boldsymbol{\beta}}_V) + \mathbf{X}_V^\top\mathbf{X}^{\circ 2}\text{vec}(\Theta) + \mathbf{X}_V^\top\boldsymbol{\varepsilon}\right) \quad (\text{D.5})$$

and

$$\widehat{\mathbf{z}}_{V^c} = \frac{1}{n\lambda_1} \left(\mathbf{X}_{V^c}^\top \mathbf{X}_V (\boldsymbol{\beta}_V - \widehat{\boldsymbol{\beta}}_V) + \mathbf{X}_{V^c}^\top \mathbf{X}^{\circ 2} \text{vec}(\boldsymbol{\Theta}) + \mathbf{X}_{V^c}^\top \boldsymbol{\varepsilon} \right). \quad (\text{D.6})$$

By (D.5) and Condition (A1), the difference $\boldsymbol{\beta}_V - \widehat{\boldsymbol{\beta}}_V$ can be expressed as

$$\boldsymbol{\beta}_V - \widehat{\boldsymbol{\beta}}_V = n\lambda_1 (\mathbf{X}_V^\top \mathbf{X}_V)^{-1} \text{sign}(\boldsymbol{\beta}_V) - (\mathbf{X}_V^\top \mathbf{X}_V)^{-1} \mathbf{X}_V \mathbf{X}^{\circ 2} \text{vec}(\boldsymbol{\Theta}) - (\mathbf{X}_V^\top \mathbf{X}_V)^{-1} \mathbf{X}_V^\top \boldsymbol{\varepsilon}. \quad (\text{D.7})$$

Replacing the $(\boldsymbol{\beta}_V - \widehat{\boldsymbol{\beta}}_V)$ in (D.6) by (D.7) yields that

$$\begin{aligned} \widehat{\mathbf{z}}_{V^c} &= \mathbf{X}_{V^c}^\top \mathbf{X}_V (\mathbf{X}_V^\top \mathbf{X}_V)^{-1} \text{sign}(\boldsymbol{\beta}_V) - \frac{\mathbf{X}_{V^c}^\top \mathbf{X}_V (\mathbf{X}_V^\top \mathbf{X}_V)^{-1} \mathbf{X}_V \mathbf{X}^{\circ 2} \text{vec}(\boldsymbol{\Theta})}{n\lambda_1} + \frac{\mathbf{X}_{V^c}^\top \mathbf{X}^{\circ 2} \text{vec}(\boldsymbol{\Theta})}{n\lambda_1} \\ &\quad - \frac{\mathbf{X}_{V^c}^\top \mathbf{X}_V (\mathbf{X}_V^\top \mathbf{X}_V)^{-1} \mathbf{X}_V^\top \boldsymbol{\varepsilon}}{n\lambda_1} + \frac{\mathbf{X}_{V^c}^\top \boldsymbol{\varepsilon}}{n\lambda_1} \\ &= \mathbf{X}_{V^c}^\top \mathbf{X}_V (\mathbf{X}_V^\top \mathbf{X}_V)^{-1} \text{sign}(\boldsymbol{\beta}_V) + \mathbf{X}_{V^c}^\top [\mathbf{I} - \mathbf{X}_V (\mathbf{X}_V^\top \mathbf{X}_V)^{-1} \mathbf{X}_V] \left(\frac{\mathbf{X}^{\circ 2} \text{vec}(\boldsymbol{\Theta})}{n\lambda_1} \right) \\ &\quad + \mathbf{X}_{V^c}^\top [\mathbf{I} - \mathbf{X}_V (\mathbf{X}_V^\top \mathbf{X}_V)^{-1} \mathbf{X}_V] \left(\frac{\boldsymbol{\varepsilon}}{n\lambda_1} \right) \\ &= \mathbf{X}_{V^c}^\top \mathbf{X}_V (\mathbf{X}_V^\top \mathbf{X}_V)^{-1} \text{sign}(\boldsymbol{\beta}_V) + \mathbf{X}_{V^c}^\top [\mathbf{I} - \mathbf{X}_V (\mathbf{X}_V^\top \mathbf{X}_V)^{-1} \mathbf{X}_V] \left(\frac{\mathbf{X}^{\circ 2} \text{vec}(\boldsymbol{\Theta}) + \boldsymbol{\varepsilon}}{n\lambda_1} \right) \\ &\triangleq \boldsymbol{\mu} + \boldsymbol{\Psi}. \end{aligned}$$

By triangle inequality,

$$\|\widehat{\mathbf{z}}_{V^c}\|_\infty \leq \|\boldsymbol{\mu}\|_\infty + \|\boldsymbol{\Psi}\|_\infty. \quad (\text{D.8})$$

With regularity condition (A2), $\|\boldsymbol{\mu}\|_\infty \leq 1 - \gamma_1$. Hence, we only have to show that $\|\boldsymbol{\Psi}\|_\infty \leq \gamma_1$ with high probability. Let $\Pi_{V^\perp}(\mathbf{X}) \triangleq \mathbf{I} - \mathbf{X}_V (\mathbf{X}_V^\top \mathbf{X}_V)^{-1} \mathbf{X}_V$ be an $n \times n$ orthogonal projection matrix, then one can show that for $r \in V^c$,

$$\boldsymbol{\Psi}_r = \mathbf{X}_r^\top \Pi_{V^\perp}(\mathbf{X}) \left(\frac{\mathbf{X}^{\circ 2} \text{vec}(\boldsymbol{\Theta}) + \boldsymbol{\varepsilon}}{n\lambda_1} \right),$$

which is the r th component in $\boldsymbol{\Psi}$, is a Gaussian random variable with mean $\xi \triangleq \mathbf{X}^{\circ 2} \text{vec}(\boldsymbol{\Theta})$ and variance at most $\frac{\sigma^2 K_{\text{clm}}^2}{n\lambda_1^2}$. Without loss of generality, let $\boldsymbol{\Psi}_{\max}$ be a normal random variable with mean $\mathbf{X}^{\circ 2} \text{vec}(\boldsymbol{\Theta})$ and possibly maximal variance $\frac{\sigma^2 K_{\text{clm}}^2}{n\lambda_1^2}$. That is, for any $r \in \{1, \dots, p\}$, $\text{Var}(\boldsymbol{\Psi}_{\max}) \geq \text{Var}(\boldsymbol{\Psi}_r)$. Applying Lemma B.1 gives that

$$P\left(|\boldsymbol{\Psi}_{\max}| \geq \gamma_1\right) \leq 2 \exp\left(-\frac{(\gamma_1 - \xi)^2 n\lambda_1^2}{2\sigma^2 K_{\text{clm}}^2}\right). \quad (\text{D.9})$$

The upper bound in (D.9) approaches to 0 as $n \rightarrow \infty$, which implies

$$P\left(|\Psi_{\max}| \leq \gamma_1\right) \geq 1 \text{ as } n \rightarrow \infty.$$

Introducing the union bound further yields that

$$P\left(\|\Psi\|_\infty \geq \gamma_1\right) \leq 2(p-k)\exp\left(-\frac{(\gamma_1 - \xi)^2 n \lambda_1^2}{2\sigma^2 K_{\text{clm}}^2}\right), \quad (\text{D.10})$$

ensuring that $\|\Psi\|_\infty \leq \gamma_1$ as $n \rightarrow \infty$. Combining $\|\mu\|_\infty \leq 1 - \gamma_1$ from regularity condition (A2) and $\|\Psi\|_\infty$ with (D.8) gives (D.3), ensuring that the sparsity recovery is guaranteed for \widehat{V} .

Part 2: *Inference for \widehat{E}*

In this part, our goal is to show

$$P(\widehat{E} = E) \rightarrow 1 \text{ as } n \rightarrow \infty.$$

The procedure in Part 1 is repeated with the target of inference changed to Θ . Let the vector of true coefficients be $\text{vec}(\Theta) = \left(\text{vec}(\Theta_E)^\top, \text{vec}(\Theta_{EC})^\top\right)^\top$ where $\text{vec}(\Theta_E)$ is a m -dimensional nonzero vector that contains the coefficients of informative interaction terms and $\text{vec}(\Theta_{EC})$ consists of those of non-informative interaction terms and is equal to $\mathbf{0}_{\binom{p}{2}-m}$. Moreover, $\text{vec}(\widehat{\Theta}) = \left(\text{vec}(\widehat{\Theta}_E)^\top, \text{vec}(\widehat{\Theta}_{EC})^\top\right)^\top$ is the corresponding estimated version of $\text{vec}(\Theta)$. Let $\widehat{\mathbf{z}}' = \left(\widehat{\mathbf{z}}'_E, \widehat{\mathbf{z}}'_{EC}\right)$ be the dual vector of $\text{vec}(\widehat{\Theta})$, where the sub-vectors $\widehat{\mathbf{z}}'_E$ and $\widehat{\mathbf{z}}'_{EC}$ are the dual vectors of $\text{vec}(\widehat{\Theta}_E)$ and $\text{vec}(\widehat{\Theta}_{EC})$, respectively, and the j th component is $\widehat{z}'_j = \text{sign}(\widehat{\theta}_j)$ if $\widehat{\theta}_j \neq 0$ and $|\widehat{z}'_j| \leq 1$ otherwise. The zero sub-gradient condition with respect to $\text{vec}(\Theta)$ is

$$-\frac{1}{n}(\mathbf{X}^{\circ 2})^\top(\mathbf{Y} - \mathbf{X}^{\circ 2}\text{vec}(\widehat{\Theta})) + \lambda_2 \widehat{\mathbf{z}}' = \mathbf{0}.$$

Similar to the procedure in Part 1, the condition

$$\|\widehat{\mathbf{z}}'_{EC}\|_\infty \leq 1 \quad (\text{D.11})$$

is sufficient to attain the sparsity for \widehat{E} and therefore is of primary attention in this part.

With the similar procedure shown in Part 1, we express the dual vector as

$$\widehat{\mathbf{z}}'_{EC} = \mathbf{X}_{EC}^{\circ 2\top} \mathbf{X}_E^{\circ 2} (\mathbf{X}_E^{\circ 2\top} \mathbf{X}_E^{\circ 2})^{-1} \text{sign}(\text{vec}(\Theta_E)) + \mathbf{X}_{EC}^{\circ 2\top} [\mathbf{I} - \mathbf{X}_E^{\circ 2} (\mathbf{X}_E^{\circ 2\top} \mathbf{X}_E^{\circ 2})^{-1} \mathbf{X}_E^{\circ 2}] \left(\frac{\mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}}{n\lambda_2} \right).$$

In a similar manner with Part 1, we can apply the triangle inequality to this vector. Then, by utilizing Condition (A2) and Lemma B.1, one can show that (D.11) holds with high probability with $n \rightarrow \infty$. Therefore, the sparsity consistency for \widehat{E} is assured as $n \rightarrow \infty$.

Part 3: Inference for \widehat{G}

As $G = (V, E)$, the statement $\widehat{G} = G$ can be viewed as $\widehat{V} = V$ and $\widehat{E} = E$. On one hand, By Bonferroni's inequality, we have

$$P(\widehat{G} = G) = P(\widehat{V} = V, \widehat{E} = E) \geq P(\widehat{V} = V) + P(\widehat{E} = E) - 1. \quad (\text{D.12})$$

On the other hand, the inclusion relationship between events implies that

$$P(\widehat{G} = G) \leq \min \left\{ P(\widehat{V} = V), P(\widehat{E} = E) \right\}. \quad (\text{D.13})$$

Combining (D.12) and (D.13) yields that (e.g. Chen and Yi, 2021)

$$P(\widehat{V} = V) + P(\widehat{E} = E) - 1 \leq P(\widehat{G} = G) \leq \min \left\{ P(\widehat{V} = V), P(\widehat{E} = E) \right\}. \quad (\text{D.14})$$

By the squeeze theorem, we have $P(\widehat{G} = G)$ as $n \rightarrow \infty$. Consequently, the sparsity for G is verified. \square

E Proof of Theorem 3.2

In this section, we consider the limiting distribution where the underlying network structure G is correctly specified by previous variable selection procedure. A new design matrix of size $n \times (1 + k + m)$ is defined as $\mathbf{X}^* = \begin{bmatrix} \mathbf{1} & \mathbf{X}_{\widehat{V}} & \mathbf{X}_{\widehat{E}} \end{bmatrix}$ and the submatrices $\mathbf{X}_{\widehat{V}}$ and $\mathbf{X}_{\widehat{E}}$ consist of the chosen main and interaction effects, respectively. To derive the asymptotic results for our estimator, we first expand our estimator in terms of Hadamard derivative to determine the dominant term. Secondly, as the dominant term consists of a sequence of estimation in different parameters, we adopt M-estimation (Stefanski and Boos, 2002) to investigate the asymptotic result of the dominant term.

Part 1: Expansion and Determining the Dominant Term

Let $U^{(a)} = F^{(a)} \circ (F^{(a)})^{-1}$, which follows the uniform distribution function as defined in

(B.2), and $\widehat{U}^{(a)} = \widehat{F}^{(a)} \circ (F^{(a)})^{-1}$ for $a = 0, 1$. By the similar derivation of (B.4) in Lemma B.2, we have that

$$\begin{aligned}
& \sqrt{n} \left\{ T(\widehat{F}^{(1)}) - T(F^{(1)}) \right\} \\
&= \sqrt{n} \rho'_U(\widehat{U}^{(1)} - U^{(1)}) + \sqrt{n} \text{Rem}(\widehat{U}^{(1)} - U^{(1)}) \\
&= \sqrt{n} \left[\rho'_U \left((\widehat{F}^{(1)} - F^{(1)}) \circ \{F^{(1)}\}^{-1} \right) \right] + \sqrt{n} \text{Rem}(\widehat{U}^{(1)} - U^{(1)}) \\
&= \sqrt{n} \left[\rho'_U \left(\left(\sum_{i=1}^n \frac{A_i}{\widehat{\pi}(\mathbf{X}_i^*)} \right)^{-1} \sum_{i=1}^n \left(\frac{A_i \delta_{Y_i}}{\widehat{\pi}(\mathbf{X}_i^*)} - F^{(1)} \right) \circ \{F^{(1)}\}^{-1} \right) \right] + \sqrt{n} \text{Rem}(\widehat{U}^{(1)} - U^{(1)}) \\
&= \sqrt{n} \left(\sum_{i=1}^n \frac{A_i}{\widehat{\pi}(\mathbf{X}_i^*)} \right)^{-1} \left(\sum_{i=1}^n \frac{A_i \phi_{F^{(1)}}(Y_i)}{\widehat{\pi}(\mathbf{X}_i^*)} \right) + \sqrt{n} \text{Rem}(\widehat{U}^{(1)} - U^{(1)}).
\end{aligned} \tag{E.1}$$

Similar derivation for $\sqrt{n} \left\{ T(\widehat{F}^{(0)}) - T(F^{(0)}) \right\}$ gives that

$$\begin{aligned}
\sqrt{n} \left\{ T(\widehat{F}^{(0)}) - T(F^{(0)}) \right\} &= \sqrt{n} \left(\sum_{i=1}^n \frac{1 - A_i}{1 - \widehat{\pi}(\mathbf{X}_i^*)} \right)^{-1} \left(\sum_{i=1}^n \frac{(1 - A_i) \phi_{F^{(0)}}(Y_i)}{1 - \widehat{\pi}(\mathbf{X}_i^*)} \right) \\
&\quad + \sqrt{n} \text{Rem}(\widehat{U}^{(0)} - U^{(0)}).
\end{aligned} \tag{E.2}$$

Combining (E.1) and (E.2) leads to that

$$\begin{aligned}
& \sqrt{n} \left[\left(T(\widehat{F}^{(1)}) - T(\widehat{F}^{(0)}) \right) - \left(T(F^{(1)}) - T(F^{(0)}) \right) \right] \\
&= \sqrt{n} \left(\sum_{i=1}^n \frac{A_i}{\widehat{\pi}(\mathbf{X}_i^*)} \right)^{-1} \left(\sum_{i=1}^n \frac{A_i \phi_{F^{(1)}}(Y_i)}{\widehat{\pi}(\mathbf{X}_i^*)} \right) \\
&\quad - \sqrt{n} \left(\sum_{i=1}^n \frac{1 - A_i}{1 - \widehat{\pi}(\mathbf{X}_i^*)} \right)^{-1} \left(\sum_{i=1}^n \frac{(1 - A_i) \phi_{F^{(0)}}(Y_i)}{1 - \widehat{\pi}(\mathbf{X}_i^*)} \right) \\
&\quad + \sqrt{n} \left(\text{Rem}(\widehat{U}^{(1)} - U^{(1)}) - \text{Rem}(\widehat{U}^{(0)} - U^{(0)}) \right).
\end{aligned} \tag{E.3}$$

Since $\text{Rem}(\widehat{U}^{(a)} - U^{(a)})$ in (E.3) converges to 0 in probability as $n \rightarrow \infty$ for $a = 0, 1$ (Fernholz, 1983), it suffices to analyze

$$\sqrt{n} \left[\left(\sum_{i=1}^n \frac{A_i}{\widehat{\pi}(\mathbf{X}_i)} \right)^{-1} \sum_{i=1}^n \frac{A_i \phi_{F^{(1)}}(Y_i)}{\widehat{\pi}(\mathbf{X}_i)} - \left(\sum_{i=1}^n \frac{1 - A_i}{1 - \widehat{\pi}(\mathbf{X}_i)} \right)^{-1} \sum_{i=1}^n \frac{(1 - A_i) \phi_{F^{(0)}}(Y_i)}{1 - \widehat{\pi}(\mathbf{X}_i)} \right]. \tag{E.4}$$

We employ M-estimation to further investigate the asymptotic variance of (E.4).

Part 2: Estimating Equations Construction

First, (14) indicates that $\hat{\boldsymbol{\eta}}$ must satisfy the following equation

$$\sum_{i=1}^n \psi_{\boldsymbol{\eta}}(A_i, \mathbf{X}_i^* | \hat{\boldsymbol{\eta}}) \triangleq \sum_{i=1}^n \left[A_i - \pi(\mathbf{X}_i^* | \hat{\boldsymbol{\eta}}) \right] \mathbf{X}_i^{*\top} = \mathbf{0}_{1+k+m}. \quad (\text{E.5})$$

Next, we consider the estimating equation for IPW-CDF under $a \in \{0, 1\}$, where are given by

$$\psi_1(Y_i, \mathbf{X}_i^*, A_i | \boldsymbol{\eta}, \kappa_1) \triangleq \frac{A_i \left(\phi_{F^{(1)}}(Y_i) - \kappa_1 \right)}{\pi(\mathbf{X}_i^* | \boldsymbol{\eta})} \quad (\text{E.6})$$

and

$$\psi_0(Y_i, \mathbf{X}_i^*, A_i | \boldsymbol{\eta}, \kappa_0) \triangleq \frac{(1 - A_i) \left(\phi_{F^{(0)}}(Y_i) - \kappa_0 \right)}{1 - \pi(\mathbf{X}_i^* | \boldsymbol{\eta})}, \quad (\text{E.7})$$

where κ_1 and κ_0 are two parameters and are used to ensure that $E[\psi_1(Y, \mathbf{X}^*, A | \boldsymbol{\eta}, \kappa_1)] = 0$ and $E[\psi_0(Y, \mathbf{X}^*, A | \boldsymbol{\eta}, \kappa_0)] = 0$. Moreover, the solution to $\sum_{i=1}^n \psi_1(Y_i, \mathbf{X}_i^*, A_i | \boldsymbol{\eta}, \kappa_1) = 0$ and $\sum_{i=1}^n \psi_0(Y_i, \mathbf{X}_i^*, A_i | \boldsymbol{\eta}, \kappa_0) = 0$ are denoted as $\hat{\kappa}_1$ and $\hat{\kappa}_0$, respectively. One may check that

$$\hat{\kappa}_1 = \left(\sum_{i=1}^n \frac{A_i}{\hat{\pi}(\mathbf{X}_i^*)} \right)^{-1} \left(\sum_{i=1}^n \frac{A_i \phi_{F^{(1)}}(Y_i)}{\hat{\pi}(\mathbf{X}_i^*)} \right)$$

and

$$\hat{\kappa}_0 = \left(\sum_{i=1}^n \frac{1 - A_i}{1 - \hat{\pi}(\mathbf{X}_i^*)} \right)^{-1} \left(\sum_{i=1}^n \frac{(1 - A_i) \phi_{F^{(0)}}(Y_i)}{1 - \hat{\pi}(\mathbf{X}_i^*)} \right).$$

Part 3: *M-Estimation*

Following the procedure in Stefanski and Boos (2002) with the estimating equations (E.5)-(E.7) in Part 2, we define $\boldsymbol{\vartheta} \triangleq (\boldsymbol{\eta}^\top, \kappa_1, \kappa_0)^\top$ and

$$\boldsymbol{\psi}(Y_i, \mathbf{X}_i^*, A_i | \boldsymbol{\vartheta}) \triangleq \begin{bmatrix} \psi_{\boldsymbol{\eta}}(A_i, \mathbf{X}_i^* | \boldsymbol{\eta}) \\ \psi_1(Y_i, \mathbf{X}_i^*, A_i | \boldsymbol{\eta}, \kappa_1) \\ \psi_0(Y_i, \mathbf{X}_i^*, A_i | \boldsymbol{\eta}, \kappa_0) \end{bmatrix} = \begin{bmatrix} \left[A_i - \pi(\mathbf{X}_i^* | \boldsymbol{\eta}) \right] \mathbf{X}_i^{*\top} \\ \frac{A_i \left(\phi_{F^{(1)}}(Y_i) - \kappa_1 \right)}{\pi(\mathbf{X}_i^* | \boldsymbol{\eta})} \\ \frac{(1 - A_i) \left(\phi_{F^{(0)}}(Y_i) - \kappa_0 \right)}{1 - \pi(\mathbf{X}_i^* | \boldsymbol{\eta})} \end{bmatrix}. \quad (\text{E.8})$$

Since $\hat{\boldsymbol{\vartheta}} \triangleq (\hat{\boldsymbol{\eta}}^\top, \hat{\kappa}_1, \hat{\kappa}_0)^\top$ is the solution of $\sum_{i=1}^n \boldsymbol{\psi}(Y_i, \mathbf{X}_i^*, A_i | \boldsymbol{\vartheta}) = \mathbf{0}_{3+k+m}$, then under suitable conditions (Stefanski and Boos, 2002), we have that

$$\sqrt{n}(\hat{\boldsymbol{\vartheta}} - \boldsymbol{\vartheta}) \xrightarrow{d} \text{N}(0, \mathbf{V}(\boldsymbol{\vartheta})) \text{ as } n \rightarrow \infty, \quad (\text{E.9})$$

where $\mathbf{V}(\boldsymbol{\vartheta}) = \mathbf{A}^{-1}(\boldsymbol{\vartheta})\mathbf{B}(\boldsymbol{\vartheta})\{\mathbf{A}^{-1}(\boldsymbol{\vartheta})\}^\top$ with the matrices

$$\begin{aligned} \mathbf{B}(\boldsymbol{\vartheta}) &= E \left[\boldsymbol{\psi}(Y_i, \mathbf{X}_i^*, A_i \mid \boldsymbol{\vartheta}) \left\{ \boldsymbol{\psi}(Y_i, \mathbf{X}_i^*, A_i \mid \boldsymbol{\vartheta}) \right\}^\top \right] \\ &= E \left[\begin{array}{c|cc} \psi_\boldsymbol{\eta} \psi_\boldsymbol{\eta}^\top & \psi_\boldsymbol{\eta} \psi_1 & \psi_\boldsymbol{\eta} \psi_0 \\ \hline \psi_1 \psi_\boldsymbol{\eta}^\top & \psi_1 \psi_1 & \psi_1 \psi_0 \\ \psi_0 \psi_\boldsymbol{\eta}^\top & \psi_0 \psi_1 & \psi_0 \psi_0 \end{array} \right] \\ &\triangleq \begin{bmatrix} \mathbf{b}_{11} & \mathbf{b}_{21}^\top \\ \mathbf{b}_{21} & \mathbf{b}_{22} \end{bmatrix} \end{aligned} \quad (\text{E.10})$$

and

$$\begin{aligned} \mathbf{A}(\boldsymbol{\vartheta}) &= -E \left[\frac{\partial}{\partial \boldsymbol{\vartheta}} \boldsymbol{\psi}(Y, \mathbf{X}^*, A \mid \boldsymbol{\vartheta}) \right] \\ &= -E \left[\begin{array}{ccc} \frac{\partial}{\partial \boldsymbol{\eta}} \boldsymbol{\psi}_\boldsymbol{\eta}(Y, \mathbf{X}^*, A \mid \boldsymbol{\eta}) & \mathbf{0}_{1+k+m} & \mathbf{0}_{1+k+m} \\ \frac{\partial}{\partial \boldsymbol{\eta}} \psi_1(Y, \mathbf{X}^*, A \mid \boldsymbol{\eta}, \kappa_1) & \frac{\partial}{\partial \kappa_1} \psi_1(Y, \mathbf{X}^*, A \mid \boldsymbol{\eta}, \kappa_1) & 0 \\ \frac{\partial}{\partial \boldsymbol{\eta}} \psi_0(Y, \mathbf{X}^*, A \mid \boldsymbol{\eta}, \kappa_0) & 0 & \frac{\partial}{\partial \kappa_0} \psi_0(Y, \mathbf{X}^*, A \mid \boldsymbol{\eta}, \kappa_0) \end{array} \right] \\ &= -E \left[\begin{array}{c|cc} \pi(\mathbf{X}^* \mid \boldsymbol{\eta}) \left(1 - \pi(\mathbf{X}^* \mid \boldsymbol{\eta}) \right) \mathbf{X}^{*\top} \mathbf{X}^* & \mathbf{0}_{1+k+m} & \mathbf{0}_{1+k+m} \\ \hline \frac{1 - \pi(\mathbf{X}^* \mid \boldsymbol{\eta})}{\pi(\mathbf{X}^* \mid \boldsymbol{\eta})} A \left(\phi_{F^{(1)}}(Y_i) - \kappa_1 \right) \mathbf{X}^* & -1 & 0 \\ -\frac{\pi(\mathbf{X}^* \mid \boldsymbol{\eta})}{1 - \pi(\mathbf{X}^* \mid \boldsymbol{\eta})} (1 - A) \left(\phi_{F^{(0)}}(Y_i) - \kappa_0 \right) \mathbf{X}^* & 0 & -1 \end{array} \right] \\ &\triangleq \begin{bmatrix} \mathbf{a}_{11} & \mathbf{0}_{(1+k+m) \times 2} \\ \mathbf{a}_{21} & \mathbf{I}_2 \end{bmatrix} \end{aligned} \quad (\text{E.11})$$

with the inverse matrix

$$\mathbf{A}^{-1}(\boldsymbol{\vartheta}) = \begin{bmatrix} \mathbf{a}_{11}^{-1} & \mathbf{0}_{(1+k+m) \times 2} \\ -\mathbf{a}_{11}^{-1} \mathbf{a}_{21} & \mathbf{I}_2 \end{bmatrix} \quad (\text{E.12})$$

and \mathbf{I}_2 being a 2×2 identity matrix. It suggests that $\mathbf{V}(\boldsymbol{\vartheta})$ can be expressed as

$$\begin{aligned}\mathbf{V}(\boldsymbol{\vartheta}) &= \mathbf{A}^{-1}(\boldsymbol{\vartheta})\mathbf{B}(\boldsymbol{\vartheta})\{\mathbf{A}^{-1}(\boldsymbol{\vartheta})\}^\top \\ &= \begin{bmatrix} \mathbf{a}_{11}^{-1} & \mathbf{0}_{(1+k+m) \times 2} \\ -\mathbf{a}_{11}^{-1}\mathbf{a}_{21} & \mathbf{I}_2 \end{bmatrix} \begin{bmatrix} \mathbf{a}_{11} & \mathbf{b}_{21}^\top \\ \mathbf{b}_{21} & \mathbf{b}_{22} \end{bmatrix} \begin{bmatrix} \mathbf{a}_{11}^{-1} & -\mathbf{a}_{21}^\top\mathbf{a}_{11}^{-1} \\ \mathbf{0}_{2 \times (1+k+m)} & \mathbf{I}_2 \end{bmatrix} \\ &= \begin{bmatrix} \mathbf{a}_{11}^{-1} & -\mathbf{a}_{11}^{-1}\mathbf{a}_{21}^\top - \mathbf{a}_{11}^{-1}\mathbf{b}_{21} \\ (-\mathbf{a}_{21} + \mathbf{b}_{21})\mathbf{a}_{11}^{-1} & (\mathbf{a}_{21} - \mathbf{b}_{21})\mathbf{a}_{11}^{-1}\mathbf{a}_{21}^\top - \mathbf{a}_{21}\mathbf{a}_{11}^{-1}\mathbf{b}_{21}^\top + \mathbf{b}_{22} \end{bmatrix}.\end{aligned}$$

Now we thoroughly investigate the components of the matrices (E.10) and (E.11). Firstly, by Lemma 7.3.11 of Casella and Berger (2002), we have that $\mathbf{a}_{11} = \mathbf{b}_{11} = E\left\{\pi(\mathbf{X}^* | \boldsymbol{\eta})\left(1 - \pi(\mathbf{X}^* | \boldsymbol{\eta})\right)\mathbf{X}^{*\top}\mathbf{X}^*\right\}$. Secondly,

$$\begin{aligned}\mathbf{a}_{21} &\triangleq \left[-E\left(\frac{\partial}{\partial \boldsymbol{\eta}}\psi_1\right) - E\left(\frac{\partial}{\partial \boldsymbol{\eta}}\psi_0\right)\right]^\top \\ &= \left[E\left\{\frac{1 - \pi(\mathbf{X}^* | \boldsymbol{\eta})}{\pi(\mathbf{X}^* | \boldsymbol{\eta})}A(\phi_{F^{(1)}}(Y) - \kappa_1)\mathbf{X}^*\right\} - E\left\{\frac{\pi(\mathbf{X}^* | \boldsymbol{\eta})}{1 - \pi(\mathbf{X}^* | \boldsymbol{\eta})}(1 - A)(\phi_{F^{(0)}}(Y) - \kappa_0)\mathbf{X}^*\right\}\right]^\top \\ &= \left[E\left\{\left(1 - \pi(\mathbf{X}^* | \boldsymbol{\eta})\right)\mathbf{X}^*(\phi_{F^{(1)}}(Y^{(1)}) - \kappa_1)\right\} - E\left\{\pi(\mathbf{X}^* | \boldsymbol{\eta})\mathbf{X}^*(\phi_{F^{(0)}}(Y^{(0)}) - \kappa_0)\right\}\right]^\top.\end{aligned}$$

For the component $\mathbf{b}_{21} \triangleq \left[E(\psi_1\psi_\boldsymbol{\eta}^\top) \ E(\psi_0\psi_\boldsymbol{\eta}^\top)\right]^\top$ in $\mathbf{B}(\boldsymbol{\vartheta})$, we observe that

$$\begin{aligned}E(\psi_1\psi_\boldsymbol{\eta}^\top) &= E\left[\frac{A(\phi_{F^{(1)}}(Y) - \kappa_1)}{\pi(\mathbf{X}^* | \boldsymbol{\eta})}\left\{A - \pi(\mathbf{X}^* | \boldsymbol{\eta})\mathbf{X}^*\right\}\right] \\ &= E\left\{\frac{A(\phi_{F^{(1)}}(Y) - \kappa_1)\mathbf{X}^*}{\pi(\mathbf{X}^* | \boldsymbol{\eta})} - A(\phi_{F^{(1)}}(Y) - \kappa_1)\mathbf{X}^*\right\} \\ &= E\left\{A(\phi_{F^{(1)}}(Y) - \kappa_1)\mathbf{X}^*\left(\frac{1}{\pi(\mathbf{X}^* | \boldsymbol{\eta})} - 1\right)\right\} \\ &= E\left\{A(\phi_{F^{(1)}}(Y) - \kappa_1)\mathbf{X}^*\left(\frac{1 - \pi(\mathbf{X}^* | \boldsymbol{\eta})}{\pi(\mathbf{X}^* | \boldsymbol{\eta})}\right)\right\} \\ &= E\left\{(\phi_{F^{(1)}}(Y^{(1)}) - \kappa_1)\mathbf{X}^*\left(1 - \pi(\mathbf{X}^* | \boldsymbol{\eta})\right)\right\} \\ &= -E\left(\frac{\partial}{\partial \boldsymbol{\eta}}\psi_1\right)\end{aligned}\tag{E.13}$$

and

$$E(\psi_0 \psi_{\boldsymbol{\eta}}^\top) = -E\left(\frac{\partial}{\partial \boldsymbol{\eta}} \psi_0\right)$$

derived by the similar step in (E.13). It suggests that $\mathbf{b}_{21} = \mathbf{a}_{21}$. Moreover,

$$\begin{aligned} \mathbf{b}_{22} &= E \begin{bmatrix} \psi_1 \psi_1 & \psi_1 \psi_0 \\ \psi_0 \psi_1 & \psi_0 \psi_0 \end{bmatrix} \\ &= \begin{bmatrix} E\left(\frac{A(\phi_{F^{(1)}}(Y) - \kappa_1)^2}{\pi(\mathbf{X}^* | \boldsymbol{\eta})^2}\right) & E\left\{\frac{A(1-A)(\phi_{F^{(1)}}(Y) - \kappa_1)(\phi_{F^{(0)}}(Y) - \kappa_0)}{\pi(\mathbf{X}^* | \boldsymbol{\eta})\{1 - \pi(\mathbf{X}^* | \boldsymbol{\eta})\}}\right\} \\ E\left\{\frac{A(1-A)(\phi_{F^{(1)}}(Y) - \kappa_1)(\phi_{F^{(0)}}(Y) - \kappa_0)}{\pi(\mathbf{X}^* | \boldsymbol{\eta})\{1 - \pi(\mathbf{X}^* | \boldsymbol{\eta})\}}\right\} & E\left[\frac{(1-A)(\phi(Y) - \kappa_0)^2}{\{1 - \pi(\mathbf{X}^* | \boldsymbol{\eta})\}^2}\right] \end{bmatrix} \\ &= \begin{bmatrix} E\left\{\frac{(\phi_{F^{(1)}}(Y^{(1)}) - \kappa_1)^2}{\pi(\mathbf{X}^* | \boldsymbol{\eta})}\right\} & 0 \\ 0 & E\left\{\frac{(\phi_{F^{(0)}}(Y^{(0)}) - \kappa_0)^2}{1 - \pi(\mathbf{X}^* | \boldsymbol{\eta})}\right\} \end{bmatrix}, \end{aligned}$$

where the non-diagonal terms are 0 since $A(1-A) = 0$.

By the property of the multivariate normal distribution, we immediately have that

$$\sqrt{n}(\widehat{\boldsymbol{\eta}} - \boldsymbol{\eta}) \xrightarrow{d} \text{N}(0, \mathcal{A}^{-1})$$

as $n \rightarrow \infty$, where $\mathcal{A} \triangleq \mathbf{a}_{11}$. This gives the result (i). On the other hand, by (E.9), we have that, as $n \rightarrow \infty$,

$$\sqrt{n} \left\{ \begin{pmatrix} \widehat{\kappa}_1 \\ \widehat{\kappa}_0 \end{pmatrix} - \begin{pmatrix} \kappa_1 \\ \kappa_0 \end{pmatrix} \right\} \xrightarrow{d} \text{N}(\mathbf{0}_2, \mathbf{v}_{22}), \quad (\text{E.14})$$

where

$$\begin{aligned} \mathbf{v}_{22} &\triangleq (\mathbf{a}_{21} - \mathbf{b}_{21})\mathbf{a}_{11}^{-1}\mathbf{a}_{21}^\top - \mathbf{a}_{21}\mathbf{a}_{11}^{-1}\mathbf{b}_{21}^\top + \mathbf{b}_{22} \\ &= \mathbf{a}_{21}\mathbf{a}_{11}^{-1}\mathbf{a}_{21}^\top - \mathbf{b}_{21}\mathbf{a}_{11}^{-1}\mathbf{a}_{21}^\top - \mathbf{a}_{21}\mathbf{a}_{11}^{-1}\mathbf{b}_{21}^\top + \mathbf{b}_{22} \\ &= \mathbf{a}_{21}\mathbf{a}_{11}^{-1}\mathbf{a}_{21}^\top - \mathbf{b}_{21}\mathbf{a}_{11}^{-1}\mathbf{a}_{21}^\top - \mathbf{a}_{21}\mathbf{a}_{11}^{-1}\mathbf{b}_{21}^\top + \mathbf{b}_{21}\mathbf{a}_{11}^{-1}\mathbf{b}_{21}^\top - \mathbf{b}_{21}\mathbf{a}_{11}^{-1}\mathbf{b}_{21}^\top + \mathbf{b}_{22} \\ &= (\mathbf{a}_{21} - \mathbf{b}_{21})\mathbf{a}_{11}^{-1}(\mathbf{a}_{21} - \mathbf{b}_{21})^\top - \mathbf{b}_{21}\mathbf{a}_{11}^{-1}\mathbf{b}_{21}^\top + \mathbf{b}_{22} \\ &= -\mathbf{b}_{21}\mathbf{a}_{11}^{-1}\mathbf{b}_{21}^\top + \mathbf{b}_{22}. \end{aligned}$$

Applying delta method to (E.14) with $\boldsymbol{\Delta} \triangleq (1, -1)^\top$, we have that, as $n \rightarrow \infty$,

$$\sqrt{n} \left[(\widehat{\kappa}_1 - \widehat{\kappa}_0) - (\kappa_1 - \kappa_0) \right] \xrightarrow{d} \text{N}\left(0, \mathcal{C} - \mathcal{B}\mathcal{A}^{-1}\mathcal{B}^\top\right), \quad (\text{E.15})$$

where $\mathcal{C} \triangleq \mathbf{\Delta}^\top \mathbf{b}_{22} \mathbf{\Delta}$ and $\mathcal{B} \triangleq \mathbf{\Delta}^\top \mathbf{b}_{21}$.

Recall that κ_1 and κ_0 are used to $E[\psi_1(Y, \mathbf{X}^*, A \mid \boldsymbol{\eta}, \kappa_1)] = 0$ and $E[\psi_0(Y, \mathbf{X}^*, A \mid \boldsymbol{\eta}, \kappa_0)] = 0$, respectively. Solving the first equation leads to

$$\begin{aligned}
\kappa_1 &= E \left\{ \frac{A \phi_{F^{(1)}}(Y)}{\pi(\mathbf{X}^*)} \right\} \\
&= E \left\{ \frac{1}{\pi(\mathbf{X}^*)} E \left(A \phi_{F^{(1)}}(Y) \mid \mathbf{X}^* \right) \right\} \\
&= E \left\{ \frac{E \left(1 \times \phi_{F^{(1)}}(Y^{(1)}) \mid \mathbf{X}^*, A = 1 \right) P(A = 1 \mid \mathbf{X}^*)}{\pi(\mathbf{X}^*)} \right. \\
&\quad \left. + \frac{E \left(0 \times \phi_{F^{(1)}}(Y^{(0)}) \mid \mathbf{X}^*, A = 0 \right) P(A = 0 \mid \mathbf{X}^*)}{\pi(\mathbf{X}^*)} \right\} \\
&= E \left\{ \frac{E \left(\phi_{F^{(1)}}(Y^{(1)}) \mid \mathbf{X}^*, A = 1 \right) \pi(\mathbf{X}^*)}{\pi(\mathbf{X}^*)} \right\} \\
&= E \left\{ E \left(\phi_{F^{(1)}}(Y^{(1)}) \mid \mathbf{X}^*, A = 1 \right) \right\} \\
&= E \left\{ E \left(\phi_{F^{(1)}}(Y^{(1)}) \mid \mathbf{X}^* \right) \right\} \\
&= E \left\{ \phi_{F^{(1)}}(Y^{(1)}) \right\} \\
&= 0,
\end{aligned}$$

where the last equality utilizes the property of influence curves (Hampel, 1974). Similar derivations yield that

$$\begin{aligned}
\kappa_0 &= E \left\{ \frac{(1-A) \phi_{F^{(0)}}(Y)}{1 - \pi(\mathbf{X}^*)} \right\} \\
&= 0.
\end{aligned}$$

Consequently, combining (E.3) with (E.15) gives that, as $n \rightarrow \infty$,

$$\sqrt{n} \left[\left(T(\widehat{F}^{(1)}) - T(F^{(0)}) \right) - \left(T(F^{(1)}) - T(F^{(0)}) \right) \right] \xrightarrow{d} \mathbf{N} \left(0, -\mathcal{B} \mathcal{A}^{-1} \mathcal{B}^\top + \mathcal{C} \right),$$

which gives the result (ii).

References

- Casella, G. and Berger, R. L. (2002). *Statistical Inference, Second Edition*. Duxbury Pacific Grove, CA.
- Chen, L.-P. and Yi, G. Y. (2021). Analysis of noisy survival data with graphical proportional hazards measurement error model. *Biometrics*, 77, 956-96.
- Fernholz, L. T. (1983). *Von Mises Calculus for Statistical Functionals*. Springer Science & Business Media.
- Hampel, F. R. (1974). The influence curve and its role in robust estimation. *Journal of the American Statistical Association*, 69(346), 383-393.
- Stefanski, L. A. and Boos, D. D. (2002). The calculus of M-estimation. *The American Statistician*, 56, 29-38.
- Serfling, R.J. (1980). *Approximation Theorems of Mathematical Statistics*. John Wiley & Sons, New York.
- Hastie, T., Tibshirani, R. and Wainwright, M. (2015). *Statistical Learning with Sparsity: The Lasso and Generalizations*. CRC press.
- Lunceford, J. K. and Davidian, M. (2004). Stratification and weighting via the propensity score in estimation of causal treatment effects: a comparative study. *Statistics in Medicine*, 23, 2937-2960
- van der Vaart, A.W. and Wellner, J.A. (1996). *Weak Convergence and Empirical Processes: With Application to Statistics*. Springer-Verlag, New York.
- Yang, E., Ravikumar, P., Allen, G. I. and Liu, Z. (2015). Graphical models via univariate exponential family distributions. *The Journal of Machine Learning Research*, 16, 3813-3847.