

Length-Constrained Directed Expander Decomposition and Length-Constrained Vertex-Capacitated Flow Shortcuts

Bernhard Haeupler*

bernhard.haeupler@inf.ethz.ch

INSAIT, Sofia University "St. Kliment Ohridski" & ETH Zürich

Thatchaphol Saranurak[‡]

thsa@umich.edu

University of Michigan

Yaowei Long[†]

yaoweil@umich.edu

University of Michigan

Shengzhe Wang

shengzhe.wang@inf.ethz.ch

ETH Zürich

Abstract

We show the existence of *length-constrained expander decomposition* in directed graphs and undirected vertex-capacitated graphs. Previously, its existence was shown only in undirected edge-capacitated graphs [HRG22, HHT24]. Along the way, we prove the multi-commodity maxflow-mincut theorems for length-constrained expansion in both directed and undirected vertex-capacitated graphs.

Based on our decomposition, we build a *length-constrained flow shortcut* for undirected vertex-capacitated graphs, which roughly speaking is a set of edges and vertices added to the graph so that every multi-commodity flow demand can be routed with approximately the same vertex-congestion *and* length, but all flow paths only contain few edges. This generalizes the shortcut for undirected edge-capacitated graphs from [HHL⁺24].

Length-constrained expander decomposition and flow shortcuts have been crucial in the recent algorithms in undirected edge-capacitated graphs [HHL⁺24, HLS24]. Our work thus serves as a foundation to generalize these concepts to directed and vertex-capacitated graphs.

*Partially funded by the Ministry of Education and Science of Bulgaria's support for INSAIT as part of the Bulgarian National Roadmap for Research Infrastructure and through the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (ERC grant agreement 949272).

[†]Part of this work was done while at INSAIT, Sofia University "St. Kliment Ohridski", Bulgaria. This work was partially funded from the Ministry of Education and Science of Bulgaria (support for INSAIT, part of the Bulgarian National Roadmap for Research Infrastructure).

[‡]Supported by NSF Grant CCF-2238138. Part of this work was done while at INSAIT, Sofia University "St. Kliment Ohridski", Bulgaria. This work was partially funded from the Ministry of Education and Science of Bulgaria (support for INSAIT, part of the Bulgarian National Roadmap for Research Infrastructure).

Contents

1	Introduction	1
1.1	Our Results	2
1.2	Our Techniques	3
2	Preliminaries	7
2.1	Directed Graphs	7
2.2	Vertex-Capacitated Graphs	8
3	Length-Constrained Directed Expansion	9
3.1	Basic Concepts of Length-Constrained Directed Expansion	9
3.2	Routing Characterization of Length-Constrained Directed Expansion	10
3.3	Length-Constrained Directed Expander Decomposition: Existence	10
3.3.1	Exponential Demand	11
3.3.2	Existential Proof of the Decomposition	16
4	Length-Constrained Vertex Expansion	19
4.1	Basic Concepts of Length-Constrained Vertex Expansion	19
4.2	Reduction to Directed Edge-Capacitated Graphs	20
4.3	Existential Proof of the Decomposition	24
4.4	Routing Characterization of Length-Constrained Vertex Expansion	25
5	Length-Constrained Vertex-Capacitated Flow Shortcuts	27
5.1	The Construction	27
5.2	Forward Mapping: Proof of Lemma 5.5	29
5.3	Backward Mapping: Proof of Lemma 5.6	31
A	Length-Constrained Directed Expansion (Continued)	37
A.1	Connection to Classic Directed Expander Decomposition	37
A.2	Routing Characterization: Proof of Theorem 3.8	38
A.3	Linkedness	41

1 Introduction

Expander decomposition found its early applications in property testing [GR98], clustering [KVV04], and approximation algorithms [CKS05] and, for the last two decades, has been the crucial ingredient in important developments of fast graph algorithms. This includes the first almost-linear time algorithms for spectral sparsifiers and Laplacian solvers [ST04], approximate max flow [KLOS14, She13, RST14], deterministic global min cut [LS21], exact max flow [CKL⁺22], as well as many almost-optimal dynamic algorithms for minimum spanning trees [NSWN17], shortest paths [CK19, BGS22], sparsifiers [BBG⁺20], k -edge-connectivity [JS22], minimum cuts [JST24, EHHL25], and more [GRST21]. Significant effort [SW19, CS19, CGL⁺20, CS20, LS21, LNPS23, HKGW23, ADK23, GPPG24, CMGS25] has then focused on constructing expander decomposition itself.

Below, we discuss two successful orthogonal generalizations of expander decomposition.

Vertex and Directed Expander Decomposition. In 2005, Chekuri, Khanna, and Shepherd [CKS05] showed that the construction of expander decomposition in undirected edge-capacitated graphs naturally extends to work in undirected vertex-capacitated graphs and applies them for approximating all-or-nothing vertex-capacitated flow problems. Later, this was extended to directed graphs, an even more general setting [CE15].¹

Since 2020, almost-linear time expander decomposition algorithms in these generalized settings have been developed [BGS20, LS22, HKGW23, SG24] and found impressive applications. For the vertex-capacitated ones, they were crucial for the fastest deterministic vertex connectivity algorithms [SY22, NSY23] and data structures for connectivity queries under vertex failures [LS22, LPS25, JPP25]. For the directed ones, they were used for dynamic algorithms in directed graphs [BGS20] and the new combinatorial approaches for exact max flow [CK24, BBST24].

Length-Constrained Expander Decomposition and Flow Shortcuts. More recently, Haeupler, Räcke, and Ghaffari [HRG22] introduced *length-constrained expanders* (LC-expanders). At a very high level, these are graphs such that any “reasonable” demand can be routed with low congestion *and* length. In contrast, normal expanders only guarantee low congestion. [HRG22] constructed LC-expander decomposition and applied it to show universally optimal distributed algorithms. In general, LC-expander decomposition is much more effective for problems that simultaneously concern length and congestion.

Based on the new decomposition, [HHL⁺24] introduced the notion of *LC-flow shortcut*², a new kind of graph augmentation. Roughly speaking, an LC-flow shortcut is a set of edges and vertices added to the graph so that every multi-commodity flow demand can be routed with approximately the same congestion *and* length, but all flow paths only have a few edges. This is formalized as follows (see Section 2 for background).

Definition 1.1 (Length-Constrained Flow Shortcut). *Given a graph $G = (V, E)$, we say an edge set E' (possibly with endpoints outside V) is a t -step flow shortcut of G with length slack λ and congestion slack κ if*

- (Forward Mapping) for every demand D routable in G with congestion 1 and length h , D is routable in $G \cup E'$ with congestion 1, length λh , and maximum step t , and
- (Backward Mapping) for every demand D on $V(G)$ routable in $G \cup E'$ with congestion 1 and length h , D is routable in G with congestion κ and length h .

¹In fact, expander decomposition was only implicit in [CKS05, CE15] as their definitions were specific to their applications. The purely graph-theoretic definition was later formalized in [BGS20].

²It was called a *low-step flow emulator* in [HHL⁺24].

In any undirected edge-capacitated graph, [HHL⁺24] showed, for any $\epsilon > 0$, the existence of a $O(1/\epsilon)$ -step LC-flow shortcut E' of size $|E'| \leq O(n^{1+O(\epsilon)} \text{polylog}(n))$ with length slack $O(1/\epsilon^3)$ and congestion slack $n^{O(\epsilon)}$.³ Combined with newly developed close-to-linear time LC-expander decomposition [HHT24, HHG25], they also obtained a close-to-linear time construction for LC-flow shortcuts albeit with worse quality.

LC-flow shortcuts have led to significant further progress. This includes the first close-to-linear time constant-approximation algorithm for minimum cost multi-commodity flow [HHL⁺24]. The dynamic but weaker version of flow shortcuts was also the key object in the first deterministic dynamic constant-approximate distance oracle with $O(n^\epsilon)$ update time [HLS24].

However, all applications of LC-expander decomposition until now are limited to undirected edge-capacitated graphs.

1.1 Our Results

To extend the reach of the expander decomposition paradigm further, the history above suggests the following research question:

Can we construct length-constrained expander decomposition and flow shortcuts beyond undirected edge-capacitated graphs?

Indeed, we answer this question affirmatively. In this paper, we focus on the existential results, but the arguments naturally give polynomial-time algorithms. For future work, we are working towards almost-linear-time constructions, which would lead to further applications for minimum cost (multi-commodity) flow in vertex-capacitated and directed graphs. Below, we discuss our contribution in more detail.

Length-Constrained Directed and Vertex Expander Decompositions. We formalize the notions of length-constrained expanders in directed graphs and in undirected vertex-capacitated graphs (Sections 3.1 and 4.1). Then, we show the existence of length-constrained expander decomposition in directed graphs (Theorem 3.9) and in undirected vertex-capacitated graphs (Theorem 4.1). Along the way, we also show that the definition of length-constrained expanders based on cuts is almost equivalent to the characterization based on multi-commodity flow (Theorems 3.8 and 4.2). This can be viewed as a version of the approximate multicommodity maxflow mincut theorem [LR99] but for length-constrained expansion in directed and vertex-capacitated graphs.

While this part does not require technical novelty, it is an important foundation for our paper and, we believe, for future work using this concept.

Length-Constrained Vertex-Capacitated Flow Shortcuts. Our main technical contribution (Theorem 5.2) is to show that, for any undirected *vertex-capacitated* graph and any $\epsilon > 0$, there exists a $2^{O(\frac{1}{\epsilon})}$ -step flow shortcut E' of size $|E'| = O(n^{1+O(\epsilon)} \text{polylog}(n))$ with length slack $O(1/\epsilon^3)$ and congestion slack $n^{O(\epsilon)}$. This generalizes the flow shortcut of [HHL⁺24] in undirected edge-capacitated graphs.

Our trade-off between size, length slack, and congestion slack matches the one of [HHL⁺24]. However, our step-bound is $2^{O(\frac{1}{\epsilon})}$ instead of $O(1/\epsilon^2)$. This is due to technical barriers unique to vertex-capacitated graphs, which also requires us to use very different analysis. We leave as a very interesting open problem if it is possible to obtain $\text{poly}(1/\epsilon)$ steps.

³The shortcut E' in [HHL⁺24] actually has $O(1/\epsilon^4)$ length slack and $O(1/\epsilon^2)$ maximum step, but this is only because they tried to ensure that all endpoints of E' are in V . Allowing endpoints outside V , one can replace their *router* with a star and improve the quality to be as we stated.

We note that obtaining similar LC-flow shortcuts on directed graphs is currently out of reach because it would give the breakthrough on *reachability shortcuts*. Given a graph $G = (V, E)$, an edge set E' is a t -step reachability shortcut of G if, for every pair of vertices $u, v \in V$, u can reach v in G if and only if u can reach v in $G \cup E'$ using at most t steps. Observe that an LC-flow shortcut in a directed graph is strictly stronger than a reachability shortcut. It is a major open problem whether there exists a $n^{o(1)}$ -step reachability shortcut of size $n^{1+o(1)}$.⁴

1.2 Our Techniques

Next, we give a technical overview of our LC-flow shortcut on vertex-capacitated graphs. We will explain how the strategy used in [HHL⁺24] fails in our setting and how we overcome the obstacle. For simplicity, here we only consider graphs with *unit capacity*. Also, we only construct a slightly weaker notion of LC-flow shortcut in the sense that, it receives an additional length parameter h and the forward mapping only guarantees that every demand routable in G with length $h' \leq h$ and congestion 1 is routable in $G \cup E'$ with length λh , congestion 1 and step t .

Preliminaries. First, we give a brief background on length-constrained expansion. A demand $D : V \times V \rightarrow \mathbb{R}_{\geq 0}$ assigns value to pair of vertices (u, v) and D is h -length if it assigns non-zero values only to vertex pairs of distance $\text{dist}_G(u, v) \leq h$. A demand D is *routable* with congestion κ and length λ if there exists a multi-commodity flow routing D with congestion κ and length λ . D *respects* a node-weighting $A : V \rightarrow \mathbb{R}_{\geq 0}$ if for each vertex u , $\sum_v D(u, v) \leq A(u)$. Let $|A| = \sum_{u \in V} A(u)$. For any $s \geq 1$, A is (h, s) -length ϕ -expanding in G if every h -length A -respecting demand is routable with length hs and congestion $O(\frac{\log n}{\phi})$.⁵ A length-constrained cut C , generally speaking, assigns to each edge an integral length increase, and $G - C$ is the graph G applied with the length increase from cut C . We informally say that G is a *length-constrained expander (LC-expander)* if a node-weighting A whose support is the whole vertex set V is expanding in G . An (h, s) -length ϕ -expander decomposition for A is a length-constrained cut C such that A is (h, s) -length ϕ -expanding in $G - C$.




Figure 1: An LC-flow shortcut of a low-diameter LC-expander.

Warm-up: Shortcutting LC-expanders. Before explaining the obstacle, we first show how to shortcut an LC-vertex expander as a warm-up. Suppose that a node-weighting A is (h, s) -length ϕ -vertex expanding in G . Say, $A := \mathbb{1}_V$ (i.e., $A(v) = 1$ for all $v \in V$).

⁴When the endpoints of E' must be in V , [Hes03, HP21, BH23] already showed that there is no $\Omega(n^{1/4})$ -step reachability shortcut of size $O(n)$. The lower bounds extend to the shortcut of size $n^{1+\epsilon}$ with a worse step bound.

⁵Our definition in the paper (Definition 3.6) is actually cut-based. This almost-equivalent flow-based definition follows from Theorem 4.2 and is more convenient in this overview.

Suppose further that G has diameter at most h . In this case, our shortcut is simply a star S connecting each original vertex v to a Steiner vertex r_S with an (hs) -length $A(v)$ -capacity edge. We can shortcut any feasible flow in G with 2 steps. An illustrative example is shown in Figure 1. The length slack is $O(s)$ since an h -length original feasible flow is mapped forward to a $(2hs)$ -length feasible flow in the star. The congestion slack is $O(\log n/\phi)$ because any feasible flow in the star induces an h -length A -respecting demand. Since A is (h, s) -length ϕ -expanding in G , we route such demand in G with congestion $O(\log n/\phi)$ and without length increasing.

In general, the diameter can be large. Thus, we can construct a *sparse neighborhood cover* to decompose the graph into clusters with diameter h , such that (1) for each vertex v , there is a cluster containing all vertices within distance h/s from v , and (2) each vertex is inside $n^{O(1/s)}$ clusters. Then, we can construct a shortcut by adding an (hs) -edge-length star on each cluster. By a similar argument, we obtain a flow shortcut graph for (h/s) -length original flows with length slack $O(s^2)$, congestion slack $O(n^{O(1/s)} \log n/\phi)$ and step 2.

So far, when we build an LC-flow shortcut for an LC-expander, the vertex-capacitated setting presents no difficulties compared to the edge-capacitated setting, because the above simple approach works in both settings. However, the differences between the two settings arise when generalizing this approach to general graphs via expander hierarchies.

Previous Approach: Shortcutting General Graphs via Boundary-Linkedness. The key idea of [HHL⁺24] is to exploit a hierarchy of *boundary-linked* LC-expander decomposition, defined as follows. Let G be an edge-unit-capacity graph. Initialize the node-weight $A_0 = \deg_G$. For each level $0 \leq i \leq d$, compute a cut $C_{i+1} \subseteq E$ of size $|C_{i+1}| \approx \phi|A_i|$ such that $A_i + \deg_{C_{i+1}}$ is (h, s) -length ϕ -expanding in $G - C_{i+1}$ where $\deg_{C_{i+1}}(v)$ counts the number of C_{i+1} -edges incident to v .⁶ The cut C_{i+1} is called the *boundary-linked* LC-expander decomposition for A_i because it gives a stronger expansion guarantee of $A_i + \deg_{C_{i+1}}$ instead of just A_i . Then, we set $A_{i+1} := \deg_{C_{i+1}}$ and continue to the next level $i + 1$. By setting $\phi = 1/(n^{O(1/s)}n^\epsilon)$, we have that $d = O(1/\epsilon)$ and $A_{d+1} = 0$.

From the above construction, we conclude that, for each i , $A_i + A_{i+1}$ is (h, s) -length ϕ -expanding in $G - C_{i+1}$. Therefore, as we have seen in the warm-up, we can add stars on the support of $A_i + A_{i+1}$ so that any flows routing h -length $(A_i + A_{i+1})$ -respecting demands in $G - C_{i+1}$ can be shortcut.

Now consider a feasible h -length flow in G . The boundary-linkedness suggests a natural *bottom-up* shortcut scheme. For each flow path P , we can think of routing P 's *head packet* and *tail packet* (initially at P 's left and right endpoints, denoted by u_0 and v_0) to the same place via shortcuts. Take the head packet as an example. Start with $u_0 \in \text{supp}(A_0) = V$. At each level $0 \leq i \leq d - 1$, let u_{i+1} be the left endpoint of the first $P \cap C_{i+1}$ -edge behind u_i . By definition, $u_{i+1} \in \text{supp}(A_{i+1})$ and P 's subpath between u_i and u_{i+1} is disjoint from C_{i+1} , so we can use star graphs at level i to route the head packet from u_i to u_{i+1} within 2 steps. In sum, each of the head and tail packets is routed from the bottom up until they reach $u_d, v_d \in \text{supp}(A_d)$, and the top-level star graphs can route them together. The total number of steps is $O(d) = O(1/\epsilon)$.⁷

The Obstacle from Vertex Cuts. The overall strategy of the above approach is to shortcut flow paths from a vertex of A_i to an endpoint of edges in C_{i+1} . This was possible since the boundary-linked expander decomposition guarantees that $A_i + \deg_{C_{i+1}}$ is expanding.

In the vertex-capacitated graph, however, the cut $C_{i+1} \subseteq V$ is now a vertex set. To follow

⁶In the actual construction, C_{i+1} assigns fractional values to edges and is called a *moving cut*, defined in Section 3.1. Here, we assume C_{i+1} is a classic edge cut for simplicity.

⁷We note that the step bound in [HHL⁺24] is $O(1/\epsilon^2)$ because they used powers of expander graphs instead of star graphs to avoid creating vertices outside G , which brought another $O(1/\epsilon)$ factor.

the same strategy, we have two natural options. We shortcut flow from a vertex of A_i to either (1) a vertex in C_{i+1} , or (2) a neighbor of C_{i+1} .

In the first case, the strategy requires that $A_i + C_{i+1}$ is expanding in $G - C_{i+1}$. This is trivially impossible because C_{i+1} is not even in the graph $G - C_{i+1}$. In the second case, let $N(C_{i+1})$ denote the neighbors of C_{i+1} that are not in C_{i+1} . The strategy requires $A_i + N(C_{i+1})$ is expanding in $G - C_{i+1}$. However, possibly $N(C_{i+1})$ is very big and has size $|N(C_{i+1})| = \Omega(n|C_{i+1}|)$. It is unlikely that expander decomposition exists to guarantee the expansion of such a large node-weighting. Even if it exists, we would set $A_{i+1} = N(C_{i+1})$ and, hence, we cannot guarantee $|A_{i+1}| \ll |A_i|$. So the number of levels of the hierarchy is unbounded.

In either option, this overall strategy fails in the vertex-capacitated graphs. At a very high level, this is because edges have two endpoints while vertices may have an unbounded number of neighbors.

Our Approach: Top-Down Analysis without Boundary-linkedness. We construct a similar hierarchy of LC-vertex expander decomposition *without* boundary-linkedness as follows. Let $G = (V, E)$ be a vertex-unit-capacity graph. Initialize node-weighting $A_0 = \mathbb{1}_V$. At each level $0 \leq i \leq d$, computes a cut $C_{i+1} \subseteq V$ such that A_i is (h, s) -length ϕ -vertex-expanding in $G - C_{i+1}$, and set $A_{i+1} := \mathbb{1}_{C_{i+1}}$. In particular, the top level d has $C_{d+1} = \emptyset$. The LC-vertex-expander decomposition guarantees $|C_{i+1}| \approx \phi|A_i|$, so the number d of levels is $O(1/\epsilon)$ by choosing proper ϕ .

Next, we construct the shortcut as follows. For each i , by the expansion of A_i , we can add stars on the support of A_i into our shortcut so that any flows routing h -length A_i -respecting demands in $G - C_{i+1}$ can be shortcut. To analyze the shortcut quality, we will no longer try to route from A_i to A_{i+1} as in the edge-capacitated setting, because we no longer have boundary-linkedness guarantee.

Our analysis is instead *top-down*. At each level i , we shortcut the current flow path as much as possible, and then the prefix and suffix that have not yet been shortcut will be deferred to lower levels as subproblems. To be more concrete, say our initial goal is to shortcut a flow path P in a feasible h -length original flow. At each level $0 \leq i \leq d$, assume we will receive a subpath P' of P with length at most h in $G - C_{i+1}$ (note that P is a valid input to the top level d because C_{d+1} is empty). We will shortcut P' using star graphs at levels up to i as follows (see Figure 2 for an illustration when $i = 1$).




Figure 2: A toy example of forward mapping given we have 2 levels in total. Crossings represent cut vertices in C_1 along the witness path $P_{u,v}$.

Step 1. Let x_i and y_i be the first and last P' -vertices in $\text{supp}(A_i)$ respectively. We can easily shortcut the subpath $P'[x_i, y_i]$ (i.e. the subpath from x_i to y_i) within 2 steps using the star graphs at level i .

Step 2. Let x'_i be the P' -vertex right before x_i and let y'_i be the P' -vertex right after y_i . We regard shortcutting the prefix $P'[u, x'_i]$ and the suffix $P'[y'_i, v]$ as two subproblems at level $i - 1$, where u, v are endpoints of P' . Note that both $P'[u, x'_i]$ and $P'[y'_i, v]$ has length at most h in $G - C_i$ because they are disjoint from C_i by definition.

Step 3. After the recursion, we obtain shortcuts for both $P'[u, x'_i]$ and $P'[y'_i, v]$. The shortcut for $\overline{P'}$ is given by concatenating shortcuts for $P'[u, x'_i]$, $P'[x_i, y_i]$ and $P'[y'_i, v]$ using two original edges (x'_i, x_i) and (y_i, y'_i) .

It is not hard to see the final step bound is $2^{O(d)} = 2^{O(1/\epsilon)}$ since the recursion has d levels and each level has two branches. We note that the actual argument is more complicated because the cuts C_i are actually moving cuts which have fractional cut values, and there is no clear partition of P' into 3 parts.

2 Preliminaries

This section includes preliminaries for directed graphs and vertex-capacitated graphs in Section 2.1 and Section 2.2. We note that in Section 3 we consider directed graphs, while in Section 4 and Section 5 we focus on vertex-capacitated graphs.

We always use n and m to denote the number of vertices and edges of the original graph in the context (for example, the original graph of each main theorem is its input graph). Throughout the paper, all input graphs will have length and capacity functions, and we assume that all lengths and capacities are *positive integers* upper bounded by $N = \text{poly}(n)$. To simplify notation, sometimes we may hide $O(\log N)$ factors in $O(\log n)$ or n^ϵ (where ϵ is a constant).

2.1 Directed Graphs

Let $G = (V, E)$ be a directed graph with $n := |V|$ vertices and $m := |E|$ edges. Let $\ell_G : E \rightarrow \mathbb{N}^+$ denote the edge length function of G . A path from vertex v to vertex w is called a (v, w) -path. For any path P , $\ell_G(P) = \sum_{e \in P} \ell_G(e)$ and let $|P|$ denote the number of edges in P (we also say P has $|P|$ steps). The *distance* between vertices v and w is $\text{dist}_G(v, w) = \min_{P: (v, w)\text{-path}} \ell_G(P)$. A *ball* of radius r around a vertex v is a $\text{ball}_G(v, r) = \{w \mid \text{dist}_G(v, w) \leq r\}$. We further define that $\text{dist}_G(v, w) = \infty$ if vertex w cannot be reached from vertex v .

Let $u_G : E \rightarrow \mathbb{R}_{>0}$ denote the edge capacity function of G . Sometimes we use $u(e)$ to represent the capacity over edge e if G is explicitly mentioned. We further define the directed degree of a vertex v in G , denoted by $\text{deg}_G^{\text{in}}(v)$ and $\text{deg}_G^{\text{out}}(v)$, as $\sum_{(w, v) \in E} u_G(w, v)$ and $\sum_{(v, w) \in E} u_G(v, w)$ respectively. Namely, they represent the capacity summation of incoming edges to v and outgoing edges from v . We use $\text{deg}_G^{\text{sum}}(v) = \text{deg}_G^{\text{in}}(v) + \text{deg}_G^{\text{out}}(v)$ to represent the capacity of all edges incident to v . We use $\text{deg}_G^{\text{min}}(v) = \min\{\text{deg}_G^{\text{in}}(v), \text{deg}_G^{\text{out}}(v)\}$ to measure the minimum of in-degree and out-degree of a vertex v .

Multicommodity Flows. A (*multicommodity*) *flow/routing* in G is a function $F : \mathcal{P} \rightarrow \mathbb{R}_{\geq 0}$ (where \mathcal{P} denotes the set of simple paths in G) that assigns each simple path P in G a flow value $F(P) \geq 0$. We define P to be a *flow path* of F if $F(P) > 0$. Further P is a (v, w) -*flow path* of F if P is both a (v, w) -path and a flow path of F . Let $\text{path}(F)$ denote the set of all paths P where $F(P) > 0$. The *value* of F is denoted by $\text{value}(F) = \sum_{P \in \text{path}(F)} F(P)$. We point out that, for each flow path $P \in \text{path}(F)$, we also regard P as a flow with only one flow path. The value of this flow P is $\text{value}(P) = F(P)$ unless otherwise stated.

The *congestion* of F on an edge e is $\text{cong}_F(e) = \frac{F(e)}{u_G(e)}$ where $F(e) = \sum_{P: P \ni e} F(P)$ denotes the total flow value of all paths passing through e . The *congestion* of F is $\text{cong}(F) = \max_{e \in E} \text{cong}_F(e)$. We define a flow as *feasible* if its congestion does not exceed one. The *step* of F is the maximum step count across all flow paths in F , given by $\text{step}(F) = \max_{P \in \text{path}(F)} |P|$. Similarly, the *length* of F is defined as the maximum length of all flow paths of F , i.e., $\text{leng}(F) = \max_{P \in \text{path}(F)} \ell(P)$. Sometimes we will refer to $\text{leng}(F, G')$ as the length of F in G' , where G' is the same graph as G except that G' may have different edge lengths.

Node-Weightings. A *node-weighting* $A : V \rightarrow \mathbb{R}_{\geq 0}$ of G assigns a non-negative value $A(v)$ to a vertex v . The *size* of A is denoted by $|A| = \sum_v A(v)$ and let $\text{supp}(A) := \{v : A(v) > 0\}$. For two node-weightings A, A' we define $\min(A, A')$ and $A + A'$ as pointwise operations, and we write $A \preceq A'$ if A is pointwise at most A' .

We further define the volume of a vertex subset $S \subseteq V$ w.r.t a node-weighting A to be $\text{vol}_A(S) = \sum_{v \in S} A(v)$. Note that the degree function is also a node-weighting, and we let $\text{vol}_G^{\text{sum}}(S) = \sum_{v \in S} \text{deg}_G^{\text{sum}}(v)$.

Demands. A *demand* $D : V \times V \rightarrow \mathbb{R}_{\geq 0}$ assigns a non-negative value $D(v, w) \geq 0$ to each

ordered pair of vertices in V to specify the units of demand $D(v, w)$ vertex v wants to send to vertex w . We note that it is always safe to assume $D(u, u) = 0$. The size of a demand is written as $|D|$ and is defined as $\sum_{v,w} D(v, w)$. For each vertex v , we let $D(v, \cdot) = \sum_w D(v, w)$ and $D(\cdot, v) = \sum_w D(w, v)$ denote the total demand starting and ending at v respectively. We say a demand D is

- *h-length constrained*: if for each $v, w \in V$ with $D(v, w) > 0$, we have $\text{dist}_G(v, w) \leq h$;
- *symmetric*: if for any pairs of $v, w \in V$, $D(v, w) = D(w, v)$;
- *sub-demand for D'* : if for any pairs of $v, w \in V$, $D(v, w) \leq D'(v, w)$;
- *A-respecting for some node-weighting A* : if for any $v \in V$, $\max\{D(v, \cdot), D(\cdot, v)\} \leq A(v)$.

The demand routed by F (or the corresponding demand of F), denoted by $\text{Dem}(F)$ or D_F , has $D_F(u, v) = \sum_{(u,v)\text{-flow path } P \in \text{path}(F)} F(P)$. We say a demand D can be routed with length h , congestion γ and step t if there is a flow F routing D with length h , congestion γ and step t .

2.2 Vertex-Capacitated Graphs

An undirected vertex-capacitated graph, denoted by $G = (V, E)$ has lengths and capacities on both edges and vertices. Formally, G has length function $\ell_G : V \cup E \rightarrow \mathbb{N}^+$ and capacity function $u_G : V \cup E \rightarrow \mathbb{N}^+$. We will use $u_{V(G)}$ to denote the restriction of u_G on V , i.e. the vertex capacity function of G . Naturally, for any simple path P in G , its length $\ell_G(P) = \sum_{\text{vertices } v \in P} \ell_G(v) + \sum_{\text{edges } e \in P} \ell_G(e)$, but its step $|P|$ is still the number of edges in P . For each vertex $v \in V$, its (capacitated) degree is $\text{deg}_G(v) = u_G(v) + \sum_{e \in E \text{ incident to } v} u_G(e)$.

Vertex-Capacitated Flows. The notation of flow naturally extends to the setting of vertex-capacitated graphs. The only difference is the definition of congestion. We note that the congestion of flow F over a vertex v is the $\text{cong}_F(v) = \frac{F(v)}{u_G(v)}$ where $F(v) = \sum_{P: P \ni v} F(P)$ denotes the total flow value of all flow paths going through v . Thus the general congestion of flow F also considers the congestion over vertices, i.e.,

$$\text{cong}_F = \max\left\{ \max_{v \in V(G)} \text{cong}_F(v), \max_{e \in E(G)} \text{cong}_F(e) \right\}$$

We further note that the step of flow F in vertex-capacitated graphs still depends on the number of edges in flow paths.

Neighborhood Covers. We will use neighborhood covers in Section 5. Given a graph G with lengths, a clustering \mathcal{S} in G is a collection of pairwise disjoint vertex sets $S_1, \dots, S_{|\mathcal{S}|}$, called clusters. A neighborhood cover \mathcal{N} with width ω and covering radius h is a collection of ω many clusterings $\mathcal{S}_1, \dots, \mathcal{S}_\omega$ such that for every node v there exists a cluster $S \in \mathcal{S}_i$ such that $\text{ball}(v, h) \subseteq S$, where $\text{ball}(v, h) = \{u \in V(G) \mid \text{dist}_G(u, v) \leq h\}$. We use $S \in \mathcal{N}$ to denote that S is a cluster in some clustering of \mathcal{N} . The clustering \mathcal{N} has diameter h_{diam} if every cluster $S \in \mathcal{N}$ has diameter at most h_{diam} i.e., $\max_{u, v \in S} \text{dist}_G(u, v) \leq h_{\text{diam}}$. We note that the shortest path between $u, v \in S$ may use vertices that are not in the cluster.

Theorem 2.1 ([Pel00]). *Given a vertex-capacitated graph G with a length parameter h and an integer $k \geq 1$, there exists a neighborhood cover \mathcal{N} with covering radius h , diameter $h_{\text{diam}} \leq (2k - 1) \cdot h$ and width $\omega = n^{O(1/k)} k$.*

3 Length-Constrained Directed Expansion

In this section, we follow the theory of length-constrained expansion and extend it to the setting of directed graphs. We start with the generalization of notations from length-constrained expanders in Section 3.1, which serves as the foundation for subsequent results. Next, we characterize length-constrained expansion in directed graphs with routing in Section 3.2, and show the existence of length-constrained directed expander decomposition in Section 3.3.

3.1 Basic Concepts of Length-Constrained Directed Expansion

The following definition of moving cuts and separation was introduced by Haeupler, Wajc and Zuzic in [HWZ20].

Definition 3.1 (Length-Constrained Cut). *An h -length moving cut $C : E \mapsto \{0, \frac{1}{h}, \frac{2}{h}, \dots, 1\}$ assigns to each edge e a fractional cut value between zero and one which is a multiple of $\frac{1}{h}$. The size of C is defined as $|C| = \sum_e u(e) \cdot C(e)$. The length increase associated with the h -length moving cut C is denoted with $\ell_{C,h}$ and defined as assigning an edge e the length increase $\ell_{C,h}(e) = h \cdot C(e)$. Any moving cut which only assigns cut values equal to either 0 or 1 is called a pure moving cut. We define the degree of a moving cut over vertex v to be $\deg_C(v) = \sum_{e \ni v} u_G(e) \cdot C(e)$.*

Definition 3.2 (h -Length Separated Demand). *For any demand D and any h -length moving cut C , we define the amount of h -length separated demand as the sum of demands between vertices that are h -length separated by C . We denote this quantity with $\text{sep}_h(C, D)$, i.e.,*

$$\text{sep}_h(C, D) = \sum_{u,v: \text{dist}_{G-C}(u,v) > h} D(u, v).$$

Definition 3.3 (h -Length Sparsity of a Cut C for Demand D). *For any demand D and any h -length moving cut C with $\text{sep}_h(C, D) > 0$, the h -length sparsity of C with respect to D is the ratio of C 's size to how much demand it h -length separates i.e.,*

$$\text{spars}_h(C, D) = \frac{|C|}{\text{sep}_h(C, D)}.$$

Above we generalize the definition of length-constrained moving cut w.r.t arbitrary directed h -length demand. However, for the definition of a directed length-constrained expander, we restrict to symmetric h -length demands.

Definition 3.4 ((h, s) -Length Sparsity of a Cut w.r.t. a Node-Weighting). *The (h, s) -length sparsity of any $h \cdot s$ -length moving cut C with respect to a node-weighting A is defined as:*

$$\text{spars}_{(h,s)}(C, A) = \min_{A\text{-respecting } h\text{-length symmetric demand } D} \text{spars}_{h \cdot s}(C, D).$$

Intuitively, $(h \cdot s)$ -length sparsity of a cut measures how much it $h \cdot s$ -length separates h -length demand w.r.t its own size. Furthermore, for a given node-weighting, we associate the sparsest cut w.r.t the node-weighting with its conductance.

Definition 3.5 ((h, s) -Length Conductance of a Node-Weighting). *The (h, s) -length conductance of a node-weighting A in a graph G is defined as the (h, s) -length sparsity of the sparsest $h \cdot s$ -length moving cut C with respect to A , i.e.,*

$$\text{cond}_{(h,s)}(A) = \min_{h \cdot s\text{-length moving cut } C} \text{spars}_{(h,s)}(C, A).$$

Definition 3.6 ((h, s) -Length ϕ -Expanding Node-Weightings). *We say a node-weighting A is (h, s) -length ϕ -expanding if the (h, s) -length conductance of A in G is at least ϕ .*

To see the connection, in Appendix A.1, we explain how our notion of length-constrained directed expansion generalizes the non-length-constrained version of directed expansion. Lastly, we give the formal definition of length-constrained directed expander decompositions as follows:

Definition 3.7 (Length-Constrained Directed Expander Decomposition). *Given a graph $G = (V, E)$, a directed (h, s) -length ϕ -expander decomposition for a node-weighting A with length slack s and cut slack κ is an $h \cdot s$ -length cut C of size at most $\kappa \cdot \phi|A|$ such that A is (h, s) -length ϕ -expanding in $G - C$.*

3.2 Routing Characterization of Length-Constrained Directed Expansion

The definition of ϕ -expanding characterizes the sparsity of moving cuts in directed graphs. With the routing characterization, we certify the notation to be meaningful and show that sparsity is closely related to demand routing.

Theorem 3.8 (Routing Characterization of Length-Constrained Directed Expanders). *Given a directed graph G and node-weighting A , for any $h \geq 1$, $\phi < 1$ and $s \geq 1$ we have:*

- *If A is (h, s) -length ϕ -expanding in G , then every A -respecting h -length symmetric demand can be routed in G with congestion at most $O(\frac{\log N}{\phi})$ and length at most $h \cdot s$.*
- *If A is not (h, s) -length ϕ -expanding in G , then some A -respecting h -length symmetric demand cannot be routed with congestion at most $\frac{1}{2\phi}$ and length at most $\frac{h \cdot s}{2}$.*

The proof idea of Theorem 3.8 is similar to the undirected case as shown in [HRG22], and for completeness, we restate and adapt the proof for the directed setting in Appendix A.2.

3.3 Length-Constrained Directed Expander Decomposition: Existence

Now, we prove the existence of length-constrained directed expander decompositions. The following theorem formally states the result:

Theorem 3.9. *For any $G = (V, E)$, a node-weighting A , $h > 1$, $\alpha \geq 1$, $\phi < 1$ and a length slack parameter $s = O(\log n)$, there is a directed (h, s) -length ϕ -expander decomposition for A with cut slack $\kappa = O(n^{O(\frac{1}{s})} \log n)$.*

The proof of Theorem 3.9 again follows the undirected-case proof in [HRG22], and we append it for completeness. The basic idea to prove the above main theorem is that we can continuously find a cut with (h, s) -sparsity less than ϕ in the graph if it is not an (h, s) -length ϕ -expander for node-weighting A . We can apply the cut to the graph and repeat the same procedure. Finally the union of those cuts will render the graph as an (h, s) -length ϕ -expander.

In order to argue the upper bound for the total cut size, we need to consider the summation of each individual cut size. However, the size of each cut depends on the sparsity associated with different demands, which adds complexity to the problem. Thus, we first introduce a special base demand, called exponential demand, in Section 3.3.1 to relate all other demands in terms of sparsity. Using this, we apply a potential argument to prove the above main theorem in Section 3.3.2.

In Appendix A.3, we will discuss *boundary-linked* LC-directed expander decomposition (also called *linked* LC-directed expander decomposition). Expander decompositions with boundary-linkedness have been shown to be very useful in the length-constrained undirected setting and the classic (i.e. non-length-constrained) setting. Hence we include this part which may lead to future applications, although it has no application in our work.

3.3.1 Exponential Demand

Exponential demand can be viewed as a worst-case demand because, for any sparse cut with a witnessing demand, it admits comparable sparsity w.r.t that same cut. Specifically, we would like to show the following lemma:

Lemma 3.10. *Given a directed graph $G = (V, E)$ and a node-weighting A , suppose that some A -respecting h -length symmetric demand D has $h \cdot s$ -length sparsity at most ϕ w.r.t. some $h \cdot s$ -length moving cut C . Then the α -exponential demand $D_{h,A}^\alpha$ has $\frac{h \cdot s}{2}$ -length sparsity at most $2^{8\alpha+1}\phi$ w.r.t the same cut C .*

We will develop the definition of the exponential demand (i.e. Definition 3.13) in the following part and start with a specific weight function base on distance.

Exponential Distance Weight For a directed graph $G = (V, E)$, and a length-bound h , we define h -length α -exponential distance weight of a vertex u w.r.t. vertex v as

$$w_h^\alpha(u, v) := \begin{cases} 1 & u = v \\ 2^{-\alpha \cdot \overline{\text{dist}}(u,v)/h} & \overline{\text{dist}}(u, v) \leq \frac{2h \log_2 n}{\alpha} \\ 0 & \text{otw.} \end{cases}$$

where $1 \leq \alpha \leq \log n$ and $\overline{\text{dist}}(u, v) = \text{dist}(u, v) + \text{dist}(v, u)$ is the round-trip distance between vertex pairs. It immediately follows that the round-trip distance is symmetric, i.e., $\overline{\text{dist}}(v, u) = \overline{\text{dist}}(u, v)$.

The following lemma serves as an introduction to the property of exponential distance weight.

Lemma 3.11. *For any graph G , length-bound h , vertices $u, v, w \in V$ the following hold for the h -length α -exponential distance weights:*

1. $\sum_{a \in V} w_h^\alpha(u, a) \leq n$.
2. $\sum_{a \in V} w_h^\alpha(u, a) \geq 1$.
3. $w_h^\alpha(u, v) \geq 2^{-\alpha \cdot \overline{\text{dist}}(u,v)/h} - 1/n^2$.
4. $2^{-\alpha \cdot \overline{\text{dist}}(w,v)/h} \cdot w_h^\alpha(u, v) - 1/n^2 \leq w_h^\alpha(u, w) \leq 2^{\alpha \cdot \overline{\text{dist}}(w,v)/h} \cdot (w_h^\alpha(u, v) + 1/n^2)$

Proof. We provide the proofs as an itemized list corresponding to the statements in the lemma.

1. Every vertex in v receives a weight $w_h^\alpha(u, a)$ of at most one and $|V| = n$.
2. Follows because $w_h^\alpha(u, u) = 1$.
3. If $\overline{\text{dist}}(u, v) < \frac{2h \log_2 n}{\alpha}$, then it is true for $w_h^\alpha(u, v) = 2^{-\alpha \cdot \overline{\text{dist}}(u,v)/h}$ and $\frac{1}{n^2} > 0$; If $\overline{\text{dist}}(u, v) \geq \frac{2h \log_2 n}{\alpha}$, then $2^{-\alpha \cdot \overline{\text{dist}}(u,v)/h} \leq \frac{1}{n^2}$ and $w_h^\alpha(u, v) = 0$, which concludes the property.
4. By triangle inequality we have $\overline{\text{dist}}(u, w) + \overline{\text{dist}}(w, v) = \text{dist}(u, w) + \text{dist}(w, u) + \text{dist}(w, v) + \text{dist}(v, w) \geq \text{dist}(u, v) + \text{dist}(v, u) = \overline{\text{dist}}(u, v)$. This further gives $\overline{\text{dist}}(u, w) \geq \overline{\text{dist}}(u, v) - \overline{\text{dist}}(w, v)$, and thus

$$2^{-\alpha \cdot \overline{\text{dist}}(u,w)/h} \leq 2^{\alpha \cdot \overline{\text{dist}}(w,v)/h} \cdot 2^{-\alpha \cdot \overline{\text{dist}}(u,v)/h}$$

We note that $w_h^\alpha(u, w) \leq 2^{-\alpha \cdot \overline{\text{dist}}(u, w)/h}$, and by Property 3, we upper bound $2^{-\alpha \cdot \overline{\text{dist}}(u, v)/h}$ by $w_h^\alpha(u, v) + 1/n^2$ to get

$$w_h^\alpha(u, w) \leq 2^{\alpha \cdot \overline{\text{dist}}(w, v)/h} \cdot (w_h^\alpha(u, v) + 1/n^2)$$

Since the choice of vertices is symmetric, by swapping vertices w and v , we can similarly have that

$$\begin{aligned} w_h^\alpha(u, v) &\leq 2^{\alpha \cdot \overline{\text{dist}}(v, w)/h} \cdot (w_h^\alpha(u, w) + 1/n^2) \\ 2^{-\alpha \cdot \overline{\text{dist}}(v, w)/h} \cdot w_h^\alpha(u, v) &\leq w_h^\alpha(u, w) + 1/n^2 \\ 2^{-\alpha \cdot \overline{\text{dist}}(v, w)/h} \cdot w_h^\alpha(u, v) - 1/n^2 &\leq w_h^\alpha(u, w) \end{aligned}$$

And note that $\overline{\text{dist}}(v, w) = \overline{\text{dist}}(w, v)$, which concludes the proof. \square

The intuition behind the distance weight is that we assign more weight to closer vertices. We let

$$w_h^\alpha(u) := \sum_{b \in V} w_h^\alpha(u, b)$$

denote the h -length α -exponential weight of u . It can be interpreted as a normalization factor of the exponential weight over vertex u , which determines the fraction of demand that u is supposed to send out to any other vertex.

Mixing Factor To explicitly denote this ratio, we define $M_h^\alpha(u, v)$ to be the h -length α -mixing factor from vertices u to v as follows:

$$M_h^\alpha(u, v) = \frac{w_h^\alpha(u, v)}{w_h^\alpha(u)}$$

It immediately follows that $0 \leq M_h^\alpha(u, v) \leq 1$.

This mixing factor instructs a unit demand between every pair of vertices according to the distance. We are interested in the overlap of the mixing factor, in particular, for close enough vertex pairs u and v , we count the summation of $\min\{M_h^\alpha(u, b), M_h^\alpha(v, b)\}$ over every vertex $b \in V$. It turns out that there exists a lower bound for the overlap with the following lemma:

Lemma 3.12. *Given a directed graph $G = (V, E)$, $1 \leq \alpha \leq \log n$, for any pair of vertices u, v where $\overline{\text{dist}}(u, v) \leq 2h$, we have*

$$\sum_{b \in V} \min\{M_h^\alpha(u, b), M_h^\alpha(v, b)\} \geq 2^{-8\alpha}$$

Proof.

$$\sum_{b \in V} \min\{M_h^\alpha(u, b), M_h^\alpha(v, b)\} = \sum_{b \in V} \min\left\{\frac{w_h^\alpha(u, b)}{w_h^\alpha(u)}, \frac{w_h^\alpha(v, b)}{w_h^\alpha(v)}\right\}$$

We note that only vertex b where $w_h^\alpha(u, b) > 0$ and $w_h^\alpha(v, b) > 0$ will contribute to the above summation otherwise the minimum will take the value 0. Then for simplicity of notation, we let

$$B_u = \left\{w \in V \mid \overline{\text{dist}}(u, w) \leq \frac{2h \log_2 n}{\alpha}\right\}$$

denote the ball centered on vertex u with the radius of $\frac{2h \log_2 n}{\alpha}$ in round-trip distance. For any vertex $v \in V \setminus B_u$, we have $w_h^\alpha(u, v) = 0$. Further we can rewrite the summation as

$$\sum_{b \in V} \min\left\{\frac{w_h^\alpha(u, b)}{w_h^\alpha(u)}, \frac{w_h^\alpha(v, b)}{w_h^\alpha(v)}\right\} = \sum_{b \in B_u \cap B_v} \min\left\{\frac{w_h^\alpha(u, b)}{w_h^\alpha(u)}, \frac{w_h^\alpha(v, b)}{w_h^\alpha(v)}\right\} \quad (1)$$

$$\geq \sum_{b \in B_u \cap B_v} \frac{\min\{w_h^\alpha(u, b), w_h^\alpha(v, b)\}}{\max\{w_h^\alpha(u), w_h^\alpha(v)\}} \quad (2)$$

For inequality (2), we use the fact that $\min\{\frac{a}{b}, \frac{c}{d}\} \geq \frac{\min\{a, c\}}{\max\{b, d\}}$. To further bound the above summation, we first build the relationship between weights over vertices u and v . For the numerator, for vertices u, v and $b \in B_u \cap B_v$, and by triangle inequality that $\overline{\text{dist}}(b, u) \geq \overline{\text{dist}}(b, v) - \overline{\text{dist}}(u, v)$, we have

$$w_h^\alpha(b, u) \leq 2^{-\alpha \cdot \overline{\text{dist}}(b, u)/h} \leq 2^{\alpha \cdot \overline{\text{dist}}(u, v)/h} \cdot 2^{-\alpha \cdot \overline{\text{dist}}(b, v)/h}$$

Note that we have $\overline{\text{dist}}_G(u, v) \leq 2h$ and $2^{-\alpha \cdot \overline{\text{dist}}(b, v)/h} = w_h^\alpha(v, b)$ since $b \in B_v$, we actually tighten the inequality and get

$$w_h^\alpha(u, b) = w_h^\alpha(b, u) \leq 2^{2\alpha} \cdot w_h^\alpha(v, b)$$

We rewrite it as

$$w_h^\alpha(v, b) \geq 2^{-2\alpha} \cdot w_h^\alpha(u, b)$$

For the denominator, from the same property and the symmetry between vertices u and v , we have

$$\begin{aligned} w_h^\alpha(v) &= \sum_{b \in B_u} w_h^\alpha(v, b) + \sum_{b \in V \setminus B_u} w_h^\alpha(v, b) \\ &\leq \sum_{b \in B_u} 2^{2\alpha} \cdot w_h^\alpha(u, b) + \sum_{b \in V \setminus B_u} 2^{2\alpha} \cdot (w_h^\alpha(u, b) + \frac{1}{n^2}) \\ &= \sum_{b \in V} 2^{2\alpha} \cdot w_h^\alpha(u, b) + \sum_{b \in V \setminus B_u} 2^{2\alpha} \cdot \frac{1}{n^2} \\ &\leq 2^{2\alpha} (w_h^\alpha(u) + \frac{1}{n}) \\ &\leq 2^{2\alpha+1} w_h^\alpha(u) \end{aligned}$$

where we use the fact that $\frac{1}{n} < w_h^\alpha(u)$.

Now combine two inequalities for the numerator and the denominator respectively we have

$$\begin{aligned} \sum_{b \in B_u \cap B_v} \frac{\min\{w_h^\alpha(u, b), w_h^\alpha(v, b)\}}{\max\{w_h^\alpha(u), w_h^\alpha(v)\}} &\geq \sum_{b \in B_u \cap B_v} \frac{\min\{w_h^\alpha(u, b), 2^{-2\alpha} \cdot w_h^\alpha(u, b)\}}{\max\{w_h^\alpha(u), 2^{2\alpha+1} w_h^\alpha(u)\}} \\ &= \frac{1}{2^{4\alpha+1}} \cdot \frac{\sum_{b \in B_u \cap B_v} w_h^\alpha(u, b)}{w_h^\alpha(u)} \\ &= \frac{1}{2^{4\alpha+1}} \cdot \frac{\sum_{b \in B_u \cap B_v} w_h^\alpha(u, b)}{\sum_{b \in B_u} w_h^\alpha(u, b)} \\ &= \frac{1}{2^{4\alpha+1}} \cdot \frac{\sum_{b \in B_u \cap B_v} w_h^\alpha(u, b)}{\sum_{b \in B_u \cap B_v} w_h^\alpha(u, b) + \sum_{b \in B_u \setminus B_v} w_h^\alpha(u, b)} \end{aligned}$$

Note that in the second equation, we rewrite the denominator since only vertices in the B_u contribute to the $w_{h,A}^\alpha(u)$. We divide it into two parts in the last equation and provide a bound between them in what follows.

For a vertex $b \in B_u \setminus B_v$, we know that it should be far away from vertex v , but close enough to vertex u . However, under the restriction that $\overline{\text{dist}}(u, v) \leq 2h$, b has to lie near the boundary of B_u , namely $\overline{\text{dist}}(u, b) \geq \frac{2h \log_2 n}{\alpha} - 2h$, and thus $w_h^\alpha(u, b) \leq 2^{2\alpha} \cdot \frac{1}{n^2}$. Consequently,

$$\sum_{b \in B_u \setminus B_v} w_h^\alpha(u, b) \leq \sum_{b \in B_u \setminus B_v} 2^{2\alpha} \cdot \frac{1}{n^2} \quad (3)$$

$$\leq 2^{2\alpha} \cdot \frac{1}{n} \quad (4)$$

$$\leq 2^{2\alpha} \cdot w_h^\alpha(u, u) \quad (5)$$

$$\leq 2^{2\alpha} \cdot \sum_{b \in B_u \cap B_v} w_h^\alpha(u, b) \quad (6)$$

In the inequality (4), we use the assumption that $|V| \leq n$. Then in the next inequality (5), we have $\frac{1}{n} \leq 1 = w_h^\alpha(u, u)$. The inequality (6) is because $u \in B_v$ (since $\alpha \leq \log_2 n$). This further concludes that

$$\frac{\sum_{b \in B_u \cap B_v} w_h^\alpha(u, b)}{\sum_{b \in B_u \cap B_v} w_h^\alpha(u, b) + \sum_{b \in B_u \setminus B_v} w_h^\alpha(u, b)} \geq \frac{1}{2^{2\alpha} + 1}$$

And finally, we are able to show that

$$\sum_{b \in V} \min\{M_h^\alpha(u, b), M_h^\alpha(v, b)\} \geq \frac{1}{2^{4\alpha+1}} \cdot \frac{1}{2^{2\alpha} + 1} \geq 2^{-8\alpha}$$

□

The above lemma shows that we can have a mixing factor with favorable overlap in the graph G . The mixing factor can be viewed as an instructor of how we send out the demand in the graph. It further helps define the exponential demand more easily as follows:

Definition 3.13 (α -Exponential Demand w.r.t h -length). *In a directed graph $G = (V, E)$, the α -exponential demand $D_{h,A}^\alpha$ w.r.t h -length and a node-weighting A is the demand defined as:*

$$D_{h,A}^\alpha(u, v) = A(u) \cdot M_h^\alpha(u, v) + A(v) \cdot M_h^\alpha(v, u)$$

For simplicity, we sometimes hide the parameter α and refer to the demand as the exponential demand.

Remark 3.14. *We note that for vertex pair $u, v \in V$ that is far away from each, i.e. $\overline{\text{dist}}(u, v) > 2h \log n / \alpha$, $D_{h,A}^\alpha(u, v)$ is zero. Since $2h \log n / \alpha > h$, the α -exponential demand is actually not an h -length demand. Furthermore, suppose there are vertices u and v such that $A(u) \ll A(v)$. From the symmetric construction, exponential demand might send more value from u than $A(u)$, thereby failing to respect node-weighting A . In conclusion, the α -exponential demand is neither guaranteed to be an h -length demand nor an A -respecting demand, but we will see in the following that it works like a worst-case demand and connects with other h -length demands in terms of sparsity.*

Corollary 3.15. *The α -exponential demand $D_{h,A}^\alpha$ is symmetric,*

$$D_{h,A}^\alpha(u, v) = D_{h,A}^\alpha(v, u)$$

As mentioned in the beginning, the reason we are interested in such an exponential demand is that it evenly mixed in the entire graph such that it can be regarded as a worst-case demand. We are now ready for the proof of the main lemma in this section.

Lemma 3.10. *Given a directed graph $G = (V, E)$ and a node-weighting A , suppose that some A -respecting h -length symmetric demand D has $h \cdot s$ -length sparsity at most ϕ w.r.t. some $h \cdot s$ -length moving cut C . Then the α -exponential demand $D_{h,A}^\alpha$ has $\frac{h \cdot s}{2}$ -length sparsity at most $2^{8\alpha+1}\phi$ w.r.t the same cut C .*

Proof. We first note that only separated demand pairs in D are involved in the sparsity w.r.t cut C , thus we restrict to a sub-demand \widehat{D} of D where $\widehat{D}(u, v) = D(u, v)$ if $\text{dist}_{G-C}(u, v) > h \cdot s$, otherwise $\widehat{D}(u, v) = 0$. It naturally follows that

$$\text{sep}_{h \cdot s}(C, D) = \text{sep}_{h \cdot s}(C, \widehat{D}) = |\widehat{D}| \geq \frac{|C|}{\phi}.$$

Further, we construct an intermediate demand D' to relate the exponential demand $D_{h,A}^\alpha$ to the fully separated demand \widehat{D} . This helps to build a connection between their sparsity w.r.t C . Intuitively, we decompose and reroute every demand pair in \widehat{D} according to the configuration and size of exponential demand $D_{h,A}^\alpha$. Specifically, for any demand $\widehat{D}(u, v)$ in \widehat{D} , for every vertex $b \in V$, we add $\widehat{D}(u, v) \cdot \min\{\frac{D_{h,A}^\alpha(u, b)}{A(u)}, \frac{D_{h,A}^\alpha(v, b)}{A(v)}\}$ to $D'(u, b)$ and $D'(b, v)$. From such a construction, the value of $D'(u, v)$ comes from non-zero demand $\widehat{D}(u, w)$ and $\widehat{D}(w, v)$, and in total, we have that

$$\begin{aligned} D'(u, v) &= \sum_{w: \widehat{D}(u, w) > 0} \widehat{D}(u, w) \cdot \min\left\{\frac{D_{h,A}^\alpha(u, v)}{A(u)}, \frac{D_{h,A}^\alpha(w, v)}{A(w)}\right\} \\ &+ \sum_{w: \widehat{D}(w, v) > 0} \widehat{D}(w, v) \cdot \min\left\{\frac{D_{h,A}^\alpha(w, u)}{A(w)}, \frac{D_{h,A}^\alpha(v, u)}{A(v)}\right\} \end{aligned}$$

In other words, demand D' depends on the fully separated demand \widehat{D} and the exponential demand. By the property that \widehat{D} is an A -respecting demand, it turns out that demand D' is a sub-demand for $2 \cdot D_{h,A}^\alpha$. To see this, we note that $\min\{a, b\} \leq a$ and $\min\{a, b\} \leq b$ for any $a, b \in \mathbb{R}$, then we have,

$$D'(u, v) \leq \sum_{w: \widehat{D}(u, w) > 0} \widehat{D}(u, w) \cdot \frac{D_{h,A}^\alpha(u, v)}{A(u)} + \sum_{w: \widehat{D}(w, v) > 0} \widehat{D}(w, v) \cdot \frac{D_{h,A}^\alpha(v, u)}{A(v)} \quad (7)$$

$$= \sum_{w: \widehat{D}(u, w) > 0} \frac{\widehat{D}(u, w)}{A(u)} \cdot D_{h,A}^\alpha(u, v) + \sum_{w: \widehat{D}(w, v) > 0} \frac{\widehat{D}(w, v)}{A(v)} \cdot D_{h,A}^\alpha(u, v) \quad (8)$$

$$\leq 2 \cdot D_{h,A}^\alpha(u, v) \quad (9)$$

We use symmetry in equation (8) where $D_{h,A}^\alpha(v, u) = D_{h,A}^\alpha(u, v)$. And for the last inequality (9), we use the fact that $\sum_{w: \widehat{D}(u, w) > 0} \widehat{D}(u, w) \leq A(u)$ since \widehat{D} is A -respecting. Since the inequality applies to every vertex pair $u, v \in V$, it confirms that D' is a sub-demand for $2 \cdot D_{h,A}^\alpha$. This directly gives that $\text{sep}_{h \cdot s/2}(C, D_{h,A}^\alpha) \geq \frac{1}{2} \text{sep}_{h \cdot s/2}(C, D')$.

Then we show that the $\text{sep}_{h \cdot s/2}(C, D')$ is at least some fraction (dependent on α) of the amount of the fully separated demand \widehat{D} w.r.t the same cut C . For each demand $\widehat{D}(u, v)$ that contributes to the $\text{sep}_{h \cdot s}(C, \widehat{D})$, we have $\text{dist}_{G-C}(u, v) > h \cdot s$. As a result, for any other vertex b , we have either $\text{dist}_{G-C}(u, b) > \frac{h \cdot s}{2}$ or $\text{dist}_{G-C}(b, v) > \frac{h \cdot s}{2}$. Then either from $D'(u, b)$ or from

$D'(b, v)$, the amount of $\widehat{D}(u, v) \cdot \min\{\frac{D_{h,A}^\alpha(u,b)}{A(u)}, \frac{D_{h,A}^\alpha(v,b)}{A(v)}\}$ is contributed to $\text{sep}_{h.s/2}(C, D')$. If we sum up all vertices b from the vertex set, in total

$$\sum_{b \in V} \widehat{D}(u, v) \cdot \min\{\frac{D_{h,A}^\alpha(u,b)}{A(u)}, \frac{D_{h,A}^\alpha(v,b)}{A(v)}\} \geq \widehat{D}(u, v) \cdot \sum_{b \in V} \min\{M_h^\alpha(u, b), M_h^\alpha(v, b)\} \quad (10)$$

$$\geq 2^{-8\alpha} \widehat{D}(u, v) \quad (11)$$

is contributed to the overall separated amount of D' for a single pair (u, v) . The reason that we can apply Lemma 3.12 to get inequality (11) is that D is a symmetric h -length demand, and thus $\text{dist}(u, v) \leq h$ and $\text{dist}(v, u) \leq h$, which means $\overline{\text{dist}}(u, v) \leq 2h$. After summing up all pairs, we have

$$\text{sep}_{h.s/2}(C, D') \geq \sum_{u,v \in V} 2^{-8\alpha} \widehat{D}(u, v) = 2^{-8\alpha} |\widehat{D}| = 2^{-8\alpha} \text{sep}_{h.s}(C, \widehat{D})$$

And finally we can conclude that

$$\text{sep}_{h.s/2}(C, D_{h,A}^\alpha) \geq \frac{1}{2} \text{sep}_{h.s/2}(C, D') \geq 2^{-8\alpha-1} \text{sep}_{h.s}(C, \widehat{D}) = 2^{-8\alpha-1} \text{sep}_{h.s}(C, D).$$

This gives the lemma. \square

The above lemma helps to relate every demand associated with a sparse cut to the exponential demand. In other words, whenever we are faced with a sparse cut, we can stick with the exponential demand for analysis by only losing a factor of $\exp(\alpha)$.

3.3.2 Existential Proof of the Decomposition

Finally we will show the existence of length-constrained expander decompositions. For the existence of the length-constrained expander decomposition for a graph G w.r.t some node-weighting A , we can find sparse cuts iteratively from the graph. Namely, if the graph G is not an expander, it is guaranteed to admit a sparse moving cut C . We can apply this cut to the graph and get a new graph $G' = G - C$. This can be done iteratively until the updated graph is already expander, or in other words, there does not exist any sparse moving cut. This gives a sequence of moving cuts, and we can combine them as a single moving cut to show the existence of expander decompositions.

We first formally describe the sequence of moving cuts as follows:

Definition 3.16 (Sequence of Moving Cuts). *Given a directed graph $G = (V, E)$, and node-weighting A , let (C_1, C_2, \dots, C_n) be a sequence of $h \cdot s$ moving cuts, let $G - \sum_{j < i} C_j$ denote the graph that is applied with cuts from C_1 to C_{i-1} . We define (C_1, C_2, \dots, C_n) as a sequence of ϕ -sparse moving cuts if and only if the (h, s) -length sparsity of C_i w.r.t A in $G - \sum_{j < i} C_j$ is at most ϕ .*

It would be less interesting if those moving cuts have very large size, thus it is important to bound the overall size of those moving cuts.

Lemma 3.17. *Let C_1, \dots, C_k be an sequence of ϕ -sparse $h \cdot s$ -length cuts for some node-weighting A in the graph G where $h > 1$, $\phi < 1$, $1 \leq \alpha \leq \log n$ and $s > \frac{4 \log_2 n}{\alpha}$, then $\sum_i |C_i| \leq (2^{8\alpha+2} \phi \ln n) \cdot |A|$.*

Proof. Let G_1 denote the initial graph G , $G_i = G - \sum_{j < i} C_j$. Inspired by the Lemma 3.10, we will introduce the exponential demand for each graph G_i w.r.t the same node-weighting A . To

avoid clutter, let w_i denote the exponential distance weight w_h^α with respect to the graph G_i . We further use D_i to denote the corresponding exponential demand $D_{h,A}^i$ w.r.t graph G_i .

In the graph G_i , by Lemma 3.10 and the fact that $\text{spars}_{(h,s)}(C, A) \leq \phi$ (meaning there exists a symmetric h -length A -respecting demand D_i^* such that $\text{spars}_{hs}(C, D_i^*) \leq \phi$), we have

$$\text{sep}_{h,s/2}^i(C_i, D_i) \geq 2^{-8\alpha-1} \cdot \frac{|C_i|}{\phi}.$$

Further, we define a potential function $P_i : V \rightarrow \mathbb{R}$ w.r.t the graph G_i . It assigns a value to each vertex u with the amount of $P_i(u) = A(u) \ln(w_i(u))$. We note the fact that $w_i(u) \geq w_i(u, u) \geq 1$, which guarantees that $P_i(u) \geq 0$ for all i and vertices u .

Start with graph G_i , each vertex u will have potential $P_i(u)$. After applying cut C_i to the graph G_i , we first get the resulting graph G_{i+1} with same node-weighting A . we have $P_{i+1}(u) = A(u) \ln(w_{i+1}(u))$. Since we only increase the length of some edges in G_i , the exponential weight can only decrease between any vertex pairs. Consequently, there is always a decrease from $P_i(u)$ to $P_{i+1}(u)$, and we have

$$P_i(u) - P_{i+1}(u) = A(u) \cdot (\ln(w_i(u)) - \ln(w_{i+1}(u))) \quad (12)$$

$$= A(u) \cdot (-\ln(1 - (1 - \frac{w_{i+1}(u)}{w_i(u)}))) \quad (13)$$

$$\geq A(u) \cdot (1 - \frac{w_{i+1}(u)}{w_i(u)}) \quad (14)$$

$$\geq A(u) \cdot (\frac{w_i(u) - w_{i+1}(u)}{w_i(u)}) \quad (15)$$

We use the fact that $-\ln(1-x) \geq x$ when $0 \leq x < 1$ for inequality (14).

In the graph G_i , the distance of each demand pair that contributes to $\text{sep}_{h,s/2}^i(C_i, D_i)$ will be at least $\frac{h \cdot s}{2} > \frac{2h \log_2 n}{\alpha}$. In other words, let (u, v) be a demand pair that contributes $D_i(u, v)$ to the separation, we have that $\text{dist}_{G_{i+1}}(u, v) > \frac{2h \log_2 n}{\alpha}$ and $\overline{\text{dist}}_{G_{i+1}}(u, v) = \overline{\text{dist}}_{G_{i+1}}(v, u) > \frac{2h \log_2 n}{\alpha}$, so $w_h^{i+1}(u, v) = w_h^{i+1}(v, u) = 0$. This allows us to further lower bound the potential reduction as follows.

$$\begin{aligned} \sum_{u \in V} (P_i(u) - P_{i+1}(v)) &\geq \sum_{u \in V} A(u) \cdot \frac{w_i(u) - w_{i+1}(u)}{w_i(u)} \\ &= \sum_{\text{ordered } (u,v) \in V \times V} A(u) \cdot \frac{w_i(u, v) - w_{i+1}(u, v)}{w_i(u)} \\ &\geq \sum_{\substack{\text{ordered } (u,v) \in V \times V \text{ s.t.} \\ \overline{\text{dist}}_{G_{i+1}}(u,v) > hs/2}} A(u) \cdot \frac{w_i(u, v)}{w_i(u)} \\ &= \frac{1}{2} \text{sep}_{h,s/2}^i(C_i, D_i), \end{aligned}$$

where the last equality is by $D_i(u, v) = A(u) \cdot \frac{w_h^i(u, v)}{w_h^i(u)} + A(v) \cdot \frac{w_h^i(v, u)}{w_h^i(v)}$ for each $u, v \in V$. As a result, the overall potential reduction is at least

$$\sum_{u \in V} P_i(u) - P_{i+1}(u) \geq \frac{1}{2} \cdot \text{sep}_{h,s/2}^i(C_i, D_i) \geq 2^{-8\alpha-2} \cdot \frac{|C_i|}{\phi}$$

Finally, we can come to a conclusion over the summation of size of all cuts.

$$\begin{aligned}
\sum_i |C_i| &\leq 2^{8\alpha+2} \phi \cdot \sum_i \sum_{u \in V} (P_i(u) - P_{i+1}(u)) \\
&\leq 2^{8\alpha+2} \phi \cdot \sum_{u \in V} P_1(u) \\
&\leq 2^{8\alpha+2} \phi \cdot \sum_{u \in V} A(u) \ln(w_1(u)) \\
&\leq 2^{8\alpha+2} \phi \cdot |A| \cdot \ln n.
\end{aligned}$$

For the last inequality, we use that $\ln(w_1(u)) \leq \ln n$. This concludes the proof. \square

The upper bound over the size of the sequence of moving cuts directly implies the existence of length-constrained directed expander decompositions.

Theorem 3.9. *For any $G = (V, E)$, a node-weighting A , $h > 1$, $\alpha \geq 1$, $\phi < 1$ and a length slack parameter $s = O(\log n)$, there is a directed (h, s) -length ϕ -expander decomposition for A with cut slack $\kappa = O(n^{O(\frac{1}{s})} \log n)$.*

Proof. From graph G , if node-weighting A is already (h, s) -length ϕ -expanding in G , then we are done, because the empty cut is a valid expander decomposition. Otherwise, there exists an $h \cdot s$ -length cut C with $h \cdot s$ sparsity strictly smaller than ϕ . We take an arbitrary cut satisfying the above condition and denote it as C_1 . It is further applied to graph G to get $G_2 = G - C_1$. With one further step, if we assume that A is still not (h, s) -length ϕ -expanding in G_2 , we can find an $h \cdot s$ -length cut C_2 similarly as above. We update $G_3 = G_2 - C_2 = G - (C_1 + C_2)$ where $C_1 + C_2$ represents that we union two cuts together by summing the cut value on each edge. W.l.o.g we can assume that $C_1 + C_2$ is still an $h \cdot s$ -length cut since it is meaningless to make the length increase of an edge larger than $h \cdot s$ when we are considering the $h \cdot s$ -length sparsity.

We repeat the above procedure until we reach some integer k where A is (h, s) -length ϕ -expanding in $G_{k+1} = G - \sum_{j \leq k} C_j$. Let $C_{\leq k}$ denote the union of all such cuts, we can assume it is an $h \cdot s$ -length cut as discussed above. Then by definition $C_{\leq k}$ is a valid expander decomposition for G and A , and Lemma 3.17 guarantees that $|C_{\leq k}| = \sum_{j \leq k} |C_j| \leq 2^{8\alpha+2} \phi \cdot |A| \cdot \ln n$ as long as $s > \frac{4 \log_2 n}{\alpha}$. This gives that $\kappa \leq 2^{8\alpha+2} \cdot \ln n = O(n^{O(\frac{1}{s})} \log n)$. \square

4 Length-Constrained Vertex Expansion

In this section we extend the theory of length-constrained expander decomposition to vertex-capacitated graphs. See Section 2.2 for preliminaries of vertex-capacitated graphs.

The basic concepts of length-constrained vertex expansion are analogous to those of length-constrained directed expansion in Section 3.1. The major difference is that now a moving cut C can assign cut values to both vertices and edges. See Section 4.1 for formal description of the basic concepts.

The main results of this section is the existence of length-constrained expander decomposition for vertex-capacitated graphs (Theorem 4.1) and the routing characterization of length-constrained vertex expanders (Theorem 4.2).

Theorem 4.1 (Existential (h, s) -length Expander Decomposition for Vertex-Capacitated Graphs). *For any vertex-capacitated graph $G_{\text{vc}} = (V_{\text{vc}}, E_{\text{vc}})$, node-weighting A_{vc} , $h > 1$, $\phi < 1$ and a length slack parameter $s = O(\log n)$, there is an (h, s) -length ϕ -expander decomposition for A with cut slack $\kappa = O(n^{O(\frac{1}{s})} \log n)$.*

Theorem 4.2 (Routing Characterization of Length-Constrained Vertex Expanders). *Given a vertex-capacitated graph G_{vc} and node-weighting A_{vc} , for any $h \geq 1$, $\phi < 1$ and $s \geq 1$ we have:*

- *If A_{vc} is (h, s) -length ϕ -expanding in G_{vc} , then every h -length A_{vc} -respecting demand can be routed in G_{vc} with congestion at most $O(\frac{\log N}{\phi})$ and dilation at most $h \cdot s$.*
- *If A_{vc} is not (h, s) -length ϕ -expanding in G_{vc} , then some h -length A_{vc} -respecting demand cannot be routed with congestion at most $\frac{1}{6\phi}$ and dilation at most $\frac{h \cdot s}{2}$.*

In Section 4.2, we introduce a key reduction that transforms vertex-capacitated graphs into directed edge-capacitated graphs, demonstrating their equivalence. This equivalence is crucial for the proofs of Theorem 4.1 and Theorem 4.2 in Section 4.3 and Section 4.4.

4.1 Basic Concepts of Length-Constrained Vertex Expansion

We start with defining concepts related to length-constrained vertex expanders.

Moving Cuts for Vertex-Capacitated Graphs. A notable distinction for vertex-capacitated graphs is that moving cuts can be applied to vertices by exerting a similar length increase on them.

An h -length moving cut $C : V \cup E \rightarrow \{0, \frac{1}{h}, \dots, 1\}$ on a vertex-capacitated graph G assigns to each edge e and each vertex v a fractional cut value between zero and one which is a multiple of $\frac{1}{h}$. The size of C is defined as $|C| = \sum_e u_G(e) \cdot C(e) + \sum_v u_G(v) \cdot C(v)$. The length increase associated with the h -length moving cut C is denoted with $\ell_{C,h}$. Generalizing from the length increase over edges, the moving cut C assigns a vertex v length increase $\ell_{C,h}(v) = h \cdot C(v)$. We similarly define the degree of the vertex moving cut over a vertex v to be $\deg_C(v) = u_G(v) \cdot C(v) + \sum_{e:e \ni v} u_G(e) \cdot C(e)$.

By applying the cut C to a vertex-capacitated graph G , the resulting graph is $G - C$ where the length of each vertex and edge increases accordingly. To distinguish between two types of moving cuts, we will clarify the type of graph where the moving cut is applied.

Remark 4.3. *From the setting of vertex capacity, we note that the undirected edge-capacitated graph is a special case for the vertex-capacitated graph. We can reduce an arbitrary undirected edge-capacitated graph to a vertex-capacitated graph by setting the length of vertices to some small*

constant and allowing arbitrarily large capacity over vertices. Then it is too expensive to have a fractional cut over any vertices. Thus if we generalize previous results to vertex-capacitated graphs, we actually build up a more general framework for length-constrained expanders and expander decompositions.

(h, s) -Length Sparsity for Vertex-Capacitated Graphs. The definition of sparsity of moving cuts, and the conductance of the node-weighting similarly generalize from the edge cut cases. It may be useful for the reader to recall the definition in Section 3.1.

We remark that for undirected vertex-capacitated graphs, the (h, s) -length sparsity is no longer restricted to symmetric demands.

$$\text{spars}_{(h,s)}(C, A) = \min_{A\text{-respecting } h\text{-length demand } D} \text{spars}_{h \cdot s}(C, D).$$

As mentioned in Section 3.1, we say that A is (h, s) -length ϕ -expanding in G if $\text{cond}_{(h,s)}(A) \geq \phi$. The definition of length-constrained expander decompositions for vertex-capacitated graphs follows.

Definition 4.4 (Length-Constrained Vertex Expander Decomposition). *Given a vertex-capacitated graph G , an (h, s) -length ϕ -expander decomposition for a node-weighting A with length slack s and cut slack κ is an $h \cdot s$ -length cut C of size at most $\kappa \cdot \phi |A|$ such that A is (h, s) -length ϕ -expanding in $G - C$.*

4.2 Reduction to Directed Edge-Capacitated Graphs




Figure 3: (Left) A vertex-capacitated graph. (Right) The corresponding directed edge-capacitated graph.

In Section 3, we presented several results concerning directed edge-capacitated graphs, including the existence of expander decompositions. It further turns out that we can reduce an undirected vertex-capacitated graph to a directed edge-capacitated graph to apply those results. This is an essential step to prove previous expansion results for vertex-capacitated graphs, and we start with a detailed introduction to the reduction.

Namely, for any vertex-capacitated graph $G_{vc} = (V_{vc}, E_{vc})$ with a node-weighting A_{vc} , one can construct a directed edge-capacitated graph $G_{ec} = (V_{ec}, E_{ec})$ with a corresponding node-weighting A_{ec} through the following reduction:

1. Let $V_{ec} = \emptyset$, $E_{ec} = \emptyset$ and $A_{ec}(v) = 0$ for all vertices at the beginning.
2. For each vertex $v \in V_{vc}$, add three vertices v_{in} , v_{mid} and v_{out} to V_{ec} ; Add (v_{in}, v_{mid}) , (v_{mid}, v_{out}) and (v_{in}, v_{out}) to E_{ec} . The length of each edge is set as $\ell_{G_{vc}}(v)$, and the capacity of each edge is set as $u_{G_{vc}}(v)$. We call such edges inner edges.

3. For each edge $e = (u, v) \in E_{vc}$, add (u_{out}, v_{in}) and (v_{out}, u_{in}) to E_{ec} . The length of them is set as $l_{G_{vc}}(e)$ and the capacity is set as $u_{G_{vc}}(e)$. We call such edges outer edges.
4. For each vertex $v \in \text{supp}(A_{vc})$, set $A_{ec}(v_{mid}) = A_{vc}(v)$.

An example is shown in Figure 3. In the right part, inner edges and outer edges are colored red and blue respectively. In what follows, we use G_{vc} to denote an arbitrary vertex-capacitated graph, and G_{ec} always refers to the edge-capacitated graph constructed from G_{vc} as described above. For simplicity, we may occasionally omit to mention G_{vc} , but whenever G_{ec} is referenced, it is understood to be based on a specific vertex-capacitated graph.

Also in the undirected setting, the direction of demands does not make a difference. Thus for simplicity, when we talk about the routing, we can stick to a symmetric demand D in both types of graphs. Namely, for any demand D in G_{vc} , we can have an equivalent symmetric demand D' by balancing the demand in both directions. And we further use the same notation D' in G_{ec} where $D'(u_{mid}, v_{mid}) = D'(v_{mid}, u_{mid}) = D'(u, v)$ whenever $D'(u, v) > 0$ for any pair of vertices $u, v \in V_{vc}$.

Reduction from Vertex-Capacitated to Directed Edge-Capacitated Graphs. As mentioned before, the motivation to construct a directed edge-capacitated graph G_{ec} that shares a similar structure as the vertex-capacitated graph G_{vc} is that we can transform problems into the directed and edge-capacitated setting which we have explored in Section 3. To apply our previous results to the vertex-capacitated case, it is essential to elucidate the connection between the two types of graphs. Specifically, we need to show the equivalence of expansion between two types of graphs in our construction as follows:

Theorem 4.5. *Let $G_{vc} = (V_{vc}, E_{ec})$ be a vertex-capacitated graph with node-weighting A_{vc} , and $G_{ec} = (V_{ec}, E_{ec})$ be the corresponding edge-capacitated graph with node-weighting A_{ec} .*

1. *If A_{vc} is (h, s) -length ϕ -expanding in G_{vc} , then A_{ec} is (h, s) -length ϕ -expanding in G_{ec} .*
2. *If A_{ec} is (h, s) -length ϕ -expanding in G_{ec} , then A_{vc} is (h, s) -length $\frac{\phi}{3}$ -expanding in G_{vc} .*

We develop the theorem gradually with some necessary lemmas concerning the equivalence of sparse cut in both types of graphs.




Figure 4: Suppose there is a path $\{u, v, w\}$ in the vertex-capacitated graph G_{vc} (Left), then we can follow it and create a path $\{u_{mid}, u_{out}, v_{in}, v_{out}, w_{in}, w_{mid}\}$ in G_{ec} (Right). We note that both paths have the same length.

We first note an important property of our construction is that the equivalence of pairwise distance is kept.

Lemma 4.6. *Given a vertex-capacitated graph $G_{vc} = (V_{vc}, E_{vc})$ and the corresponding edge-capacitated graph $G_{ec} = (V_{ec}, E_{ec})$, for any $u, v \in V_{vc}$, $\text{dist}_{G_{vc}}(u, v) = \text{dist}_{G_{ec}}(u_{mid}, v_{mid}) = \text{dist}_{G_{ec}}(v_{mid}, u_{mid})$.*

Proof. For any shortest path between u and v in G_{vc} , it is possible to give a path of the same length in G_{ec} between u_{mid} and v_{mid} by following the trajectory. For the other direction, a similar argument also applies for the shortest path between u_{mid} and v_{mid} because it always goes directly from some vertex a_{in} to a_{out} to achieve shorter length. An example is shown in Figure 4. We further note that the directed distance between u_{mid} and v_{mid} is symmetric due to the construction, and this gives the lemma. \square

As mentioned above, we then discuss the equivalence of moving cuts between vertex-capacitated graphs and directed edge-capacitated graphs. We note that it would be convenient to normalize the cut in graph G_{ec} for simplicity. Actually, in G_{ec} , for any h -length moving cut C with a fully separated demand D such that $\text{spars}_h(C, D) = \frac{|C|}{|D|} \leq \phi$, we can construct a normalized cut C' as follows:

1. For any group of vertices v_{in} , v_{mid} and v_{out} , C' cut the same length over inner edges between them by taking the maximum cut value over those edges from C , i.e.,

$$\begin{aligned} C'((v_{in}, v_{mid})) &= C'((v_{mid}, v_{out})) = C'((v_{in}, v_{out})) \\ &= \max\{C((v_{in}, v_{mid})), C((v_{mid}, v_{out})), C((v_{in}, v_{out}))\} \end{aligned}$$

2. For any pair of outer edges (u_{out}, v_{in}) and (v_{out}, u_{in}) , C' also cuts the same length over them by taking the larger cut value over two edges from C , i.e.,

$$C'((u_{out}, v_{in})) = C'((v_{out}, u_{in})) = \max\{C((u_{out}, v_{in})), C((v_{out}, u_{in}))\}$$

Cut C' is normalized in the sense that the cut value over the same group edges is the same, and thus it easily induces a cut in the base graph G_{vc} . What is more, we do not lose more than a constant factor in terms of sparsity.

Lemma 4.7. *Given the constructed graph G_{ec} and the h -length moving cut C with a fully separated demand D such that $\text{spars}_h(C, D) = \frac{|C|}{|D|} \leq \phi$, the normalized cut C' has sparsity $\text{spars}_h(C', D) \leq 3 \text{spars}_h(C, D) \leq 3\phi$.*

Proof. We first note that from our construction of C' , the size of C' only triples at most, which means $|C'| \leq 3|C|$. Further, since we only increase the cut length, the separation value w.r.t D cannot be less and $\text{sep}_h(C', D) \geq \text{sep}_h(C, D)$. This over all shows that

$$\text{spars}_h(C', D) = \frac{|C'|}{\text{sep}_h(C', D)} \leq \frac{3|C|}{\text{sep}_h(C, D)} = 3 \text{spars}_h(C, D) \leq 3\phi$$

\square

With the help of cut normalization, we further establish the equivalence of (h, s) -length sparsity between two types of graphs. In general, if a sparse cut exists in the edge-capacitated graph, we can construct a sparse normalized cut and transfer it to the vertex-capacitated graph while preserving the cut's sparsity. A similar argument also applies to the other direction.

Lemma 4.8. *Let $G_{vc} = (V_{vc}, E_{ec})$ be a vertex-capacitated graph with node-weighting A_{vc} , and $G_{ec} = (V_{ec}, E_{ec})$ be the corresponding edge-capacitated graph with node-weighting A_{ec} .*

1. *If there is an $h \cdot s$ -length moving cut C in vertex-capacitated G_{vc} such that $\text{spars}_{(h,s)}(C, A_{vc}) < \phi$, then there exists an $h \cdot s$ -length moving cut C' in G_{ec} such that $\text{spars}_{(h,s)}(C', A_{ec}) < 3\phi$.*
2. *If there is an $h \cdot s$ -length moving cut C' in edge-capacitated G_{ec} such that $\text{spars}_{(h,s)}(C', A_{ec}) < \phi$, then there exists an $h \cdot s$ -length moving cut C in G_{vc} such that $\text{spars}_{(h,s)}(C, A_{vc}) < \phi$.*

Proof. We will prove Item 1 and Item 2 separately.

Proof of Item 1. Since $\text{spars}_{(h,s)}(c, A_{\text{vc}}) < \phi$, let

$$D^* = \arg \min_{A_{\text{vc}}\text{-respecting } h\text{-length demand } D} \text{spars}_{h,s}(C, D)$$

then we have $\text{spars}_{h,s}(C, D^*) < \phi$. W.l.o.g, we can assume D^* is symmetric and fully separated. As mentioned in the reduction, this immediately gives an A_{ec} -respecting symmetric h -length demand D^* in G_{ec} as well. To see that D^* is also h -length in G_{ec} , we note that from Lemma 4.6, we have

$$\text{dist}_{G_{\text{ec}}}(u_{\text{mid}}, v_{\text{mid}}) = \text{dist}_{G_{\text{ec}}}(v_{\text{mid}}, u_{\text{mid}}) = \text{dist}_{G_{\text{vc}}}(u, v)$$

Then we construct a sparse cut C' based on C , and show that the sparsity of C' w.r.t D^* is also dependent on ϕ . For any vertex $u \in V_{\text{vc}}$ where $C(u) > 0$, we set

$$C'((u_{\text{in}}, u_{\text{mid}})) = C'((u_{\text{mid}}, u_{\text{out}})) = C'((u_{\text{in}}, u_{\text{out}})) = C(u)$$

For any $e = (u, v) \in E_{\text{vc}}$ where $C(e) > 0$, we set

$$C'((u_{\text{out}}, v_{\text{in}})) = C'((v_{\text{out}}, u_{\text{in}})) = C(e)$$

From the construction, we have that $|C'| \leq 3|C|$. Then we show that D^* is also fully separated by C' . We know that for any vertex pairs $u, v \in V_{\text{vc}}$ where $D^*(u, v) > 0$, the distance between them is bounded by h in G_{vc} but at least $h \cdot s$ after applying cut in $G_{\text{vc}} - C$, i.e., $\text{dist}_{G_{\text{vc}}}(u, v) \leq h$ and $\text{dist}_{G_{\text{vc}}-C}(u, v) > h \cdot s$. In the edge-capacitated graph G_{ec} , the key observation is that the equivalence of distance is still kept. We still have $\text{dist}_{G_{\text{vc}}-C}(u, v) = \text{dist}_{G_{\text{ec}}-C'}(u_{\text{mid}}, v_{\text{mid}})$, because we set the length increase over edges in G_{ec} same as the corresponding vertices or edges in the original graph G_{vc} . Namely, if we construct an edge-capacitated graph based on $G_{\text{vc}} - C$, we get exactly $G_{\text{ec}} - C'$. Again from Lemma 4.6, this implies that $\text{dist}_{G_{\text{ec}}}(u_{\text{mid}}, v_{\text{mid}}) \leq h$ and $\text{dist}_{G_{\text{ec}}-C'}(u_{\text{mid}}, v_{\text{mid}}) > h \cdot s$, which means that D^* is also fully separated by C' in G_{ec} . Thus we have $\text{sep}_{h,s}(C, D^*) = \text{sep}_{h,s}(C', D^*)$. We can conclude that

$$\begin{aligned} \text{spars}_{(h,s)}(C', A_{\text{ec}}) &= \min_{A_{\text{ec}}\text{-respecting } h\text{-length symmetric demand } D} \text{spars}_{h,s}(C', D) \\ &\leq \text{spars}_{h,s}(C', D^*) \\ &= \frac{|C'|}{\text{sep}_{h,s}(C', D^*)} \\ &\leq \frac{3|C|}{\text{sep}_{h,s}(C, D^*)} \\ &\leq 3\phi \end{aligned}$$

Proof of Item 2. Since $\text{spars}_{(h,s)}(C', A_{\text{ec}}) < \phi$, let

$$D^* = \arg \min_{A_{\text{ec}}\text{-respecting } h\text{-length symmetric demand } D} \text{spars}_{h,s}(C', D)$$

From Lemma 4.7, we know there exists a normalized $h \cdot s$ -length cut C_n from C' . We note that the separation from C_n is at least as large as that from cut C' , i.e.

$$\text{sep}_{h,s}(C_n, D^*) \geq \text{sep}_{h,s}(C', D^*)$$

Similarly, we construct a $h \cdot s$ -length moving cut C in the vertex-capacitated graph G_{vc} . Namely, for any vertex $u \in V_{\text{vc}}$, we set $C(u) = C_n((u_{\text{in}}, u_{\text{mid}}))$. For any edge $e = (u, v) \in E_{\text{vc}}$, we set $C(e) = C_n((u_{\text{out}}, v_{\text{in}}))$. From the property of C_n that the length increase is the same

among the same group of inner edges and outer edges, we immediately have $|C| \leq \frac{1}{2}|C_n|$. However, the construction is equivalent to that we assign the maximum value of the cut length among the same group of inner edges or the pair of outer edges from C' to the corresponding vertex or edge in G_{vc} respectively. This actually shows that $|C| \leq |C'|$, which is a tighter bound. From a similar argument as above, we can show that the demand D^* is A_{vc} -respecting and h -length in G_{vc} as well. From the construction of C and Lemma 4.6, we have $\text{seph}_{h,s}(C, D^*) = \text{seph}_{h,s}(C_n, D^*) \geq \text{seph}_{h,s}(C', D^*)$. We further have

$$\begin{aligned} \text{spars}_{(h,s)}(C, A_{vc}) &= \min_{A_{vc}\text{-respecting } h\text{-length demand } D} \text{spars}_{h,s}(C, D) \\ &\leq \text{spars}_{h,s}(C, D^*) \\ &= \frac{|C|}{\text{seph}_{h,s}(C, D^*)} \\ &\leq \frac{|C'|}{\text{seph}_{h,s}(C', D^*)} \\ &\leq \phi \end{aligned}$$

□

We summarize the equivalence of (h, s) -length sparsity with the main theorem in this section.

Theorem 4.5. *Let $G_{vc} = (V_{vc}, E_{vc})$ be a vertex-capacitated graph with node-weighting A_{vc} , and $G_{ec} = (V_{ec}, E_{ec})$ be the corresponding edge-capacitated graph with node-weighting A_{ec} .*

1. *If A_{vc} is (h, s) -length ϕ -expanding in G_{vc} , then A_{ec} is (h, s) -length ϕ -expanding in G_{ec} .*
2. *If A_{ec} is (h, s) -length ϕ -expanding in G_{ec} , then A_{vc} is (h, s) -length $\frac{\phi}{3}$ -expanding in G_{vc} .*

Proof. Given A_{vc} is (h, s) -length ϕ -expanding in G_{vc} , suppose A_{ec} is not (h, s) -length ϕ -expanding in G_{ec} , then there exists a cut C' in G_{ec} such that $\text{spars}_{(h,s)}(C', A_{ec}) < \phi$. From Lemma 4.8, we can find a cut C in G_{vc} where $\text{spars}_{(h,s)}(C, A_{vc}) < \phi$. This contradicts that A_{vc} is (h, s) -length ϕ -expanding in G_{vc} .

A similar argument applies to the other direction. □

4.3 Existential Proof of the Decomposition

In this section, we present one of the key results, namely the existence of an expander decomposition for undirected graphs with vertex capacity. Following the reduction introduced before, we apply Theorem 3.9 to the edge-capacitated graph G_{ec} to obtain the expander decomposition and then translate the cut into the vertex-capacitated graph. We still need to demonstrate that the translated cut forms a valid expander decomposition for G_{vc} , and the resulting cut has a favorable size.

Theorem 4.1 (Existential (h, s) -length Expander Decomposition for Vertex-Capacitated Graphs). *For any vertex-capacitated graph $G_{vc} = (V_{vc}, E_{vc})$, node-weighting A_{vc} , $h > 1$, $\phi < 1$ and a length slack parameter $s = O(\log n)$, there is an (h, s) -length ϕ -expander decomposition for A with cut slack $\kappa = O(n^{O(\frac{1}{s})} \log n)$.*

Proof. Given the vertex-capacitated graph G_{vc} and the node-weighting A_{vc} , we construct the corresponding edge-capacitated graph G_{ec} and node-weighting A_{ec} as defined before. Whenever A_{ec} is not (h, s) -length ϕ -expanding in G_{ec} , we can find a sparse cut in G_{ec} . We can always

normalize such a cut in G_{ec} as shown in Lemma 4.7, and apply it to the graph G_{ec} . This process can be done iteratively until the resulting graph does not admit a sparse cut. This follows the idea from Lemma 3.17 and Theorem 3.9. We note that we do not apply the normalization to the union of all sparse cuts because the normalization will change the conductance of the node-weighting in the graph applied with the cut.

Specifically, from graph $G_{\text{ec}}^{(1)} = G_{\text{ec}}$, if A_{ec} is already (h, s) -length 3ϕ -expanding in G_{ec} , then we are done. Otherwise, we take an arbitrary $h \cdot s$ -length cut C' with $h \cdot s$ sparsity strictly smaller than 3ϕ w.r.t the node-weighting A_{ec} , and from Lemma 4.7, we can construct a normalized cut $C_{\text{ec}}^{(1)}$. We note that the normalization only increases the length, thus for any A_{ec} -respecting demand D , we have $\text{sep}_{h \cdot s}(C_{\text{ec}}^{(1)}, D) \geq \text{sep}_{h \cdot s}(C', D)$. Considering the size of the normalized cut can increase by up to threefold, this gives that $\text{spars}_{h \cdot s}(C_{\text{ec}}^{(1)}, A_{\text{ec}}) \leq 9\phi$. We then apply $C_{\text{ec}}^{(1)}$ to graph G_{ec} to update $G_{\text{ec}}^{(2)} = G_{\text{ec}} - C_{\text{ec}}^{(1)}$.

We repeat the above procedure, until we reach to some integer k where A_{ec} is (h, s) -length 3ϕ -expanding in $G_{\text{ec}}^{(k+1)} = G_{\text{ec}} - \sum_{j \leq k} C_{\text{ec}}^{(j)}$. We take the union of all these cuts and set a threshold of $h \cdot s$ over the length increase of every edge and vertex. We name the resulting $h \cdot s$ -length cut C_{ec} , and by definition, C_{ec} is a valid expander decomposition for the edge-capacitated graph G_{ec} and A_{ec} . From a similar analysis from Lemma 3.17, we can demonstrate that an upper bound still exists on the summation of all these cut sizes with the potential function technique. We have $|C_{\text{ec}}| \leq O(2^{8\alpha} \log n) \cdot \phi |A_{\text{ec}}|$. What is more important is that C_{ec} is normalized.

From C_{ec} , we can construct a corresponding vertex cut C_{vc} . Namely, for vertex $u \in V_1$, we assign $C_{\text{vc}}(u) = C_{\text{ec}}((u_{\text{in}}, u_{\text{mid}}))$ which is the same among the same group of inner edges. For edge $e = (u, v) \in E_{\text{vc}}$, we assign $C_{\text{vc}}((u, v)) = C_{\text{ec}}((u_{\text{out}}, v_{\text{in}}))$ which is the same between the pair of outer edges. We apply C_{vc} to G_{vc} , and we get the resulting graph $G_{\text{vc}} - C_{\text{vc}}$. We observe that the edge-capacitated graph $G_{\text{ec}} - C_{\text{ec}}$ exactly corresponds to the graph $G_{\text{vc}} - C_{\text{vc}}$. We further note that the node-weighting A_{vc} corresponds to A_{ec} due to our construction at the beginning. Since we have shown A_{ec} to be (h, s) -length 3ϕ -expanding in $G_{\text{ec}} - C_{\text{ec}}$, from Theorem 4.5, we have that A_{vc} is ϕ -expanding in $G_{\text{vc}} - C_{\text{vc}}$. From that $|C_{\text{vc}}| \leq \frac{1}{2}|C_{\text{ec}}|$ and $|A_{\text{vc}}| = |A_{\text{ec}}|$, we also have $|C_{\text{vc}}| \leq O(2^{8\alpha} \log n) \cdot \phi |A_{\text{vc}}|$. This concludes the theorem similarly as Theorem 3.9. \square

4.4 Routing Characterization of Length-Constrained Vertex Expansion

Furthermore, due to the structural similarity introduced by the reduction, an efficient routing for a given demand in one type of graph naturally translates into a corresponding favorable routing in the other. This reduction then facilitates the transfer of routing characterizations from length-constrained directed expansion to vertex-capacitated graphs as follows:

Theorem 4.2 (Routing Characterization of Length-Constrained Vertex Expanders). *Given a vertex-capacitated graph G_{vc} and node-weighting A_{vc} , for any $h \geq 1$, $\phi < 1$ and $s \geq 1$ we have:*

- If A_{vc} is (h, s) -length ϕ -expanding in G_{vc} , then every h -length A_{vc} -respecting demand can be routed in G_{vc} with congestion at most $O(\frac{\log N}{\phi})$ and dilation at most $h \cdot s$.
- If A_{vc} is not (h, s) -length ϕ -expanding in G_{vc} , then some h -length A_{vc} -respecting demand cannot be routed with congestion at most $\frac{1}{6\phi}$ and dilation at most $\frac{h \cdot s}{2}$.

Proof. We first construct a corresponding edge-capacitated graph G_{ec} with node-weighting A_{ec} as describe before. Since A_{vc} is (h, s) -length ϕ -expanding in G , from Theorem 4.5, we have that A_{ec} is also (h, s) -length ϕ -expanding in G_{ec} . From Theorem 3.8, we have that every h -length A_{ec} -respecting symmetric demand can be routed in G_{ec} with congestion at most $O(\frac{\log N}{\phi})$ and dilation at most $h \cdot s$. We will exploit the routing in G_{ec} to route demands in the vertex-capacitated graph.

Namely, for any h -length A_{vc} -respecting demand D , we can assume it is symmetric. As mentioned before, D is also an h -length A_{ec} -respecting symmetric demand in the edge-capacitated graph. Then there exists a flow F_{ec} in G_{ec} that witnesses the routing with desired congestion and dilation. Consider any $(u_{\text{mid}}, v_{\text{mid}})$ -flow path P_{ec} of F_{ec} , we can construct a corresponding (u, v) -flow path P_{vc} by following the P_{ec} as shown in Figure 4. The observation is that for any edge in G_{ec} with u_{out} as the head vertex, it ends at some vertex v_{in} . And from the vertex v_{in} , any path must arrive at v_{out} in either one or two steps. Thus we can record groups of vertices that are passed by P_{ec} , and create the path P_{vc} by traversing vertices in G_{vc} corresponding to groups of vertices in G_{ec} one by one. In general, we construct a corresponding flow F_{vc} path in G_{vc} by following the flow path of F_{ec} . From the length and capacity property of the reduction, flow F_{vc} in G_{vc} indeed routes D with at most the same length and up to three times congestion compared to the original congestion, which is still $O(\frac{\log N}{\phi})$.

If A_{vc} is not (h, s) -length ϕ -expanding in G_{vc} , from Theorem 4.5, we have that A_{ec} cannot be (h, s) -length 3ϕ -expanding in G_{ec} . Then there exists an h -length A_{ec} -respecting demand D such that it cannot be routed with congestion at most $\frac{1}{6\phi}$ and length at most $\frac{h \cdot s}{2}$.

Similarly, we have an h -length A_{vc} -respecting demand D in G_{vc} . Suppose D can be routed in G_{vc} with congestion at most $\frac{1}{6\phi}$ and length at most $\frac{h \cdot s}{2}$, and the routing is witnessed by a flow F_{vc} . We construct a flow F_{ec} in G_{ec} by similarly following the trajectory of each flow path in F_{vc} . The resulting flow F_{ec} witnesses a routing for D with at most the same congestion and the same length. But this gives a contradiction. Thus we also find a h -length A_{vc} -respecting demand D that cannot be routed in G_{vc} with congestion at most $\frac{1}{6}\phi$ and length at most $\frac{h \cdot s}{2}$. \square

5 Length-Constrained Vertex-Capacitated Flow Shortcuts

In this section, we show the existence of LC-flow shortcuts in vertex-capacitated graphs. The Definition 5.1 below generalizes Definition 1.1 of LC-flow shortcuts in the sense that an additional length parameter h is given and the forward mapping only holds for demands routable in G with congestion 1 and length at most h .

Definition 5.1 (Length-Constrained Flow Shortcut). *Given a graph $G = (V, E)$, we say an edge set E' (possibly with endpoints outside V) is an t -step h -LC-flow shortcut of G with length slack λ and congestion slack κ if*

- (Forward Mapping) for every demand D routable in G with congestion 1 and length $h' \leq h$, D is routable in $G \cup E'$ with congestion 1, length $\lambda h'$, and maximum step t , and
- (Backward Mapping) for every demand D on $V(G)$ routable in $G \cup E'$ with congestion 1 and length h' , D is routable in G with congestion κ and length h' .

We note that in [HHL⁺24] there is an analogous but weaker definition of h -LC-flow shortcut, in which the forward mapping only guarantee that D is routable in $G \cup E'$ with length λh instead of $\lambda h'$ (and the same congestion and step). That is, the length slack in Definition 5.1 is *competitive* in the sense that it upper bounds the ratio between the lengths of the shortcut flow and the original flow. Hence, by choosing a sufficiently large h , the total length of vertices and edges in G , an h -LC-flow shortcut is automatically an LC-flow shortcut.

Theorem 5.2. *Given a vertex-capacitated graph G with parameters $\epsilon = \Omega(\frac{1}{\log n})$, there exists a t -step LC-flow shortcut E' with length slack $O(1/\epsilon^3)$, congestion slack $O(n^{O(\epsilon)} \log^3 n / \epsilon^2)$, $t = 2^{O(\frac{1}{\epsilon})}$ and size $|E'| \leq O(n^{1+O(\epsilon)} \log n / \epsilon^2)$.*

Theorem 5.2 is the main theorem of this section. In what follows, actually we will focus on constructing h -LC-flow shortcut. Setting $h = (m + n)N$ gives the LC-flow shortcut in Theorem 5.2.

5.1 The Construction

Algorithm 1 LC-FlowShortcut(G, ϵ, h)

- 1: Initialize $A_0 = u_{V(G)}$, where $u_{V(G)}$ denotes the vertex capacity function of G .
 - 2: Initialize $s = 1/\epsilon$, $\phi = 1/(n^\epsilon \kappa)$ ($\kappa = O(n^{O(1/s)} \log n)$ is the cut slack from Theorem 4.1).
 - 3: Initialize $i \leftarrow 0$.
 - 4: **while** $|A_i| > 0$ **do**
 - 5: **for** j from 1 to $\lceil \log_2 h \rceil$ **do**
 - 6: $h_j \leftarrow 2^j$, $h_{\text{cov},j} = 4h_j$, $h_{\text{diam},j} = h_{\text{cov},j} \cdot s$.
 - 7: $C_{i+1,j} \leftarrow$ an $(h_{\text{diam},j}, s)$ -length ϕ -expander decomposition of A_i in G by Theorem 4.1.
 - 8: $\mathcal{N}_{i,j} \leftarrow$ a neighborhood cover with covering radius $h_{\text{cov},j}$, diameter $h_{\text{diam},j}$ in $G - C_{i+1,j}$ by Theorem 2.1.
 - 9: $H_{i,j} = \bigcup_{S \in \mathcal{N}_{i,j}} H_S$, where H_S is the $h_j s$ -length A_i -capacitated star graph on S .
 - 10: **end for**
 - 11: $A_{i+1} = \sum_j \frac{h_{\text{diam},j} \cdot s}{h_j} \cdot \deg_{C_{i+1,j}} = 4s^2 \cdot \sum_j \deg_{C_{i+1,j}}$.
 - 12: **end while**
 - 13: Return $E' = \bigcup_{i,j} E(H_{i,j})$.
-

The construction of the flow shortcut graph G' is given by Algorithm 1. The star graphs in Line 9 are formally defined in Definition 5.3.

Definition 5.3 (Star Graphs). *Given a graph G with a node-weighting A and a length parameter h , the h -length A -capacitated star graph on some $S \subseteq V(G)$, denoted by H_S , has*

$$V(H_S) = (\text{supp}(A) \cap S) \cup \{r_S\} \text{ and } E(H_S) = \{(v, r_S) \mid v \in V(H_S) \setminus \{r_S\}\},$$

where the vertex r_S is a Steiner vertex serving as the center and $V(H_S) \setminus \{r_S\}$ are original vertices. The length and capacity of each original vertex is unchanged, while r_S has length 1 and capacity $\sum_{v \in S} A(v)$. Each edge (v, r_S) has length h and capacity $A(v)$.

In short, Algorithm 1 mainly constructs an length-constrained expander hierarchy $\{A_i, C_{i+1,j} \mid 0 \leq i \leq d, 1 \leq j \leq \lceil \log_2 h \rceil\}$, where d is the largest i such that $|A_i| > 0$. We point out that $C_{d+1,j}$ is a zero cut for all j . Then the shortcut graph G' is obtained by adding star graphs on neighborhoods of each LC-expander $G - C_{i+1,j}$.

We remark that we do LC-expander decompositions with different length parameters h_j at one level because we aim at a shortcut graph with length slack significantly smaller than its step bound. Intuitively, if we only use LC-expanders with length parameter around h to shortcut an original h -length flow path P , then inevitably each step will have length around h , which means the length slack cannot go far below the number of steps. Now, providing LC-expanders with different length parameters, when we want to shortcut a subpath of P with length h' far smaller than h , we can choose the appropriate LC-expander to obtain a shortcut with length around h' instead of h . Another benefit is that this automatically gives a competitive length slack (this is why in Definition 5.1 we define the length slack of h -LC-flow shortcut to be competitive).

We first argue the size bound of E' . Observe that, in Line 7, we have $|C_{i+1,j}| \leq \kappa \phi |A_i| \leq |A_i|/n^\epsilon$ by Theorem 4.1. In Line 8, the width of each neighborhood cover $\mathcal{N}_{i,j}$ is $\omega = n^{O(1/s)} s = n^{O(\epsilon)}/\epsilon$ by Theorem 2.1. Furthermore, because in Line 11 we have $|A_{i+1}| \leq 4s^2 \sum_{j'} |C_{i+1,j'}| \leq 4s^2 \log h |A_i|/n^\epsilon = O(\log N/(\epsilon^2 n^\epsilon)) |A_i|$, we can upper bound d by $d \leq O(\log |A_0|/\log(\epsilon^2 n^\epsilon/\log N)) = O(1/\epsilon)$. Finally, by the algorithm, we have

$$|E'| \leq O(d \log h) \cdot \omega \cdot n = O(n^{1+O(\epsilon)} \log n/\epsilon^2).$$

Next we show the quality of the shortcut. Before that, we introduce a helper lemma Lemma 5.4, which shows the demands that each $H_{i,j}$ can route within small steps.

Lemma 5.4. *For each i, j , any demand \hat{D} that is $h_{\text{cov},j}$ -length in $G - C_{i+1,j}$, A_i -respecting and $u_{V(G)}$ -respecting can be routed in $H_{i,j}$ with length $2h_{\text{diam},j}s + 1$, congestion 1 and step 2.*

Proof. Recall that $H_{i,j} = \bigcup_{S \in \mathcal{N}_{i,j}} H_S$. For each demand pair $(u, v) \in \text{supp}(\hat{D})$, we will assign it in H_S , where $S \in \mathcal{N}_{i,j}$ is an arbitrary cluster containing both u and v . Note that such an S must exist because \hat{D} is $h_{\text{cov},j}$ -length in $G - C_{i+1,j}$ and $\mathcal{N}_{i,j}$ has covering radius $h_{\text{cov},j}$ in $G - C_{i+1,j}$. Now, for each $S \in \mathcal{N}_{i,j}$, its assigned demand \hat{D}_S has $V(\hat{D}_S) \subseteq S$ and it is A_i -respecting and $u_{V(G)}$ -respecting, so \hat{D}_S can be routed in H_S with length $2h_{\text{diam},j}s + 1$, congestion 1 and step 2 by the definition of H_S . \square

Now we show that the shortcut E' constructed by Algorithm 1 has length slack $O(1/\epsilon^3)$, congestion slack $n^{O(\epsilon)}$ and step $2^{O(1/\epsilon)}$. To do this, it suffices to show the quality of the forward mapping and backward mapping, i.e. Lemma 5.5 and Lemma 5.6, whose proofs are given in Section 5.2 and Section 5.3. Let $G' = G \cup E'$ be the shortcut graph.

Lemma 5.5 (Forward Mapping). *For any feasible h -length flow F in G , there is a feasible flow F' routing $\text{Dem}(F)$ in G' with $\text{leng}(F') \leq \text{leng}(F) \cdot O(1/\epsilon^3)$ and $\text{step}(F') \leq 2^{O(1/\epsilon)}$.*

Lemma 5.6 (Backward Mapping). *For any feasible flow F' in G' such that $V(\text{Dem}(F')) \subseteq V(G)$, there is a flow F routing $\text{Dem}(F')$ in G with $\text{leng}(F) \leq \text{leng}(F')$ and $\text{cong}(F) \leq n^{O(\epsilon)}$.*

5.2 Forward Mapping: Proof of Lemma 5.5

We will employ a top-down argument. Start from the top level d . At the beginning of processing a level $i \geq 0$, we are given a feasible flow F_i with the following invariant: each flow path $P \in \text{path}(F_i)$ has $\text{len}(P, G - C_{i+1,j}) \leq h_{\text{cov},j} = 4h_j$, where j is the minimum index such that $h_j \geq \text{len}(P, G)$ (which means $h_j/2 \leq \text{len}(P, G) \leq h_j$). Initially at the top level d , we set $F_d = F$. Note that F_d satisfies the invariant above because $C_{d+1,j}$ is a zero cut for any j .

First, at the bottom level $i = 0$, we can easily shortcut every flow path in F_0 . For each star graph $H_{0,j}$, we assign it a demand $\hat{D}_{0,j}$ which sums over $\text{Dem}(P)$ for each flow path $P \in \text{path}(F_0)$ such that j is the minimum index with $h_j \geq \text{len}(P, G)$. Observe that each $\hat{D}_{0,j}$ is an $h_{\text{cov},j}$ -length in $G - C_{1,j}$ and A_0 -respecting, so it can be routed in $H_{0,j}$ with length $2h_{\text{diam},j}s + 1$, congestion 1 and step 2 by Lemma 5.4.

From now on we consider levels $i \geq 1$. When processing a level i , for each flow path $P \in \text{path}(F_i)$, we may shortcut some subpaths of P using shortcut edges in H . The subpaths of P that have not been shortcut will be added to F_{i-1} , meaning that they are deferred to lower levels to get shortcut. At the end, the final F_{i-1} should ensure the invariants above, and we proceed to the lower level $i - 1$.

Now we will explain the shortcut at a level $i \geq 1$ in detail. Fix a flow path $P \in \text{path}(F_i)$. Let j be the minimum index such that $h_j \geq \text{len}(P, G)$. We consider two cases.

Case 1: Defer. When $4s^2 \cdot \sum_{j'} C_{i,j'}(P) \leq 3$, we simply add P to F_{i-1} . We have

$$\text{len}(P, G - C_{i,j}) = \text{len}(P, G) + h_{\text{diam},j} \cdot s \cdot C_{i,j}(P) \leq 4h_j \leq h_{\text{cov},j},$$

where the inequality is by $\text{len}(P, G) \leq h_j$, $h_{\text{diam},j} = 4sh_j$ and $C_{i,j}(P) \leq 3/(4s^2)$. Therefore, in this case, the flow path added to F_{i-1} satisfies the invariant.

Case 2: Shortcut. Now suppose $4s^2 \cdot \sum_{j'} C_{i,j'}(P) > 3$. Let u and v be P 's endpoints. We say u is the left side and v the right side. For each $0 \leq k \leq |P|$, we refer to w_k as the P -vertex with k steps away from u . In particular, $w_0 = u$ and $w_{|P|} = v$.

We now define two functions $x : V(P) \rightarrow \mathbb{R}$ denoting the *budgets* of P -vertices from the left and the right respectively: for each vertex $w_k \in V(P)$,

$$x_P(w_k) = 4s^2 \sum_{j'} C_{i,j'}(w_{k-1}, w_k) + C_{i,j'}(w_k) + C_{i,j'}(w_k, w_{k+1}).$$

In particular, $x_P(w_0) = 4s^2 \sum_{j'} C_{i,j'}(w_0) + C_{i,j'}(w_0, w_1)$ and $x_P(w_{|P|}) = 4s^2 \sum_{j'} C_{i,j'}(w_{|P|-1}, w_{|P|}) + C_{i,j'}(w_{|P|})$. Let k_L be the minimum index such that $\sum_{0 \leq k \leq k_L} x_P(w_k) \geq 1$, and symmetrically let k_R be the maximum index such that $\sum_{k_R \leq k \leq |P|} x_P(w_k) \geq 1$. To avoid clutter, we let $L = \{w_0, \dots, w_{k_L}\}$ and $R = \{w_{k_R}, \dots, w_{|P|}\}$. The following Claim 5.7 says that L and R have at most one common vertex.

Claim 5.7. $k_L \leq k_R$.

Proof. By the definition of k_L and k_R , we have $\sum_{0 \leq k \leq k_L-1} x_P(w_k) < 1$ and $\sum_{k_R+1 \leq k \leq |P|} x_P(w_k) < 1$. However, $\sum_{0 \leq k \leq |P|} x_P(w_k) \geq 4s^2 \sum_{j'} C_{i,j'}(P) > 3$. This means there exists a vertex w_k with $k_L - 1 < k < k_R + 1$, which implies $k_L \leq k_R$. \square

We assign each vertex $w \in L \cup R$ a load $x'_P(w) \leq x_P(w)$, satisfying that $\sum_{w \in L} x'_P(w) = 1$ and $\sum_{w \in R} x'_P(w) = 1$. We consider the following three-phase strategy to route $F_i(P)$ flow units from u to v . *Phase 1:* the vertex u sends $F_i(P)$ flow units and each $w \in L$ receives $F_i(P)x'_P(w)$ units. *Phase 2:* Each vertex $w \in L$ sends $F_i(P)x'_P(w)$ units and each vertex $w \in R$ receives

$F_i(P)x'_P(w)$ units. *Phase 3:* Each vertex $w \in R$ sends exactly $F_i(P)x'_P(w)$ units, and the vertex v receives $F_i(P)$ units.

The First and Third Phases. Roughly speaking, for the first and third phases, we will add their corresponding original flows into F_{i-1} , meaning that they will be deferred to lower levels to get shortcut.

Regarding the first phase, recall that for each $w_k \in L$, we want to route $F_i(P)x'(w_k)$ units from u to w_k . To do this, we add into F_{i-1} a flow path $P_k = P[w_0, w_{k-1}]$ with value $F_i(P) \cdot x'(w_k)$. That is, we require the lower levels to give a shortcut that routes $F_i(P) \cdot x'(w_k)$ from u to w_{k-1} (the P -vertex one step closer to u than w_k). Then, we route $F_i(P)x'(w_k)$ units from w_{k-1} to w_k using the original edge $(w_{k-1}, w_k) \in E(G)$.

The third phase is handled in a similar way. For each $w_k \in R$, we route $F_i(P)x'(w_k)$ flow units from w_k to w_{k+1} using the original edge (w_{k-1}, w_k) , and then we add into F_{i-1} a flow path $P_k = P[w_{k+1}, v]$ with value $F_i(P) \cdot x'(w_k)$.

To proceed to the lower level $i - 1$, it remains to show that the flow paths added into F_{i-1} satisfy the invariant. Consider a flow path $P_k = P[u, w_{k-1}]$ added from the first phase (where $w_k \leq L$, i.e. $0 \leq k \leq k_L$). We want to show that $\text{leng}(P_k, G - C_{i,j_k}) \leq h_{\text{cov},j_k}$, where j_k is the minimum index such that $h_{j_k} \geq \text{leng}(P_k, G)$. By the definition of the budget function x , we have $\sum_{j'} C_{i,j'}(P_k) \leq \frac{1}{4s^2} \sum_{0 \leq k' \leq k-1} x_P(w_{k'}) < 1/(4s^2)$, where the second inequality is by $k \leq k_L$ and $\sum_{0 \leq k' \leq k_L-1} x_P(w_{k'}) < 1$ (from the definition of k_L). In particular, $C_{i,j_k}(P_k) \leq 1/(4s^2)$. Because C_{i,j_k} is an $(h_{\text{diam},j_k} s)$ -length moving cut, we have

$$\text{leng}(P_k, G - C_{i,j_k}) = \text{leng}(P_k, G) + h_{\text{diam},j_k} \cdot s \cdot C_{i,j_k}(P_k) \leq 2h_{j_k} \leq h_{\text{cov},j_k},$$

as desired, where the first inequality uses $h_{\text{diam},j_k} = 4sh_{j_k}$. By a similar argument, we can show that each flow path added from the third phase also satisfies the invariant, so we will not explain it in detail.

The Second Phase. The second phase is where the shortcut happens. We define an arbitrary (multi-commodity) demand $\hat{D}_{i,P}$ capturing this single-commodity demand. Namely, $\hat{D}_{i,P}$ satisfies that (1) $|\hat{D}_{i,P}| = F_i(P)$; (2) for each $w \in L$, $\hat{D}_{i,P}(w, \cdot) = F_i(P)x'_P(w)$; and (3) for each $w \in R$, $\hat{D}_{i,P}(\cdot, w) = F_i(P)x'_P(w)$. We will assign $\hat{D}_{i,P}$ to $H_{i,j}$, meaning that we route $\hat{D}_{i,P}$ using shortcut edges in $H_{i,j}$.

By the above assignment, for each $H_{i,j}$ at level i , its total assigned demand, denoted by $\hat{D}_{i,j}$, sums over $\hat{D}_{i,P}$ of all $P \in \text{path}(F_i)$ s.t. j is the minimum index with $h_j \geq \text{leng}(P, G)$. The following Lemma 5.8 showing that $\hat{D}_{i,j}$ can be routed with low steps, small length and congestion 1.

Lemma 5.8. *For each $H_{i,j}$ at level i , its total assigned demand $\hat{D}_{i,j}$ can be routed in $H_{i,j}$ with length $2h_{\text{diam},j}s + 1$, congestion 1 and step 2.*

Proof. First, we show that the demand $\hat{D}_i := \sum_{P \in \text{path}(F_i)} \hat{D}_{i,P}$ is A_i -respecting, which means $\hat{D}_{i,j} \preceq \hat{D}_i$ is also A_i -respecting. To see this, consider any vertex $w \in V(G)$, and we have

$$\begin{aligned} \max\{\hat{D}_i(w, \cdot), \hat{D}_i(\cdot, w)\} &\leq \sum_{P \in \text{path}(F_i)} \max\{\hat{D}_{i,P}(w, \cdot), \hat{D}_{i,P}(\cdot, w)\} \leq \sum_{P \in \text{path}(F_i) \text{ s.t. } w \in P} F_i(P)x_P(w) \\ &\leq 4s^2 \left(u_G(w) \sum_{j'} C_{i,j'}(w) + \sum_{e \in E(G) \text{ incident to } w} u_G(e) \cdot \sum_{j'} C_{i,j'}(e) \right) \\ &= A_i(w). \end{aligned}$$

Here the key step is the third inequality, which is because (1) F_i is feasible, and (2) $x_P(w)$ is the total level- i cut values of w and its incident P -edges scaled up by $4s^2$.

Next, $\hat{D}_{i,j}$ is $h_{\text{cov},j}$ -length in $G - C_{i+1,j}$ because for each demand pair $(u, v) \in \text{supp}(\hat{D}_{i,j})$, u, v is on some $P \in \text{path}(F_i)$ and $\text{len}_G(P, G - C_{i+1,j}) \leq h_{\text{cov},j}$.

Finally, since $\hat{D}_{i,j}$ is $h_{\text{cov},j}$ -length in $G - C_{i+1,j}$, A_i -respecting and trivially $u_{V(G)}$ -respecting (since \hat{D}_i can be routed in G according to the feasible flow F_i), $\hat{D}_{i,j}$ can be routed in $H_{i,j}$ with length $2h_{\text{diam},j}s + 1$, congestion 1 and step 2 by Lemma 5.4. \square

Quality of the Forward Mapping. We now show that $\text{Dem}(F)$ can be routed in $G' = G \cup \bigcup_{i,j} H_{i,j}$ with length $\text{len}_G(F) \cdot O(1/\epsilon^3)$, congestion 1 and step $2^{O(1/\epsilon)}$.

Length and Step. We first argue the length and step bounds using induction on levels. It suffices to prove the following statement for each level i : for each flow path $P \in \text{path}(F_i)$, the above forward mapping scheme maps it to a flow $\hat{F}_{i,P}$ in $G \cup \sum_{i' \leq i,j'} H_{i',j'}$ with length $20(i+1) \cdot s^2 \text{len}_G(P, G)$ and step $6 \cdot 2^{i+1} - 4$. The statement for the top level $i = d$ implies the length and step bounds after plugging in $d \leq O(1/\epsilon)$ and $s = 1/\epsilon$.

Consider the base case $i = 0$. A flow path $P \in \text{path}(F_0)$ is shortcut by $H_{0,j}$, where j is the minimum index with $h_j \geq \text{len}_G(P, G)$, which implies $h_j/2 \leq \text{len}_G(P, G)$. As we discussed above, the shortcut has length $2h_{\text{diam},j}s + 1$ and step 2. Since $2h_{\text{diam},j}s + 1 \leq 8s^2h_j + 1 \leq 20s^2 \text{len}_G(P, G)$, we complete the proof of the base case.

Consider an inductive step $i \geq 1$. Assume the statement holds for $i - 1$. For each flow path $P \in \text{path}(F_i)$, if it belongs to Case 1, the length and step of $\hat{F}_{i,P}$ can be directly bounded by the induction hypothesis, so we assume it belongs to Case 2 from now on. Recall that $\hat{F}_{i,P}$ is the concatenation of routings from three phases.

Phases 1 and 3. Consider the first-phase routing (the third-phase routing is analogous). By the induction hypothesis, its corresponding flow path $P_k = P[u, w_{k-1}]$ deferred to F_{i-1} will be mapped to a flow \hat{F}_{i-1,P_k} in the shortcut graph with length $20is^2 \cdot \text{len}_G(P[u, w_{k-1}], G)$ and step $6 \cdot 2^i - 4$. The extra edge at the end of the routing (w_{k-1}, w_k) has length $\text{len}_G(P[w_{k-1}, w_k], G)$ and step 1. In conclusion, the first-phase routing has length $\max_{w_k \in L} (20is^2 \cdot \text{len}_G(P[u, w_{k-1}], G) + \text{len}_G(P[w_{k-1}, w_k], G)) \leq 20is^2 \cdot \text{len}_G(P[u, w_{k_L}], G)$ and step $6 \cdot 2^i - 3$.

Phase 2. The second-phase routing is on $H_{i,j}$, where j satisfies $h_j/2 \leq \text{len}_G(P, G)$. For each $w_k \in L$, by Lemma 5.8 and an argument similar to the base case, the second-phase routing has length $20s^2 \text{len}_G(P, G)$ and step 2.

Finally, by Claim 5.7, $\hat{F}_{i,P}$ has length $20(i+1)s^2 \cdot \text{len}_G(P, G)$ and step $6 \cdot 2^{i+1} - 4$.

Congestion. The congestion in $\bigcup_{i,j} H_{i,j} \subseteq G'$ is 1 by Lemma 5.8. The congestion on $G \subseteq G'$ is also 1 because the flow we left in the original graph (in the second phase) is at most the feasible flow F_i .

5.3 Backward Mapping: Proof of Lemma 5.6

We first decompose the flow F' in G' into flows in subgraphs G and $H_{i,j}$ (for all i, j) of G' , denoted by F'_G and $F'_{i,j}$ respectively. Formally speaking, $F'_{i,j}$ collects, for all $P \in \text{path}(F')$, all flow subpaths in $P \cap H_{i,j}$ (the values of these subpaths are still $F'(P)$), and F'_G collects all flow subpaths in $P \cap G$ for all $P \in \text{path}(F')$.

Next, we show that $\text{Dem}(F'_{i,j})$ can be routed by a flow $F_{i,j}$ in G with length $h_{\text{diam},j} \cdot s$ and congestion $O(\omega \cdot \log N/\phi)$, where $\omega = n^{O(\epsilon)} \log n$ is the width of the neighborhood cover $\mathcal{N}_{i,j}$ and $\phi = 1/n^{O(\epsilon)}$ is the expansion of the hierarchy. Because $F'_{i,j}$ is a feasible flow in $H_{i,j}$, its demand $\text{Dem}(F'_{i,j})$ is $(\omega \cdot A_i)$ -respecting and $h_{\text{diam},j}$ -length in $G - C_{i+1,j}$ by Definition 5.3 of star graphs and the diameter of $\mathcal{N}_{i,j}$. Because A_i is $(h_{\text{diam},j}, s)$ -length expanding in $G - C_{i+1,j}$, $\text{Dem}(F'_{i,j})$ can be routed in $G - C_{i+1,j}$ with length $h_{\text{diam},j} \cdot s$ and congestion $O(\log N/\phi)$ by Theorem 4.2.

Finally, we let the flow F in G routing $\text{Dem}(F')$ be the concatenation of F'_G and all $F_{i,j}$. We have $\text{leng}(F) \leq \text{leng}(F')$ because each $F_{i,j}$ has length $h_{\text{diam},j} \cdot s \leq \text{leng}(F'_{i,j}) = 2h_{\text{diam},j}s + 1$. The congestion of F is

$$1 + (d \log h) \cdot O(\omega \cdot \log N/\phi) = n^{O(\epsilon)} \log n \log^2 N/\epsilon^2 = O(n^{O(\epsilon)} \log^3 n/\epsilon^2)$$

because F'_G has congestion 1 in G , each $F_{i,j}$ has congestion $O(\omega \cdot \log N/\phi)$, and there are totally $d \cdot \log h$ many $F_{i,j}$.

References

- [ADK23] Daniel Agassy, Dani Dorfman, and Haim Kaplan. Expander decomposition with fewer inter-cluster edges using a spectral cut player. In *50th International Colloquium on Automata, Languages, and Programming (ICALP 2023)*. Schloss-Dagstuhl-Leibniz Zentrum für Informatik, 2023.
- [BBG⁺20] Aaron Bernstein, Jan van den Brand, Maximilian Probst Gutenberg, Danupon Nanongkai, Thatchaphol Saranurak, Aaron Sidford, and He Sun. Fully-dynamic graph sparsifiers against an adaptive adversary. *arXiv preprint arXiv:2004.08432*, 2020.
- [BBST24] Aaron Bernstein, Joakim Blikstad, Thatchaphol Saranurak, and Ta-Wei Tu. Maximum flow by augmenting paths in $n^{2+o(1)}$ time. *arXiv preprint arXiv:2406.03648*, 2024.
- [BGS20] Aaron Bernstein, Maximilian Probst Gutenberg, and Thatchaphol Saranurak. Deterministic decremental reachability, scc, and shortest paths via directed expanders and congestion balancing. In *Annual Symposium on Foundations of Computer Science, FOCS*, pages 1123–1134, 2020.
- [BGS22] Aaron Bernstein, Maximilian Probst Gutenberg, and Thatchaphol Saranurak. Deterministic decremental sssp and approximate min-cost flow in almost-linear time. In *2021 IEEE 62nd Annual Symposium on Foundations of Computer Science (FOCS)*, pages 1000–1008. IEEE, 2022.
- [BH23] Greg Bodwin and Gary Hoppenworth. Folklore sampling is optimal for exact hopsets: Confirming the \sqrt{n} barrier. In *2023 IEEE 64th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 701–720. IEEE, 2023.
- [CE15] Chandra Chekuri and Alina Ene. The all-or-nothing flow problem in directed graphs with symmetric demand pairs. *Mathematical Programming*, 154:249–272, 2015.
- [CGL⁺20] Julia Chuzhoy, Yu Gao, Jason Li, Danupon Nanongkai, Richard Peng, and Thatchaphol Saranurak. A deterministic algorithm for balanced cut with applications to dynamic connectivity, flows, and beyond. In *IEEE Symposium on Foundations of Computer Science (FOCS)*, pages 1158–1167, 2020.
- [CK19] Julia Chuzhoy and Sanjeev Khanna. A new algorithm for decremental single-source shortest paths with applications to vertex-capacitated flow and cut problems. In *Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing*, pages 389–400, 2019.
- [CK24] Julia Chuzhoy and Sanjeev Khanna. Maximum bipartite matching in $n^{2+o(1)}$ time via a combinatorial algorithm. In *Proceedings of the 56th Annual ACM Symposium on Theory of Computing*, pages 83–94, 2024.
- [CKL⁺22] Li Chen, Rasmus Kyng, Yang P. Liu, Richard Peng, Maximilian Probst Gutenberg, and Sushant Sachdeva. Maximum flow and minimum-cost flow in almost-linear time. In *63rd IEEE Annual Symposium on Foundations of Computer Science, FOCS*, pages 612–623. IEEE, 2022.
- [CKS05] Chandra Chekuri, Sanjeev Khanna, and F Bruce Shepherd. Multicommodity flow, well-linked terminals, and routing problems. In *Proceedings of the thirty-seventh annual ACM symposium on Theory of computing*, pages 183–192, 2005.

- [CMGS25] Daoyuan Chen, Simon Meierhans, Maximilian Probst Gutenberg, and Thatchaphol Saranurak. Parallel and distributed expander decomposition: Simple, fast, and near-optimal. In *Proceedings of the 2025 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1705–1719. SIAM, 2025.
- [CS19] Yi-Jun Chang and Thatchaphol Saranurak. Improved distributed expander decomposition and nearly optimal triangle enumeration. In *Proceedings of the 2019 ACM Symposium on Principles of Distributed Computing*, pages 66–73, 2019.
- [CS20] Y. Chang and T. Saranurak. Deterministic distributed expander decomposition and routing with applications in distributed derandomization. In *IEEE Symposium on Foundations of Computer Science (FOCS)*, pages 377–388, full version arXiv:2007.14898, 2020.
- [EHHL25] Antoine El-Hayek, Monika Henzinger, and Jason Li. Fully dynamic approximate minimum cut in subpolynomial time per operation. In *Proceedings of the 2025 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 750–784. SIAM, 2025.
- [GPPG24] Lars Gottesbüren, Nikos Parotsidis, and Maximilian Probst Gutenberg. Practical expander decomposition. In *32nd Annual European Symposium on Algorithms (ESA 2024)*, volume 308, pages 61–1. Schloss Dagstuhl-Leibniz-Zentrum für Informatik, 2024.
- [GR98] Oded Goldreich and Dana Ron. A sublinear bipartiteness tester for bounded degree graphs. In *Proceedings of the thirtieth annual ACM symposium on Theory of computing*, pages 289–298, 1998.
- [GRST21] Gramoz Goranci, Harald Räcke, Thatchaphol Saranurak, and Zihan Tan. The expander hierarchy and its applications to dynamic graph algorithms. In *Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 2212–2228, 2021.
- [Hes03] William Hesse. Directed graphs requiring large numbers of shortcuts. In *Proceedings of the Fourteenth Annual ACM-SIAM Symposium on Discrete Algorithms, January 12-14, 2003, Baltimore, Maryland, USA*, pages 665–669. ACM/SIAM, 2003.
- [HHG25] Bernhard Haeupler, Jonas Huebotter, and Mohsen Ghaffari. A cut-matching game for constant-hop expanders. In *Proceedings of the 2025 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1651–1678. SIAM, 2025.
- [HHL⁺24] Bernhard Haeupler, D. Ellis Hershkowitz, Jason Li, Antti Roeykoe, and Thatchaphol Saranurak. Low-step multi-commodity flow emulators. In *Proceedings of the 56th Annual ACM Symposium on Theory of Computing, STOC '24*, page 71–82. ACM, June 2024.
- [HHT24] Bernhard Haeupler, D Ellis Hershkowitz, and Zihan Tan. New structures and algorithms for length-constrained expander decompositions. *arXiv preprint arXiv:2404.13446*, 2024.
- [HKGW23] Yiding Hua, Rasmus Kyng, Maximilian Probst Gutenberg, and Zihang Wu. Maintaining expander decompositions via sparse cuts. In *Proceedings of the 2023 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 48–69. SIAM, 2023.
- [HLS24] Bernhard Haeupler, Yaowei Long, and Thatchaphol Saranurak. Dynamic deterministic constant-approximate distance oracles with n^ϵ worst-case update time. *arXiv preprint arXiv:2402.18541*, 2024.

- [HP21] Shang-En Huang and Seth Pettie. Lower bounds on sparse spanners, emulators, and diameter-reducing shortcuts. *SIAM J. Discret. Math.*, 35(3):2129–2144, 2021.
- [HRG22] Bernhard Haeupler, Harald Räcke, and Mohsen Ghaffari. Hop-constrained expander decompositions, oblivious routing, and distributed universal optimality. In *Proceedings of the 54th Annual ACM SIGACT Symposium on Theory of Computing, STOC*, page 1325–1338, 2022.
- [HWZ20] Bernhard Haeupler, David Wajc, and Goran Zuzic. Network coding gaps for completion times of multiple unicasts. In *IEEE Symposium on Foundations of Computer Science (FOCS)*, pages 494–505, 2020.
- [JPP25] Yonggang Jiang, Merav Parter, and Asaf Petruschka. New oracles and labeling schemes for vertex cut queries. *arXiv preprint arXiv:2501.13596*, 2025.
- [JS22] Wenyu Jin and Xiaorui Sun. Fully dynamic st edge connectivity in subpolynomial time. In *2021 IEEE 62nd Annual Symposium on Foundations of Computer Science (FOCS)*, pages 861–872. IEEE, 2022.
- [JST24] Wenyu Jin, Xiaorui Sun, and Mikkel Thorup. Fully dynamic min-cut of superconstant size in subpolynomial time. In *Proceedings of the 2024 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 2999–3026. SIAM, 2024.
- [KLOS14] Jonathan A Kelner, Yin Tat Lee, Lorenzo Orecchia, and Aaron Sidford. An almost-linear-time algorithm for approximate max flow in undirected graphs, and its multicommodity generalizations. In *Proceedings of the twenty-fifth annual ACM-SIAM symposium on Discrete algorithms*, pages 217–226. SIAM, 2014.
- [KVV04] Ravi Kannan, Santosh Vempala, and Adrian Vetta. On clusterings: Good, bad and spectral. *Journal of the ACM (JACM)*, 51(3):497–515, 2004.
- [LNPS23] Jason Li, Danupon Nanongkai, Debmalya Panigrahi, and Thatchaphol Saranurak. Near-linear time approximations for cut problems via fair cuts. In *Proceedings of the 2023 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 240–275. SIAM, 2023.
- [LPS25] Yaowei Long, Seth Pettie, and Thatchaphol Saranurak. Connectivity labeling schemes for edge and vertex faults via expander hierarchies. In *Proceedings of the 2025 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1–47. SIAM, 2025.
- [LR99] Tom Leighton and Satish Rao. Multicommodity max-flow min-cut theorems and their use in designing approximation algorithms. *Journal of the ACM (JACM)*, 46(6):787–832, 1999.
- [LS21] Jason Li and Thatchaphol Saranurak. Deterministic weighted expander decomposition in almost-linear time. *arXiv preprint arXiv:2106.01567*, 2021.
- [LS22] Yaowei Long and Thatchaphol Saranurak. Near-optimal deterministic vertex-failure connectivity oracles. In *IEEE Symposium on Foundations of Computer Science (FOCS)*, pages 1002–1010, 2022.
- [NSWN17] Danupon Nanongkai, Thatchaphol Saranurak, and Christian Wulff-Nilsen. Dynamic minimum spanning forest with subpolynomial worst-case update time. In *Proceedings of the 58th Annual IEEE Symposium on Foundations of Computer Science*, pages 950–961, 10 2017.

- [NSY23] Chaitanya Nalam, Thatchaphol Saranurak, and Sorrachai Yingchareonthawornchai. Deterministic k -vertex connectivity in k^2 max-flows. *arXiv preprint arXiv:2308.04695*, 2023.
- [Pel00] David Peleg. *Distributed computing: a locality-sensitive approach*. SIAM, 2000.
- [RST14] Harald Räcke, Chintan Shah, and Hanjo Täubig. Computing cut-based hierarchical decompositions in almost linear time. In *Proceedings of ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 227–238, 2014.
- [SG24] Aurelio L. Sulser and Maximilian Probst Gutenberg. A simple and near-optimal algorithm for directed expander decompositions, 2024.
- [She13] Jonah Sherman. Nearly maximum flows in nearly linear time. In *2013 IEEE 54th Annual Symposium on Foundations of Computer Science*, pages 263–269. IEEE, 2013.
- [ST04] Daniel A Spielman and Shang-Hua Teng. Nearly-linear time algorithms for graph partitioning, graph sparsification, and solving linear systems. In *Proceedings of the thirty-sixth annual ACM symposium on Theory of computing*, pages 81–90, 2004.
- [SW19] Thatchaphol Saranurak and Di Wang. Expander decomposition and pruning: Faster, stronger, and simpler. In *Proceedings of ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 2616–2635, 2019.
- [SY22] Thatchaphol Saranurak and Sorrachai Yingchareonthawornchai. Deterministic small vertex connectivity in almost linear time. In *2022 IEEE 63rd Annual Symposium on Foundations of Computer Science (FOCS)*, pages 789–800. IEEE, 2022.

A Length-Constrained Directed Expansion (Continued)

A.1 Connection to Classic Directed Expander Decomposition

The classic setting of a cut in a graph is a bipartition of the vertex set. A cut is considered sparse if few edges connect the two partitions.

Definition A.1 (Classic Directed Sparse Cut). *Given a directed graph $G = (V, E)$, let $E(A, B)$ denote the set of edges with head vertices in A and tail vertices in B , a cut $(S, V \setminus S)$ where $\text{vol}_G^{\text{sum}}(S) \leq \text{vol}_G^{\text{sum}}(V \setminus S)$ is ϕ -out-sparse if $\sum_{e \in E(S, V \setminus S)} u_G(e) < \phi \cdot \text{vol}_G^{\text{sum}}(S)$. Similarly, the cut is ϕ -in-sparse if $\sum_{e \in E(V \setminus S, S)} u_G(e) < \phi \cdot \text{vol}_G^{\text{sum}}(S)$.*

In directed graphs, the standard definition of an expander only requires that the graph be free from sparse cuts in both directions. That is, a directed graph is a ϕ -expander if and only if it is strongly connected and there is no ϕ -in-sparse or ϕ -out-sparse cut. Below, we extend the above definition by interpreting an expander as the union of strongly connected components where each does not admit a sparse cut.

Definition A.2 (Classic Directed Expander). *Given a directed graph $G = (V, E)$, let $X_1, \dots, X_n \subset V$ be a partition of V and each X_i is a strongly connected component (SCC) of G . For $\phi \in (0, 1)$, a graph G is a classic directed ϕ -expander if every induced graph $G[X_i]$ has no ϕ -out-sparse cut and ϕ -in-sparse cut.*

Now we discuss the connection between classic directed expanders and length-constrained directed expanders, assuming we only consider *pure cuts* when defining LC-directed expanders. Here a pure cut are just a classic edge cuts, i.e. a subset of edges. We can also interpret a pure cut as a moving cut with cut values either 0 or 1.

We say a graph G is an (h, s) -length ϕ -expander if \deg_G^{sum} is (h, s) -length ϕ -expanding in G . Then in fact, for sufficient large h , e.g. $h = n^2 N$, G is an (h, s) -length ϕ -expander if and only if G is a classic $\Theta(\phi)$ -expander. In other words, LC-directed expanders and classic directed expanders are equivalent when we consider only pure cuts and very large h .

It is relatively easy to see that if G is not a classic ϕ -expander, then G is not an (h, s) -length $O(\phi)$ -expander. Let $(S, X \setminus S)$ for some SCC X of G (with $\text{vol}_{G[X]}^{\text{sum}}(S) \leq \text{vol}_{G[X]}^{\text{sum}}(X \setminus S)$) be the witnessing ϕ -sparse (say ϕ -out-sparse) cut. Then the pure cut $C = E(S, X \setminus S)$ is an (h, s) -length ϕ -sparse cut of G , because we can construct a \deg_G^{sum} -respecting h -length symmetric demand D with $|D| \geq \Omega(\text{vol}_{G[X]}^{\text{sum}}(S))$ by considering a bipartite graph (with multiple edges) between X and $X \setminus S$ such that each vertex $v \in X$ incident to exactly $\deg_{G[X]}^{\text{sum}}(v)$ edges and each vertex $v \in X \setminus S$ incident to at most $\deg_{G[X]}^{\text{sum}}(v)$ edges.

Now we show that if G is not an (h, s) -length ϕ -expander, then G is not a classic ϕ -expander. Let C be an (h, s) -length ϕ -sparse cut of G with witnessing demand D . Because D is h -length and symmetric, D has no inter-SCC demand pairs. This means there exists an SCC X of G such that $\text{spar}_{h,s}(C_X, D_X) \leq \phi$, where C_X and D_X are C and D restricted on $G[X]$. Let X'_1, X'_2, \dots, X'_k be the SCCs of $G[X] \setminus C_X$. We say a cut $(S, X \setminus S)$ is *valid* if

- it respects the partition $\{X'_1, X'_2, \dots, X'_k\}$, i.e. S is the union of a subset of X'_i , and
- S has no out-edges in $G \setminus C_X$.

Observe that any valid cut $(S, X \setminus S)$ has $\min\{|E(S, X \setminus S)|, |E(X \setminus S, S)|\} \leq |C_X|$. Thus it remains to find a valid cut $(S^*, X \setminus S^*)$ with $\min\{\text{vol}_{G[X]}^{\text{sum}}(S^*), \text{vol}_{G[X]}^{\text{sum}}(X \setminus S^*)\} = \Omega(|D_X|)$.

Let $d(X'_i) = \sum_{u \in X'_i, v \in X} D_X(u, v)$. By definition, $\sum_{X'_i} d(X'_i) = |D_X|$. Because D_X is $\deg_{G[X]}^{\text{sum}}$ -respecting (since D has no inter-SCC demand pairs), each X'_i has $\text{vol}_{G[X]}^{\text{sum}}(X'_i) \geq d(X'_i)$. Therefore, we just need to find a valid cut $(S^*, X \setminus S^*)$ with $\sum_{X'_i \subseteq S^*} d(X'_i) = \Omega(|D_X|)$, which is an easy exercise by considering a topological order of $\{X'_1, \dots, X'_k\}$.

A.2 Routing Characterization: Proof of Theorem 3.8

In this section, we give the proof of Theorem 3.8. We note that each direction in Theorem 3.8 can be explained with the following Theorem A.3 and Theorem A.4 respectively.

We start with the easier direction Theorem A.3, which shows that there exists a worst-case demand that cannot be routed with desirable congestion and length given the low conductance.

Theorem A.3. *In a directed graph G , suppose that for a demand D there exists a $2h$ -length moving cut C with $2h$ -length sparsity $\phi = \text{spars}_{2h}(C, D)$. Then, any h -length routing of D has congestion at least $\frac{1}{2\phi}$.*

Proof. Without loss of generality, we can assume all demand pairs $(u, v) \in \text{supp}(D)$ is $2h$ -length separated by C , i.e. $\text{sep}_{2h}(C, D) = |D|$, otherwise we consider the sub-demand of D that is $2h$ -length separated by C instead.

Let F be an h -length flow routing D in G with minimum congestion denoted by γ . For each flow path $P \in \text{path}(F)$, we have $\text{len}_G(P) \leq h$ but $\text{len}_G(P, G - C) \geq 2h$, so

$$\sum_{P \in \text{path}(F)} F(P) \cdot \sum_{e \in P} \ell_{C, 2h}(e) \geq \text{value}(F) \cdot h = |D| \cdot h.$$

On the other hand, we have

$$\begin{aligned} \sum_{P \in \text{path}(F)} F(P) \cdot \sum_{e \in P} \ell_{C, 2h}(e) &= \sum_{e \in E(G)} \ell_{C, 2h}(e) \sum_{P \in \text{path}(F) \text{ s.t. } P \ni e} F(P) \\ &\leq \sum_{e \in E(G)} \ell_{C, 2h}(e) \cdot \gamma \cdot u(e) \\ &= \gamma \cdot \sum_{e \in E(G)} 2h \cdot C(e) \cdot u(e) \\ &= \gamma \cdot 2h \cdot |C|. \end{aligned}$$

Therefore, we have $\gamma \geq \frac{|D| \cdot h}{2h \cdot |C|} = \frac{1}{2\phi}$ where we use $\frac{|C|}{|D|} = \phi$ □

On the other hand, Theorem A.4 says that if there exists a lower bound over the sparsity of some demand D , we can also construct an h -length routing of D with a bounded congestion.

Theorem A.4. *In a directed graph G , suppose that D is a demand such that every h -length moving cut C has sparsity at least ϕ for a D , i.e., $\text{spars}_h(C, D) \geq \phi$. Then, D can be routed along h -length paths with congestion at most $O(\frac{\log N}{\phi})$.*

The remainder of this section is the proof Theorem A.4. For a given h -length demand D which sends d_i amount of demand from some vertex s_i to some vertex t_i , we consider the (exponential size) concurrent multicommodity flow LP and its dual given in Figure 5. In it we denote with \mathcal{P}_i the (exponential size) set of all h -length directed paths that go from s_i to t_i :

We note that the primal LP exactly computes a routing for $z \cdot D$ while constraining the congestion to be no more than 1. It is equivalent to the inverse of the optimal h -length congestion of a fractional routing for D .

Lemma A.5. *Let $z, \{f_i(p) \mid i \in [k], p \in \mathcal{P}_i\}$ be an optimal solution for $\text{CONCURRENTFLOW}(D, h)$. Then the optimal h -length congestion for any fractional h -length routing of D is exactly $\frac{1}{z}$.*

Proof. Scale the flows by $\frac{1}{z}$. They now satisfy the demands (thanks to the first set of primal constraints) and cause at most congestion $\frac{1}{z}$ on any edge (thanks to the second set of primal constraints). Therefore the optimal h -length congestion of fractionally routing D is at most $\frac{1}{z}$.

Primal: CONCURRENTFLOW(D, h)	Dual: CUT(D, h)
maximize z	minimize $L = \sum_{e \in E} u(e) \cdot \ell_e$
subject to:	subject to:
$\forall i \in [k]: \sum_{p \in \mathcal{P}_i} f_i(p) \geq z \cdot d_i$	$\forall i \in [k], p \in \mathcal{P}_i: \sum_{e \in p} \ell_e \geq c_i$
$\forall e \in E: \sum_{p \ni e} f_i(p) \leq u(e)$	$\sum_{i \in [k]} d_i c_i \geq 1$
$\forall i \in [k], p: f_i(p) \geq 0$	$\forall e \in E: \ell_e \geq 0$
	$\forall i \in [k]: c_i \geq 0$

Figure 5: The concurrent flow LP relaxation and its dual.

On the other hand, any fractional h -length routing of D with congestion γ can be scaled down by a factor of $\frac{1}{\gamma}$ to give a feasible solution for CONCURRENTFLOW(D, h) with $z = \frac{1}{\gamma}$. Thus if the optimal h -length congestion for any fractional h -length demand is $\gamma' < \frac{1}{z}$, then we can get a feasible solution with $z' = \frac{1}{\gamma'} > z$, which contradicts that z is the optimal value. \square

Next, we show that a feasible solution to the dual LP with value L can be transformed into a h -length moving cut C for D of sparsity $\text{spars}_h(C, D) = O(L \cdot \log N)$. The intuition is to think of the dual variables c_i as indicating whether or not (or to what extent) we are separating the demands d_i while the ℓ_i variables will correspond to scaled length increases in the moving cut. The scaling is such that we are (fractionally) cutting one unit of demand (as forced by the second set of dual demands). With this intuition in mind, the following lemma [HRG22] shows that the c_i values can be "rounded" into an appropriate subset of demands to separate, together with the appropriate scaling factor.

Lemma A.6. *Given sequences $c_1, \dots, c_k, d_1, \dots, d_k \in \mathbb{R}_{\geq 0}$ with $\sum_{i \in [k]} d_i \cdot c_i \geq 1$ there exists a non-empty subset $I \subseteq [k]$ with $\min_{i \in I} c_i \geq \frac{1}{\alpha \cdot \sum_{i \in I} d_i}$ for $\alpha = 1 + \ln \left(\frac{\sum_{i \in [k]} d_i}{\min_{i \in [k]} d_i} \right)$.*

Proof. Suppose, without loss of generality, that $c_1 \geq c_2 \geq \dots \geq c_k$ and assume for the sake of contradiction that none of the sets $[1], [2], \dots, [k]$ satisfy the condition. In other words, if we let $d([i]) = \sum_{j=1}^i d_j$, then $c_i < \frac{1}{\alpha} \cdot \frac{1}{d([i])}$ for all $i \in [k]$. Multiplying both sides by d_i and summing them up, we get that $1 \leq \sum_{i=1}^k d_i c_i < \frac{1}{\alpha} \sum_{i=1}^k \frac{d_i}{d([i])}$. Reordering terms, this implies $\sum_{i=1}^k \frac{d_i}{d([i])} > \alpha$.

Define $f(x)$ as $1/d_1$ on $[0, d_1)$; $1/(d_1 + d_2)$ on $[d_1, d_1 + d_2)$; ...; $1/d([i])$ on $[d([i-1]), d([i])]$ for $i \in [k]$. Now we have

$$\int_0^{d([k])} f(x) dx = \frac{d_1}{d_1} + \frac{d_2}{d_1 + d_2} + \frac{d_3}{d_1 + d_2 + d_3} + \dots + \frac{d_k}{d([k])} > \alpha.$$

However, since $f(x) \leq 1/x$

$$\begin{aligned} \int_0^{d([k])} f(x) dx &= \int_0^{d_1} f(x) dx + \int_{d_1}^{d([k])} f(x) dx \\ &\leq 1 + \int_{d_1}^{d([k])} \frac{1}{x} dx = 1 + \ln \frac{d([k])}{d_1}. \end{aligned}$$

Hence we have $\alpha < \int_0^{d([k])} f(x) dx \leq 1 + \ln \frac{d([k])}{d_1}$, but we set $\alpha = 1 + \ln \left(\frac{\sum_{i \in [k]} d_i}{\min_{i \in [k]} d_i} \right) \geq 1 + \ln \frac{d([k])}{d_1}$, where we reach a contradiction and finish the proof. \square

Lemma A.7. *Suppose $\{\ell_e \mid e \in E\}, \{c_i \mid i \in [k]\}$ is a feasible solution for CUT(D, h). Then there exists an h -length moving cut C for D with h -length sparsity $\text{spars}_h(C, D) = O(L \cdot \log d_{\text{ratio}})$ where $L = \sum_{e \in E} u(e) \cdot \ell_e$ and $d_{\text{ratio}} = \sum_{i \in [k]} d_i / \min_{i \in [k]} d_i$.*

Proof. Applying Lemma A.6 to the c_i values results in a subset $I \subseteq [k]$ of demands with $c_{\min} = \min_{i \in I} c_i \geq \frac{1}{\alpha \cdot \sum_{i \in I} d_i}$ for $\alpha = 1 + \ln \left(\frac{\sum_{i \in [k]} d_i}{\min_{i \in [k]} d_i} \right) = O(\log d_{\text{ratio}})$. Given the scaling factor c_{\min} , for each edge e , we round down the value of $\min\{1, \frac{2\ell_e}{c_{\min}}\}$ to multiple of $1/h$, and define it to be the cut value over e from the h -length moving cut C .

We show that C successfully h -separates every demand $i \in I$. For this, we note that the first set of dual constraints guarantees that for every demand $i \in I$ and for every path $p \in \mathcal{P}_i$, the length of p , assuming every edge e has length ℓ_e , is at least c_{\min} , i.e., $\forall i \in I, p \in \mathcal{P}_i : \sum_{e \in p} \ell_e \geq c_{\min}$. Recall that edge lengths are positive integers as mentioned in Section 2, so each edge has length at least 1 in G . The length of a path $p \in \mathcal{P}_i$ under cut C is therefore at least

$$\begin{aligned} \sum_{e \in p} \ell_{G-C}(e) &= \sum_{e \in p} \ell_G(e) + h \cdot C(e) \\ &\geq \sum_{e \in p} 1 + h \cdot (\min\{1, 2\ell_e/c_{\min}\} - 1/h) \\ &\geq h \cdot \min\{1, \frac{2 \sum_{e \in p} \ell_e}{c_{\min}}\} \\ &= 2h > h \end{aligned}$$

This guarantees that C indeed h -separates every demand $i \in I$.

The amount of demands in D separated by C therefore satisfies

$$\text{sep}_h(C, D) \geq \sum_{i \in I} d_i \geq \frac{1}{\alpha \cdot c_{\min}},$$

where the last inequality comes from the construction if I guarantees that $c_{\min} \geq \frac{1}{\alpha \cdot \sum_{i \in I} d_i}$. By the definition of C , the size $|C|$ of the h -length cut C , is

$$\sum_e u(e) \cdot C(e) \leq \sum_e \frac{u(e) \cdot 2\ell_e}{c_{\min}} = \frac{\sum_e u(e) \cdot 2\ell_e}{c_{\min}} = \frac{2L}{c_{\min}}.$$

Finally, the h -length sparsity $\text{spars}_h(C, D)$ of C for D is $\frac{|C|}{\text{sep}_h(C, D)} \leq \frac{2L/c_{\min}}{1/(\alpha \cdot c_{\min})} = 2L \cdot \alpha = O(L \cdot \log d_{\text{ratio}})$, as desired. \square

The proof of Theorem A.4 follows now immediately from Lemma A.5, Lemma A.7 and strong LP duality.

Proof of Theorem A.4. First of all, D must be h -length, otherwise the zero cut C has $\text{spars}_h(C, D) = 0$. Let $d_{\min} = \min\{D(u, v) \mid \forall u, v \text{ s.t. } D(u, v) > 0\}$ and $d_{\text{ratio}} = |D|/d_{\min}$. We can assume $d_{\text{ratio}} \leq n^4 N$ by the following reasons.

- First we have $|D| \leq n^2 N / \phi$. Assume the opposite, the h -length moving cut C with $C(e) = 1$ for all $e \in E(G)$ trivially has $\text{sep}_h(C, D) = |D|$ and $|C| = \sum_{e \in E(G)} u_G(e) \leq n^2 N$ (here we use that edge capacities are at most N). This means $\text{spars}_h(C, D) = |C|/|D| < \frac{n^2 N}{n^2 N / \phi} = \phi$, a contradiction.
- When $d_{\min} \leq 1/(\phi n^2)$, we consider another demand D' which drops all demand pairs (u, v) with $D(u, v) \leq 1/(\phi n^2)$, which means $d'_{\min} \geq 1/(\phi n^2)$. Also, D' still satisfies $\text{spars}_h(C, D) \geq \phi$ for all h -length moving cuts C . Furthermore, if D' can be routed with length h and congestion $O(\log N / \phi)$, then D can also be routed with length h and congestion $O(\log N / \phi) + n^2 / (\phi n^2) = O(\log N / \phi)$ (since each edge has capacity at least 1).

Providing $d_{\text{ratio}} \leq n^4 N$, we prove the original statement by proving the counter-positive. That is, if a demand D with $d_{\text{ratio}} \leq n^4 N$ cannot be routed along h -length paths with congestion at most γ then there must exist a cut C with h -length sparsity $\text{spars}_h(C, D) = O(\frac{\log N}{\gamma})$ for D .

To see this note that, due to Lemma A.5, the value of $\text{CONCURRENTFLOW}(D, h)$ and because of strong duality also the value of the dual $\text{CUT}(D, h)$ is at most $\frac{1}{\gamma}$. Lemma A.7 now directly implies that there exist the desired h -length cut with sparsity $O(\frac{\log d_{\text{ratio}}}{\gamma}) = O(\frac{\log N}{\gamma})$ for D . \square

A.3 Linkedness

We can actually show the existence of a strengthened version of length-constrained expander decompositions called “linked” length-constrained expander decompositions. For each applied moving cut C , it will slightly increase the original node-weighting to allow more demand in the graph, which makes it more difficult for the graph to be a length-constrained expander. We define such an increase as a new node-weighting added to the original weighting:

Definition A.8 (Linked Node-Weighting). *Let C be an h -length moving cut of a graph $G = (V, E)$ and let ℓ be a positive integer divisible by h . The linked node-weighting L_C^ℓ assigns to each vertex v the value of $\ell \cdot \deg_C(v)$ where $\deg_C(v) = \sum_{e \ni v} u_G(e) \cdot C(e)$.*

Definition A.9 (Linked Length-Constrained Directed Expander Decomposition). *Given a directed graph $G = (V, E)$, a directed ℓ -linked (h, s) -length ϕ -expander decomposition for a node-weighting A with cut slack κ and length slack s is an $h \cdot s$ -length cut C of size at most $\kappa \cdot \phi |A|$ such that $A + L_C^\ell$ is (h, s) -length ϕ -expanding in $G - C$.*

We have defined the sequence of moving cuts in Definition 3.16, but now we also need to incorporate the linked node-weighting added whenever we apply the cut to the graph. We similarly describe the sequence of linked moving cuts as follows:

Definition A.10 (Sequence of Linked Moving Cuts). *Given a directed graph $G = (V, E)$, and node-weighting A , let (C_1, C_2, \dots, C_n) be a sequence of $h \cdot s$ moving cuts, let $G - \sum_{j < i} C_j$ denote the graph that is applied with cuts from C_1 to C_{i-1} and let $A + \sum_{j < i} L_{C_j}^\ell$ denote the node-weighting that is added with linked node-weighting from $L_{C_1}^\ell$ to $L_{C_{i-1}}^\ell$. We define (C_1, C_2, \dots, C_n) as a sequence of ℓ -linked ϕ -sparse moving cuts if and only if the (h, s) -length sparsity of C_i w.r.t $A + \sum_{j < i} L_{C_j}^\ell$ in $G - \sum_{j < i} C_j$ is at most ϕ .*

We again need to bound the overall size of those moving cuts. We employ a similar potential argument, but what is different that the potential will not only decrease from the cut but also increase from the added linked node-weighting. We divide the potential update into two phases to argue about the potential change more clearly, and show that the potential is overall decreasing.

Lemma A.11. *Let C_1, \dots, C_k be an sequence of ℓ -linked ϕ -sparse $h \cdot s$ -length cuts for some node-weighting A in the graph G where $h > 1$, $\phi < 1$, $\ell \leq 2^{-8\alpha-4} \cdot \frac{1}{\phi \ln n}$, $\alpha \geq 1$ and $s > \frac{4 \log_2 n}{\alpha}$, then $\sum_i |C_i| \leq (2^{8\alpha+3} \phi \ln n) \cdot |A|$.*

Proof. Let G_1 denote the initial graph G , $G_i = G - \sum_{j < i} C_j$, and let $A_i = A + \sum_{j < i} L_{C_j}^\ell$. From our assumption of the sequence of moving cuts, for every i there exists a symmetric h -length A_i -respecting demand D_i^* in the graph G_i such that the $h \cdot s$ sparsity of C_i w.r.t D_i^* is at most ϕ .

We again introduce the exponential demand for each graph G_i and the corresponding node-weighting A_i . For simplicity, let w_i denote the exponential distance weight w_h^α w.r.t G_i . We further use D_i to denote the corresponding exponential demand D_{h, A_i}^i w.r.t graph G_i and node-weighting A_i . It is worth noting that exponential demands are different from each other because

the graph G_i and the node-weighting A_i is changed for each i . Specifically, from Lemma 3.10, we have

$$\text{sep}_{h \cdot s/2}^i(C_i, D_i) \geq 2^{-8\alpha-1} \cdot \frac{|C_i|}{\phi}$$

Further, we define the same potential function $P_i : V \rightarrow \mathbb{R}$ w.r.t the graph G_i and node-weighting A_i . It assigns a value to each vertex u with the amount of $P_i(u) = A_i(u) \ln(w_i(u))$.

We divide the potential change into two phases to simplify analysis. In the first phase, we apply cut C_i to the graph G_i and then the resulting graph is G_{i+1} . The distance of some edges increases due to C_i , but the node-weighting still remains as the A_i . Then in the second phase, we added the corresponding linked node-weighting $L_{C_i}^\ell$ to A_i and get A_{i+1} . The distance between vertices in the graph G_{i+1} remains the same in this phase.

Start with graph G_i , each vertex u will have potential $P_i(u)$. After applying cut C_i to the graph G_i , we first get the resulting graph G_{i+1} with same node-weighting A_i , and let $P'_i(u)$ denote the potential of vertex u at this intermediate phase, we have $P'_i(u) = A_i(u) \ln(w_{i+1}(u))$. Since we only increase the length of some edges in G_i , the exponential weight can only decrease between any vertex pairs. Consequently, there is indeed a decrease from $P_i(u)$ to $P'_i(u)$, and we have the same result as in Lemma 3.17,

$$P_i(u) - P'_i(u) \geq A_i(u) \cdot \left(\frac{w_i(u) - w_{i+1}(u)}{w_i(u)} \right) \quad (16)$$

The same analysis gives that the overall potential reduction is still at least

$$\sum_{u \in V} P_i(u) - P'_i(u) \geq \frac{1}{2} \cdot \text{sep}_{h \cdot s/2}^i(C_i, D_{h, A_i}^i) \geq 2^{-8\alpha-2} \cdot \frac{|C_i|}{\phi}$$

In the second phase, the length of edges in graph G_{i+1} does not increase, but the node-weighting A_i is added with $L_{C_i}^\ell$ to get A_{i+1} . Thus the potential over vertex u increases from $P'_i(u)$ to $P_{i+1}(u)$, and we have

$$\begin{aligned} P_{i+1}(u) - P'_i(u) &= (A_{i+1}(u) - A_i(u)) \cdot \ln(w_h^{i+1}(u)) \\ &= \deg_{C_i}(u) \cdot \ell \cdot \ln(w_h^{i+1}(u)) \end{aligned}$$

By summing up, we have

$$\begin{aligned} \sum_{u \in V} P_{i+1}(u) - P'_i(u) &= \sum_{u \in V} \deg_{C_i}(u) \cdot \ell \cdot \ln(w_{i+1}(u)) \\ &\leq 2|C_i| \cdot \ell \cdot \ln n \\ &\leq 2^{-8\alpha-3} \cdot \frac{|C_i|}{\phi} \end{aligned}$$

We use the condition that $\ell \leq 2^{-8\alpha-4} \cdot \frac{1}{\phi \ln n}$ in second inequality. We show that the overall potential P_i decreases at least $2^{-8\alpha-2} \cdot \frac{|C_i|}{\phi}$ to P'_i , and then increases at most $2^{-8\alpha-3} \cdot \frac{|C_i|}{\phi}$ to P_{i+1} , which is equivalent to that the overall potential is monotonously decreasing as the following:

$$\sum_{u \in V} P_i(u) - P_{i+1}(u) \geq 2^{-8\alpha-3} \cdot \frac{|C_i|}{\phi}$$

where $\sum_{u \in V} P_1(u) \geq \sum_{u \in V} P_2(u) \geq \dots \geq \sum_{u \in V} P_{k+1}(u)$. By this, we can come to a similar conclusion over the summation of size of all cuts.

$$\sum_i |C_i| \leq 2^{8\alpha+3} \phi \cdot |A| \cdot \ln n$$

□

Theorem A.12. *For any $G = (V, E)$, a node-weighting A , $h > 1$, $\alpha \geq 1$, $\ell \leq 2^{-8\alpha-4} \cdot \frac{1}{\phi \ln n}$, $\phi < 1$ and a length slack parameter $s = O(\log n)$, there is a directed ℓ -linked (h, s) -length ϕ -expander decomposition for A with cut slack $\kappa = O(n^{O(\frac{1}{s})} \log n)$.*

Proof. Similar as the analysis in Theorem 3.9, we could find cut iteratively from the graph G until the resulting graph is ϕ -expander. We take the union of cuts as $C_{\leq k}$ and by definition $C_{\leq k}$ is a valid expander decomposition for G and A . Further Lemma A.11 guarantees that $|C_{\leq k}| = \sum_{j \leq k} |C_j| \leq 2^{8\alpha+3} \phi \cdot |A| \cdot \ln n$ as long as $s > \frac{4 \ln n}{\alpha}$. This gives that $\kappa \leq 2^{8\alpha+3} \cdot \ln n = O(n^{O(\frac{1}{s})} \log n)$. □