

# Steiner triple systems with high discrepancy

Lior Gishboliner \*

Stefan Glock †

Amedeo Sgueglia †

## Abstract

In this paper, we initiate the study of discrepancy questions for combinatorial designs. Specifically, we show that, for every fixed  $r \geq 3$  and  $n \equiv 1, 3 \pmod{6}$ , any  $r$ -colouring of the triples on  $[n]$  admits a Steiner triple system of order  $n$  with discrepancy  $\Omega(n^2)$ . This is not true for  $r = 2$ , but we are able to asymptotically characterise all 2-colourings which do not contain a Steiner triple system with high discrepancy. The key step in our proofs is a characterization of 3-uniform hypergraphs avoiding a certain natural type of induced subgraphs, contributing to the structural theory of hypergraphs.

## 1 Introduction

Discrepancy theory aims to answer the following question: Given a ground set  $\Omega$  and a family  $\mathcal{P} \subseteq 2^\Omega$ , does there exist an  $r$ -colouring of  $\Omega$  such that each set in  $\mathcal{P}$  contains roughly the same number of elements of each colour? Formally, let  $r \geq 2$  be an integer and  $f : \Omega \rightarrow [r]$  be an  $r$ -colouring of  $\Omega$ . For a set  $P \in \mathcal{P}$  and a colour  $c$ , let  $c(P) := \{x \in P : f(x) = c\}$ . The discrepancy of  $P$  with respect to  $f$  is defined as  $D_f(P) := r \cdot \max_{c \in [r]} \left( |c(P)| - \frac{|P|}{r} \right)$ ; the larger  $D_f(P)$  is, the less balanced is the colouring of  $P$ . The discrepancy of the family  $\mathcal{P}$  with respect to  $f$  is then defined as  $\max_P D_f(P)$ .

Here we consider discrepancy for hypergraphs. In this setting,  $\Omega$  is the edge set of a given hypergraph  $G$  and  $\mathcal{P}$  is a family of subgraphs of  $G$ . Two early results of this type are the works of Erdős and Spencer [12] on the discrepancy of cliques in the complete graph, and of Erdős, Füredi, Loeb and Sós [11] on the discrepancy of copies of a given spanning tree in the complete graph. In the last few years, there has been a renewed interest in questions of this type. For (2-uniform) graphs, there are by now discrepancy results for many different types of subgraphs, such as perfect matchings and Hamilton cycles [2, 13, 16, 17], 1-factorisations [1], spanning trees [17],  $H$ -factors [3, 8] and powers of Hamilton cycles [7]. More recently, the analogous question has been studied in the context of hypergraphs for perfect matchings and Hamilton cycles [4, 14, 15, 20, 24].

In this paper, we initiate this type of research for combinatorial designs, which form another natural class of spanning hypergraphs. One of the most intensively studied objects in design theory are Steiner triple systems, which also have connections and applications to other areas such as coding theory, geometry, graph decompositions and group theory (see e.g. [9]).

A *Steiner triple system* (STS for short) of order  $n$  is a set  $\mathcal{S}$  of 3-subsets of  $[n]$  such that every 2-subset of  $[n]$  is contained in exactly one of the triples of  $\mathcal{S}$ . Hence, we view an STS as a substructure of the complete 3-uniform hypergraph  $K_n^{(3)}$ . Due to the required symmetry condition with respect to the 2-subsets, they are much more complex than the previously studied

---

\*Department of Mathematics, University of Toronto, Canada. *Email:* [lior.gishboliner@utoronto.ca](mailto:lior.gishboliner@utoronto.ca). In the beginning of this work, LG was supported by SNSF grant 200021\_196965.

†Fakultät für Informatik und Mathematik, Universität Passau, Germany. *Email:* [stefan.glock@uni-passau.de](mailto:stefan.glock@uni-passau.de), [amedeo.sgueglia@uni-passau.de](mailto:amedeo.sgueglia@uni-passau.de). SG is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – 542321564, AS is funded by the Alexander von Humboldt Foundation.

substructures like perfect matchings and Hamilton cycles. In fact, even the existence question is highly non-trivial. By a famous result of Kirkman [23], we know that a Steiner triple system of order  $n$  exists if and only if  $n \equiv 1, 3 \pmod{6}$ . Given the multitude of results concerning discrepancy problems for spanning graphs and hypergraphs, the following question suggests itself: Assume that  $n \equiv 1, 3 \pmod{6}$ , and consider an  $r$ -colouring of the 3-subsets of  $[n]$  (corresponding to the edges of  $K_n^{(3)}$ ). Can we find a Steiner triple system of order  $n$  in which one of the colours appears significantly more often than the average?

Recall that an STS of order  $n$  has  $\binom{n}{2}/3$  triples; therefore, by significant, we mean that the discrepancy is  $\Omega(n^2)$ . In other words, we would like to have a colour appearing on at least  $(\frac{1}{r} + \Omega(1)) \frac{\binom{n}{2}}{3}$  edges of the STS. Perhaps surprisingly, the following construction gives a negative answer to this question when  $r = 2$ .

**Example 1.1.** Partition  $[n]$  into two sets  $X$  and  $Y$ , and assign colour blue to all triples touching both  $X$  and  $Y$ , and colour red to all the other triples. In an STS, every edge between  $X$  and  $Y$  must be covered by a blue triple, and every blue triple covers exactly two of these edges. Hence, in any STS, the number of blue triples is  $\frac{|X||Y|}{2}$ . Let us now choose the sizes of  $X, Y$  so that  $\frac{|X||Y|}{2}$  is roughly half of the number of edges in an STS. Take  $|X| = \lfloor \frac{3+\sqrt{3}}{6}n \rfloor$ , so  $|X| = \frac{3+\sqrt{3}}{6}n - O(1)$  and  $|Y| = n - |X| = \frac{3-\sqrt{3}}{6}n + O(1)$ . Then  $\frac{|X||Y|}{2} = \frac{1}{2} \cdot \frac{1}{3} \binom{n}{2} \pm O(n)$ , as required. Thus, in this colouring, any STS has discrepancy  $O(n)$ , i.e., is roughly balanced.

However, we can show that this is essentially the only 2-edge-colouring of  $K_n^{(3)}$  which does not contain a Steiner triple system with high discrepancy.

**Theorem 1.2.** *For every  $\eta > 0$ , there exists  $\mu > 0$  such that the following holds for all sufficiently large  $n$  with  $n \equiv 1, 3 \pmod{6}$ . Every 2-edge-colouring of  $K_n^{(3)}$  either contains a Steiner triple system with discrepancy at least  $\mu n^2$ , or it can be obtained from the colouring in Example 1.1 by switching the colour of at most  $\eta n^3$  triples.*

In contrast with the 2-colour case, when the number of colours is three or more, we can show that a Steiner triple system with high discrepancy always exists.

**Theorem 1.3.** *For all  $r \in \mathbb{N}$  with  $r \geq 3$ , there exists  $\mu > 0$  such that the following holds for all sufficiently large  $n$  with  $n \equiv 1, 3 \pmod{6}$ . Every  $r$ -edge-colouring of  $K_n^{(3)}$  contains a Steiner triple system with discrepancy at least  $\mu n^2$ .*

**Organisation.** Section 2 provides an overview of the various ideas we use to prove our main theorems and collects a list of known tools. In Section 3 we state and prove our key structure theorem for 2-edge-colourings of  $K_n^{(3)}$ . In Sections 4 and 5, we show its relevance for finding an STS with high discrepancy. Section 6 contains the proofs of Theorems 1.2 and 1.3, and Section 7 consists of concluding remarks and further problems.

## 2 Proof overview and tools

### 2.1 Overview

In order to obtain an STS with high discrepancy, we will set aside a certain number of small (edge-coloured) graphs, which we call *gadgets*. These will be used to boost the discrepancy. Let us formalise this notion by defining the gadgets we are going to use.

A *Pasch configuration* is the 6-vertex 3-graph with edges 135, 146, 236, 245 (see the left drawing in Figure 1). Let  $K_{2,2,2}^{(3)}$  denote the complete 3-partite 3-graph with parts of size 2. Observe that  $K_{2,2,2}^{(3)}$  edge-decomposes into two Pasch configurations: Indeed, suppose that the vertex set

of  $K_{2,2,2}^{(3)}$  is [6] and the three parts are  $\{1, 2\}$ ,  $\{3, 4\}$  and  $\{5, 6\}$ . Then  $\{135, 146, 236, 245\}$  and  $\{136, 145, 235, 246\}$  are two edge-disjoint Pasch configurations covering all the 8 edges of  $K_{2,2,2}^{(3)}$  (see Figure 1). The crucial property is that both Pasch configurations have the same shadow<sup>1</sup>, namely the 12 edges of  $K_{2,2,2}$ . This means that if an STS contains one of the Pasch configurations, then we can exchange it for the other Pasch configuration, while still having an STS. An edge-coloured copy  $K$  of  $K_{2,2,2}^{(3)}$  is called a *gadget* if there is a colour  $c$  such that the two Pasch configurations decomposing  $K$  have a different number of edges in colour  $c$ . We will use gadgets to manipulate the colour profile of an STS.




Figure 1: The left figure shows a Pasch configuration. The two figures together show that the edge set of the complete 3-partite 3-graph with parts  $\{1, 2\}$ ,  $\{3, 4\}$  and  $\{5, 6\}$  decomposes into two Pasch configurations:  $\{135, 146, 236, 245\}$  (in green) and  $\{136, 145, 235, 246\}$  (in brown).

Consider an  $r$ -edge-colouring of  $K_n^{(3)}$  (with  $n \equiv 1, 3 \pmod{6}$ ). If we can find a collection  $\mathcal{K}$  of  $\Omega(n^6)$  many gadgets, then we proceed as follows to show that the colouring contains an STS with high discrepancy (c.f. Theorem 4.1). We first pass to a subcollection  $\mathcal{I} \subseteq \mathcal{K}$  of gadgets whose shadows are pairwise edge-disjoint and every vertex is contained in at most a small linear number of gadgets from  $\mathcal{I}$  (this will be achieved by sampling every gadget independently with a suitable probability). Now, consider the 2-graph obtained from  $K_n$  by removing the edges which are in the shadow of a gadget in  $\mathcal{I}$ . This graph still has high minimum degree and thus its edge set can be decomposed into triangles (c.f. Theorem 2.1). Considering these triangles as edges of  $K_n^{(3)}$ , we obtain a linear hypergraph<sup>2</sup> covering almost all edges of  $K_n$ . Regardless of the colour of the edges of this linear hypergraph, we can complete it to an STS with high discrepancy by choosing, for each gadget in  $\mathcal{I}$ , the appropriate Pasch configuration.

However, there are edge-colourings of  $K_n^{(3)}$  which do not even contain a single gadget: for example, this is the case for the 2-edge-colouring described in Example 1.1, where we do not put any condition on the sizes of  $X$  and  $Y$  (and in particular for a monochromatic colouring, which corresponds to one of the sets  $X, Y$  being empty). As our main structural result (c.f. Theorem 3.1), we will show that any 2-edge-colouring having few gadgets is close to the colouring given by Example 1.1 (up to switching the two colours).

Thus, in the 2-colour case, we either find enough gadgets (and thus get an STS with high discrepancy, via the argument described above), or we get a partition  $X, Y$  as in Example 1.1. Now, if the sizes of  $X$  and  $Y$  differ significantly from those specified in Example 1.1 (i.e., if  $|X|/n$  is far from both  $\frac{3+\sqrt{3}}{6}$  and  $1 - \frac{3+\sqrt{3}}{6} = \frac{3-\sqrt{3}}{6}$ ), then significantly more than half of the edges of  $K_n^{(3)}$  have the same colour, in which case a random embedding of any given STS into  $K_n^{(3)}$  is expected to have high discrepancy (c.f. Fact 2.2). Therefore the only case where we cannot guarantee an STS with high discrepancy is when the colouring is close to the one in Example 1.1 (with the same sizes for  $X, Y$ ). This proves Theorem 1.2.

<sup>1</sup>The *shadow* of a 3-graph  $P$  is the set of all pairs of vertices which are contained in an edge of  $P$ .

<sup>2</sup>A hypergraph is *linear* if every two edges intersect in at most one vertex.

Finally, let us consider the case of  $r \geq 3$  colours. Before discussing our strategy, we make the following observation. Given an  $r$ -edge-colouring of  $K_n^{(3)}$ , consider the 2-edge-colouring of  $K_n^{(3)}$  obtained by identifying  $r - 1$  of the  $r$  colours into one colour. Observe that if a copy of  $K_{2,2,2}^{(3)}$  is a gadget for this 2-colouring, then it is also a gadget for the original colouring. Therefore we can simply refer to a gadget without specifying the colouring. To take advantage of our structure theorem for 2-colourings, we consider the 2-edge-colourings of  $K_n^{(3)}$  obtained by identifying  $(r - 1)$ -subsets of colours. If one of these 2-colourings contains many gadgets, we are done (as, by the observation above, these gadgets are gadgets for the original colouring as well). Otherwise, by the above discussion, we know the structures of each of them quite precisely. By comparing them, we will prove that all but at most two colours appear on only very few edges. Overall, when  $r \geq 3$ , we always find  $\Omega(n^6)$  gadgets unless one colour class is small (c.f. Theorem 5.1). In both cases, this implies that we can find an STS with high discrepancy: in the first case by the argument described at the beginning of this section, and in the second case by embedding any given STS into  $K_n^{(3)}$  at random.

## 2.2 Tools

A  $K_3$ -decomposition of a (2-uniform) graph  $G$  is a partition of  $E(G)$  into triangles. Thus, an STS is equivalent to a  $K_3$ -decomposition of  $K_n$ . Note that if  $G$  has a  $K_3$ -decomposition then  $|E(G)|$  is divisible by 3 and all vertices of  $G$  have even degree. A graph having these properties is called  $K_3$ -divisible. Kirkman's theorem can then be rephrased as stating that  $K_n$  admits a  $K_3$ -decomposition if and only if  $K_n$  is  $K_3$ -divisible.

As described in the proof overview, our strategy relies on the use of gadgets. To make this strategy work, it is crucial to be able to find a  $K_3$ -decomposition of the graph obtained from  $K_n$  by removing the edges in the shadows of the gadgets. A famous conjecture of Nash-Williams [26] asks for a minimum degree generalisation of Kirkman's theorem and postulates that every  $K_3$ -divisible  $n$ -vertex graph with minimum degree at least  $3n/4$  has a  $K_3$ -decomposition. This is still wide open, but the statement is known to be true if  $3/4$  is replaced by a larger constant, which suffices for our purposes. The following minimum degree version of Kirkman's theorem was proved in [6] by reducing the problem to a fractional variant (see [5] for a short proof).

**Theorem 2.1.** *For sufficiently large  $n$ , every  $K_3$ -divisible  $n$ -vertex graph  $G$  with  $\delta(G) \geq 0.91n$  has a triangle decomposition.*

Given an  $r$ -edge-coloured  $K_n^{(3)}$ , finding an STS with high discrepancy is straightforward when there is a colour used only few times. This is formalised in the following fact, for which we provide a proof for completeness.

**Fact 2.2.** *For every  $\beta > 0$  and  $r \in \mathbb{N}$  with  $r \geq 2$ , there exists  $\mu > 0$  such that the following holds for every  $n$  with  $n \equiv 1, 3 \pmod{6}$ . Let  $K_n^{(3)}$  be  $r$ -edge-coloured and suppose that there exists a colour  $c$  such that the number of edges of colour  $c$  is at most  $(\frac{1}{r} - \beta) \cdot \binom{n}{3}$ . Then there exists a Steiner triple system with discrepancy at least  $\mu n^2$ .*

*Proof.* Let  $\mu \ll \beta, 1/r$ , fix an arbitrary STS of order  $n$  and consider a random embedding into  $K_n^{(3)}$  (i.e. a random bijection from its points to  $[n]$ ). Then the expected number of triples of colour  $c$  is at most  $\frac{1}{3} \binom{n}{2} \cdot \frac{(\frac{1}{r} - \beta) \cdot \binom{n}{3}}{\binom{n}{3}} = (\frac{1}{r} - \beta) \cdot \frac{1}{3} \binom{n}{2}$ . Therefore there exists an embedding  $\mathcal{S}$  of the STS with at most  $(\frac{1}{r} - \beta) \cdot \frac{1}{3} \binom{n}{2}$  triples of colour  $c$ . By averaging over the other colours, there exists  $c' \in [r] \setminus \{c\}$  such that  $\mathcal{S}$  has at least

$$\frac{\frac{1}{3} \binom{n}{2} - (\frac{1}{r} - \beta) \cdot \frac{1}{3} \binom{n}{2}}{r - 1} \geq \frac{1}{r} \cdot \frac{1}{3} \binom{n}{2} + \mu n^2$$

triples of colour  $c'$ , as desired. □

We will use the hypergraph removal lemma, first conjectured by Erdős, Frankl and Rödl [10], then proved independently by Gowers [19] and Nagle, Rödl, Schacht and Skokan [25, 28, 29] and later by Tao [30]. More precisely, we will use the induced hypergraph removal lemma, proved by Rödl and Schacht [27].

**Lemma 2.3** (Induced hypergraph removal lemma). *For every finite family of  $k$ -graphs  $\mathcal{F}$  and every  $\alpha > 0$ , there exist  $\delta > 0$  and  $n_0 \in \mathbb{N}$  such that the following holds. Let  $\mathcal{H}$  be a  $k$ -graph on  $n \geq n_0$  vertices. If for every  $F \in \mathcal{F}$ ,  $\mathcal{H}$  contains at most  $\delta n^{v(F)}$  induced copies of  $F$ , then one can change (add or delete) at most  $\alpha n^k$  edges of  $\mathcal{H}$  so that the resulting hypergraph contains no induced copy of any member of  $\mathcal{F}$ .*

Finally, we need the following form of the Chernoff bound (see e.g. [21, Corollaries 2.3 and 2.4]).

**Lemma 2.4** (Chernoff bound). *Let  $X$  be the sum of independent Bernoulli random variables. Then for every  $\delta \in (0, 1)$  we have*

$$\mathbb{P}\left[|X - \mathbb{E}[X]| \geq \delta \mathbb{E}[X]\right] \leq 2 \exp\left(-\frac{\delta^2}{3} \mathbb{E}[X]\right).$$

Moreover, for every  $k \geq 7 \mathbb{E}[X]$  we have  $\mathbb{P}[X > k] \leq \exp(-k)$ .

### 3 Key structure theorem

In this section we establish our key structural result, characterizing 2-edge-colourings of  $K_n^{(3)}$  which have no (or only few) gadgets. Recall that a gadget is an edge-coloured copy  $K$  of  $K_{2,2,2}^{(3)}$  for which there is a colour  $c$  such that the two Pasch configurations decomposing  $K$  have a different number of edges in colour  $c$ .

**Theorem 3.1.** *For any  $\alpha > 0$ , there is  $\delta > 0$  such that for every  $n$  large enough, every 2-edge-colouring of  $K_n^{(3)}$  satisfies (at least) one of the following properties:*

- (1) *There are at least  $\delta n^6$  gadgets.*
- (2) *There is a partition  $[n] = X \cup Y$  such that after changing the colour of at most  $\alpha n^3$  edges, the following holds: All edges inside  $X$  and inside  $Y$  have the same colour, and all edges intersecting both  $X$  and  $Y$  have the other colour.*

To prove Theorem 3.1, it is enough to prove the following proposition, which makes the stronger assumption that the 2-edge-colouring of  $K_n^{(3)}$  has no gadgets at all. We will then combine this with the induced hypergraph removal lemma (Lemma 2.3) to deduce Theorem 3.1.

**Proposition 3.2.** *In every 2-edge-colouring of  $K_n^{(3)}$  with no gadgets, there is a partition  $[n] = X \cup Y$  such that after changing the colour of  $O(n^2)$  edges, the following holds: All edges inside  $X$  and inside  $Y$  have the same colour, and all edges intersecting both  $X$  and  $Y$  have the other colour.*

We now give the quick derivation of Theorem 3.1 from Proposition 3.2 and Lemma 2.3.

*Proof of Theorem 3.1.* Let  $\mathcal{F}$  be the family of 6-vertex 3-graphs  $F$  such that if the edges of  $F$  are coloured red and the non-edges of  $F$  are coloured blue, then this 2-colouring of  $K_6^{(3)}$  contains a gadget. Then a red/blue colouring of  $E(K_n^{(3)})$  has no gadgets if and only if the hypergraph  $\mathcal{H}$  of red edges is induced  $\mathcal{F}$ -free. Choose  $\delta > 0$  such that Lemma 2.3 applies with  $k = 3$  and  $\alpha/2$  in place of  $\alpha$ .

Now consider any red/blue colouring of  $E(K_n^{(3)})$ , and let  $\mathcal{H}$  be the hypergraph consisting of the red edges. If the colouring has at least  $\delta n^6$  gadgets, we are done. Otherwise,  $\mathcal{H}$  has at most  $\delta n^6$  induced copies of 3-graphs from  $\mathcal{F}$ . Hence, by Lemma 2.3, we can change the colour



of at most  $\alpha n^3/2$  edges of  $K_n^{(3)}$  to obtain a colouring having no induced copies of any  $F \in \mathcal{F}$ , and hence no gadgets. We can then apply Proposition 3.2 to this colouring. Overall, we have changed the colour of at most  $\alpha n^3/2 + O(n^2) \leq \alpha n^3$  edges, where the inequality holds since  $n$  is large enough.  $\square$

We now move on to the proof of Proposition 3.2. We will consider edge-colourings  $f: E(K_n) \rightarrow \{0, \pm 1\}$ . In such a colouring, an *unbalanced*  $C_4$  consists of 4 distinct vertices  $a, b, c, d$  and the edges  $ab, bc, cd, da$  such that  $f(ab) + f(cd) \neq f(bc) + f(ad)$ . The following lemma characterizes the structure of  $\{0, \pm 1\}$ -edge-colourings of  $K_n$  with no unbalanced  $C_4$ .

**Lemma 3.3.** *Let  $n \geq 50$ . Let  $f: E(K_n) \rightarrow \{0, \pm 1\}$  with no unbalanced  $C_4$ . Then, after deleting at most 8 vertices, one of the following holds:*

- (1) *All edges have colour 0.*
- (2) *There is a vertex-partition  $A \cup B$  such that all edges inside  $A$  have colour 1, all edges inside  $B$  have colour  $-1$ , and all edges between  $A$  and  $B$  have colour 0.*

*Proof.* Let  $M_1$  (resp.  $M_{-1}$ ) be a largest matching of edges of colour 1 (resp.  $-1$ ). We first claim that every edge inside  $V(M_1)$  is coloured with colour 1, and similarly for  $M_{-1}$ . So let  $ab, cd \in M_1$ . If one of the edges between  $\{a, b\}$  and  $\{c, d\}$  has colour different from 1, then we get an unbalanced  $C_4$ . Indeed, without loss of generality, suppose that  $f(bc) \neq 1$ . Then  $f(ab) + f(cd) = 2$  while  $f(bc) + f(ad) \leq f(ad) \leq 1$ , so  $a, b, c, d$  is an unbalanced  $C_4$ . This shows that  $V_1 := V(M_1)$  is a clique in colour 1. Similarly,  $V_{-1} := V(M_{-1})$  is a clique in colour  $-1$ . Also, by the maximality of  $M_1, M_{-1}$ , all edges inside  $V_0 := [n] \setminus (V_1 \cup V_{-1})$  have colour 0.

Next, we claim that for all distinct  $\alpha, \beta \in \{0, \pm 1\}$ , all but at most  $3n$  of the edges in  $E(V_\alpha, V_\beta)$  have colour  $\frac{\alpha+\beta}{2}$ . (Note that it could happen that  $\frac{\alpha+\beta}{2} \notin \{0, \pm 1\}$ .) Indeed, suppose otherwise, i.e., that at least  $3n + 1$  of the edges in  $E(V_\alpha, V_\beta)$  do not have this colour. Then, by using the pigeonhole principle (over the colours) and the fact that any graph with more edges than vertices has a matching of size 2, there exist two disjoint edges  $ab, cd \in E(V_\alpha, V_\beta)$  such that  $f(ab) = f(cd) =: \gamma \neq \frac{\alpha+\beta}{2}$ . Without loss of generality,  $a, d \in V_\alpha$  and  $b, c \in V_\beta$ . But now  $f(ad) + f(bc) = \alpha + \beta \neq 2\gamma = f(ab) + f(cd)$ , meaning that  $a, b, c, d$  is an unbalanced  $C_4$ . This proves our claim.

If  $\alpha = 0$  and  $\beta \in \{\pm 1\}$ , then  $\frac{\alpha+\beta}{2}$  is not an integer, so no edge can have colour  $\frac{\alpha+\beta}{2}$ . This means that  $|V_0| \cdot |V_1| \leq 3n$  and  $|V_0| \cdot |V_{-1}| \leq 3n$ . So  $|V_0|(n - |V_0|) \leq 6n$ . Hence,  $|V_0| \leq 6$  or  $n - |V_0| \leq 6$  (using  $n \geq 50$ ). In the latter case, after deleting the at most 6 vertices in  $[n] \setminus V_0$ , all remaining edges have colour 0, meaning that Item 1 in the lemma holds.

Suppose now that  $|V_0| \leq 6$ . Delete the vertices of  $V_0$ . Set  $A := V_1$  and  $B := V_{-1} \setminus V_1$ , so that  $A, B$  are disjoint. By the same argument as above, there is no matching of size 3 in  $E(A, B)$  whose edges have colours 1 or  $-1$  only. Indeed, otherwise, by the pigeonhole principle, there would be two edges between  $A$  and  $B$  with the same colour  $\gamma \neq 0$ , and this would give an unbalanced  $C_4$ . By the König–Egerváry theorem, we can delete at most 2 vertices such that in the remaining graph, all edges between  $A$  and  $B$  have colour 0. Thus, Item 2 in the lemma holds.  $\square$

We also need the following result concerning 2-edge-colourings of  $K_n^{(3)}$ . Given (not necessarily disjoint)  $A, B, C \subseteq [n]$ , we say that an edge  $abc$  is of *type*  $ABC$  if  $a \in A, b \in B, c \in C$ .

**Lemma 3.4.** *Let  $\chi$  be a 2-edge-colouring of  $K_n^{(3)}$  and let  $A, B, C \subseteq [n]$  with  $|A|, |B|, |C| \geq 360$ . Suppose that every two of the sets  $A, B, C$  are disjoint or equal. Also, suppose that there are at least  $m$  edges of type  $ABC$  in each colour. Then there exists  $Z \in \{A, B, C\}$  such that there are at least  $\frac{m \cdot |Z|}{900}$  pairs of edges  $e, e'$  of type  $ABC$  with  $\chi(e) \neq \chi(e')$ ,  $|e \cap e'| = 2$  and  $e \triangle e' \subseteq Z$ .*

*Proof.* Let the two colours be red and blue, and assume that red is the smallest colour class with respect to edges of type  $ABC$ . Every time we consider an edge, it is implicitly assumed that it is of type  $ABC$ , and we denote by  $e(A, B, C)$  the number of edges of this type. We first consider pairs of edges  $f = abc$ ,  $f' = a'b'c'$  with  $a, a' \in A$ ,  $b, b' \in B$ ,  $c, c' \in C$ ,  $\chi(f) \neq \chi(f')$  and  $f \cap f' = \emptyset$ . Then the number of such pairs is at least  $m \cdot \left( \frac{e(A, B, C)}{2} - |A||B| - |B||C| - |A||C| \right) \geq m \cdot \frac{|A||B||C|}{30}$ . Indeed, we have at least  $m$  choices to choose a red edge, and for each such choice, we then have to choose a blue edge which is vertex-disjoint from the chosen red one. There are at least  $\frac{e(A, B, C)}{2}$  blue edges, and at most  $|A||B| + |B||C| + |A||C|$  of them intersect the chosen red edge. Moreover,  $\frac{e(A, B, C)}{2} \geq \frac{|A|(|B|-1)(|C|-2)}{12} \geq \frac{|A||B||C|}{24}$  and

$$\begin{aligned} \frac{e(A, B, C)}{2} - |A||B| - |B||C| - |A||C| &\geq \frac{|A||B||C|}{24} - |A||B| - |B||C| - |A||C| \\ &\geq |A||B||C| \left( \frac{1}{24} - \frac{1}{|C|} - \frac{1}{|A|} - \frac{1}{|B|} \right) \geq \frac{|A||B||C|}{30}, \end{aligned}$$

where we used that  $|A|, |B|, |C| \geq 360$ .

Now, fix a pair  $f, f'$  as above. Consider the 4 edges  $f_1 = f = abc$ ,  $f_2 = abc'$ ,  $f_3 = ab'c'$ ,  $f_4 = f' = a'b'c'$ . Then  $|f_i \cap f_{i+1}| = 2$  for every  $1 \leq i \leq 3$ . As  $\chi(f_1) \neq \chi(f_4)$ , there must exist  $1 \leq i \leq 3$  with  $\chi(f_i) \neq \chi(f_{i+1})$ . Note that if  $i = 1$  then  $f_i \triangle f_{i+1} = \{c, c'\}$ , if  $i = 2$  then  $f_i \triangle f_{i+1} = \{b, b'\}$ , and if  $i = 3$  then  $f_i \triangle f_{i+1} = \{a, a'\}$ . Therefore, in any case,  $f_i \triangle f_{i+1} \subseteq Z$  for some  $Z \in \{A, B, C\}$ . This shows that there are at least  $\frac{2}{\binom{6}{3}} \frac{m|A||B||C|}{30} = \frac{m|A||B||C|}{900}$  vertex sets of size six which contain a pair of edges  $e, e'$  with  $\chi(e) \neq \chi(e')$ ,  $|e \cap e'| = 2$  and  $e \triangle e' \subseteq Z$  for some  $Z \in \{A, B, C\}$ . The factor of  $2 \cdot \binom{6}{3}^{-1}$  accounts for the fact that each such 6-set is counted at most  $\frac{1}{2} \cdot \binom{6}{3}$  times.

By averaging, there exists  $Z \in \{A, B, C\}$  such that for at least  $\frac{m|A||B||C|}{900}$  of the 6-sets considered above it holds that  $e \triangle e' \subseteq Z$ . Suppose that  $Z = A$ ; the other cases (i.e.,  $Z = B$  or  $Z = C$ ) are analogous. Each pair  $e, e'$  with  $|e \cap e'| = 2$  and  $e \triangle e' \subseteq A$  is contained in at most  $|B||C|$  vertex sets of size six with two vertices from each part, where we used the assumption that any two of the sets  $A, B, C$  are disjoint or equal. Hence, the number of pairs  $e, e'$  with  $\chi(e) \neq \chi(e')$ ,  $|e \cap e'| = 2$  and  $e \triangle e' \subseteq A$  is at least

$$\frac{m|A||B||C|/900}{|B||C|} = \frac{m|A|}{900}.$$

□

We are now ready to prove Proposition 3.2.

*Proof of Proposition 3.2.* We may and will assume that  $n$  is large enough. Let  $\chi : E(K_n^{(3)}) \rightarrow \{0, 1\}$  be a 2-edge-colouring of  $K_n^{(3)}$  with no gadgets. For distinct vertices  $x, y \in [n]$ , let  $f_{xy}$  be the function  $f_{xy} : \binom{[n] \setminus \{x, y\}}{2} \rightarrow \{0, \pm 1\}$  given by  $f_{xy}(uv) = \chi(xuv) - \chi(yuv)$ . Thus,  $f_{xy}$  is a colouring with colours  $0, \pm 1$  of the (2-uniform) clique on the  $n - 2$  vertices  $[n] \setminus \{x, y\}$ . Also, observe that  $f_{xy} = -f_{yx}$ . We first claim that  $f_{xy}$  has no unbalanced  $C_4$ . Indeed, let  $a, b, c, d \in [n] \setminus \{x, y\}$  be distinct vertices. Consider the copy of  $K_{2,2,2}^{(3)}$  with parts  $\{x, y\}, \{a, c\}, \{b, d\}$ . The two Pasch configurations decomposing this copy are  $P_1 = \{xab, xcd, yad, ybc\}$  and  $P_2 = \{yab, ycd, xad, xbc\}$ . Since there is no gadget, we have  $\sum_{e \in P_1} \chi(e) = \sum_{e \in P_2} \chi(e)$ . On the other hand,

$$\begin{aligned} 0 &= \sum_{e \in P_1} \chi(e) - \sum_{e \in P_2} \chi(e) = \sum_{uv \in \{ab, cd\}} (\chi(xuv) - \chi(yuv)) - \sum_{uv \in \{bc, ad\}} (\chi(xuv) - \chi(yuv)) \\ &= \sum_{uv \in \{ab, cd\}} f_{xy}(uv) - \sum_{uv \in \{bc, ad\}} f_{xy}(uv). \end{aligned}$$

So  $f_{xy}(ab) + f_{xy}(cd) = f_{xy}(bc) + f_{xy}(ad)$ . This shows that  $f_{xy}$  has no unbalanced  $C_4$ , as claimed.

By Lemma 3.3 (applied to  $f_{xy}$ ), there is a set  $S_{xy} \subseteq [n] \setminus \{x, y\}$  of size at most 8 such that after deleting  $S_{xy}$ , the colouring  $f_{xy}$  satisfies one of the following:

1. All edges have colour 0.
2. There is a vertex-partition  $A_{xy} \cup B_{xy}$  such that all edges inside  $A_{xy}$  have colour 1, all edges inside  $B_{xy}$  have colour  $-1$ , and all edges between  $A_{xy}$  and  $B_{xy}$  have colour 0.

The pair  $xy$  is called *even* if Item 1 holds and *odd* if Item 2 holds. Note that  $xy$  is even (resp. odd) if and only if  $yx$  is even (resp. odd). Also, if  $xy$  is odd, we may assume that  $A_{yx} = B_{xy}$  and  $B_{yx} = A_{xy}$ .

*Claim 1.* Let  $x, y, z \in [n]$  be distinct vertices. If  $xy$  and  $yz$  have the same parity (i.e., both are even or both are odd) then  $xz$  is even. If  $xy$  and  $yz$  have different parity then  $xz$  is odd.

*Proof.* Note that restricted to  $\binom{[n] \setminus \{x, y, z\}}{2}$ , we have  $f_{xz} = f_{xy} + f_{yz}$ . Indeed, for every distinct  $u, v \in [n] \setminus \{x, y, z\}$ , it holds that

$$f_{xz}(uv) = \chi(xuv) - \chi(zuv) = (\chi(xuv) - \chi(yuv)) + (\chi(yuv) - \chi(zuv)) = f_{xy}(uv) + f_{yz}(uv).$$

Delete the (at most 17) vertices in  $S_{xy} \cup S_{yz} \cup \{y\}$ . Suppose first that  $xy, yz$  are even. Then  $f_{xy}(uv) = f_{yz}(uv) = 0$  for any two remaining vertices  $u, v$  and  $f_{xz}(uv) = f_{xy}(uv) + f_{yz}(uv) = 0$ . This means that  $f_{xz} = f_{xy} + f_{yz}$  cannot satisfy Item 2, so it must satisfy Item 1 and hence  $xz$  is even. Suppose now that  $xy, yz$  have different parity, say  $xy$  is odd and  $yz$  is even. Then for every  $u, v \in A_{xy}$  or  $u, v \in B_{xy}$ , we have  $f_{xz}(uv) = f_{xy}(uv) + f_{yz}(uv) \in \{\pm 1\}$ . This means that  $f_{xz}$  cannot satisfy Item 1, so it must satisfy Item 2 and  $xz$  is odd. Finally, suppose that  $xy, yz$  are both odd. One of the four sets  $A_{xy} \cap A_{yz}, A_{xy} \cap B_{yz}, B_{xy} \cap A_{yz}, B_{xy} \cap B_{yz}$  has size at least  $\frac{n-17}{4}$ . For any two  $u, v$  in this set,  $f_{xy}(uv), f_{yz}(uv) \in \{\pm 1\}$  and thus  $f_{xz}(uv) \equiv 0 \pmod{2}$ . This means that  $xz$  cannot satisfy Item 2 (because in that case the edges with colour 0 form a bipartite graph), so  $xz$  is even.  $\square$

Claim 1 implies that the even pairs induce cliques (since if  $xy$  and  $yz$  are even, then  $xz$  is even) and that the number of disjoint even cliques is at most 2 (since if  $xy$  and  $yz$  are odd, then  $xz$  is even). In other words, there is a partition of  $[n]$  into  $X \cup Y$  such that every pair of vertices which is fully contained in  $X$  or fully contained in  $Y$  is even, and every other pair is odd. We will show that either  $X \cup Y$  or  $[n] \cup \emptyset$  is the desired partition.

We start by proving that almost all edges of any given type have the same colour. There are four edge types:  $XXX$  (i.e., edges entirely contained in  $X$ );  $XXY$  (i.e., edges with two vertices in  $X$  and one in  $Y$ ); and similarly  $YYX$  and  $YYY$ .

*Claim 2.* There exist colours  $c_1, c_2, c_3, c_4 \in \{0, 1\}$  such that all but at most  $10^8 |X|^2$  edges of type  $XXX$  have colour  $c_1$ , all but at most  $10^8 |X| |n|$  edges of type  $XXY$  have colour  $c_2$ , all but at most  $10^8 |Y| |n|$  edges of type  $XYX$  have colour  $c_3$ , and all but at most  $10^8 |Y|^2$  edges of type  $YYY$  have colour  $c_4$ .

*Proof.* We start with the edges of type  $XXX$  and let  $m := 10^8 |X|^2$ . First note that we can assume  $|X| \geq 360$ , as otherwise the total number of edges inside  $X$  is smaller than  $m$ . We upper-bound the number of pairs of edges  $e, e' \subseteq X$  with  $\chi(e) \neq \chi(e')$  and  $|e \cap e'| = 2$ , as follows. Write  $e := xuv, e' := x'uv$ . We first choose  $x, x'$ . Since  $xx'$  is even, if  $u, v \notin S_{xx'}$  then  $f_{xx'}(uv) = 0$  and hence  $\chi(xuv) = \chi(x'uv)$ . Therefore we must have  $u \in S_{xx'}$  or  $v \in S_{xx'}$ , and thus there are at most  $8|X|$  choices of  $uv$  (because  $|S_{xx'}| \leq 8$ ). Hence, the total number of pairs of edges  $e, e' \subseteq X$  with  $\chi(e) \neq \chi(e')$  and  $|e \cap e'| = 2$  is at most  $\binom{|X|}{2} \cdot 8|X| < \frac{m|X|}{900}$ . By Lemma 3.4 (applied on input  $A = B = C = X$ ), there must be  $c_1 \in \{0, 1\}$  such that all but at most  $m$  of the edges of type  $XXX$  have colour  $c_1$ .



We now move to edges of type  $XXY$  and let  $m' := 10^8|X|n$ . As before, we can assume  $|X|, |Y| \geq 360$ . We bound the number of pairs of edges  $e, e'$  of type  $XXY$  with  $\chi(e) \neq \chi(e')$ ,  $|e \cap e'| = 2$  and  $e \triangle e' \subseteq Y$ , as follows. Write  $e := yuv, e' := y'uv$  with  $u, v \in X$  and  $y, y' \in Y$ . We first choose  $y, y'$ . Since  $yy'$  is even, in order to have  $\chi(e) \neq \chi(e')$ , we must have  $u \in S_{yy'}$  or  $v \in S_{yy'}$ , and thus there are at most  $8|X|$  choices of  $uv$  (because  $|S_{yy'}| \leq 8$ ). Hence, the number of pairs of edges  $e, e'$  of type  $XXY$  with  $\chi(e) \neq \chi(e')$ ,  $|e \cap e'| = 2$  and  $e \triangle e' \subseteq Y$  is at most  $\binom{|Y|}{2} \cdot 8|X| < \frac{m'|Y|}{900}$ . By the same argument, the number of pairs of edges  $e, e'$  of type  $XXY$  with  $\chi(e) \neq \chi(e')$ ,  $|e \cap e'| = 2$  and  $e \triangle e' \subseteq X$  is at most  $\binom{|X|}{2} \cdot 8n < \frac{m'|X|}{900}$ . Therefore, by Lemma 3.4 (applied on input  $A = B = X$  and  $C = Y$ ), there must be  $c_2 \in \{0, 1\}$  such that all but at most  $m'$  of the edges of type  $XXY$  have colour  $c_2$ .

The two remaining types  $YYX$  and  $YYY$  are symmetric.  $\square$

If  $|X| \leq 10^{12}$ , then the number of edges touching  $X$  is at most  $|X|n^2 \leq 10^{12}n^2$ . We may then recolour these  $O(n^2)$  edges, as well as at most  $O(n^2)$  other edges (using Claim 2), to obtain a colouring for which the assertion of the proposition holds (for the partition  $X \cup Y$ ). Thus, we may assume that  $|X| > 10^{12}$  and similarly  $|Y| > 10^{12}$ .

An edge of type  $XXX$  (resp.  $XXY, XYY, YYY$ ) is called *good* if it has colour  $c_1$  (resp.  $c_2, c_3, c_4$ ). Otherwise, the edge is *bad*. Sample a set  $S$  of six distinct vertices, three from  $X$  and three from  $Y$ , uniformly at random. We claim that with positive probability, all  $\binom{6}{3} = 20$  edges inside  $S$  are good. Indeed let  $e$  be an edge of  $K_n^{(3)}$  with  $i$  vertices in  $X$  and  $3 - i$  vertices in  $Y$  (so  $0 \leq i \leq 3$ ). The probability that  $e \subseteq S$  is

$$\frac{\binom{|X|-i}{3-i}}{\binom{|X|}{3}} \cdot \frac{\binom{|Y|-(3-i)}{i}}{\binom{|Y|}{3}} \leq 729|X|^{-i}|Y|^{i-3}.$$

Using Claim 2, the expected number of bad edges  $e \subseteq S$  of type  $XXX$  is at most  $10^8|X|^2 \cdot 729|X|^{-3} < 1/4$  (where the inequality uses that  $|X| \geq 10^{12}$ ). By symmetry, the expected number of bad edges  $e \subseteq S$  of type  $YYY$  is less than  $1/4$ . Similarly, the expected number of bad edges  $e \subseteq S$  of type  $XXY$  is at most  $10^8|X|n \cdot 729|X|^{-2}|Y|^{-1} < 1/4$  (using that, since  $|X|, |Y| \geq 10^{12}$  and  $n$  is large, we have  $|X| \cdot |Y| \geq 10^{12}(n - 10^{12}) \geq 10^{12}n/2$ ). By symmetry, the same holds for edges of type  $XYY$ . By linearity of expectation, the total number of bad edges  $e \subseteq S$  is less than 1, and hence there is an outcome in which all edges in  $S$  are good.

Fix a set  $S$  for which this holds and write  $S \cap X =: \{x_1, x_2, x_3\}$  and  $S \cap Y =: \{y_1, y_2, y_3\}$ . Let  $\mathcal{P}_1 := \{x_1x_2x_3, x_1y_2y_3, y_1x_2y_3, y_1y_2x_3\}$  and  $\mathcal{P}_2 := \{x_1x_2y_3, x_1y_2x_3, y_1x_2x_3, y_1y_2y_3\}$ . Observe that  $\mathcal{P}_1$  and  $\mathcal{P}_2$  are two edge-disjoint Pasch configurations,  $\mathcal{P}_1$  has exactly one edge of colour  $c_1$  and three edges of colour  $c_3$ , while  $\mathcal{P}_2$  has exactly one edge of colour  $c_4$  and three edges of colour  $c_2$ . Moreover  $\mathcal{P}_1 \cup \mathcal{P}_2$  is a copy of  $K_{2,2,2}^{(3)}$  with parts  $\{x_i, y_i\}$  for  $i = 1, 2, 3$ . Observe that in order for this copy to not be a gadget, we must have  $c_1 = c_4$  and  $c_2 = c_3$ .

If  $c_1 \neq c_2$ , then the partition  $X \cup Y$  satisfies the assertion of the proposition. Otherwise,  $c_1 = c_2 = c_3 = c_4 =: c$ , and all but at most  $4 \cdot 10^8n^2$  edges of  $K_n^{(3)}$  have colour  $c$ , so we can take the partition  $[n] \cup \emptyset$ .  $\square$

## 4 From many gadgets to high discrepancy

In this section, we prove that if a colouring of  $E(K_n^{(3)})$  contains many gadgets, then it admits a Steiner triple system with high discrepancy.

**Theorem 4.1.** *For every  $r \in \mathbb{N}$  with  $r \geq 2$  and  $\delta > 0$ , there exist  $n_0 \in \mathbb{N}$  and  $\mu > 0$  such that for every  $n \geq n_0$  with  $n \equiv 1, 3 \pmod{6}$  the following holds. Consider an  $r$ -edge-colouring of  $K_n^{(3)}$  and assume that there exists a collection of  $\delta n^6$  gadgets. Then there is a Steiner triple system with at least  $\frac{1}{r} \cdot \frac{1}{3} \binom{n}{2} + \mu n^2$  triples of the same colour.*

*Proof.* Let  $\varepsilon := 10^{-5}$  and assume  $n$  is sufficiently large with  $n \equiv 1, 3 \pmod{6}$ . Let  $\mathcal{K}$  denote the collection of at least  $\delta n^6$  gadgets. Recall that the *shadow* of a 3-uniform hypergraph  $P$  is the (2-uniform) graph consisting of all pairs which are contained in some edge of  $P$ . Also, two graphs are *edge-disjoint* if they share no edges. We refine  $\mathcal{K}$  to a subcollection containing quadratically many gadgets, which do not use the same vertex too much, and whose shadows are pairwise edge-disjoint. We will do so by sampling the gadgets with an appropriate probability.

We first show that, for  $p := \varepsilon n^{-4}$ , there exists  $\mathcal{J} \subseteq \mathcal{K}$  such that the following hold:

(P1)  $|\mathcal{J}| \geq |\mathcal{K}|p/2$ ;

(P2) denoting by  $\mathcal{Y}$  the set of pairs of gadgets of  $\mathcal{J}$  whose shadows are not edge-disjoint, we have  $|\mathcal{Y}| < |\mathcal{K}|p/100$ ;

(P3) every vertex of  $K_n^{(3)}$  is contained in at most  $7\varepsilon n$  gadgets of  $\mathcal{J}$ .

Let  $\mathcal{J}$  be obtained from  $\mathcal{K}$  by sampling each gadget with probability  $p$ , independently. We show that with positive probability all the properties (P1), (P2) and (P3) hold. First, note that  $\mathbb{E}[|\mathcal{J}|] = |\mathcal{K}| \cdot p \geq \delta \varepsilon n^2$  and thus, by the Chernoff bound (Lemma 2.4), we have  $|\mathcal{J}| \geq |\mathcal{K}|p/2$  with probability at least 0.9. Next, observe that there are at most  $n^4$  copies of  $K_{2,2,2}^{(3)}$  in  $K_n^{(3)}$  whose shadow contains a given (2-uniform) edge, and that the shadow of  $K_{2,2,2}^{(3)}$  contains 12 edges. Therefore, given a gadget  $K$ , there are at most  $12n^4$  other gadgets whose shadow shares an edge with  $K$ . It follows that  $\mathbb{E}[|\mathcal{Y}|] \leq |\mathcal{K}| \cdot (12n^4) \cdot p^2 \leq 12|\mathcal{K}|p\varepsilon$  and, by Markov's inequality,  $\mathbb{P}[|\mathcal{Y}| \geq |\mathcal{K}|p/100] \leq 1200\varepsilon \leq 0.1$ . Finally, given  $v \in V(K_n^{(3)})$ , let  $X_v$  be the collection of gadgets of  $\mathcal{J}$  containing the vertex  $v$ . Then  $\mathbb{E}[|X_v|] \leq n^5 \cdot p = \varepsilon n$ , using that there are at most  $n^5$  copies of  $K_{2,2,2}^{(3)}$  in  $K_n^{(3)}$  containing  $v$ . Then, using the second part of Lemma 2.4, we have  $\mathbb{P}[|X_v| > 7\varepsilon n] \leq \exp(-7\varepsilon n)$ , and by a union bound over the  $n$  vertices, with probability at least 0.9, we have  $|X_v| \leq 7\varepsilon n$  for every  $v \in V(K_n^{(3)})$ . Overall, this proves the existence of a suitable  $\mathcal{J} \subseteq \mathcal{K}$ .

Let  $\mathcal{J} \subseteq \mathcal{K}$  satisfy the properties (P1)–(P3) and define  $\mathcal{I}$  by removing from  $\mathcal{J}$  all gadgets appearing in a pair in  $\mathcal{Y}$ . Then  $|\mathcal{I}| \geq |\mathcal{K}|p/3 \geq \delta \varepsilon n^2/3$ , the shadows of the gadgets in  $\mathcal{I}$  are pairwise edge-disjoint and every vertex of  $K_n^{(3)}$  is contained in at most  $7\varepsilon n$  gadgets of  $\mathcal{I}$ .

Let  $G$  be the  $n$ -vertex (2-uniform) graph obtained from  $K_n$  by removing the edges which are in the shadow of a gadget in  $\mathcal{I}$ . Since we remove 12 (2-uniform) edges per gadget,  $e(G)$  is divisible by 3. Moreover, for each gadget and each vertex  $v$  contained in the gadget, we remove exactly 4 edges adjacent to  $v$  and, since the shadows of the gadgets in  $\mathcal{I}$  are pairwise edge-disjoint, the removed edges adjacent to  $v$  are distinct for distinct gadgets. Therefore  $G$  is  $K_3$ -divisible and  $\delta(G) \geq n - 1 - 4 \cdot 7\varepsilon n \geq 0.91n$ . It follows from Theorem 2.1 that  $G$  has a triangle decomposition  $\mathcal{T}$ . Observe that the number of triples in  $\mathcal{T}$  is exactly

$$|\mathcal{T}| = \frac{\binom{n}{2} - 12|\mathcal{I}|}{3}. \quad (4.1)$$

We now show that by choosing the appropriate Pasch configuration in each gadget of  $\mathcal{I}$ , we get an STS with high discrepancy. Note that each Pasch configuration in a  $K_{2,2,2}^{(3)}$  forms a triangle decomposition of the shadow of the  $K_{2,2,2}^{(3)}$ . Hence, choosing a Pasch configuration for each gadget completes the triangle-decomposition  $\mathcal{T}$  of  $G$  to an STS.

Let  $[r]$  be the colour-set of  $E(K_n^{(3)})$ . For each gadget  $I \in \mathcal{I}$  and colour  $c \in [r]$ , let  $I_c$  be the largest number of edges in colour  $c$  among the two Pasch configurations of  $I$ . We claim that

$$\sum_{c \in [r]} I_c \geq 5. \quad (4.2)$$

Indeed, let  $P_1, P_2$  be the two Pasch configurations in  $I$ , and let  $a_c$  (resp.  $b_c$ ) be the number of edges of colour  $c$  in  $P_1$  (resp.  $P_2$ ). Then  $I_c = \max(a_c, b_c)$ , so  $I_c \geq \frac{a_c + b_c}{2}$ . Also, since  $I$  is a

gadget, there is a colour  $c^*$  with  $a_{c^*} \neq b_{c^*}$ , and so  $I_{c^*} > \frac{a_{c^*} + b_{c^*}}{2}$ . It follows that  $\sum_{c \in [r]} I_c > \frac{1}{2} \sum_{c \in [r]} (a_c + b_c) = \frac{1}{2} (e(P_1) + e(P_2)) = 4$ , implying (4.2).

For each colour  $c \in [r]$ , let  $T_c$  be the number of triples of colour  $c$  in  $\mathcal{T}$ . By choosing for each  $I \in \mathcal{I}$  the Pasch configuration with  $I_c$  edges of colour  $c$ , we get a Steiner triple system with  $S_c := T_c + \sum_{I \in \mathcal{I}} I_c$  edges in colour  $c$ . We have

$$\sum_{c \in [r]} S_c = \sum_{c \in [r]} T_c + \sum_{c \in [r]} \sum_{I \in \mathcal{I}} I_c \geq \frac{\binom{n}{2} - 12|\mathcal{I}|}{3} + 5 \cdot |\mathcal{I}| = \frac{1}{3} \binom{n}{2} + |\mathcal{I}| \geq \frac{1}{3} \binom{n}{2} + \frac{\delta \varepsilon n^2}{3},$$

where the first inequality uses (4.1) and (4.2). By averaging, there is a colour  $c \in [r]$  such that

$$S_c \geq \frac{1}{r} \cdot \left[ \frac{1}{3} \binom{n}{2} + \frac{\delta \varepsilon n^2}{3} \right],$$

which proves the theorem with  $\mu := (\delta \varepsilon)/(3r)$ .  $\square$

## 5 Edge-colourings with more than two colours

Here we use the structure theorem (Theorem 3.1) to show that for  $r \geq 3$ , if an  $r$ -edge-colouring of  $K_n^{(3)}$  does not have many gadgets, then all but at most two colour classes are tiny.

**Theorem 5.1.** *Let  $r \in \mathbb{N}$  with  $r \geq 3$  and  $\zeta > 0$ . Then there exists  $\delta > 0$  such that for every  $r$ -edge-colouring of  $K_n^{(3)}$ , (at least) one of the following properties holds:*

- (1) *There are two colours  $c^*, d^*$  such that at most  $\zeta n^3$  edges have colour different from  $c^*, d^*$ ;*
- (2) *There are at least  $\delta n^6$  gadgets.*

*Proof.* Fix a new constant  $\alpha$  such that  $\delta \ll \alpha \ll \zeta, 1/r$ . Let the colour-set be  $[r]$  and assume that Item (2) in the statement of the theorem does not hold, i.e. that  $K_n^{(3)}$  contains less than  $\delta n^6$  gadgets. For a colour  $c \in [r]$ , let  $\chi_c : E(K_n^{(3)}) \rightarrow \{c, \bar{c}\}$  be the colouring of  $K_n^{(3)}$  which identifies the colours in  $[r] \setminus \{c\}$  into one (new) colour  $\bar{c}$ . I.e.,  $\chi_c(e) = c$  if  $\chi(e) = c$  and  $\chi_c(e) = \bar{c}$  otherwise. Then  $\chi_c$  has less than  $\delta n^6$  gadgets, because a gadget in  $\chi_c$  is also a gadget in the original colouring. Hence, by Theorem 3.1, there is a partition  $[n] = X_c \cup Y_c$  such that after changing the colours of at most  $\alpha n^3$  edges, all edges inside  $X$  and  $Y$  have colour  $a_c$ , and all edges intersecting both  $X_c$  and  $Y_c$  have colour  $b_c$ , where  $\{a_c, b_c\} = \{c, \bar{c}\}$ . We call  $a_c$  the colour *inside* the parts  $X_c, Y_c$ , and  $b_c$  the colour *across* the partition  $(X_c, Y_c)$ .

In particular, by considering the Venn diagram of the partitions  $(X_c, Y_c)_{c \in [r]}$  and averaging, we get a set  $Z$  of size at least  $n/2^r$  such that  $Z \subseteq X_c$  or  $Z \subseteq Y_c$  for each  $c \in [r]$ . By averaging, there is a colour  $c^*$  such that at least  $\frac{1}{r} \binom{|Z|}{3} \geq 2\alpha n^3$  edges inside  $Z$  have colour  $c^*$ . Therefore  $c^*$  is the colour inside the part  $X_{c^*}$  and  $Y_{c^*}$ , that is  $a_{c^*} = c^*$ . If  $|X_{c^*}| \leq \zeta n/2$ , then the number of edges of colour different from  $c^*$  (in  $\chi_{c^*}$  and thus in the original colouring of  $K_n^{(3)}$ ) is at most  $|X_{c^*}|n^2 + \alpha n^3 \leq \zeta n^3/2 + \alpha n^3 < \zeta n^3$ , so Item (1) in the statement of the theorem holds (in fact, all colour classes but one are tiny). Similarly, Item (1) holds if  $|Y_{c^*}| \leq \zeta n/2$ . Therefore, from now on, we can assume  $|X_{c^*}|, |Y_{c^*}| > \zeta n/2$ .

The number of edges of colour distinct from  $c^*$  (in  $\chi_{c^*}$  and thus in the original colouring) is at least

$$\begin{aligned} \binom{|X_{c^*}|}{2} \cdot |Y_{c^*}| + |X_{c^*}| \cdot \binom{|Y_{c^*}|}{2} - \alpha n^3 &= \frac{|X_{c^*}| \cdot |Y_{c^*}| \cdot (n-2)}{2} - \alpha n^3 \\ &\geq \zeta n^3/16 - \alpha n^3 \geq \zeta n^3/20, \end{aligned}$$

where we used that  $|X_{c^*}| + |Y_{c^*}| = n$ ,  $|X_{c^*}|, |Y_{c^*}| > \zeta n/2$  and so  $|X_{c^*}| \cdot |Y_{c^*}| > \zeta n^2/4$ . By averaging, there is a colour  $d^* \in [r] \setminus \{c^*\}$  such that the number of edges of colour  $d^*$  is at least  $\zeta n^3/(20r)$ .

We now show that  $d^*$  is the colour across in  $\chi_{d^*}$ . (In fact, the same argument shows that for any  $d \in [r] \setminus \{c^*\}$ ,  $d$  is the colour across in  $\chi_d$ .) Let  $A := X_{c^*} \cap X_{d^*}$ ,  $B := X_{c^*} \cap Y_{d^*}$ ,  $C := Y_{c^*} \cap X_{d^*}$  and  $D := Y_{c^*} \cap Y_{d^*}$ . Then at least one of these four sets has size at least  $n/4$ . Without loss of generality, suppose that  $|A| \geq n/4$ . Since  $A \subseteq X_{c^*}$  and  $a_{c^*} = c^*$ , the number of edges of colour  $c^*$  inside  $A$  is at least  $\binom{|A|}{3} - \alpha n^3 \geq 2\alpha n^3$ . Since  $d^* \neq c^*$  and  $A \subseteq X_{d^*}$ , then  $d^*$  cannot be the colour inside the parts in  $\chi_{d^*}$ , as wanted.

Moreover, we must have  $|X_{d^*}|, |Y_{d^*}| > \zeta n/(5r)$  as otherwise, by a similar calculation as above, the number of edges of colour  $d^*$  would be at most  $\binom{|X_{d^*}|}{2} \cdot |Y_{d^*}| + |X_{d^*}| \cdot \binom{|Y_{d^*}|}{2} + \alpha n^3 < \zeta n^3/(20r)$ , contradicting the choice of  $d^*$ . Next, we show that the partitions  $(X_{c^*}, Y_{c^*})$  and  $(X_{d^*}, Y_{d^*})$  do not differ too much, in the following sense.

*Claim 3.* Let  $m := \zeta n/(10r)$ . We have  $|A| < m$  or  $|B| < m$ . Similarly,  $|C| < m$  or  $|D| < m$ .

*Proof.* We only prove the first statement, the second one can be proved similarly. So suppose by contradiction that  $|A|, |B| \geq m$ . As  $b_{d^*} = d^*$ , the number of the edges across  $(A, B)$  of colour  $d^*$  is at least  $\binom{|A|}{2} \cdot |B| - \alpha n^3 \geq 2\alpha n^3$ . On the other hand, as  $A \cup B \subseteq X_{c^*}$  and  $a_{c^*} = c^*$ , all but at most  $\alpha n^3$  of the edges inside  $A \cup B$  have colour  $c^*$ , a contradiction.  $\square$

As already mentioned, we can assume that  $|A| \geq n/4$ . Then, by Claim 3,  $|B| < \zeta n/(10r)$ . Since  $B \cup D = Y_{d^*}$  and  $|Y_{d^*}| > \zeta n/(5r)$ , we must have  $|D| > \zeta n/(10r)$ . Therefore, again by Claim 3, we have  $|C| < \zeta n/(10r)$ . Then  $|B \cup C| < \zeta n/(5r)$  and the number of edges touching  $B \cup C$  is at most  $|B \cup C| \cdot n^2 < \zeta n^3/(5r)$ . As  $a_{c^*} = c^*$ , all but  $\alpha n^3$  of the edges inside  $A$  and inside  $D$  have colour  $c^*$ . Also, as  $b_{d^*} = d^*$ , all but at most  $\alpha n^3$  of the edges across  $(A, D)$  have colour  $d^*$ . This means that the number of edges whose colour is neither  $c^*$  nor  $d^*$  is at most  $2\alpha n^3 + \zeta n^3/(5r) < \zeta n^3$ . In particular, Item (1) in the statement of the theorem holds.  $\square$

## 6 Proofs of main theorems

We are now ready to prove our main theorems, concerning Steiner triple systems with high discrepancy in  $r$ -edge-colourings of  $K_n^{(3)}$ . We start with the case  $r \geq 3$ .

*Proof of Theorem 1.3.* Let  $r \geq 3$  and  $1/n \ll \mu \ll \delta \ll \zeta, 1/r$ . Let  $K_n^{(3)}$  be edge-coloured in  $[r]$ . Then by Theorem 5.1, there exists a colour  $c$  such that the number of edges of colour  $c$  is at most  $\zeta n^3$ , or there are at least  $\delta n^6$  gadgets. In the first case, the result follows from Fact 2.2. In the second case, we conclude using Theorem 4.1.  $\square$

Finally, we move to the case  $r = 2$ .

*Proof of Theorem 1.2.* Given  $\eta$ , define new constants such that

$$1/n \ll \mu \ll \delta \ll \alpha \ll \beta \ll \gamma \ll \eta, 1/r.$$

Let  $K_n^{(3)}$  be 2-edge-coloured. By Theorem 3.1, there are at least  $\delta n^6$  gadgets or there is a partition  $[n] = X \cup Y$  such that after changing the colour of at most  $\alpha n^3$  edges, which we call the *bad* edges, all edges inside  $X$  or inside  $Y$  have the same colour, say red, and all edges intersecting both  $X$  and  $Y$  have the other colour, say blue. In the first case (i.e., if there are at least  $\delta n^6$  gadgets), Theorem 4.1 immediately gives an STS with discrepancy at least  $\mu n^2$ .

In the second case, we consider the sizes of  $X$  and  $Y$ . Put  $x := |X|/n$ , so  $|Y| = (1-x)n$ . Put  $z := x^3 + (1-x)^3$ . Then the number<sup>3</sup> of red edges in  $K_n^{(3)}$  is  $(z + o(1))\binom{n}{3} \pm \alpha n^3 = (z \pm 7\alpha)\binom{n}{3}$ . Hence, if  $|z - 1/2| \geq 2\beta$ , then, as  $\beta \gg \alpha$ , the number of red edges is at least  $(1/2 + \beta)\binom{n}{3}$  or at most  $(1/2 - \beta)\binom{n}{3}$ . Now Fact 2.2 gives an STS with discrepancy at least  $\mu n^2$ .

Suppose now that  $|z - 1/2| < 2\beta$ . The solutions to  $x^3 + (1-x)^3 = 1/2$  are  $x = \frac{3 \pm \sqrt{3}}{6} \in [0, 1]$ . By continuity, and as  $\beta \ll \gamma$ , we have that  $|x - \frac{3+\sqrt{3}}{6}| \leq \gamma$  or  $|x - \frac{3-\sqrt{3}}{6}| \leq \gamma$ . These two cases are symmetric (by replacing  $x$  with  $1-x$ ), so assume that  $|x - \frac{3+\sqrt{3}}{6}| \leq \gamma$ . Then, by moving  $O(\gamma n)$  vertices between  $X$  and  $Y$ , we can make sure that  $|X| = \lfloor \frac{3+\sqrt{3}}{6}n \rfloor$ . By recolouring the edges touching the moved vertices, as well as the at most  $\alpha n^3$  bad edges, we obtain that all edges inside  $X$  and inside  $Y$  are red and all edges intersecting both  $X, Y$  are blue. As  $\alpha, \gamma \ll \eta$ , this recolours at most  $\eta n^3$  edges. After this recolouring, we have the construction given by Example 1.1, as required.  $\square$

## 7 Concluding remarks

It would be interesting to extend our results to general designs. Given  $k, \ell \in \mathbb{N}$ , a  $(k, \ell)$ -Steiner system of order  $n$  is a set  $\mathcal{S}$  of  $k$ -subsets of  $[n]$  such that every  $\ell$ -subset of  $[n]$  is contained in exactly one of the sets of  $\mathcal{S}$ . Thus, an STS corresponds to  $(k, \ell) = (3, 2)$ . When general Steiner systems exist was a long-standing open problem, until finally settled by Keevash [22], who showed that they exist whenever  $n$  is sufficiently large and satisfies a necessary ‘‘divisibility condition’’. An alternative proof of Keevash’s result and a minimum degree version analogous to Theorem 2.1 was proved in [18]. In the following, we will always assume that  $n$  is such that a  $(k, \ell)$ -Steiner system of order  $n$  exists. It would be interesting to understand for which values of  $k, \ell, r$  it holds that every  $r$ -edge-colouring of  $K_n^{(k)}$  contains a  $(k, \ell)$ -Steiner system of order  $n$  with high discrepancy.

A natural starting point would be to generalize Example 1.1 as follows. Let  $k \geq 3$ , and partition  $[n]$  into  $k-1$  sets  $X_1, \dots, X_{k-1}$ . Assign colour blue to all  $k$ -sets which intersect all of the sets  $X_1, \dots, X_{k-1}$ , and colour red to all other  $k$ -sets. Every  $(k-1)$ -subset in  $X_1 \times \dots \times X_{k-1}$  must be covered by a blue  $k$ -set, and every blue  $k$ -set covers exactly two  $(k-1)$ -subsets in  $X_1 \times \dots \times X_{k-1}$ . Therefore, the number of blue  $k$ -sets in any  $(k, k-1)$ -Steiner system is always  $\frac{|X_1| \dots |X_{k-1}|}{2}$ . Unfortunately, for  $k \geq 4$ , one cannot choose the sizes of  $X_1, \dots, X_{k-1}$  such that  $\frac{|X_1| \dots |X_{k-1}|}{2}$  is roughly half the number of edges of a  $(k, k-1)$ -Steiner system. Indeed, a  $(k, k-1)$ -Steiner system has  $\binom{n}{k}$  edges, whereas the AM-GM inequality gives  $|X_1| \dots |X_{k-1}| \leq (\frac{n}{k-1})^{k-1}$ . As  $\frac{1}{(k-1)^{k-1}} < \frac{1}{(k-1)!k}$  for  $k \geq 4$ , this means that  $\frac{|X_1| \dots |X_{k-1}|}{2}$  is always strictly less than half the number of edges in a  $(k, k-1)$ -Steiner system.

This suggests the possibility that for  $k \geq 4$ , every 2-edge-colouring (and also  $r$ -edge-colouring for  $r \geq 3$ ) of  $K_n^{(k)}$  has a  $(k, k-1)$ -Steiner system with high discrepancy. One way to try and prove this would be to generalize our approach in this paper. The suitable generalization of a gadget is as follows. Consider a copy  $K$  of  $K_{2,2,\dots,2}^{(k)}$  (the complete  $k$ -partite  $k$ -graph with parts of size 2). Let  $\{a_1, b_1\}, \dots, \{a_k, b_k\}$  be the parts of  $K$ , let  $A := \{a_1, \dots, a_k\}$  and  $B := \{b_1, \dots, b_k\}$ , and let  $P_1 := \{e \in E(K) : |e \cap A| \text{ even}\}$  and  $P_2 := E(K) \setminus P_1 = \{e \in E(K) : |e \cap A| \text{ odd}\}$ . It is easy to see that each  $(k-1)$ -tuple in the shadow of  $K$  belongs to exactly one edge of  $P_1$  and exactly one edge of  $P_2$ .<sup>4</sup> We say that  $K$  is a *gadget* if there is a colour  $c$  such that  $P_1, P_2$  have a different number of edges of colour  $c$ . One can show that if a 2-edge-colouring of  $K_n^{(k)}$  has  $\Omega(n^{2k})$  gadgets, then there exists a  $(k, k-1)$ -Steiner system with high discrepancy. We wonder if the following analogue of Proposition 3.2 is true.

<sup>3</sup>We use the notation  $a = b \pm c$  to mean that  $b - c \leq a \leq b + c$ .

<sup>4</sup>In the case  $k = 3$ ,  $P_1, P_2$  are Pasch configurations.



**Problem 7.1.** Is the following true? If a 2-edge-colouring of  $K_n^{(k)}$  has no gadgets, then there is a partition  $[n] = X_1 \cup \dots \cup X_{k-1}$  such that, after changing the colour of  $o(n^k)$  edges, all edges which intersect all of the sets  $X_1, \dots, X_{k-1}$  have the same colour, and all other edges have the other colour.

## References

- [1] J. Ai, F. He, S. Im, and H. Lee, *On high discrepancy 1-factorizations of complete graphs*, arXiv:2503.17176, 2025.
- [2] J. Balogh, B. Csaba, Y. Jing, and A. Pluhár, *On the discrepancies of graphs*, Electron. J. Combin. **27** (2020), Paper No. 2.12, 14.
- [3] J. Balogh, B. Csaba, A. Pluhár, and A. Treglown, *A discrepancy version of the Hajnal–Szemerédi theorem*, Combin. Probab. Comput. **30** (2021), 444–459.
- [4] J. Balogh, A. Treglown, and C. Zárate-Guerén, *A note on color-bias perfect matchings in hypergraphs*, SIAM J. Discrete Math. **38** (2024), 2543–2552.
- [5] B. Barber, S. Glock, D. Kühn, A. Lo, R. Montgomery, and D. Osthus, *Minimalist designs*, Random Structures Algorithms **57** (2020), 47–63.
- [6] B. Barber, D. Kühn, A. Lo, and D. Osthus, *Edge-decompositions of graphs with high minimum degree*, Adv. Math. **288** (2016), 337–385.
- [7] D. Bradač, *Powers of Hamilton cycles of high discrepancy are unavoidable*, Electron. J. Combin. **29** (2022), P3.22.
- [8] D. Bradač, M. Christoph, and L. Gishboliner, *Minimum degree threshold for  $H$ -factors with high discrepancy*, Electron. J. Combin. **31** (2024), Paper No. 3.33, 82.
- [9] C. J. Colbourn and J. H. Dinitz (eds.), *Handbook of Combinatorial Designs*, 2nd ed., Discrete Math. Appl. (Boca Raton), Chapman & Hall/CRC, 2006.
- [10] P. Erdős, P. Frankl, and V. Rödl, *The asymptotic number of graphs not containing a fixed subgraph and a problem for hypergraphs having no exponent*, Graphs Combin. **2** (1986), 113–121.
- [11] P. Erdős, Z. Füredi, M. Loeb, and V. T. Sós, *Discrepancy of trees*, Studia Sci. Math. Hungar. **30** (1995), 47–57.
- [12] P. Erdős and J. H. Spencer, *Imbalances in  $k$ -colorations*, Networks **1** (1971/72), 379–385.
- [13] A. Freschi, J. Hyde, J. Lada, and A. Treglown, *A note on color-bias Hamilton cycles in dense graphs*, SIAM J. Discrete Math. **35** (2021), 970–975.
- [14] L. Gishboliner, S. Glock, P. Michaeli, and A. Sgueglia, *Defect and transference versions of the Alon–Frankl–Lovász theorem*, arXiv:2503.05089, 2025.
- [15] L. Gishboliner, S. Glock, and A. Sgueglia, *Tight Hamilton cycles with high discrepancy*, Combin. Probab. Comput., to appear.
- [16] L. Gishboliner, M. Krivelevich, and P. Michaeli, *Color-biased Hamilton cycles in random graphs*, Random Structures Algorithms **60** (2022), 289–307.
- [17] ———, *Discrepancies of spanning trees and Hamilton cycles*, J. Combin. Theory Ser. B **154** (2022), 262–291.

- [18] S. Glock, D. Kühn, A. Lo, and D. Osthus, *The existence of designs via iterative absorption: hypergraph  $F$ -designs for arbitrary  $F$* , Mem. Amer. Math. Soc. **284** (2023), monograph 1406.
- [19] W. T. Gowers, *Hypergraph regularity and the multidimensional Szemerédi theorem*, Ann. of Math. (2) **166** (2007), 897–946.
- [20] H. Hàn, R. Lang, J. P. Marciano, M. Pavez-Signé, N. Sanhueza-Matamala, A. Treglown, and C. Zárate-Guerén, *Colour-bias perfect matchings in hypergraphs*, arXiv:2408.11016, 2024.
- [21] S. Janson, T. Łuczak, and A. Ruciński, *Random graphs*, Wiley-Intersci. Ser. Discrete Math. Optim., Wiley-Interscience, 2000.
- [22] P. Keevash, *The existence of designs*, arXiv:1401.3665, 2014.
- [23] T. P. Kirkman, *On a problem in combinatorics*, Cambridge Dublin Math. J. **2** (1847), 191–204.
- [24] H. Lu, J. Ma, and S. Xie, *Discrepancies of perfect matchings in hypergraphs*, arXiv:2408.06020, 2024.
- [25] B. Nagle, V. Rödl, and M. Schacht, *The counting lemma for regular  $k$ -uniform hypergraphs*, Random Structures Algorithms **28** (2006), 113–179.
- [26] C. St. J. A. Nash-Williams, *An unsolved problem concerning decomposition of graphs into triangles*, In: Combinatorial Theory and its Applications III (P. Erdős, A. Rényi, and V.T. Sós, eds.), North Holland, 1970, pp. 1179–1183.
- [27] V. Rödl and M. Schacht, *Generalizations of the removal lemma*, Combinatorica **29** (2009), 467–501.
- [28] V. Rödl and J. Skokan, *Regularity lemma for  $k$ -uniform hypergraphs*, Random Structures Algorithms **25** (2004), 1–42.
- [29] ———, *Applications of the regularity lemma for uniform hypergraphs*, Random Structures Algorithms **28** (2006), 180–194.
- [30] T. Tao, *A variant of the hypergraph removal lemma*, J. Combin. Theory Ser. A **113** (2006), 1257–1280.