Information-theoretic subset selection of multivariate Markov chains via submodular optimization

Zheyuan Lai^{*1} and Michael C.H. Choi^{†2}

¹Department of Statistics and Data Science, National University of Singapore, Singapore ²Department of Statistics and Data Science and Yale-NUS College, National University of Singapore, Singapore

April 1, 2025

Abstract

We study the problem of optimally projecting the transition matrix of a finite ergodic multivariate Markov chain onto a lower-dimensional state space. Specifically, we seek to construct a projected Markov chain that optimizes various information-theoretic criteria under cardinality constraints. These criteria include entropy rate, information-theoretic distance to factorizability, independence, and stationarity. We formulate these tasks as best subset selection problems over multivariate Markov chains and leverage the submodular (or supermodular) structure of the objective functions to develop efficient greedy-based algorithms with theoretical guarantees. We extend our analysis to k-submodular settings and introduce a generalized version of the distorted greedy algorithm, which may be of independent interest. Finally, we illustrate the theory and algorithms through extensive numerical experiments with publicly available code on multivariate Markov chains associated with the Bernoulli-Laplace and Curie-Weiss model.

Keywords: Markov chains, submodularity, greedy algorithms, Kullback-Leibler divergence AMS 2020 subject classification: 60J10, 60J22, 90C27, 94A15, 94A17

1 Introduction

Consider a multivariate ergodic Markov chain with transition matrix P that admits a stationary distribution π on a finite product state space \mathcal{X} with $d \in \mathbb{N}$ coordinates. Given a subset $S \subseteq \{1, \ldots, d\}$, let $P^{(S)}$ denote the projected transition matrix onto S, and let Π be the transition matrix where each row is given by π . These notations are formally defined in Section 2 and 6.

A number of natural and interesting information-theoretic optimization problems arise in the context of subset selection for multivariate Markov chains. For example, among all subsets S with $|S| \leq m$, which projected transition matrix $P^{(S)}$ maximizes the entropy rate, thereby identifying the most "random" coordinates? Similarly, which choice of S minimizes the Kullback-Leibler (KL) divergence between $P^{(S)}$ and $\Pi^{(S)}$, making it closest to stationarity and highlighting subsets nearest to equilibrium? These insights have direct applications in Markov chain Monte Carlo (MCMC), where selecting and analyzing optimal subsets may lead to the design of accelerated MCMC samplers.

Model reduction problems for Markov chains have been studied previously using spectral theory [3]. While such approach is powerful, combinatorial approaches to subset selection in the context of Markov chains remain underexplored, a gap we address in this paper by leveraging submodular optimization. Our work builds on recent efforts, such as [2], and adapts a combinatorial lens to develop efficient greedy-based algorithms for finding approximate optimizers. In another related work [6], the authors study the lumping problem of Markov chains from an information-theoretic and combinatorial perspective. Our main contributions are as follows:

• Identification of new submodular functions in Markov chain theory. We prove that, under suitable assumptions, the distance to stationarity and the distance to independence of the

^{*}Email: zheyuan_lai@u.nus.edu

[†]Email: mchchoi@nus.edu.sg, corresponding author

complement set are, respectively, supermodular and submodular. This extends the line of work initiated in [2].

- Adaptation of greedy-based optimization algorithms for submodular functions in Markov chains. We apply existing greedy-based algorithms to optimize the aforementioned information-theoretic functionals. To the best of our knowledge, this is the first application of greedy optimization algorithms in the submodular optimization of Markov chains.
- Development of a generalized distorted greedy algorithm for k-submodular function maximization subjected to cardinality constraints. Building on the distorted greedy algorithm from [7], we propose a generalized version with theoretical guarantees, which is of independent interest.
- Numerical validation on structured multivariate Markov chains. We conduct extensive experiments on Markov chains derived from the Bernoulli-Laplace model and the Curie-Weiss model to evaluate the performance of the proposed greedy algorithms.

The remainder of this paper is organized as follows. Section 2.1 provides a review of submodularity and k-submodularity. Section 2.2 introduces the distorted greedy algorithm and presents our generalized version with theoretical guarantees. Section 2.3 discusses submodular functions that naturally arise in the information-theoretic study of multivariate Markov chains. We then explore optimization problems related to entropy rate (Section 3), distance to factorizability (Section 4), distance to independence (Section 5), distance to stationarity (Section 6), and distance to factorizability over a fixed set (Section 7). Finally, we illustrate the algorithms through numerical experiments in Section 8.

2 Preliminaries

2.1 Definition and properties of submodular functions

We first recall the definition of a submodular function [15]. Given a finite nonempty ground set U, a set function $f: 2^U \to \mathbb{R}$ defined on subsets of U is called **submodular** if for all $S, T \subseteq U$,

$$f(S) + f(T) \ge f(S \cap T) + f(S \cup T).$$

f is said to be **supermodular** if -f is submodular, and f is said to be **modular** if f is both submodular and supermodular.

Next, we recall a result that states the complement of a submodular function is still submodular:

Lemma 2.1. If $S \mapsto f(S)$ is submodular, then $S \mapsto f(U \setminus S)$ is submodular.

Proof. We choose $S \subseteq T \subseteq U$ and $e \in U \setminus T$, then

$$\begin{pmatrix} f(U \setminus (S \cup \{e\})) - f(U \setminus S) \end{pmatrix} - \begin{pmatrix} f(U \setminus (T \cup \{e\})) - f(U \setminus T) \end{pmatrix} \\ = \begin{pmatrix} f(U \setminus T) - f(U \setminus (T \cup \{e\})) \end{pmatrix} - \begin{pmatrix} f(U \setminus S) - f(U \setminus (S \cup \{e\})) \end{pmatrix} \ge 0$$

since $S \mapsto f(S)$ is submodular and $U \setminus T \subseteq U \setminus S$, and hence $S \mapsto f(U \setminus S)$ is submodular.

We call a submodular function $f: 2^U \to \mathbb{R}$ symmetric if $f(A) = f(U \setminus A)$ for all $A \subseteq U$.

A multivariate generalization of submodularity is known as k-submodularity [4] where $k \in \mathbb{N}$. In particular, 1-submodular function is equivalent to submodular function. Let $f : (k+1)^U \to \mathbb{R}$ be a set function. The function f is said to be k-submodular if

$$f(\mathbf{S}) + f(\mathbf{T}) \ge f(\mathbf{S} \sqcap \mathbf{T}) + f(\mathbf{S} \sqcup \mathbf{T}) \quad \forall \mathbf{S}, \mathbf{T} \in (k+1)^U,$$

where $\mathbf{S} \sqcap \mathbf{T}$ is the k-tuple whose *i*-th set is $S_i \cap T_i$ and $\mathbf{S} \sqcup \mathbf{T}$ is the k-tuple whose *i*-th set is $(S_i \cup T_i) \setminus (\bigcup_{i \neq i} (S_j \cup T_j))$. A function f is said to be k-supermodular if -f is k-submodular.

For $\mathbf{S} = (S_1, \ldots, S_k), \mathbf{T} = (T_1, \ldots, T_k) \in (k+1)^U$, we write $\mathbf{S} \preceq \mathbf{T}$ if and only if $S_i \subseteq T_i \ \forall i \in \llbracket k \rrbracket$, where $\llbracket k \rrbracket := \{1, 2, \ldots, k\}$. A function f is said to be **monotonically non-decreasing** (resp. **non-increasing**) if

$$f(\mathbf{S}) \leq (\text{resp.} \geq) f(\mathbf{T}) \quad \forall \mathbf{S} \preceq \mathbf{T}.$$

Let $\Delta_{e,i} f(\mathbf{S})$ be the marginal gain of adding e to the *i*-th set of \mathbf{S} :

$$\Delta_{e,i}f(\mathbf{S}) := f(S_1, \dots, S_i \cup \{e\}, \dots, S_k) - f(S_1, \dots, S_i, \dots, S_k)$$

Note that f being monotonically non-decreasing is equivalent to $\Delta_{e,i}f(\mathbf{S}) \geq 0$ for all $\mathbf{S} \in (k+1)^U$, $i \in [k]$, and $e \notin \operatorname{supp}(\mathbf{S})$, where we define $\operatorname{supp}(\mathbf{S}) := \bigcup_{i=1}^k S_i$. A function f is said to be **pairwise** monotonically non-decreasing (resp. non-increasing) if

$$\Delta_{e,i} f(\mathbf{S}) + \Delta_{e,j} f(\mathbf{S}) \ge (\text{resp.} \le) 0$$

for all $\mathbf{S} \in (k+1)^U$, $e \notin \operatorname{supp}(\mathbf{S})$, and $i, j \in [k]$ such that $i \neq j$. A function f is said to be **orthant** submodular (resp. orthant supermodular) if

$$\Delta_{e,i} f(\mathbf{S}) \ge (\text{resp.} \le) \Delta_{e,i} f(\mathbf{T}) \tag{1}$$

for all $i \in \llbracket k \rrbracket$ and $\mathbf{S}, \mathbf{T} \in (k+1)^U$ such that $\mathbf{S} \preceq \mathbf{T}, e \notin \operatorname{supp}(\mathbf{T})$.

The following result that we recall characterizes k-submodularity:

Theorem 2.2 (Characterization of k-submodularity (Theorem 7 of [15])). A function f is k-submodular (resp. k-supermodular) if and only if f is both orthant submodular (resp. supermodular) and pairwise monotonically non-decreasing (resp. non-increasing).

The next two results relates the sum of individually supermodular or submodular functions to k-supermodularity or k-submodularity respectively.

Lemma 2.3. Let $F: (k+1)^U \to \mathbb{R}$ defined to be

$$F(\mathbf{S}) = F(S_1, \dots, S_k) := \sum_{i=1}^k F_i(S_i)$$

be the sum of k monotonically non-increasing and supermodular functions $(F_i)_{i=1}^k$ with $F_i: 2^U \to \mathbb{R}$ for all $i \in [\![k]\!]$. Then F is k-supermodular.

Proof. Throughout this proof, let $i \neq j \in [\![k]\!]$. First, we seek to prove that F is pairwise monotonically non-increasing, in which case we aim to show $\Delta_{e,i}F(\mathbf{S}) + \Delta_{e,j}F(\mathbf{S}) \leq 0$ for $e \notin \operatorname{supp}(\mathbf{S})$:

$$\Delta_{e,i}F(\mathbf{S}) + \Delta_{e,j}F(\mathbf{S}) = (F_i(S_i \cup \{e\}) - F_i(S_i)) + (F_j(S_j \cup \{e\}) - F_j(S_i)) \le 0,$$

given that F_i, F_j are both monotonically non-increasing. Next, we seek to show that F is orthant supermodular, in which case we aim to show that $\Delta_{e,i}F(\mathbf{S}) \leq \Delta_{e,i}F(\mathbf{T})$ for any $\mathbf{S} \leq \mathbf{T}$ and $e \notin \operatorname{supp}(\mathbf{T})$:

$$\Delta_{e,i}F(\mathbf{S}) - \Delta_{e,i}F(\mathbf{T}) = (F_i(S_i \cup \{e\}) - F_i(S_i)) - (F_i(T_i \cup \{e\}) - F_i(T_i)) \le 0,$$

given that F_i is supermodular. Therefore, F is k-supermodular given that it is pairwise monotonically non-increasing and orthant supermodular using Theorem 2.2.

Corollary 2.4. Let $G: (k+1)^U \to \mathbb{R}$ defined to be

$$G(\mathbf{S}) = G(S_1, \dots, S_k) := \sum_{i=1}^k G_i(S_i)$$

be the sum of k monotonically non-decreasing and submodular functions $(G_i)_{i=1}^k$ with $G_i : 2^U \to \mathbb{R}$ for all $i \in [k]$. Then G is k-submodular.

Proof. By applying Lemma 2.3 to -G, we see that -G is k-supermodular, which is equivalent to G being k-submodular.

The next result, that we shall apply in subsequent sections, transforms a non-monotone submodular f to a monotonically non-decreasing submodular g:

Theorem 2.5 (Transform a non-monotone submodular f to a monotone submodular g (Proposition 14.18 of [10])). Let $f: 2^U \to \mathbb{R}$ be a submodular function and $\beta \in \mathbb{R}$, then $g: 2^U \to \mathbb{R}$ defined by

$$g(S) := f(S) - \beta + \sum_{e \in S} (f(U \setminus \{e\}) - f(U))$$

is submodular and monotonically non-decreasing.

We aim to prove a generalized version of Theorem 2.5, that transforms a given constrained orthant submodular function into a k-submodular function. Suppose that we are given $\mathbf{V} \in (k+1)^U$. Then, constrained to \mathbf{V} , we can transform an orthant submodular function into a k-submodular function.

Theorem 2.6. Let $f : (k+1)^U \to \mathbb{R}$ be an orthant submodular function, $\beta \in \mathbb{R}$ and $\mathbf{V} \in (k+1)^U$. then $g : (k+1)^U \preceq \mathbf{V} \to \mathbb{R}$ with

$$g(\mathbf{S}) := f(\mathbf{S}) - \beta + \sum_{i=1}^{k} \sum_{e \in S_i} \left(f(V_1, \dots, V_i \setminus \{e\}, \dots, V_k) - f(V_1, \dots, V_i, \dots, V_k) \right)$$

is k-submodular and monotonically non-decreasing.

Proof. Suppose that $\mathbf{S} \leq \mathbf{T}, i \in [\![k]\!]$, and $e \in V_i \setminus T_i$. Since f is orthant submodular, we have $\Delta_{e,i} f(\mathbf{S}) \geq \Delta_{e,i} f(\mathbf{T})$, and hence

$$\Delta_{e,i}g(\mathbf{S}) = \Delta_{e,i}f(\mathbf{S}) + f(V_1, \dots, V_i \setminus \{e\}, \dots, V_k) - f(V_1, \dots, V_i, \dots, V_k)$$

$$\geq \Delta_{e,i}f(\mathbf{T}) + \Delta_{e,i}\sum_{j=1}^k \sum_{u \in T_j} \left(f(V_1, \dots, V_j \setminus \{u\}, \dots, V_k) - f(V_1, \dots, V_j, \dots, V_k)\right)$$

$$= \Delta_{e,i}g(\mathbf{T}).$$

This gives q is orthant submodular.

To prove the orthant monotonicity, we choose $\mathbf{S} \in (k+1)^U$, $i \in [[k]]$, and $e \in V_i \setminus S_i$. From the orthant submodularity of f, since $S_i \subseteq V_i \setminus \{e\}$, we have

$$\Delta_{e,i}g(\mathbf{S}) = \Delta_{e,i}f(\mathbf{S}) - (f(V_1, \dots, V_i, \dots, V_k) - f(V_1, \dots, V_i \setminus \{e\}, \dots, V_k)) \ge 0.$$

Therefore g is monotonically non-decreasing, which implies that g is pairwise monotonically non-decreasing, and hence g is k-submodular.

In the remaining of this subsection and also the coming subsection, we recall a few classical submodular optimization algorithms.

To maximize a monotonically non-decreasing submodular function, one can apply a heuristic greedy algorithm (see Section 4 of [13]) with $(1 - e^{-1})$ -approximation guarantee. For non-monotone submodular functions, we recall a local search algorithm (see Theorem 3.4 of [5]) in Algorithm 1 that comes along with an approximation guarantee.

Algorithm 1 Local Search Algorithm [5]

Require: Ground set U with |U| = d, submodular function f, positive $\epsilon > 0$ 1: Initialize $S \leftarrow \{e\}$, where $f(\{e\})$ is the maximum over all singletons $e \in U$ 2: while $\exists a \in U \setminus S$ such that $f(S \cup \{a\}) \ge (1 + \epsilon/d^2)f(S)$ do 3: $S \leftarrow S \cup \{a\}$ 4: end while 5: if $\exists a \in S$ such that $f(S \setminus \{a\}) \ge (1 + \epsilon/d^2)f(S)$ then 6: $S \leftarrow S \setminus \{a\}$ 7: Go back to line 2 8: end if 9: Output: f(S) and $f(U \setminus S)$

Theorem 2.7 (Approximation guarantee of Algorithm 1). Algorithm 1 is a $(\frac{1}{3} - \frac{\epsilon}{d})$ -approximation algorithm for maximizing non-negative submodular functions, and $(\frac{1}{2} - \frac{\epsilon}{d})$ -approximation algorithm for maximizing non-negative submodular functions. The time complexity of Algorithm 1 is $\mathcal{O}(\frac{1}{\epsilon}d^3 \log d)$

2.2 Distorted greedy algorithms to maximize the difference of a submodular function and a modular function

In this paper, it turns out that some functions we are interested in optimizing can be written as a difference of a submodular function and a modular function. In this section, we shall consider maximizing

the difference of a monotonically non-decreasing submodular g and a modular c on the ground set U with cardinality constraint being at most $m \in \mathbb{N}$. Precisely, we consider the problem

$$\max_{S \subseteq U; \ |S| \le m} g(S) - c(S),$$

and

$$OPT = OPT(g, c, U, m) := \underset{S \subseteq U; \ |S| \le m}{\arg \max} g(S) - c(S)$$

In this setting, a distorted greedy algorithm (Algorithm 2) has been proposed along with a theoretical lower bound [7].

Algorithm 2 Distorted greedy algorithm for maximizing the difference between a monotonically non-decreasing submodular function and a modular function

Require: monotonically non-decreasing submodular g with $g(\emptyset) \ge 0$, non-negative modular c, cardinality m, ground set U

1: Initialize $S_0 \leftarrow \emptyset$ 2: for i = 0 to m - 1 do 3: $e_i \leftarrow \underset{e \in U}{\arg \max} \left\{ \left(1 - \frac{1}{m}\right)^{m - (i+1)} \left(g(S_i \cup \{e\}) - g(S_i)\right) - c(\{e\})\right\}$ 4: if $\left(1 - \frac{1}{m}\right)^{m - (i+1)} \left(g(S_i \cup \{e_i\}) - g(S_i)\right) - c(\{e_i\}) > 0$ then 5: $S_{i+1} \leftarrow S_i \cup \{e_i\}$ 6: else 7: $S_{i+1} \leftarrow S_i$ 8: end if 9: end for 10: Output: S_m .

Theorem 2.8 (Lower bound for distorted greedy algorithm). Algorithm 2 provides the following lower bound:

$$g(S_m) - c(S_m) \ge (1 - e^{-1})g(\text{OPT}) - c(\text{OPT}),$$

where S_m is the final output set.

Let $\mathbf{V} \in (k+1)^U$, and consider maximizing the difference of a monotonically non-decreasing ksubmodular g and a modular c on the ground set U with cardinality constraint being at most $m \in \mathbb{N}$. Precisely, we consider the problem

$$\max_{\mathbf{S} \preceq \mathbf{V}; \ |\operatorname{supp}(\mathbf{S})| \le m} g(\mathbf{S}) - c(\mathbf{S}), \tag{2}$$

and

$$\mathbf{OPT} = \mathbf{OPT}(g, c, U, \mathbf{V}, m) := \underset{\mathbf{S} \preceq \mathbf{V}; \ |\mathrm{supp}(\mathbf{S})| \le m}{\arg \max} g(\mathbf{S}) - c(\mathbf{S}).$$

We propose a generalized distorted greedy algorithm (Algorithm 3) for solving (2), which is of independent interest.

The rest of this section is devoted to giving a lower bound for the generalized distorted greedy algorithm. We assume that g is monotonically non-decreasing, k-submodular, $g(\emptyset) \ge 0$, while c is non-negative, modular and $c(\emptyset) = 0$.

In order to prove the lower bound for the generalized distorted greedy algorithm, we first define the distorted objective function $\Phi_i : (k+1)^U \to \mathbb{R}$, for $m \in \mathbb{N}$ and $0 \le i \le m-1$, that

$$\Phi_i(\mathbf{S}) := (1 - m^{-1})^{m-i} g(\mathbf{S}) - c(\mathbf{S})$$

We also denote $\Psi_i : (k+1)^U \times \llbracket k \rrbracket \times U \to \mathbb{R}$ that

$$\Psi_i(\mathbf{S}, j, e) := \max\{0, (1 - m^{-1})^{m - (i+1)} \Delta_{e,j} g(\mathbf{S}) - c(\{e\})\}.$$

Algorithm 3 Generalized distorted greedy algorithm for maximizing the difference of k-submodular function and a modular function

Require: k-submodular monotonically non-decreasing g with $g(\emptyset) \ge 0$, non-negative modular c with $c(\emptyset) = 0$, cardinality m, ground set U, $\mathbf{V} = (V_1, \ldots, V_k) \in (k+1)^U$.

1: Initialize
$$\mathbf{S}_{0} = (S_{0,1}, \dots, S_{0,k}) \leftarrow \emptyset$$

2: for $i = 0$ to $m - 1$ do
3: $(j^{*}, e^{*}) \leftarrow \underset{j \in [k], e \in V_{j} \setminus S_{i,j}}{\arg \max} \left\{ \left(1 - \frac{1}{m}\right)^{m - (i+1)} \Delta_{e,j} g(\mathbf{S}_{i}) - c(\{e\}) \right\}$
4: if $\left(1 - \frac{1}{m}\right)^{m - (i+1)} \Delta_{e^{*}, j^{*}} g(\mathbf{S}_{i}) - c(\{e^{*}\}) > 0$ then
5: $S_{i+1, j^{*}} \leftarrow S_{i, j^{*}} \cup \{e^{*}\}$
6: else
7: $S_{i+1, j^{*}} \leftarrow S_{i, j^{*}}$
8: end if
9: for $l \neq j^{*}$ do
10: $S_{i+1, l} \leftarrow S_{i, l}$
11: end for
12: end for
13: Output: $\mathbf{S}_{m} = (S_{m, 1}, \dots, S_{m, k}).$

Lemma 2.9. The difference of the distorted objective function of two iterations can be written as

$$\Phi_{i+1}(\mathbf{S}_{i+1}) - \Phi_i(\mathbf{S}_i) = \Psi_i(\mathbf{S}_i, j^*, e^*) + \frac{1}{m} \left(1 - \frac{1}{m}\right)^{m - (i+1)} g(\mathbf{S}_i)$$

Proof. Similar to Lemma 1 of [7], we can show

$$\begin{split} \Phi_{i+1}(\mathbf{S}_{i+1}) - \Phi_i(\mathbf{S}_i) &= \left(1 - \frac{1}{m}\right)^{m-(i+1)} g(\mathbf{S}_{i+1}) - c(\mathbf{S}_{i+1}) - \left(1 - \frac{1}{m}\right)^{m-i} g(\mathbf{S}_i) + c(\mathbf{S}_i) \\ &= \left(1 - \frac{1}{m}\right)^{m-(i+1)} g(\mathbf{S}_{i+1}) - c(\mathbf{S}_{i+1}) - \left(1 - \frac{1}{m}\right)^{m-(i+1)} \left(1 - \frac{1}{m}\right) g(\mathbf{S}_i) + c(\mathbf{S}_i) \\ &= \left(1 - \frac{1}{m}\right)^{m-(i+1)} \left(g(\mathbf{S}_{i+1}) - g(\mathbf{S}_i)\right) - \left(c(\mathbf{S}_{i+1}) - c(\mathbf{S}_i)\right) \\ &+ \frac{1}{m} \left(1 - \frac{1}{m}\right)^{m-(i+1)} g(\mathbf{S}_i). \end{split}$$

If $(1 - m^{-1})^{m - (i+1)} \Delta_{e^*, j^*} g(\mathbf{S}) - c(\{e^*\}) > 0$, then e^* is added to the solution set. In the algorithm we have $e^* \in V_{j^*} \setminus S_{i, j^*}, g(\mathbf{S}_{i+1}) - g(\mathbf{S}_i) = \Delta_{e^*, j^*} g(\mathbf{S}_i), c(\mathbf{S}_{i+1}) - c(\mathbf{S}_i) = c(\{e^*\})$, hence

$$\Phi_{i+1}(\mathbf{S}_{i+1}) - \Phi_i(\mathbf{S}_i) = \Psi_i(\mathbf{S}_i, j^*, e^*) + \frac{1}{m} \left(1 - \frac{1}{m}\right)^{m - (i+1)} g(\mathbf{S}_i)$$

If $(1 - m^{-1})^{m - (i+1)} \Delta_{e^*, j^*} g(\mathbf{S}) - c(\{e_i\}) \leq 0$, the algorithm does not add e^* into the solution set, hence $\mathbf{S}_{i+1} = \mathbf{S}_i$. In this case, we also have

$$\Phi_{i+1}(\mathbf{S}_{i+1}) - \Phi_i(\mathbf{S}_i) = 0 + \frac{1}{m} \left(1 - \frac{1}{m}\right)^{m-(i+1)} g(\mathbf{S}_i) = \Psi_i(\mathbf{S}_i, j^*, e^*) + \frac{1}{m} \left(1 - \frac{1}{m}\right)^{m-(i+1)} g(\mathbf{S}_i).$$

Summarizing these two cases, we see that

$$\Phi_{i+1}(\mathbf{S}_{i+1}) - \Phi_i(\mathbf{S}_i) = \Psi_i(\mathbf{S}_i, j^*, e^*) + \frac{1}{m} \left(1 - \frac{1}{m}\right)^{m - (i+1)} g(\mathbf{S}_i)$$

Lemma 2.10. A lower bound for Ψ_i is

$$\Psi_i(\mathbf{S}_i, j^*, e^*) \ge \frac{1}{m} \left(\left(1 - \frac{1}{m} \right)^{m - (i+1)} \left(g(\mathbf{OPT}) - g(\mathbf{S}_i) \right) - c(\mathbf{OPT}) \right)$$

Proof. For $j \in [\![k]\!]$, let

$$U_{i,j} := (V_j \setminus S_{i,j}) \cap \operatorname{OPT}_j,$$
$$U_i := \bigcup_{j=1}^k U_{i,j},$$
$$\mathbf{U}_i := (U_{i,1}, U_{i,2}, \dots, U_{i,k}),$$

and hence

$$S_{i,j} \cup U_{i,j} = S_{i,j} \cup \operatorname{OPT}_j.$$
(3)

We then have

$$\begin{split} m\Psi_{i}(\mathbf{S}_{i}, j^{*}, e^{*}) &= m \max_{j \in [\![k]\!], e \in V_{j} \setminus S_{i,j}} \left\{ 0, \left(1 - \frac{1}{m}\right)^{m - (i+1)} \Delta_{e,j} g(\mathbf{S}_{i}) - c(\{e\}) \right\} \\ &\geq |\operatorname{supp}(\mathbf{OPT})| \max_{j \in [\![k]\!], e \in U_{i,j}} \left\{ 0, \left(1 - \frac{1}{m}\right)^{m - (i+1)} \Delta_{e,j} g(\mathbf{S}_{i}) - c(\{e\}) \right\} \\ &\geq |U_{i}| \max_{j \in [\![k]\!], e \in U_{i,j}} \left\{ \left(1 - \frac{1}{m}\right)^{m - (i+1)} \Delta_{e,j} g(\mathbf{S}_{i}) - c(\{e\}) \right\} \\ &\geq \sum_{j=1}^{k} \sum_{e \in U_{i,j}} \left(\left(1 - \frac{1}{m}\right)^{m - (i+1)} \Delta_{e,j} g(\mathbf{S}_{i}) - c(\{e\}) \right) \\ &= \left(1 - \frac{1}{m}\right)^{m - (i+1)} \sum_{j=1}^{k} \sum_{e \in U_{i,j}} \Delta_{e,j} g(\mathbf{S}_{i}) - c(\mathbf{U}_{i}) \\ &\geq \left(1 - \frac{1}{m}\right)^{m - (i+1)} \sum_{j=1}^{k} \sum_{e \in U_{i,j}} \Delta_{e,j} g(\mathbf{S}_{i}) - c(\mathbf{OPT}), \end{split}$$

where the last inequality follows from the fact that c is non-negative. Then, the desired result follows if we show that

$$\sum_{j=1}^{k} \sum_{e \in U_{i,j}} \Delta_{e,j} g(\mathbf{S}_i) \ge g(\mathbf{OPT}) - g(\mathbf{S}_i).$$

Since g is orthant submodular, by Lemma 1.1 of [11], we have

$$\sum_{e \in U_{i,j}} \Delta_{e,j} g(\mathbf{S}_i) \ge g(S_{i,1}, \dots, S_{i,j-1}, S_{i,j} \cup U_{i,j}, S_{i,j+1}, \dots, S_k) - g(\mathbf{S}_i),$$

and hence it further suffices to prove

$$\sum_{j=1}^{k} g(S_{i,1}, \dots, S_{i,j-1}, S_{i,j} \cup U_{i,j}, S_{i,j+1}, \dots, S_k) \ge g(\mathbf{OPT}) + (k-1)g(\mathbf{S}_i).$$
(4)

Since g is k-submodular, then

$$g(\mathbf{X}) + g(\mathbf{Y}) \ge g(\mathbf{X} \sqcup \mathbf{Y}) + g(\mathbf{X} \sqcap \mathbf{Y}),$$

for any $\mathbf{X}, \mathbf{Y} \in (k+1)^U$. We seek to apply this definition to update each of the k coordinates by adding $(U_{i,j})_{j=1}^k$ sequentially. For the first step, we have

$$g(S_{i,1} \cup U_{i,1}, S_{i,2}, \dots, S_{i,k}) + g(S_{i,1}, S_{i,2} \cup U_{i,2}, S_{i,3}, \dots, S_{i,k})$$

$$\geq g((S_{i,1} \cup U_{i,1}) \setminus (\cup_{l \neq 1}^k S_{i,l} \cup U_{i,2}), (S_{i,2} \cup U_{i,2}) \setminus (\cup_{l \neq 2}^k S_{i,l} \cup U_{i,1}), S_{i,3}, \dots, S_{i,k}) + g(\mathbf{S}_i)$$

$$= g(S_{i,1} \cup U_{i,1}, S_{i,2} \cup U_{i,2}, S_{i,3}, \dots, S_{i,k}) + g(\mathbf{S}_i),$$

where the last equality uses the fact that with $n \in [\![k]\!]$,

$$(S_{i,n} \cup U_{i,n}) = (S_{i,n} \cup U_{i,n}) \setminus (\bigcup_{l \neq n}^k (S_{i,l} \cup U_{i,l})).$$

In the *n*-th step with $n \in [k]$, we thus have

$$g(S_{i,1} \cup U_{i,1}, \dots, S_{i,n} \cup U_{i,n}, \dots, S_{i,k}) + g(S_{i,1}, \dots, S_{i,n}, S_{i,n+1} \cup U_{i,n+1}, \dots, S_{i,k})$$

$$\geq g(S_{i,1} \cup U_{i,1}, \dots, S_{i,n+1} \cup U_{i,n+1}, \dots, S_{i,k}) + g(\mathbf{S}_i).$$

Repeating the above analysis leads to

$$\sum_{j=1}^{k} g(S_{i,1}, \dots, S_{i,j-1}, S_{i,j} \cup U_{i,j}, S_{i,j+1}, \dots, S_k) \ge g(\mathbf{S}_i \sqcup \mathbf{U}_i) + (k-1)g(\mathbf{S}_i).$$

Finally, using the assumption that g is monotonically non-decreasing and **OPT** \leq **S**_i \sqcup **U**_i in view of (3), we have

$$\sum_{j=1}^{k} g(S_{i,1}, \dots, S_{i,j-1}, S_{i,j} \cup U_{i,j}, S_{i,j+1}, \dots, S_k) \ge g(\mathbf{OPT}) + (k-1)g(\mathbf{S}_i),$$

and hence (4) holds.

Finally, we prove a lower bound for the generalized distorted greedy algorithm:

Theorem 2.11 (Lower bound for generalized distorted greedy algorithm). Algorithm 3 provides the following lower bound:

$$g(\mathbf{S}_m) - c(\mathbf{S}_m) \ge (1 - e^{-1})g(\mathbf{OPT}) - c(\mathbf{OPT}),$$

where $\mathbf{S}_m = (S_{m,1}, \dots, S_{m,k})$ is the final output set.

Proof. According to our assumptions, we have

$$\Phi_0(\mathbf{S}_0) = \left(1 - \frac{1}{m}\right)^m g(\emptyset) - c(\emptyset) \ge 0$$

and

$$\Phi_m(\mathbf{S}_m) = \left(1 - \frac{1}{m}\right)^0 g(\mathbf{S}_m) - c(\mathbf{S}_m) = g(\mathbf{S}_m) - c(\mathbf{S}_m).$$

Therefore, we have

$$g(\mathbf{S}_m) - c(\mathbf{S}_m) \ge \Phi_m(\mathbf{S}_m) - \Phi_0(\mathbf{S}_0) = \sum_{i=0}^{m-1} \Phi_{i+1}(\mathbf{S}_{i+1}) - \Phi_i(\mathbf{S}_i).$$
(5)

We apply Lemma 2.9 and 2.10 to yield

$$\Phi_{i+1}(\mathbf{S}_{i+1}) - \Phi_i(\mathbf{S}_i) = \Psi_i(\mathbf{S}_i, j^*, e^*) + \frac{1}{m} \left(1 - \frac{1}{m}\right)^{m - (i+1)} g(\mathbf{S}_i)$$
$$\geq \frac{1}{m} \left(1 - \frac{1}{m}\right)^{m - (i+1)} g(\mathbf{OPT}) - \frac{1}{m} c(\mathbf{OPT})$$

We plug the above bound into (5) to obtain

$$g(\mathbf{S}_m) - c(\operatorname{supp}(\mathbf{S}_m)) \ge \sum_{i=0}^{m-1} \left[\frac{1}{m} \left(1 - \frac{1}{m} \right)^{m-(i+1)} g(\mathbf{OPT}) - \frac{1}{m} c(\mathbf{OPT}) \right]$$
$$= \left[\frac{1}{m} \sum_{i=0}^{m-1} \left(1 - \frac{1}{m} \right)^i \right] g(\mathbf{OPT}) - c(\mathbf{OPT})$$
$$= \left(1 - \left(1 - \frac{1}{m} \right)^m \right) g(\mathbf{OPT}) - c(\mathbf{OPT})$$
$$\ge (1 - e^{-1}) g(\mathbf{OPT}) - c(\mathbf{OPT}).$$

2.3 Some submodular functions in information theory of multivariate Markov chains

Throughout this paper, we consider a finite d-dimensional state space described by $\mathcal{X} = \mathcal{X}^{(1)} \times \ldots \times \mathcal{X}^{(d)}$. For $S \subseteq \llbracket d \rrbracket$, we write $\mathcal{X}^{(S)} = \times_{i \in S} \mathcal{X}^{(i)}$. We denote by $\mathcal{L}(\mathcal{X})$ to be the set of transition matrices on \mathcal{X} , and $\mathcal{P}(\mathcal{X}) = \{\pi; \min_{x \in \mathcal{X}} \pi(x) > 0\}$ to be the set of probability masses with support on \mathcal{X} . Let $\pi \in \mathcal{P}(\mathcal{X})$ be any given probability distribution, and denote $\mathcal{L}(\pi) \subseteq \mathcal{L}(\mathcal{X})$ as a set of π -reversible transition matrices on \mathcal{X} , where a transition matrix $P \in \mathcal{L}(\mathcal{X})$ is said to be π -reversible if the detailed balance condition holds such that $\pi(x)P(x,y) = \pi(y)P(y,x)$ for all $x,y \in \mathcal{X}$. Additionally, we say that $P \in \mathcal{L}(\mathcal{X})$ is π -stationary if it satisfies $\pi = \pi P$.

We now recall the definition of the tensor product of transition matrices and probability masses, see e.g. Exercise 12.6 [12]. Define, for $M_l \in \mathcal{L}(\mathcal{X}^{(l)}), \pi_l \in \mathcal{P}(\mathcal{X}^{(l)}), x^l, y^l \in \mathcal{X}^{(l)}$ for $l \in \{i, j\}, i \neq j \in [\![d]\!]$,

$$(M_i \otimes M_j)((x^i, x^j), (y^i, y^j)) := M_i(x^i, y^i)M_j(x^j, y^j),$$

 $(\pi_i \otimes \pi_j)(x^i, x^j) := \pi_i(x^i)\pi_j(x^j).$

A transition matrix $P \in \mathcal{L}(\mathcal{X})$ is said to be in a product form if there exists $M_i \in \mathcal{L}(\mathcal{X}^{(i)})$ for $i \in \llbracket d \rrbracket$ such that $P = \bigotimes_{i=1}^{d} M_i$ can be expressed as a *d*-fold tensor product. A probability mass π is said to be in a product form if there exists $\pi_i \in \mathcal{P}(\mathcal{X}^{(i)})$ such that $\pi = \bigotimes_{i=1}^{d} \pi_i$.

We then recall the definition of leave-S-out and keep-S-in transition matrices of a given transition matrix P, see Section 2.2 of [2]. Let $\pi \in P(\mathcal{X})$, $P \in \mathcal{L}(\mathcal{X})$, and $S \subseteq \llbracket d \rrbracket$. For any $(x^{(-S)}, y^{(-S)}) \in \mathcal{X}^{(-S)} \times \mathcal{X}^{(-S)}$, we define the **leave-**S-out transition matrix to be $P_{\pi}^{(-S)}$ with entries given by

$$P_{\pi}^{(-S)}(x^{(-S)}, y^{(-S)}) := \frac{\sum_{(x^{(S)}, y^{(S)}) \in \mathcal{X}^{(S)} \times \mathcal{X}^{(S)}} \pi(x^{1}, \dots, x^{d}) P((x^{1}, \dots, x^{d}), (y^{1}, \dots, y^{d}))}{\sum_{x^{(S)} \in \mathcal{X}^{(S)}} \pi(x^{1}, \dots, x^{d})}$$

The **keep**-S-in transition matrix of P with respect to π is

$$P_{\pi}^{(S)} := P_{\pi}^{(-\llbracket d \rrbracket \setminus S)} \in \mathcal{L}(\mathcal{X}^{(S)})$$

In the special case of $S = \{i\}$ for $i \in [d]$, we write

$$P_{\pi}^{(-i)} = P_{\pi}^{(-\{i\})}, \quad P_{\pi}^{(i)} = P_{\pi}^{(\{i\})}.$$

When P is π -stationary, we omit the subscript π and write directly $P^{(-S)}, P^{(S)}$. We also apply the convention of $P^{(\emptyset)} = P^{(-\llbracket d \rrbracket)} = 1$.

We proceed to recall the Shannon entropy of a probability distribution and the entropy rate of the transition matrix, see Section 1 of [14]. For $\pi \in \mathcal{P}(\mathcal{X})$, its **Shannon entropy** is defined as

$$H(\pi) := -\sum_{x \in \mathcal{X}} \pi(x) \ln \pi(x)$$

where the standard convention of $0 \ln 0 := 0$ applies. For π -stationary $P \in \mathcal{L}(\mathcal{X})$, the **entropy rate** of P is defined as

$$H(P) := -\sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{X}} \pi(x) P(x, y) \ln P(x, y),$$

where the standard convention of $0 \ln 0 := 0$ applies.

We shall also recall the definition of **KL divergence** between Markov chains (Definition 2.1 of [2]) and the distance to independence (Definition 2.2 of [2]). For given $\pi \in \mathcal{P}(\mathcal{X})$ and transition matrices $M, L \in \mathcal{L}(\mathcal{X})$, we define the KL divergence from L to M with respect to π as

$$D_{KL}^{\pi}(M||L) := \sum_{x \in \mathcal{X}} \pi(x) \sum_{y \in \mathcal{X}} M(x,y) \ln\left(\frac{M(x,y)}{L(x,y)}\right),$$

where the convention of $0 \ln \frac{0}{a} := 0$ applies for $a \in [0, 1]$. Note that π need not be the stationary distribution of L or M. In particular, when M, L are assumed to be π -stationary, we write

$$D(M||L) := D_{KL}^{\pi}(M||L),$$

which can be interpreted as the KL divergence rate from L to M. Given $P \in \mathcal{L}(\mathcal{X})$, we define the **distance to independence** of P with respect to D_{KL}^{π} to be

$$\mathbb{I}^{\pi}(P) := \min_{L_i \in \mathcal{L}(\mathcal{X}^{(i)}), \ \forall i \in \llbracket d \rrbracket} D^{\pi}_{KL}(P \| \otimes_{i=1}^d L_i) = D^{\pi}_{KL}(P \| \otimes_{i=1}^d P^{(i)}_{\pi}).$$

We write

$$\mathbb{I}(P) = \mathbb{I}^{\pi}(P)$$

if P is π -stationary.

We recall the partition lemma for KL divergence of Markov chains (see Theorem 2.4 of [2]).

Theorem 2.12 (Partition lemma). Let $\pi \in \mathcal{P}(\mathcal{X})$, $P, L \in \mathcal{L}(\mathcal{X})$ and suppose $S \subseteq \llbracket d \rrbracket$, we have:

$$D_{KL}^{\pi}(P||L) \ge D_{KL}^{\pi^{(S)}}(P^{(S)}||L^{(S)})$$

We then recall some established results on submodularity of some functions arising in Markov chain theory as discussed in Proposition 2.6 of [2].

Theorem 2.13 (Submodularity of some functions in Markov chain theory). Let $S \subseteq \llbracket d \rrbracket$. Let $P \in \mathcal{L}(\mathcal{X})$ be a π -stationary transition matrix. We have

- (Submodularity of the entropy rate of P) The mapping $S \mapsto H(P^{(S)})$ is submodular.
- (Submodularity of the distance to $(S, \llbracket d \rrbracket \setminus S)$ -factorizability of P) The mapping $S \mapsto D(P \parallel P^{(S)} \otimes P^{(-S)})$ is submodular.
- (Supermodularity and monotonicity of the distance to independence) The mapping $S \mapsto \mathbb{I}(P^{(S)})$ is monotonically non-decreasing and supermodular.

Next, we investigate the map $S \mapsto \mathbb{I}(P^{(-S)})$, and show that it is monotonically non-increasing and supermodular.

Theorem 2.14 (Supermodularity and monotonicity of the distance to independence of $P^{(-S)}$). The mapping $S \mapsto \mathbb{I}(P^{(-S)})$ is monotonically non-increasing and supermodular.

Proof. We first prove the monotonicity. Suppose $S \subseteq T \subseteq \llbracket d \rrbracket \setminus T \subseteq \llbracket d \rrbracket \setminus S$, hence according to the partition lemma (Lemma 2.12), we have:

$$\mathbb{I}(P^{(-S)}) = D(P^{(-S)} \| \otimes_{i \in \llbracket d \rrbracket \setminus S} P^{(i)}) \ge D(P^{(-T)} \| \otimes_{i \in \llbracket d \rrbracket \setminus T} P^{(i)}) = \mathbb{I}(P^{(-T)}),$$

therefore, $S \mapsto \mathbb{I}(P^{(-S)})$ is monotonically non-increasing.

We then look into the supermodularity of this map. Since

$$\mathbb{I}(P^{(-S)}) = \sum_{i \in \llbracket d \rrbracket \backslash S} H(P^{(i)}) - H(P^{(-S)}),$$

then $\mathbb{I}(P^{(-S)})$ is supermodular because $H(P^{(-S)})$ is submodular in view of Lemma 2.1 and Lemma 2.3.

3 Submodular maximization of the entropy rate $H(P^{(S)})$

Given $P \in \mathcal{L}(\mathcal{X})$ and $m \in \mathbb{N}$, we aim to investigate the following submodular maximization problem with cardinality constraint:

$$\max_{S \subseteq \llbracket d \rrbracket; \ |S| \le m} H(P^{(S)}). \tag{6}$$

From Theorem 2.13, the map $S \mapsto H(P^{(S)})$ is submodular but generally not monotonically nondecreasing. Since the widely-used heuristic greedy algorithm is near-optimal only when the objective submodular function is monotonically non-decreasing (see Section 4 of [13]), in this regard our problem does not have a classical greedy-based approximation guarantee. On the other hand, since $H(P^{(S)}) \ge 0$ and $H(P^{(\emptyset)}) = 0$, if we consider the unconstrained maximization problem of (6), we can apply Algorithm 1 with $(\frac{1}{3} - \frac{\epsilon}{d})$ -approximation guarantee (see Theorem 2.7). Instead, we consider

$$H(P) = H(\pi \boxtimes P) - H(\pi),$$

where we define the edge measure of P with respect to π as $(\pi \boxtimes P)(x, y) := \pi(x)P(x, y)$ and $\pi \boxtimes P \in \mathcal{P}(\mathcal{X} \times \mathcal{X})$.

Then, the map

$$S \mapsto H(P^{(S)}) = H(\pi^{(S)} \boxtimes P^{(S)}) - H(\pi^{(S)})$$
(7)

can be considered as a monotonically non-decreasing submodular function $H(\pi^{(S)} \boxtimes P^{(S)})$ minus a nonnegative modular function $H(\pi^{(S)})$ if we assume π to be of product form. This fits into the setting of the distorted greedy as in Algorithm 2, and leads us to Corollary 3.1.

Corollary 3.1. Let $P \in \mathcal{L}(\mathcal{X})$ be π -stationary where π is of product form. In Algorithm 2, we take $g(S) = H(\pi^{(S)} \boxtimes P^{(S)}), c(S) = H(\pi^{(S)}), and \text{ OPT} = \arg \max_{S \subseteq \llbracket d \rrbracket; |S| \leq m} H(P^{(S)}).$ Therefore, Theorem 2.8 gives

$$H(P^{(S_m)}) \ge (1 - e^{-1})H(\pi^{(\text{OPT})} \boxtimes P^{(\text{OPT})}) - H(\pi^{(\text{OPT})})$$

where S_m is the output of Algorithm 2.

More generally for P with non-product-form π as stationary distribution, in view of Theorem 2.5, for any $\beta \in \mathbb{R}$ we have a monotonically non-decreasing submodular g given by

$$g(S) = H(P^{(S)}) - \beta + \sum_{e \in S} (H(P^{(-e)}) - H(P)),$$
(8)

and we also denote the following modular function

$$c(S) = -\beta + \sum_{e \in S} (H(P^{(-e)}) - H(P))$$

$$= -\beta + \sum_{e \in S} (D(P \| P^{(e)} \otimes P^{(-e)}) - H(P^{(e)})).$$
(9)

As $H(P^{(e)}) \leq \log |\mathcal{X}^{(e)}|, c$ is ensured to be non-negative if $\beta \leq -\sum_{i=1}^{d} \log |\mathcal{X}^{(i)}|$. Since

$$H(P^{(S)}) = g(S) - c(S),$$

we can employ Algorithm 2 to perform distorted greedy maximization with a lower bound.

Corollary 3.2. Let $P \in \mathcal{L}(\mathcal{X})$ be π -stationary. In Algorithm 2, we take g as in (8), c as in (9), $\beta \leq -\sum_{i=1}^{d} \log |\mathcal{X}^{(i)}|$, and $\text{OPT} = \arg \max_{S \subseteq \llbracket d \rrbracket; |S| \leq m} H(P^{(S)})$. Therefore, Theorem 2.8 gives

$$H(P^{(S_m)}) \ge (1 - e^{-1})g(\text{OPT}) - c(\text{OPT}),$$

where S_m is the output of Algorithm 2.

Note that the lower bound of Corollary (3.2) depends on β through g and c. If β is chosen to be too small, then the lower bound might be too loose as the right hand side might be negative.

3.1 k-submodular maximization of the entropy rate of the tensorized keep- S_i -in matrices $H(\bigotimes_{i=1}^k P^{(S_i)})$

In this subsection, we investigate the following map

$$(k+1)^{\llbracket d \rrbracket} \ni \mathbf{S} = (S_1, \dots, S_k) \mapsto f(\mathbf{S}) = H(\otimes_{i=1}^k P^{(S_i)}) = \sum_{i=1}^k H(P^{(S_i)}),$$
(10)

and consider maximization problems of the form, for given $\mathbf{V} \in (k+1)^{\llbracket d \rrbracket}$,

$$\max_{\mathbf{S} \preceq \mathbf{V}; |\text{supp}(\mathbf{S})| \le m} H(\otimes_{i=1}^{k} P^{(S_i)}).$$
(11)

In the special case of k = 1 and $\mathbf{V} = \llbracket d \rrbracket$, we recover the problem (6).

First, we consider the special case where P is π -stationary with π taking on a product form. Similar to the map (7), we re-write the map (10) as

$$\mathbf{S} \mapsto f(\mathbf{S}) = \sum_{i=1}^{k} H(\pi^{(S_i)} \boxtimes P^{(S_i)}) - \sum_{i=1}^{k} H(\pi^{(S_i)}).$$
(12)

Since $H(\pi^{(S_i)} \boxtimes P^{(S_i)})$ is monotonically non-decreasing and submodular, then by Corollary 2.4, the following function g is monotonically non-decreasing and k-submodular

$$g(\mathbf{S}) = \sum_{i=1}^{k} H(\pi^{(S_i)} \boxtimes P^{(S_i)}).$$
(13)

Since π is of product form, we denote the non-negative modular function c as

$$c(\mathbf{S}) = \sum_{i=1}^{k} H(\pi^{(S_i)}).$$
(14)

Therefore, we have

$$f(\mathbf{S}) = g(\mathbf{S}) - c(\mathbf{S}),$$

and the distorted greedy algorithm yields an approximate maximizer with a lower bound as in Theorem 2.11.

Corollary 3.3. Let $P \in \mathcal{L}(\mathcal{X})$ be π -stationary where π is of product form. In Algorithm 3, we take g as in (13) and c as in (14), and $\mathbf{OPT} = \arg \max_{\mathbf{S} \preceq \mathbf{V}; |\operatorname{supp}(\mathbf{S})| \leq m} f(\mathbf{S})$. Then by Theorem 2.11, we have the following lower bound

$$f(\mathbf{S}_m) = H(\bigotimes_{i=1}^k P^{(S_{m,i})}) \ge (1 - e^{-1})g(\mathbf{OPT}) - c(\mathbf{OPT}),$$

where $\mathbf{S}_m = (S_{m,1}, \ldots, S_{m,k})$ is the output of Algorithm 3.

In the special case where k = 1 and $\mathbf{V} = \llbracket d \rrbracket$, we recover Corollary 3.1.

Next, we investigate the case where P is π -stationary for general π which may not be of product form. We first prove an orthant submodularity result.

Lemma 3.4. The map (10) is orthant submodular.

Proof. We shall prove that $\Delta_{e,i}f(\mathbf{S}) \geq \Delta_{e,i}f(\mathbf{T})$, where we choose $\mathbf{S} \leq \mathbf{T}$ and $e \notin \operatorname{supp}(\mathbf{T})$. Given the submodularity of $S \mapsto H(P^{(S)})$, we have

$$H(P^{(S_i \cup \{e\})}) - H(P^{(S_i)}) \ge H(P^{(T_i \cup \{e\})}) - H(P^{(T_i)}),$$

which is equivalent to $\Delta_{e,i} f(\mathbf{S}) \geq \Delta_{e,i} f(\mathbf{T})$.

In view of Theorem 2.6, since the map (10) is orthant submodular, then for any $\beta \in \mathbb{R}$, if $\mathbf{S} \leq \mathbf{V}$, we have a monotonically non-decreasing k-submodular function g given by

$$g(\mathbf{S}) = \sum_{i=1}^{k} H(P^{(S_i)}) - \beta + \sum_{i=1}^{k} \sum_{e \in S_i} (H(P^{(V_i \setminus \{e\})}) - H(P^{(V_i)})),$$
(15)

and we also denote the following modular function

$$c(\mathbf{S}) = -\beta + \sum_{i=1}^{k} \sum_{e \in S_i} (H(P^{(V_i \setminus \{e\})}) - H(P^{(V_i)}))$$

$$= -\beta + \sum_{i=1}^{k} \sum_{e \in S_i} (D(P^{(V_i)} || P^{(e)} \otimes P^{(V_i \setminus \{e\})}) - H(P^{(e)})).$$
(16)

As $H(P^{(e)}) \leq \log |\mathcal{X}^{(e)}|, c$ is ensured to be non-negative if $\beta \leq -\sum_{i=1}^{k} \sum_{e \in V_i} \log |\mathcal{X}^{(e)}|$. Since

$$f(\mathbf{S}) = \sum_{i=1}^{k} H(P^{(S_i)}) = g(\mathbf{S}) - c(\mathbf{S}),$$

then we can apply Algorithm 3 to perform distorted greedy maximization with a guaranteed lower bound.

Corollary 3.5. Let $P \in \mathcal{L}(\mathcal{X})$ be π -stationary and $\mathbf{V} \in (k+1)^{\llbracket d \rrbracket}$. In Algorithm 3, we take g as in (15) and c as in (16), $\beta \leq -\sum_{i=1}^{k} \sum_{e \in V_i} \log |\mathcal{X}^{(e)}|$, and $\mathbf{OPT} = \arg \max_{\mathbf{S} \preceq \mathbf{V}; |\mathrm{supp}(\mathbf{S})| \leq m} f(\mathbf{S})$. Therefore, Theorem 2.11 gives

$$f(\mathbf{S}_m) = H(\bigotimes_{i=1}^k P^{(S_{m,i})}) \ge (1 - e^{-1})g(\mathbf{OPT}) - c(\mathbf{OPT}),$$

where $\mathbf{S}_m = (S_{m,1}, S_{m,2}, \dots, S_{m,k})$ is the output of Algorithm 3.

Note that the lower bound of Corollary 3.5 depends on β through g and c. If β is chosen to be too small, then the lower bound might be too loose as the right hand side might be negative.

4 Submodular optimization of distance to factorizability $D(P || P^{(S)} \otimes P^{(-S)})$

4.1 Submodular minimization of the distance to factorizability

For

$$2^{\llbracket d \rrbracket} \ni S \mapsto D(P \| P^{(S)} \otimes P^{(-S)})$$

we first recall that this map is submodular (see Lemma 2.13). Since $D(P||P^{(S)} \otimes P^{(-S)}) = D(P||P^{(-S)} \otimes P^{(S)})$, then this map is also symmetric. In this case, there exists an algorithm for minimizing non-negative symmetric submodular functions (see Theorem 14.25 of [10]) that gives

$$S^* \in \underset{\emptyset \neq S \subset \llbracket d \rrbracket; \ |S| \le m}{\operatorname{arg\,min}} D(P \| P^{(S)} \otimes P^{(-S)})$$

with time complexity $\mathcal{O}(d^3\theta)$. Here, θ denotes the worst case time needed to evaluate $D(P||P^{(S)} \otimes P^{(-S)})$ for any given subset S.

4.2 Submodular maximization of the distance to factorizability

Given $P \in \mathcal{L}(\mathcal{X})$ and $m \in \mathbb{N}$, we aim to investigate the following submodular maximization problem subject to a cardinality constraint

$$\max_{S \subseteq \llbracket d \rrbracket; \ |S| \le m} D(P \| P^{(S)} \otimes P^{(-S)}).$$

$$\tag{17}$$

Since $D(P||P^{(S)} \otimes P^{(-S)}) \ge 0$ and $D(P||P^{(\emptyset)} \otimes P^{(\llbracket d \rrbracket)}) = 0$, if we consider the unconstrained version of (17), we can apply Algorithm 1 with $(\frac{1}{2} - \frac{\epsilon}{d})$ -approximation guarantee (see Theorem 2.7) since $D(P||P^{(S)} \otimes P^{(-S)})$ is symmetric.

In view of Theorem 2.5, we choose $\beta = 0$ and take

$$g(S) = D(P \| P^{(S)} \otimes P^{(-S)}) + \sum_{e \in S} D(P \| P^{(-e)} \otimes P^{(e)}),$$
(18)

which is submodular and monotonically non-decreasing. In this case, we also take the modular and non-negative function c to be

$$c(S) = \sum_{e \in S} D(P \| P^{(-e)} \otimes P^{(e)}).$$

$$\tag{19}$$

Therefore,

$$D(P \| P^{(S)} \otimes P^{(-S)}) = g(S) - c(S)$$

can be expressed as the difference of a non-negative, submodular, monotonically non-decreasing function and a non-negative modular function, hence Algorithm 2 can be applied to approximately maximize $D(P||P^{(S)} \otimes P^{(-S)})$.

Corollary 4.1. Let $P \in \mathcal{L}(\mathcal{X})$ be π -stationary. In Algorithm 2, we take g as in (18) and c as in (19), and $OPT = \arg \max_{S \subseteq \llbracket d \rrbracket; \ |S| \leq m} D(P \Vert P^{(S)} \otimes P^{(-S)})$. By Theorem 2.8, we have

$$D(P \| P^{(S_m)} \otimes P^{(-S_m)}) \ge (1 - e^{-1})g(\text{OPT}) - c(\text{OPT})$$

where S_m is the final output set of Algorithm 2.

4.3 k-submodular maximization of distance to factorizability of the tensorized keep- S_i -in matrices $D(P || P^{(S_1)} \otimes \ldots \otimes P^{(S_k)} \otimes P^{(-\cup_{i=1}^k S_i)})$

In this section, we investigate the following map

$$(k+1)^{\llbracket d \rrbracket} \ni \mathbf{S} \mapsto f(\mathbf{S}) = D(P \| P^{(S_1)} \otimes \ldots \otimes P^{(S_k)} \otimes P^{(-\bigcup_{i=1}^k S_i)}), \tag{20}$$

We consider the maximization problem of the form, for given $\mathbf{V} \in (k+1)^{\llbracket d \rrbracket}$,

$$\max_{\mathbf{S} \preceq \mathbf{V}; |\operatorname{supp}(\mathbf{S})| \le m} D(P \| P^{(S_1)} \otimes \ldots \otimes P^{(S_k)} \otimes P^{(-\bigcup_{i=1}^k S_i)}).$$
(21)

In the special case of k = 1 and $\mathbf{V} = \llbracket d \rrbracket$, we recover problem (17).

Lemma 4.2. The map (20) is orthant submodular.

Proof. We shall prove that $\Delta_{e,i} f(\mathbf{S}) \geq \Delta_{e,i} f(\mathbf{T})$, where we choose $\mathbf{S} \preceq \mathbf{T}$ and $e \notin \operatorname{supp}(\mathbf{T})$. We compute that

$$\begin{aligned} \Delta_{e,i}f(\mathbf{S}) - \Delta_{e,i}f(\mathbf{T}) &= H(P^{(S_i \cup \{e\})}) - H(P^{(S_i)}) + H(P^{(-\operatorname{supp}(\mathbf{S}) \cup \{e\})}) - H(P^{(-\operatorname{supp}(\mathbf{S}))}) \\ &- H(P^{(T_i \cup \{e\})}) + H(P^{(T_i)}) - H(P^{(-\operatorname{supp}(\mathbf{T}) \cup \{e\})}) + H(P^{(-\operatorname{supp}(\mathbf{T}))}) \\ &= \left[\left(H(P^{(S_i \cup \{e\})}) - H(P^{(S_i)}) \right) - \left(H(P^{(T_i \cup \{e\})}) - H(P^{(T_i)}) \right) \right] \\ &+ \left[\left(H(P^{(-\operatorname{supp}(\mathbf{T}))}) - H(P^{(-\operatorname{supp}(\mathbf{T}) \cup \{e\})}) \right) \\ &- \left(H(P^{(\operatorname{supp}(\mathbf{S}))}) - H(P^{(-\operatorname{supp}(\mathbf{S}) \cup \{e\})}) \right) \right], \end{aligned}$$

where each of the two terms above are non-negative given the submodularity of $S \mapsto H(P^{(S)})$ (recall Theorem 2.13).

In view of Theorem 2.6, since the map (20) is orthant submodular, for any $\beta \in \mathbb{R}$, if $\mathbf{S} \leq \mathbf{V}$, we have a monotonically non-decreasing k-submodular function given by

$$g(\mathbf{S}) = f(\mathbf{S}) - \beta + \sum_{i=1}^{k} \sum_{e \in S_i} \left[D(P \| P^{(V_1)} \otimes \ldots \otimes P^{(V_i \setminus \{e\})} \otimes \ldots \otimes P^{(V_k)} \otimes P^{(-\operatorname{supp}(\mathbf{V}) \setminus \{e\})}) \right]$$

$$= D(P \| P^{(V_1)} \otimes \ldots \otimes P^{(V_i)} \otimes \ldots \otimes P^{(V_k)} \otimes P^{(-\operatorname{supp}(\mathbf{V}))}) \right]$$

$$= f(\mathbf{S}) - \beta + \sum_{i=1}^{k} \sum_{e \in S_i} \left[H(P^{(V_i \setminus \{e\})}) + H(P^{(-\operatorname{supp}(\mathbf{V}) \setminus \{e\})}) - H(P^{(V_i)}) - H(P^{(-\operatorname{supp}(\mathbf{V}))}) \right]$$

$$= f(\mathbf{S}) - \beta + \sum_{i=1}^{k} \sum_{e \in S_i} \left[D(P^{(V_i)} \| P^{(V_i \setminus \{e\})} \otimes P^{(e)}) - D(P^{(-\operatorname{supp}(\mathbf{V}) \setminus \{e\})} \| P^{(-\operatorname{supp}(\mathbf{V}))} \otimes P^{(e)}) \right],$$
(22)

and we also obtain the following modular function

$$c(\mathbf{S}) = -\beta + \sum_{i=1}^{k} \sum_{e \in S_i} \left[D(P^{(V_i)} \| P^{(V_i \setminus \{e\})} \otimes P^{(e)}) - D(P^{(-\operatorname{supp}(\mathbf{V}) \setminus \{e\})} \| P^{(-\operatorname{supp}(\mathbf{V}))} \otimes P^{(e)}) \right].$$
(23)

Thus, if we choose

$$\beta \leq -\sum_{i=1}^{k} \sum_{e \in V_i} \left(H(P^{(-\operatorname{supp}(\mathbf{V}) \setminus \{e\})}) + H(P^{(e)}) \right),$$

then c is non-negative. With these choices, f can be written as

$$f(\mathbf{S}) = D(P \| P^{(S_1)} \otimes \ldots \otimes P^{(S_k)} \otimes P^{(-\bigcup_{i=1}^k S_i)}) = g(\mathbf{S}) - c(\mathbf{S}).$$

We can then apply Algorithm 3 to perform distorted greedy maximization with a lower bound.

Corollary 4.3. Let $P \in \mathcal{L}(\mathcal{X})$ be π -stationary and $\mathbf{V} \in (k+1)^{\llbracket d \rrbracket}$. In Algorithm 3, we take g as in (22) and c as in (23). We choose

$$\beta \leq -\sum_{i=1}^{k} \sum_{e \in V_i} \left(H(P^{(-\operatorname{supp}(\mathbf{V}) \setminus \{e\})}) + H(P^{(e)}) \right),$$

and let $\mathbf{OPT} = \arg \max_{\mathbf{S} \prec \mathbf{V}; |\operatorname{supp}(\mathbf{S})| \leq m} f(\mathbf{S})$. Therefore, Theorem 2.11 gives

$$f(\mathbf{S}_m) = D(P \| P^{(S_{m,1})} \otimes \ldots \otimes P^{(S_{m,k})} \otimes P^{(-\cup_{i=1}^k S_{m,i})}) \ge (1 - e^{-1})g(\mathbf{OPT}) - c(\mathbf{OPT}),$$

where $\mathbf{S}_m = (S_{m,1}, \ldots, S_{m,k})$ is the output of Algorithm 3.

Note that the lower bound of Corollary 4.3 depends on β through g and c. If β is chosen to be too small, then the lower bound might be too loose as the right hand side might be negative.

5 Supermodular minimization of distance to independence $\mathbb{I}(P^{(S)})$

Given $P \in \mathcal{L}(\mathcal{X})$ and $d, m \geq 2$, we aim to investigate the following supermodular (recall Theorem 2.13) minimization problem

$$\min_{S \subseteq \llbracket d \rrbracket; \ |S| = m} \mathbb{I}(P^{(S)}).$$
(24)

We shall be interested in the constraint |S| = m rather than $|S| \leq m$ as in Section 3 and Section 4 because $S \mapsto \mathbb{I}(P^{(S)})$ is monotonically non-decreasing.

The supermodular minimization problem (24) is equivalent to the following submodular maximization problem

$$\max_{S \subseteq \llbracket d \rrbracket; \ |S|=m} f(S) = - \mathbb{I}(P^{(S)}) = H(P^{(S)}) - \sum_{e \in S} H(P^{(e)}).$$
(25)

Note that we restrict m to be at least 2, since we have the trivial result that $\mathbb{I}(P^{(e)}) = \mathbb{I}(P^{(\emptyset)}) = 0$ if the constraint is m = 0 or m = 1. From Theorem 2.13, f(S) is monotonically non-increasing and submodular. Therefore, the heuristic greedy algorithm (see Section 4 of [13]) cannot provide a theoretical guarantee.

In view of Theorem 2.5, for any $\beta \in \mathbb{R}$, we have a monotonically non-decreasing submodular function g given by

$$g(S) = f(S) - \beta + \sum_{e \in S} (H(P^{(-e)}) + H(P^{(e)}) - H(P))$$

$$= f(S) - \beta + \sum_{e \in S} D(P || P^{(e)} \otimes P^{(-e)}).$$
(26)

We choose $\beta = 0$ and let the following non-negative, modular function be

$$c(S) = \sum_{e \in S} D(P \| P^{(e)} \otimes P^{(-e)})$$

$$\tag{27}$$

so that f(S) = g(S) - c(S). By Theorem 2.8, we can apply Algorithm 2 to obtain a lower bound.

Corollary 5.1. Let $P \in \mathcal{L}(\mathcal{X})$ be π -stationary along with $d, m \geq 2$. In Algorithm 2, we take g as in (26), c as in (27), and $OPT = \max_{S \subseteq \llbracket d \rrbracket; |S| = m} f(S)$. By Theorem 2.8, we have the following lower bound

$$f(S_m) = -\mathbb{I}(P^{(S_m)}) \ge (1 - e^{-1})g(\text{OPT}) - c(\text{OPT}),$$

where S_m is the output of Algorithm 2.

5.1 Supermodular minimization of distance to independence of the complement set $\mathbb{I}(P^{(-S)})$

From Theorem 2.14, $\mathbb{I}(P^{(-S)})$ is monotonically non-increasing and supermodular. Given $P \in \mathcal{L}(\mathcal{X})$, $d \geq 2$, and $m \leq d-2$, we shall investigate the following optimization problem

$$\max_{S \subseteq \llbracket d \rrbracket; \ |S| \le m} f(S) = - \mathbb{I}(P^{(-S)}).$$

Note that we restrict m to be at most d-2, since we have the trivial result that $\mathbb{I}(P^{(e)}) = \mathbb{I}(P^{(\emptyset)}) = 0$ if the constraint is m = d or m = d - 1.

Since $f(S) = -\mathbb{I}(P^{(-S)})$ is monotonically non-decreasing and submodular, then we can apply the heuristic greedy algorithm (see Section 4 of [13]) that comes along with a $(1 - e^{-1})$ -approximation guarantee.

5.2 k-supermodular minimization of distance to independence of the tensorized keep- S_i -in matrices $\mathbb{I}(\bigotimes_{i=1}^k P^{(S_i)})$

In this section, we investigate the following map

$$(k+1)^{\llbracket d \rrbracket} \ni \mathbf{S} = (S_1, \dots, S_k) \mapsto \mathbb{I}(\otimes_{i=1}^k P^{(S_i)}).$$

$$(28)$$

Lemma 5.2. For $k \in \mathbb{N}$ and $\mathbf{S} \in (k+1)^{\llbracket d \rrbracket}$, we have

$$\mathbb{I}(\otimes_{i=1}^k P^{(S_i)}) = \sum_{i=1}^k \mathbb{I}(P^{(S_i)})$$

Proof. We shall prove by induction on k. When k = 1, the equality trivially holds. When k = 2, according to the chain rule of KL divergence (see Theorem 2.15 of [14]),

$$\begin{split} \mathbb{I}(P^{(S_1)} \otimes P^{(S_2)}) &= D(P^{(S_1)} \otimes P^{(S_2)} \| \otimes_{i \in S_1 \cup S_2} P^{(i)}) \\ &= D(P^{(S_1)} \| \otimes_{i \in S_1} P^{(i)}) + D(P^{(S_2)} \| \otimes_{i \in S_2} P^{(i)}) \\ &= \mathbb{I}(P^{(S_1)}) + \mathbb{I}(P^{(S_2)}). \end{split}$$

Suppose $\mathbb{I}(\bigotimes_{i=1}^{m} P^{(S_i)}) = \sum_{i=1}^{m} \mathbb{I}(P^{(S_i)})$ holds (k = m), then using the chain rule of KL divergence again (Theorem 2.15 of [14]), we have

$$\mathbb{I}(\bigotimes_{i=1}^{m+1} P^{(S_i)}) = D(\bigotimes_{i=1}^{m} P^{(S_i)} \otimes P^{(S_{m+1})} \| \bigotimes_{i \in (\bigcup_{i=1}^{m} S_i) \cup S_{m+1}} P^{(i)})
= D(\bigotimes_{i=1}^{m} P^{(S_i)} \| \bigotimes_{i \in \bigcup_{i=1}^{m} S_i} P^{(i)}) + D(P^{(S_{m+1})} \| \bigotimes_{i \in S_{m+1}} P^{(i)})
= \sum_{i=1}^{m+1} \mathbb{I}(P^{(S_i)}).$$

Lemma 5.3. The map (28) is pairwise monotonically non-decreasing. In particular, when P is non-factorizable and π -stationary, the map (28) is pairwise monotonically strictly increasing for all pairs.

Proof. Let $f(\mathbf{S}) = \mathbb{I}(\bigotimes_{i=1}^{k} P^{(S_i)})$. We shall prove that $\Delta_{e,i} f(\mathbf{S}) + \Delta_{e,j} f(\mathbf{S}) \ge 0$, where $i \neq j \in \llbracket d \rrbracket$ and $e \notin \operatorname{supp}(\mathbf{T})$. Since $\mathbb{I}(P^{(S)}) = \sum_{i \in S} H(P^{(i)}) - H(P^{(S)})$, we note that

$$\begin{aligned} \Delta_{e,i}f(\mathbf{S}) + \Delta_{e,j}f(\mathbf{S}) &= \mathbb{I}(P^{(S_i \cup \{e\})}) - \mathbb{I}(P^{(S_i)}) + \mathbb{I}(P^{(S_j \cup \{e\})}) - \mathbb{I}(P^{(S_i)}) \\ &= \left[H(P^{(e)}) + H(P^{(S_i)}) - H(P^{(S_i \cup \{e\})})\right] \\ &+ \left[H(P^{(e)}) + H(P^{(S_j)}) - H(P^{(S_j \cup \{e\})})\right] \\ &= D(P^{(S_i \cup \{e\})} \|P^{(S_i)} \otimes P^{(e)}) + D(P^{(S_j \cup \{e\})} \|P^{(S_j)} \otimes P^{(e)}), \end{aligned}$$

which is non-negative. In particular, when P is non-factorizable, it is strictly positive.

Lemma 5.4. The map (28) is orthant supermodular.

Proof. Let $f(\mathbf{S}) = \mathbb{I}(\bigotimes_{i=1}^{k} P^{(S_i)})$. For any $\mathbf{S} \preceq \mathbf{T}$, we shall prove that $\Delta_{e,i} f(\mathbf{S}) \leq \Delta_{e,i} f(\mathbf{T})$, where $i \in \llbracket d \rrbracket$ and $e \in \llbracket d \rrbracket \setminus \sup(\mathbf{T})$.

$$\begin{aligned} \Delta_{e,i}f(\mathbf{S}) - \Delta_{e,i}f(\mathbf{T}) &= \left[H(P^{(e)}) + H(P^{(S_i)}) - H(P^{(S_i \cup \{e\})}) \right] \\ &- \left[H(P^{(e)}) + H(P^{(T_i)}) - H(P^{(T_i \cup \{e\})}) \right] \\ &= \left[H(P^{(T_i \cup \{e\})}) - H(P^{(T_i)}) \right] - \left[H(P^{(S_i \cup \{e\})}) - H(P^{(S_i)}) \right] \le 0 \end{aligned}$$

where the inequality holds owing to the submodularity of $S \mapsto H(P^{(S)})$ in view of Theorem 2.13.

Collecting the previous two results, we see that, for non-factorizable P, the map (28) is not k-supermodular as k-supermodularity requires both the pairwise monotonically non-increasing property and orthant supermodularity (see Theorem 2.2).

Given $P \in \mathcal{L}(\mathcal{X})$, $d, m \geq k+1$ and $\mathbf{V} \in (k+1)^{\llbracket d \rrbracket}$, since the map (28) is orthant supermodular, we are interested in the following orthant submodular maximization problem

$$\max_{\mathbf{S} \preceq \mathbf{V}; \, |\operatorname{supp}(\mathbf{S})|=m} f(\mathbf{S}) = -\mathbb{I}(\otimes_{i=1}^k P^{(S_i)}) = -\sum_{i=1}^k \mathbb{I}(P^{(S_i)}).$$

We are restricting m to be at least k + 1 following the pigeonhole principle, as we need at least one S_i with $|S_i| > 1$. If $m \le k$, we can take either $S_i = \{e\}$ or $S_i = \emptyset$ for all $i \in [\![k]\!]$ so that the optimization problem becomes trivial.

In view of Theorem 2.6, we have a monotonically non-decreasing and k-submodular function g given by

$$g(\mathbf{S}) = f(\mathbf{S}) - \beta + \sum_{i=1}^{k} \sum_{e \in S_i} [H(P^{(V_i \setminus \{e\})}) + H(P^{(e)}) - H(P^{(V_i)})]$$
(29)
= $f(\mathbf{S}) - \beta + \sum_{i=1}^{k} \sum_{e \in S_i} D(P^{(V_i)} || P^{(V_i \setminus \{e\})} \otimes P^{(e)}).$

We take $\beta = 0$, and denote the following non-negative modular function as

$$c(\mathbf{S}) = \sum_{i=1}^{k} \sum_{e \in S_i} D(P^{(V_i)} \| P^{(V_i \setminus \{e\})} \otimes P^{(e)})$$
(30)

so that $f(\mathbf{S}) = g(\mathbf{S}) - c(\mathbf{S})$. By applying Algorithm 3, we can obtain a result with the following lower bound by Theorem 2.11.

Corollary 5.5. Let $P \in \mathcal{L}(\mathcal{X})$ be π -stationary along with $d, m \geq k+1$ and $\mathbf{V} \in (k+1)^{\llbracket d \rrbracket}$. In Algorithm 3, we take g as in (29), c as in (30), and $\mathbf{OPT} = \arg \max_{\mathbf{S} \preceq \mathbf{V}; |\operatorname{supp}(\mathbf{S})|=m} f(\mathbf{S})$, then by Theorem 2.11, we have the following lower bound

$$f(\mathbf{S}_m) = -\mathbb{I}(\bigotimes_{i=1}^k P^{(S_{m,i})}) \ge (1 - e^{-1})g(\mathbf{OPT}) - c(\mathbf{OPT}),$$

where $\mathbf{S}_m = (S_{m,1}, \ldots, S_{m,k})$ is the output of Algorithm 3.

In the special case where k = 1 and $\mathbf{V} = \llbracket d \rrbracket$, we recover Corollary 5.1.

5.3 k-supermodular minimization of distance to independence of the tensorized keep- $V_i \setminus S_i$ -in matrices $\mathbb{I}(\bigotimes_{i=1}^k P^{(V_i \setminus S_i)})$

For given $\mathbf{V} \in (k+1)^{\llbracket d \rrbracket}$, we consider the following map in view of Lemma 5.2,

$$\{\mathbf{S} \in (k+1)^{\llbracket d \rrbracket}; \ \mathbf{S} \preceq \mathbf{V}\} \ni \mathbf{S} = (S_1, \dots, S_k) \mapsto \mathbb{I}(\otimes_{i=1}^k P^{(V_i \setminus S_i)}) = \sum_{i=1}^k \mathbb{I}(P^{(V_i \setminus S_i)}).$$
(31)

We first prove a result concerning monotonicity and k-supermodularity of the map above.

Theorem 5.6. The map (31) is monotonically non-increasing and k-supermodular.

Proof. In view of Theorem 2.14, for each component S_i , we take V_i as the ground set, hence $\mathbb{I}(P^{(V_i \setminus S_i)})$ is monotonically non-increasing and supermodular. From Lemma 5.2, this function is the sum of k monotonically non-increasing and supermodular functions. From Lemma 2.3, we conclude that this map is k-supermodular and monotonically non-increasing.

Therefore, we denote the following monotonically non-decreasing, k-submodular function g as

$$g(\mathbf{S}) = -\mathbb{I}(\bigotimes_{i=1}^{k} P^{(V_i \setminus S_i)}) = -\sum_{i=1}^{k} \mathbb{I}(P^{(V_i \setminus S_i)}).$$
(32)

Given $d \ge k+1$, $m \le d-k-1$, we are interested in the following maximization problem given by

$$\max_{\mathbf{S} \preceq \mathbf{V}; |\operatorname{supp}(\mathbf{S})| \le m} g(\mathbf{S}).$$

We are restricting m by $m \leq d - k - 1$ following the pigeonhole principle, as we want $|V_i \setminus S_i| \geq 2$ for at least one i. If $m \geq d - k$, we can choose either $V_i \setminus S_i = \{e\}$ or $V_i \setminus S_i = \emptyset$ so that the optimization problem is trivial.

By taking c = 0 as a non-negative modular function, we can apply Algorithm 3 to obtain an optimization result with $(1 - e^{-1})$ -approximation guarantee.

Corollary 5.7. Let $P \in \mathcal{L}(\mathcal{X})$ be π -stationary along with $d \ge k+1$, $m \le d-k-1$ and $\mathbf{V} \in (k+1)^{\llbracket d \rrbracket}$. In Algorithm 3, we take g as in (32), c = 0 and denote

$$\mathbf{OPT} = \underset{\mathbf{S} \preceq \mathbf{V}; |\operatorname{supp}(\mathbf{S})| \le m}{\operatorname{arg\,max}} g(\mathbf{S}).$$

From Theorem 2.11, we can obtain the following lower bound

$$g(\mathbf{S}_m) \ge (1 - e^{-1})g(\mathbf{OPT}),$$

where $\mathbf{S}_m = (S_{m,1}, \ldots, S_{m,k})$ is the output of Algorithm 3.

6 Supermodular minimization of distance to stationarity $D(P^{(S)} || \Pi^{(S)})$

In this section, we investigate the following map:

$$2^{\llbracket d \rrbracket} \ni S \mapsto D(P^{(S)} \| \Pi^{(S)}), \tag{33}$$

where Π is the matrix of stationary distribution with each row of Π being π . We first show that this map is monotonically non-decreasing.

Lemma 6.1. The map (33) is monotonically non-decreasing.

Proof. We choose $S \subseteq T \subseteq \llbracket d \rrbracket$. By the partition lemma (Theorem 2.12), we have

$$D(P^{(S)} \| \Pi^{(S)}) \le D(P^{(T)} \| \Pi^{(T)}),$$

and hence this map is monotonically non-decreasing.

We are interested in the following optimization problem

$$\max_{S \subseteq \llbracket d \rrbracket; \ |S|=m} D(P^{(S)} \| \Pi^{(S)}),$$

as solving the above can help to identify coordinates which are furthest away from the equilibrium in one step.

To solve this optimization problem with a theoretical guarantee, we recall the batch greedy algorithm (Algorithm 4, see Theorem 7 of [8]).

It turns out that the theoretical guarantee depends on the supermodularity ratio and submodularity ratio of a set function f, that we shall now briefly recall. The supermodularity ratio of a non-negative set function f (Definition 6 of [8]) with respect to the set U and a cardinality constraint $m \ge 1$ is

$$\eta_{U,m} := \min_{S \subseteq U; \ T : |T| \le m, S \cap T = \emptyset} \frac{f(S \cup T) - f(S)}{\sum_{e \in T} [f(S \cup \{e\}) - f(S)]}$$

Algorithm 4 Batch greedy algorithm

Require: monotonically non-decreasing set function f; ground set U; total cardinality constraint m; number of steps l and cardinality constraints q_i such that $\sum_{i=1}^{l} q_i = m$

- 1: Initialize $S_0 = \emptyset$
- 2: for i = 1 to l do
- 3: Determine incremental gains $f(S_{i-1} \cup \{e\}) f(S_{i-1}), \forall e \in U \setminus S_{i-1}$
- 4: Find Q, comprising the elements with top- q_i incremental gains
- 5: $S_i \leftarrow S_{i-1} \cup Q$
- 6: **end for**
- 7: Output: S_l

while the submodularity ratio of f (Definition 32 of [8]) with respect to the set U and a cardinality constraint $k \ge 1$ is

$$\gamma_{U,m} := \min_{S \subseteq U; \ T: |T| \le m, S \cap T = \emptyset} \frac{\sum_{e \in T} [f(S \cup \{e\}) - f(S)]}{f(S \cup T) - f(S)}.$$

We then state the lower bound pertaining to Algorithm 4 (see Theorem 7 of [8]).

Theorem 6.2 (Lower bound for batch greedy algorithm). Let $P \in \mathcal{L}(\mathcal{X})$ be π -stationary and U be the ground set. Let f be a monotonically non-decreasing set function with $f(\emptyset) = 0$. Algorithm 4 yields the following lower bound

$$f(S_l) \ge \left(1 - \prod_{i=1}^l \left(1 - \frac{q_i \cdot \eta_{U,q_i} \cdot \gamma_{U,m}}{m}\right)\right) \max_{S \subseteq U; \ |S|=m} f(S),$$

where S_l is the output set of Algorithm 4.

Since we have a monotonically mon-decreasing map (33) with $D(P^{(\emptyset)} || \Pi^{(\emptyset)}) = 0$, we can apply the Algorithm 4 (see Theorem 7 of [8]) with the following lower bound.

Corollary 6.3. Let $P \in \mathcal{L}(\mathcal{X})$ be π -stationary and $U = \llbracket d \rrbracket$ be the ground set. Let f be (33) which is a monotonically non-decreasing set function with $f(\emptyset) = 0$. Algorithm 4 yields the following lower bound

$$f(S_l) \ge \left(1 - \prod_{i=1}^l \left(1 - \frac{q_i \cdot \eta_{U,q_i} \cdot \gamma_{U,m}}{m}\right)\right) \max_{S \subseteq \llbracket d \rrbracket; \ |S|=m} f(S),$$

where S_l is the output set of Algorithm 4.

We now consider the special case where the stationary distribution π is of product form. In this case, we can show the supermodularity of the map (33).

Lemma 6.4. The map (33) is supermodular if P is π -stationary where π is of product form. Proof.

$$\begin{split} D(P^{(S)} \| \Pi^{(S)}) &= \sum_{x^{(S)}} \sum_{y^{(S)}} \pi^{(S)}(x^{(S)}) P^{(S)}(x^{(S)}, y^{(S)}) \ln \frac{P^{(S)}(x^{(S)}, y^{(S)})}{\pi^{(S)}(y^{(S)})} \\ &= -H(P^{(S)}) - \sum_{x^{(S)}} \sum_{y^{(S)}} \pi^{(S)}(x^{(S)}) P^{(S)}(x^{(S)}, y^{(S)}) \ln \pi^{(S)}(y^{(S)}) \\ &= -H(P^{(S)}) - \sum_{y^{(S)}} \ln \pi^{(S)}(y^{(S)}) \sum_{x^{(S)}} \pi^{(S)}(x^{(S)}) P^{(S)}(x^{(S)}, y^{(S)}) \\ &= -H(P^{(S)}) + H(\pi^{(S)}). \end{split}$$

The last equation holds since P is π -stationary and hence

$$\pi^{(S)}(y^{(S)}) = \sum_{x^{(S)}} \pi^{(S)}(x^{(S)}) P^{(S)}(x^{(S)}, y^{(S)}).$$

Since the stationary distribution π is of product form, then $\pi = \bigotimes_{i=1}^{d} \pi^{(i)}$, hence $H(\pi^{(S)}) = \sum_{i \in S} H(\pi^{(i)})$, which is a modular function. Also, since $H(P^{(S)})$ is submodular, then $-H(P^{(S)})$ is supermodular. Therefore, $D(P^{(S)} || \Pi^{(S)})$ is supermodular because it is a sum of a supermodular function and a modular function.

We proceed to investigate the following optimization problem when P is π -stationary with product form π ,

$$\max_{S \subseteq [\![d]\!]; |S| \le m} f(S) = -D(P^{(S)} \| \Pi^{(S)}).$$

In view of Theorem 2.5, the following function g is monotonically non-decreasing and submodular since f is submodular:

$$g(S) = f(S) - \beta + \sum_{e \in S} (H(P^{(-e)}) - H(\pi^{(-e)}) - H(P) + H(\pi))$$

= $f(S) - \beta + \sum_{e \in S} (D(P || P^{(e)} \otimes P^{(-e)}) + D(P^{(e)} || \Pi^{(e)})).$ (34)

Choosing $\beta = 0$, we denote the following non-negative modular function as

$$c(S) = \sum_{e \in S} (D(P \| P^{(e)} \otimes P^{(-e)}) + D(P^{(e)} \| \Pi^{(e)})).$$
(35)

Since f(S) = g(S) - c(S), we apply Algorithm 2 to obtain a result with the following lower bound:

Corollary 6.5. Let $P \in \mathcal{L}(\mathcal{X})$ be π -stationary with π to be product form. In Algorithm 2, we take g as in (34), c as in (35), and $OPT = \arg \max_{S \subseteq \llbracket d \rrbracket; |S| \leq m} f(S)$. By Theorem 2.8, we have the following lower bound

$$f(S_m) = -D(P^{(S_m)} \| \Pi^{(S_m)}) \ge (1 - e^{-1})g(\text{OPT}) - c(\text{OPT}),$$

where S_m is the output set of Algorithm 2.

6.1 Supermodular minimization of distance to stationarity of the complement set $D(P^{(-S)} || \Pi^{(-S)})$

In this section, we shall investigate the following map:

$$2^{\llbracket d \rrbracket} \ni S \mapsto D(P^{(-S)} \| \Pi^{(-S)}).$$
(36)

Owing to Lemma 6.1, we first see that the map (36) is monotonically non-increasing. In addition, the map (36) is supermodular if P is π -stationary with product form π in view of Lemma 2.1 and Lemma 6.4.

We are interested in the following optimization problem

$$\max_{S \subseteq [\![d]\!]; |S| \le m} f(S) = -D(P^{(-S)} \| \Pi^{(-S)}),$$

as solving the above allows us to identify coordinates whose complement set is the closest to equilibrium in one step.

Under the assumption of product form π , as the map (36) is monotonically non-increasing and supermodular, f is monotonically non-decreasing and submodular. We apply the heuristic greedy algorithm (Section 4 of [13]) to obtain an approximate maximizer along with a $(1 - e^{-1})$ -approximation guarantee.

6.2 k-supermodular minimization of distance to stationarity of tensorized keep- S_i -in matrices $D(\bigotimes_{i=1}^k P^{(S_i)} || \bigotimes_{i=1}^k \Pi^{(S_i)})$

In this section, for given $\mathbf{V} \in (k+1)^{\llbracket d \rrbracket}$, we investigate the following map:

$$(k+1)^{\llbracket d \rrbracket} \ni \mathbf{S} = (S_1, \dots, S_k) \mapsto f(\mathbf{S}) = D(\bigotimes_{i=1}^k P^{(S_i)} \| \bigotimes_{i=1}^k \Pi^{(S_i)}).$$
(37)

We first give an orthant supermodularity result.

Lemma 6.6. The map (37) is orthant supermodular if P is π -stationary where π is of product form.

Proof. By the chain rule or tensorization property of KL divergence (see Theorem 2.15 and 2.16 of [14]), we see that

$$D(\bigotimes_{i=1}^{k} P^{(S_i)} \| \bigotimes_{i=1}^{k} \Pi^{(S_i)}) = \sum_{i=1}^{k} D(P^{(S_i)} \| \Pi^{(S_i)}).$$

We now take $\mathbf{S} \leq \mathbf{T}$ and $e \in [\![d]\!] \setminus T_i$. By (1), we aim to show that $\Delta_{e,i} f(\mathbf{S}) \leq \Delta_{e,i} f(\mathbf{T})$, which indeed holds since

$$\begin{aligned} \Delta_{e,i} f(\mathbf{S}) &= D(P^{(S_i \cup \{e\})} \| \Pi^{(S_i \cup \{e\})}) - D(P^{(S_i)} \| \Pi^{(S_i)}) \\ &\leq D(P^{(T_i \cup \{e\})} \| \Pi^{(T_i \cup \{e\})}) - D(P^{(T_i)} \| \Pi^{(T_i)}) = \Delta_{e,i} f(\mathbf{T}) \end{aligned}$$

because $S \mapsto D(P^{(S)} || \Pi^{(S)})$ is supermodular (see Theorem 6.4).

We are interested in the following optimization problem

$$\max_{\mathbf{S} \preceq \mathbf{V}; |\operatorname{supp}(\mathbf{S}_m)| \le m} -f(\mathbf{S}),$$

where f is orthant supermodular.

In view of Theorem 2.6, we have the following monotonically non-decreasing, k-submodular function g:

$$g(\mathbf{S}) = -f(\mathbf{S}) - \beta + \sum_{i=1}^{k} \sum_{e \in S_i} (D(P^{(V_i)} \| P^{(e)} \otimes P^{(V_i \setminus e)}) + D(P^{(e)} \| \Pi^{(e)})).$$
(38)

We take $\beta = 0$, and denote the non-negative modular function as

$$c(\mathbf{S}) = \sum_{i=1}^{k} \sum_{e \in S_i} (D(P^{(V_i)} \| P^{(e)} \otimes P^{(V_i \setminus e)}) + D(P^{(e)} \| \Pi^{(e)})).$$
(39)

Since $-f(\mathbf{S}) = g(\mathbf{S}) - c(\mathbf{S})$, we apply Algorithm 3 to obtain an approximate maximizer along with a lower bound.

Corollary 6.7. Let $P \in \mathcal{L}(\mathcal{X})$ be π -stationary with π be of product form and $\mathbf{V} \in (k+1)^{\llbracket d \rrbracket}$. In Algorithm 3, we take g as in (38), c as in (39), and $\mathbf{OPT} = \arg \max_{\mathbf{S} \preceq \mathbf{V}; |\operatorname{supp}(\mathbf{S}_m)| \leq m} -f(\mathbf{S})$. Then Theorem 2.11 gives the following lower bound

$$-f(\mathbf{S}_m) \ge (1 - e^{-1})g(\mathbf{OPT}) - c(\mathbf{OPT}),$$

where $\mathbf{S}_m = (S_{m,1}, \ldots, S_{m,k})$ is the output of Algorithm 3.

6.3 k-supermodular minimization of distance to stationarity of tensorized keep- $V_i \setminus S_i$ -in matrices $D(\bigotimes_{i=1}^k P^{(V_i \setminus S_i)} || \bigotimes_{i=1}^k \Pi^{(V_i \setminus S_i)})$

For given $\mathbf{V} \in (k+1)^{\llbracket d \rrbracket}$, we investigate the following map:

$$\{\mathbf{S} \in (k+1)^{\llbracket d \rrbracket}; \ \mathbf{S} \preceq \mathbf{V}\} \ni \mathbf{S} = (S_1, \dots, S_k) \mapsto D(\bigotimes_{i=1}^k P^{(V_i \setminus S_i)} \| \bigotimes_{i=1}^k \Pi^{(V_i \setminus S_i)}).$$
(40)

Theorem 6.8. The map (40) is monotonically non-increasing and k-supermodular if P is π -stationary where π is of product form.

Proof. By the chain rule or tensorization property of KL divergence (see Theorem 2.15 and 2.16 of [14]), we see that

$$D(\otimes_{i=1}^{k} P^{(V_{i} \setminus S_{i})} \| \otimes_{i=1}^{k} \Pi^{(V_{i} \setminus S_{i})}) = \sum_{i=1}^{k} D(P^{(V_{i} \setminus S_{i})} \| \Pi^{(V_{i} \setminus S_{i})})$$

which is a sum of k monotonically non-increasing and supermodular functions in view of Lemma 2.3. \Box

We are interested in the following optimization problem

$$\max_{\mathbf{S} \leq \mathbf{V}; \ |\operatorname{supp}(\mathbf{S})| \leq m} g(\mathbf{S}) = -D(\bigotimes_{i=1}^{k} P^{(V_i \setminus S_i)} \| \bigotimes_{i=1}^{k} \Pi^{(V_i \setminus S_i)}).$$
(41)

Since the map (40) is monotonically non-increasing and k-supermodular, then g is monotonically nondecreasing and k-submodular. We apply Algorithm 3 to obtain a $(1 - e^{-1})$ -approximation guarantee.

Corollary 6.9. Let $P \in \mathcal{L}(\mathcal{X})$ be π -stationary with product form π and $\mathbf{V} \in (k+1)^{\llbracket d \rrbracket}$. We take g as in (41), c = 0 and $\mathbf{OPT} = \arg \max_{\mathbf{S} \preceq \mathbf{V}; |\operatorname{supp}(\mathbf{S})| \le m} g(\mathbf{S})$. According to Theorem 2.11, we have the following lower bound for Algorithm 3

$$g(\mathbf{S}_m) \ge (1 - e^{-1})g(\mathbf{OPT}),$$

where $\mathbf{S}_m = (S_{m,1}, \dots, S_{m,k})$ is the output of Algorithm 3.

In the special case where k = 1 and $\mathbf{V} = [\![d]\!]$, the above Corollary reduces to the $(1 - e^{-1})$ -approximation guarantee as in Section 6.1.

7 Distance to factorizability over a fixed set $D(P^{(W \cup S)} || P^{(W)} \otimes P^{(S)})$

We fix a set $W \subseteq \llbracket d \rrbracket$ and investigate the following function:

$$\{S \subseteq \llbracket d \rrbracket; \ S \cap W = \emptyset\} \ni S \mapsto f(S) = D(P^{(W \cup S)} \| P^{(W)} \otimes P^{(S)}).$$

$$\tag{42}$$

We shall investigate the following optimization problem with cardinality constraint

$$\max_{S \subseteq \llbracket d \rrbracket; S \cap W = \emptyset; |S| = m} f(S).$$

We pick $S, T \subseteq \{S \subseteq \llbracket d \rrbracket; S \cap W = \emptyset\}$ with $S \subseteq T$ and compute that

$$f(S) - f(T) = [H(P^{(T \cup W)}) - H(P^{(T)})] - [H(P^{(S \cup W)}) - H(P^{(S)})] \le 0,$$

where the inequality follows from the property that $S \mapsto H(P^{(S)})$ is submodular (see Theorem 2.13). Therefore f is monotonically non-decreasing. Also, $f(\emptyset) = D(P^{(W)} || P^{(W)} \otimes P^{(\emptyset)}) = 0$. As such, we can apply Algorithm 4 (see Theorem 6.2) with a lower bound.

Corollary 7.1. Let $P \in \mathcal{L}(\mathcal{X})$ be π -stationary, $W \subseteq \llbracket d \rrbracket$, and $U = \llbracket d \rrbracket \setminus W$ be the ground set. Let f be (42) which is a monotonically non-decreasing set function with $f(\emptyset) = 0$. Algorithm 4 yields the following lower bound

$$f(S_l) \ge \left(1 - \prod_{i=1}^l \left(1 - \frac{q_i \cdot \eta_{U,q_i} \cdot \gamma_{U,m}}{m}\right)\right) \sum_{S \subseteq \llbracket d \rrbracket; S \cap W = \emptyset; |S| = m} f(S),$$

where S_l is the output set of Algorithm 4.

8 Numerical experiments¹

We conduct a case study to evaluate the numerical performance of the submodular optimization algorithms on the information-theoretic properties of multivariate Markov chains.

8.1 Experiment settings - Curie-Weiss model

We consider a discrete d-dimensional hypercube state space given by

$$\mathcal{X} = \{-1, +1\}^d.$$

¹The code is available at: https://github.com/zheyuanlai/SubmodOptMC.

Let the Hamiltonian function be that of the Curie-Weiss model (see Chapter 13 of [1]) on \mathcal{X} with interaction coefficients $\frac{1}{2^{|j-i|}}$ and external magnetic field $h \in \mathbb{R}$, that is, for $x = (x^1, \ldots, x^d) \in \mathcal{X}$,

$$\mathcal{H}(x) = -\sum_{i=1}^{d} \sum_{j=1}^{d} \frac{1}{2^{|j-i|}} x^{i} x^{j} - h \sum_{i=1}^{d} x^{i}.$$

We consider a Glauber dynamics with a simple random walk proposal targeting the Gibbs distribution at temperature $T \ge 0$. At each step we pick uniformly at random one of the *d* coordinates and flip it to the opposite sign, along with an acceptance-rejection filter, that is,

$$P(x,y) = \begin{cases} \frac{1}{d} e^{-\frac{1}{T}(\mathcal{H}(y) - \mathcal{H}(x))_{+}}, & \text{if } y = (x^{1}, x^{2}, \dots, -x^{i}, \dots, x^{d}), i \in [\![d]\!], \\ 1 - \sum_{y; \ y \neq x} P(x, y), & \text{if } x = y, \\ 0, & \text{otherwise}, \end{cases}$$

where for $m \in \mathbb{R}$ we denote $m_+ := \max\{m, 0\}$ the non-negative part of m. The stationary distribution of P is the Gibbs distribution at temperature T given by

$$\pi(x) = \frac{e^{-\frac{1}{T}\mathcal{H}(x)}}{\sum_{z \in \mathcal{X}} e^{-\frac{1}{T}\mathcal{H}(z)}}.$$

Parameters. We aim to generate a 10-dimensional Markov chain from the Curie-Weiss model. We choose d = 10, and hence the state space is of product form with $\mathcal{X} = \{-1, +1\}^{10}$. The choices of Hamiltonian function $\mathcal{H}(x)$, transition matrix P, and the stationary distribution $\pi(x)$ are detailed in Section 8.1, and we choose T = 10 as the temperature, h = 1 as the external magnetic field. For the numerical experiments of the generalized distorted greedy algorithm, we choose $\mathbf{V} = (V_1, V_2, V_3)$ where $V_1 = \{1, 2, 3, 4\}, V_2 = \{5, 6, 7\}$, and $V_3 = \{8, 9, 10\}$.

8.2 Experiment settings - Bernoulli-Laplace level model

We consider a d-dimensional Bernoulli–Laplace level model as described in Section 4.2 of [9]. Let

$$\mathcal{X} = \{x = (x^1, \dots, x^d) \in \mathbb{N}_0^d; x^1 + \dots + x^d = N\}$$

be the state space, where x^i can be interpreted as the number of "particles" of type *i* out of the total number *N*. The stationary distribution of such Markov chain, π , is given by the multivariate hypergeometric distribution described in Lemma 4.18 of [9]. Concretely, we have

$$\pi(x) = \frac{\prod_{i=1}^{d} \binom{l_i}{x^i}}{\binom{l_1+\dots+l_d}{N}}, \quad x \in \mathcal{X},$$
(43)

for some fixed parameters $l_1, \ldots, l_d \in \mathbb{N}$ representing the total number of "particles" of type *i*.

Following the spectral decomposition for reversible Markov chains (see Section 2.1 of [9] for background), the transition matrix P is written as:

$$P(x,y) = \sum_{n=0}^{N} \beta_n \phi_n(x) \phi_n(y) \pi(y),$$
(44)

where β_n are the eigenvalues and $\phi_n(x)$ is the eigenfunction.

From Definition 4.15 of [9], in the Bernoulli-Laplace level model, s is the swap size parameter satisfying

$$0 \le s \le \min\left\{N, \sum_{i=1}^d l_i - N\right\},\,$$

where we consider $\sum_{i=1}^{d} l_i > N$. From Theorem 4.19 of [9], the eigenvalues for the Bernoulli-Laplace level model are given by

$$\beta_n = \sum_{k=0}^n \binom{n}{k} \frac{(N-s)_{[n-k]} s_{[k]}}{N_{[n-k]} \left(\sum_{i=1}^d l_i - N\right)_{[k]}}, \quad 0 \le n \le N,$$

where $a_{[k]} = a(a-1)\cdots(a-k+1)$, and we apply the convention that $a_{[0]} = 1$.

In this case, we choose the eigenfunction as

$$\phi_n(x) = \left\{ \mathbf{Q_n}\left(x; N, -\sum_{i=1}^d l_i\right) \right\}_{|\mathbf{n}|=n}$$

where $\mathbf{Q}_{\mathbf{n}}$ are the multivariate Hahn polynomials for the hypergeometric distribution as defined in Proposition 2.3 of [9].

Parameters. We aim to generate a 10-dimensional Markov chain from the Bernoulli-Laplace level model. We consider the special case where s = 1 and choose d = 11, $l_1 = \ldots = l_{10} = 1$, N = 10, and $l_{11} = 10$. We let $x^{11} = N - \sum_{i=1}^{10} x^i$, and hence the state space is of product form with $\mathcal{X} = \{0, 1\}^{10}$.

The transition probabilities follow the dynamics as in (44), where particles hop between coordinates while respecting capacity constraints, and the stationary distribution π is computed as in (43). For the numerical experiments of the generalized distorted greedy algorithm, we choose $\mathbf{V} = (V_1, V_2, V_3)$ where $V_1 = \{1, 2, 3, 4\}, V_2 = \{5, 6, 7\}$, and $V_3 = \{8, 9, 10\}$.

8.3 Experiment results of Section 3

In this section, we report the numerical experiment results related to Section 3, which contains the performance of the heuristic greedy algorithm (see Section 4 of [13]), the distorted greedy algorithm (see Corollary 3.2), and the generalized distorted greedy algorithm (see Corollary 3.5) on the Bernoulli-Laplace level model (see Section 8.2) and the Curie-Weiss model (see Section 8.1). For each experiment, we conduct submodular optimization with cardinality constraint m, with m ranging from 1 to 10.

	Greedy		Distorted Greedy		
m	Subset S_m	$H(P^{(S_m)})$	Subset S_m	$H(P^{(S_m)})$	
1	{10}	0.46094	{10}	0.46094	
2	$\{3, 10\}$	0.83616	$\{1, 10\}$	0.83573	
3	$\{1, 3, 10\}$	1.17940	$\{1, 2, 5\}$	1.18116	
4	$\{1, 2, 3, 10\}$	1.49461	$\{1, 2, 3, 5\}$	1.50706	
5	$\{1, 2, 3, 4, 10\}$	1.77855	$\{1, 2, 3, 4, 5\}$	1.80193	
6	$\{1, 2, 3, 4, 5, 10\}$	2.03516	$\{1, 2, 3, 4, 5, 6\}$	2.06105	
7	$\{1, 2, 3, 4, 5, 6, 10\}$	2.25729	$\{1, 2, 3, 4, 5, 6, 7\}$	2.28328	
8	$\{1, 2, 3, 4, 5, 6, 7, 10\}$	2.43498	$\{1, 2, 3, 4, 5, 6, 7, 8\}$	2.45453	
9	$\{1, 2, 3, 4, 5, 6, 7, 8, 10\}$	2.51897	$\{1, 2, 3, 4, 5, 6, 7, 8, 10\}$	2.51897	
10	$\{1, 2, 3, 4, 5, 6, 7, 8, 10\}$	2.51897	$\{1, 2, 3, 4, 5, 6, 7, 8, 10\}$	2.51897	

Table 1: Comparison of the greedy algorithm and the distorted greedy algorithm. Entropy rate of the full chain of the Bernoulli-Laplace level model is H(P) = 1.96068.



Figure 1: Entropy rate against subset size for the three algorithms (B-L model).

Cardinality m	Subset $S_{m,1}$	Subset $S_{m,2}$	Subset $S_{m,3}$	$H(\otimes_{i=1}^{3} P^{(S_{m,i})})$
1	Ø	Ø	{10}	0.46094
2	Ø	{7}	$\{10\}$	0.90046
3	Ø	{7}	$\{8, 9\}$	1.26966
4	{4}	{7}	$\{8, 9\}$	1.70072
5	{4}	$\{5, 7\}$	$\{8, 9\}$	2.08692
6	{4}	$\{5, 6, 7\}$	$\{8, 9\}$	2.43035
7	{4}	$\{5, 6, 7\}$	$\{8, 9, 10\}$	2.71405
8	$\{3, 4\}$	$\{5, 6, 7\}$	$\{8, 9, 10\}$	3.10451
9	$\{1, 2, 4\}$	$\{5, 6, 7\}$	$\{8, 9, 10\}$	3.46267
10	$\{1, 2, 3, 4\}$	$\{5, 6, 7\}$	$\{8, 9, 10\}$	3.78968

Table 2: Performance evaluation of the generalized distorted greedy algorithm. Entropy rate of the full chain of the Bernoulli-Laplace level model is H(P) = 1.96068.

For the Bernoulli-Laplace level model, Table 1 and Figure 1a show the entropy rates of the output of the greedy algorithm and the distorted greedy algorithm (Algorithm 2); Table 2 and Figure 1b show the entropy rates of the tensorized output of the generalized distorted greedy algorithm (Algorithm 3).

	Greedy		Distorted Greedy		
m	Subset S_m	$H(P^{(S_m)})$	Subset S_m	$H(P^{(S_m)})$	
1	{1}	0.29085	{1}	0.29085	
2	$\{1, 10\}$	0.57371	$\{1, 10\}$	0.57371	
3	$\{1, 9, 10\}$	0.83933	$\{1, 9, 10\}$	0.83933	
4	$\{1, 2, 9, 10\}$	1.09570	$\{1, 2, 9, 10\}$	1.09570	
5	$\{1, 2, 6, 9, 10\}$	1.33953	$\{1, 2, 6, 9, 10\}$	1.33953	
6	$\{1, 2, 4, 6, 9, 10\}$	1.57098	$\{1, 2, 4, 6, 9, 10\}$	1.57098	
7	$\{1, 2, 4, 6, 8, 9, 10\}$	1.78757	$\{1, 2, 4, 6, 8, 9, 10\}$	1.78757	
8	$\{1, 2, 3, 4, 6, 8, 9, 10\}$	1.98500	$\{1, 2, 3, 4, 6, 7, 9, 10\}$	1.98458	
9	$\{1, 2, 3, 4, 6, 7, 8, 9, 10\}$	2.15793	$\{1, 2, 3, 4, 6, 7, 8, 9, 10\}$	2.15793	
10	$\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$	2.29109	$\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$	2.29109	

Table 3: Comparison of the greedy algorithm and the distorted greedy algorithm. Entropy rate of the full chain of the Curie-Weiss model is H(P) = 2.29109.



Figure 2: Entropy rate against subset size for the three algorithms (C-W model).

Cardinality m	Subset $S_{m,1}$	Subset $S_{m,2}$	Subset $S_{m,3}$	$H(\otimes_{i=1}^{3} P^{(S_{m,i})})$
1	{1}	Ø	Ø	0.29085
2	{1}	{7}	Ø	0.57067
3	{1}	{7}	$\{10\}$	0.86152
4	{1}	$\{5,7\}$	$\{10\}$	1.13316
5	{1}	$\{5,7\}$	$\{9, 10\}$	1.40732
6	{1}	$\{5, 6, 7\}$	$\{9, 10\}$	1.66816
7	{1}	$\{5, 6, 7\}$	$\{8, 9, 10\}$	1.93090
8	$\{1, 2\}$	$\{5, 6, 7\}$	$\{8, 9, 10\}$	2.20505
9	$\{1, 2, 4\}$	$\{5, 6, 7\}$	$\{8, 9, 10\}$	2.46832
10	$\{1, 2, 3, 4\}$	$\{5, 6, 7\}$	$\{8, 9, 10\}$	2.72011

Table 4: Performance evaluation of the generalized distorted greedy algorithm. Entropy rate of the full chain of the Curie-Weiss model is H(P) = 2.29109.

For the Curie-Weiss model, Table 3 and Figure 2a show the entropy rates of the output of the greedy algorithm and the distorted greedy algorithm (Algorithm 2); Table 4 and Figure 2b show the entropy rates of the tensorized output of the generalized distorted greedy algorithm (Algorithm 3).

Notably, in Table 1 and Figure 1a, the distorted greedy algorithm outperforms the heuristic greedy algorithm when the cardinality constraint equals to m = 3, 4, 5, 6, 7, 8. This is because, in the distorted greedy algorithm, the distortion term $(1 - \frac{1}{m})^{m-(i+1)}$ at each step is different with different cardinality constraint m, which results in possibly better or different results than the heuristic greedy algorithm. However, the distorted greedy algorithm does not necessarily select better subset than the heuristic greedy algorithm, see the example of m = 2 in Table 1 and m = 8 in Table 3.

8.4 Experiment results of Section 4

We report the numerical experiment results related to Section 4, which contains the performance of the heuristic greedy algorithm (Section 4 of [13]), the distorted greedy algorithm (Algorithm 2), and the generalized distorted greedy algorithm (Algorithm 3) on the Curie-Weiss model as detailed in Section 8.1.

		Greedy	Distorted Greedy		
m	Subset S_m	$D\left(P\ P^{(S_m)}\otimes P^{(-S_m)}\right)$	Subset S_m	$D\left(P\ P^{(S_m)}\otimes P^{(-S_m)}\right)$	
1	$\{6\}$	0.14837	$\{6\}$	0.14837	
2	$\{2, 6\}$	0.24497	$\{3, 10\}$	0.24496	
3	$\{2, 6, 9\}$	0.30927	$\{3,7\}$	0.24525	
4	$\{2, 5, 6, 9\}$	0.34590	$\{2, 7, 10\}$	0.30905	
5	$\{2, 3, 5, 6, 9\}$	0.35758	$\{2, 3, 6, 10\}$	0.34590	

Table 5: Comparison of the greedy algorithm and the distorted greedy algorithm.



Figure 3: Distance to factorizability against subset size for the three algorithms.

For the experiments related to heuristic greedy and distorted greedy algorithms, since the map $S \mapsto D(P || P^{(S)} \otimes P^{(-S)})$ is symmetric, we conduct submodular maximization with cardinality constraint m, with m only ranging from 1 to 5. The results are shown on Table 5 and Figure 3a. These results show that although the distorted greedy algorithm has a lower bound as detailed in Corollary 4.1, the performance is not guaranteed to be better than the heuristic greedy algorithm. We also conduct the generalized distorted greedy algorithm as detailed in Corollary 4.3 with cardinality constraint m ranging from 1 to 10, and the results are shown on Table 6 and Figure 3b.

m	Subset $S_{m,1}$	Subset $S_{m,2}$	Subset $S_{m,3}$	$D\left(P\ \left(\otimes_{i=1}^{3}P^{(S_{m,i})}\right)\otimes P^{(-\cup_{i=1}^{3}S_{m,i})}\right)$
1	Ø	{6}	Ø	0.14836
2	Ø	{7}	{8}	0.25388
3	{4}	{7}	{8}	0.33529
4	{4}	$\{5,7\}$	{8}	0.39056
5	$\{2,4\}$	$\{5,7\}$	{8}	0.43104
6	$\{2,4\}$	$\{5,7\}$	$\{8, 10\}$	0.45978
7	$\{2,4\}$	$\{5, 6, 7\}$	$\{8, 10\}$	0.46887
8	$\{2,4\}$	$\{5, 6, 7\}$	$\{8, 10\}$	0.46887
9	$\{2,4\}$	$\{5, 6, 7\}$	$\{8, 10\}$	0.46887
10	$\{2,4\}$	$\{5, 6, 7\}$	$\{8, 10\}$	0.46887

Table 6: Performance evaluation of the generalized distorted greedy algorithm.

We conduct similar numerical experiments on the Bernoulli-Laplace level model (see Section 8.2). Among all cardinality constraints, the greedy algorithm and the distorted greedy algorithm output $S_m = \{10\}$, and the generalized distorted greedy algorithm outputs $S_{m,1} = S_{m,2} = \emptyset$, $S_{m,3} = \{10\}$. The reason behind it is that for a 10-dimensional Markov chain, the coordinate 10 is "far" from other coordinates.

8.5 Experiment results of Section 5

We report the numerical experiment results related to Section 5, which contains the performance of the heuristic greedy algorithm (see Section 4 of [13]), the distorted greedy algorithm (see Corollary 5.1), and the generalized distorted greedy algorithm (see Corollary 5.5) on the Bernoulli-Laplace level model (see Section 8.2) and the Curie-Weiss model (see Section 8.1). For each experiment, we conduct supermodular minimization with different cardinality constraint m's.

For the Bernoulli-Laplace level model, Table 7 and Figure 4a show the distance to independence of the outputs of the greedy algorithm and the distorted greedy algorithm (Algorithm 2). We note that the distorted greedy algorithm often outperforms the greedy algorithm. Table 8 and Figure 4b show the distance to independence of the tensorized outputs of the generalized distorted greedy algorithm (Algorithm 3).

	Greedy	Distorted Greedy		
m	Subset S_m	$\left \mathbb{I}(P^{(S_m)}) \right $	Subset S_m	$\mathbb{I}(P^{(S_m)})$
2	$\{1, 10\}$	0.05140	$\{1, 2\}$	0.03406
3	$\{1, 2, 10\}$	0.13505	$\{1, 2, 3\}$	0.10318
4	$\{1, 2, 3, 10\}$	0.24989	$\{1, 2, 3, 4\}$	0.20793
5	$\{1, 2, 3, 4, 10\}$	0.39701	$\{1, 2, 3, 4, 5\}$	0.34753
6	$\{1, 2, 3, 4, 5, 10\}$	0.57523	$\{1, 2, 3, 4, 5, 6\}$	0.52441
7	$\{1, 2, 3, 4, 5, 6, 10\}$	0.78911	$\{1, 2, 3, 4, 5, 6, 7\}$	0.74171
8	$\{1, 2, 3, 4, 5, 6, 7, 10\}$	1.05094	$\{1, 2, 3, 4, 5, 6, 7, 8\}$	1.01576
9	$\{1, 2, 3, 4, 5, 6, 7, 8, 10\}$	1.41226	$\{1, 2, 3, 4, 5, 6, 7, 8, 10\}$	1.41226
10	$\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$	2.41825	$\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$	2.41825

Table 7: Comparison of the greedy algorithm and the distorted greedy algorithm (B-L model).

m	Subset $S_{m,1}$	Subset $S_{m,2}$	Subset $S_{m,3}$	$\mathbb{I}\left(\otimes_{i=1}^{3} P^{(S_{m,i})}\right)$
4	$\{1, 2\}$	$\{5\}$	{8}	0.03406
5	$\{1, 2\}$	$\{5, 6\}$	$\{8\}$	0.07999
6	$\{1, 2\}$	$\{5, 6\}$	$\{8, 9\}$	0.14286
7	$\{1, 2, 3\}$	$\{5, 6\}$	$\{8, 9\}$	0.21199
8	$\{1, 2, 3\}$	$\{5, 6, 7\}$	$\{8, 9\}$	0.30727
9	$\{1, 2, 3, 4\}$	$\{5, 6, 7\}$	$\{8, 9\}$	0.41202
10	$\{1, 2, 3, 4\}$	$\{5, 6, 7\}$	$\{8, 9, 10\}$	0.58925

Table 8: Performance evaluation of the generalized distorted greedy algorithm (B-L model).



Figure 4: Distance to independence against subset size for the three algorithms (B-L model).

	Greedy	Distorted Greedy		
m	Subset S_m	$\left \mathbb{I}(P^{(S_m)}) \right $	Subset S_m	$\mathbb{I}(P^{(S_m)})$
2	$\{4, 10\}$	0.00757	$\{1, 7\}$	0.00757
3	$\{4, 7, 10\}$	0.02350	$\{1, 6, 10\}$	0.02398
4	$\{2, 4, 7, 10\}$	0.04889	$\{1, 5, 7, 10\}$	0.04961
5	$\{2, 4, 6, 7, 10\}$	0.08592	$\{1, 3, 5, 7, 10\}$	0.08591
6	$\{2, 4, 6, 7, 8, 10\}$	0.13555	$\{1, 3, 5, 7, 8, 10\}$	0.13533
7	$\{2, 3, 4, 6, 7, 8, 10\}$	0.19989	$\{1, 3, 4, 5, 7, 8, 10\}$	0.20017
8	$\{2, 3, 4, 5, 6, 7, 8, 10\}$	0.28356	$\{1, 3, 4, 5, 6, 7, 8, 10\}$	0.28399
9	$\{2, 3, 4, 5, 6, 7, 8, 9, 10\}$	0.39102	$\{1, 3, 4, 5, 6, 7, 8, 9, 10\}$	0.39191
10	$\left\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\right\}$	0.53813	$\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$	0.53813

Table 9: Comparison of the greedy algorithm and the distorted greedy algorithm (C-W model).

m	Subset $S_{m,1}$	Subset $S_{m,2}$	Subset $S_{m,3}$	$\mathbb{I}\left(\otimes_{i=1}^{3} P^{(S_{m,i})}\right)$
4	{1}	$\{5, 7\}$	{8}	0.00778
5	$\{1, 4\}$	$\{5, 7\}$	$\{8\}$	0.01556
6	$\{1, 4\}$	$\{5, 7\}$	$\{8, 10\}$	0.02376
7	$\{1, 3, 4\}$	$\{5, 7\}$	$\{8, 10\}$	0.04172
8	$\{1, 3, 4\}$	$\{5, 6, 7\}$	$\{8, 10\}$	0.06029
9	$\{1, 3, 4\}$	$\{5, 6, 7\}$	$\{8, 9, 10\}$	0.07972
10	$\{1, 2, 3, 4\}$	$\{5, 6, 7\}$	$\{8, 9, 10\}$	0.10911

Table 10: Performance evaluation of the generalized distorted greedy algorithm (C-W model).



Figure 5: Distance to independence against subset size for the three algorithms (C-W model).

For the Curie-Weiss model, Table 9 and Figure 5a show the distance of independence of the outputs of the greedy algorithm and the distorted greedy algorithm (Algorithm 2), in which these two algorithms output similar results. Table 10 and Figure 5b show the distance of independence of the tensorized outputs of the generalized distorted greedy algorithm (Algorithm 3).

In addition, we report the numerical experiment results related to the distance to independence of the complement set, as detailed in Section 5.1 and Section 5.3. The performance of the greedy algorithm on the two models is shown in Table 11 and Figure 6a, while the performance of the generalized distorted greedy algorithm can be seen from Table 12 and Figure 6b.

	Bernoulli-Lap	lace	Curie-Weiss		
m	Subset S_m	$\mathbb{I}(P^{(-S_m)})$	Subset S_m	$\mathbb{I}(P^{(-S_m)})$	
1	{9}	1.41226	{1}	0.39102	
2	$\{9, 10\}$	1.01576	$\{1, 10\}$	0.28314	
3	$\{8, 9, 10\}$	0.74171	$\{1, 5, 10\}$	0.19981	
4	$\{7, 8, 9, 10\}$	0.52441	$\{1, 5, 7, 10\}$	0.13517	
5	$\{6, 7, 8, 9, 10\}$	0.34753	$\{1, 3, 5, 7, 10\}$	0.08523	
6	$\{5, 6, 7, 8, 9, 10\}$	0.20793	$\{1, 3, 5, 7, 8, 10\}$	0.04845	
7	$\{4, 5, 6, 7, 8, 9, 10\}$	0.10318	$\{1, 3, 4, 5, 7, 8, 10\}$	0.02304	
8	$\{3, 4, 5, 6, 7, 8, 9, 10\}$	0.03406	$\{1, 3, 4, 5, 7, 8, 9, 10\}$	0.00736	

Table 11: Performance evaluation of greedy algorithm.

	Bernoulli-Laplace				Curie-Weiss			
m	$S_{m,1}$	$S_{m,2}$	$S_{m,3}$	$\mathbb{I}\left(\otimes_{i=1}^{3} P^{(-S_{m,i})}\right)$	$S_{m,1}$	$S_{m,2}$	$S_{m,3}$	$\mathbb{I}\left(\otimes_{i=1}^{3} P^{(-S_{m,i})}\right)$
1	Ø	Ø	$\{10\}$	0.41202	${2}$	Ø	Ø	0.07972
2	$\{4\}$	Ø	$\{10\}$	0.30727	$\{2\}$	Ø	$\{9\}$	0.06029
3	{4}	$\{7\}$	{10}	0.21198	{2}	$\{6\}$	<i>{</i> 9 <i>}</i>	0.04172
4	$\{3, 4\}$	$\{7\}$	{10}	0.14286	$\{2, 3\}$	$\{6\}$	<i>{</i> 9 <i>}</i>	0.02376
5	$\{3, 4\}$	$\{7\}$	$\{9, 10\}$	0.07999	$\{2, 3\}$	$\{6\}$	$\{9, 10\}$	0.01556
6	$\{3, 4\}$	$\{5, 7\}$	$\{9, 10\}$	0.03406	$\{1, 2, 3\}$	$\{6\}$	$\{9, 10\}$	0.00778

Table 12: Performance evaluation of the generalized distorted greedy algorithm.



Figure 6: Distance to independence of the complement set against subset size.

8.6 Experiment results of Section 6

We first report the numerical experiment results related to Algorithm 4. For both the Bernoulli-Laplace level model and the Curie-Weiss model, we consider the following two configurations of the batch greedy algorithm to maximize $D(P^{(S)} || \Pi^{(S)})$ subject to the cardinality constraint m:

- Approach 1: l = m and $q_i = 1$ for $i \in [l]$;
- Approach 2: $l = \lfloor \frac{m}{2} \rfloor$, $q_i = 2$ for $i \in \lfloor l-1 \rfloor$; $q_l = 2$ if m is even, $q_l = 1$ if m is odd.

In Approach 1, we recover the heuristic greedy algorithm since we are adding one element per iteration. We compare the performance of Approach 1 and Approach 2 for both models, and the results are shown in Table 13 and Table 14. Although the stationary distribution π of the Bernoulli-Laplace level model and the Curie-Weiss model are not of product form, we still apply the heuristic distorted greedy algorithm as in Corollary 6.5, and the results are summarized in Table 15. The comparison of these algorithms on the two models is shown in Figure 7.

From these results, one can conclude that the performance of Approach 1 is slightly better than Approach 2, and the performance of the distorted greedy algorithm is the worst among the three approaches.

	Approach	ı 1	Approach 2		
m	Subset S_l	$D(P^{(S_l)} \ \Pi^{(S_l)})$	Subset S_l	$D(P^{(S_l)} \ \Pi^{(S_l)})$	
1	{1}	0.26693	{1}	0.26693	
2	$\{1, 2\}$	0.59421	$\{1, 2\}$	0.59421	
3	$\{1, 2, 7\}$	0.98856	$\{1, 2, 7\}$	0.98856	
4	$\{1, 2, 7, 10\}$	1.47330	$\{1, 2, 4, 7\}$	1.46082	
5	$\{1, 2, 7, 9, 10\}$	2.07889	$\{1, 2, 4, 7, 10\}$	2.03226	
6	$\{1, 2, 7, 8, 9, 10\}$	2.85834	$\{1, 2, 4, 7, 9, 10\}$	2.73225	
7	$\{1, 2, 6, 7, 8, 9, 10\}$	3.70196	$\{1, 2, 4, 7, 8, 9, 10\}$	3.64286	
8	$\{1, 2, 5, 6, 7, 8, 9, 10\}$	4.69790	$\{1, 2, 4, 6, 7, 8, 9, 10\}$	4.65621	
9	$\{1, 2, 4, 5, 6, 7, 8, 9, 10\}$	5.91911	$\{1, 2, 4, 5, 6, 7, 8, 9, 10\}$	5.91911	
10	$\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$	7.56130	$\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$	7.56130	

Table 13: Comparison of different configurations of the batch greedy algorithm (B-L model).

	Approach 1	L	Approach 2		
m	Subset S_l	$D(P^{(S_l)} \ \Pi^{(S_l)})$	Subset S_l	$D(P^{(S_l)} \ \Pi^{(S_l)})$	
1	$\{6\}$	0.40245	$\{6\}$	0.40245	
2	$\{3, 6\}$	0.81082	$\{5, 6\}$	0.80739	
3	$\{3, 6, 8\}$	1.22606	$\{5, 6, 8\}$	1.22234	
4	$\{3, 4, 6, 8\}$	1.64626	$\{3, 5, 6, 8\}$	1.64615	
5	$\{3, 4, 6, 8, 9\}$	2.07613	$\{2, 3, 5, 6, 8\}$	2.07601	
6	$\{2, 3, 4, 6, 8, 9\}$	2.51741	$\{2, 3, 5, 6, 8, 9\}$	2.51771	
7	$\{2, 3, 4, 5, 6, 8, 9\}$	2.97051	$\{2, 3, 4, 5, 6, 8, 9\}$	2.97051	
8	$\{1, 2, 3, 4, 6, 8, 9\}$	3.44141	$\{2, 3, 4, 5, 6, 7, 8, 9\}$	3.44085	
9	$\{1, 2, 3, 4, 6, 8, 9, 10\}$	3.93647	$\{1, 2, 3, 4, 5, 6, 7, 8, 9\}$	3.93568	
10	$\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$	4.46975	$\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$	4.46975	

Table 14: Comparison of different configurations of the batch greedy algorithm (C-W model).

	Bernoulli-Laplace l	evel model	Curie-Weiss model		
m	Subset S_m	$D(P^{(S_m)} \ \Pi^{(S_m)})$	Subset S_m	$D(P^{(S_m)} \ \Pi^{(S_m)})$	
1	{10}	0.23219	{1}	0.39435	
2	$\{1, 10\}$	0.57719	$\{1, 10\}$	0.79669	
3	$\{1, 2, 10\}$	0.98552	$\{1, 2, 10\}$	1.20915	
4	$\{1, 2, 3, 5\}$	1.45314	$\{1, 2, 9, 10\}$	1.63086	
5	$\{1, 2, 3, 4, 5\}$	1.99871	$\{1, 2, 3, 9, 10\}$	2.06307	
6	$\{1, 2, 3, 4, 5, 6\}$	2.63821	$\{1, 2, 3, 8, 9, 10\}$	2.50704	
7	$\{1, 2, 3, 4, 5, 6, 7\}$	3.39168	$\{1, 2, 3, 4, 8, 9, 10\}$	2.96498	
8	$\{1, 2, 3, 4, 5, 6, 7, 8\}$	4.30094	$\{1, 2, 3, 4, 5, 8, 9, 10\}$	3.43971	
9	$\{1, 2, 3, 4, 5, 6, 7, 8, 10\}$	5.46950	$\{1, 2, 3, 4, 5, 6, 8, 9, 10\}$	3.93647	
10	$\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$	7.56130	$\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$	4.46975	

Table 15: Performance evaluation of the distorted greedy algorithm.



Figure 7: Distance to stationarity of the output against subset size.

We then report the numerical experiment results in Section 6.2, see Table 16 and Figure 8. Note that since the stationary distributions of the Bernoulli-Laplace level model (see Section 8.2) and the Curie-Weiss model (see Section 8.1) are not of product form, these simulations are heuristic in nature, as Corollary 6.7 does not provide a theoretical guarantee in this setting.

	Bernoulli-Laplace level model			Curie-Weiss model				
m	$S_{m,1}$	$S_{m,2}$	$S_{m,3}$	Value	$S_{m,1}$	$S_{m,2}$	$S_{m,3}$	Value
1	Ø	Ø	$\{10\}$	0.23191	{1}	Ø	Ø	0.39436
2	Ø	$\{7\}$	$\{10\}$	0.48566	{1}	Ø	$\{10\}$	0.78871
3	{4}	$\{7\}$	$\{10\}$	0.74787	{1}	$\{7\}$	$\{10\}$	1.19100
4	$\{3,4\}$	$\{7\}$	$\{10\}$	1.07820	{1}	$\{7\}$	$\{9, 10\}$	1.59492
5	$\{3,4\}$	$\{5,7\}$	$\{10\}$	1.41218	$\{1, 2\}$	$\{7\}$	$\{9, 10\}$	1.99886
6	$\{3,4\}$	$\{5,7\}$	$\{8, 10\}$	1.76157	$\{1,2\}$	$\{6,7\}$	$\{9, 10\}$	2.40381
7	$\{1, 3, 4\}$	$\{5,7\}$	$\{8, 10\}$	2.15778	$\{1,2\}$	$\{5, 6, 7\}$	$\{9, 10\}$	2.81582
8	$\{1, 3, 4\}$	$\{5, 6, 7\}$	$\{8, 10\}$	2.56632	$\{1, 2, 3\}$	$\{5, 6, 7\}$	$\{9, 10\}$	3.22828
9	$\{1, 3, 4\}$	$\{5, 6, 7\}$	$\{8, 9, 10\}$	3.02745	$\{1, 2, 3\}$	$\{5, 6, 7\}$	$\{8, 9, 10\}$	3.64075
10	$\{1, 2, 3, 4\}$	$\{5, 6, 7\}$	$\{8, 9, 10\}$	3.49326	$\{1, 2, 3, 4\}$	$\{5, 6, 7\}$	$\{8, 9, 10\}$	4.06242

Table 16: Performance evaluation of Algorithm 3. "Value" refers to $D(\bigotimes_{i=1}^{3} P^{(S_{m,i})} \| \bigotimes_{i=1}^{3} \Pi^{(S_{m,i})})$.



Figure 8: Performance evaluation of the generalized distorted greedy algorithm.

We proceed to present the numerical experiment results in Section 6.1 and Section 6.3 (see Table 17, Table 18, and Figure 9). Note that since the stationary distribution π of both models is not of product form, we do not have the $(1 - e^{-1})$ -approximation guarantee.

	Bernoulli-Laplace	level model	Curie-Weiss model		
m	Subset S_m	$D(P^{(-S_m)} \ \Pi^{(-S_m)})$	Subset S_m	$D(P^{(-S_m)} \ \Pi^{(-S_m)})$	
1	{9}	5.46950	{10}	3.93568	
2	$\{9, 10\}$	4.30094	$\{9, 10\}$	3.43908	
3	$\{8, 9, 10\}$	3.39168	$\{8, 9, 10\}$	2.96487	
4	$\{7, 8, 9, 10\}$	2.63821	$\{7, 8, 9, 10\}$	2.507645	
5	$\{6, 7, 8, 9, 10\}$	1.99871	$\{6, 7, 8, 9, 10\}$	2.06420	
6	$\{4, 6, 7, 8, 9, 10\}$	1.45314	$\{5, 6, 7, 8, 9, 10\}$	1.63242	
7	$\{3, 4, 6, 7, 8, 9, 10\}$	0.98630	$\{4, 5, 6, 7, 8, 9, 10\}$	1.21075	
8	$\{1, 3, 4, 6, 7, 8, 9, 10\}$	0.58961	$\{3, 4, 5, 6, 7, 8, 9, 10\}$	0.79828	
9	$\{1, 2, 3, 4, 6, 7, 8, 9, 10\}$	0.25830	$\{2, 3, 4, 5, 6, 7, 8, 9, 10\}$	0.39435	
10	$\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$	0.00000	$\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$	0.00000	

Table 17: Performance evaluation of the greedy algorithm.

	Bernoulli-Laplace level model			Curie-Weiss model				
m	$S_{m,1}$	$S_{m,2}$	$S_{m,3}$	Value	$S_{m,1}$	$S_{m,2}$	$S_{m,3}$	Value
1	{4}	Ø	Ø	3.02668	{4}	Ø	Ø	3.64075
2	{4}	Ø	$\{9\}$	2.56554	{4}	Ø	$\{8\}$	3.22828
3	{4}	$\{6\}$	$\{9\}$	2.15700	$\{3, 4\}$	Ø	$\{8\}$	2.81582
4	$\{1,4\}$	$\{6\}$	$\{9\}$	1.76235	$\{3,4\}$	$\{5\}$	$\{8\}$	2.40381
5	$\{1,4\}$	$\{6\}$	$\{8, 9\}$	1.41297	$\{3,4\}$	$\{5, 6\}$	$\{8\}$	1.99886
6	$\{1,4\}$	$\{5, 6\}$	$\{8, 9\}$	1.07899	$\{2, 3, 4\}$	$\{5, 6\}$	$\{8\}$	1.59492
7	$\{1, 2, 4\}$	$\{5, 6\}$	$\{8, 9\}$	0.74955	$\{2, 3, 4\}$	$\{5, 6\}$	$\{8, 9\}$	1.19099
8	$\{1, 2, 3, 4\}$	$\{5, 6\}$	$\{8, 9\}$	0.48566	$\{2, 3, 4\}$	$\{5, 6, 7\}$	$\{8, 9\}$	0.78871
9	$\{1, 2, 3, 4\}$	$\{5, 6, 7\}$	$\{8, 9\}$	0.23191	$\{2, 3, 4\}$	$\{5, 6, 7\}$	$\{8, 9, 10\}$	0.39436
10	$\{1, 2, 3, 4\}$	$\{5, 6, 7\}$	$\{8, 9, 10\}$	0.00000	$\{1, 2, 3, 4\}$	$\{5, 6, 7\}$	$\{8, 9, 10\}$	0.00000

Table 18: Performance evaluation of Algorithm 3. "Value" refers to $D(\bigotimes_{i=1}^{3} P^{(V_i \setminus S_{m,i})} \| \bigotimes_{i=1}^{3} \Pi^{(V_i \setminus S_{m,i})}).$



Figure 9: Distance to stationarity of the complement set against subset size.

8.7 Experiment results of Section 7

We perform Algorithm 4 with the following configuration: $l = \lceil \frac{m}{2} \rceil$, $q_i = 2$ for $i \in [l-1]$; $q_l = 2$ if m is even, $q_l = 1$ if m is odd. We choose the fixed subset as $W = \{1, 2, 3\}$. The performance of the batch greedy algorithm on the two models is shown in Table 19 and Figure 10.

	Bernoulli-I	Laplace level model	Curie-Weiss model		
m	Subset S_l	$D(P^{(W\cup S_l)} \ P^{(W)} \otimes P^{(S_l)})$	Subset S_l	$D(P^{(W\cup S_l)} \ P^{(W)} \otimes P^{(S_l)})$	
1	{10}	0.14671	{4}	0.02751	
2	$\{9, 10\}$	0.26354	$\{4, 10\}$	0.05651	
3	$\{8, 9, 10\}$	0.37787	$\{4, 5, 10\}$	0.08919	
4	$\{7, 8, 9, 10\}$	0.49198	$\{4, 5, 9, 10\}$	0.12616	
5	$\{6, 7, 8, 9, 10\}$	0.61908	$\{4, 5, 6, 9, 10\}$	0.17028	
6	$\{5, 6, 7, 8, 9, 10\}$	0.79889	$\{4, 5, 6, 8, 9, 10\}$	0.22527	
7	$\{4, 5, 6, 7, 8, 9, 10\}$	1.06993	$\{4, 5, 6, 7, 8, 9, 10\}$	0.30491	

Table 19: Performance evaluation of the batch greedy algorithm.



Figure 10: Performance evaluation of the batch greedy algorithm.

Acknowledgements

Michael Choi acknowledges the financial support of the project "MAPLE: Mechanistic Accelerated Prediction of Protein Secondary Structure via LangEvin Monte Carlo" with grant number 22-5715-P0001 under the NUS Faculty of Science Ministry of Education Tier 1 grant Data for Science and Science for Data collaborative scheme, project NUSREC-HPC-00001 and NUSREC-CLD-00001 for NUS HPC-AI Priority Projects for Research Program, as well as the startup funding of the National University of Singapore with grant number A-0000178-01-00.

Data availability

No data was used for the research described in the article.

Declarations

Conflict of interests The authors have no relevant financial or non-financial interests to disclose.

References

- [1] Anton Bovier and Frank Den Hollander. *Metastability: a potential-theoretic approach*, volume 351. Springer, 2016.
- [2] Michael C.H. Choi, Youjia Wang, and Geoffrey Wolfer. Geometry and factorization of multivariate Markov chains with applications to the swapping algorithm. arXiv preprint arXiv:2404.12589, 2024.
- [3] Kun Deng, Prashant G. Mehta, and Sean P. Meyn. Optimal Kullback-Leibler aggregation via spectral theory of Markov chains. *IEEE Trans. Automat. Control*, 56(12):2793–2808, 2011.
- [4] Alina Ene and Huy Nguyen. Streaming algorithm for monotone k-submodular maximization with cardinality constraints. In *International Conference on Machine Learning*, pages 5944–5967. PMLR, 2022.
- [5] Uriel Feige, Vahab S Mirrokni, and Jan Vondrák. Maximizing non-monotone submodular functions. SIAM Journal on Computing, 40(4):1133–1153, 2011.
- [6] Bernhard C. Geiger and Christoph Temmel. Lumpings of Markov chains, entropy rate preservation, and higher-order lumpability. *Journal of Applied Probability*, 51(4):1114–1132, 2014.

- [7] Chris Harshaw, Moran Feldman, Justin Ward, and Amin Karbasi. Submodular maximization beyond non-negativity: Guarantees, fast algorithms, and applications. In *Proceedings of the 36th International Conference on Machine Learning*, volume 97 of *Proceedings of Machine Learning Re*search, pages 2634–2643. PMLR, 09–15 Jun 2019.
- [8] Jayanth Jagalur-Mohan and Youssef Marzouk. Batch greedy maximization of non-submodular functions: Guarantees and applications to experimental design. Journal of Machine Learning Research, 22(252):1–62, 2021.
- [9] Kshitij Khare and Hua Zhou. Rates of convergence of some multivariate Markov chains with polynomial eigenfunctions. Ann. Appl. Probab., 19(2):737–777, 2009.
- [10] Bernhard H Korte, Jens Vygen, B Korte, and J Vygen. Combinatorial optimization, volume 1. Springer, 2011.
- [11] Jon Lee, Maxim Sviridenko, and Jan Vondrák. Submodular maximization over multiple matroids via generalized exchange properties. *Mathematics of Operations Research*, 35(4):795–806, 2010.
- [12] David A Levin and Yuval Peres. Markov chains and mixing times, volume 107. American Mathematical Soc., 2017.
- [13] G. L. Nemhauser, L. A. Wolsey, and M. L. Fisher. An analysis of approximations for maximizing submodular set functions. I. *Math. Programming*, 14(3):265–294, 1978.
- [14] Yury Polyanskiy and Yihong Wu. Information Theory: From Coding to Learning. Cambridge University Press, 2025.
- [15] Justin Ward and Stanislav Živný. Maximizing k-submodular functions and beyond. ACM Trans. Algorithms, 12(4):Art. 47, 26, 2016.