

Self-Similar Bridge between Regular and Critical Regions

Vyacheslav I. Yukalov^{1,2,a}, Elizaveta P. Yukalova³, and
Didier Sornette^{4,5}

¹*Bogolubov Laboratory of Theoretical Physics,
Joint Institute for Nuclear Research, Dubna 141980, Russia*

²*Instituto de Física de São Carlos, Universidade de São Paulo,
CP 369, São Carlos 13560-970, São Paulo, Brazil*

³*Laboratory of Information Technologies,
Joint Institute for Nuclear Research, Dubna 141980, Russia*

⁴*Institute of Risk Analysis, Prediction and Management (Risks-X),
Southern University of Science and Technology, Shenzhen, People's Republic of China*

⁵*Swiss Finance Institute, c/o University of Geneva,
40 blvd. Du Pont d'Arve, CH 1211 Geneva 4, Switzerland*

Abstract

In statistical and nonlinear systems, two qualitatively distinct parameter regions are typically identified: the regular region, characterized by smooth behavior of key quantities, and the critical region, where these quantities exhibit singularities or strong fluctuations. Due to their starkly different properties, these regions are often perceived as being weakly related, if at all. However, we demonstrate that these regions are intimately connected, a relationship that can be explicitly revealed using self-similar approximation theory. This framework enables the prediction of observable quantities near the critical point based on information from the regular region and vice versa. Remarkably, the method relies solely on asymptotic expansions with respect to a parameter, regardless of whether the expansion originates in the regular or critical region. The mathematical principles of self-similar theory remain consistent across both cases. We illustrate this connection by extrapolating from the regular region to predict the existence, location, and critical indices of a critical point of an equation of state for a statistical system, even when no direct information about the critical region is available. Conversely, we explore extrapolation from the critical to the regular region in systems with discrete scale invariance, where log-periodic oscillations in observables introduce additional complexity. Our findings provide insights and solutions applicable to diverse phenomena, including material fracture, stock market crashes, and earthquake forecasting.

^acorresponding author e-mail: yukalov@theor.jinr.ru

Keywords: Critical region, Regular region, Self-similar bridge

1 Introduction

For statistical and nonlinear systems exhibiting critical phenomena, two fundamentally distinct regions of their characteristic parameters are generally recognized. The first is the region where the system displays smooth behavior of its observable quantities. The second is the critical region, characterized by sharp variations in certain observable quantities, often accompanied by oscillations and singular points. Due to these contrasting behaviors, these regions often appear unrelated, making it seem impossible to predict the behavior in one region based on knowledge of the other.

This paper argues that the regular and critical regions are deeply interconnected. When analyzing a specific observable quantity, let us remember that it retains the same defining properties across all values of the variable or parameter in question. These properties might be prominently exhibited in one region while remaining hidden, yet still present, in the other.

Critical phenomena are marked by scale invariance, which dictates the asymptotic behavior of studied quantities in the critical region [1–6]. Typically, the behavior of observables in these regions is described by renormalization group equations, which leads to power law behavior as a specific case of continuous real-valued scaling [7–14]. However, observables far outside the critical point may retain scaling properties, albeit in a hidden form.

Similarly, critical phenomena associated with discrete scaling involve complex critical exponents, whose observable signatures manifest as log-periodic oscillations superimposed on the leading power-law behavior [5, 15]. These oscillatory features may persist beyond the critical point, though they might not be readily apparent.

The possibility of log-periodic oscillations was mentioned by Harris [16]. In a work devoted to the theory of branching processes, he suggested that iterating a discrete mapping might yield log-periodic oscillations. The significance of discrete scale invariance lies in its broad applicability across various phenomena, including phase transitions and temporal critical phenomena with finite-time singularities. Discrete scale invariance, along with its characteristic log-periodic oscillations, has been documented in numerous contexts, such as models defined on hierarchical lattices [17–23], models on fractal structures [24–30] or on aperiodic structures with fractal spectrum [31–34], earthquake physics [35, 36], material fracture [37–41], financial bubbles [42–60], and many other systems [15, 61–73]. Consequently, the derived solutions connecting regular and critical regions have wide-ranging applicability.

Technically, the bridge between the regular and critical regions is constructed using self-similar approximation theory, which has been recently reviewed in [13, 74] (see also [75, 76]). This approach allows for the extrapolation of an observable quantity from one region to another, using only an asymptotic expansion available in one region. For instance, starting with an asymptotic expansion for small values of a variable $x \rightarrow 0$, the behavior of the observable at finite and even large values of x (including $x \rightarrow \infty$) can be determined. This is achieved by extracting the self-similar properties of the expansion terms and extending this self-similarity to other regions.

From a mathematical standpoint, it makes no difference whether the initial expansion is derived in the regular or the critical region, because a suitable change of variables allows one to transition from small to finite or large values. What truly matters is the overall quality of the expansion – namely, how many terms are included and how well those terms are defined.

The paper is organized as follows. In Section 2, to strengthen the completeness and persuasiveness of this paper, we first outline the fundamental principles of self-similar ap-

proximation theory and then describe the steps taken to perform the extrapolation. Section 3 offers an explicit example that verifies the capability of self-similar approximation theory to bridge the regular and critical regions for the equation of state of a statistical system. This example illustrates that, even starting from a regular region far removed from the critical point and even without prior knowledge of its existence, the theory delivers accurate quantitative predictions for both the location of the critical point and the associated critical exponent. Section 4 presents the construction of a bridge extrapolating the process that links the critical region of a system, exhibiting discrete scale invariance, with its regular region. Finally, Section 5 outlines the conclusions.

2 Self-Similar Approximation Theory

The method we employ for extrapolating asymptotic expansions from one region of the variable to another is based on self-similar approximation theory. Naturally, there is no need to repeat the full mathematical foundation of the theory here, as it has been extensively detailed in several papers and review articles [13, 74]. However, it is useful to briefly revisit the core concepts of the theory in order that the reader would understand why the self-similar approach can successfully extrapolate asymptotic expansions and bridge different regions with seemingly so diverse behavior.

Suppose we are interested in a real function $f(x)$ of a real variable $x \geq 0$, which is not known explicitly. However, we do have access to its k -th order expansion $f_k(x)$ expressed in powers of x and valid in the region where $x \rightarrow 0$. We need to extrapolate this asymptotic expansion to the region of a finite variable x . This can be done by means of self-similar approximation theory, whose main points are sketched below. We give only the basic ideas, and the detailed mathematical description can be found in the review articles [13, 74].

1. *Implantation of control parameters.* The given asymptotic series, represented by the truncated expansion $f_k(x)$, is typically divergent. To improve its convergence properties, we introduce control parameters u to obtain an expanded form $F_k(x, u)$. These control parameters can be incorporated in various ways, including through initial conditions, variable transformations, or series transformations. This incorporation induces the convergence behavior of the series in subsequent steps.
2. *Fractal transformation technique.* An efficient method of implanting control parameters is through the fractal transform

$$F_k(x, \{n_j\}) = \prod_{j=1}^k x^{-n_j} (1 + b_j x),$$

where n_j and b_j are control parameters to be defined from training conditions.

3. *Determination of control functions.* To induce the convergence of the series $\{F_k(x, u)\}$, the control parameters, in general, have to become control functions $u_k(x)$, which are defined by optimization conditions, such as the minimization of a cost functional, or training conditions so that the sequence $\{F_k(x, u_k(x))\}$ be convergent.

4. *Construction of approximation cascade.* The transition from an expanded form F_k to the next expanded form F_{k+1} can be interpreted as the temporal motion within a functional space in discrete time represented by the approximation order k . The series $\{F_k(x, u_k(x))\}$ can be reformulated as a trajectory $\{y_k(f)\}$ of a dynamical system in discrete time, called cascade. The evolution equation of the approximation cascade, in the vicinity of a fixed point, is given by the self-similar relation $y_{k+p}(f) = y_k(y_p(f))$.
5. *Embedding of cascade into flow.* In order to pass from discrete time to continuous time, the approximation cascade can be embedded into the approximation flow, whose trajectory passes through all points of the cascade trajectory and the evolution equation satisfies the self-similar relation $y(t + t', f) = y(t, (y(t', f)))$.
6. *Integration of evolution equation.* The self-similar relation can be reformulated as a Lie differential equation, which can then be integrated and analyzed to define fixed-point solutions. These solutions serve as the effective limit $f_k^*(x)$ for the approximation sequence. The effective limit $f_k^*(x)$, representing a fixed point, is termed a self-similar approximant of order k .

Let the small-variable expansion with respect to $x > 0$ be

$$f_k(x) = \sum_{n=0}^k a_n x^n \quad (x \rightarrow 0), \quad (1)$$

where $a_0 \neq 0$. And we wish to extrapolate this expansion to finite values of x , and maybe even to large values. Self-similar approximants, derived using the aforementioned procedure, are found to have the form of the factor approximants

$$f_k^*(x) = a_0 \prod_{j=1}^{N_k} (1 + A_j x)^{n_j}, \quad (2)$$

in which

$$N_k = \begin{cases} k/2, & k = 2, 4, \dots \\ (k+1)/2, & k = 3, 4, \dots \end{cases}$$

while A_j and n_j are control parameters defined by the training conditions

$$\lim_{x \rightarrow 0} \frac{1}{n!} \frac{d^n}{dx^n} f_k^*(x) = a_n. \quad (3)$$

The function $f_k^*(x)$ possesses a critical point x_c if there exists $m \in (0, k]$ for which

$$A_m = \min_j A_j < 0 \quad \text{and} \quad n_m < 0, \quad (4)$$

so that $1 + A_m x_c = 0$, and the function $f_k^*(x)$ exhibits divergence at $x = x_c$. In the critical region, the function $f_k^*(x)$ has the form

$$f_k^*(x) \simeq \frac{C_k}{(x_c - x)^\alpha}, \quad (5)$$

with the critical point and critical exponent given by

$$x_c = \frac{1}{|A_m|}, \quad \alpha = |n_m|, \quad (6)$$

the coefficient C_k being

$$C_k = a_0 \prod_{j \neq m}^{N_k} (1 + A_j x_c)^{n_j}. \quad (7)$$

Note that, if an expansion with respect to large variables, say $z \gg 1$, is given, then it can be easily transformed to an expansion over small variables by means of the variable change $z = 1/x$.

An important case that is often met is when the available information is minimal, say the expansion is

$$f(x) \simeq a_0 + a_1 x + a_2 x^2, \quad (8)$$

with nonzero a_0 and a_1 , while $a_2 \rightarrow 0$. The optimal self-similar approximant for the given minimal information can be determined by applying the techniques described above to the truncated series (8), resulting in the self-similar factor approximant.

$$f^*(x) = a_0(1 + Ax)^n, \quad (9)$$

where n is different from 0 or 1. From the training conditions (3), we have $n = a_1/a_0 A$ and $A \rightarrow 0$. Taking into account that

$$\lim_{A \rightarrow 0} (1 + Ax)^{1/A} = e^x,$$

we obtain

$$f^*(x) = a_0 \exp\left(\frac{a_1}{a_0} x\right). \quad (10)$$

With additional expansion terms for $f_k(x)$, more refined approximations for $f_k^*(x)$ can be achieved, with higher-order terms providing greater accuracy. However, given the available minimal information, the best possible result is the extrapolation given by approximant (10).

To conclude this section, let us summarize the key insight of self-similar approximation theory. Although perturbative series and their coefficients are typically derived for a specific region of a variable, they inherently contain deeper information about the full function they approximate, since the series are generated by that very function. The mathematical techniques outlined here allow us to uncover and leverage this hidden structure within asymptotic series. This explains intuitively why self-similar approximation methods can effectively extrapolate results obtained in one domain of a variable to another, broader domain.

For practical applications of the theory to specific problems, there is no need to revisit the entire underlying methodology. Instead, the essential procedure is straightforward. Consider an expansion (1) that holds in the region of small values of the variable x , which we seek to extend to finite or even large values. The solution depends on the number of terms available in the given expansion (1). In the simplest case, where only a few terms are known, the approximant is given by (10). When more terms are available, the appropriate extrapolating form is the self-similar factor approximant (2), with parameters A_j and n_j determined by the training condition (3). This condition explicitly defines all parameters A_j and n_j . If these parameters satisfy the condition (4), then a critical point exists, leading to the approximant (5). This concise formulation encapsulates everything required for the practical implementation of the general theory.

3 Bridging the Regular and Critical Regions for Statistical Systems

In statistical systems, reliable expansions are typically available for their regular regions, far from critical points, whereas the primary interest often lies in understanding the critical behavior. As a typical example, let us consider the equation of state for a fluid of hard discs of diameter σ . The equation of state is presented by the compressibility factor

$$Z = \frac{P}{\rho k_B T} \quad \left(\rho \equiv \frac{N}{V} \right), \quad (11)$$

in which P is pressure, $\rho = N/V$ is the density, and T is the temperature. By means of perturbation theory, the compressibility factor can be found as an expansion in powers of density,

$$Z = 1 + \frac{1}{\rho} \sum_{n=2}^{\infty} B_n \rho^n \quad (\rho \rightarrow 0), \quad (12)$$

where B_n are the virial coefficients [77–83]. This expansion describes the regular region of asymptotically small density $\rho \rightarrow 0$ located far outside of the critical region that is located near the closest packing density $\rho_c \sigma^2 = 2/\sqrt{3} = 1.1547$. The principal question is whether it could be possible to predict the critical behavior of the fluid knowing only the above asymptotic expansion valid solely for the regular region far outside of the critical point of the closest packing density.

It is convenient to use dimensionless quantities expressing the compressibility factor as a function of the packing fraction

$$x \equiv \frac{1}{2} \rho B_2 = \frac{\pi}{4} \rho \sigma^2. \quad (13)$$

Then the asymptotic expansion (12) reads as

$$Z_k = 1 + \sum_{n=1}^k b_n x^n \quad (x \rightarrow 0), \quad (14)$$

where the coefficients

$$b_n = 2^n \frac{B_{n+1}}{B_2^n}$$

take the values

$$\begin{aligned} b_1 = 2, \quad b_2 = 3.1280, \quad b_3 = 4.2579, \quad b_4 = 5.3369, \quad b_5 = 6.3630, \\ b_6 = 7.3519, \quad b_7 = 8.3191, \quad b_8 = 9.2721, \quad b_9 = 10.2163. \end{aligned}$$

Summing expansion (14) by means of the self-similar approximation theory using the self-similar factor approximants [13, 74, 84–86], we find the equation of state

$$Z_k^* = \frac{C_k}{(x_c - x)^\alpha}, \quad (15)$$

Table 1: Prediction for the critical point x_c and critical exponent α , based on the self-similar summation of the asymptotic expansion at $x \rightarrow 0$ where x is defined by expression (13).

k	x_c	α
2	0.887	1.773
3	0.999	2.124
4	1.138	2.957
5	1.092	2.591
6	0.998	1.831
7	0.993	1.783
8	0.994	1.794
9	0.994	1.790

in which the coefficient C_k converges to $C = 1.54$ for large k . The critical normalized density x_c and the critical exponent α for different approximation orders k are given in Table 1. The critical point converges to $x_c = 0.99$ and the critical exponent, to $\alpha = 1.79$, which is in perfect agreement with the values obtained in numerical calculations [81].

Thus, the self-similar approximation theory not only enables us to connect the critical region to the distant regular region, but it also provides precise quantitative predictions for the location of the critical point and the value of the critical exponent.

4 Bridging the Critical and Regular Regions for Systems with Discrete Scale Invariance

For systems with discrete scale invariance, the situation is the reverse of what was described in the previous section. There are quite numerous studies for the critical region and practically no information on the regular region. It is therefore necessary to start with the critical region and try to realize an extrapolation towards the regular region.

In the critical region, accomplishing the decimation procedure for lattice spin systems [8, 10, 12], it has been shown [17, 87] that, in the vicinity of the phase transition point, the existence of discrete scale invariance leads to a specific renormalization group equation for the system free energy. The latter can be considered as a function of temperature, or of an effective coupling $q = \exp(J/T)$, where J is a strength of spin interactions and T is the temperature. Using the dimensionless relative coupling $x = 1 - q/q_c$, where $q_c = \exp(J/T_c)$, with T_c being a critical temperature, consider the decimation of scale $\lambda > 1$ and the related renormalization of x

$$\hat{R}(x) = \lambda x, \quad \hat{R}^n(x) = \lambda^n x. \quad (16)$$

The corresponding free energy per lattice site has been shown [17, 87] to satisfy the equation

$$f(x) \simeq h_0(x) + \frac{1}{\mu} f[\hat{R}(x)], \quad (17)$$

related to the discrete scale invariance in the vicinity of the critical point, where x is close to zero and $\mu > 1$ is the renormalization scale for the free energy. Equation (17) is approximately valid only near a critical point, as each step of the decimation procedure typically involves the multiplication of effective couplings. The distinction between relevant and irrelevant couplings, which reduces their number, can generally be made only in the vicinity of a critical point [7–14].

By iterating the above equation, one obtains a formal solution

$$f(x) \simeq \sum_{n=0}^{\infty} \frac{1}{\mu^n} h_0[\hat{R}^n(x)] . \quad (18)$$

The existence of discrete scale invariance is related to the occurrence of periodic or chaotic maps in the renormalization procedure [17].

It has been shown [88] that the free energy satisfying the renormalization group equations (17) and (18), connected with discrete scale invariance, can be represented as

$$f(x) \simeq h_0(x) + x^\alpha \sum_{n=0}^{\infty} A_n x^{in\omega} , \quad (19)$$

where $\alpha = \ln \mu / \ln \lambda$ and $\omega = 2\pi / \ln \lambda$. It is easy to notice that using the identity

$$\operatorname{Re} \sum_{n=0}^{\infty} A_n x^{in\omega} = \sum_{n=-\infty}^{\infty} B_n x^{in\omega} ,$$

in which $B_0 = A_0$ and $B_n = B_{-n} = A_n/2$ for $n \geq 1$, the free energy (19) can be rewritten in the form

$$f(x) \simeq h_0(x) + x^\alpha \sum_{n=-\infty}^{\infty} B_n x^{in\omega} . \quad (20)$$

The appearance of complex powers leads to the arising log-periodic oscillations, since

$$x^{i\omega} = \cos(\omega \ln x) + i \sin(\omega \ln x) .$$

In order to get complex powers, it is possible to work from the beginning in the complex plane, keeping in mind that the observable quantity is real. Assuming that, in the complex plane in the vicinity of a critical point, the observable satisfies the renormalization group equation

$$\frac{dF(x)}{d \ln x} \simeq G[F(x)] , \quad (21)$$

in which $G[F(x)]$ is the Gell-Man-Low function [7,9,10,13,89,90], one can look for a solution by expanding the Gell-Man-Low function in powers of $F(x)$, as has been done in Refs. [91–95].

In all cases, one obtains the behavior of observable quantities in the vicinity of a critical point, where discrete scale invariance is present. Here, our aim is to extrapolate the critical behavior to the regular region far outside the critical vicinity. Since there exist numerous phenomena exhibiting discrete scale invariance, we shall consider the behavior of their general typical representative denoted by a dimensionless function $f(x)$. We consider this characteristic quantity $f(x)$ as a function of a dimensionless variable x . The quantity

of interest can be a thermodynamic characteristic, an asset market price, a financial index, the intensity of acoustic emissions from a loaded material undergoing damage, or its amplitude of deformation, an electric current or some other potential precursors of impending earthquakes. Without loss of generality, the function $f(x)$ can be defined to be non-negative.

In the vicinity of a critical point x_c , the studied quantity experiences a sharp increase. Depending on the considered particular case, the dimensionless variable x can represent relative pressure, relative elongation, relative temperature, relative effective coupling, relative time, or other relative system parameters,

$$x = \left\{ 1 - \frac{p}{p_c}, 1 - \frac{l}{l_c}, 1 - \frac{T}{T_c}, 1 - \frac{q}{q_c}, 1 - \frac{t}{t_c} \right\}. \quad (22)$$

In that choice of variable, the initial point equals one, while the critical point corresponds to zero,

$$x_0 = 1, \quad x_c = 0. \quad (23)$$

This choice is convenient, because then the critical region is characterized by $x \ll 1$, so that in this region it is possible to use expansions in powers of x . Thus the introduced relative variable takes values in the interval

$$0 = x_c \leq x \leq x_0 = 1, \quad (24)$$

while the normalised physical variable, varies in the interval

$$0 = 1 - x_0 \leq 1 - x \leq 1 - x_c = 1. \quad (25)$$

An observable quantity $I(x)$ can always be normalized, say defining $f(x) = I(x)/I(0)$, so that the dimensionless form for the function of interest is positive and satisfies the normalization

$$f(0) = 1, \quad f(x) \geq 0. \quad (26)$$

Our aim is to study the general situation where, in the critical region in the vicinity of a critical point, there exists discrete scaling. Then, in the critical region, the studied function $f(x)$ can be represented as

$$f(x) \simeq 1 + g(x) \quad (x \rightarrow 0), \quad (27)$$

where $g(x)$ is a function that in the critical region enjoys discrete scale invariance. When using the scale invariance relation, representation (27) becomes equivalent to expression (17). In view of the normalization condition $f(0) = 1$, we have the boundary condition

$$g(0) = 0. \quad (28)$$

In order to find the general form of the considered function $f(x)$ in the whole interval $x \in [0, 1]$ from the knowledge of the existence in the critical region $x \rightarrow 0$ of discrete scaling, it is necessary to resort to the self-similar extrapolation procedure.

The function $g(x)$ is assumed to satisfy a general type of scale invariance yielding complex exponents [5, 15]. At the same time, there is no need to extend the consideration to the complex plane, since complex exponents naturally appear after employing Fourier transformation.

The property of discrete scale invariance, arising in the vicinity of a critical point, reads as

$$g(\lambda x) = \mu g(x) . \quad (29)$$

Here the scaling parameters $\lambda > 1$ and $\mu > 1$ are assumed to be fixed. This is contrary to continuous scaling where these parameters are arbitrary. Repeated use of the scaling relation yields

$$g(\lambda_n x) = \mu_n g(x) , \quad (30)$$

where the scaling parameters $\lambda_n = \lambda^n$ and $\mu_n = \mu^n$ compose a discrete spectrum with $n = 0, \pm 1, \pm 2, \dots$

It is easy to check that the general solution of relation (29) has the form

$$g(x) = x^\alpha y(x) , \quad (31)$$

in which, independently of n ,

$$\alpha \equiv \frac{\ln \mu}{\ln \lambda} \quad (32)$$

and the function $y(x)$ can be expressed as

$$y(x) = F\left(\frac{\ln x}{\ln \lambda}\right) , \quad (33)$$

with $F(z)$ being a periodic function of period one,

$$F(z+1) = F(z) \quad \left(z \equiv \frac{\ln x}{\ln \lambda}\right) . \quad (34)$$

A periodic function can be expanded in the Fourier series

$$F(z) = \sum_{n=-\infty}^{\infty} B_n e^{i2\pi n z} , \quad (35)$$

which, using the notation of z in (34), can be rewritten in the form

$$y(x) = \sum_{n=-\infty}^{\infty} B_n e^{in\omega \ln x} , \quad (36)$$

with the log-frequency

$$\omega = \frac{2\pi}{\ln \lambda} . \quad (37)$$

The log-frequency ω encodes the existence of the preferred discrete scaling ratio λ , such that $\ln \lambda$ is the period of the oscillations in the $\ln x$ variable. Thus, λ is not a frequency (the inverse of time) in the usual sense but the preferred scaling factor. Keeping in mind a real function $F(z)$, hence real $y(x)$, gives the equality $B_n^* = B_{-n}$.

Thus for the sought quantity (27) in the critical region, we have

$$f(x) \simeq 1 + x^\alpha y(x) \quad (x \rightarrow 0) . \quad (38)$$

This expression is valid only in the vicinity of a critical point, where $x \rightarrow 0$. Therefore it is possible to consider this expression as providing the first terms of an expansion

$$f(x) = \sum_{m=0}^{\infty} a_m x^{m\alpha} y^m(x), \quad (39)$$

where $a_0 = a_1 = 1$.

A representation in the form of infinite series (39) is not merely inconvenient for practical use, but it has several principal deficiencies. Generally, such series are asymptotic and have meaning only in the limit $x \rightarrow 0$. For a finite value of x , they may not converge and even can become negative, breaking the condition of semi-positiveness of $f(x)$. Dealing with such a series requires to define its effective sum. This can be done by resorting to self-similar approximation theory delineated in Sec. 2. In the case, when just a few terms of an expansion is available and it is necessary to guarantee semi-positiveness of the sought function, the method leads to the approximant (10). This form guarantees the best self-similar extrapolation under the minimally available information.

Note that, when more information is available, and a number of terms in an expansion is sufficiently large, it is possible to extend the form (10) by repeated iteration leading to continued exponentials [96–107], which finds its justification [108] in self-similar approximation theory [13, 74]. This type of exponential summation has been successfully used in several problems [13, 40, 41, 102, 108–113].

Employing the self-similar approximant (10), the effective sum (39) can be represented by the expression

$$f(x) = \exp\{x^\alpha y(x)\}. \quad (40)$$

As is mentioned above, the used approximation is based on the self-similar approximation theory that enjoys a sound mathematical foundation and has been demonstrated to provide accurate approximations for a great number of examples.

The coefficients B_n in expansion (35) are proportional to $\mu^{-n} < 1$, which allows us to treat B_n as decreasing with n (see [17, 88]). Keeping the first three terms, we have

$$F(z) = B_0 + B_1 e^{i2\pi z} + B_1^* e^{-i2\pi z}, \quad (41)$$

which yields

$$y(x) = a + b \cos(\omega \ln x) - c \sin(\omega \ln x), \quad (42)$$

where

$$a = B_0, \quad b = 2\text{Re}B_1, \quad c = 2\text{Im}B_1.$$

Equivalently, this can be represented as

$$y(x) = a + b \cos(\omega \ln x + \varphi), \quad (43)$$

with

$$\varphi = \arctan \frac{c}{b} = \arctan \frac{\text{Im}B_1}{\text{Re}B_1}.$$

In this way, the extrapolation of the studied observable quantity from the critical region, where $x \rightarrow 0$, to arbitrary values of x reads as

$$f(x) = \exp\{x^\alpha [a + b \cos(\omega \ln x + \varphi)]\}. \quad (44)$$

Note that setting $b \rightarrow 0$ corresponds to the case of continuous scale invariance, so that function (44) reduces to

$$f_0(x) = \exp(ax^\alpha). \quad (45)$$

The extrapolated function (44) represents the quantity of interest for all x in the domain $[0, 1]$. In terms of physical behavior, we have to consider the variable $1 - x$ for studying the behavior of the function $f(x)$ from the initial point $1 - x_0 = 0$ to the critical point $1 - x_c = 1$.

In order to analyse the influence of discrete scale invariance, it is useful to compare the full function $f(x)$ with the situation when only continuous scale invariance exists, and discrete scale invariance is absent. This corresponds to function (45). In the vicinity of the critical point, the continuous scale invariance leads to the power law in the exponential.

The typical behavior of $f(x)$, as compared to $f_0(x)$, as functions of $1 - x$, in the whole range $[0, 1]$ is demonstrated in Figs. 1 to 5. The parameters are taken close to the values describing the behaviour before material ruptures [39–41] and crashes in financial markets [43–46]. The role of the parameters entering $f(x)$ is emphasized.

Figure 1 illustrates that the parameter ω controls the number of oscillations, the larger ω , the larger the number of oscillations. The phase shift φ , as is shown in Fig. 2, moves the location of the oscillations, but does not change the overall picture qualitatively. Increasing the parameter α smooths the log-periodic oscillations, and makes $f(x)$ larger, as is clear from Fig. 3. Decreasing parameter a diminishes the oscillation amplitude, as in Fig. 4. The absolute value of b influences the oscillation amplitude, making it larger by increasing $|b|$, as is evident from Fig. 5.

The function $F(z)$, according to its representation (35), contains many modes with different frequencies. The reduction to the simple form (41) is valid only when $1/\mu^n \ll 1$ for $n > 1$ so that the coefficients B_n quickly decrease with n and only the lowest mode is the most important.

In order to understand the influence of higher modes, it is necessary to study the behaviour of quantity (40) with the general form of the function

$$F(z) = B_0 + \sum_{n=1}^{\infty} (B_n e^{i2\pi n z} + B_n^* e^{-i2\pi n z}), \quad (46)$$

which leads to

$$y(x) = a + \sum_{n=1}^{\infty} b_n \cos(n\omega \ln x + \varphi_n), \quad (47)$$

with $b_n = b/\mu^{n-1}$.

Note that in Ref. [88] it is shown that two cases can happen, one is when the coefficients b_n decrease exponentially as a function of n , so that only a few lower modes survive, and the other case, when the coefficients b_n decrease slower, so that many modes are to be taken into account. These two cases correspond to very weak and strong log-periodicity, respectively. The form $b_n = b/\mu^{n-1}$, depending on the value of μ , can characterize these two cases. For $\mu > 1$, the coefficients decrease exponentially and one can limit the consideration to a few first modes, while for μ close to one a number of modes needs to be taken into account, as is shown below.

Figure 6 shows the typical behaviour of $f(x)$, when the number N of modes increases until the function behaviour ceases to change. The phases of all modes are assumed to

be equal, $\varphi_n = \varphi$. First, increasing the number of modes makes the oscillations more complex. When this number becomes close to $N = 25$, a type of mode condensation occurs, characterized by shapes resembling deep troughs. After $N = 70$, the behaviour of $f(x)$ practically does not change. Summarizing the information in Fig. 6, we see that: The log-periodic oscillations become more and more complex with increasing the number of modes N . A mode condensation effect starts as the number of modes reaches $N = 25$. Beyond $N = 70$, the function stabilizes and does not change significantly, indicating that higher modes do not essentially alter the function.

Varying the phase φ can result in the appearance of high peaks before the abrupt falls, as in Fig. 7. The phase shift affects the alignment of oscillations. Overall, phase shift does not change much the log-periodic structure of oscillations merely relocating the oscillation peaks and slightly smoothing the latter. These variations may be useful in tuning models for specific applications, e.g., for the prediction of financial crashes or material rupture.

Decreasing the absolute value of negative a makes the peaks smoother, as in Fig. 8, as compared with these in Fig. 7. Smaller negative a suppresses the peaks and troughs, compared to larger negative values. Again, as in Fig. 7, the function stabilizes at $N = 70$, which confirms that $N = 70$ is sufficient to capture full oscillation complexity.

The exponent α controls how quickly log-periodic oscillations grow. Smaller $\alpha = 0.3$ in Fig.9, as compared with $\alpha = 1$ in Fig. 8, leads to smaller amplitudes of the oscillations and slower growth, as is shown in Fig. 9. The function is stable with respect to the increasing number of modes N after $N = 70$ confirming the robustness of the self-similar approximation.

The parameter ω controls the number of log-periodic oscillations. Figure 10 is plotted for larger ω , as compared to Fig. 9, which results in more frequent oscillations. Influencing how densely spaced the oscillations are impacts how observable log-periodic patterns appear in empirical data.

Figures 11 and 12 demonstrate the influence of randomized phase effects on log-periodic oscillations. Random phases are drawn from normal distributions with different standard deviations. When phase values are randomized, oscillations become more chaotic. With increasing number of modes N oscillations become increasingly irregular. For high N , irregularity dominates, resembling rather a chaotic dynamics than a structured log-periodic pattern. Thus randomness significantly affects the structure of fluctuations, with higher standard deviations leading to less predictable and more chaotic behavior. This picture is relevant for modeling natural phenomena where oscillations are disrupted by random noise.

Concluding, these figures explore how different parameters of the model influence log-periodic oscillations. Highlighting the key points it is possible to summarize the following. Increasing the number of modes N enhances oscillation complexity. Beyond the threshold $N = 70$, further mode contributions are getting practically negligible. Varying the phase shifts φ moves the oscillations, but does not change their fundamental nature. The amplitude parameter a determines how pronounced the oscillations appear. The exponent α governs the overall growth rate of oscillations. Log-frequency ω directly influences how many oscillations appear over a given range. Phase randomness disturbs and even can destroy the standard picture of log-periodic oscillations, mimicking real-world systems where the main trends are usually superimposed by random noise.

5 Conclusion

We have considered the relation between the regular region far outside of a critical point and the critical region in the vicinity of the critical point. Despite rather different behavior of observable quantities in these regions, they are not completely disconnected, but there is a bridge between them. We put forward the idea that the connection between the regular and critical regions can be recovered on the basis of self-similarity in the variation of their asymptotic forms in these regions. It is possible, starting with an asymptotic expansion in one of the regions, to explicitly extrapolate this expansion to another region. This can be accomplished employing the self-similar approximation theory, thus creating a self-similar bridge between the regions.

From a mathematical perspective, it is important to note that it makes no difference whether the starting point is the regular region or the critical region. This is because what we need is the existence of an expansion in powers of some parameter. The theory allows for the extrapolation of any expansion, irrespectively of the nature of the parameter. Of course, it is reasonable to start from that region where there is a better defined expansion.

As an example of self-similar extrapolation from the regular to critical region, we have considered a statistical system, represented by a fluid of hard disks. We have shown that, on the basis of an expansion in the regular region, being far outside the critical region and even having no information on its existence, it is straightforward to extrapolate the equation of state to the critical point and, moreover, to predict the location of the critical point and to find the related critical exponent.

Another example we studied involves systems with discrete scale invariance. In this case, while more information is available on the system's critical behavior, the regular region remains largely unexplored. This presents the challenge of extrapolating from the critical region to areas far beyond it. This issue is common across various critical phenomena exhibiting discrete scale invariance, such as earthquakes, material fractures, stock market crashes, and other "rupture" events.

A key conclusion to highlight is that log-periodic oscillations observed near a critical point do not emerge spontaneously. Rather, they have existed well outside the critical region, though typically smaller in amplitude, less frequent, and smoother. This indicates that it is possible to identify and study log-periodic oscillations even at significant distances from the critical point. Furthermore, for certain parameter settings, these oscillations outside the critical region can exhibit quite large amplitudes. Varying the parameters can result in vastly different behaviors, providing models for a wide range of processes, including the aforementioned earthquakes, material fractures, and stock market crashes.

The examples analyzed reinforce the concept that the regular and critical regions are deeply interconnected. This relationship can be uncovered using self-similar approximation theory, which enables the construction of a self-similar bridge linking the regular and critical regions. This conclusion is broad in scope and applies to systems of various types and natures.

Statements and Declarations

Funding/Competing Interests: D.S. was partially supported by the National Natural

Science Foundation of China (Grant No. T2350710802 and No. U2039202), Shenzhen Science and Technology Innovation Commission Project (Grants No. GJHZ20210705141805017 and No. K23405006), and the Center for Computational Science and Engineering at Southern University of Science and Technology.

Financial interests: The authors have no relevant financial or non-financial interests to disclose.

Conflicts of interests: The authors declare no conflicts of interest.

Author contributions: All authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by V.I. Yukalov, E.P. Yukalova, and D. Sornette. The first draft of the manuscript was written by V.I. Yukalov and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Data Availability Statement: No Data associated in the manuscript.

References

- [1] Fisher, M.E. *The Nature of Critical Points*; University of Colorado: Boulder, 1965.
- [2] Brout, R. *Phase Transitions*; Benjamin: New York, 1965.
- [3] Stanley, H.E. *Introduction to Phase Transitions and Critical Phenomena*; Oxford University: Oxford, 1987.
- [4] Yukalov, V.I.; Shumovsky, A.S. *Lectures on Phase Transitions*; World Scientific: Singapore, 1990.
- [5] Sornette, D. *Critical Phenomena in Natural Sciences*; Springer: Berlin, 2006.
- [6] Bogolubov, N.N. *Quantum Statistical Mechanics*; World Scientific, Singapore, 2015.
- [7] Wilson, K.G.; Kogut, J. The renormalization group and the ϵ expansion. *Phys. Rep.* **1974**, *12*, 75–199.
- [8] Hu, B. Introduction to real-space renormalization group methods in critical and chaotic phenomena. *Phys. Rep.* **1982**, *91*, 233–295.
- [9] Bogolubov, N.N.; Shirkov, D.V. *Quantum Fields*; Benjamin: London, 1983.
- [10] Kadanoff, L.P. *Statistical Physics: Statics, Dynamics and Renormalization*; World Scientific: Singapore, 2000.
- [11] Ma, S.K. *Modern Theory of Critical Phenomena*; Routledge: New York, 2009.
- [12] Efrati, E.; Wang, Z.; Kolan, A.; Kadanoff, L.P. Real-space renormalization in statistical mechanics. *Rev. Mod. Phys.* **2014**, *86*, 647–669.
- [13] Yukalov, V.I. Interplay between approximation theory and renormalization group. *Phys. Part. Nucl.* **2019**, *50*, 141–209.
- [14] Dupuis, N.; Canet, L.; Eichhorn, A.; Metzner, W.; Pawłowski, J.M. Tissier, M.; Wschebor, N. The nonperturbative functional renormalization group and its applications. *Phys. Rep.* **2021**, *910*, 1–114.
- [15] Sornette, D. Discrete scale invariance and complex dimensions. *Phys. Rep.* **1998**, *297*, 239–270. (extended version at <http://xxx.lanl.gov/abs/cond-mat/9707012>)
- [16] Harris, T.E. Branching processes. *Ann. Math. Stat.*, **1948**, *19*, 474–494.
- [17] Derrida, B.; Eckmann, J.P.; Erzan, A Renormalisation groups with periodic and aperiodic orbits. *J. Phys. A* **1983**, *16*, 893–906.
- [18] Derrida, B.; De Seze, L.; Itzykson C. Fractal structure of zeros in hierarchical models. *J. Stat. Phys.* **1983**, *33*, 559–569.
- [19] Derrida, B.; Itzykson, C.; Luck, J.M. Oscillatory critical amplitudes in hierarchical models. *Comm. Math. Phys.* **1984**, *94*, 115–132.

- [20] Bessis, D.; Geronimo, J.S.; Moussa, P. Mellin transforms associated with Julia sets and physical applications. *J. Stat. Phys.* **1984**, *34*, 75–110.
- [21] Itzykson, C.; Luck, J.M. Zeroes of the partition function for statistical models on regular and hierarchical lattices. *Prog. Phys.* **1985**, *11*, 45–82.
- [22] Costin, O.; Giacomin, G. Oscillatory critical amplitudes in hierarchical models and the Harris function of branching processes. *J. Stat. Phys.* **2013**, *150*, 471–486.
- [23] Derrida, B.; Giacomin, G. Log-periodic critical amplitudes: A perturbative approach. *J. Stat. Phys.* **2014**, *154*, 286–304.
- [24] Vallejos, R.O.; Anteneodo, C. Thermodynamical fingerprints of fractal spectra. *Phys. Rev. E* **1998** *58*, 4134–4140.
- [25] Knezevic, M.; Knezevic, D. Oscillatory behavior of critical amplitudes of the Gaussian model on a hierarchical structure *Phys. Rev. E* **1999**, *60*, 3396–3398.
- [26] Lessa, J.C.; Andrade, R.F.S. Log-periodic oscillations for a uniform spin model on a fractal. *Phys. Rev. E* **2000**, *62*, 3083–3089.
- [27] Bab, M.A.; Fabricius, G.; Albano, E.V. On the occurrence of oscillatory modulations in the power law behavior of dynamic and kinetic processes in fractals. *Europhys. Lett.* **2008**, *81*, 10003.
- [28] Padilla, L.; Martin, H.O.; Iguain, J.L. Log-periodic modulation in one-dimensional random walks. *Europhys. Lett.* **2009**, *85*, 20008.
- [29] Akkermans, E.; Benichou, O.; Dunne, G.V.; Teplyaev, A.; Voituriez R. Spatial log-periodic oscillations of first-passage observables in fractals. *Phys. Rev. E* **2012**, *86*, 061125.
- [30] Dunne, G.V. Heat kernels and zeta functions on fractals. *J. Phys. A* **2012**, *45*, 374016.
- [31] Luck, J.M.; Nieuwenhuizen, T.M. A soluble quasi-crystalline magnetic model: The XY quantum spin chain. *Europhys. Lett.* **1986**, *2*, 257–266.
- [32] Karevski, D.; Turban, L. Log-periodic corrections to scaling: Exact results for aperiodic Ising quantum chains. *J. Phys. A* **1996**, *29*, 3461–3470.
- [33] Andrade, R.F.S. Detailed characterization of log-periodic oscillations for an aperiodic Ising model. *Phys. Rev. E* **2000**, *61*, 7196–7199.
- [34] Carpena, P.; Coronado, A.V.; Bernaola-Galvan, P. Thermodynamics of fractal spectra: Cantor sets and quasiperiodic sequences. *Phys. Rev. E* **2000**, *61*, 2281–2289.
- [35] Sornette, D.; Sammis, C.G. Complex critical exponents from renormalization group theory of earthquakes: Implications for earthquake predictions. *J. Phys. I France* **1995** *5*, 607–619.

- [36] Johansen, A.; Sornette, D.; Wakita, H.; Tsunogai, U.; Newman, W.I.; Saleur, H. Discrete scaling in earthquake precursory phenomena: Evidence in the Kobe earthquake, Japan. *J. Phys. I France* **1996**, *6*, 1391–1402 .
- [37] Anifrani, J.C.; Le Floc'H, C.; Sornette, D.; Souillard, B. Universal Log-periodic correction to renormalization group scaling for rupture stress prediction from acoustic emissions, *J. Phys. I France* **1995**, *5*, 631–638.
- [38] Johansen, A.; Sornette, D. Evidence of discrete scale invariance by canonical averaging. *Int. J. Mod. Phys. C* **1998**, *9*, 433–447.
- [39] Johansen, A.; Sornette, D. Critical ruptures, *Eur. Phys. J. B* **2000**, *18*, 163–181.
- [40] Moura, A.; Yukalov, V.I. Self-similar extrapolation for the law of acoustic emission before failure of heterogeneous materials. *Int. J. Fract.* **2002**, *118*, 63–68.
- [41] Yukalov, V.I.; Moura, A.; Nechad, H. Self-similar law of energy release before materials fracture. *J. Mech. Phys. Solids* **2004**, *52*, 453–465.
- [42] Sornette, D.; Johansen, A.; Bouchaud, J.-P. Stock market crashes, precursors and replicas. *J. Phys. I France* **1996**, *6*, 167–175.
- [43] Sornette, D.; Johansen, A. Large financial crashes. *Physica A* **1997**, *245*, 411–422.
- [44] Johansen, A.; Sornette, D. Modeling the stock market prior to large crashes. *Eur. Phys. J. B* **1999**, *9*, 167–174.
- [45] Johansen, A.; Sornette, D. Bubbles and anti-bubbles in Latin-American, Asian and western markets: An empirical study. *Int. J. Theor. Appl. Finance* **2001**, *4*, 853–920.
- [46] Sornette, D.; Johansen, A. Significance of log-periodic precursors to financial crashes. *Quantit. Finance* **2001**, *1*, 452–471.
- [47] Feigenbaum, J.A. A statistical analysis of log-periodic precursors to financial crashes. *Quantit. Finance* **2001**, *1*, 346–360.
- [48] Sornette, D. *Why Stock Markets Crash (Critical Events in Complex Financial Systems)*; Princeton University Press: Princeton, 2002; second print with extended preface, 2017.
- [49] Feigenbaum, J. Financial physics. *Rep. Prog. Phys.* **2003**, *66*, 1611–1649.
- [50] Clark, A. Evidence of log-periodicity in corporate bond spreads. *Physica A* **2004**, *338*, 585–595.
- [51] Bree, D.S.; Joseph, N.L. Testing for financial crashes using the log periodic power law model. *Int. Rev. Financial Anal.* **2013**, *30*, 287–297.
- [52] Fantazzini, D.; Geraskin, P. Everything you always wanted to know about log periodic power laws for bubble modelling but were afraid to ask. *Eur. J. Finance* **2013**, *19*, 366–392.

- [53] Gustavsson, M.; Leven, D.; Sjogren, H. The timing of the popping: Using the log-periodic power law model to predict the bursting of bubbles on financial markets. *Financ. Hist. Rev.* **2016**, *23*, 193–217.
- [54] Ko, B.; Song, J.W.; Chang, W. Crash forecasting in the Korean stock market based on the log-periodic structure and pattern recognition. *Physica A* **2018**, *492*, 308–323.
- [55] S. Chen; S. Zheng; H. Meersman. Testing for the burst of bubbles in dry bulk shipping market using log periodic power law model. *Maritime Busin. Rev.* **2018**, *3*, 128–144.
- [56] Jhun, J.; Palacios, P.; Weatherall, J.O. Market crashes as critical phenomena? Explanation, idealization, and universality in econophysics. *Synthese* **2018**, *195*, 4477–4505.
- [57] Dai, B.; Zhang, F.; Tarzia, D.; Ahn, K. Forecasting financial crashes: Revisit to log-periodic power law. *Complexity* **2018**, *2018*, 4237471.
- [58] Song, R.; Shu, M.; Zhu, W. The 2020 global stock market crash: Endogenous or exogenous? *Physica A* **2022**, **585**, 126425.
- [59] Shu, M.; Song, R. Detection of financial bubbles using a log-periodic power law singularity (LPPLS) model. *WIREs Comput. Statist.* **2024**, *16*, e1632.
- [60] Cepni, O.; Gupta, R.; Nel, J.; Nielsen, J. Monetary policy shocks and multi-scale positive and negative bubbles in an emerging country: The case of India. *Financ. Innov.* **2025**, *11*, 35.
- [61] Yamakov, V.; Milchev, A.; Foo, G.M.; R.B.; Pandey R.B.; Stauffer, D. Log-periodic oscillations for biased diffusion of a polymer chain in a porous medium. *Eur. Phys. J. B* **1999**, *9*, 659–667.
- [62] Sienkiewicz, J.; Fronczak, P.; Holyst, J. Log-periodic oscillations due to discrete effects in complex networks. *Phys. Rev. E* **2007**, *75*, 066102.
- [63] Faillettaz, J.; Pralong, A.; Funk, M.; Deichmann, N. Evidence of log-periodic oscillations and increasing icequake activity during the breaking-off of large ice masses. *J. Glaciol.* **2008**, *54*, 725–737.
- [64] Khamzin, A.A.; Nigmatullin, R.R.; Popov, I.I.; Zhelifonov, M.P. Log-periodic oscillations in the specific heat behaviour for self-similar Ising type spin systems. *J. Phys. Conf. Ser.* **2012**, *394*, 012008.
- [65] Bazak, B.; Barnea, N. Log-periodic oscillations in the photo response of Efimov trimers. *Few Body Syst.* **2014**, *55*, 851–856.
- [66] Thiem, S. Origin of the log-periodic oscillations in the quantum dynamics of electrons in quasiperiodic systems. *Phil. Mag.* **2015**, *95*, 1233–1243.
- [67] Rybczynski, M.; Wilk, G.; Wlodarczyk, Z. System size dependence of the log-periodic oscillations of transverse momentum spectra. *EPJ Web Conf.* **2015**, *90*, 01002.

- [68] Wang, H. et al. Discovery of log-periodic oscillations in ultraquantum topological materials. *Sci. Adv.* **2018**, *4*, NO 11.
- [69] Wang, H. et al. Log-periodic quantum magneto-oscillations and discrete-scale invariance in topological material HfTe₅. *Nat. Sci. Rev.* **2019**, *6*, 914–920.
- [70] Bhojar, P.D.; Gade, P.M. Emergence of logarithmic-periodic oscillations in contact process with topological disorder. *Phys. Rev. E* **2021**, *103*, 022115.
- [71] Banerjee, A.; Pavithran, I.; Sujith, R.I. Imprints of log-periodicity in thermoacoustic systems close to lean blowout. *Phys. Rev. E* **2023**, *107*, 024219.
- [72] Dorbath, E.; Gulzar, A.; Stock, G. Log-periodic oscillations as real-time signatures of hierarchical dynamics in proteins. *J. Chem. Phys.* **2024**, *160*, 074103.
- [73] Luck, J.M. Revisiting log-periodic oscillations. *Physica A* **2024**, *643*, 129821.
- [74] Yukalov, V.I.; Yukalova, E.P. From asymptotic series to self-similar approximants. *Physics* **2021**, *3*, 829–878.
- [75] Yukalov, V.I.; Yukalova, E.P. Self-similar extrapolation in quantum field theory. *Phys. Rev. D* **2021**, *103*, 076019.
- [76] Yukalov, V.I.; Yukalova, E.P. Strong-coupling limits induced by weak-coupling expansions. *Ann. Phys. (N.Y.)* **2024**, *467*, 169716.
- [77] Mayer, J.E.; Mayer, M. *Statistical Mechanics*; Wiley: London, 1977.
- [78] Feynman, R.P. *Statistical Mechanics*; Benjamin: London, 1982.
- [79] Hansen, J.; McDonald, I. *Theory of Simple Liquids*; Academic Press: Amsterdam, 1990.
- [80] Santos, A.; Lopez de Haro, M.; Bravo Yuste, S. An accurate and simple equation of state for hard disks. *J. Chem. Phys.* **1995**, *103*, 4622–4625.
- [81] Clisby, N.; McCoy, B.M. Ninth and tenth order virial coefficients for hard spheres in D dimensions. *J. Stat. Phys.* **2006**, *122*, 15–57.
- [82] Mulero, A.; Cachadina, I.; Solana, J.R. The equation of state of the hard-disc fluid revisited. *Mol. Phys.* **2009**, *107*, 1457–1465.
- [83] Maestre, M.A.G.; Santos, A.; Robles, M.; Lopez de Haro, M. On the relation between coefficients and the close-packing of hard disks and hard spheres. *J. Chem. Phys.* **2011**, *134*, 084502.
- [84] Yukalov, V.I.; Gluzman, S.; Sornette, D. Summation of power series by self-similar factor approximants. *Physica A* **2003**, *328*, 409–438.
- [85] Gluzman, S.; Yukalov, V.I.; Sornette, D. Self-similar factor approximants. *Phys. Rev. E* **2003**, *67*, 026109.

- [86] Yukalov, V.I.; Yukalova, E.P. Self-similar sequence transformation for critical exponents. *Phys. Lett. A* **2022**, *425*, 127899.
- [87] Niemeijer, T.; van Leeuwen, J.M. Renormalization theory for Ising-like spin systems. *Phase Transitions and Critical Phenomena*, vol. 6, pp. 425–505 Domb, C.; Green M.S. (eds.) Academic, New York, 1976.
- [88] Gluzman, S.; Sornette, D. Log-periodic route to fractal functions. *Phys. Rev. E* **2002**, *65*, 036142.
- [89] Gell-Mann, M.; Low, F.F. Quantum electrodynamics at small distances. *Phys. Rev.* **1954**, *95*, 1300–1312.
- [90] Bogolubov, N.N.; Shirkov, D.V. *Introduction to the Theory of Quantized Fields*; Wiley: New York, 1980.
- [91] Johansen, A.; Sornette, D. Financial “anti-bubbles”: Log-periodicity in gold and nikkei collapses. *Int. J. Mod. Phys. C* **1999**, *10*, 563–575.
- [92] Johansen, A.; Sornette, D. Evaluation of the quantitative prediction of a trend reversal on the Japanese stock market in 1999. *Int. J. Mod. Phys. C* **2000**, *11*, 359–364.
- [93] Sornette, D.; Zhou, W.-X. The US 2000-2002 market descent: How much longer and deeper? *Quantit. Finance* **2002**, *2*, 468–481.
- [94] Johansen, A.; Sornette, D. Finite-time singularity in the dynamics of the world population and economic indices. *Physica A* **2001**, *294*, 465–502.
- [95] Zhou, W.-X.; Sornette, D. Testing the stability of the 2000-2003 US stock market “antibubble”. *Physica A* **2005**, *348*, 428–452.
- [96] Euler, L. De formulis exponentialibus replicatis. *Acta Acad. Petropolitanae* **1777**, *1*, 38–60.
- [97] Barrow, D.F. Infinite Exponentials *Am. Math. Monthly* —bf 1936, *43*, 150–160.
- [98] Bell, E.T. The iterated exponential integers. *Ann. Math.* **1938**, *39*, 539–557.
- [99] Knoebel, R.A. Exponentials reiterated. *Am. Math. Monthly* **1981**, *88*, 235–252.
- [100] Rippon, P.J. Infinite exponentials. *Math. Gazette* **1983**, *67*, 189–196.
- [101] Bromer, N. Superexponentiation. *Math. Mag.* **1987**, *60*, 169–174.
- [102] Bender, C.M.; Vinson, J.P. Summation of power series by continued exponentials. *J. Math. Phys.* **1996**, *37*, 4103–4120.
- [103] Anderson, J. Iterated exponentials. *Am. Math. Monthly* **2004**, *111*, 668–679.
- [104] Devaney, R.L.; Josic, K.; Moreno Rocha, M.; Seal, P.; Shapiro, Y.; Frumosu, A.T. Playing catch-up with iterated exponentials. *Am. Math. Monthly* **2004**, *111*, 704–709.

- [105] Hooshmand, M.H. Ultra power and ultra exponential functions. *Integr. Transf. Spec. Funct.* **2006**, *17*, 549–558.
- [106] Marshall, A.J.; Tan, Y. A rational number of the form a^a with a irrational. *Math. Gazette* **2012**, *96*, 106–109.
- [107] Paulsen, W. Tetration for complex bases. *Adv. Comput. Math.* **2018**, *45*, 243–267.
- [108] Yukalov, V.I.; Gluzman, S. Self-similar exponential approximants. *Phys. Rev. E* **1998**, *58*, 1359–1382.
- [109] Yukalov, V.I.; Yukalova, E.P. Self-similar structures and fractal transforms in approximation theory. *Chaos Solit. Fract.* **2002**, *14*, 839–861.
- [110] Gluzman, S.; Sornette, D.; Yukalov, V.I. Reconstructing generalized exponential laws by self-similar exponential approximants. *Int. J. Mod. Phys. C* **2003**, *14*, 509–527.
- [111] Abhignan, V.; Sankaranarayanan, R. Continued functions and perturbation series: Simple tools for convergence of diverging series in $O(n)$ -symmetric φ^4 field theory at weak coupling limit. *J. Stat. Phys.* **2021**, *183*, 4.
- [112] Abhignan, V.; Sankaranarayanan, R. Continued functions and Borel-Leroy transformation: Resummation of six-loop ϵ -expansions from different universality classes. *J. Phys. A* **2021**, *54*, 425401.
- [113] Abhignan, V.; Sankaranarayanan, R. Continued functions and critical exponents: Tools for analytical continuation of divergent expressions in phase transition studies. *Eur. Phys. J. B* **2023**, *96*, 31.




Figure 1: Functions $f(x)$ (solid line) and $f_0(x)$ (dashed-dotted line) for the fixed parameters $\alpha = 1$, $a = -3$, $b = -1$, $\varphi = 0.1$, and ω being varied: (a) $\omega = 10$; (b) $\omega = 50$;




Figure 2: Functions $f(x)$ (solid blue line) for $\varphi = 0.1$, $f(x)$ (dashed-dotted green line) for $\varphi = 10$, and function $f_0(x)$ (dashed red line) are presented for the fixed parameters $\alpha = 1$, $\beta = -1$, $a = -3$, $b = -1$, and $\omega = 10$. Note that $f_0(x)$ does not depend on φ .




Figure 3: Functions $f(x)$ (oscillating solid blue line) and $f_0(x)$ (monotonic solid red line) for the fixed parameters $a = -5$, $b = -2$, $\omega = 10$, $\varphi = 0.1$, and α being varied: (a) $\alpha = 1$; (b) $\alpha = 3$;




Figure 4: Functions $f(x)$ (solid blue line) and $f_0(x)$ (dashed-dotted red line) for the fixed parameters $\alpha = 1$, $b = -1$, $\omega = 25$, $\varphi = 5$, and a being varied: (a) $a = -3$; (b) $a = -5$;




Figure 5: Functions $f(x)$ (solid blue line) and $f_0(x)$ (dashed-dotted red line) for the fixed parameters $\alpha = 1$, $a = -3$, $\omega = 10$, $\varphi = 1$, and b being varied: (a) $b = -1$; (b) $b = -2$;




Figure 6: Function $f(x)$ for the fixed parameters $a = -3$, $b = -2$, $\alpha = 1$, $\omega = 10$, $\mu = 1.1$, $\varphi = 0.1$, and N varied: (a) $N = 1$; (b) $N = 5$; (c) $N = 25$; (d) $N = 70$. Note that function $f(x)$ does not change its behaviour for $N > 70$.




Figure 7: Function $f(x)$ for the fixed parameters $a = -3$, $b = -2$, $\alpha = 1$, $\omega = 10$, $\mu = 1.1$, $\varphi = 0.5$, and N varied: (a) $N = 1$; (b) $N = 5$; (c) $N = 25$; (d) $N = 70$. Note that function $f(x)$ does not change its behaviour for $N > 70$.







Figure 8: Function $f(x)$ for the fixed parameters $b = -2$, $\alpha = 1$, $\omega = 10$, $\mu = 1.1$, $\varphi = 0.1$, $a = -1$, and N varied: (a) $N = 1$; (b) $N = 5$; (c) $N = 25$; (d) $N = 70$. Note that function $f(x)$ does not change its behaviour for $N > 70$.




Figure 9: Function $f(x)$ for the fixed parameters $a = -3$, $b = -2$, $\omega = 10$, $\mu = 1.1$, $\varphi = 0.1$, $\alpha = 0.3$, and N varied: (a) $N = 1$; (b) $N = 5$; (c) $N = 25$; (d) $N = 70$. Note that function $f(x)$ does not change its behaviour for $N > 70$.




Figure 10: Function $f(x)$ for the fixed parameters $a = -3$, $b = -2$, $\mu = 1.1$, $\varphi = 0.1$, $\alpha = 1$, $\omega = 25$, and N varied: (a) $N = 1$; (b) $N = 5$; (c) $N = 25$; (d) $N = 70$. Note that function $f(x)$ does not change its behaviour for $N > 70$.




Figure 11: Function $f(x)$ for the fixed parameters $a = -3$, $b = -2$, $\omega = 25$, $\alpha = 1$, $\mu = 1.1$, and N varied: (a) $N = 1$; (b) $N = 5$; (c) $N = 25$; (d) $N = 70$; (e) $N = 100$; (f) $N = 200$. The parameters φ_n are generated as random numbers from the normal distribution with standard deviation $\sigma = 0.1$ and zero mean.




Figure 12: Function $f(x)$ for the fixed parameters $a = -3$, $b = -2$, $\omega = 25$, $\alpha = 1$, $\mu = 1.1$, and N varied: (a) $N = 1$; (b) $N = 5$; (c) $N = 25$; (d) $N = 70$; (e) $N = 100$; (f) $N = 200$. The parameters φ_n are generated as random numbers from the normal distribution with standard deviation $\sigma = 1$ and zero mean.