

THE GRONWALL INEQUALITY

RALPH HOWARD

1. INTRODUCTION.

The Gronwall inequality as given here estimates the difference of solutions to two differential equations $y'(t) = f(t, y(t))$ and $z'(t) = g(t, z(t))$ in terms of the difference between the initial conditions for the equations and the difference between f and g . The usual version of the inequality is when $f = g$, but there is little extra work involved in proving the more general case [3].

This note started life as note/homework problem on a web page for an ordinary differential equation class in 1998. Since then it has been referenced several times: [2, 4, 1, 5, 6]. So it seemed worthwhile to put a copy on the arXiv so that there is a version more permanent than one on a personal web page. While I do not know of an explicit reference to the inequality here, I very much doubt it is original to me.

2. THE INEQUALITY

Theorem 2.1 (The Gronwall Inequality). *Let \mathbf{X} be a Banach space and $U \subset \mathbf{X}$ an open convex set in \mathbf{X} . Let $f, g: [a, b] \times U \rightarrow \mathbf{X}$ be continuous functions and let $y, z: [a, b] \rightarrow U$ satisfy the initial value problems*

$$(2.1) \quad y'(t) = f(t, y(t)), \quad y(a) = y_0,$$

$$(2.2) \quad z'(t) = g(t, z(t)), \quad z(a) = z_0.$$

Also assume there is a constant $C \geq 0$ so that

$$(2.3) \quad \|g(t, x_2) - g(t, x_1)\| \leq C\|x_2 - x_1\|$$

and a continuous function $\phi: [a, b] \rightarrow [0, \infty)$ so that

$$(2.4) \quad \|f(t, y(t)) - g(t, y(t))\| \leq \phi(t).$$

Then for $t \in [a, b]$

$$(2.5) \quad \|y(t) - z(t)\| \leq e^{C|t-a|}\|y_0 - z_0\| + e^{C|t-a|} \int_a^t e^{-C|s-a|} \phi(s) ds.$$

Date: April 1, 2025.

2020 Mathematics Subject Classification. 34A12, 34A34.

Key words and phrases. Gronwall inequality.

Remark 2.2. The inequality (2.4) is a little awkward as it involves an inequality along the solution $y(t)$ which we may not know. But we can replace (2.4) by the stronger hypothesis

$$(2.6) \quad \|f(t, x) - g(t, x)\| \leq \phi(t) \quad \text{for all } x \in U$$

(which clearly implies (2.4)) and get the same result. Of course this may mean using a choice of ϕ that is larger than is needed in (2.4). \square

Remark 2.3. Note we are not assuming f satisfies a Lipschitz condition and therefore solutions to the initial value problem $y'(t) = f(t, y(t))$, $y(a) = y_0$ need not be unique. In this case the inequality can be used to estimate $y(t)$ by comparing it to the solution to $z'(t) = g(t, z(t))$, $z(a) = y_0$ with the same initial condition. In this case the inequality becomes

$$\|y(t) - z(t)\| \leq e^{C|t-a|} \int_a^t e^{-C|s-a|} \phi(s) ds.$$

Then if we assume that the inequality (2.6) holds then and $y_1'(t) = f(t, y_1(t))$ with $y_1(a) = y_1$ then using the obvious inequality $\|y(t) - y_1(t)\| \leq \|y(t) - z(t)\| + \|z(t) - y_1(t)\|$ where $z'(t) = g(t, z(t))$, $z(a) = y_0$ we get the inequality

$$\|y(t) - y_1(t)\| \leq e^{C|t-a|} \|y_0 - y_1\| + 2e^{C|t-a|} \int_a^t e^{-C|s-a|} \phi(s) ds.$$

This gives a version of the Gronwall inequality for differential equations that do not satisfy a Lipschitz condition in terms of the Lipschitz constant of a “nearby” differential equation that does (where near is “nearby” is measured by the size of ϕ). \square

Proof of Theorem 2.5. Let $x: [a, b] \rightarrow \mathbf{X}$ be continuously differentiable. Then the map $t \mapsto \|x(t)\|$ is Lipschitz and therefore absolutely continuous. Thus it is differentiable almost everywhere, the Fundamental Theorem of Calculus applies, and it is easily checked that $\frac{d}{dt} \|x'(t)\| \leq \|x'(t)\|$. Whence, setting $x(t) = y(t) - z(t)$, using the hypothesis (2.4), and (2.3),

$$\begin{aligned} \frac{d}{dt} \|y(t) - z(t)\| &\leq \|y'(t) - z'(t)\| \\ &= \|f(t, y(t)) - g(t, z(t))\| \\ &\leq \|f(t, y(t)) - g(t, y(t))\| + \|g(t, y(t)) - g(t, z(t))\| \\ &\leq \phi(t) + C\|y(t) - z(t)\|. \end{aligned}$$

That is

$$\frac{d}{dt} \|y(t) - z(t)\| - C\|y(t) - z(t)\| \leq \phi(t).$$

Multiplication by the integrating factor e^{-Ct} yields

$$\frac{d}{dt} (e^{-Ct} \|y(t) - z(t)\|) \leq e^{-Ct} \phi(t).$$

Integrate this from a to t to get

$$e^{-Ct}\|y(t) - z(t)\| - \|y_0 - z_0\| \leq \int_0^t e^{-Cs}\phi(s) ds.$$

This is equivalent to (2.5). \square

The following is the standard form of the Gronwall inequality.

Corollary 2.4. *Let \mathbf{X} be a Banach space and $U \subset \mathbf{X}$ an open convex set in \mathbf{X} . Let $f: [a, b] \times U \rightarrow \mathbf{X}$ be a continuous function and let $y, z: [a, b] \rightarrow U$ satisfy the initial value problems*

$$(2.7) \quad y'(t) = f(t, y(t)), \quad y(a) = y_0,$$

$$(2.8) \quad z'(t) = f(t, z(t)), \quad z(a) = z_0.$$

Also assume there is a constant $C \geq 0$ so that

$$(2.9) \quad \|f(t, x_2) - f(t, x_1)\| \leq C\|x_2 - x_1\|.$$

Then for $t \in [a, b]$

$$(2.10) \quad \|y(t) - z(t)\| \leq e^{C|t-a|}\|y_0 - z_0\|.$$

Proof. In Theorem 2.1 let $f = g$. Then we can take $\phi(t) \equiv 0$ in (2.4). Then (2.5) reduces to (2.10). \square

ACKNOWLEDGEMENT

I wish to thank Josh Zahl for encouraging me put this version on the arXiv.

REFERENCES

- [1] Miles Aron, Peter Bowers, Nicole Byer, Robert Decker, Aslihan Demirkaya, and Jun Hwan Ryu, *Numerical results on existence and stability of steady state solutions for the reaction-diffusion and Klein-Gordon equations*, *Involve* **7** (2014), no. 6, 723–742. MR 3284880
- [2] Jean Bourgain and Abel Klein, *Bounds on the density of states for Schrödinger operators*, *Invent. Math.* **194** (2013), no. 1, 41–72. MR 3103255
- [3] T. H. Gronwall, *Note on the derivatives with respect to a parameter of the solutions of a system of differential equations*, *Ann. of Math. (2)* **20** (1919), no. 4, 292–296. MR 1502565
- [4] Abel Klein and C. S. Sidney Tsang, *Local behavior of solutions of the stationary Schrödinger equation with singular potentials and bounds on the density of states of Schrödinger operators*, *Comm. Partial Differential Equations* **41** (2016), no. 7, 1040–1055. MR 3528526
- [5] Rui She, Qiyu Kang, Sijie Wang, Yuán-Ruì Yáng, Kai Zhao, Yang Song, and Wee Peng Tay, *Robustmat: Neural diffusion for street landmark patch matching under challenging environments*, *IEEE Transactions on Image Processing* **32** (2023), 5550–5563.
- [6] V. Srikrishnan and Subhasis Chaudhuri, *Stabilization of parametric active contours using a tangential redistribution term*, *IEEE Transactions on Image Processing* **18** (2009), no. 8, 1859–1872.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTH CAROLINA, COLUMBIA, SC
29208

Email address: howard@math.sc.edu

URL: www.math.sc.edu/~howard