

A RATIONAL COHOMOLOGY WHICH INCLUDING THAT OF A SPIN HYPERELLIPTIC MAPPING CLASS GROUP

YAN FU, GEFEI WANG

ABSTRACT. Let $\mathfrak{G} = \mathfrak{S}_q \overleftrightarrow{\times} \mathfrak{S}_q$ be the $\mathbb{Z}/2$ -extension of the product of two symmetric groups $\mathfrak{S}_q \times \mathfrak{S}_q$. In this paper, we compute the \mathfrak{G} -invariant part of the rational cohomology of the pure braid group P_n , where $n = 2q$, denoted by $H^*(P_n)^{\mathfrak{G}}$. As is known classically, $H^*(P_n)^{\mathfrak{G}}$ includes the rational cohomology of a spin hyperelliptic mapping class group, denoted by $H^*(\mathcal{S}(\Sigma_g; c))$, where $2g + 2 = n = 2q$.

1. INTRODUCTION

Let Σ_g be a compact connected oriented surface of genus g , which admits a branched 2-fold covering $\varphi : \Sigma_g \rightarrow S^2$ with $2g + 2$ branch points. The branched covering induces a group homomorphism of the Artin braid group B_{2g+2} to the mapping class group of genus g .

A 2-Spin structure c , or spin structure c for short, on the surface Σ_g can be regarded as a map

$$c : H_1(\Sigma_g, \mathbb{Z}/2) \rightarrow \mathbb{Z}/2$$

such that $c(x + y) = c(x) + c(y) + x \cdot y$ for any $x, y \in H_1(\Sigma_g, \mathbb{Z}/2)$, where $x \cdot y$ is the intersection number. The mapping class group of genus g acts on the set of spin structures in a natural way, and we call the isotropy group at a spin structure c the spin mapping class group associated with c (cf. [5]).

We want to study unstable aspects of the rational cohomology of the spin mapping class group. As first examples, we have spin hyperelliptic mapping class groups. Let $\mathcal{S}(\Sigma_g)$ be the hyperelliptic mapping class group for a compact connected oriented surface Σ_g of genus g . We can consider the spin hyperelliptic mapping class group $\mathcal{S}(\Sigma_g; c)$ as the isotropy group of $\mathcal{S}(\Sigma_g)$ at each spin structure c . The action of the group $\mathcal{S}(\Sigma_g)$ on the set of spin structures on the surface Σ_g has $\lceil \frac{g}{2} \rceil + 1$ orbits. Hence we have $\lceil \frac{g}{2} \rceil + 1$ spin hyperelliptic mapping class groups for Σ_g . The isotropy group \mathfrak{G}_i of each orbits can be described by the direct products of two symmetric groups $\mathfrak{S}_{g+1+2i} \times \mathfrak{S}_{g+1-2i}$ or a $\mathbb{Z}/2$ -extension of direct products of two symmetric groups $\mathfrak{S}_{g+1} \overleftrightarrow{\times} \mathfrak{S}_{g+1}$ (cf. e.g., [8] Proposition 3.1, p.3.95, or [11]). Now we consider the rational cohomology groups $H^*(\mathcal{S}(\Sigma_g; c))$. From [4] and by using the Lyndon-Hochschild-Serre spectral sequence, $H^*(\mathcal{S}(\Sigma_g; c))$ is isomorphic to the \mathfrak{G}_i -invariant part of the rational cohomology group of the pure sphere braid group SP_{2g+2} , denoted by $H^*(SP_{2g+2})^{\mathfrak{G}_i}$. Moreover, for any $n \geq 2$, the natural surjection of the pure Artin braid group P_n onto the pure sphere braid group SP_n induces an

2020 *Mathematics Subject Classification.* Primary 20F36; Secondary 20J06, 57K20, 14H30.

Key words and phrases. spin structures, mapping class groups, cohomology of groups.

The second named author is partially supported by National Key R&D Program of China 2020YFA0712800 and NSFC grant No.12131009.

injection of the rational cohomology groups (*cf.* e.g., [6]). Hence the \mathfrak{S}_i -invariant part of the rational cohomology group of P_{2g+2} , denoted by $H^*(P_{2g+2})^{\mathfrak{S}_i}$, includes $H^*(\mathcal{S}(\Sigma_g; c))$ as a subalgebra.

In [12], based on the result of Lehrer and Solomon in [7], the second named author gave the set of generators of $H^*(P_n)^{\mathfrak{S}}$ and calculated $H^*(P_n)^{\mathfrak{S}}$ for the cases $\mathfrak{S} = \mathfrak{S}_{n-q} \times \mathfrak{S}_q$, $n - q \geq q$. More explicitly, as a \mathbb{Q} -vector space, we have

$$H^*(P_n)^{\mathfrak{S}} \cong \bigoplus_{\lambda} \bigoplus_{\substack{(s \otimes \zeta_{\lambda}, 1) \in (Z_{\lambda})_s = 1 \\ s \in \mathfrak{S} \setminus \mathfrak{S}_n / Z_{\lambda}}} (s \otimes \zeta_{\lambda}),$$

where λ are the $(n - *)$ partitions of n , ζ_{λ} is a 1-dimensional representation of λ , Z_{λ} is the centralizer of λ , $(Z_{\lambda})_s = sZ_{\lambda}s^{-1} \cap \mathfrak{S}$ and $s \otimes \zeta_{\lambda}$ is defined by $(s \otimes \zeta_{\lambda})(g) = \zeta_{\lambda}(s^{-1}gs)$. We give a complete invariant

$$\chi : \mathfrak{S} \setminus \mathfrak{S}_n / Z_{\lambda} \rightarrow \{\emptyset\} \sqcup \left(\prod_d (\mathbb{Z}_{\geq 0})^d / \langle c_d \rangle \right)$$

for $H^*(P_n)$, where c_d is a cycle of length d . While the case $\mathfrak{S} = \mathfrak{S}_q \overleftrightarrow{\times} \mathfrak{S}_q$, $n = 2q$, is unresolved.

In this article, we calculate $H^*(P_n)^{\mathfrak{S}}$ for the case $\mathfrak{S} = \mathfrak{S}_q \overleftrightarrow{\times} \mathfrak{S}_q$, $n = 2q$. In section 2, we give some revisions on the $(\mathfrak{S}_{n-q} \times \mathfrak{S}_q)$ -invariant part of the rational cohomology of P_n , where $n - q \geq q$. In section 3, we consider the natural surjection

$$\phi : H^*(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q} \rightarrow H^*(P_n)^{\mathfrak{S}}$$

and give the kernel of ϕ . Since we have already calculated $H^*(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q}$, we get $H^*(P_n)^{\mathfrak{S}}$. In section 4, we calculate the dimension of $H^*(P_n)^{\mathfrak{S}}$, prove that

Theorem 3.1.

$$\dim(H^*(P_n)^{\mathfrak{S}}) = \frac{\dim(H^*(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q}) + |E^P|}{2} - |K^P|.$$

Where E^P and K^P are subsets of the set of generators of $H^*(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q}$. We also give a formula to calculate the cardinality $|E^P|$ and $|K^P|$.

Acknowledgments

The second named author would like to thank Nariya Kawazumi for helpful discussions. All authors contribute equally.

2. REVISIONS ON $H^*(P_n)^{\mathfrak{S}_{n-q} \times \mathfrak{S}_q}$

For a partition $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_j > 0)$ of n , let $c_{\lambda} = c_{\lambda_1} c_{\lambda_2} \dots c_{\lambda_j}$ be an element of \mathfrak{S}_n , where

$$c_{\lambda_i} = (\lambda_1 + \lambda_2 + \dots + \lambda_{i-1} + 1, \lambda_1 + \lambda_2 + \dots + \lambda_{i-1} + 2, \dots, \lambda_1 + \lambda_2 + \dots + \lambda_i)$$

is a cycle of length λ_i . Let N_{λ} be the subgroup of \mathfrak{S}_n generated by $\{\nu_i | \lambda_i = \lambda_{i+1}\}$, where

$$\begin{aligned} \nu_i &= (\lambda_1 + \lambda_2 + \dots + \lambda_{i-1} + 1, \lambda_1 + \lambda_2 + \dots + \lambda_i + 1) \\ &\quad (\lambda_1 + \lambda_2 + \dots + \lambda_{i-1} + 2, \lambda_1 + \lambda_2 + \dots + \lambda_i + 2) \dots \\ &\quad (\lambda_1 + \lambda_2 + \dots + \lambda_i, \lambda_1 + \lambda_2 + \dots + \lambda_{i+1}), \end{aligned}$$

$C_{\lambda} = \langle c_{\lambda_1} \rangle \times \langle c_{\lambda_2} \rangle \times \dots \times \langle c_{\lambda_j} \rangle$ the direct product of cyclic groups $\langle c_{\lambda_i} \rangle$. Let Z_{λ} be the centralizer of the element c_{λ} . Then the centralizer Z_{λ} of the element c_{λ} is the semi-product of N_{λ} and C_{λ} , $Z_{\lambda} = N_{\lambda} \ltimes C_{\lambda}$.

Assume that α_λ is the character of N_λ defined by $\alpha_\lambda(\nu_i) = (-1)^{\lambda_i+1}$, $\phi_\lambda = (\phi_{\lambda_1} \otimes \phi_{\lambda_2} \otimes \cdots \otimes \phi_{\lambda_j})\varepsilon$ is the character of C_λ , where $\phi_{\lambda_i}(g_i) = e^{\frac{2\pi i}{\lambda_i}}$ and ε is the sign character of \mathfrak{S}_n . From [7], the i -th cohomology group of P_n has a $\mathbb{Q}\mathfrak{S}_n$ -module isomorphism

$$H^i(P_n) \cong \bigoplus_{\lambda} \text{Ind}_{Z_\lambda}^{\mathfrak{S}_n}(\zeta_\lambda),$$

where $\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_j > 0)$ are those j partitions of n such that $i + j = n$ and $\zeta_\lambda = \alpha_\lambda \phi_\lambda$.

From the classical linear representation theory, we have

$$H^*(P_n)^{\mathfrak{S}} \cong \bigoplus_{\lambda} \bigoplus_{\substack{(s \otimes \zeta_\lambda, 1)_{(Z_\lambda)_s} = 1 \\ s \in \mathfrak{S} \setminus \mathfrak{S}_n/Z_\lambda}} (s \otimes \zeta_\lambda),$$

where $(Z_\lambda)_s = sZ_\lambda s^{-1} \cap \mathfrak{S}$ and $s \otimes \zeta_\lambda$ is defined by $(s \otimes \zeta_\lambda)(g) = \zeta_\lambda(s^{-1}gs)$.

Let $(\mathbb{Z}/2)^n$ be the set of maps $\delta : [n] \rightarrow \mathbb{Z}/2$. Then we write

$$\delta(1, 2, \dots, n) = (\delta(1), \delta(2), \dots, \delta(n)).$$

The right \mathfrak{S}_n action on $(\mathbb{Z}/2)^n$ is defined by

$$\delta \circ \sigma(1, 2, \dots, n) = (\delta(\sigma(1)), \delta(\sigma(2)), \dots, \delta(\sigma(n))),$$

where $\sigma \in \mathfrak{S}_n$.

Definition 2.1 ([12] Definition 2.4). *We define a map*

$$\{\lambda_1 + \cdots + \lambda_{i-1} + 1, \dots, \lambda_1 + \cdots + \lambda_i\} \rightarrow \mathbb{Z}/2,$$

denoted by $\delta_{\lambda_i}^{k_i}$, by $\delta_{\lambda_i}^k(a) = (\delta \circ c_{\lambda_i}^{k-1})(a)$ and call it the i -th cycle of δ . We write

$$\delta_{\lambda_1}^{k_1} \delta_{\lambda_2}^{k_2} \cdots \delta_{\lambda_j}^{k_j} = \delta \circ (c_{\lambda_1}^{k_1-1} c_{\lambda_2}^{k_2-1} \cdots c_{\lambda_j}^{k_j-1}).$$

Hence the Z_λ -orbit of δ is

$$\{\delta_{\lambda_1}^{k_1} \delta_{\lambda_2}^{k_2} \cdots \delta_{\lambda_j}^{k_j} \circ \nu \mid \nu \in N_\lambda, 1 \leq k_i \leq \lambda_i\}.$$

If $\lambda_i = \lambda_{i+1}$ for some i , then $\nu_i \in N_\lambda$ acts on $\delta_{\lambda_1}^{k_1} \delta_{\lambda_2}^{k_2} \cdots \delta_{\lambda_j}^{k_j}$ by

$$\delta_{\lambda_1}^{k_1} \cdots \delta_{\lambda_i}^{k_i} \delta_{\lambda_{i+1}}^{k_{i+1}} \cdots \delta_{\lambda_j}^{k_j} \circ \nu_i = \delta_{\lambda_1}^{k_1} \cdots \delta_{\lambda_i}^{k_i+1} \delta_{\lambda_{i+1}}^{k_i} \cdots \delta_{\lambda_j}^{k_j}.$$

Let

$$\begin{aligned} B_i(\delta) &= \{\lambda_1 + \cdots + \lambda_{i-1} + 1 \leq i_j \leq \lambda_1 + \cdots + \lambda_i \mid \delta(i_j) = 1\} \\ &\subset \{\lambda_1 + \cdots + \lambda_{i-1} + 1, \dots, \lambda_1 + \cdots + \lambda_i\}. \end{aligned}$$

We write $B_i(\delta) = \{b_1 < \cdots < b_{d_i}\}$ with $d_i = |B_i(\delta)|$. We define an element in $(\mathbb{Z}_{\geq 0})^{d_i} / \langle c_{d_i} \rangle$, denoted by $\chi_i(\delta)$, by

$$\begin{aligned} \chi_i(\delta) &= (b_2 - b_1 - 1, \dots, b_{d_i} - b_{d_i-1} - 1, \lambda_i - b_{d_i} + b_1) \\ &= (b_3 - b_2 - 1, \dots, \lambda_i - b_{d_i} + b_1, b_2 - b_1 - 1) \\ &= \cdots, \end{aligned}$$

and call it the i -th invariant cycle of δ , while $\chi_i(\delta) = (\lambda_i - 1)$ if $|B_i(\delta)| = 1$ and $\chi_i(\delta) = \emptyset$ if $B_i(\delta) = \emptyset$. We define a subset of

$$\{\emptyset\} \sqcup \left(\prod_d (\mathbb{Z}_{\geq 0})^d / \langle c_d \rangle \right),$$

denoted by $\chi(\delta)$, by

$$\chi(\delta) = \{\chi_1(\delta), \chi_2(\delta), \dots, \chi_j(\delta)\}$$

and call it the full invariant cycle set of δ . We define $I_\lambda(q)$ by

$$I_\lambda(q) = \{\chi(\delta) \mid |B_1(\delta)| + \dots + |B_j(\delta)| = q\}.$$

Now we let $\mathfrak{G} = \mathfrak{S}_q \times \mathfrak{S}_q$, $s \in \mathfrak{G} \backslash \mathfrak{S}_n / Z_\lambda$. The complete invariant

$$\chi : \mathfrak{G} \backslash \mathfrak{S}_n / Z_\lambda \rightarrow \{\emptyset\} \sqcup \left(\prod_d (\mathbb{Z}_{\geq 0})^d / \langle c_d \rangle \right)$$

is defined by $\chi(s) = \chi(\delta_s) = \{\chi_1(\delta_s), \chi_2(\delta_s), \dots, \chi_j(\delta_s)\}$, where δ_s is defined by

$$\delta_s(i) = \begin{cases} 0, & \text{if } 1 \leq s(i) \leq n - q, \\ 1, & \text{if } n - q + 1 \leq s(i) \leq n. \end{cases}$$

Theorem 2.2 ([12] Theorem 2.5). $(s \otimes \zeta_\lambda, 1)_{(Z_\lambda)_s} = 1$ if and only if $\chi(\delta_s)$ satisfies

- (1) $1 \leq |B_i(\delta_s)| \leq \lambda_i - 1$ for every $\lambda_i \geq 3$.
- (2) If $\lambda_i \geq 3$ with $\lambda_i \equiv 0, 1, 3 \pmod{4}$, then the number of minimum elements in

$$\{(a_1, a_2, \dots, a_{d_i}), (a_2, a_3, \dots, a_{d_i}, a_1), \dots\}$$

equals 1.

- (3) If $\lambda_i \geq 2$ with $\lambda_i \equiv 2 \pmod{4}$, then the number of minimum elements in

$$\{(a_1, a_2, \dots, a_{d_i}), (a_2, a_3, \dots, a_{d_i}, a_1), \dots\}$$

is at most 2.

- (4) If there exist some $\lambda_i = \lambda_{i+1} = 2k$, then $|B_i(\delta_s)| > |B_{i+1}(\delta_s)|$ or $|B_i(\delta_s)| = |B_{i+1}(\delta_s)|$ with $\chi_i(\delta_s) < \chi_{i+1}(\delta_s)$.

Let $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_a, 1, 1, \dots)$, $\lambda_i \geq 2$, $1 \leq i \leq a$, be a $(n - *)$ partition of n . We denote by $\Lambda(n - *)$ the set of all $(n - *)$ partitions of n . For a partition $\lambda \in \Lambda(n - *)$, we consider

$$d = (d_1, d_2, \dots, d_a) \in (\mathbb{Z}_{\geq 0})^a$$

and

$$(\lambda, d) = ((\lambda_1, d_1), (\lambda_2, d_2), \dots, (\lambda_a, d_a))$$

which satisfies $d_i \geq d_{i+1}$ if $\lambda_i = \lambda_{i+1}$. If d satisfies $d_1 + d_2 + \dots + d_a \leq q$, then it is easy to see that there exists some full invariant cycle sets $\chi(\delta_s)$ such that $|B_i(\delta_s)| = d_i$. We define

$$\Pi(\lambda_i, d_i) = \{\chi_i(\delta_s) \in (\mathbb{Z}_{\geq 0})^{d_i} / \langle c_{d_i} \rangle \mid |B_i(\delta_s)| = d_i, (s \otimes \phi_{\lambda_i} \varepsilon, 1)_{(Z_\lambda)_s} = 1\}$$

and denote by $|\Pi(\lambda_i, d_i)|$ the cardinality of $\Pi(\lambda_i, d_i)$.

By the definition of χ and from Theorem 2.2, if we take

$$\bigcup_\lambda \{s \otimes \zeta_\lambda \mid (s \otimes \zeta_\lambda, 1)_{(Z_\lambda)_s} = 1\}$$

as the set of generators of $H^*(P_n)^\mathfrak{G}$, then we can get each generator $s \otimes \zeta_\lambda$ by $\chi_i(\delta_s) \in \Pi(\lambda_i, d_i)$ for $1 \leq i \leq a$ and some d_i (cf. [12] Theorem 2.6 and Theorem 3.1).

3. SOME RESULTS OF $H^*(P_n)^{\mathfrak{G}}$ WHEN $\mathfrak{G} = \mathfrak{S}_q \overleftrightarrow{\times} \mathfrak{S}_q$

In this section, we consider the $\mathbb{Z}/2$ -extension of the direct product of two symmetric groups $\mathfrak{G} = \mathfrak{S}_q \overleftrightarrow{\times} \mathfrak{S}_q$, where the $\mathbb{Z}/2$ -extension is given by

$$\sigma = (1, 2q)(2, 2q-1) \cdots (q, q+1).$$

Lemma 3.1. *For an invariant cycle*

$$\chi_i(\delta_s) = (\underbrace{0, \dots, 0}_{b_1}, \underbrace{a_1, 0, \dots, 0}_{b_2}, \dots, \underbrace{0, \dots, 0}_{b_m}, a_m),$$

we have

$$\chi_i(\delta_{\sigma s}) = (b_1 + 1, \underbrace{0, \dots, 0}_{a_1-1}, b_2 + 1, \underbrace{0, \dots, 0}_{a_2-1}, \dots, b_m + 1, \underbrace{0, \dots, 0}_{a_m-1}).$$

Furthermore, it induces a bijection between $\Pi(\lambda_i, d_i)$ and $\Pi(\lambda_i, \lambda_i - d_i)$.

Proof: By the definition of σ , we have

$$\delta_{\sigma s}(i) = \delta_{\sigma}(s(i)) = \delta(2q - s(i)) = \begin{cases} 1, & \text{if } 1 \leq s(i) \leq q, \\ 0, & \text{if } q+1 \leq s(i) \leq 2q. \end{cases}$$

On the other hand, we have $(s \otimes \zeta_{\lambda}, 1)_{(Z_{\lambda})_s} = (\sigma s \otimes \zeta_{\lambda}, 1)_{(Z_{\lambda})_{\sigma s}}$. Hence $\chi_i(\delta_{\sigma s}) \in \Pi(\lambda_i, \lambda_i - d_i)$ if and only if $\chi_i(\delta_s) \in \Pi(\lambda_i, d_i)$. These prove the Lemma. \square

We write $\overline{\chi}_i(\delta_s) = \chi_i(\delta_{\sigma s})$ and call it the dual of $\chi_i(\delta_s)$. Then σ acts on the full invariant cycle set

$$\chi(\delta_s) = \{\chi_1(\delta_s), \chi_2(\delta_s), \dots, \chi_j(\delta_s)\}$$

in a natural way by

$$\sigma \circ \chi(\delta_s) = \{\overline{\chi}_1(\delta_s), \overline{\chi}_2(\delta_s), \dots, \overline{\chi}_j(\delta_s)\}.$$

We also denote $\overline{\chi}(\delta_s) = \sigma \circ \chi(\delta_s)$ and call it the dual of $\chi(\delta_s)$.

Note that $\mathfrak{S}_q \times \mathfrak{S}_q$ is a normal subgroup of \mathfrak{G} , σ acts on $H^*(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q}$ in a natural way, hence there exists a natural surjection

$$\phi : H^*(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q} \rightarrow H^*(P_n)^{\mathfrak{G}}$$

of $\mathbb{Q}\mathfrak{G}$ -modules defined by

$$\phi(s \otimes \zeta_{\lambda}) = s \otimes \zeta_{\lambda} + \sigma(s \otimes \zeta_{\lambda}),$$

where $n = 2q$. More explicitly, we have

$$\sigma(s \otimes \zeta_{\lambda}) = \varepsilon(\lambda, s)(\sigma s \otimes \zeta_{\lambda}),$$

where $\varepsilon(\lambda, s) = \pm 1$ is depending on λ and s .

We take

$$\bigcup_{\lambda} \{s \otimes \zeta_{\lambda} | (s \otimes \zeta_{\lambda}, 1)_{(Z_{\lambda})_s} = 1\}$$

as the set of generators of $H^*(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q}$. For a fixed generator $s \otimes \zeta_{\lambda}$ with a partition λ of $H^*(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q}$, since σs is uniquely determined by s , $\phi(s \otimes \zeta_{\lambda}) \neq 0$ is a generator of $H^*(P_n)^{\mathfrak{G}}$. The kernel of ϕ is generated by

$$\bigcup_{\lambda} \{s \otimes \zeta_{\lambda} - \varepsilon(\lambda, s)(\sigma s \otimes \zeta_{\lambda}) | (s \otimes \zeta_{\lambda}, 1)_{(Z_{\lambda})_s} = 1\}.$$

Recall that λ is a $(n - *)$ partition of n . We denote by $\Lambda(n - *)$ the set of all $(n - *)$ partitions of n . For a partition

$$\lambda = (\underbrace{\lambda_1, \dots, \lambda_1}_{|(\lambda_1)|}, \underbrace{\lambda_2, \dots, \lambda_2}_{|(\lambda_2)|}, \dots, \underbrace{\lambda_a, \dots, \lambda_a}_{|(\lambda_a)|}, 1, \dots, 1) \in \Lambda(n - *),$$

we write

$$\lambda_i \in \{\lambda_1, \lambda_2, \dots, \lambda_a\}$$

and

$$|(\lambda_i)| = \#\{1 \leq j \leq |\lambda_1| + \dots + |\lambda_a| \mid \lambda_j = \lambda_i\}.$$

We consider

$$d = (d_1, d_2, \dots, d_{|(\lambda_1)| + \dots + |(\lambda_a)|}) \in (\mathbb{Z}_{\geq 0})^{|(\lambda_1)| + \dots + |(\lambda_a)|}$$

and

$$(\lambda, d) = ((\lambda_1, d_1), (\lambda_2, d_2), \dots, (\lambda_{|(\lambda_1)| + \dots + |(\lambda_a)|}, d_{|(\lambda_1)| + \dots + |(\lambda_a)|}))$$

which satisfies $d_i \geq d_{i+1}$ if $\lambda_i = \lambda_{i+1}$. For a fixed λ , we denote

$$E = \left\{ (\lambda, d) \left| \begin{array}{l} \lambda \in \Lambda(n - *), d_1 + \dots + d_{|(\lambda_1)| + \dots + |(\lambda_a)|} \leq q, \\ \exists 0 \leq k_i \leq \lfloor \frac{\lambda_i}{2} \rfloor, 1 \leq i \leq a, \text{ s.t.} \\ d_{|(\lambda_1)| + \dots + |(\lambda_{i-1})| + j} = \lambda_i - d_{|(\lambda_1)| + \dots + 2k_i - j + 1}, \\ d_{|(\lambda_1)| + \dots + |(\lambda_{i-1})| + j} > \frac{\lambda_i}{2}, \\ 1 \leq i \leq a, 1 \leq j \leq k_i, \\ d_{|(\lambda_1)| + \dots + |(\lambda_{i-1})| + j} = \frac{\lambda_i}{2}, \\ 2k_i + 1 \leq j \leq |(\lambda_i)| \end{array} \right. \right\}.$$

Definition 3.2. *Let*

$$\lambda = (\underbrace{\lambda_1, \dots, \lambda_1}_{|(\lambda_1)|}, \underbrace{\lambda_2, \dots, \lambda_2}_{|(\lambda_2)|}, \dots, \underbrace{\lambda_a, \dots, \lambda_a}_{|(\lambda_a)|}, 1, \dots, 1)$$

be a partition of n . We define a subset of the set of generators of

$$\bigotimes_{|(\lambda_1)|} H^{\lambda_1 - 1}(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q} \otimes \dots \otimes \bigotimes_{|(\lambda_a)|} H^{\lambda_a - 1}(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q}$$

denoted by E^P , by

$$E^P = \bigcup_{(\lambda, d) \in E} \left\{ s \otimes \zeta_\lambda \left| \begin{array}{l} (s \otimes \zeta_\lambda, 1)_{(Z_\lambda)_s} = 1, \\ \overline{\chi_{|(\lambda_1)| + \dots + |(\lambda_{i-1})| + j}(\delta_s)} = \chi_{|(\lambda_1)| + \dots + |(\lambda_{i-1})| + 2k_i - j + 1}(\delta_s), \\ \chi_{|(\lambda_1)| + \dots + |(\lambda_{i-1})| + j}(\delta_s) < \overline{\chi_{|(\lambda_1)| + \dots + |(\lambda_{i-1})| + j}(\delta_s)}, \\ 1 \leq i \leq a, 1 \leq j \leq k_i, \\ \chi_{|(\lambda_1)| + \dots + |(\lambda_{i-1})| + j}(\delta_s) < \chi_{|(\lambda_1)| + \dots + |(\lambda_{i-1})| + j + 1}(\delta_s), \\ 1 \leq i \leq a, 1 \leq j \leq k_i - 1, \\ \overline{\chi_{|(\lambda_1)| + \dots + |(\lambda_{i-1})| + j}(\delta_s)} = \overline{\chi_{|(\lambda_1)| + \dots + |(\lambda_{i-1})| + j}(\delta_s)}, \\ \chi_{|(\lambda_1)| + \dots + |(\lambda_{i-1})| + j}(\delta_s) < \chi_{|(\lambda_1)| + \dots + |(\lambda_{i-1})| + j + 1}(\delta_s), \\ 1 \leq i \leq a, 2k_i < j \leq |(\lambda_i)| \end{array} \right. \right\}.$$

It is easy to see that if $s \otimes \zeta_\lambda \in E^P$, then $\pm s \otimes \zeta_\lambda$ is a generator of $H^*(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q}$, where the sign is depending on the lexicographic order of $\chi(\delta_s)$. Hence with respect to the sign, we also say E^P is a subset of the set of the generators of $H^*(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q}$. We will use E^P to compute $H^*(P_n)^{\mathfrak{S}}$.

Lemma 3.3. *Let $s \otimes \zeta_\lambda$ be a generator of $H^*(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q}$. $s \otimes \zeta_\lambda = \sigma s \otimes \zeta_\lambda$ if and only if $s \otimes \zeta_\lambda \in E^P$.*

Proof: Assume that $s \otimes \zeta_\lambda \in E^P$, we consider it as an element in

$$\bigotimes_{|(\lambda_1)|} H^{\lambda_1-1}(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q} \otimes \dots \otimes \bigotimes_{|(\lambda_a)|} H^{\lambda_a-1}(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q}.$$

By the definition, the invariant cycles associated with $\sigma \circ (s \otimes \zeta_\lambda) = \pm(\sigma s) \otimes \zeta_\lambda$ satisfies

$$\begin{aligned} \overline{\chi_{|(\lambda_1)|+\dots+|(\lambda_{i-1})|+j}(\delta_{\sigma s})} &= \chi_{|(\lambda_1)|+\dots+|(\lambda_{i-1})|+j}(\delta_s) \\ &= \chi_{|(\lambda_1)|+\dots+|(\lambda_{i-1})|+2k_i-j+1}(\delta_{\sigma s}) = \overline{\chi_{|(\lambda_1)|+\dots+|(\lambda_{i-1})|+2k_i-j+1}(\delta_s)}, \\ \chi_{|(\lambda_1)|+\dots+|(\lambda_{i-1})|+j}(\delta_s) &< \overline{\chi_{|(\lambda_1)|+\dots+|(\lambda_{i-1})|+j}(\delta_s)}, \\ 1 \leq i \leq a, 1 \leq j \leq k_i, \\ \chi_{|(\lambda_1)|+\dots+|(\lambda_{i-1})|+j}(\delta_s) &> \chi_{|(\lambda_1)|+\dots+|(\lambda_{i-1})|+j+1}(\delta_s), \\ 1 \leq i \leq a, k_i \leq j \leq 2k_i - 1, \\ \chi_{|(\lambda_1)|+\dots+|(\lambda_{i-1})|+j}(\delta_s) &= \overline{\chi_{|(\lambda_1)|+\dots+|(\lambda_{i-1})|+j}(\delta_s)}, \\ \chi_{|(\lambda_1)|+\dots+|(\lambda_{i-1})|+j}(\delta_s) &> \chi_{|(\lambda_1)|+\dots+|(\lambda_{i-1})|+j+1}(\delta_s), \\ 1 \leq i \leq a, 2k_i < j \leq |(\lambda_i)|. \end{aligned}$$

Hence as an element in

$$\bigotimes_{|(\lambda_1)|} H^{\lambda_1-1}(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q} \otimes \dots \otimes \bigotimes_{|(\lambda_a)|} H^{\lambda_a-1}(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q},$$

we have $\pm(\sigma s) \otimes \zeta_\lambda = s \otimes \zeta_\lambda$. Thus as a generator of $H^*(P_n)^{\mathfrak{S}_p \times \mathfrak{S}_p}$, we have $s \otimes \zeta_\lambda = \sigma s \otimes \zeta_\lambda$.

Conversely, if $s \otimes \zeta_\lambda = \sigma s \otimes \zeta_\lambda$, then we have $s \otimes \zeta_\lambda \in E^P$ obviously. \square

Lemma 3.4. *If $s \otimes \zeta_\lambda \in E^P$, then*

$$\varepsilon(\lambda, s) = \prod_{\lambda_i \in \{\lambda_1, \lambda_2, \dots, \lambda_a\}} (-1)^{\frac{|(\lambda_i)|(\lambda_i-1)(\lambda_i-2)}{2} + (\lambda_i-1)^2 k_i (2k_i-1)}.$$

Proof: By the definition, we get

$$\varepsilon(\lambda, s) = \prod_{\lambda_i \in \{\lambda_1, \lambda_2, \dots, \lambda_a\}} (-1)^{\frac{|(\lambda_i)|(\lambda_i-1)(\lambda_i-2)}{2} + (\lambda_i-1)^2 k_i (2k_i-1)}$$

by the direct computation. \square

Now consider the natural surjection

$$\phi : H^*(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q} \rightarrow H^*(P_n)^{\mathfrak{S}}.$$

Theorem 3.5. *The none trivial elements in the kernel of ϕ restricted by E^P is generated by*

$$K^P = \bigcup_{(\lambda, d) \in K} \{s \otimes \zeta_\lambda\} \subset E^P,$$

where $K \subset E$ is the set of all (λ, d) such that : there exists odd number of $1 \leq i \leq a$, such that λ_i and k_i satisfy one of:

- (1) If $k_i \equiv 0 \pmod{2}$, then $\lambda_i \equiv 0, 3 \pmod{4}$ with $|(\lambda_i)| \equiv 1, 3 \pmod{4}$.
- (2) If $k_i \equiv 1 \pmod{2}$, then $\lambda_i \equiv 2 \pmod{4}$.
- (3) If $k_i \equiv 1 \pmod{2}$, then $\lambda_i \equiv 0 \pmod{4}$ with $|(\lambda_i)| \equiv 0, 2 \pmod{4}$.
- (4) If $k_i \equiv 1 \pmod{2}$, then $\lambda_i \equiv 3 \pmod{4}$ with $|(\lambda_i)| \equiv 1, 3 \pmod{4}$.

Proof: Assume that $(s \otimes \zeta_\lambda, 1)_{(Z_\lambda)_s} = 1$. We will show that $\phi(s \otimes \zeta_\lambda) \neq 0$ when $s \otimes \zeta_\lambda \notin E^P$ at first. From Lemma 3.3, conversely, we have $s \otimes \zeta_\lambda \neq \pm \sigma s \otimes \zeta_\lambda$. Hence $\phi(s \otimes \zeta_\lambda) \neq 0$.

Now assume that $s \otimes \zeta_\lambda \notin E^P$, then from Lemma 3.4 we have

$$\varepsilon(\lambda, s) = \prod_{\lambda_i \in \{\lambda_1, \lambda_2, \dots, \lambda_a\}} (-1)^{\frac{|\lambda_i|(\lambda_i-1)(\lambda_i-2)}{2} + (\lambda_i-1)^2 k_i (2k_i-1)}.$$

We will calculate $\varepsilon(\lambda, s) = -1$. For each λ_i , we have

$$(-1)^{\frac{|\lambda_i|(\lambda_i-1)(\lambda_i-2)}{2} + (\lambda_i-1)^2 k_i (2k_i-1)} = -1$$

if and only if λ_i and k_i satisfy one of

- (1) If $k_i \equiv 0 \pmod 2$, then $\lambda_i \equiv 0, 3$ with $|\lambda_i| \equiv 1, 3 \pmod 4$.
- (2) If $k_i \equiv 1 \pmod 2$, then $\lambda_i \equiv 2 \pmod 4$.
- (3) If $k_i \equiv 1 \pmod 2$, then $\lambda_i \equiv 0 \pmod 4$ with $|\lambda_i| \equiv 0, 2 \pmod 4$.
- (4) If $k_i \equiv 1 \pmod 2$, then $\lambda_i \equiv 3 \pmod 4$ with $|\lambda_i| \equiv 1, 3 \pmod 4$.

These prove the theorem. \square

4. THE DIMENSION OF $H^*(P_n)^{\mathfrak{S}}$

For a fixed λ , we denote

$$\begin{aligned} E_1(\lambda, d) &= \{1 \leq i \leq |\lambda_1| + \dots + |\lambda_a| \mid \lambda_i \equiv 1 \pmod 2, d_i > \frac{\lambda_i}{2}\}, \\ E_{21}(\lambda, d) &= \{1 \leq i \leq |\lambda_1| + \dots + |\lambda_a| \mid \lambda_i \equiv 0 \pmod 2, d_i > \frac{\lambda_i}{2}\}, \\ E_{22}(\lambda, d) &= \{1 \leq i \leq |\lambda_1| + \dots + |\lambda_a| \mid \lambda_i \equiv 0 \pmod 2, d_i = \frac{\lambda_i}{2}\}, \\ |(\lambda_i, d_i)| &= \#\{1 \leq j \leq a \mid (\lambda_j, d_j) = (\lambda_i, d_i)\}, \\ \Pi(\lambda_i, \frac{\lambda_i}{2}) &= \left\{ \chi_i(\delta_s) \mid \begin{array}{l} (s \otimes \phi_{\lambda_i}, 1)_{(Z_\lambda)_s} = 1, \\ \chi_i(\delta_s) = \chi_i(\delta_s) \end{array} \right\}. \end{aligned}$$

We denote by $|E^P|$ and $|K^P|$ the cardinality of E^P and K^P respectively. For $1 \leq b \leq |(\lambda_i, d_i)|$, we also define simply

$$\overline{K(b)} = \{(k_1, k_2, \dots, k_b) \in (\mathbb{Z}_{>0})^b \mid k_1 + k_2 + \dots + k_b = |(\lambda_i, d_i)|\}$$

for $\overline{K(\lambda_i, d_i, b)}$.

Theorem 4.1.

$$\dim(H^*(P_n)^{\mathfrak{S}}) = \frac{\dim(H^*(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q}) + |E^P|}{2} - |K^P|.$$

Proof: By the definition of ϕ , we can calculate $\dim(H^*(P_n)^{\mathfrak{S}})$ directly. For a fixed λ , we assume that $(s \otimes \zeta_\lambda, 1)_{(Z_\lambda)_s} = 1$.

If $s \otimes \zeta_\lambda \notin E^P$, then we have $s \otimes \zeta_\lambda \neq \sigma s \otimes \zeta_\lambda$ with $\phi(s \otimes \zeta_\lambda) = \pm \phi(\sigma s \otimes \zeta_\lambda)$. The kernel of ϕ associated with this part is generated by

$$\{s \otimes \zeta_\lambda - \varepsilon(\lambda, s)(\sigma s \otimes \zeta_\lambda) \mid (s \otimes \zeta_\lambda, 1)_{(Z_\lambda)_s} = 1\}.$$

Hence the dimension of $H^*(P_n)^{\mathfrak{S}}$ associated with this part is

$$\frac{\dim(H^*(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q}) - |E^P|}{2}.$$

If $s \otimes \zeta_\lambda \in E^P$, then from Theorem 3.5, K^P is a set of generators of the kernel of ϕ associated with this part. Hence the dimension of $H^*(P_n)^{\mathfrak{S}}$ associated with this part is

$$|E^P \setminus K^P| = |E^P| - |K^P|.$$

Thus we have

$$\begin{aligned} \dim(H^*(P_n)^{\mathfrak{S}}) &= \frac{\dim(H^*(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q}) - |E^P|}{2} + |E^P| - |K^P| \\ &= \frac{\dim(H^*(P_n)^{\mathfrak{S}_q \times \mathfrak{S}_q}) + |E^P|}{2} - |K^P|. \end{aligned}$$

These prove the theorem. \square

Theorem 4.2.

$$|E^P| = \sum_{(\lambda, d) \in E} \left(\prod_{i \in E_1(\lambda, d)} \left(\sum_{b=1}^{|\lambda_i, d_i|} |K(b)| \binom{|\Pi(\lambda_i, d_i)|}{b} \right) \right. \\ \times \prod_{i \in E_{21}(\lambda, d)} \left(\frac{|\Pi(\lambda_i, d_i)|}{|(\lambda_i, d_i)|} \right) \\ \times \prod_{i \in E_{22}(\lambda, d)} \left(\frac{|\Pi(\lambda_i, \frac{\lambda_i}{2})| - |\Pi(\lambda_i, \frac{\lambda_i}{2})|}{2k_i} \right) \left(\frac{|\overline{\Pi(\lambda_i, \frac{\lambda_i}{2})}|}{|(\lambda_i, \frac{\lambda_i}{2})| - 2k_i} \right) \Bigg)$$

and

$$|K^P| = \sum_{(\lambda, d) \in K} \left(\prod_{i \in E_1(\lambda, d)} \left(\sum_{b=1}^{|\lambda_i, d_i|} |K(b)| \binom{|\Pi(\lambda_i, d_i)|}{b} \right) \right. \\ \times \prod_{i \in E_{21}(\lambda, d)} \left(\frac{|\Pi(\lambda_i, d_i)|}{|(\lambda_i, d_i)|} \right) \\ \times \prod_{i \in E_{22}(\lambda, d)} \left(\frac{|\Pi(\lambda_i, \frac{\lambda_i}{2})| - |\Pi(\lambda_i, \frac{\lambda_i}{2})|}{2k_i} \right) \left(\frac{|\overline{\Pi(\lambda_i, \frac{\lambda_i}{2})}|}{|(\lambda_i, \frac{\lambda_i}{2})| - 2k_i} \right) \Bigg).$$

Proof: Similar with the proof of Theorem 3.1 in [12], since $s \otimes \zeta_\lambda \in E^P$ is only depending on each first k_i and last $|(\lambda_i)| - 2k_i$ invariant cycles associated with each $\lambda_i \in \{\lambda_1, \dots, \lambda_a\}$ and $d_{|(\lambda_1)| + \dots + |(\lambda_{i-1})| + j}$, $1 \leq j \leq k_i$, we get these formulas immediately. \square

From Theorem 3.3 in [12], we already get the cardinality of $\Pi(\lambda_i, d_i)$. Now we compute the cardinality of $\overline{\Pi(\lambda_i, \frac{\lambda_i}{2})}$. We write $\overline{\Pi(2d, \bar{d})}$ for some $d \in \mathbb{Z}^+$ respectively.

Lemma 4.3. $\chi_i(\delta_s) = \overline{\chi_i}(\delta_s)$ if and only if there exists some divisors $k \geq 1$ of d , such that $\delta_s(j) = \delta_s(j + \frac{d}{k}) + 1$, where

$$\lambda_1 + \dots + \lambda_{i-1} + 1 \leq j \leq \lambda_1 + \dots + \lambda_{i-1} + \frac{2k-1}{k}d.$$

Moreover, $\chi_i(\delta_s) \in \overline{\Pi(2d, \bar{d})}$ if and only if $\delta_s(j) = \delta_s(j + d) + 1$ with $\delta_s(j) \neq \delta_s(j + \frac{d}{k}) + 1$ for every divisor $k \geq 2$ of d .

Proof: By the definition of σ , we have $\overline{\chi_i}(\delta_s) = \delta_{\sigma s}(j) = \delta_s(j) + 1$ for every $1 \leq j \leq 2q$.

Assume that $\chi_i(\delta_s) = \overline{\chi_i}(\delta_s)$, then there exists some divisors $k \geq 1$ of d , such that $c_{2d}^{\frac{d}{k}} \circ \delta_{\sigma s} = \delta_s$. Hence we have $\delta_s(j) = \delta_s(j + \frac{d}{k}) + 1$.

On the other hand, if there exists a divisor $k \geq 1$ of d , such that $\delta_s(j) = \delta_s(j + \frac{d}{k}) + 1$, then we have $c_{2d}^{\frac{d}{k}} \circ \delta_{\sigma s} = \delta_s$. Hence we have $\chi_i(\delta_s) = \overline{\chi_i}(\delta_s)$. Thus the first part of this Lemma follows.

Note that if $\chi_i(\delta_s)$ satisfies $\delta_s(j) = \delta_s(j + \frac{d}{k}) + 1$ for a divisor $k \geq 2$ of d , then $c_{2d}^{\frac{2d}{k}} \circ \delta_s = c_{2d}^{\frac{d}{k}} \circ \delta_{\sigma s} = \delta_s$. Since $2d \equiv 0, 2 \pmod{4}$, we have $\chi_i(\delta_s) \notin \overline{\Pi(2d, d)}$ for such $\chi_i(\delta_s)$. By the conversation, we complete our proof. \square

Theorem 4.4. *Let $d = 2^{k_0} p_1^{k_1} \cdots p_m^{k_m}$, where p_i are some odd primes for $1 \leq i \leq m$, then*

$$|\overline{\Pi(2d, d)}| = \frac{1}{2d} (2^d + \sum_{j=1}^m (-1)^j 2^{\frac{d}{p_{i_1} \cdots p_{i_j}}}).$$

Proof: For δ_s , if $\delta_s(j) = \delta_s(j + \frac{d}{2^k}) + 1$ for some $1 \leq k \leq k_0$, then we have $\delta_s(j) = \delta_s(j + d)$. Hence we don't need to consider this case.

We write

$$A_0 = \{\delta_s | \delta_s(j) = \delta_s(j+d)+1 \text{ with } \delta_s(j) \neq \delta_s(j+\frac{d}{k})+1 \text{ for every divisor } k \geq 2 \text{ of } d\}$$

and

$$A_i = \{\delta_s | \delta_s(j) = \delta_s(j + \frac{d}{p_i}) + 1\}.$$

Then from Lemma 4.3,

$$(\bigcup_{i=1}^m A_i) / \langle c_{2d} \rangle$$

is the set of all $\chi_i(\delta_s)$ such that $\chi_i(\delta_s) = \overline{\chi_i}(\delta_s)$,

$$A_0 \subset \bigcup_{i=1}^m A_i,$$

$$A_0 \cap A_i = \emptyset, 1 \leq i \leq m \text{ and } A_0 / \langle c_{2d} \rangle \simeq \overline{\Pi(2d, d)}.$$

Note that

$$|\bigcup_{i=1}^m A_i| = 2^d = |\bigcup_{i=0}^m A_i|,$$

by using the inclusion-exclusion principle, we have

$$|A_0| = |\bigcup_{i=1}^m A_i| - \sum_{j=1}^m \sum_{1 \leq i_1 \leq \cdots \leq i_j \leq m} (-1)^j |A_{i_1} \cap \cdots \cap A_{i_j}|.$$

Since

$$|A_{i_1} \cap \cdots \cap A_{i_j}| = 2^{\frac{d}{p_{i_1} \cdots p_{i_j}}},$$

we have

$$|A_0| = 2^d + \sum_{j=1}^m (-1)^j 2^{\frac{d}{p_{i_1} \cdots p_{i_j}}}.$$

Note that for every $\delta_s \in A_0$ and for every $1 \leq j \leq 2d-1$, we have $c_{2d}^j \circ \delta_s \neq \delta_s$. Hence we have

$$|\overline{\Pi(2d, d)}| = \frac{1}{2d} (2^d + \sum_{j=1}^m (-1)^j 2^{\frac{d}{p_{i_1} \cdots p_{i_j}}}),$$

the theorem follows. \square

REFERENCES

- [1] V. I. Arnold, A remark on the ramification of hyperelliptic integrals as functions of parameters, *Funct. Anal. Appl.*, **2**(1968), 187–189.
- [2] V. I. Arnold, The cohomology ring of the colored braids group, *Math. Notes*, **5**(1969), 138–140.
- [3] Joan S. Birman, *Braids, Links, and Mapping Class Groups*, Annals of Math. Studies **82**(1975).
- [4] Joan S. Birman, Hugh M. Hilden, On the mapping class groups of closed surfaces as covering spaces, *Advances in the Theory of Riemann Surfaces*, Annals of Math. Studies **66**(1971), 81–115.
- [5] D. Johnson, Spin structures and quadratic forms on surfaces, *J. London Math. Soc.*, **22**(1980), no. 2, 365–373.
- [6] N. Kawazumi, On the homotopy type of the moduli space of n -point sets of P^1 , *J. Fac. Sci. Univ. Tokyo*, **37**(1990), 263–297.
- [7] G. I. Lehrer, L. Solomon, On the action of the symmetric group on the cohomology of the complement of its reflecting hyperplanes, *J. Algebra*, **104**(1986), 410–424.
- [8] D. Mumford, *Tata Lectures on Theta*, II, Birkhäuser, Boston-Basel-Stuttgart, 1984.
- [9] O. Randal-Williams, The Picard group of the moduli space of r -Spin Riemann surfaces, *Adv. Math.*, **231**(2012), no. 1, 482–515.
- [10] O. Randal-Williams, Homology of the moduli spaces and mapping class groups of framed, r -Spin and Pin surfaces, *J. Topol.*, **7**(2014), no. 1, 155–186.
- [11] G. Wang, The artin braid group actions on the set of spin structures on a surface, *Hokkaido Math. J.*, **52**(2023), no. 3, 427–462.
- [12] G. Wang, On the rational cohomology of spin hyperelliptic mapping class groups, preprint (2024), arXiv:2401.03110

YAN FU, CHERN INSTITUTE OF MATHEMATICS, NANKAI UNIVERSITY, TIANJIN, P. R. CHINA
Email address: 1120230012@mail.nankai.edu.cn

GEFEI WANG, SCHOOL OF MATHEMATICAL SCIENCE, PEKING UNIVERSITY, BEIJING, P. R. CHINA
Email address: wanggefei@math.pku.edu.cn