



Testing for integer integration in functional time series

Won-Ki Seo* 

School of Economics
University of Sydney

Han Lin Shang 

Department of Actuarial Studies and Business Analytics
Macquarie University

Abstract

We develop a statistical testing procedure to examine whether the curve-valued time series of interest is integrated of order d for an integer $d \geq 0$. The proposed procedure can distinguish between integer-integrated time series and fractionally-integrated ones, and it has broad applicability in practice. Monte Carlo simulation experiments show that the proposed testing procedure performs reasonably well. We apply our methodology to Canadian yield curve data and French sub-national age-specific mortality data. We find evidence that these time series are mostly integrated of order one, while some have fractional orders exceeding or falling below one.

Keywords: Functional time series; fractional integration; sequential testing; yield curves; age-specific mortality rates

*The R code used in this paper is available at the author's website.

1 Introduction

Recent advances in computing and storage technologies have facilitated the presence of functional data with a graphical representation of curves (Ramsay and Silverman, 2005), images (Sørensen et al., 2013), and shapes (Srivastava and Klassen, 2016). Unlike conventional multivariate analysis, each datum in functional data analysis consists of random functions, which are realizations of a stochastic process commonly residing in a Hilbert space. In many scientific fields ranging from genomics to finance, analysing functional data has had a significant impact on statistical methods and thinking, permanently changing how we display, model and forecast high-frequency data. For an overview of functional data analysis, Ramsay and Silverman (2005) presented several state-of-the-art statistical techniques while Ferraty and Vieu (2006) listed a range of nonparametric techniques. Hsing and Eubank (2015) provided theoretical foundations with an introduction to linear operators, while Mateu and Giraldo (2022) studied spatially-dependent functional data.

Koner and Staicu (2023) presents a survey of second-generation functional data, including the analysis of temporally dependent functional data. For an overview of functional time series, Kokoszka and Reimherr (2017) introduced several state-of-the-art statistical techniques. Functional time series analysis balances functional data and time series analyses. Similar to its univariate or multivariate counterparts, a temporal dependence structure exists in functional observations. It arises in many fields, including biology (Beran et al., 2023), transportation (Klepsch et al., 2017), environmental science (Damon and Guillas, 2002), finance (Kokoszka and Zhang, 2012) and demography (Chiou and Müller, 2009). Depending on whether the continuum is also a time variable, functional time series can be classified into two groups.

For analysing functional time series, a growing body of literature has developed statistical methods and theory. The bulk of which assumes stationarity over the temporal domain, i.e., short-range dependence. These include the functional autoregressive model (Bosq, 2000; Kokoszka and Reimherr, 2013), linear regression with dependent functional variables (Seong and Seo, 2025), nonparametric functional regression (Ferraty and Vieu, 2006), functional factor model (Bathia et al., 2010), functional moving average model (Klepsch and Klüppelberg, 2017), and functional autoregressive moving average model (Klepsch et al., 2017).

On the other hand, there have been many notable developments in long memory (see, e.g., Li et al., 2020; Sabzikar and Kokoszka, 2023; Baek et al., 2024) and cointegration (see, e.g., Chang et al., 2016b, 2017) of functional time series. The long-memory curve time series describes processes with greater persistence than short-range dependent ones. In the stationary case, autocovariance decays very slowly, and the spectral density is unbounded,

typically at zero frequency. [Chang et al. \(2016b\)](#) considered cointegration for functional time series and developed statistical methods based on functional principal component analysis. [Beare et al. \(2017\)](#) and [Seo and Beare \(2019\)](#) extended the Granger-Johansen representation theorem for nonstationary functional time series taking values in a Hilbert space and a Bayes Hilbert space, respectively; more recent extensions in this area include [Beare and Seo \(2020\)](#), [Franchi and Paruolo \(2020\)](#), and [Seo \(2023\)](#).

A research gap in the literature concerning stationary curve time series is how to determine if the time series of interest is $I(0)$ (i.e., short-range dependent) or $I(d)$ for $d \in (-0.5, 0.5) \setminus \{0\}$ (i.e., fractionally-integrated). To fill this gap, we offer our solution by presenting a functional version of the KPSS-type test, which has been widely used for examining the stationarity of scalar- or vector-valued time series. A similar gap exists in the literature on nonstationary curve-valued time series, where it is often assumed that such a time series is $I(1)$ or integrated of any integer order, and hence, appropriately differenced sequences are assumed to be stationary $I(0)$ sequences. We thus extend our proposed test to assess the widespread assumption of integer integration for nonstationary time series. A useful testing procedure based on sequential applications of our proposed tests is also provided; it not only assesses the statistical plausibility of integer integration but also enables practitioners to identify an admissible range for the integration order of functional time series.

The remainder of the paper is outlined as follows. In [Section 2](#), we define notation and introduce fractionally-integrated functional time series. [Section 3](#) outlines assumptions and develops the proposed tests for examining integer integration; in [Section 3.2](#), we test the presence of short-range dependence against dependence implied by fractional integration in a stationary functional time series. Its extension to the nonstationary case is documented in [Section 3.3](#). Some extensions of the proposed tests, including a sequential testing procedure, are considered in [Section 3.4](#). In [Section 4](#) we present a series of Monte Carlo simulation studies. In [Section 5](#), we apply our hypothesis tests to Canadian yield curve data and French sub-national age-specific mortality rates. Concluding remarks are given in [Section 6](#).

2 Preliminary

2.1 Notation

We let \mathcal{H} be a Hilbert space of square-integrable functions on a compact interval, equipped with inner product $\langle \cdot, \cdot \rangle$ and norm $\|\cdot\|$. Let $\mathcal{L}_{\mathcal{H}}$ denote the space of continuous linear operators equipped with the usual operator norm $\|\cdot\|_{\mathcal{L}_{\mathcal{H}}}$. For any $A \in \mathcal{L}_{\mathcal{H}}$, $\text{ran } A$ denotes the range of

A and $\text{rank } A$ denotes the rank of A (i.e. the dimension of $\text{ran } A$). The adjoint A^* of A is the unique element of $\mathcal{L}_{\mathcal{H}}$ satisfying that $\langle Ah_1, h_2 \rangle = \langle h_1, A^*h_2 \rangle$ for all $h_1, h_2 \in \mathcal{H}$. If $A = A^*$, it is said to be self-adjoint. We say that A is nonnegative (resp. positive) if $\langle Ah, h \rangle \geq 0$ (resp. $\langle Ah, h \rangle > 0$) for all nonzero $h \in \mathcal{H}$. An element $A \in \mathcal{L}_{\mathcal{H}}$ is a compact operator if $A = \sum_{j=1}^{\infty} a_j h_{1j} \otimes h_{2j}$ for some orthonormal bases $\{h_{1j}\}_{j \geq 1}$ and $\{h_{2j}\}_{j \geq 1}$ and some sequence of real numbers $\{a_j\}_{j \geq 1}$ tending to zero, where \otimes denotes the tensor product on \mathcal{H} . If A is nonnegative, self-adjoint and compact, we may assume that $h_{1j} = h_{2j}$ and $a_j \geq 0$, and in this case $\{a_j\}_{j \geq 1}$ become the eigenvalues of A .

A \mathcal{H} -valued random element, say U , is a measurable map from the underlying probability space to \mathcal{H} . We say that U is square-integrable if $\mathbb{E}[\|U\|^2] < \infty$, and for such a random element, we may define its expectation $\mathbb{E}[U]$ (satisfying $\mathbb{E}[\langle U, h \rangle] = \langle \mathbb{E}[U], h \rangle$ for any $h \in \mathcal{H}$) and covariance operator $C_U := \mathbb{E}[(U - \mathbb{E}[U]) \otimes (U - \mathbb{E}[U])]$.

2.2 Fractionally-integrated functional time series

We consider a \mathcal{H} -valued, possibly nonstationary, time series Y_t whose d -th (fractional) difference $\Delta^d(Y_t - Y_0)$ can be made stationary I(0) time series under a suitable initial condition. Hereafter, such a time series is conveniently called a Fractionally-Integrated (FI) Hilbert-valued Time Series (HTS). The formal definition of the I(0) property in this paper is given as follows:

Definition 1. *The sequence Z_t is I(0) if the following are satisfied: (i) for each $t \geq 1$, Z_t allows the representation $Z_t = \sum_{j=0}^{\infty} A_j \eta_{t-j}$ for a mean-zero independently and identically distributed (iid) sequence η_t with covariance C_{η} and a sequence of bounded linear operator $\{A_j\}_{j \geq 1} \subset \mathcal{L}_{\mathcal{H}}$ with $\sum_{j=0}^{\infty} \|A_j\|_{\mathcal{L}_{\mathcal{H}}} < \infty$; (ii) its long-run covariance $(\sum_{j=0}^{\infty} A_j)C_{\eta}(\sum_{j=0}^{\infty} A_j)^*$ is a nonzero operator.*

The I(0) property is adopted from recent articles (e.g., [Beare et al., 2017](#); [Beare and Seo, 2020](#); [Seo, 2023](#)) concerning integer-integrated functional time series. As will be discussed in detail later, however, the meaning of $\Delta^d(Y_t - Y_0)$ being I(0) is more general and differs in the present paper. This difference arises not only because d is allowed to be a fraction, but also, more importantly, because we require only that the highest memory of the time series $\Delta^d(Y_t - Y_0)$ be zero, while allowing it to contain anti-persistent components, provided that stationarity is preserved. Specifically, there may be a projection map P such that $\Delta^{\tilde{d}}P(Y_t - Y_0)$ is I(0) for some $\tilde{d} \in (-1/2, 0]$ in our setup. This treatment is not only more realistic but also necessary for our subsequent analysis in developing a novel test for integer-order integration

of functional time series.

We are interested in examining integer-order integration hypotheses, i.e., $Y_t \sim I(d)$, where

$$H_0 : d \text{ is an integer} \quad \text{against} \quad H_1 : d \text{ is a fraction.}$$

Testing the above hypotheses is important for practitioners because it helps determine an important characteristic of functional time series and identify a suitable statistical model.

The cases with $d = 0$ and 1 may be the most empirically relevant, and thus, we will focus on these cases subsequently. Moreover, because of technical reasons, we restrict the range of admissible values of d as $d > -1/2$. To sum up, we will mainly examine if d is an integer contained in the interval $(-1/2, 2)$; however, our test to be developed can be extended to allow the case with $d \geq 2$ in an obvious way, which will be discussed in Section 3 (see Remark 3.3).

3 Testing integer integration

3.1 Assumptions

We will consider the time series of functional observations $\mathbf{Y}_0 = \{Y_t - Y_0\}_{t=1}^T$. For some nonnegative integer $\delta \geq 0$, we hereafter let $\mathbf{Y}_\delta = \{Y_{\delta,t}\}_{t=\delta+1}^T$ be the δ -th difference of \mathbf{Y}_0 , i.e.,

$$Y_{\delta,t} = \Delta^\delta(Y_t - Y_0), \quad t \geq \delta + 1, \quad (3.1)$$

where Δ^0 is understood as the identity operator. For our asymptotic analysis, we will assume that the time series \mathbf{Y}_0 satisfies some regularity conditions, which are introduced below:

Assumption F. *The time series $\mathbf{Y}_0 = \{Y_{0,t}\}_{t \geq 1}$ is a FIHTS satisfying the following properties:*

- (i) *For some $K > 0$, there exist a \mathcal{H} -valued sequence $\{X_t\}_{t \geq 1}$ and orthogonal projections P_1, \dots, P_K such that $\sum_{j=1}^K P_j = I_{\mathcal{H}}$ (the identity map on \mathcal{H}) and $Y_{0,t} = \sum_{j=1}^K P_j X_t$ for $t \geq 1$. Moreover, for each $j = 1 \dots, K$ and*

$$d_1 > d_2 > \dots > d_K > -1/2,$$

there exists a stationary $I(0)$ sequence $\{v_{j,t}\}_{t \in \mathbb{Z}}$ satisfying the following:

$$v_{1,t} = \Delta^{d_1} P_1 X_t = \sum_{k=0}^{\infty} \psi_{1,k} \varepsilon_{t-k}, \quad (3.2)$$

\vdots

$$v_{K,t} = \Delta^{d_K} P_K X_t = \sum_{k=0}^{\infty} \psi_{K,k} \varepsilon_{t-k}, \quad (3.3)$$

where $\{\psi_{j,k}\}_{j \geq 1}$ is a sequence in $\mathcal{L}_{\mathcal{H}}$ with $\sum_{k=0}^{\infty} k \|\psi_{j,k}\|_{\mathcal{L}_{\mathcal{H}}} < \infty$ for all j , and ε_t is an iid copies of ε with mean zero and positive definite covariance C_{ε} .

(ii) $\mathbb{E}[\|\varepsilon_t\|^p] < \infty$ for some $p > \max\{4, 2/(2d_K + 1)\}$.

(iii) $\mathcal{H}_1 := \text{ran } P_1$ is finite dimensional, i.e., $\dim(\mathcal{H}_1) = \text{rank } P_1 < \infty$.

(iv) $d_1 \in (-1/2, 2)$.

Some comments on Assumption **F** are in order. We first note that \mathbf{Y}_0 is a FIHTS with the dominant subspace $\mathcal{H}_1 := \text{ran } P_1$ (see e.g., Li et al., 2020, 2023; Seo and Shang, 2024). The case where \mathcal{H}_1 is a finite dimensional subspace has been considered in the literature on FIHTS, and thus Assumption **F(iii)** does not seem to be restrictive. In our theoretical development, finite dimensionality of \mathcal{H}_1 is only used to easily obtain a stochastic bound of a certain operator (see e.g., (A.8)). It should also be noted that we do not require finite dimensionality of the other subspaces (i.e., $\text{ran } P_2, \dots, \text{ran } P_K$), and thus the time series described by Assumptions **F(i)-(iii)** is still quite general than those considered in the recent literature. If $d_1 = 0$, the time series X_t defined in Assumption **F** is $I(0)$ in the sense of Definition 1. That is, the time series basically allows potential memory reduction on \mathcal{H} and thus our statistical inference needs to be based on elements in \mathcal{H}_1 ; if we use $(I_{\mathcal{H}} - P_1)Z_t$, then we may mistakenly conclude that its memory is negative even if \mathbf{Y}_0 is $I(0)$ and, due to this nature, it is important to base our statistical inference on the dominant subspace \mathcal{H}_1 . Assumption **F(iv)** is introduced to simplify the subsequent discussion. Under this assumption, the order of integer integration is either 0 or 1. This assumption does not place any practical restriction given that examples of $I(d)$ processes with $d \geq 2$ appear to be scarce and thus they have been seldom discussed in the literature on function time series analysis. Moreover, even if we mainly develop our theoretical results under this assumption, it is straightforward to extend the subsequent results to the case with $d_1 \in (-1/2, m)$ for some other integer $m > 2$ (see Remark 3.3).

It will be convenient to introduce some additional notation. For any time series $\mathbf{Z} = \{Z_t\}_{t=t_0}^T$ (which will be set to $\mathbf{Y}_0, \mathbf{Y}_1$ or their projections in the subsequent discussion), where the realizations are given for $t = t_0, \dots, T$ (e.g., $t_0 = 1$ if $\mathbf{Z} = \mathbf{Y}_0$ while $t_0 = 2$ if $\mathbf{Z} = \mathbf{Y}_1$), we define

$$\mathcal{K}_T(\mathbf{Z}) := \sum_{t=t_0}^T \left(\sum_{s=t_0}^t Z_s \otimes \sum_{s=t_0}^t Z_s \right).$$

We also let

$$\Lambda_T(\mathbf{Z}) := T^{-1} \sum_{s=-q}^q w(s, q) \sum_{t_0 \leq t, t-s \leq T} Z_t \otimes Z_{t-s}, \quad (3.4)$$

where $w(s, q) = 1 - |s|/(q+1)$, which corresponds to the Bartlett kernel weight function. $\mathcal{K}_T(\mathbf{Z})$ (resp. $\Lambda_T(\mathbf{Z})$) is the unnormalized sample covariance (long-run covariance) operator of $\sum_{s=t_0}^t Z_s$ (resp. Z_t).

3.2 Testing I(0)-ness against FI alternatives

We first consider testing I(0)-ness (i.e., $d_1 = 0$) against stationary FI alternatives (i.e., the cases where $d_1 \in (-1/2, 1/2) \setminus \{0\}$), and then this will be later extended to the case accommodating nonstationary FI alternatives (i.e., the cases where $d_1 \geq 1/2$). The test is constructed from the time series \mathbf{Y}_0 .

Under Assumption **F**, we know that $v_{1,t} = \Delta^{d_1} P_1 X_t$ is a stationary I(0) sequence. We let Λ_{v_1} denote the population long-run covariance operator of $v_{1,t}$, i.e.,

$$\Lambda_{v_1} = \sum_{s=-\infty}^{\infty} \mathbb{E} [v_{1,t} \otimes v_{1,t-s}].$$

Let \bar{h} be any unit-norm element of the dominant subspace \mathcal{H}_1 . We first establish the following preliminary result, which helps in understanding the practical challenges involved in developing a test for examining integer integration: below and hereafter, we let W denote the standard Brownian motion on $[0, 1]$ and let \int represent \int_0^1 for notational simplicity.

Proposition 3.1 (An infeasible test). *Suppose that $-1/2 < d_1 < 1/2$ and Assumption **F** is satisfied. Then,*

$$T^{-2} \langle \mathcal{K}_T(\mathbf{Y}_0) \bar{h}, \bar{h} \rangle \xrightarrow{d} c_D \int W(r)^2 dr \quad \text{if } d_1 = 0,$$

where $c_D = \langle \Lambda_{v_1} \bar{h}, \bar{h} \rangle$. On the other hand,

$$\begin{aligned} T^{-2} \langle \mathcal{K}_T(\mathbf{Y}_0) \bar{h}, \bar{h} \rangle &\xrightarrow{p} 0 \quad \text{if } d_1 \in (-1/2, 0), \\ T^{-2} \langle \mathcal{K}_T(\mathbf{Y}_0) \bar{h}, \bar{h} \rangle &\xrightarrow{p} \infty \quad \text{if } d_1 \in (0, 1/2). \end{aligned}$$

From Proposition 3.1, we know that the statistic $\langle \mathcal{K}_T(\mathbf{Y}_0) \bar{h}, \bar{h} \rangle$ has discriminating power for the null hypothesis of I(0)-ness against stationary FI alternatives. However, in practice, Proposition 3.1 cannot directly be used for statistical inference since c_D and \bar{h} , which is an element of \mathcal{H}_1 , are unknown. Given that \bar{h} is an element of the dominant subspace, one might attempt to replace \bar{h} with a reasonable estimator \bar{h}_T , which can be obtained by estimating the dominant subspace of \mathcal{H}_1 using some existing methods (see, e.g., Li et al., 2020), and then employing a standard long-run covariance estimator, if available, to replace $c_D = \langle \Lambda_{v_1} \bar{h}, \bar{h} \rangle$ with its feasible counterpart. In fact, our estimator is obtained in a broadly similar manner. However, it should be noted that the asymptotic properties of the estimator of \bar{h} will inevitably and crucially depend on the value of d_1 and possibly on the other memory parameters, d_2, \dots, d_K (it is difficult to believe that a reasonable estimator, independent of the memory properties, exists). As a consequence, it is also not straightforward at all to replace c_D with its feasible counterpart while preserving discriminating power against FI alternatives for any nonzero values of d_1 .

In order to have a feasible and asymptotically valid test, we require additional conditions for the time series introduced in Assumption F. We will hereafter assume the following conditions for the stationary component of \mathbf{Y}_0 :

Assumption S. Let $d_S := \max\{d_j : d_j < 1/2\}$ and let $P_S = \sum_{d_j \leq d_S} P_j$ in Assumption F. The stationary sequence of $P_S X_t$ satisfies the following: (i) $T^{-1} \sum_{t=1}^T P_S X_t = O_p(T^{-1/2+d_S})$; (ii) for any $\ell \geq 0$, $T^{-1} \sum_{t=\ell+1}^T P_S X_t \otimes P_S X_{t-\ell} \xrightarrow{p} C_{S,\ell} := \mathbb{E}[P_S X_t \otimes P_S X_{t-\ell}]$ and (iii) $C_{S,0}$ is positive definite on \mathcal{H}_1 .

Some comments on the above are in order. Note that the time series $\{P_S X_t\}_{t \geq 1}$ is a stationary process, and in this case, assuming condition (ii) does not appear to be restrictive. For example, Salish and Gleim (2019) establish (i) and (ii) for FIHTS (with long-range dependence) under nonrestrictive conditions. Assumption S along with Assumption F will be maintained throughout this section.

We next show that a feasible version of the test can be obtained from $\mathcal{K}_T(\mathbf{Y}_0)$ and its dominant eigenvector \bar{h}_T , corresponding to the maximal eigenvalue. Specifically, consider the

test statistic

$$V_0 = T^{-2} \frac{\langle \mathcal{K}_T(\mathbf{Y}_0) \bar{h}_T, \bar{h}_T \rangle}{\langle \Lambda_T(\mathbf{Y}_0) \bar{h}_T, \bar{h}_T \rangle}.$$

The above statistic is used to examine the null of I(0)-ness against FI alternatives. V_0 may be viewed as a ratio of two different variances (associated with $\mathcal{K}_T(\mathbf{Y}_0)$ and $\Lambda_T(\mathbf{Y}_0)$), and thus it may be called a variance-ratio statistic. The dominant eigenvector \bar{h}_T can easily be computed from the standard functional principal component analysis (FPCA), and thus the test statistic is easy-to-compute.

We will show that the test based on V_0 is consistent against stationary FI alternatives under the following additional assumptions: below, we let $\mathbf{X}_1 = \{P_1 X_t\}_{t=1}^T$, where X_t and P_1 are detailed in Assumption **F** (note that the subscript 1 is associated with P_1 and is used differently from that of \mathbf{Y}_1).

Assumption A1. $d_1 = d_S = \max\{d_j : d_j < 1/2\}$, and furthermore, the following hold: (i) $q^{-2d_1} \langle \Lambda_T(\mathbf{X}_1)h, h \rangle \xrightarrow{P} c_{j,h} > 0$ for any $h \in \mathcal{H}_1$ with $\|h\| = 1$; (ii) $q \rightarrow \infty$ and $q = o(T^v)$ for $v \in (0, 1/2)$; if $d_1 < 0$, it is further assumed that $v < |(d_1 - d_2)/2d_1|$.

Assumption **A1(i)** is a high-level assumption employed to conveniently obtain a feasible version of the test. Under Assumptions **F** and **S**, primitive sufficient conditions for this can be found in e.g., [Abadir et al. \(2009\)](#), as detailed in Remark **3.1**. Requirements for the lag-truncation parameter q are detailed by Assumption **A1(ii)**. A choice of q for the case when $d_1 \geq 0$ is common in practical studies, necessitating the sample long-run covariance estimation, and thus does not place any practical restriction. The requirement for the case $d_1 < 0$ is introduced for mathematical convenience, related to anti-persistent components; combined with Assumption **A1(i)**, this assumption facilitates extending our asymptotic results developed for persistent FIHTS (with $d_1 > 0$) to the case with anti-persistent FIHTS (with $d_1 < 0$); see our proof of Proposition **3.2**. This specifically requires that q increases at a sufficiently slow rate depending on the values of d_1 and d_2 . A theoretical choice satisfying this condition regardless of the values of d_1 and d_2 is $q = O(\log T)$. We conjecture that this assumption is not essential for our theoretical results and can be relaxed by developing appropriate asymptotic results for anti-persistent functional time series. However, investigating this issue requires a comprehensive study of its own, and thus we do not further pursue it in the present paper.

Remark 3.1. If $d_1 < 1/2$ (and thus $d_1 = d_S$), for any $h \in \mathcal{H}_j$ with $\|h\| = 1$, $\langle \Lambda_T(\mathbf{X}_1)h, h \rangle$ is the sample long-run covariance estimator of a fourth-order stationary I(d_1) sequence (Assumption **F**). From Theorem 2.2 of [Abadir et al. \(2009\)](#), we know that, under certain

non-restrictive conditions on the spectral density and autocovariances of such a time series (detailed in equation (1.1) and Assumption M in their paper), $\langle q^{-2d_1} \Lambda_T(\mathbf{X}_1)h, h \rangle$ converges to a positive constant depending on d_1 and h , under a requirement on q implied by Assumption A1(ii) (see Section A.3 of the Appendix for more details). In this case, Assumption A1(i) is clearly satisfied.

We next develop a feasible test of the null of I(0)-ness against stationary FI alternatives.

Proposition 3.2 (Feasible test of I(0)-ness). *Suppose that $-1/2 < d_1 < 1/2$ and Assumptions F, S and A1 are satisfied. Then,*

$$V_0 \xrightarrow{p \text{ or } d} \begin{cases} 0 & \text{if } d_1 \in (-1/2, 0), \\ \int W(r)^2 dr & \text{if } d_1 = 0, \\ \infty & \text{if } d_1 \in (0, 1/2). \end{cases}$$

Proposition 3.2 shows that the V_0 -test has power against stationary FI alternatives with $d_1 \in (-1/2, 1/2) \setminus \{0\}$. Given this result, it appears to be reasonable to conjecture that this test also has power against nonstationary FI alternatives concerning more persistent FIHTS, and this is investigated in the following:

Proposition 3.3. *Suppose that $d_1 \geq 1/2$ and Assumptions F and S hold. If $d_1 = 1/2$, we further assume that, for any $h \in \mathcal{H}_1$ and $t = 1, \dots, T$,*

$$\langle P_1 X_t, h \rangle = O_p(\sqrt{\log T}), \quad (3.5)$$

where $P_1 X_t$ is defined in Assumption F. Then, $V_0 \xrightarrow{p} \infty$.

In fact, based on some key asymptotic results used in the proof of Proposition 3.2, it is not difficult to show that the test statistic V_0 diverges to infinity under nonstationary FI alternatives with $d_1 > 1/2$. To conveniently address the particular case of weak nonstationarity ($d_1 = 1/2$), Proposition 3.3 imposes condition (3.5). This condition does not seem restrictive and is expected to hold generally given that $\langle P_1 X_t, h \rangle$ can be understood as the partial sum process $\sum_{s=1}^t x_s$, where $\{x_s\}_{s \geq 1}$ is an I(-1/2) time series, and $\sum_{s=1}^T x_s = O_p(\sqrt{\log T})$ holds under standard regularity conditions for such a time series; see, e.g., Liu (1998, Lemma 2.1) and Tanaka (2017, Ch. 12).

Propositions 3.2 and 3.3 give us a consistent test for examining if $d_1 \in (-1/2, 0)$, $\{0\}$, or $(0, \infty)$. Rejection in the upper (resp. lower) tail of the V_0 -test means that the time series is fractionally-integrated with persistence (resp. anti-persistence).

3.3 Testing I(1)-ness against FI alternatives

Suppose that the V_0 -test proposed in Section 3.2 rejects the null hypothesis of I(0)-ness in the upper tail and thus $d_1 > 0$ is concluded. Practitioners may then be interested in testing the null of I(1)-ness against FI alternatives as the next step. Considering the properties of an I(1) time series, it may be appropriate to apply the test considered in Section 3.2 to the first-differenced observations $\mathbf{Y}_1 = \{Y_{1,t}\}_{t=2}^T$ (see (3.1)). Specifically, define

$$V_1 = T^{-2} \frac{\langle \mathcal{K}_T(\mathbf{Y}_1) \bar{h}_T, \bar{h}_T \rangle}{\langle \Lambda_T(\mathbf{Y}_1) \bar{h}_T, \bar{h}_T \rangle}, \quad (3.6)$$

where \bar{h}_T is the dominant eigenvector which we used in Section 3.2. We will demonstrate that the test based on the above statistic can be used to examine I(1)-ness against FI alternatives, not only as a test applied after the V_0 -test in Section 3.2, but also as a standalone test for examining I(1)-ness against I(d)-ness for any other possible d in the admissible range $(-1/2, 2)$ (see Remark 3.2 to appear).

We investigate the asymptotic properties of V_1 . To this end, for the time series defined in Assumption F, we let $\Delta \mathbf{X}_1 = \{P_1 \Delta X_t\}_{t=2}^T$ and employ the following conditions:

Assumption A2. $d_1 > 1/2$ and the following hold: (i) $q^{-2(d_1-1)} \langle \Lambda_T(\Delta \mathbf{X}_1) h, h \rangle \xrightarrow{p} c_{j,h} > 0$ for any $h \in \mathcal{H}_1$ with $\|h\| = 1$; (ii) if $d_1 = 3/2$, for any $h \in \mathcal{H}_1$, $\langle P_1 \Delta X_t, h \rangle = O_p(\sqrt{\log T})$; (iii) $q \rightarrow \infty$ and $q = o(T^v)$ for $v \in (0, 1/2)$; if $d_1 \in (1/2, 1)$, $v < |(d_1 - d_2)/2(d_1 - 1)|$.

The above conditions are obvious adaptations of Assumption A1 and (3.5) for nonstationary FIHTS with $d_1 > 1/2$. We first show that the V_1 -test is consistent against nonstationary FI alternatives.

Proposition 3.4. *Suppose that $d_1 > 1/2$ and Assumptions F, S and A2 are satisfied. Then, the following hold:*

$$V_1 \xrightarrow{p \text{ or } d} \begin{cases} 0 & \text{if } d_1 \in (1/2, 1), \\ \int W(r)^2 dr & \text{if } d_1 = 1, \\ \infty & \text{if } d_1 > 1. \end{cases}$$

That is, the V_1 -test can distinguish the null of I(1)-ness from nonstationary FI alternatives. Moreover, we can determine which of the following intervals includes the memory parameter d_1 : $(1/2, 1)$, $\{1\}$, and $(1, 2)$.

Even if the V_1 -test, described in Proposition 3.4, can be used to examine I(1)-ness, there is an unsatisfactory property as a test used after rejection of the V_0 -test in the upper tail:

according to Propositions 3.2 and 3.3, rejection of the V_0 -test means that the memory parameter of \mathbf{Y}_0 is contained in the interval $(0, \infty)$, but rejection of the V_1 -test in the lower tail in Proposition 3.4 means that the memory of \mathbf{Y}_0 is included in the interval $(1/2, 1)$, not $(0, 1)$ containing all fractions between 0 and 1. Thus it would be useful if we extend the V_1 -test given in Proposition 3.4 so that the test has a discriminating power when \mathbf{Y}_0 is FIHTS in Assumption F with $d_1 \in [0, 1/2]$. We hereafter investigate this issue. Given that $\mathbf{Y}_0 \sim I(d_1)$ implies that $\Delta X_t \sim I(d_1 - 1)$, this obviously requires a detailed investigation of statistical properties of an $I(d_1 - 1)$ process. A complicated case occurs particularly when $d_1 = 1/2$, and, to deal with this case conveniently, we assume the following condition for the subsequent discussion:

Assumption A3. *If $d_1 = 1/2$, the following hold: (i) $\langle P_1 X_t, h \rangle = O_p(\sqrt{\log T})$; (ii) $q \rightarrow \infty$ and $q = o(T^v)$ for $v \in (0, 1/2)$ and there exists a divergent sequence $d_T = o(T/\log T)$ such that $d_T \langle \Lambda_T(\Delta \mathbf{X}_1)h, h \rangle$ is bounded away from zero with probability approaching one for every $h \in \mathcal{H}_1$.*

Assumption A3(i) is a repetition of the requirement for the case with $d_1 = 1/2$ given in Proposition 3.3. Assumption A3(ii) seems to be a fairly high-level condition, but it is not restrictive. For example, Cho et al. (2015, Theorem 6) show that Gaussian $I(-1/2)$ processes generally satisfy this condition with $d_T = q/\log q$, which is of much smaller asymptotic order than $T/\log T$ under $q = o(T^{1/2})$. We thus believe that some deviations from the conditions employed by Cho et al. (2015) can also be accommodated under Assumption A3. The desired extension of the V_1 -test in Proposition 3.4 is given as a direct consequence of our next result:

Proposition 3.5. *Suppose that $0 \leq d_1 \leq 1/2$ and Assumptions F, S and A3 hold. Then $V_1 \rightarrow_p 0$.*

Remark 3.2. Proposition 3.5 can be extended to the case with $-1/2 < d_1 \leq 1/2$ without requiring further assumptions. In fact, our proof of Proposition 3.5 given in the Appendix is provided to accommodate this more general case. This means that the V_1 -test is a standalone test, which can be used to examine $I(1)$ -ness against $I(d)$ for any other values of d in the admissible range $(-1/2, 2)$ (see Assumption F(iv)).

Remark 3.3 (V_2 -test and its further extension). Due to simplicity and empirical relevance, we focus in this paper on the case with $d_1 \in (-1/2, 2)$. However, the V_1 -test can be extended to another test, say the V_2 -test, using the second-differenced time series of $Y_{2,t}$ (see (3.1)), for examining $I(2)$ -ness against FI alternatives. The extension from the V_1 -test to the V_2 -test

is quite obvious and similar to that from the V_0 -test to the V_1 -test, which is described by Propositions 3.4 and 3.5. In a similar manner, the V_k -test for an integer $k \geq 3$ can also be considered.

3.4 Extensions

3.4.1 Sequential testing procedure

In this section, we consider a combined procedure of the two tests developed in Sections 3.2 and 3.3 as a way to assess the statistical plausibility of the null hypothesis of integer integration. Let η_α be the $\alpha\%$ quantile of $\int W(r)^2 dr$ for $\alpha \in (0, 1/2)$. Consider the following test function:

$$D_{\text{seq}} = \mathbb{1}\{\eta_\alpha < V_0 < \eta_{1-\alpha}\} + \mathbb{1}\{V_0 > \eta_{1-\alpha}\}\mathbb{1}\{\eta_\alpha < V_1 < \eta_{1-\alpha}\}, \quad (3.7)$$

where $\mathbb{1}\{\cdot\}$ denotes the binary indicator function and $D_{\text{seq}} = 1$ is understood as acceptance of the null hypothesis of integer integration. If $d_1 = 0$, we find that $\mathbb{P}(\eta_\alpha < V_0 < \eta_{1-\alpha}) \rightarrow 1 - 2\alpha$ while $\mathbb{P}(V_0 > \eta_{1-\alpha}) \rightarrow 0$. On the other hand, if $d_1 = 1$, $\mathbb{P}(V_0 > \eta_{1-\alpha}) \rightarrow 1$ and $\mathbb{P}(\eta_\alpha < V_1 < \eta_{1-\alpha}) \rightarrow 1 - 2\alpha$. Thus under either of these cases, $\mathbb{P}\{D_{\text{seq}} = 1\} \rightarrow 1 - 2\alpha$. If d_1 is a fraction in $(-1/2, 2)$, then either of the following holds:

$$\mathbb{P}(V_0 < \eta_\alpha) \rightarrow 1, \quad \mathbb{P}(V_0 > \eta_{1-\alpha} \text{ and } V_1 < \eta_\alpha) \rightarrow 1 \quad \text{or} \quad \mathbb{P}(V_0 > \eta_{1-\alpha} \text{ and } V_1 > \eta_{1-\alpha}) \rightarrow 1.$$

Therefore, $\mathbb{P}\{D_{\text{seq}} = 0\} \rightarrow 1$. Therefore, (3.7) constitutes a consistent decision rule to distinguish the null hypothesis of integer integration against fractional integration with significant level 2α . These results are summarized below:

Corollary 3.1. *Suppose that all the assumptions for the V_0 - and V_1 -tests in Sections 3.2 and 3.3 are satisfied for $d_1 \in (-1/2, 2)$. Then, under the null of integer integration of order 0 or 1,*

$$\mathbb{P}\{D_{\text{seq}} = 1\} \rightarrow 1 - 2\alpha.$$

Under the alternative of fractional integration of order $d \in (-1/2, 2) \setminus \{0, 1\}$,

$$\mathbb{P}\{D_{\text{seq}} = 0\} \rightarrow 1.$$

Remark 3.4. For simplicity, we only considered FIHTS with the memory parameter $d_1 \in (-1/2, 2)$. However, we may extend the testing procedure developed in Corollary 3.1 to accommodate integrated time series of a higher order by sequentially adding the V_k -test for $k = 2, 3, \dots, M$ for a finite integer M , obtained from extending the V_1 -test (see Remark

3.3). As illustrated by Corollary 3.1, sequential applications of these tests do not distort the asymptotic size and are consistent. For example, if we consider the case with $d_1 \in (-1/2, 3)$, we can additionally apply the V_2 -test described in Remark 3.3, and in this case, the relevant test function will become

$$D_{\text{seq}} = \mathbb{1}\{\eta_\alpha < V_0 < \eta_{1-\alpha}\} + \mathbb{1}\{V_0 > \eta_{1-\alpha}\}\mathbb{1}\{\eta_\alpha < V_1 < \eta_{1-\alpha}\} \\ + \mathbb{1}\{V_0 > \eta_{1-\alpha}\}\mathbb{1}\{V_1 > \eta_{1-\alpha}\}\mathbb{1}\{\eta_\alpha < V_2 < \eta_{1-\alpha}\}.$$

As shown above, the combination of the V_0 - and V_1 -tests can not only be used to examine integer integration, but also enables us to determine which of the following subsets of $(-1/2, 2)$ contains the true memory parameter:

$$\mathcal{I}_1 = (-1/2, 0), \quad \mathcal{I}_2 = \{0\}, \quad \mathcal{I}_3 = (0, 1), \quad \mathcal{I}_4 = \{1\}, \quad \mathcal{I}_5 = (1, 2).$$

That is, by first applying the V_0 -test, we may determine if $d_1 \in \mathcal{I}_1$ (when the test is rejected in the lower tail) or $d_1 \in \mathcal{I}_2$ (when the test is not rejected) or $d_1 \in \mathcal{I}_3 \cup \mathcal{I}_4 \cup \mathcal{I}_5$ (when the test is rejected in the upper tail). If the latter is concluded from the V_0 -test, then we may apply the V_1 -test to determine if $d_1 \in \mathcal{I}_3$ (when the test is rejected in the lower tail) or $d_1 \in \mathcal{I}_4$ (when the test is not rejected) or $d_1 \in \mathcal{I}_5$ (when the test is rejected in the upper tail). This can be a useful information in implementing some statistical methods on the memory parameter requiring appropriate bounds of d_1 as an input, such as the local Whittle estimator of Li et al. (2021), Li et al. (2023), and Seo and Shang (2024) in the functional time series setup.

If the time series of interest is anticipated to be similar to an I(1) process as in many empirical examples, a reversed procedure may be more practically relevant. Specifically, we may first apply the V_1 -test to determine if $d_1 \in \mathcal{I}_5$ (when the test is rejected in the upper tail) or $d_1 \in \mathcal{I}_4$ (when the test is not rejected) or $d_1 \in \mathcal{I}_1 \cup \mathcal{I}_2 \cup \mathcal{I}_3$ (when the test is rejected in the lower tail). If the latter is concluded from the V_1 -test, we may apply the V_0 -test to determine if $d_1 \in \mathcal{I}_3$ (when the test is rejected in the upper tail) or $d_1 \in \mathcal{I}_2$ (when the test is not rejected) or $d_1 \in \mathcal{I}_1$ (when the test is rejected in the lower tail). This may be understood as a reversed version of the aforementioned testing procedure.

3.4.2 Deterministic terms

In the previous sections, we have used the initialized variable $Y_t - Y_0$, which is assumed to have zero mean. We in this section consider the case with an unknown intercept, where observations are given by $\tilde{\mathbf{Y}}_0 = \mu + \mathbf{Y}_0 = \{\mu + Y_{0,t}\}_{t=1}^T$, and consider a testing procedure that

is robust to the existence of an unknown intercept. We hereafter define

$$\tilde{\mathcal{K}}_T(\mathbf{Z}) := \sum_{t=t_0}^T \left(\sum_{s=t_0}^t (Z_s - \bar{Z}_T) \right) \otimes \left(\sum_{s=t_0}^t (Z_s - \bar{Z}_T) \right),$$

where \bar{Z}_T is the sample mean of Z_t over $t = t_0, \dots, T$. We also let

$$\tilde{\Lambda}_T(\mathbf{Z}) := T^{-1} \sum_{s=-q}^q w(s, q) \sum_{t_0 \leq t, t-s \leq T} (Z_t - \bar{Z}_T) \otimes (Z_{t-s} - \bar{Z}_T),$$

where $w(s, q) = 1 - |s|/(q+1)$. We then consider the test statistic given as follows:

$$\tilde{V}_0 = T^{-2} \frac{\langle \tilde{\mathcal{K}}_T(\tilde{\mathbf{Y}}_0) \tilde{h}_T, \tilde{h}_T \rangle}{\langle \tilde{\Lambda}_T(\tilde{\mathbf{Y}}_0) \tilde{h}_T, \tilde{h}_T \rangle},$$

where \tilde{h}_T is the dominant eigenvector of $\tilde{\mathcal{K}}_T(\mathbf{Y}_0)$.

We can similarly define \tilde{V}_1 as in (3.6). However, since we use the first-differenced sequence for this test, the test statistic does not change from V_1 in (3.6), except that \bar{h}_T is replaced by \tilde{h}_T . As will be discussed in our proof of Proposition 3.6, under Assumption F, \tilde{h}_T satisfies all the necessary properties required for \bar{h}_T , which are used to establish the asymptotic properties of the test. Hence, the two tests have the same limiting behaviors under the null and alternative hypotheses. We thus focus only on the limiting behavior of \tilde{V}_0 in this section.

Proposition 3.6. *Suppose that $-1/2 < d_1 < 1/2$, Assumptions F and S hold, and Assumption A1 is satisfied when $\Lambda_T(\mathbf{X}_1)$ is replaced by $\tilde{\Lambda}_T(\mathbf{X}_1)$. Then,*

$$\tilde{V}_0 \xrightarrow{p \text{ or } d} \begin{cases} 0 & \text{if } d_1 \in (-1/2, 0), \\ \int \tilde{W}(r)^2 dr & \text{if } d_1 = 0, \\ \infty & \text{if } d_1 \in (0, 1/2), \end{cases}$$

where \tilde{W} is the standard Brownian bridge on $[0, 1]$, given by $\tilde{W}(r) = W(r) - rW(1)$. If $d_1 \geq 1/2$, Assumptions F and S hold, and (3.5) is satisfied when $d_1 = 1/2$, then $\tilde{V}_0 \xrightarrow{p} \infty$.

Remark 3.5. We discuss primitive sufficient conditions for Assumption A1(i) to be satisfied when $\Lambda_T(\mathbf{X}_1)$ is replaced by $\tilde{\Lambda}_T(\mathbf{X}_1)$ in Section A.3 of the Appendix.

4 Simulation studies

4.1 Simulation data generating process

We provide simulation results for the proposed test. The simulation data generating process (DGP) considered in this section is similar to the curve-valued processes considered by [Seo and Shang \(2024\)](#). In our subsequent experiments, d_1 is controlled to be a specified value in $[-0.45, 0.45]$, and depending on this value, we generate FIHTS so that it exhibits various degrees of memory reduction on \mathcal{H} .

Let $\{e_k\}_{k \geq 1}$ be an orthonormal basis of \mathcal{H} , where $\{e_k\}_{k=1}^{r_1}$ is an orthonormal set, spanning the dominant subspace $\mathcal{H}_1 = \text{ran } P_1$ of dimension $r_1 > 0$. We next define $\mathcal{H}_2 = \text{ran } P_2$ as the $(r_2 - r_1)$ -dimensional subspace spanned by $\{e_k\}_{k=r_1+1}^{r_2}$, and successively define $\mathcal{H}_j = \text{ran } P_j$ as the span of $\{e_k\}_{k=r_{j-1}+1}^{r_j}$ for $j = 1, \dots, K$ and $r_1 < r_2 < \dots < r_K$.

For each j , we let $\Delta^{d_j} P_j X_t$ in [\(3.2\)](#) – [\(3.3\)](#) be generated as follows:

$$\Delta^{d_j} P_j X_t = \sum_{\ell=r_{j-1}+1}^{r_j} a_{\ell,t} \sigma_\ell e_\ell, \quad a_{\ell,t} \sim \text{ARMA}(1, 1), \quad (4.1)$$

where the AR coefficient ϕ_ℓ and MA coefficient θ_ℓ , characterizing the ARMA(1, 1) process $\{a_{\ell,t}\}$, are determined as follows in each simulation run:

$$\phi_\ell \sim U[-\mathbf{b}, \mathbf{b}] \quad \text{and} \quad \theta_\ell \sim U[-\mathbf{b}, \mathbf{b}],$$

for some $\mathbf{b} > 0$ without dependence on any other variables. In the subsequent simulation experiments, we consider the cases with $\mathbf{b} = 0.15$ and $\mathbf{b} = 0.6$, and obviously, the former case is closer to white noise than the latter. Using [\(4.1\)](#) and an appropriate anti-differencing operation, we construct $P_j X_t$ and compute $Y_{0,t}$ as $Y_{0,t} = \sum_{j=1}^K P_j X_t$ for large enough $K > 0$. σ_ℓ in [\(4.1\)](#) is a sequence decaying to zero as ℓ gets larger, this is introduced to ensure the required summability condition for the resulting time series to be \mathcal{H} -valued (see e.g. [Nielsen et al., 2023a](#)); we specifically let $\sigma_\ell = \ell^{-2}$ so that the variances of $a_{\ell,t}$ for $\ell \geq 1$ become summable.

For each simulation run where d_1 is specified, we let d_2, \dots, d_K and the dimensions of $\text{ran } P_j$, $p_j = r_j - r_{j-1}$ (for $j \geq 1$ with $r_0 = 0$), be randomly determined. By doing so, we can evaluate the average performance of the proposed test to some degree for various values of those parameters. Specifically, in our simulation concerning the stationary FIHTS (i.e., the case where $d_1 \in (-1/2, 1/2)$), we let $p_j \sim [U(1, 4)]$, where $[a]$ denotes the integer part of a . The memory parameters d_2, \dots, d_K are determined as the ordered (from the largest to

smallest) realizations $\{\bar{d}_j\}_{j=1}^K$ from uniform random variables given by

$$\bar{d}_j \sim \begin{cases} U((d_1 - 0.2) \vee -0.5, (d_1 - 0.1) \vee -0.49) & \text{if } j = 2, \\ U((d_1 - 0.5) \vee -0.5, (d_1 - 0.2) \vee -0.49) & \text{if } j \geq 3, \end{cases} \quad (4.2)$$

where $a \vee b = \max\{a, b\}$ for $a, b \in \mathbb{R}$. Since we consider the case where $d_1 \in [-0.45, 0.45]$, d_2 (i.e., the largest \bar{d}_j) obtained as above satisfies that $0.04 < |d_1 - d_2| < 0.2$, and hence the simulation DGP always allows a moderate memory reduction, which may be regarded as a realistic assumption in many empirical applications.

In the cases where nonstationary FIHTS are considered, we let $d_1 \geq 0.55$, and let $p_1(= r_1)$, $p_2(= r_2 - r_1)$ and d_2 be determined as follows:

$$p_1 \sim [U(1, 4)], \quad p_2 \sim [U(1, 4)], \quad d_2 \sim U((d_1 - 0.2) \vee 0.5, (d_1 - 0.1) \vee 0.51). \quad (4.3)$$

We then let $P_1 X_t$ and $P_2 X_t$ constitute the nonstationary part X_t^N of the time series, with $\Delta^{d_1} P_1 X_t$ and $\Delta^{d_2} P_2 X_t$ being generated as in (4.1). The time series used in the analysis is obtained by adding a stationary component X_t^S to this nonstationary time series; specifically we let the stationary part X_t^S be generated exactly as in the above case concerning stationary FIHTS with $d_1 = 0.25$.

We let the basis system $\{e_k\}_{k \geq 1}$ vary to some degree across different simulation runs (see e.g., [Aue et al., 2018](#)). Specifically, we let $\{e_k\}_{k \geq 1}$ be the Fourier basis functions of which the first five basis functions are randomly permuted, so that the dominant parts of the time series take values in different subspaces across different simulation runs. This is also to avoid that the performance of the test is evaluated with specific shapes of the dominant components of the generated functional observations.

4.2 Simulation results

We first examine the sizes and correct rejection rates of the proposed V_0 - and V_1 -tests in various scenarios regarding the integration order. The correct rejection rate is slightly different from the power of the test. Specifically, it measures the relative frequency that the null is rejected at lower (resp. upper) tail when $d_1 < 0$ (resp. $d_1 > 0$) for the V_0 -test. For the V_1 -test, it measures the relative frequency that the test is rejected at lower (resp. upper) tail when $d_1 < 1$ (resp. $d_1 > 1$). Considering that our testing results may also be used to conjecture a reasonable range for d_1 , the correct rejection rate represents a performance measure as the relative frequency of the correct decision being made by the proposed test.

Table 1 shows the simulated sizes and correct rejection rates of the V_0 - and V_1 -tests in various scenarios. It may be noticeable that the V_0 -test (resp. V_1 -test) performs better in the case where $d_1 > 0$ (resp. $d_1 > 1$) compared to the case where $d_1 < 0$ (resp. $d_1 < 1$). This may be partly due to the restriction on admissible values of d_j in the DGP in each case. As described in (4.2) and (4.3), in the simulation DGP concerning the stationary (resp. nonstationary) case, the second most persistent $I(d_2)$ component is set to a stationary (resp. nonstationary) process, and thus d_2 needs to satisfy $d_2 > -1/2$ (resp. $d_2 > 1/2$). If d_1 is lower than the specified integer value under the null in each case, the gap between d_1 and d_2 tends to be smaller, and overall d_1, \dots, d_K tend to become more homogeneous (and hence less distinguishable from the data) compared to the case where d_1 is greater than the specified integer value.

Table 1: Size and correct rejection rates

(a) V_0 -test for $H_0 : d_1 = 0$

		d_1	-0.45	-0.3	-0.15	0	0.15	0.3	0.45
		T							
b = 0.15	250		0.832	0.470	0.128	0.031	0.264	0.659	0.946
	500		0.942	0.654	0.195	0.038	0.322	0.762	0.968
	750		0.979	0.757	0.248	0.036	0.360	0.804	0.984
	1000		0.994	0.812	0.282	0.041	0.392	0.834	0.991
b = 0.60	250		0.781	0.471	0.155	0.062	0.281	0.674	0.943
	500		0.910	0.643	0.220	0.064	0.342	0.762	0.968
	750		0.952	0.736	0.269	0.062	0.378	0.802	0.981
	1000		0.978	0.791	0.306	0.064	0.398	0.841	0.990

(b) V_1 -test for $H_0 : d_1 = 1$

		d_1	0.55	0.7	0.85	1	1.15	1.3	1.45
		T							
b = 0.15	250		0.800	0.460	0.117	0.032	0.280	0.676	0.945
	500		0.935	0.658	0.168	0.035	0.358	0.790	0.971
	750		0.975	0.757	0.206	0.028	0.404	0.834	0.981
	1000		0.990	0.809	0.238	0.033	0.432	0.862	0.987
b = 0.60	250		0.752	0.465	0.148	0.062	0.294	0.687	0.939
	500		0.901	0.634	0.206	0.064	0.373	0.789	0.970
	750		0.955	0.730	0.235	0.058	0.411	0.830	0.981
	1000		0.974	0.788	0.264	0.054	0.431	0.857	0.985

Notes: In the case with either $d_1 = 0$ or 1, the size of the test is reported; the nominal size is 5%. Except for these cases, the correct rejection rates are reported. The number of Monte Carlo replications in each case is 2000.

Overall, the V_0 - and V_1 -tests perform better as the sample size T gets larger, and both

tests have good size control. The correct rejection rates also increase as d_1 moves away from the hypothesized integer value for both tests, which may well be expected from the theoretical results presented earlier. The parameter \mathbf{b} , which measures the tendency of departure of the underlying ARMA process $\{a_{\ell,t}\}$ from white noise, seems to affect the performance of the tests, but the difference is moderate for the considered values of \mathbf{b} .

We next examine the performance of the sequential testing procedure discussed in Section 3.4.1. We first investigate how accurately the testing procedure identifies integer integration (the cases where $d = 0$ or 1) in the simulation DGP. We also assess its performance in the case of fractional integration by computing the relative frequency of correctly identifying the interval that includes the fractional order. The simulation results are presented in Table 2, along with additional details of the Monte Carlo experiments. Overall, the testing procedure performs reasonably well, even with a moderate sample size of $T = 250$, and its performance improves clearly as the sample size increases.

Table 2: Relative frequencies of correct determination of integer/fractional integration

(a) Case with integer integration					
	Case	$T = 250$	$T = 500$	$T = 750$	$T = 1000$
$\mathbf{b} = 0.15$	$d_1 \in \{0, 1\}$	0.970	0.960	0.959	0.957
	$d_1 = 0$	0.969	0.959	0.959	0.952
	$d_1 = 1$	0.950	0.955	0.957	0.962
$\mathbf{b} = 0.60$	$d_1 \in \{0, 1\}$	0.947	0.941	0.936	0.934
	$d_1 = 0$	0.948	0.935	0.934	0.931
	$d_1 = 1$	0.924	0.941	0.937	0.938
(b) Case with fractional integration					
	Case	$T = 250$	$T = 500$	$T = 750$	$T = 1000$
$\mathbf{b} = 0.15$	$d_1 < 0$	0.520	0.669	0.756	0.794
	$d_1 \in (0, 1)$	0.552	0.714	0.770	0.794
	$d_1 > 1$	0.715	0.757	0.800	0.827
$\mathbf{b} = 0.60$	$d_1 < 0$	0.533	0.657	0.718	0.754
	$d_1 \in (0, 1)$	0.563	0.702	0.749	0.776
	$d_1 > 1$	0.724	0.767	0.788	0.806

Notes: In the top panel, d_1 is set to 0 or 1 in the simulation DGP, and the case $d_1 \in \{0, 1\}$ reports the relative frequency of the cases where $d_1 = 0$ or 1 is concluded by the testing procedure for each sample size; the total number of Monte Carlo replications is set to 2000. Within the same simulation results, the case $d_1 = 0$ (resp. $d_1 = 1$) reports the frequency of correctly determining d_1 by the testing procedure, relative to the frequency of the cases where $d_1 = 0$ (resp. $d_1 = 1$) in the simulation DGP. The bottom panel reports the frequency of correctly identifying the subset \mathcal{I}_j for each sample size, relative to the frequency of the cases where $d_1 \in \mathcal{I}_j$ in the simulation DGP. In each simulation run concerning this case, d_1 is randomly chosen; specifically, d_1 is determined by one of the following four uniform random variables, each of which can occur with a probability 1/4: $U(-0.485, -0.15)$, $U(0.15, 0.5)$, $U(0.5, 0.85)$, or $U(1.15, 1.5)$. The number of Monte Carlo replications in each case is 2000.

We additionally examine the finite sample performance of the \tilde{V}_0 -test, which is discussed in Section 3.4.2. In this simulation experiment, we added a nonzero mean μ_ℓ to each of the ARMA process $a_{\ell,t}$ in 4.1, and μ_ℓ is drawn from $N(0, 1)$ in each simulation, without dependence of any other variables. The simulation results are reported in Table 3. The results are qualitatively similar to those of Table 1.

Table 3: Size and correct rejection rates for the \tilde{V}_0 -test for $H_0 : d_1 = 0$

		d_1						
T		-0.45	-0.3	-0.15	0	0.15	0.3	0.45
$b = 0.15$	$T = 250$	0.824	0.478	0.136	0.044	0.206	0.474	0.743
	$T = 500$	0.966	0.724	0.224	0.042	0.268	0.624	0.896
	$T = 750$	0.993	0.842	0.298	0.035	0.316	0.723	0.943
	$T = 1000$	0.998	0.896	0.345	0.036	0.345	0.780	0.963
$b = 0.60$	$T = 250$	0.775	0.476	0.172	0.085	0.208	0.491	0.762
	$T = 500$	0.939	0.689	0.248	0.077	0.288	0.650	0.902
	$T = 750$	0.976	0.782	0.314	0.082	0.328	0.739	0.946
	$T = 1000$	0.991	0.846	0.350	0.083	0.360	0.790	0.961

Notes: The nominal size is 5%. The number of Monte Carlo replications in each case is 2000.

5 Empirical applications

5.1 Canadian yield curves for zero-coupon bonds

We apply our proposed test to the end-of-day Canadian yield curves for the period spanning 04 July 2022 to 11 September 2024; the data used in this section is publicly available at <https://www.bankofcanada.ca/rates/interest-rates/bond-yield-curves/>. Except for the days with missing observations, the total number of observations is 545. Each observation corresponds to zero-coupon bond yields at 120 regularly spaced maturities ranging from 0.25 to 30 (years). In Figure 1, we present a rainbow plot of the Canadian yield curve data by maturities. A similar dataset is used by Seo and Shang (2024), although their analysis employs a long-span, monthly frequency time series, whereas the analysis here considers a daily frequency.

In some recent articles, a time series of yield curves is often mentioned as an example of nonstationary functional time series (see e.g., Li et al., 2023; Nielsen et al., 2023a; Seo and Shang, 2024), and empirical evidence suggests that such a time series behaves similarly to an I(1) process. We examine this in detail using our methodology.

Specifically, we apply the sequential testing procedure based on the V_0 - and V_1 -tests to the yield curve data. The test results are reported in the first two columns of Table 4. As




Figure 1: Visualization of Canadian yield curves

reported in the table, the V_0 -test is rejected in the upper tail with a p -value nearly equal to zero, meaning that $I(0)$ stationarity is strongly rejected against a higher order integration. On the other hand, the null hypothesis of $I(1)$ -ness is not rejected at 5% significance level, from which we conclude that the time series of interest is $I(1)$. Given that our testing procedure can easily be extended to examine a higher order integration (see Remark 3.3), one may want to confirm this test result by implementing the V_2 -test and checking if the null of $I(2)$ -ness is rejected in the lower tail. The test result is reported in the third column of Table 4, and the null of $I(2)$ -ness is strongly rejected against a lower order integration.

Table 4: Testing results for Canadian yield curves

	V_0 -test	V_1 -test	V_2 -test
Statistic	58.06	1.82	0.004
p -value	< 0.001	0.08	< 0.001
Test result	Rejection in the upper tail	Accept	Rejection in the lower tail

Notes : The p -values in the second row are derived from 200,000 Monte Carlo replications of the approximate limiting distribution. The third row presents the test results at the 5% significance level, where the null hypothesis is rejected if the test statistic falls below the 2.5th percentile ($q_{0.025}$) or exceeds the 97.5th percentile ($q_{0.975}$) of the limiting distribution, approximately 0.045 and 2.126, respectively. The reported p -value is computed as $2 \min\{\mathbb{P}(V_j > q_{0.975}), \mathbb{P}(V_j < q_{0.025})\}$, considering each of the considered tests is two-sided.

If the considered tests are implemented at the 10% significance level, then the V_1 -test rejects the null hypothesis of $I(1)$ integration as well, and thus the testing procedure concludes that this time series is fractionally-integrated of order $d \in (1, 2)$. Overall, our test results confirm the nonstationarity of the Canadian yield curve time series with an integration order of $d = 1$ or $d \in (1, 2)$.

5.2 French sub-national mortality rates

We next apply our proposed tests to age- and sex-specific French sub-regional mortality data from 1901 to 2021, with the data sourced from the French Human Mortality Database at <https://frdata.org/en/french-human-mortality-database/> (see Bonnet, 2020, for data description). Depending on the territorial units for statistics, there exist NUT1, NUT2 and NUT3 ranging from the largest to the smallest units. For illustration, we consider the middle level, i.e., NUT2, which has 22 regions. The number of samples is 121 for all regions except one (Alsace), which allows records with many missing or overly imputed entries from 1901 to 1920; for that region, we use data from 1921 to 2021. As is common in the literature on proportional rate data, we use the logit transformation of the mortality rate in the following analysis (see e.g., Cairns et al., 2011). The logit transformation ensures the transformed data are adequately defined. Mortality rates for each year are observed for ages ranging from 0 to 110 (and older) for both genders over time. We treat the logit transformation of the mortality rates at various ages as functional observations, as in Hyndman and Ullah (2007) and Shang and Hyndman (2017), before applying our proposed tests.

In Figure 2a, we display a rainbow plot of the logit-transformed age-specific female mortality for the region Île de France, along with its functional autocorrelation (ACF) plot in Figure 2b. From the rainbow plot, we observe a decreasing trend in age-specific mortality rates over the years. By examining the functional autocorrelation plot of Mestre et al. (2021), we notice a strong persistence exhibited in the data set.




Figure 2: Graphical displays of the logit-transformed age-specific female mortality rates in region Île de France.

Age-specific mortality rates are often considered to be $I(d)$ functional time series of integer integration order d (mostly $d = 1$). As in Section 5.1, we assess this using the V_0 -, V_1 - and V_2 -tests, covering 0, 1 and 2 as possible integer integration orders.

For each region, we apply our testing procedure to identify the order of integration of age-specific mortality rates for each gender, and the test results are reported in Table 5. For most regions, our testing procedure concludes that the considered time series is $I(1)$, which may be understood as evidence supporting the popularity of using the random walk with drift as the forecasting technique in Lee and Carter’s (1992) model. For some regions, the test results suggest fractional integration orders exceeding one (Île de France and Nord-Pas-de-Calais for female data) or falling below one (five regions for male data). However, there is no case where $I(0)$ -ness is supported in the considered dataset. This finding is consistent with the existing empirical evidence of nonstationarity or long-range dependence in a time series of age-specific mortality rates.

Table 5: Testing results for 22 sub-national age- and gender-specific logit-mortality rates

No.	Region	d_1 for female data	d_1 for male data
1	Île de France	$\in (1, 2)$	1
2	Centre-Val de Loire	1	1
3	Bourgogne	1	1
4	Franche-Comté	1	1
5	Basse-Normandie	1	1
6	Haute-Normandie	1	1
7	Nord-Pas-de-Calais	$\in (1, 2)$	$\in (0, 1)$
8	Picardie	1	$\in (0, 1)$
9	Alsace	1	$\in (0, 1)$
10	Champagne-Ardenne	1	1
11	Lorraine	1	1
12	Pays de la Loire	1	1
13	Bretagne	1	1
14	Aquitaine	1	1
15	Limousin	1	$\in (0, 1)$
16	Poitou-Charentes	1	1
17	Languedoc-Roussillon	1	1
18	Midi-Pyrénées	1	1
19	Auvergne	1	1
20	Rhône-Alpes	1	1
21	Provence-Alpes-Côte d’Azur	1	1
22	Corse	1	$\in (0, 1)$

Notes : For the first and seventh regions of female data, we conducted the V_2 -test and found that the tests are rejected in the lower tail, and concluded that $d_1 \in (1, 2)$.

6 Concluding remarks

In this paper, statistical tests for examining if a functional time series of interest is integer-integrated or fractionally-integrated are proposed, and these are extended to a sequential testing procedure, which can be used to identify the integration order or an interval including it in a unified framework. Given recent interest in curve-valued time series exhibiting strong persistence, we believe that the proposed methods are useful for practitioners. To illustrate our methodology and its practical relevance, we apply the proposed tests to the Canadian yield curve data and the French sub-national age-specific mortality data. The results overall support the widespread assumption of $I(1)$ for such time series in the literature, although evidence of fractional integration is also observed in certain cases.

There are a few ways in which the methodology presented here can be useful in practice, and we briefly mention two. First, our hypothesis testing procedure can assist in identifying the admissible range of fractional integration orders of functional time series, which, in turn, may help in estimating the fractional order, a crucial component for statistical inference on such time series; see e.g., [Li et al. \(2021, 2023\)](#) concerning the local Whittle estimator of the fractional order for functional time series. Second, to illustrate, we considered yield curve and mortality datasets; however, the developed testing procedure can also be applied to various economic/statistical functional time series, which are often assumed to be integer-integrated, and can provide statistical evidence either in support of or against this assumption in favor of a fractional integration order. Some existing examples in the recent literature include earning densities ([Chang et al., 2016a](#)) and their transformations ([Seo, 2024](#)), age-specific employment rates ([Nielsen et al., 2023a](#)), and high-frequency financial time series, such as cumulative intraday returns ([Gabrys et al., 2010](#)) and intraday records of a volatility index ([Shang et al., 2019](#)).

Acknowledgment

This research was supported by the Australian Research Council Future Fellowship (grant no. FT240100338).

A Appendix

A.1 Preliminary results

We collect some preliminary results, which will be repeatedly used in the subsequent discussion, as follows:

Lemma A.1. *Suppose that Assumption F holds. Then the following hold:*

- (i) $\|\mathcal{K}_T(\mathbf{Y}_0)\|_{\mathcal{L}_{\mathcal{H}}}$ and $\|P_1\mathcal{K}_T(\mathbf{Y}_0)P_1\|_{\mathcal{L}_{\mathcal{H}}}$ are $O_p(T^{2+2d_1})$.
- (ii) The dominant eigenvector \bar{h}_T of $\mathcal{K}_T(\mathbf{Y}_0)$ converges to a random vector \tilde{h} taking values in \mathcal{H}_1 in the sense that $\|\bar{h}_T - \text{sgn}(\langle \bar{h}_T, \tilde{h} \rangle)\bar{h}_T\| \xrightarrow{p} 0$. More specifically, $\|\bar{h}_T - \text{sgn}(\langle \bar{h}_T, \tilde{h} \rangle)\tilde{h}\| = O_p(T^{-(d_1-d_2)})$.

Proof of Lemma A.1. We first show (i). Since $d_1 > -1/2$, under Assumption F, $\sum_{s=1}^t X_s$ is a Type-II fractionally-integrated process of integration order $\tilde{d}_1 = 1 + d_1 > 1/2$. Then from Theorem 3.1 of Li et al. (2023) and similar arguments used in its proof, we deduce that $T^{-2\tilde{d}_1}\langle \mathcal{K}_T(\mathbf{Y}_0)v_1, v_2 \rangle$ converges to a well defined limit. Particularly if $v_1, v_2 \in \text{ran } P_1$, it converges to an almost surely nonzero (random) constant, but if either of v_1 or v_2 is contained in $[\text{ran } P_1]^\perp$, it decays to zero. From these results, we find that $\|\mathcal{K}_T(\mathbf{Y}_0)\|_{\mathcal{L}_{\mathcal{H}}} = O_p(T^{2\tilde{d}_1})$ and $\|P_1\mathcal{K}_T(\mathbf{Y}_0)P_1\|_{\mathcal{L}_{\mathcal{H}}} = O_p(T^{2\tilde{d}_1})$, from which the desire results immediately follow.

Given that $\sum_{s=1}^t P_j X_s$ is a Type-II fractionally-integrated process of integration order $\tilde{d}_j = d_j + 1 > 1/2$, we deduce from similar arguments used in Proposition 2.1 and Theorem 3.1 of Li et al. (2023) that $\sum_{s=1}^t P_j X_s = O_p(T^{\tilde{d}_j-1/2})$ uniformly in $t = 1, \dots, T$, and thus find that $\|P_j\mathcal{K}_T(\mathbf{Y}_0)P_k\|_{\mathcal{L}_{\mathcal{H}}} = O_p(T^{\tilde{d}_j+\tilde{d}_k})$ for all $j, k \in \{1, \dots, K\}$. Combining this result with (i) and the fact that $d_1 > d_2 > \dots > d_K$, we find that

$$\|T^{-2\tilde{d}_1}(\mathcal{K}_T(\mathbf{Y}_0) - P_1\mathcal{K}_T(\mathbf{Y}_0)P_1)\|_{\mathcal{L}_{\mathcal{H}}} = O_p(T^{-(d_1-d_2)}).$$

From nearly identical arguments used in the proof of Proposition 3.2 of Chang et al. (2016b), we further find that (a) the dominant eigenvector \bar{h}_T of $\mathcal{K}_T(\mathbf{Y}_0)$ converges to the dominant eigenvector \tilde{h} of $P_1\mathcal{K}_T P_1$, which is random (and also dependent on T) but takes values in $\mathcal{H}_1 = \text{ran } P_1$ with probability one and (b) $\|\bar{h}_T - \text{sgn}(\langle \bar{h}_T, \tilde{h} \rangle)\tilde{h}\| = O_p(T^{-(d_1-d_2)})$. \square

A.2 Proofs

To reduce notational burden, we hereafter let

$$\widehat{\mathcal{K}}_0 = \mathcal{K}_T(\mathbf{Y}_0), \quad \widehat{\mathcal{K}}_1 = \mathcal{K}_T(\mathbf{Y}_1), \quad \widehat{\Lambda}_0 = \Lambda_T(\mathbf{Y}_0), \quad \widehat{\Lambda}_1 = \Lambda_T(\mathbf{Y}_1).$$

Proof of Proposition 3.1. From Lemma A.1, we know that $T^{-2}\widehat{\mathcal{K}}_0 = T^{-2}P_1\widehat{\mathcal{K}}_0P_1 + o_p(1)$. This implies that

$$T^{-2}\langle \widehat{\mathcal{K}}_0\bar{h}, \bar{h} \rangle = T^{-2}\langle P_1\widehat{\mathcal{K}}_0P_1\bar{h}, \bar{h} \rangle + o_p(1).$$

Since $d_1 = 0$, we find that

$$T^{-2}P_1\widehat{\mathcal{K}}_0P_1 = \frac{1}{T} \sum_{t=1}^T \left(\frac{1}{\sqrt{T}} \sum_{s=1}^t P_1 v_{1,s} \right) \otimes \left(\frac{1}{\sqrt{T}} \sum_{s=1}^t P_1 v_{1,s} \right).$$

When $d_1 = 0$ and the summability condition imposed on $\{\psi_{1,j}\}_{j \geq 0}$ is given, $\{P_1 v_{1,t}\}_{t \geq 1}$ becomes the so-called L^4 - m -approximable sequence (Proposition 2.1 of Hörmann and Kokoszka, 2010). We then deduce from Theorem 1.1 of Berkes et al. (2013) and Theorem 2.1 of Horváth et al. (2014) that $\sup_{0 \leq r \leq 1} \|T^{-1/2} \sum_{t=1}^{\lfloor Tr \rfloor} P_1 v_{1,t} - \mathcal{W}(r)\| = o_p(1)$, where $\mathcal{W}(r)$ is a \mathcal{H} -valued Brownian motion whose covariance operator is given by $P_1 \Lambda_{v_1} P_1$. Moreover, using similar arguments used in the proof of Lemma 1 of Nielsen et al. (2023b), we may also assume that $\|T^{-2}P_1\widehat{\mathcal{K}}_0P_1 - \int \mathcal{W}(r) \otimes \mathcal{W}(r) dr\|_{\mathcal{L}_{\mathcal{H}}} \xrightarrow{p} 0$. Given that $P_1 \Lambda_{v_1} P_1$ is the covariance operator of \mathcal{W} and $\bar{h} \in \mathcal{H}_1$, we find that

$$T^{-2}\langle \widehat{\mathcal{K}}_0\bar{h}, \bar{h} \rangle \xrightarrow{d} \int \langle \mathcal{W}(r), \bar{h} \rangle^2 \stackrel{d}{=} \langle \Lambda_{v_1} \bar{h}, \bar{h} \rangle \int W(r)^2 dr. \quad (\text{A.1})$$

From (A.1), the desired result is established.

Now suppose that $d_1 < 0$. Then, we know from Lemma A.1(i) that $T^{-2}\widehat{\mathcal{K}}_0 = o_p(1)$, from which we find that $T^{-2}\langle \widehat{\mathcal{K}}_0\bar{h}, \bar{h} \rangle \xrightarrow{p} 0$. On the other hand, in the case where $d_1 > 0$, we may deduce from Theorem 3.1 of Li et al. (2023) and Lemma A.1(i) that $T^{-2-2d_1}\langle \widehat{\mathcal{K}}_0\bar{h}, \bar{h} \rangle$ converges to a well defined nondegenerate limit, from which, $T^{-2}\langle \widehat{\mathcal{K}}_0\bar{h}, \bar{h} \rangle \xrightarrow{p} \infty$ is deduced as desired. \square

Proof of Proposition 3.2. Noting that $\widehat{\Lambda}_0$ is self-adjoint nonnegative definite, we deduce from Assumptions F, S and A1 that $q^{-2d_1}\|\widehat{\Lambda}_0\|_{\mathcal{L}_{\mathcal{H}}} = O_p(1)$. Since $\|\bar{h}_T - \text{sgn}(\langle \bar{h}_T, \bar{h} \rangle)\bar{h}\| \xrightarrow{p} 0$ (see Lemma A.1), we find that $\langle \widehat{\Lambda}_0\bar{h}_T, \bar{h}_T \rangle = \langle P_1\widehat{\Lambda}_0P_1\bar{h}, \bar{h} \rangle + o_p(1)$ when $d_1 = 0$. Moreover, under the summability condition on $\{\psi_{1,j}\}_{j \geq 1}$ given by Assumption F, $\{P_1 X_t\}_{t \geq 1}$ is the so-called L^4 - m -approximable sequence and $P_1\widehat{\Lambda}_0P_1$ is its sample long-run covariance. From Theorem 2

of Horváth et al. (2013), it is deduced that $\|P_1\widehat{\Lambda}_0P_1 - \Lambda_{v_1}\|_{\mathcal{L}\mathcal{H}} \xrightarrow{p} 0$. We thus find that

$$\langle \widehat{\Lambda}_0\bar{h}_T, \bar{h}_T \rangle \xrightarrow{p} \langle \Lambda_{v_1}\bar{h}, \bar{h} \rangle. \quad (\text{A.2})$$

On the other hand, we note that

$$\langle T^{-2}\widehat{\mathcal{K}}_0\bar{h}_T, \bar{h}_T \rangle = \langle T^{-2}P_1\widehat{\mathcal{K}}_0P_1\bar{h}, \bar{h} \rangle + o_p(1) \xrightarrow{d} \int \langle \mathcal{W}(r), \bar{h} \rangle^2 \stackrel{d}{=} \langle \Lambda_{v_1}\bar{h}, \bar{h} \rangle \int W(r)^2 dr. \quad (\text{A.3})$$

Thus the desired result when $d_1 = 0$ follows from (A.2) and (A.3).

We then consider the case where $d_1 \neq 0$. V_0 can be written as

$$V_0 = ((T/q)^{-2d_1}q^{-2d_1}\langle \widehat{\Lambda}_0\bar{h}_T, \bar{h}_T \rangle)^{-1}\langle T^{-2-2d_1}\widehat{\mathcal{K}}_0\bar{h}_T, \bar{h}_T \rangle,$$

where $T^{-2-2d_1}\langle \widehat{\mathcal{K}}_0\bar{h}_T, \bar{h}_T \rangle$ can be shown to converge to a well defined nondegenerate limit under the employed assumptions using similar arguments used in Theorem 3.1 of Li et al. (2023). Using the results given in Lemma A.1, we deduce that $q^{-2d_1}\langle \widehat{\Lambda}_0\bar{h}_T, \bar{h}_T \rangle = q^{-2d_1}\langle P_1\widehat{\Lambda}_0P_1\bar{h}, \bar{h} \rangle + O_p(q^{-2d_1}T^{-(d_1-d_2)})$. From this result and Assumptions F, S and A1, it is deduced that $q^{-2d_1}\langle \widehat{\Lambda}_0\bar{h}_T, \bar{h}_T \rangle = q^{-2d_1}\langle P_1\widehat{\Lambda}_0P_1\bar{h}, \bar{h} \rangle + o_p(1) \xrightarrow{p} c_{1,h}$ holds under either of (a) $d_1 > 0$ or (b) $d_1 < 0$ (since $O_p(q^{-2d_1}T^{-(d_1-d_2)}) = o(1)$ under Assumption A1(ii)), where $c_{1,h}$ is a positive random constant whose randomness results from that \bar{h} is random. Combining this result with the fact that $T/q \rightarrow \infty$, we find that $(T/q)^{-2d_1}q^{-2d_1}\langle \widehat{\Lambda}_0\bar{h}_T, \bar{h}_T \rangle \xrightarrow{p} 0$ if $d_1 > 0$ (and thus $V_0 \xrightarrow{p} \infty$) while $(T/q)^{-2d_1}q^{-2d_1}\langle \widehat{\Lambda}_0\bar{h}_T, \bar{h}_T \rangle \xrightarrow{p} \infty$ if $d_1 < 0$ (and thus $V_0 \xrightarrow{p} 0$). \square

Proof of Proposition 3.3. Note that the test statistic can be written as

$$(T^{-2d_1}\langle \widehat{\Lambda}_0\bar{h}_T, \bar{h}_T \rangle)^{-1}\langle T^{-2-2d_1}\widehat{\mathcal{K}}_0\bar{h}_T, \bar{h}_T \rangle. \quad (\text{A.4})$$

As discussed in our proof of Proposition 3.2, $T^{-2-2d_1}\langle \widehat{\mathcal{K}}_0\bar{h}_T, \bar{h}_T \rangle$ converges to a well defined nondegenerate limit. From Assumptions F, S, similar arguments used in the proof of Theorem 3.1 of Li et al. (2023) and the condition (3.5), it can be deduced that $T^{-2d_1}\sum_{t=1}^T X_t \otimes X_t = T^{-2d_1}\sum_{t=1}^T P_1X_t \otimes P_1X_t + o_p(1) = O_p(1)$ if $d_1 > 1/2$ while $(T \log T)^{-1}\sum_{t=1}^T X_t \otimes X_t = (T \log T)^{-1}\sum_{t=1}^T P_1X_t \otimes P_1X_t + o_p(1) = O_p(1)$ if $d_1 = 1/2$. From the construction of $\widehat{\Lambda}_0$, we may also deduce that $\widehat{\Lambda}_0 = O_p(qT^{2d_1-1})$ if $d_1 > 1/2$ and $\widehat{\Lambda}_0 = O_p(q \log T)$ if $d_1 = 1/2$; that is, $\widehat{\Lambda}_0 = O_p(q \max\{T^{2d_1-1}, \log T\})$ for any $d_1 \geq 1/2$. This in turn implies that $T^{-2d_1}\langle \widehat{\Lambda}_0\bar{h}_T, \bar{h}_T \rangle = O_p(\max\{qT^{-1}, qT^{-2d_1} \log T\})$, which converges to zero given that $d_1 \geq 1/2$ and $T/q \rightarrow \infty$ (Assumption A1(ii)) hold. From these findings, we know that the test statistic in (A.4) diverges to infinity. \square

Proof of Proposition 3.4. In the case where $d_1 = 1$, we find from only a slight modification of the arguments used in our proof of Proposition 3.2 that

$$T^{-2} \langle \widehat{\mathcal{K}}_1 \bar{h}_T, \bar{h}_T \rangle \xrightarrow{d} \int \langle \mathcal{W}(r), \bar{h} \rangle^2 \stackrel{d}{=} \langle \Lambda_{v_1} \bar{h}, \bar{h} \rangle \int W(r)^2 dr.$$

Since $\|\bar{h}_T - \text{sgn}(\langle \bar{h}_T, \bar{h} \rangle) \bar{h}\| \xrightarrow{p} 0$, we note that $\langle \widehat{\Lambda}_1 \bar{h}_T, \bar{h}_T \rangle = \langle P_1 \widehat{\Lambda}_1 P_1 \bar{h}, \bar{h} \rangle + o_p(1)$ (see Lemma A.1). Note that $P_1 \widehat{\Lambda}_1 P_1$ is the sample long-run covariance of $\{P_1 \Delta X_t\}_{t \geq 1}$, which is L^4 - m -approximable under the conditions employed in Assumption F, and thus $\|P_1 \widehat{\Lambda}_1 P_1 - \Lambda_{v_1}\|_{\mathcal{L}_{\mathcal{H}}} \xrightarrow{p} 0$ (Horváth et al., 2013, Theorem 2). From these results, the desired result when $d_1 = 1$ is obtained.

We next consider the case where $d_1 \in (1/2, 3/2) \setminus \{1\}$. We write the test statistic as follows:

$$V_1 = ((T/q)^{-2(d_1-1)} q^{-2(d_1-1)} \langle \widehat{\Lambda}_1 \bar{h}_T, \bar{h}_T \rangle)^{-1} \langle T^{-2-2(d_1-1)} \widehat{\mathcal{K}}_1 \bar{h}_T, \bar{h}_T \rangle,$$

where $\langle T^{-2-2(d_1-1)} \widehat{\mathcal{K}}_1 \bar{h}_T, \bar{h}_T \rangle$ converges to a well defined nondegenerate limit as in the previous proofs. We observe that $\|\bar{h}_T - \text{sgn}(\langle \bar{h}_T, \bar{h} \rangle) \bar{h}\| = O_p(T^{-(d_1-d_2)})$, and thus $q^{-2(d_1-1)} \langle \widehat{\Lambda}_1 \bar{h}_T, \bar{h}_T \rangle = q^{-2(d_1-1)} \langle P_1 \Lambda_1 P_1 \bar{h}, \bar{h} \rangle + O_p(q^{-2(d_1-1)} T^{-(d_1-d_2)})$ (see Lemma A.1). From similar arguments used in our proof of Proposition 3.1, we find that $(T/q)^{-2(d_1-1)} q^{-2(d_1-1)} \langle \widehat{\Lambda}_1 \bar{h}_T, \bar{h}_T \rangle \xrightarrow{p} 0$ if $d_1 \in (1, 3/2)$ while $(T/q)^{-2(d_1-1)} q^{-2(d_1-1)} \langle \widehat{\Lambda}_1 \bar{h}_T, \bar{h}_T \rangle \xrightarrow{p} \infty$ if $d_1 \in (1/2, 1)$, from which the desired results follow.

In the case where $d_1 \geq 3/2$, it can be shown that $V_1 \xrightarrow{p} \infty$ using nearly identical arguments used in our proof of Proposition 3.3. \square

Proof of Proposition 3.5. First consider the case with $d_1 \in (-1/2, 1/2)$. Then the test statistic can be written as

$$(T \langle \widehat{\Lambda}_1 \bar{h}_T, \bar{h}_T \rangle)^{-1} \langle T^{-1} \widehat{\mathcal{K}}_1 \bar{h}_T, \bar{h}_T \rangle. \quad (\text{A.5})$$

Let $y_t = \langle X_t, \bar{h}_T \rangle$. We find that

$$\langle T^{-1} \widehat{\mathcal{K}}_1 \bar{h}_T, \bar{h}_T \rangle = \frac{1}{T} \sum_{t=2}^T (y_t - y_1)^2 = \frac{1}{T} \sum_{t=2}^T y_t^2 + y_1^2 + o_p(1) = \langle \widehat{C}_0 \bar{h}_T, \bar{h}_T \rangle + y_1^2 + o_p(1), \quad (\text{A.6})$$

where $\widehat{C}_0 = T^{-1} \sum_{t=2}^T X_t \otimes X_t$. On the other hand, from similar algebra used in the proof of

Theorem 4 of (Cho et al., 2015, pp. 234–5), we find that

$$\begin{aligned} \langle \widehat{\Lambda}_1 \bar{h}_T, \bar{h}_T \rangle &= O_p(qT^{-1}) + \left(\frac{1}{T} \sum_{t=2}^{T-1} (2 - 2w(1, q)) y_t^2 \right) \\ &\quad + \sum_{r=2}^q \left(\frac{1}{T} \sum_{t=r+1}^{T-1} 2(2w(r-1, q) - w(r-2, q) - w(r, q)) y_t y_{t-r+1} \right) \\ &\quad + \left(\frac{1}{T} \sum_{t=2}^{T-1} 2(2w(q, q) - w(q-1, q)) y_t y_{t-q} \right) - \frac{1}{T} \sum_{t=q+2}^T 2w(q, q) y_t y_{t-q-1}. \end{aligned}$$

We observe, as in Cho et al. (2015), that $w(r-1, q) - w(r-2, q) - w(r, q) = 0$ for every $r = 1, \dots, q$ and also find that $2w(q, q) - w(q-1, q) = 0$. From a little bit of algebra, we find that

$$\begin{aligned} q \langle \widehat{\Lambda}_1 \bar{h}_T, \bar{h}_T \rangle &= O_p(q^2 T^{-1}) + \frac{q}{T} \sum_{t=2}^{T-1} (2/(q+1)) y_t^2 - \frac{q}{T} \sum_{t=q+2}^T (2/(q+1)) y_t y_{t-q-1} \\ &= O_p(q^2 T^{-1}) + \frac{2}{T} \sum_{t=2}^{T-1} y_t^2 - \frac{2}{T} \sum_{t=q+2}^T y_t y_{t-q-1} = o_p(1) + 2 \langle \widehat{C}_0 - \widehat{C}_{q+1}, \bar{h}_T, \bar{h}_T \rangle. \end{aligned} \tag{A.7}$$

Under Assumption S, we find that $\langle \widehat{C}_0 - \widehat{C}_{q+1}, \bar{h}_T, \bar{h}_T \rangle = \langle C_0 - C_{q+1}, \bar{h}_T, \bar{h}_T \rangle + o_p(1) = \langle C_0, \bar{h}, \bar{h} \rangle + o_p(1)$ for large enough T and q , where \bar{h} is a unit-norm vector taking values in \mathcal{H}_1 (see Lemma A.1). Since C_0 is positive definite on a finite dimensional subspace \mathcal{H}_1 , we may write $P_1 C_0 P_1 = \sum_{j=1}^{\dim(\mathcal{H}_1)} a_j f_j \otimes f_j$ with $a_{\dim(\mathcal{H}_1)} > 0$. We then observe that

$$\langle C_0, \bar{h}, \bar{h} \rangle \geq \min_{h \in \mathcal{H}_1, \|h\|=1} \langle C_0, h, h \rangle \geq a_{\dim(\mathcal{H}_1)}. \tag{A.8}$$

Combining this result with (A.5), (A.7) and the fact that $T/q \rightarrow \infty$, we find that

$$T \langle \widehat{\Lambda}_1 \bar{h}_T, \bar{h}_T \rangle \xrightarrow{p} \infty \tag{A.9}$$

From (A.5), (A.6) and (A.9), we find the test statistic converges to zero if $-1/2 < d_1 < 1/2$.

To establish the desired result for the case with $d_1 = 1/2$, we write the inverse of the test statistic as follows:

$$V_1^{-1} = d_T \langle \widehat{\Lambda}_1 \bar{h}_T, \bar{h}_T \rangle (d_T \langle T^{-2} \widehat{\mathcal{K}}_1 \bar{h}_T, \bar{h}_T \rangle)^{-1}. \tag{A.10}$$

We first observe that

$$\begin{aligned} \langle (T \log T)^{-1} \widehat{\mathcal{K}}_1 \bar{h}_T, \bar{h}_T \rangle &= \frac{1}{T \log T} \sum_{t=2}^T (y_t - y_1)^2 \\ &= \frac{1}{T \log T} \sum_{t=2}^T y_t^2 - \frac{2}{T \log T} y_1 \sum_{t=2}^T y_t + \frac{1}{\log T} y_1^2. \end{aligned} \quad (\text{A.11})$$

Note that under Assumption **A3** we have $(T \log T)^{-1} \sum_{t=1}^T y_t = O_p(1/\sqrt{\log T})$, and thus, for large enough q and T ,

$$\frac{1}{T \log T} y_1 \sum_{t=2}^T y_t + \frac{1}{\log T} y_1^2 = O_p(1/\sqrt{\log T}), \quad (\text{A.12})$$

and

$$\frac{1}{T \log T} \sum_{t=2}^T y_t^2 = \frac{1}{T \log T} \sum_{t=1}^T \langle X_t, \bar{h}_T \rangle^2 = \frac{1}{\log T} \langle P_1 \widehat{\mathcal{C}}_0 P_1 \bar{h}, \bar{h} \rangle + o_p(1) = O_p(1). \quad (\text{A.13})$$

Combining (A.11)-(A.13), we find that $d_T T^{-2} \widehat{\mathcal{K}}_1 = O_p(d_T T^{-1} \log T)$. Since $\widehat{\Lambda}_1$ is self-adjoint nonnegative definite, we find from Assumptions **A3** that $d_T \|\widehat{\Lambda}_1\|_{\mathcal{L}_{\mathcal{H}}} = O_p(1)$ and thus $d_T \langle \widehat{\Lambda}_1 \bar{h}_T, \bar{h}_T \rangle = d_T \langle P_1 \widehat{\Lambda}_1 P_1 \bar{h}, \bar{h} \rangle + o_p(1)$, which is bounded away from zero for any $h \in \mathcal{H}_1$ with probability approaching one. Since $d_T T^{-1} \log T \rightarrow 0$, we thus find that the inverse statistic considered in (A.10) diverges to infinity, implying that $V_1 \xrightarrow{p} 0$. \square

Proof of Proposition 3.6. Let $\widetilde{\Lambda}_0 = \widetilde{\Lambda}(\mathbf{Y}_0)$ and $\widetilde{\mathcal{K}}_0 = \widetilde{\mathcal{K}}(\mathbf{Y}_0)$ for notational simplicity.

With slight modifications of our proofs of Lemma **A.1**, the following can be shown: (a) $\|\widetilde{\mathcal{K}}_0\|_{\mathcal{L}_{\mathcal{H}}} = O_p(T^{2+2d_1})$, (b) $\|P_1 \widetilde{\mathcal{K}}_0 P_1\|_{\mathcal{L}_{\mathcal{H}}} = O_p(T^{2+2d_1})$, (c) \widetilde{h}_T converges to a random vector \widetilde{h} taking values in \mathcal{H}_1 and (d) $\|\widetilde{h}_T - \text{sgn}(\langle \widetilde{h}_T, \widetilde{h} \rangle) \widetilde{h}\| = O_p(T^{-(d_1-d_2)})$.

First consider the case where $d_1 \in (-1/2, 1/2)$. We know from Assumptions **F** and **S** that, for d_1 in the considered range, $T^{-1} \sum_{s=1}^T P_1 X_s = O_p(T^{d_1-1/2})$, and hence $T^{-1} \sum_{s=1}^T X_s = O_p(T^{d_1-1/2})$. Using this result, we find that

$$\begin{aligned} \widetilde{\Lambda}_0 &= T^{-1} \sum_{s=-q}^q w(s, q) \sum_{t:1 \leq t, t-s \leq T} \{X_t \otimes X_{t-s} - \bar{X}_T \otimes X_{t-s} - X_t \otimes \bar{X}_T + \bar{X}_T \otimes \bar{X}_T\}, \\ &= T^{-1} \sum_{s=-q}^q w(s, q) \sum_{t:1 \leq t, t-s \leq T} X_t \otimes X_{t-s} + O_p(qT^{2d_1-1}). \end{aligned}$$

Thus,

$$q^{-2d_1} \widetilde{\Lambda}_0 = q^{-2d_1} \widehat{\Lambda}_0 + O_p((T/q)^{2d_1-1}). \quad (\text{A.14})$$

Since $T/q \rightarrow \infty$, $O_p((T/q)^{2d_1-1}) = o_p(1)$ for any $d_1 \in (-1/2, 1/2)$, we have $q^{-2d_1}\tilde{\Lambda}_0 = q^{-2d_1}\hat{\Lambda}_0 + o_p(1)$. From similar arguments and the existing asymptotic results used in our proof of Proposition 3.1, we deduce that

$$\langle T^{-2}\tilde{\mathcal{K}}_0\tilde{h}_T, \tilde{h}_T \rangle = \langle T^{-2}P_1\tilde{\mathcal{K}}_0P_1\tilde{h}_T, \tilde{h}_T \rangle + o_p(1) \xrightarrow{d} \langle \Lambda_{v_1}\tilde{h}, \tilde{h} \rangle \int \tilde{W}(r)^2 dr.$$

Then the desired result when $d_1 = 0$ immediately follows from that $\langle \tilde{\Lambda}_0\tilde{h}_T, \tilde{h}_T \rangle = \langle \hat{\Lambda}_0\tilde{h}_T, \tilde{h}_T \rangle + o_p(1)$, as shown above, and that $\langle \hat{\Lambda}_0\tilde{h}_T, \tilde{h}_T \rangle \xrightarrow{p} \langle \Lambda_{v_1}\tilde{h}, \tilde{h} \rangle$ as we earlier established in our proof of Proposition 3.2.

In the case where $d_1 \in (-1/2, 1/2) \setminus \{0\}$, we write the test statistic as follows:

$$((T/q)^{-2d_1}q^{-2d_1}\langle \tilde{\Lambda}_0\tilde{h}_T, \tilde{h}_T \rangle)^{-1}\langle T^{-2-2d_1}\tilde{\mathcal{K}}_0\tilde{h}_T, \tilde{h}_T \rangle.$$

With some obvious modification of our proofs of Proposition 3.2, we find that $\langle T^{-2-2d_1}\tilde{\mathcal{K}}_0\tilde{h}_T, \tilde{h}_T \rangle$ converges to a well defined nondegenerate limit. Moreover, $(T/q)^{-2d_1}q^{-2d_1}\langle \tilde{\Lambda}_0\tilde{h}_T, \tilde{h}_T \rangle = (T/q)^{-2d_1}q^{-2d_1}\langle \hat{\Lambda}_0\tilde{h}_T, \tilde{h}_T \rangle + o_p(1)$ (see (A.14) and note that $T/q \rightarrow \infty$), which converges to zero (resp. diverges to infinity) if $d_1 \in (0, 1/2)$ (resp. $d_1 \in (-1/2, 0)$), as earlier shown in our proof of Proposition 3.2. From these results, the desired results when $d_1 \in (-1/2, 1/2) \setminus \{0\}$ follow.

To show that $\tilde{V}_0 \xrightarrow{p} \infty$ when $d_1 \geq 1/2$, we use nearly identical arguments used in our proof of Proposition 3.3, noting that, under the employed assumptions, $T^{-2d_1}\sum_{t=1}^T(X_t - \bar{X}_T) \otimes (X_t - \bar{X}_T) = O_p(1)$ if $d_1 > 1/2$ while $(T \log T)^{-1}\sum_{t=1}^T(X_t - \bar{X}_T) \otimes (X_t - \bar{X}_T) = O_p(1)$ if $d_1 = 1/2$.

□

A.3 Detailed discussion on Remarks 3.1 and 3.5

A set of primitive sufficient conditions for Assumption A1(i) (and also a similar condition used in Proposition 3.6) can be deduced from Abadir et al. (2009). For notational simplicity, let $x_{j,t} := \langle X_t, h_j \rangle$ for $h_j \in \mathcal{H}_j$ (see Assumption F) and let f_j be the spectral density of $x_{j,t}$. Suppose that

$$f_j(\omega) = a_j|\omega|^{-2d_j} + o(|\omega|^{-2d_j}) \quad \text{as } \omega \rightarrow 0, \quad (\text{A.15})$$

where $a_j > 0$ and $d_j \in (-1/2, 1/2)$ as considered in Assumption S. If $d_j < 0$, the following is further assumed:

$$f_j(\omega) \leq C|\omega|^{-2d_j} \quad \text{for } \omega \in [-\pi, \pi]. \quad (\text{A.16})$$

Define the lag- s autocovariance of $x_{j,t}$ as $\gamma_j(s)$ (i.e., $\gamma_j(s) = Cov(x_{j,t}, x_{j,t+s})$) and let $\kappa(s, u, r)$ be the forth-order cumulant defined by $\kappa_j(s, u, r) = \mathbb{E}[x_t x_{t+s} x_{t+u} x_{t+r}] - [\gamma_j(s)\gamma_j(u-r) + \gamma_j(u)\gamma_j(s-r) + \gamma_j(r)\gamma_j(s-u)]$. As in Assumption M of [Abadir et al. \(2009\)](#), we consider the following conditions:

$$\gamma_j(s) \sim cs^{2d_j-1} \quad \text{if } d_j \neq 0, \quad \sum_{s=-\infty}^{\infty} |\gamma_j(s)| < \infty \quad \text{if } d_j = 0, \quad (\text{A.17})$$

and

$$\sum_{s,u,r=-\infty}^{\infty} |\kappa_j(s, u, r)| \leq C \quad \text{if } d_j < 0, \quad \sup_s \sum_{u,r=-n}^n |\kappa_j(s, u, r)| \leq Cn^{2d_j} \quad \text{if } d_j \geq 0. \quad (\text{A.18})$$

Then from Theorem 2.2 of [Abadir et al. \(2009\)](#), Assumptions **F**, the condition $q = o(T^{1/2})$ and [\(A.15\)–\(A.18\)](#), we find that both $q^{-2d_j} \langle \widehat{\Lambda}_0 h_j, h_j \rangle$ and $q^{-2d_j} \langle \widetilde{\Lambda}_0 h_j, h_j \rangle$ converge to $a_j g(d_j)$, where

$$g(d_j) = \begin{cases} \frac{2\Gamma(1-2d_j) \sin(\pi d_j)}{d_j(1+2d_j)} & \text{if } d_j \neq 0, \\ 2\pi & \text{if } d_j = 0, \end{cases} \quad (\text{A.19})$$

where $\Gamma(\cdot)$ is the gamma function.

References

- Abadir, K. M., Distaso, W., and Giraitis, L. (2009). Two estimators of the long-run variance: Beyond short memory. *Journal of Econometrics*, 150(1):56–70.
- Aue, A., Rice, G., and Sönmez, O. (2018). Detecting and dating structural breaks in functional data without dimension reduction. *Journal of the Royal Statistical Society: Series B*, 80:509–529.
- Baek, C., Kokoszka, P., and Meng, X. (2024). Test of change point versus long-range dependence in functional time series. *Journal of Time Series Analysis*, 45:497–512.
- Bathia, N., Yao, Q., and Ziegelmann, F. (2010). Identifying the finite dimensionality of curve time series. *The Annals of Statistics*, 38:3352–3386.
- Beare, B. K., Seo, J., and Seo, W.-K. (2017). Cointegrated linear processes in Hilbert space. *Journal of Time Series Analysis*, 38(6):1010–1027.

- Beare, B. K. and Seo, W.-K. (2020). Representation of I(1) and I(2) autoregressive Hilbertian processes. *Econometric Theory*, 36(5):773–802.
- Beran, J., Näscher, J., Farquharson, F., Kustermann, M., Kabitz, H.-J., and Walterspacher, S. (2023). On seasonal functional modeling under strong dependence, with applications to mechanically ventilated breathing activity. *Journal of Statistical Planning and Inference*, 222:38–65.
- Berkes, I., Horváth, L., and Rice, G. (2013). Weak invariance principles for sums of dependent random functions. *Stochastic Processes and their Applications*, 123(2):385–403.
- Bonnet, F. (2020). Computations of French lifetables by department, 1901–2014. *Demographic Research*, 42:741–762.
- Bosq, D. (2000). *Linear Processes in Function Spaces*. Springer-Verlag, New York.
- Cairns, A. J., Blake, D., Dowd, K., Coughlan, G. D., Epstein, D., and Khalaf-Allah, M. (2011). Mortality density forecasts: An analysis of six stochastic mortality models. *Insurance: Mathematics and Economics*, 48(3):355–367.
- Chang, Y., Hu, B., and Park, J. Y. (2016a). On the error correction model for functional time series with unit roots. Mimeo, Indiana University, URL: <https://www.econbohu.com/wp-content/uploads/2023/09/fecm.pdf>.
- Chang, Y., Hu, B., and Park, J. Y. (2017). Econometric analysis of persistent functional dynamics. Technical report, Centre interuniversitaire de recherche en économie quantitative. URL: <https://cireqmontreal.com/wp-content/uploads/2017/03/park1.pdf>.
- Chang, Y., Kim, C. S., and Park, J. Y. (2016b). Nonstationarity in time series of state densities. *Journal of Econometrics*, 192(1):152–167.
- Chiou, J.-M. and Müller, H.-G. (2009). Modeling hazard rates as functional data for the analysis of cohort lifetables and mortality forecasting. *Journal of the American Statistical Association: Applications and Case Studies*, 104(486):572–585.
- Cho, C.-K., Amsler, C., and Schmidt, P. (2015). A test of the null of integer integration against the alternative of fractional integration. *Journal of Econometrics*, 187(1):217–237.
- Damon, J. and Guillas, S. (2002). The inclusion of exogenous variables in functional autoregressive ozone forecasting. *Environmetrics*, 13(7):759–774.

- Ferraty, F. and Vieu, P. (2006). *Nonparametric Functional Data Analysis: Theory and Practice*. Springer, New York.
- Franchi, M. and Paruolo, P. (2020). Cointegration in functional autoregressive processes. *Econometric Theory*, 36(5):803–839.
- Gabrys, R., Horváth, L., and Kokoszka, P. (2010). Tests for error correlation in the functional linear model. *Journal of the American Statistical Association: Theory and Methods*, 105(491):1113–1125.
- Hörmann, S. and Kokoszka, P. (2010). Weakly dependent functional data. *The Annals of Statistics*, 38(3):1845–1884.
- Horváth, L., Kokoszka, P., and Reeder, R. (2013). Estimation of the mean of functional time series and a two-sample problem. *Journal of the Royal Statistical Society: Series B*, 75(1):103–122.
- Horváth, L., Kokoszka, P., and Rice, G. (2014). Testing stationarity of functional time series. *Journal of Econometrics*, 179(1):66–82.
- Hsing, T. and Eubank, R. L. (2015). *Theoretical Foundations of Functional Data Analysis, with An Introduction to Linear Operators*. Wiley series in probability and statistics. Wiley, Chichester, West Sussex, United Kingdom.
- Hyndman, R. J. and Ullah, M. S. (2007). Robust forecasting of mortality and fertility rates: A functional data approach. *Computational Statistics & Data Analysis*, 51(10):4942–4956.
- Klepsch, J. and Klüppelberg, C. (2017). An innovations algorithm for the prediction of functional linear processes. *Journal of Multivariate Analysis*, 155:252–271.
- Klepsch, J., Klüppelberg, C., and Wei, T. (2017). Prediction of functional ARMA processes with an application to traffic data. *Econometrics and Statistics*, 1:128–149.
- Kokoszka, P. and Reimherr, M. (2013). Determining the order of the functional autoregressive model. *Journal of Time Series Analysis*, 34(1):116–129.
- Kokoszka, P. and Reimherr, M. (2017). *Introduction to Functional Data Analysis*. CRC Press, Boca Raton.
- Kokoszka, P. and Zhang, X. (2012). Functional prediction of intraday cumulative returns. *Statistical Modelling*, 12:377–398.

- Koner, S. and Staicu, A.-M. (2023). Second-generation functional data. *The Annual Review of Statistics and its Application*, 10:547–572.
- Lee, R. D. and Carter, L. R. (1992). Modeling and forecasting U.S. mortality. *Journal of the American Statistical Association: Applications & Case Studies*, 87(419):659–671.
- Li, D., Robinson, P., and Shang, H. L. (2023). Nonstationary fractionally integrated functional time series. *Bernoulli*, 29(2):1505–1526.
- Li, D., Robinson, P. M., and Shang, H. L. (2020). Long-range dependent curve time series. *Journal of the American Statistical Association: Theory and Methods*, 115(530):957–971.
- Li, D., Robinson, P. M., and Shang, H. L. (2021). Local Whittle estimation of long-range dependence for functional time series. *Journal of Time Series Analysis*, 42(5-6):685–695.
- Liu, M. (1998). Asymptotics of nonstationary fractional integrated series. *Econometric Theory*, 14(5):641–662.
- Mateu, J. and Giraldo, R., editors (2022). *Geostatistical Functional Data Analysis*. Wiley series in probability and statistics. Wiley, Hoboken, NJ.
- Mestre, G., Portela, J., Rice, G., Muñoz San Roque, A., and Alonso, E. (2021). Functional time series model identification and diagnosis by means of auto- and partial autocorrelation analysis. *Computational Statistics & Data Analysis*, 155:107108.
- Nielsen, M. Ø., Seo, W.-K., and Seong, D. (2023a). Inference on common trends in functional time series. Technical report, arXiv. URL: <https://arxiv.org/abs/2312.00590>.
- Nielsen, M. O., Seo, W.-K., and Seong, D. (2023b). Inference on the dimension of the nonstationary subspace in functional time series. *Econometric Theory*, 39(3):443–480.
- Ramsay, J. O. and Silverman, B. W. (2005). *Functional Data Analysis*. Springer, New York, 2nd edition.
- Sabzikar, F. and Kokoszka, P. (2023). Tempered functional time series. *Journal of Time Series Analysis*, 44(3):280–293.
- Salish, N. and Gleim, A. (2019). A moment-based notion of time dependence for functional time series. *Journal of Econometrics*, 212(2):377–392.

- Seo, W.-K. (2023). Cointegration and representation of cointegrated autoregressive processes in Banach spaces. *Econometric Theory*, 39(4):1–52.
- Seo, W.-K. (2024). Functional principal component analysis for cointegrated functional time series. *Journal of Time Series Analysis*, 45(2):320–330.
- Seo, W.-K. and Beare, B. K. (2019). Cointegrated linear processes in Bayes Hilbert space. *Statistics & Probability Letters*, 147:90–95.
- Seo, W.-K. and Shang, H. L. (2024). Fractionally integrated curve time series with cointegration. *Electronic Journal of Statistics*, 18(2):3858–3902.
- Seong, D. and Seo, W.-K. (2025). Functional instrumental variable regression with an application to estimating the impact of immigration on native wages. *Econometric Theory*, in press.
- Shang, H. L. and Hyndman, R. J. (2017). Grouped functional time series forecasting: An application to age-specific mortality rates. *Journal of Computational and Graphical Statistics*, 26(2):330–343.
- Shang, H. L., Yang, Y., and Kearney, F. (2019). Intraday forecasts of a volatility index: Functional time series methods with dynamic updating. *Annals of Operations Research*, 282(1):331–354.
- Sørensen, H., Goldsmith, J., and Sangalli, L. M. (2013). An introduction with medical applications to functional data analysis. *Statistics in Medicine*, 32(30):5222–5240.
- Srivastava, A. and Klassen, E. P. (2016). *Functional and Shape Data Analysis*. Springer, New York.
- Tanaka, K. (2017). *Time Series Analysis: Nonstationary and Noninvertible Distribution Theory*. John Wiley & Sons, Inc., Hoboken, New Jersey, 2nd edition.