

# Deviation Between Team-Optimal Solution and Nash Equilibrium in Flow Assignment Problems

Gehui Xu<sup>1</sup>, Ting Bai<sup>2</sup>, Andreas A. Malikopoulos<sup>2</sup>, and Thomas Parisini<sup>3</sup>

**Abstract**—We investigate the relationship between the team-optimal solution and the Nash equilibrium (NE) to assess the impact of strategy deviation on team performance. As a working use case, we focus on a class of flow assignment problems in which each source node acts as a cooperating decision maker (DM) within a team that minimizes the team cost based on the team-optimal strategy. In practice, some selfish DMs may prioritize their own marginal cost and deviate from NE strategies, thus potentially degrading the overall performance. To quantify this deviation, we explore the deviation bound between the team-optimal solution and the NE in two specific scenarios: (i) when the team-optimal solution is unique and (ii) when multiple solutions do exist. This helps DMs analyze the factors influencing the deviation and adopting the NE strategy within a tolerable range. Furthermore, in the special case of a potential game model, we establish the consistency between the team-optimal solution and the NE. Once the consistency condition is satisfied, the strategy deviation does not alter the total cost, and DMs do not face a strategic trade-off. Finally, we validate our theoretical analysis through some simulation studies.

## I. INTRODUCTION

Routing in networks for data transmission and logistics management has been widely studied across various domains, including wireless communication [1], road traffic [2], and freight transportation [3]. In particular, increasing attention has been given to a class of flow assignment problems [4]–[6]. These problems aim to determine the optimal allocation of traffic across multiple available routes from each source node to the destination node to achieve a desired objective, such as minimizing the total transmission cost of the entire network.

Team theory provides a well-established framework for modeling flow assignment problems from a cooperative perspective [4], [5], [7]. In this framework, source nodes act as decision makers (DMs) within a team organization. Based on available information, such as measured input flow values and network topology, each DM allocates its received flows across multiple paths leading to the destination. While each DM independently decides its routing strategy, they share

the common objective of minimizing the total cost (i.e., the team cost) which is influenced by their joint decisions [4], [5], [7], [8]. The information structure in these problems can be categorized as *static* or *dynamic* [9]. In static team problems, the information available to each DM is not affected by the decisions of others [4], [5]. In contrast, dynamic team problems mean that the information of at least one DM is affected by the decisions of others [10]. The solution to both static and dynamic team problems is characterized by the team-optimal solution [7], [9], [11].

However and inevitably, there may be selfish DMs in a team who prioritize their own costs thus being reluctant to adopt the team-optimal strategy [12], [13]. This is because achieving the team optimum often requires DMs to sacrifice their own benefits, such as tolerating longer transport time in logistics to reduce overall network costs [14] or consuming additional energy in wireless networks to assist data transmission [1]. Additionally, due to limited computing and communication resources, DMs may find it difficult to access complete network information needed to minimize the team’s total cost and instead make decisions based on local information. As a result, the flow assignment problem could also be formulated as a non-cooperative game [12], [13]. In this context, DMs focus on minimizing their individual costs based on the Nash equilibrium (NE) [15], which represents a stable strategy profile where no DM can improve its profit by changing its strategy, given the strategies of others.

As selfish DMs prioritize minimizing their individual costs rather than the team’s overall cost, deviating from the team-optimal strategy to the NE strategy may compromise the overall team performance. However, it is worth mentioning that if the team-optimal solution and the NE are of little difference, then such a strategy deviation will not significantly impact the collective interests of all DMs, as the total cost increase remains acceptable. Moreover, in certain special cases where the NE coincides with the team-optimal solution, regardless of whether the DMs deviate in their strategies, the total costs remain unchanged.

Therefore, the main objective of this preliminary paper is to explore the relationship between the team-optimal solutions and the NE to evaluate how strategy deviations impact the team’s overall cost. For the sake of clarity, we focus on a simple flow assignment problem and formulate it as a static team problem and a non-cooperative game problem, respectively. We first provide an upper bound on the deviation between the team-optimal solution and the NE in two cases: (i) when the team-optimal solution is unique and (ii) when multiple solutions exist. Next, we establish the consistency between the

\*This work was supported in part by NSF under Grants CNS-2401007, CMMI-2348381, IIS-2415478, and in part by MathWorks.

<sup>1</sup>G. Xu is with the Department of Electrical and Electronic Engineering, Imperial College London, SW7 2AZ London, U.K. E-mail: g.xu@imperial.ac.uk

<sup>2</sup>T. Bai and A. A. Malikopoulos are with the Information and Decision Science Lab, School of Civil & Environmental Engineering, Cornell University, Ithaca, New York, U.S.A. E-mails: {tingbai, amaliko}@cornell.edu

<sup>3</sup>Thomas Parisini is with the Dept. of Electrical and Electronic Engineering, Imperial College London, London SW7 2AZ, UK, and also with the Dept. of Electronic Systems, Aalborg University, Denmark and with the Dept. of Engineering and Architecture, University of Trieste, Italy. E-mail: t.parisini@imperial.ac.uk

team-optimal solution and the NE in a specific non-cooperative game setting, namely, a potential game. Finally, we present simulation studies on two logistics transportation examples to validate our theoretical analysis.

## II. FLOW ASSIGNMENT PROBLEM

In this section, we begin with a routing example in a logistics transportation system. Next, we present the mathematical formulation of the flow assignment problem, modeling it both as a static cooperative team problem and a non-cooperative game.

### A. A freight logistics problem

As a motivating example, consider a freight logistics problem where multiple independent shippers share a transportation network [3]. The shippers need to transport goods to a common destination (e.g., a warehouse or a distribution center) through multiple available transportation paths, which are connected by multiple edges. The transportation cost of each edge depends on the total volume of goods transported by all shippers using that edge. We assume all shippers belong to the same carrier and collectively minimize the total transportation cost of the carrier.

### B. Cooperative team decision-making problem

Next, we introduce the mathematical formulation of the flow assignment problem in a general form. Team theory [4], [5], [7], [9] is a mathematical formalism that can be used to model this problem. It was developed to provide a rigorous mathematical framework for cooperating members in which all members have the same objective, yet different information. In this context, we consider a “team” consisting of a number of members that cooperate to achieve a common objective. The underlying structure to model a team decision problem consists of (1) a number of members of the team; (2) the decisions of each member; (3) the information available to each member, which may be different; (4) an global objective, which is the same for all members; and (5) the existence, or not, of communication between team members.

In this problem, the routing nodes have decision-making capabilities and are considered cooperating members of a team organization. Specifically, we model the network as a directed graph  $\mathcal{G}(\mathcal{V}, \mathcal{E})$ , where  $\mathcal{V} = \{1, \dots, M\}$ ,  $M \in \mathbb{N}$ , is the node set and  $\mathcal{E} \subseteq \mathcal{V} \times \mathcal{V}$  is the edge set. There are no cycles in the network graph. Moreover, there are  $N$  ( $N < M$ ,  $N \in \mathbb{N}$ ) source nodes and a single destination node. Each source node corresponds to a shipper. Let  $\mathcal{N} = \{1, \dots, N\}$  denote the set of source nodes. The routing decisions are made only at the source nodes by the DMs. For illustration, in Fig. 1 we show a flow assignment problem with two DMs.

For each DM  $i \in \mathcal{N}$ , let  $r_i \in \mathbb{R}$  be the flow entering DM  $i$ , which represents the amount of goods received by shipper  $i$  from an external supplier. Each DM  $i$  has  $P_i \in \mathbb{N}$  available paths to the destination, connected by multiple edges. Let  $L = |\mathcal{E}| \in \mathbb{N}$  be the number of edges in the network. We introduce an indicator function  $\sigma(l, k, i)$ , which takes the value




Fig. 1. Flow assignment problem with two source nodes. Source nodes 1 and 2 in blue represent DM<sub>1</sub> and DM<sub>2</sub>. Node 3 in gray serves as a transmitting node, and node 4 in red is the destination. The parameters  $\rho_1, \dots, \rho_5$  represent the weighting coefficients.

of 1 if edge  $l$  (for  $l = 1, \dots, L$ ) belongs to  $k$ -th path (for  $k = 1, \dots, P_i$ ) from DM  $i$  to the destination, and 0 otherwise. Each DM  $i$  makes a routing decision  $u_i = \text{col}\{u_i^k\}_{k=1}^{P_i} \in \mathbb{R}^{P_i}$  to allocate the external flow  $r_i$  among the  $P_i$  available paths. Here,  $u_i^k \in \mathbb{R}$  denotes the portion of external flow  $r_i$  assigned to path  $k$ , representing the amount of goods allocated to transport route  $k$ . Note that the value of  $u_i^k$  is the same for all links that belong to path  $k$ . The local strategy set of DM  $i$  is defined by

$$\Xi_i = \left\{ u_i \in \mathbb{R}^{P_i} : \sum_{k=1}^{P_i} u_i^k = r_i, u_i^k \geq 0, \text{ for } k = 1, \dots, P_i \right\}.$$

Let  $\Xi = \prod_{i=1}^N \Xi_i \subseteq \mathbb{R}^{\sum_{i=1}^N P_i}$  be the joint strategy set for all DMs,  $\mathbf{u} = \text{col}\{u_i\}_{i=1}^N \in \Xi$  be the strategy profile for all DMs, and  $\mathbf{u}_{-i} = \text{col}\{u_1, \dots, u_{i-1}, u_{i+1}, \dots, u_N\}$  be the strategy profile for all DMs except DM  $i$ .

All DMs share a common objective of minimizing the total cost of the entire network. The cost is determined based on edge costs, which in turn depend on the total flow allocated by all DMs. Let

$$z_l(\mathbf{u}) = \sum_{i=1}^N \sum_{k=1}^{P_i} u_i^k \sigma(l, k, i)$$

be the total flow function on edge  $l$ , and let  $f_l(z_l)$  be the transmission cost function on edge  $l$ , which is commonly considered to be quadratic, i.e.,

$$f_l(z_l(\mathbf{u})) = z_l^2(\mathbf{u}) = \left[ \sum_{i=1}^N \sum_{k=1}^{P_i} u_i^k \sigma(l, k, i) \right]^2.$$

Consider  $\rho_l > 0$  as the weighting coefficient related to edge  $l$ , representing the transportation cost per unit of goods on that edge. Then the total cost function of the entire network is

$$\mathcal{C}(\mathbf{u}) = \sum_{l=1}^L \rho_l \cdot f_l(z_l(\mathbf{u})). \quad (1)$$

In this context, each DM independently determines its routing strategy  $u_i$  and collaborates in minimizing  $\mathcal{C}$ . The team decision-making problem is formulated as follows:

$$\begin{aligned} & \min_{u_1, \dots, u_N} \mathcal{C}(u_1, \dots, u_N) \\ & \text{s. t. } u_i \in \Xi_i, i \in \mathcal{N}. \end{aligned} \quad (2)$$

Since  $r_i$  for  $i \in \mathcal{N}$  are independent parameters, each DM receives this information and makes decisions independently.

This indicates that the information received by each DM is not influenced by the decisions of others [4], [5], [11]. Thus, the team problem under consideration follows a static information structure with independent available information [11]. Referring to [4], [5], we have the following assumption for the decision-making of DMs.

**Assumption 1** For each DM  $i \in \mathcal{N}$ , the routing decision  $u_i$  is determined based on the complete knowledge of the network topology  $\mathcal{G}$  and the input flow  $r_i$ .

The corresponding outcome is described by a team-optimal solution [4], [5], [11], formally defined next.

**Definition 1 (Team-optimal solution)** A strategy profile  $\mathbf{u}^* = \text{col}\{u_1^*, u_2^*, \dots, u_N^*\} \in \Xi$  is a team-optimal solution if

$$\mathcal{C}(\mathbf{u}^*) \leq \mathcal{C}(\mathbf{u}), \forall \mathbf{u} \in \Xi.$$

The team decision-making problem can be analyzed from the viewpoint of game theory. All DMs, who are considered players in a game, share the same preferences and seek to minimize the shared cost function  $\mathcal{C}$  [15]. For the sake of clarity, we will refer to “player” as “DM” throughout the remainder of our exposition. A fundamental distinction exists between the team-optimal solution and the widely recognized concept of NE [16]. Namely, a strategy profile  $\mathbf{u}^*$  is called an NE if it satisfies the following conditions

$$\begin{aligned} \mathcal{C}(u_1^*, \mathbf{u}_{-1}^*) &\leq \mathcal{C}(u_1, \mathbf{u}_{-1}^*), \forall u_1 \in \Xi_1, \\ &\vdots \\ \mathcal{C}(u_N^*, \mathbf{u}_{-N}^*) &\leq \mathcal{C}(u_N, \mathbf{u}_{-N}^*), \forall u_N \in \Xi_N. \end{aligned}$$

NE characterizes an outcome where no DM has an incentive to unilaterally deviate from their chosen strategy, provided that the strategies of all others remain unchanged. This equilibrium concept is also referred to as person-by-person optimality in the team problem [11]. Clearly, a team-optimal strategy is always an NE, but the converse does not necessarily hold. In the following sections, we introduce distinct notations to further elucidate the differences between the team-optimal solution and the NE.

### C. Non-cooperative game

In practice, DMs may prioritize minimizing their own costs due to self-interest rather than cooperatively optimizing the overall network performance. Next, we reformulate the flow assignment problem as a non-cooperative game.

Let  $\{\mathcal{N}, \{\Xi_i\}_{i \in \mathcal{N}}, \{\mathcal{C}_i\}_{i \in \mathcal{N}}\}$  be a non-cooperative game, where  $\mathcal{C}_i(u_i, \mathbf{u}_{-i}) : \mathbb{R}^{\sum_{i=1}^N P_i} \rightarrow \mathbb{R}$  denotes DM  $i$ 's payoff function, representing its selfish interest; it is associated with the transmission cost of its allocated flow across the network. Let  $g_l(z_l(\mathbf{u}))$  be the unit flow cost function, which is determined by all DMs using edge  $l$ . Typically,  $g_l(z_l(\mathbf{u}))$  is considered to be in a linear form [12], i.e.,

$$g_l(z_l(\mathbf{u})) = a_l + b_l z_l(\mathbf{u}) = a_l + b_l \sum_{i=1}^N \sum_{k=1}^{P_i} u_i^k \sigma(l, k, i),$$

where  $a_l \geq 0$  is the fixed cost and  $b_l > 0$  is the congestion cost. In a non-cooperative setting, each DM only cares about its own proportional share on edge  $l$ , given by  $(\sum_{k=1}^{P_i} u_i^k \sigma(l, k, i)) \cdot g_l(z_l(\mathbf{u}))$ . Then the individual payoff  $\mathcal{C}_i(u_i, \mathbf{u}_{-i})$  for each DM  $i$  is

$$\mathcal{C}_i(u_i, \mathbf{u}_{-i}) = \sum_{l=1}^L \rho_l \omega(l, i) \left[ \sum_{k=1}^{P_i} u_i^k \sigma(l, k, i) \right] g_l(z_l(\mathbf{u})), \quad (3)$$

where  $\omega(l, i)$  is an indicator function which simply gives 1 if edge  $l$  related with DM  $i$ , and is 0 otherwise. The individual objective of each DM is to have a minimal personal transmission cost. Given  $\mathbf{u}_{-i}$ , DM  $i$  aims to solve

$$\min_{u_i \in \Xi_i} \mathcal{C}_i(u_i, \mathbf{u}_{-i}). \quad (4)$$

The outcome of this non-cooperative game is characterized by the NE [16].

**Definition 2 (NE)** A profile  $\mathbf{u}^\diamond = \text{col}\{u_1^\diamond, \dots, u_N^\diamond\} \in \Xi$  is said to be an NE of game  $\{\mathcal{N}, \{\Xi_i\}_{i \in \mathcal{N}}, \{\mathcal{C}_i\}_{i \in \mathcal{N}}\}$  if

$$\mathcal{C}_i(u_i^\diamond, \mathbf{u}_{-i}^\diamond) \leq \mathcal{C}_i(u_i, \mathbf{u}_{-i}^\diamond), \forall i \in \mathcal{N}, \forall u_i \in \Xi_i.$$

Define  $G(\mathbf{u}) = \nabla_{\mathbf{u}} \mathcal{C}(\mathbf{u}) = \text{col}\{\nabla_{u_1} \mathcal{C}(\cdot, u_{-1}), \dots, \nabla_{u_N} \mathcal{C}(\cdot, u_{-N})\}$  as the gradient map of (1) and  $F(\mathbf{u}) = \text{col}\{\nabla_{u_1} \mathcal{C}_1(\cdot, u_{-1}), \dots, \nabla_{u_N} \mathcal{C}_N(\cdot, u_{-N})\}$  as the pseudo-gradient map of (3). We impose the following assumption.

### Assumption 2

- (1) The function  $\mathcal{C}(\mathbf{u})$  is convex and continuously differentiable in  $\mathbf{u}$ . For each DM  $i \in \mathcal{N}$ , the function  $\mathcal{C}_i(u_i, \mathbf{u}_{-i})$  is convex and continuously differentiable in  $u_i$ .
- (2) The operators  $G(\mathbf{u})$  and  $F(\mathbf{u})$  are monotone.

Assumption 2 can guarantee the existence of a team-optimal solution and an NE by using variational inequality [17]. Also, Assumption 2 does not restrict the uniqueness of the team-optimal solution and NE.

The following results verify the existence of a team-optimal solution and an NE [17].

**Lemma 1** Given Assumption 2, there exists a team-optimal solution  $\mathbf{u}^*$  in the team decision-making problem.

**Lemma 2** Given Assumption 2, there exists an NE  $\mathbf{u}^\diamond$  in game  $\{\mathcal{N}, \{\Xi_i\}_{i \in \mathcal{N}}, \{\mathcal{C}_i\}_{i \in \mathcal{N}}\}$ .

In prior efforts [4], [5], DMs adopt team-optimal strategies based on Assumption 2. However, this assumption may not always hold in practice. DMs may prioritize their own interests, being reluctant to compromise their performance or share information. Consequently, some DMs may deviate from the NE strategy, adopting the NE strategy to minimize their own costs, which could potentially reduce overall team performance. Such a strategy deviation may compromise the collective performance of the team. For example, in the freight transportation case, some shippers select the cheapest route for themselves, resulting in congestion and increased costs for others, which causes the total system cost to exceed that of the team-optimal solution.

Nevertheless, if the team-optimal solution and the NE are nearly identical, such a strategy deviation has minimal

impact. In special cases where they coincide, total costs remain unchanged regardless of individual strategy shifts.

To this end, we seek to address the following questions in this paper:

- Does there exist a deviation bound between the team-optimal solution and the NE?
- In which scenario is the team-optimal solution and the NE consistent?

### III. DEVIATION BOUND BETWEEN TEAM-OPTIMAL SOLUTION AND NE

In practice, team-optimal solutions and the NE may not be the same. Next, we explore the deviation bound between a unique team-optimal solution and a unique NE. Based on the convexity and continuity condition in Assumption 2, the team-optimal solution and the NE are equivalent to the first-order stationary points of their respective problems. This allows us to study their deviation based on gradient information. Let us denote the gradient difference

$$\theta(\mathbf{u}) = G(\mathbf{u}) - F(\mathbf{u})$$

as a perturbation term. Additionally, we define the compact set  $\Theta = \{\mathbf{u} \in \Xi : \|\mathbf{u} - \mathbf{u}^*\| < q\}$ , where  $q$  is a positive constant. The following result provides a deviation bound. The proof is presented in Appendix A.

**Theorem 1** *Given Assumption 2, suppose that there exist positive constants  $\kappa_1, \kappa_2, \nu_1$  and  $\nu_2$  such that  $G(\mathbf{u})$  is  $\kappa_1$ -strongly monotone and  $\nu_1$ -Lipschitz continuous, while  $F(\mathbf{u})$  is  $\kappa_2$ -strongly monotone and  $\nu_2$ -Lipschitz continuous. If the perturbation term  $\theta(\mathbf{u})$  satisfies*

$$\|\theta(\mathbf{u})\| \leq \delta < \omega \cdot s \cdot q$$

for all  $\mathbf{u} \in \Theta$  and some constant  $0 < s < 1$ , then

$$\|\mathbf{u}^* - \mathbf{u}^\diamond\| \leq \frac{\gamma\delta}{\omega s},$$

where

$$0 < \gamma < \min\left\{\frac{2\kappa_1}{\nu_1^2}, \frac{2\kappa_2}{\nu_2^2}\right\}, \quad \omega = \frac{2\gamma\kappa_1 - \gamma^2\nu_1^2}{2 + \gamma\nu_1}.$$

By Theorem 1, since  $\kappa_1, \kappa_2, \nu_1, \nu_2$  and  $s$  are constants, the deviation  $\|\mathbf{u}^* - \mathbf{u}^\diamond\|$  is influenced by a term proportional to  $\|\theta(\mathbf{u})\|$ . Since  $\Xi$  is compact and  $\theta(\mathbf{u})$  is continuous, there exists a constant  $\delta$  such that  $\theta(\mathbf{u}) \leq \delta$ . Given any such  $\delta$ , we can always select  $q$  large enough to ensure  $\|\theta(\mathbf{u})\| \leq \delta < \omega s q$ , ensuring that the deviation  $\|\mathbf{u}^* - \mathbf{u}^\diamond\|$  remains bounded. Therefore, the deviations from the team-optimal solution do not grow unbounded under small perturbations. If  $\theta(\mathbf{u})$  is sufficiently small, then  $\|\mathbf{u}^* - \mathbf{u}^\diamond\|$  remains small. This implies that DMs can tolerably adopt the NE strategy, as the resulting cost difference remains within an acceptable range.

Next, we relax the assumptions on  $G(\mathbf{u})$  and  $F(\mathbf{u})$  in Theorem 1 and consider the case where the solution is not unique. To quantify the deviation between solution sets, we employ the Hausdorff metric [18], which is one of the most common metrics for measuring the distance between two sets. The Hausdorff metric considers the greatest distance that a

point in one set needs to travel to reach the other [18]. The Hausdorff metric of two sets  $A, B \subseteq \mathbb{R}^{\sum_{i=1}^N P_i}$  is defined as

$$H(A, B) = \max\left\{\sup_{a \in A} \inf_{b \in B} \|a - b\|, \sup_{b \in B} \inf_{a \in A} \|a - b\|\right\}.$$

Let  $\Upsilon_{TO}$  be the set of team-optimal strategy profile  $\mathbf{u}^*$  and  $\Upsilon_{NE}$  be the set of NE strategy profile  $\mathbf{u}^\diamond$ . Let  $\Lambda_{TO}$  be the image set of  $F(\mathbf{u}^*)$  and  $\Lambda_{NE}$  be the image set of  $F(\mathbf{u}^\diamond)$ . We characterize the deviation of the NE set from the team-optimal solution set based on differences in values within the gradient space. The proof is in Appendix B.

**Theorem 2** *Given Assumption 2, suppose that there exists a constant  $\kappa_2 > 0$  such that  $F(\mathbf{u})$  is  $\kappa_2$ -strongly monotone. Then if  $H(\Lambda_{TO}, \Lambda_{NE}) < \eta$ ,*

$$H(\Upsilon_{TO}, \Upsilon_{NE}) < \kappa_2 \eta.$$

In Theorem 2, the upper bound on the Hausdorff metric  $H(\Upsilon_{TO}, \Upsilon_{NE})$  is mainly affected by the Hausdorff metric  $H(\Lambda_{TO}, \Lambda_{NE})$ . Particularly, when the NE  $\mathbf{u}^\diamond$  is an interior point of the strategy set  $\Xi$ , the set  $\Lambda_{NE}$  is the zero set, then the Hausdorff metric on the gradient space becomes  $H(\Lambda_{TO}, \Lambda_{NE}) = \sup_{\mathbf{u}^* \in \Lambda_{TO}} \|F(\mathbf{u}^*)\|$ . Regarding this maximal value as a perturbation, a lower perturbation yields a lower bound. Moreover, the Hausdorff metric characterizes the maximum deviation between two sets in the worst case. If the bound is sufficiently small, then even at the farthest point, the NE set remains close to the team-optimal solution set.

### IV. COINCIDENCE OF TEAM-OPTIMAL SOLUTION AND NE

In this section, we consider a special type of non-cooperative game, namely the potential game, and investigate the coincidence relationship between the team-optimal solution and the NE.

We define the potential game  $\{\mathcal{N}, \{\Xi_i\}_{i \in \mathcal{N}}, \{\tilde{C}_i\}_{i \in \mathcal{N}}\}$ . The first two tuples are the same as the game introduced in Section II-C, but the payoff function  $\tilde{C}_i(u_i, \mathbf{u}_{-i})$  of each DM differs. Here,  $\tilde{C}_i(u_i, \mathbf{u}_{-i})$  is defined as each DM's marginal contribution to the overall transmission cost, i.e.,

$$\tilde{C}_i(u_i, \mathbf{u}_{-i}) = \sum_{l=1}^L \rho_l \omega(l, i) \left( \sum_{k=1}^{P_i} f_l(z_l(u_i, \mathbf{u}_{-i})) \right). \quad (5)$$

Then, we introduce the concept of potential game [19].

**Definition 3 (Potential game)** *A game  $\{\mathcal{N}, \{\Xi_i\}_{i \in \mathcal{N}}, \{\tilde{C}_i\}_{i \in \mathcal{N}}\}$  is a potential game if there exists a potential function  $\Phi$  such that*

$$\Phi(u'_i, \mathbf{u}_{-i}) - \Phi(u_i, \mathbf{u}_{-i}) = \tilde{C}_i(u'_i, \mathbf{u}_{-i}) - \tilde{C}_i(u_i, \mathbf{u}_{-i}), \quad (6)$$

for every  $\mathbf{u} \in \Xi$ ,  $i \in \mathcal{N}$ , and unilateral deviation  $u'_i \in \Xi_i$ .

It follows from Definition 3 that any unilateral deviation from a strategy profile results in the same change in both individual payoffs and a unified potential function.

The following result verifies that  $\mathcal{C}(\mathbf{u})$  in (1) is a potential function. The proof is presented in Appendix C.

**Proposition 1** *Given the function  $\mathcal{C}$  and payoffs  $\tilde{C}_i$  for each DM  $i \in \mathcal{N}$ , the game  $\{\mathcal{N}, \{\Xi_i\}_{i \in \mathcal{N}}, \{\tilde{C}_i\}_{i \in \mathcal{N}}\}$  is a potential game.*

The following result provides a consistent relationship between the team-optimal solution and the NE. The proof is presented in Appendix D.

**Proposition 2** *Under potential game, if  $\mathbf{u}^*$  is a team-optimal solution minimizing  $\mathcal{C}$ , then  $\mathbf{u}^*$  is an NE  $\mathbf{u}^\diamond$  of the game  $\{\mathcal{N}, \{\Xi_i\}_{i \in \mathcal{N}}, \{\tilde{\mathcal{C}}_i\}_{i \in \mathcal{N}}\}$ . Moreover, if Assumption 2 holds, then  $\mathbf{u}^\diamond$  is also a team-optimal solution  $\mathbf{u}^*$ .*

Proposition 2 establishes the consistency between the team-optimal solution and the NE. When the condition in Proposition 2 is satisfied, the Hausdorff metric between the team-optimal solution set and the NE set is zero. This indicates that the cost for the whole team remains unchanged. Even if some DMs choose the NE strategy, the remaining DMs can still follow the team-optimal strategies.

## V. SIMULATIONS STUDIES

In this section, we illustrate the relationship between the team-optimal solution and the NE in two logistics transportation examples.




Fig. 2. Logistics network with four DMs.

Consider a logistics transportation example with four shippers acting as DMs. As shown in Fig. 2, each shipper has two available paths. Shipper 1 can choose between paths 1-8 and 1-7-8. Shipper 2 can choose between paths 2-8 and 2-5-8. Shipper 3 can choose between paths 3-5-8 and 3-6-8. Shipper 4 can choose between paths 4-6-8 and 4-7-8. Each shipper needs to transport a specific amount of goods. According to the general load capacity limits, take  $r_1 = 10$  tons,  $r_2 = 15$  tons,  $r_3 = 8$  tons, and  $r_4 = 12$  tons. The transportation network consists of 11 edges with congestion cost coefficients  $\rho_l$ , representing the transportation cost per unit flow, measured in  $\$/\text{ton}$ . The specific values of  $\rho_l$  are listed in Case 1 of Table I, where the short-distance transportation costs ( $\rho_4, \rho_5, \rho_6, \rho_7$ ) are generally lower than the long-distance transportation costs ( $\rho_1, \rho_2, \rho_3, \rho_8, \rho_9, \rho_{10}, \rho_{11}$ ). Besides, we set the transmission cost  $f_l(z_l)$  on edge  $l$  as quadratic, and the cost per unit traffic  $g_l(z_l)$  as linear, where  $a_l = 0$  and  $b_l = 1$ .

We first focus on the deviation bound between the unique team-optimal solution and the unique NE. From (1) and (3), the strong monotonicity constants  $\kappa_1, \kappa_2$  and Lipschitz continuity constants  $\nu_1, \nu_2$  of gradients  $G(\mathbf{u})$  and  $F(\mathbf{u})$  are mainly influenced by  $\rho_l$  for  $l = 1, \dots, 11$ . This further affects the deviation bound  $\frac{\gamma\delta}{\omega_s}$  in Theorem 1. Based on Case 1, we randomly select two weight coefficients,  $\rho_1$  and




Fig. 3. Deviation between the team-optimal solution and NE in different settings.

$\rho_7$ , representing the long-distance path cost and the short-distance path cost, respectively. We then observe how changes in these parameters influence the deviation bound in Fig. 3. The horizontal axis represents the value of  $\delta$  in Theorem 1. The vertical axis represents the difference between the total transmission cost  $\mathcal{C}(\mathbf{u}^*)$  and  $\mathcal{C}(\mathbf{u}^\diamond)$ , corresponding to the value of  $\frac{\gamma\delta}{\omega_s}$ . Fig. 3 shows that as the long-distance path cost  $\rho_1$  increases, the performance gap in total cost decreases. In contrast, an increase in the short-distance path cost  $\rho_7$  enlarges this gap. Moreover, Fig. 3 demonstrates that a smaller value of  $\delta$  leads to a reduced cost difference. This implies that when the NE is close to the team-optimal solution, the overall operational efficiency of the transportation network remains nearly unchanged.




Fig. 4. Deviation between team-optimal solution and NE in different scenarios.

Furthermore, in Table I, we randomly generate four additional scenarios with different values of  $\rho_1, \dots, \rho_{11}$ . The short-distance transportation costs ( $\rho_4, \rho_5, \rho_6, \rho_7$ ) are set between 5 and 10  $\$/\text{ton}^2$ , while the long-distance transportation costs ( $\rho_1, \rho_2, \rho_3, \rho_8, \rho_9, \rho_{10}, \rho_{11}$ ) are set between 8 and 20  $\$/\text{ton}^2$ . Fig. 4 presents the bound  $\frac{\gamma\delta}{\omega_s}$  for these cases. In Fig. 4, the deviation  $\|\mathbf{u}^* - \mathbf{u}^\diamond\|$  remains consistently within the corresponding upper bound, and the difference between them is relatively small.

Next, let us consider another transportation network, shown in Fig. 5. Based on this network, we study the relationship between the team-optimal solution set and the NE set in both the non-cooperative game and the potential game. In this example, DM<sub>1</sub> has three available paths, DM<sub>2</sub> has two, DM<sub>3</sub> has three, and DM<sub>4</sub> has two. Let  $\rho_1 = 10, \rho_2 = 8, \rho_3 = 6, \rho_4 = 7, \rho_5 = 8, \rho_6 = 10, \rho_7 = 5, \rho_8 = 8, \rho_9 = 7, \rho_{10} = 6,$

TABLE I  
FIVE DIFFERENT TRAFFIC SCENARIOS.

	$\rho_1$	$\rho_2$	$\rho_3$	$\rho_4$	$\rho_5$	$\rho_6$	$\rho_7$	$\rho_8$	$\rho_9$	$\rho_{10}$	$\rho_{11}$
Case 1	10	11	15	8	6	7	5	13	12	11	9
Case 2	12	9	14	7	6	5	6	14	10	8	12
Case 3	18	12	17	5	7	6	8	10	8	9	10
Case 4	14	10	16	6	5	8	7	12	9	10	11
Case 5	16	13	12	6	5	6	7	15	11	8	13




Fig. 5. Another logistics network with four DMs.




Fig. 6. Team-optimal solution and NE in different settings. Any team-optimal solution is an NE in the potential game.

$\rho_{11} = 10$ . We compute the team-optimal solution via (2), the NE in the non-cooperative game via (3), and the NE in the potential game via (5). To visualize the strategy space of all DMs, we map it to the strategy space of  $u_2^1, u_1^2, u_1^3$ . As shown in Fig. 6, the strategy profile  $\mathbf{u}^*$  that satisfies the constraints  $\mathbb{E}$  and the condition  $u_2^{1*} + u_1^{2*} + u_1^{3*} = 11.7931$  is a team-optimal solution. In Fig. 6(a), a team-optimal solution is not an NE in the non-cooperative game setting. In Fig. 6(b), any team-optimal solution is an NE in the potential game setting.

In this example, the individual cost for DM<sub>2</sub> under the non-cooperative game setting with the NE strategy is lower than that under the team-optimal solution, which motivates DM<sub>2</sub> to deviate. Moreover, this deviation has the greatest impact on the difference between the team-optimal solution set and the NE set. Therefore, we consider the following three strategy profiles: (i) all DMs choose the team-optimal strategy; (ii) DM<sub>2</sub> takes the NE strategy, while the others adopt the team-optimal strategies; (iii) all DMs choose the NE strategies. Fig. 7 investigates how different strategy profiles affect the total cost  $\mathcal{C}$ . In a general non-cooperative game, the misalignment between individual objectives and the collective goal leads to a discrepancy between the NE and the team-optimal solution. The ideal scenario occurs when all




Fig. 7. Total cost  $\mathcal{C}$  corresponding to different equilibria in different game settings.

DMs adopt the team-optimal strategies. However, when DM<sub>2</sub> deviates from the NE strategy, the resulting strategy profile does not form an equilibrium, increasing the total cost. When all DMs adopt the NE strategy, the cost decreases relatively. This suggests that the remaining DMs need to make a trade-off between the team-optimal strategy and the NE strategy. In the potential game setting, individual interests align with the collective objective, making the NE identical to the team-optimal solution. Therefore, even if DM<sub>2</sub> deviates, the total cost remains unchanged, allowing the remaining DMs to still adopt the team-optimal strategies.

## VI. CONCLUDING REMARKS

In this paper, we addressed a flow assignment problem and investigated the relationship between the team-optimal solutions and the NE to evaluate the impact of unilateral deviations on team performance. We studied the upper bound on the deviation between the team-optimal solution and the NE in cases where the team-optimal solution is unique, as well as in cases where multiple solutions exist. Furthermore, we established the consistency between the team-optimal solution and the NE in a potential game. Finally, we presented the applicability of our results in logistics examples.

A potential direction of future research could consider investigating these questions in the case of random flow assignments.

## APPENDIX

### A. Proof of Theorem 1

Before delving into the details, we present our proof strategy. We establish a deviation bound between the team-optimal

solution and the NE using gradient information. To accomplish this, we propose two projected gradient algorithms for finding the team-optimal solution and the NE. Subsequently, we treat the gradient difference as a perturbation term and derive the deviation bound based on the stability of perturbed systems.

Let us consider the following projected gradient algorithm for seeking the team-optimal solution, i.e, for  $i \in \mathcal{N}$ ,

$$\dot{u}_i = \Pi_{\Xi_i}(u_i - \gamma \nabla_{u_i} \mathcal{C}(\mathbf{u})) - u_i, \quad (7)$$

Here,  $\Pi_{\Xi_i} : \mathbb{R}^{P_i} \rightarrow \Xi_i$  is the projection map, defined as  $\Pi_{\Xi_i}(x) = \operatorname{argmin}_{y \in \Xi_i} \|x - y\|$ , and  $\gamma$  is the stepsize satisfying  $0 < \gamma < \min\{\frac{2\kappa_1}{\nu_1^2}, \frac{2\kappa_2}{\nu_2^2}\}$ . The compact form of (7) can be written as

$$\dot{\mathbf{u}} = \Pi_{\Xi}(\mathbf{u} - \gamma G(\mathbf{u})) - \mathbf{u}. \quad (8)$$

It is clear that the equilibrium of dynamics (8) is  $\mathbf{u}^*$ . Then we show that (8) converges at an exponential rate. Consider the candidate Lyapunov function  $V(\mathbf{u}) = \frac{1}{2}\|\mathbf{u} - \mathbf{u}^*\|^2$ . Its time derivative along the trajectory of (8) is

$$\begin{aligned} \dot{V}(\mathbf{u}) &= -(\mathbf{u} - \mathbf{u}^*)^T (\mathbf{u} - \Pi_{\Xi}(\mathbf{u} - \gamma G(\mathbf{u}))) \\ &= -(\mathbf{u} - \mathbf{u}^*)^T (\mathbf{u} - \Pi_{\Xi}(\mathbf{u} - \gamma G(\mathbf{u})) \\ &\quad - (\mathbf{u}^* - \Pi_{\Xi}(\mathbf{u}^* - \gamma G(\mathbf{u}^*)))) \\ &= -\|\mathbf{u} - \mathbf{u}^*\|^2 + (\mathbf{u} - \mathbf{u}^*)^T (\Pi_{\Xi}(\mathbf{u} - \gamma G(\mathbf{u})) \\ &\quad - \Pi_{\Xi}(\mathbf{u}^* - \gamma G(\mathbf{u}^*))) \\ &\leq -\|\mathbf{u} - \mathbf{u}^*\|(\|\mathbf{u} - \mathbf{u}^*\| - \|\mathbf{u} - \gamma G(\mathbf{u}) - \mathbf{u}^* - \gamma G(\mathbf{u}^*)\|). \end{aligned}$$

Then since

$$\begin{aligned} &\|\mathbf{u} - \mathbf{u}^*\| - \|\mathbf{u} - \gamma G(\mathbf{u}) - \mathbf{u}^* - \gamma G(\mathbf{u}^*)\| \\ &= \frac{\|\mathbf{u} - \mathbf{u}^*\| - \|\mathbf{u} - \gamma G(\mathbf{u}) - \mathbf{u}^* - \gamma G(\mathbf{u}^*)\|^2}{\|\mathbf{u} - \mathbf{u}^*\| + \|\mathbf{u} - \gamma G(\mathbf{u}) - \mathbf{u}^* - \gamma G(\mathbf{u}^*)\|} \\ &\geq \frac{2\gamma(\mathbf{u} - \mathbf{u}^*)^T (G(\mathbf{u}) - G(\mathbf{u}^*)) - \gamma^2 \|G(\mathbf{u}) - G(\mathbf{u}^*)\|^2}{(2 + \gamma \cdot \nu_1)\|\mathbf{u} - \mathbf{u}^*\|} \\ &\geq \frac{2\gamma \cdot \kappa_1 - \gamma^2 \cdot \kappa_1^2}{2 + \gamma \cdot \nu_1} \|\mathbf{u} - \mathbf{u}^*\| \\ &= \omega \|\mathbf{u} - \mathbf{u}^*\|, \end{aligned}$$

we have

$$\dot{V}(\mathbf{u}) \leq -\omega \|\mathbf{u} - \mathbf{u}^*\|^2 \leq -2\omega V(\mathbf{u}).$$

This represents the exponential convergence rate.

Similarly, consider the NE-seeking algorithm for DM  $i$

$$\dot{u}_i = \Pi_{\Xi_i}(\mathbf{u} - \gamma \nabla_{u_i} \mathcal{C}_i(u_i, \mathbf{u}_{-i})) - u_i.$$

The compact form of the NE-seeking dynamics is

$$\dot{\mathbf{u}} = \Pi_{\Xi}(\mathbf{u} - \gamma F(\mathbf{u})) - \mathbf{u} \quad (9)$$

We can similarly prove that (9) converge to  $\mathbf{u}^\diamond$  with an exponential convergence rate. Then, let us denote

$$\beta(\mathbf{u}) = \Pi_{\Xi}(\mathbf{u} - \gamma F(\mathbf{u})) - \Pi_{\Xi}(\mathbf{u} - \gamma G(\mathbf{u})).$$

We rewrite the NE-seeking dynamics (9) as

$$\begin{aligned} \dot{\mathbf{u}} &= \Pi_{\Xi}(\mathbf{u} - \gamma G(\mathbf{u})) - \mathbf{u} + \Pi_{\Xi}(\mathbf{u} - \gamma F(\mathbf{u})) - \Pi_{\Xi}(\mathbf{u} - \gamma G(\mathbf{u})) \\ &= \Pi_{\Xi}(\mathbf{u} - \gamma G(\mathbf{u})) - \mathbf{u} + \beta(\mathbf{u}), \end{aligned} \quad (10)$$

which can be regarded a perturbed system of (8). Moreover,

since  $\Xi$  is compact, there exists a constant  $\delta$  such that

$$\begin{aligned} \|\beta(\mathbf{u})\| &= \|\Pi_{\Xi}(\mathbf{u} - \gamma F(\mathbf{u})) - \Pi_{\Xi}(\mathbf{u} - \gamma G(\mathbf{u}))\| \\ &\leq \|\gamma F(\mathbf{u}) - \gamma G(\mathbf{u})\| \\ &\leq \gamma \|F(\mathbf{u}) - G(\mathbf{u})\| \\ &\leq \gamma \delta. \end{aligned}$$

Then the Lyapunov function  $V(\mathbf{u})$  along the trajectories of (10) satisfies

$$\begin{aligned} \dot{V}(\mathbf{u}) &\leq -\omega \|\mathbf{u} - \mathbf{u}^*\|^2 + \left\| \frac{\partial V}{\partial \mathbf{u}} \right\| \|\beta(\mathbf{u})\| \\ &\leq -\omega \|\mathbf{u} - \mathbf{u}^*\|^2 + \gamma \delta \|\mathbf{u} - \mathbf{u}^*\| \\ &= -(1-s)\omega \|\mathbf{u} - \mathbf{u}^*\|^2 - s\omega \|\mathbf{u} - \mathbf{u}^*\|^2 + \gamma \delta \|\mathbf{u} - \mathbf{u}^*\| \\ &\leq -(1-s)\omega \|\mathbf{u} - \mathbf{u}^*\|^2, \quad \forall \|\mathbf{u} - \mathbf{u}^*\| \geq \gamma \delta / s\omega. \end{aligned}$$

Referring to [20, Theorem 4.18], when  $V(\mathbf{u}) \geq \frac{1}{2}(\gamma \delta / s\omega)^2$ , we have  $\|\mathbf{u} - \mathbf{u}^*\| \geq \gamma \delta / s\omega$  and  $\dot{V}(\mathbf{u}) \leq -2(1-s)\omega V(\mathbf{u})$ . This implies

$$\begin{aligned} \|\mathbf{u} - \mathbf{u}^*\|^2 &\leq \sqrt{2V(\mathbf{u}(t_0))} e^{-(1-s)\omega(t-t_0)} \\ &\leq e^{-(1-s)\omega(t-t_0)} \|\mathbf{u}(t_0) - \mathbf{u}^*\|^2 \end{aligned}$$

over the interval  $[t_0, t_0 + T)$  for some finite  $T$ . For  $t \geq t_0 + T$ , we have

$$\|\mathbf{u} - \mathbf{u}^*\| \leq \sqrt{2V(\mathbf{u})} \leq \gamma \delta / s\omega,$$

and the proof is complete.  $\square$

## B. Proof of Theorem 2

It follows from  $F$  being  $\kappa_2$ -strongly monotone in  $\mathbf{u}$ , we have

$$\kappa_2 \|\mathbf{u}^* - \mathbf{u}^\diamond\| \leq \|F(\mathbf{u}^*) - F(\mathbf{u}^\diamond)\|.$$

Then, following the definition of  $H(\Lambda_{TO}, \Lambda_{NE})$ , we get

$$\begin{aligned} &H(\Lambda_{TO}, \Lambda_{NE}) \\ &= \max \left\{ \sup_{F(\mathbf{u}^*) \in \Lambda_{TO}} \inf_{F(\mathbf{u}^\diamond) \in \Lambda_{NE}} \|F(\mathbf{u}^*) - F(\mathbf{u}^\diamond)\|, \right. \\ &\quad \left. \sup_{F(\mathbf{u}^\diamond) \in \Lambda_{NE}} \inf_{F(\mathbf{u}^*) \in \Lambda_{TO}} \|F(\mathbf{u}^*) - F(\mathbf{u}^\diamond)\| \right\} \\ &\geq \max \left\{ \sup_{F(\mathbf{u}^*) \in \Lambda_{TO}} \inf_{F(\mathbf{u}^\diamond) \in \Lambda_{NE}} \kappa_1 \|\mathbf{u}^* - \mathbf{u}^\diamond\|, \right. \\ &\quad \left. \sup_{F(\mathbf{u}^*) \in \Lambda_{TO}} \inf_{F(\mathbf{u}^\diamond) \in \Lambda_{NE}} \kappa_2 \|\mathbf{u}^* - \mathbf{u}^\diamond\| \right\} \end{aligned}$$

Therefore,

$$\begin{aligned} &H(\Upsilon_{TO}, \Upsilon_{NE}) \\ &= \max \left\{ \sup_{\mathbf{u}^* \in \Upsilon_{TO}} \inf_{\mathbf{u}^\diamond \in \Upsilon_{NE}} \|\mathbf{u}^* - \mathbf{u}^\diamond\|, \right. \\ &\quad \left. \sup_{\mathbf{u}^\diamond \in \Upsilon_{NE}} \inf_{\mathbf{u}^* \in \Upsilon_{TO}} \|\mathbf{u}^* - \mathbf{u}^\diamond\| \right\} \\ &\leq \frac{H(\Lambda_{TO}, \Lambda_{NE})}{\kappa_2} \\ &\leq \frac{\eta}{\kappa_2}. \end{aligned}$$

This completes the proof.  $\square$

### C. Proof of Proposition 1

Given that the payoff functions  $\tilde{C}_i$  are continuously differentiable, it follows from [19] that the condition in Definition 2 is equivalent to

$$\nabla_{u_i} \tilde{C}_i(u_i, \mathbf{u}_{-i}) = \nabla_{u_i} \mathcal{C}(u_i, \mathbf{u}_{-i}) \quad \text{for } i \in \mathcal{N}.$$

By computing the partial derivatives of  $\tilde{C}_i$  and  $\mathcal{C}$ , respectively, we can verify that  $\{\mathcal{N}, \{\Xi_i\}_{i \in \mathcal{N}}, \{\tilde{C}_i\}_{i \in \mathcal{N}}\}$  is a potential game.  $\square$

### D. Proof of Proposition 2

Since  $\{\mathcal{N}, \{\Xi_i\}_{i \in \mathcal{N}}, \{\tilde{C}_i\}_{i \in \mathcal{N}}\}$  is a potential game, if  $\mathbf{u}^*$  is a team-optimal solution that satisfies

$$\mathcal{C}(\mathbf{u}_i^*, \mathbf{u}_{-i}^*) \leq \mathcal{C}(u_i, \mathbf{u}_{-i}), \forall \mathbf{u} \in \Xi.$$

Then, for each  $i \in \mathcal{N}$ , let  $\mathbf{u}$  be the tuple whose  $j$ -th subvector is equal to  $\mathbf{u}_j^*$  for  $j \neq i$  and  $i$ -th subvector is equal to  $u_i$ , where  $u_i$  is an arbitrary element of the set  $\Xi_i$ . We have

$$\mathcal{C}(\mathbf{u}_i^*, \mathbf{u}_{-i}^*) \leq \mathcal{C}(u_i, \mathbf{u}_{-i}^*), \forall i \in \mathcal{N}, \forall u_i \in \Xi_i.$$

This implies  $\mathbf{u}^*$  is an NE that satisfies

$$\tilde{C}_i(\mathbf{u}_i^*, \mathbf{u}_{-i}^*) \leq \tilde{C}_i(u_i, \mathbf{u}_{-i}^*), \forall i \in \mathcal{N}, \forall u_i \in \Xi_i.$$

On the other hand, denote  $\tilde{F}(\mathbf{u}) = \text{col}\{\nabla_{u_1} \tilde{C}_1(\cdot, u_{-1}), \dots, \nabla_{u_N} \tilde{C}_N(\cdot, u_{-N})\}$  as the pseudo-gradient of potential game  $\{\mathcal{N}, \{\Xi_i\}_{i \in \mathcal{N}}, \{\tilde{C}_i\}_{i \in \mathcal{N}}\}$ . Since  $\nabla_{u_i} \tilde{C}_i(u_i, \mathbf{u}_{-i}) = \nabla_{u_i} \mathcal{C}(u_i, \mathbf{u}_{-i})$ , we have  $\tilde{F}(\mathbf{u}) = G(\mathbf{u})$ . Then, based on Assumption 2, the existence of NE  $\mathbf{u}^\diamond$  is guaranteed. Moreover, since  $\Xi_i$  is compact and convex,  $\tilde{C}_i(u_i, \mathbf{u}_{-i})$  is convex and continuously differentiable in  $u_i$ , by convexity and the minimum principle,  $\mathbf{u}^\diamond$  satisfies

$$(u_i - u_i^\diamond)^T \nabla \tilde{C}_i(\mathbf{u}^\diamond) \geq 0, \forall i \in \mathcal{N}, \forall u_i \in \Xi_i.$$

Then by concatenating these inequalities,

$$(\mathbf{u} - \mathbf{u}^\diamond)^T \tilde{F}(\mathbf{u}^\diamond) \geq 0, \forall \mathbf{u} \in \Xi,$$

which implies

$$(\mathbf{u} - \mathbf{u}^\diamond)^T G(\mathbf{u}^\diamond) \geq 0, \forall \mathbf{u} \in \Xi,$$

Thus,  $\mathbf{u}^\diamond$  is also a team-optimal solution that satisfies

$$\mathcal{C}(\mathbf{u}^\diamond) \leq \mathcal{C}(\mathbf{u}), \forall \mathbf{u} \in \Xi. \quad \square$$

## REFERENCES

- [1] A. Orda, R. Rom, and N. Shimkin, "Competitive routing in multiuser communication networks," *IEEE/ACM Transactions on Networking*, vol. 1, no. 5, pp. 510–521, 1993.
- [2] T. Bai, A. Li, G. Xu, C. G. Cassandras, and A. A. Malikopoulos, "Routing guidance for emerging transportation systems with improved dynamic trip equity," *64th IEEE Conference on Decision and Control (in review)*, *arXiv preprint arXiv:2503.12601*, 2025.
- [3] T. G. Crainic, "Service network design in freight transportation," *European Journal of Operational Research*, vol. 122, no. 2, pp. 272–288, 2000.
- [4] M. Aicardi, F. Davoli, R. Minciardi, and R. Zoppoli, "Decentralized strategies for input flow routing in computer communication networks," *IFAC Proceedings Volumes*, vol. 22, no. 10, pp. 257–262, 1989.
- [5] —, "Decentralized routing, teams and neural networks in communications," in *29th IEEE Conference on Decision and Control (CDC)*. IEEE, 1990, pp. 2386–2390.
- [6] K.-C. Leung and V. O. Li, "Flow assignment and packet scheduling for multipath networks," in *Seamless Interconnection for Universal Services. Global Telecommunications Conference. GLOBECOM'99.(Cat. No. 99CH37042)*, vol. 1. IEEE, 1999, pp. 246–250.
- [7] R. Radner, "Team decision problems," *The Annals of Mathematical Statistics*, vol. 33, no. 3, pp. 857–881, 1962.
- [8] A. Dave and A. A. Malikopoulos, "A dynamic program for a team of two agents with nested information," in *2021 IEEE Conference on Decision and Control (CDC)*. IEEE, 2021, pp. 3768–3773.
- [9] A. A. Malikopoulos, "On team decision problems with nonclassical information structures," *IEEE Transactions on Automatic Control*, vol. 68, no. 7, pp. 3915–3930, 2023.
- [10] A. Dave, N. Venkatesh, and A. A. Malikopoulos, "Decentralized control of two agents with nested accessible information," in *2022 American Control Conference (ACC)*. IEEE, 2022, pp. 3423–3430.
- [11] R. Zoppoli, M. Sanguineti, G. Gnecco, T. Parisini *et al.*, *Neural Approximations for Optimal Control and Decision*. Springer, 2020.
- [12] E. Altman, Y. Hayel, and H. Kameda, "Evolutionary dynamics and potential games in non-cooperative routing," in *2007 5th International Symposium on Modeling and Optimization in Mobile, Ad Hoc and Wireless Networks and Workshops*. IEEE, 2007, pp. 1–5.
- [13] A. Haurie and P. Marcotte, "On the relationship between Nash—Cournot and Wardrop equilibria," *Networks*, vol. 15, no. 3, pp. 295–308, 1985.
- [14] S. Shivshankar and A. Jamalipour, "An evolutionary game theory-based approach to cooperation in vanets under different network conditions," *IEEE Transactions on Vehicular Technology*, vol. 64, no. 5, pp. 2015–2022, 2014.
- [15] T. Başar and G. J. Olsder, *Dynamic Noncooperative Game Theory*. SIAM, 1998.
- [16] J. Nash, "Non-cooperative games," *Annals of Mathematics*, pp. 286–295, 1951.
- [17] F. Facchinei and J.-S. Pang, *Finite-Dimensional Variational Inequalities and Complementarity Problems*. Springer, 2003.
- [18] E. M. Bronstein, "Approximation of convex sets by polytopes," *Journal of Mathematical Sciences*, vol. 153, no. 6, pp. 727–762, 2008.
- [19] D. Monderer and L. S. Shapley, "Potential games," *Games and Economic Behavior*, vol. 14, no. 1, pp. 124–143, 1996.
- [20] H. K. Khalil and J. W. Grizzle, *Nonlinear Systems*. Prentice hall Upper Saddle River, NJ, 2002, vol. 3.