

Wasserstein KL-divergence for Gaussian distributions

Adwait Datar¹[0000-0002-4085-9675] and Nihat Ay^{1,2}[0000-0002-8527-2579]

¹ Institute for Data Science Foundations, Hamburg University of Technology, 21073 Hamburg, Germany

² Santa Fe Institute, Santa Fe, NM 87501, USA
{adwait.datar,nihat.ay}@tuhh.de

Abstract. We introduce a new version of the KL-divergence for Gaussian distributions which is based on Wasserstein geometry and referred to as WKL-divergence. We show that this version is consistent with the geometry of the sample space \mathbb{R}^n . In particular, we can evaluate the WKL-divergence of the Dirac measures concentrated in two points which turns out to be proportional to the squared distance between these points.

Keywords: Wasserstein geometry · Kullback-Leibler divergence · Gaussian distributions · Otto metric.

1 Introduction

Classical information geometry is agnostic to the geometry of the sample space, which is chosen to be \mathbb{R}^n in this article. This sample space is equipped with the standard inner product and the corresponding distance function. The Kullback-Leibler divergence (KL-divergence), defined for probability measures on \mathbb{R}^n , represents a fundamental quantity in information geometry [1,4]. It is, however, not coupled with the standard geometry of \mathbb{R}^n . For instance, if we consider two Dirac measures concentrated in distinct points, then their KL-divergence will be infinite and therefore not sensitive to the actual distance between these points. Building on previous work [3,5], a new canonical divergence has been introduced in [2] which is based on Wasserstein geometry [6] and, in particular, the Otto metric [8]. Consider two probability measures μ and ν on \mathbb{R}^n which we assume to have a positive density with respect to the Lebesgue measure. In order to define the new canonical divergence, ν has to have a particular structure in terms of a smooth function $f : \mathbb{R}^n \rightarrow \mathbb{R}$. More precisely, we consider the gradient field $\text{grad } f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and the corresponding flow $\varphi_t : \mathbb{R}^n \rightarrow \mathbb{R}^n$, $t \in \mathbb{R}$, which solves the ordinary differential equation $\frac{d}{dt}\varphi_t(x) = \text{grad } f(\varphi_t(x))$, $\varphi_0(x) = x$. With this, we assume that ν is the image of μ with respect to the flow after a unit of time, that is $\nu = (\varphi_1)_*(\mu)$. In these terms, the new canonical divergence is defined as follows:

$$D_{\text{WKL}}(\mu\|\nu) := \int_{\mathbb{R}^n} \int_0^1 (f \circ \varphi_1 - f \circ \varphi_t) dt d\mu. \quad (1)$$

This represents the Wasserstein version of the classical KL-divergence. We therefore refer to it as the *Wasserstein KL-divergence*, abbreviated by WKL. Clearly, the outlined definition of the WKL-divergence is rather implicit and requires the knowledge about a potential function f that induces the transport of μ to ν in terms of the gradient flow of f . In the present article, we restrict attention to Gaussian distributions where this construction can be made explicit. As a result, we provide an explicit formula for the WKL-divergence in terms of the means μ_0, μ_1 and covariance matrices Σ_0, Σ_1 of the respective Gaussian distributions $\mathcal{N}(\mu_0, \Sigma_0)$ and $\mathcal{N}(\mu_1, \Sigma_1)$ on \mathbb{R}^n . We compare the WKL-divergence with the classical KL-divergence and show that it is indeed nicely coupled with the geometry of the sample space, that is \mathbb{R}^n .

Notation

We use I to denote the identity matrix of appropriate size. The transpose of a matrix A is represented by A^T , and its Moore–Penrose pseudoinverse is denoted by A^\dagger . The projection matrix onto the null space of A is denoted by $A^\perp = I - AA^\dagger$. The gradient of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ evaluated at a point x is the vector $\text{grad}f(x) \in \mathbb{R}^n$. We use $A \succ 0$ ($A \succeq 0$) to indicate that A is symmetric positive definite (semi-definite). The matrix exponential is denoted by e^A and for any $A \succ 0$, the matrix logarithm and symmetric positive definite square root of A is denoted by $\log(A)$ and \sqrt{A} (or $A^{\frac{1}{2}}$), respectively. Finally, the Frobenius norm of A is represented by $\|A\|_F$ and $\text{Tr}(A)$ denotes the trace of A .

2 Main Results

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be the quadratic potential function

$$f(x) = \frac{1}{2}x^T Ax + b^T x \quad (2)$$

where $A \in \mathbb{R}^{n \times n}$ is symmetric³ and $b \in \mathbb{R}^n$. Consider the gradient flow dynamics given by

$$\dot{x}(t) = \text{grad}f(x(t)), \quad x(0) = x_0 \quad (3)$$

and let $\varphi_t : x_0 \mapsto x(t)$ denote its flow map. Our first Lemma provides a formula for the inner integral on the right hand side of (1) in terms of the A and b that define the potential function f .

Lemma 1. *Consider the gradient flow dynamics (3) with a quadratic function f given in (2) and let $M = 2Ae^{2A} - e^{2A} + I$. The following identity holds for all $x_0 \in \mathbb{R}^n$:*

$$\frac{\int_0^1 (f \circ \varphi_1 - f \circ \varphi_t)(x_0) dt}{\int_0^1 (f \circ \varphi_1 - f \circ \varphi_t)(x_0) dt} = \frac{1}{4}(x_0 + A^\dagger b)^T M (x_0 + A^\dagger b) + \frac{1}{2}b^T A^\perp b. \quad (4)$$

³ Note that the assumption on the symmetry of A is without loss of generality since replacing A by $\frac{1}{2}(A + A^T)$ keeps the function unchanged.

Proof. The gradient flow dynamics (3) can be explicitly solved to obtain

$$\begin{aligned} x(t) &= e^{At}x_0 + \left(\int_0^t e^{A(t-\tau)} d\tau \right) b = e^{At} \underbrace{(x_0 + A^\dagger b)}_y + t(I - AA^\dagger)b - A^\dagger b \\ &= e^{At}y + tA^\perp b - A^\dagger b, \end{aligned}$$

where we have defined $y = x_0 + A^\dagger b$ for convenience and have used properties⁴ of A^\perp , A^\dagger and e^A . We now compute $f \circ \varphi_t$ as

$$f(x(t)) = \frac{1}{2}x(t)^T Ax(t) + b^T x(t) = \frac{1}{2}(y^T Ae^{2At}y - b^T A^\dagger b) + tb^T A^\perp b + y^T A^\perp b$$

Therefore,

$$(f \circ \varphi_1)(x_0) - (f \circ \varphi_t)(x_0) = \frac{1}{2}y^T A (e^{2A} - e^{2At}) y + (1-t)b^T A^\perp b.$$

Integrating with respect to time, we get,

$$\begin{aligned} \int_0^1 (f \circ \varphi_1 - f \circ \varphi_t)(x_0) dt &= \int_0^1 \left(\frac{1}{2}y^T A (e^{2A} - e^{2At}) y + (1-t)b^T A^\perp b \right) dt \\ &= \frac{1}{4}y^T (2Ae^{2A} - e^{2A} + I) y + \frac{1}{2}b^T A^\perp b. \end{aligned}$$

Plugging in the definitions of y and M , we get the desired identity. \square

The next lemma considers two Gaussian distributions μ and ν and provides a function f with the desired property that ν is the image of μ with respect to its gradient flow after a unit of time.

Lemma 2. *Let $\mu = \mathcal{N}(\mu_0, \Sigma_0)$ and $\nu = \mathcal{N}(\mu_1, \Sigma_1)$ be two Gaussian distributions with $\Sigma_0 \succ 0$ and $\Sigma_1 \succ 0$. Define the quadratic function f of the form (2) with*

$$A = \log \left(\Sigma_0^{-\frac{1}{2}} \left(\Sigma_0^{\frac{1}{2}} \Sigma_1 \Sigma_0^{\frac{1}{2}} \right)^{\frac{1}{2}} \Sigma_0^{-\frac{1}{2}} \right), \quad (5)$$

$$b = ((e^A - I)A^\dagger + A^\perp)^{-1} (\mu_1 - e^A \mu_0). \quad (6)$$

Then, under the gradient flow dynamics (3), $x(0) \sim \mu$ implies $x(1) \sim \nu$. Furthermore, the function f achieving this property is unique within the set of function of the form (2).

Proof. We have seen in the proof of Lemma 1, that dynamics (3) can be solved to obtain

$$x(t) = e^{At}x_0 + (e^{At}A^\dagger + tA^\perp - A^\dagger) b.$$

⁴ We use the fact that A^\perp , A^\dagger and e^A can be simultaneously diagonalized by orthogonal matrices owing to the symmetry of A which leads to a number of useful properties such as commutativity. These are used throughout the paper.

Using properties of Gaussian random variables, it can be shown that

$$x_0 \sim \mathcal{N}(\mu_0, \Sigma_0) \implies x(t) \sim \mathcal{N}\left(\underbrace{e^{At}\mu_0 + (e^{At}A^\dagger + tA^\perp - A^\dagger)b}_{\mu_t}, \underbrace{e^{At}\Sigma_0e^{At}}_{\Sigma_t}\right).$$

Observe that with the prescribed A and b given in (5) and (6), we indeed get

$$\Sigma_1 = e^A \Sigma_0 e^A, \quad (7)$$

$$\mu_1 = e^A \mu_0 + ((e^A - I)A^\dagger + A^\perp)b. \quad (8)$$

Note that with a change of variable $X = e^A$, (7) reduces to the matrix equation $X\Sigma_0X = \Sigma_1$ which is a special case of the algebraic Riccati equation and has been extensively studied in control theory (see [7] for example). Since $e^A \succ 0$ for any symmetric A , we are interested in the positive definite solutions X to the matrix equation $X\Sigma_0X = \Sigma_1$. Under the constraints that $\Sigma_0 \succ 0$ and $\Sigma_1 \succ 0$, uniqueness of the positive definite solution follows from [7, Theorem 5] proving the uniqueness of the solution A to (7). Uniqueness of b is obtained immediately since $((e^A - I)A^\dagger + A^\perp)$ is non-singular. This proves the final statement. \square

We now present the main result of the paper which provides a formula for the WKL-divergence between two Gaussian distributions.

Theorem 1. *Let $\mu = \mathcal{N}(\mu_0, \Sigma_0)$ and $\nu = \mathcal{N}(\mu_1, \Sigma_1)$ be two Gaussian distributions with $\Sigma_0 \succ 0$ and $\Sigma_1 \succ 0$. Then the following identity holds*

$$D_{\text{WKL}}(\mu\|\nu) = \frac{1}{4}\text{Tr}(\Sigma_0 - \Sigma_1 + \Sigma_0 R^2 \log(R^2)) + \frac{1}{4} \left\| \sqrt{Q + 2\log(R)^\perp}(\mu_1 - \mu_0) \right\|^2$$

where

$$R = \Sigma_0^{-\frac{1}{2}} \left(\Sigma_0^{\frac{1}{2}} \Sigma_1 \Sigma_0^{\frac{1}{2}} \right)^{\frac{1}{2}} \Sigma_0^{-\frac{1}{2}} \quad \text{and} \quad Q = (R - I)^\dagger (\log(R^2)R^2 - R^2 + I) (R - I)^\dagger.$$

Furthermore, if $\Sigma_0 \Sigma_1 = \Sigma_1 \Sigma_0$, then we get the following simplification:

$$D_{\text{WKL}}(\mu\|\nu) = \frac{1}{4} \left(\left\| \sqrt{Q}(\sqrt{\Sigma_1} - \sqrt{\Sigma_0}) \right\|_F^2 + \left\| \sqrt{Q + 2\log(R)^\perp}(\mu_1 - \mu_0) \right\|^2 \right).$$

Proof. Note that the outer integral in the right-hand-side of (1) corresponds to taking an expectation with respect to μ . Using Lemma 2, we define A and b according to (5) and (6) to obtain the property that $x(0) \sim \mu$ implies $x(1) \sim \nu$. Using Lemma 1 along with the well-known properties of the expectation and trace operators⁵, we get that

$$\begin{aligned} D_{\text{WKL}}(\mu\|\nu) &= \int_{\mathbb{R}^n} \int_0^1 (f \circ \varphi_1 - f \circ \varphi_t)(x_0) dt d\mu(x_0) \\ &= \mathbb{E} \left[\frac{1}{4} (x_0 + A^\dagger b)^T M (x_0 + A^\dagger b) + \frac{1}{2} b^T A^\perp b \right] \\ &= \frac{1}{4} (\text{Trace}(M \Sigma_0) + (\mu_0 + A^\dagger b)^T M (\mu_0 + A^\dagger b)) + \frac{1}{2} b^T A^\perp b \quad (9) \end{aligned}$$

⁵ We mainly use the linearity and the cyclic property of the trace operators

where A and b are given by (5) and (6). We now substitute the expression (6) for b into (9) term by term. Let us first compute and simplify $\mu_0 + A^\dagger b$

$$\begin{aligned}\mu_0 + A^\dagger b &= \mu_0 + A^\dagger (A^\dagger(e^A - I) + A^\perp)^{-1} (\mu_1 - e^A \mu_0) \\ &= \mu_0 + (e^A - I)^\dagger (\mu_1 - e^A \mu_0) = (e^A - I)^\perp \mu_0 + (e^A - I)^\dagger (\mu_1 - \mu_0)\end{aligned}$$

Using the fact that $M(e^A - I)^\perp = 0$, we get that

$$(\mu_0 + A^\dagger b)^T M (\mu_0 + A^\dagger b) = \left\| \sqrt{Q} (\mu_1 - \mu_0) \right\|^2. \quad (10)$$

Observe that $M \succeq 0$ which makes the symmetric positive definite matrix square-root well-defined. Similarly, since $A^\perp (A^\dagger(e^A - I) + A^\perp)^{-1} = A^\perp$ and $A^\perp e^A = A^\perp$, the last term $\frac{1}{2} b^T A^\perp b$ can be computed as

$$\begin{aligned}b^T A^\perp b &= (\mu_1 - e^A \mu_0)^T A^\perp (\mu_1 - e^A \mu_0) = (\mu_1 - \mu_0)^T A^\perp (\mu_1 - \mu_0) \\ &= \left\| \log(R)^\perp (\mu_1 - \mu_0) \right\|^2.\end{aligned} \quad (11)$$

Finally, the cyclic property of the trace operator gives us

$$\begin{aligned}\text{Tr}(M \Sigma_0) &= \text{Tr}((\log(R^2) R^2 - R^2 + I) \Sigma_0) \\ &= \text{Tr}(\log(R^2) R^2 \Sigma_0 - \Sigma_1 + \Sigma_0).\end{aligned} \quad (12)$$

Plugging in (10), (11) and (12) in (9) gives the desired result. Finally, when $\Sigma_0 \Sigma_1 = \Sigma_1 \Sigma_0$, the first term can be reformulated using $M(e^A - I)^\perp = 0$ as

$$\begin{aligned}\text{Tr}(M \Sigma_0) &= \text{Tr}((e^A - I)(e^A - I)^\dagger M (e^A - I)^\dagger (e^A - I) \Sigma_0) \\ &= \left\| \sqrt{Q} (\sqrt{\Sigma_1} - \sqrt{\Sigma_0}) \right\|_F^2\end{aligned}$$

□

2.1 Univariate Gaussian Distributions

We now take a closer look at the simple case of univariate Gaussian distributions, i.e., $n = 1$. The main formula is given in the following corollary which is a direct application of Theorem 1.

Corollary 1. *Let $\mu = \mathcal{N}(\mu_0, \sigma_0^2)$ and $\nu = \mathcal{N}(\mu_1, \sigma_1^2)$ be two univariate Gaussian distributions with $\sigma_0 > 0$ and $\sigma_1 > 0$. Then the following identity holds*

$$D_{\text{WKL}}(\mu || \nu) = \begin{cases} \frac{\sigma_0^2 - \sigma_1^2 + \sigma_1^2 \log\left(\frac{\sigma_1^2}{\sigma_0^2}\right)}{4(\sigma_1 - \sigma_0)^2} \left((\sigma_1 - \sigma_0)^2 + (\mu_1 - \mu_0)^2 \right) & \text{if } \sigma_0 \neq \sigma_1, \\ \frac{1}{2} (\mu_1 - \mu_0)^2 & \text{if } \sigma_0 = \sigma_1. \end{cases}$$

Furthermore, we have that

$$\lim_{\sigma_1 \rightarrow \sigma_0} \frac{\sigma_0^2 - \sigma_1^2 + \sigma_1^2 \log\left(\frac{\sigma_1^2}{\sigma_0^2}\right)}{4(\sigma_1 - \sigma_0)^2} \left((\sigma_1 - \sigma_0)^2 + (\mu_1 - \mu_0)^2 \right) = \frac{1}{2} (\mu_1 - \mu_0)^2.$$




Fig. 1. Left figure shows the variation in D_{WKL} and D_{KL} as σ shrink to 0. Right figure shows the local variation in D_{WKL} and D_{KL} as the σ varies around $\sigma_{\text{opt}} \in \{1, 3\}$.

Proof. Substituting $A \leftarrow a$, $\Sigma_0 \leftarrow \sigma_0^2$, $\Sigma_1 \leftarrow \sigma_1^2$, we get

$$R = \frac{\sigma_1}{\sigma_0} \quad \text{and} \quad Q = \left(\sigma_0^2 - \sigma_1^2 + \sigma_1^2 \log \left(\frac{\sigma_1^2}{\sigma_0^2} \right) \right) \left((\sigma_1 - \sigma_0)^2 \right)^\dagger.$$

This leads to

$$D_{\text{WKL}}(\mu \parallel \nu) = \frac{Q}{4} \left((\sigma_1 - \sigma_0)^2 + (\mu_1 - \mu_0)^2 \right) + \frac{1}{2} \log(R)^\perp (\mu_1 - \mu_0)^2$$

which gives the desired result. The final limit can be shown via successive applications of L'Hopital's rule. \square

Let $\mu = \mathcal{N}(\mu_0, \sigma^2)$ and $\nu = \mathcal{N}(\mu_1, \sigma^2)$ be two univariate Gaussian distributions with equal variance σ^2 and possibly different means μ_0 and μ_1 . Let us now compare the WKL-divergence and the classical KL-divergence between these distributions which can be computed as

$$D_{\text{WKL}}(\mu \parallel \nu) = \frac{(\mu_1 - \mu_0)^2}{2} \quad \text{and} \quad D_{\text{KL}}(\nu \parallel \mu) = \frac{(\mu_1 - \mu_0)^2}{2\sigma^2}.$$

Note that since our WKL-divergence corresponds to the dual of the KL-divergence [2], a direct comparison requires reversing the order of the distributions. It is clear from this example that as $\sigma \rightarrow 0$, the KL-divergence diverges to infinity whereas the WKL-divergence remains constant since it depends solely on the distance between the means. This is illustrated in Figure 1 (left) where we set $\mu_0 = 0$, $\mu_1 = 2$ and the standard deviations $\sigma_0 = \sigma_1 = \sigma$ is varied. This illustrates the coupling of the divergence with the geometry of the sample space which in this case is \mathbb{R} . This allows us to approximate the divergence between two Dirac measures such that it is proportional to the distance in the sample space.

Let $\mu = \mathcal{N}(\mu, \sigma_{\text{opt}}^2)$ and $\nu = \mathcal{N}(\mu, \sigma^2)$ be two univariate Gaussian distributions with equal means μ . We now look at the local curvature of the KL-divergence and the WKL-divergence around an optimum which has a strong influence on performance of optimization algorithms. The curvature corresponds




Fig. 2. Surface plots for $D_{\text{WKL}}(\mu\|\nu)$ (left) and $D_{\text{KL}}(\nu\|\mu)$ (right) where $\mu = \mathcal{N}(0, \sigma_0^2)$ and $\nu = \mathcal{N}(1, \sigma_1^2)$

to the hessian evaluated at the optimum and it can be shown that

$$\left. \frac{\partial^2}{\partial \sigma^2} D_{\text{WKL}}(\mu\|\nu) \right|_{\sigma=\sigma_{\text{opt}}} = 1 \quad \text{and} \quad \left. \frac{\partial^2}{\partial \sigma^2} D_{\text{KL}}(\nu\|\mu) \right|_{\sigma=\sigma_{\text{opt}}} = \frac{2}{\sigma_{\text{opt}}^2}.$$

Observe that local curvature of the WKL-divergence is independent of σ_{opt} whereas the local curvature of the KL-divergence blows up to infinity as σ_{opt} approaches zero. Figure 1 (right) plots the divergences for $\mu = 0$ and $\sigma_{\text{opt}} \in \{3, 6\}$ and it can be observed that the local curvature of D_{WKL} is independent of σ_{opt} whereas the local curvature of D_{KL} is high for $\sigma_{\text{opt}} = 3$ than that for $\sigma_{\text{opt}} = 6$.

2.2 Discussion on Continuity of D_{WKL}

Note that Theorem 1 implies $D_{\text{WKL}}(\mathcal{N}(\mu_0, \Sigma_0)\|\mathcal{N}(\mu_1, \Sigma_0)) = \frac{1}{2} \|(\mu_1 - \mu_0)\|^2$. In analogy with the continuity statement of Corollary 1, it can be shown that if the sequence $\Sigma_{(1)}, \Sigma_{(2)}, \dots$ satisfies $\|\Sigma_{(i)} - \Sigma_0\|_F \rightarrow 0$, then

$$D_{\text{WKL}}(\mathcal{N}(\mu_0, \Sigma_0)\|\mathcal{N}(\mu_1, \Sigma_{(i)})) \rightarrow D_{\text{WKL}}(\mathcal{N}(\mu_0, \Sigma_0)\|\mathcal{N}(\mu_1, \Sigma_0)).$$

This involves diagonalizing the involved symmetric matrices and applying the L'Hopital's rule for each eigenvalue sequence separately. Thus, D_{WKL} is continuous at $\Sigma_0 = \Sigma_1$. Since the value is independent of Σ_0 , it allows us to take limit as Σ_0 tends to 0. This gives us a way to approximate the divergence between Dirac measures concentrated at μ_0 and μ_1 and provides a finite value that is proportional to the squared distance $\|\mu_0 - \mu_1\|^2$. In contrast, the KL-divergence for $\Sigma_0 = \Sigma_1 = \Sigma$ gives

$$D_{\text{KL}}(\nu\|\mu) = \frac{1}{2} \left\| \Sigma^{-1/2}(\mu_1 - \mu_0) \right\|^2$$

which diverges to ∞ as Σ tends to 0. Figure 2 shows the surface plots for the KL-divergence and the WKL-divergence for univariate Gaussian distributions. In contrast to KL-divergence, the WKL-divergence does not blow up at the origin.

3 Outlook

Building on [2], we have presented a novel WKL-divergence for multivariate Gaussian distributions. A promising direction for future research is to explore its information theoretic applications such as maximum likelihood estimation. Comparing its empirical performance with existing divergences in real-world machine learning tasks also remains an exciting avenue.

Acknowledgments. Nihat Ay would like to thank Felix Otto for his helpful comments and suggestions.

Disclosure of Interests. The authors have no competing interests to declare that are relevant to the content of this article.

References

1. Amari, S.i., Nagaoka, H.: Methods of information geometry, vol. 191. American Mathematical Soc. (2000)
2. Ay, N.: Information geometry of the otto metric. *Information Geometry* pp. 1–24 (2024)
3. Ay, N., Amari, S.i.: A novel approach to canonical divergences within information geometry. *Entropy* **17**(12), 8111–8129 (2015)
4. Ay, N., Jost, J., Van Lê, H., Schwachhöfer, L.: *Information geometry*, vol. 64. Springer (2017)
5. Felice, D., Ay, N.: Towards a canonical divergence within information geometry. *Information geometry* **4**(1), 65–130 (2021)
6. Khan, G., Zhang, J.: When optimal transport meets information geometry. *Information Geometry* **5**(1), 47–78 (2022)
7. Kučera, V.: A review of the matrix riccati equation. *Kybernetika* **9**(1), 42–61 (1973)
8. Otto, F.: The geometry of dissipative evolution equations: the porous medium equation. *Comm. Partial Differential Equations* **26**, 101–174 (2001)