

# ASYMPTOTICALLY DISTRIBUTION-FREE GOODNESS-OF-FIT TESTING FOR POINT PROCESSES

JUSTIN BAARS, S. UMUT CAN, AND ROGER J. A. LAEVEN

**ABSTRACT.** Consider an observation of a multivariate temporal point process  $N$  with law  $\mathcal{P}$  on the time interval  $[0, T]$ . To test the null hypothesis that  $\mathcal{P}$  belongs to a given parametric family, we construct a convergent compensated counting process to which we apply an innovation martingale transformation. We prove that the resulting process converges weakly to a standard Wiener process. Consequently, taking a suitable functional of this process yields an asymptotically distribution-free goodness-of-fit test for point processes. For several standard tests based on the increments of this transformed process, we establish consistency under alternative hypotheses. Finally, we assess the performance of the proposed testing procedure through a Monte Carlo simulation study and illustrate its practical utility with two real-data examples.

**KEYWORDS.** Point processes  $\circ$  goodness-of-fit testing  $\circ$  innovation martingale transformation  $\circ$  functional central limit theorem  $\circ$  Hawkes processes  $\circ$  Monte Carlo simulation

**AFFILIATIONS.** JB, UC and RL are with the Dept. of Quantitative Economics, University of Amsterdam. RL is also with EURANDOM, Eindhoven University of Technology, and with CENTER, Tilburg University. The research of JB and RL is funded in part by the Netherlands Organization for Scientific Research under an NWO VICI grant (2020–2027).

**EMAIL.** J.R.Baars@uva.nl, S.U.Can@uva.nl, R.J.A.Laeven@uva.nl.

*Date:* April 1, 2025.

## 1. INTRODUCTION

Consider a  $d$ -variate temporal point process  $N = (N^{(1)}, \dots, N^{(d)})$  depending on time  $t \in \mathbb{R}$ . Suppose that  $N$  is modeled through a parametric null hypothesis  $\mathcal{F}_\Theta := \{N_\theta : \theta \in \Theta\}$ , where  $\Theta \subset \mathbb{R}^m$  is a finite-dimensional parameter space. Under the null hypothesis  $H_0$ , it holds that  $N \stackrel{d}{=} N_{\theta_0}$  for some  $\theta_0 \in \Theta$ . In practice,  $\theta_0$  is typically unknown, and must be estimated using an observation  $\omega_T = \{t_1, \dots, t_{n(T)}\}$  of the point process on the time interval  $[0, T]$ , leading to an estimated parameter vector  $\hat{\theta}_T$ . If maximum likelihood estimation is employed, the statistical properties of  $\hat{\theta}_T$  are well-established [37, 42, 43].

The present work is concerned with goodness-of-fit testing for temporal point processes. Given an observation  $\omega_T$  of the point process on the time interval  $[0, T]$ , we are interested in testing the hypothesis  $H_0 : N \stackrel{d}{=} N_\theta$  for some  $\theta \in \Theta$  against the alternative  $H_1 : N \not\stackrel{d}{=} N_\theta$  for all  $\theta \in \Theta$ . In goodness-of-fit problems, the test statistic  $S$  often has a distribution that is challenging to calculate and that may depend on the null hypothesis or on the true parameter, fragmenting the theory. As a result, one has to resort to *ad hoc* bootstrap methods to estimate critical values for each specific application. To address this issue, a test statistic, or test process, is needed that, under the null hypothesis, has a distribution that is independent of both  $\mathcal{F}_\Theta$  and the true parameter  $\theta_0$ .

For temporal point processes, a common approach is to apply a time transformation  $(\Lambda(t_i))_{i \in [n(T)]}$ , where  $[n] := \{1, \dots, n\}$  and  $\Lambda(t) = \int_0^t \lambda(s) ds$  is the *compensator* of the process, i.e., the time-integrated conditional intensity. According to the random time change theorem [14], Theorem 7.4.I, under the law  $\mathcal{P}_{\theta_0}$

of  $N_{\theta_0}$  the transformed arrival times  $\Lambda_{\theta_0}(t_i)$  follow a Poisson process of unit rate. Since  $\theta_0$  is unknown, typically the goodness-of-fit of a candidate model is assessed by testing whether the transformed interarrival times  $\Lambda_{\hat{\theta}_T}(t_i) - \Lambda_{\hat{\theta}_T}(t_{i-1})$  are standard exponential. In such tests, the estimation uncertainty  $\hat{\theta}_T \neq \theta_0$ , which causes a discrepancy between  $\Lambda_{\hat{\theta}_T}$  and  $\Lambda_{\theta_0}$ , is generally ignored; see [3, 10, 11, 19, 31, 32, 34, 49, 50].

In recent decades, various goodness-of-fit testing procedures for temporal point processes have been proposed, including random thinning [38], random superposition [7], super-thinning [12], and random-time-change-based methods [13, 45]. However, those procedures rely on results that hold only under the true distribution  $\mathcal{P}_{\theta_0}$ . In practice, since  $\theta_0$  is unknown and typically replaced by the estimator  $\hat{\theta}_T$ , such tests fail to account for the uncertainty introduced by estimation. To the best of our knowledge, no existing goodness-of-fit test for general temporal point processes explicitly addresses this challenge, leaving a significant gap in the literature.

Ignoring estimation uncertainty has important consequences: the distribution of the test statistic under the null hypothesis still depends on the specific model class  $\mathcal{F}_{\Theta}$  and the true parameter  $\theta_0$ , even in the limit as  $T \rightarrow \infty$ . There is no theoretical justification for ignoring the estimation uncertainty  $\hat{\theta}_T \neq \theta_0$ . In fact, even if the data is generated under the law  $\mathcal{P}_{\theta_0}$ , the estimator  $\hat{\theta}_T$  will, by construction, fit the observed data better than the true parameter  $\theta_0$ . Consequently, for a test statistic  $S(\omega_T, \theta)$  depending on both the observation  $\omega_T$  and the parameter  $\theta$ , it is natural to expect that under  $\mathcal{P}_{\theta_0}$ , the value of  $S(\omega_T, \hat{\theta}_T)$  will appear ‘less extreme’ than  $S(\omega_T, \theta_0)$ . Ignoring estimation uncertainty leads to undersized tests with reduced power. In applied work, it is common to fit a model to data and then attempt to demonstrate that the model fits well by using a goodness-of-fit test that fails to reject the null hypothesis. Ignoring estimation uncertainty in this context leads to overly conservative tests (i.e., with too low power), potentially giving a false sense of confidence in an otherwise imperfect model.

In this paper, we address the limitations of existing goodness-of-fit procedures by developing an asymptotically distribution-free testing method for a broad class of point processes. Specifically, based on the observation  $\omega_T$  we construct a test *process* on a bounded interval  $[0, \tau]$ , which converges weakly to a standard Wiener process, as  $T \rightarrow \infty$ . Consequently, the limiting process has an asymptotic distribution that is independent of both the null hypothesis and the true parameter  $\theta_0$ . Functionals of this test process can be employed as a test statistic. Since the weak limit of the test process is a standard Wiener process that does not depend on the model class  $\mathcal{F}_{\Theta}$  or the true parameter  $\theta_0$ , the asymptotic distributions of these functionals are also independent of  $\mathcal{F}_{\Theta}$  and  $\theta_0$ . This means that their critical values only need to be tabulated once, making them applicable across a wide range of testing problems — an important practical advantage.

To derive the asymptotically distribution-free test process, we first establish a functional central limit theorem (FCLT) for the counting process  $N_{\theta_0}$  compensated by  $\Lambda_{\hat{\theta}_T}$ . The resulting limit is the sum of a Wiener process and a linear drift with a random coefficient, which still depends on the true parameter  $\theta_0$  and the model class  $\mathcal{F}_{\Theta}$ . However, the specific structure of this limit enables the application of an *innovation martingale transformation*  $\mathcal{T}_{\theta_0}$ , originally introduced by Khmaladze in [24, 25, 26]. After transformation, we obtain a process that, as we formally establish, converges to a standard Wiener process, as  $T \rightarrow \infty$ .

The remainder of this paper is organized as follows. In Section 2, we introduce our setup and state the assumptions on point processes. Section 3 is dedicated to deriving a functional central limit theorem (FCLT) for the compensated *empirical* counting process. This is accomplished by first proving FCLTs for both the compensated counting process and the difference between the true and empirical compensators. Along the way, we derive a Cramér-Wold device for stochastic processes, which, to the best of our knowledge, cannot be found in the literature and may be of independent interest. We use this result to prove an FCLT for

multivariate martingales, which is also noteworthy in its own right. We then proceed to the key step of our approach: applying the innovation martingale transformation. We show that the difference between the true transformation and its empirical counterpart becomes negligible as  $T \rightarrow \infty$ , resulting in an asymptotically distribution-free test process.

In Section 4, we focus on goodness-of-fit testing for point processes. Section 4.1 explains how to construct asymptotically distribution-free test statistics from the previously derived test process. Section 4.2 discusses the bias inherent in random-time-change-based methods. In Section 5, we demonstrate that, under any reasonable alternative, standard goodness-of-fit tests such as the Kolmogorov-Smirnov, Anderson-Darling, and Cramér-von Mises tests, based on the increments of our test process, have power converging to 1, ensuring consistency under alternatives.

In Section 6, we evaluate the performance of our proposed testing procedure through simulations under a parametric null hypothesis, and compare it to methods that ignore estimation uncertainty. The results show that by explicitly accounting for this uncertainty, our procedure delivers more reliable and robust inferences. In Section 7, we further illustrate its practical advantages over classical methods through two real-data applications, where our testing procedure leads to different conclusions than tests that disregard estimation uncertainty. Finally, in Section 8 we describe how to extend our work to a spatiotemporal setup.

**Appendix.** Several technical assumptions are collected in the appendix.

## 2. MODEL DESCRIPTION AND ASSUMPTIONS

Let  $N = (N^{(1)}, \dots, N^{(d)})$  be a  $d$ -variate point process on  $\mathbb{R}$ , i.e., a random discrete  $[d]$ -marked subset of  $\mathbb{R}$ . Denote the law of  $N$  by  $\mathcal{P}$ . A realization of  $N$  consists of event times  $\omega := \{\dots, t_{-2}, t_{-1}, t_0, t_1, t_2, \dots\} \subset \mathbb{R}$  along with coordinates  $c_i \in [d]$ ; here,  $t_i < t_{i+1}$  and  $t_0 < 0 \leq t_1$ . The realization on  $[0, T]$  is then denoted by  $\omega_T = \omega \cap [0, T] = \{t_1, \dots, t_{n(T)}\}$ , where  $n(T)$  denotes the number of events in the interval  $[0, T]$ . We assume that  $\omega$  has no limit point.

We are mostly interested in the situation in which  $N \stackrel{d}{=} N_{\theta_0}$ , where  $N_{\theta_0}$  belongs to the parametric family  $\mathcal{F}_{\Theta} := \{N_{\theta} : \theta \in \Theta\}$ , with  $\Theta \subset \mathbb{R}^m$ . Here,  $\stackrel{d}{=}$  denotes equality in distribution. Denote the law of  $N_{\theta}$  by  $\mathcal{P}_{\theta}$ . We use this framework to test the parametric null hypothesis

$$H_0 : N \stackrel{d}{=} N_{\theta} \text{ for some } \theta \in \Theta, \quad (1)$$

using an observation  $\omega_T$  of  $N$  on the time interval  $[0, T]$ .

We start with assumptions on  $N_{\theta}$  and on  $\Theta$ .

### Assumptions A.

- (i)  $\Theta \subset \mathbb{R}^m$  is open, convex and bounded.
- (ii) For each  $\theta \in \Theta$ ,  $N_{\theta}$  is a stationary and ergodic point process, absolutely continuous w.r.t. the unit Poisson process on  $[0, T]$ , for each  $T \in (0, \infty)$ .
- (iii) For each  $\theta \in \Theta$ ,  $N_{\theta}$  is an orderly point process, i.e.,  $\mathbb{P}(N_{\theta}[0, \delta] \geq 2) = o(\delta)$ , as  $\delta \downarrow 0$ .

By Assumptions A, it follows that, for each  $\theta \in \Theta$ , the probabilistic behavior of  $N_{\theta}$  can be characterized by a conditional intensity function  $\lambda_{\theta} : \mathbb{R}_+ \rightarrow \mathbb{R}_+^d : t \mapsto \lambda_{\theta}(t)$ ; see [14], Proposition 7.3.IV. Denote its components by  $\lambda_{\theta}^{(k)}$ ,  $k \in [d]$ . Let

$$\mathcal{H}_t := \sigma(N_{\theta}(s) : s \in (-\infty, t]).$$

Then  $\lambda_\theta$  can be taken to be any  $(\mathcal{H}_t)_{t \in \mathbb{R}}$ -predictable function such that

$$\lambda_\theta(t) = \lim_{\Delta t \downarrow 0} \frac{1}{\Delta t} \mathbb{P}(N_\theta[t, t + \Delta t] > 0 | \mathcal{H}_t). \quad (2)$$

We assume that the point process  $N = N_\theta$  is specified through a conditional intensity function satisfying the following assumptions.

**Assumptions B.**

- (i) For each  $\theta \in \Theta$ ,  $\lambda_\theta$  is  $(\mathcal{H}_t)_{t \in \mathbb{R}}$ -predictable.
- (ii) The model  $\mathcal{F}_\Theta$  is identifiable:  $\lambda_{\theta_1}(0) = \lambda_{\theta_2}(0)$  a.s. if and only if  $\theta_1 = \theta_2$ .
- (iii)  $\lambda_\theta(0) \in C^1(\Theta)$  a.s., with for any  $\theta \in \Theta$ ,  $\partial \lambda_\theta^{(k)}(0) / \partial \theta_i \in L^2(\mathbb{P})$  for all  $k \in [d]$  and  $i \in [m]$ .
- (iv) For each  $\theta \in \Theta$ ,  $\lambda_\theta(0) > 0$  a.s.
- (v) For each  $\theta \in \Theta$ , there exists  $\Lambda_0 \in L^1(\mathbb{P})$  and a neighborhood  $U = U(\theta) \subset \Theta$  of  $\theta$  such that for all  $\theta' \in U$ ,  $\|\lambda_{\theta'}(0, \omega)\|_\infty \leq \Lambda_0(\omega)$ .
- (vi) Let  $D\lambda_\theta(t) = (\partial \lambda_\theta^{(k)}(t) / \partial \theta_i)_{k \in [d], i \in [m]} \in \mathbb{R}^{d \times m}$  be the total derivative of  $\theta \mapsto \lambda_\theta(t)$ . Assume that

$$\alpha_\theta := \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T D\lambda_\theta(t) dt \quad (3)$$

exists a.s. for  $\theta = \theta_0$ .

**Remark 1.** By stationarity, i.e., Assumption A(ii), Assumptions B can be stated without reference to  $t$ .

**Remark 2.** Assumption B(vi) is satisfied if the following condition holds: there exists  $\Lambda_1 \in L^1(\mathbb{P})$  and a neighborhood  $U = U(\theta_0) \subset \Theta$  of  $\theta_0$  such that for all  $\theta' \in U$ ,  $\|D\lambda_{\theta'}(0)\| \leq \Lambda_1$ . This follows by ergodicity (Assumption A(ii)) together with the dominated convergence theorem. In this case, using stationarity of  $\lambda_\theta$ , it follows that  $\alpha_\theta = \mathbb{E}[D\lambda_\theta(0)]$ .

Suppose that we have some estimator  $\hat{\theta}_T \in \Theta$  for  $\theta$  based on the observation  $\omega_T \subset [0, T]$  of  $N$ . We assume that this estimator satisfies a central limit theorem, as follows.

**Assumption C.** There exists an  $m$ -variate random vector  $Z$  such that under  $\mathcal{P}_{\theta_0}$  it holds that

$$\sqrt{T}(\theta_0 - \hat{\theta}_T) \xrightarrow{d} Z, \quad \text{as } T \rightarrow \infty. \quad (4)$$

Assumption C is satisfied with  $Z \sim \mathcal{N}(0, I(\theta_0)^{-1})$  (where  $I(\theta)$  is defined in Assumption C.1(ii) below) when  $\hat{\theta}_T$  is the maximum likelihood estimator and if next to Assumptions A–B one grants Assumptions C.1 below. This follows from a multivariate generalization of [37], Theorem 5; see, e.g., [42], Theorem 4.

**Assumptions C.1.**

- (i) The log-likelihood function  $L_T(\theta)$  at time  $T$  has a unique maximum a.s. Here  $L_T(\theta)$  is defined by

$$L_T(\theta) = \sum_{k=1}^d \left( \int_0^T \log \lambda_\theta^{(k)}(t) dN(t) - \int_0^T \lambda_\theta^{(k)}(t) dt \right). \quad (5)$$

- (ii) For each  $\theta \in \Theta$  and for all  $i, j \in [m]$ ,  $k \in [d]$ ,  $\frac{1}{\lambda_\theta^{(k)}} \frac{\partial \lambda_\theta^{(k)}}{\partial \theta_i} \frac{\partial \lambda_\theta^{(k)}}{\partial \theta_j} \in L^2(\mathbb{P})$ . Assume that for each  $\theta \in \Theta$  the Fisher information matrix  $I(\theta) \in \mathbb{R}^{m \times m}$  with elements

$$I_{ij}(\theta) = \sum_{k=1}^d \mathbb{E} \left[ \frac{1}{\lambda_\theta^{(k)}} \frac{\partial \lambda_\theta^{(k)}}{\partial \theta_i} \frac{\partial \lambda_\theta^{(k)}}{\partial \theta_j} \right] \quad (6)$$

is nonsingular.

- (iii)  $\lambda_\theta \in C^3(\Theta)$  with  $\frac{\partial^2 \lambda_\theta^{(k)}}{\partial \theta_i \partial \theta_j} \in L^2(\mathbb{P})$  for all  $k \in [d]$ ,  $i, j \in [m]$ .
- (iv) See appendix.

**Remark 3.** Assumptions A, B, and C.1 are satisfied by Poisson, Hawkes, Wold, and delayed renewal processes; see [37]. Furthermore, our setup and assumptions allow for marked point processes.

In Section 3, we derive a limit of a certain transformation of the stochastic process  $N$ . We need an appropriate framework for stochastic process convergence. The transformation of the next section maps the time interval  $[0, T]$  to  $[0, 1]$ , hence it suffices to work on the latter interval. On the space  $D([0, 1], \mathbb{R}^d)$  of right-continuous functions  $f : [0, 1] \rightarrow \mathbb{R}^d$  we consider multiple topologies. We consider the strong  $SU$  and the weak  $WU$  uniform topologies and the strong  $SJ_1$  and weak  $WJ_1$  Skorokhod  $J_1$  topologies. The strong topologies are defined by treating  $\mathbb{R}^d$  as the range space, while the weak topologies use the fact that  $D([0, 1], \mathbb{R}^d) = D([0, 1], \mathbb{R})^d$  in the sense of bijection, and consider a product topology on this space. As the name suggests, in general the strong topology is stronger than the weak topology. However, as we will prove later, the strong and product uniform topologies coincide, i.e.,  $SU = WU$ . See [53] for more details.

### 3. ASYMPTOTICALLY DISTRIBUTION-FREE TEST PROCESS

In this section, we derive a distribution-free limit process from a transformation of the process  $N_{\theta_0}$ , meaning that the limit is in fact independent of the true parameter  $\theta_0$ , as well as of the model  $\mathcal{F}_\Theta$ . As indicated in the Introduction, this allows us to overcome a crucial shortcoming of goodness-of-fit tests for point processes known from the literature. Our plan of action consists of four steps.

- (1) We consider the  $d$ -dimensional compensated empirical process

$$\hat{\eta}^{(T)} : u \mapsto \frac{1}{\sqrt{T}} \left( N_{\theta_0}(uT) - \int_0^{uT} \lambda_{\hat{\theta}_T}(s) ds \right), \quad (7)$$

for  $u \in [0, 1]$ , which we decompose into

$$\hat{\eta}^{(T)} : u \mapsto \frac{1}{\sqrt{T}} \left( N_{\theta_0}(uT) - \int_0^{uT} \lambda_{\theta_0}(s) ds \right) + \frac{1}{\sqrt{T}} \left( \int_0^{uT} \lambda_{\theta_0}(s) ds - \int_0^{uT} \lambda_{\hat{\theta}_T}(s) ds \right). \quad (8)$$

The first step consists of applying a martingale functional central limit theorem (martingale FCLT) to the first term of (8), in order to obtain a Gaussian limit.

- (2) We show that the second term of (8) converges to a deterministic function of  $u$  multiplied by some random variable (which is, in particular, independent of  $u$ ).
- (3) From steps 1 and 2, we obtain a limit  $\hat{\eta}_{\theta_0}$  of (8) which is still dependent on the true parameter  $\theta_0$ , as well as on the model  $\mathcal{F}_\Theta$ . However, the particular shape of the limit allows us to apply an *innovation martingale transformation*  $\mathcal{T}_{\theta_0}$ , first discussed by Khmaladze in [24, 25, 26]. After transformation,  $\mathcal{T}_{\theta_0}(\hat{\eta}_{\theta_0})$  is a standard Wiener process. This transformation  $\mathcal{T}_{\theta_0}$  still depends on the true, but unknown parameter  $\theta_0$ .
- (4) Finally, we show that the difference between  $\mathcal{T}_{\theta_0}(\hat{\eta}_{\theta_0})$  and its empirical counterpart  $\mathcal{T}_{\hat{\theta}_T}(\hat{\eta}^{(T)})$  is small for large  $T$ , hence  $\mathcal{T}_{\hat{\theta}_T}(\hat{\eta}^{(T)})$  converges to a standard Wiener process.

As we will see, due to the orthogonal structure of the martingales underlying a multivariate point process, with some work, the analysis in the multivariate case can be reduced to that of univariate case.

**3.1. Step 1: FCLT for the compensated process.** For our first step, we need a multivariate analog of [6], Theorem 18.3. In order to prove this, we apply the following lemma.

**Lemma 1** (Cramér-Wold device for stochastic processes). *Let  $(W_n)_{n \in \mathbb{N}} \subset D([0, 1], \mathbb{R}^d)$  be a sequence of  $d$ -dimensional stochastic processes such that*

$$\text{for all } c \in \mathbb{R}^d : \quad c^\top W_n \xrightarrow{d} c^\top W, \quad (9)$$

*on  $D([0, 1], \mathbb{R})$  equipped with the Skorokhod  $J_1$  topology, and where  $W$  is a Wiener process on  $\mathbb{R}^d$ . Then it holds that  $W_n \xrightarrow{d} W$  on  $D([0, 1], \mathbb{R}^d)$  equipped with the strong Skorokhod  $J_1$  topology.*

*Proof.* By the standard Cramér-Wold device, the condition (9) implies that the finite-dimensional distributions of  $W_n$  converge to those of  $W$ .

Next, condition (9) implies that the one-dimensional marginals converge to a continuous limit, meaning that we have weak convergence in  $D([0, 1], \mathbb{R})$  equipped with the uniform topology. Because of this weak convergence, we have tightness (w.r.t. the uniform topology) of the one-dimensional marginals. In turn, by [53], Theorem 11.6.7, this implies tightness of the processes  $W_n$  themselves; this tightness is w.r.t. the product topology on  $D([0, 1], \mathbb{R})^d$  obtained out of uniform topologies on  $D([0, 1], \mathbb{R})$ . Note that here the  $\|\cdot\|_\infty$ -norm on  $\mathbb{R}^d$  is used.

Using Prohorov's theorem, those two conditions imply that  $W_n \xrightarrow{d} W$  on  $D([0, 1], \mathbb{R})^d$  equipped with the product of uniform topologies,  $WU$ . We now prove that this is equal to the strong uniform topology on  $D([0, 1], \mathbb{R}^d)$ ,  $SU$ .

Indeed, the product topology  $WU$  is generated by the subbase consisting of the sets

$$\pi_i^{-1} \{x_i \in D([0, 1], \mathbb{R}) : \|x_i - y_i\|_\infty < \epsilon\}, \quad y_i \in D([0, 1], \mathbb{R}), \epsilon > 0, i \in [d],$$

where  $\pi_i$  is the projection operator onto the  $i$ th coordinate. Since

$$\pi_i^{-1} \{x_i \in D([0, 1], \mathbb{R}) : \|x_i - y_i\|_\infty < \epsilon\} = \bigcup_{\substack{z \in D([0, 1], \mathbb{R}^d) \\ z_i = y_i}} \{x \in D([0, 1], \mathbb{R}^d) : \|x - z\|_\infty < \epsilon\}$$

lies in the uniform topology on  $D([0, 1], \mathbb{R}^d)$ , it follows that  $WU \subset SU$ . Next, note that  $SU$  on  $D([0, 1], \mathbb{R}^d)$  is generated by the sets

$$\{x \in D([0, 1], \mathbb{R}^d) : \|x - y\|_\infty < \epsilon\}, \quad y \in D([0, 1], \mathbb{R}^d), \epsilon > 0.$$

Hence, the inclusion  $WU \supset SU$  follows by the equality

$$\{x \in D([0, 1], \mathbb{R}^d) : \|x - y\|_\infty < \epsilon\} = \bigcap_{i \in [d]} \pi_i^{-1} \{x_i \in D([0, 1], \mathbb{R}) : \|x_i - y_i\|_\infty < \epsilon\}.$$

To conclude, we have  $W_n \xrightarrow{d} W$  on  $D([0, 1], \mathbb{R})^d$  equipped with the strong uniform topology  $SU$ . This implies convergence in the  $SJ_1$  topology.  $\square$

**Remark 4.** *As is evident from the proof, Lemma 1 allows for generalizations to stochastic processes for which all the finite-dimensional distributions are Borel probability measures.*

The following result is a multivariate extension of [6], Theorem 18.3.

**Lemma 2** (FCLT for multivariate martingales). *Consider a stationary and ergodic two-sided  $d$ -variate martingale difference sequence  $(\xi_n)_{n \in \mathbb{Z}}$ , where  $\xi_n$  takes values in  $\mathbb{R}^d$  and satisfies  $\mathbb{E}[\xi_n | \mathcal{F}_{n-1}] = 0$ , with  $\mathcal{F}_n := \sigma(\xi_k : k \leq n)$ . Write  $\Xi := (\Xi_{ij})_{i, j \in [d]} := (\mathbb{E}[\xi^{(i)} \xi^{(j)}])_{i, j \in [d]} \in \mathbb{R}^{d \times d}$ , where  $\xi^{(i)}$  denotes the  $i$ th*

coordinate of the  $d$ -dimensional generic random variable  $\xi$ . The stochastic process  $t \mapsto X_t^n := \sum_{k \leq nt} \xi_k / \sqrt{n}$  converges weakly on  $D([0, 1], \mathbb{R}^d)$  equipped with the strong Skorokhod  $J_1$  topology to a  $d$ -variate Wiener process  $W$  with covariance matrix  $u\Xi$ .

*Proof.* By Lemma 1, it suffices to show that for arbitrary  $c \in \mathbb{R}^d$  it holds that  $c^\top X^n \xrightarrow{d} c^\top W$  on  $D[0, 1]$  equipped with the Skorokhod  $J_1$  topology.

To this end, note that  $(c^\top \xi_n)_{n \in \mathbb{Z}}$  is a univariate martingale difference sequence w.r.t.  $(\mathcal{F}_n)_{n \in \mathbb{Z}}$ , with second moment

$$\mathbb{E} \left[ (c^\top \xi_k)^2 \right] = \sum_{i=1}^n \sum_{j=1}^n c_i c_j \mathbb{E} \left[ \xi_n^{(i)} \xi_n^{(j)} \right] = \sum_{i=1}^n \sum_{j=1}^n c_i c_j \Xi_{ij} = c^\top \Xi c \geq 0.$$

The limit  $c^\top X^n \xrightarrow{d} c^\top W$  in the case  $\mathbb{E} \left[ (c^\top \xi_k)^2 \right] = 0$  is trivial, hence we may assume from now on that  $\mathbb{E} \left[ (c^\top \xi_k)^2 \right] > 0$ . Then by [6], Theorem 18.3, it follows that  $c^\top X^n$  converges weakly on  $D([0, 1], \mathbb{R})$  equipped with the Skorokhod  $J_1$  topology to a Wiener process with covariance  $u c^\top \Xi c$ , i.e., to a limit  $c^\top W$ , where  $W$  is a  $d$ -variate Wiener process with covariance  $u\Xi$ .  $\square$

The next result establishes Step 1.

**Theorem 1.** *Grant Assumptions A. Let  $\lambda_{\theta_0}$  be the conditional intensity of the point process  $N_{\theta_0} \in \mathcal{F}_{\Theta}$ . Then it holds that*

$$\left( \frac{1}{\sqrt{T}} \left( N_{\theta_0}(uT) - \int_0^{uT} \lambda_{\theta_0}(s) \, ds \right) \right)_{u \in [0, 1]} \xrightarrow{d} (W_{\theta_0}(u))_{u \in [0, 1]}, \quad (10)$$

as  $T \rightarrow \infty$ , weakly on  $D([0, 1], \mathbb{R}^d)$  equipped with the strong Skorokhod  $J_1$  topology, where  $W_{\theta_0}$  is a  $d$ -variate Wiener process with covariance

$$u \cdot \text{diag} \left( \mathbb{E}[N_{\theta_0}^{(1)}[0, 1]], \dots, \mathbb{E}[N_{\theta_0}^{(d)}[0, 1]] \right).$$

*Proof.* The result follows by an application of Lemma 2 to  $\xi_n = N_{\theta_0}[n, n+1) - \int_n^{n+1} \lambda_{\theta_0}(s) \, ds$ , combined with an estimate of continuous-time quantities by discrete-time quantities analogous to [54], eqn. (2.19). We use the calculation

$$\mathbb{E} \left[ \left( N_{\theta_0}^{(i)}(1) - \int_0^1 \lambda_{\theta_0}^{(i)}(s) \, ds \right) \left( N_{\theta_0}^{(j)}(1) - \int_0^1 \lambda_{\theta_0}^{(j)}(s) \, ds \right) \right] = \begin{cases} \mathbb{E} \left[ N_{\theta_0}^{(i)}[0, 1] \right] & \text{if } i = j; \\ 0 & \text{if } i \neq j, \end{cases}$$

which follows from [14], Proposition 14.1.VIII.  $\square$

**3.2. Step 2: Limit theorem for difference between real and empirical compensator.** For the second step of our procedure, we analyze the second term of (8).

**Theorem 2.** *Grant Assumptions A–C. Let  $\alpha_{\theta_0} \in \mathbb{R}^{d \times m}$  be as in Assumption B(vi) and let  $Z$  be as in Assumption C. Then we have, under the law  $\mathcal{P}_{\theta_0}$  of  $N_{\theta_0} \in \mathcal{F}_{\Theta}$ ,*

$$\left( \frac{1}{\sqrt{T}} \int_0^{uT} \left( \lambda_{\theta_0}(s) - \lambda_{\hat{\theta}_T}(s) \right) \, ds \right)_{u \in [0, 1]} \rightarrow (u \alpha_{\theta_0} Z)_{u \in [0, 1]}, \quad (11)$$

as  $T \rightarrow \infty$ , weakly on  $D([0, 1], \mathbb{R}^d)$  equipped with the strong Skorokhod  $J_1$  topology.

*Proof.* Our proof is structured as follows. Assumptions C grants the CLT  $\sqrt{T}(\theta_0 - \hat{\theta}_T) \xrightarrow{d} Z$ . This allows us to perform a Taylor expansion on the integrand of the prelimit from (11):  $\lambda_{\theta_0}(s) - \lambda_{\hat{\theta}_T}(s) = D\lambda_{\theta_0}(s)(\theta_0 - \hat{\theta}_T) + o_{\mathbb{P}}(|\theta_0 - \hat{\theta}_T|)$ . We use this to rewrite the prelimit from (11) to

$$\frac{1}{\sqrt{T}} \int_0^{uT} (\lambda_{\theta_0}(s) - \lambda_{\hat{\theta}_T}(s)) ds = u \frac{1}{uT} \int_0^{uT} D\lambda_{\theta_0}(s) ds \sqrt{T}(\theta_0 - \hat{\theta}_T) + o_{\mathbb{P}}(1). \quad (12)$$

We prove the result by showing that the  $i$ th coordinate of the right-hand side (12) converges uniformly in probability to the  $i$ th coordinate of the right-hand side of (11). From this, convergence in the Skorokhod topology follows.

Denote the  $k$ th row of  $\alpha_{\theta_0}$  by  $\alpha_{\theta_0}^{(k)}$ , i.e.,

$$\alpha_{\theta_0}^{(k)} := \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \frac{\partial \lambda_{\theta}^{(k)}(s)}{\partial \theta} ds \in \mathbb{R}^m.$$

We need to show that, for each  $k \in [d]$ ,

$$u \frac{1}{uT} \int_0^{uT} \frac{\partial \lambda_{\theta}^{(k)}(s)}{\partial \theta} ds \sqrt{T}(\theta_0 - \hat{\theta}_T) \rightarrow u \alpha_{\theta_0}^{(k)} Z, \quad (13)$$

uniformly in probability. The second factor  $\sqrt{T}(\theta_0 - \hat{\theta}_T)$  is not dependent on  $u$  and converges to  $Z$ , by Assumptions C, hence uniform convergence in probability  $\sqrt{T}(\theta_0 - \hat{\theta}_T) \rightarrow Z$  is immediate. Indeed, by the Skorokhod representation theorem, there exists a version of  $\sqrt{T}(\theta_0 - \hat{\theta}_T)$  which converges *in probability* to a random variable with distribution  $Z$ .

For the first factor

$$u \frac{1}{uT} \int_0^{uT} \frac{\partial \lambda_{\theta}^{(k)}(s)}{\partial \theta} ds$$

note that by Assumption B(iii),  $\|\partial \lambda_{\theta}^{(k)}(0)/\partial \theta\|$  has finite expectation, hence by stationarity (Assumption A(ii)), the same holds for  $\|\partial \lambda_{\theta}^{(k)}(s)/\partial \theta\|$ ,  $s \geq 0$ . In fact, it follows that  $\mathbb{E}\|\partial \lambda_{\theta}^{(k)}(s)/\partial \theta\|$  is independent of time  $s$ . By Fubini's theorem, we infer that

$$\mathbb{E} \left[ \frac{1}{T} \int_0^{uT} \left\| \frac{\partial \lambda_{\theta}^{(k)}(s)}{\partial \theta} \right\| ds \right] = \frac{1}{T} \int_0^{uT} \mathbb{E} \left\| \frac{\partial \lambda_{\theta}^{(k)}(s)}{\partial \theta} \right\| ds = u \mathbb{E} \left\| \frac{\partial \lambda_{\theta}^{(k)}(0)}{\partial \theta} \right\|.$$

Fix  $\delta, \epsilon > 0$ . For  $u \in [0, 1]$ ,  $T > 0$ , an application of Markov's inequality yields that with probability of at least  $1 - \delta/2$  it holds for all  $u' \in [0, u]$  that

$$\frac{1}{T} \int_0^{u'T} \left\| \frac{\partial \lambda_{\theta}^{(k)}(s)}{\partial \theta} \right\| ds \leq \frac{2}{\delta} u \mathbb{E} \left\| \frac{\partial \lambda_{\theta}^{(k)}(0)}{\partial \theta} \right\|.$$

Note that this bound is independent of  $T > 0$ . Hence, selecting

$$u^{(k)} = \frac{\epsilon}{4} \left( \delta \left( \mathbb{E} \left\| \frac{\partial \lambda_{\theta}^{(k)}(0)}{\partial \theta} \right\| \right) \right)^{-1} \wedge \frac{1}{\|\alpha_{\theta_0}\|}$$

shows that with probability of at least  $1 - \delta/2$

$$\sup_{u \in [0, u^{(k)}]} \left\| \frac{1}{T} \int_0^{uT} \frac{\partial \lambda_{\theta}^{(k)}(s)}{\partial \theta} ds - u \alpha_{\theta_0}^{(k)} \right\| \leq \epsilon.$$

Given  $u^{(k)}$ , we use Assumption B(vi) to select  $T^{(k)}$  such that for  $T \geq T^{(k)}$  and  $u \in [u^{(k)}, 1]$  it holds with probability of at least  $1 - \delta/2$  that

$$\left\| u \left( \frac{1}{uT} \int_0^{uT} \frac{\partial \lambda_{\theta}^{(k)}(s)}{\partial \theta} ds - \alpha_{\theta_0}^{(k)} \right) \right\| \leq \epsilon.$$



We have shown that on an event of probability at least  $1 - \delta$ , for  $T \geq T^{(k)}$  it holds that

$$\sup_{u \in [0,1]} \left\| \frac{1}{T} \int_0^{uT} \frac{\partial \lambda_{\theta}^{(k)}(s)}{\partial \theta} ds - u \alpha_{\theta_0}^{(k)} \right\| \leq \epsilon.$$

In other words,  $\frac{1}{T} \int_0^{uT} \frac{\partial \lambda_{\theta}^{(k)}(s)}{\partial \theta} ds$  converges uniformly in probability to  $u \alpha_{\theta_0}^{(k)}$ .

Next, by a simple union bound similar to the one used in the proof of [51], Theorem 2.7(vi), it follows that

$$\left( \frac{1}{T} \int_0^{uT} \frac{\partial \lambda_{\theta}^{(k)}(s)}{\partial \theta} ds, \sqrt{T}(\theta_0 - \hat{\theta}_T) \right) \rightarrow \left( u \alpha_{\theta_0}^{(k)}, Z \right),$$

uniformly in probability. By an application of the continuous mapping theorem (to the function  $g(x, y) = x \cdot y$ ) it follows that (13) holds, uniformly over  $u \in [0, 1]$ , in probability.

We extend to (11) by selecting  $u^* = \min_{k \in [d]} u^{(k)}$  and  $T^* = \max_{k \in [d]} T^{(k)}$ .

Note that uniform convergence in probability amounts to uniform convergence on a set of probability  $1 - \delta/2$ . Therefore, conditional on a high probability set, convergence in the Skorokhod  $J_1$  topology holds, hence the tightness conditions for the  $J_1$  topology hold. Conditional on a set of probability  $1 - \delta/2$ , the probabilities appearing in the tightness conditions can be bounded by  $\delta/2$ . Therefore, they can even be bounded by  $\delta$ . Hence, uniform convergence in probability implies convergence in the Skorokhod  $J_1$  topology.  $\square$

**Remark 5.** *Since the limit  $u \mapsto u \alpha_{\theta_0} Z$  is continuous, it follows that (11) even holds on  $D([0, 1], \mathbb{R}^d)$  equipped with the uniform topology.*

Combining Theorem 1 with Theorem 2, we have the following limit result for the empirical process  $\hat{\eta}^{(T)}$  defined in (7).

**Corollary 1.** *Grant Assumptions A–C. Let  $W_{\theta_0}$  be a  $d$ -variate Wiener process with covariance*

$$u \cdot \text{diag} \left( \mathbb{E}[N_{\theta_0}^{(1)}[0, 1]], \dots, \mathbb{E}[N_{\theta_0}^{(d)}[0, 1]] \right).$$

*Then it holds under the law  $\mathcal{P}_{\theta_0}$  of  $N_{\theta_0} \in \mathcal{F}_{\Theta}$  that*

$$\left( \frac{1}{\sqrt{T}} \left( N_{\theta_0}(uT) - \int_0^{uT} \lambda_{\hat{\theta}_T}(s) ds \right) \right)_{u \in [0,1]} \xrightarrow{d} (W_{\theta_0}(u) + u \alpha_{\theta_0} Z)_{u \in [0,1]} =: (\hat{\eta}_{\theta_0}(u))_{u \in [0,1]}, \quad (14)$$

*as  $T \rightarrow \infty$ , weakly on  $D([0, 1], \mathbb{R}^d)$  equipped with the strong Skorokhod  $J_1$  topology.*

**3.3. Step 3: Transforming  $\hat{\eta}_{\theta_0}$  into a standard Wiener process.** The limiting result (14) derived in Step 2 still depends on the true but unknown parameter, through the covariance of  $W_{\theta_0}$  and the linear stochastic drift term  $u \alpha_{\theta_0} Z$ . Due to this dependency on the true parameter, we cannot directly use this limit to construct an asymptotically valid testing procedure for all  $\theta \in \Theta$ , causing the theory to become fragmented.

To address this issue, we apply a suitable *innovation martingale transform* to the limit process, which results in a new limit process that is independent of the true parameter  $\theta_0$  and of the specific family  $\mathcal{F}_{\Theta}$  we are testing for. Variants of innovation martingale transformations have been applied to various statistical problems in the literature over the last couple of decades; see [15, 16, 27, 28, 29, 30, 35, 36, 47].

An innovation martingale transformation has also been applied to processes involving populations of size  $n$  over fixed time horizons  $[0, T]$ , where an asymptotically distribution-free test process has been derived in the limit as  $n \rightarrow \infty$  [4, 48]. However, that large- $n$  setup differs fundamentally from the large- $T$  setup considered in this work. In the large- $n$  regime, one considers a system consisting of many interacting particles over a fixed time horizon. The limiting behavior as  $n \rightarrow \infty$  is typically governed by an LLN or CLT, where

randomness is averaged out due to aggregation across the population. The resulting asymptotics are then driven by scaling properties of the population-level process. By contrast, the large- $T$  regime concerns a single realization of a process evolving over an increasingly long time horizon. Here, the asymptotics are dictated by the temporal structure of the process rather than by population averaging. Growth in  $T$  does not inherently reduce stochasticity but instead reveals long-term dependencies and fluctuation behavior that are absent in the large- $n$  setting. As a result, techniques that rely on concentration due to population size do not carry over directly, and different probabilistic tools are required to understand the limiting behavior.

**Theorem 3.** *Grant Assumptions A–C. Let  $\mu_{\theta_0}^{(k)} := \mathbb{E}[N_{\theta_0}^{(k)}[0, 1]]$ . For  $\theta \in \Theta$ , let  $\mathcal{T}_\theta : D([0, 1], \mathbb{R}^d) \rightarrow D([0, 1], \mathbb{R}^d)$  be the transformation*

$$\mathcal{T}_\theta(\eta)(B) := \left\{ \frac{1}{\sqrt{\mu_{\theta_0}^{(k)}}} \left( \eta^{(k)}(B) - \int_B \frac{\eta^{(k)}(1) - \eta^{(k)}(v)}{1 - v} dv \right) \right\}_{k \in [d]}, \quad B \in \mathcal{B}[0, 1]. \quad (15)$$

Then it holds under the law  $\mathcal{P}_{\theta_0}$  of  $N_{\theta_0} \in \mathcal{F}_\Theta$  that  $\mathcal{T}_{\theta_0}(\hat{\eta}_{\theta_0})$  is a standard  $d$ -dimensional Wiener process, where  $\hat{\eta}_{\theta_0}$  is defined in (14).

*Proof.* Each coordinate of the limit  $\hat{\eta}_{\theta_0}$  found in Corollary 1 is of the general form treated in [8], Section 3. Fix a coordinate  $k \in [d]$ . We can specify [8], Theorem 3.1, to univariate time frames by taking functions independent of the second coordinate. We identify the objects appearing in that theorem. We take the scanning family  $A_v = [0, v]$  on  $[0, 1]$ . We have  $R(u) = \mu_{\theta_0}^{(k)} u$  and

$$Q(B) = \int_B dQ(u) = \int_B du = \text{Leb}(B) = \int_B q(u) dR(u) = \mu_{\theta_0}^{(k)} \int_B q(u) du,$$

hence  $q(u) = 1/\mu_{\theta_0}^{(k)}$ . Next, we have

$$I(A_v^c) = \int_{A_v^c} q(u)^2 dR(u) = \int_v^1 \frac{1}{(\mu_{\theta_0}^{(k)})^2} \mu_{\theta_0} du = \frac{1 - v}{\mu_{\theta_0}^{(k)}}.$$

Note that  $Q(B \cap A_{dv}) = Q(B \cap \{dv\}) = \mathbf{1}\{v \in B\} dv$ . Then [8], Theorem 3.1, gives that

$$\eta_{\theta_0}^{(k)}(B) - \int_B \frac{\eta_{\theta_0}^{(k)}(1) - \eta_{\theta_0}^{(k)}(v)}{1 - v} dv \quad (16)$$

is a Wiener process with variance  $\mu_{\theta_0}^{(k)} u$ , implying that  $\mathcal{T}_{\theta_0}(\hat{\eta}_{\theta_0})$  is a standard  $d$ -dimensional Wiener process.

Indeed,  $\mathcal{T}_{\theta_0}(\hat{\eta}_{\theta_0})$  is a  $d$ -dimensional Gaussian process with zero mean, hence its probabilistic behavior is characterized by its covariance structure  $C_{ij}(u) := \text{Cov}(\mathcal{T}_{\theta_0}(\hat{\eta}_{\theta_0})_i(u), \mathcal{T}_{\theta_0}(\hat{\eta}_{\theta_0})_j(u))$ ,  $i, j \in [d]$ ,  $u \in [0, 1]$ . Using linearity of  $\mathcal{T}_{\theta_0}^{(i)}$  and  $\mathcal{T}_{\theta_0}^{(j)}$  in  $\eta$  together with the definition on  $\hat{\eta}_{\theta_0}$  given in (14), we expand

$$\begin{aligned} C_{ij}(u) &= \text{Cov} \left( \mathcal{T}_{\theta_0}^{(i)}(W_{\theta_0}^{(i)}(u)), \mathcal{T}_{\theta_0}^{(j)}(W_{\theta_0}^{(j)}(u)) \right) + \text{Cov} \left( \mathcal{T}_{\theta_0}^{(i)}(W_{\theta_0}^{(i)}(u)), \mathcal{T}_{\theta_0}^{(j)}(u\alpha_{\theta_0}^{(j)} Z) \right) \\ &\quad + \text{Cov} \left( \mathcal{T}_{\theta_0}^{(i)}(u\alpha_{\theta_0}^{(i)} Z), \mathcal{T}_{\theta_0}^{(j)}(W_{\theta_0}^{(j)}(u)) \right) + \text{Cov} \left( \mathcal{T}_{\theta_0}^{(i)}(u\alpha_{\theta_0}^{(i)} Z), \mathcal{T}_{\theta_0}^{(j)}(u\alpha_{\theta_0}^{(j)} Z) \right). \end{aligned}$$

For  $i \neq j$ , the first term vanishes because of independence between  $W_{\theta_0}^{(i)}$  and  $W_{\theta_0}^{(j)}$ , while the remaining three terms vanish because  $\mathcal{T}_{\theta_0}^{(i)}(u\alpha_{\theta_0}^{(k)} Z) \equiv 0$  for each  $k \in [d]$ . Therefore,  $C_{ij}(u) \equiv 0$ .  $\square$

**3.4. Step 4: Replacing  $\theta_0$  by  $\hat{\theta}_T$ .** Although the limit process from Theorem 3 does not depend on the true parameter, the transformation  $\mathcal{T}_{\theta_0}$  and the process  $(\hat{\eta}_{\theta_0})$  still do. Hence, in practical applications  $\mathcal{T}_{\theta_0}(\hat{\eta}_{\theta_0})$  cannot be used. Instead, the best one could do is to replace it by  $\mathcal{T}_{\hat{\theta}_T}(\hat{\eta}^{(T)})$  where  $\hat{\eta}^{(T)}$  is the empirical process defined in (7). Here, we show that the difference between  $\mathcal{T}_{\hat{\theta}_T}(\hat{\eta}^{(T)})$  and  $\mathcal{T}_{\theta_0}(\hat{\eta}_{\theta_0})$  vanishes in the limit as  $T \rightarrow \infty$ . Since  $\mathcal{T}_{\theta_0}(\hat{\eta}_{\theta_0})$  is a standard  $d$ -dimensional Wiener process, this provides us with

an asymptotically distribution-free limit process. Also, the quantity  $\mathcal{T}_{\hat{\theta}_T}(\hat{\eta}^{(T)})$  can be calculated explicitly without knowledge of  $\theta_0$ .

**Theorem 4.** *Grant Assumptions A–C. Select some  $\tau \in (0, 1)$ . Then under the law  $\mathcal{P}_{\theta_0}$  of  $N_{\theta_0} \in \mathcal{F}_{\Theta}$ ,  $\mathcal{T}_{\hat{\theta}_T}(\hat{\eta}^{(T)})$  converges weakly on  $D([0, \tau], \mathbb{R}^d)$  equipped with the strong Skorokhod  $J_1$ -topology to a standard Wiener process, as  $T \rightarrow \infty$ .*

*Proof.* The proof consists of two parts. First, we show that  $\mu_{\hat{\theta}_T} \rightarrow \mu_{\theta_0}$  as  $T \rightarrow \infty$ . Second, we show that

$$\tilde{\mathcal{T}} := \text{diag} \left( \sqrt{\mu_{\theta_0}^{(1)}}, \dots, \sqrt{\mu_{\theta_0}^{(d)}} \right) \mathcal{T}_{\theta}$$

satisfies  $\tilde{\mathcal{T}}(\hat{\eta}^{(T)}) \rightarrow \tilde{\mathcal{T}}(\hat{\eta}_{\theta_0})$  uniformly on  $[0, \tau] \ni u$  in probability as  $T \rightarrow \infty$ ; note that  $\tilde{\mathcal{T}}$  could also be defined through the formula

$$\tilde{\mathcal{T}}(\eta)(B) := \left\{ \eta^{(k)}(B) - \int_B \frac{\eta^{(k)}(1) - \eta^{(k)}(v)}{1 - v} dv \right\}_{k \in [d]}, \quad B \in \mathcal{B}[0, 1].$$

For the first part, by stationarity (Assumption A(ii)) it holds that  $\mu_{\theta_0}^{(k)} = \mathbb{E}[\lambda_{\theta_0}^{(k)}(0)]$ . By Assumption C, it holds that  $\hat{\theta}_T \rightarrow \theta_0$  in probability, as  $T \rightarrow \infty$ . By passing to a subsequence, this convergence can be taken a.s. By Assumption B(iii) and the continuous mapping theorem,  $\lambda_{\hat{\theta}_T}^{(k)}(0) \rightarrow \lambda_{\theta_0}^{(k)}(0)$  a.s., as  $T \rightarrow \infty$ . By Assumption B(v) and dominated convergence, it follows that  $\mu_{\hat{\theta}_T}^{(k)} \rightarrow \mu_{\theta_0}^{(k)}$  as  $T \rightarrow \infty$ .

By Assumption B(iii), it holds that  $\mu_{\theta_0}^{(k)} > 0$ , hence to prove the theorem it suffices to prove the second part. To this end, we apply the Skorokhod representation theorem to the result of Corollary 1 to obtain a probability space supporting probabilistic equivalent versions of  $\hat{\eta}^{(T)}$  and  $\hat{\eta}_{\theta_0}$  satisfying

$$\sup_{u \in [0, \tau]} \left\| \hat{\eta}^{(T)}(u) - \hat{\eta}_{\theta_0}(u) \right\|_{\infty} \rightarrow 0 \quad \text{a.s.} \quad (17)$$

We will work on this probability space.

To prove that  $\tilde{\mathcal{T}}(\hat{\eta}^{(T)}) \rightarrow \tilde{\mathcal{T}}(\hat{\eta}_{\theta_0})$  uniformly on  $[0, \tau] \ni u$  in probability, it suffices to prove for  $k \in [d]$  that

$$\sup_{u \in [0, \tau]} \left| \int_0^u \frac{\Delta_k^{(T)}(1) - \Delta_k^{(T)}(v)}{1 - v} dv \right| \rightarrow 0 \quad (18)$$

in probability, as  $T \rightarrow \infty$ , i.e., to prove ucp convergence. Here  $\Delta^{(T)} := \hat{\eta}^{(T)} - \hat{\eta}_{\theta_0}$  denotes the difference between the empirical process from Step 2 and its limit. Since  $\frac{1}{1-v}$  is bounded on  $[0, \tau]$ , we simply infer the convergence from (18) from (17).  $\square$

#### 4. GOODNESS-OF-FIT TESTING PROCEDURES

**4.1. Asymptotically correct goodness-of-fit test based on the test process.** Given a sample  $\omega_T$  on  $[0, T]$  of a point process  $T$ , we are interested in testing the parametric hypothesis

$$H_0 : N \stackrel{d}{=} N_{\theta} \text{ for some } \theta \in \Theta. \quad (19)$$

In a practical situation, the sample  $\omega_T$  would be used to calculate an estimator  $\hat{\theta}_T$  for  $\theta$ , after which one asks whether  $\mathcal{F}_{\Theta} = \{N_{\theta} : \theta \in \Theta\}$  is indeed a suitable model for the data. To answer this question, we use Theorem 4 to construct a goodness-of-fit test for testing the null hypothesis  $H_0$  given in (19). The ideas of this subsection will be implemented in Sections 6–7. In Section 5, we consider consistency under alternatives for this goodness-of-fit test.

In Section 3, we proved in Theorem 4 that for any  $\tau \in (0, 1)$ , the process  $\mathcal{T}_{\hat{\theta}_T}(\hat{\eta}^{(T)})$  converges weakly on  $D([0, \tau], \mathbb{R}^d)$  to a standard Wiener process  $W$ . The process  $\mathcal{T}_{\hat{\theta}_T}(\hat{\eta}^{(T)})$  only depends on the sample  $\omega_T$ . Therefore, it can be calculated explicitly. Now suppose  $\mathcal{K}$  is a test statistic depending continuously on  $\mathbb{R}^d$ -valued stochastic processes on  $[0, \tau]$ . Then  $\mathcal{K}(\mathcal{T}_{\hat{\theta}_T}(\hat{\eta}^{(T)}))$  converges in distribution to  $\mathcal{K}(W)$ .

The limit process  $W$  from Theorem 4 is a standard Wiener process on  $[0, \tau]$ . In particular, for any  $n \in \mathbb{N}$ , the random variables

$$Z_i := \sqrt{\frac{n}{\tau}} (W(i\tau/n) - W((i-1)\tau/n)), \quad i \in [n], \quad (20)$$

compose an i.i.d. sample of size  $n$  from the  $d$ -dimensional standard normal distribution.

Suppose that one replaces  $Z_i$  by  $\hat{Z}_i^{(T)}$  based on  $\hat{W}^{(T)} := \mathcal{T}_{\hat{\theta}_T}(\hat{\eta}^{(T)})$  instead of  $W$ ; that is,

$$\hat{Z}_i^{(T)} := \sqrt{\frac{n}{\tau}} (\hat{W}^{(T)}(i\tau/n) - \hat{W}^{(T)}((i-1)\tau/n)), \quad i \in [n]. \quad (21)$$

By the preceding arguments, it follows that  $(\hat{Z}_i^{(T)})_{i \in [n]}$  converges to a sample of standard normal random variables, as  $T \rightarrow \infty$  in such a way that  $T/n = T/n(T) \rightarrow \infty$ . Hence, replacing parametric null (19) by

$$H_0^{(T)} : (\hat{Z}_1^{(T)}, \dots, \hat{Z}_{n(T)}^{(T)}) \text{ is an i.i.d. sample from the } \mathcal{N}(0, I_d)\text{-distribution,} \quad (22)$$

where  $I_d$  denotes the  $d \times d$  identity matrix, leads to asymptotically correct tests, i.e., tests having the correct rejection rates under  $H_0$ , asymptotically, as  $T \rightarrow \infty$ . The hypothesis (22) can be evaluated using any normality test: well-known examples include (multivariate versions of) the Kolmogorov-Smirnov, Anderson-Darling and Cramér-von Mises tests.

In order to obtain high power under alternatives, we want to take a large sample size  $n = n(T)$ . However, for fixed  $T$ , a lower value of this hyperparameter  $n$  amounts to more data being used to calculate  $\hat{Z}_i^{(T)}$ , hence better approximations to standard normals, yielding a more robust testing procedure. In other words, there is a trade-off between small and large  $n$ . One way to find a middle ground is by taking  $n = \text{ceil}(c\sqrt{T})$  for some  $c > 0$ , where  $\text{ceil}(\cdot)$  denotes the ceiling function. A lower choice of  $c$  prioritizes the robustness of the testing procedure, while a higher  $c$  prioritizes power. For this choice of the hyperparameter  $n$ , given a sample  $\omega_T$  of a point process  $N$  on  $[0, T]$ , one may test (19) using Algorithm 1.

In the present situation, a ‘naive’ testing procedure ignoring estimation uncertainty can be composed by simply treating the (standardized) empirical process

$$\tilde{W}^{(T)} := \text{diag} \left( \sqrt{\mu_{\theta_0}^{(1)}}, \dots, \sqrt{\mu_{\theta_0}^{(d)}} \right)^{-1} \hat{\eta}^{(T)} \quad (23)$$

as a standard Wiener process, e.g., by testing the random variables

$$\tilde{Z}_i^{(T)} := \sqrt{\frac{n}{\tau}} (\tilde{W}^{(T)}(i\tau/n) - \tilde{W}^{(T)}((i-1)\tau/n)), \quad i \in [n]. \quad (24)$$

for normality. This amounts to leaving out step (iii) of Algorithm 1. In Section 6, we will compare this ‘naive’ testing procedure empirically to our asymptotically correct testing procedure, and we show empirically that it is generally a bad idea to ignore the estimation uncertainty  $\hat{\theta}_T \neq \theta_0$ .

**Remark 6.** We know from Corollary 1 that  $\tilde{W}^{(T)}$  converges to a Wiener process plus a random linear drift term. The magnitude of the bias of  $\tilde{Z}_i^{(T)}$  caused by this drift is then of order  $1/\sqrt{n}$ . Since test statistics based on a functional of the empirical process — such as the Kolmogorov-Smirnov, Anderson-Darling and Cramér-von Mises tests — converge at rate  $\sqrt{n(T)} \asymp \sqrt{T}$ , the bias of the ‘naive’ testing procedure cannot be mitigated through a choice of the hyperparameter  $n$ .

**Algorithm 1** Asymptotically correct goodness-of-fit test based on transformed empirical process

**Input:** A realization  $\omega_T$  of a point process  $N$  on  $[0, T]$  and hyperparameters  $\tau \in (0, 1)$  and  $n = n(T)$ ; e.g., use  $n = \text{ceil}(c\sqrt{T})$  for some  $c > 0$

**Output:** Outcome of a goodness-of-fit test

(i) Estimate  $\theta$  using an estimator  $\hat{\theta}_T$  satisfying Assumption C; e.g., use the MLE

(ii) Compute the compensated empirical process  $\hat{\eta}^{(T)}$ :

$$\hat{\eta}^{(T)} : u \mapsto \frac{1}{\sqrt{T}} \left( N(uT) - \int_0^{uT} \lambda_{\hat{\theta}_T}(s) ds \right)$$

(iii) For  $\mu_{\hat{\theta}_T}^{(k)} := \mathbb{E}[N_{\hat{\theta}_T}^{(k)}[0, 1]] \approx N^{(k)}(T)/T$ , compute the transformed process  $\hat{W}^{(T)} := \mathcal{T}_{\hat{\theta}_T}(\hat{\eta}^{(T)})$  using

$$\mathcal{T}_{\hat{\theta}_T}(\hat{\eta}^{(T)})(u) := \left\{ \frac{1}{\sqrt{\mu_{\hat{\theta}_T}^{(k)}}} \left( (\hat{\eta}^{(T)})^{(k)}(u) - \int_0^u \frac{(\hat{\eta}^{(T)})^{(k)}(1) - (\hat{\eta}^{(T)})^{(k)}(v)}{1-v} dv \right) \right\}_{k \in [d]}$$

(iv) For  $i \in [n]$ , compute:

$$\hat{Z}_i^{(T)} := \sqrt{\frac{n}{\tau}} \left( \hat{W}^{(T)}(i\tau/n) - \hat{W}^{(T)}((i-1)\tau/n) \right)$$

(v) Perform an (asymptotically) exact normality test on the sample  $(\hat{Z}_i^{(T)})_{i \in [n]}$ ; e.g., use the Kolmogorov-Smirnov, Anderson-Darling or Cramér-von Mises test

**4.2. Random-time-change-based goodness-of-fit testing.** Typically, when goodness-of-fit tests are conducted in the context of univariate point processes, one considers transformed event times  $\Lambda_{\hat{\theta}_T}(t_i)$ , where  $\Lambda_{\theta}(t) = \int_0^t \lambda_{\theta}(t) dt$  is the *compensator* of the point process, i.e., the time-integrated conditional intensity. By the random time change theorem [14], Theorem 7.4.I, under  $\mathcal{P}_{\theta_0}$  the times  $\Lambda_{\theta_0}(t_i)$  are distributed according to a unit Poisson process. Hence, a goodness-of-fit test is performed by testing whether  $\Lambda_{\hat{\theta}_T}(t_i) - \Lambda_{\hat{\theta}_T}(t_{i-1})$  are standard exponential.<sup>1</sup> However, the estimation uncertainty  $\hat{\theta}_T \neq \theta_0$  leading to a difference between  $\Lambda_{\hat{\theta}_T}$  and  $\Lambda_{\theta_0}$  is typically ignored; see [3, 10, 11, 19, 31, 32, 34, 49, 50]. Even though the goodness-of-fit test outlined in Algorithm 1 relies on different ideas than this test based on the random time change theorem, it is, of course, possible to compare the outcomes of both tests.

We now consider the effect of estimation uncertainty. By Theorem 2 with  $u = t_i/T = O(1/T)$ , holding for fixed  $t_i$  it follows that, as  $T \rightarrow \infty$ ,

$$\Lambda_{\hat{\theta}_T}(t_i) - \Lambda_{\hat{\theta}_T}(t_{i-1}) = \Lambda_{\theta_0}(t_i) - \Lambda_{\theta_0}(t_{i-1}) - \frac{1}{\sqrt{T}}(t_i - t_{i-1})\alpha_{\theta_0}Z + o_{\mathbb{P}}(1/\sqrt{T}). \quad (25)$$

By the random time change theorem, this means that the transformed interarrival times  $\Lambda_{\hat{\theta}_T}(t_i) - \Lambda_{\hat{\theta}_T}(t_{i-1})$  approximately equal standard exponential random variables plus some bias term proportional to the interarrival time  $t_i - t_{i-1}$ . This bias term leads to an incorrect goodness-of-fit testing procedure, assessing whether the times  $(\Lambda_{\hat{\theta}_T}(t_i))$  compose a realization of a unit Poisson process. In fact, when one uses a test statistic that is a functional of the empirical process — such as the Kolmogorov-Smirnov, Anderson-Darling and Cramér-von Mises tests — to test the transformed interarrival times, the convergence of the tests statistics occurs at rate  $\sqrt{n(T)} \asymp \sqrt{T}$ . Therefore, the deviations  $\frac{1}{\sqrt{T}}(t_i - t_{i-1})\alpha_{\theta_0}Z + o_{\mathbb{P}}(1/\sqrt{T}) = O_{\mathbb{P}}(1/\sqrt{T})$  in (25) from standard exponentials are in general *not* negligible.

<sup>1</sup>By a multivariate version of the random time change theorem, [14], Theorem 14.6.IV, this testing procedure allows for a multivariate generalization.

If a consistent estimate of  $\alpha_{\theta_0}Z$  could be obtained from the data, one could modify (25) to develop an asymptotically distribution-free goodness-of-fit test based on the random time change theorem. However, this is not a straightforward task.

### 5. CONSISTENCY UNDER $H_1$ OF THE GOODNESS-OF-FIT TEST BASED ON THE TEST PROCESS

In step (iii) of Algorithm 1, we transform the compensated empirical process  $\hat{\eta}^{(T)}$  to  $\hat{W}^{(T)} := \mathcal{T}_{\hat{\theta}_T}(\hat{\eta}^{(T)})$  using the innovation martingale transform defined in (7). This transformed process converges to a process independent of the true parameter  $\theta_0$ , as well as of the model  $\mathcal{F}_{\Theta}$ , hence leads to asymptotically correct goodness-of-fit tests. However, we notice that due to the transformation information may be lost, see [24]. In other words, it may be possible that a test based on the transformed process  $\hat{W}^{(T)}$  is incapable of detecting certain deviations from the null that other tests might be able to detect. In this subsection, we discuss the consistency under  $H_1$  of a goodness-of-fit testing procedure outlined in Algorithm 1, where for the normality test in step (v) we consider, for example, a Kolmogorov-Smirnov, an Anderson-Darling or a Cramér-von Mises test. Those tests are consistent under alternatives, meaning that if one tests a simple null  $H_0 : X \stackrel{d}{=} Y$ , then for a sample of  $Z \stackrel{d}{\neq} Y$ , the power of the test converges to 1.

**Theorem 5.** *Grant Assumptions A–B and Assumptions C.1. Suppose that we observe a realization of a stationary and ergodic point process  $N$  having law  $\mathcal{P}$  and intensity  $\lambda \notin \mathcal{L}_{\Theta} := \{\lambda_{\theta} : \theta \in \Theta\}$ ; i.e.,  $N$  does not belong to the null (19). Fit  $N$  to the parametric hypothesis  $H_0$  by calculating the maximum likelihood estimator  $\hat{\theta}_T$ . If  $\lambda$  does not satisfy*

$$\lambda \stackrel{d}{=} \lambda_{\theta} + c \quad \text{for some } \theta \in \Theta \quad \text{and some } c \in \mathbb{R}^d,$$

*and if one tests the null (19) using Algorithm 1, using a goodness-of-fit test that is consistent under alternatives at rate  $\sqrt{n(T)}$  in step (v), then the power of the goodness-of-fit test given in Algorithm 1 converges to 1.*

*Proof.* The maximum likelihood estimator  $\hat{\theta}_T$  is consistent in probability for the ‘least-false’ estimator  $\theta^* \in \Theta$  maximizing the expected likelihood  $\mathbb{E}_{\mathcal{P}}[L_T(\theta)]$ ; see [20], Section 2.2. In particular, a CLT  $\sqrt{T}(\theta^* - \hat{\theta}_T) \xrightarrow{d} Z$ , as  $T \rightarrow \infty$ , still holds; cf. Assumption C.1.

Select  $\tau \in (0, 1)$  as in Theorem 4 and work on  $[0, \tau] \ni u$ . Recall that  $\hat{W}^{(T)} := \mathcal{T}_{\hat{\theta}_T}(\hat{\eta}^{(T)})$ , where

$$\begin{aligned} \hat{\eta}^{(T)}(u) &= \frac{1}{\sqrt{T}} \left( N(uT) - \int_0^{uT} \lambda(s) ds \right) + \frac{1}{\sqrt{T}} \left( \int_0^{uT} \lambda(s) ds - \int_0^{uT} \lambda_{\theta^*}(s) ds \right) \\ &\quad + \frac{1}{\sqrt{T}} \left( \int_0^{uT} \lambda_{\theta^*}(s) ds - \int_0^{uT} \lambda_{\hat{\theta}_T}(s) ds \right). \end{aligned} \quad (26)$$

By the CLT  $\sqrt{T}(\theta^* - \hat{\theta}_T) \xrightarrow{d} Z$  and an argument analogous to the proof of Theorem 2, the third term on the right-hand side of (26) converges to a shift term linear in  $u$ , with random coefficient, which is annihilated by the innovation martingale transformation  $\mathcal{T}_{\hat{\theta}_T}$ . Next, the transformation (through  $\mathcal{T}_{\hat{\theta}_T}$ ) of the first term on the right-hand side of (26) converges weakly on  $D([0, \tau], \mathbb{R}^d)$  to a standard Wiener process  $W_1$ , as  $T \rightarrow \infty$ .

In order to be able to detect the deviation from the null  $\lambda \notin \mathcal{L}_{\Theta} := \{\lambda_{\theta} : \theta \in \Theta\}$ , the second term of (26) should be a non-zero stochastic process after transformation through  $\mathcal{T}_{\hat{\theta}_T}$ . Note that transformation of the second term

$$\mathcal{T}_{\hat{\theta}_T} \left( u \mapsto \frac{1}{\sqrt{T}} \left( \int_0^{uT} \lambda(s) ds - \int_0^{uT} \lambda_{\hat{\theta}_T}(s) ds \right) \right) \quad (27)$$

has, after a change of variables for the second integral,  $k$ th component equal to

$$\frac{1}{\sqrt{T}\sqrt{\mu_{\hat{\theta}_T}^{(k)}}} \left( \int_0^{uT} g_k(s) ds - \int_0^{uT} \frac{1}{T-v} \int_v^T g_k(s) ds dv \right) = \frac{1}{\sqrt{T}\sqrt{\mu_{\hat{\theta}_T}^{(k)}}} \int_0^{uT} \left( g_k(s) - \bar{g}_k^T(s) \right) ds, \quad (28)$$

where  $g_k(s) := \lambda^{(k)}(s) - \lambda_{\hat{\theta}_T}^{(k)}(s)$  and  $\bar{g}_k^T(s) := (T-s)^{-1} \int_s^T g_k(t) dt$ .

Note that by ergodicity of  $N$  (Assumption A(ii)) and by the a.s. continuity of the integral, it holds that  $\sup_{s \in [0, \tau T]} |\bar{g}_k^T(s) - \mathbb{E}[g_k(0)]| \rightarrow 0$ , a.s., as  $T \rightarrow \infty$ . Furthermore, we have  $\bar{g}_k^T(s) = \mathbb{E}[g_k(0)] + \mathcal{O}_{\mathbb{P}}(1/\sqrt{T})$  uniformly on  $[0, \tau T] \ni s$ .

Using [22], Theorem VIII.3.79, with  $p = q = 2$  it follows that

$$\mathcal{T}_{\hat{\theta}_T} \left( u \mapsto \frac{1}{\sqrt{T}} \left( \int_0^{uT} \lambda(s) ds - \int_0^{uT} \lambda_{\hat{\theta}_T}(s) ds \right) \right) \xrightarrow{d} W_2(u) + u\bar{Z},$$

weakly on  $D([0, \tau], \mathbb{R}^d)$  equipped with the strong Skorokhod  $J_1$ -topology, where  $\bar{Z}$  is a  $d$ -variate random vector, and where  $W_2$  is a  $d$ -variate Wiener process with covariance

$$u \mapsto u \cdot \lim_{T \rightarrow \infty} \left( 2 \int_0^{\tau T} \mathbb{E} \left[ \frac{g_i(0) - \bar{g}_i^T(0)}{\mu_{\hat{\theta}_T}^{(i)}} \frac{g_j(t) - \bar{g}_j^T(t)}{\mu_{\hat{\theta}_T}^{(j)}} \right] dt \right)_{i,j \in [d]}.$$

Note that  $\bar{Z} + W_2$  is a non-zero stochastic process if there is some  $k \in [d]$  such that  $\text{Var}(g_k(0)) \neq 0$ , hence if  $\lambda$  is not equal (in  $L^1(\mathbb{P})$ ) to  $\lambda_{\theta} + c$  for some  $\theta \in \Theta$  and some  $c \in \mathbb{R}^d$ .

Now consider the asymptotically correct goodness-of-fit testing procedure outlined in Algorithm 1, selecting  $n(T)$  such that  $T/n(T) \rightarrow \infty$  and  $n(T) \rightarrow \infty$  as  $T \rightarrow \infty$ , where we base the test on the transformed process  $\hat{W}^{(T)} := \mathcal{T}_{\hat{\theta}_T}(\hat{\eta}^{(T)})$  under misspecification  $\lambda \notin \mathcal{L}_{\Theta} := \{\lambda_{\theta} : \theta \in \Theta\}$ . Then in step (iv) of that testing procedure, it holds that the sample  $(\hat{Z}_i^{(T)})_{i \in [n(T)]}$  converges in distribution to a sample of shifted normally distributed random vectors, with a shift of order  $1/\sqrt{n(T)}$ , while the sample size  $n(T)$  diverges, as  $T \rightarrow \infty$ . Consequently, when one applies a goodness-of-fit test that is consistent under alternatives at rate  $\sqrt{n(T)}$  (such as the Kolmogorov-Smirnov, Anderson-Darling and Cramér-von Mises tests) in step (v) of that testing procedure, the power under the alternative  $N$  converges to 1.  $\square$

**Remark 7.** *An interesting follow-up question is whether the conclusion of Theorem 5 remains valid under a root- $T$  alternative. That is, consider a family of alternative models  $\check{N}_T$  defined on  $[0, T]$  with a conditional intensity function of the form  $\check{\lambda}_T = \lambda_{\theta} + \check{\lambda}/\sqrt{T}$ , where  $\theta \in \Theta$  and where  $\check{\lambda}$  is the conditional intensity of some other point process on  $[0, T]$ .*

*Following the proof of Theorem 5, the statistics  $\hat{Z}_i^{(T)}$  again converge to a sample of standard normal variables, but with a perturbation of order  $1/n(T)$  in both their variance and mean. This implies that the shift in mean does not contribute to the power of the test. In contrast, the variance perturbation can yield a test of non-trivial power, unless the dependency structure of the processes  $W_1$  and  $W_2$  appearing in the proof of Theorem 5 enforces their sum to be a standard Wiener process. Determining the precise conditions under which this occurs requires a detailed understanding of their dependence, which appears to be a hard problem. Nonetheless, this suggests the conjecture that for a generic root- $T$  alternative, the induced variance perturbation leads to a test with non-trivial power.*

The proof of Theorem 5 implies that if the test from Section 4.1 is not consistent under an ergodic alternative with intensity  $\lambda$ , then it holds for some  $c \in \mathbb{R}^d$  that  $\lambda(t) - \lambda_{\theta^*}(t) \rightarrow c$ , as  $t \rightarrow \infty$ . Under a stationary alternative, this is only possible if  $\lambda \stackrel{d}{=} \lambda_{\theta^*} + c$  to begin with. By choosing a model ‘closed under adding

constant intensities' (or: a model with a parametric intensity specification including a linear parametric term), i.e., a model such that

$$\{\lambda_\theta + c : c \in \mathbb{R}^d, \lambda_\theta(s) + c \geq 0 \text{ for all } s \geq 0 \text{ a.s.}, \theta \in \Theta\} \subset \mathcal{L}_\Theta,$$

we have consistency under any *stationary* deviation from  $H_0$ . Note that this implies that it is a good idea to add a linear parametric term to the conditional intensity, if one wishes to ensure consistency under  $H_1$ . If our model is not 'closed under adding constant intensities', then the only stationary deviations from the null that our test might fail to detect are point processes having an intensity that is a member of

$$\{\lambda_\theta + c : c \in \mathbb{R}^d \setminus \{0\}, \lambda_\theta(s) + c \geq 0 \text{ for all } s \geq 0 \text{ a.s.}, \theta \in \Theta\}.$$

On the other hand, even for a null model 'closed under adding constant intensities', it is still possible that our test fails to detect an ergodic, but nonstationary deviation  $N$  from the null, but where  $\lambda \rightarrow \lambda_\theta$  a.s., for some  $\theta \in \Theta$ . However, one might argue that any such model is in fact 'close' to the null, since it converges to a null model as it converges to stationarity.

## 6. SIMULATIONS FOR PARAMETRIC HAWKES NULL HYPOTHESES

In this section, we investigate the behavior of the goodness-of-fit tests outlined in Section 4, by considering parametric null hypotheses consisting of linear self-exciting Hawkes processes, whose parameters we estimate from simulated sample paths of point processes. The Hawkes process was introduced in 1971, see [17, 18], and see e.g., [2, 5, 21] for more recent advances. In particular, we consider rejection rates both under simulations of point processes which are a member of the null, and of processes which are not. We compare the asymptotically exact testing procedure outlined in Algorithm 1 to 'naive' testing procedures ignoring estimation uncertainty  $\hat{\theta}_T \neq \theta_0$ , and we argue that our asymptotically exact testing procedure has an advantage over the test based on just the compensated empirical process and over the random-time-change-based test described in Section 4.2.

For the parameter space  $\Theta = (0, 10)^3$  satisfying Assumption A(i), consider the parametric null hypothesis

$$H_0^{\text{Exp}} : N \stackrel{d}{=} N_\theta^{\text{Exp}} \text{ for some } \theta \in \{(\mu, \alpha, \beta) \in \Theta : \alpha < \beta\}, \quad (29)$$

where  $N_\theta^{\text{Exp}} = N_{\mu, \alpha, \beta}^{\text{Exp}}$  is a univariate linear exponential Hawkes process, i.e., a process having intensity

$$\lambda_\theta^{\text{Exp}}(t) = \lambda_{\mu, \alpha, \beta}^{\text{Exp}}(t) = \mu + \sum_{0 \leq t_i < t} \alpha e^{-\beta(t-t_i)}, \quad (30)$$

where  $t_i$  denote the event times of  $N_\theta^{\text{Exp}}$ .

Also, consider the similar parametric null hypothesis

$$H_0^{\text{PL}} : N \stackrel{d}{=} N_\theta^{\text{PL}} \text{ for some } \theta \in \{(\mu, \alpha, \beta) \in \Theta : \alpha < \beta\}, \quad (31)$$

where  $N_\theta^{\text{PL}} = N_{\mu, \alpha, \beta}^{\text{PL}}$  is a univariate linear power-law Hawkes process, i.e., a process having intensity

$$\lambda_\theta^{\text{PL}}(t) = \lambda_{\mu, \alpha, \beta}^{\text{PL}}(t) = \mu + \sum_{0 \leq t_i < t} \alpha (1 + t - t_i)^{-(\beta+1)}, \quad (32)$$

where  $t_i$  denote the event times of  $N_\theta^{\text{PL}}$ .

For both hypotheses, we simulate realizations on  $[0, T]$  for  $T = 5,000$  and  $T = 50,000$ . We set  $n = \text{ceil}(\sqrt{T}/4)$  for both the asymptotically correct transformation-based testing procedure and the 'naive' testing procedure outlined in Section 4.1.



**TABLE 1.** Using the parametric null hypothesis  $H_0^{\text{Exp}}$ ,  $T = 5,000$  and  $n = \text{ceil}(\sqrt{T}/4)$ , we conduct 500 simulations for one model contained in  $H_0$  and five models not contained in  $H_0$ . We report the number of rejections  $R_{0.01}$ ;  $R_{0.05}$ ;  $R_{0.20}$  out of 500 tests, using significance levels 0.01; 0.05; 0.20, using both the transformation-based testing procedure and the ‘naive’ testing procedure outlined in Algorithm 1, using Kolmogorov-Smirnov (KS), Cramér-von Mises (CvM) and Anderson-Darling (AD) normality tests in step (v) thereof.

Test	Transformation-based testing procedure			‘Naive’ testing procedure		
	KS	CvM	AD	KS	CvM	AD
$N^{\text{ExpH}}$	6; 25; 95	4; 25; 96	3; 22; 98	0; 0; 9	0; 0; 6	0; 0; 7
$N^{\text{PLH}}$	14; 40; 106	15; 41; 113	16; 48; 135	0; 0; 10	0; 0; 8	0; 3; 23
$N_{1,2,2}^{\text{SN}}$	2; 17; 95	2; 16; 93	2; 13; 89	0; 0; 5	0; 0; 4	0; 0; 7
$N_{1/5,10,2}^{\text{SN}}$	4; 47; 214	0; 30; 244	0; 28; 251	0; 6; 81	0; 3; 129	0; 4; 154
$N^{\text{Periodic}}$	29; 221; 470	20; 305; 490	10; 324; 496	4; 156; 451	9; 256; 496	5; 281; 497
$N^{\text{SC}}$	500; 500; 500	500; 500; 500	500; 500; 500	500; 500; 500	500; 500; 500	500; 500; 500

**Remark 8.** The choice of the hyperparameter  $n$  is motivated in Section 4.1. It turns out that  $c = 1/4$  yields the right rejection rates for various choices of  $N_{\mu,\alpha,\beta}^{\text{Exp}}$ . However, one should be careful with selecting models with  $\alpha$  close to  $\beta$ , since, for fixed  $T$ , for such parameters  $\hat{W}^{(T)}$  might behave more erratically than a Brownian motion. This is a property inherent to the Hawkes process, not to our testing procedure; see, e.g., [23].

Using the transformation-based and the ‘naive’ testing procedures, we perform goodness-of-fit tests for 500 simulated realizations of the following processes:

- (1)  $N^{\text{ExpH}} := N_{1/2,1,2}^{\text{Exp}}$ .
- (2)  $N^{\text{PLH}} := N_{1/2,1,2}^{\text{PL}}$ .
- (3)  $N_{1,2,2}^{\text{SN}}$ , which is a shot-noise process  $N_{\mu,\alpha,\beta}^{\text{SN}}$  with parameters  $(\mu, \alpha, \beta) = (1, 2, 2)$ . Here,  $N_{\mu,\alpha,\beta}^{\text{SN}}$  has intensity

$$\lambda_{\mu,\alpha,\beta}^{\text{SN}}(t) = \sum_{0 \leq t_i < t} \alpha e^{-\beta(t-t_i)}, \quad (33)$$

where the  $t_i$  are exogenous ‘shots’, arriving according to an independent Poisson process of rate  $\mu$ .

- (4)  $N_{1/5,10,2}^{\text{SN}}$ .
- (5)  $N^{\text{Periodic}}$ , which is a time-inhomogeneous Poisson process  $N_{\mu,\alpha,\beta,\gamma}^{\text{Periodic}}$  with parameters  $(\mu, \alpha, \beta, \gamma) = (5/4, 1, 1/5, 0)$ . Here,  $N_{\mu,\alpha,\beta,\gamma}^{\text{Periodic}}$  has intensity

$$\lambda_{\mu,\alpha,\beta,\gamma}^{\text{Periodic}}(t) = \mu + \alpha \sin(\beta(t - \gamma)), \quad \mu \geq \alpha. \quad (34)$$

- (6)  $N^{\text{SC}}$ , which is a self-correcting point process  $N_{\mu,\alpha,\beta}^{\text{SC}}$  with  $(\mu, \alpha, \beta) = (1, 1/2, \log(2))$ . Here, letting  $N(t_-) = N_{\mu,\alpha,\beta}^{\text{SC}}([0, t))$ , the process  $N_{\mu,\alpha,\beta}^{\text{SC}}$  has intensity

$$\lambda_{\mu,\alpha,\beta}^{\text{SC}}(t) = \mu e^{\beta t} \alpha^{N(t_-)}. \quad (35)$$

For the simulations under  $H_0^{\text{Exp}}$ , we consider the ‘small’ time horizon  $T = 5,000$  in Table 1 and the ‘large’ time horizon  $T = 50,000$  in Table 2. The processes (1)–(6) average around 1 event per time unit, hence those time frames correspond to samples of around 5,000 and 50,000 events.

First, note that for the model  $N^{\text{ExpH}}$  (which belongs to  $H_0^{\text{Exp}}$ ), the rejection rates for the transformation-based testing procedure align well with the nominal significance levels. Specifically, we observe approximately 5, 25, and 100 rejections out of 500 at significance levels of 0.01, 0.05, and 0.20, respectively. This confirms

**TABLE 2.** We report the number of rejections  $R_{0.01}; R_{0.05}; R_{0.20}$  out of 500, using significance levels 0.01; 0.05; 0.20 in an experiment analogous to the one conducted for Table 1, but now with parametric null hypothesis  $H_0^{\text{Exp}}$  and  $T = 50,000$ .


Test	Transformation-based testing procedure			'Naive' testing procedure		
	KS	CvM	AD	KS	CvM	AD
$N^{\text{ExpH}}$	6; 25; 110	5; 20; 107	4; 25; 108	0; 0; 9	0; 0; 5	0; 0; 6
$N^{\text{PLH}}$	9; 31; 101	8; 30; 113	9; 37; 144	0; 0; 11	0; 0; 9	0; 2; 28
$N_{1,2,2}^{\text{SN}}$	6; 20; 91	5; 21; 96	4; 21; 96	0; 0; 6	0; 0; 4	0; 0; 6
$N_{1/5,10,2}^{\text{SN}}$	83; 258; 464	68; 337; 488	143; 418; 499	24; 166; 441	43; 278; 478	104; 396; 494
$N^{\text{Periodic}}$	495; 500; 500	499; 500; 500	500; 500; 500	494; 500; 500	500; 500; 500	500; 500; 500
$N^{\text{SC}}$	500; 500; 500	500; 500; 500	500; 500; 500	500; 500; 500	500; 500; 500	500; 500; 500

the asymptotic correctness of the transformation-based procedure, as stated in Section 4.1. In contrast, the ‘naive’ testing procedure — based on untransformed empirical processes and ignoring the estimation uncertainty ( $\hat{\theta}_T \neq \theta_0$ ) — produces undersized tests. For  $N^{\text{ExpH}}$ , the number of rejections at significance levels of 0.01, 0.05, and 0.20 is far below the expected counts of 5, 25, and 100, respectively. This demonstrates that ignoring estimation uncertainty leads to unreliable tests. Finally, the power of the transformation-based procedure against alternative hypotheses appears to be higher than that of the ‘naive’ procedure. However, we emphasize that this need not hold in general.

Suppose that in step (v) of the testing procedure described in Algorithm 1, we apply the Kolmogorov-Smirnov normality test. We can then compare the simulated Kolmogorov-Smirnov test statistics for  $N^{\text{ExpH}} \in H_0^{\text{Exp}}$  with their theoretical null distribution given by the Kolmogorov distribution, using a Q-Q plot. Under the null, the Q-Q plot should be centered around the line  $y = x$ . We investigate this for  $H_0^{\text{Exp}}$  at both  $T = 5,000$  and  $T = 50,000$ , comparing the transformation-based and ‘naive’ testing procedures (see Figure 1). As expected from Sections 3 and 4.1, the Q-Q plots for the transformation-based procedure align closely with the line  $y = x$  (see Figures 1(a) and (c)). Moreover, the deviation from  $y = x$  is smaller at  $T = 50,000$  compared to  $T = 5,000$ , which aligns with the asymptotic nature of our test as  $T \rightarrow \infty$ . By contrast, the ‘naive’ testing procedure yields different results. Since the standardized empirical process in (23) corresponds to a Brownian motion with a linear drift and a random coefficient, the random variables  $\tilde{Z}_i^{(T)}$  defined in (24) are normally distributed, but not standard normal. This is reflected in the Q-Q plots for the ‘naive’ procedure (see Figures 1(b) and (d)), where the points form an approximately straight line but deviate from  $y = x$ .

Now, consider the alternative hypotheses. Detecting deviations from the null  $H_0^{\text{Exp}}$  varies in difficulty. As it turns out, the processes  $N^{\text{PLH}}$  and  $N_{1,2,2}^{\text{SN}}$  are challenging to distinguish from the null, while detecting  $N_{1/5,10,2}^{\text{SN}}$  is more feasible, and identifying  $N^{\text{Periodic}}$  is relatively easy. For  $N^{\text{SC}}$ , the deviation from the null is straightforward to detect.

These differences can be explained as follows. First,  $N^{\text{PLH}}$  is a Hawkes process similar to those in  $H_0^{\text{Exp}}$ , but with a power-law decay for the excitation kernel instead of exponential decay, leading to long memory. Detecting such differences in kernel shape is inherently difficult. Second, while the shot-noise process  $N_{1,2,2}^{\text{SN}}$  exhibits clustering behavior like the Hawkes process, it lacks self-excitation. Since its shot intensity  $\mu = 1$  is high compared to  $\beta = 2$ , clusters frequently overlap, making it hard to distinguish between exogenous and endogenous clustering. In contrast,  $N_{1/5,10,2}^{\text{SN}}$  has a lower shot intensity than  $N_{1,2,2}^{\text{SN}}$ , resulting in more clearly separated clusters. Since cluster initiation in the shot-noise process is exogenous, we expect the null to be more easily rejected for  $N_{1/5,10,2}^{\text{SN}}$  than for  $N_{1,2,2}^{\text{SN}}$ . Now, consider the easier alternatives. The periodic Poisson



**FIGURE 1.** We consider 500 simulations of  $N^{\text{ExpH}}$  both for  $T = 5,000$  and  $T = 50,000$ , and both for the transformation-based and the 'naive' testing procedures outlined in Section 4.1. We test for standard normal increments using the Kolmogorov-Smirnov test. We draw Q-Q plots for the empirical distribution of the test statistics vs. the theoretical limit distribution, i.e., the *Kolmogorov distribution*. The dashed red line is a straight line fitted to the scatterplot, while the line  $y = x$  is drawn solid blue.

process  $N^{\text{Periodic}}$  produces peaks in intensity, leading to concentrated event groups that resemble clusters. For small  $T$ , detecting this may be challenging due to the limited number of observed 'clusters'. However, the regularity of these clusters should reveal that the process is not an overdispersed one. Finally, since the self-correcting process  $N^{\text{SC}}$  models behavior opposite to that of a Hawkes process, rejecting the null under this alternative should be quite easy.

Although Theorem 5 ensures that the power of the Kolmogorov-Smirnov, Anderson-Darling, and Cramér-von Mises tests converges to 1 as  $T \rightarrow \infty$  under any alternative, the previous paragraph suggests that, in practice, detecting the deviations  $N^{\text{PLH}}$  and  $N_{1,2,2}^{\text{SN}}$  requires a sample  $\omega_T$  over a *large* time frame  $[0, T]$ . To investigate this, we run 100 computationally intensive simulations for  $H_0^{\text{Exp}}$  versus  $N^{\text{PLH}}$  and  $N_{1,2,2}^{\text{SN}}$  for large values of  $T$ . The rejection rates for  $N^{\text{PLH}}$  are well above the significance levels at  $T = 5 \cdot 10^5$ , while for  $N_{1,2,2}^{\text{SN}}$ , we need  $T = 5 \cdot 10^6$  to observe similar behavior. These rejection rates are summarized in Table 3.

**TABLE 3.** We report the number of rejections  $R_{0.01}$ ;  $R_{0.05}$ ;  $R_{0.20}$  out of 100, using significance levels 0.01; 0.05; 0.20 in an experiment analogous to the one conducted for Table 1, using the parametric null hypothesis  $H_0^{\text{Exp}}$  and the alternative models  $N^{\text{PLH}}$  and  $N_{1,2,2}^{\text{SN}}$  for large time frames with  $T = 5 \cdot 10^5$  and  $T = 5 \cdot 10^6$ , respectively.

Test	Transformation-based testing procedure			'Naive' testing procedure		
	KS	CvM	AD	KS	CvM	AD
$N^{\text{PLH}}, T = 5 \cdot 10^5$	3; 10; 35	2; 16; 40	3; 20; 41	0; 3; 11	0; 1; 13	0; 5; 19
$N_{1,2,2}^{\text{SN}}, T = 5 \cdot 10^6$	10; 17; 40	10; 18; 41	15; 23; 42	5; 10; 29	4; 5; 21	4; 9; 22

**TABLE 4.** We report the number of rejections  $R_{0.01}$ ;  $R_{0.05}$ ;  $R_{0.20}$  out of 500, using significance levels 0.01; 0.05; 0.20 in an experiment analogous to the one conducted for Table 1, but now with parametric null hypothesis  $H_0^{\text{PL}}$  and  $T = 5,000$ .

Test	Transformation-based testing procedure			'Naive' testing procedure		
	KS	CvM	AD	KS	CvM	AD
$N^{\text{PLH}}$	3; 23; 100	4; 20; 97	6; 20; 100	0; 0; 9	0; 0; 3	0; 0; 10
$N^{\text{ExpH}}$	7; 22; 92	6; 18; 90	4; 14; 86	0; 0; 4	0; 0; 2	0; 0; 3
$N_{1,2,2}^{\text{SN}}$	9; 26; 96	8; 29; 93	5; 25; 92	0; 1; 3	0; 0; 5	0; 0; 6
$N_{1/5,10,2}^{\text{SN}}$	8; 47; 243	4; 43; 298	1; 41; 304	0; 5; 140	0; 11; 204	0; 14; 245
$N^{\text{Periodic}}$	15; 131; 417	10; 169; 460	3; 173; 475	0; 46; 350	1; 98; 431	0; 111; 451
$N^{\text{SC}}$	500; 500; 500	500; 500; 500	500; 500; 500	500; 500; 500	500; 500; 500	500; 500; 500

**TABLE 5.** We report the number of rejections  $R_{0.01}$ ;  $R_{0.05}$ ;  $R_{0.20}$  out of 500, using significance levels 0.01; 0.05; 0.20 in an experiment analogous to the one conducted for Table 1, but now with parametric null hypothesis  $H_0^{\text{PL}}$  and  $T = 50,000$ .

Test	Transformation-based testing procedure			'Naive' testing procedure		
	KS	CvM	AD	KS	CvM	AD
$N^{\text{PLH}}$	4; 24; 99	2; 19; 101	2; 18; 100	0; 0; 4	0; 0; 3	0; 0; 6
$N^{\text{ExpH}}$	3; 28; 115	4; 24; 110	2; 21; 110	0; 0; 10	0; 0; 9	0; 0; 9
$N_{1,2,2}^{\text{SN}}$	3; 22; 96	3; 22; 85	2; 20; 87	0; 0; 16	0; 0; 7	0; 0; 15
$N_{1/5,10,2}^{\text{SN}}$	104; 334; 489	123; 412; 498	244; 469; 500	43; 269; 481	87; 378; 496	204; 468; 498
$N^{\text{Periodic}}$	431; 498; 500	487; 500; 500	499; 500; 500	409; 499; 500	493; 500; 500	499; 500; 500
$N^{\text{SC}}$	500; 500; 500	500; 500; 500	500; 500; 500	500; 500; 500	500; 500; 500	500; 500; 500

We repeat our experiments for  $H_0^{\text{PL}} \ni N^{\text{PLH}}$  consisting of univariate power-law Hawkes processes. We report our findings for  $T = 5,000$  and  $T = 50,000$  in Table 4 and Table 5, respectively. The observed rejection rates are comparable to the ones in Tables 1 and 2 for  $H_0^{\text{Exp}}$ . This confirms the results of Sections 3–5 once more. Furthermore, it is possible to draw Q-Q plots analogous to Figure 1, yielding comparable results.

**Remark 9.** *Our results hold as well for null hypotheses different than the self-exciting ones considered here. For example, for a null hypothesis consisting of periodic Poisson processes  $N_{\mu,\alpha,\beta,\gamma}^{\text{Periodic}}$ , we run 500 simulations under  $N^{\text{Periodic}}$ , with  $T = 5,000$  and using the Anderson-Darling normality test. Using significance levels 0.01, 0.05 and 0.20, this yields 10, 32 and 104 rejections when using the transformation-based testing procedure, and 0, 0 and 5 rejections when using the 'naive' testing procedure, respectively.*

**TABLE 6.** We report the number of rejections  $R_{0.01}$ ;  $R_{0.05}$ ;  $R_{0.20}$  out of 500, using significance levels 0.01; 0.05; 0.20 in an experiment similar to the one conducted for Table 1, but now using the testing procedure described in Section 4.2, with  $T = 5,000$ .

Test	$H_0^{\text{Exp}}$			$H_0^{\text{PL}}$		
	KS	CvM	AD	KS	CvM	AD
$N^{\text{ExpH}}$	0; 0; 3	0; 0; 0	0; 0; 1	0; 5; 91	0; 1; 78	0; 9; 121
$N^{\text{PLH}}$	0; 4; 49	0; 0; 36	0; 3; 59	0; 0; 1	0; 0; 1	0; 0; 2
$N_{1,2,2}^{\text{SN}}$	0; 5; 71	0; 0; 31	0; 0; 23	3; 52; 255	0; 8; 195	0; 8; 164
$N_{1/5,10,2}^{\text{SN}}$	3; 66; 239	0; 13; 212	5; 51; 271	0; 0; 7	0; 0; 2	0; 1; 9
$N^{\text{Periodic}}$	44; 165; 349	26; 163; 391	43; 193; 406	500; 500; 500	500; 500; 500	500; 500; 500
$N^{\text{SC}}$	500; 500; 500	500; 500; 500	500; 500; 500	500; 500; 500	500; 500; 500	500; 500; 500

Note that the Kolmogorov-Smirnov, Anderson-Darling and Cramér-von Mises tests all yield goodness-of-fit tests of the right size when applied in step (v) of the transformation-based testing procedure outlined in Algorithm 1. Since in our simulations the power under alternatives is highest for Anderson-Darling and lowest for Kolmogorov-Smirnov, we advice to use the former in applications. We will do so in the next section.

We conclude by examining the random-time-change-based test discussed in Section 4.2, which, like the ‘naive’ testing procedure in this section, ignores estimation uncertainty. We fit the same six processes as before to both  $H_0^{\text{Exp}}$  and  $H_0^{\text{PL}}$ , using time frame  $[0, T]$  for  $T = 5,000$ , and then perform the goodness-of-fit test outlined in Section 4.2. The results are summarized in Table 6. We observe that the random-time-change-based test, similar to the ‘naive’ procedure, appears undersized. Additionally, the power under any alternative seems to be lower than that of the asymptotically exact test presented in Section 4.1. Thus, the random-time-change-based test shares the shortcomings of the ‘naive’ testing procedure.

## 7. DATA ANALYSIS

In this section, we illustrate the testing procedure outlined in Algorithm 1 by applications to real-world data. In particular, we compare the goodness-of-fit tests conducted in [39] and [46] to our asymptotically exact test.

**7.1. Temporal ETAS model.** First, we apply the testing procedure from Algorithm 1 to earthquake data used in [39], the seminal paper that established a framework for applying Omori’s law and the Gutenberg-Richter law to earthquake data. Building on this temporal model, subsequent years saw the development of the more advanced spatiotemporal ETAS (Epidemic Type Aftershock Sequence) model, which has become a cornerstone in earthquake modeling. For further developments and applications, see, e.g., [1, 40, 44, 52].

We consider the earthquake data set used in [39], Table 1, consisting of shallow earthquakes of less than 100 kilometers depth, of magnitude at least 6.0 on the Richter scale, in the Off Tohoku district east of Japan. This is the polygonal region with vertices at (42°N, 142°E), (39°N, 142°E), (38°N, 141°E), (35°N, 140.5°E), (35°N, 144°E), (39°N, 146°E), and (42°N, 146°E); see [39], Fig. 1 for a map. This region is one of the most seismically active areas in Japan. The data set spans the period between 1885 and 1980, and consists of 483 earthquakes satisfying the depth and magnitude conditions.

The earthquake dynamics are modeled by marked point processes, with time stamps  $t_i$  corresponding to earthquake occurrences and marks  $m_i$  corresponding to earthquake magnitudes, assumed to be independent

of  $t_i$ . More specifically, [39] considers marked Hawkes processes with conditional intensity functions

$$\lambda(t) = \mu + \sum_{t_i < t} c(m_i)g(t - t_i). \quad (36)$$

Based on the Akaike information criterion, the model with  $g(t) = K/(t+c)$  and  $c(m) = e^{\beta(m-M)}$  is selected. Here  $M = 6.0$  is the cut-off magnitude. The decay kernel  $g(t)$  and the mark function  $c(m)$  are theoretically motivated by Omori's law and the Gutenberg-Richter law, respectively. This gives the conditional intensity function

$$\lambda_\theta(t) = \lambda_{\mu,K,c,\beta}(t) = \mu + \sum_{t_i < t} e^{\beta(m_i-M)} \frac{K}{t - t_i + c}. \quad (37)$$

From [39], Table 3, the maximum likelihood estimators for the parameters are


$$\hat{\theta} = (\hat{\mu}, \hat{K}, \hat{c}, \hat{\beta}) = (.00536, .017284, .01959, 1.61385).$$

Having selected this 'best' model, it is noted that it is still possible that there exists a more suitable model. To assess the suitability of the model (37), [39] considers the transformed event times  $(\tau_i)_{i \in [483]}$ , where  $\tau_i = \Lambda_{\hat{\theta}}(t_i)$ , and where  $\Lambda_\theta(t) = \int_0^t \lambda_\theta(s) ds$  is the *compensator* of the point process. By the random time change theorem, [14], Theorem 7.4.I, it follows that the transformed times  $(\Lambda_{\theta_0}(t_i))_{i \in [483]}$  using the *true* parameter  $\theta_0$  form a realization of a stationary Poisson process of unit intensity. Therefore, it is noted in [39] that if  $\lambda_{\hat{\theta}}$  is a good approximation of  $\lambda_{\theta_0}$ , then  $(\tau_i)_{i \in [483]}$  is expected to behave like a stationary Poisson process of unit intensity. However, from Section 4.2 we know that even if  $\lambda_{\hat{\theta}}$  is a good approximation of  $\lambda_{\theta_0}$ , the difference between  $(\Lambda_{\hat{\theta}}(t_i))_{i \in [483]}$  and  $(\Lambda_{\theta_0}(t_i))_{i \in [483]}$  is not necessarily neglectable: ignoring estimation uncertainty might lead to inaccurate tests.

In [39], the null hypothesis that  $(\tau_i)_{i \in [483]}$  is a realization of a Poisson process of unit intensity is tested with the aid of a Kolmogorov-Smirnov test; see [39], Figure 9. One could use the transformed times  $(\tau_i)_{i \in [483]}$  to construct the interarrival times  $(x_i)_{i \in [482]}$ , where  $x_i := \tau_{i+1} - \tau_i$  are standard exponentially distributed under the null. Following [39], we perform a Kolmogorov-Smirnov test for the null hypothesis stating that  $(x_i)_{i \in [482]}$  is a sample from a standard exponential distribution. This yields a  $p$ -value of .5756. Hence, for any reasonable significance level, the null is not rejected, indicating a seemingly good fit of the model to the data.

We examine the situation in greater detail, accounting for the estimation uncertainty in  $\hat{\theta}$  by performing the asymptotically exact transformation-based test introduced in Algorithm 1. Specifically, the transformed empirical process is calculated using (15) as outlined in step (iii) of the procedure, and its path is shown in Figure 2. Under the null hypothesis that the true intensity satisfies (37), this figure should resemble a standard Brownian motion. Performing the transformation-based testing procedure involves selecting a hyperparameter  $n$  and a normality test for step (v) of the procedure. Unlike the setup in Section 4.1, where there is approximately one event per time unit (days), here the choice  $n = \text{ceil}(\sqrt{N[0, T]}/4) = 6$  instead of  $n = \text{ceil}(\sqrt{T}/4)$  is more appropriate. Using the Anderson-Darling normality test suggested at the end of Section 6, we obtain a  $p$ -value of 0.0568 (and 0.2475 for the 'naive' testing procedure).

The relatively low  $p$ -value and the unexpected jumps in Figure 2 raise concerns about the adequacy of the model's fit. While the  $p$ -value does not lead to rejection at a typical significance level such as  $\alpha = 0.05$ , it suggests that the model may not fully capture the underlying dynamics of the data. This is consistent with the expectation that the temporal ETAS model might overlook certain features better accounted for in the spatiotemporal version. As we have seen, a test based on random time change, ignoring estimation uncertainty, yields a much higher  $p$ -value, indicating a seemingly good fit. These discrepancies highlight



**FIGURE 2.** Path of the transformed empirical process under the null hypothesis that the intensity of earthquake occurrences satisfies (37). The plot exhibits deviations from the expected behavior of a standard Brownian motion, including unexpected jumps, suggesting potential inadequacies in the temporal ETAS model's fit to the data.

that ignoring estimation uncertainty could lead to inaccurate procedures with unwarranted confidence in an otherwise imperfect model.


**7.2. A recursive point process model.** More recently, in 2019, a variant of the Hawkes process was introduced in [46], having the property that the expected offspring size corresponding to an event is inversely related to the conditional intensity at the time that event occurred. This 'recursive point process model' is based on the idea that when a disease occurs infrequently within a population, as seen in the initial phases of an outbreak, individuals with the disease are likely to transmit it at a higher rate. In contrast, as the disease becomes more widespread, the transmission rate diminishes due to the implementation of preventive measures and the growing proportion of individuals with prior exposure. The model is described as recursive because the conditional intensity at any given moment depends on the productivity of earlier points, which itself is influenced by the conditional intensity at those same points.

This recursive model is specified through its conditional intensity

$$\lambda(t) = \mu + \sum_{t_i < t} H(\lambda(t_i))g(t - t_i), \quad (38)$$

with  $\mu > 0$ , where  $H : (0, \infty) \rightarrow [0, \infty)$  is assumed to be a non-increasing function and where  $g : [0, \infty) \rightarrow [0, \infty)$  is a density function. Note that this model is self-exciting, and can be seen as a variant of the classical Hawkes process. A parametric specification is chosen where  $H$  satisfies a power law and where  $g$  is an exponential density. More specifically,  $H(x) = \kappa/x^\alpha$  and  $g(u) = \beta \exp(-\beta u)$ , yielding a parametric model depending on  $\theta = (\mu, \kappa, \beta, \alpha)$ , where all parameters are assumed to be positive.

In [46], Section 7, this model is fitted to data consisting of recorded cases of Rocky Mountain Spotted Fever (RMSF) in California from 1960 to 2011. The freely accessible data were obtained from Project Tycho ([www.tycho.pitt.edu](http://www.tycho.pitt.edu)). Today, an updated version of this data set is available on Project Tycho, which includes the original data set of 67 disease counts, as well as 2569 additional disease counts. As in [46],



**FIGURE 3.** For both the small (a) and the large (b) data sets we fit the model for the data, after which we transform arrival times and we test for standard exponentiality of the interarrival times. Deviations from  $y = x$  are evident, especially for the larger data set.

Section 7, since the data are recorded with a weekly temporal resolution, the onset time for each individual case was randomly assigned from a uniform distribution within the corresponding 7-day interval. In this section, we replicate the model fitting from [46], Section 7, yielding the following parameter estimates for the small data set:

$$(\hat{\mu}_S, \hat{\kappa}_S, \hat{\beta}_S, \hat{\alpha}_S) = (0.000233, 0.0409, 0.00315, 0.600), \quad (39)$$

and for the large data set, consisting of  $67 + 2569 = 2636$  disease counts:

$$(\hat{\mu}_L, \hat{\kappa}_L, \hat{\beta}_L, \hat{\alpha}_L) = (0.00216, 0.793, 0.0606, 0.192). \quad (40)$$

In [46], Section 7, after fitting the (small) data set to the model, the fit is assessed using super-thinned residuals. This method is motivated by principles similar to those behind the random-time-change-based test described in Section 4.1. Specifically, both methods ignore estimation uncertainty, which can lead to inaccurate results. In this section, instead of using super-thinned residuals, we apply a classical random-time-change-based goodness-of-fit test as outlined in Section 4.2. After fitting the model, we transform the interarrival times  $t_{i+1} - t_i$ , using the estimated compensator, into  $\Lambda_{\hat{\theta}}(t_{i+1}) - \Lambda_{\hat{\theta}}(t_i)$ , and test whether the transformed interarrival times follow a standard exponential distribution. The corresponding QQ-plots, along with the reference line  $y = x$  (around which the interarrival times should be centered under the null hypothesis), are shown in Figure 3. For the small data set, a Kolmogorov-Smirnov test yields a  $p$ -value of 0.1632, consistent with [46], Figure 3, where a slight deviation from the straight line  $y = x$  is observed in Figure 3(A), but within the Kolmogorov-Smirnov confidence bounds, so we do not reject the null at the 0.05 significance level. For the large data set, the deviation from the straight line is more pronounced, leading to a rejection of the null; the  $p$ -value is 0.000.

Next, we account for the estimation uncertainty in  $\hat{\theta}$  by performing the asymptotically exact transformation-based test described in Algorithm 1. For the small sample, we select the hyperparameter  $n = 8$ , as  $n = \text{ceil}(\sqrt{N[0, T]}/4) = 2$  normals is very few. For the large sample, we use  $n = \text{ceil}(\sqrt{N[0, T]}/4) = 13$ . Following the procedure, we apply an Anderson-Darling test for normality in step (v), as suggested in Section 6. For the small sample, the  $p$ -value is 0.2506 (compared to 0.5948 for the ‘naive’ testing procedure). For the large sample, the  $p$ -value is 0.0382 (compared to 0.3790 for the ‘naive’ procedure). The results of these tests are consistent with the conclusions of the random-time-change-based tests above, at  $\alpha = .05$ . While



the fit in Figure 3(A) may seem reasonably good, concerns arise for large data sets where rejection might be driven by estimation uncertainty. The less-than-perfect fit indicated by the random-time-change-based test for the large sample is confirmed by rejection in our asymptotically exact test, which accounts for estimation uncertainty.

Possible objections are that the rejection based on the full data set may result from a trend in the data (e.g., due to global warming affecting tick activity) or that the large sample size enables the detection of very small deviations from the model. To investigate this, we select a subset of the data consisting of 203 infections recorded between 1970 and 1977. This yields the following parameter estimates:

$$(\hat{\mu}_{70-77}, \hat{\kappa}_{70-77}, \hat{\beta}_{70-77}, \hat{\alpha}_{70-77}) = (0.00191, 0.753, 0.0537, 0.196). \quad (41)$$

A goodness-of-fit test based on the random-time-change method (see Section 4.2) results in a  $p$ -value of 0.0725, meaning that we would not reject the null hypothesis at a typical significance level of  $\alpha = 0.05$ . Next, using  $n = \text{ceil}(\sqrt{N[0, T]}/4) = 4$ , an Anderson-Darling test for normality in step (v) of Algorithm 1 yields a  $p$ -value of 0.0069 (compared to 0.4257 for the ‘naive’ testing procedure). Hence, here one would reject the null at a typical significance level of  $\alpha = 0.05$ . This demonstrates, using real data, that our testing procedure may have higher power than approaches that ignore estimation uncertainty, a phenomenon we already observed in Section 6.

It is important to note that the recursive model defined in (38) is self-exciting, making it a plausible choice for modeling diseases that spread directly between members of a population. However, RMSF is transmitted to humans via ticks and is not contagious between humans. Although tick bites might exhibit temporal clustering, such as spikes during favorable conditions like good hiking weather, we do not expect one tick bite to trigger another. Instead, tick bites likely follow a shot-noise type of intensity, with external factors driving exposure likelihood. As such, it is not surprising that the recursive model does not perfectly fit RMSF infection data. Given the challenges outlined in Section 6 regarding separating shot-noise from self-exciting dynamics, it is not surprising that neither the random-time-change-based test nor our asymptotically exact test rejects the null hypothesis when applied to the small sample of just 67 observations.

This recursive model, however, may be particularly well-suited for diseases like COVID-19, influenza, and Ebola, which are transmissible between humans. In such scenarios, a higher number of infections (hence, a higher conditional intensity) typically triggers stricter preventive measures by policymakers or individuals, thereby decreasing the expected number of secondary infections (or the so-called ‘ $R$ -number’). This dynamic is accounted for by the recursive model but is not captured by the traditional self-exciting model.

## 8. CONCLUDING REMARKS

In this work, we introduced an asymptotically distribution-free test process for point processes, which can be used to construct asymptotically distribution-free goodness-of-fit tests. Such tests are straightforward to implement and address the limitations of tests that ignore estimation uncertainty, which are often undersized and lack power.

Several extensions and follow-up questions arise.

- Although our primary focus has been on  $d$ -dimensional point processes for  $d \in \mathbb{N}$ , it is worth noting that our results extend naturally to spatiotemporal point processes defined on an infinite spatial domain  $\mathcal{X}$ , without introducing significant challenges. In this case, one would work with a conditional intensity function  $\lambda_\theta(s, t)$  and analyze space-integrated quantities, assuming the regularity conditions required for the central limit theorem  $\sqrt{T}(\theta_0 - \hat{\theta}_T) \xrightarrow{d} Z$ , see [43]. Additionally,

if  $\alpha_\theta := \lim_{T \rightarrow \infty} \int_0^T \int_{\mathcal{X}} \frac{\partial \lambda_\theta(s,t)}{\partial \theta} ds dt$  exists almost surely for  $\theta = \theta_0$ , one can derive a version of Corollary 1. This results in a Wiener process with variance  $\mathbb{E}[N_{\theta_0}([0, 1], \mathcal{X})]$  and drift  $\alpha_{\theta_0} Z$ , which is in the suitable form for applying an innovation martingale transformation. Therefore, we can proceed as in Section 3.3 and beyond. This framework allows for the construction of an asymptotically distribution-free goodness-of-fit test for spatiotemporal point processes.

- As noted in Section 4.2, if a consistent estimate of  $\alpha_{\theta_0} Z$  could be obtained from the data, it would be possible to modify (25) to develop an asymptotically distribution-free random-time-change-based goodness-of-fit test. However, solving this problem is far from straightforward due to the dependence of  $\alpha_{\theta_0} Z$  on the unobservable quantity  $\Lambda_{\theta_0}(t_i) - \Lambda_{\theta_0}(t_{i-1})$  in (25). Despite this challenge, this remains an intriguing direction for future research.

#### REFERENCES

- [1] G. ADELIO and M. CHIODI (2015). FLP estimation of semi-parametric models for space–time point processes and diagnostic tools. *Spatial Statistics* **14**, pp. 119–132.
- [2] Y. AİT-SAHALIA, J. A. CACHO-DIAZ, and R. J. A. LAEVEN (2015). Modeling financial contagion using mutually exciting jump processes. *Journal of Financial Economics* **117**, pp. 585–606.
- [3] H. ALBRECHER, M. BLADT, D. KORTSCHAK, F. PRETTENTHALER, and T. SWIERCZYNSKI (2019). Flood occurrence change-point analysis in the paleoflood record from Lake Mondsee (NE Alps). *Global and Planetary Change* **178**, pp. 65–76.
- [4] P. K. ANDERSEN, Ø. BORGAN, R. D. GILL, and N. KEIDING (1993). *Statistical Models Based on Counting Processes*. Springer Series in Statistics, Springer-Verlag, New York.
- [5] E. BACRY, S. DELATTRE, M. HOFFMANN, and J. F. MUZY (2013). Some limit theorems for Hawkes processes and application to financial statistics. *Stochastic Processes and their Applications* **123**, pp. 2475–2499.
- [6] P. BILLINGSLEY (1999). *Convergence of Probability Measures*. John Wiley & Sons.
- [7] P. BRÉMAUD (1981). *Point Processes and Queues: Martingale Dynamics*. Springer-Verlag, New York.
- [8] S. U. CAN, J. H. J. EINMAHL, E. V. KHMALADZE and R. J. A. LAEVEN (2015). Asymptotically distribution-free goodness-of-fit testing for tail copulas. *The Annals of Statistics* **43**, pp. 878–902.
- [9] S. U. CAN, J. H. J. EINMAHL and R. J. A. LAEVEN (2020). Goodness-of-fit testing for copulas: A distribution-free approach. *Bernoulli* **26**, pp. 3163–3190.
- [10] V. CHAVEZ-DEMOULIN and J. A. MCGILL (2012). High-frequency financial data modeling using Hawkes processes. *Journal of Banking and Finance* **36**, pp. 3415–3426.
- [11] F. CHEN and W. H. TAN (2018). Marked self-exciting point process modelling of information diffusion on Twitter. *The Annals of Applied Statistics* **12**, pp. 2175–2196.
- [12] R. A. CLEMENTS, F. P. SCHOENBERG, and A. VEEN (2013). Evaluation of space–time point process models using super-thinning. *Environmetrics* **23**(7), pp. 606–616.
- [13] D. VERE-JONES and F. P. SCHOENBERG (2004). Rescaling marked point processes. *Australian and New Zealand Journal of Statistics* **46**(1), pp. 133–143.
- [14] D. J. DALEY and D. VERE-JONES (2003). *An Introduction to the Theory of Point Processes*, Vol I and II, 2nd ed. Springer-Verlag, New York.
- [15] M. A. DELGADO, J. HIDALGO, and C. VELASCO (2005). Distribution-free goodness-of-fit tests for linear processes. *Annals of Statistics* **33**, pp. 2568–2609.
- [16] H. DETTE and B. HETZLER (2009). Khmaladze transformation of integrated variance processes with applications to goodness-of-fit testing. *Mathematical Methods of Statistics* **18**, pp. 97–116.
- [17] A. G. HAWKES (1971). Spectra of some self-exciting and mutually exciting point processes. *Biometrika* **58**, pp. 83–90.
- [18] A. G. HAWKES and D. OAKES (1974). A cluster process representation of a self-exciting process. *Journal of Applied Probability* **11**, pp. 493–503.
- [19] P. EMBRECHTS, T. LINIGER and L. LIN (2011). Multivariate Hawkes processes: an application to financial data. *Journal of Applied Probability* **48**, pp. 367–378.
- [20] N. L. HJORT (1986). Bayes estimators and asymptotic efficiency in parametric counting process models. *Scandinavian Journal of Statistics* **13**, pp. 63–85.
- [21] M. IKEFUJI, R. J. A. LAEVEN, J. R. MAGNUS and Y. YUE (2022). Earthquake risk embedded in property prices: Evidence from five Japanese cities. *Journal of the American Statistical Association* **117**, pp. 82–93.

- [22] J. JACOD and A. SHIRYAEV (2003). *Limit Theorems for Stochastic Processes*. Springer-Verlag.
- [23] T. JAISSON and M. ROSENBAUM (2015). Limit theorems for nearly unstable Hawkes processes. *The Annals of Applied Probability* **25** pp. 600–631.
- [24] E. V. KHMALADZE (1981). A martingale approach in the theory of goodness-of-fit tests. *Teoriya Veroyatnostei i ee Primeneniya* **26**, pp. 246–265.
- [25] E. V. KHMALADZE (1988). An innovation approach to goodness-of-fit tests in  $\mathbb{R}^m$ . *Annals of Statistics* **16**, pp. 1503–1516.
- [26] E. V. KHMALADZE (1993). Goodness of fit problem and scanning innovation martingales. *Annals of Statistics* **21**, pp. 798–829.
- [27] E. V. KHMALADZE and H. L. KOUL (2004). Martingale transforms goodness-of-fit tests in regression models. *Annals of Statistics* **32**, pp. 995–1034.
- [28] E. V. KHMALADZE and H. L. KOUL (2009). Goodness-of-fit problem for errors in nonparametric regression: Distribution-free approach. *Annals of Statistics* **37**, pp. 3165–3185.
- [29] R. KOENKER and Z. XIAO (2002). Inference on the quantile regression process. *Econometrica* **70**, pp. 1583–1612.
- [30] R. KOENKER and Z. XIAO (2006). Quantile autoregression. *Journal of the American Statistical Association* **101**, pp. 980–990.
- [31] T. J. KWAN, F. CHEN and W. T. M. DUNSMUIR (2023). Alternative asymptotic inference theory for a nonstationary Hawkes process *Journal of Statistical Planning and Inference* **227**, pp. 75–90.
- [32] M. LALLOUACHE and D. CHALLET (2016). The limits of statistical significance of Hawkes processes fitted to financial data. *Quantitative Finance* **16**, pp. 1–11.
- [33] M. LIFSHITS (2012). *Lectures on Gaussian Processes*, SpringerBriefs in Mathematics, Heidelberg.
- [34] R. LUO, V. KRISHNAMURTHY and E. BLASCH (2022). Hawkes process modeling of block arrivals in Bitcoin Blockchain. Preprint. Available at <https://arxiv.org/abs/2203.16666>.
- [35] I. W. MCKEAGUE, A. M. NIKABADZE, and Y. Q. SUN (1995). An omnibus test for independence of a survival time from a covariate. *Annals of Statistics* **23**, pp. 450–475.
- [36] A. NIKABADZE and W. STUTE (1997). Model checks under random censorship. *Statistics and Probability Letters* **32**, pp. 249–259.
- [37] Y. OGATA (1978). The asymptotic behaviour of maximum likelihood estimators for stationary point processes. *Annals of the Institute of Statistical Mathematics* **30**, pp. 243–261.
- [38] Y. OGATA (1981). On Lewis’ simulation method for point processes. *IEEE Transactions on Information Theory* **IT-27**, pp. 23–31.
- [39] Y. OGATA (1988). Statistical models for earthquake occurrences and residual analysis for point processes. *Journal of the American Statistical Association* **83**, pp. 9–27.
- [40] Y. OGATA (1998). Space-time point-process models for earthquake occurrences. *Annals of the Institute of Statistical Mathematics* **50**, pp. 379–402.
- [41] T. OZAKI (1979). Maximum likelihood estimation of Hawkes’ self-exciting point processes. *Annals of the Institute of Statistical Mathematics* **31**, pp. 145–155.
- [42] M. L. PURI and P. D. TUAN (1986). Maximum likelihood estimation for stationary point processes. *Proceedings of the National Academy of Sciences* **83**, pp. 541–545.
- [43] S. L. RATHBUN (1996). Asymptotic properties of the maximum likelihood estimator for spatio-temporal point processes. *Journal of Statistical Planning and Inference* **51**, pp. 55–74.
- [44] A. REINHART (2018). A review of self-exciting spatio-temporal point processes and their applications. *Statistical Science* **33**, pp. 299–318.
- [45] F. P. SCHOENBERG (1999). Transforming spatial point processes into Poisson processes. *Stochastic Processes and Their Applications* **81**, pp. 155–164.
- [46] F. P. SCHOENBERG, M. HOFFMANN, and R. J. HARRIGAN (2019). A recursive point process model for infectious diseases. *Annals of the Institute of Statistical Mathematics* **71**, pp. 1271–1287.
- [47] W. STUTE, S. THIES, and L.-X. ZHU (1998). Model checks for regression: An innovation process approach. *Annals of Statistics* **26**, pp. 1916–1934.
- [48] Y. SUN, R. C. TIWARI, and J. N. ZALKIKAR (2001). Goodness of fit tests for multivariate counting process models with applications. *Scandinavian Journal of Statistics* **28**(1), pp. 241–256.
- [49] I. M. TOKE and F. POMPONIO (2012). Modelling trades-through in a limit order book using Hawkes processes. *Economics – The Open-Access, Open-Assessment E-Journal* **6**, pp. 1–23.
- [50] H. J. T. UNWIN, I. ROUTLEDGE, S. FLAXMAN, M.-A. RIZOIU, S. LAI, J. COHEN, D. J. WEISS, S. MISHRA, and S. BHATT (2021). Using Hawkes Processes to model imported and local malaria cases in near-elimination settings. *PLoS Computational Biology* **17**(4), e1008830.

- [51] A. W. VAN DER VAART (1998). *Asymptotic Statistics*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press.
- [52] A. VEEN and F. P. SCHOENBERG (2008). Estimation of space–time Branching process models in seismology using an EM–type algorithm. *Journal of the American Statistical Association* **103**, pp. 614–624.
- [53] W. WHITT (2002). *Stochastic-Process Limits: An Introduction to Stochastic-Process Limits and Their Application to Queues*. Springer Series in Operations Research.
- [54] L. ZHU (2013). Central limit theorem for nonlinear Hawkes processes. *Journal of Applied Probability* **50**, pp. 760–771.

## APPENDIX

For the following set of assumptions, let  $\mathcal{H}_{0,t} := \sigma(N_\theta(s) : s \in ([0, t]))$ , and let  $\lambda_\theta^*$  be any  $(\mathcal{H}_t)_{t \in \mathbb{R}^-}$ -predictable function such that

$$\lambda_\theta^*(t) = \lim_{\Delta t \downarrow 0} \frac{1}{\Delta t} \mathbb{P}(N_\theta[t, t + \Delta t] > 0 | \mathcal{H}_{0,t}).$$

### Assumptions C.1(v).

(a)  $\mathbb{E}[\sup_{\delta \in (0,1]} \delta^{-1} (N[0, \delta])^2] < \infty$ .

(b) For each  $\theta \in \Theta$ , there exists a neighborhood  $U = U(\theta) \subset \Theta$  of  $\theta$  such that if

$$H(t, \omega) := \max_{\substack{i,j,k \in [m] \\ \ell \in [d]}} \sup_{\theta' \in U} \left| \frac{\partial^3 \lambda_{\theta'}^{(\ell)}}{\partial \theta_i \partial \theta_j \partial \theta_k} \right|;$$

$$G(t, \omega) := \max_{\substack{i,j,k \in [m] \\ \ell \in [d]}} \sup_{\theta' \in U} \left| \frac{\partial^3 \log \lambda_{\theta'}^{(\ell)}}{\partial \theta_i \partial \theta_j \partial \theta_k} \right|,$$

then  $\mathbb{E}[H(0, \omega)] < \infty$  and  $\mathbb{E}[\|\lambda_{\theta_0}(0, \omega)\|^2 G(0, \omega)^2] < \infty$ .

(c) For each  $\theta \in \Theta$ ,  $i, j \in [m]$ , the following tend to zero in probability as  $t \rightarrow \infty$ :

$$\lambda_\theta - \lambda_\theta^*, \quad \frac{\partial \lambda_\theta}{\partial \theta_i} - \frac{\partial \lambda_\theta^*}{\partial \theta_i} \quad \text{and} \quad \frac{\partial^2 \lambda_\theta}{\partial \theta_i \partial \theta_j} - \frac{\partial^2 \lambda_\theta^*}{\partial \theta_i \partial \theta_j}.$$

(d) For each  $\theta \in \Theta$ , there is some  $\alpha > 0$  such that the following have finite, uniformly (w.r.t.  $t$ ) bounded  $(2 + \alpha)$ th moments

$$\frac{\lambda_\theta}{\lambda_\theta^*}, \quad \frac{1}{\lambda_\theta^*} \frac{\partial \lambda_\theta^*}{\partial \theta_i} \frac{\partial \lambda_\theta^*}{\partial \theta_j} \quad \text{and} \quad \frac{\partial^2 \lambda_\theta^*}{\partial \theta_i \partial \theta_j}, \quad i, j \in [m].$$

(e) For each  $\theta \in \Theta$ ,  $i \in [m]$ , as  $T \rightarrow \infty$

$$\mathbb{E} \left[ \frac{1}{\sqrt{T}} \int_0^T \left| \frac{\partial \lambda_\theta}{\partial \theta_i} - \frac{\partial \lambda_\theta^*}{\partial \theta_i} \right| dt \right] \rightarrow 0;$$

$$\mathbb{E} \left[ \frac{1}{\sqrt{T}} \int_0^T |\lambda_\theta - \lambda_\theta^*| \left| \frac{1}{\lambda_\theta^*} \frac{\partial \lambda_\theta^*}{\partial \theta_i} \right| dt \right] \rightarrow 0.$$

(f) For each  $\theta \in \Theta$ , there exists a neighborhood  $U = U(\theta) \subset \Theta$  of  $\theta$  such that

$$\max_{\substack{i,j,k \in [m] \\ \ell \in [d]}} \sup_{\theta' \in U} \left| \frac{\partial^3 \lambda_{\theta'}^{(\ell)}(t)}{\partial \theta_i \partial \theta_j \partial \theta_k} - \frac{\partial^3 (\lambda_{\theta'}^*)^{(\ell)}(t)}{\partial \theta_i \partial \theta_j \partial \theta_k} \right| \xrightarrow{\mathbb{P}} 0 \text{ as } t \rightarrow \infty.$$

(g) For each  $\theta \in \Theta$ , there exists a neighborhood  $U = U(\theta) \subset \Theta$  of  $\theta$  and some  $\alpha > 0$  such that

$$\max_{\substack{i,j,k \in [m] \\ \ell \in [d]}} \sup_{\theta' \in U} \frac{\partial^3 (\lambda_{\theta'}^*)^{(\ell)}}{\partial \theta_i \partial \theta_j \partial \theta_k} \quad \text{and} \quad \max_{\substack{i,j,k \in [m] \\ \ell \in [d]}} \sup_{\theta' \in U} \frac{\partial^3 \log (\lambda_{\theta'}^*)^{(\ell)}}{\partial \theta_i \partial \theta_j \partial \theta_k}$$

have finite, uniformly (w.r.t.  $t$ ) bounded  $(2 + \alpha)$ th moments.