

Finite sums associated with some polynomial identities

Kunle Adegoke

Department of Physics and Engineering Physics
Obafemi Awolowo University
220005 Ile-Ife, Nigeria
adegoke00@gmail.com

Robert Frontczak

Independent Researcher
72762 Reutlingen, Germany
robert.frontczak@web.de

Karol Gryszka

Institute of Mathematics
University of the National Education Commission, Krakow
Podchorążych 2, 30-084 Kraków, Poland
karol.gryszka@uken.krakow.pl

Abstract

In this paper, we present a general framework for the derivation of interesting finite combinatorial sums starting with certain classes of polynomial identities. The sums that can be derived involve products of binomial coefficients and also harmonic numbers and squared harmonic numbers. We apply the framework to discuss combinatorial sums associated with some prominent polynomial identities from the recent past.

2020 Mathematics Subject Classification: 05A10, 05A19, 11B83.

Keywords: Polynomial identity, Combinatorial identity, Harmonic number, Binomial coefficient.

1 Motivation and Preliminaries

Harmonic numbers $(H_n)_{n \geq 0}$ are defined by

$$H_0 = 0 \quad \text{and for } n \geq 1 : \quad H_n = \sum_{k=1}^n \frac{1}{k}.$$

Due to their ubiquitous connections with combinatorics and number theory harmonic numbers and generalized harmonic numbers are very popular research objects. The mathematical literature is extremely rich, still growing constantly (see the papers of Batir [5, 6, 7, 8], Boyadzhiev [9, 10], Choi [12, 13, 14], Chu [15, 16, 17], Genčev [22], Jin and Du [24], Sofo and Srivastava [29], Wang and Wei [32], Wei et al. [33], Wei and Wang [34] or Xu [35], to name a few recent and important articles in the subject).

In 2009, Boyadzhiev [9] studied binomial sums with harmonic numbers using the Euler transform. His main result is the following identity valid for $n \geq 1$ and all $t \in \mathbb{C}$

$$\sum_{k=1}^n \binom{n}{k} H_k t^k = (1+t)^n \left(H_n - \sum_{k=1}^n \frac{1}{k(1+t)^k} \right). \quad (1)$$

We have learned recently, however, that Boyadzhiev's identity (1) was discovered earlier by Donald Knuth. Knuth published it in his book [25, Page 77, Eq. (10)] in 1997. So, Boyadzhiev actually only rediscovered this identity, which will be referred to as the Knuth-Boyadzhiev (KB) identity. Frontczak [20] obtained a complement of the KB identity given in (1) and derived

$$\sum_{k=0}^n \binom{n}{k} H_k t^k = ((1+t)^n - 1) H_n - t \sum_{k=0}^{n-1} H_{n-1-k} (1+t)^k. \quad (2)$$

In addition, Frontczak [21] also derived an analogue formula involving the skew-harmonic numbers. It is also notable that the KB identity is a special case of a binomial sum identity involving multiple harmonic-like numbers derived by Adegoke and Frontczak [3]. Batir [7] and Batir and Sofo [8] did research in the same direction. Among other things, they derived in [8] the identity

$$\sum_{k=1}^n \binom{n}{k} A_k t^k = (1+t)^{n-1} \sum_{k=0}^{n-1} \left(\sum_{j=0}^k \binom{k}{j} a_{j+1} t^{j+1} \right) \frac{1}{(1+t)^k}, \quad (3)$$

where $(a_n)_{n \geq 1}$ is any sequence in \mathbb{C} , and $A_n = a_1 + a_2 + \dots + a_n$ is its n th partial sum with $A_0 = 0$. Identity (3) is another generalization of the KB result and contains identity (1) as a special case for $a_n = 1/n$. We also note that Frontczak's identity (2) is stated wrongly in [8]. In [7] Batir proved a generalization of (3) involving generalized binomial coefficients.

In this paper, we present a general framework (outlined in Section 3) for deriving various finite combinatorial sums involving (central) binomial coefficients, harmonic numbers, odd harmonic numbers, and squared harmonic numbers. The framework can be succinctly described as a set of transformations applied to a given polynomial identity (like (1) or (3)) which leads to many associated combinatorial sums. For example, we will demonstrate that the finite sum

$$\sum_{k=1}^n (-1)^{k-1} \frac{\binom{n}{k}}{\binom{k+r}{s}} H_k = \frac{s}{r-s+1} \left(\sum_{k=1}^n \frac{1}{k \binom{n-k+r}{r-s+1}} - \frac{H_n}{\binom{n+r}{r-s+1}} \right), \quad (4)$$

which holds for all $s, r \in \mathbb{C} \setminus \mathbb{Z}^-$ such that $s \neq 0$ and $r - s \notin \mathbb{Z}^-$, is a consequence of (1). Identity (4) contains several interesting sums as direct descendents and special cases. Furthermore, we will derive some presumably new identities involving binomial coefficients, such as the following identity

$$\sum_{k=0}^n (-1)^k \frac{2^{2k}}{2k+1} \frac{\binom{n}{k}}{\binom{2k}{k}} = \frac{1}{2n+1}$$

which results from applying the general framework to Chebyshev polynomials of the second kind.

We apply the framework to discuss combinatorial sums associated with several prominent polynomial identities from the recent past. It is important to note that we do not explicitly formulate every possible result based on these identities to avoid repetition of some overly complex expression. For additional polynomial identities to which our general framework can be applied, as well as more combinatorial sums that can be derived, we refer the reader to Section 6.

2 Definitions and required identities

Harmonic numbers H_s and odd harmonic numbers O_s are defined for $0 \neq s \in \mathbb{C} \setminus \mathbb{Z}^-$ by the recurrence relations

$$H_s = H_{s-1} + \frac{1}{s} \quad \text{and} \quad O_s = O_{s-1} + \frac{1}{2s-1},$$

with $H_0 = 0$ and $O_0 = 0$. One way to generalize this definition is to consider harmonic numbers $H_s^{(m)}$ and odd harmonic numbers $O_s^{(m)}$ of order $m \in \mathbb{C}$ that are defined by

$$H_s^{(m)} = H_{s-1}^{(m)} + \frac{1}{s^m} \quad \text{and} \quad O_s^{(m)} = O_{s-1}^{(m)} + \frac{1}{(2s-1)^m},$$

with $H_0^{(m)} = 0$ and $O_0^{(m)} = 0$ so that $H_s = H_s^{(1)}$ and $O_s = O_s^{(1)}$. The recurrence relations imply that if $s = n$ is a non-negative integer, then

$$H_n^{(m)} = \sum_{j=1}^n \frac{1}{j^m} \quad \text{and} \quad O_n^{(m)} = \sum_{j=1}^n \frac{1}{(2j-1)^m}.$$

In particular, we have the connection between harmonic numbers and the psi (or digamma) function [30]

$$H_z = \psi(z+1) + \gamma, \tag{5}$$

where γ is the Euler-Mascheroni constant.

Lemma 1. For $u, v \in \mathbb{C} \setminus \mathbb{Z}^-$, we have

$$\int_0^1 x^u (1-x)^v dx = \frac{1}{\binom{u+v+1}{u+1}(u+1)}. \quad (6)$$

Proof. The identity (6) is the well-known Beta function integral. \square

Lemma 2. If n is an integer, then

$$\begin{aligned} H_{n-1/2} &= 2O_n - 2 \ln 2 \\ H_{n-1/2} - H_{-1/2} &= 2O_n, \\ H_{n-1/2} - H_{1/2} &= 2(O_n - 1), \\ H_{n+1/2} - H_{-1/2} &= 2O_{n+1}, \\ H_{n+1/2} - H_{1/2} &= 2(O_{n+1} - 1), \\ H_{n+1/2} - H_{n-1/2} &= \frac{2}{2n+1}, \\ H_{n-1/2} - H_{-3/2} &= 2(O_n - 1), \\ H_{n+1/2} - H_{-3/2} &= 2(O_{n+1} - 1). \end{aligned} \quad (7)$$

Proof. Use (5) as the definition of the harmonic numbers in conjunction with the known result for the digamma function at half-integer arguments [30, Eq. (51)], namely,

$$\psi(n+1/2) = -\gamma - 2 \ln 2 + 2 \sum_{k=1}^n \frac{1}{2k-1}.$$

\square

Lemma 3. We have

$$\binom{r+1/2}{s} = \frac{\binom{2r+1}{2s} \binom{2s}{s}}{\binom{r}{s} 2^{2s}}, \quad (8)$$

$$\binom{1/2}{r} = (-1)^{r+1} \frac{\binom{2r}{r}}{2^{2r} (2r-1)},$$

$$\binom{r-1/2}{s} = \binom{2r}{r} \binom{r}{s} \frac{1}{\binom{2(r-s)}{r-s} 2^{2s}}, \quad (9)$$

$$\binom{-1/2}{r} = (-1)^r \binom{2r}{r} 2^{-2r}, \quad (10)$$

$$\binom{r}{s+1/2} = \frac{1}{\pi} \frac{2^{2r+2}}{s+1} \binom{r}{s} \binom{2(r-s)}{r-s}^{-1} \binom{2(s+1)}{s+1}^{-1}.$$

Proof. These consequences of the generalized binomial coefficients, are easy to derive using the Gamma function and some of its special values. At any rate they are entries in Gould's book [23]. \square

3 General results for arbitrary polynomial identities

In this section, we introduce a set of general results that are applicable for a large family of polynomial identities. The polynomial identities under consideration are of the general form

$$\sum_{k=l_1}^{u_1} f(k)W_k(t) = \sum_{k=l_2}^{u_2} g(k)V_k(t), \quad (11)$$

where W_k, V_k are polynomials belonging to the following set (we skip the dependence on k and n to simplify the notation)

$$BP = \{t^k, t^{n-k}, (1-t)^k, (1-t)^{n-k}, t^k(1-t)^{n-k}, t^{n-k}(1-t)^k\},$$

l_1, l_2, u_1, u_2 are non-negative integers, $f(k)$ and $g(k)$ are sequences and t is a complex variable. These will be our standing assumptions for the rest of the paper. To deduce the results in topic, we perform a series of basic operations in order, which we call *the (general) algorithm*. We describe that in details in the following example.

Example 4. Suppose that the polynomial identity contains on one of the sides the following sum:

$$\sum_{k=l_1}^{u_1} f(k)(1-t)^k. \quad (12)$$

We now multiply that by $(1-t)^r t^{s-1}$ to obtain

$$\sum_{k=l_1}^{u_1} f(k)(1-t)^{k+r} t^{s-1}.$$

The next step is to integrate termwise from 0 to 1 and use the Beta integral representation (6), which gives

$$\sum_{k=l_1}^{u_1} f(k) \frac{1}{s \cdot \binom{k+r+s}{s}}.$$

The final usual step is to substitute $r \mapsto r-s$ and multiply the equation by s to eliminate an unwanted factor in the denominator. This gives the final form of the identity without harmonic numbers

$$\sum_{k=l_1}^{u_1} f(k) \frac{1}{\binom{k+r}{s}} \quad (13)$$

of (12). Note that (12) is just a part of some, unspecified, polynomial identity and a set of operations described above is performed on both sides of that identity.

We can now perform one more operation to "generate" harmonic numbers from (13). We differentiate that with respect to s . In case of (13) this gives the following sum:

$$\sum_{k=l_1}^{u_1} f(k) \frac{H_s - H_{k+r-s}}{\binom{k+r}{s}} \quad (14)$$

where $s, r \in \mathbb{C} \setminus \mathbb{Z}^-$, $s \neq 0$ and $r - s \notin \mathbb{Z}^-$, which are again the standing assumptions for these kind of terms. We can also differentiate with respect to r , which gives

$$\sum_{k=l_1}^{u_1} f(k) \frac{H_{k+r-s} - H_{k+r}}{\binom{k+r}{s}}. \quad (15)$$

The above example allows us to transform any identity (11) with polynomials in the set BP to an identity involving binomial coefficients and harmonic numbers.

The polynomial identities under consideration that can be applied to the described method are of the following form (which we will refer to as *standard forms*)

$$\sum_{k=l_1}^{u_1} f(k)t^k = \sum_{k=l_2}^{u_2} g(k)t^k, \quad (16)$$

$$\sum_{k=l_1}^{u_1} f(k)t^{n-k} = \sum_{k=l_2}^{u_2} g(k)t^k, \quad (17)$$

$$\sum_{k=l_1}^{u_1} f(k)(1-t)^k = \sum_{k=l_2}^{u_2} g(k)t^k, \quad (18)$$

$$\sum_{k=l_1}^{u_1} f(k)(1-t)^{n-k} = \sum_{k=l_2}^{u_2} g(k)t^k, \quad (19)$$

$$\sum_{k=l_1}^{u_1} f(k)t^k(1-t)^{n-k} = \sum_{k=l_2}^{u_2} g(k)t^k, \quad (20)$$

$$\sum_{k=l_1}^{u_1} f(k)t^{n-k}(1-t)^k = \sum_{k=l_2}^{u_2} g(k)t^k. \quad (21)$$

We note that for example

$$\sum_{k=l_1}^{u_1} f(k)(1+t)^k = \sum_{k=l_2}^{u_2} g(k)t^k$$

is not on the above list, but substitution $t \mapsto -t$ shows it is of the form (18). All standard forms have their right-hand-side equal and it is just a matter of simple changes to work with any possible identity in the BP set.

Let us explicitly formulate the results for identities (18) and (20), but the very similar results (with obvious and necessary changes) can be derived and presented for the remaining cases.

Lemma 5. *Let an arbitrary polynomial identity have the following form:*

$$\sum_{k=l_1}^{u_1} f(k)(1-t)^k = \sum_{k=l_2}^{u_2} g(k)t^k. \quad (22)$$

Let $r, s \in \mathbb{C} \setminus \mathbb{Z}^-$ are such that $s \neq 0$ and $r - s \notin \mathbb{Z}^-$. Then the following identity holds:

$$\sum_{k=l_1}^{u_1} f(k) \frac{1}{\binom{k+r}{s}} = \sum_{k=l_2}^{u_2} g(k) \frac{s}{r-s+1} \frac{1}{\binom{k+r}{r-s+1}} \quad (23)$$

Furthermore, if $s, r \in \mathbb{C} \setminus \mathbb{Z}^-$ such that $s \neq 0$ and $r - s \notin \mathbb{Z}^-$, then

$$\sum_{k=l_1}^{u_1} f(k) \frac{H_s - H_{k+r-s}}{\binom{k+r}{s}} = \sum_{k=l_2}^{u_2} g(k) \frac{s+1}{(r-s+1)^2} \frac{1}{\binom{k+r}{r-s+1}} + \sum_{k=l_2}^{u_2} g(k) \frac{s}{r-s+1} \frac{H_{k+s-1} - H_{r-s+1}}{\binom{k+r}{r-s+1}}. \quad (24)$$

In the above lemma, we differentiated with respect to s to obtain (24). In the next lemma, we differentiated with respect to r to obtain (27).

Lemma 6. Let an arbitrary polynomial identity have the following form:

$$\sum_{k=l_1}^{u_1} f(k) t^k (1-t)^{n-k} = \sum_{k=l_2}^{u_2} g(k) t^k. \quad (25)$$

Let $r, s \in \mathbb{C} \setminus \mathbb{Z}^-$ are such that $s \neq 0$ and $r - s \notin \mathbb{Z}^-$. Then the following identity holds:

$$\sum_{k=l_1}^{u_1} \frac{f(k)}{(k+s) \binom{n+r}{k+s}} = \sum_{k=l_1}^{u_1} \frac{g(k)}{(k+s) \binom{k+r}{k+s}} \quad (26)$$

Furthermore, if $s, r \in \mathbb{C} \setminus \mathbb{Z}^-$ such that $s \neq 0$ and $r - s \notin \mathbb{Z}^-$, then

$$\sum_{k=l_1}^{u_1} f(k) \frac{H_{n-k+r-s} - H_{n+r}}{(k+s) \binom{n+r}{k+s}} = \sum_{k=l_1}^{u_1} g(k) \frac{H_{r-s} - H_{k+r}}{(k+s) \binom{k+r}{k+s}} \quad (27)$$

More identities that are special cases of the above and general ones can be deduced. For example, we have the following.

Corollary 7. Under the assumptions of Lemma 5, we have

$$\sum_{k=l_1}^{u_1} \frac{f(k)}{\binom{k+s}{s}} = \sum_{k=l_2}^{u_2} \frac{g(k)s}{k+s}, \quad \sum_{k=l_1}^{u_1} \frac{f(k)}{k+1} = \sum_{k=l_2}^{u_2} \frac{g(k)}{k+1},$$

and

$$\begin{aligned} \sum_{k=l_1}^{u_1} f(k) \frac{H_s - H_k}{\binom{k+s}{s}} &= \sum_{k=l_2}^{u_2} g(k) \frac{s+1}{k+s} + \sum_{k=l_2}^{u_2} g(k) s \frac{H_{k+s-1} - 1}{k+s}, \\ \sum_{k=l_1}^{u_1} f(k) \frac{1 - H_k}{k+1} &= \sum_{k=l_2}^{u_2} g(k) \frac{2}{k+1} + \sum_{k=l_2}^{u_2} g(k) \frac{H_k - 1}{k+1}, \end{aligned}$$

Proof. Substitute $r = s$ in Lemma 5 to obtain the first identity in each set. Then, take $s = 1$ to obtain the second one. \square

Similar set of results can be made after applying Lemma 6.

Corollary 8. *Under the assumptions of Lemma 6, we have*

$$\sum_{k=l_1}^{u_1} \frac{f(k)}{(k+s)\binom{n+s}{k+s}} = \sum_{k=l_2}^{u_2} \frac{g(k)}{k+s}, \quad \sum_{k=l_1}^{u_1} \frac{f(k)}{(k+1)\binom{n+1}{k+1}} = \sum_{k=l_2}^{u_2} \frac{g(k)}{k+1},$$

and

$$\begin{aligned} \sum_{k=l_1}^{u_1} f(k) \frac{H_{n-k} - H_{n+s}}{(k+s)\binom{n+s}{k+s}} &= - \sum_{k=l_2}^{u_2} g(k) \frac{H_{k+s}}{k+s}, \\ \sum_{k=l_1}^{u_1} f(k) \frac{H_{n-k} - H_{n+1}}{(k+1)\binom{n+1}{k+1}} &= - \sum_{k=l_2}^{u_2} g(k) \frac{H_{k+1}}{k+1}, \end{aligned}$$

The aim of the next sections is to generate identities of the above kind using some initial polynomial identity.

3.1 A simple application

To give the reader an idea of how the general identities could be use and how universal they can be, let us consider the binomial identity

$$(1-t)^n = \sum_{k=0}^n (-1)^k \binom{n}{k} t^k$$

Here, we use Corollary 7 with $s = 1$, $l_1 = u_1 = n$, $l_2 = 0$, $u_2 = n$, $f(n) = 1$ and $g(k) = (-1)^k \binom{n}{k}$. This gives, after some simple algebraic manipulation, the following well-known identities:

$$\frac{1}{n+1} = \sum_{k=0}^n (-1)^k \binom{n}{k} \frac{1}{k+1}, \quad (28)$$

$$\frac{H_n}{n+1} = \sum_{k=0}^n (-1)^{k+1} \binom{n}{k} \frac{H_{k+1}}{k+1}. \quad (29)$$

To see the application of Corollary 8, let us consider

$$\sum_{k=0}^n t^k (1+t)^{n-k} = (t+1)^{n+1} - t^{n+1} = \sum_{k=0}^n \binom{n+1}{k} t^k.$$

From that we get:

$$\sum_{k=0}^n \frac{(-1)^k}{(k+1) \binom{n+1}{k+1}} = \sum_{k=0}^n \frac{(-1)^k}{k+1} \binom{n+1}{k},$$

$$\sum_{k=0}^n (-1)^k \frac{H_{n-k} - H_{n+1}}{(k+1) \binom{n+1}{k+1}} = \sum_{k=0}^n (-1)^{k+1} \binom{n+1}{k} \frac{H_{k+1}}{k+1}.$$

The first identity may look curious, but in fact both sides of it are 0, which follows directly from (28).

4 Applications

4.1 Sums associated with polynomial identities of Knuth-Boyadzhiev and Frontczak

Recall that the identity of Knuth and Boyadzhiev (1) can be stated as

$$\sum_{k=1}^n (-1)^{k-1} \binom{n}{k} H_k t^k = -(1-t)^n H_n + \sum_{k=1}^n \frac{(1-t)^{n-k}}{k}, \quad (\text{A})$$

whereas Frontczak's compliment (2) has the form

$$\sum_{k=0}^n (-1)^k \binom{n}{k} H_k t^k = ((1-t)^n - 1) H_n + t \sum_{k=0}^{n-1} H_{n-1-k} (1-t)^k. \quad (\text{B})$$

These forms are members of the polynomial family introduced in the previous section. The next results are therefore immediate.

Theorem 9. *If n is a non-negative integer, then*

$$\sum_{k=0}^n (-1)^k \binom{n}{k} \frac{(2k+1)}{2^{2k}} \binom{2k}{k} H_{k+1} = \frac{\binom{2n}{n} H_{n+1}}{2^{2n} n+1} - \frac{2}{n+1} \sum_{k=1}^n \frac{\binom{2(n-k)}{n-k}}{2^{2(n-k)} k}, \quad (30)$$

and, more generally, if $s, r \in \mathbb{C} \setminus \mathbb{Z}^-$ such that $s \neq 0$ and $r-s \notin \mathbb{Z}^-$, then

$$\sum_{k=1}^n (-1)^{k-1} \frac{\binom{n}{k}}{\binom{k+r}{s}} H_k = \frac{s}{r-s+1} \left(\sum_{k=1}^n \frac{1}{k \binom{n-k+r}{r-s+1}} - \frac{H_n}{\binom{n+r}{r-s+1}} \right). \quad (31)$$

Proof. To obtain (31), we apply the general algorithm to (A). Identity (30) is obtained by setting $r = -1$, $s = -1/2$ in (31). Note that we used

$$\lim_{x \rightarrow 0} \frac{1}{\binom{x-1}{1/2}} = 0.$$

□

Corollary 10. For any $r \in \mathbb{C} \setminus \mathbb{Z}^-$ we have

$$\sum_{k=1}^n (-1)^{k-1} \frac{\binom{n}{k}}{\binom{k+r}{r}} H_k = \frac{r}{n+r} (H_{n+r} - H_r) + \frac{n}{(n+r)^2}. \quad (32)$$

Proof. Set $s = r$ in (31). This gives

$$\sum_{k=1}^n (-1)^{k-1} \frac{\binom{n}{k}}{\binom{k+r}{r}} H_k = r \sum_{k=1}^n \frac{1}{k(n-k+r)} - \frac{r}{n+r} H_n.$$

But

$$\begin{aligned} \sum_{k=1}^n \frac{1}{k(n-k+r)} &= \frac{1}{n+r} \sum_{k=1}^n \left(\frac{1}{k} + \frac{1}{n-k+r} \right) \\ &= \frac{1}{n+r} (H_n + H_{n-1+r} - H_{r-1}) \end{aligned}$$

and hence

$$\sum_{k=1}^n (-1)^{k-1} \frac{\binom{n}{k}}{\binom{k+r}{r}} H_k = \frac{r}{n+r} (H_{n-1+r} - H_{r-1}).$$

□

Remark. Corollary 10 is Corollary 1.2 in Chu's paper [16] and also Theorem 7 in [4].

Theorem 11. If n is a non-negative integer, then

$$\begin{aligned} \sum_{k=0}^n (-1)^k \frac{2k+1}{2^{2k}(k+1)} \binom{n}{k} \binom{2k}{k} H_{k+1} O_{k+1} &= -\frac{\binom{2n}{n}}{2^{2n}(n+1)} H_{n+1} (O_n - 1) \\ &+ \frac{1}{n+1} \sum_{k=1}^n \frac{\binom{2(n-k)}{n-k}}{2^{2(n-k)} k} (O_{n-k} - 1), \end{aligned} \quad (33)$$

and, more generally, if $s, r \in \mathbb{C} \setminus \mathbb{Z}^-$ such that $s \neq 0$ and $r-s \notin \mathbb{Z}^-$, then

$$\begin{aligned} \sum_{k=1}^n (-1)^{k-1} \binom{n}{k} H_k \frac{H_{k+r-s} - H_s}{\binom{k+r}{s}} \\ = -\frac{r+1}{(r-s+1)^2} \sum_{k=1}^n \frac{1}{k \binom{n-k+r}{r-s+1}} - \frac{s}{r-s+1} \sum_{k=1}^n \frac{H_{n-k+s-1} - H_{r-s+1}}{k \binom{n-k+r}{r-s+1}} \\ + \frac{(r+1) H_n}{(r-s+1)^2 \binom{n+r}{r-s+1}} - \frac{s H_n (H_{r-s+1} - H_{n+s-1})}{(r-s+1) \binom{n+r}{r-s+1}}. \end{aligned} \quad (34)$$

Proof. Differentiate (31) with respect to s . □

Corollary 12. For any nonzero $r \in \mathbb{C} \setminus \mathbb{Z}^-$ we have

$$\begin{aligned} \sum_{k=1}^n (-1)^{k-1} \frac{\binom{n}{k}}{\binom{k+r}{r}} H_k^2 &= \frac{1}{n+r} (H_{n+r} - H_r) (rH_r - 1) + \frac{n}{(n+r)^2} \left(H_r - \frac{1}{r} \right) \\ &\quad + \frac{r}{n+r} H_n H_{n+r-1} - r \sum_{k=1}^n \frac{H_{n-k+r-1}}{k(n-k+r)}. \end{aligned} \quad (35)$$

In particular,

$$\sum_{k=1}^n (-1)^{k-1} \frac{\binom{n}{k}}{k+1} H_k^2 = \frac{1}{2(n+1)} (3H_n^{(2)} - H_n^2). \quad (36)$$

Proof. Set $s = r$ in (34) and simplify using Corollary 10. The special case follows upon setting $r = 1$ in conjunction with

$$\sum_{k=1}^n \frac{H_{n-k}}{k} = H_n^2 - H_n^{(2)},$$

and

$$\sum_{k=1}^n \frac{H_{n-k}}{n-k+1} = \frac{1}{2} (H_n^2 - H_n^{(2)}).$$

□

Remark. Identity (36) is Equation (2.22) in Theorem 2 of Choi [13]. There is also the identity (corresponding to $r = 0$ in the above relation)

$$\sum_{k=1}^n (-1)^k \binom{n}{k} H_k^2 = \frac{H_n}{n} - \frac{2}{n^2}. \quad (37)$$

This identity has been obtained by Wang [31] by the method of Riordan arrays, and rediscovered by Boyadzhiev [11].

A different expression for the sum (31) can be derived from Frontczak's identity (B).

Theorem 13. If $s, r \in \mathbb{C} \setminus \mathbb{Z}^-$ such that $s \neq 0$ and $r - s \notin \mathbb{Z}^-$, then

$$\sum_{k=0}^n (-1)^k \frac{\binom{n}{k}}{\binom{k+r}{s}} H_k = H_n \left(\frac{s}{r-s+1} \frac{1}{\binom{n+r}{r-s+1}} - \frac{1}{\binom{n}{s}} \right) + s \sum_{k=0}^{n-1} \frac{H_{n-1-k}}{(k+s)\binom{r+k+1}{r-s+1}}. \quad (38)$$

Proof. Apply the general algorithm to (B) and simplify. □

Corollary 14. *If $s, r \in \mathbb{C} \setminus \mathbb{Z}^-$ such that $s \neq 0$ and $r - s \notin \mathbb{Z}^-$, then*

$$\sum_{k=0}^{n-1} \frac{H_{n-1-k}}{(k+s)\binom{r+k+1}{r-s+1}} = \frac{H_n}{s\binom{r}{s}} - \frac{1}{r-s+1} \sum_{k=0}^n \frac{1}{k\binom{n-k+r}{r-s+1}}. \quad (39)$$

In particular, we have for all $s \geq 1$

$$\sum_{k=0}^n \frac{H_{n-k}}{(k+s)(k+1+s)} = \frac{n+1}{s(n+1+s)} H_{n+1} - \frac{1}{n+1+s} (H_{n+s} - H_{s-1}) \quad (40)$$

and

$$\begin{aligned} \sum_{k=0}^n \frac{H_{n-k}}{(k+s)(k+1+s)(k+2+s)} &= \frac{1}{2} \left(\frac{H_{n+1}}{s(s+1)} - \frac{1}{n+1+s} (H_{n+1} + H_{n+s} - H_{s-1}) \right. \\ &\quad \left. + \frac{1}{n+2+s} (H_{n+1} + H_{n+1+s} - H_s) \right). \end{aligned} \quad (41)$$

Proof. Compare (38) with (31). The first particular case follows by setting $r = s$, replacing n by $n + 1$, and using the same evaluation for the sum as in the proof of Corollary 10. To get the sum (41) set $r = s + 1$, replace n by $n + 1$, and make use of the partial fraction decomposition

$$\frac{1}{k(n-k+s)(n-k+s+1)} = \frac{1}{k(n-k+s)} - \frac{1}{k(n-k+s+1)}.$$

□

Theorem 15. *If $s, r \in \mathbb{C} \setminus \mathbb{Z}$ are such that $s \neq 0$ and $r - s \notin \mathbb{Z}$, then*

$$\begin{aligned} \sum_{k=0}^n (-1)^k \binom{n}{k} \frac{H_s - H_{k+r-s}}{\binom{k+r}{s}} H_k &= H_n \left(\frac{r+1}{(r-s+1)^2} \frac{1}{\binom{n+r}{r-s+1}} + \frac{s}{r-s+1} \frac{H_{n+s-1} - H_{r-s+1}}{\binom{n+r}{r-s+1}} \right) \\ &\quad + H_n \frac{H_{r-s} - H_s}{\binom{r}{s}} \\ &\quad + \sum_{k=0}^{n-1} H_{n-1-k} \frac{s(k+s)(H_{k+s} - H_{r-s+1}) + k}{(k+s)^2 \binom{n+r}{r-s+1}}. \end{aligned}$$

Proof. Differentiate (38) with respect to s .

□

Corollary 16. *We have*

$$\sum_{k=0}^n (-1)^k \binom{n}{k} \frac{H_k^2}{\binom{k+r}{r}} = H_n H_r - H_n \left(\frac{1}{n+r} + \frac{r}{n+r} H_{n+r-1} \right) - \frac{r}{n+r} (H_{n-1+r} - H_{r-1})$$

$$\begin{aligned}
& -\frac{r}{n+r} \sum_{k=0}^{n-1} \frac{H_{n-1-k} H_{k+r}}{k+r} + \frac{r}{n+r} \sum_{k=0}^{n-1} \frac{H_{n-k-1}}{k+r} \\
& -\frac{1}{n+r} \sum_{k=0}^{n-1} \frac{k H_{n+k}}{(k+r)^2}
\end{aligned}$$

and in particular,

$$\begin{aligned}
\sum_{k=0}^n (-1)^k \binom{n}{k} \frac{H_k^2}{k+1} &= \frac{nH_n - H_n - H^2}{n+1} + \frac{1}{n+1} (H_n^2 + H_n^{(2)}) \\
& - \frac{1}{n+1} \sum_{k=1}^n \frac{H_{n-k} H_k}{k} - \frac{1}{n+1} \sum_{k=0}^{n-1} \frac{k H_{n+k}}{(k+1)^2}.
\end{aligned}$$

Proof. Set $s = r$ in ... to obtain

$$\begin{aligned}
\sum_{k=0}^n (-1)^k \binom{n}{k} \frac{H_r - H_k}{\binom{k+r}{r}} H_k &= H_n \left(\frac{r+1}{n+r} + \frac{r}{n+r} (H_{n+r-1} - 1) \right) \\
& - H_n H_r \\
& + \sum_{k=0}^{n-1} H_{n-1-k} \frac{r(k+r)(H_{k+r} - 1) + k}{(k+r)^2(n+r)}.
\end{aligned}$$

Use Corollary 10 to the left-hand side and rearrange. This gives the main identity. Particular case follows upon setting $r = 1$. \square

4.2 Sums derived from polynomial identities involving H_n^2 and $H_n^{(2)}$

Next, we recall the following two identities.

Theorem 17. *Let $n \in \mathbb{N}$ and $t \in \mathbb{C}$. Then we have*

$$\sum_{k=0}^n \binom{n}{k} H_k^2 t^k = H_n^2 (1+t)^n - \sum_{k=1}^n \frac{H_n - 2H_{k-1} + H_{n-k}}{k} (1+t)^{n-k} \quad (\text{C})$$

and

$$\sum_{k=0}^n \binom{n}{k} H_k^{(2)} t^k = H_n^{(2)} (1+t)^n - \sum_{k=1}^n \frac{H_n - H_{n-k}}{k} (1+t)^{n-k}. \quad (\text{D})$$

Proof. Identity (C) is a restatement of Sofo's and Batir's result from [8]

$$\sum_{k=0}^n \binom{n}{k} H_k^2 t^k = (1+t)^n \left(H_n^2 - \sum_{k=1}^n \frac{H_n - 2H_k + H_{n-k}}{k(1+t)^k} - 2 \sum_{k=1}^n \frac{1}{k^2(1+t)^k} \right),$$

while (D) follows from combining this identity with

$$\sum_{k=0}^n \binom{n}{k} \left(H_k^2 - H_k^{(2)} \right) t^k = (H_n^2 - H_n^{(2)}) (1+t)^n + 2 \sum_{k=1}^n \frac{H_{k-1} - H_{n-k}}{k} (1+t)^{n-k},$$

which comes from [3]. Identity (D) is also stated as Identity 24 in Choi's paper [12]. \square

Using the general algorithm (as in Theorems 9 and 13) we can deduce the following results.

Theorem 18. *If $s, r \in \mathbb{C} \setminus \mathbb{Z}^-$ such that $s \neq 0$ and $r - s \notin \mathbb{Z}^-$, then*

$$\sum_{k=0}^n (-1)^k \binom{n}{k+r} H_k^{(2)} = \frac{s}{r-s+1} \left(\frac{H_n^{(2)}}{\binom{n+r}{r-s+1}} - \sum_{k=1}^n \frac{H_n - H_{n-k}}{k \binom{n-k+r}{r-s+1}} \right). \quad (42)$$

Corollary 19. *For all $r \geq 1$ we have*

$$\sum_{k=0}^n (-1)^{k+1} \binom{n}{k+r} H_k^{(2)} = \frac{r}{n+r} \left(H_n (H_{n-1+r} - H_{r-1}) - \sum_{k=0}^{n-1} \frac{H_k}{k+r} \right). \quad (43)$$

In particular,

$$\sum_{k=0}^n (-1)^{k+1} \binom{n}{k} \frac{H_k^{(2)}}{k+1} = \frac{1}{2(n+1)} (H_n^2 + H_n^{(2)}) \quad (44)$$

and

$$\sum_{k=0}^n (-1)^k \binom{n}{k} \frac{H_k^{(2)}}{(k+1)(k+2)} = \frac{1}{n+1} \left(\frac{1}{2} (H_{n+1}^2 - H_{n+1}^{(2)}) + H_n - H_n H_{n+1} - \frac{n}{n+1} \right). \quad (45)$$

Remark. Choi [13] states in Theorem 2, Equation (2.23), the result

$$\sum_{k=1}^n (-1)^{k+1} \binom{n}{k} \frac{H_k^{(2)}}{k+1} = \frac{1}{2(n+1)} (5H_n^{(2)} - 3H_n^2),$$

which is not correct.

4.3 Sums associated with an identity of Dattoli et al.

In this subsection we work with the following polynomial identity due to Dattoli et al. [19]:

$$\sum_{k=0}^n \binom{n}{k} (-1)^k \frac{1}{k+2} t^k (1+t)^{n-k} = \sum_{k=0}^n \binom{n}{k} \frac{t^k}{(k+1)(k+2)}. \quad (E)$$

Using the general algorithm again (as in previous sections) we can transform (E) into our next result.

Theorem 20. If $s, r \in \mathbb{C} \setminus \mathbb{Z}^-$ such that $s \neq 0$ and $r - s \notin \mathbb{Z}^-$, then

$$\sum_{k=0}^n \binom{n}{k} \frac{1}{(k+2)(k+s)\binom{n+r}{k+s}} = \sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(k+1)(k+2)(k+s)\binom{k+r}{k+s}}. \quad (46)$$

Corollary 21. For all $r \geq 1$ we have

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(k+1)(k+2)(k+r)} = \frac{1}{(n+1)\cdots(n+r)} \sum_{k=0}^n \frac{\prod_{j=1}^{r-1}(k+j)}{k+2}, \quad (47)$$

where the product equals 1 for $r = 1$. In particular, we have the identities

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(k+1)^2(k+2)} = \frac{H_{n+2} - 1}{n+1}, \quad (48)$$

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(k+1)(k+2)^2} = \frac{n+2 - H_{n+2}}{(n+1)(n+2)}, \quad (49)$$

and

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(k+1)(k+2)(k+3)} = \frac{1}{2(n+3)}. \quad (50)$$

Proof. Set $r = s$ in Theorem 20 and simplify using

$$\binom{n+r}{k+r} = \frac{(n+r)\cdots(n+1)}{(k+r)\cdots(k+1)} \binom{n}{k}.$$

The particular examples correspond to $r = 1$, $r = 2$ and $r = 3$, respectively, where it was used that

$$\sum_{k=0}^n \frac{1}{k+2} = H_{n+2} - 1,$$

$$\sum_{k=0}^n \frac{k+1}{k+2} = n+2 - H_{n+2},$$

and

$$\sum_{k=0}^n (k+1) = \frac{(n+1)(n+2)}{2}.$$

□

Theorem 22. If $s, r \in \mathbb{C} \setminus \mathbb{Z}^-$ such that $s \neq 0$ and $r - s \notin \mathbb{Z}^-$, then

$$\begin{aligned} & \sum_{k=0}^n \binom{n}{k} \frac{H_{n-k+r-s}}{(k+2)(k+s)\binom{n+r}{k+s}} - H_{n+r} \sum_{k=0}^n \binom{n}{k} \frac{1}{(k+2)(k+s)\binom{n+r}{k+s}} \\ &= H_{r-s} \sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(k+1)(k+2)(k+s)\binom{k+r}{k+s}} - \sum_{k=0}^n \binom{n}{k} \frac{(-1)^k H_{k+r}}{(k+1)(k+2)(k+s)\binom{k+r}{k+s}}. \end{aligned} \quad (51)$$

Proof. Differentiate (46) with respect to r . □

Corollary 23. For all $r \geq 1$ we have

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k H_{k+r}}{(k+1)(k+2)(k+r)} = \frac{1}{(n+1) \cdots (n+r)} \left(H_{n+r} \sum_{k=0}^n \frac{\prod_{j=1}^{r-1} (k+j)}{k+2} - \sum_{k=0}^n \frac{H_{n-k}}{k+2} \prod_{j=1}^{r-1} (k+j) \right). \quad (52)$$

In particular, we have the identities

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k H_{k+1}}{(k+1)^2(k+2)} = \frac{1}{n+1} \left(H_{n+1}(H_{n+2} - 1) - \sum_{k=0}^n \frac{H_{n-k}}{k+2} \right), \quad (53)$$

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k H_{k+2}}{(k+1)(k+2)^2} = \frac{1}{n+1} - \frac{1}{(n+1)(n+2)} \left(H_{n+2}^2 - H_{n+1} - \sum_{k=0}^n \frac{H_{n-k}}{k+2} \right), \quad (54)$$

and

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k H_{k+3}}{(k+1)(k+2)(k+3)} = \frac{1}{2(n+3)} \left(H_{n+3} - H_{n+1} + \frac{3n+4}{2(n+2)} \right). \quad (55)$$

Proof. Set $r = s$ in Theorem 22 and simplify. The particular examples correspond to $r = 1$, $r = 2$ and $r = 3$, respectively. Note also that we used

$$\sum_{k=0}^n H_{n-k} = \sum_{k=0}^n H_k = (n+1)(H_{n+1} - 1)$$

as well as

$$\sum_{k=0}^n k H_k = \frac{1}{4} n(n+1)(2H_{n+1} - 1). \quad \square$$

Theorem 24. If $s, r \in \mathbb{C} \setminus \mathbb{Z}^-$ such that $s \neq 0$ and $r - s \notin \mathbb{Z}^-$, then

$$\begin{aligned} & \sum_{k=0}^n \binom{n}{k} \frac{H_{k+s} - H_{n-k+r-s}}{(k+2)(k+s) \binom{n+r}{k+s}} - \sum_{k=0}^n \binom{n}{k} \frac{1}{(k+2)(k+s)^2 \binom{n+r}{k+s}} \\ &= \sum_{k=0}^n \binom{n}{k} \frac{(-1)^k (H_{k+s} - H_{r-s})}{(k+1)(k+2)(k+s) \binom{k+r}{k+s}} - \sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(k+1)(k+2)(k+s)^2 \binom{k+r}{k+s}}. \end{aligned} \quad (56)$$

Proof. Differentiate (46) with respect to s . □

Corollary 25. For all $r \geq 1$ we have

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(k+1)(k+2)(k+r)^2} = \frac{1}{(n+1)\cdots(n+r)} \sum_{k=0}^n \left(H_{n+r} - H_{k+r} + \frac{1}{k+r} \right) \frac{\prod_{j=1}^{r-1} (k+j)}{k+2}. \quad (57)$$

In particular, we have the identities

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(k+1)^3(k+2)} = \frac{1}{n+2} + \frac{H_{n+1}(H_{n+2}-1)}{n+1} - \frac{H_{n+2}^2 - H_{n+2}^{(2)}}{2(n+1)}, \quad (58)$$

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(k+1)(k+2)^3} = \frac{1}{n+1} - \frac{H_{n+2}^2 + H_{n+2}^{(2)}}{2(n+1)(n+2)}, \quad (59)$$

and

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(k+1)(k+2)(k+3)^2} = \frac{H_{n+3}}{(n+1)(n+2)(n+3)} + \frac{n-2}{4(n+1)(n+2)}. \quad (60)$$

Proof. Set $r = s$ in Theorem 24. This gives

$$\begin{aligned} & \frac{1}{(n+1)\cdots(n+r)} \left(\sum_{k=0}^n (H_{k+r} - H_{n-k}) \frac{(k+1)\cdots(k+r-1)}{k+2} - \sum_{k=0}^n \frac{(k+1)\cdots(k+r-1)}{k+2} \right) \\ &= \sum_{k=0}^n \binom{n}{k} \frac{(-1)^k H_{k+r}}{(k+1)(k+2)(k+r)} - \sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(k+1)(k+2)(k+r)^2}. \end{aligned}$$

Now, simplify using (52). The particular examples correspond to $r = 1$, $r = 2$ and $r = 3$, respectively, after some lengthy calculations. Here we used

$$\sum_{k=0}^n \frac{H_{k+1}}{k+2} = \frac{1}{2} \left(H_{n+2}^2 - H_{n+2}^{(2)} \right),$$

$$\sum_{k=0}^n \frac{1}{(k+1)(k+2)} = \frac{n+1}{n+2},$$

$$\sum_{k=0}^n \frac{H_{k+2}}{k+2} = \frac{1}{2} \left(H_{n+2}^2 + H_{n+2}^{(2)} \right) - 1,$$

and

$$\sum_{k=0}^n H_{k+3}(k+1) = \frac{1}{4} (2(n-1)(n+4)H_{n+4} - n^2 + n + 24),$$

all being standard sums. □

4.4 Sums associated with an identity for the Chebyshev polynomials of the second kind

This application deals with sums that are derived using an identity for the Chebyshev polynomials of the second kind $U_n(t)$ defined by

$$U_0(t) = 1, \quad U_1(t) = 2t, \quad U_{n+1}(t) = 2tU_n(t) - U_{n-1}(t).$$

From [2, Lemma 41] we have that

$$\sum_{k=0}^n (-1)^k 2^{2k} \binom{n+k}{n-k} t^{2k} = (-1)^n U_{2n}(t).$$

Using the representation

$$U_n(t) = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n+1}{2k+1} (t^2 - 1)^k t^{n-2k},$$

we immediately get the identity

$$\sum_{k=0}^n (-1)^k 2^{2k} \binom{n+k}{n-k} t^k = \sum_{k=0}^n (-1)^{n-k} \binom{2n+1}{2k+1} (1-t)^k t^{n-k} \quad (\text{F})$$

and the general algorithm can be applied to transform (F) into our next results.

Theorem 26. *If $s, r \in \mathbb{C} \setminus \mathbb{Z}^-$ such that $s \neq 0$ and $r - s \notin \mathbb{Z}^-$, then*

$$\sum_{k=0}^n (-1)^k 2^{2k} \frac{\binom{n+k}{n-k}}{(k+s) \binom{k+r}{k+s}} = \sum_{k=0}^n (-1)^{n-k} \frac{\binom{2n+1}{2k+1}}{(n-k+s) \binom{n+r}{n-k+s}}. \quad (61)$$

Corollary 27. *For all $r \geq 1$ we have*

$$\sum_{k=0}^n (-1)^k 2^{2k} \frac{\binom{n+k}{n-k}}{k+r} = \sum_{k=0}^n (-1)^{n-k} \frac{\binom{2n+1}{2k+1}}{(n-k+r) \binom{n+r}{k}}. \quad (62)$$

In particular, we have the identity

$$\sum_{k=0}^n (-1)^{n-k} \frac{\binom{2n+1}{2k+1}}{\binom{n}{k}} = \frac{(-1)^n (2n+1) - 1}{2n} \quad (n \geq 1). \quad (63)$$

Proof. Set $r = s$ in Theorem 26 and simplify. The particular example corresponds to $r = 1$ in conjunction to

$$\sum_{k=0}^n (-1)^k 2^{2k} \frac{\binom{n+k}{n-k}}{k+1} = \frac{(-1)^n (2n+1) - 1}{2n(n+1)},$$

where the last expression is true since

$$\sum_{k=0}^n (-1)^{n-k} 2^{2k} \frac{\binom{n+k}{n-k}}{k+1} = \int_0^1 U_{2n}(\sqrt{t}) dt = \frac{2n - \cos(\pi n) + 1}{2n(n+1)}.$$

□

Corollary 28. *We have*

$$\sum_{k=0}^n (-1)^k \frac{2^{2k}}{2k+1} \frac{\binom{n}{k}}{\binom{2k}{k}} = \frac{1}{2n+1} \quad (64)$$

and

$$\sum_{k=0}^n (-1)^{k+1} 2^{2k} \frac{k}{2k+1} \frac{\binom{n}{k}}{\binom{2k}{k}} = \frac{2n}{(2n+1)(2n-1)}. \quad (65)$$

Proof. Set $r = s = 1/2$ in Theorem 26. Using (8) this choice gives

$$\sum_{k=0}^n (-1)^k 2^{2k} \frac{\binom{n+k}{n-k}}{2k+1} = \sum_{k=0}^n (-1)^{n-k} \frac{2^{2k}}{2(n-k)+1} \frac{\binom{n}{k} \binom{2n+1}{2k+1}}{\binom{2k}{k} \binom{2n+1}{2k}}. \quad (66)$$

The binomial coefficients can be simplified to

$$\frac{\binom{2n+1}{2k+1}}{\binom{2n+1}{2k}} = \frac{2(n-k)+1}{2k+1}.$$

The final step in the proof is to use

$$\sum_{k=0}^n (-1)^k 2^{2k} \frac{\binom{n+k}{n-k}}{2k+1} = \frac{(-1)^n}{2n+1},$$

which follows from

$$\sum_{k=0}^n (-1)^{n-k} 2^{2k} \frac{\binom{n+k}{n-k}}{2k+1} = \int_0^1 U_{2n}(t) dt = \frac{\sin(\pi n) + 1}{2n+1}.$$

This completes the proof of the first identity. The second follows as a combination with

$$\sum_{k=0}^n (-1)^{k+1} 2^{2k} \frac{\binom{n}{k}}{\binom{2k}{k}} = \frac{1}{2n-1}.$$

□

Corollary 29. *We have*

$$\sum_{k=0}^n (-1)^{n-k} \frac{2^{2k}}{2(n-k)+3} \frac{\binom{n+1}{k} \binom{2n+1}{2k+1}}{\binom{2k}{k} \binom{2n+3}{2k}} = \frac{(-1)^n (4n^2 + 4n - 1)}{(2n-1)(2n+1)(2n+3)}. \quad (67)$$

Proof. Set $r = s = 3/2$ in Theorem 26 and use (8) again. The left hand side can be evaluated as

$$\sum_{k=0}^n (-1)^{n-k} 2^{2k} \frac{\binom{n+k}{n-k}}{2k+3} = \int_0^1 t^2 U_{2n}(t) dt = \frac{4n^2 + 4n - 1}{(2n-1)(2n+1)(2n+3)}.$$

□

Corollary 30. *We have*

$$\sum_{k=0}^n (-1)^{n-k} k \frac{\binom{2n+1}{2k+1}}{\binom{n}{k}} = \begin{cases} 1, & \text{if } n = 1; \\ \frac{(-1)^n (2n+1)(2n-1) - (2n^2+1)}{4(n-1)n}, & \text{if } n \geq 2. \end{cases} \quad (68)$$

Proof. Set $r = 2$ in (62). After simplification of the binomial coefficients this results in the relation

$$\sum_{k=0}^n (-1)^{n-k} 2^{2k} \frac{\binom{n+k}{n-k}}{k+2} = \frac{1}{(n+2)(n+1)} \sum_{k=0}^n (-1)^{n-k} (n+1-k) \frac{\binom{2n+1}{2k+1}}{\binom{n}{k}}.$$

The left hand side equals

$$\sum_{k=0}^n (-1)^{n-k} 2^{2k} \frac{\binom{n+k}{n-k}}{k+2} = 2 \int_0^1 t^3 U_{2n}(t) dt = \frac{(2n+1)(2n(n+1)-3) + 3(-1)^n}{4(n-1)n(n+1)(n+2)}$$

and hence

$$\sum_{k=0}^n (-1)^{n-k} (n+1-k) \frac{\binom{2n+1}{2k+1}}{\binom{n}{k}} = \begin{cases} -5, & \text{if } n = 1; \\ \frac{(-1)^n (2n+1)(2n(n+1)-3) + 3}{4(n-1)n}, & \text{if } n \geq 2. \end{cases}$$

The final step is to split this sum into two parts, use the sum identity from (63) and simplify. □

Remark. Similar combinatorial sums were studied recently by Chu and Guo [18]. They have proved, among other things, the evaluations

$$\sum_{k=0}^n (-1)^k \frac{\binom{n+2}{2k+1}}{\binom{n}{k}} = 2 \quad \text{and} \quad \sum_{k=0}^n (-1)^k \frac{\binom{n+2}{2k}}{\binom{n}{k}} = -\frac{2}{n}.$$

Theorem 31. *If $s, r \in \mathbb{C} \setminus \mathbb{Z}^-$ such that $s \neq 0$ and $r - s \notin \mathbb{Z}^-$, then*

$$\begin{aligned} & \sum_{k=0}^n (-1)^{k+1} 2^{2k} \frac{\binom{n+k}{n-k}}{(k+s) \binom{k+r}{k+s}} H_{k+r} \\ &= \sum_{k=0}^n (-1)^{n-k} \frac{\binom{2n+1}{2k+1}}{(n-k+s) \binom{n+r}{n-k+s}} (H_{k+r-s} - H_{n+r} - H_{r-s}). \end{aligned} \quad (69)$$

Proof. Differentiate (61) with respect to r and simplify. \square

Corollary 32. *We have*

$$\sum_{k=0}^n (-1)^{k+1} 2^{2k} \frac{\binom{n+k}{n-k}}{k+r} H_{k+r} = \sum_{k=0}^n (-1)^{n-k} \frac{\binom{2n+1}{2k+1}}{(n-k+r) \binom{n+r}{k}} (H_k - H_{n+r})$$

and

$$\sum_{k=0}^n (-1)^{k+1} \frac{2^{2k+1}}{2k+1} \binom{n+k}{n-k} O_{k+1} = \frac{2(-1)^{n+1}}{2n+1} O_{n+1} + \sum_{k=0}^n (-1)^{n-k} \frac{2^{2k}}{2k+1} \frac{\binom{n}{k}}{\binom{2k}{k}} H_k.$$

Proof. Set $r = s$ in (69). The particular case follows from setting $r = \frac{1}{2}$, which yields with (8)

$$\sum_{k=0}^n (-1)^{k+1} 2^{2k} \frac{\binom{n+k}{n-k}}{2k+1} (2O_{k+1} - 2 \ln 2) = \sum_{k=0}^n (-1)^{n-k} \frac{2^{2k}}{2k+1} \frac{\binom{n}{k}}{\binom{2k}{k}} (H_k - 2O_{n+1} + 2 \ln 2).$$

We use (66) to remove logarithms. Finally, we use (64) to simplify the right-hand side and the result follows. \square

5 Additional results

In this section, we present an approach that is closely related to the general algorithm. First, we recall the following Lemma from [1], which shows that the standard form (18) has additional insights to offer.

Lemma 33 ([1, Corollary 3]). *Let an arbitrary polynomial identity have the following form:*

$$\sum_{k=s}^w f(k)(1+t)^k = \sum_{k=m}^r g(k)t^k, \quad (70)$$

where m, w, r and s are non-negative integers, $f(k)$ and $g(k)$ are sequences, and t is a complex variable. Let v be an arbitrary real number. Then

$$\sum_{k=s}^w \frac{f(k)}{2^k} \binom{2k+v}{(2k+v)/2} \binom{k+v}{v/2}^{-1} = \sum_{k=\lfloor (m+1)/2 \rfloor}^{\lfloor r/2 \rfloor} \frac{g(2k)}{2^{2k}} \binom{2k}{k} \binom{(2k+v)/2}{v/2}^{-1}, \quad (71)$$

and

$$\sum_{k=m}^r \frac{g(k)(-1)^k}{2^k} \binom{2k+v}{(2k+v)/2} \binom{k+v}{v/2}^{-1} = \sum_{k=\lfloor (s+1)/2 \rfloor}^{\lfloor w/2 \rfloor} \frac{f(2k)}{2^{2k}} \binom{2k}{k} \binom{(2k+v)/2}{v/2}^{-1}. \quad (72)$$

In particular,

$$\sum_{k=s}^w \frac{f(k)}{2^k} \binom{2k}{k} = \sum_{k=\lfloor (m+1)/2 \rfloor}^{\lfloor r/2 \rfloor} \frac{g(2k)}{2^{2k}} \binom{2k}{k}, \quad (73)$$

and

$$\sum_{k=m}^r \frac{g(k)(-1)^k}{2^k} \binom{2k}{k} = \sum_{k=\lfloor (s+1)/2 \rfloor}^{\lfloor w/2 \rfloor} \frac{f(2k)}{2^{2k}} \binom{2k}{k}. \quad (74)$$

Lemma 34 ([1, Theorem 8]). *Let the polynomial identity be given as in (70). Let u and v be arbitrary complex numbers such that $\Re u > -1$ and $\Re v > -1$. Then*

$$\begin{aligned} & \sum_{k=s}^n \frac{f(k)}{2^{2k}} \binom{v}{v/2} \binom{2k+u}{(2k+u)/2} \binom{(2k+u+v)/2}{v/2}^{-1} \\ &= \sum_{k=m}^r \frac{(-1)^k g(k)}{2^{2k}} \binom{u}{u/2} \binom{2k+v}{(2k+v)/2} \binom{(2k+u+v)/2}{u/2}^{-1}. \end{aligned} \quad (75)$$

In particular,

$$\sum_{k=s}^n \frac{f(k)}{2^{2k}} \binom{2k}{k} = \sum_{k=m}^r \frac{(-1)^k g(k)}{2^{2k}} \binom{2k}{k}. \quad (76)$$

We can now write more identities, for instance from (A), (C) and (D).

Identity (A) can be written in the standard form (70) with

$$f(k) = \frac{\delta_{nk}(H_k + 1) - 1}{n - k + \delta_{nk}}, \quad g(k) = \binom{n}{k} H_k, \quad w = n = r, \quad s = m = 0, \quad (77)$$

where here and throughout this paper δ_{ij} is Kronecker's delta symbol having the value 1 when $i = j$ and zero otherwise. Identity (C) can be written in the standard form (70) with

$$f(k) = \delta_{nk} H_n^2 - (1 - \delta_{nk}) \frac{(H_n - 2H_{n-k-1+\delta_{nk}} + H_k)}{n - k + \delta_{nk}}, \quad (78)$$

$$g(k) = \binom{n}{k} H_k^2, \quad (79)$$

We now formulate the result for (D). Similar results can be obtained for other, just mentioned identities.

Theorem 35. *We have*

$$\sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{2k} 2^{n-2k} \binom{2k}{k} H_{2k}^{(2)} = \binom{2n}{n} H_n^{(2)} - \sum_{k=0}^{n-1} \binom{2k}{k} 2^{n-k} \frac{H_n - H_k}{n - k}, \quad (80)$$

$$\sum_{k=0}^n (-1)^k \binom{n}{k} 2^{-k} \binom{2k}{k} H_k^{(2)} = - \sum_{k=0}^{\lfloor n/2 \rfloor - 1} \binom{2k}{k} 2^{-2k} \frac{H_n - H_{2k}}{n - 2k} + \begin{cases} 2^{-n} \binom{n}{n/2} H_n^{(2)}, & \text{if } n \text{ is even;} \\ -2^{-n+1} \binom{n-1}{(n-1)/2} \frac{1}{n}, & \text{if } n \text{ is odd;} \end{cases} \quad (81)$$

and more generally, if n is a non-negative integer and v is a real number, then

$$\begin{aligned} & \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{2k} 2^{n-2k} \binom{2k}{k} \binom{(2k+v)/2}{v/2}^{-1} H_{2k}^{(2)} \\ &= \binom{2n+v}{(2n+v)/2} \binom{n+v}{v/2}^{-1} H_n^{(2)} \\ & \quad - \sum_{k=0}^{n-1} \binom{2k+v}{(2k+v)/2} 2^{n-k} \binom{k+v}{v/2}^{-1} \frac{H_n - H_k}{n-k}, \end{aligned} \quad (82)$$

and

$$\begin{aligned} & \sum_{k=0}^n (-1)^k \binom{n}{k} 2^{-k} \binom{2k+v}{(2k+v)/2} \binom{k+v}{v/2}^{-1} H_k^{(2)} \\ &= - \sum_{k=0}^{\lfloor n/2 \rfloor - 1} \binom{2k}{k} 2^{-2k} \binom{(2k+v)/2}{v/2}^{-1} \frac{H_n - H_{2k}}{n-2k} \\ & \quad + \begin{cases} \binom{n}{n/2} 2^{-n} \binom{(n+v)/2}{v/2}^{-1} H_n^{(2)}, & \text{if } n \text{ is even;} \\ -\binom{n-1}{(n-1)/2} 2^{-n+1} \binom{(n-1+v)/2}{v/2}^{-1} \frac{1}{n}, & \text{if } n \text{ is odd.} \end{cases} \end{aligned} \quad (83)$$

Proof. Identities (82) and (83) now follow from (71) and (72). \square

Theorem 36. *We have*

$$\sum_{k=0}^n (-1)^k \binom{n}{k} 2^{-2k} \binom{2k}{k} H_k^{(2)} = \binom{2n}{n} 2^{-2n} H_n^{(2)} - \sum_{k=0}^{n-1} \frac{H_n - H_k}{n-k} \binom{2k}{k} 2^{-2k}, \quad (84)$$

and more generally, if n is a non-negative integer and u and v are complex numbers such that $\Re u > -1$ and $\Re v > -1$, then

$$\begin{aligned} & \sum_{k=0}^n (-1)^k \binom{n}{k} 2^{-2k} \binom{2k+v}{(2k+v)/2} \binom{(2k+u+v)/2}{u/2}^{-1} H_k^{(2)} \\ &= \binom{v}{v/2} \binom{u}{u/2}^{-1} \binom{2n+u}{(2n+u)/2} \binom{(2n+u+v)/2}{v/2}^{-1} 2^{-2n} H_n^{(2)} \\ & \quad - \binom{v}{v/2} \binom{u}{u/2}^{-1} \sum_{k=0}^{n-1} \frac{H_n - H_k}{n-k} \binom{2k+u}{(2k+u)/2} 2^{-2k} \binom{(2k+u+v)/2}{v/2}^{-1}. \end{aligned} \quad (85)$$

Proof. Use identity (75) with the parameters and variables given in (77). Particular cases follow after setting $u = v$ and then replacing u by $2u$ and then setting $u = 0$. \square

6 Conclusion

In this article, we have introduced a general framework (algorithm) to study combinatorial sums associated with certain classes of polynomial identities. The sums involve products of binomial coefficients and also harmonic numbers. To underline its usefulness we have shown applications based on some prominent polynomial identities from the recent past. This is only the starting point and many new families of sums can be studied. We invite the readers to apply the framework to sums associated with

- the identity of Simons [28]

$$\sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} t^k = \sum_{k=0}^n (-1)^{n+k} \binom{n}{k} \binom{n+k}{k} (1+t)^k$$

- the Narayana polynomials [26, 27]

$$\sum_{k=0}^n \frac{1}{n} \binom{n}{k-1} \binom{n}{k} t^k = \frac{1}{n+1} \sum_{k=0}^n (-1)^k \binom{n+1}{k} \binom{2n-k}{n} (1-t)^k$$

- an identity involving central binomial coefficients [23]

$$\sum_{k=0}^n 2^{2(n-k)} \binom{n}{k} \binom{2k}{k} t^k = \sum_{k=0}^n \binom{2k}{k} \binom{2(n-k)}{n-k} (1+t)^k$$

and many more.

References

- [1] K. Adegoke, A short proof of Knuth's old sum, preprint, 2024, <https://www.preprints.org/manuscript/202411.1249/v3>
- [2] K. Adegoke, R. Frontczak and T. Goy, Binomial Fibonacci sums from Chebyshev polynomials, *J. Integer Seq.*, **26** (2023), Article 23.9.6.
- [3] K. Adegoke and R. Frontczak, Combinatorial identities with multiple harmonic-like numbers, *AppliedMath*, **4** (2024), 986–998.
- [4] K. Adegoke and R. Frontczak, Some notes on an identity of Frisch, *Open J. Math. Sci.*, **8** (2024), 216–226.

- [5] N. Batir, On some combinatorial identities and harmonic sums, *Int. J. Number Theory*, **13** (2017), 1695–1709.
- [6] N. Batir, Combinatorial identities involving harmonic numbers, *Integers*, **20** (2020), #A25.
- [7] N. Batir, Finite binomial sum identities with harmonic numbers, *J. Integer Seq.*, **24** (2021), Article 21.4.3.
- [8] N. Batir and A. Sofo, A unified treatment of certain classes of combinatorial identities, *J. Integer Seq.*, **24** (2021), Article 21.3.2.
- [9] K. N. Boyadzhiev, Harmonic number identities via Euler’s transform, *J. Integer Seq.*, **12** (2009), Article 09.6.1.
- [10] K. N. Boyadzhiev, Series transformation formulas of Euler type, Hadamard product of series, and harmonic number identities, *Indian J. Pure Appl. Math.* **42** (2011), 371–386.
- [11] K. N. Boyadzhiev, Binomial transform and the backward difference, *Adv. Appl. Discrete Math.*, **18** (2014), 43–63.
- [12] J. Choi, Certain summation formulas involving harmonic numbers and generalized harmonic numbers, *Appl. Math. Comput.* **218** (2011), 734–740.
- [13] J. Choi, Finite summation formulas involving binomial coefficients, harmonic numbers and generalized harmonic numbers, *J. Inequal. Appl.* (2013), Article 49, 11 pages.
- [14] J. Choi and H. M. Srivastava, Some summation formulas involving harmonic numbers and generalized harmonic numbers, *Math. Comp. Mod.* **54** (2011), 2220–2234.
- [15] W. Chu, Partial-fraction decompositions and harmonic number identities, *J. Combin. Math. Combin. Comput.* **60** (2007), 139–153.
- [16] W. Chu, Summation formulae involving harmonic numbers, *Filomat* **26** (1) (2012), 143–152.
- [17] W. Chu and A. M. Fu, Dougall-Dixon formula and harmonic number identities, *Ramanujan J.* **18** (2009), 11–31.
- [18] W. Chu and D. Guo, Alternating sums of binomial quotients, *Math. Commun.* **27**, (2022), 203–213.
- [19] G. Dattoli, S. Licciardi and R. M. Pidotella, Inverse derivative operator and umbral methods for the harmonic numbers and telescopic series study, *Symmetry* **13**, 2021, 781.
- [20] R. Frontczak, Harmonic sums via Euler’s transform: Complementing the approach of Boyadzhiev, *J. Integer Seq.*, **23** (2020), Article 20.3.2.

- [21] R. Frontczak, Binomial sums with skew-harmonic numbers, *Palest. J. Math.*, **10** (2021), 756–763.
- [22] M. Genčev, Binomial sums involving harmonic numbers, *Math. Slovaca*, **61** (2011), 215–226.
- [23] H. W. Gould, *Combinatorial Identities, A Standardized Set of Tables Listing 500 Binomial Coefficient Summations, revised edition*. Published by the author, Morgantown, WV, 1972.
- [24] H. T. Jin and D. K. Du, Abel’s lemma and identities on harmonic numbers, *Integers*, **15** (2015), #A22.
- [25] D. E. Knuth, *The Art of Computer Programming, Volume 1, Fundamental Algorithms, Third Edition*, 1997.
- [26] T. Mansour and Y. Sun, Dyck paths and partial Bell polynomials, *Austr. J. Combin.* **42** (2008), 285–297.
- [27] T. Mansour and Y. Sun, Identities involving Narayana polynomials and Catalan numbers, *Discrete Math.* **309** (2009), 4079–4088.
- [28] S. Simons, A curious identity, *Math. Gaz.* **85** (2001), 296–298.
- [29] A. Sofo and H. M. Srivastava, Identities for the harmonic numbers and binomial coefficients, *Ramanujan J.* **25** (2011), 93–113.
- [30] H. M. Srivastava and J. Choi, *Series Associated with the Zeta and Related Functions*, Springer Science+Media, B.V., 2001.
- [31] W. Wang, Riordan arrays and harmonic number identities, *Comput. Math. Appl.* **60** (2010), 1494–1509.
- [32] J. Wang and C. Wei, Derivative operator and summation formulae involving generalized harmonic numbers, *J. Math. Anal. Appl.* 434 (1) (2016), 315–341.
- [33] C. Wei, D. Gong and Q. Yan, Telescoping method, derivative operators and harmonic number identities, *Integral Transforms Spec. Funct.* 25 (3) (2014), 203–214.
- [34] C. Wei and X. Wang, Summation formulas involving generalized harmonic numbers, *J. Difference Equ. Appl.*, **22** (2016), 1554–1567.
- [35] C. Xu, Identities for the shifted harmonic numbers and binomial coefficients, *Filomat* **31:19** (2017), 6071–6086.