

# Elementary Proof of the Completeness of OEIS A051221 Below 2000

Seichi Azuma

seazuma1@gmail.com

## Abstract

The OEIS sequence A051221 consists of nonnegative integers of the form  $10^x - y^2$ . The known values are those less than or equal to 2000 with  $x \leq 7$ , and it is conjectured that no new values in this range appear for  $x \geq 8$ . In this paper, we give an elementary proof that this is indeed the case by using Pell-type equations.

## 1 Introduction

The sequence A051221 in the OEIS<sup>1</sup> is defined as the set of all nonnegative integers that can be expressed in the form  $10^x - y^2$  for integers  $x \geq 0$  and  $y \geq 0$ . Currently, the sequence consists of

$$0, 1, 6, 9, 10, 19, 36, 39, \dots, 1999.$$

The current list was obtained by computing the values in the set

$$T(X) = \bigcup_{x=1}^X \{10^x - y^2 \mid y \in \mathbb{Z}\} \cap [0, 2000],$$

and stopping when  $T(X+1) = T(X)$ . Since this equality holds at  $X = 7$ , the currently known elements of the sequence correspond to values of  $x \leq 7$ . However, this does not immediately imply that new values cannot appear for  $x \geq 9$ . Intuitively, it is unlikely that a square is very close to a power of 10, but proving this rigorously is nontrivial. In this paper, we present an elementary proof that  $T(X) = T(7)$  for all  $X \geq 7$ , thereby confirming that the known list is complete within the range  $[0, 2000]$ .

Generally, equation  $a^x - y^2 = c$  is known as a generalized Lebesgue-Ramanujan-Nagell equation. A comprehensive survey by Maohua Le and Gökhan Soydan [1] summarizes the known results, including the finiteness of integer solutions  $(x, y)$  for fixed  $a$  and  $c$ . For specific cases, Christian Hercher and Karl Fegert provided an elementary proof using Pell-type equations for expressions of this form [2]. Our method follows a similar method.

---

<sup>1</sup><https://oeis.org/A051221>

## 2 Reformulation and Strategy

Note that when  $x$  is even, the smallest possible positive value of the form  $10^x - y^2$  is

$$10^x - (10^{x/2} - 1)^2 = 2 \cdot 10^{x/2} - 1,$$

which exceeds 2000 whenever  $x/2 \geq 4$ , that is,  $x \geq 8$ . Therefore, it is sufficient to show that no additional values of the form  $10^x - y^2$  can appear for odd  $x \geq 9$ .

We fix an integer  $c \in [0, 2000]$  that is not known to be in the sequence. Let  $x = 2u + 1$ . Then the equation becomes

$$10 \cdot (10^u)^2 - y^2 = c.$$

This is known as Pell-type equation and can be analyzed via quadratic fields. As shown in the next section, any solution  $10t^2 - s^2 = c$  can be expressed with integers  $a, b, K$  satisfying:

$$s + t\sqrt{10} = (a + b\sqrt{10})(19 + 6\sqrt{10})^K, \quad |a| \leq 3\sqrt{c}, \quad \sqrt{c/10} \leq |b| \leq \sqrt{c}, \quad 10b^2 - a^2 = c.$$

For each  $c$ , the pair  $(a, b)$  that satisfies the condition is finite, so our approach is to show that  $t$  cannot be the form of  $10^u$  for any integer  $k$  for each  $(a, b)$ . We consider the sequence of integers  $t_k$  defined by

$$(a + b\sqrt{10})(19 + 6\sqrt{10})^k = s_k + t_k\sqrt{10},$$

and show the impossibility of  $t_k = 10^u$  for any  $k$  and  $u \geq 4$  by showing that  $t_k \equiv 0 \pmod{10^4}$  and  $t_k \equiv 10^m \pmod{p}$  do not hold simultaneously for any  $m$  for some  $p$ . Note that negating  $a, b$  results in negating  $s_k, t_k$  and that negating  $b, k$  corresponds to taking the conjugate, which negates  $t_k$ . Hence, it is sufficient to consider only nonnegative  $a$  and  $b$ , and to prove the impossibility of  $t_k = \pm 10^u$ .

## 3 Main Lemma

The key step in our argument is the following lemma.

**Lemma 3.1.** *Let  $c$  be a fixed positive integer. Suppose that there exist integers  $s, t$  that satisfy the equation*

$$10t^2 - s^2 = c.$$

*Then there exist integers  $a, b, K$  that satisfy*

$$s + t\sqrt{10} = (a + b\sqrt{10})(19 + 6\sqrt{10})^K, \quad 0 \leq |a| \leq 3\sqrt{c}, \quad \sqrt{\frac{c}{10}} \leq |b| \leq \sqrt{c}, \quad 10b^2 - a^2 = c.$$

*Proof.* We first assume  $s + t\sqrt{10} > 0$ . We define  $K$  as

$$K = \left\lfloor \frac{\log(s + t\sqrt{10}) - \log \sqrt{c}}{\log(19 + 6\sqrt{10})} + \frac{1}{2} \right\rfloor.$$

This implies

$$(19 + 6\sqrt{10})^{k-1/2} \leq \frac{s + t\sqrt{10}}{\sqrt{c}} \leq (19 + 6\sqrt{10})^{k+1/2}.$$

We then define  $a, b$  by

$$a + b\sqrt{10} = \frac{s + t\sqrt{10}}{(19 + 6\sqrt{10})^k}.$$

Since  $19 + 6\sqrt{10}$  is a unit in  $\mathbb{Z}[\sqrt{10}]$ ,  $a$  and  $b$  are integers, and we have

$$10b^2 - a^2 = 10t^2 - s^2 = c.$$

Note that  $\sqrt{19 + 6\sqrt{10}} = 3 + \sqrt{10}$ , so the inequality above implies

$$\sqrt{c}(-3 + \sqrt{10}) \leq a + b\sqrt{10} \leq \sqrt{c}(3 + \sqrt{10}).$$

Now consider the set of real numbers of the form  $a + b\sqrt{10}$  satisfying this inequality. These values correspond to the points  $(a, b)$  in the hyperbola

$$10b^2 - a^2 = c$$

within a bounded region. By analyzing the endpoints, we have

$$-3\sqrt{c} \leq a \leq 3\sqrt{c}, \quad \sqrt{c/10} \leq b \leq \sqrt{c},$$

since the smallest value of  $b$  occurs when  $a = 0$ , which gives  $10b^2 = c$ .

When  $s + t\sqrt{10} < 0$ , we apply the same discussion to  $-s - t\sqrt{10}$  and conclude similarly that

$$-3\sqrt{c} \leq a \leq 3\sqrt{c}, \quad \sqrt{c/10} \leq -b \leq \sqrt{c}.$$

□

## 4 Example

Let  $c = 31$ , which is not in the known list of values. Then, the only integer pair  $(a, b) = (3, 2)$  satisfies the conditions

$$0 \leq a \leq 3\sqrt{c}, \quad \sqrt{\frac{c}{10}} \leq b \leq \sqrt{c}, \quad 10b^2 - a^2 = c.$$

Define the sequence of integers  $t_k$  by

$$(a + b\sqrt{10})(19 + 6\sqrt{10})^k = s_k + t_k\sqrt{10}.$$

The sequence  $\{t_k\}$  satisfies the recurrence:

$$t_0 = b = 2, \quad t_1 = 6a + 19b = 56, \quad t_{k+2} = 38t_{k+1} - t_k.$$

Let  $N = 10^4$  and  $p = 160001$ . The sequences  $(t_k \bmod N)$  and  $(t_k \bmod p)$  are linear recurrence sequences modulo fixed moduli and, therefore, eventually periodic. In this case,

these sequences have a common period 40000, and we find that  $t_k \equiv 0 \pmod{N}$  occurs precisely when  $k \equiv 3309 \pmod{5000}$ . By computing  $t_k \pmod{p}$  at those values of  $k$ , we have

$$t_k \equiv 4354, 121626, 16949, 146265, 155647, 38439, 143052, 13736 \pmod{p}.$$

We also observe that the set of residues  $\{\pm 10^m \pmod{p}\}$  forms a multiplicative subgroup of order 1250 in  $\mathbb{F}_p^\times$ , and none of the eight residues above belongs to this subgroup.

Therefore, we conclude that  $t_k = \pm 10^u$  does not occur for any  $k$  and  $u \geq 4$ , which confirms that  $c = 31$  cannot appear in the form  $10^x - y^2$  for any  $x \geq 9$ .

## 5 Whole Computation

We now describe the complete computation that verifies the completeness of the sequence A051221 within the interval  $[0, 2000]$ .

For every integer  $c \in [0, 2000]$  that is not contained in the known list of the form  $10^x - y^2$  for  $x \leq 7$ , we enumerate all integer pairs  $(a, b)$  satisfying the following conditions:

$$0 \leq a \leq 3\sqrt{c}, \quad \sqrt{\frac{c}{10}} \leq b \leq \sqrt{c}, \quad 10b^2 - a^2 = c.$$

For each valid pair  $(a, b)$ , we define the sequence  $t_k$  by

$$t_0 = b, \quad t_1 = 6a + 19b, \quad t_{k+2} = 38t_{k+1} - t_k.$$

We compute  $t_k \pmod{N}$  and  $t_k \pmod{p}$  for

$$N = 10^4, \quad p = 160001,$$

and verify whether  $t_k \equiv 0 \pmod{N}$  and  $t_k \equiv \pm 10^m \pmod{p}$  can hold simultaneously as in the previous example.

This method successfully excludes most cases except for the following five cases:

$$(a, b) = (22, 8), (38, 16), (68, 24), (67, 24), (3, 12).$$

We handle these remaining cases taking  $p = 1601$  instead of  $p = 160001$ , where the multiplicative subgroup  $\{\pm 10^m \pmod{p}\}$  has size 200. For each of these five cases, we confirm that none of the  $t_k$  values congruent with 0 (mod  $N$ ) modulo  $p$  belong to the subgroup  $\{\pm 10^m \pmod{p}\}$ , thus eliminating all possibilities. This completes the proof that no new values of the form  $10^x - y^2$  appear in the interval  $[0, 2000]$  for  $x \geq 9$ . All computations were performed using Google Colab, and the corresponding notebook is available at:

[https://colab.research.google.com/drive/1c77020BIBM1\\_UMuH4MmnhKy0POM9LBC-?usp=sharing](https://colab.research.google.com/drive/1c77020BIBM1_UMuH4MmnhKy0POM9LBC-?usp=sharing)

## References

- [1] M. Le and G. Soydan, *A brief survey on the generalized Lebesgue–Ramanujan–Nagell equation*, *Surv. Math. Appl.* **15** (2020), 473–523. ISSN 1842-6298 (electronic), 1843-7265 (print).
- [2] C. Hercher and K. Fegert, *Triangular numbers with a single repeated digit*, *J. Integer Seq.* **28** (2025), Article 25.2.1.