
Can Diffusion Models Disentangle? A Theoretical Perspective

Liming Wang¹ Muhammad Jehanzeb Mirza¹
 Yishu Gong² Yuan Gong³ Jiaqi Zhang⁴ Brian H. Tracey² Katerina Placek² Marco Vilela² James R. Glass¹

Abstract

This paper presents a novel theoretical framework for understanding how diffusion models can learn disentangled representations. Within this framework, we establish identifiability conditions for general disentangled latent variable models, analyze training dynamics, and derive sample complexity bounds for disentangled latent subspace models. To validate our theory, we conduct disentanglement experiments across diverse tasks and modalities, including subspace recovery in latent subspace Gaussian mixture models, image colorization, image denoising, and voice conversion for speech classification. Additionally, our experiments show that training strategies inspired by our theory, such as style guidance regularization, consistently enhance disentanglement performance.

1. Introduction

Diffusion models (DMs) (Sohl-Dickstein et al., 2015; Song & Ermon, 2019; Ho et al., 2020) are generative models capable of approximating probability distributions by learning noisy versions of their scores (Fisher, 1935). While such approaches enjoy both empirical successes (e.g., (Ramesh et al., 2022)) and theoretical guarantees (Chen et al., 2023b; Pabbaraju et al., 2023), they tend to represent the latent structure of the underlying distribution *implicitly*. However, in learning tasks such as controllable generation, it is useful to represent the task-specific latent structure *explicitly* in the generative model to reflect the inductive biases of the problems. One approach, known as *conditional diffusion models* (CDMs), achieves this goal with DMs by labeling such variables and conditioning the model on these labels (Wu et al., 2023a;

Yang et al., 2023; Hudson et al., 2024). However, it remains unclear whether and when CDMs can learn an explicit representation that captures the conditional dependency relations between the variables, especially when some of them are unlabeled. For example, if the latent factors are (approximately) independent, it is desirable to have a *disentangled* representation with decomposable parts for each variable. Learning such a representation is called *disentanglement*. An intriguing theoretical question then arises: *what are the fundamental limits for CDMs to learn disentangled representations?* A theory capable of answering this question can potentially lead to more powerful, compositional generative models for a wider range of applications in self-supervised representation learning, causal inference and computational creativity.

To answer the question, we focus on one canonical example of the disentanglement problem — content-style disentanglement. The choice is justified on three grounds. First, the task involves a simple latent variable model but captures the essence of the disentanglement problem. Second, it is practically useful, as certain data, such as those from under-represented minorities and subjects, have no data labeled with both content and style to explicitly disentangle factors in the data. Given only unpaired data with style labels for training, a two-variable disentanglement system tries to change the style of the observed data, without modifying the content during inference. Such a problem is at the core of applications such as voice conversion (VC) and image editing with *non-parallel* data. To generate samples with new combinations of content and style, the model has to learn a representation that disentangles the content variable from the style during training. Lastly, DM-based disentanglement recently revolutionized various fields such as VC (e.g., (Popov et al., 2022; Choi et al., 2023; Seed Team, 2024)) and image editing (e.g., (Wu et al., 2023b; Yang et al., 2023)) and provided new opportunities for generating realistic synthetic data for classification tasks. However it is not fully understood whether and how such models perform disentanglement and improve downstream performance of classification systems.

This paper makes three key contributions:

1. **Theoretical identifiability:** We prove that a diffusion model (DM) trained with an information-regularized score-matching objective can identify disentangled

¹Computer Science and Artificial Intelligence Laboratory, Massachusetts Institute of Technology, USA ²Takeda Development Center Americas, USA ³xAI, USA ⁴Broad Institute, Massachusetts Institute of Technology, USA. Correspondence to: Liming Wang <limingw@csail.mit.edu>.

latent variables for general two-variable latent variable models, provided the score functions satisfy mild Lipschitz continuity conditions (Section 3.1).

2. **Novel training objective:** We propose and analyze a style-guided score-matching objective, specifically designed for DM disentanglement using gradient-based methods (Section 3.3). For a class of probability distributions known as latent subspace models, our theoretical analysis establishes the optimality of this objective and provides insights into its sample complexity as a function of the sample dimension, noise schedules, and Lipschitz properties of the score functions.
3. **Empirical validation:** We validate our theory through extensive experiments on synthetic and real-world disentanglement tasks (Section 4), including subspace recovery for Gaussian mixture models, image colorization, image denoising, and voice conversion for speech classification. Our results demonstrate the practical utility of our theory, showing that training strategies inspired by it — such as data augmentation and style guidance regularization — consistently improve performance across all tasks.

2. Disentanglement

In a two-variable disentanglement problem, a learner is given a random observation (sample) $X \sim p$ with support $\mathcal{X} \subseteq \mathbb{R}^{d_X}$, which is determined by two underlying random variables: the *content* Z with support $\mathcal{Z} \subseteq \mathbb{R}^{d_Z}$, and the *style* G with support $\mathcal{G} \subseteq \mathbb{R}^{d_G}$. The generative process is modeled as:

$$X := \alpha \bar{X} + N := \alpha \bar{\psi}(Z, G) + N, \quad (1)$$

where $\alpha := \sqrt{1 - \delta^2}$ and $N \sim \mathcal{N}(0, \delta^2)$ represents independent Gaussian white noise with a small deviation $\delta > 0$. Here, N is independent of Z and G . For example, in a voice conversion task, the content variable $Z \sim p_Z$ represents the semantic information of the speech, while the style variable $G \sim p_G$ encodes speaker identity, accent and emotional characteristics. To make the problem well-posed, we assume that the style variable G is *observable* by the learner. Notably, the definitions of content and style variables are symmetric, differing only their observability. Consequently, the framework is equally applicable when the content variable Z is observable instead. We constrain N to be Gaussian because our primary interest lies in the limiting case as $\delta \rightarrow 0$. This limiting scenario cannot be analyzed directly due to the divergence of certain information-theoretic quantities.

Further, we make the following independence assumptions.

Assumption 2.1. The generative process in Equation 1 possesses the following statistical properties:

1. *Disentanglement:* $Z \perp\!\!\!\perp G$;

2. *Conditional disentanglement:* $Z - X - G$ forms a Markov chain.

3. *Injectivity:* There are injective mappings $z: \mathcal{X} \mapsto \mathcal{Z}$ and $g: \mathcal{X} \mapsto \mathcal{G}$ such that

$$Z = z(\bar{X}), G = g(\bar{X});$$

4. *Invertibility:* There exists a mapping $\psi(\cdot, \cdot)$ such that $X = \psi(z(X), g(X))$.

The task of *disentanglement* is then to recover Z given X and G . To evaluate the success of disentanglement, we use the following definitions.

Definition 2.2. (ϵ -disentanglement) Two variables X and Y are ϵ -disentangled if there exists $\epsilon > 0$ such that the mutual information $I(X; Y) := \mathbb{E}_{p_{XY}(x, y)} \log \frac{p_{XY}(x, y)}{p_X(x)p_Y(y)} \leq \epsilon$.

For applications such as voice conversion and image editing, we are also interested in the notion of (ϵ, ϕ, p) -editability.

Definition 2.3. (ϵ, ϕ, p) -editability) Given a function ϕ and a probability measure p , two random variables X and Y are (ϵ, ϕ, p) -*editable* if given two of their independent, identically distributed (i.i.d) copies (X_1, Y_1) and (X_2, Y_2) , and let $Z_1 := \phi(X_1, Y_1)$ and $Z_{12} = \phi(X_1, Y_2)$,

1. $d_{\text{TV}}(p_{Z_1}, p) \leq \epsilon$;
2. $d_{\text{TV}}(p_{Z_{12}}, p_{Z_1}) \leq \epsilon$.

where $d_{\text{TV}}(p, q) := \frac{1}{2} \int_{\mathcal{X}} |p(x) - q(x)| dx$ is the total variation distance between p and q . Further, X and Y are simply (ϵ, ϕ) -*editable* if $p \equiv p_{Z_1}$.

For instance, the content Z and style G defined by Eq. 1 and Assumption 2.1 are (ϵ, ψ) -*editable* by definition, since for any i.i.d copies (Z_1, G_1) and (Z_2, G_2) , $\bar{X}' := \psi(Z_1, G_2)$ satisfies $p_{\bar{X}'}(x) = p_Z(z(x))p_G(g(x)) = p_{\bar{X}}(x), \forall x$. This notion is also useful for analyzing content distortion, particularly in verifying whether samples X can be generated by modifying style G while preserving content Z : Given a content variable Z and a learned generative model $\hat{\psi}$, if Z and G are $(\epsilon, \hat{\psi}, p_{X|Z})$ -*editable*, then it is possible to generate samples X from an ϵ -closed conditional distribution of $p_{X|Z}$ with preserved content Z and modified style G .

3. Diffusion model-based disentanglement

3.1. Unsupervised disentanglement and editability

One approach to disentanglement is to learn the generative process in Eq. 1 using *diffusion models* (DM) (Song & Ermon, 2019), as illustrated in Figure 1. The training of a DM has two main stages: noising and denoising. In the *noising* stage,




Figure 1. Graphical model of the diffusion-based disentanglement process. Solid circles represent observed variables, dashed circles are latent variables, and square boxes denote learnable components. In the forward process, independent latent content Z and observed style G generate a clean sample \bar{X} , which is then corrupted by Brownian motion. In the reverse process, the corrupted observation X_T is restored using a learnable score network s_θ , conditioned on the style G and a learnable bottleneck variable \hat{Z} .

the model progressively corrupts the input data by adding noise according to a stochastic differential equation (SDE):

$$dX_t = f(X_t, t)dt + \nu(t)dB_t, X_0 \sim p, \quad (2)$$

for some *parameter-free* functions f and ν and a truncated Brownian motion $\{B_t\}_{t \in [0, T]}$. A common choice sets $f(X_t, t) := -X_t$ and $\nu(t) \equiv \sqrt{2}$, yielding the Ornstein–Uhlenbeck (OU) process

$$dX_t = -X_t dt + \sqrt{2}dB_t, X_0 \sim p, \quad (3)$$

whose stationary distribution is standard Gaussian and which exhibits favorable convergence properties (Chen et al., 2023b;a). Our framework readily extends to alternate choices of (f, ν) corresponding to different noise schedulers.

Let p_t denote the distribution of X_t and $p_{t|s}$ the conditional distribution of X_t given X_s . Define the time-reversed process as $X_t^\leftarrow := X_{T-t}$. In the *denoising* stage, the model learns to reconstruct the observation X from its corrupted versions X_t^\leftarrow 's at some time $t \geq t_0$, where $t_0 > 0$ is a small threshold. This is achieved by simulating the *reverse process* of Eq. 3, conditioned on the style variable G and a *bottleneck variable* $\hat{Z} = z_\phi(X_{t_0})$, where z_ϕ is a learnable transformation with parameterized by ϕ :

$$dX_t^\leftarrow = [X_t^\leftarrow + 2\nabla_x \log p_{T-t}(X_t^\leftarrow | \hat{Z}, G)]dt + \sqrt{2}dB_t^\leftarrow, \quad X_0^\leftarrow \sim p_T. \quad (4)$$

Classical theory in SDE ensures that the reverse process Eq. 4 has p as its stationary distribution (Anderson, 1982). To simulate this reverse process, the learner needs to estimate the *gradient* of the conditional probability distribution

$p_t(X_t | \hat{Z}, G)$ with respect to X_t for all X_t . To achieve this, the learner employs a *parameterized score function* $s_\theta : \mathcal{X} \times \mathcal{Z} \times \mathcal{G} \times [0, T] \mapsto \mathbb{R}^{d_x}$ and trains it to approximate the true score function via a learning objective known as *score matching*:

$$L(\theta, \phi) := \mathbb{E}_{t, p_t} \|s_\theta(X_t, z_\phi(X), G, t) - \nabla_x \log p_t(X_t | z_\phi(X), G)\|^2, \quad (5)$$

where the time distribution is chosen as $p(t) \equiv \frac{\mathbb{1}_{t_1 \leq t \leq T}}{T - t_1}$ for some $t_1 > t_0$. While the true score function $\nabla_x \log p_t(X_t | \hat{Z}, G)$ is not directly accessible, this objective is equivalent to the following *conditional* score matching objective based on the gradient $\nabla_x \log p_{t|0}(X_t | X)$'s:

$$\begin{aligned} L_c(\theta, \phi) &:= \mathbb{E}_{t, p_t} \|s_\theta(X_t, z_\phi(X_{t_0}), G, t) - \nabla_x \log p_{t|0}(X_t | X)\|^2 \\ &= \mathbb{E}_{t, p_t} \left\| s_\theta(X_t, z_\phi(X_{t_0}), G, t) + \frac{N_t}{\sigma(t)} \right\|^2, \end{aligned} \quad (6)$$

where N_t is a standard Gaussian vector independent of X and the variance function is given by $\sigma(t) := \sqrt{1 - \exp(-2t)}$. To achieve DM-based disentanglement, we introduce a *regularized* score matching objective with hyperparameters $\gamma, \rho > 0$ and the rectified linear unit (ReLU) function $(x)_+ := \max\{x, 0\}$:

$$L_c^{\gamma, \rho}(\theta, \phi) := L_c(\theta, \phi) + \gamma(I(z_\phi(X_{t_0}); X) - \rho)_+. \quad (7)$$

During inference, DM generates new samples \hat{X}_t^\leftarrow by simulating the *estimated reverse process*:

$$\begin{aligned} d\hat{X}_t^\leftarrow &= [\hat{X}_t^\leftarrow + 2s_\theta(\hat{X}_{k\eta}^\leftarrow, \hat{Z}, G, T - k\eta)]dt + \sqrt{2}dB_t^\leftarrow, \\ \hat{X}_0^\leftarrow &\sim p_T, t \in [k\eta, (k+1)\eta]. \end{aligned} \quad (8)$$

Define $\hat{X}_t := \hat{X}_{T-t}^\leftarrow$, $\hat{X} := \hat{X}_{t_0}$ and introduce the parameterized *maximum likelihood (ML) estimator* ψ_θ as

$$\psi_\theta(\hat{Z}, G) := \mathbb{E}[\hat{X} | \hat{Z}, G]. \quad (9)$$

A key distinction of DM-based disentanglement from autoencoder-based approaches (Qian et al., 2019) is the involvement of multiple bottleneck variables: the time-dependent, task independent X_t 's and the time-dependent, task-dependent \hat{Z} . We argue that both are essential for the disentanglement to work *in general*, except in special cases discussed later. First, the presence of X_t is crucial because, without it, the score function cannot approximate the gradient effectively due to the absence of a time-dependent variable. On the other hand, without \hat{Z} , the sequence X_t 's alone generally fails to capture the content information of X . In fact, as $T \rightarrow \infty$, the mutual information between X and X_T , conditioned on G , vanishes:

$$I(X; X_T | G) \propto -D_{\text{KL}}(p_{T|0}(\cdot | G) || \mathcal{N}(0, I_{d_x})) \rightarrow 0,$$

by the property of the OU process (Chen et al., 2023b). As a result, even with a well-trained score function $s_\theta(X_t, \hat{Z}, G, t) = s_\theta(X_t, G, t) \approx \nabla \log p_t(\cdot|G)$, the mutual information between the samples X' from the reverse process and X degrades, following the data processing inequality,

$$I(X'; X|G) \leq I(X_T; X|G) \xrightarrow{T \rightarrow \infty} 0. \quad (10)$$

However, when both X_t 's and \hat{Z} are present, and \hat{Z} minimizes the regularized score matching objective Eq. 7, we prove that \hat{Z} and G are ϵ -disentangled, where ϵ depends on the quality of the score function s_θ and the mutual information between the bottleneck \hat{Z} and X , $I(\hat{Z}; X_t)$. To this end, we make the following assumptions.

Assumption 3.1. The observation X is sub-gaussian with second moment: $\mathbb{E}\|X\|^2 =: \sigma_X^2 d_X$.

Assumption 3.2. The score functions for $p_{\hat{X}}(\cdot)$, $p_{Z|X}(\cdot|x)$ and $p_{G|X}(\cdot|x)$ are λ_s -Lipschitz for any $x \in \mathcal{X}$:

$$\begin{aligned} \|\nabla_x \log p_{\hat{X}}(x') - \nabla_x \log p_{\hat{X}}(x)\| &\leq \lambda_s \|x' - x\|, \\ \|\nabla_z \log p_{Z|X}(z'|x) - \nabla_z \log p_{Z|X}(z|x)\| &\leq \lambda_s \|z' - z\|, \\ \|\nabla_g \log p_{G|X}(g'|x) - \nabla_g \log p_{G|X}(g|x)\| &\leq \lambda_s \|g' - g\|, \\ \forall (x, x', z, z', g, g') &\in \mathcal{X}^2 \times \mathcal{Z}^2 \times \mathcal{G}^2. \end{aligned}$$

Assumption 3.3. The content function $z(\cdot)$ is λ_z -Lipschitz and the style function $g(\cdot)$ is λ_g -Lipschitz.

Assumption 3.4. The function $\psi(\cdot, \cdot)$ is λ_ψ -Lipschitz in both arguments with $\lambda_\psi(\lambda_z + \lambda_g) = 1 + O(\delta)$.

Note that Assumption 3.1 and Assumption 3.2 are analogous to those in (Chen et al., 2023b). Our result relies crucially on the following lemma.

Lemma 3.5. *There exists (θ_1, ϕ_1) such that for*

$$\begin{aligned} \delta &= O\left(\frac{\epsilon^2}{\lambda_s^2 \lambda_z^2 \sigma_X^2 d_X^4}\right) < \min\left\{\frac{1}{2}, \frac{1}{\sqrt{d_X}}\right\}, \\ t_0 &= -\log(1 - \delta^2)^{1/2}, t_1 = -\log(1 - \delta)^{1/2}, \end{aligned}$$

the followings hold:

1. *The mutual information between \hat{Z} and X obeys*

$$I(z_{\phi_1}(X_{t_0}); X) = I(Z; X) + O\left(\frac{\epsilon^2}{\lambda_s \lambda_z \sigma_X d_X^3}\right);$$
2. *The conditional score matching loss satisfies for some*

$$C_2 = 1 + O(\delta),$$

$$\begin{aligned} L_c(\theta_1, \phi_1) &\leq \frac{C_2 \delta^2 d_X (e^{2T} - e^{2t_1})}{2(T - t_1)(e^{2t_1} - 1)(e^{2T} - 1)} \\ &= O\left(\frac{\epsilon^2}{\lambda_s^2 \lambda_z^2 \sigma_X^2 d_X^3 T}\right). \end{aligned}$$

Using Lemma 3.5, we prove the following theorem.

Theorem 3.6. *Suppose Assumption 2.1-3.4 hold, and (θ^*, ϕ^*) be a minimizer of $L_c^{\gamma, \rho}$ defined in Eq. 7 with*

$$\rho = I(Z; X) + O(\epsilon^2), \gamma = \frac{C\epsilon}{2T}, T = \Omega(\log(1/\epsilon))$$

for some constant $C > 0$ and δ, t_0, t_1 as in Lemma 3.5. Then the bottleneck $z_{\phi^*}(X_{t_0})$ and the style G are $O(\epsilon)$ -disentangled.

We prove that \hat{Z} and G are $(\epsilon, \psi_{\theta^*}, p)$ -editable for some optimal θ^* under additional assumptions in Appendix C.

Theorem 3.7. *Under Assumption 2.1-3.2 and Assumption C.1-C.2, let δ, t_0 and (θ^*, ϕ^*) be chosen as in Theorem 3.6, and set the diffusion step size $\eta := O\left(\frac{\epsilon}{\lambda_\theta^2 d_X T} + \frac{\sqrt{\epsilon}}{\lambda_\theta \sigma_X \sqrt{T}}\right)$, the bottleneck $\hat{Z} := z_{\phi^*}(X_{t_0})$ and the style G are $(O(\sqrt{\epsilon}), \psi_{\theta^*}, p)$ -editable.*

3.2. Disentanglement with Data Augmentation

A key challenge with an unsupervised disentangled model is the issue of *content distortion*: While the generated samples from the ML estimator $\hat{\psi}$ preserve the overall distribution of the original data, they do not necessarily retain the *content* of individual samples. For content preservation, the content variable Z needs to be predictive of the bottleneck variable \hat{Z} independently of the style G . The condition implies that the conditional mutual information $I(\hat{Z}; G|Z) \approx 0$. However, disentanglement alone only enforces $I(\hat{Z}; G) \approx 0$, which does not necessarily imply conditional independence from G given Z . As result, when \hat{Z} and G are $(\epsilon, \hat{\psi}, p)$ -editable, $\hat{Z}|Z$ and G are not necessarily $(\epsilon, \hat{\psi}, p_{X|Z})$ -editable. A concrete example illustrating this phenomenon is included in Appendix D. Previous study (Qian et al., 2019) appears to overlook this issue, likely due to the empirical success of their methods. This suggests that commonly used datasets, such as speech and images, may possess inherent inductive biases that facilitate disentanglement, warranting further theoretical investigation.

To mitigate this issue, one potential approach is to incorporate *data augmentation*. In particular, suppose we can leverage an inductive bias to generate a synthetic sample of the form $X^c := \alpha \bar{\psi}(Z, G^c) + N'$ such that G and $G^c|Z$ are ϵ -disentangled and N' and N are independent for some $\epsilon > 0$. Given such synthetic samples, we can enhance training by introducing the following *mix-and-match loss*:

$$\mathcal{L}_{\text{mm}}(\theta, \phi) = \mathbb{E}_{t, p_t} \left\| s_\theta(X_t, \hat{z}_\phi(X_{t_0}^c), G) + \frac{N_t}{\sigma(t)} \right\|^2, \quad (11)$$

where $X_{t_0}^c := X^c + N_{t_0}$. By optimizing Eq. 11, we can show that $\hat{Z}|Z$ and G are $O(\epsilon_1)$ -disentangled and $(O(\epsilon_2), \hat{\psi}, p_{X|Z})$ -editable for some arbitrarily small $\epsilon_1, \epsilon_2 > 0$.

Theorem 3.8. *Under Assumption 2.1-C.2, suppose $I(G; G^c|Z) \leq \epsilon$, and let δ, t_0, t_1 be chosen as in Theorem 3.6,*




Figure 2. **The dual-encoder score network for disentangling content and style of latent subspace models.** The upper pathway processes the corrupted observation X_t to compute a content score function $s_Z(X_t, t)$, while the lower pathway processes the style G_t to compute a style score function $s_G(G_t, t)$. The two scores are combined along with the positional encoding of the time step t to produce the final score function $s_\theta(X_t, t)$.

η be chosen as in Theorem 3.7 and (θ^*, ϕ^*) be the optimal solution to Eq. 11, then $z_{\phi^*}(X_{t_0}^c)$ and G are $O(\epsilon)$ -disentangled and $(O(\sqrt{\epsilon}), \psi_{\theta^*}, p_{X|Z})$ -editable.

3.3. Disentanglement with latent subspace model

One class of latent variable models that avoids content distortion issues is the *latent subspace model* (LSM). This approach has been widely used in classical disentanglement models in speech processing, such as the *i-vector* model (Dehak et al., 2011) and has been shown to approximately hold for various self-supervised learning representations (Liu et al., 2023). These representations have been applied to disentanglement tasks in recent works (Baas et al., 2023; Choi et al., 2023; Huang et al., 2024).

Definition 3.9. A latent subspace model samples via

$$Z \sim p_Z, G \sim \alpha, X = A_Z Z + A_G G, \quad (12)$$

where $A_Z \in \mathbb{R}^{d_X \times d_Z}$, $A_G \in \mathbb{R}^{d_X \times d_G}$ are orthogonal matrices and the subspaces for the content and speaker are orthogonal and span the whole space, i.e., $R(A_Z)^\perp = R(A_G)$ with $d_Z + d_G = d_X$, where $R(A)$ is the column space of matrix A . Further, let X_t be the noisy feature variable at time t of the diffusion process and define $Z_t := A_Z^\top X_t =: z(X_t)$, $G_t := A_G^\top X_t =: g(X_t)$.

A useful property of LSM is that its score is *decomposable*:

$$\nabla_x \log p_t(X_t) = A_Z \nabla_z \log p_{Z_t}(z(X_t)) + A_G \nabla_g \log p_{G_t}(g(X_t)). \quad (13)$$

This is a generalization of the subspace model used in (Chen et al., 2023a), where they assume $p_{G_t} = \mathcal{N}(\mathbf{0}_{d_G}, I_{d_G})$. Another useful property of LSM is that the distance between the estimated and true subspaces captures the level of disentanglement, since for $\hat{Z} := P_U X$, $U^\top A_G = 0$ implies $I(\hat{Z}; G) = 0$. Further, disentanglement (and similarly editability) for (\hat{Z}, G) automatically implies disentanglement

(editability) for $(\hat{Z}|Z, G)$, since if $U^\top A_G = 0$,

$$I(\hat{Z}; G|Z) = I(P_U N_{t_0} + P_U G; G) = I(P_U N_{t_0}; G) = 0.$$

Therefore, LSM does not suffer from content distortion.

For LSM, we prove that the model is able to learn a disentangled representation *without* conditioned on functions of X , i.e., $s_\theta(X_t, \hat{Z}(X), G, t) = s'_\theta(X_t, G, t)$. To this end, we consider the following *regularized* score matching loss:

$$L^{\lambda_r}(s_Z, s_G, U, V) := 2L_0(s_Z, s_G, U, V) + \frac{1}{2}L_b(s_Z, s_G, U, V) + \lambda_r L_r(s_Z, U), \quad (14)$$

where

$$\begin{aligned} L_0 &:= \mathbb{E}_{t, p_t(x)} \|\tilde{U} \tilde{s}(x, t) - \nabla_x \log p_t(x)\|_2^2 \\ L_b &:= \mathbb{E}_t \|\tilde{U}^\top \tilde{U} - \mathbb{E}_{p_t(x)} \tilde{s}(x, t)^\top \tilde{s}(x, t)\|_2^2 \\ L_r &:= 2\mathbb{E}_{t, p_t(x)} \|V s_G(g(x), t) - \nabla_x \log p(x)\|_2^2 + \\ &\quad \frac{1}{2} \mathbb{E}_t \|V^\top V - \mathbb{E}_{p_t(x)} s_G(g(x), t) s_G(g(x), t)^\top\|_2^2, \end{aligned}$$

where $\tilde{U} := [U, V]$, $\tilde{s}(x, t) := [s_Z(x, t)^\top, s_G(x, t)^\top]^\top$ and $\lambda_r > 0$ is the *style guidance weight*. L_r is the *style guidance regularization* loss and can be viewed as a way to limit the mutual information between the bottleneck variable and the observation $I(\hat{Z}; X)$, by encouraging maximal contribution from the style variable G in the score matching process. As shown in Figure 2, the estimated score functions along the content and style subspaces, s_Z and s_G respectively are parameterized by two *two-layer ReLU neural networks*:

$$s_Z(x, t) := \frac{1}{\sqrt{d_H}} \sum_{j=1}^{d_H} \theta_Z^{(2), j} (\theta_Z^{(1), j})^\top [x^\top, \text{PE}(t)]^\top, \quad (15)$$

$$s_G(g, t) := \frac{1}{\sqrt{d_H}} \sum_{j=1}^{d_H} \theta_G^{(2), j} (\theta_G^{(1), j})^\top [g^\top, \text{PE}(t)]^\top. \quad (16)$$

with parameters $\theta_Z := (\theta_Z^{(1)}, \theta_Z^{(2)}) \in \mathbb{R}^{(d_X + d_T) \times d_H} \times \mathbb{R}^{d_Z \times d_H}$ and $\theta_G := (\theta_G^{(1)}, \theta_G^{(2)}) \in \mathbb{R}^{(d_G + d_T) \times d_H} \times \mathbb{R}^{d_G \times d_H}$. $\text{PE} : \mathbb{R}_+ \mapsto \mathbb{R}^{d_T}$ denotes the *position encoding* of the time variable, and L_b is the balancing loss commonly used in the low-rank matrix factorization literature (e.g., (Ge et al., 2017)). The inclusion of PE is not redundant: our theoretical analysis indicates that convergence of the gradient-based method requires PE of different time steps to be incoherent. This property is approximately satisfied by sinusoidal-based PE. Further, we demonstrate that the LSM objective Eq. 14 recovers the true style and speaker subspaces, validating its effectiveness in disentanglement tasks.

Theorem 3.10. *For the linear subspace model 3.9 and the objective in Eq. 14, then any minimizer (U^*, V^*) of Eq. 14 satisfy $R(U^*) = R(A_Z)$ and $R(V^*) = R(A_G)$.*

Further, we analyze the training dynamics of gradient-based methods for LSM disentanglement in the *infinite-width* regime as $d_T, d_H \rightarrow \infty$. For the gradient flow equations

$$[\dot{U}, \dot{V}] = [-\nabla_U L^{\lambda_r}(s_Z, s_G, U, V), -\nabla_V L^{\lambda_r}(s_Z, s_G, U, V)] \quad (17)$$

$$[\dot{\theta}_Z, \dot{\theta}_G] = [-\nabla_{\theta_Z} L^{\lambda_r}(s_Z, s_G, U, V), -\nabla_{\theta_G} L^{\lambda_r}(s_Z, s_G, U, V)], \quad (18)$$

Our analysis leads to the following theorem.

Theorem 3.11. *Suppose $\min\{d_T, d_H\} \rightarrow \infty$, and the neural network weights are initialized by standard Gaussians. Further, choose $\text{PE}(\cdot)$ such that $\text{PE}(t)$'s are bounded and linearly independent for all $t \in [t_0, T]$. Then for $\lambda_r = 3$, the system of gradient flow equations in Eq. 17-18 converges to a critical point (\hat{U}, \hat{V}) such that $R(\hat{U}) = R(A_Z), R(\hat{V}) = R(A_G)$.*

Further, our framework can be extended to the finite-sample regime. Consider the following empirical objective:

$$\hat{L}^{\lambda_r}(\tilde{s}, \tilde{U}) = 2\hat{L}_0(\tilde{s}, \tilde{U}) + \frac{1}{2}\hat{L}_b(\tilde{s}, \tilde{U}) + \lambda_r \hat{L}_r(s_Z, U), \quad (19)$$

where $s^*(x, t) := \nabla_x \log p_t(x)$ and

$$\begin{aligned} \hat{L}_0 &:= \frac{1}{n} \sum_{i=1}^n \|\tilde{U} \tilde{s}(x^i, t^i) - s^*(x^i, t^i)\|_2^2, \\ \hat{L}_b &:= \left\| \tilde{U}^\top \tilde{U} - \frac{1}{n} \sum_{i=1}^n \tilde{s}(x^i, t^i) \tilde{s}(x^i, t^i)^\top \right\|, \\ \hat{L}_r &:= \frac{2}{n} \sum_{i=1}^n \|V s_G(g(x^i), t^i) - \nabla_x \log p_{t^i}(x^i)\|_2^2 + \\ &\quad \frac{1}{2} \left\| V^\top V - \frac{1}{n} \sum_{i=1}^n s_G(g(x^i), t^i) s_G(g(x^i), t^i)^\top \right\|^2. \end{aligned}$$

By solving the gradient flow equations Eq. 17-18 with \hat{L} in place of \hat{L} , our analysis yields the following sample complexity bound.

Theorem 3.12. *Under Assumption 3.1-3.2, let $\min\{d_T, d_H\} \rightarrow \infty$, and the neural network weights are initialized by standard Gaussians. Further, choose $\text{PE}(\cdot)$ such that $\text{PE}(t)$'s are bounded and linearly independent for all $t \in [t_0, T]$. Then for $\lambda_r = 3$, the empirical version of the system of gradient flow equations in Eq. 17-18 converges to a critical point $(\hat{s}_Z, \hat{s}_G, \hat{U}, \hat{V})$ such that with probability at least $1 - O(\frac{1}{n})$, the following holds:*

1. *The population objective*

$$L_0(\hat{s}_Z, \hat{s}_G, \hat{U}, \hat{V}) = O\left(\sqrt{\frac{d_X^5 \log^3 n}{n}}\right),$$

2. *The content are style subspace are recovered as*

$$\begin{aligned} \|P_{\hat{U}} - A_Z A_Z^\top\|_F^2 &= O\left(\frac{1}{\sigma_{d_Z}^2(s_Z^*)} \sqrt{\frac{d_X^5 \log^3 n}{n}}\right), \\ \|P_{\hat{V}} - A_G A_G^\top\|_F^2 &= O\left(\frac{1}{\sigma_{d_G}^2(s_G^*)} \sqrt{\frac{d_X^5 \log^3 n}{n}}\right), \end{aligned}$$

where $\sigma_i(s)$ is the i -th largest singular value of the operator s .

3. *The bottleneck \hat{Z} and style G are $O\left(\frac{d_X^{5/4} \log^{3/4} n}{\sigma_{d_Z}(s_Z^*) n^{1/4}}\right)$ -disentangled;*

4. *Set $t_0 := \frac{1}{2} \log \frac{n^{1/8}}{n^{1/8} - \log^{3/2} n} > 0$, and let $\hat{Z}_{t_0} := \hat{Z} + \sigma(t_0) N_{t_0}$, then $\hat{Z}_{t_0}|Z$ and $A_G G_{t_0}$ are $(\epsilon', \hat{\psi}, p)$ -editable, with $\hat{\psi}(z, g) := \hat{z} + P_{\hat{V}} g$ for sufficiently large n , and $\epsilon' := O\left(\frac{d_X^{5/8} \log^{3/8} n}{\min\{\sigma_{d_Z}^{1/2}(s_Z^*), \sigma_{d_G}^{1/2}(s_G^*)\} n^{1/32}}\right)$.*

While our bounds depend on the total dimension d_X , our model can be generalized to represent content and style variables in a lower-dimensional subspace. This can be achieved by incorporating a residual connection in the score network in Figure 2. Further, our analysis primarily focuses on the unconditional score matching objective. However, it can be readily extended to the conditional score matching objective by accounting for the finite-sample noise introduced by the cross terms $\langle s_\theta(x_t, t), \nabla_{x_t} \log \frac{p_t(x_t)}{p_{t_0}(x_t|x)} \rangle$. Finally, while our analysis is centered on infinite-width, two-layer neural networks, the methodology can potentially be extended to finite-width and deeper networks as in (Du et al., 2019).

4. Experiments

This section presents the empirical evaluation of our theory. First, we conduct disentanglement experiments on synthetic datasets generated by Gaussian mixture models (GMM). Next, to demonstrate the applicability of our theory to real-world scenarios, we perform disentanglement tasks such as colorization and denoising on two standard image datasets, MNIST (Deng, 2012) and CIFAR10 (Krizhevsky et al., 2009). Finally, to assess the generalizability of our theory across modalities, we tackle the voice conversion adaptation task, a canonical disentanglement task, to enhance speech emotion classification on the IEMOCAP (Busso et al., 2008) dataset.

Implementation details: For the GMM dataset, we train a wide, two-layer MLP to align with our theoretical framework. For image and speech data, we employ score networks based on U-Nets (Ronneberger et al., 2015), widely adopted for

DMs (Song et al., 2021). For images, the model is trained with the following regularized score-matching objective:

$$L_c^{\lambda_r}(\theta, \phi) := \mathbb{E}_{t, p_t} \left\| s_\theta(X_t, \hat{Z}, G, t) + \frac{N_t}{\sigma(t)} \right\|^2 + \lambda_r \mathbb{E}_{t, p_t} \left\| s_\theta(X_t, \mathbf{0}_{d_Z}, G, t) + \frac{N_t}{\sigma(t)} \right\|^2, \quad (20)$$

where the regularizer is inspired by the regularizer L_r used in the LSM objective Eq. 14 and thus we refer to λ_r also as the style guidance weight and the weighted loss as the style guidance loss. For speech data, we utilize an existing DM-based VC approach (Popov et al., 2022) in a plug-and-play manner. More implementation details on the model architecture, training, and inference can be found in Appendix I.




Figure 3. **LSGMM disentanglement results with a two-layer multilayer perceptron (MLP) score function.** The subspace recovery error is normalized between $[0, 1]$. In four random trials for each noise scheduler, DM consistently recovers (error < 0.1) the correct content subspace and achieves disentanglement with sufficiently large λ_r and sample size.

Latent subspace GMM disentanglement: First, we conduct subspace recovery experiments on *latent subspace GMMs* (LSGMM), a class of LSMs where each subspace follows a GMM. Further, Figure 3 shows the subspace recovery error as a function of the style guidance weight λ_r and the sample size n . Consistent with the predictions of Theorem 3.11, the LSGMM achieves the smallest subspace reconstruction error when the style guidance weight is sufficiently large, and the result is consistent across different noise schedulers. Moreover, since all score networks are wide, two-layer MLPs trained using gradient-based methods, these results provide empirical support for Theorem 3.11. Additionally, Figure 3 reveals a sublinear decay rate of the subspace recovery error as the sample size increases, aligning with the prediction of Theorem 3.12.

Image disentanglement: Next, we evaluate our theoretical findings on image data from MNIST and CIFAR-10. For MNIST, we perform the task of *colorization*, where the goal is to transform MNIST images into RGB while preserving the digit shape and applying a specified background color. This is a disentanglement problem where the digit shape is the

content and the background color is the style. The results are visualized in Figure 4. As shown in Figure 4b, without any regularization, the model fails to achieve disentanglement and instead copies the input without any meaningful colorization. In contrast, as demonstrated in Figure 4c, incorporating the regularizer enables the model to effectively separate color from shape, allowing for successful colorization.

For CIFAR10, we conduct an image denoising experiment, which is a disentanglement task with the clean image as the content and the independent noise as the style. In this setup, we introduce a random color shift as the noise, though our approach can, in principle, be extended to other types of independent noise. As shown in Figure 5, the model exhibits the same trend observed in the MNIST experiment, demonstrating the effectiveness of the style guidance loss, as predicted by Theorem 3.12. Quantitative results for both datasets using different regularization weights in four different metrics can be found in Appendix I.2.

Speech disentanglement: To further assess the practical utility of our theoretical framework, we apply DM-based disentanglement to speech data, focusing on *voice conversation adaptation* (VCA). The style variable corresponds to the speaker identity and the content represents a target characteristic of the speech signal, such as the emotion state and health condition of the speaker. The task of VCA is then to learn a disentangled representation to improve speech classification tasks under distribution shift in the style variable. Such a task serves as a probe into the speech representation to measure its degree of disentanglement, as more disentangled representation tends to be more robust against domain shift in style.

We focus our attention on VCA for speech emotion recognition on IEMOCAP, though the same approach has shown beneficial effect for speech biomarker impairment classification such as Alzheimer detection (ADReSS (Luz et al., 2020)) and Amyotrophic Lateral Sclerosis (ALS) severity classification (ALS-TDI (Vieira et al., 2022)), as included in Appendix I.3.

Figure 6 depicts the classification performance as a function of the number of target speakers used to perform speaker-emotion disentanglement on IEMOCAP. For each target speaker number, we randomly select 4 speaker combinations. We compare classifiers based on DM-based disentangled representation (Diff-VC (Popov et al., 2022)) with: 1) classifiers without using any disentangled representation (No VC); 2) those using other disentanglement methods such as pitch shifting, KNN-VC (Baas et al., 2023) and TriAAN-VC (Park et al., 2023). We found that Diff-VC performs the best, reaching an average of 97.2% accuracy, which is 25.7% better than the no-VC classifier and 36.1% than the pitch shifting adaptation. This result demonstrates that DM can indeed achieve approximate disentanglement for speech data, as predicted by Theorem 3.6, and that approximate disentanglement can benefit speech classification tasks. We also observe




Figure 4. **Disentanglement results on MNIST.** The content is the gray-scale image and the style is the background color of the image. The task is to colorize a gray-scale image without observing paired gray-scale and colored images during training. The training objective is Eq. 19 without the balancing loss and with the LSM score functions replaced by the score function forms in the general case, where the style-only score function is simply the same score function with the content embedding masked out. 4b and 4c suggest that successful disentanglement is achieved with a sufficiently large style guidance weight $\lambda_r = 3$. More examples can be found in Appendix I.2.




Figure 5. **Disentanglement results on CIFAR10.** The content is the clean image and the style is the random corruption on the image. The task is to restore the clean image without observing clean images during training. As shown in 5b and 5c, disentanglement is achieved with a sufficiently large style guidance weight $\lambda_r = 3$. More examples can be found in Appendix I.2.




Figure 6. **Speech emotion recognition results on IEMOCAP as a probing task for speech disentanglement.** DM-based disentanglement between the content (emotion) and style (speaker) is shown to be superior in disentanglement than other methods. Further, augmentation techniques such as using multiple speaker further improve disentanglement performance.

a steady trend of improvement in classification accuracy as the number of target speakers increases. One interpretation of this result is that having multiple speakers serves as a form of data augmentation, which improves disentanglement by reducing content distortion, confirming Theorem 3.8.

5. Related works

Disentangled representation learning: The concept of disentanglement we adopt is first defined explicitly as a generalization of statistical independence (Tishby et al., 1999) based on mutual information, though other definitions exist, e.g., (Higgins et al., 2018). Disentanglement is a crucial concept for deep learning in fields such as representation learning (Bengio et al., 2013; Schmidhuber, 1992; Tschanen et al., 2018), voice conversion (Qian et al., 2019; Wang et al., 2021a; Popov et al., 2022) and image editing (Wu et al., 2023b; Yang et al., 2023; Motamed et al., 2024). Neural network-based architectures have been proposed to learn disentangled representation (Hsu et al., 2017; Higgins et al., 2017; Kim & Mnih, 2018; Chen et al., 2016; Wu et al., 2023a; Yang et al., 2023; Hudson et al., 2024) among others, though theoretical understanding of such models remain limited. To understand the learnability of disentangled representation, (Locatello et al., 2019) proved a no-free-lunch theorem on disentanglement inspired by classical results in independent component analysis (Comon, 1994). Motivated by the task of VC, (Qian et al., 2019) proves that for the content-style latent variable model, disentanglement is indeed possible when the style variable is observed, a result our theory extends to DM-based approaches and noisy, continuous content and style variables.

Diffusion model theory: Early works on DMs focus on their ability to learn general data distributions, under different assumptions on statistical properties of the data distribution such as log-Sobolev inequality (Lee et al., 2022), and bounded moments (Block et al., 2020; Chen et al., 2023b) and score approximation accuracy in terms of L^∞ -accuracy (Bortoli et al., 2021) and L^2 -accuracy (Lee et al., 2022; Chen et al., 2023b). Others attempt to understand the benefit of DM over maximum-likelihood-based generative models (Pabbaraju et al., 2023). More recent works have started to analyze the ability of DM to learn latent low-dimensional subspace (Chen et al., 2023a) and manifold structure (Bortoli et al., 2022). Further, (Fu et al., 2024) studies the convergence properties of CDMs for a variety of latent variable learning tasks and the role of classifier-free guidance in such tasks.

6. Conclusion

In this work, we present a theoretical framework to understand the capacity of diffusion models to disentangle latent variables. By addressing key theoretical questions regarding identifiability, training dynamics, and sample complexity, our framework provides a comprehensive foundation for diffusion model-based disentanglement. We validate our theory through synthetic experiments and evaluations on image and speech datasets, demonstrating strong empirical support.

Impact Statement

This paper presents work whose goal is to advance the field of Machine Learning. There are many potential societal consequences of our work, none which we feel must be specifically highlighted here.

References

- Anderson, B. D. Reverse-time diffusion equation models. *Stochastic Processes and their Applications*, 12(3):313–326, 1982. ISSN 0304-4149. doi: 10.1016/0304-4149(82)90051-5. URL <https://www.sciencedirect.com/science/article/pii/0304414982900515>.
- Baas, M., van Niekerk, B., and Kamper, H. Voice conversion with just nearest neighbors. In *Interspeech*, 2023.
- Baevski, A., Zhou, H., Mohamed, A., and Auli, M. wav2vec 2.0: A framework for self-supervised learning of speech representations. In *Neural Information Processing System*, 2020.
- Barron, A. R. Universal approximation bounds for superpositions of a sigmoidal function. *IEEE Transactions on Information Theory*, 39(3):930–945, 1993. doi: 10.1109/18.256500.
- Bengio, Y., Courville, A., and Vincent, P. Representation learning: A review and new perspectives. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 2013. URL <https://arxiv.org/pdf/1206.5538>.
- Block, A., Mroueh, Y., and Rakhlin, A. Generative modeling with denoising auto-encoders and langevin sampling. *arXiv preprint arXiv:2002.00107*, 2020. URL <https://arxiv.org/abs/2002.00107>.
- Bortoli, V. D., Thornton, J., Heng, J., and Doucet, A. Diffusion schrödinger bridge with applications to score-based generative modeling. In Ranzato, M., Beygelzimer, A., Dauphin, Y., Liang, P., and Vaughan, J. W. (eds.), *Neural Information Processing System*, volume 34, pp. 17695–17709. Curran Associates, Inc., 2021.
- Bortoli, V. D., Thornton, A., Hutchinson, J. H., Mathieu, J. A., Teh, Y., and Doucet, A. Convergence of denoising diffusion models under the manifold hypothesis. *Transactions on Machine Learning Research (TMLR)*, 2022. URL <https://openreview.net/forum?id=MhK5aXo3gB>.
- Busso, C., Bulut, M., Lee, C.-C., Kazemzadeh, A., Mower, E., Kim, S., Chang, J.-N., Lee, S., and Narayanan, S. Iemocap: Interactive emotional dyadic motion capture database. In *Language Resources and Evaluation (LREC)*, pp. 335–339, 2008.
- Caron, M., Mairal, J., Touvron, H., Misra, I., Bojanowski, P., Jégou, H., and Joulin, A. Emerging properties in self-supervised vision transformers. In *Proceedings of the IEEE/CVF International Conference on Computer Vision (ICCV)*, pp. 9650–9660, 2021. URL <https://arxiv.org/abs/2104.14294>.
- Chae, M. and Walker, S. G. Wasserstein upper bounds of the total variation for smooth densities. *Statistics & Probability Letters*, 163:108771, 2020. ISSN 0167-7152. doi: 10.1016/j.spl.2020.108771. URL <https://www.sciencedirect.com/science/article/pii/S0167715220300742>.
- Chen, M., Huang, K., Zhao, T., and Wang, M. Score approximation, estimation and distribution recovery of diffusion models on low-dimensional data. In *International Conference on Machine Learning*, 2023a. URL <https://openreview.net/pdf?id=KB4mLiouEX>.
- Chen, S., Chewi, S., Li, J., Li, Y., Salim, A., and Zhang, A. Sampling is as easy as learning the score: theory for diffusion models with minimal data assumptions. In *The Eleventh International Conference on Learning Representations*, 2023b. URL https://openreview.net/forum?id=zyLVMgsZOU_.

- Chen, X., Duan, Y., Houthoofd, R., Schulman, J., Sutskever, I., and Abbeel, P. Infogan: Interpretable representation learning by information maximizing generative adversarial nets. In *Neural Information Processing System*, NIPS'16, pp. 2180–2188, Red Hook, NY, USA, 2016. Curran Associates Inc. URL <https://arxiv.org/abs/1606.03657>.
- Choi, H.-Y., Lee, S.-H., and Lee, S.-W. Diff-HierVC: Diffusion-based Hierarchical Voice Conversion with Robust Pitch Generation and Masked Prior for Zero-shot Speaker Adaptation. In *Proc. INTERSPEECH 2023*, pp. 2283–2287, 2023. doi: 10.21437/Interspeech.2023-817.
- Comon, P. Independent component analysis, a new concept? *Signal processing*, 36(3):287–314, 1994. URL <https://core.ac.uk/download/pdf/52776748.pdf>.
- Dehak, N., Kenny, P. J., Dehak, R., Dumouchel, P., and Ouellet, P. Front-end factor analysis for speaker verification. In *IEEE Transactions on Audio, Speech, and Language Processing*, volume 19, pp. 788–798. IEEE, 2011. doi: 10.1109/TASL.2010.2064307.
- Deng, L. The mnist database of handwritten digit images for machine learning research. *IEEE Signal Processing Magazine*, 29(6):141–142, 2012.
- Du, S., Lee, J., Li, H., Wang, L., and Zhai, X. Gradient descent finds global minima of deep neural networks. In Chaudhuri, K. and Salakhutdinov, R. (eds.), *International Conference on Machine Learning*, volume 97 of *Proceedings of Machine Learning Research*, pp. 1675–1685. PMLR, 09–15 Jun 2019. URL <https://proceedings.mlr.press/v97/du19c.html>.
- Fisher, R. A. The detection of linkage with ‘dominant’ abnormalities. *Annals of Eugenics*, 6(2):187–201, 1935.
- Fu, H., Yang, Z., Wang, M., and Chen, M. Unveil conditional diffusion models with classifier-free guidance: A sharp statistical theory. *ArXiv*, abs/2403.11968, 2024. URL <https://api.semanticscholar.org/CorpusID:268532529>.
- Ge, R., Jin, C., and Zheng, Y. No spurious local minima in nonconvex low rank problems: A unified geometric analysis. In Precup, D. and Teh, Y. W. (eds.), *Proceedings of the 34th International Conference on Machine Learning*, volume 70 of *Proceedings of Machine Learning Research*, pp. 1233–1242. PMLR, 06–11 Aug 2017. URL <https://proceedings.mlr.press/v70/ge17a.html>.
- Girsanov, I. V. On transforming a certain class of stochastic processes by absolutely continuous substitution of measures. *Theory of Probability & Its Applications*, 5(3): 285–301, 1960. doi: 10.1137/1105034.
- Hajek, B. and Raginsky, M. ECE 543: Statistical Learning Theory. <http://maxim.ece.illinois.edu/teaching/SLT/>, 2021. Accessed: 2025-01-25.
- Higgins, I., Matthey, L., Pal, A., Burgess, C., Glorot, X., Botvinick, M., Mohamed, S., and Lerchner, A. beta-VAE: Learning basic visual concepts with a constrained variational framework. In *International Conference on Learning Representations*, 2017. URL <https://openreview.net/forum?id=Sy2fzU9gl>.
- Higgins, I., Amos, D., Pfau, D., Racaniere, S., Matthey, L., Rezende, D., and Lerchner, A. Towards a definition of disentangled representations. *arXiv preprint arXiv:1812.02230*, 2018. URL <https://arxiv.org/pdf/1812.02230>.
- Ho, J., Jain, A., and Abbeel, P. Denoising diffusion probabilistic models. In *Neural Information Processing System*, volume 33, pp. 6840–6851, 2020.
- Hsu, W.-N., Zhang, Y., and Glass, J. Unsupervised learning of disentangled and interpretable representations from sequential data. In *Advances in Neural Information Processing Systems*, pp. 1878–1889, 2017. URL <https://arxiv.org/abs/1709.07902>.
- Huang, F., Zeng, K., and Zhu, W. DiffVC+: Improving diffusion-based voice conversion for speaker anonymization. In *Interspeech*, 2024.
- Hudson, D. A., Zoran, D., Malinowski, M., Lampinen, A. K., Jaegle, A., McClelland, J. L., Matthey, L., Hill, F., and Lerchner, A. Soda: Bottleneck diffusion models for representation learning. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR)*, pp. to appear, June 2024. URL https://openaccess.thecvf.com/content/CVPR2024/papers/Hudson_SODA_Bottleneck_Diffusion_Models_for_Representation_Learning_CVPR_2024_paper.pdf.
- Jacot, A., Gabriel, F., and Hongler, C. Neural tangent kernel: Convergence and generalization in neural networks. In *Neural Information Processing System*, 2018.
- Kim, H. and Mnih, A. Disentangling by factorising. In Dy, J. and Krause, A. (eds.), *International Conference on Machine Learning*, volume 80 of *Proceedings of Machine Learning Research*, pp. 2649–2658. PMLR, 10–15 Jul 2018. URL <https://proceedings.mlr.press/v80/kim18b.html>.
- Kingma, D. P. and Ba, J. Adam: A method for stochastic optimization. In *International Conference on Learning Representations*, 2015. URL <https://arxiv.org/abs/1412.6980>.

- Krizhevsky, A., Nair, V., and Hinton, G. Learning multiple layers of features from tiny images. <http://www.cs.toronto.edu/~kriz/cifar.html>, 2009. Accessed: 2025-01-25.
- Lee, H., Lu, J., and Tan, Y. Convergence for score-based generative modeling with polynomial complexity. In Oh, A. H., Agarwal, A., Belgrave, D., and Cho, K. (eds.), *Neural Information Processing System*, 2022.
- Lee, J. D., Simchowitz, M., Jordan, M. I., and Recht, B. Gradient descent only converges to minimizers. In *Conference on Learning Theory*, 2016.
- Liu, O. D., Tang, H., and Goldwater, S. Self-supervised predictive coding models encode speaker and phonetic information in orthogonal subspaces. In *Interspeech*, 2023.
- Locatello, F., Bauer, S., Lucic, M., Raetsch, G., Gelly, S., Schölkopf, B., and Bachem, O. Challenging common assumptions in the unsupervised learning of disentangled representations. In Chaudhuri, K. and Salakhutdinov, R. (eds.), *Proceedings of the 36th International Conference on Machine Learning*, volume 97 of *Proceedings of Machine Learning Research*, pp. 4114–4124. PMLR, Jun 2019. URL <http://proceedings.mlr.press/v97/locatello19a.html>.
- Luz, S., Haider, F., de la Fuente, S., and Davida Fromm, B. M. Alzheimer’s dementia recognition through spontaneous speech: The adress challenge. In *Proceedings of Interspeech 2020*, pp. 2172–2176, 2020.
- Ma, Z., Chen, M., Zhang, H., Zheng, Z., Chen, W., Li, X., Ye, J., Chen, X., and Hain, T. Emobox: Multilingual multi-corpus speech emotion recognition toolkit and benchmark. In *Interspeech*, 2024.
- Motamed, S., Paudel, D. P., and Gool, L. V. Lego: Learning to disentangle and invert personalized concepts beyond object appearance in text-to-image diffusion models. In *Proceedings of the European Conference on Computer Vision (ECCV)*, 2024. doi: 10.1007/978-3-031-72633-0_7. URL https://www.ecva.net/papers/eccv_2024/papers_ECCV/papers/02309.pdf.
- Nichol, A. and Dhariwal, P. Improved denoising diffusion probabilistic models. In *International Conference on Machine Learning*, 2021.
- Pabbaraju, C., Rohatgi, D., Sevekari, A., Lee, H., Moitra, A., and Risteski, A. Provable benefits of score matching. In *Neural Information Processing System*, 2023. URL <https://openreview.net/forum?id=waXoG35kbb>.
- Park, H. J., Yang, S. W., Kim, J. S., Shin, W., and Han, S. W. Triaan-vc: Triple adaptive attention normalization for any-to-any voice conversion. In *ICASSP*, 2023.
- Paszke, A., Gross, S., Massa, F., Lerer, A., Bradbury, J., Gregory, T., Lin, Z., Gimelshein, N., Antiga, L., Desmaison, A., Kopf, A., Yang, E., DeVito, Z., Raison, M., Tejani, A., Chilamkurthy, S., Steiner, B., Fang, L., Bai, J., and Chintala, S. Pytorch: An imperative style, high-performance deep learning library. In *Advances in Neural Information Processing Systems 32*, pp. 8024–8035. Curran Associates, Inc., 2019.
- Popov, V., Vovk, I., Gogoryan, V., Sadekova, T., Kudinov, M. S., and Wei, J. Diffusion-based voice conversion with fast maximum likelihood sampling scheme. In *International Conference on Learning Representations*, 2022. URL <https://openreview.net/forum?id=8c50f-DoWAu>.
- Qian, K., Zhang, Y., Chang, S., Yang, X., and Hasegawa-Johnson, M. AutoVC: Zero-shot voice style transfer with only autoencoder loss. In *International Conference on Machine Learning*, pp. 5210–5219, 2019. URL <http://proceedings.mlr.press/v97/qian19c.html>.
- Radford, A., Kim, J. W., Xu, T., Brockman, G., McLeavey, C., and Sutskever, I. Robust speech recognition via large-scale weak supervision. In *International Conference on Machine Learning*, 2023.
- Ramesh, A., Dhariwal, P., Nichol, A., Chu, C., and Chen, M. Hierarchical text-conditional image generation with clip latents. *arXiv preprint arXiv:2204.06125*, 2022.
- Ravanelli, M., Parcollet, T., Moumen, A., de Langen, S., Subakan, C., Plantinga, P., Wang, Y., Mousavi, P., Libera, L. D., Ploujnikov, A., Paissan, F., Borra, D., Zaiem, S., Zhao, Z., Zhang, S., Karakasidis, G., Yeh, S.-L., Champion, P., Rouhe, A., Braun, R., Mai, F., Zuluaga-Gomez, J., Mousavi, S. M., Nautsch, A., Liu, X., Sagar, S., Duret, J., Mdhaftar, S., Laperriere, G., Rouvier, M., Mori, R. D., and Esteve, Y. Open-source conversational ai with SpeechBrain 1.0, 2024. URL <https://arxiv.org/abs/2407.00463>.
- Ronneberger, O., Fischer, P., and Brox, T. U-net: Convolutional networks for biomedical image segmentation. In *Medical Image Computing and Computer-Assisted Intervention (MICCAI)*, 2015.
- Schmidhuber, J. Learning factorial codes by predictability minimization. *Neural Computation*, 4(6):863–879, 1992.
- Seed Team, B. Seed-TTS: A family of high-quality versatile speech generation models. *arXiv preprint arXiv:2406.02430*, 2024.
- Sohl-Dickstein, J., Weiss, E., Maheswaranathan, N., and Ganguli, S. Deep unsupervised learning using nonequilibrium

- thermodynamics. In Bach, F. and Blei, D. (eds.), *Proceedings of the 32nd International Conference on Machine Learning*, volume 37 of *Proceedings of Machine Learning Research*, pp. 2256–2265, Lille, France, 7 2015. PMLR.
- Song, Y. and Ermon, S. Generative modeling by estimating gradients of the data distribution. In *Neural Information Processing System*, 2019. URL <https://arxiv.org/pdf/1907.05600>.
- Song, Y. and Ermon, S. Improved techniques for training score-based generative models. In *Neural Information Processing System*, 2020. URL <https://arxiv.org/pdf/2006.09011>.
- Song, Y., Sohl-Dickstein, J., Kingma, D. P., Kumar, A., Ermon, S., and Poole, B. Score-based generative modeling through stochastic differential equations. In *International Conference on Learning Representations*, 2021. URL <https://openreview.net/forum?id=PXTIG12RRHS>.
- Tishby, N., Pereira, F. C., and Bialek, W. The information bottleneck method. In *Proceedings of the 37th annual Allerton conference on communication, control, and computing*, pp. 368–377. University of Illinois Press, 1999.
- Tschannen, M., Bachem, O., and Lucic, M. Recent advances in autoencoder-based representation learning. *arXiv preprint arXiv:1812.05069*, 2018. URL <https://arxiv.org/pdf/1812.05069>.
- Vershynin, R. *High-Dimensional Probability—An Introduction with Applications in Data Science*. Cambridge University Press, Sept. 2018. doi: <https://doi.org/10.1017/9781108231596>.
- Vieira, F., Venugopalan, S., Premasiri, A., et al. A machine-learning based objective measure for ALS disease severity. *NPJ Digital Medicine*, 5:45, 2022. doi: 10.1038/s41746-022-00588-8. URL <https://doi.org/10.1038/s41746-022-00588-8>.
- Wang, L., Gong, Y., Dawalatabad, N., Vilela, M., Placek, K., Tracey, B., Gong, Y., Premasiri, A., Vieira, F., and Glass, J. Automatic prediction of amyotrophic lateral sclerosis progression using longitudinal speech transformer. In *Interspeech*, 2024.
- Wang, W. et al. Vqmivc: Vector quantization and mutual information-based unsupervised speech representation disentanglement for one-shot voice conversion. In *Interspeech*, 2021a. URL <https://arxiv.org/pdf/2106.10132>.
- Wang, Y., Boumadane, A., and Heba, A. A fine-tuned wav2vec 2.0/hubert benchmark for speech emotion recognition, speaker verification and spoken language understanding. *arXiv preprint arXiv:2111.02735*, 2021b.
- Wu, Q., Liu, Y., Zhao, H., Kale, A., Bui, T., Yu, T., Lin, Z., Zhang, Y., and Chang, S. Uncovering the disentanglement capability in text-to-image diffusion models. In *CVPR*, pp. 1900–1910, June 2023a. URL https://openaccess.thecvf.com/content/CVPR2023/html/Wu_Uncovering_the_Disentanglement_Capability_in_Text-to-Image_Diffusion_Models_CVPR_2023_paper.html.
- Wu, Q., Liu, Y., Zhao, H., Kale, A., Bui, T., Yu, T., Lin, Z., Zhang, Y., and Chang, S. Uncovering the disentanglement capability in text-to-image diffusion models. In *CVPR*, pp. 1900–1910, 2023b. doi: 10.1109/CVPR52729.2023.00189. URL https://openaccess.thecvf.com/content/CVPR2023/papers/Wu_Uncovering_the_Disentanglement_Capability_in_Text-to-Image_Diffusion_Models_CVPR_2023_paper.pdf.
- Yang, T., Wang, Y., Lu, Y., and Zheng, N. Disdiff: Unsupervised disentanglement of diffusion probabilistic models. In *Neural Information Processing System*, 2023. URL <https://openreview.net/forum?id=3ofe0lpwQP>.
- Zhang, R., Isola, P., Efros, A. A., Shechtman, E., and Wang, O. The unreasonable effectiveness of deep features as a perceptual metric. In *CVPR*, 2018.

A. Overview of the Appendix

The appendix is organized as follows:

- Section B: Proof of Theorem 3.6. The proof relies mainly on Lemma 3.5, which is proved in Section B.2. Lemma 3.5 in turn relies on Lemma B.1, B.2 and B.3 proved in Section B.3, B.4 and B.5 respectively. Lemma B.1 is further based on Lemma B.4 proved in Section B.6;
- Section C: Proof of Theorem 3.7. The proof relies on Lemma C.3, C.4 and C.5 proved in Section C.1, C.2 and C.3 respectively;
- Section D: an example of continuous random variables \hat{Z} and G such that (\hat{Z}, G) are $(\epsilon, \hat{\psi}, p)$ -editable but $(\hat{Z}|Z, G)$ are not $(\epsilon, \hat{\psi}, p_{X|Z})$ -editable.
- Section E: Proof of Theorem 3.8;
- Section F: Proof of Theorem 3.10;
- Section G: Proof of Theorem 3.11; The proof utilizes Lemma G.1, G.2 and G.3 proved in Section G.1, G.2 and G.3 respectively. Lemma G.3 in turn makes use of Lemma G.4 proved in Section G.4.
- Section H: Proof of Theorem 3.12. The proof relies on Lemma H.1 proved in Section H.1. Lemma H.1 further requires Lemma H.2 and H.3 proved in Section H.2 and H.3 respectively.
- Section I: Experimental details. The setups of GMM experiments are described in Section I.1, followed by the image experiments in Section I.2 and the speech experiments in Section I.3.

B. Proof of Theorem 3.6

B.1. Main proof

First, we assume Lemma 3.5 to be true and defer its proof to Section B.2. Therefore, by the property of (θ_1, ϕ_1) and the optimality of (θ^*, ϕ^*) ,

$$\begin{aligned} L_c^{\gamma, \rho}(\theta^*, \phi^*) &= L_c(\theta^*, \phi^*) + \gamma(I(z_{\phi^*}(X_{t_0}); X) - \rho)_+ \\ &\leq L_c(\theta_1, \phi_1) + \gamma I(z_{\phi_1}(X_{t_0}); X) \\ &\leq \frac{C_2 \delta d_X}{2(T-t_1)} + C_1 \gamma \delta + \gamma(I(Z; X) - \rho)_+, \end{aligned}$$

for some $C_1 = O(\lambda_s \lambda_z \sigma_X d_X)$ and $C_2 := \lambda_{\psi}^2 (\lambda_z + \lambda_g)^2 (1 + \delta^2 \sigma_X^2)$. Since both L_c and $I(Z; X)$ are nonnegative, this implies

$$\begin{aligned} L_c(\theta^*, \phi^*) &\leq \frac{C_2 \delta d_X}{2(T-t_1)} + C_1 \gamma \delta + \gamma(I(Z; X) - \rho)_+, \\ I(z_{\phi^*}(X_{t_0}); X) &\leq I(Z; X) + \frac{C_2 \delta d_X}{2\gamma(T-t_1)} + C_1 \delta. \end{aligned}$$

Choose $\gamma = \frac{C_2 \sqrt{\delta}}{2C_1(T-t_1)}$ and $\rho = O(\delta)$, then we have

$$\begin{aligned} L_c(\theta^*, \phi^*) &\leq \frac{C'_2 \delta d_X}{2(T-t_1)}, \\ I(z_{\phi^*}(X_{t_0}); X) &\leq I(Z; X) + C'_1 \sqrt{\delta} d_X, \end{aligned}$$

for some $C'_2 = 1 + O(\delta)$ and $C'_1 = O(\lambda_s \lambda_z \sigma_X d_X)$.

Let $h(X) := -\int_{\mathcal{X}} p(x) \log p(x) dx$ denote the differential entropy of continuous random variable X . By Assumption 2.1 and the definition of \hat{Z} , we have both $G - X - X_{t_0} - Z$ and $G - \bar{X} - X - \hat{Z}$ form Markov chains, and therefore

$$\begin{aligned} I(Z, G; X) &= h(Z, G) - h(Z, G|X) \\ &= h(Z) - h(Z|X) + h(G) - h(G|X) \\ &= I(Z; X) + I(G; X), \\ I(X_T, \hat{Z}, G; X) &= I(\hat{Z}, G; X) + I(X_T; X|\hat{Z}, G) \\ &= h(\hat{Z}, G) - h(\hat{Z}|X) - h(G|X) + I(X_T; X|\hat{Z}, G) \\ &= I(\hat{Z}; X) + I(G; X) - I(\hat{Z}; G) + I(X_T; X|\hat{Z}, G). \end{aligned}$$

As a result,

$$\begin{aligned} I(\hat{Z}; G) &= I(\hat{Z}; X) + I(G; X) + I(X_T; X|\hat{Z}, G) - I(X_T, \hat{Z}, G; X) \\ &= (I(Z, G; X) - I(X_T, \hat{Z}, G; X)) + I(X_T; X|\hat{Z}, G) + (I(\hat{Z}; X) - I(Z; X)) \\ &= (I(Z, G; X) - I(X_T, \hat{Z}, G; X)) + I(X_T; X|\hat{Z}, G) + C'_1 \sqrt{\delta} d_X, \end{aligned} \quad (21)$$

where the third equality uses item 1. To bound the first bracketed term of the RHS, use the definition $I(Z, G; X)$:

$$I(Z, G; X) = h(X) - h(N) = h(X) - \frac{1}{2} \log(2\pi e \delta^2 d_X),$$

and apply the maximum entropy inequality on $I(X_T, z(X_{t_0}), G; X)$:

$$\begin{aligned} I(X_T, z(X_{t_0}), G; X) &\geq h(X) - \frac{1}{2} \log 2\pi e \mathbb{E} \left\| (e^T - e^{-T}) s_{\theta^*}(X_T, \hat{Z}, G, T) + e^T X_T - X \right\|^2 \\ &\geq h(X) - \frac{1}{2} \log 2\pi e^{2T+1} (1 - e^{-2T})^2 \lim_{t_1 \rightarrow T} L_c(\theta^*, \phi^*) \\ &\geq h(X) - \frac{1}{2} \log 2\pi e C_2 \delta^2 d_X = I(Z, G; X) - \frac{1}{2} \log C_2. \end{aligned}$$

The last inequality uses the optimality of (θ^*, ϕ^*) and therefore $s_{\theta}(x, z, g, t)$'s needs to achieve minimal loss at any $t \in [t_1, T]$, which is upper-bounded by item 2 of Lemma 3.5 as

$$\lim_{t_1 \rightarrow T} L_c(\theta_1, \phi_1) \leq \lim_{t_1 \rightarrow T} \frac{C_2 \delta^2 d_X (e^{2T} - e^{2t_1})}{2(T - t_1)(e^{2t_1} - 1)(e^{2T} - 1)} = \frac{C_2 \delta^2 d_X}{e^{2T} (1 - e^{-2T})^2}.$$

To bound the second term of the RHS of Eq. 21, notice that

$$\begin{aligned} I(X_T; X|\hat{Z}, G) &= h(X_T|\hat{Z}, G) - h(X_T|X, \hat{Z}, G) \\ &\leq h(X_T) - h(X_T|X_{t_0}) \\ &= I(X_T; X_{t_0}) = \mathbb{E}_{p_{X_{t_0}}} D_{\text{KL}}(p_{X_T|X_{t_0}} \| p_{X_T}) \leq C_3 \exp(-T), \end{aligned}$$

for some $C_3 > 0$, where the first inequality uses the facts that $h(X_T) \leq h(X_T|Y)$ for any Y and the data processing inequality along the Markov chain $\hat{Z}, G - X_{t_0} - X_T$. The last inequality uses the convergence property of the OU process (Chen et al., 2023b).

Combining the bounds and choose $T := \Omega(\log \frac{1}{\delta})$, we conclude that

$$\begin{aligned} I(z_{\phi}(X_{t_0}); G) &\leq \frac{1}{2} \log C'_2 + C_3 \exp(-T) + C'_1 \sqrt{\delta} d_X \\ &= O\left(\lambda_s \lambda_z \sigma_X \sqrt{\delta} d_X^2 + C_3 \delta + \frac{1}{2} \log \lambda_{\psi}^2 (\lambda_z + \lambda_g)^2 (1 + \delta^2 \sigma_X^2)\right) = O(\lambda_s \lambda_z \sigma_X d_X^2 \sqrt{\delta}), \end{aligned}$$

where the last inequality uses the assumption that $\lambda_{\psi}(\lambda_z + \lambda_g) = 1 + O(\delta)$.

B.2. Proof of Lemma 3.5

To prove the lemma, we need the following technical lemmas, whose proofs are deferred to Section B.3, B.4 and B.5 respectively.

Lemma B.1. *Suppose random variable $Y = \alpha X + N \sim p_Y$, for independent random variables $X \sim p_X$ and $N \sim \mathcal{N}(0, \sigma^2 I_d)$, where $\alpha := \sqrt{1 - \sigma^2}$. Then, for any distribution $q(x|z)$ with an L -Lipschitz score function in x for any $z \in \mathcal{X}$, then the following inequality holds:*

$$\mathbb{E}_{p_{XY Z}(x,y,z)} \log \frac{q(y|z)}{q(x|z)} \leq CL[\sigma^2(1 + \sigma^2)\mathbb{E}\|X\|^2 + \sigma^2 d + \sigma \sqrt{\mathbb{E}\|X\|^2 d}],$$

for some constant $C > 0$ independent of σ , $\mathbb{E}\|X\|^2$ and d .

Lemma B.2. *For $\delta < 1/2$ and $\alpha := \sqrt{1 - \delta^2}$, the KL divergence between two Gaussian distributions $\mathcal{N}(\alpha\mu, \delta^2 I_d)$ and $\mathcal{N}(\frac{1}{\alpha}\mu, \frac{\delta^2}{\alpha^2} I_d)$ is upper-bounded as*

$$D_{\text{KL}}\left(\mathcal{N}(\alpha\mu, \delta^2 I_d) \left\| \mathcal{N}\left(\frac{1}{\alpha}\mu, \frac{\delta^2}{\alpha^2} I_d\right)\right.\right) \leq \left(\frac{\|\mu\|^2}{2} + \frac{d}{6}\right)\delta^2.$$

Lemma B.3. *Suppose random variable $Y = \alpha X + N \sim p_Y$, for independent random variables $X \sim p_X$ and $N \sim \mathcal{N}(0, \sigma^2 I_d)$, where $\alpha := \sqrt{1 - \sigma^2}$. Further, suppose $p_X(x) > 0$ for any $x \in \mathbb{R}^d$ and the score function of p_X is L -Lipschitz, then the following holds*

$$\max\{D_{\text{KL}}(p_X \| p_Y), D_{\text{KL}}(p_Y \| p_X)\} \leq CL(\mathbb{E}\|X\|^2)^{1/2} \delta d,$$

for some constant $C > 0$.

Assuming Lemma B.1-B.3, choose $t_0 := \frac{1}{2} \log \frac{1}{1 - \delta^2}$. Then by the property of the OU process,

$$X_{t_0} = \alpha X + \delta N_{t_0}.$$

Next, set ϕ_1 such that $z_{\phi_1}(x) = z(x)$ and define

$$\begin{aligned} q(z|x) &:= p_{z(X_{t_0})|X}(z|x), \\ q(z) &:= p_{z(X_{t_0})}(z), \\ p(z|x) &:= p_{Z|X}(z|x), \\ p(z) &:= p_Z(z), \end{aligned}$$

then by definition,

$$\begin{aligned} &I(z(X_{t_0}); X) - I(Z; X) \\ &= \mathbb{E}_{\bar{X}, X, X_{t_0}} \log \frac{q(z(X_{t_0})|X)p(Z)}{q(X_{t_0})p(z(\bar{X})|X)} \\ &= \mathbb{E}_{\bar{X}, X, X_{t_0}} \log \frac{q(z(X_{t_0})|X)}{p(z(\bar{X})|X)} + \mathbb{E}_{\bar{X}, X_{t_0}} \log \frac{p(Z)}{q(z(X_{t_0}))}. \end{aligned}$$

For the first term on the RHS, notice that

$$\begin{aligned} &\mathbb{E}_{\bar{X}, X, X_{t_0}} \log \frac{q(z(X_{t_0})|X)}{p(z(\bar{X})|X)} \\ &= \mathbb{E}_{p_{X X_{t_0}}(x, x')} \log \frac{q(z(x')|x)}{p(z(x')|x)} + \mathbb{E}_{p_{\bar{X} X X_{t_0}}(\bar{x}, x, x')} \log \frac{p(z(x')|x)}{p(z(\bar{x})|x)} \\ &= \mathbb{E}_{p_X} D_{\text{KL}}(p_{z(X_{t_0})|X} \| p_{Z|X}) + \mathbb{E}_{p_{\bar{X} X X_{t_0}}(\bar{x}, x, x')} \log \frac{p(z(x')|x)}{p(z(\bar{x})|x)} \\ &= \mathbb{E}_{p_X} D_{\text{KL}}(p_{z(X_{t_0})|X} \| p_{Z|X}) + O(\lambda_s \lambda_z \sigma_X d_X \delta), \end{aligned} \tag{22}$$

where the last equality uses Lemma B.1 with $L := \lambda_s \lambda_z$ and $\sigma := \sqrt{1 - \alpha^4} = O(\delta)$, and the fact that $\delta < 1/\sqrt{d_X}$ and thus the term involving $\delta\sqrt{d_X}$ dominates those involving $\delta^2 d_X$ in the bound. To bound the KL term, apply data processing inequality,

$$\begin{aligned}
 & \mathbb{E}_{p_X} D_{\text{KL}}(p_{z(X_{t_0})|X} \| p_{Z|X}) \\
 & \leq \mathbb{E}_{p_X} D_{\text{KL}}(p_{X_{t_0}|X} \| p_{\bar{X}|X}) \\
 & = \mathbb{E}_{p_{X_{t_0}}(x,y)} \log \frac{\mathcal{N}(y|\alpha_{t_0}x, \delta^2 I_{d_X})}{p_{\bar{X}}(y) \mathcal{N}(x|\alpha y, \delta^2 I_{d_X}) / p_X(x)} \\
 & = \mathbb{E}_{p_{X_{t_0}}(x,y)} \log \frac{p_X(x)}{p_{\bar{X}}(y)} + O((d_X + \mathbb{E}\|X\|)\delta^2) \\
 & = D_{\text{KL}}(p_X \| p_{\bar{X}}) + \mathbb{E}_{p_{X_{t_0}}(x,y)} \log \frac{p_{\bar{X}}(x)}{p_{\bar{X}}(y)} + O((d_X + \mathbb{E}\|X\|)\delta^2),
 \end{aligned}$$

where the second-to-last equality uses Lemma B.2:

$$\begin{aligned}
 & \mathbb{E}_{p_{X_{t_0}}(x,y)} \log \frac{\mathcal{N}(y|\alpha x, \delta^2 I_{d_X})}{\mathcal{N}(x|\alpha y, \delta^2 I_{d_X})} \\
 & = \mathbb{E}_{p_{X_{t_0}}(x,y)} \log \frac{\mathcal{N}(y|\alpha_{t_0}x, \delta_{t_0}^2 I_{d_X})}{\mathcal{N}(y|\frac{1}{\alpha}x, \frac{\delta^2}{\alpha^2} I_{d_X})} + d_X \log(1/\alpha) \\
 & = O((d_X + \mathbb{E}\|X\|)\delta^2).
 \end{aligned}$$

To proceed, notice further that

$$\begin{aligned}
 & D_{\text{KL}}(p_X \| p_{\bar{X}}) + \mathbb{E}_{p_{X_{t_0}}(x,y)} \log \frac{p_{\bar{X}}(x)}{p_{\bar{X}}(y)} \\
 & = D_{\text{KL}}(p_X \| p_{\bar{X}}) + O(\lambda_s \sigma d_X \delta) \\
 & = O(\lambda_s d_X \sigma_X \delta),
 \end{aligned}$$

where the first equality uses Lemma B.1 with $\delta < 1/\sqrt{d_X}$, and the second inequality uses Lemma B.3. Combining this with Eq. 22, the first term is upper-bounded as

$$\mathbb{E}_{\bar{X}, X, X_{t_0}} \log \frac{q(z(X_{t_0})|X)}{p(z(\bar{X})|X)} = O((\lambda_s + \lambda_s \lambda_z) d_X \sigma_X \delta). \quad (23)$$

For the second term, notice that by Assumption 3.2 and the fact that a weighted sum preserves Lipschitzness, $\nabla_z \log p(z) = \mathbb{E}_{p(x|z)} \nabla_z \log p(z|x)$ is λ_s -Lipschitz, and therefore

$$\begin{aligned}
 & \mathbb{E}_{\bar{X}, X_{t_0}} \log \frac{p(z(\bar{X}))}{q(z(X_{t_0}))} \\
 & = \mathbb{E}_{p_{X_{t_0}}(x)} \log \frac{p(z(x))}{q(z(x))} + \mathbb{E}_{p_{\bar{X}|X_{t_0}}(x,y)} \log \frac{p(z(x))}{p(z(y))} \\
 & = -D_{\text{KL}}(p_{z(X_{t_0})} \| p_Z) + \mathbb{E}_{p_{\bar{X}|X_{t_0}}(x,y)} \log \frac{p(z(x))}{p(z(y))} \\
 & = O(\lambda_s \lambda_z d_X \sigma_X \delta),
 \end{aligned} \quad (24)$$

where the last equality use the non-negativity of KL divergence and Lemma B.1 on $p(z)$ with $\delta < 1/\sqrt{d_X}$. Combining Eq. 23 and Eq. 24 yields item 1 of the theorem:

$$I(z(X_{t_0}); X) - I(Z; X) \leq C_1 \delta = O\left(\frac{\epsilon^2}{\lambda_s \lambda_z \sigma_X d_X^3}\right),$$

where $C_1 := O(\lambda_s \lambda_z \sigma_X d_X)$.

To prove item 2, set θ_1 so that the score function

$$s_{\theta_1}(X_t, z_\phi(X_{t_0}), G, t) = \frac{\exp(-t)\psi(z_\phi(X_{t_0}), G) - X_t}{1 - \exp(-2t)}.$$

Then by the conditional disentanglement property in Assumption 2.1 and set $t_1 := \frac{1}{2} \log \frac{1}{1-\delta}$, the loss L_c becomes

$$\begin{aligned} L_c(\theta_1, \phi_1) &= \frac{1}{T-t_1} \int_{t_1}^T \frac{\exp(-2t) \mathbb{E} \|\psi(z(X_{t_0}), G) - X\|^2}{(1 - \exp(-2t))^2} dt \\ &=: \frac{\tilde{L}_c(\theta_1, \phi_1)}{T-t_1} \int_{e^{t_1}}^{e^T} \frac{d\tau}{2(\tau-1)^2} \\ &= \frac{\tilde{L}_c(\theta_1, \phi_1)}{2(T-t_1)} \left(\frac{1}{e^{2t_1}-1} - \frac{1}{e^{2T}-1} \right) \leq \frac{\tilde{L}_c(\theta_1, \phi_1)(e^{2T} - e^{2t_1})}{2(T-t_1)(e^{2t_1}-1)(e^{2T}-1)}, \end{aligned}$$

Further, notice that

$$\begin{aligned} \tilde{L}_c(\theta_1, \phi_1) &:= \mathbb{E} \|\psi(z(X_{t_0}), G) - X\|^2 \\ &= \mathbb{E} \|\psi(z(X_{t_0}), G) - \psi(z(X), g(X))\|^2 \\ &= \lambda_\psi^2 \mathbb{E} (\lambda_z \|X_{t_0} - X\| + \lambda_g \|X - \bar{X}\|)^2 \\ &\leq \lambda_\psi^2 (\lambda_z + \lambda_g)^2 (1 + \delta^2 \sigma_X^2) \delta^2 d_X \\ &=: C_2 \delta^2 d_X, \end{aligned}$$

where $C_2 := \lambda_\psi^2 (\lambda_z + \lambda_g)^2 (1 + \delta^2 \sigma_X^2)$. The first inequality uses Cauchy-Schwarz inequality, and the second inequality uses the fact that $\psi(z(X), G)$ is the MMSE estimator and $Z - z(X) - X - G$ forms a Markov chain. The last inequality uses the fact that

$$\begin{aligned} \mathbb{E} \|X_{t_0} - X\|^2 &= \mathbb{E} \|X - \bar{X}\|^2 \\ &= \mathbb{E} \|(\alpha-1)X + N_t\|^2 \\ &= (1 - \sqrt{1-\delta^2})^2 \mathbb{E} \|X\|^2 + \mathbb{E} \|N_{t_0}\|^2 \\ &\leq \delta^4 \sigma_X^2 d_X + \delta^2 d_X. \end{aligned}$$

As a result, item 2 follows from

$$L_c(\theta_1, \phi_1) \leq \frac{C_2 \delta^2 d_X (e^{2T} - e^{2t_1})}{2(T-t_1)(e^{2t_1}-1)(e^{2T}-1)} \leq \frac{C_2 \delta d_X}{2(T-t_1)} = O\left(\frac{\epsilon^2}{\lambda_s^2 \lambda_z^2 \sigma_X^2 d_X^3 T}\right)$$

with the choice of t_1 and δ and the fact that $0 < \delta < \min\{T, 1\}$.

B.3. Proof of Lemma B.1

To begin, we make use the following lemma proved in Section B.6.

Lemma B.4. *Suppose the score function $s_q(x|z) := \nabla_x \log q(x|z)$ of the probability density $q(x|z)$ is L -Lipschitz as a function of x , then the following inequality holds:*

$$\log \frac{q(y|z)}{q(x|z)} \leq (L\|x\| + \|s_q(\mathbf{0}_d|z)\|) \|y-x\| + \frac{L\|y-x\|^2}{2}.$$

Set $s_q(x|z) := \nabla_x \log q(x|z)$, then by Lemma B.4,

$$\begin{aligned} &\mathbb{E}_{p_{XY}Z(x,y,z)} \log \frac{q(y|z)}{q(x|z)} \\ &\leq \mathbb{E}_{p_{XY}Z(x,y,z)} (L\|x\| + \|s_q(\mathbf{0}_d|z)\|) \|y-x\| + \frac{L}{2} \mathbb{E}_{p_{XY}(x,y)} \|y-x\|^2. \end{aligned}$$

For the first term of the RHS, notice that

$$\begin{aligned}
 & \mathbb{E}_{p_{XYZ}(x,y,z)}(L\|x\| + \|s_q(\mathbf{0}_d|z)\|)\|y-x\| \\
 & \leq \sqrt{\mathbb{E}(L\|X\| + \|s_q(\mathbf{0}_d|Z)\|)^2} \mathbb{E}\|Y-X\|^2 \\
 & = \sqrt{\mathbb{E}(L\|X\| + \|s_q(\mathbf{0}_d|Z)\|)^2} \sqrt{(1-\alpha)^2 \mathbb{E}\|X\|^2 + \sigma^2 d} \\
 & \leq \sqrt{2L^2 \mathbb{E}\|X\|^2 + C_1} \sqrt{(1-\alpha)^2 \mathbb{E}\|X\|^2 + \sigma^2 d} \\
 & \leq C_2 L(\sigma^2 \mathbb{E}\|X\|^2 + \sigma \sqrt{\mathbb{E}\|X\|^2 d}),
 \end{aligned}$$

where $C_1 := 2\mathbb{E}\|s_q(\mathbf{0}_d|Z)\|^2$ and C_2 large enough. To bound the second term of the RHS, notice that

$$\begin{aligned}
 \frac{L}{2} \mathbb{E}_{p_{XY}(x,y)} \|y-x\|^2 &= \frac{L}{2} [(1-\alpha)^2 \mathbb{E}\|X\|^2 + \mathbb{E}\|N\|^2] \\
 &\leq \frac{L\sigma^2}{2} (\sigma^2 \mathbb{E}\|X\|^2 + d).
 \end{aligned}$$

Combining the two terms yields

$$\mathbb{E}_{p_{XYZ}(x,y,z)} \log \frac{q(y|z)}{q(x|z)} \leq CL[\sigma^2(1+\sigma^2)\mathbb{E}\|X\|^2 + \sigma^2 d + \sigma \sqrt{\mathbb{E}\|X\|^2 d}],$$

for some $C > 0$ large enough.

B.4. Proof of Lemma B.2

Use the formula for the KL divergence between Gaussians:

$$\begin{aligned}
 & D_{\text{KL}}(\mathcal{N}(\alpha\mu, \delta^2 I_d) \|\mathcal{N}(\mu/\alpha, (\delta/\alpha)^2 I_d)) \\
 &= \left(\frac{\delta^2}{(\delta/\alpha)^2} - 1 \right) d/2 + \frac{\|\alpha\mu - \mu/\alpha\|^2}{2(\delta/\alpha)^2} + d \log \frac{1}{\alpha} \\
 &= -\frac{d}{2} \delta^2 + \frac{\delta^2 \|\mu\|^2}{2} + d \log \frac{1}{\alpha} \\
 &= \frac{(\|\mu\|^2 - d)\delta^2}{2} + \frac{d}{2} \log \left(1 + \frac{\delta^2}{1-\delta^2} \right) \\
 &\leq \frac{(\|\mu\|^2 - d)\delta^2}{2} + \frac{2d\delta^2}{3} = \left(\frac{\|\mu\|^2}{2} + \frac{d}{6} \right) \delta^2.
 \end{aligned}$$

B.5. Proof of Lemma B.3

By Jensen's inequality,

$$\begin{aligned}
 & D_{\text{KL}}(p_Y \| p_X) \\
 &= \mathbb{E}_{\int p_X(x) \mathcal{N}(y|\alpha x, \sigma^2 I_d) dx} \log \frac{\int p_X(x) \mathcal{N}(y|\alpha x, \sigma^2 I_d) dx}{p_X(y)} \\
 &\leq \mathbb{E}_{p_X(x) \mathcal{N}(y|\alpha x, \sigma^2 I_d)} \log \frac{p_X(x) \mathcal{N}(y|\alpha x, \sigma^2 I_d)}{p_X(y) \mathcal{N}(x|y/\alpha, (\sigma/\alpha)^2 I_d)} \\
 &= \mathbb{E}_{p_X(x) \mathcal{N}(y|\alpha x, \sigma^2 I_d)} \log \frac{p_X(x)}{p_X(y)} + d \log \alpha \\
 &\leq \mathbb{E}_{p_X(x) \mathcal{N}(y|\alpha x, \sigma^2 I_d)} \log \frac{p_X(x)}{p_X(y)} - \frac{\sigma^2 d}{2} \\
 &\leq CL(\sigma^2 \mathbb{E}\|X\|^2 + \sigma^2 d + \sigma \sqrt{d} \mathbb{E}\|X\|),
 \end{aligned}$$

for some $C > 0$, where the last inequality uses Lemma B.1.

Similarly, apply Jensen's inequality and Lemma B.1,

$$\begin{aligned}
 D_{\text{KL}}(p_X \| p_Y) &= \mathbb{E}_{p_X(x)} \log \frac{p_X(x)}{\int p_X(y) \mathcal{N}(x|\alpha y, \sigma^2 I_d) dy} \\
 &\leq \mathbb{E}_{p_X(x) \mathcal{N}(y|x/\alpha, (\sigma/\alpha)^2 I_d)} \log \frac{p_X(x) \mathcal{N}(y|x/\alpha, (\sigma/\alpha)^2 I_d)}{p_X(y) \mathcal{N}(x|\alpha y, \sigma^2 I_d)} \\
 &= \mathbb{E}_{p_X(x) \mathcal{N}(y|x/\alpha, (\sigma/\alpha)^2 I_d)} \log \frac{p_X(x)}{p_X(y)} + d \log \alpha \\
 &\leq CL(\sigma^2 \mathbb{E} \|X\|^2 + \sigma^2 d + \sigma \sqrt{d} \mathbb{E} \|X\|).
 \end{aligned}$$

B.6. Proof of Lemma B.4

by the Lipschitz property of the score function of s_q , for any $(x, y) \in \mathcal{X}^2$,

$$\begin{aligned}
 \log \frac{q(y|z)}{q(x|z)} &\leq |\langle s_q(x|z), y-x \rangle| + \frac{L \|y-x\|^2}{2} \\
 &\leq \|s_q(x|z)\| \|y-x\| + \frac{L \|y-x\|^2}{2}.
 \end{aligned}$$

Apply the Lipschitz property of s_q again,

$$\log \frac{q(y|z)}{q(x|z)} \leq (L \|x\| + \|s_q(\mathbf{0}_d|z)\|) \|y-x\| + \frac{L \|y-x\|^2}{2}. \quad (25)$$

C. Proof of Theorem 3.7

To prove this theorem, we need two additional assumptions below.

Assumption C.1. For any $\theta \in \Theta, \phi \in \Phi$, the estimated score function is of the form $s_\theta(x, z, g, t) := \frac{\bar{s}_\theta(z, g, t) - x}{1 - \exp(-2t)}$, where $\bar{s}_\theta(\cdot, \cdot, \cdot)$ is λ_Θ -Lipschitz in all arguments, and the score function for $p_{z_\phi(X_{t_0}), G|X}(z, g|x)$ is λ_Φ -Lipschitz in z and g for any $x \in \mathcal{X}$.

We further make an assumption for ψ_θ similar to Assumption 3.2 for ψ . The assumption ensures that the estimated mapping ψ_θ is sufficiently smooth.

Assumption C.2. For any $\theta \in \Theta$, the function ψ_θ is invertible and twice-differentiable with bounded first and second-order derivatives:

$$\begin{aligned}
 \|J_{[z^\top, g^\top]^\top} \psi_\theta(z, g)\| &\in [l_\psi, \lambda_\psi], \\
 \sum_j^{d_X} \|\nabla_{[z^\top, g^\top]^\top}^2 [\psi_\theta(z, g)]_j\|^2 &\leq H_\psi, \forall (z, g) \in \mathcal{Z} \times \mathcal{G},
 \end{aligned}$$

for some $0 < l_\psi \leq \lambda_\psi$ and $H_\psi > 0$, where $J_x f(x)$ is the Jacobian matrix of function f with respect to vector x and $[v]_i$ denotes the i -th coordinate of vector v .

Assumption C.1 ensures the function class for the bottleneck and the score function are Lipschitz, analogous to the setting in (Chen et al., 2023a). For Assumption C.2, note that the invertibility of ψ_θ is not strictly necessary and can be relaxed to injectivity by partitioning the domain. Further, Assumption C.1 can be relaxed to only require $s_\theta(\cdot)$ to be Lipschitz if we use a weaker notion of editability by allowing stochastic processes in the definition instead of the deterministic relation $\psi_\theta(\hat{Z}, G)$.

Again, we start by introducing a couple helpful lemmas and postponing their proofs to Section C.1, C.2 and C.3.

Lemma C.3. (Novikov's condition) *The following bounds hold with*

$$\begin{aligned}
 \mathbb{E}_{X, (X_t^\leftarrow)_t} \exp \left(\frac{1}{2} \int_0^{T-t} \|s_\theta(X_t^\leftarrow, \hat{Z}, G, T-t) - \nabla_x \log p_{T-t_0}(X_t^\leftarrow | X)\|^2 dt \right) &< \infty \\
 \mathbb{E}_{(X_t^\leftarrow)_t} \exp \left(\frac{1}{2} \int_0^{T-t} \|\mathbb{E}[s_\theta(X_t^\leftarrow, \hat{Z}, G, T-t) | X_t^\leftarrow] - \nabla_x \log p_{T-t_0}(X_t^\leftarrow)\|^2 dt \right) &< \infty.
 \end{aligned}$$

Lemma C.4. (moment bounds) *The conditional variance and the second moment of \hat{X} are bounded:*

$$\begin{aligned}\text{Var}(\hat{X}|X) &:= \mathbb{E}\|\hat{X} - \mathbb{E}[\hat{X}|X]\|^2 \leq C_1 \delta d_X, \\ \mathbb{E}\|\hat{X}\|^2 &\leq C_2 \sigma_X^2 d_X,\end{aligned}$$

for some constant $C_1, C_2 > 0$.

Lemma C.5. (Theorem 2.1 of (Chae & Walker, 2020)) *For joint probability density p_{XY} such that its marginals have bounded gradients:*

$$\max\left\{\sup_{x \in \mathcal{X}} \|\nabla_x p_X(x)\|, \sup_{y \in \mathcal{Y}} \|\nabla_y p_Y(y)\|\right\} < \infty,$$

the following inequalities hold:

$$d_{\text{TV}}(p_X, p_Y) \leq C \mathcal{W}_1^{1/2}(p_X, p_Y) \leq C (\mathbb{E}_{(X,Y) \sim p_{XY}} \|X - Y\|^2)^{1/4},$$

for some scalar constant $C > 0$, where $\mathcal{W}_r(p, q) := (\int_{\pi \in \Pi(p, q)} \|x - y\|^r d\pi(x, y))^{1/r}$ is the Wasserstein- r distance and $\Pi(p, q)$ is the set of distributions with marginals p and q .

To prove the first condition of editability, first define $\hat{\psi} := \psi_{\theta^*}$ and $\Psi := \hat{\psi}(\hat{Z}, G)$ and use triangle inequality:

$$d_{\text{TV}}(p_\Psi, p) \leq d_{\text{TV}}(p_\Psi, p_{\hat{X}}) + d_{\text{TV}}(p_{\hat{X}}, p_{X_{t_0}}) + d_{\text{TV}}(p_{X_{t_0}}, p). \quad (26)$$

To bound the first term on the RHS, we apply Lemma C.5 to $p_{\Psi, \hat{X}}$. To this end, we first check that p_Ψ and $p_{\hat{X}}$ have bounded gradients. Indeed, notice that

$$\begin{aligned}\|\nabla_x p_{\hat{X}}(x)\| &= \|p_{\hat{X}}(x) \nabla_x \log p_{\hat{X}}(x)\| \\ &\leq \mathbb{E}_{p_{\hat{X}}(x)} \|\nabla_x \log p_{\hat{X}}(x)\| \\ &= \mathbb{E}_{p_{\hat{X}}(x)} \|\mathbb{E}_{\hat{Z}, G} [s_\theta(\hat{X}, \hat{Z}, G, t_0)]\| \\ &\leq \mathbb{E}_{p_{\hat{X}}(x)} [\|\mathbb{E}_{\hat{Z}, G} [s_\theta(\hat{X}, \hat{Z}, G, t_0) - s_\theta(\mathbf{0}_{d_X}, \hat{Z}, G, t_0)]\| + \|\mathbb{E}_{\hat{Z}, G} s_\theta(\mathbf{0}_{d_X}, \hat{Z}, G, t_0)\|] \\ &\leq \lambda_\Theta \mathbb{E}_{p_{\hat{X}}(x)} \|\hat{X}\| + C \\ &\leq \lambda_\Theta C_1 \sigma_X d_X^{1/2} + C < \infty,\end{aligned}$$

for some constant $C_1 > 0$ and $C := \sup_{(z, g) \in \mathcal{Z} \times \mathcal{G}} \|s_\theta(\mathbf{0}_{d_X}, z, g, t_0)\|$. The third-to-last equality uses Assumption C.1 and the second-to-last inequality uses Lemma C.4.

Further, define $v := [\hat{z}^\top, g^\top]^\top \in \mathcal{V}$, notice that by Assumption C.2,

$$\begin{aligned}\|\nabla_x p_\Psi(x)\| &= \|p_\Psi(x) \nabla_x \log p_\Psi(x)\| \\ &\leq \mathbb{E}_{p_\Psi(x)} \|\nabla_x \hat{\psi}^{-1}(x) \nabla_{[\hat{z}^\top, g^\top]^\top} \log p_{\hat{Z}, G}(\hat{\psi}^{-1}(x))\| + \|\nabla_x \log \det(J_x \hat{\psi}^{-1}(x))\| \\ &\leq \mathbb{E}_{p_\Psi(x)} \frac{\|\nabla_{[\hat{z}^\top, g^\top]^\top} \log p_{\hat{Z}, G}(\hat{\psi}^{-1}(x))\|}{\inf_{v \in \mathcal{V}} \|\nabla_v \hat{\psi}(v)\|} + \|\nabla_x \log \det(J_x \hat{\psi}^{-1}(x))\| \\ &\leq \frac{1}{l_\psi} (\lambda_\Phi \mathbb{E}_{p_\Psi(x)} \|\psi^{-1}(x)\| + C_2) + \|\nabla_x \log \det(J_x \hat{\psi}^{-1}(x))\| \\ &\leq \frac{\lambda_\Phi}{l_\psi \lambda_\psi} \mathbb{E}_{p_\Psi(x)} \|x\| + C_3 + \|\nabla_x \log \det(J_x \hat{\psi}^{-1}(x))\|,\end{aligned}$$

where $J_x f$ denotes the Jacobian matrix of vector function f with respect to vector x , $C_2 := \|\nabla_v \log p_{\hat{Z}, G}(\mathbf{0}_{d_Z}, \mathbf{0}_{d_G})\|$ and $C_3 := \frac{C_2}{l_\psi} + \frac{\lambda_s}{l_\psi \lambda_\psi} \|\psi^{-1}(\mathbf{0}_{d_X})\|$. To bound the last term of the RHS, use chain rule and the identity

$\frac{d\det(A(t))}{dt} = \det(A(t))\text{Tr}(A(t)^{-1}\frac{dA(t)}{dt})$ for invertible matrix A ,

$$\begin{aligned}
 & \|\nabla_x \log \det(J_x \hat{\psi}^{-1}(x))\|^2 \\
 & \leq \sum_{i=1}^{d_X} \left(\frac{\partial}{\partial x_i} \log \det(J_x \hat{\psi}^{-1}(x)) \right)^2 \\
 & \leq \sum_{i=1}^{d_X} \left(\frac{\partial}{\partial x_i} \log \det(J_v \hat{\psi}(\psi^{-1}(x))) \right)^2 \\
 & = \sum_{i=1}^{d_X} \left(\text{Tr} \left(J_v^{-1} \hat{\psi}(\hat{\psi}^{-1}(x))^\top \frac{\partial J_v \hat{\psi}(\hat{\psi}^{-1}(x))}{\partial x_i} \right) \right)^2 \\
 & \leq \frac{1}{\|J_v \hat{\psi}(\hat{\psi}^{-1}(x))\|^2} \sum_{i=1}^{d_X} \left\| \frac{\partial J_v \hat{\psi}(\hat{\psi}^{-1}(x))}{\partial x_i} \right\|^2 \\
 & \leq \frac{1}{\|J_v \hat{\psi}(\hat{\psi}^{-1}(x))\|^4} \sum_{j=1}^{d_Z+d_G} \|\nabla_v^2 \hat{\psi}[\hat{\psi}^{-1}(x)]_j\|^2 \\
 & \leq \frac{H_\psi^2}{l_\psi^4} < \infty.
 \end{aligned}$$

As a result of Lemma C.5,

$$d_{\text{TV}}(p_\Psi, p_{\hat{X}}) = O((\mathbb{E}\|\hat{X} - \mathbb{E}[\hat{X}|X]\|^2)^{1/4}) = O(\delta^{1/4} d_X^{1/4}).$$

where the first equality uses Lemma C.5 and the second equality uses Lemma C.4.

To bound the second term, by Lemma C.3, Novikov's condition holds for s_{θ^*} , therefore Girsanov's theorem (Chen et al., 2023b) and Theorem 3.6 imply that

$$\begin{aligned}
 d_{\text{TV}}(p_{\hat{X}|X}, p_{X_{t_1}|X}) &= \mathbb{E}_{p(X)} d_{\text{TV}}(p_{\hat{X}|X}, \mathcal{N}(X, \delta^2 I_{d_X})) \\
 &= O\left((\sqrt{d_X} \eta + \sigma_X^2 \eta) \lambda_\Theta \sqrt{T} + \sqrt{L_c(\theta^*, \phi^*) T}\right) \\
 &= O\left((\sqrt{d_X} \eta + \sigma_X^2 \eta) \lambda_\Theta \sqrt{T} + \sqrt{\delta d_X}\right)
 \end{aligned}$$

Further, by Jensen's inequality,

$$\begin{aligned}
 d_{\text{TV}}(p_{\hat{X}}, p_{X_{t_1}}) &\leq d_{\text{TV}}(p_{\hat{X}|X}, p_{X_{t_1}|X}) \\
 &= O\left((\sqrt{d_X} \eta + \sigma_X^2 \eta) \lambda_\Theta \sqrt{T} + \sqrt{\delta d_X}\right).
 \end{aligned}$$

Finally, for the last term on the RHS of Eq. 26, we use Pinsker's inequality and Lemma B.3 to conclude that

$$\begin{aligned}
 d_{\text{TV}}(p_{X_{t_1}}, p) &\leq \sqrt{\frac{D_{\text{KL}}(p||p_{X_{t_1}})}{2 \log 2}} \\
 &= O(\sqrt{\lambda_s d_X \sigma_X \delta^{1/2}}).
 \end{aligned}$$

Combining bounds for the terms, we obtain

$$\begin{aligned}
 d_{\text{TV}}(p_\Psi, p) &= O(\delta^{1/4} d_X^{1/4} + (\sqrt{d_X} \eta + \sigma_X^2 \eta) \lambda_\Theta \sqrt{T} + \sqrt{\delta d_X} + \sqrt{\lambda_s \sigma_X \delta^{1/2} d_X}) \\
 &= O((\sqrt{d_X} \eta + \sigma_X^2 \eta) \lambda_\Theta \sqrt{T} + \sqrt{\lambda_s \sigma_X \delta^{1/2} d_X}).
 \end{aligned}$$

Set the diffusion step size $\eta = \frac{\epsilon}{\lambda_s^2 d_X T} + \frac{\sqrt{\epsilon}}{\lambda_\Theta \sigma_X^2 \sqrt{T}}$ and $\delta = \frac{\epsilon^2}{\lambda_s \lambda_z \sigma_X d_X}$ yields item 1.

To prove the second condition for editability, consider i.i.d copies (\hat{Z}_1, G_1) and (\hat{Z}_2, G_2) sampled from $p_{\hat{Z}G}$, and apply Theorem 3.6 with Pinsker's inequality

$$d_{\text{TV}}(p_{\hat{Z}_1 G_1}, p_{\hat{Z}_2 G_2}) \leq \sqrt{\frac{D_{\text{KL}}(p_{\hat{Z}G} \| p_{\hat{Z}G})}{2 \log 2}} = O(\sqrt{\lambda_s \lambda_z \sigma_X d_X^2 \delta^{1/2}})$$

Further, applying data processing inequality yields

$$d_{\text{TV}}(p_{\hat{\psi}(\hat{Z}_1, G_1)}, p_{\hat{\psi}(\hat{Z}_2, G_2)}) \leq d_{\text{TV}}(p_{\hat{Z}_1 G_1}, p_{\hat{Z}_2 G_2}) = O(\sqrt{\lambda_s \lambda_z \sigma_X d_X^2 \delta^{1/2}}),$$

which implies (\hat{Z}, G) is $(O(\sqrt{\lambda_s \lambda_z \sigma_X d_X^2 \delta^{1/2}}), \hat{\psi})$ -editable. Setting $\delta := \frac{\epsilon^2}{\lambda_s^2 \lambda_z^2 \sigma_X^2 d_X^4}$ proves item 2 of the theorem.

C.1. Proof of Lemma C.3

We start by proving the first bound. First, by the equivalence of the forward and reverse process,

$$\begin{aligned} & \mathbb{E}_{X, (X_t^\leftarrow)_t} \exp \left(\frac{1}{2} \int_0^{T-t_1} \|s_\theta(X_t^\leftarrow, \hat{Z}, G, T-t) - \nabla_x \log p_{T-t_1}(X_t^\leftarrow | X)\|^2 dt \right) \\ &= \mathbb{E}_{p(X)} \mathbb{E}_{t, p_t(\cdot | X)} \exp \left(\frac{1}{2} \int_{t_1}^T \|s_\theta(X_t, \hat{Z}, G, t) - \nabla_x \log p_t(X_t | X)\|^2 dt \right) \end{aligned} \quad (27)$$

$$= \mathbb{E}_{p(X)} \mathbb{E}_{t, p_t(\cdot | X)} \exp \left(\frac{1}{2} \int_{t_1}^T \left\| \frac{\bar{s}_\theta(\hat{Z}, G, t) - X_t}{1 - e^{-2t}} + \frac{X_t - e^{-t} X}{1 - e^{-2t}} \right\|^2 dt \right) \quad (28)$$

$$= \mathbb{E}_{p(X)} \mathbb{E}_{t, p_t(\cdot | X)} \exp \left(\frac{1}{2} \int_{t_1}^T \frac{\|\bar{s}_\theta(\hat{Z}, G, t) - e^{-t} X\|^2}{(1 - e^{-2t})^2} dt \right), \quad (29)$$

where the second-to-last equality uses Assumption C.1 and the closed-form formula of the conditional score function of an OU process. Now, let $\sigma(t) := \sqrt{1 - e^{-2t}}$ and $\lambda := 2\lambda_\Theta(1 + \lambda_z + \lambda_s)$, and by Assumption C.1 again, we have

$$\begin{aligned} & \mathbb{E} \exp \left(\frac{1}{2} \int_{t_1}^T \frac{\|\bar{s}_\theta(\hat{Z}, G, t) - e^{-t} X\|^2}{\sigma(t)^4} dt \right) \\ & \lesssim \mathbb{E} \exp \left(\frac{1}{2} \int_{t_1}^T \frac{\lambda^2 (\|X\|^2 + \|X_{t_0}\|^2 + \|\bar{X}\|^2 + t^2)}{\sigma(t)^4} dt \right) \\ & = \mathbb{E} \exp \left(\frac{\lambda^2 (\|X\|^2 + \|X_{t_0}\|^2 + \|\bar{X}\|^2)}{2} \int_{t_1}^T \frac{dt}{\sigma(t)^4} + \int_{t_1}^T \frac{\lambda^2 t^2 dt}{2\sigma(t)^4} \right) \\ & = \mathbb{E} \exp \left(\frac{\lambda^2 (\|X\|^2 + \|X_{t_0}\|^2 + \|\bar{X}\|^2)}{2} \int_{t_1}^T \frac{dt}{\sigma(t)^4} + \int_{t_1}^T \frac{\lambda^2 t^2 dt}{2\sigma(t)^4} \right) \\ & \leq \mathbb{E} \exp \left(\frac{\lambda^2 T (\|X\|^2 + \|X_{t_0}\|^2 + \|\bar{X}\|^2)}{\delta^2} + \frac{\lambda^2 T^3}{2\delta^2} \right) < \infty, \end{aligned}$$

where the last inequality uses the fact that the concatenated vector $[X^\top, X_{t_0}^\top, \bar{X}^\top]^\top$ is sub-gaussian and the second-to-last inequality uses the bound

$$\int_{t_1}^T \frac{dt}{\sigma(t)^4} \leq \frac{T}{\sigma(t_1)^4} = \frac{T}{(1 - e^{-2t_1})^2} = \frac{T}{\delta^2}.$$

For the second inequality, we apply Jensen's inequality to upper-bound it by the first bound:

$$\begin{aligned}
 & \mathbb{E}_{t,p_t} \exp \left(\frac{1}{2} \int_{t_1}^T \left\| \mathbb{E}[s_\theta(X_t, \hat{Z}, G, t) | X_t] - \nabla_x \log p_t(X_t) \right\|^2 dt \right) \\
 &= \mathbb{E}_{t,p_t} \exp \left(\frac{1}{2} \int_{t_1}^T \left\| \mathbb{E}[s_\theta(X_t, \hat{Z}, G, t) - \nabla_x \log p_t(X_t | X) | X_t] \right\|^2 dt \right) \\
 &\leq \mathbb{E}_{t,p_t(X_t)} \mathbb{E}_{p_{0|t}(X|X_t)} \exp \left(\frac{1}{2} \int_{t_1}^T \left\| s_\theta(X_t, \hat{Z}, G, t) - \nabla_x \log p_t(X_t | X) \right\|^2 dt \right) < \infty,
 \end{aligned}$$

where first equality uses the property of the score function:

$$\mathbb{E}[\nabla_x \log p_t(X_t | X) | X_t] = \nabla_x \log p_t(X_t) + \mathbb{E}[\nabla_x \log p_t(X | X_t) | X_t] = \nabla_x \log p_t(X_t).$$

C.2. Proof of Lemma C.4

Let $\mathbb{P}_{[0,T]}$ be the measure on $B_{[0,T]}^{\leftarrow} := \{B_t^{\leftarrow}\}_{t \in [0,T]}$ and $\frac{d\mathbb{P}}{d\mathbb{Q}}$ be the Radon–Nikodym derivative between \mathbb{P} and \mathbb{Q} . Further, define another measure $\mathbb{Q}_{[0,T]}$ as

$$\frac{d\mathbb{Q}_t}{d\mathbb{P}_t} = \exp \left(\int_0^t b_s dB_s^{\leftarrow} - \frac{1}{2} \int_0^t \|b_s\|^2 ds \right),$$

where $b_t := s_\theta(X_t, \hat{Z}, G, t) - \nabla_x \log p_t(X_t | X) = \frac{\bar{s}_\theta(\hat{Z}, G, t) - X}{\sigma(t)^2}$.

By Lemma C.3, the Novikov condition holds and we can apply Girsanov's theorem (Girsanov, 1960) on \mathbb{Q} , which implies if we change the measure of B_t from \mathbb{P}_t to \mathbb{Q}_t , then there exists a Brownian motion β_t such that

$$B_t = \beta_t + \int_0^t \theta_\tau d\tau,$$

and thus under \mathbb{Q}_t ,

$$\begin{aligned}
 dX_t^{\leftarrow} &= (X_t^{\leftarrow} + 2\nabla_x \log p_{T-t}(X_t^{\leftarrow} | X)) dt + \sqrt{2} dB_t^{\leftarrow} \\
 &= \left(X_t^{\leftarrow} + 2s_\theta(X_t, \hat{Z}, G, t) \right) dt + \sqrt{2} d\beta_t.
 \end{aligned}$$

Therefore, the discrepancy in distributions of X_t^{\leftarrow} 's between the estimated and true reverse process is captured by the discrepancy between measures $\mathbb{P}_{[0,T]}$ and $\mathbb{Q}_{[0,T]}$ for the driving process $B_{[0,T]}^{\leftarrow}$.

To assess such the difference between the two measures, notice that

$$\begin{aligned}
 & \mathbb{E}_{\mathbb{Q}_t} [\exp \langle r, B_t^{\leftarrow} \rangle | X] \\
 &= \mathbb{E}_{\mathbb{P}_t} \left[\exp \left(\langle r, B_t^{\leftarrow} \rangle + \int_0^t \langle b_\tau, dB_\tau^{\leftarrow} \rangle - \frac{1}{2} \int_0^t \|b_\tau\|^2 d\tau \right) \middle| X \right] \\
 &= \mathbb{E}_{\mathbb{P}_t} \left[\exp \left[\int_0^t \left\langle r + \frac{\bar{s}_\theta(\hat{Z}, G, \tau) - X}{\sigma(\tau)^2}, dB_\tau^{\leftarrow} \right\rangle - \int_0^t \frac{1}{2\sigma(\tau)^4} \|\bar{s}_\theta(\hat{Z}, G, \tau) - X\|^2 d\tau \right] \middle| X \right] \\
 &= \exp \left(\frac{1}{2} \int_0^t \left\| r + \frac{\bar{s}_\theta(\hat{Z}, G, \tau) - X}{\sigma(\tau)^2} \right\|^2 - \frac{1}{2\sigma(\tau)^4} \|\bar{s}_\theta(\hat{Z}, G, \tau) - X\|^2 d\tau \right) \\
 &= \exp \left(\frac{r^2 t}{2} + \int_0^t \left\langle r, \frac{\bar{s}_\theta(\hat{Z}, G, \tau) - X}{\sigma(\tau)^2} \right\rangle d\tau \right) = \mathbb{E}_{\mathcal{N} \left(\int_0^t \frac{\bar{s}_\theta(\hat{Z}, G, \tau) - X}{\sigma(\tau)^2} d\tau, t I_{d_X} \right)} [\exp \langle r, B_t^{\leftarrow} \rangle | X],
 \end{aligned}$$

which implies that B_t remains a Gaussian random variable under \mathbb{Q}_t with the same variance as that under \mathbb{P}_t . Therefore, we conclude that

$$\begin{aligned}\text{Var}(\hat{X}|X) &= \mathbb{E}_{X_{[0,T]} \sim \mathbb{P}_{[0,T]}, X_{[0,t_1]}^{\leftarrow} | X_T \sim \mathbb{Q}_{[t_1,T]}} \|\hat{X} - \mathbb{E}[\hat{X}|X]\|^2 \\ &= \mathbb{E}_{X_{[0,T]} \sim \mathbb{P}_{[0,T]}, X_{[0,t_1]}^{\leftarrow} | X_T \sim \mathbb{P}_{[t_1,T]}} \|\hat{X} - \mathbb{E}[\hat{X}|X]\|^2 = \delta d_X.\end{aligned}$$

To bound the second moment, define measure $\mathbb{Q}'_{[0,T]}$ on $B_{[0,T]}^{\leftarrow}$

$$\frac{d\mathbb{Q}'_t}{d\mathbb{P}_t} = \exp\left(\int_0^t b'_s dB_s^{\leftarrow} - \frac{1}{2} \int \|b'_s\|^2 ds\right),$$

where $b'_t := \mathbb{E}[s_\theta(X_t, \hat{Z}, G, t) | X_t] - \nabla_x \log p_t(X_t)$. By Lemma C.3, Novikov condition holds for $\mathbb{E}[s_\theta(X_t, \hat{Z}, G, t) | X_t]$, and therefore by Girsanov's theorem and Cauchy-Schwarz inequality,

$$\begin{aligned}\mathbb{E}_{\mathbb{Q}'} \|\hat{X}\|^2 &= \mathbb{E}_{\mathbb{P}} \frac{d\mathbb{Q}'}{d\mathbb{P}} \|\hat{X}\|^2 \\ &\leq \mathbb{E}_{\mathbb{P}}^{1/2} \|\hat{X}\|^4 \mathbb{E}_{\mathbb{P}'}^{1/2} \left(\frac{d\mathbb{Q}'}{d\mathbb{P}}\right)^2 \\ &\leq C \sigma_X^2 d_X \mathbb{E}_{\mathbb{P}}^{1/2} \left(\frac{d\mathbb{Q}'}{d\mathbb{P}}\right)^2 \\ &= C \sigma_X^2 d_X \mathbb{E}_{\mathbb{P}}^{1/2} \exp\left(\int_{t_1}^T \|\mathbb{E}[s_\theta(X_t, \hat{Z}, G, t) | X_t] - \nabla_x \log p_t(X_t)\|^2 dt\right) \\ &\leq C_2 \sigma_X^2 d_X,\end{aligned}$$

for $C, C_2 > 0$, where the second-to-last inequality uses Girsanov's theorem as follows:

$$\begin{aligned}\mathbb{E}_{\mathbb{P}} \left(\frac{d\mathbb{Q}'}{d\mathbb{P}}\right)^2 &= \mathbb{E}_{\mathbb{P}} \exp\left(2 \int_{t_1}^T b'_t dB_t - \int_{t_1}^T \|b'_t\|^2 dt\right) \\ &= \mathbb{E}_{\mathbb{P}} \exp\left(\int_{t_1}^T \|b'_t\|^2 dt\right).\end{aligned}$$

C.3. Proof of Lemma C.5

We provide a sketch of the proof and refer the readers to (Chae & Walker, 2020) for more details. The main strategy is to upper bound $d_{\text{TV}}(p, q)$ by a lower bound of $\mathcal{W}_2(p, q)$ in terms of a metric more closely related to d_{TV} called the *Lévy-Prokhorov metric* defined as

$$d_P(p, q) := \inf\{\epsilon > 0 | p(A) < q(A + B^\epsilon) + \epsilon \text{ and } q(A) < p(A + B^\epsilon) + \epsilon, \forall A \in \mathcal{B}(\mathcal{X})\},$$

where B^ϵ is the ϵ -ball, the set $A + B$ is the Minkowski sum of A, B and $\mathcal{B}(\mathcal{X})$ is the Borel sigma algebra on the sample space \mathcal{X} . This is indeed a lower bound of \mathcal{W}_2 (and \mathcal{W}_r in general) since for any $A : p(A) < q(A)$,

$$p(A + B^\epsilon) \geq \Pr_{X \sim p, Y \sim q} [Y \in A, \|X - Y\| \leq \epsilon] \quad (30)$$

$$\geq q(A) - \Pr_{X \sim p, Y \sim q} [\|X - Y\| > \epsilon] \quad (31)$$

$$\geq q(A) - \frac{\mathcal{W}_2^2(p, q)}{\epsilon}. \quad (32)$$

Therefore, $\epsilon = \mathcal{W}_2(p, q)$ is sufficient for $p(A + B^\epsilon) > q(A)$ and thus $\mathcal{W}_2(p, q) \geq d_P$ by the definition of d_P .

To upper bound d_{TV} , notice that

$$\begin{aligned} d_{\text{TV}}(p,q) &= \sup_{A \subseteq \mathcal{B}(\mathcal{X})} p(A) - q(A) \\ &:= p(A^*) - q(A^*) \\ &\leq q(A^* + B^{d_P(p,q)}) + d_P(p,q) - q(A^*) \\ &\leq q((A^* + B^{d_P(p,q)}) \setminus A^*) + d_P(p,q). \end{aligned}$$

Finally, by the smoothness properties of q , we can upper-bound the first term on the RHS and obtain an upper-bound of d_{TV} in terms of d_P , which in terms provides an upper-bound of $d_{\text{TV}}(p,q)$ in terms of $\mathcal{W}_2(p,q)$.

D. Content distortion example

Consider the following example.

Example 1. Under the same independence relations in Assumption 2.1 with $t_0 = t_1 = 0$, let $X := [X(1), X(2)] = [Z + \Xi(1), G + \Xi(2)]$ with $Z \sim \mathcal{N}(0,1)$, $G \sim \text{Unif}\{-1,1\}$, $\Xi(1), \Xi(2) \sim \mathcal{N}(0,\epsilon)$. Further, let the inputs to the score function be $\hat{Z} = X(1)\text{sign}(G)$. Then we have \hat{Z} and G are independent since for any $z \in \mathbb{R}, g < 0$,

$$\begin{aligned} \Pr[\hat{Z} \geq z | G = g] &= \Pr[Z + \Xi(1) \leq -z | G = g] \\ &= \Pr[-Z - \Xi(1) \geq z] \\ &= Q\left(\frac{z}{1+\epsilon^2}\right) = \Pr[\hat{Z} \geq z], \end{aligned}$$

where $Q(\cdot)$ is the tail distribution of a standard Gaussian variable and the last equality uses the symmetry $X(1) \stackrel{d.}{=} -X(1)$. Therefore, $I(\hat{Z}; G) = 0$ which automatically implies \hat{Z}, G are (ϵ, ψ) -editable for any ϵ, ψ . It remains to show that $d_{\text{TV}}(p_\Psi, p) \xrightarrow{\epsilon \rightarrow 0} 0$ for the ML estimator $\Psi := \hat{\psi}(\hat{Z}, G) := \mathbb{E}[X | \hat{Z}, G]$. To this end, notice that

$$\Psi := \hat{\psi}(\hat{Z}, G) = \mathbb{E}[X | Z + \Xi(1), G] = \left[\frac{1}{1+\epsilon} (Z + \Xi(1)), G \right] = \left[\frac{1}{1+\epsilon} \hat{Z} \text{sign}(G), G \right],$$

and use Pinsker's inequality, we have

$$\begin{aligned} d_{\text{TV}}(p_\Psi, p) &\leq \sqrt{\frac{D_{\text{KL}}(p_\Psi || p)}{2 \log 2}} \\ &= \sqrt{\frac{\frac{\epsilon}{2} + \frac{1}{2} \log(1+\epsilon)}{2 \log 2}} \leq \sqrt{\frac{\epsilon}{2 \log 2}} \xrightarrow{\epsilon \rightarrow 0} 0. \end{aligned}$$

Therefore, \hat{Z}, G is $(\sqrt{\epsilon/2 \log 2}, \hat{\psi}, p)$ -editable.

However, we will show next that this does not imply $\hat{Z} | Z$ and G are $(c, \hat{\psi}, p_{X|Z})$ -editable for some constant $c \gg 0$. In view of the facts that

$$\begin{aligned} \text{sign}(G) &\sim \text{Unif}\{-1,1\}, \\ \hat{Z} &\sim \mathcal{N}(0, 1+\epsilon), \\ \hat{Z} | Z &\sim \frac{1}{2} \mathcal{N}(-Z, \epsilon) + \frac{1}{2} \mathcal{N}(Z, \epsilon), \\ \hat{Z} | Z, G &\sim \mathcal{N}(0, \epsilon), \end{aligned}$$

and consider i.i.d copies $(\hat{Z}_1 | Z, G_1) = ((Z + \Xi_1(1))\text{sign}(G_1), G_1)$ and $(\hat{Z}_2 | Z, G_2) = ((Z + \Xi_2(1))\text{sign}(G_2), G_2)$, then we have

$$\begin{aligned} \Psi_1 | Z &:= \hat{\psi}(\hat{Z}_1, G_1) | Z = \left[\frac{1}{1+\epsilon} (Z + \Xi_1(1)), G_1 \right] \\ \Psi_{12} | Z &:= \hat{\psi}(\hat{Z}_1, G_2) | Z = \left[\frac{1}{1+\epsilon} (Z + \Xi_1(1))\text{sign}(G_1 G_2), G_2 \right], \end{aligned}$$

As a result, by the variational characterization of d_{TV} :

$$\begin{aligned} d_{\text{TV}}(p_{\Psi_1|Z}, p_{\Psi_{12}|Z}) &= \sup_{|f| \leq 1/2} \mathbb{E}_{p_{\Psi_1|Z}} f(\Psi_1) - \mathbb{E}_{p_{\Psi_{12}|Z}} f(\Psi_{12}) \geq \\ &= \mathbb{E} \left[\frac{\Psi_1(1)}{2|Z|/(1+\epsilon) + 2\epsilon} \mathbb{1}_{|\Psi_1(1) - Z/(1+\epsilon)| \leq \epsilon} \middle| Z \right] - \mathbb{E} \left[\frac{\Psi_{12}(1)}{2|Z|/(1+\epsilon) + 2\epsilon} \mathbb{1}_{|\Psi_{12} - Z/(1+\epsilon)| \leq \epsilon} \middle| Z \right] = \\ &= \frac{(1-2Q(1))(1-2Q(1)(1-2Q(1)))Z}{2|Z|/(1+\epsilon) + 2\epsilon} \xrightarrow{\epsilon \rightarrow 0} (1/2 - Q(1))(1-2Q(1)(1-2Q(1))) \approx 0.27 =: c. \end{aligned}$$

Therefore, $(\hat{Z}|Z, G)$ is not $(0.27, \hat{\psi}, p_{X|Z})$ -editable.

E. Proof of Theorem 3.8

By data processing inequality on the Markov chain $z_{\phi^*}(X^c)|Z - X^c|Z - G|Z$,

$$\begin{aligned} I(z_{\phi^*}(X^c); G|Z) &\leq I(X^c; G|Z) \\ &\leq I(Z, G^c; G|Z) = I(G^c; G|Z) \leq \epsilon. \end{aligned} \tag{33}$$

Therefore, $(\hat{Z}|Z, G)$ is ϵ -disentangled.

Further, use similar techniques as in Theorem 3.6, we can prove that

$$\tilde{L}_{\text{mm}}(\theta^*, \phi^*) = O\left(\frac{\delta d_X}{T - t_1}\right).$$

To prove the first condition of editability, let $\Psi := \psi_{\theta^*}(\hat{Z}, G)$ and $\hat{Z} := z_{\phi^*}(X_{t_0}^c)$, notice that

$$\begin{aligned} d_{\text{TV}}(p_{\Psi|Z}, p_{X|Z}) &\leq d_{\text{TV}}(p_{\Psi|Z}, p_{\hat{X}|Z}) + d_{\text{TV}}(p_{\hat{X}|Z}, p_{X_{t_0}|Z}) + d_{\text{TV}}(p_{X_{t_0}|Z}, p_{X|Z}) \\ &\leq d_{\text{TV}}(p_{\Psi|Z}, p_{\hat{X}|Z}) + d_{\text{TV}}(p_{\hat{X}|X}, p_{X_{t_0}|X}) + d_{\text{TV}}(p_{X_{t_0}|\bar{X}}, p_{X|\bar{X}}), \end{aligned}$$

where the first inequality uses Jensen's inequality and the second inequality uses data processing inequality with the Markov chain $Z - \bar{X} - X - X_{t_0}$. Using Lemma C.5, we can show that the first term is bounded as

$$d_{\text{TV}}(p_{\Psi|Z}, p_{\hat{X}|Z}) = O\left(\delta^{1/4} d_X^{1/4}\right).$$

Check Novikov's condition and apply Girsanov's theorem as in Theorem 3.7, we can prove that

$$\begin{aligned} d_{\text{TV}}(p_{\hat{X}|X}, p_{X_{t_1}|X}) &= O\left(\left(\sqrt{d_X} \eta + \sigma_X^2 \eta\right) \lambda_{\Theta} \sqrt{T} + \sqrt{\tilde{L}_{\text{mm}}(\theta^*, \phi^*) T}\right) \\ &= O\left(\left(\sqrt{d_X} \eta + \sigma_X^2 \eta\right) \lambda_{\Theta} \sqrt{T} + \sqrt{\delta d_X}\right). \end{aligned}$$

Again similar to Theorem 3.7, we can bound the third term as

$$d_{\text{TV}}(p_{X_{t_1}|\bar{X}}, p_{X|\bar{X}}) = O\left(\sqrt{\lambda_s d_X \sigma_X \delta^{1/2}}\right).$$

Choosing the same step size η as in Theorem 3.7 and combining the terms yields

$$d_{\text{TV}}(p_{\Psi|Z}, p_{X|Z}) = O\left(\sqrt{\lambda_s \sigma_X \delta^{1/2} d_X}\right).$$

Set $\delta = \frac{\epsilon^2}{\lambda_s^2 \lambda_z^2 \sigma_X^2 d_X^4}$ yields the first condition of editability.

To prove the second condition, consider i.i.d copies $(\hat{Z}_1|Z, G_1)$ and $(\hat{Z}_2|Z, G_2)$ sampled from $p_{\hat{Z}|Z}$ and apply Eq. 33 with Pinsker's inequality:

$$d_{\text{TV}}(p_{\psi_{\theta^*}(\hat{Z}_1, G_1)|Z}, p_{\psi_{\theta^*}(\hat{Z}_2, G_2)|Z}) \leq d_{\text{TV}}(p_{\hat{Z}_1|Z}, p_{\hat{Z}_2|Z}) \leq \sqrt{\frac{D_{\text{KL}}(p_{\hat{Z}_1|Z} \| p_{\hat{Z}_2|Z})}{2 \log 2}} = O(\sqrt{\epsilon}).$$

As a result, we prove that $(\hat{Z}|Z, G)$ is $(O(\sqrt{\epsilon}), \psi_{\theta^*}, p_{X|Z})$ -editable.

F. Proof of Theorem 3.10

Let $s_Z^*(x) := \nabla_z \log p_{Z_t}(z(x))$ and $s_G^*(g) := \nabla_g \log p_{G_t}(g)$ and P_A be the projection matrix onto $R(A)$, then for any (s_Z, s_G, U, V) ,

$$\begin{aligned} L_0(s_Z, s_G, U, V) &:= \mathbb{E}_{t, p_t(x)} \|(P_{A_Z}(Us_Z(x, t) + Vs_G(g(x), t)) - A_Z s_Z^*(x, t)) + (P_{A_G}(Us_Z(x, t) + Vs_G(g(x), t)) - A_G s_G^*(g(x), t))\|^2 \\ &= \mathbb{E}_{t, p_t(x, t)} \left[\|P_{A_Z}(Us_Z(x, t) + Vs_G(g(x), t)) - A_Z s_Z^*(x, t)\|^2 + \right. \\ &\quad \left. \|P_{A_G}(Us_Z(x, t) + Vs_G(g(x), t)) - A_G s_G^*(g(x), t)\|^2 \right] \geq 0 = L_0(s_Z^*, s_G^*, A_Z, A_G). \end{aligned} \quad (34)$$

To analyze the equality condition, notice by the fact that minimizing over a larger set leads to smaller loss and the independence between Z_t and G_t ,

$$\begin{aligned} \mathbb{E}_{t, p_t(x)} \|P_{A_Z}(Us_Z(x, t) + Vs_G(g(x), t)) - A_Z s_Z^*(x, t)\|^2 &\geq \mathbb{E}_{t, p_t(x)} \|\mathbb{E}_{p_t(x)}[A_Z s_Z^*(x, t) | Us_Z(x, t) + Vs_G(g(x), t)] - A_Z s_Z^*(x, t)\|^2 \\ &\geq \mathbb{E}_{t, p_t(x)} \|\mathbb{E}_{p_t(x)}[A_Z s_Z^*(x, t) | Us_Z(x, t), Vs_G(g(x), t)] - A_Z s_Z^*(x, t)\|^2 \\ &= \mathbb{E}_{t, p_t(x)} \|\mathbb{E}_{p_t(x)}[A_Z s_Z^*(x, t) | Us_Z(x, t)] - A_Z s_Z^*(x, t)\|^2, \end{aligned} \quad (35)$$

with equality if and only if

$$\begin{aligned} \mathbb{E}_{p_t(x)}(s_Z^*(x, t) - P_{A_Z} Vs_G(g(x), t)) s_G(g(x), t)^\top &= 0, \forall t \in [0, T] \\ \Rightarrow P_{A_Z} Vs_G(g(x), t) &= 0, \text{a.s.}, \forall x, t. \end{aligned}$$

by the orthogonality principle. As a result, the equality of Eq. 35 is achieved if and only if

$$P_{A_Z} Us_Z(x, t) = A_Z s_Z^*(x, t), \quad (36)$$

$$P_{A_Z} Vs_G(g(x), t) = 0, \quad (37)$$

$$P_{A_G} Us_Z(x, t) + Vs_G(g(x), t) = A_G s_G^*(g(x), t), \text{a.s.}, \forall x, t. \quad (38)$$

Now, we turn our attention to the regularizer L_r and notice that for any (s_Z, U) that satisfies Eq. 36-38,

$$\begin{aligned} L_r(s_G, V) &\geq \mathbb{E}_{t, p_t(x)} \|Vs_G(g(x), t) - \nabla_x \log p_t(x)\|^2 = \mathbb{E}_{t, p_t(x)} \|Us_Z(x, t)\|^2 \\ &\geq \mathbb{E}_{t, p_t(x)} \|P_{A_Z} Us_Z(x, t)\|^2 = \mathbb{E}_{t, p_t(x)} \|A_Z s_Z^*(x, t)\|^2, \end{aligned}$$

where both equalities are achieved if and only if

$$\begin{aligned} V^\top V &= \mathbb{E} s_G(g, t) s_G(g, t)^\top, \\ \|P_{A_G} Us_Z(x, t)\| &= 0, \text{a.s.}, \forall x, t. \end{aligned}$$

Finally, we shall show that for any (A_Z, A_G, s_Z^*, s_G^*) , there exists some optimal solution (U, V, s_Z, s_G) such that

$$Us_Z = A_Z s_Z^*, Vs_G = A_G s_G^*, \quad (39)$$

$$V^\top V = \mathbb{E} s_G(g, t) s_G(g, t)^\top, L_b(s_Z, s_G, U, V) = 0. \quad (40)$$

To this end, consider the SVD of the operator $A_Z s_Z^*(x, t) + A_G s_G^*(g(x), t)$ as

$$\begin{aligned} \forall x, t, \quad A_Z s_Z^*(x, t) + A_G s_G^*(g(x), t) &= [\Phi_Z, \Phi_G] \begin{bmatrix} \Sigma_Z & \mathbf{0}_{d_Z \times d_G} \\ \mathbf{0}_{d_G \times d_Z} & \Sigma_G \end{bmatrix} \begin{bmatrix} \Psi_Z(x, t) \\ \Psi_G(g(x), t) \end{bmatrix} \\ \text{s.t.} \quad \Sigma_Z &= \text{diag}(\sigma_1(s_Z^*), \dots, \sigma_{d_Z}(s_Z^*)), \Sigma_G = \text{diag}(\sigma_1(s_G^*), \dots, \sigma_{d_G}(s_G^*)), \\ &\begin{bmatrix} \Phi_Z^\top \\ \Phi_G^\top \end{bmatrix} [\Phi_Z, \Phi_G] = \mathbb{E} \begin{bmatrix} \Psi_Z(x, t) \\ \Psi_G(g(x), t) \end{bmatrix} [\Psi_Z(x, t)^\top, \Psi_G(g(x), t)^\top] = I_{d_X} \end{aligned}$$

Set $U := \Phi_Z \Sigma_Z^{1/2}, s_Z^*(x, t) = \Sigma_Z^{1/2} \Psi_Z(x, t), V := \Phi_G \Sigma_G^{1/2}, s_G^*(x, t) = \Sigma_G^{1/2} \Psi_G(x, t)$, we have (U, V, s_Z, s_G) satisfies Eq. 39. Further, notice that

$$\begin{aligned} U^\top U &= \Sigma_Z^{1/2} \Phi_Z^\top \Phi_Z \Sigma_Z^{1/2} = \Sigma_Z = \Sigma_Z^{1/2} \mathbb{E} \Psi_Z(x, t) \Psi_Z(x, t)^\top \Sigma_Z^{1/2} = \mathbb{E} s_Z(x, t) s_Z(x, t)^\top, \\ U^\top V &= \Sigma_Z^{1/2} \Phi_Z^\top \Phi_G \Sigma_G^{1/2} = 0 = \Sigma_Z^{1/2} \mathbb{E} \Psi_Z(x, t) \Psi_G(x, t)^\top \Sigma_G^{1/2} = \mathbb{E} s_Z(x, t) s_G(g(x), t)^\top, \\ V^\top V &= \Sigma_G^{1/2} \Phi_G^\top \Phi_G \Sigma_G^{1/2} = \Sigma_G = \Sigma_G^{1/2} \mathbb{E} \Psi_G(g, t) \Psi_G(g, t)^\top \Sigma_G^{1/2} = \mathbb{E} s_G(g, t) s_G(g, t)^\top. \end{aligned}$$

Therefore, (U, V, s_Z, s_G) also satisfies Eq. 40.

G. Proof of Theorem 3.11

Define matrices

$$\begin{aligned} W(x, t) &:= \begin{bmatrix} U & V \\ s_Z(x, t)^\top & s_G(g(x), t)^\top \end{bmatrix} =: \begin{bmatrix} \tilde{U} \\ s(x, t)^\top \end{bmatrix}, \\ \tilde{W}(x, t) &:= \begin{bmatrix} W(x, t) \\ \mathbf{0}_{(d_X+1) \times d_Z}, \sqrt{\frac{\lambda_r}{3}} W_G(x, t) \end{bmatrix}, \\ W^*(x, t) &:= \begin{bmatrix} A_Z & A_G \\ s_Z^*(x, t)^\top & s_G^*(g(x), t)^\top \end{bmatrix} =: \begin{bmatrix} A \\ s^*(x, t)^\top \end{bmatrix}, \\ \tilde{W}^*(x, t) &:= \begin{bmatrix} W^*(x, t) \\ \mathbf{0}_{(d_X+1) \times d_Z}, \sqrt{\frac{\lambda_r}{3}} W_G^*(x, t) \end{bmatrix}, \\ N(x, t) &:= W(x, t) W(x, t)^\top, N^*(x, t) := W^*(x, t) W^*(x, t)^\top, \\ \tilde{N}(x, t) &:= \tilde{W}(x, t) \tilde{W}(x, t)^\top, \tilde{N}^*(x, t) := \tilde{W}^*(x, t) \tilde{W}^*(x, t)^\top. \end{aligned}$$

Further, define the *direction of improvement* of $W(x, t)$'s and $\tilde{W}(x, t)$'s respectively as

$$\begin{aligned} \Delta(x, t) &:= W(x, t) - W^*(x, t) R, \\ \tilde{\Delta}(x, t) &:= \tilde{W}(x, t) - \tilde{W}^*(x, t) R, \\ R &:= \underset{R: R^\top R = R R^\top = I_{d_X}}{\operatorname{argmin}} \mathbb{E}_{t, p_t(x)} [\|\tilde{W}(x, t) - \tilde{W}^*(x, t) R\|^2] \\ &:= \underset{R: R^\top R = R R^\top = I_{d_X}}{\operatorname{argmin}} \mathbb{E}_{t, p_t(x)} \left[\|W(x, t) - W^*(x, t) R\|^2 + \frac{\lambda_r}{3} \|W_G(x, t) - W_G^*(x, t) R_G\|^2 \right], \end{aligned}$$

where for any matrices M , define M_Z to be its first d_Z columns and M_G to be its $(d_Z + 1)$ -th through d_X -th columns.

For any set of matrices $\{C(y, t)\}_{y \in \mathcal{Y}, t \in [0, T]}$ and probability measures $q_t(y)$'s, define random matrix \mathbf{C} as

$$\mathbf{C} = C(y, t) \quad \text{w.p.} \quad p(t) q_t(y),$$

where $p(t)$ is some fixed distribution of the diffusion time t . Further, define a blockwise representation of \mathbf{C} as

$$\mathbf{C} = \begin{bmatrix} \mathbf{C}_0 & \mathbf{c}_2 \\ \mathbf{c}_1 & c_{11} \end{bmatrix},$$

where \mathbf{C}_0 is \mathbf{C} deleting the last row and column.

For a pair of random matrices $(\mathbf{C}_1, \mathbf{C}_2)$, define

$$\mathbf{C}_1 \mathbf{C}_2 = C_1(y, t) C_2(y', t) \quad \text{w.p.} \quad p(t) q_t(y) q_t(y').$$

Next, let $[\mathbf{A}, \mathbf{B}]_{\mathcal{K}}$ denote the *bilinear form* between random matrices (\mathbf{A}, \mathbf{B}) weighted by the operator \mathcal{K} , and $\langle \mathbf{A}, \mathbf{B} \rangle_{\mathcal{K}} = [\mathbf{A}, \mathbf{B}]_{\mathcal{K}}$ be an *inner product* between random matrices \mathbf{A} and \mathbf{B} if \mathcal{K} is positive definite. Then we define the

following bilinear forms between random matrices (\mathbf{A}, \mathbf{B}) with weight operators \mathcal{I}, \mathcal{G} and \mathcal{H}_0 respectively as

$$\begin{aligned}\langle \mathbf{A}, \mathbf{B} \rangle_{\mathcal{I}} &:= \mathbb{E}_{t, q_t(y)} \langle A(y, t), B(y, t) \rangle \\ [\mathbf{A}, \mathbf{B}]_{\mathcal{G}} &:= \mathbb{E}_{t, q_t(y)} [\langle A_0(y, t), B_0(y, t) \rangle + a_{11}(y, t)b_{11}(y, t) - \\ &\quad \langle a_1(y, t), b_1(y, t) \rangle - \langle a_2(y, t), b_2(y, t) \rangle], \\ [\mathbf{A}, \mathbf{B}]_{\mathcal{H}_0} &:= \mathbb{E}_{t, q_t(y)} [\langle a_1(y, t), b_1(y, t) \rangle + \langle a_2(y, t), b_2(y, t) \rangle].\end{aligned}$$

It can be verified that $\langle \cdot, \cdot \rangle_{\mathcal{I}}$ is indeed an inner product satisfying properties such as conjugate symmetry, linearity in the first argument and positive definiteness, and therefore we can define the norm $\|\mathbf{A}\|_{\mathcal{I}} := \langle \mathbf{A}, \mathbf{A} \rangle_{\mathcal{I}}$. In addition, it can be checked that $[\cdot, \cdot]_{\mathcal{G}}$ and $[\cdot, \cdot]_{\mathcal{H}_0}$ are conjugate symmetric and linear in the first argument. An important relation we use repeatedly later is the fact that

$$2[\mathbf{C}, \mathbf{C}]_{\mathcal{H}_0} + [\mathbf{C}, \mathbf{C}]_{\mathcal{G}} = [\mathbf{C}, \mathbf{C}]_{2\mathcal{H}_0 + \mathcal{G}} = \|\mathbf{C}\|_{\mathcal{I}}^2, \quad (41)$$

since

$$[\mathbf{C}, \mathbf{C}]_{2\mathcal{H}_0 + \mathcal{G}} = 2(\|\mathbf{c}_1\|_{\mathcal{I}}^2 + \|\mathbf{c}_2\|_{\mathcal{I}}^2) + \|\mathbf{C}_0\|_{\mathcal{I}}^2 + \|c_{11}\|_{\mathcal{I}}^2 - \|\mathbf{c}_1\|_{\mathcal{I}}^2 - \|\mathbf{c}_2\|_{\mathcal{I}}^2 = \|\mathbf{C}\|_{\mathcal{I}}^2.$$

Under these definitions, we prove the following Lemma in Section G.1.

Lemma G.1. *Let $\tilde{L}^{\lambda_r}(\mathbf{W}) := L^{\lambda_r}(s_Z, s_G, U, V)$ with $\lambda_r = 3$, and \mathbf{W} be an approximate critical point of $\tilde{L}(\mathbf{W})$ so that $\langle \nabla \tilde{L}(\mathbf{W}), \Delta' \rangle_{\mathcal{I}} \leq \epsilon \|\Delta'\|_{\mathcal{I}}$ for any random matrix Δ' . Then the following holds*

$$[\Delta, \Delta]_{\nabla^2 \tilde{L}^{\lambda_r}(\mathbf{W})} \leq \|\tilde{\Delta} \tilde{\Delta}^\top\|_{\mathcal{I}}^2 - 3\|\tilde{\mathbf{N}} - \tilde{\mathbf{N}}^*\|_{\mathcal{I}}^2 + \epsilon \|\Delta\|_{\mathcal{I}}, \quad (42)$$

where $\nabla^2 f(\mathbf{W})$ denotes the Hessian operator of the functional f at \mathbf{W} .

To proceed, we use the following lemma for random matrices analogous to Lemma 40 and 41 in (Ge et al., 2017) and defer its proof to Section G.2.

Lemma G.2. *If $\mathbb{E}_{p(\mathbf{X})}[U(\mathbf{X})^\top Y(\mathbf{X})]$ is a positive semi-definite (PSD) matrix, then for independent, identically distributed random variables \mathbf{X}, \mathbf{X}' ,*

$$\begin{aligned}\mathbb{E}\|U(\mathbf{X})U(\mathbf{X}')^\top - Y(\mathbf{X})Y(\mathbf{X}')^\top\|^2 \geq \\ \max\left\{\frac{1}{2}\mathbb{E}\|(U(\mathbf{X}) - Y(\mathbf{X}))(U(\mathbf{X}') - Y(\mathbf{X}'))^\top\|^2, 2(\sqrt{2} - 1)\mathbb{E}\|(U(\mathbf{X}) - Y(\mathbf{X}))U(\mathbf{X}')^\top\|^2\right\}.\end{aligned}$$

Further, we prove in Section G.3 the following lemma showing that gradient descent converges to a local optimum of the objective \tilde{L} .

Lemma G.3. *The gradient flow equation Eq. 17-Eq. 18 converges in probability to a solution $(\hat{U}, \hat{V}, \hat{\theta}_Z, \hat{\theta}_G)$ such that for some $\epsilon > 0$ and random matrix $\Delta'(t)$:*

$$\begin{aligned}\langle \nabla \tilde{L}^{\lambda_r}(\mathbf{W}), \Delta' \rangle_{\mathcal{I}} &\leq \epsilon, \\ [\Delta', \Delta']_{\nabla^2 \tilde{L}^{\lambda_r}} &\geq 0.\end{aligned}$$

To prove the theorem, first consider the singular value decomposition $\mathbb{E}_{t, p_t(x)} \tilde{W}^*(x, t)^\top \tilde{W}(x, t) =: \Phi \Sigma \Psi^\top$, and by the definition of the direction of improvement,

$$\begin{aligned}R &:= \operatorname{argmin}_{RR^\top = R^\top R = I_{d_X}} \|\tilde{\mathbf{W}} - \tilde{\mathbf{W}}^* R\|_{\mathcal{I}}^2 \\ &= \operatorname{argmax}_{RR^\top = R^\top R = I_{d_X}} \langle \tilde{\mathbf{W}}, \tilde{\mathbf{W}}^* R \rangle_{\mathcal{I}} = \operatorname{argmax}_{RR^\top = R^\top R = I_{d_X}} \langle \mathbb{E}_{t, p_t(x)} \tilde{W}^*(x, t)^\top \tilde{W}(x, t), R \rangle \\ &= \operatorname{argmax}_{RR^\top = R^\top R = I_{d_X}} \langle \Sigma, \Phi^\top R \Psi \rangle = \Phi \Psi^\top,\end{aligned}$$

where the last equality holds since $R' := \Phi^\top R\Psi$ is orthogonal, $|R'_{ii}| \leq 1$ and

$$\langle \Sigma, R' \rangle \leq \sum_{i=1}^{d_x} \Sigma_{ii},$$

with equality iff $R' = I_{d_x}$. As a result,

$$\mathbb{E}_{t, p_t(x)} \tilde{W}^\top(x, t) \tilde{W}^*(x, t) R = \Psi \Sigma \Psi^\top = \mathbb{E}_{p_t(x), t} R^\top \tilde{W}^{*\top}(x, t) \tilde{W}(x, t)$$

is PSD. Applying Lemma G.2 on $U(\mathbf{X}) = \tilde{\mathbf{W}}^* R$ and $Y(\mathbf{X}) = \tilde{\mathbf{W}}$ yields

$$\begin{aligned} & \|\tilde{\Delta} \tilde{\Delta}^\top\|_{\mathcal{I}}^2 - 3\|\tilde{\mathbf{N}} - \tilde{\mathbf{N}}^*\|_{\mathcal{I}}^2 \\ & \leq 2\|\tilde{\mathbf{N}} - \tilde{\mathbf{N}}^*\|_{\mathcal{I}}^2 - 3\|\tilde{\mathbf{N}} - \tilde{\mathbf{N}}^*\|_{\mathcal{I}}^2 = -\|\tilde{\mathbf{N}} - \tilde{\mathbf{N}}^*\|_{\mathcal{I}}^2 \leq -2(\sqrt{2}-1)\|\tilde{\Delta}\|_{\mathcal{I}}^2 \leq -0.8\|\tilde{\Delta}\|_{\mathcal{I}}^2. \end{aligned}$$

Combining this with Lemma G.1 yields

$$[\Delta, \Delta]_{\nabla^2 \tilde{L}^{\lambda_r}} \leq -0.8\|\tilde{\Delta}\|_{\mathcal{I}}^2,$$

Therefore, we have the LHS to be positive only if $\|\tilde{\Delta}\|_{\mathcal{I}} = 0$. Set $\epsilon = 0$ and applying Theorem 3.10 yields $R(U) = R(A_Z), R(V) = R(A_G)$.

G.1. Proof of Lemma G.1

It can be verified that

$$\begin{aligned} L_0(s_Z, s_G, U, V) &= \frac{1}{2}[\mathbf{N} - \mathbf{N}^*, \mathbf{N} - \mathbf{N}^*]_{\mathcal{H}_0}^2 =: \tilde{L}_0(\mathbf{W}), \\ L_b(s_Z, s_G, U, V) &= [\mathbf{N}, \mathbf{N}]_{\mathcal{G}}^2 =: \tilde{L}_b(\mathbf{W}). \end{aligned}$$

Define $\tilde{L}_r(\mathbf{W}) := L_r(s_G, U)$, then we have

$$\tilde{L}^{\lambda_r}(\mathbf{W}) = 2\tilde{L}_0(\mathbf{W}) + \frac{1}{2}\tilde{L}_b(\mathbf{W}) + \lambda_r \tilde{L}_r(\mathbf{W}) = [\mathbf{N} - \mathbf{N}^*, \mathbf{N} - \mathbf{N}^*]_{\mathcal{H}_0}^2 + \frac{1}{2}[\mathbf{N}, \mathbf{N}]_{\mathcal{G}}^2 + \lambda_r \tilde{L}_r(\mathbf{W}).$$

Then, consider the Fréchet derivative of $\tilde{L}_t(\mathbf{W})$ along $\Delta(t)$, it can be shown that

$$\begin{aligned} \langle \nabla \tilde{L}^{\lambda_r}(\mathbf{W}), \Delta \rangle_{\mathcal{I}} &= [\mathbf{N} - \mathbf{N}^*, \Delta \mathbf{W}^\top + \mathbf{W} \Delta^\top]_{2\mathcal{H}_0} + [\mathbf{N}, \Delta \mathbf{W}^\top + \mathbf{W} \Delta^\top]_{\mathcal{G}} + \lambda_r \langle \nabla \tilde{L}_r(\mathbf{W}), \Delta \rangle_{\mathcal{I}} \\ &= [\mathbf{N} - \mathbf{N}^*, \Delta \mathbf{W}^\top + \mathbf{W} \Delta^\top]_{2\mathcal{H}_0 + \mathcal{G}} + [\mathbf{N}^*, \Delta \mathbf{W}^\top + \mathbf{W} \Delta^\top]_{\mathcal{G}} + \lambda_r \langle \nabla \tilde{L}_r(\mathbf{W}), \Delta \rangle_{\mathcal{I}} \\ &= [\mathbf{N} - \mathbf{N}^*, \Delta \mathbf{W}^\top + \mathbf{W} \Delta^\top]_{2\mathcal{H}_0 + \mathcal{G}} + 2[\mathbf{N}^*, \mathbf{N}]_{\mathcal{G}} + \lambda_r \langle \nabla \tilde{L}_r(\mathbf{W}), \Delta \rangle_{\mathcal{I}}, \quad (43) \end{aligned}$$

where the last equality uses the fact that $[\mathbf{N}^*, \mathbf{W}^* \mathbf{W}^\top]_{\mathcal{G}} = [\mathbf{N}^*, \mathbf{W} \mathbf{W}^{*\top}]_{\mathcal{G}} = 0$. To see this, notice that

$$\begin{aligned} [\mathbf{N}^*, \mathbf{W}^* \mathbf{W}^\top]_{\mathcal{G}} &= \langle A A^\top, \tilde{U} A^\top \rangle + \mathbb{E}_{t, p_t(x) p_t(x')} s^*(x, t)^\top s^*(x', t) s(x, t)^\top s^*(x', t) \\ &\quad - \mathbb{E}_{t, p_t(x)} s^*(x, t)^\top A^\top \tilde{U} s^*(x, t) - \mathbb{E}_{t, p_t(x)} s^*(x, t)^\top A^\top A s(x, t) \\ &= \text{Tr}(A(A^\top A - \mathbb{E}_{t, p_t(x)}[s^*(x, t) s^*(x, t)]) \tilde{U}) + \\ &\quad \mathbb{E}_{t, p_t(x)}[s^*(x, t)^\top (\mathbb{E}_{p_t(x', t)}[s^*(x', t) s^*(x', t)^\top] - A^\top A) s(x, t)] = 0, \end{aligned}$$

where the last inequality uses the fact that $\tilde{L}_b(\mathbf{W}^*) = 0$ and therefore

$$\begin{aligned} U^{*\top} U^* &= \mathbb{E}_{t, p_t(x)} s_Z^*(x, t) s_Z^*(x, t)^\top, \\ V^{*\top} V^* &= \mathbb{E}_{t, p_t(x)} s_G^*(x, t) s_G^*(x, t)^\top. \end{aligned}$$

Similarly, we can show that $[\mathbf{N}^*, \mathbf{W} \mathbf{W}^{*\top}]_{\mathcal{G}} = 0$.

Now, consider the Hessian of \tilde{L} along Δ by taking the Fréchet derivative of Eq. 43,

$$\begin{aligned} [\Delta, \Delta]_{\nabla^2 \tilde{L}^{\lambda_r}} &= [\Delta \mathbf{W}^\top + \mathbf{W} \Delta^\top, \Delta \mathbf{W}^\top + \mathbf{W} \Delta^\top]_{2\mathcal{H}_0 + \mathcal{G}}^2 + 2[\mathbf{N} - \mathbf{N}^*, \Delta \Delta^\top]_{2\mathcal{H}_0 + \mathcal{G}} + \\ &\quad 2[\mathbf{N}^*, \mathbf{W} \Delta^\top + \Delta \mathbf{W}^\top]_{\mathcal{G}} + \lambda_r [\Delta, \Delta]_{\nabla^2 \tilde{L}_r(\mathbf{W})} \\ &= \|\Delta \mathbf{W}^\top + \mathbf{W} \Delta^\top\|_{\mathcal{I}}^2 + 2[\mathbf{N} - \mathbf{N}^*, \Delta \Delta^\top]_{2\mathcal{H}_0 + \mathcal{G}} + 2[\mathbf{N}^*, \mathbf{W} \Delta^\top + \Delta \mathbf{W}^\top]_{\mathcal{G}} + \lambda_r [\Delta, \Delta]_{\nabla^2 \tilde{L}_r(\mathbf{W})}, \end{aligned}$$

where we use Eq. 41 in the last equality. For the first term of the right-hand side, by the choice of Δ ,

$$\begin{aligned} \|\Delta \mathbf{W}^\top + \mathbf{W} \Delta^\top\|_{\mathcal{I}}^2 &= \|\mathbf{N} - \mathbf{N}^* + \Delta \Delta^\top\|_{\mathcal{I}}^2 \\ &= \|\Delta \Delta^\top\|_{\mathcal{I}}^2 + 2\langle \mathbf{N} - \mathbf{N}^*, \Delta \mathbf{W}^\top + \mathbf{W} \Delta^\top \rangle_{\mathcal{I}} - \|\mathbf{N} - \mathbf{N}^*\|_{\mathcal{I}}^2 \\ &= \|\Delta \Delta^\top\|_{\mathcal{I}}^2 + 2\langle \nabla \tilde{L}^{\lambda_r}(\mathbf{W}), \Delta \rangle_{\mathcal{I}} - 4\langle \mathbf{N}^*, \mathbf{N} \rangle_{\mathcal{G}} - 2\lambda_r \langle \nabla \tilde{L}_r(\mathbf{W}), \Delta \rangle_{\mathcal{I}} - \|\mathbf{N} - \mathbf{N}^*\|_{\mathcal{I}}^2. \end{aligned}$$

For the second term,

$$\begin{aligned} [\mathbf{N} - \mathbf{N}^*, \Delta \Delta^\top]_{2\mathcal{H}_0 + \mathcal{G}} &= [\mathbf{N} - \mathbf{N}^*, \mathbf{W} \Delta^\top + \Delta \mathbf{W}^\top]_{2\mathcal{H}_0 + \mathcal{G}} - [\mathbf{N} - \mathbf{N}^*, \mathbf{N} - \mathbf{N}^*]_{2\mathcal{H}_0 + \mathcal{G}} \\ &= \langle \nabla \tilde{L}^{\lambda_r}(\mathbf{W}), \Delta \rangle_{\mathcal{I}} - 2\langle \mathbf{N}^*, \mathbf{N} \rangle_{\mathcal{G}} - \lambda_r \langle \nabla \tilde{L}_r(\mathbf{W}), \Delta \rangle_{\mathcal{I}} - \|\mathbf{N} - \mathbf{N}^*\|_{\mathcal{I}}^2. \end{aligned} \quad (44)$$

Combined with the fact that

$$[\mathbf{N}^*, \mathbf{W} \Delta^\top + \Delta \mathbf{W}^\top]_{\mathcal{G}} = 2[\mathbf{N}^*, \mathbf{N}]_{\mathcal{G}},$$

we obtain

$$\begin{aligned} [\Delta, \Delta]_{\nabla^2 \tilde{L}^{\lambda_r}} &= \|\Delta \Delta^\top\|_{\mathcal{I}}^2 + 4\langle \nabla \tilde{L}^{\lambda_r}(\mathbf{W}), \Delta \rangle_{\mathcal{I}} - 3\|\mathbf{N} - \mathbf{N}^*\|_{\mathcal{I}}^2 \\ &\quad - 4[\mathbf{N}, \mathbf{N}^*]_{\mathcal{G}} - 4\lambda_r \langle \nabla \tilde{L}_r(\mathbf{W}), \Delta \rangle_{\mathcal{I}} + \lambda_r [\Delta, \Delta]_{\nabla^2 \tilde{L}_r(\mathbf{W})} \\ &\leq \|\Delta \Delta^\top\|_{\mathcal{I}}^2 - 3\|\mathbf{N} - \mathbf{N}^*\|_{\mathcal{I}}^2 + 4\langle \nabla \tilde{L}^{\lambda_r}(\mathbf{W}), \Delta \rangle_{\mathcal{I}} - 4\lambda_r \langle \nabla \tilde{L}_r(\mathbf{W}), \Delta \rangle_{\mathcal{I}} + \lambda_r [\Delta, \Delta]_{\nabla^2 \tilde{L}_r(\mathbf{W})} \\ &\leq \|\Delta \Delta^\top\|_{\mathcal{I}}^2 - 3\|\mathbf{N} - \mathbf{N}^*\|_{\mathcal{I}}^2 + 4\epsilon \|\Delta\|_{\mathcal{I}} - 4\lambda_r \langle \nabla \tilde{L}_r(\mathbf{W}), \Delta \rangle_{\mathcal{I}} + \lambda_r [\Delta, \Delta]_{\nabla^2 \tilde{L}_r(\mathbf{W})}, \end{aligned} \quad (45)$$

where the first inequality uses the fact that

$$\begin{aligned} [\mathbf{N}, \mathbf{N}^*]_{\mathcal{G}} &= \langle UU^\top, AA^\top \rangle + \mathbb{E}_{t, p_t(x)p_t(x')} s^*(x, t)^\top s^*(x', t) s(x, t)^\top x(x', t) - 2\mathbb{E}_{t, p_t(x)} s(x, t) U^\top A s^*(x', t) \\ &= \|U^\top A - \mathbb{E}_{t, p_t(x)} s(x, t) s^*(x, t)^\top\|^2 \geq 0, \end{aligned}$$

and the second inequality uses the condition that \mathbf{W} is an approximate critical point of $\tilde{L}^{\lambda_r}(\mathbf{W})$. It remains to bound the \tilde{L}_r related terms. Define $\mathbf{N}^Z := \mathbf{W}_Z \mathbf{W}_Z^\top$, $\mathbf{N}^G := \mathbf{W}_G \mathbf{W}_G^\top$ and similarly \mathbf{N}^{Z*} and \mathbf{N}^{G*} . Notice that

$$\begin{aligned} \tilde{L}_r(\mathbf{W}) &= [\mathbf{N}^G - \mathbf{N}^*, \mathbf{N}^G - \mathbf{N}^*]_{\mathcal{H}_0} + \frac{1}{2} [\mathbf{N}^G, \mathbf{N}^G]_{\mathcal{G}}^2 \\ &= [\mathbf{N}^G - \mathbf{N}^{G*}, \mathbf{N}^G - \mathbf{N}^{G*}]_{\mathcal{H}_0}^2 - 2[\mathbf{N}^G - \mathbf{N}^{G*}, \mathbf{N}^{Z*}]_{\mathcal{H}_0} + [\mathbf{N}^{Z*}, \mathbf{N}^{Z*}]_{\mathcal{H}_0} + \frac{1}{2} [\mathbf{N}^G, \mathbf{N}^G]_{\mathcal{G}} \\ &= [\mathbf{N}^G - \mathbf{N}^{G*}, \mathbf{N}^G - \mathbf{N}^{G*}]_{\mathcal{H}_0} + \frac{1}{2} [\mathbf{N}^G, \mathbf{N}^G]_{\mathcal{G}} = 2\tilde{L}_0(\mathbf{W}_G) + \frac{1}{2} \tilde{L}_b(\mathbf{W}_G) = \tilde{L}^0(\mathbf{W}_G), \end{aligned}$$

where the second-to-last equality uses the fact that

$$\begin{aligned} [\mathbf{N}^G - \mathbf{N}^{G*}, \mathbf{N}^{Z*}]_{\mathcal{H}_0} &= 2\mathbb{E}_{t, p_t(x)} \langle V s_G(g(x), t) - A_G s_G^*(g(x), t), A_Z s_Z^*(z(x), t) \rangle \\ &= 2\mathbb{E}_{t, p_t(x)} \langle V s_G(g(x), t), A_Z s_Z^*(z(x), t) \rangle \\ &= 2\langle V \mathbb{E}_{t, p_t(x)} s_G(g(x), t), A_Z \mathbb{E}_{t, p_t(x)} s_Z^*(z(x), t) \rangle = 0, \end{aligned}$$

where the second-to-last equality uses the independence of the content variable Z and the style variable G and the last equality uses the property of the score function:

$$\mathbb{E}_{p_Z} s_Z^*(z) = \mathbb{E}_{p_Z} \nabla_z \log p_Z(z) = \int \nabla_z p_Z(z) dz = \nabla_z \int p_Z(z) dz = 0.$$

Therefore, we can apply Eq. 45 with $\lambda_r = 0$ to obtain

$$[\Delta, \Delta]_{\tilde{L}_r(\mathbf{W})} = \langle \Delta_G, \Delta_G \rangle_{\tilde{L}^0(\mathbf{W}_G)} \quad (46)$$

$$\leq \|\Delta_G \Delta_G^\top\|_{\mathcal{I}}^2 - 3\|\mathbf{N}^G - \mathbf{N}^{G*}\|_{\mathcal{I}}^2 + 4\langle \nabla \tilde{L}_r(\mathbf{W}), \Delta \rangle_{\mathcal{I}}. \quad (47)$$

Plugging Eq. 46 into Eq. 45 yields

$$\begin{aligned} [\Delta, \Delta]_{\tilde{L}^{\lambda_r}(\mathbf{W})} &\leq \|\Delta \Delta^\top\|_{\mathcal{I}}^2 + \lambda_r \|\Delta_G \Delta_G^\top\|_{\mathcal{I}} - 3(\|\mathbf{N} - \mathbf{N}^*\|_{\mathcal{I}}^2 + \lambda_r \|\mathbf{N}^G - \mathbf{N}^{G*}\|_{\mathcal{I}}^2) + \epsilon \|\Delta\|_{\mathcal{I}} \\ &= \|\tilde{\Delta} \tilde{\Delta}\|_{\mathcal{I}}^2 - 3\|\tilde{\mathbf{N}} - \tilde{\mathbf{N}}^*\|_{\mathcal{I}}^2 + \epsilon \|\Delta\|_{\mathcal{I}}, \end{aligned}$$

where the last equality uses the definition of $\mathbf{N}, \mathbf{N}^*, \Delta$ and Δ^* . For example, for \mathbf{N} , we have

$$\begin{aligned} \|\tilde{\mathbf{N}}\|_{\mathcal{I}}^2 &= \left\| \begin{bmatrix} \mathbf{W}\mathbf{W}^\top & \sqrt{\frac{\lambda_r}{3}} \mathbf{W}_G \mathbf{W}_G^\top \\ \sqrt{\frac{\lambda_r}{3}} \mathbf{W}_G \mathbf{W}_G^\top & \frac{\lambda_r}{3} \mathbf{W}_G \mathbf{W}_G^\top \end{bmatrix} \right\|_{\mathcal{I}}^2 = \|\mathbf{N}\|_{\mathcal{I}}^2 + \lambda_r \|\mathbf{N}^G\|_{\mathcal{I}}^2 + \left(\frac{\lambda_r^2}{9} - \frac{\lambda_r}{3} \right) \|\mathbf{N}^G\|_{\mathcal{I}}^2 \\ &= \|\mathbf{N}\|_{\mathcal{I}}^2 + 3\|\mathbf{N}^G\|_{\mathcal{I}}^2, \end{aligned}$$

where in the last equality we set $\lambda_r = 3$.

G.2. Proof of Lemma G.2

First, let $\Delta(\mathbf{X}) := U(\mathbf{X}) - Y(\mathbf{X})$, by definition,

$$\mathbb{E} \Delta(\mathbf{X})^\top U(\mathbf{X}) = \mathbb{E} U(\mathbf{X})^\top U(\mathbf{X}) - \mathbb{E} Y(\mathbf{X})^\top U(\mathbf{X}) = \mathbb{E} U(\mathbf{X})^\top \Delta(\mathbf{X}),$$

and

$$\mathbb{E} \|U(\mathbf{X})U(\mathbf{X}')^\top - Y(\mathbf{X})Y(\mathbf{X}')^\top\|^2 = \mathbb{E} \|\Delta(\mathbf{X})U(\mathbf{X}')^\top + U(\mathbf{X})\Delta(\mathbf{X}')^\top - \Delta(\mathbf{X})\Delta(\mathbf{X}')^\top\|^2.$$

Expanding the square norm,

$$\begin{aligned} &\mathbb{E} \|\Delta(\mathbf{X})U(\mathbf{X}')^\top + U(\mathbf{X})\Delta(\mathbf{X}')^\top - \Delta(\mathbf{X})\Delta(\mathbf{X}')^\top\|^2 \\ &= 2\mathbb{E} \|\Delta(\mathbf{X})U(\mathbf{X}')^\top\|^2 + \mathbb{E} \|\Delta(\mathbf{X})\Delta(\mathbf{X}')^\top\|^2 + 2\langle \mathbb{E} \Delta(\mathbf{X})^\top U(\mathbf{X}), \mathbb{E} U(\mathbf{X})^\top \Delta(\mathbf{X}) \rangle - \\ &\quad 2\langle \mathbb{E} \Delta(\mathbf{X})U(\mathbf{X}')^\top + U(\mathbf{X})\Delta(\mathbf{X}')^\top, \mathbb{E} \Delta(\mathbf{X})\Delta(\mathbf{X}')^\top \rangle \\ &= \mathbb{E} \|\Delta(\mathbf{X})\Delta(\mathbf{X}')^\top\|^2 + 2\langle \mathbb{E} \Delta(\mathbf{X})^\top U(\mathbf{X}), \mathbb{E} \Delta(\mathbf{X})^\top \Delta(\mathbf{X}) \rangle + 2\|\mathbb{E} \Delta(\mathbf{X})^\top U(\mathbf{X})\|^2 - \\ &\quad 4\langle \mathbb{E} \Delta(\mathbf{X})^\top U(\mathbf{X}), \mathbb{E} \Delta(\mathbf{X})^\top \Delta(\mathbf{X}) \rangle \\ &= \frac{1}{2} \|\mathbb{E} \Delta(\mathbf{X})^\top \Delta(\mathbf{X})\|^2 + 2\langle \mathbb{E} U(\mathbf{X})^\top Y, \mathbb{E} \Delta(\mathbf{X})^\top \Delta(\mathbf{X}) \rangle + \\ &\quad \|\sqrt{2}U(\mathbf{X})^\top \Delta(\mathbf{X}) - \frac{1}{\sqrt{2}}\Delta(\mathbf{X})^\top \Delta(\mathbf{X})\|^2 \\ &\geq \frac{1}{2} \|\mathbb{E} \Delta(\mathbf{X})^\top \Delta(\mathbf{X})\|^2 = \frac{1}{2} \mathbb{E} \|(U(\mathbf{X}) - Y(\mathbf{X}))(U(\mathbf{X}') - Y(\mathbf{X}'))^\top\|^2, \end{aligned}$$

where the second equality uses the symmetry of $\mathbb{E} U(\mathbf{X})^\top \Delta(\mathbf{X})$ the last inequality uses the PSD of $\mathbb{E} U(\mathbf{X})^\top Y(\mathbf{X})$.

Similarly,

$$\begin{aligned} &\mathbb{E} \|U(\mathbf{X})U(\mathbf{X}')^\top - Y(\mathbf{X})Y(\mathbf{X}')^\top\|^2 \\ &= 2\mathbb{E} \|\Delta(\mathbf{X})U(\mathbf{X}')^\top\|^2 + \mathbb{E} \|\Delta(\mathbf{X})\Delta(\mathbf{X}')^\top\|^2 + 2\|\mathbb{E} \Delta(\mathbf{X})^\top U(\mathbf{X})\|^2 - \\ &\quad 4\langle \mathbb{E} \Delta(\mathbf{X})^\top U(\mathbf{X}), \mathbb{E} \Delta(\mathbf{X})^\top \Delta(\mathbf{X}) \rangle \\ &= (2\sqrt{2} - 2)\mathbb{E} \|U(\mathbf{X})\Delta(\mathbf{X}')^\top\|^2 + (4 - 2\sqrt{2})\langle \mathbb{E} U(\mathbf{X})^\top Y(\mathbf{X}), \mathbb{E} \Delta(\mathbf{X})^\top \Delta(\mathbf{X}) \rangle + \\ &\quad \|\sqrt{2}\mathbb{E} U(\mathbf{X})^\top \Delta(\mathbf{X}) - \mathbb{E} \Delta(\mathbf{X})^\top \Delta(\mathbf{X})\|^2 \\ &\geq (2\sqrt{2} - 2)\mathbb{E} \|U(\mathbf{X})\Delta(\mathbf{X}')^\top\|^2 = 2(\sqrt{2} - 1)\mathbb{E} \|(U(\mathbf{X}) - Y(\mathbf{X}))U(\mathbf{X}')^\top\|^2, \end{aligned}$$

where the last inequality uses the PSD of $\mathbb{E} U(\mathbf{X})^\top Y(\mathbf{X})$.

G.3. Proof of Lemma G.3

The proof relies on the following lemma.

Lemma G.4. Define the neural tangent kernels (NTKs) for the score functions s_Z and s_G as

$$\begin{aligned} K_Z(x, t, x', t') &:= J_{\text{vec}(\theta_Z)} s_Z(x, t)^\top J_{\text{vec}(\theta_Z)} s_Z(x', t'), \\ K_G(x, t, x', t') &:= J_{\text{vec}(\theta_G)} s_G(g(x), t)^\top J_{\text{vec}(\theta_G)} s_G(g(x'), t'), \forall (x, t, x', t') \in \mathcal{X} \times [t_0, T] \times \mathcal{X} \times [t_0, T], \end{aligned}$$

where $\text{vec}(\theta)$ denotes the flattened version of the parameter θ . Further, suppose the parameters are initialized randomly as $\theta_Z(0), \theta_G(0) \sim \mathcal{N}(0, I)$. Then as $d_H \rightarrow \infty$, the NTKs K_Z and K_G converge to some kernels K_Z^* and K_G^* fixed during training. Further, define the minimal eigenvalues of operator $K : \mathcal{X} \times [t_0, T] \times \mathcal{X} \times [t_0, T] \mapsto \mathbb{R}^d$ as

$$\lambda_{\min}(K) := \inf_{v: \mathbb{E}_{t, p(x)} \|v(t, x)\|^2 = 1} \mathbb{E}_{t, t', p(x)p(x')} v(x, t)^\top K(x, t, x', t') v(x', t'),$$

then the minimal eigenvalues of K_Z^* and K_G^* satisfy $\min\{\lambda_{\min}(K_Z^*), \lambda_{\min}(K_G^*)\} > 0$.

Define the random gradient of loss L with respect to random matrix \mathbf{Y} with probability density q_t as

$$\nabla_{\mathbf{Y}} L(\mathbf{Y}) = \nabla_{Y(x, t)} L(\mathbf{Y}) \quad \text{w.p. } p(t)q_t(x).$$

By the definition of the gradient flow equations in Eq. 17-18, we have

$$\begin{aligned} \dot{\tilde{L}}(\mathbf{W}) &= \langle \nabla_{\mathbf{W}} \tilde{L}, \dot{\mathbf{W}} \rangle_{\mathcal{I}} \\ &= -\|\nabla_U \tilde{L}(\mathbf{W})\|^2 - \|\nabla_V \tilde{L}(\mathbf{W})\|^2 - \|\mathbb{E}_{t, p_t} J_{\theta_Z} s_Z \nabla_{s_Z} \tilde{L}(\mathbf{W})\|^2 - \|\mathbb{E}_{t, p_t} J_{\theta_G} s_G \nabla_{s_G} \tilde{L}(\mathbf{W})\|^2, \end{aligned}$$

where the first two terms of the RHS by the property of the gradient flow, vanishes if and only if the gradients $\nabla_U \tilde{L}(\mathbf{W})$ and $\nabla_V \tilde{L}(\mathbf{W})$ become 0. For the third term of the RHS, notice that

$$\begin{aligned} &\|\mathbb{E}_{t, p_t} J_{\theta_Z} s_Z \nabla_{s_Z} \tilde{L}(\mathbf{W})\|^2 \\ &= \mathbb{E}_{t, t', p_t(x)p_{t'}(x')} \nabla_{s_Z(x, t)} \tilde{L}(\mathbf{W})^\top J_{\theta_Z} s_Z(x, t)^\top J_{\theta_Z} s_Z(x', t')^\top \nabla_{s_Z(x', t')} \tilde{L}(\mathbf{W}) \\ &= \mathbb{E}_{t, t', p_t(x)p_{t'}(x')} \nabla_{s_Z(x, t)} \tilde{L}(\mathbf{W})^\top K_Z(x, t, x', t') \nabla_{s_Z(x', t')} \tilde{L}(\mathbf{W}) \\ &\geq \lambda_{\min}(K_Z) \int \|p(t)p_t(x) \nabla_{s_Z(x, t)} \tilde{L}(\mathbf{W})\|^2 dx dt. \end{aligned}$$

By Lemma G.4, $\lambda_{\min}(K_Z) \xrightarrow{d_H \rightarrow \infty} \lambda_{\min}(K_Z^*) > 0$ and thus the term vanishes if and only if

$$\|p(t)p_t(x) \mathbb{E}_{t, p_t(x)} \nabla_{s_Z(x, t)} \tilde{L}(\mathbf{W})\|^2 = 0, \forall x, t.$$

Similarly, we can show that the gradient flow converges only if

$$\|p(t)p_t(x) \mathbb{E}_{t, p_t(x)} \nabla_{s_G(g(x), t)} \tilde{L}(\mathbf{W})\|^2 = 0, \forall x, t.$$

For the second order condition, we use the well-known result that gradient descent (with small noise) is able to escape saddle points almost surely (Lee et al., 2016).

G.4. Proof of Lemma G.4

Due to the symmetry in their forms, it suffices to prove the statement for s_Z and K_Z and we omit the subscript when the context is clear. Large of large number and the standard theory on neural tangent kernel (Jacot et al., 2018) yields

$$K_Z \xrightarrow{d_H \rightarrow \infty} \mathbb{E}_{\text{vec}(\theta_Z) \sim \mathcal{N}(0, I)} J_{\text{vec}(\theta_Z)} s_Z(x, t)^\top J_{\text{vec}(\theta_Z)} s_Z(x', t'), \quad (48)$$

which stays fixed during training.

Define $\tilde{x} := [x^\top, \text{PE}(t)]^\top$ and $a(\tilde{x}) := (\theta^{(1)\top} \tilde{x})_+$. Later, we will slightly abuse the notation to represent (x, t) when \tilde{x} appears in the true score functions. By definition,

$$\begin{aligned} K_Z^{(2)}(x, t, x', t') &= \mathbb{E}_{\mathcal{N}(0, I)} J_{\text{vec}(\theta_Z^{(2)})} s_Z(\tilde{x})^\top J_{\text{vec}(\theta_Z^{(2)})} s_Z(\tilde{x}') \\ &= \mathbb{E}_{\mathcal{N}(0, I)} J_{\text{vec}(\theta_Z^{(2)})} \theta_Z^{(2)} a(\tilde{x})^\top J_{\text{vec}(\theta_Z^{(2)})} \theta_Z^{(2)} a(\tilde{x}') \\ &= \mathbb{E}_{\mathcal{N}(0, I)} (a(\tilde{x})^\top \otimes I_{d_Z}) (a(\tilde{x}') \otimes I_{d_X}) = \mathbb{E}_{\mathcal{N}(0, I)} a(\tilde{x})^\top a(\tilde{x}') I_{d_Z}, \end{aligned}$$

where \otimes is the Kronecker product. Notice that by definition $K_Z^{(2)} \succeq 0$. Similarly,

$$\begin{aligned} K_Z^{(1)}(x, t, x', t') &:= \mathbb{E}_{\mathcal{N}(0, I)} J_{\text{vec}(\theta_Z^{(1)})} s_Z(\tilde{x})^\top J_{\text{vec}(\theta_Z^{(1)})} s_Z(\tilde{x}') \\ &= \mathbb{E}_{\mathcal{N}(0, I)} J_{a(\tilde{x})} s_Z(\tilde{x})^\top J_{\theta_Z^{(1)}} a(\tilde{x})^\top J_{\theta_Z^{(1)}} a(\tilde{x}') J_{a(\tilde{x}')} s_Z(\tilde{x}') \\ &=: (\tilde{x}^\top \tilde{x}') \mathbb{E}_{\mathcal{N}(0, I)} \theta_Z^{(2)} S^2(\tilde{x}, \tilde{x}') \theta_Z^{(2)\top}, \\ &= (\tilde{x}^\top \tilde{x}') \mathbb{E}_{\mathcal{N}(0, I)} S(\tilde{x}, \tilde{x}'), \end{aligned}$$

where we use the independence between $(\theta_Z^{(1)}, \theta_Z^{(2)})$ to cancel out $\theta_Z^{(1)}$, and

$$S_{ij}(\tilde{x}, \tilde{x}') := \begin{cases} \mathbb{1}[\theta^{(1), j\top} \tilde{x} \geq 0, \theta^{(1), j\top} \tilde{x}' \geq 0], & \text{if } i = j, \\ 0, & \text{otherwise.} \end{cases}$$

By assumption, we have $\theta^{(1), j\top} \tilde{x}$'s are zero-mean Gaussians and thus

$$\Pr[\theta^{(1), j\top} \tilde{x} \geq 0, \theta^{(1), j\top} \tilde{x}' \geq 0] = \frac{1}{4} + \frac{1}{2\pi} \arcsin\left(\frac{\tilde{x}^\top \tilde{x}'}{\|\tilde{x}\| \|\tilde{x}'\|}\right),$$

where we apply the formula for bivariate Gaussian variable (N_1, N_2) : $\Pr[N_1 \geq 0, N_2 \geq 0] = \frac{1}{4} + \arcsin(\rho)$ where $\rho := \frac{\text{Cov}(N_1, N_2)}{\sqrt{\text{Var}(N_1)\text{Var}(N_2)}}$.

As a result,

$$K_Z^{(1)}(\tilde{x}, \tilde{x}') := \frac{1}{4} \tilde{x}^\top \tilde{x}' \left[1 + \frac{2}{\pi} \arcsin\left(\frac{\tilde{x}^\top \tilde{x}'}{\|\tilde{x}\| \|\tilde{x}'\|}\right) \right] I_{d_Z}.$$

Notice that $K_Z^{(1)}$ is a positive definite operator since for any finite set of distinct samples $\tilde{X} := [\tilde{x}_1, \dots, \tilde{x}_n]$ with $t_i \neq t_j$ for all i, j , the matrix

$$K := \{K_Z(\tilde{x}_i, \tilde{x}_j)\}_{ij} \succ \frac{1}{4} \tilde{X}^\top \tilde{X} \otimes I_{d_Z} \succ 0,$$

where we use $\|\tilde{x}\| > 0$ and $\text{PE}(t_i)$'s are linearly independent from the condition of the lemma, and thus their Gram matrix $\tilde{X}^\top \tilde{X} \succ 0$. Consequently, $K_Z = K_Z^{(1)} + K_Z^{(2)} \succ 0$ and therefore its minimal eigenvalue is positive.

H. Proof of Theorem 3.12

For i.i.d samples $[x^1, \dots, x^n]$, let the Hilbert space spanned by the NTKs $K_Z(x, t, \cdot, \cdot)$'s and $K_G(x, t, \cdot, \cdot)$'s be \mathcal{H}_{K_Z} and \mathcal{H}_{K_G} respectively. Further, for any $f(x, t) := \int_{t_0}^T \int_{\mathcal{X}} K(x, t, x', t') c(x', t') dx' dt' \in \mathcal{H}_K, c(x, t) \in \mathbb{R}^d, \forall x, t$, define the *NTK norm* as

$$\|f\|_K := \langle f, f \rangle_K := \sqrt{\mathbb{E}_{t, t', p_t(x) p_t(x')} c(x, t)^\top K(x, t, x', t') c(x', t)}. \quad (49)$$

Further, define the *subspace score matching losses* as

$$\begin{aligned} L_Z(U, s_Z) &:= \mathbb{E}_{t, p_t(x)} \|U s_Z(x, t) - A_Z s_Z^*(x, t)\|^2 =: \mathbb{E}_{t, p_t(x)} \ell_Z(x, t; U, s_Z) \\ L_G(V, s_G) &:= \mathbb{E}_{t, p_t(x)} \|V s_G(g(x), t) - A_G s_G^*(x, t)\|^2 =: \mathbb{E}_{t, p_t(x)} \ell_G(x, t; V, s_G), \end{aligned}$$

and their empirical versions as

$$\begin{aligned}\hat{L}_Z(U, s_Z) &:= \frac{1}{n} \sum_{i=1}^n \ell_Z(\tilde{x}^i; U, s_Z) \\ \hat{L}_G(V, s_G) &:= \frac{1}{n} \sum_{i=1}^n \ell_G(\tilde{x}^i; V, s_G).\end{aligned}$$

To prove the theorem, we make use of the following lemma proved in Section H.1.

Lemma H.1. (generalization error bound) *Let $\min\{d_T, d_H\} \rightarrow \infty$, and for i.i.d samples $[x^1, \dots, x^n]$, let the reproducing kernel Hilbert space (RKHS) spanned by the NTKs $K_Z(x, t, \cdot, \cdot)$'s and $K_G(x, t, \cdot, \cdot)$'s be \mathcal{H}_{K_Z} and \mathcal{H}_{K_G} respectively, and denote $K_{Z,(x,t)} := K_Z(x, t, \cdot, \cdot)$. Further, define the function classes $\mathcal{S}_Z, \mathcal{U}, \mathcal{S}_G, \mathcal{V}$ as*

$$\begin{aligned}\mathcal{S}_Z &:= \left\{ f = \sum_{i=1}^N c_i K_{Z, \tilde{x}_i} \mid \forall N \in \mathbb{N}, \|f\|_K \leq C_Z \sqrt{\frac{\sigma_1(s_Z^*) d_Z^{1/2}}{\lambda_{\min}(K_Z^*)}}, \forall [\tilde{x}_1, \dots, \tilde{x}_N] \in (\mathcal{X} \times [t_0, T])^N \right\}, \\ \mathcal{U} &:= \left\{ U : \|U\|_F \leq C_Z \sqrt{\sigma_1(s_Z^*) d_Z^{1/2}} \right\}, \\ \mathcal{S}_G &:= \left\{ f = \sum_{i=1}^N c_i K_{G, \tilde{x}_i} \mid \forall N \in \mathbb{N}, \|f\|_K \leq C_G \sqrt{\frac{\sigma_1(s_G^*) d_G^{1/2}}{\lambda_{\min}(K_G^*)}}, \forall [\tilde{x}_1, \dots, \tilde{x}_N] \in (\mathcal{X} \times [t_0, T])^N \right\}, \\ \mathcal{V} &:= \left\{ V : \|V\|_F \leq C_G \sqrt{\sigma_1(s_G^*) d_G^{1/2}} \right\},\end{aligned}$$

where $\sigma_i(A)$ is the i -th largest singular value of the operator A . Then with probability at least $1 - O(\frac{1}{n})$,

$$\begin{aligned}L_Z(\hat{U}, \hat{s}_Z) &\leq \min_{(U, s_Z) \in \mathcal{U} \times \mathcal{S}_Z} L_Z(U, s_Z) + O\left(\sqrt{\frac{d_X^5 \log^3 n}{n}}\right), \\ L_G(\hat{V}, \hat{s}_G) &\leq \min_{(V, s_G) \in \mathcal{V} \times \mathcal{S}_G} L_G(V, s_G) + O\left(\sqrt{\frac{d_X^5 \log^3 n}{n}}\right).\end{aligned}$$

Now we are ready to prove the theorem. First, by the universal approximation theorem of neural networks (e.g. (Barron, 1993)) and the Lipschitzness of the true score functions, for C_Z, C_G large enough, $s_Z^* \in \mathcal{S}_Z$ and $s_G^* \in \mathcal{S}_G$ and therefore

$$\begin{aligned}\min_{(U, s_Z) \in \mathcal{U} \times \mathcal{S}_Z} L_Z(U, s_Z) &= L_Z(A_Z, s_Z^*) = 0, \\ \min_{(V, s_G) \in \mathcal{V} \times \mathcal{S}_G} L_G(V, s_G) &= L_G(A_G, s_G^*) = 0.\end{aligned}$$

As a result, we have with probability at least $1 - O(\frac{1}{n})$,

$$L_Z(\hat{U}, \hat{s}_Z) \leq \epsilon(n), \quad L_G(\hat{V}, \hat{s}_G) \leq \epsilon(n),$$

where $\epsilon(n) = O\left(\sqrt{\frac{d_X^5 \log^3 n}{n}}\right)$. Therefore, we prove item 1 by noticing

$$L_0(\hat{U}, \hat{V}, \hat{s}_Z, \hat{s}_G) \leq 2L_Z(\hat{U}, \hat{s}_Z) + 2L_G(\hat{V}, \hat{s}_G) \leq 2\epsilon(n) = O\left(\sqrt{\frac{d_X^5 \log^3 n}{n}}\right).$$

We proceed to prove item 2. Let $Q_Z \in \mathbb{R}^{d_X \times d_Z}, Q_G \in \mathbb{R}^{d_X \times d_G}$ be orthogonal matrices such that

$$P_{R(\hat{U})} = Q_Z Q_Z^\top, P_{R(\hat{V})} = Q_G Q_G^\top,$$

then we have $\hat{U}\hat{s}_Z := Q_Z Q_Z^\top \hat{U}\hat{s}_Z$ and $\hat{V}\hat{s}_G := Q_G Q_G^\top \hat{V}\hat{s}_G$,

Then applying Lemma 7 of (Chen et al., 2023a) yields with probability at least $1 - 4\delta$,

$$\begin{aligned}\|P_{R(\hat{U})} - A_Z A_Z^\top\|^2 &\leq \frac{\epsilon(n)}{\sigma_{d_Z}(s_Z^*)^2} \\ \|P_{R(\hat{V})} - A_G A_G^\top\|^2 &\leq \frac{\epsilon(n)}{\sigma_{d_G}(s_G^*)^2}.\end{aligned}$$

Finally, for item 3, define $Z_{\parallel} := P_U A_Z Z$ and $Z_{\perp} := P_U A_G G$, where $P_U := P_{R(U)}$ is the projection matrix onto the $R(U)$. By definition of \hat{Z} , we have

$$\hat{Z} = P_U A_Z Z + P_U A_G G = Z_{\parallel} + Z_{\perp}.$$

Since Z_{\parallel} is a function of Z , we have $Z_{\parallel} \perp\!\!\!\perp G$. In addition, we have $\hat{Z} - Z_{\perp} - G$ forms a Markov chain, and therefore by data processing inequality,

$$\begin{aligned}I(\hat{Z}; G) &\leq I(\hat{Z}; Z_{\perp}) = I(Z_{\parallel} + Z_{\perp}; Z_{\perp}) \\ &= \mathbb{E}_{p_{Z_{\perp}}(z, y)} \log \frac{p_{\hat{Z}Z_{\perp}}(z, y)}{p_{\hat{Z}}(z)p_{Z_{\perp}}(y)} \\ &= \mathbb{E}_{p_{\hat{Z}Z_{\perp}}(z, y)} \log \frac{p_{Z_{\parallel}}(z-y)}{p_{\hat{Z}}(z)} \\ &= \mathbb{E}_{p_{\hat{Z}Z_{\perp}}(z, y)} \log \frac{p_{Z_{\parallel}}(z)}{p_{\hat{Z}}(z)} + \mathbb{E}_{p_{\hat{Z}Z_{\perp}}(z, y)} \log \frac{p_{Z_{\parallel}}(z-y)}{p_{Z_{\parallel}}(z)} \\ &= \mathbb{E}_{p_{\hat{Z}}(z)} \log \frac{p_{Z_{\parallel}}(z)}{p_{\hat{Z}}(z)} + \mathbb{E}_{p_{\hat{Z}Z_{\perp}}(z, y)} \log \frac{p_{Z_{\parallel}}(z-y)}{p_{Z_{\parallel}}(z)} \\ &= -D_{\text{KL}}(p_{\hat{Z}} \| p_{Z_{\parallel}}) + \mathbb{E}_{p_{\hat{Z}Z_{\perp}}(z, z-y)} \log \frac{p_{Z_{\parallel}}(z-y)}{p_{Z_{\parallel}}(z)} \\ &\leq \mathbb{E}_{p_{\hat{Z}Z_{\perp}}(z, z-y)} \log \frac{p_{Z_{\parallel}}(z-y)}{p_{Z_{\parallel}}(z)} = \mathbb{E}_{p_{Z_{\perp}}(y)p_{Z_{\parallel}}(z-y)} \log \frac{p_{Z_{\parallel}}(z-y)}{p_{Z_{\parallel}}(z)} \\ &= O\left(\mathbb{E}_{p_{Z_{\parallel}}(z-y)p_{Z_{\perp}}(y)} \|z-y\| \|y\| + \mathbb{E}_{p_{Z_{\perp}}(y)} \|y\|^2\right) = O(\|P_U A_G\| + \|P_U A_G\|^2),\end{aligned}$$

where the last inequality uses the nonnegativity of KL divergence and the second-to-last equality combines Lemma B.4 with the fact that $Z_{\parallel} = P_U A_Z Z$ is a 1-Lipschitz function of Z and therefore its score function along the column space of $P_U A_Z$ is λ_s -Lipschitz. To upper bound $\|P_U A_G\|^2$, we utilize item 2 by noticing that

$$\|P_U A_G\|^2 = \frac{1}{2} (\|P_U - A_Z A_Z^\top\|^2 + \text{rank}(U) - d_Z) \leq \frac{\epsilon(n)}{2\sigma_{d_Z}(s_Z^*)^2},$$

where the last inequality uses the fact that $\text{rank}(U) \leq d_Z$. As a result,

$$I(\hat{Z}; G) = O\left(\frac{\sqrt{\epsilon(n)}}{\sigma_{d_Z}(s_Z^*)}\right) = O\left(\frac{d_X^{5/4} \log^{3/4} n}{\sigma_{d_Z}(s_Z^*) n^{1/4}}\right).$$

Similarly, to prove item 4, let $f_1(Z) := P_U A_Z Z$ and $f_2(G) := P_U A_G G$, choose $t_0 := \frac{1}{2} \log \frac{1}{1-\delta^2}$ for some $\delta > 0$, and define

$$\hat{Z}_{t_0} := \hat{Z} + \delta N_{t_0} = P_U A_Z Z_{t_0} + P_U A_G G_{t_0} = f_1(Z_{t_0}) + f_2(G_{t_0}).$$

Then we have

$$\begin{aligned}
 & I(\hat{Z}_{t_0}; G|Z) \\
 &= \mathbb{E}_{p_{\hat{Z}_{t_0}|Z}} \log \frac{p_{\hat{Z}_{t_0}|G,Z}(\hat{z}|g,z)}{p_{\hat{Z}_{t_0}|Z}(\hat{z}|z)} \\
 &= \mathbb{E}_{p_{\hat{Z}_{t_0}|Z}} \log \frac{p_{\delta N_{t_0}}(\hat{z} - f_1(z) - f_2(g))}{p_{f_2(g) + \delta N_{t_0}}(\hat{z} - f_1(z))} \\
 &= -D_{\text{KL}}(p_{\delta N_{t_0}} \| p_{f_2(g) + \delta N_{t_0}}) + \mathbb{E}_{p_{\hat{Z}_{t_0}|Z}} \log \frac{p_{\delta N_{t_0}}(\hat{z} - f_1(z) - f_2(g))}{p_{\delta N_{t_0}}(\hat{z} - f_1(z))} \\
 &\leq \frac{1}{\delta^2} O\left(\mathbb{E}_{p_{\hat{Z}_{t_0}|Z}} \|\hat{z} - f_1(z)\| \|f_2(g)\| + \mathbb{E}_{p_{G}(g)} \|f_2(g)\|^2\right) \\
 &= O\left(\frac{1}{\delta^2} \|P_U A_G\|\right) = O\left(\frac{d_X^{5/4} \log^{3/4} n}{\delta^2 \sigma_{d_Z}(s_Z^*) n^{1/4}}\right),
 \end{aligned}$$

where the inequality uses Lemma B.4. By Pinsker's inequality, and choose $\delta := \frac{\log^{3/4} n}{n^{1/16}}$ this implies that the second condition of editability holds with

$$\epsilon = O\left(\frac{d_X^{5/8} \log^{3/4} n}{\sigma_{d_Z}(s_Z^*)^{1/2} n^{1/16}}\right).$$

To check the first condition holds, we employ Lemma C.5. To this end, we first check that $p_{\hat{\psi}(\hat{Z}_{t_0}, A_G G_{t_0})}$ has bounded gradient. Since we know $p_{X_{t_0}}$ has bounded gradient by a similar argument in Theorem 3.7, and

$$\hat{\psi}(\hat{Z}_{t_0}, A_G G_{t_0}) := (P_U A_Z A_Z^\top + P_V A_G A_G^\top) X_{t_0}$$

is a linear function of X_{t_0} , it suffices to show that $[P_U A_Z, P_V A_G]$ is invertible. This is the case with high probability since by item 2,

$$\|P_U A_Z A_Z^\top P_U + P_V A_G A_G^\top P_V - I_{d_X}\| \leq \|P_U - A_Z A_Z^\top\|^2 + \|P_V - A_G A_G^\top\|^2 = O\left(\frac{\log^{3/2} n}{n^{1/2}}\right)$$

with high probability. Therefore, by matrix perturbation inequality,

$$\sigma_{d_X}^2([P_U A_Z, P_V A_G]) \geq 1 - O\left(\frac{\log^{3/2} n}{n^{1/2}}\right) > 0$$

for sufficiently large n . Therefore, applying Lemma C.5,

$$\begin{aligned}
 & d_{\text{TV}}(p_{\hat{\psi}(\hat{Z}_{t_0}, A_G G_{t_0})}, p_X) \\
 &\leq O\left(\mathbb{E}\|\hat{\psi}(\hat{Z}_{t_0}, G_{t_0}) - X\|^2\right)^{1/4} \\
 &= O\left(\mathbb{E}\|(P_U A_Z A_Z^\top - A_Z A_Z^\top)X\|^2 + \mathbb{E}\|(P_V A_G A_G^\top - A_G A_G^\top)X\|^2 + \sigma(t_0)^2 \mathbb{E}\|(P_U A_Z A_Z^\top + P_V A_G A_G^\top)N_{t_0}\|^2\right)^{1/4} \\
 &= O\left(\frac{d_X^{5/2} \log^{3/2} n}{\min\{\sigma_{d_Z}^2(s_Z^*), \sigma_{d_G}^2(s_G^*)\} n^{1/2}} + \frac{\log^{3/2} n}{n^{1/8}}\right)^{1/4} = O\left(\frac{d_X^{5/8} \log^{3/8} n}{\min\{\sigma_{d_Z}^{1/2}(s_Z^*), \sigma_{d_G}^{1/2}(s_G^*)\} n^{1/32}}\right).
 \end{aligned}$$

Combining the two conditions, we have $\hat{Z}_{t_0}|Z$ and $A_G G_{t_0}$ are $\left(O\left(\frac{d_X^{5/8} \log^{3/8} n}{\min\{\sigma_{d_Z}^{1/2}(s_Z^*), \sigma_{d_G}^{1/2}(s_G^*)\} n^{1/32}}\right), \hat{\psi}, p\right)$ -editable.

H.1. Proof of Lemma H.1

As $d_H, d_T \rightarrow \infty$, we have $K_Z = K_Z^*, K_G = K_G^*$. First, we prove the following lemmas in Section H.2 and H.3 respectively.

Lemma H.2. Suppose $n > \max\{d_Z, d_G\}$, and $\min\{d_T, d_H\} \rightarrow \infty$, the following holds for the empirical risk minimizer $(\hat{s}_Z, \hat{s}_G, \hat{U}, \hat{V})$ with probability at least $1 - 2\exp(-n)$:

$$\begin{aligned}\hat{L}_Z(\hat{U}, \hat{s}_U) &= \hat{L}_G(\hat{V}, \hat{s}_V) = 0, \\ \|\hat{s}_Z\|_{K_Z} &\leq C_Z \sqrt{\frac{\sigma_1(s_Z^*) d_Z^{1/2}}{\lambda_{\min}(K_Z^*)}}, \\ \|\hat{s}_G\|_{K_G} &\leq C_G \sqrt{\frac{\sigma_1(s_G^*) d_G^{1/2}}{\lambda_{\min}(K_G^*)}},\end{aligned}$$

for some constant $C_Z, C_G > 0$.

Lemma H.3. For the NTK of the estimated content and style score functions K_Z and K_G defined in Lemma G.3 and for any $(x, t) \in \mathcal{X} \times [t_0, T]$, the following holds:

$$\begin{aligned}\|K(x, t, x, t)\| &\leq \frac{3}{2}(\|x\|^2 + \|PE(t)\|^2) \\ \|K_G(x, t, x, t)\| &\leq \frac{3}{2}(\lambda_g^2 \|x\|^2 + \|PE(t)\|^2).\end{aligned}$$

Then we make use of the following properties of sub-gaussian random variables from Lemma 16 of (Chen et al., 2023a).

Lemma H.4. Consider a probability density function $p(x) \leq \exp(-C\|x\|_2^2/2)$ for $x \in \mathbb{R}^d$ and constant $C > 0$. Let R be a fixed radius. Then the following holds

$$\begin{aligned}\int_{\|x\| > R} p(x) dx &\leq \frac{2d\pi^{d/2}}{C\Gamma(d/2+1)} R^{d-2} \exp(-CR^2/2), \\ \int_{\|x\| > R} \|x\|^2 p(x) dx &\leq \frac{2d\pi^{d/2}}{C\Gamma(d/2+1)} R^d \exp(-CR^2/2).\end{aligned}$$

Define $\rho_Z := C_Z \sqrt{\frac{\sigma_1(s_Z^*) d_Z^{1/2}}{\lambda_{\min}(K_Z^*)}}$, and without loss of generality, assume $\sup_{t \in [t_0, T]} \|PE(t)\| \leq T$. To prove lemma H.1, we first bound the Rademacher average of \mathcal{S}_Z as

$$\begin{aligned}\mathcal{R}_n(\mathcal{S}_Z) &= \mathbb{E}_{\epsilon^n} \sup_{s_Z \in \mathcal{S}_Z(\tilde{x}^{1:n})} \frac{1}{n} \sum_{i=1}^n \langle \epsilon_i, s_Z(\tilde{x}^i) \rangle \\ &= \frac{1}{n} \mathbb{E}_{\epsilon^n} \sup_{c \in \mathcal{C}_Z(\tilde{x}^{1:n})} \sum_{i=1}^n \sum_{j=1}^{d_Z} \epsilon_{ij} \langle s_{Z,j}, K_{Z, \tilde{x}^i, j} \rangle_{K_Z} \\ &= \frac{1}{n} \mathbb{E}_{\epsilon^n} \sup_{c \in \mathcal{C}_Z(\tilde{x}^{1:n})} \sum_{j=1}^{d_Z} \left\langle s_{Z,j}, \sum_{i=1}^n \epsilon_{ij} K_{Z, \tilde{x}^i, j} \right\rangle_{K_Z} \\ &\leq \frac{\rho_Z}{n} \mathbb{E}_{\epsilon^n} \sqrt{\sum_{j=1}^{d_Z} \left\| \sum_{i=1}^n \epsilon_{ij} K_{Z, \tilde{x}^i, j} \right\|_{K_Z}^2} \\ &\leq \frac{\rho_Z}{n} \sqrt{\mathbb{E}_{\epsilon^n} \sum_{j=1}^{d_Z} \left\| \sum_{i=1}^n \epsilon_{ij} K_{Z, \tilde{x}^i, j} \right\|_{K_Z}^2} \\ &= \frac{\rho_Z}{n} \sqrt{\sum_{i=1}^n p(t^i) p_{t^i}(x^i) \mathbb{E}_{\epsilon_i} \epsilon_i^\top K_Z(\tilde{x}^i, \tilde{x}^i) \epsilon_i} \\ &\leq \frac{\rho_Z}{n} \sqrt{\sum_{i=1}^n p(t^i) p_{t^i}(x^i) \text{Tr}(K_Z(\tilde{x}^i, \tilde{x}^i))}.\end{aligned}$$

Averaging over \tilde{x}^n and applying Cauchy-Schwarz inequality, we can bound the Rademacher complexity of the data-dependent function class \mathcal{S}_Z as

$$\begin{aligned}\mathbb{E}_{\tilde{x}^n} \mathcal{R}_n(\mathcal{S}_Z) &= \mathbb{E}_{\tilde{x}^n} \frac{\rho_Z}{n} \sqrt{\sum_{i=1}^n p^{(t^i)} p_{t^i}(x^i) \text{Tr}(K_Z(\tilde{x}^i, \tilde{x}^i))} \\ &\leq \frac{\rho_Z}{\sqrt{n}} \sqrt{\mathbb{E}_{t, p_t(x) p_t(x)} \text{Tr}(K_Z(\tilde{x}, \tilde{x}))} =: \frac{\rho_Z C_{K_Z}}{\sqrt{n}},\end{aligned}$$

where $C_{K_Z} := \sqrt{\mathbb{E}_{t, p_t(x) p_t(x)} \text{Tr}(K_Z(\tilde{x}, \tilde{x}))} = O(d_X)$.

To proceed, let the content subspace matrix and score function class learned by solving Eq. 17-Eq. 18 given training data $[\tilde{x}^1, \dots, \tilde{x}^n]$ be $\mathcal{U}(\tilde{x}^{1:n})$ and $\mathcal{S}_Z(\tilde{x}^{1:n})$ respectively, then by Lemma G.2, with probability at least $1 - 2\exp(-\Omega(n))$, the event

$$\mathcal{E} := \mathcal{S}_Z(\tilde{x}^{1:n}) \subseteq \mathcal{S}_Z \quad (50)$$

will occur, which implies with the same probability bound, the empirical risk minimizer over $\mathcal{S}_Z(x^{1:n})$ satisfies

$$\hat{L}_Z(\hat{U}, \hat{s}_Z) = \min_{(U, s_Z) \in \mathcal{U}(\tilde{x}^{1:n}) \times \mathcal{S}_Z(\tilde{x}^{1:n})} \frac{1}{n} \sum_{i=1}^n \ell_Z(\tilde{x}^i; U, s_Z) = \min_{U, s_Z \in \mathcal{S}_Z} \frac{1}{n} \sum_{i=1}^n \ell_Z(\tilde{x}^i; U, s_Z) = 0.$$

Therefore, let an empirical risk minimizer of L_Z over \mathcal{S}_Z be \hat{U}', \hat{s}'_Z , then for any $\epsilon > 0$, we can bound the generalization error probability as

$$\begin{aligned}&\Pr \left[L_Z(\hat{U}, \hat{s}_Z) \geq \min_{(U, s_Z) \in \mathcal{U} \times \mathcal{S}_Z} L_Z(U, s_Z) + \epsilon \right] \\ &\leq \Pr(\mathcal{E}) + \Pr \left[L_Z(\hat{U}', \hat{s}'_Z) \geq \min_{(U, s_Z) \in \mathcal{U} \times \mathcal{S}_Z} L_Z(U, s_Z) + \epsilon \right] \\ &\leq \Pr \left[L_Z(\hat{U}', \hat{s}'_Z) \geq \min_{(U, s_Z) \in \mathcal{U} \times \mathcal{S}_Z} L_Z(U, s_Z) + \epsilon \right] + 2\exp(-\Omega(n)).\end{aligned} \quad (51)$$

Next, since the squared loss ℓ_Z is not Lipschitz with respect to x , we apply a truncation argument on L_Z . To this end, define the truncated version of L_Z as

$$L_Z^{\text{trunc}}(U, s_Z) := \mathbb{E}_{t, p_t} \ell(x, t; U, s_Z) \mathbb{1}_{\|x\| \leq R}$$

for some radius $R > 0$. Similarly we can define its empirical version as L_Z^{trunc} . Then L_Z admits the following decomposition:

$$\begin{aligned}&L_Z(\hat{U}', \hat{s}'_Z) - L_Z(A_Z, s_Z^*) \\ &= L_Z^{\text{trunc}}(\hat{U}', \hat{s}'_Z) - L_Z^{\text{trunc}}(A_Z, s_Z^*) + L_Z(\hat{U}, \hat{s}_Z) - L_Z^{\text{trunc}}(\hat{U}', \hat{s}'_Z) + L_Z^{\text{trunc}}(A_Z, s_Z^*) - L_Z(A_Z, s_Z^*) \\ &\leq \underbrace{L_Z^{\text{trunc}}(\hat{U}', \hat{s}'_Z) - L_Z^{\text{trunc}}(A_Z, s_Z^*)}_{(i)} + \underbrace{L_Z(\hat{U}, \hat{s}_Z) - L_Z^{\text{trunc}}(\hat{U}', \hat{s}'_Z)}_{(ii)}\end{aligned}$$

To bound (i), notice that for $s_Z \in \mathcal{S}_Z$, with balanced weight $U \in \mathcal{U}_Z$,

$$\begin{aligned}&|\ell(x', t'; U, s_Z) \mathbb{1}_{\|x'\| \leq R} - \ell(x, t; U, s_Z) \mathbb{1}_{\|x\| \leq R}| \\ &\leq \sup_{x \in \mathcal{X}, t \in [t_0, T]} \sup_{(U, s_Z) \in \mathcal{U} \times \mathcal{S}_Z} |\ell(x, t; U, s_Z) \mathbb{1}_{\|x\| \leq R}| \\ &\leq \sup_{x \in \mathcal{X}, t \in [t_0, T]} \sup_{(U, s_Z) \in \mathcal{U} \times \mathcal{S}_Z} 2(\|U s_Z(x, t)\|^2 + \|s_Z^*(x, t)\|^2) \mathbb{1}_{\|x\| \leq R} \\ &= \sup_{x \in \mathcal{X}, t \in [t_0, T]} \sup_{(U, s_Z) \in \mathcal{U} \times \mathcal{S}_Z} 2\|U s_Z(x, t)\|^2 \mathbb{1}_{\|x\| \leq R} + 2\lambda_s^2(R^2 + T^2) \\ &= (3\rho_Z^4 \lambda_{\min}(K_Z^*) + 2\lambda_s^2)(R^2 + T^2) =: B,\end{aligned}$$

where the second-to-last inequality uses the Lipschitz property of s_Z^* :

$$\|s_Z^*(x, t)\|^2 \mathbb{1}_{\|x\| \leq R} \leq \lambda_s^2 \|\tilde{x}\|^2 \mathbb{1}_{\|x\| \leq R} \leq \lambda_s^2(R^2 + T^2), \quad (52)$$

where the last inequality uses Lemma H.3 as follows:

$$\begin{aligned}
 \|U s_Z(x, t)\| &\leq \|U\| \|s_Z(x, t)\| \\
 &\leq \rho_Z \sqrt{\lambda_{\min}(K_Z)} \sqrt{\sum_{j=1}^{d_Z} \langle s_{Z,j}, K_{Z,x,t,j} \rangle_{K_Z}^2} \\
 &\leq \rho_Z \sqrt{\lambda_{\min}(K_Z)} \|s_Z\|_{K_Z} \|K_{Z,x,t}\|_{K_Z} \\
 &\leq \rho_Z^2 \sqrt{\lambda_{\min}(K_Z)} \|K_Z(x, t, x, t)\| \leq \rho_Z^2 \sqrt{\frac{3}{2} \lambda_{\min}(K_Z)} \|\tilde{x}\|.
 \end{aligned} \tag{53}$$

Using Eq. 52-53, we can show that

$$\begin{aligned}
 \|U s_Z(\tilde{x}^i) - A_Z s_Z^*(\tilde{x}^i)\| \mathbb{1}_{\|\tilde{x}^i\| \leq R} &\leq \|U s_Z(\tilde{x}^i)\| \mathbb{1}_{\|\tilde{x}^i\| \leq R} + \|A_Z s_Z^*(\tilde{x}^i)\| \mathbb{1}_{\|\tilde{x}^i\| \leq R} \\
 &\leq (\rho_Z^2 \sqrt{3 \lambda_{\min}(K_Z^*)/2} + \lambda_s) \sqrt{R^2 + T^2}.
 \end{aligned}$$

Let $\lambda_{\text{trunc}} := 2(\rho_Z^2 \sqrt{3 \lambda_{\min}(K_Z^*)/2} + \lambda_s) \sqrt{R^2 + T^2}$ and use the inequality for $\max\{|s|, |t|\} \leq \lambda$,

$$|s^2 - t^2| < 2\lambda|s - t|,$$

we deduce that

$$|\ell(\tilde{x}^i; U', s'_Z) - \ell(\tilde{x}^i; U, s_Z)| \mathbb{1}_{\|\tilde{x}^i\| \leq R} \leq \lambda_{\text{trunc}} \left| \|U' s'_Z(\tilde{x}^i) - A_Z s_Z^*(\tilde{x}^i)\| - \|U s_Z(\tilde{x}^i) - A_Z s_Z^*(\tilde{x}^i)\| \right|,$$

and thus $\ell(\tilde{x}^i; U, s_Z)$ is a λ_{trunc} -Lipschitz function of $\|U s_Z(\tilde{x}^i) - A_Z s_Z^*(\tilde{x}^i)\|$. Denote

$$\begin{aligned}
 \mathcal{U} \mathcal{S}_Z &:= \{U s_Z : U \in \mathcal{U}, s_Z \in \mathcal{S}_Z\}, \\
 \ell \circ \mathcal{U} \mathcal{S}_Z(\tilde{x}^n) &:= \{(\ell(\tilde{x}^1; U, s_Z), \dots, \ell(\tilde{x}^n; U, s_Z)) : U s_Z \in \mathcal{U} \mathcal{S}_Z\},
 \end{aligned}$$

and apply the standard generalization bound for Lipschitz function on L_Z^{trunc} (see, e.g., Theorem 9.1 of (Hajek & Raginsky, 2021)) to conclude that with probability at least $1 - \delta$,

$$\begin{aligned}
 (i) &\leq 2\mathbb{E}_{\tilde{x}^n} \mathcal{R}_n(\ell \circ \mathcal{U} \mathcal{S}_Z(\tilde{x}^n)) + \frac{B}{2} \sqrt{\frac{8 \log(1/\delta)}{n}} \\
 &\leq 2\lambda_{\text{trunc}} \left(\mathbb{E} \mathcal{R}_n(\mathcal{U} \mathcal{S}_Z) + \frac{\lambda_s \sqrt{R^2 + T^2}}{\sqrt{n}} \right) + \frac{B}{2} \sqrt{\frac{8 \log(1/\delta)}{n}} \\
 &\leq 2\lambda_{\text{trunc}} \frac{\rho_Z^2 C_{K_Z} + \lambda_s \sqrt{R^2 + T^2}}{\sqrt{n}} + \frac{B}{2} \sqrt{\frac{8 \log(1/\delta)}{n}} \\
 &= O\left(\frac{\lambda_s \rho_Z^4 (R^2 + T^2)}{\sqrt{n}} + \lambda_s \rho_Z^4 (R^2 + T^2) \sqrt{\frac{\log 1/\delta}{n}} \right) = O\left(\frac{\lambda_s d_Z T^2 R^2}{\sqrt{n}} + \lambda_s d_Z T^2 R^2 \sqrt{\frac{\log 1/\delta}{n}} \right),
 \end{aligned}$$

where the second inequality uses triangle inequality within the definition of Rademacher complexity followed by the bound in Eq. 52, and the third inequality uses the contraction principle of Rademacher complexity (Hajek & Raginsky, 2021).

To bound (ii), notice that

$$\begin{aligned}
 (ii) &= \mathbb{E}_{t,p_t(x)} \ell_Z(x,t; \hat{U}', \hat{s}'_Z) \mathbb{1}_{\|x\|>R} \\
 &\leq 2\mathbb{E}_{t,p_t(x)} (\|\hat{U}' \hat{s}'_Z(x,t)\|^2 + \|s^*_Z(x,t)\|^2) \mathbb{1}_{\|x\|>R} \\
 &\leq 2\mathbb{E}_{t,p_t(x)} \left[\|\hat{U}' \hat{s}'_Z(x,t)\|^2 + \lambda_s^2 (\lambda_z^2 \|x\|^2 + t^2) \right] \mathbb{1}_{\|x\|>R} \\
 &\leq 2\mathbb{E}_{t,p_t(x)} \left[\frac{3}{2} \rho_Z^4 \lambda_{\min}(K_Z) d_Z (\|x\|^2 + \|\text{PE}(t)\|^2) + \lambda_s^2 (\lambda_z^2 \|x\|^2 + t^2) \right] \mathbb{1}_{\|x\|>R} \\
 &= (3\rho_Z^4 \lambda_{\min}(K_Z) d_Z + 2\lambda_s^2 \lambda_z^2) \mathbb{E}_{t,p_t(x)} \|x\|^2 \mathbb{1}_{\|x\|>R} + \mathbb{E}_{t,p_t(x)} \left(\lambda_s^2 t^2 + \frac{3}{2} \|\text{PE}(t)\|^2 \right) \mathbb{1}_{\|x\|>R} \\
 &= (3\rho_Z^4 \lambda_{\min}(K_Z) d_Z + 2\lambda_s^2 \lambda_z^2) \frac{\sigma_X^2 d_X 2^{-d_X/2+1} R^{d_X}}{\Gamma(d_X/2+1)} \exp(-R^2/2\sigma_X^2) + \\
 &\quad \left(\frac{3}{2} \rho_Z^4 \lambda_{\min}(K_Z) d_Z \mathbb{E}_t \|\text{PE}(t)\|^2 + \frac{\lambda_s^2 (T^3 - t_0^3)}{3(T-t_0)} \right) \frac{\sigma_X^2 d_X 2^{-d_X/2+1}}{\Gamma(d_X/2+1)} R^{d_X-2} \exp(-R^2/2\sigma_X^2) \\
 &= O\left(\frac{\lambda_s^2 T^2 \sigma_X^2 d_X^3 2^{-d_X/2+1}}{\Gamma(d_X/2+1)} R^{d_X} \exp(-R^2/2\sigma_X^2) \right).
 \end{aligned}$$

where the second-to-last inequality uses Eq. 52 and the last inequality applies Eq. 53.

Take $R := O(\sqrt{d_X \log d_X + \log n})$ such that (ii) $\leq \frac{\lambda_s^2 d_X T^2}{n}$, and then combining (i) and (ii) yields with probability at least $1 - \delta$,

$$L_Z(\hat{U}', \hat{s}'_Z) - L_Z(A_Z, s^*_Z) \leq O\left(\frac{1 + \sqrt{\log 1/\delta}}{\sqrt{n}} (d_X^{5/2} + \log n) \right). \quad (54)$$

Setting $\delta := \frac{1}{n}$ and combining Eq. 51 and Eq. 54 yields with probability at least $1 - O(\frac{1}{n})$,

$$L_Z(\hat{U}, \hat{s}_Z) - L_Z(A_Z, s^*_Z) \leq O\left(\frac{1 + \sqrt{\log n}}{\sqrt{n}} (d_X^{5/2} + \log n) \right) = O\left(\sqrt{\frac{d_X^5 \log^3 n}{n}} \right).$$

Combining this bound with Eq. 51 yields the desired bound. And similarly, we can prove the bound for \hat{s}_G .

H.2. Proof of Lemma H.2

As $d_H, d_T \rightarrow \infty$, we have $K_Z = K_Z^*, K_G = K_G^*$. By the optimality of \hat{s}_Z and \hat{s}_G and Theorem 3.11, we have

$$\begin{aligned}
 \hat{U} \hat{s}_Z(x^i, t^i) &= A_Z s^*_Z(x^i, t^i), \\
 \hat{V} \hat{s}_G(g(x^i), t^i) &= A_G s^*_G(g(x^i), t^i), \\
 \frac{1}{n} \sum_{i=1}^n \hat{s}_Z(x^i, t^i) \hat{s}_Z(x^i, t^i) &= \hat{U}^\top \hat{U}, \\
 \frac{1}{n} \sum_{i=1}^n \hat{s}_G(g(x^i), t^i) \hat{s}_G(g(x^i), t^i) &= \hat{V}^\top \hat{V}, \forall i = 1, \dots, n.
 \end{aligned} \quad (55)$$

The first two equalities immediately yield

$$\hat{L}_Z(\hat{U}, \hat{s}_U) = \hat{L}_G(\hat{V}, \hat{s}_V) = 0.$$

To prove the last two inequalities in the lemma, let $\mathbb{E}_{\hat{p}_n}[f(x, t)] := \frac{1}{n} \sum_{i=1}^n f(x^i, t^i)$. Then Eq. 55 implies that

$$\begin{aligned}
 &\mathbb{E}_{\hat{p}_n} \|A_Z s^*_Z(x, t)\|^2 \\
 &= \mathbb{E}_{\hat{p}_n} \text{Tr}(\hat{s}_Z^\top(x, t) \hat{U}^\top \hat{U} \hat{s}_Z(x, t)) = \text{Tr}\left(\mathbb{E}_{\hat{p}_n} \hat{s}_Z(x, t) \hat{s}_Z^\top(x, t) \hat{U}^\top \hat{U} \right) \\
 &= \text{Tr}\left(\mathbb{E}_{\hat{p}_n} \hat{s}_Z(x, t) \hat{s}_Z^\top(x, t) \mathbb{E}_{\hat{p}_n} \hat{s}_Z(x, t) \hat{s}_Z^\top(x, t) \right) = \|\mathbb{E}_{\hat{p}_n} \hat{s}_Z(x, t) \hat{s}_Z^\top(x, t)\|^2 \geq (\mathbb{E}_{\hat{p}_n} \|\hat{s}_Z(x, t)\|)^2,
 \end{aligned} \quad (56)$$

Let $\Sigma_n(s) := \mathbb{E}_{\hat{p}_n} s(x,t) s(x,t)^\top$ and $\lambda_i(M)$ be the i -th eigenvalue of the matrix M , then we have

$$\begin{aligned} \mathbb{E}_{\hat{p}_n} \|\hat{s}_Z(x,t)\|^2 &\leq \sqrt{\mathbb{E}_{\hat{p}_n} \|A_Z s_Z^*(x,t)\|^2} = \sqrt{\mathbb{E}_{\hat{p}_n} \|s_Z^*(x,t)\|^2} \\ &= \sqrt{\sum_{i=1}^{d_Z} \lambda_i(\Sigma_n(s_Z^*))} \leq \sqrt{\|\Sigma_n(s_Z^*)\|_{\text{op}} d_Z}. \end{aligned} \quad (57)$$

Further, suppose $\hat{s}_Z(\tilde{x}) = \frac{1}{n} \sum_{i=1}^n K_Z(\tilde{x}, \tilde{x}^i) c_Z(\tilde{x}^i)$ and define

$$\bar{K}_Z := \begin{bmatrix} K_Z(\tilde{x}^1, \tilde{x}^1) & \cdots & K_Z(\tilde{x}^1, \tilde{x}^n) \\ \vdots & \ddots & \vdots \\ K_Z(\tilde{x}^n, \tilde{x}^1) & \cdots & K_Z(\tilde{x}^n, \tilde{x}^n) \end{bmatrix}, \bar{c}_Z := [c_Z(\tilde{x}^1)^\top, \dots, c_Z(\tilde{x}^n)^\top]^\top,$$

then we have

$$\begin{aligned} \|\hat{s}_Z\|_{\bar{K}_Z}^2 &= \frac{1}{n} \bar{c}_Z^\top \bar{K}_Z \bar{c}_Z \leq \frac{\bar{c}_Z^\top \bar{K}_Z^2 \bar{c}_Z}{n \lambda_n(\bar{K}_Z)} = \frac{1}{n \lambda_n(\bar{K}_Z)} \sum_{j=1}^n \left\| \sum_{i=1}^n \bar{K}_Z(\tilde{x}^i, \tilde{x}^j) c_Z(\tilde{x}^j) \right\|^2 \\ &= \frac{1}{\lambda_n(\bar{K}_Z)} \mathbb{E}_{\hat{p}_n} \|\hat{s}_Z(\tilde{x})\|^2 \leq \frac{1}{\lambda_{\min}(K_Z^*)} \mathbb{E}_{\hat{p}_n} \|\hat{s}_Z(\tilde{x})\|^2 \leq \frac{\sqrt{\|\Sigma_n(s_Z^*)\|_{\text{op}} d_Z}}{\lambda_{\min}(K_Z^*)}. \end{aligned}$$

It remains to prove the concentration of the operator norm $\|\Sigma_n(s_Z^*)\|_{\text{op}}$ around $\|\Sigma_\infty(s_Z^*)\|_{\text{op}} =: \|\Sigma(s_Z^*)\|_{\text{op}}$. To this end, we use the assumptions that $s_Z^*(x,t)$ is λ_s -Lipschitz and (z,t) is sub-gaussian with sub-gaussian norm at most $\sigma_Z^2 d_Z + T^2$ to conclude that $s_Z^*(x,t)$ is sub-gaussian with sub-gaussian norm at most $\lambda_s(\sigma_Z^2 d_Z + T^2)$. Then Theorem 4.7.1 of (Vershynin, 2018) yields with probability at least $1 - 2\exp(-u)$,

$$\|\Sigma_n(s_Z^*) - \Sigma(s_Z^*)\|_{\text{op}} \leq C \left(\sqrt{\frac{r+u}{n}} + \frac{r+u}{n} \right) \|\Sigma(s_Z^*)\|_{\text{op}}, \quad (58)$$

where $r := \frac{\text{Tr}(\Sigma(s_Z^*))}{\|\Sigma(s_Z^*)\|_{\text{op}}}$ is the stable rank of $\Sigma(s_Z^*)$. As a result, with probability at least $1 - 2\exp(-n)$,

$$\|\hat{s}_Z\|_{K_Z} \leq \frac{C_Z \|\Sigma(s_Z^*)\|_{\text{op}}^{1/4} d_Z^{1/4}}{\lambda_{\min}(K_Z^*)^{1/2}} = C_Z \sqrt{\frac{\sigma_1(s_Z^*) d_Z^{1/2}}{\lambda_{\min}(K_Z^*)}},$$

where $C_Z := 1 + C^{1/2}(\sqrt{r+1} + r + 1)^{1/2}$ suffices.

Finally, using a similar argument we can show that

$$\|\hat{s}_G\|_{K_G} \leq C_G \sqrt{\frac{\sigma_1(s_G^*) d_G^{1/2}}{\lambda_{\min}(K_G^*)}}$$

with probability at least $1 - 2\exp(-n)$.

H.3. Proof of Lemma H.3

As $d_H, d_T \rightarrow \infty$, we have $K_Z = K_Z^*, K_G = K_G^*$. By definition of the NTKs,

$$\begin{aligned} \|K_Z(x, t, x, t)\| &= \left\| \mathbb{E}_{\mathcal{N}(0, I)} \|a(x, t)\|^2 I_{d_Z} + \frac{1}{2} \|\tilde{x}\|^2 I_{d_Z} \right\| \\ &= \left\| \mathbb{E}_{\mathcal{N}(0, I)} \|a(x, t)\|^2 + \frac{1}{2} \|\tilde{x}\|^2 \right\| d_Z \\ &= \left\| \lim_{d_H \rightarrow \infty} \frac{1}{d_H} \sum_{j=1}^{d_H} \mathbb{E}_{\mathcal{N}(0, I)} (\tilde{x}^\top \Theta_j^{(1)})_+ (\Theta_j^{(1)\top} \tilde{x})_+ + \frac{1}{2} \|\tilde{x}\|^2 \right\| d_Z \\ &\leq \left\| \lim_{d_H \rightarrow \infty} \frac{1}{d_H} \sum_{j=1}^{d_H} \mathbb{E}_{\mathcal{N}(0, I)} (\tilde{x}^\top \Theta_j^{(1)} \Theta_j^{(1)\top} \tilde{x})_+ + \frac{1}{2} \|\tilde{x}\|^2 \right\| d_Z = \left\| \|\tilde{x}\|^2 + \frac{1}{2} \|\tilde{x}\|^2 \right\| = \frac{3}{2} \|\tilde{x}\|^2. \end{aligned}$$

Similarly, we can prove the bound for K_G .

I. Experiment details

I.1. Latent subspace GMM disentanglement

For the synthetic disentanglement experiment, we choose the subspace dimension to be $d_Z = d_G = 5$ and sample the content variable via $Z \sim \frac{1}{2} \mathcal{N}(\mu_1^Z, \sigma_0^2 \mathbf{I}_{d_Z}) + \frac{1}{2} \mathcal{N}(\mu_2^Z, \sigma_0^2 \mathbf{I}_{d_Z})$ and the style variable via $G \sim \frac{1}{2} \mathcal{N}(\mu_1^G, \sigma_0^2 \mathbf{I}_{d_G}) + \frac{1}{2} \mathcal{N}(\mu_2^G, \sigma_0^2 \mathbf{I}_{d_G})$, where $\sigma_0 = 0.1$. In this way, we generate i.i.d 4000 samples for training.

We follow the network architecture shown in Figure 2 with $d_H = 512$. For the time embedding $\text{PE}(\cdot)$, we opt for a *Gaussian Fourier projection layer* to encode temporal information between $(0, 1]$ defined as:

$$\text{PE}(t) := \begin{bmatrix} \sin(2\pi\Omega t) \\ \cos(2\pi\Omega t) \end{bmatrix},$$

where $\Omega \sim \mathcal{N}(\mathbf{0}_{512}, 9000 \mathbf{I}_{512})$. We train the models for 10,000 steps with an Adam (Kingma & Ba, 2015) optimizer with

Name	$\alpha(t)$	$\sigma^2(t)$
VE	1	$\frac{25^{2t} - 1}{2 \log 25}$
VP	$e^{-0.05t - 4.975t^2}$	$1 - e^{-0.1t - 9.95t^2}$
sub-VP	$e^{-0.05t - 4.975t^2}$	$(1 - e^{-0.1t - 9.95t^2})^2$
VP (cosine)	$e^{-\frac{t}{2} - \frac{1}{\pi} \sin(\frac{t}{2})}$	$1 - e^{-t - \frac{2}{\pi} \sin(\frac{t}{2})}$

Table 1. The default noise schedule hyperparameters for the synthetic data experiments. Continuous time ($t \in [10^{-5}, 1]$) is used in the expression.

learning rate 10^{-5} and batch size equal to the entire training set. To ensure convergence, we pretrained the speaker score network \hat{s}^G . We experiment with various noise schedulers, including the variance exploding (VE), vanilla variance preserving (VP) (Ho et al., 2020), sub-VP (Song et al., 2021) and cosine VP (Nichol & Dhariwal, 2021). The detailed schedule hyperparameters are listed in Table 1 and are chosen based on rules of thumbs in (Song & Ermon, 2020; Song et al., 2021).

I.2. Image disentanglement

For all experiments, we use a Gaussian Fourier projection layer to encode temporal information between $(0, 1]$. Further, the U-Net architecture used as the score function for MNIST and CIFAR10 are shown in Table 3 and Table 4 respectively. To capture the content information of the image, we use the 16×16 feature map from a pretrained `vit-small-patch16-224-dino` variant of the DINO model (Caron et al., 2021). The feature map is then resampled to the same size as the image.

For both datasets, we train the DM using an Adam optimizer (Kingma & Ba, 2015) with a batch size of 128 and a learning rate 10^{-4} for 50 epochs. A VE schedule is used during conditional score matching. During inference, we use probability

Table 2. **Quantitative results for image disentanglement on MNIST and CIFAR10 test sets.** MSE, PSNR and SSIM stands for mean squared error, peak signal-to-noise ratio and structural similarity respectively between the generated and target samples. LPIPS (Zhang et al., 2018) is a perceptual metric based on features from deep image classifiers. The results are averaged over two random trials.

	MNIST				CIFAR10			
	MSE(\downarrow)	PSNR(\uparrow)	SSIM(\uparrow)	LPIPS(\downarrow)	MSE(\downarrow)	PSNR(\uparrow)	SSIM(\uparrow)	LPIPS(\downarrow)
$\lambda_r = 0$	0.19 \pm 0.01	5.2 \pm 0.2	0.42 \pm 0.04	0.30 \pm 0.02	0.44 \pm 0.03	3.5 \pm 0.2	0.05 \pm 0.00	0.62 \pm 0.01
$\lambda_r = 0.03$	0.07 \pm 0.04	9.6 \pm 2.3	0.53 \pm 0.06	0.18 \pm 0.04	0.35 \pm 0.02	4.4 \pm 0.1	0.05 \pm 0.00	0.61 \pm 0.02
$\lambda_r = 0.3$	0.01 \pm 0.00	16.0 \pm 0.3	0.66 \pm 0.00	0.1 \pm 0.00	0.18 \pm 0.06	7.3 \pm 0.4	0.06 \pm 0.00	0.53 \pm 0.02
$\lambda_r = 3$	0.01 \pm 0.00	17.3 \pm 0.3	0.66 \pm 0.01	0.1 \pm 0.00	0.11 \pm 0.01	9.3 \pm 0.2	0.07 \pm 0.00	0.49 \pm 0.00

flow (Song et al., 2021) with 500 steps to perform sampling. For the CIFAR10 denoising experiment, we feed a all-zero matrix as the noise map. All models are implemented in Pytorch (Paszke et al., 2019) on two A5000 GPUs. The training time is approximately an hour for both datasets and the inference is approximately 10 seconds for 64 samples.

Full quantitative results on MNIST and CIFAR10 are shown in Table 2

I.3. Speech disentanglement

The dataset statistics are shown in Table 5. The overall results for realistic datasets are shown in Table 6

For the IEMOCAP dataset, we use a system available on SpeechBrain (Ravanelli et al., 2024) that finetunes on the wav2vec 2.0 backbone (Baevski et al., 2020) with a multi-layer perceptron classifier (MLP) (Wang et al., 2021b). The classifier is trained using Adam optimizer for 30 epochs with a batch size of 4 and a learning rate of 10^{-4} for the MLP and the 10^{-5} learning rate for wav2vec 2.0 weights. The system is then evaluated using the standard classification accuracy metric and 5-fold cross validation (Busso et al., 2008; Ma et al., 2024). For each fold, we use all 8 speakers from the training set as target speakers.

On the ALS and ADReSS, we use whisper-medium (Radford et al., 2023) features, as they have shown to be the most effective for speech impairment classification (Wang et al., 2024). To avoid unfair comparison, We concatenate hidden representations over *all* layers of the whisper-medium encoder rather than selecting a particular layer and perform mean pooling over the frame-level features. For both datasets, we follow the standard splits used in previous works (Vieira et al., 2022) to have no overlaps between speaker in the training and test sets. And for both datasets, we use the 15 most frequent speakers from the training set as target speakers for the VC to achieve maximize conversion quality via better speaker representation.

We apply the VCs in mostly a zero-shot, plug-and-play fashion, and leave finetuning to specific datasets for future works. For the Diff-VC, we use the publicly available score network and vocoder checkpoints trained on LibriTTS and adopt the original inference hyperparameter settings for all experiments. Similarly, we use the pretrained models and for other VC models. Further, we use a maximum of 120 second speech from the target speaker to compute the target speaker embedding for all models except KNN-VC, where we use all the target speech as the pool for nearest neighbor search. We also compare VC adaptation with common data augmentation technique such as pitch shifting, where we shift the pitch of all the speech utterances to equally spaced pitch levels over the $F0$ range of the training speech data with levels equal to the number of target speakers and train separate classifiers for each level as in the case of using VC adaptation.

For ALS severity classification as shown in Table 6, KNN-VC achieves the best performance among the VCs, reaching 65% macro-F1 with 15 target speakers and hard majority voting, compared to 54.9% when training without VC adaptation and 61.7% with pitch shifting. For cognitive impairment detection as shown in Table 6, TriAAN-VC performed the best followed by the KNN-VC method, both achieved 83.3% macro-F1 with soft majority voting, which is 12.7% better than methods without VC adaptation and 14.5% and 6.2% better than the pitch shifting adaptation using hard and soft majority voting respectively. On IEMOCAP, we found that Diff-VC performs the best, reaching an average of 97.2% accuracy, which is 25.7% better than the no-VC classifier and 36.1% than the pitch shifting adaptation. Though a phenomenon out of the scope of predictions by our theory, we hypothesized that such “specialization” of the VC methods is due to the different level of generalization ability of different VCs to latent variables *other than* the speaker identity, such as recording conditions and health conditions of the speaker. For instance, Diff-VC does not perform well on ALS compared to KNN-VC, probably due to the domain mismatch between the health conditions of its training set, which contains little pathological speech, compared to KNN-VC which uses the WavLM representation trained on much larger speech dataset with diverse speech.




Figure 7. More image colorization results on MNIST

As to the advantage of hard vs. soft voting, we observe different trends across different datasets and VC methods. On ALS-TDI, hard voting works better than soft voting by 8.4% and 16% for the best two methods Diff-VC and KNN-VC, though worse by 3.9% and 1.3% for pitch shifting and Diff-VC. On IEMOCAP, the gap between soft and hard voting is negligible, with soft majority voting shows a 0.1%-0.7% edge over hard majority voting across VC methods. On ADReSS, we found soft voting methods to be better than hard voting for all the VC methods by 4.1%–6.2%, while worse for the pitch shifting method by 8.3% (68.8% vs. 77.1%). Since soft voting uses a random classifier for voting, it tends to perform well when the model is “confidently” correct and “hesitantly” wrong, as it puts more weights on confident classifiers than hesitant ones. This suggests that the average confidence score estimated in terms of the classifier posteriors on incorrect examples will be high for classifier ensembles that perform well with hard voting than soft voting.

Table 7 8 9 show the complete results for the realistic dataset experiments.



Figure 8. More image denoising results on CIFAR10

Table 3. **U-Net score network used in the MNIST colorization experiment.** The input is $9 \times 28 \times 28$ created by stacking the image, the projected and resized DINO feature map to $3 \times 28 \times 28$ and the background color vector broadcasted to $3 \times 28 \times 28$. $a + b$ denotes that component a accepts hidden embedding from the previous layer and component b accepts the time embedding, and the outputs of a and b are added with proper broadcasting. “Conv2d” refers to 2-D convolutional layer, “GroupNorm” stands for group normalization layer, and “ConvTrans2d” stands for 2-D transposed convolutional layer.

MNIST U-Net
$384 \times 3 \times 1 \times 1$ Conv2d with stride 1 (DINO projection layer)
$9 \times 32 \times 3 \times 3$ Conv2d with stride 2 + 1024×32 Linear
GroupNorm with 4 groups Swish activation
$32 \times 64 \times 3 \times 3$ Conv2d with stride 2 + 1024×64 Linear
GroupNorm with 32 groups Swish activation
$64 \times 128 \times 3 \times 3$ Conv2d with stride 2 + 1024×128 Linear
GroupNorm with 32 groups Swish activation
$128 \times 256 \times 3 \times 3$ Conv2d with stride 2 + 1024×256 Linear
GroupNorm with 32 groups Swish activation
$256 \times 128 \times 4 \times 4$ ConvTrans2d with stride 2 + 1024×128 Linear
GroupNorm with 32 groups Swish activation
$256 \times 64 \times 4 \times 4$ ConvTrans2d with stride 2 + 1024×64 Linear
GroupNorm with 32 groups Swish activation
$128 \times 32 \times 4 \times 4$ ConvTrans2d with stride 2 + 1024×32 Linear
GroupNorm with 32 groups Swish activation
$64 \times 3 \times 3 \times 3$ ConvTrans2d with stride 1

Table 4. **U-Net score network used in the CIFAR10 denoising experiment.** The input is $12 \times 32 \times 32$ created by stacking the image, the projected and resized DINO feature map to $3 \times 32 \times 32$ and the noise map broadcasted to $3 \times 32 \times 32$. A dual encoder with separate U-Nets for the content and the style variables is used and $a + b + c$ denotes that component a accepts the content embedding from the previous layer, component b accepts the previous style embedding and c accepts the time embedding. Further, the outputs of a, c and the outputs of b, c are added separately with proper broadcasting. $a + b$ here denotes that a accepts the previous content embedding, b accepts the previous style embedding and the two outputs are added.

CIFAR10 U-Net	
384 × 6 × 1 × 1 Conv2d with stride 1 (DINO projection layer)	
6 × 32 × 3 × 3 + 3 × 32 × 3 × 3 Conv2d with stride 2 + 1024 × 32 Linear	
GroupNorm with 4 groups Swish activation	
32 × 64 × 3 × 3 + 32 × 64 × 3 × 3 Conv2d with stride 2 + 1024 × 64 Linear	
GroupNorm with 32 groups Swish activation	
64 × 128 × 3 × 3 + 64 × 128 × 3 × 3 Conv2d with stride 2 + 1024 × 128 Linear	
GroupNorm with 32 groups Swish activation	
128 × 256 × 3 × 3 + 128 × 256 × 3 × 3 Conv2d with stride 2 + 1024 × 256 Linear	
GroupNorm with 32 groups Swish activation	
256 × 128 × 4 × 4 + 256 × 128 × 4 × 4 ConvTrans2d with stride 2 + 1024 × 128 Linear	
GroupNorm with 32 groups Swish activation	
256 × 64 × 4 × 4 + 256 × 64 × 4 × 4 ConvTrans2d with stride 2 + 1024 × 64 Linear	
GroupNorm with 32 groups Swish activation	
128 × 32 × 4 × 4 + 128 × 32 × 4 × 4 ConvTrans2d with stride 2 + 1024 × 32 Linear	
GroupNorm with 32 groups Swish activation	
64 × 3 × 3 × 3 + 64 × 3 × 3 × 3 ConvTrans2d with stride 1	

Table 5. **Datasets and VC-adapted classifiers used during realistic data experiments**

	$ \mathcal{Y} $	Feature	Classifier	#Classifiers	Reference	VC	DM-based	Reference
IEMOCAP	4	wav2vec 2.0 base	MLP	8	(Busso et al., 2008)	TriAAN-VC	No	(Park et al., 2023)
ADReSS	2	whisper-medium	SVM	15	(Luz et al., 2020)	KNN-VC	No	(Baas et al., 2023)
ALS-TDI	5	whisper-medium	SVM	15	(Vieira et al., 2022)	Diff-VC	Yes	(Popov et al., 2022)

Table 6. Overall results on realistic datasets. All metrics are between 0-100. A: single (average); B: single (best); MV: majority vote; SMV: soft majority vote.

VC type	Impairment								Emotion			
	ALS-TDI, F1 \uparrow				ADReSS, F1 \uparrow				IEMOCAP, Acc. (5-fold) \uparrow			
	A	B	MV	SMV	A	B	MV	SMV	A	B	MV	SMV
No VC	54.9	54.9	54.9	54.9	70.6	70.6	70.6	70.6	71.5	71.5	71.5	71.5
Pitch shifting	55.8	60.3	57.6	<u>61.5</u>	71.2	77.1	77.1	68.8	60.6	55.1	61.1	61.1
KNN-VC	<u>55.8</u>	<u>61.7</u>	<u>64.8</u>	49.9	71.5	<u>79.2</u>	<u>79.2</u>	<u>83.3</u>	70.4	69.3	71.4	71.5
TriAAN-VC	55.7	60.7	61.7	53.3	<u>72.4</u>	75.0	77.1	<u>83.3</u>	65.1	64.1	66.8	67.2
Diff-VC	47.0	51.2	50.3	49.2	65.6	69.4	66.7	70.8	<u>87.0</u>	<u>94.3</u>	<u>96.5</u>	<u>97.2</u>

Table 7. Emotion recognition results on IEMOCAP. 8 speakers in the training set of each fold are used as target speakers.

VC type	Voting type	Accuracy					
		1	2	3	4	5	Avg.
No VC	-	72.6	76.6	68.9	68.9	70.3	71.5
Pitch shifting	single (best)	64.0	65.3	57.4	57.7	58.6	60.6
	single (avg)	61.3	62.1	50.2	48.9	53.0	55.1
	majority	61.2	65.3	58.5	57.8	62.5	61.1
	soft majority	60.8	65.4	57.8	57.5	61.5	61.1
KNN-VC	single (best)	71.2	75.4	68.3	71.9	69.1	70.4
	single (avg)	69.6	72.6	67.0	69.9	67.4	69.3
	majority	70.3	75.6	68.5	72.8	70.0	71.4
	soft majority	70.3	76.1	68.5	72.8	69.9	71.5
TriAAN-VC	single (best)	65.5	66.9	63.0	67.5	62.6	65.1
	single (avg)	64.6	66.3	61.1	65.6	63.1	64.1
	majority	66.9	69.0	63.9	67.9	66.5	66.8
	soft majority	66.6	69.9	63.6	68.8	67.0	67.2
Diff-VC	single (avg)	87.0	88.3	86.2	87.6	86.1	87.0
	single (best)	94.4	94.8	94.2	95.2	92.9	94.3
	majority	97.5	96.7	95.2	98.1	94.9	96.5
	soft majority	97.7	97.6	96.3	98.7	95.6	97.2

Table 8. Alzheimer detection results on ADReSS

VC type	Voting type	Precision	Recall	F1	Accuracy
No VC	-	71.4	70.8	70.6	70.8
Pitch shifting	single (avg)	71.8	71.4	71.4	71.2
	single (best)	77.1	77.1	77.1	77.1
	majority	77.1	77.1	77.1	77.1
	soft majority	68.8	68.8	68.7	68.8
KNN-VC	single (avg)	71.8	71.5	71.4	71.5
	single (best)	80.0	79.2	79.1	79.2
	majority	79.4	79.2	79.1	79.2
	soft majority	83.6	83.3	83.3	83.3
TriAAN-VC	single (avg)	72.5	72.4	72.4	72.4
	single (best)	75.2	75.0	75.0	75.0
	majority	77.5	77.1	77.0	77.1
	soft majority	83.3	83.3	83.3	83.3
Diff-VC	single (avg)	65.7	65.4	65.4	65.6
	single (best)	69.4	69.4	69.4	69.4
	majority	66.7	66.7	66.7	66.7
	soft majority	72.2	70.8	70.4	70.8

Table 9. ALS severity classification results on ALS-TDI with a whisper-medium+SVM classifier

VC type	Voting type	Precision \uparrow	Recall \uparrow	F1 \uparrow
No VC	-	59.8	53.7	54.9
Pitch shifting	single (avg.)	60.5	54.1	55.8
	single (best)	67.4	57.7	60.3
	majority	73.0	54.9	57.6
	soft majority	68.4	59.0	61.5
KNN-VC	single (avg.)	58.5	54.6	55.8
	single (best)	65.7	59.5	61.7
	majority	67.9	62.9	64.8
	soft majority	51.1	49.6	49.9
TriAAN-VC	single (avg.)	60.2	54.5	55.7
	single (best)	68.0	58.2	60.7
	majority	69.9	59.1	61.7
	soft majority	54.1	52.8	53.3
Diff-VC	single (avg.)	48.2	47.0	47.0
	single (best)	53.0	51.0	51.2
	majority	49.8	50.9	50.3
	soft majority	50.1	48.8	49.2