## Semiparametric Counterfactual Regression

Kwangho Kim\*

Department of Statistics, Korea University

kwanghk@korea.ac.kr

#### Abstract

We study counterfactual regression, which aims to map input features to outcomes under hypothetical scenarios that differ from those observed in the data. This is particularly useful for decision-making when adapting to sudden shifts in treatment patterns is essential. We propose a doubly robust-style estimator for counterfactual regression within a generalizable framework that accommodates a broad class of risk functions and flexible constraints, drawing on tools from semiparametric theory and stochastic optimization. Our approach uses incremental interventions to enhance adaptability while maintaining consistency with standard methods. We formulate the target estimand as the optimal solution to a stochastic optimization problem and develop an efficient estimation strategy, where we can leverage rapid development of modern optimization algorithms. We go on to analyze the rates of convergence and characterize the asymptotic distributions. Our analysis shows that the proposed estimators can achieve  $\sqrt{n}$ -consistency and asymptotic normality for a broad class of problems. Numerical illustrations highlight their effectiveness in adapting to unseen counterfactual scenarios while maintaining parametric convergence rates.

*Keywords:* Causal Inference; Counterfactual Prediction; Incremental Intervention; Dataset shift; Transportability.

<sup>\*</sup>The author thanks Edward Kennedy and Alec McClean for very helpful discussions. R scripts for various examples, motivating illustrations, and simulation studies are available on GitHub at https://github.com/kwangho-joshua-kim/counterfactual-prediction.

## 1 Introduction

Counterfactuals (or potential outcomes) describe how certain aspects of the world would have been different under hypothetical scenarios. They are widely employed in causal inference to explore questions like "what would have happened if X had occurred", even when X never actually took place. In fact, causal inference tasks can be understood as comparing outcomes, or, more generally, functionals, across two or more counterfactual distributions.

Recently, counterfactuals have proven useful for predicting outcomes in deployment settings that differ significantly from the training phase. This is commonly referred to as *counterfactual prediction*. In counterfactual prediction, the goal is to map input features to outcomes under hypothetical scenarios that may differ substantially from the observed world (e.g., what if treatment access were limited to 50% of individuals compared to current practice?). This can be particularly useful to inform decision-making in clinical practice, where we need to address abrupt shifts in treatment patterns (e.g., Hernán et al. 2019; Dickerman and Hernán 2020; van Geloven et al. 2020; Dickerman et al. 2022). The recent strikes of junior doctors in South Korea may serve as a representative example (Lancet Editorial 2024; Park 2024). Moreover, certain interventions in the training data may be undesirable or incompatible with the target deployment setting, requiring their removal via prediction under a hypothetical intervention that completely excludes them (e.g., Sperrin et al. 2018). This is especially relevant in clinical research when predicting risk as to treatments initiated after baseline (Schulam and Saria 2017; van Geloven et al. 2020; Lin et al. 2021).

More generally, counterfactual prediction may be used when transferring a model from a source population to a similar target population that differs only in post-baseline treatment patterns. Therefore, counterfactual prediction is closely related to domain adaptation and out-of-distribution generalization, especially when distributional shifts result from specific interventions. In principle, retraining the model with data collected from the new setting is preferred, but due to cost or as an interim solution, one may instead seek to adapt the existing model to approximate outcomes under a hypothetical intervention that aligns with treatment patterns in the target setting.

Counterfactual prediction presents challenges absent in standard prediction tasks, as the data required to construct predictive models are inherently unobservable in their entirety. Surprisingly, while recent advances in predictive modeling have significantly enhanced causal inference, the reverse, leveraging causal inference to address challenges in predictive modeling, has received much less attention. Counterfactual prediction is closely related to the estimation of the conditional average treatment effect (CATE), which is represented as the contrast between two distinct counterfactual regression functions. However, unlike counter-

factual prediction, efficient estimation of the CATE can exploit a structure much simpler than that of individual counterfactual regression functions (Kennedy 2020). Also, in multivalued treatment settings, estimating each counterfactual regression function separately is often more practical than computing all relative-effect combinations.

Some recent progress has been made for counterfactual prediction beyond its role within the CATE estimation framework. To address covariate distributional shifts induced by varying treatments, several studies have adopted direct modeling, or plug-in, approaches grounded in the parametric g-formula. (e.g., Li et al. 2016; Schulam and Saria 2017; Nguyen et al. 2020; Lin et al. 2021; Dickerman et al. 2022). Coston, Kennedy and Chouldechova (2020); Coston, Mishler, Kennedy and Chouldechova (2020) introduced a nonparametric method for counterfactual prediction and associated risk, addressing runtime confounding and fairness. Other approaches utilize representation learning techniques to construct a common representation space that balances the source and target domains (e.g., Johansson et al. 2016; Shalit et al. 2017; Hassanpour and Greiner 2019a,0). While previous work has focused primarily on deterministic interventions, McClean et al. (2024) develop nonparametric methods for estimating conditional effects, defined under a specific class of stochastic interventions known as incremental interventions. However, a generic learning framework to efficiently and flexibly predict counterfactual outcomes still remains absent.

Our work aims to fill this gap in the literature. We propose doubly robust-style estimators for counterfactual regression that can accommodate a broad class of risk functions and flexible constraints (e.g., shape, fairness), leveraging techniques at the intersection of semiparametric theory and stochastic optimization. As in McClean et al. (2024), our hypothetical interventions, or treatment policies, are formulated using incremental interventions (Kennedy 2019), which shift the propensity score rather than assigning treatments to fixed values, while still allowing recovery of standard deterministic interventions as a special case. Our target estimand approximates the regression function by projecting it onto a finite-dimensional model space, guided by a specific class of risk and constraints. We show that the target parameters can be framed as optimal solutions to a stochastic program. Subsequently, we propose an estimation strategy by formulating and solving an appropriate approximating program, which is derived by characterizing the efficient influence functions of each coefficient in the true stochastic optimization problem. We go on to derive a closed-form expression for the asymptotic distribution of our estimator, and show that the proposed estimator can be  $\sqrt{n}$ -consistent and asymptotically normal under relatively weak regularity conditions. Experiments demonstrate that the proposed estimator not only effectively adapts to unseen counterfactual scenarios but also attains parametric rates of convergence.

## 2 Framework

### 2.1 Setup

Suppose that we have access to an i.i.d. sample  $(Z_1, ..., Z_n)$  of n tuples  $Z = (Y, A, X) \sim \mathbb{P}$ , where  $Y \in \mathcal{Y}$  represents the outcome,  $A \in \{0, 1\}$  denotes a binary intervention, and  $X \in \mathcal{X} \subseteq \mathbb{R}^p$  comprises observed covariates. Here, we assume A is binary for simplicity, but in principle it can be multi-valued. In general, we consider  $\mathcal{Y}$  as a subset of Euclidean space, though it may also be discrete. We let  $Y^a$  denote the counterfactual (or potential) outcome that would be observed under treatment assignment  $A = a, a \in \mathcal{A}$ . Throughout, we refer to a *factual prediction* as a mapping from a set of predictors to the observable outcome Y.

In contrast to previous approaches to counterfactual prediction discussed in Section 1, we employ incremental interventions to generate our target counterfactual scenarios with greater flexibility. *Incremental interventions* are a specialized type of stochastic intervention that shifts the exposure propensity of each unit by a predetermined amount (Kennedy 2019). Specifically, let  $\pi(X) = \mathbb{P}(A = 1 \mid X)$  denote the observational propensity score. Then the incremental intervention replaces  $\pi$  with the distribution defined by

$$q(X;\delta,\pi) = \frac{\delta\pi(X)}{\delta\pi(X) + 1 - \pi(X)} \quad \text{for} \quad 0 < \delta < \infty,$$

where  $\delta$  is a user-specified increment parameter, indicating how the intervention changes the odds of receiving treatment; if  $\delta > 1$  (< 1), then the intervention increases (decreases) the odds. This corresponds to a counterfactual scenario in which individuals receive a hypothetical treatment  $A^*$ , where  $A^* \mid X \sim \text{Bernoulli}\{q(X;\delta,\pi)\}$ . There are at least two key reasons why incremental interventions are particularly appealing. First, they more accurately capture treatment variations that may naturally occur in practice. More importantly, they completely eliminate the need for the stringent positivity assumption. Their extensions have also been explored in studies on dropout (Kim et al. 2021), conditional effects (McClean et al. 2024), and continuous exposures (Schindl et al. 2024), among others.

Let  $Q(a|x;\delta)$  denote the *intervention distribution*, the probability distribution of treatment assignment A = a for the incremental intervention associated with  $q(X;\delta,\pi)$ , i.e.,  $dQ(a|X;\delta) = q(X;\delta,\pi)^a \{1 - q(X;\delta,\pi)\}^{1-a}$ . Let  $Q(\delta)$  represent draws from  $Q(a|x;\delta)$ . In our counterfactual prediction tasks, the primary quantity of interest is the counterfactual outcome  $Y^{Q(\delta)}$ . Throughout, we assume *consistency*,  $Y = Y^a$  if A = a, and *no unmeasured confounding*,  $A \perp Y^a \mid X$ . Kennedy (2019) showed that under the assumptions of consistency and no unmeasured confounding, the mean counterfactual outcome under  $Q(a|x;\delta)$  can be identified as

$$\mathbb{E}(Y^{Q(\delta)}) = \int_{\mathcal{A}} \int_{\mathcal{X}} \mathbb{E}(Y \mid X = x, A = a) \, dQ(a \mid x; \delta) \, d\mathbb{P}(x) = \mathbb{E}\left[\frac{\delta\pi(X)\mu_1(X) + \{1 - \pi(X)\}\mu_0(X)}{\delta\pi(X) + \{1 - \pi(X)\}}\right] = \mathbb{E}\left[\frac{Y(\delta A + 1 - A)}{\delta\pi(X) + \{1 - \pi(X)\}}\right], \quad (1)$$

where  $\mu_a(X) = \mathbb{E}(Y \mid X, A = a)$ . However, in counterfactual regression, our goal is to obtain an accurate prediction of  $Y^{Q(\delta)}$  given X, rather than to estimate the marginal mean  $\mathbb{E}(Y^{Q(\delta)})$ . We allow  $\delta = 0$  and  $\delta = \infty$ , which correspond to the deterministic interventions A = 0 and A = 1, respectively. Note that this requires the *positivity assumption*,  $\varepsilon < \pi(X) < 1 - \varepsilon$  a.s. for some  $\varepsilon > 0$ , in addition to the assumptions of consistency and no unmeasured confounding.

**Notation.** We introduce some notation used in this paper. For any fixed vector v and matrix M, we let  $||v||_2$  and  $||M||_F$  denote the Euclidean norm and the Frobenius norm, respectively.  $|| \cdot ||_2$  is understood as the spectral norm when it is used with a matrix. Let  $\mathbb{P}_n$  denote the empirical measure over  $(Z_1, ..., Z_n)$ . In addition, we use  $||f||_{2,\mathbb{P}}$  to denote the  $L_2(\mathbb{P})$  norm of f defined by  $||f||_{2,\mathbb{P}} = [\int f(z)^2 d\mathbb{P}(z)]^{1/2}$ . Lastly, we let  $s^*(\mathsf{P})$  denote the set of optimal solutions of an optimization program  $\mathsf{P}$ , and define  $\mathsf{dist}(x, \mathcal{S}) = \inf \{||x - y||_2 : y \in \mathcal{S}\}$  as the distance from a point x to a set S.

### 2.2 Motivating Illustration

Consider a simple data-generating process where the covariates  $X \sim \text{Uniform}[0, 10]$  and,

$$A \sim \text{Bernoulli}(\text{expit}(2.8 - 0.3X)), \quad Y \sim N(1 + 0.75X + 0.5X^2 - 10A, X),$$

where  $expit(\cdot)$  denotes the inverse logit function. Under this model, approximately 75.4% of individuals initiate treatment; however, as X increases, the likelihood of treatment initiation decreases. Moreover, individuals who are less likely to receive treatment tend to exhibit greater variability in their treatment effects.

Suppose that we are concerned with the expected outcome under a hypothetical treatment policy governed by the intervention distribution  $Q(a|x; \delta)$ . This scenario may represent a counterfactual inquiry: for example, given the baseline predictors, what would be the risk of death if access to surgery were restricted to only a randomly selected x% of individuals? (Dickerman and Hernán 2020). Thus, our objective is to predict  $Y^{Q(\delta)}$  from X. When the shift is substantial (i.e., when  $\delta$  deviates significantly from 1), the factual prediction model is expected to perform poorly. A small  $\delta(\delta \ll 1)$  can correspond to a scenario in which



Figure 1: (a) Density of the intervention distribution  $q(X; \delta, \pi)$  for varying values of  $\delta$ ; (b) and (c) display the densities of the true counterfactual outcomes and the predicted outcomes from factual and proposed counterfactual regression methods for  $\delta = 0.1$  and  $\delta = 0.01$ , respectively. The factual regression exhibits a notable distributional discrepancy from the true counterfactual outcomes, resulting in substantial estimation errors.

physicians suddenly go on strike (e.g., Park 2024). Here, we posit that no new data are available for model adaptation or retraining.

We consider a counterfactual scenario in which the odds of receiving treatment are substantially reduced by a factor of 0.1 and 0.01. We suppose that the propensity score  $\pi$  is unknown and is estimated using logistic regression. Training and test datasets of equal size n = 1000 are generated, where in the test data, A is sampled from Bernoulli $\{q(X; \delta, \pi)\}$ . The factual regression corresponds to the outcome regression model  $\mu_a$ , estimated via ordinary least squares with higher-order polynomial terms, where the treatment variable A is imputed using  $\hat{\pi}$ . For counterfactual regression, we apply the proposed method with the  $L_2$  loss without any constraints (Example 2C). We assume that both the outcome and propensity score models are correctly specified in their parametric forms.

This simple example illustrates an interesting, yet presumably common phenomenon. As  $\delta$  decreases, the probability of receiving treatment declines sharply compared to the original propensity score  $\pi$ , particularly for small values of X (Figure 1a). Figures 1b and c show that the factual regression fails to generalize to these counterfactual scenarios, while the proposed counterfactual regression successfully adapts. The outcomes predicted by the factual regression exhibit a substantial distributional discrepancy from the true counterfactual outcomes. The discrepancy becomes even more pronounced as  $\delta$  decreases from 0.1 to 0.01, whereas the proposed estimator consistently generalizes well across varying values of  $\delta$ . A similar phenomenon is observed even when more complex nonparametric models are used

for the outcome regression, or when the parametric models are misspecified.

### 2.3 Estimand

Learning tasks are generally characterized through risk functions, and across domains, a wide variety of nontraditional criteria are often employed to design and evaluate machine learning algorithms (Wang et al. 2020). Given a loss function L, we define the risk of a regression model  $f : \mathcal{X} \to \mathbb{R}$  as  $\mathcal{R}\left(f(X), Y^{Q(\delta)}\right) = \mathbb{E}\left\{L\left(f(X), Y^{Q(\delta)}\right)\right\}$ . For instance, when the  $L_2$  loss (or squared error loss) is used without any model constraints, the function f that minimizes  $\mathcal{R}(f(X), Y^{Q(\delta)})$  is given by the conditional mean  $\mathbb{E}\{Y^{Q(\delta)} \mid X\}$ , which corresponds to the conditional incremental effects studied in McClean et al. (2024). We consider some specified finite-dimensional parametric model  $\{f(x; \beta) : \beta \in \mathbb{R}^k\}$  as the regression model for  $Y^{Q(\delta)}$ . Our target parameter can be formulated as the optimal solution to the following stochastic program:

$$\underset{\beta \in \mathbb{R}^k}{\text{minimize}} \quad \mathcal{R}\left(f(X;\beta), Y^{Q(\delta)}\right) + \lambda \rho(\beta) \quad \text{subject to} \quad \beta \in \mathcal{C},$$
(2)

where the constraint set is given by  $C = \{\beta : g_j(\beta) \leq 0, j = 1, ..., r\}$  for deterministic functions  $g_j : \mathbb{R}^k \to \mathbb{R}$ . Here,  $\rho(\cdot)$  is a pre-specified penalty function, and  $\lambda \geq 0$  is the associated tuning parameter. The solution to the above program corresponds to the coefficients of the best-fitting regression function for  $Y^{Q(\delta)}$  within the class  $\{f(x;\beta) : \beta \in \mathbb{R}^k\}$ , subject to a set of constraints C. Since we make no assumptions about the true functional relationship between  $Y^{Q(\delta)}$  and X, all our results are formally nonparametric. The above projection approach, where a model is not assumed to be correct but is instead used solely to define approximations, has been widely employed in statistics and causal inference (see, for example, Kennedy et al. 2021, Section 3.1.1).

**Remark 2.1.** Another advantage of our framework is that it accommodates a general setting where only a subset of covariates  $V \subseteq X$  is available at the time of prediction. This flexibility allows for scenarios like runtime confounding, where certain factors used by decision-makers are recorded in the training data (for constructing nuisance estimates) but are unavailable during prediction. For further details, see, for example, Coston, Kennedy and Chouldechova (2020); Kim, Mishler and Zubizarreta (2022).

**Models.** We present a few illustrative examples of model classes for  $f(x;\beta)$ . An archetypal example is a class of generalized linear models, where the counterfactual outcome has a distribution that is in the exponential family. More generally, one may consider a functional

aggregation or linear ensemble in the following form:

$$f(X;\beta) = \sum_{j=1}^{k} \beta_j b_j(X), \qquad (3)$$

with a set of known basis functions  $\{b_j\}$ . Examples of such basis functions include orthogonal polynomials, wavelets, and splines. Aggregated predictors of this form have been extensively studied and broadly applied across a wide range of disciplines (e.g., Tsybakov 2003; Nemirovski et al. 2009; Polley and Van der Laan 2010).

**Constraints.** We often seek to incorporate various constraints into our regression models. A few examples that may be included in the constraint set C are presented below.

**Example 1A** (Constrained regression). There is a growing demand for the use of constrained regression in various scientific contexts. For instance, a set of linear inequality constraints  $C\beta \leq d$ , where  $C \in \mathbb{R}^{r \times k}$ ,  $d \in \mathbb{R}^r$ , can be employed to enforce shape constraints (James et al. 2013; Gaines et al. 2018), ensure structural consistency (Liew 1976), or maintain physical fidelity in calibration models (Schwendinger et al. 2024). The same type of linear inequality constraints can be utilized to target specific sub-populations in website advertising (James et al. 2019), design marketing mix models (Chen et al. 2021), and account for the compositional nature of microbiome data (Lu et al. 2019). More general constraints, where coefficients may depend on X, are also employed in interpretable machine learning (e.g., Zeng et al. 2017).

The functions  $\{g_j\}$  in the constraint set  $\mathcal{C}$  do not need to be known or fixed a priori. If  $\{g_j\}$  depend on  $\mathbb{P}$  or  $Y^{Q(\delta)}$ , they must also be estimated from the data. Some examples are given below.

**Example 1B** (Algorithmic fairness). Data-driven policy making can inadvertently result in discriminatory treatment of sensitive groups (e.g., gender, race). To mitigate such biases, fairness metrics impose constraints on the joint distribution of outcomes and sensitive features (e.g., Hardt et al. 2016; Corbett-Davies et al. 2017; Barocas et al. 2023). To evaluate unfairness within each subgroup, we utilize the fairness function  $uf: \mathcal{Y} \times \mathcal{X} \times \{0,1\}^2 \to \mathbb{R}$ , which accommodates a broad range of (counterfactual) fairness measures (e.g., Mishler and Kennedy 2021; Kim and Zubizarreta 2023). Let  $F \in \mathcal{X}$  be a binary sensitive feature. Using the fairness function, our fairness criterion can be expressed at the population level as:

$$|\mathbb{E}\left\{uf(Z)f(X;\beta)\right\}| \le \epsilon,\tag{4}$$

where  $\epsilon \geq 0$  represents the acceptable fairness threshold. For instance, the criteria of statis-

tical parity and conditional statistical parity, which are among the most widely recognized fairness measures, can be implemented by defining the fairness functions as:

$$uf(Z) = \frac{1-F}{\mathbb{E}(1-F)} - \frac{F}{\mathbb{E}(F)},\tag{5}$$

$$uf(Z) = \frac{(1-F)\mathbb{1}\{L=l\}}{\mathbb{E}[(1-F)\mathbb{1}\{L=l\}]} - \frac{F\mathbb{1}\{L=l\}}{\mathbb{E}[F\mathbb{1}\{L=l\}]},$$
(6)

where L is some function of X and represents a legitimate factor used to specify conditional parity, leading to

$$\begin{split} |\mathbb{E}\left\{f(X;\beta) \mid F=0\right\} - \mathbb{E}\left\{f(X;\beta) \mid F=1\right\}| &\leq \epsilon, \\ |\mathbb{E}\left\{f(X;\beta) \mid F=0, L=l\right\} - \mathbb{E}\left\{f(X;\beta) \mid F=1, L=l\right\}| &\leq \epsilon, \end{split}$$

respectively. These conditions require the regression model to be marginally (or conditionally) independent of the given sensitive feature.

**Example 1C** (Balance for the positive class). In many practical settings, such as criminal justice and healthcare policy, it is important to ensure that a model's decisions remain independent of individuals who exhibit a certain characteristic (e.g., a risk score exceeding a threshold), even if the model can only identify them imperfectly (e.g., Kleinberg et al. 2016). In this case, with  $Y^{Q(\delta)}$  viewed as a risk score, one may wish to incorporate the following balancing condition into the modeling framework:

$$uf(Z) = \frac{(1-S)\mathbb{1}\left\{Y^{Q(\delta)} > 0\right\}}{\mathbb{E}\left[(1-S)\mathbb{1}\left\{Y^{Q(\delta)} > 0\right\}\right]} - \frac{S\mathbb{1}\left\{Y^{Q(\delta)} > 0\right\}}{\mathbb{E}\left[S\mathbb{1}\left\{Y^{Q(\delta)} > 0\right\}\right]},\tag{7}$$

which leads to the fairness criterion

$$\left| \mathbb{E}\left\{ f(X;\beta) \mid F = 0, Y^{Q(\delta)} > 0 \right\} - \mathbb{E}\left\{ f(X;\beta) \mid F = 1, Y^{Q(\delta)} > 0 \right\} \right| \le \epsilon.$$

**Challenges.** A key challenge stems from the counterfactual nature of the functionals involved in the optimization problem (2). The two most widely used approaches in stochastic programming are stochastic approximation (SA) and sample average approximation (SAA) (e.g., Nemirovski et al. 2009; Shapiro et al. 2014). Such standard methods typically assume that each unknown component is of the form  $\mathbb{E}[H(\beta,\xi)]$ , where *H* is a fixed, known, real-valued function, and  $\xi$  is a random vector whose distribution is  $\mathbb{P}$ . Additionally, the following assumptions are commonly made: i) an iid sample of realizations,  $\xi_1, \xi_2, \ldots, \xi_n$ from the random vector  $\xi$  can be generated; ii) for each pair  $(\beta, \xi)$ , the value  $H(\beta, \xi)$  or its stochastic subgradients are accessible. In such settings, sample mean constructions or subgradient descent methods are readily applicable. However, these standard methods are not directly applicable to our framework, as the optimization problem involves functionals of unobserved counterfactuals, whose identification relies on unknown nuisance functions. Moreover, they cannot leverage efficient estimators for counterfactual components, such as semiparametric estimators with cross-fitting (Chernozhukov et al. 2017; Newey and Robins 2018). Therefore, more general approaches are required beyond the standard SA and SAA frameworks.

## 3 Estimation

In this section, we develop an estimation strategy for the target parameter defined as the solution to the optimization problem (2). Our estimation procedure can be summarized as the following three-step sequencing:

- **Step 1.** Derive the efficient influence function for each coefficient that requires estimation in the target stochastic program (2).
- **Step 2.** Construct the efficient semiparametric estimators for the coefficients in (2) based on the efficient influence functions derived in Step 1, and utilize these estimators to formulate the corresponding approximating program for (2).
- **Step 3.** Solve an optimization problem equivalent to the approximating program defined in Step 2.

We provide a more detailed discussion about each step in the proposed strategy. Our goal in Step 1 is to find the efficient influence function (EIF) for each unknown component in our target program (2). The EIF enables construction of the efficient semiparametric estimator by de-biasing generic plug-in estimators, where we may achieve local minimax lower bounds (Bickel et al. 1993; van der Vaart 2002; Tsiatis 2006; Kennedy 2016). This also yields desirable properties for our main estimator in Step 3, such as double robustness or general second-order bias, which allows us to relax nonparametric conditions on nuisance function estimation.

The EIF for  $\mathbb{E}\{Y^{Q(\delta)}\}\$  can be useful in specifying the EIF for various functionals involving  $Y^{Q(\delta)}$ . For convenience, we let  $\mu_a(X) = \mathbb{E}[Y \mid X, A = a], \pi_a(X) = \mathbb{P}(A = a \mid X), a \in \{0, 1\},$  and  $\eta_Q = (\mu_a, \pi_a)$ . By Kennedy (2019, Theorem 2), the uncentered EIF for  $\mathbb{E}\{Y^{Q(\delta)}\}\$  is given as

$$\varphi_Q(Z;\eta_Q,\delta) = \frac{\delta\pi_1(X)\phi_1(Z) + \pi_0(X)\phi_0(Z)}{\delta\pi_1(X) + \pi_0(X)} + \frac{\delta\Delta(X)\{A - \pi_1(X)\}}{\{\delta\pi_1(X) + \pi_0(X)\}^2},\tag{8}$$

where  $\Delta(X) = \mu_1(X) - \mu_0(X)$ , and  $\phi_a(Z) = \frac{\mathbb{1}(A=a)}{\pi_a(X)} \{Y - \mu_a(X)\} + \mu_a(X)$ . For instance, it can be deduced that for an arbitrary fixed real-valued function  $\tau : \mathcal{X} \to \mathbb{R}$ , the uncentered EIF for the parameter  $\mathbb{E}\{Y^{Q(\delta)}\tau(X)\}$  is simply given by  $\varphi_Q(Z;\eta_Q,\delta)\tau(X)$  (Kim, Kennedy and Zubizarreta 2022, Lemma A.1). More practical and relevant examples will be provided shortly.

Once the EIF is specified, in Step 2, we construct an efficient semiparametric estimator for each coefficient in (2), essentially as a sample average of the EIF with the estimated nuisance components, i.e., *one-step estimator*. To our knowledge, there are two main approaches for constructing such efficient estimators; one is based on empirical process conditions, and the other is to use sample splitting. One may assume that the function class for the EIF and corresponding estimators are not too complex (e.g., Donsker or low-entropy type conditions), but this would limit the flexibility of the nuisance estimators. To avoid this, alternatively, we can use sample splitting (or cross-fitting) to allow for arbitrarily complex nuisance estimators. Both approaches can be viewed as ways to avoid using the same data twice, one for constructing relevant nuisance components, and the other for de-biasing, which can introduce a threat of overfitting (Chernozhukov et al. 2018). Here, we are agnostic about which approach will be used. We refer the interested readers to Kennedy (2016,0) and references therein.

Subsequently, we formulate an approximating (sample-based) program of (2) by substituting all the unknown coefficients with their estimators. Let  $\varphi_{\mathcal{R}}(Z; \eta_{\mathcal{R}}, \delta, \beta)$  be the uncentered EIF for  $\mathcal{R}\left(f(X;\beta), Y^{Q(\delta)}\right)$  with the corresponding nuisance functions  $\eta_{\mathcal{R}}$ , so that  $\mathcal{R}\left(f(X;\beta), Y^{Q(\delta)}\right) = \mathbb{E}\{\varphi_{\mathcal{R}}(Z;\eta_{\mathcal{R}},\delta,\beta)\}$ . Then, the semiparametric estimator for our risk function is given by  $\mathbb{P}_n\{\varphi_{\mathcal{R}}(Z;\hat{\eta}_{\mathcal{R}},\delta,\beta)\}$ , provided that  $\hat{\eta}_{\mathcal{R}}$  converges to  $\eta_{\mathcal{R}}$  at sufficiently fast rates. As discussed above,  $\hat{\eta}_{\mathcal{R}}$  can be constructed on the same sample if we are willing to rely on appropriate empirical process conditions, or, alternatively, on a separate, independent sample using sample splitting. The approximating program can be generically formulated as follows:

$$\underset{\beta \in \mathbb{R}^k}{\text{minimize}} \quad \mathbb{P}_n \left\{ \varphi_{\mathcal{R}}(Z; \hat{\eta}_{\mathcal{R}}, \delta, \beta) \right\} + \lambda \rho(\beta) \quad \text{subject to} \quad \beta \in \widehat{\mathcal{C}}, \tag{9}$$

where  $\eta_{\mathcal{R}}$  is a set of the relevant nuisance functions, and  $\widehat{\mathcal{C}}$  represents the estimated feasible set  $\{\beta : \widehat{g}_j(\beta) \leq 0, j = 1, \ldots, r\}$ . Here, each  $\widehat{g}_j$  is the EIF-based semiparametric estimator, constructed as discussed above.

It can be shown, using standard arguments from the semiparametric literature, that each stochastic component of the approximating program (9) serves as a  $\sqrt{n}$ -consistent, asymptotically normal, and efficient estimator for its counterpart in (2) (Kennedy 2022, Section

4). For example, when  $\mathcal{R}$  is smooth, we have that  $\forall \beta, \delta$ ,

$$\mathbb{P}_{n}\left\{\varphi_{\mathcal{R}}(Z;\hat{\eta}_{\mathcal{R}},\delta,\beta)\right\} - \mathcal{R}\left(f(X;\beta),Y^{Q(\delta)}\right) \xrightarrow{d} N\left(0,\operatorname{var}(\varphi_{\mathcal{R}}(Z;\eta_{\mathcal{R}},\delta,\beta))\right),\tag{10}$$

if the second-order von Mises remainder term  $\mathbb{P}_n \{\varphi_{\mathcal{R}}(Z; \hat{\eta}_{\mathcal{R}}, \delta, \beta)\} - \mathbb{E}\{\varphi_{\mathcal{R}}(Z; \eta_{\mathcal{R}}, \delta, \beta)\} + \int \varphi_{\mathcal{R}}(z; \hat{\eta}_{\mathcal{R}}, \delta, \beta) d\mathbb{P}(z)$  vanishes at a rate faster than  $\sqrt{n}$ , along with appropriate empirical process conditions.

Finally, in Step 3, the approximating program derived in Step 2 (or its equivalent reformulation) can be solved using various off-the-shelf optimization solvers. An important benefit of the proposed approach is that, depending on the problem, one can leverage rapid development of modern optimization algorithms (e.g., Boyd et al. 2011; Bertsekas 2015; Jain et al. 2017). Finally, our proposed estimator for counterfactual regression is  $f(X; \hat{\beta})$ , where  $\hat{\beta}$  is the optimal solution to the approximating program from Step 3.

In what follows, we illustrate the proposed estimation procedure through examples.

**Example 2A** (Cross-entropy loss). The proposed framework is also applicable to a classification task. For simplicity, assume  $\mathcal{Y} = \{0, 1\}$ . Then we may consider the cross-entropy loss  $L(y, \hat{y}) = -y \log \hat{y} - (1 - y) \log(1 - \hat{y})$ . For a generalized linear model  $f(X; \beta)$ , the corresponding risk is defined by  $-\mathbb{E}\left\{Y^{Q(\delta)} \log f(X; \beta) + (1 - Y^{Q(\delta)}) \log(1 - f(X; \beta))\right\}$ , yielding the EIF

$$-\mathbb{E}\left\{\varphi_Q(Z;\eta_Q,\delta)\log f(X;\beta) + (1-\varphi_Q(Z;\eta_Q,\delta))\log(1-f(X;\beta))\right\}.$$

This leads to the following approximating program:

$$\underset{\beta \in \mathbb{R}^k}{\text{minimize}} \quad -\mathbb{P}_n\left\{\varphi_Q(Z; \hat{\eta}_Q, \delta) \log f(X; \beta) + (1 - \varphi_Q(Z; \hat{\eta}_Q, \delta)) \log(1 - f(X; \beta))\right\}.$$

Kim, Kennedy and Zubizarreta (2022) studied counterfactual classification within the framework developed in this work, specifically focusing on the case of a deterministic intervention ( $\delta = 0$ ) and deterministic constraints, where C is known and fixed.

**Example 2B** (Mean squared logarithmic loss). Consider a generalized linear model  $f(X;\beta)$ , and the mean squared logarithmic loss defined by  $L(y,\hat{y}) = \{\log(1+y) - \log(1+\hat{y})\}^2$  with the  $L_2$  penalty function. This loss function is known to be particularly useful when the data exhibit a wide range of values (Hodson et al. 2021). In this case, the EIF for the risk  $\mathbb{E}[\{\log(Y^{Q(\delta)}+1) - \log(f(X;\beta)+1)\}^2]$  is given by

$$\log^{2} (f(X;\beta)+1) - 2 \left\{ \log (f(X;\beta)+1) \varphi'_{Q}(Z;\eta'_{Q},\delta) \right\} + \varphi''_{Q}(Z;\eta''_{Q},\delta),$$

where  $\varphi'_Q(Z; \eta'_Q, \delta)$  ( $\varphi''_Q(Z; \eta''_Q, \delta)$ ) is the same as  $\varphi_Q$  in (8) except that  $\mu_a(X)$  and  $\phi_a(Z)$  are replaced by  $\mu'_a(X) = \mathbb{E}[\log(Y+1) \mid X, A = a]$  ( $\mu''_a(X) = \mathbb{E}[\log^2(Y+1) \mid X, A = a]$ ) and  $\phi'_a(Z) = \frac{\mathbb{I}(A=a)}{\pi_a(X)} \{\log(Y+1) - \mu'_a(X)\} + \mu'_a(X)$  ( $\phi''_a(Z) = \frac{\mathbb{I}(A=a)}{\pi_a(X)} \{\log^2(Y+1) - \mu''_a(X)\} + \mu''_a(X))$ , respectively, with  $\eta'_Q = (\mu'_a, \pi_a)$  ( $\eta''_Q = (\mu''_a, \pi_a)$ ). Then our approximating problem is equivalent to:

$$\underset{\beta \in \mathbb{R}^k}{\text{minimize}} \quad \frac{1}{2} \mathbb{P}_n \left\{ \log^2 \left( f(X;\beta) + 1 \right) \right\} - \mathbb{P}_n \left\{ \log \left( f(X;\beta) + 1 \right) \varphi_Q'(Z;\hat{\eta}_Q',\delta) \right\} + \lambda \|\beta\|_2^2 \right\}$$

**Example 2C** ( $L_2$  loss with fairness criterion). One of the most commonly used loss functions is the squared error loss, or  $L_2$  loss:  $L = (y - \hat{y})^2$ . Consider the linear aggregation form (3), with  $L_2$  loss and  $L_2$  penalty. Then, the risk and the EIF are given by  $\mathbb{E}\{Y^{Q(\delta)} - b(X)^{\top}\beta\}^2$ and

$$\varphi_Q'(Z;\eta_Q',\delta) - 2\beta^\top \varphi_Q(Z;\eta_Q,\delta)b(X) + \beta^\top \left\{ b(X)b(X)^\top \right\} \beta,$$

respectively, where  $\varphi'_Q(Z; \eta'_Q, \delta)$  is defined analogously to  $\varphi_Q$  except that  $\mu_a(X)$  and  $\phi_a(Z)$ are replaced by  $\mu'_a(X) = \mathbb{E}[Y^2 \mid X, A = a]$  and  $\phi'_a(Z) = \frac{\mathbb{I}(A=a)}{\pi_a(X)} \{Y^2 - \mu'_a(X)\} + \mu'_a(X)$ , with  $\eta'_Q = (\mu'_a, \pi_a)$ . Also, suppose that the fairness criterion of statistical parity (5) is required on the regression function with respect to a sensitive variable  $F \in \mathcal{X}$ . Then our approximating program is equivalent to the following quadratic optimization problem:

$$\begin{array}{ll} \underset{\beta \in \mathbb{R}^{k}}{\text{minimize}} & \frac{1}{2} \beta^{\top} \mathbb{P}_{n} \left\{ b(X) b(X)^{\top} \right\} \beta - \beta^{\top} \mathbb{P}_{n} \{ \varphi_{Q}(Z; \widehat{\eta}_{Q}, \delta) b(X) \} + \lambda \|\beta\|_{2}^{2} \\ \text{subject to} & \left| \mathbb{P}_{n} \left[ \left\{ \frac{(1-F)}{\mathbb{P}_{n}(1-F)} - \frac{F}{\mathbb{P}_{n}(F)} \right\} b(X)^{\top} \right] \beta \right| \leq \epsilon. \end{array}$$

One can add other fairness criteria on top of the statistical parity. When  $g_j$  depends on the counterfactual outcome  $Y^{Q(\delta)}$  as a non-smooth function (e.g., Example 1C), additional structural conditions, such as the margin condition (e.g., Kim and Zubizarreta 2023), or techniques like undersmoothing (e.g., Kennedy et al. 2017), may be required to ensure  $\sqrt{n}$ consistency and asymptotic normality.

## 4 Asymptotic Analysis

Here, we study the rates of convergence and the limiting distribution of our proposed estimator. This analysis reduces to studying the behavior of the optimal solution estimators for the target program defined in (2). We may express a generic form of (2) as

$$\underset{\beta \in \mathbb{R}^k}{\text{minimize }} \psi(\beta) \quad \text{subject to} \quad g_j(\beta) \le 0, \ j = 1, \dots, r, \tag{P}_g)$$

for some deterministic real-valued functions  $\psi$ ,  $g_j$ . The corresponding approximating program takes the form:

$$\underset{\beta \in \mathbb{R}^k}{\text{minimize }} \widehat{\psi}(\beta) \quad \text{subject to} \quad \widehat{g}_j(\beta) \le 0, \ j = 1, \dots, r, \qquad (\widehat{\mathsf{P}}_g)$$

where we let the random functions  $\widehat{\Psi}$ ,  $\widehat{g}_j$  denote estimators for  $\Psi$ ,  $g_j$ , respectively. In our setting, we have  $\Psi(\beta) = \mathbb{E} \{ \varphi_{\mathcal{R}}(Z; \eta_{\mathcal{R}}, \delta, \beta) \} + \lambda \rho(\beta)$  and  $\widehat{\Psi}(\beta) = \mathbb{P}_n \{ \varphi_{\mathcal{R}}(Z; \widehat{\eta}_{\mathcal{R}}, \delta, \beta) \} + \lambda \rho(\beta)$ .

As discussed in Section 2.3, the specific structure of our counterfactual coefficients introduces challenges that hinder the direct application of standard methods from the stochastic optimization literature. To address these challenges, we first introduce the classical results of Shapiro (1993), which apply to a broad class of stochastic programs, including the one considered in this study.

Let  $\beta^*$  and  $\hat{\beta}$  denote the optimal solutions of the true program  $(\mathsf{P}_g)$  and the approximating program  $(\hat{\mathsf{P}}_g)$ , respectively, i.e.,  $\beta^* \in \mathsf{s}^*(\mathsf{P}_g)$  and  $\hat{\beta} \in \mathsf{s}^*(\hat{\mathsf{P}}_g)$ . Assuming that the functions  $\psi, g_j$  are continuously differentiable, the set  $\{\beta : g_j(\beta) \leq 0, j = 1, \ldots, r\}$  is convex, and that a constraint qualification is satisfied, Shapiro (1993) analyzed the asymptotic behavior of  $\hat{\beta}$  around  $\beta^*$ , using the generalized equations approach. While this approach imposes less restrictive differentiability assumptions compared to the SAA and SA methods, it requires the consistency of our solution estimators to be ensured a priori, as stated below.

**Assumption 1.** Let  $\gamma^*$  and  $\hat{\gamma}$  denote the Lagrange multiplier vectors associated with the optimal solutions  $\beta^*$  and  $\hat{\beta}$ , respectively. We require  $\hat{\beta} \xrightarrow{p} \beta^*$  and  $\hat{\gamma} \xrightarrow{p} \gamma^*$ .

Although numerous results establish regularity conditions under which Assumption 1 holds by construction for conventional SAA and SA methods (e.g., Dupacová and Wets 1988; Shapiro 2000), verifying this condition in our context is not straightforward. Moreover, obtaining a closed-form expression for the limiting distribution of the optimal solution estimators necessitates computing the directional derivatives of the solution to the associated generalized equation.

In our work, we narrow our focus to specialized yet sufficiently general forms of our target problem where the above issues can be successfully addressed without introducing additional regularity conditions. Specifically, we first examine the smooth optimization problem with a fixed feasible set. We then consider a scenario in which the objective function exhibits separable stochastic components, and all constraints are linear.

### 4.1 Smooth functions with fixed feasible set.

We consider the case where both  $\mathsf{P}_g$  and  $\widehat{\mathsf{P}}_g$  share a common set of deterministic constraints  $\mathcal{C} = \{\beta : g_j(\beta) \leq 0, j = 1, \dots, r\}$ :

$$\underset{\beta \in \mathbb{R}^k}{\text{minimize }} \Psi(\beta) \quad \text{subject to} \quad g_j(\beta) \le 0, \ j = 1, \dots, r, \tag{P}_f)$$

$$\underset{\beta \in \mathbb{R}^k}{\text{minimize }} \widehat{\Psi}(\beta) \quad \text{subject to} \quad g_j(\beta) \le 0, \ j = 1, \dots, r.$$

This type of problem has been studied by Kim, Mishler and Zubizarreta (2022) in the context of classification tasks. Assume that  $\psi$ ,  $g_j$  are twice differentiable with respect to  $\beta$ , and that the Hessian matrix of  $\psi$  is positive definite at  $\beta^* \in s^*(\mathsf{P}_f)$ . If (10) holds, it follows that

$$\nabla_{\beta}\widehat{\psi}(\beta^*) - \nabla_{\beta}\psi(\beta^*) \xrightarrow{d} N\left(0, \nabla^2_{\beta}\psi(\beta^*) \operatorname{var}\left(\varphi_{\mathcal{R}}(Z; \eta_{\mathcal{R}}, \delta, \beta^*)\right) \nabla^2_{\beta}\psi(\beta^*)^{\top}\right).$$
(11)

Next, for any feasible point  $\bar{\beta} \in \mathcal{C}$  in  $\mathsf{P}_f$ , we let

$$L(\bar{\beta},\bar{\gamma}) = \psi(\bar{\beta}) + \sum_{j \in J_0(\bar{\beta})} \bar{\gamma}_j g_j(\bar{\beta})$$

denote the Lagrangian function with multipliers  $\bar{\gamma}_j \geq 0$ , and define the *active index set* by

$$J_0(\bar{\beta}) = \{ 1 \le j \le r : g_j(\bar{\beta}) = 0 \}.$$

We now introduce some standard regularity conditions for our analysis.

**Definition 4.1** (LICQ). Linear independence constraint qualification (LICQ) is satisfied at  $\bar{\beta} \in C$ , if the vectors  $\nabla_{\beta}g_{i}(\bar{\beta})$ ,  $j \in J_{0}(\bar{\beta})$  are linearly independent.

**Definition 4.2** (SC). Strict Complementarity (SC) is satisfied at  $\overline{\beta}$ , if the Karush-Kuhn-Tucker (KKT) condition

$$\nabla_{\beta} L(\bar{\beta}, \bar{\gamma}) = 0,$$

is satisfied such that  $\bar{\gamma}_j > 0, \forall j \in J_0(\bar{\beta}).$ 

LICQ is arguably one of the most widely used constraint qualifications that ensure the necessity of the first-order KKT conditions at optimal solution points. SC requires that if the

*j*-th inequality constraint is active, then the corresponding dual variable is strictly positive. Consequently, exactly one of  $\bar{\gamma}_j$  and  $g_j(\bar{\beta})$  is zero for each  $1 \leq j \leq r$ . SC is commonly employed in nonlinear programming, especially in the context of parametric optimization (e.g., Still 2018). We require that LICQ and SC hold at each optimal solution of  $\mathsf{P}_f$ .

Assumption 2. LICQ and SC hold at  $\beta^* \in s^*(\mathsf{P}_f)$  with the associated multipliers  $\gamma^*$ .

If we assume that the optimal solution  $\beta^*$  is unique and LICQ holds at  $\beta^*$ , then the corresponding multipliers  $\gamma^*$  are determined uniquely (Wachsmuth 2013). We also require the second-order growth condition as follows.

**Assumption 3.** There exists a neighborhood W of  $\beta^* \in s^*(\mathsf{P}_f)$ , and a constant  $\kappa > 0$  such that for all  $\beta \in W \cap \mathcal{C}$ ,

$$\Psi(\beta) \ge \Psi(\beta^*) + \kappa \|\beta - \beta^*\|_2^2.$$

Assumption 3 guarantees that each optimal solution is locally isolated, and is a standard condition in nonlinear programming (e.g., Shapiro et al. 2014; Still 2018). Assumption 3 can be ensured by various forms of second-order sufficient conditions. In our case, it holds, for example, if  $\varsigma^{\top} \nabla_{\beta}^2 g_j(\beta^*) \varsigma \geq 0$ , for any  $\varsigma \in \{\varsigma : \nabla_{\beta} g_j(\beta^*) = 0, j \in J_0(\beta^*)\}$ .

Now, given (11), the closed-form expression for the limiting distribution of  $\hat{\beta}$  is derived using the result of Kim, Mishler and Zubizarreta (2022, Lemma B.2), as formally stated below.

**Theorem 4.1.** Assume that  $P_f$  has a unique optimal solution  $\beta^*$  (i.e.,  $s^*(P_f)$  is a singleton), that (10) holds, and that Assumptions 2, 3 are satisfied. Then,

$$n^{1/2} \left( \widehat{\beta} - \beta^* \right) \xrightarrow{d} \begin{bmatrix} \nabla_{\beta}^2 \psi(\beta^*) + \sum_j \gamma_j^* \nabla_{\beta}^2 g_j(\beta^*) & \mathsf{B}(\beta^*) \\ \mathsf{B}^\top(\beta^*) & 0 \end{bmatrix}^{-1} \begin{bmatrix} \mathbf{1} \\ \mathbf{0} \end{bmatrix} N \left( 0, \Sigma_{\beta^*} \right),$$

where  $\mathsf{B} = \left[ \nabla_{\beta} g_j(\beta^*)^{\top}, \ j \in J_0(\beta^*) \right]$  and  $\Sigma_{\beta^*} = \nabla_{\beta}^2 \psi(\beta^*) var(\varphi_{\mathcal{R}}(Z; \eta_{\mathcal{R}}, \delta, \beta^*)) \nabla_{\beta}^2 \psi(\beta^*)^{\top}.$ 

Theorem 4.1 gives conditions under which  $\hat{\beta}$  is  $\sqrt{n}$ -consistent and asymptotically normal, without requiring the a-priori consistency in Assumption 1. However, restricting the analysis to fixed feasible sets limits the applicability to a broader range of regression problems (e.g., fairness).

# 4.2 Objective function with separable stochastic components and linear constraints.

Here, we analyze another specialization of  $\mathsf{P}_g$ , wherein a closed-form expression for the asymptotic distribution can be obtained without resorting to deterministic constraints or requiring the a-priori consistency condition. For a finite-dimensional set  $\mathbb{T}$  in Euclidean space, consider a statistical functional  $T : \mathbb{P} \to \mathbb{T}$  and a twice continuously differentiable real-valued function  $h : \mathbb{R}^k \times \mathbb{T} \to \mathbb{R}$ . Additionally, let  $C \in \mathbb{R}^{r \times k}$  and  $d \in \mathbb{R}^k$ , where the elements of C and d may depend on  $\mathbb{P}$ . Therefore,  $T(\mathbb{P}), C$ , and d are unknown and must be estimated. On the other hand, we assume that the function h is deterministic and known, and that all stochastic components are separable from h, meaning that any dependence on  $\mathbb{P}$  is confined to (a transformation of)  $T(\mathbb{P})$ , rather than being embedded within the functional form of h.  $T(\mathbb{P})$  can be viewed as a set of (identified) counterfactual components of interest. Our objective is to estimate the optimal solution of the following stochastic program:

$$\underset{\beta \in \mathbb{R}^k}{\text{minimize } h(\beta, T) \quad \text{subject to} \quad C\beta \le d, \qquad (\mathsf{P}_{sl})$$

where we write  $T(\mathbb{P}) \equiv T$ . Since the true program  $(\mathsf{P}_{sl})$  is not directly solvable, we compute an optimal solution of the following approximating program:

$$\underset{\beta \in \mathbb{R}^k}{\text{minimize } h(\beta, \widehat{T}) \quad \text{subject to} \quad \widehat{C}\beta \leq \widehat{d}.$$
 ( $\widehat{\mathsf{P}}_{sl}$ )

Thus, we allow for varying feasible sets, i.e., the feasible sets of programs  $\mathsf{P}_{sl}$  and  $\widehat{\mathsf{P}}_{sl}$  are not required to coincide as a fixed deterministic set, with each set defined by linear constraints. Note that all the cases in Examples 1A - 1C can be formulated as linear constraints. A representative example for h is the case where  $L_2$  loss is used with the aggregated predictor defined in (3), as in Example 2C.

We will employ the exact counterparts of the regularity assumptions utilized in the previous subsection. A sufficient second-order condition for Assumption 3 can be specified as

$$\varsigma^{\top} \nabla^2_{\beta} h(\beta^*, T) \varsigma > 0, \quad \forall \varsigma \in \{\varsigma \in \mathbb{R}^k \mid \nabla_{\beta} h(\beta^*, T)^{\top} \varsigma \le 0, C_j^{\top} \varsigma \le 0, j \in J_0(\beta^*)\} \setminus \{0\},$$

for  $\beta^* \in s^*(\mathsf{P}_{sl})$  and  $J_0(\beta^*) = \{1 \leq j \leq r : C_j^\top \beta^* - d_j = 0\}$ , which, in particular, holds if  $\nabla_{\beta}^2 h(\beta^*, T)$  is positive definite. We also require the following mild consistency condition, with no requirement on rates of convergence.

Assumption 4.  $\max \left\{ \|\hat{T} - T\|_2, \|\hat{C} - C\|_F, \|\hat{d} - d\|_2 \right\} = o_{\mathbb{P}}(1).$ 

The consistency properties of  $\hat{\beta} \in s^*(\hat{\mathsf{P}}_{sl})$  can be established under relatively weak assump-

tions, as formalized in the following theorem.

**Theorem 4.2.** Under Assumptions 3 and 4, it follows that for  $\hat{\beta} \in s^*(\hat{P}_{sl})$ ,

$$dist(\widehat{\beta}, \mathsf{s}^*(\mathsf{P}_{sl})) = O_{\mathbb{P}}\left(\max\left\{\|\widehat{T} - T\|_2, \|\widehat{C} - C\|_F, \|\widehat{d} - d\|_2\right\}\right).$$

The proof of Theorem 4.2 is presented in Appendix A.1. Theorem 4.2 suggests that the convergence rates for estimating the optimal solutions of  $\mathsf{P}_{sl}$  are essentially determined by the rates at which each stochastic component is estimated. The result in Theorem 4.2 follows from the local Lipschitz stability property of optimal solutions in general nonlinear parametric optimization.

Characterizing the asymptotic distribution of  $\hat{\beta}$  requires stronger assumptions than those needed for establishing consistency. First, we impose the following rate condition, which is stronger than Assumption 4.

# Assumption 5. $\max\left\{\|\widehat{T} - T\|_2, \|\widehat{C} - C\|_F, \|\widehat{d} - d\|_2\right\} = O_{\mathbb{P}}(n^{-1/2}).$

Next, as before, we require Assumption 2 to hold for our target program  $\mathsf{P}_{sl}$ . For  $\beta^* \in \mathsf{s}^*(\mathsf{P}_{sl})$ , LICQ holds if the vectors  $\{C_j^\top : j \in J_0(\beta^*)\}$  are linearly independent, and SC holds if the KKT conditions

$$\nabla_{\beta}h(\beta^*,T) + \sum_{j \in J_0(\beta^*)} \gamma_j^* C_j^\top \beta^* = 0, \quad \operatorname{diag}(\gamma^*)(C\beta^* - d) = 0$$

are satisfied such that  $\gamma_j^* > 0, \forall j \in J_0(\beta^*)$ .

For our analysis, we utilize a generalized delta method for directionally differentiable mappings (e.g., Shapiro 1993, 2000). To this end, we require the following technical condition.

Assumption 6. For  $|J_0(\beta^*)| \times k$  matrix  $C_{ac} = [C_j, j \in J_0(\beta^*)]$  and some random vector  $\Upsilon_{\beta^*} \in \mathbb{R}^{k+|J_0(\beta^*)|}$ ,

$$n^{1/2} \begin{bmatrix} \nabla_{\beta} h(\beta^*, \widehat{T}) - \nabla_{\beta} h(\beta^*, T) + \sum_{j} \gamma_{j \in J_0(\beta^*)}^* \left\{ \widehat{C}_j - C_j \right\} \\ -(\widehat{\mathsf{C}}_{ac} - \mathsf{C}_{ac})\beta^* \end{bmatrix} \xrightarrow{d} \Upsilon_{\beta^*}$$

Assumption 6 states that, on the active set, the KKT multipliers of the approximating program  $\hat{\mathsf{P}}_{sl}$  jointly converge in distribution to those of the true program  $\mathsf{P}_{sl}$  at  $\sqrt{n}$  rates. Similar conditions are used in the analysis of parametric programs (e.g., Shapiro 1990). Since inference on optimal solution estimators is typically conducted using bootstrap methods, the

case where  $\Upsilon_{\beta^*}$  follows a multivariate normal distribution is particularly important. The absence of this property undermines the consistency of the bootstrap for solution estimators (Fang and Santos 2019). It is straightforward to see that if (10) holds, then Assumptions 5 and 6 are satisfied, with  $\Upsilon_{\beta^*}$  following a multivariate normal. In the next theorem, we provide a closed-form expression for the asymptotic distribution of our proposed estimator, characterized as the optimal solution to  $\widehat{\mathsf{P}}_{sl}$ .

**Theorem 4.3.** Assume that  $\mathsf{P}_{sl}$  has a unique optimal solution  $\beta^*$ , and that Assumptions 2, 3, 5, 6 are satisfied. Then, for  $\widehat{\beta} \in \mathsf{s}^*(\widehat{\mathsf{P}}_{sl})$ , we have that

$$n^{\frac{1}{2}} \left( \widehat{\beta} - \beta^* \right) = \begin{bmatrix} \nabla_{\beta}^2 h(\beta^*, T) & \mathsf{C}_{ac}^{\mathsf{T}} \\ \mathsf{C}_{ac} & 0 \end{bmatrix}^{-1} \begin{bmatrix} \mathbf{1} \\ \operatorname{diag}(\gamma_{ac}^*) \mathbf{1} \end{bmatrix}^{\mathsf{T}} \Upsilon_{\beta^*} + o_{\mathbb{P}}(1), \tag{12}$$

where  $C_{ac}, \Upsilon_{\beta^*}$  are defined according to Assumption 6, and  $\gamma_{ac}^* = [\gamma_j^*, j \in J_0(\beta^*)]$ .

We provide the proof of Theorem 4.3 in Appendix A.2. To derive (12), we calculate the directional derivative of the optimal solutions in the relevant parametric program using an appropriate version of the implicit function theorem (Dontchev and Rockafellar 2009). We apply Theorem 4.3 to Examples 2B and 2C, which leads to the following corollaries.

**Corollary 4.4.** Consider the mean squared logarithmic loss with the  $L_2$  penalty term as in Example 2B. Assume that f is twice continuously differentiable with respect to  $\beta$ ,  $||1/\hat{\pi}_a||_{\infty} < \infty$ ,  $||\hat{\mu}'_a - \mu'_a||_{2,\mathbb{P}} + ||\hat{\mu}''_a - \mu''_a||_{2,\mathbb{P}} + ||\hat{\pi}_a - \pi_a||_{2,\mathbb{P}} = o_{\mathbb{P}}(1)$ , and  $||\hat{\pi}_a - \pi_a||_{2,\mathbb{P}}(||\hat{\mu}'_a - \mu'_a||_{2,\mathbb{P}} + ||\hat{\mu}''_a - \mu''_a||_{2,\mathbb{P}}) = o_{\mathbb{P}}(n^{-\frac{1}{2}})$ . Then,  $\hat{\beta}$  is  $\sqrt{n}$ -consistent and asymptotically normal for  $\beta^*$ .

**Corollary 4.5.** Consider the  $L_2$  loss with the  $L_2$  penalty term, the aggregated predictors (3), and the statistical parity as in Example 2C. Assume that  $\|1/\hat{\pi}_a\|_{\infty} < \infty$ ,  $\|\hat{\mu}_a - \mu_a\|_{2,\mathbb{P}} + \|\hat{\pi}_a - \pi_a\|_{2,\mathbb{P}} = o_{\mathbb{P}}(1)$ , and  $\|\hat{\pi}_a - \pi_a\|_{2,\mathbb{P}} \|\hat{\mu}_a - \mu_a\|_{2,\mathbb{P}} = o_{\mathbb{P}}(n^{-\frac{1}{2}})$ . Then,  $\hat{\beta}$  is  $\sqrt{n}$ -consistent and asymptotically normal for  $\beta^*$ .

Corollary 4.5 can be extended to general fairness criteria (4), provided that the corresponding coefficients are  $\sqrt{n}$ -estimable.

## 5 Experiments

We conduct a simulation study to assess the performance of the proposed estimators, emphasizing the validation of the theoretical results established in Section 4. We focus on the canonical setting described in Example 2C, using  $L_2$  loss and statistical parity as the fairness criterion, with a threshold of  $\epsilon = 0.1$ . The data-generating process is specified as follows:  $F \sim$  Bernoulli(0.5),  $X_1 \sim$  Uniform[0, 10],  $X_2 \sim N(4F - 2, 2)$ ,  $\pi(X) = \exp(2.5 - 0.3X_1 - FX_2)$ ,



Figure 2: RMSE versus nuisance convergence rates for (a)  $\delta = 0.1$  and (b)  $\delta = 0.01$ . The proposed estimators consistently attain convergence rates faster than those of the nuisance components.

and  $\mu_A(X) = 0.5\sqrt{X_1} + 2X_2 - 5A$  where  $X = (X_1, X_2)$ . We construct the nuisance estimators as  $\hat{\pi}(X) = \text{expit} \{ \text{logit}(\pi(X)) + \epsilon_{\pi} \}$  and  $\hat{\mu}_a(X) = \mu_a(X) + \epsilon_{\mu}$ , where  $\epsilon_{\pi}, \epsilon_{\mu} \sim N(n^{-r}, n^{-2r})$ . These choices guarantee that  $\|\hat{\pi}_a - \pi_a\|_{2,\mathbb{P}} = O(n^{-r})$  and  $\|\hat{\mu}_a - \mu_a\|_{2,\mathbb{P}} = O(n^{-r})$ , where we vary  $r \in (0, 0.5)$  for different scenarios. This simulation setup allows us to study how the proposed semiparametric estimators perform under different convergence rates of the nuisance components.

For each pair of  $(n, \delta)$ , where  $n \in \{500, 1000, 5000\}$ ,  $\delta \in \{0.1, 0.01\}$ , we generate data and compute nuisance estimates as described above, with  $r \in \{0.05 + 0.025k : k = 0, ..., 18\}$ . We then compute the coefficients of the proposed estimator  $\hat{\beta}$  by solving the corresponding constrained quadratic program. Each scenario was replicated 500 times, and the root-meansquare error (RMSE) is computed with respect to the true  $\beta^*$ . The results are shown in Figure 2. The proposed estimators achieve convergence rates faster than those of the nuisance components and rapidly approach the parametric rate of  $O(n^{-1/2})$  particularly for larger sample sizes, thereby supporting the theoretical results in the previous section.

## 6 Discussion

We have developed a general framework for counterfactual regression under incremental interventions. We believe this work offers new insights with potential relevance beyond causal inference, particularly in related areas of the data science community such as domain adaptation, out-of-distribution generalization, and transfer learning. There are several promising directions for future research. One natural extension is to accommodate continuous or timevarying treatments, which frequently arise in practice. Another involves generalizing the framework to allow for more complex non-linear objective functions and constraints, though such extensions may require additional regularity conditions. Finally, applying the proposed methods in the context of optimal treatment regimes or conditional effect estimation is also of interest, as it may provide new guidance on how treatment policies can be more effectively tailored under varying circumstances.

## References

- Barocas, S., Hardt, M. and Narayanan, A. (2023), Fairness and Machine Learning: Limitations and Opportunities, MIT Press. URL: http://www.fairmlbook.org
- Bertsekas, D. (2015), Convex optimization algorithms, Athena Scientific.
- Bickel, P. J., Klaassen, C. A., Bickel, P. J., Ritov, Y., Klaassen, J., Wellner, J. A. and Ritov, Y. (1993), Efficient and adaptive estimation for semiparametric models, Vol. 4, Springer.
- Boyd, S., Parikh, N., Chu, E., Peleato, B., Eckstein, J. et al. (2011), 'Distributed optimization and statistical learning via the alternating direction method of multipliers', *Foundations and Trends® in Machine learning* **3**(1), 1–122.
- Chen, H., Zhang, M., Han, L. and Lim, A. (2021), 'Hierarchical marketing mix models with sign constraints', *Journal of Applied Statistics* **48**(13-15), 2944–2960.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., Hansen, C. and Newey, W. (2017), 'Double/debiased/neyman machine learning of treatment effects', American Economic Review 107(5), 261–65.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., Hansen, C., Newey, W. and Robins, J. (2018), 'Double/debiased machine learning for treatment and structural parameters', *The Econometrics Journal* 21(1), C1–C68.
- Corbett-Davies, S., Pierson, E., Feller, A., Goel, S. and Huq, A. (2017), Algorithmic decision making and the cost of fairness, *in* 'Proceedings of the 23rd acm sigkdd international conference on knowledge discovery and data mining', pp. 797–806.
- Coston, A., Kennedy, E. and Chouldechova, A. (2020), Counterfactual predictions under runtime confounding, *in* 'Advances in neural information processing systems', Vol. 33, pp. 4150–4162.
- Coston, A., Mishler, A., Kennedy, E. H. and Chouldechova, A. (2020), Counterfactual risk assessments, evaluation, and fairness, *in* 'Proceedings of the 2020 Conference on Fairness, Accountability, and Transparency', pp. 582–593.
- Dickerman, B. A., Dahabreh, I. J., Cantos, K. V., Logan, R. W., Lodi, S., Rentsch, C. T., Justice, A. C. and Hernán, M. A. (2022), 'Predicting counterfactual risks under hypothetical treatment strategies: an application to hiv', *European journal of epidemiology* 37(4), 367–376.

- Dickerman, B. A. and Hernán, M. A. (2020), 'Counterfactual prediction is not only for causal inference', *European Journal of Epidemiology* **35**(7), 615–617.
- Dontchev, A. L. and Rockafellar, R. T. (2009), *Implicit functions and solution mappings*, Vol. 543, Springer.
- Dupacová, J. and Wets, R. (1988), 'Asymptotic behavior of statistical estimators and of optimal solutions of stochastic optimization problems', *The annals of statistics* 16(4), 1517– 1549.
- Faigle, U., Kern, W. and Still, G. (2013), Algorithmic principles of mathematical programming, Vol. 24, Springer Science & Business Media.
- Fang, Z. and Santos, A. (2019), 'Inference on directionally differentiable functions', The Review of Economic Studies 86(1), 377–412.
- Gaines, B. R., Kim, J. and Zhou, H. (2018), 'Algorithms for fitting the constrained lasso', Journal of Computational and Graphical Statistics 27(4), 861–871.
- Hardt, M., Price, E. and Srebro, N. (2016), 'Equality of opportunity in supervised learning', Advances in neural information processing systems 29.
- Hassanpour, N. and Greiner, R. (2019*a*), Counterfactual regression with importance sampling weights., *in* 'IJCAI', Macao, pp. 5880–5887.
- Hassanpour, N. and Greiner, R. (2019b), Learning disentangled representations for counterfactual regression, *in* 'International Conference on Learning Representations'.
- Hernán, M. A., Hsu, J. and Healy, B. (2019), 'A second chance to get causal inference right: a classification of data science tasks', *Chance* **32**(1), 42–49.
- Hodson, T. O., Over, T. M. and Foks, S. S. (2021), 'Mean squared error, deconstructed', Journal of Advances in Modeling Earth Systems 13(12), e2021MS002681.
- Jain, P., Kar, P. et al. (2017), 'Non-convex optimization for machine learning', *Foundations* and *Trends® in Machine Learning* **10**(3-4), 142–363.
- James, G. M., Paulson, C. and Rusmevichientong, P. (2013), 'Penalized and constrained regression', Unpublished manuscript, http://www-bcf. usc. edu/gareth/research/Research. html.
- James, G. M., Paulson, C. and Rusmevichientong, P. (2019), 'Penalized and constrained optimization: an application to high-dimensional website advertising', *Journal of the American Statistical Association*.

- Johansson, F., Shalit, U. and Sontag, D. (2016), Learning representations for counterfactual inference, *in* 'International conference on machine learning', PMLR, pp. 3020–3029.
- Kennedy, E. H. (2016), Semiparametric theory and empirical processes in causal inference, in 'Statistical causal inferences and their applications in public health research', Springer, pp. 141–167.
- Kennedy, E. H. (2019), 'Nonparametric causal effects based on incremental propensity score interventions', *Journal of the American Statistical Association* **114**(526), 645–656.
- Kennedy, E. H. (2020), 'Optimal doubly robust estimation of heterogeneous causal effects', arXiv preprint arXiv:2004.14497.
- Kennedy, E. H. (2022), 'Semiparametric doubly robust targeted double machine learning: a review', *arXiv preprint arXiv:2203.06469*.
- Kennedy, E. H., Balakrishnan, S. and Wasserman, L. (2021), 'Semiparametric counterfactual density estimation', arXiv preprint arXiv:2102.12034.
- Kennedy, E. H., Ma, Z., McHugh, M. D. and Small, D. S. (2017), 'Non-parametric methods for doubly robust estimation of continuous treatment effects', *Journal of the Royal Statistical Society Series B: Statistical Methodology* 79(4), 1229–1245.
- Kim, K., Kennedy, E. H. and Naimi, A. I. (2021), 'Incremental intervention effects in studies with dropout and many timepoints#', Journal of Causal Inference 9(1), 302–344.
- Kim, K., Kennedy, E. and Zubizarreta, J. (2022), 'Doubly robust counterfactual classification', Advances in Neural Information Processing Systems 35, 34831–34845.
- Kim, K., Mishler, A. and Zubizarreta, J. R. (2022), 'Counterfactual mean-variance optimization', arXiv preprint arXiv:2209.09538.
- Kim, K. and Zubizarreta, J. R. (2023), Fair and robust estimation of heterogeneous treatment effects for policy learning, *in* 'International Conference on Machine Learning', PMLR, pp. 16997–17014.
- Kleinberg, J., Mullainathan, S. and Raghavan, M. (2016), 'Inherent trade-offs in the fair determination of risk scores', arXiv preprint arXiv:1609.05807.
- Lancet Editorial, T. L. R. H.-W. (2024), 'Junior doctor strikes in south korea: more doctors are needed?', *The Lancet Regional Health: Western Pacific* 44, 101056.
- Li, J., Zhao, L., Tian, L., Cai, T., Claggett, B., Callegaro, A., Dizier, B., Spiessens, B., Ulloa-Montoya, F. and Wei, L.-J. (2016), 'A predictive enrichment procedure to identify

potential responders to a new therapy for randomized, comparative controlled clinical studies', *Biometrics* 72(3), 877–887.

- Liew, C. K. (1976), 'Inequality constrained least-squares estimation', Journal of the American Statistical Association 71(355), 746–751.
- Lin, L., Sperrin, M., Jenkins, D. A., Martin, G. P. and Peek, N. (2021), 'A scoping review of causal methods enabling predictions under hypothetical interventions', *Diagnostic and* prognostic research 5(1), 1–16.
- Lu, J., Shi, P. and Li, H. (2019), 'Generalized linear models with linear constraints for microbiome compositional data', *Biometrics* **75**(1), 235–244.
- McClean, A., Branson, Z. and Kennedy, E. H. (2024), 'Nonparametric estimation of conditional incremental effects', *Journal of Causal Inference* **12**(1), 20230024.
- Mishler, A. and Kennedy, E. (2021), 'Fade: Fair double ensemble learning for observable and counterfactual outcomes', *arXiv preprint arXiv:2109.00173*.
- Nemirovski, A., Juditsky, A., Lan, G. and Shapiro, A. (2009), 'Robust stochastic approximation approach to stochastic programming', *SIAM Journal on optimization* **19**(4), 1574– 1609.
- Newey, W. K. and Robins, J. R. (2018), 'Cross-fitting and fast remainder rates for semiparametric estimation', *arXiv preprint arXiv:1801.09138*.
- Nguyen, T.-L., Collins, G. S., Landais, P. and Le Manach, Y. (2020), 'Counterfactual clinical prediction models could help to infer individualized treatment effects in randomized controlled trials—an illustration with the international stroke trial', *Journal of clinical epidemiology* **125**, 47–56.
- Park, H. (2024), 'Junior doctors' strike in south korea: systemic barriers undermine medical practice', bmj 385.
- Polley, E. C. and Van der Laan, M. J. (2010), 'Super learner in prediction'.
- Schindl, K., Shen, S. and Kennedy, E. H. (2024), 'Incremental effects for continuous exposures', arXiv preprint arXiv:2409.11967.
- Schulam, P. and Saria, S. (2017), 'Reliable decision support using counterfactual models', Advances in Neural Information Processing Systems 30, 1697–1708.
- Schwendinger, B., Schwendinger, F. and Vana, L. (2024), 'Holistic generalized linear models', Journal of Statistical Software 108, 1–49.

- Shalit, U., Johansson, F. D. and Sontag, D. (2017), Estimating individual treatment effect: generalization bounds and algorithms, *in* 'International conference on machine learning', PMLR, pp. 3076–3085.
- Shapiro, A. (1990), 'On differential stability in stochastic programming', Mathematical Programming 47, 107–116.
- Shapiro, A. (1993), 'Asymptotic behavior of optimal solutions in stochastic programming', Mathematics of Operations Research 18(4), 829–845.
- Shapiro, A. (2000), Statistical inference of stochastic optimization problems, *in* 'Probabilistic constrained optimization', Springer, pp. 282–307.
- Shapiro, A., Dentcheva, D. and Ruszczyński, A. (2014), Lectures on stochastic programming: modeling and theory, SIAM.
- Sperrin, M., Martin, G. P., Pate, A., Van Staa, T., Peek, N. and Buchan, I. (2018), 'Using marginal structural models to adjust for treatment drop-in when developing clinical prediction models', *Statistics in medicine* 37(28), 4142–4154.
- Still, G. (2018), 'Lectures on parametric optimization: An introduction', *Optimization On*line.
- Tsiatis, A. A. (2006), Semiparametric theory and missing data, Vol. 4, Springer.
- Tsybakov, A. B. (2003), Optimal rates of aggregation, in 'Learning Theory and Kernel Machines: 16th Annual Conference on Learning Theory and 7th Kernel Workshop, COLT/Kernel 2003, Washington, DC, USA, August 24-27, 2003. Proceedings', Springer, pp. 303–313.
- van der Vaart, A. W. (2002), Semiparametric statistics, *in* 'Lectures on probability theory and statistics (Saint-Flour, 1999)', Springer, pp. 331–457.
- van Geloven, N., Swanson, S. A., Ramspek, C. L., Luijken, K., van Diepen, M., Morris, T. P., Groenwold, R. H., van Houwelingen, H. C., Putter, H. and le Cessie, S. (2020), 'Prediction meets causal inference: the role of treatment in clinical prediction models', *European journal of epidemiology* 35(7), 619–630.
- Wachsmuth, G. (2013), 'On licq and the uniqueness of lagrange multipliers', *Operations Research Letters* **41**(1), 78–80.
- Wang, Q., Ma, Y., Zhao, K. and Tian, Y. (2020), 'A comprehensive survey of loss functions in machine learning', Annals of Data Science pp. 1–26.

Zeng, J., Ustun, B. and Rudin, C. (2017), 'Interpretable classification models for recidivism prediction', Journal of the Royal Statistical Society Series A: Statistics in Society 180(3), 689–722.

### APPENDIX

## A Proofs

**Extra notation.** First, we introduce some extra notation used throughout in the proofs. We let  $\langle M_1, M_2 \rangle \coloneqq tr\left(M_1^\top M_2\right)/k$  for  $k \times k$  matrices  $M_1, M_2$  (so  $||M_1||_F = \sqrt{\langle M_1, M_1 \rangle}$ ). We let  $\mathbb{B}_{\delta}(\bar{z})$  denote the open ball with radius  $\delta > 0$  around the point  $\bar{z}$  with  $|| \cdot ||_2$  (unless otherwise mentioned), i.e.,  $\mathbb{B}_{\delta}(\bar{z}) = \{z \mid ||z - \bar{z}||_2 < \delta\}$ . We use  $C^r(\mathbb{S})$  to denote a set of functions that are r times continuously differentiable on  $\mathbb{S}$ . Further, we let  $o_{x \to \alpha}(1)$  denote some function of x that converges to 0 when  $x \to \alpha$ , i.e., a function  $h_{\alpha}(x)$  such that  $\lim_{x \to \alpha} ||h_{\alpha}(x)|| = 0$ . This is to distinguish from typical little-o (o) asymptotic notation with respect to n in our proofs. Finally, for  $\theta \in \mathbb{R}^p$  and any vector-valued function  $H : \mathbb{R}^p \to \mathbb{R}^q$  whose first-order partial derivatives exist, we denote its Jacobian matrix with respect to  $\theta$  by  $J_{\theta}H \in \mathbb{R}^{q \times p}$ .

### A.1 Proof of Theorem 4.2

First, we discuss the notion of Lipschitz stability of local minimizers in general nonlinear parametric optimization. Let  $\Xi \subset \mathbb{R}^q$  be some finite-dimensional open parameter set. For  $\theta \in \Xi$ , consider a parametric program

$$\begin{array}{ll} \underset{x}{\operatorname{minimize}} & f(x,\theta) \\ \text{subject to} & x \in \mathcal{S}_q(\theta) = \{ x \mid g_j(x,\theta) \le 0, j \in J \}, \end{array}$$

where  $f, g_j \in C^2(\mathbb{R}^k \times \Xi)$ . Let  $x^*(\theta) \in s^*(\mathsf{P}(\theta))$ , a local minimizer of  $\mathsf{P}(\theta)$  with the corresponding parameter  $\theta$ . Next, we define the Lipschitz stability property of local minimizers.

**Definition A.1** (Lipschitz stability of local minimizer).  $x_0 \equiv x^*(\theta_0)$  is called Lipschitz stable if there exist some constants  $L, \epsilon > 0$  and a local minimizer  $x(\theta)$  of  $\mathsf{P}(\theta)$  such that

$$||x(\theta) - x_0||_2 \le L ||\theta_0 - \theta||_2 \quad \forall \theta \in \mathbb{B}_{\delta}(\theta_0).$$

See, for example, Still (2018), Chapter 6 for more details for Lipschitz and other types of stability result of local minimizers in smooth  $(C^2)$  nonlinear parametric optimization.

Assume that the parameter  $\theta$  unknown and is to be estimated from data with an estimator  $\hat{\theta}$ . The next lemma show that if the Lipschitz stability result holds for each local minimizer of  $\mathsf{P}(\theta)$  then the estimation error of the solutions is bounded by that of the parameters.

**Lemma A.1.** Suppose a local minimizer  $x^* \equiv x^*(\theta)$  of  $\mathsf{P}(\theta)$  is Lipschitz stable. Provided

that  $\hat{\theta}$  converges in probability to  $\theta$ , we have

$$dist(x^*, \mathsf{s}^*(\mathsf{P}(\widehat{\theta}))) = O_{\mathbb{P}}\left(\left\|\widehat{\theta} - \theta\right\|_2\right)$$

*Proof.* By definition of Lipschitz stability, there exist constants  $L, \epsilon > 0$  and a local solution  $x(\hat{\theta})$  of  $\mathsf{P}(\hat{\theta})$  such that

$$||x(\widehat{\theta}) - x^*||_2 \le L ||\theta - \widehat{\theta}||_2, \quad \forall \widehat{\theta} \in \mathbb{B}_{\delta}(\theta).$$

Given that  $\hat{\theta} \xrightarrow{p} \theta$ , we have

$$\begin{aligned} \|\widehat{x} - x^*\|_2 &= \|\widehat{x} - x^*\|_2 \mathbb{1}\left\{\widehat{\theta} \in \mathbb{B}_{\delta}(\theta)\right\} + \|\widehat{x} - x^*\|_2 \mathbb{1}\left\{\widehat{\theta} \notin \mathbb{B}_{\delta}(\theta)\right\} \\ &= \|\widehat{x} - x^*\|_2 + \left(\mathbb{1}\left\{\widehat{\theta} \in \mathbb{B}_{\delta}(\theta)\right\} - 1\right)\|\widehat{x} - x^*\|_2 + \|\widehat{x} - x^*\|_2 \mathbb{1}\left\{\widehat{\theta} \notin \mathbb{B}_{\delta}(\theta)\right\} \\ &= O\left(\|\theta - \widehat{\theta}\|_2\right) + o_{\mathbb{P}}\left(\|\widehat{x} - x^*\|_2\right),\end{aligned}$$

where the last equality follows by  $\mathbb{1}\left\{\widehat{\theta} \notin \mathbb{B}_{\delta}(\theta)\right\} = o_{\mathbb{P}}(1)$  and  $\mathbb{1}\left\{\widehat{\theta} \in \mathbb{B}_{\delta}(\theta)\right\} - 1 = o_{\mathbb{P}}(1)$ . Hence we have  $\|\widehat{x} - x^*\|_{2}$ 

$$\frac{\|\widehat{x} - x^*\|_2}{\|\widehat{\theta} - \theta\|_2} = O_{\mathbb{P}}(1)$$

and the result follows by the fact that  $dist(x^*, s^*(\mathsf{P}(\widehat{\theta}))) \leq \|\widehat{x} - x^*\|_2$ .

The proof of Theorem 4.2 follows directly from the following argument.

*Proof.*  $\mathsf{P}_{sl}$  can be viewed as a special case of the parametric program  $\mathsf{P}(\theta)$ , where the parameter  $\theta$  includes all the varying components (T, C, d). Owing to the linearity of the constraints, the strong second-order condition (Still 2018, Definition 6.1) is satisfied for each optimal solution under Assumption 3. Hence, by Theorem 6.4 of Still (2018),  $\beta^* \in \mathsf{s}^*(\mathsf{P}_{sl})$  is Lipschitz stable. The remainder of the proof follows directly from Lemma A.1.

### A.2 Proof of Theorem 4.3

*Proof.* For notational simplicity, we write  $h(\beta, T) \equiv h(\beta)$ ,  $h(\beta, \hat{T}) \equiv \hat{h}(\beta)$ . Let  $\gamma^*$  and  $\hat{\gamma}$  be the Lagrange multipliers associated with the optimal solutions of  $\mathsf{P}_{sl}$  and  $\hat{\mathsf{P}}_{sl}$ , respectively. First, we aim to show that

$$\|\widehat{\gamma} - \gamma\|_{2} = O_{\mathbb{P}}\left(\|\widehat{\beta} - \beta^{*}\|_{2} + n^{-1/2}\right).$$
(13)

Since the feasible sets of  $\mathsf{P}_{sl}$  and  $\widehat{\mathsf{P}}_{sl}$  are defined solely by linear inequalities, the KKT

optimality conditions hold for both problems (Faigle et al. 2013, Theorem 5.4). Then, by the complementary slackness, for each  $j \in \{1, ..., r\}$ , we have  $\gamma_j^*(C_j\beta^* - d_j) = 0$  and  $\hat{\gamma}_j(C_j\hat{\beta} - \hat{d}_j) = 0$ . Consequently,

$$\gamma_j^*(C_j\beta^* - d_j) - \hat{\gamma}_j(\hat{C}_j\hat{\beta} - \hat{d}_j) = (\gamma_j^* - \hat{\gamma}_j)(C_j\beta^* - d_j) + \hat{\gamma}_j\left\{(C_j\beta^* - d_j) - (\hat{C}_j\hat{\beta} - \hat{d}_j)\right\}$$
$$= (\gamma_j^* - \hat{\gamma}_j)(C_j\beta^* - d_j) + \hat{\gamma}_jC_j(\beta^* - \hat{\beta}) + \hat{\gamma}_j\left\{(C_j - \hat{C}_j)\hat{\beta} + \hat{d}_j - d_j\right\}$$
$$= 0.$$

Recall that  $J_0(\beta^*)$  is the active index set for  $\mathsf{P}_{sl}$ . Then, from the above

$$j \notin J_0(\beta^*) \Rightarrow \hat{\gamma}_j - \gamma_j^* = O\left(\|\hat{\beta} - \beta^*\|_2 + \|\hat{C}_j - C_j\|_2 + |\hat{d}_j - d_j|\right).$$
(14)

On the other hand, by combining the stationarity conditions (or dual conditions) from both  $\mathsf{P}_{sl}$  and  $\widehat{\mathsf{P}}_{sl}$ , and subsequently adding and subtracting terms, it follows that:

$$\begin{split} 0 &= \nabla_{\beta} \widehat{h}(\widehat{\beta}) - \nabla_{\beta} h(\beta^{*}) + \sum_{j \in J_{0}(\beta^{*})} \left\{ \widehat{\gamma}_{j} \widehat{C}_{j}^{\top} - \gamma_{j}^{*} C_{j}^{\top} \right\} \\ &= \nabla_{\beta} \widehat{h}(\widehat{\beta}) - \nabla_{\beta} \widehat{h}(\beta^{*}) + \nabla_{\beta} \widehat{h}(\beta^{*}) - \nabla_{\beta} h(\beta^{*}) + \sum_{j \in J_{0}(\beta^{*})} \left\{ (\widehat{\gamma}_{j} - \gamma_{j}^{*}) C_{j}^{\top} + \widehat{\gamma}_{j} (\widehat{C}_{j}^{\top} - C_{j}^{\top}) \right\} \\ &= (\widehat{\beta} - \beta^{*})^{\top} \nabla_{\beta}^{2} \widehat{h}(\beta^{*}) + (\widehat{\beta} - \beta^{*}) o_{\widehat{\beta} \to \beta^{*}}(1) + \nabla_{\beta} \widehat{h}(\beta^{*}) - \nabla_{\beta} h(\beta^{*}) \\ &+ \sum_{j \in J_{0}(\beta^{*})} \left\{ (\widehat{\gamma}_{j} - \gamma_{j}^{*}) C_{j}^{\top} + \widehat{\gamma}_{j} (\widehat{C}_{j}^{\top} - C_{j}^{\top}) \right\}, \end{split}$$

where the last equality follows by Taylor's theorem. Since we have assumed non-zero rows in C, by the Cauchy–Schwarz and triangle inequalities we obtain

$$j \in J_0(\beta^*) \Rightarrow |\hat{\gamma}_j - \gamma_j^*| = O\left(\|\hat{\beta} - \beta^*\|_2 + \|\nabla_\beta \hat{h}(\beta^*) - \nabla_\beta h(\beta^*)\|_2 + \|\hat{C}_j - C_j\|_2\right).$$
(15)

Combining the results from 14 and 15, we obtain:

$$\begin{aligned} \|\widehat{\gamma} - \gamma\|_{2} &= O\left(\|\widehat{\beta} - \beta^{*}\|_{2} + \|\nabla_{\beta}\widehat{h}(\beta^{*}) - \nabla_{\beta}h(\beta^{*})\|_{2} + \sum_{j=1}^{r} \|\widehat{C}_{j} - C_{j}\|_{2} + \sum_{j=1}^{r} |\widehat{d}_{j} - d_{j}|\right) \\ &= O\left(\|\widehat{\beta} - \beta^{*}\|_{2} + \|\nabla_{\beta}\widehat{h}(\beta^{*}) - \nabla_{\beta}h(\beta^{*})\|_{2} + \|\widehat{C} - C\|_{F} + \|\widehat{d} - d\|_{2}\right),\end{aligned}$$

and thus under Assumption 5, we obtain (13).

Next, consider the following perturbed parametrized program  $\mathsf{P}_{\xi}$ :

$$\begin{array}{ll} \underset{\beta \in \mathbb{R}^k}{\text{minimize}} & h(\beta) + \beta^\top \xi_1 \\ \text{subject to} & C\beta - d - \xi_2 \le 0, \end{array}$$
 (P<sub>\xi</sub>)

for a parameter  $\xi = (\xi_1, \xi_2) \in \mathbb{R}^k \times \mathbb{R}^r$ . ( $\mathsf{P}_{\xi}$ ) can be viewed as a perturbed program of ( $\mathsf{P}_{sl}$ ); for  $\xi = 0$ , ( $\mathsf{P}_{\xi}$ ) coincides with the program ( $\mathsf{P}_{sl}$ ). Let  $\bar{\beta}(\xi)$  denote the solution of the program  $\mathsf{P}_{\xi}$ . Clearly, we get  $\bar{\beta}(0) = \beta^*$ .

By Theorem 4.2 and Assumption 5, we obtain  $\hat{\beta} \xrightarrow{p} \beta^*$ . Furthermore, from (13), it follows that  $\hat{\gamma} \xrightarrow{p} \gamma^*$ , thereby ensuring that the consistency conditions Assumption 1 are satisfied. Also, by the positive definiteness of  $\nabla^2_{\beta} f(\beta^*)$  and the fact that we have only linear constraints, the uniform version of the quadratic growth condition also holds at  $\bar{\beta}(\xi)$  (see Shapiro (1993, Assumption A3)). Hence, given that the SC condition holds (Assumption 2), by Shapiro (1993, Theorem 3.1), we have

$$\widehat{\beta} = \overline{\beta}(\zeta) + o_{\mathbb{P}}(n^{-1/2}), \tag{16}$$

where

$$\zeta = \begin{bmatrix} \nabla_{\beta} \hat{h}(\beta^{*}) - \nabla_{\beta} h(\beta^{*}) - \sum_{j \in J_{0}(\beta^{*})} \gamma_{j}^{*} \left\{ \hat{C}_{j}^{\top} - C_{j}^{\top} \right\} \\ -(\hat{\mathsf{C}}_{ac} - \mathsf{C}_{ac})\beta^{*} \end{bmatrix} \equiv \begin{bmatrix} \zeta_{1} \\ \zeta_{2} \end{bmatrix}.$$

If  $\bar{\beta}(\xi)$  is differentiable at  $\xi = 0$  in the sense of Frechet, we have

$$\beta(\xi) - \beta^* = D_0\beta(\xi) + o(\|\xi\|),$$

where the mapping  $D_0\bar{\beta}(\cdot): \mathbb{R}^k \to \mathbb{R}^k$  is the directional derivative of  $\bar{\beta}(\cdot)$  at  $\xi = 0$ . Thus in this case, letting  $\xi = \zeta$  yields

$$n^{1/2}\left(\widehat{\beta}-\beta^*\right)=D_0\overline{\beta}(n^{1/2}\zeta)+o_{\mathbb{P}}(1),$$

which follows by (16) and that  $\bar{\beta}(0) = \beta^*$ .

Now we show that such mapping  $D_0\bar{\beta}(\cdot)$  exists and is indeed linear. To this end, we will show that  $\bar{\beta}(\xi)$  is locally totally differentiable at  $\xi = 0$ , followed by the implicit function theorem. Define a vector-valued function  $H \in \mathbb{R}^{k+|J_0(\beta^*)|}$  by

$$H(x,\xi,\gamma) = \begin{pmatrix} \nabla_{\beta}h(\beta) + C^{\top}\gamma + \xi_1 \\ \operatorname{diag}(\gamma)(C\beta - d - \xi_2) \end{pmatrix}.$$

Let the solution of  $H(x,\xi,\gamma) = 0$  be  $\bar{\beta}(\xi), \bar{\gamma}(\xi)$ . By virtue of the SC condition,  $\bar{\beta}(\xi), \bar{\gamma}(\xi)$ satisfies the KKT conditions for  $(\mathsf{P}_{\xi})$ . Then by the classical implicit function theorem (e.g., Dontchev and Rockafellar 2009, Theorem 1B.1), there exists a neighborhood  $\mathbb{B}_{\bar{r}}(0)$ , for some  $\bar{r} > 0$ , of  $\xi = 0$  such that  $\bar{\beta}(\xi)$  and its total derivative exist for  $\forall \xi \in \mathbb{B}_{\bar{r}}(0)$ . In particular, the derivative at  $\xi = 0$  is computed by

$$\nabla_{\boldsymbol{\xi}}\bar{\boldsymbol{\beta}}(0) = -\boldsymbol{J}_{\boldsymbol{\beta},\boldsymbol{\gamma}}H(\bar{\boldsymbol{\beta}}(0),0,\bar{\boldsymbol{\gamma}}(0))^{-1}\left[\boldsymbol{J}_{\boldsymbol{\xi}}H(\bar{\boldsymbol{\beta}}(0),0,\bar{\boldsymbol{\gamma}}(0))\right],$$

where in our case  $\bar{\beta}(0) = \beta^*, \bar{\gamma}(0) = \gamma^*$ , and thus

$$\boldsymbol{J}_{\boldsymbol{\beta},\boldsymbol{\gamma}}H(\bar{\boldsymbol{\beta}}(0),0,\bar{\boldsymbol{\gamma}}(0)) = \begin{bmatrix} \nabla_{\boldsymbol{\beta}}^{2}h(\boldsymbol{\beta}^{*}) & \mathsf{C}_{ac} \\ \mathsf{C}_{ac}^{\top} & \mathbf{0} \end{bmatrix},$$

and

$$oldsymbol{J}_{\xi}H(ar{eta}(0),0,ar{\gamma}(0)) = egin{bmatrix} oldsymbol{1} \ -\operatorname{diag}(\gamma)oldsymbol{1} \end{bmatrix}.$$

Here the inverse of  $\mathbf{J}_{\beta,\gamma}H(\bar{\beta}(0), 0, \bar{\gamma}(0))$  always exists under the LICQ condition (Assumption 2). Therefore we obtain that

$$D_0\bar{\beta}(n^{1/2}\zeta) = \begin{bmatrix} \nabla_{\beta}^2 h(\beta^*) & \mathsf{C}_{ac} \\ \mathsf{C}_{ac}^\top & \mathbf{0} \end{bmatrix}^{-1} \begin{bmatrix} \mathbf{1} \\ -\operatorname{diag}(\gamma_j^*)\mathbf{1} \end{bmatrix} n^{1/2}\zeta.$$

Since we assume that

$$n^{1/2}\zeta = n^{1/2} \begin{bmatrix} \nabla_{\beta} \hat{h}(\beta^*) - \nabla_{\beta} h(\beta^*) - \sum_{j \in J_0(\beta^*)} \gamma_j^* \left\{ \hat{C}_j^\top - C_j^\top \right\} \\ -(\hat{\mathsf{C}}_{ac} - \mathsf{C}_{ac})\beta^* \end{bmatrix} \xrightarrow{d} \Upsilon_{\beta^*},$$

for any random variable  $\Upsilon,$  by Slutsky's theorem we finally get the desired result

$$n^{1/2} \left( \widehat{eta} - eta^* 
ight) \xrightarrow{d} \begin{bmatrix} 
abla^2_{eta} h(eta^*) & \mathbf{C}_{ac} \\ \mathbf{C}^{ op}_{ac} & \mathbf{0} \end{bmatrix}^{-1} \begin{bmatrix} \mathbf{1} \\ -\operatorname{diag}(\gamma^*_j) \mathbf{1} \end{bmatrix} \Upsilon_{eta^*}.$$

L		
L		
L		