

A computational transition for detecting multivariate shuffled linear regression by low-degree polynomials

Zhangsong Li
Peking University

April 7, 2025

Abstract

In this paper, we study the problem of multivariate shuffled linear regression, where the correspondence between predictors and responses in a linear model is obfuscated by a latent permutation. Specifically, we investigate the model $Y = \frac{1}{\sqrt{1+\sigma^2}}(\Pi_* X Q_* + \sigma Z)$, where X is an $n * d$ standard Gaussian design matrix, Z is an $n * m$ Gaussian noise matrix, Π_* is an unknown $n * n$ permutation matrix, and Q_* is an unknown $d * m$ on the Grassmanian manifold satisfying $Q_*^\top Q_* = \mathbb{I}_m$.

Consider the hypothesis testing problem of distinguishing this model from the case where X and Y are independent Gaussian random matrices of sizes $n * d$ and $n * m$, respectively. Our results reveal a phase transition phenomenon in the performance of low-degree polynomial algorithms for this task. (1) When $m = o(d)$, we show that all degree- D polynomials fail to distinguish these two models even when $\sigma = 0$, provided with $D^4 = o(\frac{d}{m})$. (2) When $m = d$ and $\sigma = \omega(1)$, we show that all degree- D polynomials fail to distinguish these two models provided with $D = o(\sigma)$. (3) When $m = d$ and $\sigma = o(1)$, we show that there exists a constant-degree polynomial that strongly distinguish these two models. These results establish a smooth transition in the effectiveness of low-degree polynomial algorithms for this problem, highlighting the interplay between the dimensions m and d , the noise level σ , and the computational complexity of the testing task.

1 Introduction

In this paper, we study the multivariate shuffled linear regression problem, which is rigorously defined as follows.

Definition 1.1. *Consider the following linear model*

$$Y = \frac{1}{\sqrt{1+\sigma^2}}(\Pi_* X Q_* + \sigma Z), \quad (1.1)$$

here $X \in \mathbb{R}^{n*d}$ is the design matrix, $Q_* \in \mathbb{R}^{d*m}$ is the unknown regression matrix, Π_* is an unknown permutation matrix that shuffles the rows of X , and $Z \in \mathbb{R}^{n*m}$ is the observation noise. We focus on the Bayesian setting where $X \in \mathbb{R}^{n*d}$, $Z \in \mathbb{R}^{n*m}$ are two independent random matrices with i.i.d. standard normal entries, Π_* is a uniform $n * n$ permutation matrix, $Q_* \in \mathbb{R}^{d*m}$ is sampled from a prior distribution ν supported on all $d * m$ matrices and X, Z, Π_*, Q_* are independent.

Two basic problems regarding this model are as follows: (1) the *detection* problem, i.e., testing this model against two independent Gaussian matrices; (2) the *estimation* problem, i.e., recovering Π_* and Q_* from the observations (X, Y) . If Π_* is known, (1.1) reduces to standard linear

regression. Otherwise, this problem is known as shuffled regression [PWC17, APZ17+, LWX24], unlabeled sensing [UHV18, ZSL22, ZL20], or linear regression with permuted/mismatched data [SBD19, MW23, SDL20], as the correspondence between the predictors (the rows of X) and the responses (the rows of Y) is lost. Thus, it is a significantly more challenging problem as one needs to jointly estimate the permutation Π_* and the regression coefficients Q_* . This problem has attracted considerable theoretical and practical interest due to its applications in areas such as robotics, data integration, and de-anonymization; for further details, we refer the readers to [UHV18, Section 1] and [ZL20, Section 1.1]. In this work we are interested in the algorithmic aspect of the detection problem, where the goal is to solve problem (1) using *efficient algorithms* (i.e., algorithms with polynomial running time). Our results can be informally summarized as follows:

Theorem 1.2 (Informal). *Suppose $m \leq d$ and ν is the Haar measure over the $d * m$ Grassmanian manifold $\{Q \in \mathbb{R}^{d * m} : Q^\top Q = \mathbb{I}_m\}$. Then the following results hold:*

- (1) *when $m = o(d)$, there is evidence suggesting that all algorithms based on degree- D polynomials fails to distinguish (X, Y) with two independent Gaussian matrices even with $\sigma = 0$, provided that $D^4 = o(\frac{d}{m})$;*
- (2) *when $m = d$ and $\sigma = \omega(1)$, there is evidence suggesting that all algorithms based on degree- D polynomials fails to distinguish (X, Y) with two independent Gaussian matrices provided that $D = o(\sigma)$;*
- (3) *when $m = d$ and $\sigma = o(1)$, there is an efficient algorithm based on degree- $O(1)$ polynomial that strongly distinguish (X, Y) with two independent Gaussian matrices.*

Remark 1.3. *In view of the widely accepted assumption that the failure of degree- $O(\log n)$ polynomials are evidences for computational hardness and the close connection between detection and estimation, we tend to feel that the estimation problem is computationally hard when $d/m \geq \text{poly}(\log n)$ or when $d = m, \sigma \geq \text{poly}(\log n)$.*

In addition, we note that in the literature the shuffled linear regression problem (1.1) is usually stated in the non-Bayesian setting, where Q_ is just an arbitrary unknown $d * m$ matrix. However, since in the singular value decomposition $Q_* = U_* \Lambda_* V_*$ where $V_* \in O(m)$ and Λ_* is an $m * m$ diagonal matrix, assuming V_* and Λ_* is known to us will only simplify the problem (note that we can replace the observation Y with $Y V_*^\top \Lambda_*^-$ where Λ_*^- is the pseudo-inverse of Λ_*). Thus, our hardness result (1) covers the most widely studied non-Bayesian setting and suggests that estimation is hard in this setting in the similar parameter regime.*

1.1 Discussions

Information-computation gap and low-degree polynomial framework. A special case in our result is the single-variate setting $m = 1$, where $Q_* \in \mathbb{R}^d$ is sampled uniformly from the unit sphere \mathbb{S}^{d-1} . In this setting, the authors of [LWX24] determined the information threshold for recovering Π_* given (X, Y) when $d = o(n)$, which roughly speaking is $\sigma = O(n^{-2})$ for exact recovery and $\sigma = O(n^{-1})$ for almost-exact recovery. In addition, in a recent forthcoming work [GWX25+] the authors determined the information threshold for the detection problem up to a $\text{poly}(d)$ factor when $d = o(\frac{n}{\log n})$, conjectured to be $\sigma^2 = O(\frac{1}{d \log d})$.¹ However, when $d = \omega(\log n)$

¹To be more precise, in [GWX25+] the authors show that on the one hand, when $d = o(\frac{\log n}{\log \log n})$ the informational detection threshold is $\sigma^2 = o(\frac{1}{d \log d})$, and an efficient algorithm achieves the information threshold. On the other hand, for general $d = o(\frac{n}{\log n})$, detection is informationally possible if $\sigma^2 = O(\frac{1}{(d \log d)^\alpha})$ and informationally impossible if $\sigma^2 = \omega(\frac{1}{d \log d})$. They also conjecture that $\sigma^2 = O(\frac{1}{d \log d})$ is the true information threshold for detection.

currently no known polynomial-time algorithms that solve the detection or estimation problem except in the noiseless case $\sigma = 0$ [HSS17, AHSS17]. This suggests the presence of an *information-computation gap*, a phenomenon commonly observed in high-dimensional statistical inference tasks [ZK16, RSS19, KWB22, Gam21]. Since at this point it seems rather elusive to prove hardness on the detection problem for a typical instance under the assumption of $P \neq NP$, we can only hope to prove hardness under even (much) stronger hypothesis.

For this purpose, the *low-degree polynomial framework* emerged as a powerful tool to provide *evidences* for computational lower bounds. Indeed, it has been proved that the class of low-degree polynomial algorithms is a useful proxy for computationally efficient algorithms, in the sense that the best-known polynomial-time algorithms for a wide variety of high-dimensional inference problems are captured by the low-degree class such as spectral methods, approximate message passing and small subgraph counts; see e.g., [Hop18, SW22, KWB22]. Furthermore, it is conjectured in [Hop18] that the failure of degree- D polynomial algorithms implies the failure of all “robust” algorithms with running time $n^{\tilde{O}(D)}$ (here \tilde{O} means having at most this order up to a $\text{poly}(\log n)$ factor). Consequently, the failure of degree- $O(\text{poly}(\log n))$ polynomial algorithms to solve a problem is often interpreted as evidence of its intrinsic average-case hardness, or at least it is expected that breaking such impossibility results would require a major breakthrough in algorithms. Thus, our results provide further evidence supporting the existence of an information-computation gap in this problem when $d \geq \text{poly}(\log n)$.

Lattice-based algorithm “beats” low-degree polynomials. Although in the high-dimensional single-variate case $m = 1, d = \text{poly}(\log n)$ we provide evidence that all algorithms based on low-degree polynomials fail to achieve the detection task even in the noiseless case $\sigma = 0$, we note that there is a lattice-based polynomial-time algorithm solving the detection and estimation tasks when $\sigma = 0$ [HSS17, AHSS17]. Thus, the shuffled regression problem is a rare example where a lattice-based algorithm outperforms the family of low-degree polynomial algorithms. A similar phenomenon has been observed in the problem of clustering a variant of Gaussian mixtures, as discovered in [ZSWB22].

However, as discussed in [ZSWB22], this phenomenon does not necessary contradict the low-degree conjecture proposed in [Hop18]. Indeed, the lattice-based algorithms in both [ZSWB22] and [HSS17, AHSS17] are highly sensitive to the specifics of the model, and rely on the observations being “noiseless” in some sense. In fact, these algorithms appear to fail as long as the noise level satisfies $\sigma = \exp(-\text{poly}(n))$. Thus, we tend to feel that our low-degree hardness result actually suggests that all “robust” algorithms fail to solve the *noisy* detection/estimation problem even when the noise σ is polynomially small.

Procrustes matching and geometric graph matching. Another special case in our results which may be of particular interest is the setting where $m = d$ and Q_* is uniformly sampled from the family of $d \times d$ orthogonal matrices. This problem, known as *Procrustes matching*, has numerous applications in fields such as natural language processing and computer vision [RCB97, MDK+16, DL17, GJB19].

In addition, it was shown in [WWXY22] that this problem is further equivalent to a geometric variant of the random graph matching problems [WWXY22, GL24+, EGMM24]. More precisely, consider the Gaussian model

$$Y = \Pi_* X + \sigma Z, \quad \text{where } X, Y, Z \in \mathbb{R}^{n \times d}$$

such that X_i, Z_i being i.i.d. $\mathcal{N}(0, \mathbb{I}_d)$ vectors. The observation is two correlated Wishart matrices $A = XX^\top$ and $B = YY^\top$. In the low-dimensional regime where $d = O(\log n)$, the information

threshold for estimating Π_* was determined in [WWXY22] and an efficient algorithm approaching the information threshold up to a poly(d) factor was found in [GL24+]. In contrast, the related detection problem was less understood, especially in the high-dimensional regime $d = \omega(\log n)$. Our work shows the computation transition phenomenon for this detection problem under the low-degree polynomial framework, suggesting that (roughly speaking) $\sigma = O(1)$ is the separation of the computational “easy” and “hard” regime.

Open problems. While we have characterized the computation transition for low-degree polynomial algorithms in the detection problem, the information-theoretic threshold for this detection problem in multivariate case remains largely open. In the forthcoming work [GWX25+], inspired by the analysis of the broken sample problem [DCK20, KN22, JWX25+] the authors shows that in the single-variate case, the detection threshold is given by $\sigma^2 = O(\frac{1}{d \log d})$ when $d = o(\frac{\log n}{\log \log n})$ and it lies in the range $O(\frac{1}{d^2 (\log d)^2}) \ll \sigma^2 \ll O(\frac{1}{d \log d})$ when $\frac{\log n}{\log \log n} \ll d \ll \frac{n}{\log n}$. Extending these results to the multivariate setting presents an intriguing open problem.

Another interesting question left open by our work is whether we could provide stronger evidence on the information-computation gaps emerged in this detection problem when $d/m \geq \text{poly}(\log n)$. For instance, it would be interesting if we could establish the sum-of-squares lower bound [GJJ+20, JPRX23] or statistical query (SQ) lower bound [BBH+21, DKPZ23] for this problem. In addition, we note that in the problem of clustering a variant of Gaussian mixtures mentioned above, the hardness of the noisy problem can be shown by reducing to the standard assumption [MR09, Conjecture 1.2] from lattice-based cryptography that certain worst-case lattice problems are hard against quantum algorithms [BRST21]. However, it remains unclear whether similar reductions exist for our problem.

1.2 Notations

We record in this subsection some notation conventions. Denote μ to be the uniform distribution over the set of all permutation matrices \mathfrak{S}_n and denote ν to be the Haar distribution over the $d * m$ Grassmanian manifold. In addition, let \mathbb{S}^{d-1} be the unit d -dimensional sphere. For a matrix or a vector M , we will use M^\top to denote its transpose. Denote $O(m)$ to be the set of all $m * m$ orthogonal matrices. For a $k * k$ matrix $M = (m_{ij})_{k * k}$, let $\det(M)$ and $\text{tr}(M)$ be the determinant and trace of M , respectively. Denote $M \succ 0$ if M is positive definite and $M \succeq 0$ if M is semi-positive definite. In addition, if M is symmetric we let $\varsigma_1(M) \geq \varsigma_2(M) \geq \dots \geq \varsigma_k(M)$ be the eigenvalues of M . Denote by $\text{rank}(M)$ the rank of the matrix M . For two $k * l$ matrices M_1 and M_2 , we define their inner product to be

$$\langle M_1, M_2 \rangle := \sum_{i=1}^k \sum_{j=1}^l M_1(i, j) M_2(i, j).$$

We also define the Frobenius norm, operator norm, and ∞ -norm of M respectively by

$$\|M\|_F = \text{tr}(MM^\top) = \langle M, M \rangle^{\frac{1}{2}}, \quad \|M\|_{\text{op}} = \varsigma_1(MM^\top)^{\frac{1}{2}}, \quad \|M\|_\infty = \max_{\substack{1 \leq i \leq k \\ 1 \leq j \leq l}} |M_{i,j}|$$

where $\text{tr}(\cdot)$ is the trace for a squared matrix. We will use \mathbb{I}_k to denote the $k * k$ identity matrix (and we drop the subscript if the dimension is clear from the context). Similarly, we denote $\mathbb{O}_{k * l}$ the $k * l$ zero matrix and denote $\mathbb{J}_{k * l}$ the $k * l$ matrix with all entries being 1. We will abbreviate $\mathbb{O}_k = \mathbb{O}_{k * 1}$ and $\mathbb{J}_k = \mathbb{J}_{k * 1}$. The indicator function of sets A is denoted by $\mathbf{1}_A$.

For any $\alpha = (\alpha_1, \dots, \alpha_k) \in \mathbb{N}^k$, define $|\alpha| = \sum_{1 \leq i \leq k} \alpha_i$ and $\alpha! = \alpha_1! \dots \alpha_k!$. In addition, for any $c \in \mathbb{N}$ and $x \in \mathbb{R}^k$, define $c\alpha = (c\alpha_1, \dots, c\alpha_k)$ and $x^\alpha = x_1^{\alpha_1} \dots x_k^{\alpha_k}$. For $m \in \mathbb{N}$ and $\alpha \in \mathbb{N}^k$ with $|\alpha| = m$, denote $\binom{m}{\alpha} = \frac{m!}{\alpha_1! \dots \alpha_d!}$. We denote $\mathbf{A} \in (\mathbb{N}^k)^{\otimes n}$, if $\mathbf{A} = (\alpha_i : i \in [n])$ such that $\alpha_i \in \mathbb{N}^k$. For two $\mathbf{A}, \mathbf{A}' \in (\mathbb{N}^k)^{\otimes n}$ with $\mathbf{A} = (\alpha_i : i \in [n])$ and $\mathbf{A}' = (\alpha'_i : i \in [n])$, we say $\mathbf{A} \cong \mathbf{A}'$ if and only if there exists a bijection $\pi : [n] \rightarrow [n]$ such that $\alpha_i = \alpha'_{\pi(i)}$ for all $i \in [n]$. Denote $\text{Aut}(\mathbf{A})$ to be the number of bijections $\pi : [n] \rightarrow [n]$ such that $\alpha_i = \alpha_{\pi(i)}$ for all $i \in [n]$. For a set A , we use both $|A|$ and $\#A$ to denote its cardinality.

For any two positive sequences $\{a_n\}$ and $\{b_n\}$, we write equivalently $a_n = O(b_n)$, $b_n = \Omega(a_n)$, $a_n \lesssim b_n$ and $b_n \gtrsim a_n$ if there exists a positive absolute constant c such that $a_n/b_n \leq c$ holds for all n . We write $a_n = o(b_n)$, $b_n = \omega(a_n)$, $a_n \ll b_n$, and $b_n \gg a_n$ if $a_n/b_n \rightarrow 0$ as $n \rightarrow \infty$. We write $a_n = \Theta(b_n)$ if both $a_n = O(b_n)$ and $a_n = \Omega(b_n)$ hold.

2 The low-degree polynomial framework

In this section we provide the formal statement of Theorem 1.2. More precisely, we will consider the following hypothesis testing problem, where

- under the null hypothesis \mathcal{H}_0 , we let $X \in \mathbb{R}^{n \times d}$ and $Y \in \mathbb{R}^{n \times m}$ be two independent Gaussian matrices with i.i.d. standard normal entries, respectively;
- under the alternative hypothesis \mathcal{H}_1 , we let (X, Z, Π_*, Q_*) be independent random matrices with $X \in \mathbb{R}^{n \times d}$, $Z \in \mathbb{R}^{n \times m}$ having i.i.d. standard normal entries, Π_* an $n \times n$ uniform permutation matrix and Q_* sampled from ν , and let $Y = \frac{1}{\sqrt{1+\sigma^2}}(\Pi_* X Q_* + \sigma Z)$.

We establish evidences the computational hardness of this hypothesis testing problem by focusing on a specific class of algorithms known as *low-degree polynomials*. Inspired by the sum-of-squares hierarchy, the low-degree polynomial method offers a promising approach for deriving computational lower bounds in high-dimensional inference problems. At its core, this method analyzes algorithms that rely on evaluating collections of polynomials with moderate degrees. The study of this category of algorithms is motivated by the examination of high-dimensional hypothesis testing problems [BHK+19, Hop18, HKP+17, HS17], with an extensive overview provided in [KWB22]. A key strength of the low-degree polynomial approach is its ability to deliver tight hardness results for a wide range of problems. Notable examples include detection problems such as planted clique, planted dense subgraph, community detection, sparse PCA, correlated random graphs (see [HS17, HKP+17, Hop18, KWB22, SW22, DMW23+, BKW20, DKW+22, MW25, DDL23+, KMW24, CDGL24+]), optimization problems such as maximal independent sets in sparse random graphs [GJW20, Wein22], and constraint satisfaction problems such as random k -SAT [BH22]. In the remaining of this paper, we will focus on applying this framework in the context of hypothesis testing problems.

More precisely, denote $\mathbb{P} = \mathbb{P}_n$ to be the law of (X, Y) under \mathcal{H}_1 and $\mathbb{Q} = \mathbb{Q}_n$ to be the law of (X, Y) under \mathcal{H}_0 . In addition, let $\mathbb{R}[X, Y]_{\leq D}$ denote the set of multivariate polynomials in the entries of (X, Y) with degree at most D . With a slight abuse of notation, we will often say “a polynomial” to mean a sequence of polynomials $f = f_n \in \mathbb{R}[X, Y]_{\leq D}$, one for each problem size n ; the degree $D = D_n$ of such a polynomial may scale with n . To probe the computational threshold for testing between two sequences of probability measures \mathbb{P} and \mathbb{Q} , we consider the following notions of strong separation and weak separation defined in [BAH+22, Definition 1.6].

Definition 2.1. Let $f \in \mathbb{R}[X, Y]_{\leq D}$ be a polynomial.

- We say f strongly separates \mathbb{P}_n and \mathbb{Q}_n if as $n \rightarrow \infty$

$$\sqrt{\max \{ \text{Var}_{\mathbb{P}}(f(A, B)), \text{Var}_{\mathbb{Q}}(f(A, B)) \}} = o(|\mathbb{E}_{\mathbb{P}}[f(A, B)] - \mathbb{E}_{\mathbb{Q}}[f(A, B)]|);$$

- We say f weakly separates \mathbb{P}_n and \mathbb{Q}_n if as $n \rightarrow \infty$

$$\sqrt{\max \{ \text{Var}_{\mathbb{P}}(f(A, B)), \text{Var}_{\mathbb{Q}}(f(A, B)) \}} = O(|\mathbb{E}_{\mathbb{P}}[f(A, B)] - \mathbb{E}_{\mathbb{Q}}[f(A, B)]|).$$

See [BAH+22] for a detailed discussion of why these conditions are natural for hypothesis testing. In particular, according to Chebyshev's inequality, strong separation implies that we can threshold $f(A, B)$ to test \mathbb{P} against \mathbb{Q} with vanishing type-I and type-II errors. Now we can state our rigorous statement justifying Theorem 1.2.

Theorem 2.2. (1) Suppose $1 \leq m = o(d)$ and $D^4 = o(\frac{d}{m})$. Then no degree- D polynomial weakly separates \mathbb{P} and \mathbb{Q} .

(2) Suppose $m = d, \sigma = \omega(1)$ and $D = o(\sigma^{-1})$. Then no degree- D polynomial weakly separates \mathbb{P} and \mathbb{Q} .

(3) Suppose $m = d$ and $\sigma = o(1)$. Then there is a degree- $O(1)$ polynomial that strongly separates \mathbb{P} and \mathbb{Q} .

3 Proof of Theorem 2.2

In this section we prove Theorem 2.2 formally. In Subsection 3.1 we provide a universal bound for the low-degree advantage between \mathbb{P} and \mathbb{Q} for all $d, m \in \mathbb{N}$. In Subsections 3.2 we prove Item (1) in the single-variate as preparations. In Subsections 3.3, 3.4 and 3.5 we prove Items (1), (2) and (3) in Theorem 2.2 respectively.

3.1 Universal bound on low-degree advantage

We begin with the classical framework for proving the failure of weak separation of polynomials, as introduced in [HS17].

Definition 3.1. For a quadruple (n, d, m, D) with d, m, D possibly depend on n , we define the low-degree advantage by

$$\text{Adv}_{\leq D}(\mathbb{P}, \mathbb{Q}) := \sup_{f \in \mathbb{R}[X, Y]_{\leq D}} \frac{\mathbb{E}_{\mathbb{P}}[f]}{\sqrt{\mathbb{E}_{\mathbb{Q}}[f^2]}}. \quad (3.1)$$

It is clear (see [BAH+22, Definition 1.8]) that $\text{Adv}_{\leq D}(\mathbb{P}, \mathbb{Q}) = 1 + o(1)$ as $n \rightarrow \infty$ implies that no polynomial in $\mathbb{R}[X, Y]_{\leq D}$ weakly separates \mathbb{P} and \mathbb{Q} .

The rest of this section is devoted to proving $\text{Adv}_{\leq D}(\mathbb{P}, \mathbb{Q}) = 1 + o(1)$ under various settings. The following polynomials will play a fundamental role in our analysis.

Definition 3.2. For all $m \in \mathbb{N}$, define the Hermite polynomials by

$$H_0(z) = 1, \quad H_1(z) = z, \quad H_{m+1}(z) = zH_m(z) - mH_{m-1}(z). \quad (3.2)$$

In addition, define $\mathcal{H}_m(z) = \frac{1}{\sqrt{m!}} H_m(z)$, and for all $x \in \mathbb{R}^d$ and $\alpha \in \mathbb{N}^d$ define

$$\mathcal{H}_\alpha(x) := \prod_{i=1}^d \mathcal{H}_{\alpha_i}(x_i). \quad (3.3)$$

For all $y \in \mathbb{R}^m$ and $\beta \in \mathbb{N}^m$, define $\mathcal{H}_\beta(y)$ in the similar manner. For all $\mathbf{A} = (\alpha_i : i \in [n]) \in (\mathbb{N}^d)^{\otimes n}$ and $\mathbf{B} = (\beta_j : j \in [n]) \in (\mathbb{N}^m)^{\otimes n}$ such that $\alpha_i \in \mathbb{N}^d, \beta_j \in \mathbb{N}^m$, define

$$\phi_{\mathbf{A}, \mathbf{B}}(X, Y) = \prod_{1 \leq i \leq n} \mathcal{H}_{\alpha_i}(X_i) \prod_{1 \leq j \leq n} \mathcal{H}_{\beta_j}(Y_j). \quad (3.4)$$

Define $|\mathbf{A}| = \sum_{1 \leq i \leq n} |\alpha_i|$ and $|\mathbf{B}| = \sum_{1 \leq j \leq n} |\beta_j|$. It is well known (see, e.g., [Sze39]) that $\{\phi_{\mathbf{A}, \mathbf{B}} : |\mathbf{A}| + |\mathbf{B}| \leq D\}$ forms a standard orthogonal basis of $\mathbb{R}[X, Y]_{\leq D}$ under the measure \mathbb{Q} , i.e., we have

$$\mathbb{E}_{\mathbb{Q}}[\phi_{\mathbf{A}, \mathbf{B}} \phi_{\mathbf{A}', \mathbf{B}'}] = \mathbf{1}_{\{(\mathbf{A}, \mathbf{B}) = (\mathbf{A}', \mathbf{B}')\}}. \quad (3.5)$$

Consequently, we can explicitly express the low-degree advantage $\text{Adv}_{\leq D}(\mathbb{P}, \mathbb{Q})$ in terms of this basis, as in the next lemma.

Lemma 3.3. *Recall (3.1). For any $n, D \geq 1$, it holds that*

$$\text{Adv}_{\leq D}(\mathbb{P}, \mathbb{Q}) = \left(\sum_{(\mathbf{A}, \mathbf{B}) : |\mathbf{A}| + |\mathbf{B}| \leq D} \mathbb{E}_{\mathbb{P}}[\phi_{\mathbf{A}, \mathbf{B}}(X, Y)]^2 \right)^{1/2}. \quad (3.6)$$

Proof. For any $f \in \mathbb{R}[X, Y]_{\leq D}$, it can be uniquely expressed as

$$f = \sum_{(\mathbf{A}, \mathbf{B}) : |\mathbf{A}| + |\mathbf{B}| \leq D} C_{\mathbf{A}, \mathbf{B}} \phi_{\mathbf{A}, \mathbf{B}},$$

where $C_{\mathbf{A}, \mathbf{B}}$'s are real constants. Applying Cauchy-Schwartz inequality one gets

$$\frac{\mathbb{E}_{\mathbb{P}}[f]}{\sqrt{\mathbb{E}_{\mathbb{Q}}[f^2]}} = \frac{\sum_{(\mathbf{A}, \mathbf{B}) : |\mathbf{A}| + |\mathbf{B}| \leq D} C_{\mathbf{A}, \mathbf{B}} \mathbb{E}_{\mathbb{P}}[\phi_{\mathbf{A}, \mathbf{B}}]}{\sqrt{\sum_{(\mathbf{A}, \mathbf{B}) : |\mathbf{A}| + |\mathbf{B}| \leq D} C_{\mathbf{A}, \mathbf{B}}^2}} \leq \left(\sum_{(\mathbf{A}, \mathbf{B}) : |\mathbf{A}| + |\mathbf{B}| \leq D} (\mathbb{E}_{\mathbb{P}}[\phi_{\mathbf{A}, \mathbf{B}}])^2 \right)^{1/2},$$

with equality holds if and only if $C_{\mathbf{A}, \mathbf{B}} \propto \mathbb{E}_{\mathbb{P}}[\phi_{\mathbf{A}, \mathbf{B}}]$. \square

The rest part of this section is devoted to bounding the right hand side of (3.6) under various assumptions in Theorem 2.2. For $\alpha \in \mathbb{N}^d, \beta \in \mathbb{N}^m$ and $Q \in \mathbb{R}^{d \times m}$, denote

$$\Lambda_{\alpha, \beta}(Q) := \mathbb{E}_U \left[\mathcal{H}_\alpha(U) \mathcal{H}_\beta \left(\frac{1}{\sqrt{1 + \sigma^2}} (UQ + \sigma V) \right) \right], \text{ where } U \perp V \text{ and } U, V \sim \mathcal{N}(0, \mathbb{I}_d). \quad (3.7)$$

We first bound the right hand side of (3.6) by $\Lambda_{\alpha, \beta}(Q)$'s.

Lemma 3.4. *We have*

$$\sum_{|\mathbf{A}| + |\mathbf{B}| \leq D} \mathbb{E}_{\mathbb{P}}[\phi_{\mathbf{A}, \mathbf{B}}(X, Y)]^2 \leq \sum_{0 \leq k \leq D} \sum_{\substack{(\alpha_1, \dots, \alpha_k) : \alpha_i \in \mathbb{N}^d \\ (\beta_1, \dots, \beta_k) : \beta_i \in \mathbb{N}^m \\ 0 < |\alpha_i|, |\beta_j| \leq D}} \mathbb{E}_{Q \sim \nu} \left[\prod_{1 \leq i \leq k} \Lambda_{\alpha_i, \beta_i}(Q) \right]^2. \quad (3.8)$$

Proof. For all $\mathbf{A} \in (\mathbb{N}^m)^{\otimes n}$ and $\mathbf{B} \in (\mathbb{N}^d)^{\otimes n}$, we can characterize (\mathbf{A}, \mathbf{B}) with two subsets $S, K \subset [n]$ and two ordered families of integer vectors $(\alpha_1, \dots, \alpha_{|S|}), (\beta_1, \dots, \beta_{|K|})$ where $\alpha_i \in \mathbb{N}^d, \beta_j \in \mathbb{N}^m$ and $\alpha_i \neq \mathbb{O}_d$ and $\beta_j \neq \mathbb{O}_m$. Also denote $\mathbb{P}_\pi = \mathbb{P}(\cdot \mid \pi_* = \pi)$. Note that when $\pi(S) \neq K$, suppose that there exists $j \in S \setminus \pi^{-1}(K)$, then

$$\begin{aligned} \mathbb{E}_{\mathbb{P}_\pi}[\phi_{\mathbf{A}, \mathbf{B}}] &= \mathbb{E}_{Q \sim \nu, X_i \sim \mathcal{N}(0, \mathbb{I}_d)} \left[\prod_{1 \leq i \leq n} \mathcal{H}_{\alpha_i}(X_i) \mathcal{H}_{\beta_i}(X_i Q) \right] \\ &= \mathbb{E}_{Q \sim \nu} \left[\mathbb{E}_X \left[\mathcal{H}_{\alpha_j}(X_j) \right] \mathbb{E}_X \left[\prod_{i \neq j} \mathcal{H}_{\alpha_i}(X_i) \mathcal{H}_{\beta_i}(X_i Q) \right] \right] = 0. \end{aligned}$$

Similarly, when there exists $j \in K \setminus \pi(S)$ we can check that $\mathbb{E}_{\mathbb{P}_\pi}[\phi_{\mathbf{A}, \mathbf{B}}] = 0$. If $\pi(S) = K$, let σ to be the restriction of π on S , we then have

$$\mathbb{E}_{\mathbb{P}_\pi}[\phi_{\mathbf{A}, \mathbf{B}} \mid Q] = \prod_{1 \leq i \leq |S|} \Lambda_{\alpha_i, \beta_{\sigma(i)}}(Q).$$

Thus, we have shown that

$$\mathbb{E}_{\mathbb{P}}[\phi_{\mathbf{A}, \mathbf{B}}] = \begin{cases} 0, & |S| \neq |K|; \\ \frac{(n-k)!}{n!} \sum_{\sigma \in \mathfrak{S}_k} \mathbb{E}_{Q \sim \nu} \left[\prod_{1 \leq i \leq k} \Lambda_{\alpha_i, \beta_{\sigma(i)}}(Q) \right], & |S| = |K| = k. \end{cases}$$

Thus, we have the left hand side of (3.8) is bounded by

$$\begin{aligned} & \sum_{0 \leq k \leq D} \sum_{\substack{S, K \subset [n] \\ |S|=|K|=k}} \sum_{\substack{\alpha_1, \dots, \alpha_k \in \mathbb{N}^d \\ \beta_1, \dots, \beta_k \in \mathbb{N}^m \\ 0 < |\alpha_i|, |\beta_j| \leq D}} \left(\frac{(n-k)!}{n!} \sum_{\sigma \in \mathfrak{S}_k} \mathbb{E}_{Q \sim \nu} \left[\prod_{1 \leq i \leq k} \Lambda_{\alpha_i, \beta_{\sigma(i)}}(Q) \right] \right)^2 \\ & \leq \sum_{0 \leq k \leq D} \sum_{\substack{\alpha_1, \dots, \alpha_k \in \mathbb{N}^d \\ \beta_1, \dots, \beta_k \in \mathbb{N}^m \\ 0 < |\alpha_i|, |\beta_j| \leq D}} \left(\frac{1}{k!} \sum_{\sigma \in \mathfrak{S}_k} \mathbb{E}_{Q \sim \nu} \left[\prod_{1 \leq i \leq k} \Lambda_{\alpha_i, \beta_{\sigma(i)}}(Q) \right] \right)^2. \end{aligned} \quad (3.9)$$

Using Cauchy-Schwartz inequality, we have (3.9) is bounded by

$$\begin{aligned} & \sum_{0 \leq k \leq D} \sum_{\substack{\alpha_1, \dots, \alpha_k \in \mathbb{N}^d \\ \beta_1, \dots, \beta_k \in \mathbb{N}^m \\ 0 < |\alpha_i|, |\beta_j| \leq D}} \frac{1}{k!} \sum_{\sigma \in \mathfrak{S}_k} \mathbb{E}_{Q \sim \nu} \left[\prod_{1 \leq i \leq k} \Lambda_{\alpha_i, \beta_{\sigma(i)}}(Q) \right]^2 \\ & = \sum_{0 \leq k \leq D} \sum_{\substack{\alpha_1, \dots, \alpha_k \in \mathbb{N}^d \\ \beta_1, \dots, \beta_k \in \mathbb{N}^m \\ 0 < |\alpha_i|, |\beta_j| \leq D}} \mathbb{E}_{Q \sim \nu} \left[\prod_{1 \leq i \leq k} \Lambda_{\alpha_i, \beta_i}(Q) \right]^2, \end{aligned}$$

where in equality we use that fix any $\sigma \in \mathfrak{S}_k$ there is a one-to-one correspondence between $(\beta_1, \dots, \beta_k)$ and $(\beta_{\sigma(1)}, \dots, \beta_{\sigma(k)})$. This completes our proof. \square

3.2 Proof of Item (1) in single-variate case

To get a feeling on how we bound the right hand side of (3.8), we first prove Item (1) of Theorem 2.2 in single-variate case (with a milder assumption on D)

$$m = 1, \sigma = 0, D^6 = o(d).$$

Based on Lemma 3.4, we see that

$$\text{Adv}_{\leq D}(\mathbb{P}, \mathbb{Q})^2 \leq \sum_{0 \leq k \leq D} \sum_{\substack{\alpha_1, \dots, \alpha_k \in \mathbb{N}^d \\ \beta_1, \dots, \beta_k \in \mathbb{N} \\ 0 < |\alpha_i|, |\beta_j| \leq D}} \mathbb{E}_{Q \sim \nu} \left[\prod_{1 \leq i \leq k} \Lambda_{\alpha_i, \beta_i}(Q) \right]^2, \quad (3.10)$$

where

$$\Lambda_{\alpha, \beta}(Q) = \mathbb{E}_{U \sim \mathcal{N}(0, \mathbb{I}_d)} \left[\mathcal{H}_\alpha(U) \mathcal{H}_\beta(\langle U, Q \rangle) \right].$$

In this case, we can calculate $\Lambda_{\alpha, \beta}(Q)$ for each $\alpha \in \mathbb{N}^d$ and $\beta \in \mathbb{N}$ directly, as incorporated in the next lemma.

Lemma 3.5. *We have*

$$\Lambda_{\alpha, \beta}(Q) = \mathbf{1}_{\{|\alpha|=\beta\}} \cdot \mathfrak{M}(\alpha; \beta) Q^\alpha, \quad (3.11)$$

where

$$\mathfrak{M}(\alpha; \beta) = \sqrt{\frac{\alpha!}{\beta!}} \cdot \binom{\beta}{\alpha}. \quad (3.12)$$

Proof. By direct calculation, we have that

$$\begin{aligned} \Lambda_{\alpha, \beta}(Q) &\stackrel{\text{Lemma A.1}}{=} \sum_{\gamma \in \mathbb{N}^d: |\gamma|=\beta} \mathfrak{M}(\gamma; \beta) \mathbb{E}_{U \sim \mathcal{N}(0, \mathbb{I}_d)} \left[Q^\gamma \cdot \mathcal{H}_\gamma(U) \mathcal{H}_\alpha(U) \right] \\ &= \sum_{\gamma \in \mathbb{N}^d: |\gamma|=\beta} \mathbf{1}_{\{\gamma=\alpha\}} \cdot \mathfrak{M}(\gamma; \beta) Q^\gamma = \mathbf{1}_{\{|\alpha|=\beta\}} \cdot \mathfrak{M}(\alpha; \beta) Q^\alpha, \end{aligned}$$

as desired. □

We can now finish the proof of Theorem 2.2, Item (1).

Proof of Theorem 2.2, Item (1). Using Lemma 3.5, we get that

$$\begin{aligned} &\sum_{\substack{\alpha_1, \dots, \alpha_k \in \mathbb{N}^d; \beta_1, \dots, \beta_k \in \mathbb{N} \\ 0 < |\alpha_i|, |\beta_j| \leq D}} \mathbb{E}_{Q \sim \nu} \left[\prod_{1 \leq i \leq k} \Lambda_{\alpha_i, \beta_i}(Q) \right]^2 \\ &= \sum_{\substack{\alpha_1, \dots, \alpha_k \in \mathbb{N}^d; \beta_1, \dots, \beta_k \in \mathbb{N} \\ 0 < |\alpha_i|, |\beta_j| \leq D}} \mathbb{E}_{Q \sim \nu} \left[\prod_{1 \leq i \leq k} \left(\mathbf{1}_{\{|\alpha_i|=\beta_i\}} \cdot \mathfrak{M}(\alpha_i; \beta_i) Q^{\alpha_i} \right) \right]^2 \\ &= \sum_{\substack{\alpha_1, \dots, \alpha_k \in \mathbb{N}^d \\ \alpha_i \neq \mathbf{0}_d, |\alpha_i| \leq D}} \mathfrak{M}(\alpha_1; |\alpha_1|)^2 \dots \mathfrak{M}(\alpha_k; |\alpha_k|)^2 \mathbb{E}_{Q \sim \nu} \left[Q^{\alpha_1 + \dots + \alpha_k} \right]^2 \\ &\stackrel{(3.12)}{\leq} \sum_{\gamma \neq \mathbf{0}_d, |\gamma| \leq D} \mathbb{E}_{Q \sim \nu} \left[Q^\gamma \right]^2 \sum_{\substack{\alpha_1 + \dots + \alpha_k = \gamma \\ \alpha_i \neq \mathbf{0}, |\alpha_i| \leq D}} \binom{|\alpha_1|}{\alpha_1} \dots \binom{|\alpha_k|}{\alpha_k}. \end{aligned} \quad (3.13)$$

Using Lemma A.4, we see that

$$\begin{aligned}
(3.13) &= \sum_{\gamma \neq \mathbb{O}_d, |\gamma| \leq D/2} \mathbb{E}_{Q \sim \nu} [Q^{2\gamma}]^2 \sum_{\substack{\alpha_1 + \dots + \alpha_k = 2\gamma \\ \alpha_i \neq \mathbb{O}_d, |\alpha_i| \leq D}} \binom{|\alpha_1|}{\alpha_1} \dots \binom{|\alpha_k|}{\alpha_k} \\
&= \sum_{\gamma \neq \mathbb{O}_d, |\gamma| \leq D/2} \left(\frac{\Gamma(\frac{d}{2})(2\gamma_1 - 1)!! \dots (2\gamma_d - 1)!!}{\Gamma(\frac{d}{2} + \gamma_1 + \dots + \gamma_d) 2^{\gamma_1 + \dots + \gamma_d}} \right)^2 \sum_{\substack{\alpha_1 + \dots + \alpha_k = 2\gamma \\ \alpha_i \neq \mathbb{O}_d, |\alpha_i| \leq D}} \binom{|\alpha_1|}{\alpha_1} \dots \binom{|\alpha_k|}{\alpha_k} \\
&\leq \sum_{\gamma \neq \mathbb{O}_d, |\gamma| \leq D/2} d^{-2|\gamma|} D^{2|\gamma|} \sum_{\substack{\alpha_1 + \dots + \alpha_k = 2\gamma \\ \alpha_i \neq \mathbb{O}_d, |\alpha_i| \leq D}} \binom{|\alpha_1|}{\alpha_1} \dots \binom{|\alpha_k|}{\alpha_k}, \tag{3.14}
\end{aligned}$$

where the inequality follows from

$$\left((2\gamma_1 - 1)!! \dots (2\gamma_d - 1)!! \right)^2 \leq D^{2(\gamma_1 + \dots + \gamma_d)} = D^{2|\gamma|}$$

for $|\gamma| \leq D/2$. In addition, we have that

$$\begin{aligned}
\sum_{\substack{\alpha_1 + \dots + \alpha_k = 2\gamma \\ \alpha_i \neq \mathbb{O}_d, |\alpha_i| \leq D}} \binom{|\alpha_1|}{\alpha_1} \dots \binom{|\alpha_k|}{\alpha_k} &\leq \sum_{\substack{\alpha_1 + \dots + \alpha_k = 2\gamma \\ \alpha_i \neq \mathbb{O}_d, |\alpha_i| \leq D}} |\alpha_1|! \dots |\alpha_k|! \\
&\leq D^{|\alpha_1| + \dots + |\alpha_k|} \cdot \#\{\alpha_1 + \dots + \alpha_k = 2\gamma : \alpha_i \neq \mathbb{O}_d, |\alpha_i| \leq D\} \\
&\leq D^{2|\gamma|} \cdot k^{2|\gamma|}.
\end{aligned}$$

Plugging this estimation into (3.14), we get that

$$\begin{aligned}
(3.14) &\leq \sum_{\gamma \neq \mathbb{O}_d, |\gamma| \leq D/2} d^{-2|\gamma|} (D^4 k^2)^{|\gamma|} \leq \sum_{k=1}^{D/2} d^{-2k} (D^4 k^2)^k \cdot \#\{\gamma \neq \mathbb{O}_d : |\gamma| = k\} \\
&\leq \sum_{k=1}^{D/2} d^{-2k} (D^4 k^2)^k \cdot \binom{d+k}{k} \leq \sum_{k=1}^{D/2} d^{-k} (D^4 k)^k = [1 + o(1)] \cdot \frac{D^4 k}{d},
\end{aligned}$$

where the last transition follows from $D^4 k \leq D^5 = o(d)$. Thus, we get from (3.10) that

$$\text{Adv}_{\leq D}(\mathbb{P}, \mathbb{Q})^2 \leq 1 + \sum_{1 \leq k \leq D/2} [1 + o(1)] \cdot \frac{D^4 k}{d} = [1 + o(1)] \cdot \frac{D^6}{d} = 1 + o(1),$$

as desired. \square

3.3 Proof of Item (1) for general dimensions

In this subsection we extend our argument to the general case that $1 \leq m = o(d)$ and $D^4 = o(\frac{d}{m})$. Still, based on (3.8), it suffices to show that for $1 \leq k \leq D$ we have

$$\sum_{\substack{(\alpha_1, \dots, \alpha_k) : \alpha_i \in \mathbb{N}^d \setminus \{\mathbb{O}_d\}, |\alpha_i| \leq D \\ (\beta_1, \dots, \beta_k) : \beta_i \in \mathbb{N}^m \setminus \{\mathbb{O}_m\}, |\beta_i| \leq D}} \mathbb{E}_{Q \sim \nu} \left[\prod_{1 \leq i \leq k} \Lambda_{\alpha_i, \beta_i}(Q) \right]^2 = o(D^{-1}). \tag{3.15}$$

However, in this case it seems rather complicated to calculate $\Lambda_{\alpha, \beta}(Q)$ directly. Thus, we will take an alternative approach by first relating the left hand side of (3.15) to an easier hypothesis testing problem.

Definition 3.6. Consider the following hypothesis testing problem, where our observation are two vectors $X = (X_1, \dots, X_k)^\top \in \mathbb{R}^{k*d}$, $Y = (Y_1, \dots, Y_k)^\top \in \mathbb{R}^{k*m}$ such that:

- under the null hypothesis $\overline{\mathcal{H}}_0$, we sample $X_i \sim \mathcal{N}(0, \mathbb{I}_d)$ and $Y_i \sim \mathcal{N}(0, \mathbb{I}_m)$ independently.
- under the alternative hypothesis $\overline{\mathcal{H}}_1$, we first sample $X_1, \dots, X_k \sim \mathcal{N}(0, \mathbb{I}_d)$ and $Q \sim \nu$ independently, and then define $Y = XQ$.

In addition, we define \mathbb{P} to be the law of (X, Y) under $\overline{\mathcal{H}}_1$ and let \mathbb{Q} to be the law of (X, Y) under $\overline{\mathcal{H}}_0$, respectively.

Our next lemma shows that we can bound $\Lambda(\alpha, \beta)$ by the chi-square norm between \mathbb{P} and \mathbb{Q} .

Lemma 3.7. We have

$$\sum_{\substack{(\alpha_1, \dots, \alpha_k): \alpha_i \in \mathbb{N}^d \setminus \{\mathbb{O}_d\}, |\alpha_i| \leq D \\ (\beta_1, \dots, \beta_k): \beta_i \in \mathbb{N}^m \setminus \{\mathbb{O}_m\}, |\beta_i| \leq D}} \mathbb{E}_{Q \sim \nu} \left[\prod_{1 \leq i \leq k} \Lambda_{\alpha_i, \beta_i}(Q) \right]^2 \leq \mathbb{E}_{(X, Y) \sim \mathbb{Q}} \left[\left(\frac{d\mathbb{P}}{d\mathbb{Q}}(X, Y) \right)^2 \right] - 1.$$

Proof. Denote $L(X, Y) = \frac{d\mathbb{P}}{d\mathbb{Q}}(X, Y)$. In addition, for $\alpha = (\alpha_1, \dots, \alpha_k)$ and $\beta = (\beta_1, \dots, \beta_k)$ we write

$$\kappa(\alpha; \beta) = \mathbb{E}_{Q \sim \nu} \left[\prod_{1 \leq i \leq k} \Lambda_{\alpha_i, \beta_i}(Q) \right] = \mathbb{E}_{(X, Y) \sim \mathbb{P}} \left[\prod_{1 \leq i \leq k} \mathcal{H}_{\alpha_i}(X) \mathcal{H}_{\beta_i}(Y) \right].$$

Then we have

$$\begin{aligned} & \sum_{\substack{(\alpha_1, \dots, \alpha_k): \alpha_i \in \mathbb{N}^d, |\alpha_i| \leq D \\ (\beta_1, \dots, \beta_k): \beta_i \in \mathbb{N}^m, |\beta_i| \leq D}} \kappa(\alpha; \beta)^2 \stackrel{(3.7)}{=} \mathbb{E}_{(X, Y) \sim \mathbb{P}} \left[\sum_{\substack{(\alpha_1, \dots, \alpha_k): \alpha_i \in \mathbb{N}^d, |\alpha_i| \leq D \\ (\beta_1, \dots, \beta_k): \beta_i \in \mathbb{N}^m, |\beta_i| \leq D}} \kappa(\alpha; \beta) \mathcal{H}_{\alpha_i}(X) \mathcal{H}_{\beta_i}(Y) \right] \\ &= \mathbb{E}_{(X, Y) \sim \mathbb{Q}} \left[\sum_{\substack{(\alpha_1, \dots, \alpha_k): \alpha_i \in \mathbb{N}^d, |\alpha_i| \leq D \\ (\beta_1, \dots, \beta_k): \beta_i \in \mathbb{N}^m, |\beta_i| \leq D}} \kappa(\alpha; \beta) L(X, Y) \mathcal{H}_{\alpha_i}(X) \mathcal{H}_{\beta_i}(Y) \right] \\ &\leq \mathbb{E}_{(X, Y) \sim \mathbb{Q}} \left[L(X, Y)^2 \right]^{\frac{1}{2}} \mathbb{E}_{(X, Y) \sim \mathbb{Q}} \left[\left(\sum_{\substack{(\alpha_1, \dots, \alpha_k): \alpha_i \in \mathbb{N}^d, |\alpha_i| \leq D \\ (\beta_1, \dots, \beta_k): \beta_i \in \mathbb{N}^m, |\beta_i| \leq D}} \kappa(\alpha; \beta) \mathcal{H}_{\alpha_i}(X) \mathcal{H}_{\beta_i}(Y) \right)^2 \right]^{\frac{1}{2}} \\ &= \mathbb{E}_{(X, Y) \sim \mathbb{Q}} \left[L(X, Y)^2 \right]^{\frac{1}{2}} \left(\sum_{\substack{(\alpha_1, \dots, \alpha_k): \alpha_i \in \mathbb{N}^d, |\alpha_i| \leq D \\ (\beta_1, \dots, \beta_k): \beta_i \in \mathbb{N}^m, |\beta_i| \leq D}} \kappa(\alpha; \beta)^2 \right)^{\frac{1}{2}}, \end{aligned}$$

where the last equality follows from (3.5). This yields the desired result since $\kappa(\mathbb{O}_d, \mathbb{O}_m) = 1$. \square

We can now prove (3.15), thus finishing the proof of Theorem 2.2, Item (2).

Proof of (3.15). We will calculate $L(X, Y)$ directly. We divide our proof into two cases.

Case 1: $k \leq m$ and $k^4 = o(d/m)$. Note that given $Y = B$ and $XX^\top = A$, the conditional distribution of X under both \mathbb{P} and \mathbb{Q} is the uniform distribution over

$$\left\{ X \in \mathbb{R}^{k*d} : XX^\top = A \right\}.$$

Thus, we have

$$\mathbf{L}(X, Y) = \mathbf{L}(XX^\top, Y).$$

Note that XX^\top is a $k * k$ Wishart matrix with dimension d , it is well-known that the density of XX^\top is given by (see, e.g., [Wis28] and [BDER16])

$$f_{XX^\top}(A) = \mathbf{1}_{\{A \succeq 0\}} \cdot \omega(d, k) \det(A)^{\frac{1}{2}(d-k-1)} e^{-\frac{1}{2} \text{tr}(A)}, \quad (3.16)$$

where $\omega(\cdot, \cdot)$ is the Wishart constant defined by

$$\frac{1}{\omega(s, t)} = \pi^{t(t-1)/4} 2^{st/2} \prod_{j=1}^t \Gamma\left(\frac{s-j+1}{2}\right). \quad (3.17)$$

In addition, we claim that the conditional distribution of Y given $\{XX^\top = A\}$ is given by

$$Y \mid \{XX^\top = A\} \stackrel{d}{=} A^{\frac{1}{2}} \cdot \mathbf{Z}_{k*m}, \quad (3.18)$$

where \mathbf{Z}_{k*m} is the upper-left $k * m$ submatrix of a uniform $d * d$ orthogonal matrix $\mathbf{\Gamma}_d$. In fact, conditioned on the spectral decomposition of X

$$X = \begin{pmatrix} A^{\frac{1}{2}} & \mathbb{O}_{k*(d-k)} \end{pmatrix} U \text{ where } U \in O(d),$$

we have

$$Y = XQ = \begin{pmatrix} A^{\frac{1}{2}} & \mathbb{O}_{k*(d-k)} \end{pmatrix} UQ$$

and thus (3.18) follows from UQ is uniformly distributed over $O(d)$. Applying Lemma A.5, we see that the conditional density of Y given $\{XX^\top = A\}$ is

$$f(Y \mid XX^\top = A) = \frac{\omega(d-m, k)}{(2\pi)^{km/2} \omega(d, k)} \det(\mathbb{I}_m - Y^\top A^{-1} Y)^{(d-k-m-1)/2} \det(A)^{-m/2} \cdot \chi(A^{-1/2} Y),$$

where $\chi(A)$ is the indicator function that all the eigenvalues of $A^\top A$ lies in $[0, 1]$. In addition, note that under \mathbb{Q} , the density of Y is independent of x and is given by

$$\frac{1}{(2\pi)^{km/2}} e^{-\text{tr}(Y^\top Y)/2}. \quad (3.19)$$

Thus, we have

$$\mathbf{L}(A, Y) = \frac{\omega(d-m, k)}{\omega(d, k)} \cdot e^{\frac{\text{tr}(Y^\top Y)}{2}} \det(\mathbb{I}_m - Y^\top A^{-1} Y)^{\frac{d-k-m-1}{2}} \det(A)^{-m/2} \cdot \chi(A^{-1/2} Y). \quad (3.20)$$

Plugging (3.20) into the formula

$$\mathbb{E}_{\mathbb{Q}}[\mathbf{L}(XX^\top, Y)^2] = \int_{A \succeq 0} \int_Y \mathbf{L}(A, Y)^2 \cdot \frac{e^{-\text{tr}(Y^\top Y)/2}}{(2\pi)^{km/2}} \cdot \omega(d, k) \det(A)^{\frac{1}{2}(d-k-1)} e^{-\frac{1}{2} \text{tr}(A)} dA dY, \quad (3.21)$$

we obtain that $\mathbb{E}_{\mathbb{Q}}[\mathbf{L}(X, Y)^2]$ equals

$$\int_{A \succeq 0} \int_Y \frac{\omega(d-m, k)^2 \det(\mathbb{I}_m - Y^\top A^{-1} Y)^{d-k-m-1} \det(A)^{\frac{1}{2}(d-k-2m-1)} \chi(A^{-1/2} Y)}{\omega(d, k) (2\pi)^{\frac{km}{2}} e^{\frac{1}{2}(\text{tr}(A) - \text{tr}(Y^\top Y))}} dA dY. \quad (3.22)$$

By substituting Y with $A^{1/2}Z$, we get that

$$(3.22) = \int_{A \geq 0} \int_Z \frac{\omega(d-m, k)^2 \det(\mathbb{I}_m - Z^\top Z)^{d-k-m-1} \det(A)^{\frac{1}{2}(d-k-m-1)} \chi(Z)}{\omega(d, k) (2\pi)^{\frac{km}{2}} e^{\frac{1}{2}(\text{tr}(A) - \text{tr}(Z^\top AZ))}} d\text{Ad}Z. \quad (3.23)$$

Note that

$$\text{tr}(A) - \text{tr}(Z^\top AZ) = \text{tr}(A(\mathbb{I}_k - ZZ^\top)),$$

by substituting A with $(\mathbb{I}_k - ZZ^\top)^{-1/2} B (\mathbb{I}_k - ZZ^\top)^{-1/2}$, we get that (note that $\det(\mathbb{I}_m - Z^\top Z) = \det(\mathbb{I}_k - ZZ^\top)$)

$$(3.23) = \int_{B \geq 0} \int_Z \frac{\omega(d-m, k)^2 \det(\mathbb{I}_m - Z^\top Z)^{\frac{1}{2}(d-3k-m-1)} \det(B)^{\frac{1}{2}(d-k-m-1)} \chi(Z)}{\omega(d, k) (2\pi)^{\frac{km}{2}} e^{\frac{1}{2}\text{tr}(B)}} dB dZ. \quad (3.24)$$

Note that

$$\begin{aligned} \int_{B \succ 0} e^{-\frac{1}{2}\text{tr}(B)} \det(B)^{\frac{1}{2}(d-k-m-1)} dB &= \frac{1}{\omega(d-m, k)}, \\ \int_Z \det(\mathbb{I}_m - Z^\top Z)^{\frac{1}{2}(d-3k-m-1)} \chi(Z) dZ &= \frac{(2\pi)^{mk/2} \omega(d-2k, k)}{\omega(d-2k-m, k)}. \end{aligned}$$

We then get that

$$\begin{aligned} (3.24) &= \frac{(2\pi)^{mk/2} \omega(d-2k, k)}{\omega(d-2k-m, k)} \cdot \frac{1}{\omega(d-m, k)} \cdot \frac{\omega(d-m, k)^2}{\omega(d, k) (2\pi)^{km/2}} \\ &= \frac{\omega(d-m, k)}{\omega(d-m-2k, k)} \cdot \frac{\omega(d-2k, k)}{\omega(d, k)}. \end{aligned} \quad (3.25)$$

Applying Lemma A.3, we get that (recall the assumption $k^4 \leq D^4 = o(d/m)$)

$$(3.25) = [1 + o(1)] \cdot \left(\frac{d}{d-m}\right)^{k^2} = [1 + o(1)] \cdot \left(1 + \frac{mk^2}{d}\right) = 1 + o(D^{-1}).$$

Case 2: $m^4 \leq k^4 = o(d/m)$. In this case, again we have

$$\mathbf{L}(X, Y) = \mathbf{L}(XX^\top, Y)$$

and the density of XX^\top is given by (3.16). Using Lemma A.5, we see that the conditional density of Y given $\{XX^\top = A\}$ is

$$f(Y \mid XX^\top = A) = \frac{\omega(d-k, m)}{(2\pi)^{km/2} \omega(d, m)} \det(\mathbb{I}_k - A^{-\frac{1}{2}} Y Y^\top A^{-\frac{1}{2}})^{\frac{d-k-m-1}{2}} \det(A)^{-\frac{m}{2}} \cdot \chi(Y A^{-1/2}).$$

Since the density of Y under \mathbf{Q} is given by (3.19), we get that

$$\mathbf{L}(A, Y) = \frac{\omega(d-k, m)}{\omega(d, m)} \cdot e^{\frac{\text{tr}(Y^\top Y)}{2}} \det(\mathbb{I}_k - A^{-\frac{1}{2}} Y Y^\top A^{-\frac{1}{2}})^{\frac{d-k-m-1}{2}} \det(A)^{-\frac{m}{2}} \cdot \chi(Y A^{-1/2}).$$

Plugging this result into (3.21), we obtain that $\mathbb{E}_{\mathbf{Q}}[\mathbf{L}(X, Y)^2]$ equals

$$\int_{A \succ 0} \int_Y \frac{\omega(d-k, m)^2 \omega(d, k) \det(\mathbb{I}_k - A^{-\frac{1}{2}} Y Y^\top A^{-\frac{1}{2}})^{d-k-m-1} \det(A)^{\frac{1}{2}(d-k-2m-1)} \chi(Y A^{-1/2})}{\omega(d, m)^2 (2\pi)^{\frac{km}{2}} e^{\frac{1}{2}(\text{tr}(A) - \text{tr}(Y^\top Y))}} d\text{Ad}Y. \quad (3.26)$$

Substituting $Y = A^{1/2}Z$, we get that

$$(3.26) = \int_{A \succ 0} \int_Z \frac{\omega(d-k, m)^2 \omega(d, k) \det(\mathbb{I}_k - ZZ^\top)^{d-k-m-1} \det(A)^{\frac{1}{2}(d-k-m-1)} \chi(Z)}{\omega(d, m)^2 (2\pi)^{\frac{km}{2}} e^{\frac{1}{2}(\text{tr}(A) - \text{tr}(AZZ^\top))}} d\text{Ad}Z. \quad (3.27)$$

Substituting $A = (\mathbb{I}_k - ZZ^\top)^{-1/2} B (\mathbb{I}_k - ZZ^\top)^{-1/2}$, we get that

$$(3.27) = \int_{B \succ 0} \int_Z \frac{\omega(d-k, m)^2 \omega(d, k) \det(\mathbb{I}_k - ZZ^\top)^{d-3k-m-1} \det(B)^{\frac{1}{2}(d-k-m-1)} \chi(Z)}{\omega(d, m)^2 (2\pi)^{\frac{km}{2}} e^{\frac{1}{2} \text{tr}(B)}} dB dZ. \quad (3.28)$$

Note that

$$\begin{aligned} \int_{B \succ 0} e^{-\frac{1}{2} \text{tr}(B)} \det(B)^{\frac{1}{2}(d-k-m-1)} dB &= \frac{1}{\omega(d-m, k)}, \\ \int_Z \det(\mathbb{I}_k - Z^\top Z)^{\frac{1}{2}(d-3k-m-1)} \chi(Z) dZ &= \frac{(2\pi)^{mk/2} \omega(d-2k, m)}{\omega(d-3k, m)}. \end{aligned}$$

We then get that

$$\begin{aligned} (3.28) &= \frac{\omega(d-k, m)^2}{\omega(d, m)^2} \cdot \frac{\omega(d, k)}{\omega(d-m, k)} \cdot \frac{\omega(d-2k, m)}{\omega(d-3k, m)} \\ &= [1 + o(1)] \cdot d^{mk} \cdot (d-m)^{-mk/2} \cdot (d-3k)^{-mk/2} \\ &= [1 + o(1)] \cdot \left(\frac{d^2}{(d-m)(d-3k)} \right)^{mk/2} = 1 + o(mk^2/d) = 1 + o(D^{-1}), \end{aligned}$$

where in the second equality we use Lemma A.3 and in the last inequality we use the assumption $m^4 \leq k^4 = o(d/m)$. Combining the two cases leads to the desired result. \square

3.4 Proof of Item (2)

In this subsection we discuss the case that $m = d, \sigma = \omega(1)$ and $D = o(\sigma)$. Still, based on (3.8), it suffices to show that for $1 \leq k \leq D$ we have

$$\sum_{\substack{(\alpha_1, \dots, \alpha_k): \alpha_i \in \mathbb{N}^d \\ (\beta_1, \dots, \beta_k): \beta_i \in \mathbb{N}^m \\ 0 < |\alpha_i|, |\beta_j| \leq D}} \mathbb{E}_{Q \sim \nu} \left[\prod_{1 \leq i \leq k} \Lambda_{\alpha_i, \beta_i}(Q) \right]^2 = o(D^{-1}). \quad (3.29)$$

We will adopt the similar approach as in Subsection 3.3 by relating the left hand side of (3.29) to a simpler hypothesis testing problem. Consider the following hypothesis testing problem, where our observation are two vectors $X = (X_1, \dots, X_k)^\top \in \mathbb{R}^{k*d}, Y = (Y_1, \dots, Y_k)^\top \in \mathbb{R}^{k*d}$ such that:

- under the null hypothesis $\overline{\mathcal{H}}_0$, we sample $X_i \sim \mathcal{N}(0, \mathbb{I}_d)$ and $Y_i \sim \mathcal{N}(0, \mathbb{I}_d)$ independently.
- under the alternative hypothesis $\overline{\mathcal{H}}_1$, we first sample $X_1, \dots, X_k \sim \mathcal{N}(0, \mathbb{I}_d)$ and $Q \sim \nu$ independently (recall that now ν is the Haar measure over $O(d)$), and then define

$$Y = \frac{1}{\sqrt{1+\sigma^2}} (XQ + \sigma Z).$$

In addition, we define \mathbf{P} to be the law of (X, Y) under $\overline{\mathcal{H}}_1$ and let \mathbf{Q} to be the law of (X, Y) under $\overline{\mathcal{H}}_0$, respectively. Similar as in Lemma 3.7, we get that

$$\sum_{\substack{(\alpha_1, \dots, \alpha_k): \alpha_i \in \mathbb{N}^d \\ (\beta_1, \dots, \beta_k): \beta_i \in \mathbb{N}^m \\ 0 < |\alpha_i|, |\beta_j| \leq D}} \mathbb{E}_{Q \sim \nu} \left[\prod_{1 \leq i \leq k} \Lambda_{\alpha_i, \beta_i}(Q) \right]^2 \leq \mathbb{E}_{(X, Y) \sim \mathbf{Q}} \left[\left(\frac{d\mathbf{P}}{d\mathbf{Q}}(X, Y) \right)^2 \right] - 1. \quad (3.30)$$

The rest part of this subsection is devoted to the direct calculation of the right hand side of (3.30). Clearly, the probability density function of (X, Y) under \mathbf{Q} is given by

$$f_{\mathbf{Q}}(X, Y) = \frac{e^{-\|X\|_{\mathbb{F}}^2/2}}{(2\pi)^{kd/2}} \cdot \frac{e^{-\|Y\|_{\mathbb{F}}^2/2}}{(2\pi)^{kd/2}}.$$

In addition, the probability density function of (X, Y) under \mathbf{P} is given by

$$f_{\mathbf{P}}(X, Y) = \frac{e^{-\|X\|_{\mathbb{F}}^2/2}}{(2\pi)^{kd/2}} \cdot \int_{O(d)} \frac{e^{-\frac{\sigma^2}{2(1+\sigma^2)} \left\| Y - \frac{XQ}{\sqrt{1+\sigma^2}} \right\|_{\mathbb{F}}^2}}{(2\pi\sigma^2/(1+\sigma^2))^{kd}} dQ.$$

Thus, we get that

$$\mathbf{L}(X, Y) = \left(\frac{1+\sigma^2}{\sigma^2} \right)^{kd/2} e^{-(\|X\|_{\mathbb{F}}^2 + \|Y\|_{\mathbb{F}}^2)/2\sigma^2} \cdot \int_{O(d)} e^{\sqrt{1+\sigma^2} \langle X, YQ \rangle / \sigma^2} dQ.$$

Thus, we have

$$\mathbf{L}(X, Y)^2 = \left(\frac{1+\sigma^2}{\sigma^2} \right)^{kd} e^{-(\|X\|_{\mathbb{F}}^2 + \|Y\|_{\mathbb{F}}^2)/\sigma^2} \cdot \int_{O(d) \times O(d)} e^{\sqrt{1+\sigma^2} \langle X, Y(Q_1+Q_2) \rangle / \sigma^2} dQ_1 dQ_2.$$

We then have

$$\begin{aligned} & \mathbb{E}_{(X, Y) \sim \mathbf{Q}} \left[\mathbf{L}(X, Y)^2 \right] \\ &= \left(\frac{1+\sigma^2}{\sigma^2} \right)^{kd} \int_{O(d) \times O(d)} \mathbb{E}_{(X, Y) \sim \mathbf{Q}} \left[e^{-(\|X\|_{\mathbb{F}}^2 + \|Y\|_{\mathbb{F}}^2)/\sigma^2} \cdot e^{\sqrt{1+\sigma^2} \langle X, Y(Q_1+Q_2) \rangle / \sigma^2} \right] dQ_1 dQ_2 \\ &= \left(\frac{1+\sigma^2}{\sigma^2} \right)^{kd} \left(\frac{\sigma^2}{2+\sigma^2} \right)^{kd/2} \int_{O(d) \times O(d)} \mathbb{E}_{Y \sim \mathbf{Q}} \left[e^{-\|Y\|_{\mathbb{F}}^2/\sigma^2} \cdot e^{(1+\sigma^2) \|Y(Q_1+Q_2)\|_{\mathbb{F}}^2/2\sigma^2(2+\sigma^2)} \right] dQ_1 dQ_2 \\ &= \left(\frac{1+\sigma^2}{\sigma^2} \right)^{kd} \left(\frac{\sigma^2}{2+\sigma^2} \right)^{kd/2} \int_{O(d)} \mathbb{E}_{Y \sim \mathbf{Q}} \left[e^{-\|Y\|_{\mathbb{F}}^2/\sigma^2} \cdot e^{(1+\sigma^2) (\|Y\|_{\mathbb{F}}^2 + \langle Y, YQ \rangle) / \sigma^2(2+\sigma^2)} \right] dQ \\ &= \left(\frac{1+\sigma^2}{\sigma^2} \right)^{kd} \left(\frac{\sigma^2}{2+\sigma^2} \right)^{kd/2} \int_{O(d)} \mathbb{E}_{Y \sim \mathbf{Q}} \left[e^{-\|Y\|_{\mathbb{F}}^2/\sigma^2(\sigma^2+2)} \cdot e^{(1+\sigma^2) \langle Y, YQ \rangle / \sigma^2(2+\sigma^2)} \right] dQ, \end{aligned} \quad (3.31)$$

where the second equality follows from Lemma A.6 and the third equality follows from the fact that $Q_1 Q_2^\top \sim \nu$ for $Q_1 \perp Q_2$ and $Q_1, Q_2 \sim \nu$. Using Lemma A.7 with

$$A = A(Q) = \frac{1}{\sigma^2(\sigma^2+2)} \mathbb{I}_d - \frac{\sigma^2+1}{2\sigma^2(\sigma^2+2)} (Q + Q^\top),$$

we get that

$$(3.31) = \left(\frac{1+\sigma^2}{\sigma^2} \right)^{kd} \left(\frac{\sigma^2}{2+\sigma^2} \right)^{kd/2} \int_{O(d)} \det(\mathbb{I}_d + 2A(Q))^{-k/2} dQ. \quad (3.32)$$

Note that (denote $\epsilon = \epsilon(\sigma) = (1 + \sigma^2)^{-1} = o(1)$)

$$\begin{aligned} \det(\mathbb{I}_d + 2A(Q)) &= \det\left(\frac{1+(\sigma^2+1)^2}{\sigma^2(\sigma^2+2)}\mathbb{I}_d - \frac{\sigma^2+1}{\sigma^2(\sigma^2+2)}(Q + Q^\top)\right) \\ &= \left(\frac{(\sigma^2+1)^2}{\sigma^2(\sigma^2+2)}\right)^d \cdot \det\left((1 + \epsilon^2)\mathbb{I}_d - \epsilon(Q + Q^\top)\right) \\ &= \left(\frac{(\sigma^2+1)^2}{\sigma^2(\sigma^2+2)}\right)^d \cdot \det\left((\mathbb{I}_d + \epsilon Q)(\mathbb{I}_d + \epsilon Q^\top)\right) = \frac{(\sigma^2+1)^2}{\sigma^2(\sigma^2+2)} \cdot \det(\mathbb{I}_d + \epsilon Q)^2. \end{aligned}$$

Thus, we have

$$(3.32) = \int_{O(d)} \det\left(\mathbb{I}_d + \frac{1}{1+\sigma^2}Q\right)^{-k} dQ = 1 + O\left(\frac{k}{1+\sigma^2}\right) = 1 + o(D^{-1}), \quad (3.33)$$

where the second equality follows from Lemma A.8 and the last equality follows from $k \leq D$ and $D = o(\sigma)$. This leads to (3.29).

3.5 Proof of Item (3)

In this subsection we prove Item (3) in Theorem 2.2. Denote

$$f(X, Y) = \left(\|Y\|_{\mathbb{F}}^2 - \|X\|_{\mathbb{F}}^2\right)^2. \quad (3.34)$$

It suffices to show the following estimation.

Lemma 3.8. *Assuming $d = m$ and $\sigma = o(1)$, then we have*

- (1) $\mathbb{E}_{\mathbb{P}}[f] = o(nd)$ and $\mathbb{E}_{\mathbb{Q}}[f] = \Theta(nd)$.
- (2) $\text{Var}_{\mathbb{P}}[f], \text{Var}_{\mathbb{Q}}[f] = o(n^2d^2)$.

Proof. Note that under \mathbb{P} we have $Y = (XQ + \sigma Z)/\sqrt{1 + \sigma^2}$, thus

$$\|Y\|_{\mathbb{F}}^2 - \|X\|_{\mathbb{F}}^2 = \frac{1}{1+\sigma^2}\|XQ + \sigma Z\|_{\mathbb{F}}^2 - \|X\|_{\mathbb{F}}^2 = \frac{\sigma^2}{1+\sigma^2}(\|Z\|_{\mathbb{F}}^2 - \|X\|_{\mathbb{F}}^2) + \frac{2\sigma}{1+\sigma^2}\langle X, Z \rangle.$$

Thus, we have

$$\begin{aligned} \mathbb{E}_{\mathbb{P}}[f^2] &= \mathbb{E}_{\mathbb{P}}\left[\left(\|Y\|_{\mathbb{F}}^2 - \|X\|_{\mathbb{F}}^2\right)^4\right] \leq 2^4 \cdot \left(\mathbb{E}_{\mathbb{P}}\left[\sigma^4(\|Z\|_{\mathbb{F}}^2 - \|X\|_{\mathbb{F}}^2)^4\right] + \mathbb{E}_{\mathbb{P}}\left[\sigma^2\langle X, Z \rangle^4\right]\right) \\ &\leq O(1) \cdot \sigma^2 \cdot n^2d^2 = o(n^2d^2), \end{aligned} \quad (3.35)$$

where the second inequality follows from $\|Z\|_{\mathbb{F}}^2 - \|X\|_{\mathbb{F}}^2$ and $\langle X, Z \rangle$ are sums of nd independent random variables with finite fourth moment. In addition, under \mathbb{Q} we have X, Y are independent $n * d$ Gaussian random matrices with i.i.d. $\mathcal{N}(0, 1)$ entries. Thus, we have

$$\mathbb{E}_{\mathbb{Q}}[f] = \mathbb{E}_{\mathbb{Q}}\left[\left(\sum_{1 \leq i \leq n} \sum_{1 \leq j \leq d} (Y_{i,j}^2 - X_{i,j}^2)\right)^2\right] = 4nd, \quad (3.36)$$

$$\mathbb{E}_{\mathbb{Q}}[f^2] = \mathbb{E}_{\mathbb{Q}}\left[\left(\sum_{1 \leq i \leq n} \sum_{1 \leq j \leq d} (Y_{i,j}^2 - X_{i,j}^2)\right)^4\right] = 16n^2d^2 + O(nd). \quad (3.37)$$

Combining (3.35), (3.36) and (3.37) yields the desired result. \square

Acknowledgment. The author thanks Jian Ding for helpful discussions. The author thanks Shuyang Gong for helpful discussions on the forthcoming manuscript [GWX25+]. The author thanks Hang Du for pointing him to the reference [MM13]. This work is partially supported by National Key R&D program of China (No. 2023YFA1010103) and NSFC Key Program (Project No. 12231002).

A Preliminary results

A.1 Preliminary results on Hermite polynomials

Lemma A.1. For $x, y \in \mathbb{R}^d$ with $\|y\| = 1$ and $m \in \mathbb{N}$, we have

$$\mathcal{H}_m(\langle x, y \rangle) = \sum_{\alpha \in \mathbb{N}^d: |\alpha|=m} \sqrt{\frac{\alpha!}{m!}} \binom{m}{\alpha} y^\alpha \mathcal{H}_\alpha(x). \quad (\text{A.1})$$

Proof. Recall Definition 3.2. It was shown in [MOS13] that

$$e^{tx - \frac{t^2}{2}} = \sum_{n=0}^{\infty} \frac{H_n(x)}{n!} \cdot t^n.$$

Thus, we see that

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{H_n(\langle x, y \rangle)}{n!} \cdot t^n &= e^{\langle x, y \rangle t - \frac{t^2}{2}} = e^{\langle x, y \rangle t - \frac{\|y\|^2 t^2}{2}} = \prod_{i=1}^d e^{x_i y_i t - \frac{y_i^2 t^2}{2}} \\ &= \prod_{i=1}^d \left(\sum_{n=0}^{\infty} \frac{H_n(x_i)}{n!} \cdot (y_i t)^n \right) = \sum_{n=0}^{\infty} t^n \sum_{\alpha \in \mathbb{N}^d: |\alpha|=n} \frac{y^\alpha H_\alpha(x)}{\alpha!}. \end{aligned}$$

Thus, we see that

$$H_n(\langle x, y \rangle) = \sum_{\alpha \in \mathbb{N}^d: |\alpha|=n} \binom{n}{\alpha} y^\alpha H_\alpha(x).$$

Combined with Definition 3.2, we get the desired result. \square

A.2 Preliminary results on Wishart constant

The following result can be found in [Gau59].

Lemma A.2 (Gautschi's inequality). for all $x \in \mathbb{R}_{\geq 0}$ we have $x^{1-s} \leq \frac{\Gamma(x+1)}{\Gamma(x+s)} \leq (x+1)^{1-s}$. In particular, we have when $x \rightarrow \infty$

$$\frac{\Gamma(x + \frac{1}{2})}{\Gamma(x)} = [1 + O(\frac{1}{x})] \cdot \sqrt{x}.$$

Based on Lemma A.2, the following result provides a crude bound on $\omega(s, t)$.

Lemma A.3. Recall the definition of $\omega(s, t)$ in (3.17). For $k^3 = o(d)$ we have

$$\frac{\omega(d, k)}{\omega(d-2k, k)} = [1 + o(1)] \cdot d^{k^2} \quad \text{and} \quad \frac{\omega(d-2k, k)}{\omega(d, k)} = [1 + o(1)] \cdot d^{-k^2}.$$

In addition, for $k^3, m^3 = o(d)$ we have

$$\frac{\omega(d, m)}{\omega(d-k, m)} = [1 + o(1)] \cdot d^{km/2} \quad \text{and} \quad \frac{\omega(d-k, m)}{\omega(d, m)} = [1 + o(1)] \cdot d^{-km/2}.$$

Proof. Using (3.17), we get that

$$\begin{aligned}\frac{\omega(d, k)}{\omega(d - 2k, k)} &= 2^{k^2} \prod_{j=1}^k \frac{\Gamma(\frac{d-j+1}{2})}{\Gamma(\frac{d-2k-j+1}{2})} = 2^{k^2} \prod_{j=1}^k \prod_{i=1}^k \left(\frac{d-j+1}{2} - i \right) \\ &= \prod_{j=1}^k \prod_{i=1}^k \left([1 + O(k/d)] \cdot d \right) = [1 + o(1)] \cdot d^{k^2},\end{aligned}$$

where in the last equality we use $k^3 = o(d)$. Similarly we can get that

$$\frac{\omega(d - 2k, k)}{\omega(d, k)} = [1 + o(1)] \cdot d^{-k^2}.$$

In addition, using (3.17) again we have

$$\begin{aligned}\frac{\omega(d, m)}{\omega(d - k, m)} &= 2^{km/2} \cdot \prod_{j=1}^m \frac{\Gamma(\frac{d-j+1}{2})}{\Gamma(\frac{d-k-j+1}{2})} = 2^{km/2} \prod_{j=1}^m \prod_{i=1}^k \frac{\Gamma(\frac{d-i-j+2}{2})}{\Gamma(\frac{d-i-j+1}{2})} \\ &= 2^{km/2} \prod_{j=1}^m \prod_{i=1}^k \left(\left(1 + O\left(\frac{2}{d-i-j+1}\right) \right) \cdot \sqrt{(d-i-j+1)/2} \right) = [1 + o(1)] \cdot d^{km/2},\end{aligned}$$

where in the last equality we use $k^3, m^3 = o(d)$. Similarly we can get that

$$\frac{\omega(d - k, m)}{\omega(d, m)} = [1 + o(1)] \cdot d^{-km/2}. \quad \square$$

A.3 Preliminary results in probability

Lemma A.4. *For all $\{\gamma_i \in \mathbb{N} : 1 \leq i \leq d\}$ we have*

$$\mathbb{E}_{Q \sim \text{Unif}(\mathbb{S}^{d-1})} \left[Q_1^{\gamma_1} \dots Q_d^{\gamma_d} \right] = \mathbf{1}_{\{\gamma_1, \dots, \gamma_d \text{ is even}\}} \cdot \frac{\Gamma(\frac{d}{2})(\gamma_1-1)!\dots(\gamma_d-1)!}{\Gamma(\frac{d+\gamma_1+\dots+\gamma_d}{2})2^{\frac{\gamma_1+\dots+\gamma_d}{2}}}.$$

Proof. Denote $X = (X_1, \dots, X_d) \sim \mathcal{N}(0, \mathbb{I}_d)$. We the have $\|X\| \perp \frac{X}{\|X\|}$ and $\frac{X}{\|X\|} \sim \text{Unif}(\mathbb{S}^{d-1})$. Thus, we have

$$\begin{aligned}& \mathbb{E}_{Q \sim \text{Unif}(\mathbb{S}^{d-1})} \left[Q_1^{\gamma_1} \dots Q_d^{\gamma_d} \right] \cdot \mathbb{E} \left[\|X\|^{\gamma_1+\dots+\gamma_d} \right] \\ &= \mathbb{E} \left[\left(\frac{X_1}{\|X\|} \right)^{\gamma_1} \dots \left(\frac{X_d}{\|X\|} \right)^{\gamma_d} \right] \cdot \mathbb{E} \left[\|X\|^{\gamma_1+\dots+\gamma_d} \right] \\ &= \mathbb{E} \left[X_1^{\gamma_1} \dots X_d^{\gamma_d} \right] = \mathbf{1}_{\{\gamma_1, \dots, \gamma_d \text{ is even}\}} \cdot \prod_{1 \leq i \leq d} (\gamma_i - 1)!!.\end{aligned}$$

In addition, since the law of $\|X\|^2$ is a Gamma distribution with d sample and parameter $\frac{1}{2}$, we have

$$\mathbb{E} \left[\|X\|^{\gamma_1+\dots+\gamma_d} \right] = \frac{2^{\frac{\gamma_1+\dots+\gamma_d}{2}} \Gamma(\frac{d+\gamma_1+\dots+\gamma_d}{2})}{\Gamma(\frac{d}{2})}.$$

Thus, we see that the desired result holds. \square

Lemma A.5 ([JM19], Lemma 2.1). Let Γ_d be an $d \times d$ random matrix which is uniformly distributed on the orthogonal group $O(n)$ and let \mathbf{Z}_n be the upper-left $p \times q$ submatrix of Γ_n . If $p + q \leq n$ and $p \geq q$ then the joint density function of entries of \mathbf{Z}_n is given by

$$f(Z) = \frac{\omega(d-p, q)}{\omega(d, q)(2\pi)^{pq/2}} \cdot \text{Det}(\mathbb{I}_q - Z^\top Z)^{(n-p-q-1)/2} \chi_0(Z^\top Z),$$

where $\chi_0(Z^\top Z)$ is the indicator function of the set that all q eigenvalues of $Z^\top Z$ are in $[0, 1]$. When $p < q$, the density of \mathbf{Z}_n is obtained by interchanging p and q in the above Wishart constant.

Lemma A.6. Let Z be an $d \times m$ matrix with entries are i.i.d. $\mathcal{N}(0, 1)$ random variables. Then for all $\lambda > 0$ and $A \in \mathbb{R}^{d \times m}$ we have

$$\mathbb{E} \left[e^{-\lambda \|Z\|_{\mathbb{F}}^2 + \langle A, Z \rangle} \right] = (1 + 2\lambda)^{-dm/2} \cdot e^{\|A\|_{\mathbb{F}}^2 / (2(1+2\lambda))}.$$

Proof. Note that

$$\mathbb{E}_{X \sim \mathcal{N}(0,1)} \left[e^{-\lambda x^2 + ax} \right] = \frac{1}{\sqrt{1+2\lambda}} e^{\frac{a^2}{2(1+2\lambda)}},$$

we then have

$$\mathbb{E} \left[e^{-\lambda \|Z\|_{\mathbb{F}}^2 + \langle A, Z \rangle} \right] = \prod_{\substack{1 \leq i \leq d \\ 1 \leq j \leq m}} \frac{1}{\sqrt{1+2\lambda}} e^{\frac{a_{i,j}^2}{2(1+2\lambda)}} = (1 + 2\lambda)^{-dm/2} \cdot e^{\|A\|_{\mathbb{F}}^2 / (2(1+2\lambda))}. \quad \square$$

Lemma A.7. Let $k < d$ and let Z be an $d \times k$ matrix with entries are i.i.d. $\mathcal{N}(0, 1)$ random variables. Then for any $d \times d$ semidefinite matrix A we have

$$\mathbb{E} \left[e^{-\text{tr}(Z^\top AZ)} \right] = \det(\mathbb{I}_d + 2A)^{-k/2}.$$

Proof. Note that for all $U \in O(d)$ we have

$$\mathbb{E} \left[e^{-\text{tr}(Z^\top AZ)} \right] = \mathbb{E} \left[e^{-\text{tr}(Z^\top U^\top AUZ)} \right].$$

Thus, without losing of generality we may assume that $A = \text{Diag}(a_1, \dots, a_d)$ is a diagonal matrix. In this case, we get from direct calculation that

$$\mathbb{E} \left[e^{-\text{tr}(Z^\top AZ)} \right] = \prod_{1 \leq i \leq d} \mathbb{E} \left[e^{-a_i \sum_{j=1}^k Z_{i,j}^2} \right] = \prod_{1 \leq i \leq d} (1 + 2a_i)^{-k/2} = \det(\mathbb{I}_d + 2A)^{-k/2}. \quad \square$$

Lemma A.8. Denote $\nu = \nu_d$ to be the Haar measure over $O(d)$. For all $\epsilon k = O(1)$ and $0 \leq \epsilon \leq 1 - \Omega(1)$, we have as $d \rightarrow \infty$

$$\mathbb{E}_{Q \sim \nu} \left[\det(\mathbb{I}_d + \epsilon Q)^k \right] = 1 + O(\epsilon k).$$

Proof. Clearly it suffices to show that when $\epsilon k = \Theta(1)$ we have $\mathbb{E}_{Q \sim \nu} \left[\det(\mathbb{I}_d + \epsilon Q)^k \right] = +O(1)$, and the general cases follows from the simple inequality $\mathbb{E}[X] \leq \mathbb{E}[X^m]^{\frac{1}{m}}$ for $m \geq 1$ and $X \geq 0$. Denote $\{e^{i\theta_1}, \dots, e^{i\theta_d} : \theta_1, \dots, \theta_d \in [0, 2\pi]\}$ to be the spectrum of Q . In addition, denote \bar{F}_d to be the empirical distribution of $\theta_1, \dots, \theta_d$. We then have

$$\det(\mathbb{I}_d + \epsilon Q)^k = \exp \left(k \sum_{\ell=1}^d \log(1 + \epsilon e^{i\theta_\ell}) \right) = \exp(d \cdot W_d),$$

where

$$W_d = \int_0^{2\pi} k \log(1 + \epsilon e^{it}) d\overline{F}_d(t).$$

Denote $\varphi(t) = k \log(1 + \epsilon e^{it})$. Note that when $\epsilon k = O(1)$ and $0 \leq \epsilon \leq 1 - \Omega(1)$ we have $|\varphi'(t)| \leq \frac{\epsilon k}{1-\epsilon} = \Theta(1)$ and thus φ is 1-Lipchitz. It was well known (see, e.g., [AGZ10, Theorem 4.4.27] and [MM13, Theorem 2.1]) that \overline{F}_d weakly converges to F , the uniform distribution over $[0, 2\pi]$. In addition, we have the following bound for the tail probability of W_d :

$$\mathbb{P}\left(|W_d| \geq t\right) = \mathbb{P}\left(\left|W_d - k \int_0^{2\pi} \log(1 + \epsilon e^{it}) dF_d(t)\right| \geq t\right) \leq 2e^{-\Theta(1) \cdot d^2 t^2},$$

where the equality follows from

$$\int_0^{2\pi} \log(1 + \epsilon e^{it}) dF_d(t) = 0 \text{ for all } \epsilon \in [0, 1]$$

and the inequality follows from [MM13, Corollary 2.4]. Thus, we have

$$\mathbb{E}_{Q \sim \nu} \left[\det(\mathbb{I}_d + \epsilon Q)^k \right] \leq 2 \int_0^\infty d e^{dt} \cdot e^{-\Theta(1) \cdot d^2 t^2} dt = e^{O(1)} = O(1). \quad \square$$

References

- [APZ17+] Abubakar Abid, Ada Poon, and James Zou. Linear regression with shuffled labels. arXiv preprint, [arXiv:1705.01342](https://arxiv.org/abs/1705.01342).
- [AGZ10] Greg W. Anderson, Alice Guionnet, and Ofer Zeitouni. An introduction to random matrices. Cambridge Studies in Advanced Mathematics, volume 118. Cambridge University Press, 2010.
- [AHSS17] Alexandr Andoni, Daniel Hsu, Kevin Shi, and Xiaorui Sun. Correspondence retrieval. In *Proceedings of the 30th Conference on Learning Theory (COLT)*, pages 105–126. PMLR, 2017.
- [BKW20] Afonso S. Bandeira, Dmitriy Kunisky, and Alexander S. Wein. Computational hardness of certifying bounds on constrained PCA problems. In *11th Innovations in Theoretical Computer Science Conference (ITCS)*, Schloss Dagstuhl-Leibniz-Zentrum für Informatik, 2020.
- [BAH+22] Afonso S. Bandeira, Ahmed E. Alaoui, Samuel B. Hopkins, Tselil Schramm, Alexander S. Wein, and Ilias Zadik. The Franz-Parisi criterion and computational trade-offs in high dimensional statistics. In *Advances in Neural Information Processing Systems (NIPS)*, volume 35, pages 33831–33844. Curran Associates, Inc., 2022.
- [BHK+19] Boaz Barak, Samuel B. Hopkins, Jonathan Kelner, Pravesh K. Kothari, Ankur Moitra, and Aaron Potechin. A nearly tight sum-of-squares lower bound for the planted clique problem. In *SIAM Journal on Computing*, 48(2):687–735, 2019.
- [BBH+21] Matthew Brennan, Guy Bresler, Sam Hopkins, Jerry Li, and Tselil Schramm. Statistical query algorithms and low-degree tests are almost equivalent. In *Proceedings of the 34th Conference on Learning Theory (COLT)*, pages 774–774. PMLR, 2021.
- [BH22] Guy Bresler and Brice Huang. The algorithmic phase transition of random k -SAT for low degree polynomials. In *IEEE 62nd Annual Symposium on Foundations of Computer Science (FOCS)*, pages 298–309. IEEE, 2022.
- [BRST21] Joan Bruna, Oded Regev, Min Jae Song, and Yi Tang. Continuous LWE. In *Proceedings of the 53rd Annual ACM Symposium on Theory of Computing (STOC)*, pages 694–707. ACM, 2021.

- [BDER16] Sébastien Bubeck, Jian Ding, Ronen Eldan, and Miklós Z. Rácz. Testing for high-dimensional geometry in random graphs. In *Random Structures and Algorithms*, 49(3):503–532, 2016.
- [CDGL24+] Guanyi Chen, Jian Ding, Shuyang Gong, and Zhangsong Li. A computational transition for detecting correlated stochastic block models by low-degree polynomials. arXiv preprint, [arXiv:2409.00966](https://arxiv.org/abs/2409.00966).
- [DCK20] Osman E. Dai, Daniel Cullina, and Negar Kiyavash. Achievability of nearly-exact alignment for correlated Gaussian databases. In *Proceedings of the IEEE International Symposium on Information Theory (ISIT)*, pages 1230–1235. IEEE, 2020.
- [DMW23+] Abhishek Dhawan, Cheng Mao, and Alexander S. Wein. Detection of dense subhypergraphs by low-degree polynomials. to appear in *Random Structures and Algorithms*.
- [DKPZ23] Ilias Diakonikolas, Daniel M. Kane, Thanasis Pittas, and Nikos Zarifis. SQ lower bounds for learning mixtures of separated and bounded covariance Gaussians. In *Proceedings of the 36th Annual Conference on Learning Theory (COLT)*, pages 2319–2349. PMLR, 2023.
- [DDL23+] Jian Ding, Hang Du, and Zhangsong Li. Low-degree hardness of detection for correlated Erdős-Rényi Graphs. to appear in *Annals of Statistics*.
- [DKW+22] Yunzi Ding, Dmitriy Kunisky, Alexander S. Wein, and Afonso S. Bandeira. Subexponential-time algorithms for sparse PCA. In *Foundations of Computational Mathematics*, 22(1):1–50, 2022.
- [DL17] Nadav Dym and Yaron Lipman. Exact recovery with symmetries for Procrustes matching. In *SIAM Journal on Optimization*, 27(3):1513–1530, 2017.
- [EGMM24] Mathieu Even, Luca Ganassali, Jakob Maier, and Laurent Massoulié. Aligning embeddings and geometric random graphs: Informational results and computational approaches for the Procrustes-Wasserstein problem. In *Advances in Neural Information Processing Systems (NIPS)*, volume 37, pages 70730–70764. Curran Associates, Inc., 2024.
- [Gam21] David Gamarnik. The overlap gap property: A topological barrier to optimizing over random structures. In *Proceedings of the National Academy of Sciences of the United States of America*, 118(41):e2108492118, 2021.
- [GJW20] David Gamarnik, Aukosh Jagannath and Alexander S. Wein. Low-degree hardness of random optimization problems. In *IEEE 61st Annual Symposium on Foundations of Computer Science (FOCS)*, pages 131–140. IEEE, 2020.
- [Gau59] Walter Gautschi. Some elementary inequalities relating to the Gamma and incomplete Gamma function. In *Journal of Mathematics and Physics*, 38:77–81, 1959.
- [GJJ+20] Mrinalkanti Ghosh, Fernando G. Jeronimo, Chris Jones, Aaron Potechin, and Goutham Rajendran. Sum-of-squares lower bounds for Sherrington-Kirkpatrick via planted affine planes. In *IEEE 61st Annual Symposium on Foundations of Computer Science (FOCS)*, pages 954–965. IEEE, 2020.
- [GL24+] Shuyang Gong and Zhangsong Li. The Umeyama algorithm for matching correlated Gaussian geometric models in the low-dimensional regime. arXiv preprint, [arXiv:2402.15095](https://arxiv.org/abs/2402.15095).
- [GWX25+] Shuyang Gong, Yihong Wu, and Jiaming Xu. The broken-sample problem revisited II: Detecting hidden linear dependencies. Forthcoming, 2025.
- [GJB19] Edouard Grave, Armand Joulin, and Quentin Berthet. Unsupervised alignment of embeddings with Wasserstein Procrustes. In *Proceedings of the 22nd International Conference on Artificial Intelligence and Statistics (AISTATS)*, pages 1880–1890. PMLR, 2019.
- [Hop18] Samuel B. Hopkins. Statistical inference and the sum of squares method. PhD thesis, Cornell University, 2018.
- [HKP+17] Samuel B. Hopkins, Pravesh K. Kothari, Aaron Potechin, Prasad Raghavendra, Tselil Schramm, and David Steurer. The power of sum-of-squares for detecting hidden structures. In *IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 720–731. IEEE, 2017.

- [HS17] Samuel B. Hopkins and David Steurer. Efficient Bayesian estimation from few samples: community detection and related problems. In *IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 379–390. IEEE, 2017.
- [HSS17] Daniel J. Hsu, Kevin Shi, and Xiaorui Sun. Linear regression without correspondence. In *Advances in Neural Information Processing Systems (NIPS)*, volume 30, pages 1531–1540. Curran Associates, Inc., 2017.
- [JM19] Tiefeng Jiang and Yutao Ma. Distances between random orthogonal matrices and independent normals. In *Transactions of the American Mathematical Society*, 372(3):1509–1553, 2019.
- [JWX25+] Simiao Jiao, Yihong Wu, and Jiaming Xu. The broken sample problem revisited: Proof of a conjecture by Bai-Hsing and high-dimensional extensions. arXiv preprint, [arXiv:2503.14619](https://arxiv.org/abs/2503.14619).
- [JPRX23] Chris Jones, Aaron Potetchin, Goutham Rajendran, and Jeff Xu. Sum-of-squares lower bounds for densest k -subgraph. In *Proceedings of the 55th Annual ACM Symposium on Theory of Computing (STOC)*, pages 84–95. ACM, 2023.
- [KMW24] Dmitriy Kunisky, Cristopher Moore, and Alexander S. Wein. Tensor cumulants for statistical inference on invariant distributions. In *IEEE 65th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 1007–1026. IEEE, 2024.
- [KN22] Dmitriy Kunisky and Jonathan Niles-Weed. Strong recovery of geometric planted matchings. In *Proceedings of the 33th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 834–876. SIAM, 2022.
- [KWB22] Dmitriy Kunisky, Alexander S. Wein, and Afonso S. Bandeira. Notes on computational hardness of hypothesis testing: Predictions using the low-degree likelihood ratio. In *Mathematical Analysis, its Applications and Computation: ISAAC*, pages 1–50. Springer, 2022.
- [LWX24] Leon Lufkin, Yihong Wu, and Jiaming Xu. Sharp information-theoretic thresholds for shuffled linear regression. In *Proceedings of the IEEE International Symposium on Information Theory (ISIT)*, pages 363–367. IEEE, 2024.
- [MOS13] Wilhelm Magnus, Fritz Oberhettinger, and Raj Pal Soni. Formulas and theorems for the special functions of mathematical physics. Springer Science & Business Media, 2013.
- [MDK+16] Haggai Maron, Nadav Dym, Itay Kezurer, Shahar Kovalsky, and Yaron Lipman. Point registration via efficient convex relaxation. In *ACM Transactions on Graphics*, 35(4):1–12, 2016.
- [MW23] Rahul Mazumder and Haoyue Wang. Linear regression with partially mismatched data: Local search with theoretical guarantees. In *Mathematical Programming*, 197(2):1265–1303, 2023.
- [MM13] Elizabeth S. Meckes and Mark W. Meckes. Concentration and convergence rates for spectral measures of random matrices. In *Probability Theory and Related Fields*, 156(1-2):145–164, 2013.
- [MR09] Daniele Micciancio and Oded Regev. Lattice-based cryptography. Springer Berlin Heidelberg, 2009.
- [MW25] Andrea Montanari and Alexander S. Wein. Equivalence of approximate message passing and low-degree polynomials in rank-one matrix estimation. In *Probability Theory and Related Fields*, 191(1-2):181–233, 2025.
- [PWC17] Ashwin Pananjady, Martin J. Wainwright, and Thomas A. Courtade. Linear regression with shuffled data: Statistical and computational limits of permutation recovery. In *IEEE Transactions on Information Theory*, 64(5):3286–3300, 2017.
- [RSS19] Prasad Raghavendra, Tselil Schramm, and David Steurer. High-dimensional estimation via sum-of-squares proofs. In *Proceedings of the International Congress of Mathematicians (ICM 2018)*, pages 3389–3423, 2019.
- [RCB97] Anand Rangarajan, Haili Chui, and Fred L. Bookstein. The softassign Procrustes matching algorithm. In *Biennial International Conference on Information Processing in Medical Imaging*, pages 29–42. Springer, 1997.

- [SW22] Tselil Schramm and Alexander S. Wein. Computational barriers to estimation from low-degree polynomials. In *Annals of Statistics*, 50(3):1833–1858, 2022.
- [SBD19] Martin Slawski and Emanuel Ben-David. Linear regression with sparsely permuted data. In *Electronic Journal of Statistics*, 13(1):1–36, 2019.
- [SDL20] Martin Slawski, Emanuel Ben-David, and Ping Li. Two-stage approach to multivariate linear regression with sparsely mismatched data. In *Journal of Machine Learning Research*, 21(1):8422–8463, 2020.
- [Sze39] Gabor Szegő. Orthogonal polynomials. American Mathematical Society, 1939.
- [UHV18] Jayakrishnan Unnikrishnan, Saeid Haghghatshoar, and Martin Vetterli. Unlabeled sensing with random linear measurements. In *IEEE Transactions on Information Theory*, 64(5):3237–3253, 2018.
- [WWXY22] Haoyu Wang, Yihong Wu, Jiaming Xu, and Israel Yoloh. Random graph matching in geometric models: the case of complete graphs. In *Proceedings of 35th Conference on Learning Theory (COLT)*, pages 3441–3488. PMLR, 2022.
- [Wein22] Alexander S. Wein. Optimal low-degree hardness of maximum independent set. In *Mathematical Statistics and Learning*, 4(3):221–251, 2022.
- [Wis28] John Wishart. The generalized product moment distribution in samples from a normal multivariate population. In *Biometrika*, 20A(1-2):32–52, 1928.
- [ZSWB22] Ilias Zadik, Min Jae Song, Alexander S. Wein, and Joan Bruna. Lattice-based methods surpass sum-of-squares in clustering. In *Proceedings of 35th Conference on Learning Theory (COLT)*, pages 1247–1248. PMLR, 2022.
- [ZK16] Lenka Zdeborová and Florent Krzakala. Statistical physics of inference: thresholds and algorithms. In *Advances in Physics*, 65(5):453–552, 2016.
- [ZL20] Hang Zhang and Ping Li. Optimal estimator for unlabeled linear regression. In *Proceedings of the 37th International Conference on Machine Learning (ICML)*, pages 11153–11162. PMLR, 2020.
- [ZSL22] Hang Zhang, Martin Slawski, and Ping Li. The benefits of diversity: Permutation recovery from multiple measurement vectors in unlabeled sensing. In *IEEE Transactions on Information Theory*, 68(4):2509–2529, 2022.