

REMARKS ON MINIMAL HYPERSURFACES IN GRADIENT SHRINKING RICCI SOLITONS

YUKAI SUN AND GUANGRUI ZHU

ABSTRACT. In this paper, we prove that any compact 2-sided smooth stable minimal hypersurface in gradient Ricci soliton (M^n, g, f) with scalar curvature $R \geq (n-1)\lambda$ must have vanished second fundamental form and vanished normal Ricci curvature. For shrinking gradient Ricci solitons with scalar curvature $R \geq (n-1)\lambda$, the existence of an area-minimizing hypersurface would imply M is splitting.

1. INTRODUCTION

A compact(without boundary) minimal hypersurface Σ in a Riemannian manifold (M, g) is stable if for any $\phi \in C^\infty(\Sigma)$ such that

$$\int_{\Sigma} |\nabla^{\Sigma} \phi|^2 - (\text{Ric}(\nu, \nu) + |A|^2)\phi^2 \geq 0,$$

where ∇^{Σ} is the connection on Σ induced from (M, g) , Ric is the Ricci curvature of M , ν is the outwards pointing unit normal vector of Σ and A is the fundamental form of Σ .

Schoen-Yau [10] obtained a classification for the stable minimal hypersurface in complete three-dimensional manifolds with nonnegative scalar curvature. Chodosh-Li-Stryker[2] showed that the stable minimal hypersurface Σ is totally geodesic and $\text{Ric}(\nu, \nu) = 0$ in four-dimensional manifolds with scalar curvature bounded from below by a positive constant, nonnegative sectional curvature and weakly bounded geometry. Hong-Yan [8] also found that the stable minimal hypersurface Σ is totally geodesic and $\text{Ric}(\nu, \nu) = 0$ in five-dimensional Riemannian manifolds under some curvature conditions.

We aim to study the compact minimal hypersurface in gradient Ricci solitons with scalar curvature bounded from below. First, we revisit the definition of the gradient Ricci soliton.

Definition 1.1. A gradient Ricci soliton, denoted by (M, g, f) , is a smooth n -dimensional complete Riemannian manifold (M, g) equipped with a smooth function $f : M \rightarrow \mathbb{R}$, satisfying

$$(1.1) \quad \text{Ric} + \text{Hess } f = \lambda g$$

for some $\lambda \in \mathbb{R}$, where Ric is the Ricci tensor and $\text{Hess } f$ is the Hessian of f .

If f is not a constant function, we call the gradient Ricci soliton (M, g, f) is nontrivial.

When $\lambda > 0$, $\lambda = 0$ or $\lambda < 0$, the gradient Ricci soliton (M, g, f) is referred to as a shrinking, steady, or expanding gradient Ricci soliton, respectively. The gradient Ricci soliton can be viewed as a special structure on a Riemannian manifold that relates the Ricci curvature and metric by a function and is of great importance in the study of singularities in the Ricci flow [3, 6]. Therefore, we can ask whether

2020 *Mathematics Subject Classification.* 53C24, 53C21.

Key words and phrases. minimal surface, gradient Ricci soliton, scalar curvature, μ -bubble.

the compact stable minimal hypersurface has a vanishing fundamental form and vanishing normal Ricci curvature under the assumption of a lower bounded scalar curvature. Indeed, we can demonstrate the following theorem:

Theorem 1.2. Any compact 2-sided stable minimal hypersurface is totally geodesic with vanishing normal Ricci curvature in a nontrivial shrinking gradient Ricci soliton (M^n, g, f) with scalar curvature $R \geq (n-1)\lambda$ for $3 \leq n \leq 7$.

If $n > 7$, the minimal hypersurfaces may not be smooth. However, if $n \leq 7$, all minimal hypersurfaces are smooth [11]. Therefore, we consider the dimension of M between 3 and 7.

We can also obtain an analogous result for steady and expanding gradient Ricci solitons under a similar condition as in Theorem 1.2. However, we find that for steady and expanding gradient Ricci solitons, Munteanu-Wang[9] has proved that:

Theorem 1.3 (Theorem 3.1[9]). Let (M^n, g, f) be a steady Ricci soliton. Assume that there exists a smooth compact embedded minimal hypersurface Σ in M . Then (M, g) splits isometrically as a direct product $\Sigma \times \mathbb{R}$.

Theorem 1.4 (Theorem 3.2[9]). Let (M^n, g, f) be an expanding Ricci soliton with scalar curvature $R \geq \lambda(n-1)$. Assume that there exists a smooth compact embedded minimal hypersurface Σ in M . Then (M, g) splits isometrically as a direct product $\Sigma \times \mathbb{R}$.

Now we consider the compact area-minimizing hypersurface. We recall the definition of area-minimizing hypersurface.

Definition 1.5. Let Σ be a compact hypersurface in a Riemannian manifold (M, g) . Suppose that the homology class $[\Sigma] \in H_{n-1}(M, \mathbb{Z})$ of Σ is non-zero, i.e., $[\Sigma] \neq 0$ in $H_{n-1}(M, \mathbb{Z})$. Then Σ is called an area-minimizing hypersurface if

$$\text{Area}(\Sigma) = \inf_{\hat{\Sigma} \in [\Sigma]} \{\text{Area}(\hat{\Sigma})\},$$

where $\text{Area}(\Sigma)$ denotes the area of Σ .

Bray-Brendle-Neves[1], Hao-Shi-Sun[7], and Zhu[12] have shown that the existence of a compact area-minimizing hypersurface implies a local splitting property under some proper conditions. Motivated by their work, we have the following results for shrinking Ricci solitons.

Theorem 1.6. For $3 \leq n \leq 7$, let (M^n, g, f) be a nontrivial shrinking gradient Ricci soliton with $R \geq (n-1)\lambda$. If there exists a compact 2-side area-minimizing hypersurface Σ in M , then Σ must be an Einstein manifold, and M is isometric to the Riemannian product $\Sigma \times \mathbb{R}$.

The proof of Theorem 1.6 is using the μ -bubble method as was done in [12]. One key step for the local splitting is to carefully choose the function μ in equation (3.6).

The paper is organized as follows. In section 2, we give some facts about the gradient Ricci soliton and μ -bubble. In section 3, we prove theorem 1.2 and theorem 1.6.

Acknowledgements: Yukai Sun is partially funded by the National Key R&D Program of China Grant 2020YFA0712800.

2. BASIC FACT ABOUT GRADIENT RICCI SOLITON AND μ -BUBBLE

In this section, we revisit some key facts concerning gradient Ricci solitons and μ -bubbles.

The following lemma is well known for gradient Ricci soliton. For completeness, we give a proof.

Lemma 2.1. For gradient Ricci solitons (M, g, f) , We have the following equations

$$(2.1) \quad R + \Delta f = \lambda n,$$

$$(2.2) \quad R + |\nabla f|^2 - 2\lambda f = 0 \ (\lambda \neq 0).$$

Proof. We compute in a local normal coordinate $\{\partial_i\}_{i=1}^n$. Taking trace for formula (1.1), we get the formula (2.1).

Since

$$\nabla_i \nabla_j \nabla_k f - \nabla_j \nabla_i \nabla_k f = -R_{ijkl} \nabla_l f,$$

and

$$\nabla_j \nabla_k f = \lambda g_{jk} - R_{jk}, \quad \nabla_i \nabla_k f = \lambda g_{ik} - R_{ik},$$

we have

$$(2.3) \quad \nabla_j R_{ik} - \nabla_i R_{jk} = -R_{ijkl} \nabla_l f.$$

Taking trace about k, j for (2.3) and use the contracted second Bianchi identity $2\nabla_j R_{ij} = \nabla_i R$, we have

$$\nabla_j R_{ij} - \nabla_i R = -R_{il} \nabla_l f$$

and

$$\nabla_i R = 2R_{il} \nabla_l f.$$

We compute

$$\begin{aligned} \nabla_i (R + |\nabla f|^2 - 2\lambda f) &= \nabla_i R + 2\nabla_i \nabla_l f \nabla_l f - 2\lambda \nabla_i f \\ &= 2(R_{il} + \nabla_i \nabla_l f) \nabla_l f - 2\lambda \nabla_i f \\ &= 2\lambda g_{il} \nabla_l f - 2\lambda \nabla_i f \\ &= 0. \end{aligned}$$

So $R + |\nabla f|^2 - 2\lambda f = \text{constant}$. We can add another constant C to f to make the constant zero without changing the equation for $\lambda \neq 0$. Thus, we get the equation (2.2). \square

Next we recall the basic knowledge of μ -bubble. A Riemannian band (M^n, g) is a compact connected orientable smooth manifold with a metric g and nonempty boundary ∂M such that

$$\partial M = \partial_- M \cup \partial_+ M, \quad \partial_- M \neq \emptyset, \quad \partial_+ M \neq \emptyset, \quad \partial_- M \cap \partial_+ M = \emptyset.$$

On the Riemannian band (M^n, g) , let $\overset{\circ}{M}$ denote the interior of M and h be a smooth function either on $\overset{\circ}{M}$ or on M . A Caccioppoli set $\bar{\Omega} \subset M$ is chosen such that its boundary $\partial \bar{\Omega} \subset \overset{\circ}{M}$ is smooth and $\partial_- M \subset \bar{\Omega}$. Let $\partial^* \Omega$ denote the reduced boundary of the Caccioppoli set Ω . We consider the following functional

$$(2.4) \quad E(\Omega) = \int_{\partial^* \Omega} d\mathcal{H}^{n-1} - \int_{\Omega} (\chi_{\Omega} - \chi_{\bar{\Omega}}) h d\mathcal{H}^n$$

for $\Omega \in \mathcal{C}$, where \mathcal{C} is defined as

$$\mathcal{C} = \{\Omega \mid \text{all Caccioppoli sets } \Omega \subset M \text{ and } \Omega \Delta \bar{\Omega} \Subset \overset{\circ}{M}\},$$

here \mathcal{H}^n denotes n -dimensional Hausdorff measure. We usually omit the measure \mathcal{H}^n in the integral for simplicity. Then the μ -bubbles are critical points of the functional $E(\Omega)$.

From Proposition 2.1 in [13] and Section 5.1 in [5], we have the following existence result of the μ -bubble.

Lemma 2.2 (Existence of μ -bubble). For a Riemannian band (M^n, g) with $3 \leq n \leq 7$, if either $h \in C^\infty(\overset{\circ}{M})$ with $h \rightarrow \pm\infty$ on $\partial_{\mp}M$, or $h \in C^\infty(M)$ with

$$h|_{\partial_-M} > H_{\partial_-M}, \quad h|_{\partial_+M} < H_{\partial_+M}$$

where H_{∂_-M} is the mean curvature of ∂_-M with respect to the inward normal and H_{∂_+M} is the mean curvature of ∂_+M with respect to the outward normal. Then there exists an $\Omega \in \mathcal{C}$ with smooth boundary such that

$$E(\Omega) = \inf_{\Omega' \in \mathcal{C}} E(\Omega').$$

In the following, we discuss the first and second variations of the μ -bubble.

Lemma 2.3 (First and second variation of μ -bubble). Let Ω_t be a smooth 1-parameter family of region in \mathcal{C} with $\Omega_0 = \Omega$ and normal speed ϕ at $t = 0$, then

$$(2.5) \quad \frac{d}{dt}E(\Omega_t)|_{t=0} = \int_{\partial\Omega} (H - h)\phi,$$

where H is the mean curvature of $\partial\Omega$. In particular, a μ -bubble Ω satisfies

$$H = h.$$

For the μ -bubble Ω , we have

$$(2.6) \quad \frac{d^2}{dt^2}E(\Omega_t)|_{t=0} = \int_{\partial\Omega} [-\Delta_\Sigma\phi - |A|^2\phi - \text{Ric}(\nu, \nu)\phi - \langle \nabla h, \nu \rangle \phi] \phi \geq 0.$$

where ν is the outwards pointing unit normal vector on $\partial\Omega$.

3. PROOF OF THEOREMS

In this section, we prove our results.

We first recall the properties of the minimal hypersurface. If there is a compact minimal hypersurface Σ in gradient Ricci soliton (M^n, g, f) . Then the mean curvature H of Σ is zero. By the stable condition and the second variation of Σ , there is

$$(3.1) \quad \int_{\Sigma} [|\nabla^\Sigma\phi|^2 - (\text{Ric}(\nu, \nu) + |A|^2)\phi^2] \geq 0,$$

where $\phi \in C^\infty(\Sigma)$, ν is the outwards pointing unit normal vector on Σ and A is the second fundamental of Σ . On Σ , we also have

$$(3.2) \quad \text{Hess } f(\nu, \nu) = \Delta f - \Delta_\Sigma f - H\langle \nu, \nabla f \rangle,$$

where Δ_Σ is the Laplace operator on Σ .

Proof of Theorem 1.2. Assume that there is a compact stable minimal hypersurface Σ in M . Since

$$\text{Ric}(\nu, \nu) + \text{Hess } f(\nu, \nu) = \lambda,$$

by (3.1), we have

$$(3.3) \quad \int_{\Sigma} [|\nabla^\Sigma\phi|^2 - (\lambda - \text{Hess } f(\nu, \nu) + |A|^2)\phi^2] \geq 0,$$

combining with equations (3.2) and (2.1), there is

$$(3.4) \quad \int_{\Sigma} [|\nabla^\Sigma\phi|^2 + (\lambda(n-1) - R - |A|^2 - \Delta_\Sigma f)\phi^2] \geq 0.$$

Taking $\phi \equiv 1$, we obtain

$$(3.5) \quad \int_{\Sigma} (\lambda(n-1) - R - |A|^2) \geq 0.$$

Therefore, if $R \geq \lambda(n-1)$, we have $A \equiv 0$, $R = (n-1)\lambda$ and $\Delta f = \lambda$ on Σ . Hence,

$$\int_{\Sigma} \text{Ric}(\nu, \nu) = \int_{\Sigma} (\lambda - \text{Hess } f(\nu, \nu)) = 0.$$

By equation (3.1),

$$\int_{\Sigma} [|\nabla^{\Sigma} \phi|^2 - \text{Ric}(\nu, \nu)\phi^2] \geq 0.$$

Since the operator $-\Delta_{\Sigma} - \text{Ric}(\nu, \nu)$ is nonnegative, its first eigenvalue λ_1 is nonnegative. By Rayleigh's principle,

$$\lambda_1 = \inf_{0 \neq \phi \in H_1^2(\Sigma)} \frac{\int_{\Sigma} [|\nabla^{\Sigma} \phi|^2 - \text{Ric}(\nu, \nu)\phi^2]}{\int_{\Sigma} \phi^2} \leq \frac{\int_{\Sigma} [-\text{Ric}(\nu, \nu)]}{\int_{\Sigma}} = 0.$$

Thus $\lambda_1 = 0$, i.e.

$$-\Delta_{\Sigma} 1 - \text{Ric}(\nu, \nu)1 = 0.$$

Consequently, $\text{Ric}(\nu, \nu) = 0$, $\text{Hess } f(\nu, \nu) = \lambda$ and $\Delta_{\Sigma} f = 0$ on Σ . We get f is a constant on Σ . \square

Before proof of Theorem 1.6, we need the following lemma which was proved in Proposition 3.2[1] or Lemma 3.3[12].

Lemma 3.1. Let Σ be an area minimizing hypersurface in M with vanished second fundamental form and vanished normal Ricci curvature, then we can construct a local foliation $\{\Sigma_t\}_{-\epsilon \leq t \leq \epsilon}$ in M such that Σ_t are of constant mean curvature and $\Sigma_0 = \Sigma$.

Proof of Theorem 1.6. By Theorem 1.2 and Lemma 3.1, we can construct a local foliation $\{\Sigma_t\}_{-\epsilon \leq t \leq \epsilon}$ in M such that Σ_t are of constant mean curvature and $\Sigma_0 = \Sigma$, let H_t denote the mean curvature of Σ_t . We claim that $H_t \equiv 0$ for any $t \in (-\epsilon, \epsilon)$. It suffices to prove this for nonnegative t , as the argument for the negative side is analogous. For contradiction, suppose that the assertion is not true. then there exists a $t_0 > 0$ such that $H_{t_0} = \delta > 0$, since otherwise $H_t \leq 0$ and the area-minimizing property of Σ will imply $H_t \equiv 0$. Let Ω_{t_0} denote the region enclosed by Σ and Σ_{t_0} i.e.,

$$\Omega_{t_0} := \cup_{t \in [0, t_0]} \Sigma_t.$$

Consider the following functional

$$(3.6) \quad E(\Omega) = \int_{\partial^* \Omega \setminus \Sigma} d\mathcal{H}^{n-1} - \int_{\Omega} h(\chi_{\Omega} - \chi_{\bar{\Omega}}) d\mathcal{H}^n, h = \delta e^{-f},$$

for $\Omega, \bar{\Omega} \in \mathcal{C}$ and $\Sigma \subset \bar{\Omega}$, where

$$\mathcal{C} = \{\Omega \mid \text{all Caccioppoli sets } \Omega \subset \Omega_{t_0} \text{ and } \Omega \Delta \bar{\Omega} \Subset \overset{\circ}{\Omega}_{t_0}\}.$$

From equation (2.2), we have $0 < h < \delta$, by Lemma 2.2, there exists a μ -bubble $\hat{\Omega}$ such that $\hat{\Sigma} = \partial \hat{\Omega} \setminus \Sigma$ is a smooth two sided hypersurface disjoint from Σ and Σ_{t_0} . Let \hat{H} and \hat{A} denote the mean curvature and the second fundamental form of the $\hat{\Sigma}$, respectively. Then we have $\hat{H} = h$ and

$$\int_{\hat{\Sigma}} [|\nabla^{\hat{\Sigma}} \phi|^2 - (\text{Ric}(\hat{\nu}, \hat{\nu}) + |\hat{A}|^2 + \langle \nabla h, \hat{\nu} \rangle) \phi^2] \geq 0,$$

where $\hat{\nu}$ is the outwards pointing unit normal vector on $\hat{\Sigma}$. Taking $\phi = 1$ and combining with equations (2.1) and (3.2), we obtain

$$\int_{\hat{\Sigma}} [(n-1)\lambda - R - h\langle \nabla f, \hat{\nu} \rangle - \langle \nabla h, \hat{\nu} \rangle - |\hat{A}|^2] \geq 0,$$

we also have

$$(3.7) \quad \int_{\hat{\Sigma}} \left(\lambda(n-1) - R - |\hat{A}|^2 \right) \geq 0.$$

Hence, $|\hat{A}| \equiv 0$, which leads to a contradiction.

Since H_t vanishes for any $t \in (-\epsilon, \epsilon)$, there holds $\text{Area}(\Sigma_t) = \text{Area}(\Sigma)$, which yields that Σ_t is also area-minimizing, by Theorem 1.2, Σ_t is totally geodesic and has vanished normal Ricci curvature.

In the following, we establish the existence of a local isometry $\Phi : \Sigma \times \mathbb{R} \rightarrow M$. Let $\tilde{\nu}_t$ and \tilde{V}_t denote the outwards pointing unit normal vector field and normal variation vector field of the foliation $\{\Sigma_t\}_{-\epsilon \leq t \leq \epsilon}$, respectively. Assume $\Phi : \Sigma \times (-\epsilon, \epsilon) \rightarrow M$ is the flow generated by \tilde{V}_t . It follows that Φ is an embedding in a small neighborhood of Σ and the pull-back of the metric g is

$$\Phi^*(g) = \psi^2 dt^2 + \Phi_t^*(g_t),$$

where $\psi = \langle \tilde{V}_t, \tilde{\nu}_t \rangle > 0$ is the lapse function (see definition in (3.3) of [1]) and g_t is the induced metric on Σ_t from g . Since Σ_t is totally geodesic, we have $\partial_t \Phi_t^*(g_t) = 2\psi A_{\Sigma_t} = 0$, which implies that $\Phi_t^*(g_t) = g|_{\Sigma}$. Additionally, the stability condition yields $-\Delta_{\Sigma_t} \psi - |A_{\Sigma_t}|^2 \psi - \text{Ric}(\nu_t, \nu_t) \psi = -\Delta_{\Sigma_t} \psi = 0$, then $\psi(\cdot, t) \equiv \text{constant}$. Let

$$r(\cdot, t) = \int_0^t \psi(\cdot, s) ds,$$

then

$$\Phi^*(g) = dr^2 + g|_{\Sigma}.$$

Hence, $\Phi : \Sigma \times (-\epsilon, \epsilon) \rightarrow M$ is local a isometry. Through a continuous argument as Proposition 3.8 in [1], we conclude that there exists a local isometry $\Phi : \Sigma \times \mathbb{R} \rightarrow M$. Moreover, Φ is a Riemannian covering map. Since M has constant scalar curvature $R \equiv (n-1)\lambda$, by Theorem 10 in [4], M is isometric to a product of an Einstein manifold with \mathbb{R}^k . We conclude Σ is an Einstein manifold and M is isometric to $\Sigma \times \mathbb{R}$. \square

We now give an example of the gradient Ricci soliton which is the rigid case in Theorem 1.6

Example 3.2. $(\mathbb{S}^{n-1} \times \mathbb{R}, g, f)$ with $g = g_{\mathbb{S}^{n-1}} + dt^2$, $f(x, t) = \frac{\lambda t^2 + (n-1)}{2}$ and $\lambda = n-2$.

At the end of this paper, we would like to ask the following question:

Question 3.3. For $3 \leq n \leq 7$ and $2 \leq k \leq (n-2)$, let (M^n, g, f) be a gradient Ricci soliton with $R \geq (n-k)\lambda$. If there exists a compact 2-side area-minimizing surface Σ of codimension k in M , do we have Σ must be an Einstein manifold and M is isometric to the Riemannian product $\Sigma \times \mathbb{R}^k$?

REFERENCES

1. Hubert Bray, Simon Brendle, and Andre Neves, *Rigidity of area-minimizing two-spheres in three-manifolds*, Comm. Anal. Geom. **18** (2010), no. 4, 821–830. MR 2765731
2. Otis Chodosh, Chao Li, and Douglas Stryker, *Complete stable minimal hypersurfaces in positively curved 4-manifolds*, 2024.
3. Bennett Chow, *Ricci solitons in low dimensions*, Graduate Studies in Mathematics, vol. 235, American Mathematical Society, Providence, RI, [2023] ©2023. MR 4624811
4. Manuel Fernández-López and Eduardo García-Río, *On gradient Ricci solitons with constant scalar curvature*, Proc. Amer. Math. Soc. **144** (2016), no. 1, 369–378. MR 3415603
5. Misha Gromov, *Four lectures on scalar curvature*, Perspectives in scalar curvature. Vol. 1, World Sci. Publ., Hackensack, NJ, [2023] ©2023, pp. 1–514. MR 4577903
6. Richard S. Hamilton, *The formation of singularities in the Ricci flow*, Surveys in differential geometry, Vol. II (Cambridge, MA, 1993), Int. Press, Cambridge, MA, 1995, pp. 7–136. MR 1375255

7. Tianze Hao, Yuguang Shi, and Yukai Sun, *Llarull type theorems on complete manifolds with positive scalar curvature*, Trans. Amer. Math. Soc. **377** (2024), no. 10, 7403–7420. MR 4855340
8. Han Hong and Zetian Yan, *Rigidity and nonexistence of cmc hypersurfaces in 5-manifolds*, 2024.
9. Ovidiu Munteanu and Jiaping Wang, *A minkowski type inequality for manifolds with positive spectrum*, 2023.
10. Richard Schoen and Shing Tung Yau, *Complete three-dimensional manifolds with positive Ricci curvature and scalar curvature*, Seminar on Differential Geometry, Ann. of Math. Stud., vol. No. 102, Princeton Univ. Press, Princeton, NJ, 1982, pp. 209–228. MR 645740
11. Leon Simon, *Lectures on geometric measure theory*, Proceedings of the Centre for Mathematical Analysis, Australian National University, vol. 3, Australian National University, Centre for Mathematical Analysis, Canberra, 1983. MR 756417
12. Jintian Zhu, *Rigidity of area-minimizing 2-spheres in n -manifolds with positive scalar curvature*, Proc. Amer. Math. Soc. **148** (2020), no. 8, 3479–3489. MR 4108854
13. ———, *Width estimate and doubly warped product*, Trans. Amer. Math. Soc. **374** (2021), no. 2, 1497–1511. MR 4196400

KEY LABORATORY OF PURE AND APPLIED MATHEMATICS, SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, BEIJING, 100871, P. R. CHINA
Email address: 52195500003@stu.ecnu.edu.cn

SCHOOL OF MATHEMATICAL SCIENCES, EAST CHINA NORMAL UNIVERSITY, SHANGHAI, 200241, P. R. CHINA
Email address: 52275500052@stu.ecnu.edu.cn