

PERVERSE SHEAVES ON THE SEMI-INFINITE FLAG VARIETY AND REPRESENTATIONS OF THE FROBENIUS KERNEL

EMILIEN ZABETH

ABSTRACT. We study a category of Iwahori-equivariant modular perverse sheaves on some avatar of the semi-infinite flag variety, by adapting the work of Arkhipov-Bezrukavnikov-Braverman-Gaitsgory-Mirković. We then construct a functor between the latter category and the category of graded representations of the Frobenius kernel, and conjecture that this functor is an equivalence.

CONTENTS

1. Introduction	1
2. Conventions	6
3. Perverse sheaves on affine Grassmannians	9
4. Drinfeld’s compactifications	23
5. Perverse sheaves on the semi-infinite flag variety	29
6. Convolution	38
7. Iwahori-Whittaker objects	47
8. The convolution functor	50
9. Strategy of proof for Conjecture 8.1	54
References	58

1. INTRODUCTION

Let \mathbb{k} be a finite field of characteristic ℓ . In this paper, we study a category of modular (with coefficients in \mathbb{k}) perverse sheaves $\mathrm{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}})$ on the semi-infinite flag variety associated with a reductive algebraic group G defined over an algebraically closed field \mathbb{F} of prime characteristic $p \neq \ell$. Under certain conditions on ℓ and G , we construct an exact functor between the latter category and the extended principal block $\mathrm{Rep}_{[0]}(\check{\mathbf{G}}_1\check{\mathbf{T}})$ of the category of $\check{\mathbf{G}}_1\check{\mathbf{T}}$ -modules, where $\check{\mathbf{G}}_1$ is the Frobenius kernel of $\check{\mathbf{G}}$, the split reductive group over \mathbb{k} whose Frobenius twist coincides with the Langlands dual group of G over \mathbb{k} , and $\check{\mathbf{T}} \subset \check{\mathbf{G}}$ is a maximal torus. This functor is expected to be an equivalence, and we give a condition under which it is. Here, the semi-infinite flag variety $\mathcal{Fl}^{\frac{\infty}{2}}$ is constructed using the Drinfeld compactification space $\overline{\mathrm{Bun}}_{N^-}$, and our proof follows closely the steps taken in [ABB⁺05], where the authors work with coefficients of characteristic zero. Namely, we construct a functor between $\mathrm{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}})$ and the category of modules over the “regular perverse sheaf” – which is known to be equivalent to $\mathrm{Rep}_{[0]}(\check{\mathbf{G}}_1\check{\mathbf{T}})$ thanks to the recent work of Achar-Riche [AR22]. The key result in the construction is the fact that the IC-sheaf on the stack $\overline{\mathrm{Bun}}_{N^-}$ satisfies the Hecke eigensheaf property with respect to $\mathrm{Rep}(\check{\mathbf{G}}^{(1)})$ (see Proposition 6.5). This result was known for characteristic zero coefficients and used in the context of the geometric Langlands program (see [BFGM02]), but had not been observed yet (to the knowledge of the author) in the case of positive characteristic.

1.1. Representations of the Frobenius kernel. We fix a Borel subgroup $B \subset G$ and a maximal torus $T \subset B$. Let $\text{Fr} : \check{\mathbf{G}} \rightarrow \check{\mathbf{G}}^{(1)}$ denote the Frobenius endomorphism, and $\check{\mathbf{G}}_1$ be its kernel. If we let $L(\lambda)$ be the simple $\check{\mathbf{G}}$ -module associated with a dominant character λ of $\check{\mathbf{T}}$, then $L(\lambda)$ remains simple as a $\check{\mathbf{G}}_1$ -module when λ is ℓ -restricted, and any simple $\check{\mathbf{G}}_1$ -module is obtained as such a restriction. Thus (thanks to Steinberg’s tensor product formula), the understanding of simple $\check{\mathbf{G}}_1$ -modules is equivalent to that of simple $\check{\mathbf{G}}$ -modules (under the assumption that the derived subgroup of $\check{\mathbf{G}}$ is simply connected).

For technical reasons, it is often more convenient to work with $\check{\mathbf{G}}_1\check{\mathbf{T}}$ -modules. Simple $\check{\mathbf{G}}_1\check{\mathbf{T}}$ -modules $\{\widehat{L}(\lambda), \lambda \in X^*(\check{\mathbf{T}})\}$ are labeled by the lattice of characters $X^*(\check{\mathbf{T}})$, and remain simple when viewed as $\check{\mathbf{G}}_1$ -modules. For each $\lambda \in X^*(\check{\mathbf{T}})$, the object $\widehat{L}(\lambda)$ is defined as the socle of the baby co-Verma module $\widehat{Z}'(\lambda)$, and coincides with the head of the baby Verma module $\widehat{Z}(\lambda)$.

A central question in the theory of modular representations of algebraic groups is to understand how baby Verma modules decompose in the basis of simple modules in the Grothendieck group of $\text{Rep}(\check{\mathbf{G}}_1\check{\mathbf{T}})$, as one would then get character formulas for simple modules. Recent progress have been obtained by Riche-Williamson in [RW21], where the authors obtain a formula – proved in the case where $\ell \geq 2h - 1$, where h is the Coxeter number of $\check{\mathbf{G}}$ – involving the ℓ -canonical basis of the affine Hecke algebra. Their proof does not rely on a geometric model for the category $\text{Rep}(\check{\mathbf{G}}_1\check{\mathbf{T}})$, but rather on the geometric Satake equivalence, which gives a geometric model for the category of $\check{\mathbf{G}}$ -modules. The goal of this paper is to obtain a geometric model for the category of $\check{\mathbf{G}}_1\check{\mathbf{T}}$ -modules (or rather the principal block of this category), with the hope that one might be able to use new techniques afforded by the geometry for the understanding of the characters of simple modules.

We will in fact restrict our attention to the extended principal block $\text{Rep}_{[0]}(\check{\mathbf{G}}_1\check{\mathbf{T}})$ of the category of $\check{\mathbf{G}}_1\check{\mathbf{T}}$ -modules, whose understanding is sufficient for the study of the general case (we assume for this that $\ell > h$). This is the Serre subcategory of $\text{Rep}(\check{\mathbf{G}}_1\check{\mathbf{T}})$ generated by the family

$$\{\widehat{L}(w \bullet_\ell 0) \mid w \in W_{\text{ext}}\},$$

where \bullet_ℓ denotes the “dot” action of the extended affine Weyl group $W_{\text{ext}} := W \ltimes X^*(\check{\mathbf{T}})$ on the affine space $X^*(\check{\mathbf{T}}) \otimes_{\mathbb{Z}} \mathbb{R}$.

Remark 1.1. In view of [RW22], it would be desirable to obtain a geometric model for the whole category $\text{Rep}(\check{\mathbf{G}}_1\check{\mathbf{T}})$, instead of merely the principal block. Indeed, assume that there exists a space X (for instance an ind-algebraic stack over \mathbb{F}), endowed with an action of the multiplicative group \mathbb{G}_m and an equivalence of categories

$$(1.1) \quad \text{Perv}(X) \xrightarrow{\sim} \text{Rep}(\check{\mathbf{G}}_1\check{\mathbf{T}}),$$

(where the category $\text{Perv}(X)$ might involve an equivariance condition or “factorization” property, cf. §5.1). Then, following the philosophy of *loc. cit.*, the characters of projective indecomposable $\check{\mathbf{G}}_1\check{\mathbf{T}}$ -modules (which are the analogues of indecomposable tilting $\check{\mathbf{G}}$ -modules) might be expressed in terms of the stalks of indecomposable parity sheaves on the connected components of the fixed points X^{μ_ℓ} , where $\mu_\ell \subset \mathbb{G}_m$ is the subgroup of ℓ -th roots of unity. To the knowledge of the author, there is no conjecture in the literature concerning a space X realizing the equivalence (1.1).

1.2. The geometric Satake equivalence and Finkelberg-Mirković conjecture. The present work builds on several other celebrated equivalences of categories. First, we have an equivalence of monoidal categories (whose first complete proof was obtained in [MV04])

$$\text{Sat} : (\text{Perv}_{G[[t]]}(\text{Gr}), \star) \xrightarrow{\sim} (\text{Rep}_{\mathbb{k}}(\check{\mathbf{G}}^{(1)}), \otimes_{\mathbb{k}}),$$

where the left-hand side denotes the *Satake category* (equipped with a convolution product), consisting of perverse (étale) sheaves on the affine Grassmannian Gr , with coefficients in \mathbb{k} and which respect an equivariance condition for the left action of the positive loop group $G[[t]]$ on Gr . This result is called the *geometric Satake equivalence*, and is ubiquitous in this paper.

Next, denote by $\mathrm{Rep}_{[0]}(\check{\mathbf{G}})$ the extended principal block of the category $\mathrm{Rep}(\check{\mathbf{G}})$, defined as the Serre subcategory generated by the family $\{L(\lambda), \lambda \in (W_{\mathrm{ext}} \bullet_{\ell} 0) \cap \mathbf{Y}^+\}$. We let $\mathrm{Perv}_{I_u}(\mathrm{Gr})$ denote the category of perverse sheaves on Gr with coefficients in \mathbb{k} , equivariant with respect to the Iwahori subgroup I_u associated with B . The latter category admits a right action of the monoidal category $(\mathrm{Perv}_{G[[t]]}(\mathrm{Gr}), \star)$. Below, the functor sw^* denotes a certain auto-equivalence of the category $\mathrm{Perv}_{G[[t]]}(\mathrm{Gr})$ (see §3.3 for the definition), and W_{ext}^S a certain subset of W_{ext} labelling the simple objects $\{L_w, w \in W_{\mathrm{ext}}^S\}$ of $\mathrm{Perv}_{I_u}(\mathrm{Gr})$. The following statement is known as the Finkelberg-Mirković conjecture (cf. [FM99, §1.5]). It was recently proved by Bezrukavnikov-Riche (it is still conjectured to hold when $\ell > h$).

Theorem 1.2. *Assume that $\ell > h$, and that $\ell \neq 19$ (resp. $\ell \neq 31$) if $\check{\mathbf{G}}$ has a component of type E_7 (resp. E_8). There exists an equivalence of highest weight categories*

$$\mathrm{FM} : \mathrm{Perv}_{I_u}(\mathrm{Gr}) \xrightarrow{\sim} \mathrm{Rep}_{[0]}(\check{\mathbf{G}})$$

which satisfies

$$\mathrm{FM}(L_w) \simeq L(w^{-1} \bullet_{\ell} 0) \quad \forall w \in W_{\mathrm{ext}}^S.$$

Moreover, for any $\mathcal{F} \in \mathrm{Perv}_{I_u}(\mathrm{Gr})$, $\mathcal{G} \in \mathrm{Perv}_{G[[t]]}(\mathrm{Gr})$, there exists a bifunctorial isomorphism

$$\mathrm{FM}(\mathcal{F} \star \mathcal{G}) \simeq \mathrm{FM}(\mathcal{F}) \otimes \mathrm{Fr}^*(\mathrm{Sat}(\mathrm{sw}^* \mathcal{G})).$$

1.3. Results of Achar-Riche. In their recent work [AR22], Achar and Riche give an incarnation of $\mathrm{Rep}_{[0]}(\check{\mathbf{G}}_1 \check{\mathbf{T}})$ in terms of perverse sheaves. More precisely, they define a category $\mathrm{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ consisting of \mathbf{Y} -graded ind-objects of the category $\mathrm{Perv}_{I_u}(\mathrm{Gr})$, endowed with a right-action of the “regular perverse sheaf” \mathcal{R} . We refer to §3 for a detailed review of this construction. In particular, they prove that the category $\mathrm{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ is equivalent to $\mathrm{Rep}_{[0]}(\check{\mathbf{G}}_1 \check{\mathbf{T}})$ under the assumptions of Theorem 1.2 (cf. Theorem 3.27). This result will be important for us, since it allows the theory of perverse sheaves to enter the picture.

1.4. Main statement. The main part of this paper is dedicated to the definition and study of an “artificial” category of perverse sheaves on the semi-infinite flag variety \mathcal{Fl}^{∞} , equivariant with respect to the unipotent radical of the Iwahori subgroup I_u . The idea of considering perverse sheaves on \mathcal{Fl}^{∞} goes back to Lusztig in the 80’s (see [Lus, §11]). It was then defined as the quotient

$$G(\mathbb{F}((t)))/T \cdot N(\mathbb{F}((t))),$$

where t is a formal variable. The above set can be endowed with a structure of ind-scheme (see [FM99, §4]), on which the group I_u acts. The orbits are labelled by the affine Weyl group and their closure relations are given by Lusztig’s periodic order (these are the first requirements for the category of I_u -equivariant perverse sheaves on it to be equivalent to $\mathrm{Rep}_{[0]}(\check{\mathbf{G}}_1 \check{\mathbf{T}})$); the problem is that the closures of these orbits are schemes of infinite type (so, in particular, are not varieties). The usual methods are therefore not suited to define I_u -equivariant perverse sheaves on such a space.

To overcome this difficulty, a general idea is to replace a local problem by a global problem: namely, replacing the formal disc $\mathrm{Spec}(\mathbb{F}[[t]])$ by a smooth complete curve X (for technical simplicity, we will take X to be \mathbb{P}^1 in the main body text). The formal disc is

seen as the formal neighborhood of a fixed point x . This point of view led the authors of [FM99] and [FFKM99] to consider perverse sheaves on spaces of Drinfeld compactifications (or more precisely on quasimap spaces), and was taken up in [ABB⁺05] to establish an equivalence of categories between $\text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}})$ and the extended principal block of the category of graded modules for the small quantum group (associated with $\check{\mathbf{G}}$) at an ℓ -th root of unity – the latter category being the “quantum analogue” of $\text{Rep}_{[0]}(\check{\mathbf{G}}_1\check{\mathbf{T}})$. In these previous references, the authors worked with sheaves whose field of coefficients is of characteristic zero. We check that the constructions of [ABB⁺05] still make sense for a field of positive characteristic. Let us give a brief overview of these constructions¹.

We denote by \mathbf{X} (resp. \mathbf{Y}) the lattice of characters (resp. cocharacters) of T , and by $\mathbf{X}^+ \subset \mathbf{X}$ the subset of dominant characters determined by B . Let also B^- be the Borel subgroup which is opposite to B with respect to T , and N (resp. N^-) be the unipotent radical of B (resp. of N^-). Fix a smooth complete curve X over \mathbb{F} , a closed point $x \in X$, and let Bun_G (resp. Bun_{N^-}) denote the stack of principal G -bundles (resp. principal N^- -bundles) over X . The space $\overline{\text{Bun}}_{N^-}$ is an algebraic stack locally of finite type over \mathbb{F} which is a “compactification” of the canonical projection $\overline{\text{Bun}}_{N^-} \rightarrow \text{Bun}_G$, whose construction is recalled in §4.1. For any test scheme S , a point of $\overline{\text{Bun}}_{N^-}(S)$ consists of a pair $(\mathcal{F}_G, (\kappa_{\lambda}^-)_{\lambda \in \mathbf{X}^+})$, where $\mathcal{F}_G \in \text{Bun}_G(S)$ and $\kappa_{\lambda}^- : \mathcal{V}_{\mathcal{F}_G}^{\lambda} \rightarrow \mathcal{O}_{X \times S}$ is a morphism of coherent sheaves whose dual is injective, such that the collection (κ_{λ}^-) satisfies the Plücker relations.

In order to get a space which is stratified by the extended affine Weyl group, we need to introduce some variants of the previous stack. First, we consider the ind-algebraic stack ${}_{\infty}\overline{\text{Bun}}_{N^-}$, where the morphism κ_{λ}^- is allowed to have a pole at x for all λ . For any $\nu \in \mathbf{Y}$, we let ${}_{\leq \nu}\overline{\text{Bun}}_{N^-}$ be the variant where each κ_{λ}^- is now allowed to have a pole of order $\leq \langle \nu, \lambda \rangle$ at x (in particular, ${}_{\infty}\overline{\text{Bun}}_{N^-}$ is the direct limit of the stacks ${}_{\leq \nu}\overline{\text{Bun}}_{N^-}$ for $\nu \in \mathbf{Y}$), and ${}_{\nu}\overline{\text{Bun}}_{N^-}$ be its open substack where we impose that the pole is of order exactly $\langle \nu, \lambda \rangle$ at x . Next, in order to make the Weyl group appear in the stratification, we consider the stacks ${}^1_{\infty}\overline{\text{Bun}}_{N^-}$, ${}^1_{\leq \nu}\overline{\text{Bun}}_{N^-}$ and ${}^1_{\nu}\overline{\text{Bun}}_{N^-}$, where we add a structure of level one at x for \mathcal{F}_G (namely, an isomorphism between G and the fiber of \mathcal{F}_G at x). In particular, the group G acts on the latter three stacks, and we denote by ${}^1_{\infty}\overline{\text{Bun}}_{N^-}$ the quotient stack $N \backslash {}^1_{\infty}\overline{\text{Bun}}_{N^-}$. We have an isomorphism

$${}^1_{\nu}\overline{\text{Bun}}_{N^-} \simeq G \times^{N^-} {}^1_{\nu}\mathcal{N}$$

where ${}^1_{\nu}\mathcal{N}$ is the canonical N^- -torsor over ${}_{\nu}\overline{\text{Bun}}_{N^-}$ (cf. §4.4). The Bruhat decomposition of G/B^- induces a “stratification” (each stratum is not necessarily smooth, since ${}_{\nu}\overline{\text{Bun}}_{N^-}$ is not) of ${}^1_{\nu}\overline{\text{Bun}}_{N^-}$ by Schubert strata. For $x := wt_{\nu} \in W_{\text{ext}}$, we denote by

$$\text{IC}_x^{\frac{\infty}{2}}, \quad \Delta_x^{\frac{\infty}{2}}, \quad \nabla_x^{\frac{\infty}{2}}$$

the objects of $\text{Perv}_T({}^1_{\infty}\overline{\text{Bun}}_{N^-})$ (where the index T stands for T -equivariant) respectively defined as the intermediate extension, the !-extension and the *-extension of the IC-sheaf on the Schubert stratum in ${}^1_{\nu}\overline{\text{Bun}}_{N^-}$ associated with w . In particular, these objects are equivariant with respect to N and each $\text{IC}_x^{\frac{\infty}{2}}$ is a simple object of $\text{Perv}({}^1_{\infty}\overline{\text{Bun}}_{N^-})$.

In the theorem below, $\text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}})$ denotes a certain full subcategory of the category of T -equivariant perverse sheaves $\text{Perv}_T({}^1_{\infty}\overline{\text{Bun}}_{N^-})$ (see §5.1). The simple objects of $\text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}})$ are the IC-sheaves $\{\text{IC}_x^{\frac{\infty}{2}}, x \in W_{\text{ext}}\}$. For any $\mathcal{S} \in \text{Perv}_{I_u}(\text{Gr})$, the object $\mathcal{S} \star \mathcal{IC}_0^{\frac{\infty}{2}}$ denotes the object of $\text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}})$ obtained by the convolution process (§6.3),

¹To simplify exposition, we assume in this part that G is simply connected.

and we put

$$\widetilde{\text{For}}_0 : \text{Perv}_{I_u}(\text{Gr}) \rightarrow \text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}}), \mathcal{S} \mapsto \mathcal{S} \star \mathcal{IC}_0^{\frac{\infty}{2}}.$$

Moreover, the monoidal category $(\text{Perv}_{G[[t]]}(\text{Gr}), \star)$ acts on the left on $\text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}})$. Finally, recall that the forgetful functor induces a functor

$$\text{For}_0 : \text{Rep}_{[0]}(\check{\mathbf{G}}) \rightarrow \text{Rep}_{[0]}(\check{\mathbf{G}}_1 \check{\mathbf{T}}).$$

The following conjecture has its roots in [FM99, §1.5].

Conjecture 1.1. *Assume that $\ell \geq h$. There exists an equivalence of categories*

$$\widetilde{\text{FM}} : \text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}}) \xrightarrow{\sim} \text{Rep}_{[0]}(\check{\mathbf{G}}_1 \check{\mathbf{T}})$$

such that

$$\widetilde{\text{FM}} \circ \widetilde{\text{For}}_0 \simeq \text{For}_0 \circ \widetilde{\text{FM}},$$

satisfying

$$\widetilde{\text{FM}}(\mathcal{IC}_x^{\frac{\infty}{2}}) \simeq \widehat{\mathbf{L}}(x^{-1} \cdot_{\ell} 0), \quad \widetilde{\text{FM}}(\nabla_x^{\frac{\infty}{2}}) \simeq \widehat{\mathbf{Z}}'(x^{-1} \cdot_{\ell} 0), \quad \widetilde{\text{FM}}(\Delta_x^{\frac{\infty}{2}}) \simeq \widehat{\mathbf{Z}}(x^{-1} \cdot_{\ell} 0)$$

for all $x \in W_{\text{ext}}$ and

$$\widetilde{\text{FM}}(\mathcal{G} \star \mathcal{F}) \simeq \widetilde{\text{FM}}(\mathcal{F}) \otimes \text{Fr}^*(\text{Sat}(\text{sw}^* \mathcal{G}))$$

functorially in $\mathcal{F} \in \text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}})$, $\mathcal{G} \in \text{Perv}_{G[[t]]}(\text{Gr})$.

Remark 1.3. (1) The strategy to prove the above conjecture is to construct a functor from $\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ to $\text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}})$.

(2) The objects of $\text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}})$ are perverse sheaves on ${}^{I_u}\overline{\text{Bun}}_{N^-}$ satisfying a certain ‘‘factorization property’’. This property implies that the objects are constructible with respect to the Schubert strata (see Remark 5.19) and that the simple objects are labelled by W_{ext} . From this perspective, one can think of the factorization property as a replacement for the I_u -equivariance on $\mathcal{Fl}^{\frac{\infty}{2}}$.

1.5. What is done in this paper. We study the category $\text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}})$, and relate it to the category $\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$. For our construction to work, we need to make the following hypothesis.

Conjecture 1.2. *Let \mathbb{k} be either a finite field or a finite extension of \mathbb{Q}_{ℓ} . For any $i \in \mathbb{Z}$ and closed point $x \in \overline{\text{Bun}}_{N^-}$, the dimension of the \mathbb{k} -vector space $\mathbf{H}^i((\mathcal{IC}_{\overline{\text{Bun}}_{N^-}})_x)$ does not depend on the characteristic of \mathbb{k} .*

Here is our main result.

Theorem 1.4. *Fix ℓ , and assume that Conjecture 1.2 holds. Then there is an exact functor*

$$\text{Conv}^{\text{Hecke}} : \text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R}) \rightarrow \text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}}),$$

which commutes with Verdier duality, and induces a bijection between the isomorphism classes of simple objects.

Conjecture 1.2 is expected to hold for a general reductive group G , but there is no proof available at the moment.

Remark 1.5. In an upcoming work, Achar-Riche-Dhillon should give a proof of Conjecture 1.2 (with minor restrictions on ℓ).

Below (§5.3), we add the following observation, well-known by experts.

Proposition 1.6. *Conjecture 1.2 holds when $G = \text{SL}_n$.*

We conjecture that $\text{Conv}^{\text{Hecke}}$ is an equivalence of categories. In §9, we give a strategy of proof for this conjecture, which follows the same steps as the analogue statement of [ABB⁺05]. We plan to achieve the proof of this conjecture in future work.

1.6. Overview. After having introduced the necessary conventions in §2, we recall the main results of [AR22] in §3. Namely, we explain the construction of the category $\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$, which consists of certain ind-objects in the category $\text{Perv}_{I_u}(\text{Gr})$. Thanks to *loc. cit.* together with [BR24], this category is equivalent to $\text{Rep}_{[0]}(\check{\mathbf{G}}_1\check{\mathbf{T}})$ under certain hypothesis. In §4, we recall the constructions of the main geometric objects, namely the Drinfeld compactification and Zastava spaces. In §5, we check that the definitions of the categories of perverse sheaves on the semi-infinite flag variety from [ABB⁺05] still make sense in the modular setting. The main result of this section is that the dimensions of the stalks of the IC-sheaf \mathcal{IC}_0^{∞} on $\overline{\text{Bun}}_{N^-}$ do not depend on the characteristic (see Lemma 5.8) when $G = \text{SL}_n$. This allows us to take back the main constructions of *loc. cit.*. In particular, we exhibit an action of the monoidal category $(\text{Perv}_{G[[t]]}(\text{Gr}), \star)$ on $\text{Perv}_{G[[t]]}(\mathcal{Fl}^{\infty})$, see Proposition 6.5. This property plays a crucial role in §8, where we are able to construct a functor $\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R}) \rightarrow \text{Perv}_{I_u}(\mathcal{Fl}^{\infty})$ which is exact, sends simple objects to simple objects, and commutes with Verdier duality and convolution.

1.7. Acknowledgments. This work was conducted during PhD studies at the Université Clermont Auvergne, under the supervision of Simon Riche and Geordie Williamson. I especially want to thank S.R. for many rereadings of earlier versions.

2. CONVENTIONS

2.1. The group G . We let \mathbb{k} be a finite field of characteristic $\ell > 0$, \mathbb{F} be an algebraically closed field of prime characteristic $p \neq \ell$, and $\check{\mathbf{G}}$ be a split connected reductive group over \mathbb{k} . We denote by G the reductive group defined over \mathbb{F} , such that the Langlands dual group $G_{\mathbb{k}}^{\vee}$ over \mathbb{k} is the Frobenius twist $\check{\mathbf{G}}^{(1)}$ of $\check{\mathbf{G}}$. We let $T \subset B \subset G$ be a maximal torus and Borel subgroup, $W = N_G(T)/T$ (where $N_G(T)$ is the normalizer of T in G) denote the associated Weyl group, and $\check{\mathfrak{R}}_s \subset \check{\mathfrak{R}}_+ \subset \check{\mathfrak{R}}$ (resp. $\mathfrak{R}_s \subset \mathfrak{R}_+ \subset \mathfrak{R}$) denote the sets of simple coroots, positive² coroots and coroots (resp. simple roots, positive roots and roots) determined by B (i.e. the positive roots consist of the T weights in $\text{Lie}(B)$). We let B^- be the Borel subgroup which is opposite to B (i.e. such that $B \cap B^- = T$), put $N = [B, B]$, $N^- = [B^-, B^-]$.

Denote by $\mathbf{Y} := X_*(T)$ (resp. $\mathbf{X} := X^*(T)$) the coweight lattice (resp. weight lattice) of G , let \mathbf{Y}^{pos} (resp. \mathbf{X}^{pos}) be the sub-semigroup spanned by $\check{\mathfrak{R}}_s$ (resp. by the positive roots), and \mathbf{Y}^+ (resp. \mathbf{X}^+) be the subset of dominant coweights (resp. dominant weights) determined by B . We also define the set of strictly dominant coweights

$$\mathbf{Y}^{++} = \{\lambda \in \mathbf{Y} \mid \langle \alpha, \lambda \rangle > 0 \forall \alpha \in \check{\mathfrak{R}}_+\},$$

where $\langle \cdot, \cdot \rangle : \mathbf{X} \times \mathbf{Y} \rightarrow \mathbb{Z}$ is the usual perfect pairing. Denote by $W_{\text{ext}} := W \ltimes \mathbf{Y}$ the extended affine Weyl group. For $(w, \lambda) \in W \times \mathbf{Y}$, we will denote by $w\lambda$ the corresponding element of W_{ext} ; we will simply denote by λ the element $e\lambda$, where $e \in W$ is the identity. For $\lambda_1, \lambda_2 \in \mathbf{Y}$, we will write $\lambda_1 \leq \lambda_2$ if $\lambda_2 - \lambda_1 \in \mathbf{Y}^{\text{pos}}$.

We assume that the quotient $\mathbf{X}/\mathbb{Z}\check{\mathfrak{R}}$ of \mathbf{X} by the subgroup spanned by the roots is torsion-free. This means that the morphism of groups

$$\mathbf{Y} \rightarrow \text{Hom}_{\mathbb{Z}}(\mathbb{Z}\check{\mathfrak{R}}, \mathbb{Z})$$

induced by restriction is surjective, and is equivalent to requiring that the scheme-theoretic center of G be a torus. In particular, there exists $\zeta \in \mathbf{Y}$ such that $\langle \alpha, \zeta \rangle = 1$ for all $\alpha \in \check{\mathfrak{R}}_s$.

We let $\check{\mathbf{T}} \subset \check{\mathbf{B}} \subset \check{\mathbf{G}}$ be the maximal torus and Borel subgroup such that $T_{\mathbb{k}}^{\vee} \simeq \check{\mathbf{T}}^{(1)}$ and $B_{\mathbb{k}}^{\vee} \simeq \check{\mathbf{B}}^{(1)}$, where the superscript (1) denotes the Frobenius twist, and $T_{\mathbb{k}}^{\vee}$ (resp. $B_{\mathbb{k}}^{\vee}$) denotes the split maximal torus (resp. Borel subgroup) of the Langlands dual group of

²We take opposite conventions compared with [AR22].

G over \mathbb{k} corresponding to T (resp. to B). In the sequel, we will identify the character lattice $X^*(\check{\mathbf{T}})$ with \mathbf{Y} , in such a way that the morphism of \mathbb{Z} -modules

$$X^*(\check{\mathbf{T}}^{(1)}) \rightarrow \mathbf{Y}$$

induced by the Frobenius morphism $\mathrm{Fr}|_{\check{\mathbf{T}}} : \check{\mathbf{T}} \rightarrow \check{\mathbf{T}}^{(1)}$ identifies with $\lambda \mapsto \ell \cdot \lambda$.

For any group scheme H over \mathbb{k} , we will denote by $\mathrm{Rep}(H)$ the category of finite dimensional representations of H over \mathbb{k} . For any $V \in \mathrm{Rep}(\check{\mathbf{G}}^{(1)})$ and $\mu \in \mathbf{Y}$, we will denote by $V(\mu)$ the corresponding μ -weight space.

For all $\lambda \in \mathbf{X}^+$, we let V^λ be the Weyl module of G associated with the weight λ , and fix^3 a highest weight vector $v_\lambda \in V^\lambda$. For $\lambda_1, \lambda_2 \in \mathbf{X}^+$, there is a uniquely defined G -equivariant morphism

$$(2.1) \quad V^{\lambda_1 + \lambda_2} \rightarrow V^{\lambda_1} \otimes V^{\lambda_2}$$

sending $v_{\lambda_1 + \lambda_2}$ to $v_{\lambda_1} \otimes v_{\lambda_2}$. If \mathcal{F}_G is a G -bundle and \mathcal{F}_T is a T -bundle on some scheme S and $\lambda \in \Lambda$, we will denote by

$$\mathcal{V}_{\mathcal{F}_G}^\lambda := \mathcal{F}_G \times^G V^\lambda, \quad \mathcal{L}_{\mathcal{F}_T}^\lambda := \mathcal{F}_T \times^T \mathbb{A}_{\mathbb{F}}^1$$

the induced vector bundle and line bundle respectively on S , where T acts on the affine line $\mathbb{A}_{\mathbb{F}}^1$ via the character λ in the second twisted product.

2.2. Stacks. In this note, we will study algebraic stacks and schemes defined over \mathbb{F} . The stacks will be attached to the smooth complete curve X and to G . For any group scheme H over \mathbb{F} , we will denote by Bun_H the stack of principal H -bundles over X . This means that Bun_H is a functor from the category of \mathbb{F} -schemes to the category of groupoids, such that for any test scheme S , $\mathrm{Bun}_H(S)$ consists of the category of principal H -bundles over $X \times S$. We will denote by $\mathcal{F}_H^0 := H \times X \times S$ the trivial H -bundle. For any morphism of schemes $S' \rightarrow S$, the functor $\mathrm{Bun}_H(S) \rightarrow \mathrm{Bun}_H(S')$ is obtained by pullback.

In the sequel, we will work with variants of this stack, known as Drinfeld's compactifications. When introducing such a stack \mathcal{X} , we will say that it classifies *something*, where something will be the category $\mathcal{X}(\mathbb{F})$. More precisely, S will always be a test \mathbb{F} -scheme, and the groupoid $\mathcal{X}(S)$ will consist in the data of some vector bundles over $X \times S$ together with some morphisms of sheaves between them, and the functors $\mathcal{X}(S) \rightarrow \mathcal{X}(S')$ will be obtained by pullback for any morphism of \mathbb{F} -schemes $S' \rightarrow S$.

When defining locally closed substacks of a given stack, we will often only work with the field-valued points. This is justified by the following facts. For \mathcal{X} an algebraic stack, we denote by $|\mathcal{X}|$ its set of points, endowed with the Zariski topology. Recall that by definition, $|\mathcal{X}|$ consists of the set $\bigsqcup_{(\mathbb{L}, f)} \mathrm{Ob}(\mathcal{X}(\mathrm{Spec}(\mathbb{L}))) / \sim$, where (\mathbb{L}, f) runs through the maps $\mathrm{Spec}(\mathbb{L}) \rightarrow \mathrm{Spec}(\mathbb{F})$, with \mathbb{L} a field. Two points $x_{(\mathbb{L}, f)}, x'_{(\mathbb{L}', f')}$ are equivalent if there is a common field extension \mathbb{L}'' of \mathbb{L} and \mathbb{L}' such that both points become isomorphic when pullbacked to $\mathcal{X}(\mathrm{Spec}(\mathbb{L}''))$. The opens of $|\mathcal{X}|$ are the subsets $|U|$, where $U \hookrightarrow \mathcal{X}$ is an open substack. When \mathcal{X} is a scheme, we recover the underlying topological space. If $p : X \rightarrow \mathcal{X}$ is a presentation, with X a scheme, then the induced morphism $|X| \rightarrow |\mathcal{X}|$ is surjective, continuous and open ([Sta22, § 100.4]). If $T \subset |\mathcal{X}|$ is a locally closed subset, then there exists a unique reduced locally closed substack $\mathcal{Z} \subset \mathcal{X}$ such that $T = |\mathcal{Z}|$ as subsets of $|\mathcal{X}|$ ([Sta22, Remark 100.10.5]).

³There is a canonical choice for such a vector, see the proof of [Jan03, Lemma II.2.13].

2.3. Group arc schemes. For any closed point $x \in X$, we denote by $\hat{\mathcal{O}}_x$ the completion of the local ring of X at x , let \mathcal{K}_x be the fraction field of $\hat{\mathcal{O}}_x$, and put $D_x := \text{Spec}(\hat{\mathcal{O}}_x)$. If t denotes a local coordinate on X at x , then we get isomorphisms $\hat{\mathcal{O}}_x \simeq \mathbb{F}[[t]]$, $\mathcal{K}_x \simeq \mathbb{F}((t))$. For any \mathbb{F} -algebra R , we consider the completed tensor products $R \hat{\otimes} \hat{\mathcal{O}}_x$ and $R \hat{\otimes} \mathcal{K}_x$, so that we get identifications

$$R \hat{\otimes} \hat{\mathcal{O}}_x \simeq R[[t]] \quad R \hat{\otimes} \mathcal{K}_x \simeq R((t)).$$

For an affine group scheme H over \mathbb{F} , we will denote by $H[[t]]$, resp. $H((t))$, resp. $H[t^{-1}]$, the functor defined on points via

$$H[[t]](R) := H(R \hat{\otimes} \hat{\mathcal{O}}_x), \quad H((t)) := H(R \hat{\otimes} \mathcal{K}_x), \quad H[t^{-1}] := H(R[t^{-1}]).$$

It is a classical fact that $H[[t]]$ is a group scheme and $H((t))$ a group ind-scheme. By [Zhu16, §2.3], $H[t^{-1}]$ is a group ind-scheme.

We fix once and for all a closed point $x \in X$. For $k \in \mathbb{Z}_{\geq 0}$, we let G_k be the group scheme representing the functor $R \mapsto G(R[t]/(t^k))$, and G^k be the congruence subgroup defined as the kernel of the canonical morphism $G[[t]] \rightarrow G_k$. Notice that $G^0 = G[[t]]$.

Finally, we define the Iwahori subgroup $I \subset G[[t]]$ (resp. $I_u \subset I$, resp. $I^- \subset G[[t]]$, resp. $I_u^- \subset I$) as the pre-image of B (resp. of N , resp. of B^- , resp. of N^-) through the map $G[[t]] \rightarrow G$, $t \mapsto 0$.

2.4. Perverse sheaves on stacks. This paper will be concerned with defining various categories of perverse sheaves on algebraic stacks defined over \mathbb{F} . Since our stacks will be locally of finite type, we will freely use the formalism developed in [LO09]. For an algebraic stack \mathcal{X} locally of finite type, we will denote by $\mathbf{D}(\mathcal{X})$ the derived category of bounded complexes of \mathbb{k} -sheaves, whose cohomology is constructible. We will also denote by $({}^p\mathbf{D}(\mathcal{X})^{\leq 0}, {}^p\mathbf{D}(\mathcal{X})^{\geq 0})$ the perverse t -structure, and by $\text{Perv}(\mathcal{X})$ its heart. It follows from the constructions that, if \mathcal{X} is an \mathbb{F} -scheme of finite type acted on by a connected \mathbb{F} -algebraic group K , then the constructible equivariant derived category $\mathbf{D}_K^b(\mathcal{X})$ of \mathbb{k} -sheaves – in the sense of Bernstein-Lunts [BL94] – coincides with $\mathbf{D}(K \backslash \mathcal{X})$, where $K \backslash \mathcal{X}$ denotes the quotient stack. Likewise, the perverse t -structures coincide. We will often write $\mathbf{D}_K(\mathcal{X})$ (resp. $\text{Perv}_K(\mathcal{X})$) instead of $\mathbf{D}(K \backslash \mathcal{X})$ (resp. $\text{Perv}(K \backslash \mathcal{X})$).

In the sequel, we will often have the following situations. We are given a system of \mathbb{F} -stacks locally of finite type $(\mathcal{X}_n)_{n \in \mathbb{N}}$ together with closed embeddings $\mathcal{X}_m \hookrightarrow \mathcal{X}_n$ for $m \leq n$, and denote by $\mathcal{X} := \varinjlim_n \mathcal{X}_n$ the associated ind-stack. Then for any $m \leq n$, the $*$ -pushforward through the morphism $\mathcal{X}_m \hookrightarrow \mathcal{X}_n$ induces a fully-faithful and t -exact functor $\mathbf{D}(\mathcal{X}_m) \rightarrow \mathbf{D}(\mathcal{X}_n)$. We let $\mathbf{D}(\mathcal{X})$ and $\text{Perv}(\mathcal{X})$ be the categories defined as direct limits of the systems $(\mathbf{D}(\mathcal{X}_n))_{n \geq 0}$ and $(\text{Perv}(\mathcal{X}_n))_{n \geq 0}$ respectively. Finally, assume that each \mathcal{X}_n is a scheme (so that \mathcal{X} is an ind-scheme)⁴ and that \mathcal{X} is acted on by a group scheme $K \subset G[[t]]$ such that, for all $n \geq 0$, each \mathcal{X}_n is stable under the action of K , and there exists $u_n \in \mathbb{Z}_{\geq 0}$ such that the subgroup $K^{u_n} := K \cap G^{u_n}$ acts trivially on \mathcal{X}_n . Then we put $\mathbf{D}(K \backslash \mathcal{X}_n) := \mathbf{D}(K[t/t^{u_n}] \backslash \mathcal{X}_n)$, where $K[t/t^{u_n}]$ is defined as the algebraic group K/K^{u_n} . Using the fact that, for any $m \geq u_n$, the kernel of the canonical morphism $K[t/t^m] \rightarrow K[t/t^{u_n}]$ is unipotent, one can show that the previous definition does not depend on the choice of u_n (cf. the appendix of [BR18] for a detailed treatment in the analytic case). For $m \leq n$, pushforward induces a fully faithful and t -exact functor $\mathbf{D}(K \backslash \mathcal{X}_m) \rightarrow \mathbf{D}(K \backslash \mathcal{X}_n)$. We define the categories $\mathbf{D}(K \backslash \mathcal{X})$ and $\text{Perv}(K \backslash \mathcal{X})$ as direct limits.

⁴The cases we have in mind are when \mathcal{X} is either the affine Grassmannian or the affine flag variety.

2.5. Ind-objects. Our constructions will require to pass to categories of ind-objects. An ind-object of a (locally small) category \mathcal{C} is a formal filtered colimit of objects of \mathcal{C} , where “formal” means that the colimit is taken in the category of presheaves $\text{Fun}(\mathcal{C}^{\text{op}}, \text{Sets})$. The inductive limits in the latter category will be denoted by \varinjlim . For a complete account of these constructions, we refer the reader to [KS06, Chapter 6]. We will denote by $\text{Ind}(\mathcal{C})$ the full subcategory of $\text{Fun}(\mathcal{C}^{\text{op}}, \text{Sets})$ consisting of ind-objects of \mathcal{C} . If \mathcal{C} is abelian, then so is $\text{Ind}(\mathcal{C})$. The natural functor $\mathcal{C} \rightarrow \text{Ind}(\mathcal{C})$ is fully faithful. We will use the fact that a functor $F : \mathcal{C} \rightarrow \mathcal{G}$ between two categories induces a functor $\text{Ind}(\mathcal{C}) \rightarrow \text{Ind}(\mathcal{G})$, which we will still denote by F .

3. PERVERSE SHEAVES ON AFFINE GRASSMANNIANS

3.1. The extended affine Weyl group. We state a few well known combinatorial facts about the extended affine Weyl group. A detailed treatment of this topic can be found in [AR21, §2] (notice that we take opposite conventions with respect to *loc. cit.*, so that our positive roots correspond to the negative roots in *loc. cit.*).

Recall that the affine Weyl group $W_{\text{aff}} := W \ltimes \mathbb{Z}\check{\mathfrak{A}}$ is a subgroup of W_{ext} , which is endowed with a structure of Coxeter group. Namely, each root $\alpha \in \check{\mathfrak{A}}$ defines a reflection $s_\alpha \in W$, and we will denote by S the set of simple reflections (i.e. associated with a simple root) of the finite Weyl group, which is known to generate W . It is also well known (cf. for instance [Jan03, II.6.3]) that the set of simple reflections

$$S_{\text{aff}} := \{(s, 0), s \in S_0\} \cup \{(s_\beta, -\beta^\vee)\},$$

where β^\vee runs through the set of largest short coroots of the irreducible components of $\check{\mathfrak{A}}^\vee$, generates W_{aff} . The set S_{aff} is then a Coxeter generating system for W_{aff} . We will denote by $\ell : W_{\text{aff}} \rightarrow \mathbb{Z}_{\geq 0}$ the associated length function, and by \leq the associated Bruhat order. The length function and Bruhat order can be extended to the group W_{ext} . We let W_{ext}^S denote the subset of W_{ext} consisting of elements w whose length is minimal in wW .

The group W_{ext} acts naturally on the vector space $E := \mathbf{Y} \otimes_{\mathbb{Z}} \mathbb{R}$ via the action

$$wt_\mu \cdot \lambda := w(\lambda + \mu)$$

for any $w \in W$, $\mu \in \mathbf{Y}$, $\lambda \in \mathbf{Y} \otimes_{\mathbb{Z}} \mathbb{R}$. This action defines a hyperplane arrangement in E , and thus a set of alcoves (which are the connected components of the complement of E by the set of hyperplanes), on which W_{ext} acts. The *fundamental alcove* is the alcove defined as

$$\mathfrak{A}_{\text{fund}} := \{v \in E \mid \forall \beta \in \check{\mathfrak{A}}_+, -1 < \langle \beta, v \rangle < 0\},$$

and is a fundamental domain for the action of W_{ext} on E . If we set

$$\mathcal{C} := \{v \in E \mid \forall \beta \in \check{\mathfrak{A}}_+, \langle \beta, v \rangle < 0\},$$

then we have

$$(3.1) \quad W_{\text{ext}}^S = \{w \in W_{\text{ext}} \mid w^{-1} \cdot (\mathfrak{A}_{\text{fund}}) \subset \mathcal{C}\}.$$

We next define the *fundamental box*

$$\Pi := \{v \in E \mid \forall \beta \in \check{\mathfrak{A}}_s, -1 < \langle \beta, v \rangle < 0\},$$

together with the set of restricted elements

$$W_{\text{ext}}^{\text{res}} = \{w \in W_{\text{ext}} \mid w^{-1} \cdot (\mathfrak{A}_{\text{fund}}) \subset \Pi\}.$$

Since any alcove inside \mathcal{C} is included in a subset of the form $\lambda + \Pi$ for a unique $\lambda \in -\mathbf{Y}^+$, we have

$$W_{\text{ext}}^S = \{wt_\lambda, w \in W_{\text{ext}}^{\text{res}}, \lambda \in \mathbf{Y}^+\}.$$

Lastly, we will need the following property.

Lemma 3.1. *For any $w \in W_{\text{ext}}^S$ and $\lambda \in \mathbf{Y}^+$ we have $\ell(wt_\lambda w_0) = \ell(w) + \ell(t_\lambda w_0)$.*

Proof. Notice that we have $wt_\lambda w_0 = ww_0w_0t_\lambda w_0 = ww_0t_{w_0(\lambda)}$. Since $w_0(\lambda) \in -\mathbf{Y}^+$ (and we have chosen opposite conventions for the choice of the fundamental alcove compared [AR22]), the result then follows from [AR22, Lemma 2.7]. \square

3.2. Coset representatives. We introduce the set

$${}^S W_{\text{ext}}^S \subset W_{\text{ext}},$$

consisting of elements $w \in W_{\text{ext}}$ such that w is minimal in Ww and $vw \in W_{\text{ext}}^S$ for all $v \in W$ (see [AR21, Lemma 2.4] for equivalent definitions). In particular, we have the inclusion ${}^S W_{\text{ext}}^S \subset W_{\text{ext}}^S$.

For any $\lambda \in \mathbf{Y}$, we denote by w_λ^R the minimal element in $t_\lambda W$.

Lemma 3.2. [AR21, Lemma 2.5] *Let $\lambda \in \mathbf{Y}$. We have $w_\lambda^R \in {}^S W_{\text{ext}}^S$ iff $w_0(\lambda) \in \mathbf{Y}^{++}$. Moreover, in this case we have $w_\lambda^R = w_0 t_{w_0(\lambda)}$.*

Corollary 3.3. *We have a bijection*

$$\mathbf{Y}^{++} \rightarrow {}^S W_{\text{ext}}^S, \quad w_0(\lambda) \mapsto w_0 t_{w_0(\lambda)}.$$

Proof. This follows from the fact that the map $\mathbf{Y} \rightarrow W_{\text{ext}}^S$, $\lambda \rightarrow w_\lambda^R$ is a bijection, together with Lemma 3.2. \square

Finally, if we put

$${}^S W_{\text{ext}}^{\text{res}} := {}^S W_{\text{ext}}^S \cap W_{\text{ext}}^{\text{res}},$$

then as above we have

$${}^S W_{\text{ext}}^S = \{wt_\lambda, w \in {}^S W_{\text{ext}}^{\text{res}}, \lambda \in \mathbf{Y}^+\}.$$

3.3. Perverse sheaves on flag varieties. We start by recalling a few well-known facts about the affine Grassmannian and affine flag variety, which can be found for instance in [Fal03]. We consider the \mathbb{F} -ind-schemes Gr and Fl defined as the fppf quotients

$$\text{Gr} := G((t))/G[[t]], \quad \text{Fl} := G((t))/I.$$

For any $\lambda \in \mathbf{Y}$, we denote by $t^\lambda \in T((t))$ the image of t through $\lambda : \mathbb{G}_m((t)) \rightarrow T((t))$. For any $v \in W$, we choose a lift $\dot{v} \in N_G(T)$ and, for any element $w := vt_\lambda \in W_{\text{ext}}$, we set $\dot{w} := \dot{v}t^\lambda \in G((t))$. We define Fl_w as the I -orbit of the image of \dot{w} under the canonical projection $G((t)) \rightarrow \text{Fl}$. Notice that Fl_w is also the I_u -orbit of that same element. It is well known that each Fl_w is an affine \mathbb{F} -space of dimension $\ell(w)$, and that $\overline{\text{Fl}}_v \subset \overline{\text{Fl}}_w \Leftrightarrow v \leq w$ for all $v, w \in W_{\text{ext}}$. We have a decomposition of the associated reduced ind-scheme

$$(\text{Fl})_{\text{red}} = \bigsqcup_{w \in W_{\text{ext}}} \text{Fl}_w.$$

Likewise, for any $w \in W_{\text{ext}}$, we denote by Gr_w the I -orbit (which coincides with the I_u -orbit) of the image of \dot{w} under the canonical projection $G((t)) \rightarrow \text{Gr}$. Each Gr_w is an affine space over \mathbb{F} , of dimension $\ell(w)$ when $w \in W_{\text{ext}}^S$. From the fact that the canonical projection $\text{Fl} \rightarrow \text{Gr}$ is I -equivariant, one can deduce the stratification

$$(\text{Gr})_{\text{red}} = \bigsqcup_{w \in W_{\text{ext}}^S} \text{Gr}_w, \quad \text{and } \overline{\text{Gr}}_v \subset \overline{\text{Gr}}_w \Leftrightarrow v \leq w \quad \forall v, w \in W_{\text{ext}}^S.$$

Finally, for any $\lambda \in \mathbf{Y}$, we let $\text{Gr}^\lambda := G[[t]] \cdot [t^\lambda]$ be the $G[[t]]$ -orbit of the image $[t^\lambda]$ of t^λ in Gr , and let $j^\lambda : \text{Gr}^\lambda \hookrightarrow \text{Gr}$ denote the locally closed immersion. We have

$$\dim(\text{Gr}^\lambda) = \langle 2\rho, \lambda \rangle.$$

Using the Cartan decomposition, one can show that the action of $G[[t]]$ on Gr induces the following equality

$$(\text{Gr})_{\text{red}} = \bigsqcup_{\lambda \in \mathbf{Y}^+} \text{Gr}^\lambda.$$

Each orbit Gr^λ is a smooth \mathbb{F} -scheme of finite type and

$$\overline{\text{Gr}}^\lambda \subset \overline{\text{Gr}}^\mu \Leftrightarrow \lambda \leq \mu \quad \forall \lambda, \mu \in \mathbf{Y}^+.$$

Next, we define three categories of perverse sheaves which will be constructible with respect to the above stratifications. Each time, the formalism recalled in the second paragraph of §2.4 applies.

First notice that, for each $\lambda \in \mathbf{Y}^+$, there exists a large enough k such that G^k acts trivially on $\overline{\text{Gr}}^\lambda$. So as recalled in §2.4, we can define the $G[[t]]$ -equivariant categories $\text{D}_{G[[t]]}(\text{Gr})$ and $\text{Perv}_{G[[t]]}(\text{Gr})$. The latter category is a *highest weight category*, with weight poset \mathbf{Y}^+ . To each $\mu \in \mathbf{Y}^+$, we associate the standard, costandard and simple objects

$$\mathcal{I}_!^\mu := {}^p\tau^{\geq 0}(j_!^\mu \underline{\mathbb{k}}_{\text{Gr}^\mu}[\langle 2\rho, \mu \rangle]), \quad \mathcal{I}_*^\mu := {}^p\tau^{\leq 0}(j_*^\mu \underline{\mathbb{k}}_{\text{Gr}^\mu}[\langle 2\rho, \mu \rangle]), \quad \mathcal{IC}^\mu := j_{!*}^\mu \underline{\mathbb{k}}_{\text{Gr}^\mu}[\langle 2\rho, \mu \rangle].$$

The category $\text{Perv}_{I_u}(\text{Gr})$ is also a highest weight category, with weight poset W_{ext}^S . For each $w \in W_{\text{ext}}^S$, we will denote by Δ_w, ∇_w the corresponding standard and costandard objects. They are respectively defined as the $!$ -pushforward and $*$ -pushforward of the constant sheaf $\underline{\mathbb{k}}_{\text{Gr}_w}[\ell(w)]$ under the inclusion $\text{Gr}_w \hookrightarrow \text{Gr}$ (note that these objects are indeed perverse since the locally closed inclusion is affine). The simple object associated with w will be denoted by L_w : it is defined as the intersection cohomology complex associated with Gr_w (and coincides with the image of the canonical morphism $\Delta_w \rightarrow \nabla_w$).

We introduce notation for the category $\text{Perv}_{I_u}(\text{Fl})$. For $w \in W_{\text{ext}}$, we let \mathcal{D}_w and \mathcal{N}_w be the associated standard and costandard perverse sheaves, defined respectively as the $!$ -pushforward and $*$ -pushforward of the constant sheaf $\underline{\mathbb{k}}_{\text{Fl}_w}[\ell(w)]$ under the inclusion $\text{Fl}_w \hookrightarrow \text{Fl}$. We will denote by $\mathcal{L}_w \in \text{Perv}_{I_u}(\text{Fl})$ the intersection cohomology complex associated with Fl_w (it coincides with the image of the canonical morphism $\mathcal{D}_w \rightarrow \mathcal{N}_w$). The family $(\mathcal{L}_w, w \in W_{\text{ext}})$ are then representatives for the isomorphism classes of the simple objects in $\text{Perv}_{I_u}(\text{Fl})$. Likewise, the standard and costandard objects of the category $\text{Perv}_I(\text{Fl})$ are labelled by W_{ext} , and we will still denote by $\mathcal{D}_w, \mathcal{N}_w$ the corresponding objects associated with w .

3.4. Convolution. We now briefly recall the construction of convolution products defined on the above categories, which will be used profusely in the sequel. We start with the bifunctor

$$(3.2) \quad (-) \star (-) : \text{D}_{G[[t]]}(\text{Gr}) \times \text{D}_{G[[t]]}(\text{Gr}) \rightarrow \text{D}_{G[[t]]}(\text{Gr}),$$

which will endow $\text{D}_{G[[t]]}(\text{Gr})$ with a structure of monoidal category (cf. [BG02, §3.2.2]). For that, we consider the diagram

$$(3.3) \quad \text{Gr} \times \text{Gr} \xleftarrow{p} G((t)) \times \text{Gr} \xrightarrow{q} G((t)) \times^{G[[t]]} \text{Gr} \xrightarrow{m} \text{Gr},$$

where p and q are the canonical projections, and m is the multiplication morphism. For $(\mathcal{F}, \mathcal{G}) \in \text{D}_{G[[t]]}(\text{Gr}) \times \text{D}_{G[[t]]}(\text{Gr})$, $\mathcal{F} \star \mathcal{G}$ is defined as $m_*(\mathcal{F} \boxtimes \mathcal{G})$, where $\mathcal{F} \boxtimes \mathcal{G}$ is the image of $p^*(\mathcal{F} \boxtimes \mathcal{G})$ through the inverse of the equivalence of categories

$$q^* : \text{D}_{G[[t]] \times G[[t]]}(G((t)) \times \text{Gr}) \rightarrow \text{D}_{G[[t]]}(G((t)) \times^{G[[t]]} \text{Gr}).$$

It is a classical result (cf. for instance [BR18, §6.3]) that the convolution product is t -exact, i.e. that $\mathcal{F} \star \mathcal{F}' \in \text{Perv}_{G[[t]]}(\text{Gr})$ when $\mathcal{F}, \mathcal{F}' \in \text{Perv}_{G[[t]]}(\text{Gr})$. For any subgroup scheme $K \subset G[[t]]$, a similar construction endows the category $\text{D}_K(\text{Gr})$ with a right action of the monoidal category $\text{D}_{G[[t]]}(\text{Gr})$.

In the sequel, we will also need the convolution bifunctor

$$(-) \star^I (-) : D_{I_u}(\text{Fl}) \times D_I(\text{Gr}) \rightarrow D_{I_u}(\text{Gr}).$$

The definition is similar to that (3.2), but using the diagram

$$\text{Fl} \times \text{Gr} \xleftarrow{p} G((t)) \times \text{Gr} \xrightarrow{q} G((t)) \times^I \text{Gr} \xrightarrow{m} \text{Gr}$$

instead of (3.3). We have the following standard properties, which are stated in [AR22, Lemma 3.1].

Lemma 3.4. (1) *For any $w \in W_{\text{ext}}$, there exist canonical isomorphisms*

$$\mathcal{D}_w \star^I \mathcal{N}_{w^{-1}} \simeq \mathcal{D}_e \simeq \mathcal{N}_w \star^I \mathcal{D}_{w^{-1}}.$$

(2) *For $w, y \in W_{\text{ext}}$ such that $\ell(wy) = \ell(w) + \ell(y)$ and both wy and y belong to W_{ext}^S , there exist canonical isomorphisms*

$$\mathcal{D}_w \star^I \Delta_y \simeq \Delta_{wy} \quad \mathcal{N}_w \star^I \nabla_y \simeq \nabla_{wy}.$$

3.5. The geometric Satake equivalence and Finkelberg-Mirković conjecture. Recall that $G_{\mathbb{k}}^{\vee}$ is $\check{\mathbf{G}}^{(1)}$. The following celebrated result will be used profusely in the sequel.

Theorem 3.5 ([MV04]). *The convolution product \star makes $\text{Perv}_{G[[t]]}(\text{Gr})$ a monoidal category, and there is an equivalence of monoidal categories*

$$\text{Sat} : (\text{Perv}_{G[[t]]}(\text{Gr}), \star) \xrightarrow{\sim} (\text{Rep}_{\mathbb{k}}(\check{\mathbf{G}}^{(1)}), \otimes_{\mathbb{k}}).$$

This equivalence respects the highest weight structures. This means that \mathcal{I}_1^{μ} , \mathcal{I}_*^{μ} and \mathcal{IC}^{μ} respectively correspond – via the functor Sat – to the Weyl module $M^{(1)}(\mu)$, the induced module $N^{(1)}(\mu)$, and the simple module $L^{(1)}(\mu)$ of highest weight μ .

We let $\text{sw} : \text{Gr} \rightarrow \text{Gr}$ be the isomorphism induced by the inversion map $G((t)) \rightarrow G((t))$, $g \mapsto g^{-1}$, so that pulling back induces an equivalence of categories

$$\text{sw}^* : \text{Perv}_{G[[t]]}(\text{Gr}) \rightarrow \text{Perv}_{G[[t]]}(\text{Gr}).$$

Proposition 3.6. [FS24, VI.12.1] *The auto-equivalence sw^* corresponds– via Sat – to the Chevalley involution on $\text{Rep}_{\mathbb{k}}(\check{\mathbf{G}}^{(1)})$. In particular, for any $\mathcal{G} \in \text{Perv}_{G[[t]]}(\text{Gr})$ and $\mu \in \mathbf{Y}$, we have an isomorphism*

$$\text{Sat}(\text{sw}^* \mathcal{G})(-\mu) \simeq \text{Sat}(\mathcal{G})(w_0(\mu)).$$

The following lemma will be useful in the sequel.

Lemma 3.7. *Let $r \in \mathbb{Z}_{\geq 0}$ and put $\mu_r := (\ell^r - 1)\zeta$. The canonical morphisms*

$$\mathcal{I}_1^{\mu_r} \rightarrow \mathcal{IC}^{\mu_r} \rightarrow \mathcal{I}_*^{\mu_r}$$

are isomorphisms. In particular, the dimensions of the stalks of \mathcal{IC}^{μ_r} are independant of $\text{char}(\mathbb{k})$.

Proof. This follows from the geometric Satake equivalence together with [Jan03, II.3.19, (4)]. Let us propose a geometric proof. In [Zab22], we gave a description of the blocks of the category $\text{Perv}_{G[[t]]}(\text{Gr})$. From this description, we see that if \mathcal{IC}^{λ} is a composition factor of $\mathcal{I}_1^{\mu_r}$ (or of $\mathcal{I}_*^{\mu_r}$), then we must have $\lambda \in W_{\text{aff}} \bullet_{\ell^r} \mu_r$, and $\lambda \leq \mu_r$. But μ_r is minimal in $(W_{\text{aff}} \bullet_{\ell^r} \mu_r) \cap \mathbf{Y}^+$, so that $\lambda = \mu_r$. We deduce that $\mathcal{I}_1^{\mu_r}$ is simple, concluding the proof of the first assertion.

Finally, the last assertion follows from the fact that, when $\text{char}(\mathbb{k}) = 0$, the category $\text{Perv}_{G[[t]]}(\text{Gr})$ is known to be semi-simple, so that the canonical morphisms from the assertion are also isomorphisms. But the stalks of $\mathcal{I}_1^{\mu_r}$ are clearly independant of the characteristic of \mathbb{k} . \square

\mathbf{A}_n ($n \geq 1$)	\mathbf{B}_n ($n \geq 2$)	\mathbf{C}_n ($n \geq 3$)	\mathbf{C}_n ($n \geq 4$)	\mathbf{E}_6	\mathbf{E}_7	\mathbf{E}_8	\mathbf{F}_4	\mathbf{G}_2
$n + 1$	$2n$	$2n$	$2n - 2$	12	19	31	12	6

 FIGURE 1. Bounds on ℓ

The category $\text{Perv}_{I_u}(\text{Gr})$ also has an interpretation in terms of representation theory. We let $\text{Rep}_{[0]}(\check{\mathbf{G}})$ denote the extended principal block of the category $\text{Rep}(\check{\mathbf{G}})$, defined as the Serre subcategory generated by the family $\{L(\lambda), \lambda \in (W_{\text{ext}} \bullet_{\ell} 0) \cap \mathbf{Y}^+\}$. Notice that, for any $w \in W_{\text{ext}}^S$, the weight $w^{-1} \bullet_{\ell} 0$ is dominant by (3.1). We also denote by $\text{Fr} : \check{\mathbf{G}} \rightarrow \check{\mathbf{G}}^{(1)}$ the Frobenius morphism. The following statement is known as the Finkelberg-Mirković conjecture (cf. [FM99, §1.5]). A proof was recently obtained in [BR24].

Theorem 3.8. *Assume that, for any indecomposable constituent in the root system of $(\check{\mathbf{G}}, \check{\mathbf{T}})$, ℓ is greater than the bound in Figure 1. There exists an equivalence of highest weight categories*

$$\text{FM} : \text{Perv}_{I_u}(\text{Gr}) \xrightarrow{\sim} \text{Rep}_0(\check{\mathbf{G}})$$

which satisfies

$$\text{FM}(L_w) \simeq L(w^{-1} \bullet_{\ell} 0) \quad \forall w \in W_{\text{ext}}^S.$$

Moreover, for any $\mathcal{F} \in \text{Perv}_{I_u}(\text{Gr})$, $\mathcal{G} \in \text{Perv}_{G[[t]]}(\text{Gr})$, there exists a bifunctorial isomorphism

$$\text{FM}(\mathcal{F} \star \mathcal{G}) \simeq \text{FM}(\mathcal{F}) \otimes \text{Fr}^*(\text{Sat}(\text{sw}^* \mathcal{G})).$$

Remark 3.9. The above theorem takes a more natural form in [BR24], as the category $\text{Perv}_{I_u}(\text{Gr})$ is replaced by the category of equivariant perverse sheaves on the *opposite affine Grassmannian* $\text{Gr}^{\text{op}} := G[[t]] \backslash G((t))$. For our purposes, it is however more convenient to work with the usual affine Grassmannian.

3.6. Iwahori-Whittaker model for the Satake equivalence. Fix a primitive p -th root of unity $\zeta \in \mathbb{k}$, and consider the Artin-Schreier map $\text{AS} : \mathbb{G}_a \rightarrow \mathbb{G}_a$ determined by the map of rings $x \mapsto x^p - x$. This morphism is a Galois cover of group $\mathbb{Z}/p\mathbb{Z}$, so determines a continuous group morphism $\pi_1(\mathbb{G}_a, 0) \rightarrow \mathbb{Z}/p\mathbb{Z}$, where $\pi_1(\mathbb{G}_a, 0)$ is the étale fundamental group of \mathbb{G}_a with geometric base point 0. The composition of this map with the morphism $\mathbb{Z}/p\mathbb{Z} \rightarrow \mathbb{k}^{\times}$ (induced by ζ) yields a continuous representation of the fundamental group, and thus a local system on \mathbb{G}_a of rank one. We denote this local system by \mathcal{L}_{AS} . We also let $\psi : N^- \rightarrow \mathbb{G}_a$ be a morphism of \mathbb{F} -algebraic groups which restricts to a non-zero morphism on each subgroup $N_{-\alpha}$ for all $\alpha \in \mathfrak{R}_s$, where $N_{-\alpha} \simeq \mathbb{G}_a$ is the root subgroup associated with $-\alpha$.

Define $\chi : I_u^- \rightarrow \mathbb{G}_a$ to be the composition of ψ with the evaluation map. For any $Y \subset \text{Gr}$ which is a locally closed finite union of I_u^- -orbits, one can show (cf. [Cia21, Lemma 3.2]) that the I_u^- -action on Y factors through a quotient group of finite type J such that χ factors through $\chi_J : J \rightarrow \mathbb{G}_a$; we can then consider the $(J, \chi_J^* \mathcal{L}_{\text{AS}})$ -equivariant derived category of étale \mathbb{k} -sheaves $D_{J, \chi_J^* \mathcal{L}_{\text{AS}}}(Y)$. This category is by definition the subcategory of $D(Y)$ consisting of constructible complexes of étale \mathbb{k} -sheaves \mathcal{F} such that there exists an isomorphism $a^* \mathcal{F} \simeq \chi_J^* \mathcal{L}_{\text{AS}} \boxtimes \mathcal{F}$, where $a : J \times Y \rightarrow Y$ is the action map; this definition does not depend on the choice of J . We then define the category $D_{\mathcal{TW}}(\text{Gr})$ as a direct limit of the categories $D_{J, \chi_J^* \mathcal{L}_{\text{AS}}}(Y)$, indexed by finite and closed unions of I_u^- -orbits Y which are ordered by inclusion. Notice that, since the transition maps are push-forwards of closed immersions, they are fully faithful functors, and one can see this limit as an increasing union of categories.

The category $D_{\mathcal{IW}}(\mathrm{Gr})$ admits a canonical perverse t -structure, and we will denote by $\mathrm{Perv}_{\mathcal{IW}}(\mathrm{Gr})$ its heart. Since the $I_{\mathfrak{u}}^-$ orbits are affine \mathbb{F} -schemes, the category $\mathrm{Perv}_{\mathcal{IW}}(\mathrm{Gr})$ admits a transparent highest weight structure (in the sense of [Ric16, 3.7]).

Note that the same construction (simply replacing Gr with Fl) allows one to define the category $D_{\mathcal{IW}}(\mathrm{Fl})$.

For each $\lambda \in \mathbf{Y}$, let us denote by $Y_\lambda \subset \mathrm{Gr}$ the $I_{\mathfrak{u}}^-$ -orbit associated with $w_0(\lambda)$ and by $j^\lambda : Y_\lambda \hookrightarrow \mathrm{Gr}$ the locally closed immersion. One can show that an orbit Y_λ supports a non-zero Iwahori-Whittaker local system iff λ is strictly dominant (i.e if $\lambda \in \zeta + \mathbf{Y}_+$), and that in this case there exists exactly one (up to isomorphism) such local system of rank one on Y_λ , which we will denote by $\mathcal{L}_{\mathrm{AS}}^\lambda$. Thus, the weight poset is given by $\mathbf{Y}^{++} = \zeta + \mathbf{Y}^+$, and the standard, costandard and simple objects associated with some $\lambda \in \mathbf{Y}^{++}$ are respectively given by

$$\Delta_\lambda^{\mathcal{IW}} := j_!^\lambda \mathcal{L}_{\mathrm{AS}}^\lambda[\dim(Y_\lambda)], \quad \nabla_\lambda^{\mathcal{IW}} := j_*^\lambda \mathcal{L}_{\mathrm{AS}}^\lambda[\dim(Y_\lambda)], \quad L_\lambda^{\mathcal{IW}},$$

where $L_\lambda^{\mathcal{IW}}$ is obtained as the image of the canonical morphism $\Delta_\lambda^{\mathcal{IW}} \rightarrow \nabla_\lambda^{\mathcal{IW}}$. Moreover, the usual convolution product (cf. for instance [BGM⁺19, §2.4]) endows $D_{\mathcal{IW}}(\mathrm{Gr})$ with a right action

$$(-) \star (-) : D_{\mathcal{IW}}(\mathrm{Gr}) \times D_{G[[t]]}(\mathrm{Gr}) \rightarrow D_{\mathcal{IW}}(\mathrm{Gr})$$

of the monoidal category $D_{G[[t]]}(\mathrm{Gr})$.

The following result – which gives another incarnation of the category $\mathrm{Rep}_{\mathbb{k}}(G^\vee)$ – was proved in [ABB⁺05, §2.2.1] in the case where $\mathrm{char}(\mathbb{k}) = 0$, and generalized by the authors of [BGM⁺19] to the case of positive characteristic.

Theorem 3.10. *The functor $\mathcal{F} \mapsto \nabla_\zeta^{\mathcal{IW}} \star \mathcal{F}$ induces an equivalence of categories*

$$\mathrm{Perv}_{G[[t]]}(\mathrm{Gr}) \xrightarrow{\sim} \mathrm{Perv}_{\mathcal{IW}}(\mathrm{Gr}).$$

In the sequel, we will work with the t -exact “averaging functor”

$$\begin{aligned} \mathrm{Av}_{I_{\mathfrak{u}}, \psi} : D_{I_{\mathfrak{u}}}(\mathrm{Gr}) &\rightarrow D_{\mathcal{IW}}(\mathrm{Gr}), \\ \mathrm{Av}_{I_{\mathfrak{u}}, \psi} : D_{I_{\mathfrak{u}}}(\mathrm{Fl}) &\rightarrow D_{\mathcal{IW}}(\mathrm{Fl}). \end{aligned}$$

We refer to [AR21, §3.6-3.8] for the construction of this functor.

Remark 3.11. Compared with [BGM⁺19], we work with the opposite Iwahori subgroup $I_{\mathfrak{u}}^-$ for the equivariance condition in $\mathrm{Perv}_{\mathcal{IW}}(\mathrm{Gr})$, as opposed to $I_{\mathfrak{u}}$ in *loc. cit.*. This condition is imposed by our choice to work with $D_{I_{\mathfrak{u}}}(\mathrm{Gr})$ (as opposed to $D_{I_{\mathfrak{u}}^-}(\mathrm{Gr})$).

3.7. Representations of Frobenius kernels. We define the group schemes

$$\check{\mathbf{G}}_1 := \ker(\mathrm{Fr}), \quad \check{\mathbf{B}}_1 := \ker(\mathrm{Fr}|_{\check{\mathbf{B}}}).$$

Recall (§2.1) that we have identified the character lattice $X^*(\check{\mathbf{T}})$ with \mathbf{Y} . For any $\lambda \in \mathbf{Y}$, we denote by \mathbb{k}_λ the one-dimensional $\check{\mathbf{B}}_1 \check{\mathbf{T}}$ -module induced by λ , and define the associated baby co-Verma module:

$$\widehat{Z}'(\lambda) := \mathrm{ind}_{\check{\mathbf{B}}_1 \check{\mathbf{T}}}^{\check{\mathbf{G}}_1 \check{\mathbf{T}}} \mathbb{k}_\lambda.$$

One can show (cf. [Jan03, Part II, Proposition 9.6]) that each $\widehat{Z}'(\lambda)$ has a simple socle – denoted by $\widehat{L}(\lambda)$ – and that any simple $\check{\mathbf{G}}_1 \check{\mathbf{T}}$ -module is isomorphic to exactly one such $\widehat{L}(\lambda)$. As in [Jan03, Proposition 9.6], we define the baby Verma module

$$\widehat{Z}(\lambda) := \mathrm{coind}_{\check{\mathbf{B}}_1 \check{\mathbf{T}}}^{\check{\mathbf{G}}_1 \check{\mathbf{T}}} \mathbb{k}_\lambda.$$

In the sequel, we will work with twisted modules. For any $w \in W$, we choose a lift $\check{w} \in N_{\check{\mathbf{G}}}(\check{\mathbf{T}})$. Since \check{w} normalises $\check{\mathbf{T}}$ and $\check{\mathbf{G}}_1$ (because the latter is a normal subgroup of

$\check{\mathbf{G}}$), it also normalises $\check{\mathbf{G}}_1\check{\mathbf{T}}$. Therefore we can twist any $\check{\mathbf{G}}_1\check{\mathbf{T}}$ -module M by \dot{w} (cf. [Jan03, §I.2.15]). The resulting module will be denoted by wM .

3.8. Representations of Frobenius kernels through representations of reductive groups.

We briefly recall how one can see $\check{\mathbf{G}}_1\check{\mathbf{T}}$ -modules as $\check{\mathbf{G}}$ -modules with extra structure (cf. [AG03] for the original reference treating the case of quantum groups, and [AR22, §4.1] for our present setting). This identification comes from the following equivalences of categories

$$(3.4) \quad \text{Rep}(\check{\mathbf{G}}_1\check{\mathbf{T}}) \simeq \text{Coh}^{\check{\mathbf{G}}_1\check{\mathbf{T}}}(\text{pt}) \simeq \text{Coh}^{\check{\mathbf{G}} \times \check{\mathbf{T}}^{(1)}}(\check{\mathbf{G}} \times \check{\mathbf{T}}^{(1)}/\check{\mathbf{G}}_1\check{\mathbf{T}}),$$

where $\check{\mathbf{G}}_1\check{\mathbf{T}}$ is seen as a subgroup of $\check{\mathbf{G}} \times \check{\mathbf{T}}^{(1)}$ through the morphism $g \mapsto (g, \text{Fr}(g))$. In fact, the map $(g, t) \mapsto \text{Fr}(g)t^{-1}$ induces an isomorphism

$$\check{\mathbf{G}} \times \check{\mathbf{T}}^{(1)}/\check{\mathbf{G}}_1\check{\mathbf{T}} \simeq \check{\mathbf{G}}^{(1)}.$$

We deduce that the category of $\check{\mathbf{G}}_1\check{\mathbf{T}}$ -modules is equivalent, via (3.4), to the category of $\mathcal{O}(\check{\mathbf{G}}^{(1)})$ -modules of finite type, which are $\check{\mathbf{G}}$ -equivariant and endowed with a \mathbf{Y} -grading compatible with the \mathbf{Y} -grading of $\mathcal{O}(\check{\mathbf{G}}^{(1)})$. Here $\mathcal{O}(\check{\mathbf{G}}^{(1)})$ is endowed with the left regular $\check{\mathbf{G}}$ -representation structure, and the grading is given by

$$\mathcal{O}(\check{\mathbf{G}}^{(1)})_\lambda = \text{Ind}_{\check{\mathbf{T}}^{(1)}}^{\check{\mathbf{G}}^{(1)}}(-\lambda) \quad \forall \lambda \in X^*(\check{\mathbf{T}}^{(1)}) = \mathbf{Y}.$$

The equivalence (3.4) takes a $\check{\mathbf{G}}_1\check{\mathbf{T}}$ -module M to the \mathbf{Y} -graded object whose component associated with $\lambda \in \mathbf{Y}$ is

$$\text{Ind}_{\check{\mathbf{G}}_1\check{\mathbf{T}}}^{\check{\mathbf{G}}} (M \otimes \text{Fr}^*(\mathbb{k}_{\check{\mathbf{T}}}(-\lambda))).$$

It follows from [AR22, Lemma 4.2] that, for $\lambda \in X^*(\check{\mathbf{T}}^{(1)})$, $\mathcal{O}(\check{\mathbf{G}}^{(1)})_\lambda$ is an ind-object of $\text{Rep}(\check{\mathbf{G}}^{(1)})$.

Finally, we spell out how to express twists by elements of W in the category on the right-hand side of (3.4). Let $M \in \text{Rep}(\check{\mathbf{G}}_1\check{\mathbf{T}})$, $w \in W$, and denote by \widetilde{M} the corresponding \mathbf{Y} -graded, $\check{\mathbf{G}}$ -equivariant, $\mathcal{O}(\check{\mathbf{G}}^{(1)})$ -module of finite type. We let ${}^w\widetilde{M}$ be the object such that, for any $\lambda \in \mathbf{Y}$, we have

$${}^w\widetilde{M}_\lambda := M_{w(\lambda)},$$

and such that the structure of $\mathcal{O}(\check{\mathbf{G}}^{(1)})$ -module is obtained by twisting by w the corresponding action on M .

Proposition 3.12. *For any $w \in W$, and $M \in \text{Rep}(\check{\mathbf{G}}_1\check{\mathbf{T}})$, the image of the representation wM through the equivalence (3.4) is canonically isomorphic to ${}^w\widetilde{M}$.*

Proof. Notice that we have canonical isomorphisms

$$\begin{aligned} \text{Ind}_{\check{\mathbf{G}}_1\check{\mathbf{T}}}^{\check{\mathbf{G}}} ({}^wM \otimes \text{Fr}^*(\mathbb{k}_{\check{\mathbf{T}}}(-\lambda))) &\simeq \text{Ind}_{\check{\mathbf{G}}_1\check{\mathbf{T}}}^{\check{\mathbf{G}}} ({}^w(M \otimes \text{Fr}^*(\mathbb{k}_{\check{\mathbf{T}}}(-w(\lambda)))) \\ &\simeq {}^w \left(\text{Ind}_{\check{\mathbf{G}}_1\check{\mathbf{T}}}^{\check{\mathbf{G}}} (M \otimes \text{Fr}^*(\mathbb{k}_{\check{\mathbf{T}}}(-w(\lambda)))) \right), \end{aligned}$$

where the last isomorphism follows from equation (4) in [Jan03, §I.3.5]. Finally, left-translation by \dot{w} induces an isomorphism of $\check{\mathbf{G}}$ -modules

$$\text{Ind}_{\check{\mathbf{G}}_1\check{\mathbf{T}}}^{\check{\mathbf{G}}} (M \otimes \text{Fr}^*(\mathbb{k}_{\check{\mathbf{T}}}(-w(\lambda)))) \simeq {}^w \left(\text{Ind}_{\check{\mathbf{G}}_1\check{\mathbf{T}}}^{\check{\mathbf{G}}} (M \otimes \text{Fr}^*(\mathbb{k}_{\check{\mathbf{T}}}(-w(\lambda)))) \right).$$

One easily deduces the desired result. \square

3.9. The regular perverse sheaf. Let us define the “regular perverse sheaf” \mathcal{R} , following [AR22, §5.2]. For any $\lambda \in \mathbf{Y}^+$, we have by construction a canonical morphism

$$\mathcal{I}_!^\lambda \rightarrow \mathcal{I}_*^\lambda.$$

Since the category $(\text{Perv}_{G[[t]]}(\text{Gr}), \star)$ is *rigid* with unit object \mathcal{IC}^0 and $\mathcal{I}_*^{-w_0(\lambda)} \simeq (\mathcal{I}_!^\lambda)^\vee$ (these facts can be proved directly or using the geometric Satake equivalence), the above morphism induces a morphism

$$(3.5) \quad \mathcal{IC}^0 \rightarrow \mathcal{I}_*^\lambda \star \mathcal{I}_*^{-w_0(\lambda)}.$$

Also notice that, for any $\delta, \delta' \in \mathbf{Y}^+$, the perverse sheaf $\mathcal{I}_*^\delta \star \mathcal{I}_*^{\delta'}$ is supported on $\overline{\text{Gr}}^{\delta+\delta'}$, and its restriction to $\text{Gr}^{\delta+\delta'}$ is $\underline{\mathbb{k}}_{\text{Gr}^{\delta+\delta'}}[\langle 2\rho, \delta + \delta' \rangle]$. Thus, adjunction yields a canonical morphism

$$(3.6) \quad \mathcal{I}_*^\delta \star \mathcal{I}_*^{\delta'} \rightarrow \mathcal{I}_*^{\delta+\delta'}.$$

Let $\mu \in \mathbf{Y}$. We put

$$\mathcal{R}_\mu := \varinjlim_{\lambda \in \mathbf{Y}^+ \cap (-w_0(\mu) + \mathbf{Y}^+)} \mathcal{I}_*^{w_0(\mu)+\lambda} \star \mathcal{I}_*^{-w_0(\lambda)}.$$

The transition morphisms are given by

$$\mathcal{I}_*^{w_0(\mu)+\lambda} \star \mathcal{I}_*^{-w_0(\lambda)} \rightarrow \mathcal{I}_*^{w_0(\mu)+\lambda} \star (\mathcal{I}_*^\nu \star \mathcal{I}_*^{-w_0(\nu)}) \star \mathcal{I}_*^{-w_0(\lambda)} \rightarrow \mathcal{I}_*^{w_0(\mu)+\lambda+\nu} \star \mathcal{I}_*^{-w_0(\lambda+\nu)},$$

where the first, resp. second, morphism is given by (3.5), resp. (3.6).

Proposition 3.13. *For all $\mu \in \mathbf{Y}$, we have an isomorphism of ind-objects*

$$\text{Sat}(\text{sw}^*(\mathcal{R}_\mu)) \simeq \mathcal{O}(\check{\mathbf{G}}^{(1)})_\mu.$$

Proof. This follows from the description of $\mathcal{O}(\check{\mathbf{G}}^{(1)})_\mu$ as an ind-object of $\text{Rep}(\check{\mathbf{G}}^{(1)})$ ([AR22, Lemma 4.2]), together with the fact that $\text{sw}^*(\mathcal{I}_*^\lambda) \simeq \mathcal{I}_*^{-w_0(\lambda)}$ for all $\lambda \in \mathbf{Y}^+$ thanks to Proposition 3.6. \square

We then define the \mathbf{Y} -graded ind-object

$$\mathcal{R} := \bigoplus_{\mu \in \mathbf{Y}} \mathcal{R}_\mu.$$

The morphism (3.5) defines a unit map $\eta : \mathcal{IC}^0 \rightarrow \mathcal{R}_0$, and one can define multiplication morphisms $\mathcal{R}_\mu \star \mathcal{R}_{\mu'} \rightarrow \mathcal{R}_{\mu+\mu'}$ for all $\mu, \mu' \in \mathbf{Y}$, in a similar way to the definition of the transition morphisms above (see [AR22, (5.5)]). These morphisms make \mathcal{R} an algebra object, called the *regular perverse sheaf*.

3.10. Modules over the regular perverse sheaf. Here we follow [AR22, §5.4] (here we only consider the case $A = \emptyset$, in the notation of *loc. cit.*). By definition, a \mathbf{Y} -graded right \mathcal{R} -module

$$\mathcal{F} = \bigoplus_{\lambda \in \mathbf{Y}} \mathcal{F}_\lambda$$

is a \mathbf{Y} -graded ind-object in the category $\text{Perv}_{I_u}(\text{Gr})$, along with a collection of morphisms

$$\mathcal{F}_\mu \star \mathcal{R}_\lambda \rightarrow \mathcal{F}_{\mu+\lambda}$$

for all λ, μ , satisfying obvious associativity and unit axioms. We let

$$\text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$$

denote the category of \mathbf{Y} -graded right \mathcal{R} -modules (which is a full subcategory of the category of \mathbf{Y} -graded ind-objects in $\text{Perv}_{I_u}(\text{Gr})$). This category is abelian. For any $\nu \in \mathbf{Y}$, there is a *shift-of-grading* endofunctor $\langle \nu \rangle$ defined by

$$(\mathcal{F}\langle \nu \rangle)_\lambda := \mathcal{F}_{\lambda-\nu} \quad \forall \lambda \in \mathbf{Y}.$$

Next, we define a functor

$$\Phi : \text{Perv}_{I_u}(\text{Gr}) \rightarrow \text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R}), \quad \mathcal{F} \mapsto \bigoplus_{\mu \in \mathbf{Y}} \mathcal{F} \star \mathcal{R}_\mu.$$

This functor is exact (by exactness of the convolution product) and faithful, since it doesn't kill any nonzero object. Indeed, since the unit map η is injective and \star is exact, we get an injective map $\mathcal{F} \simeq \mathcal{F} \star \mathcal{IC}^0 \rightarrow \mathcal{F} \star \mathcal{R}_0$.

Proposition 3.14. [AR22, Lemma 5.1] *Let $\mathcal{F} \in \text{Perv}_{I_u}(\text{Gr})$, $\mathcal{G} \in \text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$. There is a natural isomorphism*

$$\text{Hom}_{\text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})}(\Phi(\mathcal{F}), \mathcal{G}) \simeq \text{Hom}_{\text{Perv}_{I_u}(\text{Gr})}(\mathcal{F}, \mathcal{G}_0).$$

Since the above result will be used a lot in §8, we recall its proof.

Proof. A morphism $\alpha : \Phi(\mathcal{F}) \rightarrow \mathcal{G}$ is sent to the composition of the morphism $\mathcal{F} \rightarrow \mathcal{F} \star \mathcal{R}_0$ with the degree-zero part $\alpha_0 : \mathcal{F} \star \mathcal{R}_0 = \Phi(\mathcal{F})_0 \rightarrow \mathcal{G}_0$.

Conversely, consider a morphism $\beta : \mathcal{F} \rightarrow \mathcal{G}_0$. For any $\lambda \in \mathbf{Y}$, we construct a morphism $\Phi(\mathcal{F})_\lambda \rightarrow \mathcal{G}_\lambda$ as the composition

$$\mathcal{F} \star \mathcal{R}_\lambda \xrightarrow{\beta \star \text{id}} \mathcal{G}_0 \star \mathcal{R}_\lambda \rightarrow \mathcal{G}_\lambda,$$

where the last morphism comes from the \mathcal{R} -module structure of \mathcal{G} .

One can easily check that the two functors that we have constructed are inverse to each other. \square

An object of $\text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ is called a *free graded \mathcal{R} -module of finite type* if it is isomorphic to a direct sum of shifts of objects of the form $\Phi(\mathcal{F})$, for $\mathcal{F} \in \text{Perv}_{I_u}(\text{Gr})$. An object of $\text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ is said to be *finitely generated* if it is a quotient of a free graded \mathcal{R} -module.

We now describe the simple objects of the abelian category $\text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$. Let $w \in W_{\text{ext}}$. By construction of the set $W_{\text{ext}}^{\text{res}}$, there exist $y \in W_{\text{ext}}^{\text{res}}$, $\lambda \in \mathbf{Y}$ such that $w = yt_\lambda$. We then put

$$\widehat{\mathcal{L}}_w := \Phi(L_y)\langle -\lambda \rangle \in \text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$$

(here the choice of a couple $(y, \lambda) \in W_{\text{ext}}^{\text{res}} \times \mathbf{Y}$ such that $yt_\lambda = w$ is non-unique, but a different choice (y', λ') yields an isomorphic object $\Phi(L_{y'})\langle -\lambda' \rangle$ thanks to the comments in [AR22, §5.5]). We thus have the equality $\widehat{\mathcal{L}}_{yt_\lambda} = \widehat{\mathcal{L}}_y\langle -\lambda \rangle$ for all $y \in W_{\text{ext}}^{\text{res}}$, $\lambda \in \mathbf{Y}$.

Theorem 3.15. [AR22, Theorem 5.6] *The assignment $w \mapsto \widehat{\mathcal{L}}_w$ induces a bijection between W_{ext} and the set of isomorphism classes of simple objects of $\text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$.*

We will denote by $\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ the full subcategory of $\text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ consisting of finite length objects.

Proposition 3.16. [AR22, Theorem 7.9] *An object of $\text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ is finitely generated iff it is of finite length.*

We end this subsection with the crucial *Hecke eigensheaf property* of objects of the category $\text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$.

Proposition 3.17. *Let $\mathcal{F} \in \text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$, $\mathcal{G} \in \text{Rep}(G_{\mathbf{k}}^{\vee})$. We have canonical isomorphisms*

$$\mathcal{F} \star \mathcal{G} \simeq \bigoplus_{\nu \in \mathbf{Y}} \text{Sat}(\text{sw}^* \mathcal{G})(\nu) \otimes \mathcal{F}\langle \nu \rangle \simeq \bigoplus_{\nu \in \mathbf{Y}} \text{Sat}(\mathcal{G})(-w_0(\nu)) \otimes \mathcal{F}\langle \nu \rangle.$$

Proof. Consider the canonical morphism $\mathcal{F} \rightarrow \mathcal{F} \star \mathcal{R}_0$ induced by the unit map. Convolving with \mathcal{G} we get a map $\mathcal{F} \star \mathcal{G} \rightarrow \mathcal{F} \star \mathcal{R}_0 \star \mathcal{G}$. But one can show (by applying the equivalence $(\text{Sat} \circ \text{sw}^*)^{-1}$ and using Proposition 3.13, cf. the proof of [AR22, Lemma 5.2]) that we have a canonical isomorphism

$$\mathcal{R}_0 \star \mathcal{G} \simeq \bigoplus_{\nu} \text{Sat}(\text{sw}^* \mathcal{G})(\nu) \otimes \mathcal{R}_{-\nu}.$$

So we obtain a morphism

$$\mathcal{F} \star \mathcal{G} \rightarrow \bigoplus_{\nu} \text{Sat}(\text{sw}^* \mathcal{G})(\nu) \otimes \mathcal{F} \star \mathcal{R}_{-\nu}.$$

Composing the above morphism with the morphism $\bigoplus_{\nu} \mathcal{F} \star \mathcal{R}_{-\nu} \rightarrow \bigoplus_{\nu} \mathcal{F}\langle \nu \rangle$ coming from the \mathcal{R} -module structure of \mathcal{F} , we obtain a map

$$\mathcal{F} \star \mathcal{G} \rightarrow \bigoplus_{\nu \in \mathbf{Y}} \text{Sat}(\text{sw}^* \mathcal{G})(\nu) \otimes \mathcal{F}\langle \nu \rangle.$$

This morphism is functorial in \mathcal{F} and \mathcal{G} , and satisfies an obvious compatibility with the tensor product and convolution products (cf. [ABB⁺05, §1.3.6]). Therefore, one can easily check (as in the proof of [AG03, Proposition 2.3]) that this morphism is in fact an isomorphism. \square

3.11. Verdier duality. We now recall the definition of the Verdier duality operation on $\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ (cf. [AR22, §9.1]). The goal is to explain the construction of an exact and involutive anti-equivalence $\mathbb{D} : \text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R}) \rightarrow \text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ which satisfies

$$(3.7) \quad \mathbb{D}(\Phi(\mathcal{F})\langle \lambda \rangle) \simeq \Phi(\mathbb{D}(\mathcal{F}))\langle \lambda \rangle \quad \forall \mathcal{F} \in \text{Perv}_{I_u}(\text{Gr}), \lambda \in \mathbf{Y}.$$

In fact, (3.7) allows one to define \mathbb{D} for free graded \mathcal{R} -modules of finite type. This functor can then be extended (although this is not completely obvious) to the whole category $\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$.

We let $\text{Free}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ denote the additive \mathbf{k} -linear category whose objects are formal finite direct sums

$$\bigoplus_{j \in J} (\mathcal{F}_j, \lambda_j),$$

where J is a finite set, and $\mathcal{F}_j \in \text{Perv}_{I_u}(\text{Gr})$, $\lambda_j \in \mathbf{Y}$ for all j . The morphisms are defined by

$$\begin{aligned} \text{Hom}_{\text{Free}_{I_u}^{\mathbf{Y}}(\mathcal{R})} \left(\bigoplus_{j \in J} (\mathcal{F}_j, \lambda_j), \bigoplus_{i \in I} (\mathcal{F}'_i, \lambda_i) \right) &:= \text{Hom}_{\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})} \left(\bigoplus_{j \in J} \Phi(\mathcal{F}_j)\langle \lambda_j \rangle, \bigoplus_{i \in I} \Phi(\mathcal{F}'_i)\langle \lambda_i \rangle \right) \\ &\simeq \bigoplus_{j, i} \text{Hom}_{\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})} (\Phi(\mathcal{F}_j), \Phi(\mathcal{F}'_i)\langle \lambda_i - \lambda_j \rangle) \end{aligned}$$

for an object $\bigoplus_{i \in I} (\mathcal{F}'_i, \lambda_i)$ of $\text{Free}_{I_u}^{\mathbf{Y}}(\mathcal{R})$. We define the exact and fully faithful functor

$$\Phi_2 : \text{Free}_{I_u}^{\mathbf{Y}}(\mathcal{R}) \rightarrow \text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R}), \quad \bigoplus_{j \in J} (\mathcal{F}_j, \lambda_j) \rightarrow \bigoplus_{j \in J} \Phi(\mathcal{F}_j)\langle \lambda_j \rangle.$$

By construction, the essential image of Φ_2 coincides with the free graded \mathcal{R} -modules.

We now define a functor

$$\mathbb{D}_{\text{fr}} : \text{Free}_{I_u}^{\mathbf{Y}}(\mathcal{R}) \rightarrow \text{Free}_{I_u}^{\mathbf{Y}}(\mathcal{R})^{\text{op}},$$

where $(-)^{\text{op}}$ means that we take the opposite category. On objects, it sends $\bigoplus_{j \in J} (\mathcal{F}_j, \lambda_j)$ to $\bigoplus_{j \in J} (\mathbb{D}(\mathcal{F}_j), \lambda_j)$. To describe how it behaves on morphisms, it is enough to treat the case of a morphism between two objects of the form $(\mathcal{F}, \lambda), (\mathcal{G}, \mu)$ by additivity. We will construct a bifunctorial isomorphism

$$\text{Hom}_{\text{Free}_{I_u}^{\mathbf{Y}}(\mathcal{R})}((\mathcal{F}, \lambda), (\mathcal{G}, \mu)) \rightarrow \text{Hom}_{\text{Free}_{I_u}^{\mathbf{Y}}(\mathcal{R})}((\mathbb{D}(\mathcal{G}), \mu), (\mathbb{D}(\mathcal{F}), \lambda)).$$

By definition, the space $\text{Hom}_{\text{Free}_{I_u}^{\mathbf{Y}}(\mathcal{R})}((\mathcal{F}, \lambda), (\mathcal{G}, \mu))$ coincides with the first space below

$$\begin{aligned} \text{Hom}_{\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})}(\Phi(\mathcal{F}), \Phi(\mathcal{G})\langle \mu - \lambda \rangle) &\simeq \text{Hom}_{\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})}(\mathcal{F}, (\Phi(\mathcal{G})\langle \mu - \lambda \rangle)_0) \\ &= \text{Hom}_{\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})}(\mathcal{F}, \mathcal{G} \star \mathcal{R}_{\lambda - \mu}) \\ &= \varinjlim_{\nu} \text{Hom}_{\text{Perv}_{I_u}(\text{Gr})}(\mathcal{F}, \mathcal{G} \star \mathcal{I}_*^{w_0(\lambda - \mu) + \nu} \star \mathcal{I}_*^{-w_0(\nu)}), \end{aligned}$$

where $\nu \in \mathbf{Y}^+ \cap (-w_0(\lambda - \mu) + \mathbf{Y}^+)$. But for any such ν , we have the following isomorphisms

$$\begin{aligned} \text{Hom}_{\text{Perv}_{I_u}(\text{Gr})}(\mathcal{F}, \mathcal{G} \star \mathcal{I}_*^{w_0(\mu - \lambda) + \nu} \star \mathcal{I}_*^{-w_0(\nu)}) &\simeq \text{Hom}_{\text{Perv}_{I_u}(\text{Gr})}(\mathcal{F} \star \mathcal{I}_!^{\lambda - \mu - w_0(\nu)} \star \mathcal{I}_!^{\nu}, \mathcal{G}) \\ &\simeq \text{Hom}_{\text{Perv}_{I_u}(\text{Gr})}(\mathbb{D}(\mathcal{G}), \mathbb{D}(\mathcal{F}) \star \mathcal{I}_*^{\lambda - \mu - w_0(\nu)} \star \mathcal{I}_*^{\nu}), \end{aligned}$$

where the last isomorphism is due to the fact that Verdier duality commutes with the convolution product and sends standard objects to costandard objects. These isomorphisms yield an isomorphism between $\text{Hom}_{\text{Free}_{I_u}^{\mathbf{Y}}(\mathcal{R})}((\mathcal{F}, \lambda), (\mathcal{G}, \mu))$ and

$$(3.8) \quad \varinjlim_{\nu} \text{Hom}_{\text{Perv}_{I_u}(\text{Gr})}(\mathbb{D}(\mathcal{G}), \mathbb{D}(\mathcal{F}) \star \mathcal{I}_*^{\lambda - \mu - w_0(\nu)} \star \mathcal{I}_*^{\nu}).$$

Setting $\nu' := -w_0(\mu - \lambda) + \nu$, we obtain an isomorphism between $\varinjlim_{\nu} \mathcal{I}_*^{\lambda - \mu - w_0(\nu)} \star \mathcal{I}_*^{\nu}$ and $\mathcal{R}_{\mu - \lambda}$. And finally between (3.8) and

$$\begin{aligned} \text{Hom}_{\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})}(\mathbb{D}(\mathcal{G}), (\Phi(\mathbb{D}(\mathcal{F}))\langle \lambda - \mu \rangle)_0) &\simeq \text{Hom}_{\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})}(\Phi(\mathbb{D}(\mathcal{G})), \mathbb{D}(\Phi(\mathcal{F}))\langle \lambda - \mu \rangle) \\ &= \text{Hom}_{\text{Free}_{I_u}^{\mathbf{Y}}(\mathcal{R})}((\mathbb{D}(\mathcal{G}), \mu), (\mathbb{D}(\mathcal{F}), \lambda)). \end{aligned}$$

Lemma 3.18. [AR22, Lemma 9.2] *The functor \mathbb{D}_{fr} is exact.*

We will admit the following fact, which concludes the desired construction and whose full proof is given in [AR22, §9.1].

Proposition 3.19. *There exists an exact and involutive anti-equivalence $\mathbb{D} : \text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R}) \rightarrow \text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ which coincides with (the functor induced by) \mathbb{D}_{fr} on the subcategory of free graded \mathcal{R} -modules.*

Remark 3.20. We can get an “explicit” description of \mathbb{D} . Indeed, any object $\mathcal{F} \in \text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ can be written as the cokernel of a morphism $\alpha : \mathcal{F}_1 \rightarrow \mathcal{F}_2$ between two free graded \mathcal{R} -modules thanks to Proposition 3.16. The object $\mathbb{D}(\mathcal{F})$ then coincides with the kernel of the morphism $\mathbb{D}(\alpha) : \mathbb{D}(\mathcal{F}_2) \rightarrow \mathbb{D}(\mathcal{F}_1)$, whose construction was explained above.

From (3.7) and the definition of the object $\widehat{\mathcal{L}}_w$ (together with the fact that Verdier duality preserves simple objects of the category $\text{Perv}_{I_u}(\text{Gr})$, since those are IC-sheaves), we get that

$$(3.9) \quad \mathbb{D}(\widehat{\mathcal{L}}_w) \simeq \widehat{\mathcal{L}}_w \quad \forall w \in W_{\text{ext}}.$$

3.12. Baby Verma and co-Verma modules. In this subsection we follow [AR22, §5.6], defining the analogue of baby co-Verma modules as ind-objects of $\text{Perv}_{I_u}(\text{Gr})$. For any $\mu \in \mathbf{Y}^+$, the orbit Gr_{t_μ} is open inside Gr^μ (recall that our choice of Borel subgroup B corresponds to the opposite Borel subgroup in [AR22]). So by adjunction, we obtain a canonical morphism

$$(3.10) \quad \mathcal{I}_*^\mu \rightarrow \nabla_\mu.$$

Thanks to Lemma 3.4, we get a canonical isomorphism

$$\mathcal{N}_w \star^I \nabla_\mu \simeq \nabla_{wt_\mu},$$

for all $w \in W_{\text{ext}}^S$, $\mu \in \mathbf{Y}^+$ (notice that $\ell(wt_\mu w_0) = \ell(w) + \ell(t_\mu w_0)$ by Lemma 3.1. since $w \in W_{\text{ext}}^S$). On the other hand, if we denote by $\pi : \text{Fl} \rightarrow \text{Gr}$ the canonical projection, we easily obtain isomorphisms

$$(3.11) \quad \mathcal{N}_w \star^I \mathcal{I}_*^\mu \simeq (\pi_* \mathcal{N}_w) \star \mathcal{I}_*^\mu \simeq \nabla_w \star \mathcal{I}_*^\mu,$$

where the first isomorphism follows from [BGM⁺19, Lemma 2.5], and the second one from the fact that $\pi_* \mathcal{N}_w \simeq \nabla_w$ (see [AR22, (3.4)]).

Thus, we get a canonical morphism

$$(3.12) \quad \nabla_w \star \mathcal{I}_*^\mu \rightarrow \nabla_{wt_\mu}$$

by convolving (3.10) with \mathcal{N}_w .

Next, for $w \in W_{\text{ext}}$, $\mu \in \mathbf{Y}$, we put

$$(\widehat{\mathcal{Z}}'_w)_\mu = \varinjlim_\lambda \nabla_{wt_{\mu+\lambda}} \star \mathcal{I}_*^{-w_0(\lambda)},$$

where λ runs over the elements of \mathbf{Y}^+ such that $wt_{\mu+\lambda} \in W_{\text{ext}}^S$. The transition morphisms in the above inductive limit are given by

$$\nabla_{wt_{\mu+\lambda}} \star \mathcal{I}_*^{-w_0(\lambda)} \rightarrow \nabla_{wt_{\mu+\lambda}} \star \mathcal{I}_*^\nu \star \mathcal{I}_*^{-w_0(\nu)} \star \mathcal{I}_*^{-w_0(\lambda)} \rightarrow \nabla_{wt_{\mu+\lambda+\nu}} \star \mathcal{I}_*^{-w_0(\lambda+\nu)}$$

where the first, resp. second, morphism is obtained via (3.5), resp. (3.6) and (3.12). We then put

$$\widehat{\mathcal{Z}}'_w := \bigoplus_{\mu \in \mathbf{Y}} (\widehat{\mathcal{Z}}'_w)_\mu.$$

The morphisms $(\widehat{\mathcal{Z}}'_w)_\mu \star \mathcal{R}_{\mu'} \rightarrow (\widehat{\mathcal{Z}}'_w)_{\mu+\mu'}$ giving the structure of \mathcal{R} -module are obtained thanks to the morphisms

$$\begin{aligned} (\nabla_{wt_{\mu+\lambda}} \star \mathcal{I}_*^{-w_0(\lambda)}) \star (\mathcal{I}_*^{w_0(\mu')+\lambda'} \star \mathcal{I}_*^{-w_0(\lambda')}) &\rightarrow (\nabla_{wt_{\mu+\lambda}} \star \mathcal{I}_*^{w_0(\mu')+\lambda'} \star \mathcal{I}_*^{-w_0(\lambda'+\lambda)}) \\ &\rightarrow \nabla_{wt_{\mu+\mu'+(\lambda+\lambda')}} \star \mathcal{I}_*^{-w_0(\lambda'+\lambda)}, \end{aligned}$$

where the first morphism is obtained by the commutativity constraint and (3.5), and the second one follows from (3.12).

The next result allows to recover all the co-Verma modules by acting on the one associated with the trivial element 0. This will later allow us to reduce our considerations to the case of $\widehat{\mathcal{Z}}'_0$.

Proposition 3.21. *For any $\gamma \in \mathbf{Y}^+$, $\nu \in \mathbf{Y}$, we have a canonical isomorphism*

$$\mathcal{N}_\gamma \star^I \widehat{\mathcal{Z}}'_\nu \simeq \widehat{\mathcal{Z}}'_{\gamma+\nu}$$

Proof. Recall that, for any $\mu \in \mathbf{Y}$, we have

$$(\widehat{\mathcal{Z}}'_\nu)_\mu = \varinjlim_{\lambda \in \mathbf{Y}^+ - \mu - \nu} \nabla_{\nu+\mu+\lambda} \star \mathcal{I}_*^{-w_0(\lambda)}.$$

Since $\ell(t_\gamma t_{\nu+\mu+\lambda}) = \ell(t_\gamma) + \ell(t_{\mu+\lambda})$, we can apply Lemma 3.4 to get a canonical isomorphism

$$\mathcal{N}_\gamma \star^I \nabla_{\nu+\mu+\lambda} \simeq \nabla_{\gamma+\nu+\mu+\lambda}.$$

Thus, we have a canonical isomorphism

$$\mathcal{N}_\gamma \star^I (\nabla_{\nu+\mu+\lambda} \star \mathcal{I}_*^{-w_0(\lambda)}) \simeq \nabla_{\gamma+\nu+\mu+\lambda} \star \mathcal{I}_*^{-w_0(\lambda)}$$

whenever γ and $\nu + \mu + \lambda$ are dominant. Therefore we obtain a canonical isomorphism

$$\mathcal{N}_\gamma \star^I (\widehat{\mathcal{Z}}'_\nu)_\mu \simeq \varinjlim_\lambda \nabla_{\gamma+\nu+\mu+\lambda} \star \mathcal{I}_*^{-w_0(\lambda)},$$

where λ runs through the set of elements of \mathbf{Y}^+ such that $\gamma + \nu + \mu + \lambda$ is dominant. The fact that the transition morphisms in the above limit coincide with the transition morphisms appearing in the definition of $(\widehat{\mathcal{Z}}'_{\gamma+\nu})_\mu$ follows from the construction of the morphism (3.12). \square

Similarly, one can check the following result.

Proposition 3.22. *For any $w, w' \in W$ such that $\ell(w w' w_0) = \ell(w) + \ell(w' w_0)$ and $\lambda \in \mathbf{Y}$, we have a canonical isomorphism*

$$\mathcal{N}_w \star^I \widehat{\mathcal{Z}}'_{w't_\lambda} \simeq \widehat{\mathcal{Z}}'_{w w' t_\lambda}.$$

Proposition 3.23. [AR22, Corollary 6.9] *For any $w \in W_{\text{ext}}$ the object $\widehat{\mathcal{Z}}'_w$ has finite length and a simple socle, isomorphic to $\widehat{\mathcal{L}}_w$. In particular, $\widehat{\mathcal{Z}}'_w$ is indecomposable.*

We are thus allowed to define the analogue of the baby Verma module

$$\widehat{\mathcal{Z}}_w := \mathbb{D}(\widehat{\mathcal{Z}}'_w) \in \text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$$

for all $w \in W_{\text{ext}}$. Since \mathbb{D} is an (exact) equivalence of categories sending $\widehat{\mathcal{L}}_w$ to $\widehat{\mathcal{L}}_w$, we deduce from Proposition 3.23 that $\widehat{\mathcal{Z}}_w$ is indecomposable and has a simple head, isomorphic to $\widehat{\mathcal{L}}_w$.

Proposition 3.24. [AR22, Proposition 9.25] *For $w, w' \in W_{\text{ext}}$ we have*

$$\text{Ext}_{\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})}^i(\widehat{\mathcal{Z}}_w, \widehat{\mathcal{Z}}_{w'}) = \begin{cases} \mathbb{k} & \text{if } i = 0 \text{ and } w = w', \\ 0 & \text{otherwise.} \end{cases}$$

3.13. Averaging functor. We denote by $\text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ the version of $\text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ where the category $\text{Perv}_{I_u}(\text{Gr})$ is replaced by $\text{Perv}_{I_u}^{\mathcal{W}}(\text{Gr})$. Namely, an object \mathcal{F} of $\text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ is a \mathbf{Y} -graded ind-object in $\text{Perv}_{I_u}^{\mathcal{W}}(\text{Gr})$, endowed with a right-action of \mathcal{R} respecting the grading on \mathcal{R} and \mathcal{F} . There is an exact and faithful functor

$$\Phi^{\mathcal{W}} : \text{Perv}_{I_u}^{\mathcal{W}}(\text{Gr}) \rightarrow \text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R}), \quad \mathcal{F} \mapsto \bigoplus_{\mu \in \mathbf{Y}} \mathcal{F} \star \mathcal{R}_\mu.$$

The above functor satisfies the same type of properties as Φ . Namely, for any $w \in {}^S W_{\text{ext}}$, written as $w = y t_\lambda$ for some $(y, \lambda) \in {}^S W_{\text{ext}}^{\text{res}} \times \mathbf{Y}$, we define

$$\widehat{\mathcal{L}}_w^{\mathcal{W}} := \Phi^{\mathcal{W}}(\mathbb{L}_y^{\mathcal{W}})\langle -\lambda \rangle \in \text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R}),$$

where y is seen as an element of \mathbf{Y}^{++} through the bijection of Corollary 3.3. Then the above object is simple, and the assignment $w \mapsto \widehat{\mathcal{L}}_w^{\mathcal{W}}$ induces a bijection between ${}^S W_{\text{ext}}$ and the set of isomorphism classes of objects of $\text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$. We denote by $\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ the subcategory of objects of finite length. As in §3.12, one can define the baby co-Verma module $\widehat{\mathcal{Z}}_w^{\mathcal{W}}$ attached to an element $w \in {}^S W_{\text{ext}}$, and show that this object admits a unique

simple socle, isomorphic to $\widehat{\mathcal{L}}_w^{\mathcal{I}\mathcal{W}}$. Using the geometric Satake equivalence together with Theorem 3.10 (and the identifications obtained in §3.8), one can check that the category $\text{Mod}_{\mathcal{I}\mathcal{W}}^{\mathbf{Y}}(\mathcal{R})$ identifies with the category of \mathbf{Y} -graded ind-objects of $\text{Rep}(\check{\mathbf{G}}^{(1)})$ endowed with a graded action of $\mathcal{O}(\check{\mathbf{G}}^{(1)})$. We therefore obtain the following result.

Proposition 3.25. [AR22, Proposition 5.7] *The category $\text{Mod}_{\mathcal{I}\mathcal{W}}^{\mathbf{Y}}(\mathcal{R})$ (resp. $\text{mod}_{\mathcal{I}\mathcal{W}}^{\mathbf{Y}}(\mathcal{R})$) is equivalent to the category of algebraic $T_{\mathbb{k}}^{\vee}$ -modules (resp. finite dimensional algebraic $T_{\mathbb{k}}^{\vee}$ -modules). In particular, the category $\text{mod}_{\mathcal{I}\mathcal{W}}^{\mathbf{Y}}(\mathcal{R})$ is semi-simple, and we have isomorphisms*

$$\widehat{\mathcal{Z}}_w^{\mathcal{I}\mathcal{W}} \simeq \widehat{\mathcal{L}}_w^{\mathcal{I}\mathcal{W}} \quad \forall w \in {}^S W_{\text{ext}}.$$

Recall the averaging functor

$$\text{Av}_{I_u, \psi} : \text{Perv}_{I_u}(\text{Gr}) \rightarrow \text{Perv}_{\mathcal{I}\mathcal{W}}(\text{Gr}).$$

This functor induces an exact functor between the abelian categories

$$\text{Av}_{I_u, \psi} : \text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R}) \rightarrow \text{mod}_{\mathcal{I}\mathcal{W}}^{\mathbf{Y}}(\mathcal{R}),$$

and the latter functor commutes with Φ , in the sense that we have a functorial isomorphism

$$\Phi^{\mathcal{I}\mathcal{W}}(\text{Av}_{I_u, \psi}(\mathcal{F})) \simeq \text{Av}_{I_u, \psi}(\Phi(\mathcal{F}))$$

for any $\mathcal{F} \in \text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$. We now describe the behaviour of the averaging functor with respect to costandard (and therefore also standard) objects.

Proposition 3.26. [AR22, Lemma 6.3] *Let $w \in W_{\text{ext}}$, and write it as $w = vu$, with $v \in W$, $u \in {}^S W_{\text{ext}}$. We have*

$$\text{Av}_{I_u, \psi}(\widehat{\mathcal{Z}}'_w) \simeq \widehat{\mathcal{Z}}'_u.$$

3.14. Representations of the Frobenius kernel in terms of \mathcal{R} -modules. The following statement is a consequence of the main results of [AR22], together with the proof of the Finkelberg-Mirković-conjecture in [BR24].

Theorem 3.27. [AR22, Proposition 1.3] *Assume that, for any indecomposable constituent in the root system of $(\check{\mathbf{G}}, \check{\mathbf{T}})$, ℓ is greater than the bound in Figure 1. Then there exists an equivalence of categories*

$$\text{FM}_{\text{Frob}} : \text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R}) \xrightarrow{\sim} \text{Rep}_{[0]}(\check{\mathbf{G}}_1 \check{\mathbf{T}})$$

such that

$$\text{FM}_{\text{Frob}} \circ \Phi \simeq \text{FM} \circ \text{For}_0$$

which satisfies

$$\text{FM}_{\text{Frob}}(\widehat{\mathcal{L}}_w) \simeq \widehat{\mathcal{L}}(w^{-1} \cdot_{\ell} 0), \quad \text{FM}_{\text{Frob}}(\widehat{\mathcal{Z}}'_w) \simeq \widehat{\mathcal{Z}}'(w^{-1} \cdot_{\ell} 0), \quad \text{FM}_{\text{Frob}}(\widehat{\mathcal{Z}}_w) \simeq \widehat{\mathcal{Z}}(w^{-1} \cdot_{\ell} 0)$$

for any $w \in W_{\text{ext}}$, and such that

$$\text{FM}_{\text{Frob}}(\mathcal{F} \star \mathcal{G}) \simeq \text{FM}_{\text{Frob}}(\mathcal{F}) \otimes \text{Fr}^*(\text{Sat}(\text{sw}^* \mathcal{G}))$$

functorially in $\mathcal{F} \in \text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$, $\mathcal{G} \in \text{Perv}_{G[[t]]}(\text{Gr})$.

The next result follows from the constructions of baby Verma and co-Verma modules in terms of coinduced and induced representations respectively.

Proposition 3.28. *There is an isomorphism of $\check{\mathbf{G}}_1 \check{\mathbf{T}}$ -modules*

$$\widehat{\mathcal{Z}}(0) \simeq {}^{w_0} \widehat{\mathcal{Z}}'((w_0 t_{-2\zeta})^{-1} \cdot_{\ell} 0).$$

Proof. We want to establish an isomorphism $\widehat{Z}(0) \simeq {}^{w_0}\widehat{Z}'(2(\ell-1)\zeta)$, or equivalently ${}^{w_0}\widehat{Z}(0) \simeq \widehat{Z}'(2(\ell-1)\zeta)$. By [Jan03, Lemma 9.2, Part II], we have an isomorphism

$$\widehat{Z}(0) \simeq \text{ind}_{\mathbf{B}_1^+ \tilde{\mathbf{T}}}^{\tilde{\mathbf{G}}_1 \tilde{\mathbf{T}}}(-2(\ell-1)\zeta).$$

We deduce that

$${}^{w_0}\widehat{Z}(0) \simeq \text{ind}_{w_0(\mathbf{B}_1^+ \tilde{\mathbf{T}})}^{\tilde{\mathbf{G}}_1 \tilde{\mathbf{T}}}(w_0(-2(\ell-1)\zeta)) = \text{ind}_{\mathbf{B}_1^+ \tilde{\mathbf{T}}}^{\tilde{\mathbf{G}}_1 \tilde{\mathbf{T}}}(2(\ell-1)\zeta) = \widehat{Z}'(2(\ell-1)\zeta),$$

where the first isomorphism is due to [Jan03, (4)-I.3.5]. \square

Remark 3.29. In fact, one can easily deduce from the previous proposition how to express any baby Verma module as a twisted baby co-Verma module (in the same fashion as [ABB⁺05, Corollary 3.2.11]), but only the above case will be useful for us in the sequel. The point is that it allows one to express baby Verma modules as ind-objects of the category $\text{Perv}_{I_u}(\text{Gr})$.

Let $\mathcal{F} \in \text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$, and $w \in W$. We define the object ${}^w\mathcal{F} \in \text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ by putting

$$({}^w\mathcal{F})_\lambda := \mathcal{F}_{w(\lambda)} \quad \forall \lambda \in \mathbf{Y}$$

and defining the right- \mathcal{R} action by twisting by w the corresponding action on \mathcal{F} (the twist of \mathcal{R} by w can be defined via the geometric Satake equivalence).

Corollary 3.30. *Assume that, for any indecomposable constituent in the root system of $(\tilde{\mathbf{G}}, \tilde{\mathbf{T}})$, ℓ is greater than the bound in Figure 1. We have an isomorphism*

$$\widehat{\mathcal{Z}}_0 \simeq {}^{w_0}(\widehat{\mathcal{Z}}'_{w_0 t_{-2\zeta}}).$$

Proof. Thanks to the assumptions, we may use Theorem 3.27 to prove this isomorphism. We have isomorphisms

$$\text{FM}_{\text{Frob}}({}^{w_0}(\widehat{\mathcal{Z}}'_{w_0 t_{-2\zeta}})) \simeq {}^{w_0}\text{FM}_{\text{Frob}}(\widehat{\mathcal{Z}}'_{w_0 t_{-2\zeta}}) \simeq {}^{w_0}\widehat{Z}'((w_0 t_{-2\zeta})^{-1} \cdot \ell 0)$$

thanks to Proposition 3.12 and Theorem 3.27. We can then conclude thanks to Corollary 3.30. \square

4. DRINFELD'S COMPACTIFICATIONS

4.1. Drinfeld's compactification and variants. We put $X := \mathbb{P}^1$. Thanks to [ABB⁺05, §4.1.1], there exists a reductive group \tilde{G} over \mathbb{F} whose derived subgroup $[\tilde{G}, \tilde{G}]$ is simply connected, endowed with a central isogeny $\varphi : \tilde{G} \rightarrow G$ such that $\ker(\varphi)$ is connected. We fix such a pair (\tilde{G}, φ) and consider the maximal torus and Borel subgroup $\tilde{T} := \varphi^{-1}(T)$, $\tilde{B} := \varphi^{-1}(B)$. We also let \tilde{N}^- be the unipotent radical of the Borel subgroup which is opposite to \tilde{B} with respect to \tilde{T} . Notice that, if we denote by $\tilde{\mathbf{X}}^+ \subset \tilde{\mathbf{X}}$ the set of dominant characters and characters associated with the pair (\tilde{T}, \tilde{B}) , then φ induces a morphism $\mathbf{X} \rightarrow \tilde{\mathbf{X}}$ that sends \mathbf{X}^+ to $\tilde{\mathbf{X}}^+$.

We have a diagram

$$\begin{array}{ccc} \text{Bun}_B & \xrightarrow{\mathfrak{p}^+} & \text{Bun}_T \\ \mathfrak{q}^+ \downarrow & & \\ & & \text{Bun}_G \end{array}$$

defined by sending a B -bundle \mathcal{F}_B to $\mathfrak{q}^+(\mathcal{F}_B) := \mathcal{F}_B \times^B G$ (resp. $\mathfrak{p}^+(\mathcal{F}_B) := \mathcal{F}_B \times^B T$). There is an important description of the stack Bun_B , called the Plücker description (cf. [BG02, §1.2.1]). Namely, when the derived subgroup $[G, G]$ is simply connected, Bun_B classifies the data of a principal G -bundle \mathcal{F}_G over $X \times S$, endowed with a B -reduction,

i.e. a principal T -bundle $\mathcal{F}_T \in \text{Bun}_T(S)$ together with a collection of morphisms of vector bundles

$$\kappa^\lambda : \mathcal{L}_{\mathcal{F}_T}^\lambda \rightarrow \mathcal{V}_{\mathcal{F}_G}^\lambda \quad \forall \lambda \in \mathbf{X}^+,$$

which are required to satisfy the Plücker relations. This means that, for $\lambda_1, \lambda_2 \in \mathbf{X}^+$, the morphism

$$\mathcal{L}_{\mathcal{F}_T}^{\lambda_1+\lambda_2} \xrightarrow{\kappa^{\lambda_1+\lambda_2}} \mathcal{V}_{\mathcal{F}_G}^{\lambda_1+\lambda_2} \rightarrow \mathcal{V}_{\mathcal{F}_G}^{\lambda_1} \otimes \mathcal{V}_{\mathcal{F}_G}^{\lambda_2}$$

(where the second arrow is induced by (2.1)) must coincide with the composition

$$\mathcal{L}_{\mathcal{F}_T}^{\lambda_1+\lambda_2} \simeq \mathcal{L}_{\mathcal{F}_T}^{\lambda_1} \otimes \mathcal{L}_{\mathcal{F}_T}^{\lambda_2} \xrightarrow{\kappa^{\lambda_1} \otimes \kappa^{\lambda_2}} \mathcal{V}_{\mathcal{F}_G}^{\lambda_1} \otimes \mathcal{V}_{\mathcal{F}_G}^{\lambda_2}.$$

One can then define the Drinfeld compactification $\overline{\text{Bun}}_B$. Since we want to work with a general group G , we first need to define a larger space $\overline{\text{Bun}}'_B(G)$. It is the functor classifying the data of a principal G -bundle \mathcal{F}_G , endowed with a *degenerate* B -reduction, i.e. a principal T -bundle \mathcal{F}_T together with a collection of morphisms

$$\kappa^\lambda : \mathcal{L}_{\mathcal{F}_T}^\lambda \rightarrow \mathcal{V}_{\mathcal{F}_G}^\lambda \quad \forall \lambda \in \mathbf{X}^+,$$

satisfying the Plücker relations, but which are no longer required to be morphisms of vector bundles. We merely ask that, for every geometric point s of S , the restriction $\kappa^\lambda|_{X \times_s}$ is injective. When $[G, G]$ is simply connected, $\overline{\text{Bun}}'_B(G)$ is the sought for Drinfeld compactification. But in general, $\overline{\text{Bun}}_B$ will be a closed substack of $\overline{\text{Bun}}'_B(G)$.

As in the previous paragraph, one can define the space $\overline{\text{Bun}}'_B(\tilde{G})$ attached to \tilde{G} . There is a morphism of stacks $\overline{\text{Bun}}'_B(\tilde{G}) \rightarrow \overline{\text{Bun}}'_B(G)$, obtained by inducing the bundles via the isogeny $\tilde{G} \rightarrow G$ (the Plücker relations for \tilde{G} induce Plücker relations for G since $\mathbf{X}^+ \subset \tilde{\mathbf{X}}^+$). The same arguments as in the proof of [ABB⁺05, Lemma 4.1.2] can be used to prove the next result.

Proposition 4.1. *The morphism of stacks $\overline{\text{Bun}}'_B(\tilde{G}) \rightarrow \overline{\text{Bun}}'_B(G)$ is a closed embedding, and its image does not depend on the choice of (\tilde{G}, φ) .*

We define $\overline{\text{Bun}}_B$ as the image of the above morphism inside $\overline{\text{Bun}}'_B(G)$. One can show that $\overline{\text{Bun}}_B$ is an algebraic stack in which Bun_B is a dense open substack, and that the canonical projection $\mathfrak{p}^+ : \overline{\text{Bun}}_B \rightarrow \text{Bun}_G$ is proper ([BG02, Proposition 1.2.2]). Notice that the isomorphism $B \simeq B^-$ (obtained by conjugation by w_0) yields an isomorphism of stacks between Bun_B and Bun_{B^-} (resp. $\overline{\text{Bun}}_B$ and $\overline{\text{Bun}}_{B^-}$). We denote by \mathfrak{q} (resp. $\bar{\mathfrak{q}}$) the canonical projection $\text{Bun}_{B^-} \rightarrow \text{Bun}_G$ (resp. $\overline{\text{Bun}}_{B^-} \rightarrow \text{Bun}_G$), and by \mathfrak{p} (resp. $\bar{\mathfrak{p}}$) the canonical projection $\text{Bun}_{B^-} \rightarrow \text{Bun}_T$ (resp. $\overline{\text{Bun}}_{B^-} \rightarrow \text{Bun}_T$).

One can easily check that the connected components of $|\text{Bun}_T|$ are parametrized by \mathbf{Y} : the connected component associated with μ consists in the T -bundles \mathcal{F}_T such that the line bundle $\mathcal{L}_{\mathcal{F}_T}^\lambda$ has degree $-\langle \lambda, \mu \rangle$ for all $\lambda \in \mathbf{X}^+$. Likewise, we have a decomposition into connected components

$$\text{Bun}_B = \bigsqcup_{\mu \in \mathbf{Y}} \text{Bun}_B^\mu,$$

where Bun_B^μ is the preimage through \mathfrak{p}^+ of the connected component of Bun_T associated with μ . We will denote by $\overline{\text{Bun}}_{B^-}^\theta$ the preimage of the connected component associated with $\theta \in \mathbf{Y}$ through $\bar{\mathfrak{p}}$.

Next, we define the stack $\overline{\text{Bun}}_{N^-}$, which is a “compactification” of the canonical morphism $\text{Bun}_{N^-} \rightarrow \text{Bun}_G$ (more precisely, there is a canonical action of T on $\overline{\text{Bun}}_{N^-}$

such that the induced morphism $\overline{\text{Bun}}_{N^-}/T \rightarrow \text{Bun}_G$ is representable and proper, cf. [FGV01, Lemma 2.2.4]). By definition, $\overline{\text{Bun}}_{N^-}$ sits in the Cartesian diagram

$$\begin{array}{ccc} \overline{\text{Bun}}_{N^-} & \longrightarrow & \overline{\text{Bun}}_{B^-} \\ \downarrow & & \downarrow \bar{q} \\ pt & \longrightarrow & \text{Bun}_T, \end{array}$$

where the bottom horizontal arrow is the inclusion of the trivial bundle. Let us spell out this definition. As before, we first need to define a larger space $\overline{\text{Bun}}'_{N^-}(G)$. It classifies the data of a principal G -bundle $\mathcal{F}_G \in \text{Bun}_G(S)$, together with a collection of morphisms of sheaves

$$\kappa^- := (\kappa^{\lambda, -} : \mathcal{V}_{\mathcal{F}_G}^\lambda \rightarrow \mathcal{O}_{X \times S})_{\lambda \in \mathbf{X}^+}$$

satisfying the Plücker relations, and such that the dual of $\kappa^{\lambda, -}|_{X \times s}$ is injective for every geometric point s (for technical reasons which will appear in §4.3, we consider here dual morphisms in the definition). One can also define the analogue stack $\overline{\text{Bun}}'_{\tilde{N}^-}(\tilde{G})$ associated with \tilde{G} , which is an algebraic stack locally of finite type over \mathbb{F} . Consider the natural morphism of stacks $\overline{\text{Bun}}'_{\tilde{N}^-}(\tilde{G}) \rightarrow \overline{\text{Bun}}'_{N^-}(G)$, which is a closed embedding ([ABB⁺05, Lemma 4.1.2]), and define $\overline{\text{Bun}}_{N^-}$ as the image of the latter morphism.

Now, we introduce some variations of $\overline{\text{Bun}}_{N^-}$ (as found in [ABB⁺05, §4.1]). Recall that we have fixed a closed point $x \in X$. For any $\nu \in \mathbf{Y}$, the stack $\leq_\nu \overline{\text{Bun}}'_{N^-}(G)$ (respectively $\leq_\nu \overline{\text{Bun}}_{N^-}$) classifies the data of a principal G -bundle \mathcal{F}_G , together with a collection of morphisms of sheaves

$$\kappa_\nu^- := (\kappa_\nu^{\lambda, -} : \mathcal{V}_{\mathcal{F}_G}^\lambda \rightarrow \mathcal{O}_{X \times S}((x \times S) \cdot \langle \nu, \lambda \rangle))_{\lambda \in \mathbf{X}^+}$$

satisfying the Plücker relations (resp. and such that the divisor of zeros of $\kappa_\nu^{\lambda, -}$ does not contain x for all λ). The stack $\leq_\nu \overline{\text{Bun}}'_{N^-}(G)$ is thus open in $\leq_\nu \overline{\text{Bun}}'_{N^-}(G)$. We also define $\leq_\nu \overline{\text{Bun}}_{N^-}$ to be the smooth open substack of $\leq_\nu \overline{\text{Bun}}'_{N^-}$, where the morphisms of sheaves are required to be morphisms of vector bundles. Next, choose a cocharacter $\nu' \in X_*(\tilde{T})$ which is sent to ν via the isogeny φ . Then one can define the version $\leq_{\nu'} \overline{\text{Bun}}'_{\tilde{N}^-}(\tilde{G})$ attached to \tilde{G} and ν' , and the same arguments as the ones used to prove Proposition 4.1 show that the natural morphism $\leq_{\nu'} \overline{\text{Bun}}'_{\tilde{N}^-}(\tilde{G}) \rightarrow \leq_\nu \overline{\text{Bun}}'_{N^-}(G)$ is a closed embedding. Moreover, its image is easily seen not to depend on the choice of ν' . So we let $\leq_\nu \overline{\text{Bun}}_{N^-}$ be the image of the latter morphism, and put

$$\leq_\nu \overline{\text{Bun}}_{N^-} := \leq_\nu \overline{\text{Bun}}'_{N^-}(G) \cap \leq_\nu \overline{\text{Bun}}_{N^-}, \quad \leq_\nu \overline{\text{Bun}}_{N^-} := \leq_\nu \overline{\text{Bun}}_{N^-}(G) \cap \leq_\nu \overline{\text{Bun}}'_{N^-}.$$

By construction we have

$$(4.1) \quad \leq_\nu \overline{\text{Bun}}_{N^-} = \leq_\nu \overline{\text{Bun}}_{N^-} - \bigcup_{\nu' < \nu} \leq_{\nu'} \overline{\text{Bun}}_{N^-}.$$

Note that all the previous stacks admit a natural action of T : an element t multiplies each $\kappa_\nu^{\lambda, -}$ by $\lambda(t)$, for all $\nu \in \mathbf{Y}$, $\lambda \in \mathbf{X}^+$.

Proposition 4.2. *For any $\nu \in \mathbf{Y}$, there is a canonical morphism $\leq_\nu \overline{\text{Bun}}_{N^-} \rightarrow \overline{\text{Bun}}_{B^-}$ which is smooth.*

Proof. By construction it is equivalent to construct a canonical morphism $\leq_\nu \overline{\text{Bun}}'_{\tilde{N}^-}(\tilde{G}) \rightarrow \overline{\text{Bun}}'_{\tilde{B}^-}(\tilde{G})$ and to check that it is smooth. First recall that there is an isomorphism of stacks

$$\text{Bun}_{\tilde{T}} \simeq (pt/\tilde{T}) \times \text{Pic}(\mathbb{P}^1)^r,$$

where $r := \text{rank}(\tilde{T})$, pt/\tilde{T} is the stack of principal \tilde{T} -bundles, and $\text{Pic}(\mathbb{P}^1)$ is the Picard scheme of \mathbb{P}^1 . Now, we have a Cartesian square (see [FGV01, (2.4)])

$$\begin{array}{ccc} \leq_{\nu} \overline{\text{Bun}}'_{N^-}(\tilde{G}) & \longrightarrow & \overline{\text{Bun}}'_{\tilde{B}^-}(\tilde{G}) \\ \downarrow & & \downarrow \bar{q} \\ pt & \longrightarrow & \text{Bun}_{\tilde{T}}, \end{array}$$

where the bottom arrow is determined by the maps

$$pt \rightarrow pt/\tilde{T}, \quad pt \rightarrow \{\nu\} \subset \mathbf{Y} \simeq \text{Pic}(\mathbb{P}^1)^r.$$

But $pt \rightarrow pt/\tilde{T}$ is smooth and $\text{Pic}(\mathbb{P}^1) \simeq \mathbb{Z}$ is discrete, so the bottom arrow is smooth. \square

We will also need the following fact in the sequel, which follows from [FGV01, Corollary 3.2.4] and says that ${}_{\nu}\text{Bun}_{N^-}$ is topologically “contractible”.

Proposition 4.3. *For any $\nu \in \mathbf{Y}$, the cohomology group*

$$H^k({}_{\nu}\text{Bun}_{N^-}, \mathbb{k}_{\nu\text{Bun}_{N^-}})$$

vanishes when $k \neq 0$, and is isomorphic to \mathbb{k} when $k = 0$.

Proof. By [FGV01, Corollary 3.2.4], there exists a unipotent group N' and a N' -bundle \mathcal{N} over the stack ${}_{\nu}\text{Bun}_{N^-}$. Since N' is unipotent, the forgetful functor

$$D({}_{\nu}\text{Bun}_{N^-}) \simeq D_{N'}(\mathcal{N}) \rightarrow D(\mathcal{N})$$

is fully faithful. Therefore we have an isomorphism

$$H^k({}_{\nu}\text{Bun}_{N^-}) \simeq H^k(\mathcal{N})$$

for all k . By *loc. cit.*, the stack \mathcal{N} is isomorphic to a tower of affine spaces. This concludes the proof. \square

For $\nu_1 \leq \nu_2 \in \mathbf{Y}$, we have a natural closed embedding

$$\leq_{\nu_1} \overline{\text{Bun}}'_{N^-}(G) \hookrightarrow \leq_{\nu_2} \overline{\text{Bun}}'_{N^-}(G),$$

which induces a closed embedding

$$\leq_{\nu_1} \overline{\text{Bun}}_{N^-} \hookrightarrow \leq_{\nu_2} \overline{\text{Bun}}_{N^-}.$$

We define ${}_{\infty}\overline{\text{Bun}}_{N^-}$ as the direct limit (in the category of ind-algebraic stacks over \mathbb{F})

$$\varinjlim_{\nu \in \mathbf{Y}} (\leq_{\nu} \overline{\text{Bun}}_{N^-}).$$

We denote by $\tilde{i}_{<\nu}$, resp. \tilde{i}_{ν} , resp. i_{ν} , the embedding of $\leq_{\nu} \overline{\text{Bun}}_{N^-}$, resp. ${}_{\nu} \overline{\text{Bun}}_{N^-}$, resp. ${}_{\nu}\text{Bun}_{N^-}$, into ${}_{\infty}\overline{\text{Bun}}_{N^-}$.

We will denote by ${}^1\text{Bun}_G$ the stack sending a test scheme S to the groupoid consisting in the data of a principal G -bundle \mathcal{F}_G over $X \times S$, together with a structure of level 1 at x for \mathcal{F}_G . Recall that a structure of level one at x is an isomorphism of sheaves $\mathcal{F}_G|_{x \times S} \simeq \mathcal{F}_G^0|_{x \times S}$. We will denote by ${}^1_{\infty}\overline{\text{Bun}}_{N^-}$ the fibered product ${}_{\infty}\overline{\text{Bun}}_{N^-} \times_{\text{Bun}_G} {}^1\text{Bun}_G$. Similarly, we will let ${}^1_{\leq_{\nu}}\overline{\text{Bun}}_{N^-}$, ${}^1_{\nu}\overline{\text{Bun}}_{N^-}$, ${}^1_{\nu}\text{Bun}_{N^-}$ be the corresponding stacks obtained by base change, and still denote by $\tilde{i}_{<\nu}$, \tilde{i}_{ν} , i_{ν} the inclusion of these stacks in ${}^1_{\infty}\overline{\text{Bun}}_{N^-}$. In the sequel, we will consider the diagonal T -action on these stacks, induced by the previously defined T -action together with the action induced by modifying the structure of level 1.

Remark 4.4. In [ABB⁺05], there are versions of the above stacks which are labelled by a non-negative integer k , to indicate a structure of level k at x for \mathcal{F}_G . Here, we only treat the cases where $k \in \{0, 1\}$. In the sequel we will sometimes adopt the notation ${}^0_\infty \overline{\text{Bun}}_{N^-} := {}_\infty \overline{\text{Bun}}_{N^-}$.

4.2. Colored effective divisors. Given $\mu = \sum_{i \in \check{\mathfrak{X}}_s} n_i i \in \mathbf{Y}^{pos}$, we consider as in [FM99] the space $X^\mu := \prod_{i \in \check{\mathfrak{X}}_s} X^{(n_i)}$ of colored effective divisors of multidegree μ , where $X^{(n_i)}$ denotes the n_i -th symmetric power of X . We will write an element $D = (D_i)_{i \in \check{\mathfrak{X}}_s} \in X^\mu$ as $D = \sum_{i \in \check{\mathfrak{X}}_s} i \cdot D_i$ and, for any $\lambda \in \Lambda$, define the divisor $\langle \lambda, D \rangle := \sum_{i \in \check{\mathfrak{X}}_s} \langle \lambda, i \rangle D_i$. A partition of μ is a decomposition $\mu = \gamma_1 + \cdots + \gamma_m$ into a sum of non-zero elements $\gamma_1, \dots, \gamma_m$ of \mathbf{Y}^{pos} , which can be seen as an element $\{\{\gamma_1, \dots, \gamma_m\}\}$ of the m -th symmetric power of \mathbf{Y}^{pos} . We will denote by $\mathfrak{A}(\mu)$ the set of partitions of μ .

For each $\Gamma = \{\{\gamma_1, \dots, \gamma_m\}\} \in \mathfrak{A}(\mu)$, we will define the locally closed stratum $X^\mu_\Gamma \subset X^\mu$. First, we define ${}'X^\mu_\Gamma$ as the subset of $X^{\gamma_1} \times \cdots \times X^{\gamma_m}$ consisting of tuples (x_1, \dots, x_m) such that

- (1) for each k , all the components of x_k in X^{γ_k} are equal, meaning that if one writes $\gamma_k = \sum_{i \in \check{\mathfrak{X}}_s} n'_i i$, $x_k = (x_k^j)_{j \in I}$, then there exists some $y_k \in X$ such that

$$x_k^j = \underbrace{(y_k, \dots, y_k)}_{n'_i} \in X^{(n'_i)} \quad \forall i \in \check{\mathfrak{X}}_s.$$

- (2) we have $y_i \neq y_j$ for $i \neq j$.

The locally closed subvariety $X^\mu_\Gamma \subset X^\mu$ is then defined as the image of the canonical morphism ${}'X^\mu_\Gamma \rightarrow X^\mu$, obtained by quotienting via the action of the symmetric group. We thus obtain a stratification $X^\mu = \sqcup_{\Gamma \in \mathfrak{A}(\mu)} X^\mu_\Gamma$.

We also need to introduce the moduli space description of X^μ (cf. [BFGM02, §2.1]). By definition, X^μ classifies the data of a principal T -bundle \mathcal{F}_T , together with a collection of non-zero morphisms of sheaves $d_\lambda : \mathcal{L}_{\mathcal{F}_T}^\lambda \rightarrow \mathcal{O}_{X \times S}$ for all $\lambda \in \mathbf{X}^+$, such that for every geometric point s of S , the divisor of zeros of $d_\lambda|_{X \times s}$ has degree $\langle \lambda, \mu \rangle$. Similarly we define ${}_{\leq \nu} X^\mu$, where this time the second data is a collection of non-zero morphisms of sheaves $\mathcal{L}_{\mathcal{F}_T}^\lambda \rightarrow \mathcal{O}_{X \times s}(x \cdot \langle \lambda, \nu \rangle)$, such that the divisor of zeros of the restriction to $X \times s$ has degree $\langle \lambda, \mu \rangle$. We then define the ind-scheme ${}_\infty X^\mu$ as a direct limit.

4.3. Zastava spaces. For $\mu \in \mathbf{Y}$, we define various Zastava spaces of degree μ , as displayed in [ABB⁺05, §4.1.4]. We let ${}_{\leq \nu} Z^\mu$ (resp. ${}^1_{\leq \nu} Z^\mu$) be the open substack of

$${}_{\leq \nu} \overline{\text{Bun}}_{N^-} \times_{\text{Bun}_G} \text{Bun}_B^\mu \quad (\text{resp. } {}^1_{\leq \nu} \overline{\text{Bun}}_{N^-} \times_{\text{Bun}_G} \text{Bun}_B^\mu)$$

defined by the condition that the reductions to N^- and B are transversal. This means that for all $\lambda \in \mathbf{X}^+$, the composition

$$\kappa_\nu^{\lambda, -} \circ \kappa^\lambda : \mathcal{L}_{\mathcal{F}_T}^\lambda \rightarrow \mathcal{O}_X(x \cdot \langle \nu, \lambda \rangle)$$

is non-zero. We also define ${}_\nu Z^\mu$ (resp. ${}^1_\nu Z^\mu$) by replacing $\leq \nu$ with ν in the previous definition, and put $Z^\mu := {}_{\leq 0} Z^\mu$. For $\nu_1 \leq \nu_2$, we have a canonical closed embedding ${}_{\leq \nu_1} Z^\mu \hookrightarrow {}_{\leq \nu_2} Z^\mu$ (resp. ${}^1_{\leq \nu_1} Z^\mu \hookrightarrow {}^1_{\leq \nu_2} Z^\mu$), so that we can let ${}_\infty Z^\mu$ (resp. ${}^1_\infty Z^\mu$) be the direct limit (in the category of ind-algebraic stacks over \mathbb{F})

$$\varinjlim_{\nu \in \mathbf{Y}} {}_{\leq \nu} Z^\mu \quad (\text{resp. } \varinjlim_{\nu \in \mathbf{Y}} {}^1_{\leq \nu} Z^\mu).$$

In the sequel, we will often have statements involving ${}^k_\nu Z^\mu$, for $\nu \in \{\nu, \leq \nu, \infty\}$, $k \in \{0, 1\}$, making the convenient identification ${}^0_\nu Z^\mu := {}_\nu Z^\mu$. Finally, we will denote by ${}^k \mathbf{p}^\mu : {}^k_\infty Z^\mu \rightarrow {}^k_\infty \overline{\text{Bun}}_{N^-}$ the canonical projection.

One of the main features of these spaces is their factorization property, which we now explain. First notice that, thanks to the moduli space description of ${}_{\leq \nu} X^\mu$ (§4.2), we have canonical projections ${}_{\leq \nu} \mathfrak{s}^\mu$ and ${}_\infty \mathfrak{s}^\mu$ from ${}_{\leq \nu} Z^\mu$ and ${}_\infty Z^\mu$ to ${}_{\leq \nu} X^\mu$ and ${}_\infty X^\mu$ respectively. We will denote by ${}^1_\infty \mathfrak{s}^\mu : {}^1_\infty Z^\mu \rightarrow {}_\infty X^\mu$ the composition of ${}_\infty \mathfrak{s}^\mu$ with the canonical projection ${}^1_\infty Z^\mu \rightarrow {}_\infty Z^\mu$, and similarly for ${}^1_{\leq \nu} \mathfrak{s}^\mu$. We put $\overset{\circ}{X} := X - \{x\}$, and let

$$\overset{\circ}{X}^\mu$$

be the corresponding open subscheme of X^μ . For $\mu_1, \mu_2 \in \mathbf{Y}$, we denote by

$$(\overset{\circ}{X}^{\mu_1} \times X^{\mu_2})_{\text{disj}}$$

the open substack of the product $\overset{\circ}{X}^{\mu_1} \times X^{\mu_2}$ consisting of pairs (x_1, x_2) such that x_1 and x_2 have no point in common.

Proposition 4.5. [BFGM02, Lemma 4.1.5] *For $\mu_1 + \mu_2 = \mu$ and $k \in \{0, 1\}$, there exists a canonical isomorphism of stacks*

$$(4.2) \quad {}^k_\infty Z^\mu \times_{{}_\infty X^\mu} (\overset{\circ}{X}^{\mu_1} \times X^{\mu_2})_{\text{disj}} \simeq (Z^{\mu_1} \times {}^k_\infty Z^{\mu_2}) \times_{(X^{\mu_1} \times {}_\infty X^{\mu_2})} (\overset{\circ}{X}^{\mu_1} \times X^{\mu_2})_{\text{disj}}.$$

The following result is used implicitly in [ABB⁺05].

Proposition 4.6. [BFGM02, §3.6] *Let $\nu \in \mathbf{Y}$, and U be an open substack of finite type of ${}^k_{\leq \nu} \overline{\text{Bun}}_{N^-}$ (resp. ${}^k_\nu \overline{\text{Bun}}_{N^-}$). There exists $\mu \in \mathbf{Y}$ large enough so that U belongs to the image of the canonical projection ${}^k_{\leq \nu} Z^\mu \rightarrow {}^k_{\leq \nu} \overline{\text{Bun}}_{N^-}$ (resp. ${}^k_\nu Z^\mu \rightarrow {}^k_\nu \overline{\text{Bun}}_{N^-}$).*

4.4. Canonical torsors. Let $(\mathcal{F}_G, \kappa_0)$ be an object of ${}^1_0 \overline{\text{Bun}}_{N^-}(S)$. By definition, the restriction $(\mathcal{F}_G|_{\{x\}}, \kappa_0|_{\{x\}})$ from X to $\{x\}$ produces an N^- -bundle over S . We introduce the stack ${}^1_0 \mathcal{N}$ as classifying the triples $(\mathcal{F}_G, \kappa_0, \Phi)$, where Φ is a trivialization of the N^- -bundle $(\mathcal{F}_G|_{\{x\}}, \kappa_0|_{\{x\}})$. The canonical projection ${}^1_0 \mathcal{N} \rightarrow {}^1_0 \overline{\text{Bun}}_{N^-}$ makes ${}^1_0 \mathcal{N}$ an N^- -torsor over ${}^1_0 \overline{\text{Bun}}_{N^-}$. A similar construction (cf. [FGV01, §3.2.6]) produces an N^- -torsor ${}^1_\nu \mathcal{N}$ over ${}^1_\nu \overline{\text{Bun}}_{N^-}$. It follows directly from the constructions (and [ABB⁺05, (27)]) that we have a canonical isomorphism

$$(4.3) \quad {}^1_\nu \overline{\text{Bun}}_{N^-} \simeq G \times^{N^-} {}^1_\nu \mathcal{N},$$

and that the T -action on ${}^1_\nu \overline{\text{Bun}}_{N^-}$ corresponds to the diagonal action on the right-hand side. The above isomorphism induces morphisms

$$(4.4) \quad \overline{\text{ev}}_\nu : {}^1_\nu \overline{\text{Bun}}_{N^-} \rightarrow G/N^-, \quad \text{ev}_\nu : {}^1_\nu \text{Bun}_{N^-} \rightarrow G/N^-.$$

where ev_ν is obtained by restricting $\overline{\text{ev}}_\nu$.

Let $\mu \in \mathbf{Y}$. The isomorphism (4.3) is compatible with the definition of the Zastava space ${}^1_\nu Z^\mu$. Namely, if we let ${}^1_\nu \tilde{\mathcal{N}}$ be the open substack of

$${}^1_\nu \mathcal{N} \times_{\text{Bun}_G} \text{Bun}_B^\mu$$

defined by the transversality condition (cf. §4.3), then we have an isomorphism

$$(4.5) \quad {}^1_\nu Z^\mu \simeq G \times^{N^-} {}^1_\nu \tilde{\mathcal{N}}.$$

Let $\bar{x}' := \{x'_1, \dots, x'_m\}$ be a collection of points of $X - \{x\}$, and denote by ${}_\infty \overline{\text{Bun}}_{N^-}^{\text{n.z.}\bar{x}'}$ the open substack of ${}_\infty \overline{\text{Bun}}_{N^-}$ defined by the condition that the divisor of zeros of the morphisms κ^λ is disjoint from \bar{x}' , for all $\lambda \in \mathbf{X}^+$. For $i = 1, \dots, m$, we denote by t'_i a local coordinate on X at x'_i . Using the same ideas as in the previous paragraph, one can then construct an $\prod_{i=1}^m N^-[[t'_i]]$ -torsor $\mathcal{N}^{\bar{x}'}$ over ${}_\infty \overline{\text{Bun}}_{N^-}^{\text{n.z.}\bar{x}'}$. Following the arguments of [FGV01, Lemma 3.2.7], one can show that the action of $\prod_{i=1}^m N^-[[t'_i]]$ on $\mathcal{N}^{\bar{x}'}$ extends to an action of $\prod_{i=1}^m N^-((t'_i))$.

We will denote by ${}^1_\infty\overline{\text{Bun}}_{N^-}^{\text{n.z.}\overline{x}'}$, ${}^1\mathcal{N}^{\overline{x}'}$ the corresponding stacks obtained by base change through the map ${}^1\text{Bun}_G \rightarrow \text{Bun}_G$.

5. PERVERSE SHEAVES ON THE SEMI-INFINITE FLAG VARIETY

In this section, we recall the construction of certain categories of perverse sheaves on (some avatar of) the semi-infinite flag manifold. This construction comes from [ABB⁺05] in the case where $\text{char}(\mathbb{k}) = 0$, but the definitions also make sense for arbitrary characteristic.

5.1. Main definition. Let $\mu \in \mathbf{Y}$ and $k \in \{0, 1\}$. Notice that, by [BFGM02, Lemma 3.7], the morphism $\text{Bun}_B^\mu \rightarrow \text{Bun}_G$ is smooth whenever $\langle \alpha, \mu \rangle > -2$ for all positive root α (or more generally when $\langle \alpha, \mu \rangle > 2g - 2$ for all positive root α when X is a general smooth projective curve of genus g). Thus, the morphism ${}^k\mathfrak{p}^\mu : {}^k_\infty Z^\mu \rightarrow {}^k_\infty\overline{\text{Bun}}_{N^-}$ from §4.3 is smooth when μ satisfies the previous condition. For all $k \in \{0, 1\}$, $\mu_1, \mu_2 \in \mathbf{Y}^{\text{pos}}$ such that $\mu = \mu_1 + \mu_2$, with $\langle \alpha, \mu_i \rangle > -2$, for $i = 1, 2$ and all positive root α , we let ${}^k s_{1,2}$ (resp. ${}^k t_{1,2}$) denote the open inclusion of the left-hand side (resp. right-hand side) of (4.2) into ${}^k_\infty Z^\mu$ (resp. $Z^{\mu_1} \times {}^k_\infty Z^{\mu_2}$). Also notice that, since the group $\prod_{i=1}^m N^-((t'_i))$ is an ind-pro-unipotent group scheme, one can define the equivariance condition for perverse sheaves on ${}^k\mathcal{N}^{\overline{x}'}$ with respect to the action of $\prod_{i=1}^m N^-((t'_i))$ (by writting the latter group as a direct limit of closed subgroup schemes, and imposing the equivariance condition for each subgroup, cf. for instance [Gai18, §1.2.1]).

We now define the category $\text{Perv}_{G^k}(\mathcal{Fl}^{\frac{\infty}{2}})$. By definition, it is the full subcategory of $\text{Perv}_T({}^k_\infty\overline{\text{Bun}}_{N^-})$ consisting of objects \mathcal{F} satisfying the following two conditions:

- (1) (Factorization) For $\mu = \mu_1 + \mu_2$ with $\langle \alpha, \mu_i \rangle > 2g - 2$ for all positive root α , the complex $({}^k s_{1,2})^*({}^k\mathfrak{p}^\mu)^*\mathcal{F}$ is isomorphic (up to a shift by the corresponding relative dimensions), under the isomorphism (4.2), to the complex

$$({}^k t_{1,2})^*(\text{IC}_{Z^{\mu_1}} \boxtimes ({}^k\mathfrak{p}^{\mu_2})^*\mathcal{F}).$$

- (2) (Equivariance) For all m -tuple \overline{x}' of $X - \{x\}$, the pullback of \mathcal{F} under the composition

$${}^k\mathcal{N}^{\overline{x}'} \rightarrow {}^k_\infty\overline{\text{Bun}}_{N^-}^{\text{n.z.}\overline{x}'} \rightarrow {}^k_\infty\overline{\text{Bun}}_{N^-}$$

is $\prod_{i=1}^m N^-((t'_i))$ -equivariant.

Remark 5.1. Compared with the definition of [ABB⁺05, §4.2.1], condition (3) is missing. When $k = 1$ (which is our main case of interest), this condition is automatic (see Remark 4.2.2 in *loc. cit.*). When $k = 0$, we shall see that condition (3) is also automatic, since the obtained category turns out to be – with or without condition (3) – the Serre subcategory of $\text{Perv}_T({}^k_\infty\overline{\text{Bun}}_{N^-})$ generated by the IC-sheaves on each $\leq_\nu \overline{\text{Bun}}_{N^-}$, for $\nu \in \mathbf{Y}$.

We omit the proofs of the next two results, as one can copy the arguments of *loc. cit.*.

Proposition 5.2. [ABB⁺05, Proposition 4.2.8] *The equivariance condition is implied by the factorization condition.*

Proposition 5.3. [ABB⁺05, Corollary 4.2.9] *Let $k \in \{0, 1\}$. As a subcategory of the category $\text{Perv}_T({}^k_\infty\overline{\text{Bun}}_{N^-})$, $\text{Perv}_{G^k}(\mathcal{Fl}^{\frac{\infty}{2}})$ is stable under extensions.*

5.2. Simple objects. We will now see that the structure of the category $\text{Perv}_{G[[t]]}(\mathcal{Fl}^{\frac{\infty}{2}})$ does *not* depend on the characteristic of \mathbb{k} (at least when Corollary 5.10 holds). Let $\nu \in \mathbf{Y}$, and $\mathcal{F} \in \text{Perv}_T(G/N^-)$ (the action of T is induced by the right action on G by translation). The isomorphism (4.3) allows us to consider the external twisted tensor product

$$(5.1) \quad \mathcal{F} \widetilde{\boxtimes} \text{IC}_{\nu\overline{\text{Bun}}_{N^-}} \in D_T({}^1_\nu\overline{\text{Bun}}_{N^-}),$$

defined as the perversely shifted pullback of $\mathcal{F} \boxtimes \mathrm{IC}_{\nu \overline{\mathrm{Bun}}_{N^-}} \in \mathrm{Perv}_T(G/N^- \times \nu \overline{\mathrm{Bun}}_{N^-})$ through the canonical projection $G \times^{N^-} {}^1\nu\mathcal{N} \rightarrow G/N^- \times \nu \overline{\mathrm{Bun}}_{N^-}$. In particular, the object (5.1) is perverse.

Proposition 5.4. [ABB⁺05, Proposition 4.2.4]

(1) For any $\mathcal{F} \in \mathrm{Perv}_T(G/N^-)$ and $\nu \in \mathbf{Y}$, the object

$$(\tilde{i}_\nu)_! (\mathcal{F} \boxtimes \mathrm{IC}_{\nu \overline{\mathrm{Bun}}_{N^-}}) \quad (\text{resp. } (\tilde{i}_\nu)_! (\mathrm{IC}_{\nu \overline{\mathrm{Bun}}_{N^-}}))$$

is an object of $\mathrm{Perv}_{G^1}(\mathcal{Fl}^{\frac{\infty}{2}})$ (resp. of $\mathrm{Perv}_{G[[t]]}(\mathcal{Fl}^{\frac{\infty}{2}})$).

(2) For any $\mathcal{G} \in \mathrm{Perv}_{G^1}(\mathcal{Fl}^{\frac{\infty}{2}})$ (resp. $\mathcal{G} \in \mathrm{Perv}_{G[[t]]}(\mathcal{Fl}^{\frac{\infty}{2}})$) and $\nu \in \mathbf{Y}$, the perverse cohomologies of $(\tilde{i}_\nu)^* \mathcal{G}$ are of the form

$$\mathcal{F} \boxtimes \mathrm{IC}_{\nu \overline{\mathrm{Bun}}_{N^-}} \quad (\text{resp. } \bigoplus_{i=1}^m \mathrm{IC}_{\nu \overline{\mathrm{Bun}}_{N^-}})$$

for some $\mathcal{F} \in \mathrm{Perv}_T(G/N^-)$ (resp. $m \geq 1$).

Proof. Let $\mu, \nu \in \mathbf{Y}$, with $\langle \alpha, \mu \rangle > 2g - 2$ for all positive root α . First notice that, by considering the base change of ${}^k\mathfrak{p}^\mu : {}^k_\infty Z^\mu \rightarrow {}^k_\infty \overline{\mathrm{Bun}}_{N^-}$ (resp. ${}^k\mathcal{N}^{\mathfrak{x}'} \rightarrow {}^k_\infty \overline{\mathrm{Bun}}_{N^-}$) through ${}^k_\nu \overline{\mathrm{Bun}}_{N^-} \rightarrow {}^k_\infty \overline{\mathrm{Bun}}_{N^-}$, one can define the factorization and equivariance conditions for objects of $\mathrm{Perv}_T({}^k_\nu \overline{\mathrm{Bun}}_{N^-})$. The same observation applies for objects of $\mathrm{Perv}_{G^1}({}^1_\nu \overline{\mathrm{Bun}}_{N^-})$ (this will be useful at the end of this proof). We denote by $\mathrm{Perv}_{G^1}({}^1_\nu \overline{\mathrm{Bun}}_{N^-})$ (resp. $\mathrm{Perv}_{G[[t]]}({}^1_\nu \overline{\mathrm{Bun}}_{N^-})$) the category of objects satisfying these two conditions. Remark that the functor $(\tilde{i}_\nu)_!$ sends $\mathrm{Perv}_{G^1}({}^1_\nu \overline{\mathrm{Bun}}_{N^-})$ (resp. $\mathrm{Perv}_{G[[t]]}({}^1_\nu \overline{\mathrm{Bun}}_{N^-})$) to $\mathrm{Perv}_{G^1}(\mathcal{Fl}^{\frac{\infty}{2}})$ (resp. $\mathrm{Perv}_{G[[t]]}(\mathcal{Fl}^{\frac{\infty}{2}})$). So we only need to check that $\mathcal{F} \boxtimes \mathrm{IC}_{\nu \overline{\mathrm{Bun}}_{N^-}}$ (resp. $\mathrm{IC}_{\nu \overline{\mathrm{Bun}}_{N^-}}$) satisfies the desired properties to prove the first point.

The fact that $\mathcal{F} \boxtimes \mathrm{IC}_{\nu \overline{\mathrm{Bun}}_{N^-}}$ (resp. $\mathrm{IC}_{\nu \overline{\mathrm{Bun}}_{N^-}}$) satisfies the equivariance condition is obvious. The factorization condition follows from the fact that the projection ${}^k_\nu Z^\mu \rightarrow {}^k_\nu \overline{\mathrm{Bun}}_{N^-}$ is smooth by the assumption on μ . Indeed, when $k = 0$, we deduce that the pullback of $\mathrm{IC}_{\nu \overline{\mathrm{Bun}}_{N^-}}$ through the morphism ${}^0_\nu Z^\mu \rightarrow {}^0_\nu \overline{\mathrm{Bun}}_{N^-}$ is an *IC*-sheaf, so the factorization property follows from the fact that the external tensor product of two *IC*-sheaves is an *IC*-sheaf (cf. [Ach21, Lemma 3.3.14]). Similarly when $k = 1$, the pullback of $\mathcal{F} \boxtimes \mathrm{IC}_{\nu \overline{\mathrm{Bun}}_{N^-}}$ through ${}^1_\nu Z^\mu \rightarrow {}^1_\nu \overline{\mathrm{Bun}}_{N^-}$ identifies with $\mathcal{F} \boxtimes \mathrm{IC}_{\nu Z^\mu}$, so that the factorization property again follows from the fact that any *IC*-sheaf verifies it.

Now we prove the second point. The factorization and equivariance properties are compatible with respect to restriction to $\nu \overline{\mathrm{Bun}}_{N^-}$. So it suffices to check that the functor $\mathcal{F} \mapsto \mathcal{F} \boxtimes \mathrm{IC}_{\nu \overline{\mathrm{Bun}}_{N^-}}$ defines equivalences of categories

$$\begin{aligned} \mathrm{Perv}(pt) &\xrightarrow{\sim} \mathrm{Perv}_{G[[t]]}({}^1_\nu \overline{\mathrm{Bun}}_{N^-}) \\ \mathrm{Perv}_T(G/N^-) &\xrightarrow{\sim} \mathrm{Perv}_{G^1}({}^1_\nu \overline{\mathrm{Bun}}_{N^-}). \end{aligned}$$

We only deal with the second case, since the same arguments can be adapted for the first one. First, we claim that every object $\mathcal{F}' \in \mathrm{Perv}_{G^1}({}^1_\nu \overline{\mathrm{Bun}}_{N^-})$ is the Goresky-Macpherson extension of its restriction to ${}^1_\nu \overline{\mathrm{Bun}}_{N^-}$. Indeed, assume that it is not, then we can find an open subscheme of finite type U inside the support of \mathcal{F}' such that either $*$ or $!$ -restriction of $\mathcal{F}'|_U$ to $({}^1_\nu \overline{\mathrm{Bun}}_{N^-} - {}^1_\nu \overline{\mathrm{Bun}}_{N^-}) \cap U$ does *not* lives in negative (resp. positive) perverse cohomological degrees (cf. for instance the proof of [Ach21, Lemma 3.3.3]). But thanks to Proposition 4.6, we can find $\mu_1 \in \mathbf{Y}$ large enough so that U belongs to the image of ${}^1_\nu Z^{\mu_1} \rightarrow {}^1_\nu \overline{\mathrm{Bun}}_{N^-}$. Therefore, using the identification (4.2), we deduce that we can find

$\mu_2 \in \mathbf{Y}$ large enough (for instance $\mu_2 = \mu_1$) such that, for $\mu = \mu_1 + \mu_2$, either $*$ or $!$ -restriction (of the pullback of $\mathcal{F}'|_U$ through the map ${}^1_\nu Z^{\mu_1 + \mu_2} \rightarrow {}^1_\nu \overline{\text{Bun}}_{N^-}$) on

$$((Z^{\mu_1} - \overset{\circ}{Z}^{\mu_1}) \times {}^1_\nu Z^{\mu_2}) \times_{(X^{\mu_1} \times_\infty X^{\mu_2})} (\overset{\circ}{X}^{\mu_1} \times X^{\mu_2})_{\text{disj}}$$

does not live in negative (resp. positive) perverse cohomological degrees. However, this contradicts the factorization property.

So we are now reduced (thanks to the fact that Goresky-Macpherson extension is a fully faithful functor) to showing that the functor $\mathcal{F} \mapsto \mathcal{F} \boxtimes \text{IC}_{\nu \text{Bun}_{N^-}}$ induces an equivalence of categories

$$\text{Perv}_T(G/N^-) \xrightarrow{\sim} \text{Perv}_{G^1}({}^1_\nu \text{Bun}_{N^-}).$$

Since ${}^1_\nu \text{Bun}_{N^-}$ is smooth, this functor coincides with the pullback via $\text{ev}_\nu : {}^1_\nu \text{Bun}_{N^-} \rightarrow G/N^-$, so that full faithfulness follows from the fact that ev_ν is smooth surjective with connected fibers. To prove the essential surjectivity, the argument of *loc. cit.* is the following. Let $\mathcal{F}' \in \text{Perv}_{G^1}({}^1_\nu \text{Bun}_{N^-})$. For any non-empty collection of points \bar{x}' distinct from x , the pullback of \mathcal{F}' through the map

$${}^1_\nu \text{Bun}_{N^-} \times_{{}^1_\nu \overline{\text{Bun}}_{N^-}} {}^1\mathcal{N}^{\bar{x}'} \rightarrow {}^1_\nu \text{Bun}_{N^-}$$

is $\prod_{i=1}^m N^-((t'_i))$ -equivariant by definition. The assertion then follows from the fact that the latter ind-group-scheme acts transitively along the fibers of the map obtained by composition

$${}^1_\nu \text{Bun}_{N^-} \times_{{}^1_\nu \overline{\text{Bun}}_{N^-}} {}^1\mathcal{N}^{\bar{x}'} \rightarrow G/N^-.$$

□

Remark 5.5. Since the action of T on G/N^- is free, we have an equivalence of categories $\text{Perv}_T(G/N^-) \simeq \text{Perv}(G/B^-)$. So the previous proof has constructed an equivalence of categories

$$\text{Perv}(G/B^-) \rightarrow \text{Perv}_{G^1}({}^1_\nu \overline{\text{Bun}}_{N^-}).$$

As a corollary of the proof of the previous proposition, we obtain:

Corollary 5.6. *Let $\mathcal{F} \in \text{Perv}_{G[[t]]}(\mathcal{Fl}^{\frac{\infty}{2}})$ and $\nu \in \mathbf{Y}$. The cohomology groups of $(i_\nu)^* \mathcal{F}$ are constant sheaves.*

Proof. We will prove by induction on m that the cohomology groups of ${}^p\tau^{\leq m}((i_\nu)^* \mathcal{F})$ are constant sheaves, starting at some negative enough integer such that ${}^p\tau^{\leq m}((i_\nu)^* \mathcal{F}) = 0$. So assume that there exists m such that the cohomology groups of ${}^p\tau^{\leq m}((i_\nu)^* \mathcal{F})$ are constant sheaves. We have a distinguished triangle

$${}^p\tau^{\leq m}((i_\nu)^* \mathcal{F}) \rightarrow {}^p\tau^{\leq m+1}((i_\nu)^* \mathcal{F}) \rightarrow {}^pH^{m+1}((i_\nu)^* \mathcal{F}) \xrightarrow{+1}.$$

By the second point of Proposition 5.4, ${}^pH^{m+1}((i_\nu)^* \mathcal{F})$ is a constant sheaf. From this together with Proposition 4.3, we deduce that the cohomology groups of ${}^p\tau^{\leq m+1}((i_\nu)^* \mathcal{F})$ are constant sheaves. □

Corollary 5.7. [ABB⁺05, Proposition 4.2.4] *Let $k \in \{0, 1\}$. All the simple objects of $\text{Perv}_{G^k}(\mathcal{Fl}^{\frac{\infty}{2}})$ can be written as in the first point of Proposition 5.4, with \mathcal{F} an irreducible object of $\text{Perv}_T(G/N^-)$.*

Proof. We only treat the case $k = 0$, since the arguments are easily adapted to the case $k = 1$. By the first point of Proposition 5.4, objects that are written in the form of *loc. cit.* are indeed objects of $\text{Perv}_{G[[t]]}(\mathcal{Fl}^{\frac{\infty}{2}})$. Conversely, let \mathcal{F} be a simple object of $\text{Perv}_{G[[t]]}(\mathcal{Fl}^{\frac{\infty}{2}})$, and denote by Z its support. We deduce from (4.1) that there exists a

unique $\nu \in \mathbf{Y}$ such that ${}_{\nu}\overline{\text{Bun}}_{N^-} \cap Z$ is open in Z . Since \mathcal{F} is an IC-sheaf (because it is a simple object), we deduce that it must be isomorphic to $(\tilde{i}_{\nu})_{!*}((\tilde{i}_{\nu})^*\mathcal{F})$. But we know that

$$(\tilde{i}_{\nu})^*\mathcal{F} \simeq \bigoplus_{i=1}^m \text{IC}_{{}_{\nu}\overline{\text{Bun}}_{N^-}}$$

for some $m \geq 1$. Since \mathcal{F} is simple we must have $m = 1$. \square

5.3. Stalks of simple objects. The goal of this subsection is to prove that, when $G = \text{SL}_n$, the results of [BFGM02] concerning the description of the stalks of $\text{IC}_{\overline{\text{Bun}}_{B^-}}$ do not depend on the characteristic of \mathbb{k} .

For $\mu \in \mathbf{Y}^{pos}$, we let ${}^{\mu}\text{Bun}_{B^-}$ be a stack fibered over X^{μ} , whose fiber over $D \in X^{\mu}$ consists of the triples $(\mathcal{F}_G, \mathcal{F}_T, \kappa) \in \overline{\text{Bun}}_{B^-}$ such that the morphism

$$\kappa^{\lambda} : \mathcal{V}_{\mathcal{F}_G} \rightarrow \mathcal{L}_{\mathcal{F}_T}^{\lambda}$$

factors through the canonical morphism $\mathcal{L}_{\mathcal{F}_T}^{\lambda}(-\langle \lambda, D \rangle) \rightarrow \mathcal{L}_{\mathcal{F}_T}^{\lambda}$, with the resulting morphism $\mathcal{V}_{\mathcal{F}_G} \rightarrow \mathcal{L}_{\mathcal{F}_T}^{\lambda}(-\langle \lambda, D \rangle)$ being a morphism of vector bundles. We also define a morphism

$$\varphi^{\mu} : \text{Bun}_{B^-} \times X^{\mu} \rightarrow \overline{\text{Bun}}_{B^-},$$

sending $((\mathcal{F}_G, \mathcal{F}_T, \kappa), D)$ to $(\mathcal{F}_G, \mathcal{F}_T(D), \tilde{\kappa})$, where $\tilde{\kappa}^{\lambda}$ is defined as the composition

$$\mathcal{V}_{\mathcal{F}_G} \xrightarrow{\kappa^{\lambda}} \mathcal{L}_{\mathcal{F}_T}^{\lambda} \rightarrow \mathcal{L}_{\mathcal{F}_T}^{\lambda}(\langle \lambda, D \rangle).$$

This morphism φ^{μ} is representable by a locally closed immersion ([BG02, Proposition 6.1.2]), and its image coincides with ${}^{\mu}\text{Bun}_{B^-}$. For $\Gamma \in \mathfrak{A}(\mu)$, we let

$$\varphi_{\Gamma}^{\mu} : \text{Bun}_{B^-} \times X_{\Gamma}^{\mu} \rightarrow \overline{\text{Bun}}_{B^-}$$

denote the locally closed immersion obtained by composing φ^{μ} with the inclusion of $\text{Bun}_{B^-} \times X_{\Gamma}^{\mu}$ into $\text{Bun}_{B^-} \times X^{\mu}$. Also notice that, for $\theta \in \mathbf{Y}^{pos}$ such that $\theta \leq \mu$, restriction of φ^{μ} yields a morphism

$$\varphi_{\mu, \theta} : \text{Bun}_{B^-}^{\theta - \mu} \times X^{\mu} \rightarrow \overline{\text{Bun}}_{B^-}^{\theta},$$

which is representable by a locally closed immersion. By Proposition 6.1.3 of *loc. cit.*, we obtain a stratification

$$(5.2) \quad \overline{\text{Bun}}_{B^-}^{\theta} = \bigsqcup_{\mu \in \mathbf{Y}^{pos}, \mu \leq \theta, \Gamma \in \mathfrak{A}(\mu)} \text{Bun}_{B^-}^{\theta - \mu} \times X_{\Gamma}^{\mu}.$$

Conjecture 5.1. *For any $i \in \mathbb{Z}$ and closed point $x \in \overline{\text{Bun}}_{B^-}$, the dimension of the \mathbb{k} -vector space $H^i((\text{IC}_{\overline{\text{Bun}}_{B^-}})_x)$ does not depend on the characteristic of \mathbb{k} .*

Lemma 5.8. *If $G = \text{SL}_n$, then Conjecture 5.1 holds.*

The proof of this lemma is postponed to §5.4, and will follow from the facts that:

- (1) The stack $\overline{\text{Bun}}_{B^-}$ admits a small resolution of singularities π when $G = \text{SL}_n$.
- (2) The cohomology of the fibers of π is independent of the characteristic.

Indeed, the first point implies that the stalk of $\text{IC}_{\overline{\text{Bun}}_{B^-}}$ at a point coincides with the cohomology of the fiber of this point through π , and the second point allows to conclude.

Proposition 5.9. *Assume that Conjecture 5.1 holds. Let $\mu \in \mathbf{Y}^{pos}$ and $\Gamma \in \mathfrak{A}(\mu)$. Then the complex $(\varphi_{\Gamma}^{\mu})^! \text{IC}_{\overline{\text{Bun}}_{B^-}} \in D(\text{Bun}_{B^-} \times X_{\Gamma}^{\mu})$ lives in perverse cohomological degrees ≥ 2 .*

Proof. In the case where $\text{char}(\mathbb{k}) = 0$, the description of

$$(\varphi_\Gamma^\mu)^* \text{IC}_{\overline{\text{Bun}}_{B^-}} \in \text{D}(\text{Bun}_{B^-} \times X_\Gamma^\mu)$$

follows from [BFGM02, Theorem 7.3]. It shows in particular that $(\varphi_\Gamma^\mu)^* \text{IC}_{\overline{\text{Bun}}_{B^-}}$ lives in perverse cohomological degrees ≤ -2 . Applying Verdier duality, we deduce that $(\varphi_\Gamma^\mu)^\dagger \text{IC}_{\overline{\text{Bun}}_{B^-}}$ lives in perverse cohomological degrees ≥ 2 when $\text{char}(\mathbb{k}) = 0$. Thus, the case of positive characteristic follows from Lemma 5.8. \square

For all $\nu \in \mathbf{Y}$, $\mu \in \mathbf{Y}^{\text{pos}}$ and $\Gamma \in \mathfrak{A}(\mu)$, we can similarly define a locally closed immersion

$$\nu\varphi_\Gamma^\mu : \nu\text{Bun}_{N^-} \times X_\Gamma^\mu \rightarrow \leq_\nu \overline{\text{Bun}}_{N^-},$$

which is obtained by base change of the morphism φ_Γ^μ through the morphism $\leq_\nu \overline{\text{Bun}}_{N^-} \rightarrow \overline{\text{Bun}}_{B^-}$.

Corollary 5.10. *Assume that Conjecture 5.1 holds. Let $\nu \in \mathbf{Y}$, $\mu \in \mathbf{Y}^{\text{pos}}$ and $\Gamma \in \mathfrak{A}(\mu)$. Then the complex $(\nu\varphi_\Gamma^\mu)^\dagger \text{IC}_{\leq_\nu \overline{\text{Bun}}_{N^-}} \in \text{D}(\nu\text{Bun}_{N^-} \times X_\Gamma^\mu)$ lives in perverse cohomological degrees ≥ 2 .*

Proof. This follows from Proposition 5.9 together with the fact that the morphism

$$\leq_\nu \overline{\text{Bun}}_{N^-} \rightarrow \overline{\text{Bun}}_{B^-}$$

is smooth (Remark 4.2), so that the perversely-shifted pullback of the IC-sheaf is the IC-sheaf. \square

5.4. Laumon's compactification. Recall the classical fact that there is an equivalence of categories between the category of rank- n vector bundle \mathcal{M} over X , endowed with an isomorphism $\wedge^n \mathcal{M} \simeq \mathcal{O}_X$, and the category of principal SL_n -bundles over X , defined on objects by sending a sheaf \mathcal{M} to the relative spectrum of the dual sheaf \mathcal{M}^\vee .

In this subsection, we prove Lemma 5.8. When G is SL_n , Laumon introduced another compactification of the morphism $\text{Bun}_{B^-} \rightarrow \text{Bun}_G$. Namely, ${}^L \overline{\text{Bun}}_{B^-}$ is the smooth algebraic stack classifying the data of a rank n vector bundle \mathcal{M} over X together with an isomorphism $\wedge^n \mathcal{M} \simeq \mathcal{O}_X$ and a complete flag $0 = \mathcal{M}_0 \subset \mathcal{M}_1 \subset \cdots \subset \mathcal{M}_n = \mathcal{M}$ of locally free subsheaves, where each inclusion $\mathcal{M}_i \subset \mathcal{M}_{i+1}$ is merely an inclusion of subsheaves, and not necessarily of vector bundles (cf. [Lau90, §3.3.1], the version where the quotients $\mathcal{M}_{i+1}/\mathcal{M}_i$ are vector bundles is isomorphic to Bun_{B^-}). There is then a natural morphism $\pi : {}^L \overline{\text{Bun}}_{B^-} \rightarrow \overline{\text{Bun}}_{B^-}$, sending a complete flag $(\mathcal{M}_i)_{0 \leq i \leq n}$ to the point

$$(\mathcal{F}_G := (\mathcal{M}, \wedge^n \mathcal{M} \simeq \mathcal{O}_X), \mathcal{F}_T, \kappa)$$

of $\overline{\text{Bun}}_{B^-}$ defined as follows: \mathcal{F}_T is the T -bundle $(\det(\mathcal{M}_{i+1}/\mathcal{M}_i))_{0 \leq i \leq n-1}$, and $(\kappa^\lambda)_{\lambda \in \mathbf{X}^+}$ is the collection of morphisms determined by the inclusions of sheaves

$$\mathcal{L}_{\mathcal{F}_T}^{\varpi_k} := \wedge^k \mathcal{M}_k \subset \wedge^k \mathcal{M} = \mathcal{V}_{\mathcal{M}}^{\varpi_k} \quad \forall 0 \leq k \leq n.$$

The connected components of ${}^L \overline{\text{Bun}}_{B^-}$ are labelled by \mathbf{Y} . We will denote by ${}^L \overline{\text{Bun}}_{B^-}^\theta$ the connected component associated with an element $\theta \in \mathbf{Y}$. By construction, ${}^L \overline{\text{Bun}}_{B^-}^\theta$ is the pre-image of $\overline{\text{Bun}}_{B^-}^\theta$ through π , and we will denote by

$$\pi_\theta : {}^L \overline{\text{Bun}}_{B^-}^\theta \rightarrow \overline{\text{Bun}}_{B^-}^\theta$$

the restriction of π . Thanks to [Lau90, Lemme 4.2.3], the fibers of π_θ are projective smooth schemes over \mathbb{F} . We need one last preliminary result.

Lemma 5.11. *The trivial bundle \mathcal{F}_G^0 is open inside Bun_G .*

Proof. Recall that we have assumed that $X = \mathbb{P}^1$. Thanks to [Zhu16, Theorem 2.3.7], we have an isomorphism of \mathbb{F} -stacks

$$[G[t^{-1}] \backslash \text{Gr}] \simeq \text{Bun}_G,$$

where we take the quotient stack on the left-hand side. The trivial bundle \mathcal{F}_G^0 corresponds to the $G[t^{-1}]$ -orbit associated with the class of the neutral element in Gr . Thanks to [Zhu16, Proposition 2.3.3], the latter orbit is open in Gr . \square

Finally, recall the definition of a "pseudoaffine" variety over \mathbb{F} (cf. [Kuz96, §2.3]). Those are defined by induction on the dimension. A pseudo affine space of dimension one is \mathbb{A}^1 . A variety Y is pseudo affine if it admits a locally trivial fibration $Y \rightarrow B$, with pseudo affine fibers and pseudo affine B . We are now ready to prove Lemma 5.8:

Proof. Let $\theta \in \mathbf{Y}^{pos}$. First, let us justify why the stalk of $\text{IC}_{\overline{\text{Bun}}_{B^-}^\theta}$ at a point $(\mathcal{F}_G, \mathcal{F}_T, \kappa) \in \overline{\text{Bun}}_{B^-}^\theta$ depends only on the type of defect of that point, i.e. to which stratum of (5.2) it belongs. Similarly to §4.3 (and as in [BFGM02]), one can define the Zastava space

$$'Z^{\mu-\theta} \subset \overline{\text{Bun}}_{B^-}^\theta \times_{\text{Bun}_G} \text{Bun}_B^\mu$$

as an open substack of the right-hand side, for any $\mu \in \mathbf{Y}^{pos}$. The Zastava spaces defined in this way satisfy an analogous factorization property (see for instance [BFGM02, §7]). The canonical projection $'Z^{\mu-\theta} \rightarrow \overline{\text{Bun}}_{B^-}^\theta$ is smooth for μ large enough, and any open substack of finite type of $\overline{\text{Bun}}_{B^-}^\theta$ belongs to its image for μ large enough (cf. [BFGM02, §3.6]). Moreover, the stratification by type of defect (5.2) of $\overline{\text{Bun}}_{B^-}^\theta$ induces a stratification of $'Z^{\mu-\theta}$. Therefore, it is enough to prove that the dimension of the cohomology of the stalk of $\text{IC}_{'Z^{\mu-\theta}}$ is constant on every stratum. Thanks to the factorization property of $'Z^{\mu-\theta}$, we can see that it is enough to check that the stalk of $\text{IC}_{\overline{\text{Bun}}_{B^-}^\theta}$ at a point of $\text{Bun}_{B^-}^{\theta-\delta}$ does not depend on the point, for $0 \leq \delta \leq \theta$. Here $\text{Bun}_{B^-}^{\theta-\delta}$ is seen as included in $\text{Bun}_{B^-}^{\theta-\delta} \times X_{\{\{\theta-\delta\}\}}^{\theta-\delta}$ via the inclusion of $\{x\}$ into $X_{\{\{\theta-\delta\}\}}^{\theta-\delta}$. But the same arguments as in the proof of Corollary 5.6 show that the restriction of $\text{IC}_{\overline{\text{Bun}}_{B^-}^\theta}$ to $\text{Bun}_{B^-}^{\theta-\delta}$ has constant stalks.

Therefore it is enough to consider points of the form $\varphi := (\mathcal{F}_G^0, \mathcal{F}_T, \kappa) \in \overline{\text{Bun}}_{B^-}^\theta$, in which case the result follows from [Kuz96]. Indeed, in this case φ belongs to the quasimap space $\mathcal{Q}^\theta := (\mathfrak{p}^\theta)^{-1}(\mathcal{F}_G^0)$, and $(\pi_\theta)^{-1}(\{\varphi\})$ coincides with the pre-image of $\{\varphi\}$ under the small resolution of singularities ${}^L\mathcal{Q}^\theta \rightarrow \mathcal{Q}^\theta$ from *loc. cit.*, where ${}^L\mathcal{Q}^\theta$ is Laumon's compactification of the space of maps of degree θ ([Kuz96, §1.4]). Thus, the stalk of $\text{IC}_{\mathcal{Q}^\theta}$ at φ coincides with the étale cohomology $H^\bullet((\pi_\theta)^{-1}(\{\varphi\}), \mathbf{k})$. The scheme of finite type $(\pi_\theta)^{-1}(\{\varphi\})$ is stratified by pseudoaffine spaces by [Kuz96, Corollary 2.4.4], so that $H^\bullet((\pi_\theta)^{-1}(\{\varphi\}), \mathbf{k})$ is easily seen to be independent of $\text{char}(\mathbf{k})$. So the dimension of the cohomology of the stalk of $\text{IC}_{\mathcal{Q}^\theta}$ at φ is independent of $\text{char}(\mathbf{k})$. But \mathcal{Q}^θ is open in $\overline{\text{Bun}}_{B^-}^\theta$ thanks to Lemma 5.11, from which we deduce the the stalk of $\text{IC}_{\overline{\text{Bun}}_{B^-}^\theta}$ at φ coincides with the stalk of $\text{IC}_{\mathcal{Q}^\theta}$ at φ . \square

Remark 5.12. It is stated in [BG02, §0.2.1] that the morphism π is a small resolution of singularities, so that computing the stalks of $\text{IC}_{\overline{\text{Bun}}_{B^-}^\theta}$ amounts to computing the cohomology of the fibers of π . We have just explained why the cohomology of those fibers is independent of $\text{char}(\mathbf{k})$.

5.5. Semi-simplicity of the spherical category. For $\nu \in \mathbf{Y}$, we put

$$\mathcal{IC}_\nu^\infty := (\bar{i}_\nu)_! (\mathbf{IC}_{\nu \overline{\text{Bun}}_{N^-}}).$$

By Proposition 5.4, the set $\{\mathcal{IC}_\nu^\infty, \nu \in \mathbf{Y}\}$ coincides with the set of isomorphism classes of simple objects of the category $\text{Perv}_{G[[t]]}(\mathcal{Fl}^{\frac{\infty}{2}})$.

Lemma 5.13. *Assume that Conjecture 5.1 holds. Let $\nu_1, \nu_2 \in \mathbf{Y}$, with $\nu_1 < \nu_2$. The complex*

$$i_{\nu_1}^! \mathcal{IC}_{\nu_2}^\infty \in D(\nu_1 \text{Bun}_{N^-})$$

lives in perverse cohomological degrees ≥ 2 .

Proof. We want to apply Corollary 5.10. For that, we just need to notice that the inclusion $\nu_1 \text{Bun}_{N^-} \subset \leq_{\nu_2} \overline{\text{Bun}}_{N^-}$ factors through the locally closed inclusion

$$\nu_1 \text{Bun}_{N^-} \times X_{\{\nu_2 - \nu_1\}}^{\nu_2 - \nu_1} \hookrightarrow \leq_{\nu_2} \overline{\text{Bun}}_{N^-}$$

from §5.3, where the inclusion $\nu_1 \text{Bun}_{N^-} \hookrightarrow \nu_1 \text{Bun}_{N^-} \times X_{\{\nu_2 - \nu_1\}}^{\nu_2 - \nu_1}$ is induced by the diagonal embedding of $\{x\}$ into $X_{\{\nu_2 - \nu_1\}}^{\nu_2 - \nu_1}$. \square

The previous Lemma allows us to take back the arguments of [ABB⁺05].

Theorem 5.14 ([ABB⁺05, Proposition 4.3.2]). *Assume that Conjecture 5.1 holds. The category $\text{Perv}_{G[[t]]}(\mathcal{Fl}^{\frac{\infty}{2}})$ is semi-simple.*

Proof. Let $\nu_1, \nu_2 \in \mathbf{Y}$. We want to prove that

$$\text{Ext}_{\infty \overline{\text{Bun}}_{N^-}}^1(\mathcal{IC}_{\nu_1}^\infty, \mathcal{IC}_{\nu_2}^\infty) = 0.$$

Arguing as in the second case of [BR18, Proof of Proposition 4.4], we see that the support of one sheaf must be included in the other for the space $\text{Ext}_{\infty \overline{\text{Bun}}_{N^-}}^1(\mathcal{IC}_{\nu_1}^\infty, \mathcal{IC}_{\nu_2}^\infty)$ to be non-trivial. Applying Verdier duality, we may and will assume that $\leq_{\nu_1} \overline{\text{Bun}}_{N^-} \subset \leq_{\nu_2} \overline{\text{Bun}}_{N^-}$, which means that $\nu_1 \leq \nu_2$.

Next, consider the stack

$$\mathcal{X} := \leq_{\nu_2} \overline{\text{Bun}}_{N^-} - (\leq_{\nu_1} \overline{\text{Bun}}_{N^-} - \nu_1 \text{Bun}_{N^-}),$$

and denote by $j : \mathcal{X} \hookrightarrow \leq_{\nu_2} \overline{\text{Bun}}_{N^-}$ the inclusion. Since \mathcal{X} contains the open substack $\nu_2 \overline{\text{Bun}}_{N^-}$, the complex $\mathcal{IC}_{\nu_2}^\infty$ coincides with the image of the IC-sheaf on \mathcal{X} through the functor $j_{!*}$. On the other hand, if we let j' denote the inclusion $\nu_1 \text{Bun}_{N^-} \hookrightarrow \mathcal{X}$, then $\mathcal{IC}_{\nu_1}^\infty$ coincides with the image of $j'_{!*}(\mathbf{IC}_{\nu_1 \text{Bun}_{N^-}})$ through $j_{!*}$, since $j_{!*} j'_{!*} \simeq (j \circ j')_{!*}$. Applying [FGV01, Lemma 6.1.4], we get an inclusion

$$\text{Ext}_{\leq_{\nu_2} \overline{\text{Bun}}_{N^-}}^1(\mathcal{IC}_{\nu_1}^\infty, \mathcal{IC}_{\nu_2}^\infty) \hookrightarrow \text{Ext}_{\mathcal{X}}^1(j'_{!*}(\mathbf{IC}_{\nu_1 \text{Bun}_{N^-}}), \mathbf{IC}_{\mathcal{X}}).$$

Since $\nu_1 \text{Bun}_{N^-}$ is closed in \mathcal{X} , the latter Ext^1 is isomorphic to

$$(5.3) \quad \text{Ext}_{\nu_1 \text{Bun}_{N^-}}^1(\mathbf{IC}_{\nu_1 \text{Bun}_{N^-}}, i_{\nu_1}^! \mathcal{IC}_{\nu_2}^\infty).$$

If $\nu_1 = \nu_2$, then the above space is $\{0\}$ since $\nu_1 \text{Bun}_{N^-}$ is contractible (Proposition 4.3). If $\nu_1 < \nu_2$, then the complex $i_{\nu_1}^! \mathcal{IC}_{\nu_2}^\infty$ lives in perverse cohomological degrees ≥ 2 thanks to Lemma 5.13. \square

The next result will be useful in §7.1.

Corollary 5.15. *Let $\nu, \nu'' \in \mathbf{Y}$ with $\nu'' \leq \nu$, $r \in \mathbb{Z}_{\geq 0}$, and put $\mu_r := (\ell^r - 1)\zeta$. For r large enough, there is an isomorphism of complexes of vector spaces*

$$\mathcal{IC}_{\nu}^{\infty} \big|_{\nu'' \text{Bun}_{N^-}} \simeq \mathcal{IC}^{\mu_r} \big|_{\text{Gr}^{\mu_r - (\nu - \nu'')}} ,$$

where $\mathcal{IC}_{\nu}^{\infty} \big|_{\nu'' \text{Bun}_{N^-}}$ (resp. $\mathcal{IC}^{\mu_r} \big|_{\text{Gr}^{\mu_r - (\nu - \nu'')}}$) is meant to denote the stalk of $\mathcal{IC}_{\nu}^{\infty}$ (resp. of \mathcal{IC}^{μ_r}) at any point of $\nu'' \text{Bun}_{N^-}$ (resp. of $\text{Gr}^{\mu_r - (\nu - \nu'')}$)

Proof. Since the dimensions of the stalks of $\mathcal{IC}_{\nu}^{\infty}$ and \mathcal{IC}^{μ_r} do not depend on the characteristic of \mathbb{k} , the result follows from the end of the proof of [ABB⁺05, Theorem 5.3.5]. Let us briefly recall the arguments.

On the one hand, the stalks of $\mathcal{IC}_{\nu}^{\infty}$ do not depend on $\text{char}(\mathbb{k})$ thanks to Lemma 5.8, so that the generating function of $\mathcal{IC}_{\nu}^{\infty} \big|_{\nu'' \text{Bun}_{N^-}}$ is given by the q -analogue of Kostant's partition function $\mathcal{K}^{\nu - \nu''}(t)$ associated with $\nu - \nu''$ thanks to [FFKM99, §3] (or the main result of [Kuz96]).

On the other hand, the generating function of $\mathcal{IC}^{\mu_r} \big|_{\text{Gr}^{\mu_r - (\nu - \nu'')}}$ does not depend on $\text{char}(\mathbb{k})$ thanks to Lemma 3.7. The latter coincides with $\mathcal{K}^{\mu_r - (\mu_r - (\nu - \nu''))}(t) = \mathcal{K}^{\nu - \nu''}(t)$ when r is large enough (so that μ_r is dominant enough) thanks to [Lus81, §§9,11]. \square

5.6. Iwahori-equivariant objects. We now define the main category of interest (following [ABB⁺05, §4.4.3]). Let $\nu \in \mathbf{Y}$, and denote by ${}^I_{\nu} \overline{\text{Bun}}_{N^-}$ and ${}^I_{\infty} \overline{\text{Bun}}_{N^-}$ (resp. ${}^I_{\nu} \overline{\text{Bun}}_{N^-}$ and ${}^I_{\infty} \overline{\text{Bun}}_{N^-}$) the quotient stacks of ${}^1 \overline{\text{Bun}}_{N^-}$ and ${}^1_{\infty} \overline{\text{Bun}}_{N^-}$ by N (resp. by B). We let $\text{Perv}_{I_u}(\mathcal{Fl}^{\infty})$ (resp. $\text{Perv}_I(\mathcal{Fl}^{\infty})$) be the subcategory of

$$\text{Perv}({}^I_{\infty} \overline{\text{Bun}}_{N^-}) \quad (\text{resp. } \text{Perv}({}^I_{\infty} \overline{\text{Bun}}_{N^-}))$$

consisting of objects whose image through the forgetful functor to $\text{Perv}({}^1_{\infty} \overline{\text{Bun}}_{N^-})$ belongs to $\text{Perv}_{G^1}(\mathcal{Fl}^{\infty})$.

By abuse of notation, we still denote by $\overline{e\nu}_{\nu} : {}^I_{\nu} \overline{\text{Bun}}_{N^-} \rightarrow N \backslash G/B^-$ the morphism obtained by composing the map ${}^1 \overline{\text{Bun}}_{N^-} \rightarrow G/N^-$ (cf. (4.4)) with the canonical projection $G/N^- \rightarrow G/B^-$, and then quotienting by N . For $w \in W$, we let

$$N_w := N \backslash (N \cdot w \cdot B^-) / B^-$$

be the Schubert cell associated with w . For $x \in W_{\text{ext}}$, written as $x = wt_{\nu}$, we let ${}^I_x \overline{\text{Bun}}_{N^-}$ be the preimage of N_w through $\overline{e\nu}_{\nu}$, and $\tilde{i}_x : {}^I_x \overline{\text{Bun}}_{N^-} \hookrightarrow {}^I_{\infty} \overline{\text{Bun}}_{N^-}$ denote the locally closed inclusion. Let $\text{IC}_{w, G/B^-}$, $\tilde{\nabla}_w$, $\tilde{\Delta}_w \in \text{Perv}_N(G/B^-)$, denote respectively the Goresky-MacPherson extension of the constant sheaf on N_w , the costandard (i.e. $!$ -extension of the constant sheaf) and standard (i.e. $*$ -extension of the constant sheaf) objects associated with w . Thanks to Proposition 5.4, we see that the simple objects of $\text{Perv}_{I_u}(\mathcal{Fl}^{\infty})$ consist of the intersection cohomology sheaves $\text{IC}_x^{\infty} \in \text{Perv}({}^I_{\infty} \overline{\text{Bun}}_{N^-})$ associated with ${}^I_x \overline{\text{Bun}}_{N^-}$, for $x \in W_{\text{ext}}$. More precisely, we have

$$\text{IC}_x^{\infty} := (\tilde{i}_{\nu})_{!*} \left(\text{IC}_{w, G/B^-} \boxtimes \text{IC}_{\nu \overline{\text{Bun}}_{N^-}} \right), \quad x = wt_{\nu}.$$

Consider the functor $F : {}^I_{\infty} \overline{\text{Bun}}_{N^-} \rightarrow {}^{\infty} \overline{\text{Bun}}_{N^-}$ induced by forgetting the structure of level 1, and put

$$\text{For}_{I_u}^{G[[t]]} := F^*[\dim(G) - \dim(N)] : D({}^{\infty} \overline{\text{Bun}}_{N^-}) \rightarrow D({}^I_{\infty} \overline{\text{Bun}}_{N^-}).$$

Similarly, one defines the functor

$$\text{For}_I^{G[[t]]} : D({}^{\infty} \overline{\text{Bun}}_{N^-}) \rightarrow D({}^I_{\infty} \overline{\text{Bun}}_{N^-}).$$

The next result will be used profusely in the sequel.

Proposition 5.16. *The functors $\text{For}_{I_u}^{G[[t]]}, \text{For}_I^{G[[t]]}$ induce fully faithful functors*

$$\text{For}_{I_u}^{G[[t]]} : \text{Perv}_{G[[t]]}(\mathcal{F}l_{\frac{\infty}{2}}) \rightarrow \text{Perv}_{I_u}(\mathcal{F}l_{\frac{\infty}{2}}), \quad \text{For}_I^{G[[t]]} : \text{Perv}_{G[[t]]}(\mathcal{F}l_{\frac{\infty}{2}}) \rightarrow \text{Perv}_I(\mathcal{F}l_{\frac{\infty}{2}})$$

which satisfy

$$\text{For}_{I_u}^{G[[t]]}(\mathcal{I}C_{\nu}^{\frac{\infty}{2}}) \simeq \text{IC}_{\nu}^{\frac{\infty}{2}}, \quad \text{For}_I^{G[[t]]}(\mathcal{I}C_{\nu}^{\frac{\infty}{2}}) \simeq \text{IC}_{\nu}^{\frac{\infty}{2}}$$

for all $\nu \in \mathbf{Y}$, where $\text{IC}_{\nu}^{\frac{\infty}{2}}$ is seen as an object of $\text{Perv}_I(\mathcal{F}l_{\frac{\infty}{2}})$ in the second isomorphism.

Proof. The morphism F is induced by the morphism $\text{For} : {}^1_{\infty}\overline{\text{Bun}}_{N^-} \rightarrow {}^{\infty}\overline{\text{Bun}}_{N^-}$ forgetting the structure of level one. The latter is a G -torsor, so that F is smooth of relative dimension $\dim(G) - \dim(N)$. Since

$$F^{-1}({}_{\nu}\overline{\text{Bun}}_{N^-}) \simeq {}^I_{\nu}\overline{\text{Bun}}_{N^-}$$

for all $\nu \in \mathbf{Y}$ we have an isomorphism

$$\text{For}_{I_u}^{G[[t]]}(\mathcal{I}C_{\nu}^{\frac{\infty}{2}}) \simeq \text{IC}({}^I_{\nu}\overline{\text{Bun}}_{N^-}).$$

But $\text{IC}({}^I_{\nu}\overline{\text{Bun}}_{N^-}) \simeq \text{IC}_{\nu}^{\frac{\infty}{2}}$ since the N -orbit of the trivial element is dense in G/B^- . The full faithfulness then follows from the semi-simplicity of $\text{Perv}_{G[[t]]}(\mathcal{F}l_{\frac{\infty}{2}})$. The proof of the assertions concerning $\text{For}_I^{G[[t]]}$ is similar. \square

We also let $\nabla_x^{\frac{\infty}{2}}, \Delta_x^{\frac{\infty}{2}}$ denote the corresponding costandard and standard objects. Namely we have

$$\nabla_x^{\frac{\infty}{2}} := (\bar{i}_{\nu})_* \left(\tilde{\nabla}_w \tilde{\boxtimes} \text{IC}_{\nu}\overline{\text{Bun}}_{N^-} \right), \quad \Delta_x^{\frac{\infty}{2}} := (\bar{i}_{\nu})! \left(\tilde{\Delta}_w \tilde{\boxtimes} \text{IC}_{\nu}\overline{\text{Bun}}_{N^-} \right).$$

Note that the objects $\nabla_x^{\frac{\infty}{2}}, \Delta_x^{\frac{\infty}{2}}$ admit canonical lifts to the category $\text{D}({}^I_{\infty}\overline{\text{Bun}}_{N^-})$. We will denote the corresponding objects by $\nabla_x^{\frac{\infty}{2}}, \Delta_x^{\frac{\infty}{2}} \in \text{D}({}^I_{\infty}\overline{\text{Bun}}_{N^-})$ for all $x \in W_{\text{ext}}$. Thus we have

$$\text{For}_{I_u}^I(\Delta_x^{\frac{\infty}{2}}) \simeq \Delta_x^{\frac{\infty}{2}}, \quad \text{For}_{I_u}^I(\nabla_x^{\frac{\infty}{2}}) \simeq \nabla_x^{\frac{\infty}{2}}$$

(where $\text{For}_{I_u}^I : \text{D}({}^I_{\infty}\overline{\text{Bun}}_{N^-}) \rightarrow \text{D}({}^I_{\infty}\overline{\text{Bun}}_{N^-})$ denotes the forgetful functor).

By construction we have isomorphisms in $\text{D}({}^I_{\infty}\overline{\text{Bun}}_{N^-})$:

$$\nabla_x^{\frac{\infty}{2}} \simeq (\bar{i}_x)_*(\text{IC}_{\frac{x}{I_u}}\overline{\text{Bun}}_{N^-}), \quad \Delta_x^{\frac{\infty}{2}} \simeq (\bar{i}_x)!(\text{IC}_{\frac{x}{I_u}}\overline{\text{Bun}}_{N^-}).$$

Moreover, thanks to the proof of [ABB⁺05, Corollary 4.4.5], we know that the morphism \bar{i}_x is affine, so that Δ_x and ∇_x are perverse. We can then reuse the arguments of [ABB⁺05, Corollary 4.4.7] to prove the following result.

Proposition 5.17. *For all $x, x' \in W_{\text{ext}}$, $i \in \{0, 1, 2\}$, we have*

$$\text{Ext}_{\text{Perv}_{I_u}(\mathcal{F}l_{\frac{\infty}{2}})}^i(\Delta_x^{\frac{\infty}{2}}, \nabla_{x'}^{\frac{\infty}{2}}) = \begin{cases} \mathbb{k} & \text{if } i = 0 \text{ and } x = x', \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Let $\nu \in \mathbf{Y}$, $\mathcal{F}' \in \text{Perv}_{G^1}({}^1_{\nu}\overline{\text{Bun}}_{N^-})$, and $\mathcal{F} \in \text{Perv}_{G^1}({}^1_{\infty}\overline{\text{Bun}}_{N^-})$. We first claim that, if $(\bar{i}_{\nu})_!(\mathcal{F}')$ is a perverse sheaf, then the canonical morphism

$$(5.4) \quad \text{Ext}_{\text{D}_T({}^1_{\infty}\overline{\text{Bun}}_{N^-})}^1((\bar{i}_{\nu})_!(\mathcal{F}'), \mathcal{F}) \rightarrow \text{Ext}_{\text{D}_T({}^1_{\infty}\overline{\text{Bun}}_{N^-})}^1((i_{\nu})_!(\mathcal{F}'|_{{}^1_{\nu}\overline{\text{Bun}}_{N^-}}), \mathcal{F})$$

is an isomorphism. This follows from the same arguments as in [ABB⁺05, Proposition 4.2.14]. The five lemma then implies that the analogue morphism (5.4) for Ext^2 is injective.

But we have:

$$\begin{aligned} \mathrm{Ext}_{D_T(I_\infty \overline{\mathrm{Bun}}_{N^-})}^i((i_x)_!(\mathbb{k}[\ell(x)]), \nabla_{x'}^{\frac{\infty}{2}}) &\simeq \mathrm{Ext}_{D_T(I_x^u \mathrm{Bun}_{N^-})}^i(\mathbb{k}[\ell(x)], (i_x)^! \nabla_{x'}^{\frac{\infty}{2}}) \\ &\simeq H_T^{i-\ell(x)}(I_x^u \mathrm{Bun}_{N^-}, (i_x)^! \nabla_{x'}^{\frac{\infty}{2}}). \\ &\simeq \begin{cases} \mathbb{k} & \text{if } i = 0 \text{ and } x = x', \\ 0 & \text{otherwise,} \end{cases} \end{aligned}$$

where the last isomorphism is due to the fact that, by definition of $I_x^u \mathrm{Bun}_{N^-}$ together with Proposition 4.3, the k -th cohomology group of the quotient $T \setminus I_x^u \mathrm{Bun}_{N^-}$ vanishes is when $k \neq 0$, and is isomorphic to \mathbb{k} otherwise. \square

Corollary 5.18. *For any $w \in W_{\mathrm{ext}}$, the object $\nabla_w^{\frac{\infty}{2}}$ has a simple socle, isomorphic to $\mathrm{IC}_w^{\frac{\infty}{2}}$.*

Proof. Let $w' \in W_{\mathrm{ext}}$ be such that we have a non-zero morphism $\mathrm{IC}_{w'}^{\frac{\infty}{2}} \rightarrow \nabla_w^{\frac{\infty}{2}}$. Composing this morphism with the surjective morphism $\Delta_{w'}^{\frac{\infty}{2}} \rightarrow \mathrm{IC}_{w'}^{\frac{\infty}{2}}$, we obtain a non-zero morphism $\Delta_{w'}^{\frac{\infty}{2}} \rightarrow \nabla_w^{\frac{\infty}{2}}$. By Proposition 5.17, we must have $w' = w$. \square

Remark 5.19. The same comments as in [ABB⁺05, Remark 4.4.7] apply here. Namely, one deduces from Proposition 5.17 that the Ext group between a standard and costandard object vanishes for all $i > 0$. Moreover, for any $\mathcal{F} \in \mathrm{Perv}_{I_u}(\mathcal{F}l^{\frac{\infty}{2}})$, we have an isomorphism

$$\mathrm{Ext}_{\mathrm{Perv}_{I_u}(\mathcal{F}l^{\frac{\infty}{2}})}^i(\Delta_x^{\frac{\infty}{2}}, \mathcal{F}) \simeq H^i(I_x^u \mathrm{Bun}_{N^-}, i_x^! \mathcal{F}),$$

and one deduces from the second point of Proposition 5.4 that the cohomology groups of $i_x^! \mathcal{F}$ are local systems. Since $I_x^u \mathrm{Bun}_{N^-}$ is contractible according to *loc. cit.*, those cohomology groups are in fact constant sheaves.

Remark 5.20. Notice that the group G acts on ${}^1_\infty \overline{\mathrm{Bun}}_{N^-}$. This allows us to define a convolution product

$$(-) \star (-) : D_N(G/B) \times D(I_\infty \overline{\mathrm{Bun}}_{N^-}) \rightarrow D(I_\infty \overline{\mathrm{Bun}}_{N^-}).$$

Namely, for $(\mathcal{F}, \mathcal{G}) \in D_N(G/B) \times D(I_\infty \overline{\mathrm{Bun}}_{N^-})$, one can form the twisted external tensor product $\mathcal{F} \tilde{\boxtimes} \mathcal{G} \in D_N(G \times {}^B 1_\infty \overline{\mathrm{Bun}}_{N^-})$, and define $\mathcal{F} \star \mathcal{G}$ to be the pushforward of $\mathcal{F} \tilde{\boxtimes} \mathcal{G}$ through the morphism $G \times {}^B 1_\infty \overline{\mathrm{Bun}}_{N^-} \rightarrow {}^1_\infty \overline{\mathrm{Bun}}_{N^-}$ induced by the action of G . One can then easily check that, for all $w_1, w_2 \in W$, $\lambda \in \mathbf{Y}$ such that $\ell(w_1 w_2 w_0) = \ell(w_1) + \ell(w_2 w_0)$, we have

$$(5.5) \quad \mathcal{N}_{w_1} \star \nabla_{w_2 t_\lambda}^{\frac{\infty}{2}} \simeq \nabla_{w_1 w_2 t_\lambda}^{\frac{\infty}{2}} \quad \mathcal{D}_{w_1} \star \Delta_{w_2 t_\lambda}^{\frac{\infty}{2}} \simeq \Delta_{w_1 w_2 t_\lambda}^{\frac{\infty}{2}},$$

where \mathcal{N}_{w_1} , resp. \mathcal{D}_{w_1} , is seen as the costandard object (resp. standard object) associated with w in $D_N(G/B)$ through the fully faithful functor $D_N(G/B) \rightarrow D_{I_u}(\mathrm{Fl})$ induced by pullback through the canonical projection $\mathrm{Fl} \rightarrow G/B$.

6. CONVOLUTION

6.1. Global affine Grassmannian. Recall (§3.3) that the affine Grassmannian Gr is an ind-scheme which can be defined as the fppf-quotient $G((t))/G[[t]]$. From this definition, one can show that for any test \mathbb{F} -algebra R , $\mathrm{Gr}(\mathrm{Spec}(R))$ is the set of isomorphism classes of pairs (\mathcal{F}_G, β) , where \mathcal{F}_G is a G -torsor over $\mathrm{Spec}(R[[t]])$ and β a trivialization over $\mathrm{Spec}(R((t)))$. We will need the global interpretation of Gr – due to Beauville and Laszlo [BL95] – which can be found for instance in [Zhu16, §1.4]. Namely, the result of *loc. cit.* shows that Gr classifies the data of (\mathcal{F}_G, β) , where $\mathcal{F}_G \in \mathrm{Bun}_G(S)$ and β is a trivialization on $(X - \{x\}) \times S$.

For each coweight $\lambda \in \mathbf{Y}$, one can give the following description of the Schubert variety $\overline{\text{Gr}}^\lambda$, as in [BG02, §3.2.1]. It is the closed subscheme of Gr classifying those (\mathcal{F}_G, β) such that, for each $\mu \in \mathbf{X}^+$ and G -module V whose weights are $\leq \mu$, we have

$$\mathcal{V}_{\mathcal{F}_G^0}(-\langle \lambda, \mu \rangle \cdot (x \times S)) \subset \mathcal{V}_{\mathcal{F}_G}$$

(or in other words, the meromorphic map $\beta_V : \mathcal{V}_{\mathcal{F}_G^0} \rightarrow \mathcal{V}_{\mathcal{F}_G}$ induced by β has a pole of order at most $\langle \lambda, \mu \rangle$ at x). Notice that, by passing to the dual representation, the above condition is equivalent to $\mathcal{V}_{\mathcal{F}_G} \subset \mathcal{V}_{\mathcal{F}_G^0}(-\langle w_0(\lambda), \mu \rangle \cdot (x \times S))$ for all V . Each Schubert variety is a finite dimensional projective scheme which is stable under the $G[[t]]$ -action. For all $\lambda, \lambda' \in \mathbf{Y}$, we have

$$\overline{\text{Gr}}^\lambda \subset \overline{\text{Gr}}^{\lambda'} \iff \lambda \leq \lambda'$$

and we put $\text{Gr}^\lambda := \overline{\text{Gr}}^\lambda - (\cup_{\mu < \lambda} \overline{\text{Gr}}^\mu)$. Then $\text{Gr}^\lambda(\mathbb{F})$ coincides with $G[[t]] \cdot [\lambda]$.

6.2. Hecke stacks. Next, we denote by \mathcal{H}_G the Hecke stack for G at x . It classifies the data $(\mathcal{F}_1, \mathcal{F}_2, \beta)$ of two bundles $\mathcal{F}_1, \mathcal{F}_2 \in \text{Bun}_G(S)$ together with an isomorphism

$$\beta : \mathcal{F}_1|_{(X-\{x\}) \times S} \simeq \mathcal{F}_2|_{(X-\{x\}) \times S}.$$

Recall that ${}^1\text{Bun}_G$ was introduced at the end of §4.1, and let ${}^{\text{Lu}}\text{Bun}_G$ denote the quotient stack of ${}^1\text{Bun}_G$ by N . For each coweight $\lambda \in \mathbf{Y}$, we will denote by $\overline{\mathcal{H}}_G^\lambda$ the closed substack of \mathcal{H}_G classifying those $(\mathcal{F}_1, \mathcal{F}_2, \beta)$ such that, for each $\mu \in \Lambda_+$ and G -module V whose weights are $\leq \mu$, we have

$$\mathcal{V}_{\mathcal{F}_1}(-\langle \lambda, \mu \rangle \cdot (x \times S)) \subset \mathcal{V}_{\mathcal{F}_2} \subset \mathcal{V}_{\mathcal{F}_1}(-\langle w_0(\lambda), \mu \rangle \cdot (x \times S)).$$

As above, the second inclusion is implied by the first one.

Consider the diagram

$$\text{Bun}_G \xleftarrow{\overleftarrow{h}} \mathcal{H}_G \xrightarrow{\overrightarrow{h}} \text{Bun}_G$$

defined by $\overleftarrow{h}(\mathcal{F}_1, \mathcal{F}_2) = \mathcal{F}_1$, $\overrightarrow{h}(\mathcal{F}_1, \mathcal{F}_2) = \mathcal{F}_2$. Let \mathfrak{G} denote the canonical $G[[t]]$ -torsor over Bun_G associated with x . By definition, \mathfrak{G} classifies the data of a G -bundle \mathcal{F}_G over X together with a trivialization $\mathcal{F}_G|_{D_x} \simeq \mathcal{F}_G^0|_{D_x}$. The projection \overleftarrow{h} (and also \overrightarrow{h}) realizes \mathcal{H}_G as a fibration over Bun_G with typical fiber Gr (cf. [BG02, §3.2.4]):

$$\mathcal{H}_G \simeq \text{Gr} \times^{G[[t]]} \mathfrak{G},$$

where $G[[t]]$ acts on the right on \mathfrak{G} , and by left translation on Gr .

Under this identification we have, $\overline{\mathcal{H}}_G^\lambda \simeq \overline{\text{Gr}}^\lambda \times^{G[[t]]} \mathfrak{G}$ (and $\overline{\mathcal{H}}_G^{-w_0(\lambda)} \simeq \overline{\text{Gr}}^\lambda \times^{G[[t]]} \mathfrak{G}$ if we project via \overrightarrow{h}).

For $k_1, k_2 \in \{0, 1\}$, we let ${}^{k_1, k_2}\mathcal{H}_G$ denote the base change of \mathcal{H}_G through the morphism

$${}^{k_1}\text{Bun}_G \times {}^{k_2}\text{Bun}_G \rightarrow \text{Bun}_G \times \text{Bun}_G$$

(where ${}^0\text{Bun}_G := \text{Bun}_G$). We still denote by \overleftarrow{h} and \overrightarrow{h} the canonical projections from ${}^{k_1, k_2}\mathcal{H}_G$ to ${}^{k_1}\text{Bun}_G$ and ${}^{k_2}\text{Bun}_G$ respectively. If we let \mathfrak{G}_x^k denote the canonical G^k -torsor over ${}^k\text{Bun}_G$ associated with x for $k \in \{0, 1\}$, then the projection \overrightarrow{h} realizes ${}^{k_1, k_2}\mathcal{H}_G$ as a fibration over ${}^{k_2}\text{Bun}_G$ with typical fiber $G((t))/G^{k_1}$ ([ABB⁺05, §5.1.1]). This means that we get a canonical isomorphism

$$(6.1) \quad {}^{k_1, k_2}\mathcal{H}_G \simeq G((t))/G^{k_1} \times^{G^{k_2}} \mathfrak{G}^{k_2},$$

where $\mathfrak{G}^0 := \mathfrak{G}$. Similarly, let us denote by ${}^{I_u,0}\mathcal{H}_G$ the Hecke stack obtained by replacing ${}^{k_1}\overline{\text{Bun}}_G$ with ${}^{I_u}\overline{\text{Bun}}_G$ in the previous construction, and letting $k_2 = 0$. Via \overrightarrow{h} , we then get a canonical isomorphism

$$(6.2) \quad {}^{I_u,0}\mathcal{H}_G \simeq G((t))/I_u \times^{G[[t]]} \mathfrak{G}.$$

Moreover, we obtain the following isomorphism via \overleftarrow{h} :

$$(6.3) \quad {}^{I_u,0}\mathcal{H}_G \simeq \text{Gr} \times^{G[[t]]} {}^{I_u}\mathfrak{G},$$

where ${}^{I_u}\mathfrak{G}$ is the canonical $G[[t]]$ -torsor over ${}^{I_u}\overline{\text{Bun}}_G$.

We then consider the following diagram, in which both squares are Cartesian

$$\begin{array}{ccccc} {}^{k_1}\overline{\text{Bun}}_{N^-} & \xleftarrow{\overleftarrow{h'}} & {}^{k_1,k_2}\mathcal{H}_{G,N^-} & \xrightarrow{\overrightarrow{h'}} & {}^{k_2}\overline{\text{Bun}}_{N^-} \\ \downarrow \text{p} & & \downarrow & & \downarrow \text{p} \\ {}^{k_1}\text{Bun}_G & \xleftarrow{\overleftarrow{h}} & {}^{k_1,k_2}\mathcal{H}_G & \xrightarrow{\overrightarrow{h}} & {}^{k_2}\text{Bun}_G. \end{array}$$

Similarly we have the following diagram

$$\begin{array}{ccccc} {}^{I_u}\overline{\text{Bun}}_{N^-} & \xleftarrow{\overleftarrow{h'}} & {}^{I_u,0}\mathcal{H}_{G,N^-} & \xrightarrow{\overrightarrow{h'}} & \overline{\text{Bun}}_{N^-} \\ \downarrow \text{p} & & \downarrow & & \downarrow \text{p} \\ {}^{I_u}\text{Bun}_G & \xleftarrow{\overleftarrow{h}} & {}^{I_u,0}\mathcal{H}_G & \xrightarrow{\overrightarrow{h}} & \text{Bun}_G. \end{array}$$

Now, the previous diagrams together with the identifications (6.1) and (6.2) yield the following canonical isomorphisms:

$$(6.4) \quad {}^{0,0}\mathcal{H}_{G,N^-} \simeq G((t))/G[[t]] \times^{G[[t]]} \tilde{\mathfrak{G}}, \quad {}^{I_u,0}\mathcal{H}_{G,N^-} \simeq G((t))/I_u \times^{G[[t]]} \tilde{\mathfrak{G}},$$

where $\tilde{\mathfrak{G}}$ denotes the canonical $G[[t]]$ -torsor over $\overline{\text{Bun}}_{N^-}$. The identification (6.3) also yield the isomorphism

$${}^{I_u,0}\mathcal{H}_{G,N^-} \simeq \text{Gr} \times^{G[[t]]} {}^{I_u}\tilde{\mathfrak{G}},$$

where ${}^{I_u}\tilde{\mathfrak{G}}$ denotes the canonical $G[[t]]$ -torsor over ${}^{I_u}\overline{\text{Bun}}_{N^-}$. For any $\lambda \in \mathbf{Y}$, we define the locally closed substack

$${}^{I_u,0}\mathcal{H}_{G,N^-}^\lambda := \text{Gr}^\lambda \times^{G[[t]]} {}^{I_u}\tilde{\mathfrak{G}}.$$

6.3. Definition of convolution. Recall ([ABB⁺05, §5.1.1]) that the inversion map $\text{sw} : G((t)) \rightarrow G((t))$, $g \mapsto g^{-1}$ induces an equivalence of categories

$$\mathcal{S} \mapsto \text{sw}^*(\mathcal{S}) : \text{D}_{G^{k_1}}(G((t))/G^{k_2}) \rightarrow \text{D}_{G^{k_2}}(G((t))/G^{k_1}).$$

We are now ready to define the convolution product for any $\mathcal{S} \in \text{D}_{G[[t]]}(\text{Gr})$:

$$\mathcal{F} \mapsto \mathcal{S} \star \mathcal{F} : \text{D}(\overline{\text{Bun}}_{N^-}) \rightarrow \text{D}(\overline{\text{Bun}}_{N^-}), \quad \mathcal{F} \mapsto (\overleftarrow{h'})_*(\text{sw}^*(\mathcal{S}) \boxtimes \mathcal{F}),$$

where $\text{sw}^*(\mathcal{S}) \boxtimes \mathcal{F}$ is seen as an object of $\text{D}({}^{0,0}\mathcal{H}_{G,N^-})$ thanks to (6.4). And respectively for any $\mathcal{S} \in \text{D}_{I_u}(\text{Gr})$:

$$\mathcal{F} \mapsto \mathcal{S} \star \mathcal{F} : \text{D}(\overline{\text{Bun}}_{N^-}) \rightarrow \text{D}({}^{I_u}\overline{\text{Bun}}_{N^-}), \quad \mathcal{F} \mapsto (\overleftarrow{h'})_*(\text{sw}^*(\mathcal{S}) \boxtimes \mathcal{F}).$$

Similarly, one can define convolution functors

$$\begin{aligned} D_I(\mathrm{Gr}) \times D(\infty \overline{\mathrm{Bun}}_{N^-}) &\rightarrow D(\infty^I \overline{\mathrm{Bun}}_{N^-}), (\mathcal{S}, \mathcal{F}) \mapsto \mathcal{S} \star \mathcal{F}, \\ D_{I_u}(\mathrm{Fl}) \times D(\infty^I \overline{\mathrm{Bun}}_{N^-}) &\rightarrow D(\infty^{I_u} \overline{\mathrm{Bun}}_{N^-}), (\mathcal{S}, \mathcal{F}) \mapsto \mathcal{S} \star \mathcal{F}, \\ D_I(\mathrm{Fl}) \times D(\infty^I \overline{\mathrm{Bun}}_{N^-}) &\rightarrow D(\infty^I \overline{\mathrm{Bun}}_{N^-}), (\mathcal{S}, \mathcal{F}) \mapsto \mathcal{S} \star \mathcal{F}. \end{aligned}$$

The next result follows easily from the constructions.

Lemma 6.1. *For any $\mathcal{F} \in D_I(\mathrm{Fl})$, $\mathcal{G} \in D(\infty \overline{\mathrm{Bun}}_{N^-})$, there exists a canonical isomorphism in $D(\infty^I \overline{\mathrm{Bun}}_{N^-})$:*

$$\mathcal{F} \star \mathrm{For}_I^{G[[t]]}(\mathcal{G}) \simeq \pi_*(\mathcal{F}) \star \mathcal{G},$$

where we recall that $\pi : \mathrm{Fl} \rightarrow \mathrm{Gr}$ is the canonical projection.

In the cases we consider, the convolution product commutes with Verdier duality.

Proposition 6.2. *The morphism $\overleftarrow{h'}$ from ${}^{0,0}\mathcal{H}_{G,N^-}$ (resp. ${}^{I_u,0}\mathcal{H}_{G,N^-}$) to $\infty \overline{\mathrm{Bun}}_{N^-}$ (resp. $\infty^{I_u} \overline{\mathrm{Bun}}_{N^-}$) is ind-proper.*

Proof. This follows from the fact that $\overleftarrow{h'}$ realizes ${}^{0,0}\mathcal{H}_{G,N^-}$ (resp. ${}^{I_u,0}\mathcal{H}_{G,N^-}$) as a fibration over $\infty \overline{\mathrm{Bun}}_{N^-}$ (resp. $\infty^{I_u} \overline{\mathrm{Bun}}_{N^-}$) with typical fiber an ind-projective ind-scheme (namely the affine Grassmannian in the first case, and the affine flag variety in the second case). \square

Corollary 6.3. *For any $\mathcal{F} \in D_{G[[t]]}(\mathcal{Fl}^{\infty})$ and $\mathcal{S} \in D_{G[[t]]}(\mathrm{Gr})$ (resp. $\mathcal{S} \in D_{I_u}(\mathrm{Gr})$), we have*

$$\mathbb{D}(\mathcal{S} \star \mathcal{F}) = \mathbb{D}(\mathcal{S}) \star \mathbb{D}(\mathcal{F}).$$

6.4. Exactness of convolution. We will explain the proof of the following key property.

Theorem 6.4. [ABB⁺05, Theorem 5.2.2] *For any $\nu \in \mathbf{Y}$, the convolution functor*

$$\mathcal{S} \mapsto \mathcal{S} \star \mathcal{IC}_{\nu}^{\infty}$$

from $D_{G[[t]]}(\mathrm{Gr})$ (resp. $D_{I_u}(\mathrm{Gr})$) to $D_{G[[t]]}(\mathcal{Fl}^{\infty})$ (resp. $D_{I_u}(\mathcal{Fl}^{\infty})$) is t -exact.

Proof. Thanks to Corollary 6.3, it is enough to prove that the convolution functor is right t -exact. Let $\mathcal{S} \in {}^p D_{G[[t]]}(\mathrm{Gr})^{\leq 0}$ (resp. $\mathcal{S} \in {}^p D_{I_u}(\mathrm{Gr})^{\leq 0}$) and $\mu \in \mathbf{Y}$. We need to prove that $\overleftarrow{i}_{\mu}^*(\mathcal{S} \star \mathcal{IC}_{\nu}^{\infty})$ lives in non-positive cohomological degrees.

The preimage $(\overleftarrow{h'})^{-1}(\mu \overline{\mathrm{Bun}}_{N^-})$ decomposes into the locally closed substacks

$$X_{\mu',\lambda} := (\overleftarrow{h'})^{-1}(\mu \overline{\mathrm{Bun}}_{N^-}) \cap (\overrightarrow{h'})^{-1}(\mu' \overline{\mathrm{Bun}}_{N^-}) \cap {}^{?,0}\mathcal{H}_{G,N^-}^{\lambda}, \quad \mu' \in \mathbf{Y}, \lambda \in \mathbf{Y}^+,$$

where $? = 0$ (resp. $? = I_u$). In order to conclude, it will be enough to show that:

- (1) The dimension of the fibers of $\overleftarrow{h'}|_{X_{\mu',\lambda}}$ is bounded by $\langle \mu' - \mu + \lambda, \rho \rangle$.
- (2) The restriction of $\mathrm{sw}^*(\mathcal{S}) \boxtimes \mathcal{IC}_{\nu}$ to $X_{\mu',\lambda}$ lives in perverse cohomological degrees less than $-\langle \mu' - \mu + \lambda, \rho \rangle$.

For the first point, the argument is that $\overleftarrow{h'}|_{X_{\mu',\lambda}}$ realizes $X_{\mu',\lambda}$ as the following fibration over ${}_{\mu}^? \overline{\mathrm{Bun}}_{N^-}$:

$$(\mathrm{Gr}^{\lambda} \cap N^-(t)) \cdot (\mu - \mu') \times^{N^-[[t]]} {}_{\mu'}^? \mathcal{N},$$

where, in the case $? = I_u$, the stack ${}_{\mu}^{I_u} \mathcal{N}$ is the $N^-[[t]]$ -torsor ${}_{\mu}^1 \mathcal{N}/N$ over ${}_{\mu}^{I_u} \overline{\mathrm{Bun}}_{N^-}$.

Let us explain the proof of the second point. Let $p_?$ be the identity $\mathrm{Gr} \rightarrow \mathrm{Gr}$ when $? = 0$, and the canonical projection $G((t))/I_u \rightarrow \mathrm{Gr}$ when $? = I_u$. Projecting via $\overrightarrow{h'}$, the locally closed substack $X_{\mu', \lambda}$ identifies with

$$p_?^{-1}(\mathrm{Gr}^{-w_0(\lambda)} \cap N^-((t)) \cdot (\mu' - \mu)) \times^{N^-[[t]]} \mu' \mathcal{N},$$

so that the restriction of $\mathrm{sw}^*(\mathcal{S}) \widetilde{\boxtimes} \mathrm{IC}_\nu$ to $X_{\mu', \lambda}$ identifies with

$$\mathrm{sw}^*(\mathcal{S})|_{p_?^{-1}(\mathrm{Gr}^{-w_0(\lambda)} \cap N^-((t)) \cdot (\mu' - \mu))} \widetilde{\boxtimes} \mathcal{IC}_\nu^{\infty} |_{\mu' \overline{\mathrm{Bun}}_{N^-}}.$$

It is a property of intersection cohomology that the complex $\mathcal{IC}_\nu^{\infty} |_{\mu' \overline{\mathrm{Bun}}_{N^-}}$ lives in negative (and in particular non-positive) perverse cohomological degrees. So we only need to check that the restriction of $\mathrm{sw}^*(\mathcal{S})$ to $p_?^{-1}(\mathrm{Gr}^{-w_0(\lambda)} \cap N^-((t)) \cdot (\mu' - \mu))$ lives in perverse cohomological degrees $\leq -\langle \mu' - \mu + \lambda, \rho \rangle$. Since $*$ -restriction is right t -exact, the complex $\mathrm{sw}^*(\mathcal{S})|_{p_?^{-1}(\mathrm{Gr}^{-w_0(\lambda)})}$ lives in non-positive perverse cohomological degrees. When $? = 0$, the complex $\mathrm{sw}^*(\mathcal{S})|_{\mathrm{Gr}^{-w_0(\lambda)}}$ is constant since it is $G[[t]]$ -equivariant and $\mathrm{Gr}^{-w_0(\lambda)}$ is an orbit. So $\mathrm{sw}^*(\mathcal{S})|_{\mathrm{Gr}^{-w_0(\lambda)} \cap N^-((t)) \cdot (\mu' - \mu)}$ lives in perverse cohomological degrees less than

$$-\mathrm{codim}(\mathrm{Gr}^{-w_0(\lambda)} \cap N^-((t)) \cdot (\mu' - \mu), \mathrm{Gr}^{-w_0(\lambda)}) \leq -\langle \mu' - \mu + \lambda, \rho \rangle.$$

Finally, assume that $? = I_u$. Let H be the stabilizer of $t^{-w_0(\lambda)}$ in $G[[t]]$ and n be a large enough integer such that we have an isomorphism $G_n/H \simeq \mathrm{Gr}^{-w_0(\lambda)}$ (where G_n was defined in §2.3). Then we have an isomorphism of \mathbb{F} -schemes

$$p_{I_u}^{-1}(\mathrm{Gr}^{-w_0(\lambda)}) \simeq G_n \times^H G[[t]]/I_u \simeq G_n \times^H G/U.$$

Since $\mathrm{sw}^*(\mathcal{S})|_{p_{I_u}^{-1}(\mathrm{Gr}^{-w_0(\lambda)})}$ is $G[[t]]$ -equivariant, we deduce that it is of the form

$$\underline{\mathbb{k}}_{\mathrm{Gr}^{-w_0(\lambda)}}[d] \widetilde{\boxtimes} \mathcal{F}$$

for some $\mathcal{F} \in \mathrm{D}(G/U)$ and integer d . Since $\mathrm{sw}^*(\mathcal{S})|_{p_?^{-1}(\mathrm{Gr}^{-w_0(\lambda)})}$ lives in non-positive perverse cohomological degrees, we must have $d \geq \dim(\mathrm{Gr}^{-w_0(\lambda)})$ and $\mathcal{F} \in {}^p\mathrm{D}(G/U)^{\leq 0}$. The identification

$$\mathrm{sw}^*(\mathcal{S})|_{p_{I_u}^{-1}(\mathrm{Gr}^{-w_0(\lambda)} \cap N^-((t)) \cdot (\mu' - \mu))} \simeq \underline{\mathbb{k}}_{\mathrm{Gr}^{-w_0(\lambda)} \cap N^-((t)) \cdot (\mu' - \mu)}[d] \widetilde{\boxtimes} \mathcal{F}$$

then allows to conclude. \square

6.5. Convolution in the spherical case. From here and until the end, we assume that Conjecture 5.1 holds. The present subsection – which is copied from [ABB⁺05, §5.2.3] – settles some results which will be essential when defining the functor from the Hecke category to $\mathrm{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}})$. For any $V \in \mathrm{Rep}(G_{\mathbb{k}}^{\vee})$ and coweight $\mu \in \mathbf{Y}$, we will denote by $V(\mu)$ the associated μ -weight space. For any morphism of \mathbb{F} -schemes of finite type $i : X \rightarrow Y$ and $\mathcal{F} \in \mathrm{D}(Y)$, we put

$$H_Y^k(X, \mathcal{F}) := H^k(Y, i^! \mathcal{F})$$

for all $k \in \mathbb{Z}$. Recall that, by [MV04], we have a canonical isomorphism of complexes of vector spaces for any $\mathcal{G} \in \mathrm{Perv}_{G_{\mathbb{k}}^{\vee}[[t]]}(\mathrm{Gr})$

$$(6.5) \quad H_{N^-((t)) \cdot \mu}^{\bullet}(\mathrm{Gr}, \mathcal{G}) \simeq H_{N^-((t)) \cdot \mu}^{\langle 2\rho, \mu \rangle}(\mathrm{Gr}, \mathcal{G}) \simeq \mathrm{Sat}(\mathcal{G})(\mu).$$

Proposition 6.5. [ABB⁺05, Proposition 5.2.4] *For any $\mathcal{G} \in \mathrm{Perv}_{G[[t]]}(\mathrm{Gr})$ and $\nu \in \mathbf{Y}$, there exist canonical isomorphisms*

$$\mathcal{G} \star \mathcal{IC}_\nu^{\frac{\infty}{2}} \simeq \bigoplus_{\mu \in \mathbf{Y}} \mathcal{IC}_{\nu+\mu}^{\frac{\infty}{2}} \otimes_{\mathbb{k}} \mathrm{Sat}(\mathrm{sw}^* \mathcal{G})(-\mu) \simeq \bigoplus_{\mu \in \mathbf{Y}} \mathcal{IC}_{\nu+\mu}^{\frac{\infty}{2}} \otimes_{\mathbb{k}} \mathrm{Sat}(\mathcal{G})(w_0(\mu)).$$

Moreover, for any $\mathcal{V}, \mathcal{U} \in \text{Perv}_{G[[t]]}(\text{Gr})$, the following diagram commutes

$$(6.6) \quad \begin{array}{ccc} (\mathcal{U} \star \mathcal{V}) \star \mathcal{IC}_{\nu}^{\infty} & \xrightarrow{\sim} & \oplus_{\mu'} (\mathcal{U} \star \mathcal{IC}_{\nu+\mu'}^{\infty}) \otimes \text{Sat}(\mathcal{V})(\mu') \\ \downarrow \sim & & \downarrow \sim \\ \oplus_{\mu} \mathcal{IC}_{\nu+\mu}^{\infty} \otimes (\text{Sat}(\mathcal{U}) \otimes \text{Sat}(\mathcal{V}))(\mu') & \xrightarrow{\sim} & \oplus_{\mu'+\mu''} \mathcal{IC}_{\nu+\mu'+\mu''}^{\infty} \otimes \text{Sat}(\mathcal{U})(\mu'') \otimes \text{Sat}(\mathcal{V})(\mu'). \end{array}$$

Proof. Since the convolution product is t -exact (Theorem 6.4) and the category $\text{Perv}_{G[[t]]}(\mathcal{Fl}^{\infty})$ is semi-simple (Theorem 5.14), we know that there exist vector spaces L_{μ} such that

$$(6.7) \quad \mathcal{G} \star \mathcal{IC}_{\nu}^{\infty} \simeq \bigoplus_{\mu \in \mathbf{Y}} \mathcal{IC}_{\nu+\mu}^{\infty} \otimes_{\mathbb{k}} L_{\mu}.$$

Thanks to Lemma 5.13 (or more easily by the usual theory of perverse sheaves), we have an isomorphism of vector spaces between L_{μ} and

$$(6.8) \quad \text{Hom}(\text{IC}_{\nu+\mu \text{Bun}_{N^-}}, (i_{\nu+\mu})^!(\mathcal{G} \star \mathcal{IC}_{\nu}^{\infty})).$$

Let λ be a dominant cocharacter such that the support of $\text{sw}^*(\mathcal{G})$ is included in $\overline{\text{Gr}}^{-w_0(\lambda)}$. By the proper base change theorem, we have an isomorphism

$$(i_{\nu+\mu})^!(\mathcal{G} \star \mathcal{IC}_{\nu}^{\infty}) \simeq (h')^{\leftarrow}{}_* (\tilde{i}_{\nu+\mu})^!(\text{sw}^*(\mathcal{G}) \boxtimes \mathcal{IC}_{\nu}^{\infty}),$$

where $\tilde{i}^{\nu+\mu}$ is the locally closed inclusion

$$Y := (h')^{\leftarrow}{}^{-1}(\nu+\mu \text{Bun}_{N^-}) \cap (h')^{\rightarrow}{}^{-1}(\leq_{\nu} \overline{\text{Bun}}_{N^-}) \cap {}^{0,0}\overline{\mathcal{H}}_{G,N^-}^{\lambda} \subset (h')^{\leftarrow}{}^{-1}(\leq_{\nu} \overline{\text{Bun}}_{N^-}) \cap {}^{0,0}\overline{\mathcal{H}}_{G,N^-}^{\lambda}$$

(by abuse of notation, we still denote by $\overleftarrow{h'}$ the restriction of $\overleftarrow{h'}$ to $Y \subset {}^{0,0}\mathcal{H}_{G,N^-}$).

For $\beta \leq \lambda$, $\beta \in \mathbf{Y}^+$, denote by Y_{β} the locally closed substack of Y defined by replacing ${}^{0,0}\overline{\mathcal{H}}_{G,N^-}^{\lambda}$ with ${}^{0,0}\mathcal{H}_{G,N^-}^{\beta}$ in the definition of Y , by $j_{\beta} : Y_{\beta} \hookrightarrow {}^{0,0}\mathcal{H}_{G,N^-}$ the locally closed inclusion, and by $(h'_{\beta})^{\leftarrow}{}_{*}$ the restriction of $\overleftarrow{h'}$ to Y_{β} . Then we have the finite stratification $Y = \cup_{\beta} Y_{\beta}$, and projecting via $\overrightarrow{h'}$ induces an isomorphism

$$Y_{\beta} \simeq \left(\text{Gr}^{-w_0(\beta)} \cap N^-(\langle t \rangle) \cdot (-\mu) \right) \times^{N^-[[t]]} \overset{\circ}{\nu}\mathcal{N},$$

where $\overset{\circ}{\nu}\mathcal{N}$ denotes the preimage of νBun_{N^-} under the canonical projection $\nu \mathcal{N} \rightarrow \overline{\nu \text{Bun}}_{N^-}$. Let us denote by $j : \text{Gr}^{-w_0(\beta)} \cap N^-(\langle t \rangle) \cdot (-\mu) \hookrightarrow \text{Gr}$ the locally closed embedding. Now, notice that for all β , we have the isomorphism

$$(h'_{\beta})^{\leftarrow}{}_{*}(j_{\beta})^!(\text{sw}^*(\mathcal{G}) \boxtimes \mathcal{IC}_{\nu}^{\infty}) \simeq (h'_{\beta})^{\leftarrow}{}_{*}(j^!(\text{sw}^*(\mathcal{G})) \boxtimes i'_{\nu} \mathcal{IC}_{\nu}^{\infty}).$$

By adjunction, we have the isomorphism

$$(6.9) \quad \begin{aligned} & \text{Hom}(\text{IC}_{\nu+\mu \text{Bun}_{N^-}}, (h'_{\beta})^{\leftarrow}{}_{*}(j^!(\text{sw}^*(\mathcal{G})) \boxtimes i'_{\nu} \mathcal{IC}_{\nu}^{\infty})) \\ & \simeq \text{Hom}((h'_{\beta})^{\leftarrow}{}^{*}(\text{IC}_{\nu+\mu \text{Bun}_{N^-}}), (j^!(\text{sw}^*(\mathcal{G})) \boxtimes i'_{\nu} \mathcal{IC}_{\nu}^{\infty})) \end{aligned}$$

Since $\nu+\mu \text{Bun}_{N^-}$ is smooth, $\text{IC}_{\nu+\mu \text{Bun}_{N^-}}$ is the constant perverse sheaf of rank one. Moreover the relative dimension of $\nu+\mu \text{Bun}_{N^-}$ compared with νBun_{N^-} is $2\langle \rho, \mu \rangle$. Therefore,

the pullback $(h'_\beta)^*(\overleftarrow{\mathrm{IC}}_{\nu+\mu\mathrm{Bun}_{N^-}})$ is isomorphic to $\mathbb{k}_{Y'_\beta}[2\langle\rho, \mu\rangle]\widetilde{\boxtimes}\mathrm{IC}_{\nu\mathrm{Bun}_{N^-}}$, where we have put $Y'_\beta := \mathrm{Gr}^{-w_0(\beta)} \cap N^-((t)) \cdot (-\mu)$. Thus, (6.9) coincides with

$$\mathrm{Hom}(\mathbb{k}_{Y'_\beta}[2\langle\rho, \mu\rangle], j^!(\mathrm{sw}^*(\mathcal{G}))) \simeq H_{Y'_\beta}^{-2\langle\rho, \mu\rangle}(\mathrm{Gr}, \mathrm{sw}^*(\mathcal{G})).$$

Now, using distinguished triangles associated with open and closed immersions together with an induction on the number of strata of Y , we deduce that (6.8) coincides with

$$H_{N^-((t)) \cdot (-\mu)}^{-2\langle\rho, \mu\rangle}(\mathrm{Gr}, \mathrm{sw}^*(\mathcal{G})).$$

Finally, the above vector space is canonically isomorphic to $\mathrm{Sat}(\mathrm{sw}^*(\mathcal{G}))(-\mu)$ thanks to (6.5). By Proposition 3.6, the latter vector space is isomorphic to $\mathrm{Sat}(\mathcal{G})(w_0(\mu))$.

The commutativity of the diagrams follows from the fact that the geometric Satake equivalence is a tensor functor. \square

6.6. Action of the regular perverse sheaf. We define the \mathbf{Y} -graded object of $\mathrm{Perv}_{G[[t]]}(\mathcal{F}l_{\frac{\infty}{2}})$:

$$\mathcal{IC}_{\frac{\infty}{2}} := \bigoplus_{\lambda \in \mathbf{Y}} \mathcal{IC}_{\lambda}^{\frac{\infty}{2}}.$$

For any $\lambda, \mu \in \mathbf{Y}$, we will define a morphism

$$H_{\mu, \lambda} : \mathcal{R}_\mu \star \mathcal{IC}_{\lambda}^{\frac{\infty}{2}} \rightarrow \mathcal{IC}_{\lambda+\mu}^{\frac{\infty}{2}}.$$

Recall (§3.9) that $\mathcal{R}_\mu := \varinjlim_{\delta} \mathcal{I}_*^{w_0(\mu)+\delta} \star \mathcal{I}_*^{-w_0(\delta)} \in \mathrm{Ind}(\mathrm{Perv}_{G[[t]]}(\mathrm{Gr}))$. For any $\delta \in \mathbf{Y}^+ \cap (-w_0(\mu) + \mathbf{Y}^+)$, we define

$$\begin{aligned} H_{\mu, \lambda}^\delta : (\mathcal{I}_*^{w_0(\mu)+\delta} \star \mathcal{I}_*^{-w_0(\delta)}) \star \mathcal{IC}_{\lambda}^{\frac{\infty}{2}} &\xrightarrow[\mathrm{Prop.6.5}]{\sim} \bigoplus_{\mu'} \mathcal{I}_*^{w_0(\mu)+\delta} \star \mathcal{IC}_{\lambda+\mu'}^{\frac{\infty}{2}} \otimes_{\mathbb{k}} \mathrm{Sat}(\mathcal{I}_*^{-w_0(\delta)})(w_0(\mu')) \\ &\rightarrow \mathcal{I}_*^{w_0(\mu)+\delta} \star \mathcal{IC}_{\lambda-w_0(\delta)}^{\frac{\infty}{2}} \\ &\xrightarrow[\mathrm{Prop.6.5}]{\sim} \bigoplus_{\mu'} \mathcal{IC}_{\lambda-w_0(\delta)+\mu'}^{\frac{\infty}{2}} \otimes_{\mathbb{k}} \mathrm{Sat}(\mathcal{I}_*^{w_0(\mu)+\delta})(w_0(\mu')) \\ &\rightarrow \mathcal{IC}_{\lambda+\mu}^{\frac{\infty}{2}}, \end{aligned}$$

where the second (resp. last) morphism is obtained by projecting onto the 1-dimensional $(-\delta)$ -weight space (resp. $(w_0(\mu) + \delta)$ -weight space), identified with \mathbb{k} via the weight vector $\dot{w}_0 \cdot v_{-w_0(\delta)}$ (resp. $v_{w_0(\mu)+\delta}$). Thanks to the commutativity of the diagram (6.6), the collection of morphisms $(H_{\mu, \lambda}^\delta)_{\delta \in \mathbf{Y}^+ \cap (-w_0(\mu) + \mathbf{Y}^+)}$ is compatible with the transition morphisms in the limit defining \mathcal{R}_μ . So we obtain the desired morphism $H_{\mu, \lambda}$. Moreover, the commutativity of (6.6) implies that the morphisms $(H_{\mu, \lambda})_{\mu, \lambda \in \mathbf{Y}}$ satisfy obvious associativity and unit axioms. Namely:

Proposition 6.6. *The morphisms $(H_{\mu, \lambda})_{\mu, \lambda \in \mathbf{Y}}$ endow $\mathcal{IC}_{\frac{\infty}{2}}$ with a structure of left \mathcal{R} -module.*

6.7. Action of convolution on standard and simple objects.

Proposition 6.7. [ABB⁺05, Proposition 5.4.2] *For $\lambda \in \mathbf{Y}^+$ and $\nu \in \mathbf{Y}$, there are canonical isomorphisms*

$$\mathcal{D}_\lambda \star \Delta_\nu^{\frac{\infty}{2}} \simeq \Delta_{\nu+\lambda}^{\frac{\infty}{2}}, \quad \mathcal{N}_\lambda \star \nabla_\nu^{\frac{\infty}{2}} \simeq \nabla_{\nu+\lambda}^{\frac{\infty}{2}}.$$

Proof. The second isomorphism will follow from the first one by applying Verdier duality together with Corollary 6.3. So we only need to prove that

$$\mathcal{D}_\lambda \star \Delta_\nu^{\frac{\infty}{2}} \simeq \Delta_{\nu+\lambda}^{\frac{\infty}{2}}.$$

One can check (same proof as in [ABB⁺05, Proposition 5.1.5]) that the perverse cohomologies of $\mathcal{D}_\lambda \star \Delta_\nu^{\frac{\infty}{2}}$ are objects of $\text{Perv}_I(\mathcal{F}l^{\frac{\infty}{2}})$. Moreover, we can deduce from the second point of Proposition 5.4 that, for all $w \in W$, $\delta \in \mathbf{Y}$, the perverse cohomologies of the restriction of an object of $\text{Perv}_I(\mathcal{F}l^{\frac{\infty}{2}})$ to ${}^I_{wt_\delta}\text{Bun}_{N^-}$ is of the form $\mathcal{F}\tilde{\boxtimes}\text{IC}_{\nu\overline{\text{Bun}}_{N^-}}$, for some $\mathcal{F} \in \text{Perv}_N(G/B^-)$. Thus, to prove that $\mathcal{D}_\lambda \star \Delta_\nu^{\frac{\infty}{2}} \simeq \Delta_{\nu+\lambda}^{\frac{\infty}{2}}$, it suffices to check that the stalk of $\mathcal{D}_\lambda \star \Delta_\nu^{\frac{\infty}{2}}$ at any point of ${}^I_{wt_\delta}\overline{\text{Bun}}_{N^-}$ is zero when $wt_\delta \neq t_{\nu+\lambda}$, and is the (perversly shifted) constant sheaf otherwise. The computation of the latter stalk can then be conducted as in the proof of [ABB⁺05, Proposition 5.4.2], where the authors show that it amounts to computing the cohomology of an empty set when $wt_\delta \neq t_{\nu+\lambda}$, and of a point scheme otherwise. \square

Next, we exhibit the commutativity of certain diagrams (cf. [ABB⁺05, (49),(50)]), which will turn out to be useful in §8. Below, those diagrams will involve the following canonical morphisms. For any $\mu, \lambda \in \mathbf{Y}^+$, $\nu \in \mathbf{Y}$, we have:

$$(6.10) \quad \nabla_\lambda \star \mathcal{IC}_\nu^{\frac{\infty}{2}} \xrightarrow{\sim} \mathcal{N}_\lambda \star \text{IC}_\nu^{\frac{\infty}{2}} \rightarrow \mathcal{N}_\lambda \star \nabla_\nu^{\frac{\infty}{2}},$$

where the first isomorphism follows from Lemma 6.1. Moreover, we will denote by

$$b_\lambda : \mathcal{I}_*^\lambda \rightarrow \nabla_\lambda$$

the morphism established in (3.10), and by

$$a_\nu : \mathcal{IC}_\nu^{\frac{\infty}{2}} \rightarrow \nabla_\nu^{\frac{\infty}{2}}$$

the canonical morphism in $\text{Perv}_{I_u}(\mathcal{F}l^{\frac{\infty}{2}})$, where we have identified $\mathcal{IC}_\nu^{\frac{\infty}{2}}$ with $\text{IC}_\nu^{\frac{\infty}{2}}$ through the functor $\text{For}_{I_u}^{G[[t]]}$ (cf. Proposition 5.16).

Proposition 6.8. *For any $\mu, \lambda \in \mathbf{Y}^+$, $\nu \in \mathbf{Y}$, the following diagrams are commutative:*

$$(6.11) \quad \begin{array}{ccc} \mathcal{I}_*^\lambda \star \mathcal{IC}_\nu^{\frac{\infty}{2}} & \xrightarrow{b_\lambda \star \text{id}} & \nabla_\lambda \star \mathcal{IC}_\nu^{\frac{\infty}{2}} \xrightarrow{(6.10)} \mathcal{N}_\lambda \star \nabla_\nu^{\frac{\infty}{2}} \\ \text{Prop. 6.5} \downarrow \sim & & \text{Prop. 6.7} \downarrow \sim \\ \oplus_{\nu'} \mathcal{IC}_{\nu+\nu'}^{\frac{\infty}{2}} \otimes \text{Sat}(\mathcal{I}_*^\lambda)(w_0(\nu')) & \longrightarrow & \mathcal{IC}_{\nu+\lambda}^{\frac{\infty}{2}} \xrightarrow{a_{\nu+\lambda}} \nabla_{\nu+\lambda}^{\frac{\infty}{2}} \end{array}$$

$$(6.12) \quad \begin{array}{ccc} \nabla_\lambda \star \mathcal{IC}_\nu^{\frac{\infty}{2}} & \longrightarrow & \nabla_\lambda \star \mathcal{I}_*^\mu \star \mathcal{I}_*^{-w_0(\mu)} \star \mathcal{IC}_\nu^{\frac{\infty}{2}} \\ \downarrow (6.10) & & \downarrow \text{Prop. 6.5} \\ \mathcal{N}_\lambda \star \nabla_\nu^{\frac{\infty}{2}} & & \nabla_\lambda \star \mathcal{I}_*^\mu \star \mathcal{IC}_{\nu-\mu}^{\frac{\infty}{2}} \\ \downarrow \sim & & \downarrow (\text{id} \star b_\mu) \star \text{id} \\ \nabla_{\lambda+\nu}^{\frac{\infty}{2}} & \longleftarrow & \nabla_{\lambda+\mu} \star \mathcal{IC}_{\nu-\mu}^{\frac{\infty}{2}}, \end{array}$$

where the bottom arrow $\nabla_{\lambda+\mu} \star \mathcal{IC}_{\nu-\mu}^{\frac{\infty}{2}} \rightarrow \nabla_{\lambda+\nu}^{\frac{\infty}{2}}$ is the composition

$$\nabla_{\lambda+\mu} \star \mathcal{IC}_{\nu-\mu}^{\frac{\infty}{2}} \xrightarrow{(6.10)} \mathcal{N}_{\lambda+\mu} \star \nabla_{\nu-\mu}^{\frac{\infty}{2}} \xrightarrow[\text{Prop. 6.7}]{\sim} \nabla_{\lambda+\nu}^{\frac{\infty}{2}}.$$

Proof. Notice that, for any $\nu \in \mathbf{Y}$, the space of morphisms between $\mathrm{IC}_{\nu}^{\infty}$ and ∇_{ν}^{∞} is one dimensional and canonical. The fact that the two morphisms $\mathcal{I}_*^{\lambda} \star \mathrm{IC}_{\nu}^{\infty} \rightarrow \nabla_{\nu+\lambda}^{\infty}$ (resp. $\nabla_{\lambda} \star \mathrm{IC}_{\nu}^{\infty} \rightarrow \nabla_{\nu+\lambda}^{\infty}$) considered in the first (resp. second) diagram coincide then follows from the construction of the isomorphisms of propositions 6.5 and 6.7. \square

From Proposition 6.7 we deduce (by replacing B with B^- in the arguments) that we have a canonical isomorphism for $\lambda \in \mathbf{Y}^+$, $\nu' \in \mathbf{Y}$

$$\mathcal{D}_{-\lambda} \star \Delta_{w_0 t_{\nu'}}^{\infty} \simeq \Delta_{w_0 t_{\nu'} - w_0(\lambda)}^{\infty}.$$

Thanks to the first point of Lemma 3.4, we deduce that there is a canonical isomorphism

$$(6.13) \quad \mathcal{N}_{\lambda} \star \Delta_{w_0 t_{\nu'}}^{\infty} \simeq \Delta_{w_0 t_{\nu'} + w_0(\lambda)}^{\infty}.$$

At the present moment, we are not able to prove the following result (which follows from [ABB⁺05, Lemma 5.4.5] in the case of \mathbb{Q}_{ℓ} -sheaves), but hope to come back to this question in future work.

Conjecture 6.1. *For any $\nu \in \mathbf{Y}$, there exists a non-zero morphism*

$$\tilde{a}_{\nu} : \mathrm{IC}_{\nu}^{\infty} \rightarrow \Delta_{w_0 t_{\nu} + 2\zeta}^{\infty}$$

such that the following diagrams commute for all $\nu \in \mathbf{Y}$, $\lambda, \mu \in \mathbf{Y}^+$:

$$(6.14) \quad \begin{array}{ccccc} \mathcal{I}_*^{\lambda} \star \mathrm{IC}_{\nu}^{\infty} & \xrightarrow{b_{\lambda} \star \mathrm{id}} & \nabla_{\lambda} \star \mathrm{IC}_{\nu}^{\infty} & \longrightarrow & \mathcal{N}_{\lambda} \star \Delta_{w_0 t_{\nu} + 2\zeta}^{\infty} \\ \downarrow \text{Prop. 6.5} \sim & & & & \downarrow \text{Prop. 6.7} \sim \\ \oplus_{\nu'} \mathrm{IC}_{\nu+\nu'}^{\infty} \otimes \mathrm{Sat}(\mathcal{I}_*^{\lambda})(w_0(\nu')) & \longrightarrow & \mathrm{IC}_{\nu+w_0(\lambda)}^{\infty} & \xrightarrow{\tilde{a}_{\nu+w_0(\lambda)}} & \Delta_{w_0 t_{\nu} + 2\zeta + w_0(\lambda)}^{\infty} \end{array}$$

$$(6.15) \quad \begin{array}{ccc} \nabla_{\lambda} \star \mathrm{IC}_{\nu}^{\infty} & \longrightarrow & \nabla_{\lambda} \star \mathcal{I}_*^{\mu} \star \mathcal{I}_*^{-w_0(\mu)} \star \mathrm{IC}_{\nu}^{\infty} \\ \downarrow & & \downarrow \text{Prop. 6.5} \\ \mathcal{N}_{\lambda} \star \Delta_{w_0 t_{\nu} + 2\zeta}^{\infty} & & \nabla_{\lambda} \star \mathcal{I}_*^{\mu} \star \mathrm{IC}_{\nu-\mu}^{\infty} \\ \downarrow \text{(6.13)} \sim & & \downarrow \\ \Delta_{w_0 t_{\nu} + 2\zeta + w_0(\lambda)}^{\infty} & \longleftarrow & \nabla_{\lambda+\mu} \star \mathrm{IC}_{\nu-\mu}^{\infty}. \end{array}$$

Theorem 6.9. [ABB⁺05, Theorem 5.3.2] *For any $w \in W_{\mathrm{ext}}^{\mathrm{res}}$ and $\nu \in \mathbf{Y}$, we have*

$$\mathrm{L}_w \star \mathrm{IC}_{\nu}^{\infty} \simeq \mathrm{IC}_{w t_{\nu}}^{\infty}.$$

Proof. By definition we have

$$\mathrm{L}_w \star \mathrm{IC}_{\nu}^{\infty} := (h')^{\leftarrow} \star (\mathrm{sw}^*(\mathrm{L}_w) \boxtimes \mathrm{IC}_{\nu}^{\infty}),$$

where $h' : I^0 \mathcal{H}_{G, N^-} \rightarrow I^{\infty} \overline{\mathrm{Bun}}_{N^-}$. Denote by X (resp. Y) the closed substack of $I^0 \mathcal{H}_{G, N^-}$ (resp. of $I^{\infty} \overline{\mathrm{Bun}}_{N^-}$) defined as the support of $\mathrm{sw}^*(\mathrm{L}_w) \boxtimes \mathrm{IC}_{\nu}^{\infty}$ (resp. of $\mathrm{IC}_{w t_{\nu}}^{\infty}$). Consider the stratifications

$$X = \bigsqcup_{(u, \mu) \in W_{\mathrm{ext}} \times \mathbf{Y}} \mathcal{O}_u \times^{N^-[[\ell]]} I_{\mu} \mathcal{N}, \quad Y = \bigsqcup_{vt_{\lambda} \in W_{\mathrm{ext}}} I_{vt_{\lambda}} \overline{\mathrm{Bun}}_{N^-},$$

where $\mathbb{O}_u \subset \text{Fl}$ denotes the $G[[t]]$ -orbit associated with an element $u \in W_{\text{ext}}$. By the proof of [ABB⁺05, Theorem 5.3.2], the morphism

$$\overleftarrow{h}'|_{\overleftarrow{h}'^{-1}(Y)} : X \rightarrow Y$$

is stratified small with respect to the dense open subset ${}^I_{w\nu} \overline{\text{Bun}}_{N^-} \subset Y$. Since \overleftarrow{h}' is moreover proper and $\text{sw}^*(L_w) \widetilde{\boxtimes} \mathcal{IC}_{\nu}^{\infty}$ coincides with the IC-sheaf on X , we deduce that $L_w \star \mathcal{IC}_{\nu}^{\infty}$ coincides with the IC-sheaf on Y (see for instance [Ach21, Proposition 3.8.10]). \square

7. IWAHORI-WHITTAKER OBJECTS

7.1. Iwahori-Whittaker for the semi-infinite flag variety. For any \mathbb{F} -stack locally of finite type \mathcal{X} acted on by N^- , the category $\text{D}_{(N^-, \psi)}(\mathcal{X})$ is defined as the full subcategory of $\text{D}(\mathcal{X})$ consisting of objects \mathcal{F} such that there exists an isomorphism $a^* \mathcal{F} \simeq \psi^* \mathcal{L}_{\text{AS}} \boxtimes \mathcal{F}$, where $a : N^- \times \mathcal{X} \rightarrow \mathcal{X}$ is the morphism defining the action (and recall that $\psi : N^- \rightarrow \mathbb{G}_a$ was defined in §3.6). This category is endowed with a natural perverse t -structure, whose heart will be denoted by $\text{Perv}_{(N^-, \psi)}(\mathcal{X})$, and which is sent to $\text{Perv}(\mathcal{X})$ by the forgetful functor $\text{For}_{(N^-, \psi)} : \text{D}_{(N^-, \psi)}(\mathcal{X}) \rightarrow \text{D}(\mathcal{X})$.

First consider the category $\text{Perv}_{(N^-, \psi)}(G/B^-)$. It is a classical fact that the only N^- -orbit supporting a non-zero Whittaker local system is the open dense orbit, i.e. the orbit associated with w_0 . Thus, the category $\text{Perv}_{(N^-, \psi)}(G/B^-)$ contains a unique simple object, denoted by ψ_{G/B^-} , defined as the Goresky-Macpherson extension of the rank-one Whittaker local system on this orbit. Next, we define the category $\text{Perv}_{(I_u^-, \psi)}(\mathcal{Fl}^{\frac{\infty}{2}})$ as the full subcategory of $\text{Perv}_{(N^-, \psi)}({}^1_{\infty} \overline{\text{Bun}}_{N^-})$ consisting of T -equivariant perverse sheaves satisfying the factorization and equivariance conditions (cf. §5.1). The same arguments as the one used in the proof of Proposition 5.4 show that the simple objects of $\text{Perv}_{(I_u^-, \psi)}(\mathcal{Fl}^{\frac{\infty}{2}})$ are of the form

$$\text{IC}_{\nu}^{\psi} := (\tilde{i}_{\nu})_* \left(\psi_{G/B^-} \widetilde{\boxtimes} \text{IC}_{\nu \overline{\text{Bun}}_{N^-}} \right), \quad \nu \in \mathbf{Y}.$$

We have the convolution functor

$$\text{Perv}_{\mathcal{IW}}(\text{Gr}) \times \text{Perv}_{G[[t]]}(\mathcal{Fl}^{\frac{\infty}{2}}) \rightarrow \text{Perv}_{(I_u^-, \psi)}(\mathcal{Fl}^{\frac{\infty}{2}}).$$

Recall (§2.1) that we have picked $\zeta \in \mathbf{Y}$ such that $\langle \alpha, \zeta \rangle = 1$ for all $\alpha \in \mathfrak{R}_s$. The proof of the next result is the same as that of Theorem 6.9.

Proposition 7.1. [ABB⁺05, Theorem 5.3.5] *For any $\nu \in \mathbf{Y}$, we have*

$$L_{\zeta}^{\mathcal{IW}} \star \mathcal{IC}_{\nu}^{\infty} \simeq \text{IC}_{\nu+\zeta}^{\psi}.$$

For any $\nu \in \mathbf{Y}$, we have the corresponding standard and costandard objects

$$\nabla_{\nu}^{\psi} := (\tilde{i}_{\nu})_* \left(\psi_{G/B^-} \widetilde{\boxtimes} \text{IC}_{\nu \overline{\text{Bun}}_{N^-}} \right), \quad \Delta_{\nu}^{\psi} := (\tilde{i}_{\nu})! \left(\psi_{G/B^-} \widetilde{\boxtimes} \text{IC}_{\nu \overline{\text{Bun}}_{N^-}} \right).$$

By the proof of [ABB⁺05, Corollary 4.4.5], we know that the inclusion of ${}^I_u \overline{\text{Bun}}_{N^-}$ into ${}^{\leq \nu} \overline{\text{Bun}}_{N^-}$ is affine, so that both objects above are perverse sheaves. They are then easily seen to be objects of $\text{Perv}_{(I_u^-, \psi)}(\mathcal{Fl}^{\frac{\infty}{2}})$.

Theorem 7.2. [ABB⁺05, Theorem 4.4.10] *For any $\nu \in \mathbf{Y}$, the canonical morphisms*

$$\Delta_{\nu}^{\psi} \rightarrow \text{IC}_{\nu}^{\psi} \rightarrow \nabla_{\nu}^{\psi}$$

are isomorphisms.

Proof. By the arguments of *loc. cit.*, we are reduced to showing that the Euler characteristic of the stalk of $\mathrm{IC}_{\nu+\zeta}^{\psi}$ at any point of the locally closed substack $\mathrm{ev}_{\nu'}^{-1}(N^- \cdot w_0)$, for $\nu' < \nu + \zeta$, is zero. By Proposition 7.1, this is equivalent to computing the stalk of $L_{\zeta}^{\mathcal{I}\mathcal{W}} \star \mathcal{I}\mathcal{C}_{\nu}^{\infty}$. Using similar arguments as in the proof of Proposition 6.5, this stalk identifies with the extension of a finite number of certain complexes of \mathbb{k} -vector spaces $K_{\nu''}$, $\nu'' < \nu$. Each $K_{\nu''}$ is defined as the stalk of the direct image under

$$\overleftarrow{h}' : (\overleftarrow{h}')^{-1}(1_{\nu'} \overline{\mathrm{Bun}}_{N^-}) \cap (\overrightarrow{h}')^{-1}(\nu'' \overline{\mathrm{Bun}}_{N^-}) \cap {}^{1,0}\overline{\mathcal{H}}_{G,N^-}^{\zeta} \rightarrow 1_{\nu'} \overline{\mathrm{Bun}}_{N^-}$$

of the restriction of $L_{\zeta}^{\mathcal{I}\mathcal{W}} \widetilde{\boxtimes} \mathcal{I}\mathcal{C}_{\nu}^{\infty}$ to the above locally closed substack. Moreover, it follows from the constructions that the projection \overrightarrow{h}' identifies that same locally closed substack with

$$(N((t)) \cdot w_0(\nu' - \nu'') \cap I^- \cdot (w_0(\zeta))) \times^{N^-[[t]]} \nu'' \mathcal{N}.$$

Therefore

$$\begin{aligned} K_{\nu''} &\simeq H_c^{\bullet}(N((t)) \cdot w_0(\nu' - \nu'') \cap I^- \cdot (w_0(\zeta)), \mathcal{L} \otimes_{\mathbb{k}} \mathcal{I}\mathcal{C}_{\nu}^{\infty}|_{\nu'' \mathrm{Bun}_{N^-}}) \\ &\simeq H_c^{\bullet}(N((t)) \cdot w_0(\nu' - \nu'') \cap I^- \cdot (w_0(\zeta)), \mathcal{L}) \otimes_{\mathbb{k}} \mathcal{I}\mathcal{C}_{\nu}^{\infty}|_{\nu'' \mathrm{Bun}_{N^-}} \end{aligned}$$

where \mathcal{L} denotes the rank-one Iwahori-Whittaker local system on $I^- \cdot (w_0(\zeta))$, and $\mathcal{I}\mathcal{C}_{\nu}^{\infty}|_{\nu'' \mathrm{Bun}_{N^-}}$ is meant to be the stalk of $\mathcal{I}\mathcal{C}_{\nu}^{\infty}$ at any point of $\nu'' \mathrm{Bun}_{N^-}$.

On the other hand, let us put $\mu_r := (\ell^r - 1)\zeta$ for any integer $r \geq 0$. The canonical morphisms

$$\mathcal{I}_1^{\mu_r} \rightarrow \mathcal{I}\mathcal{C}^{\mu_r} \rightarrow \mathcal{I}_*^{\mu_r}$$

are isomorphisms for any r thanks to Lemma 3.7, so that (by applying the functor $\Delta_{\zeta}^{\mathcal{I}\mathcal{W}} \star (-)$) the canonical morphisms

$$\Delta_{\mu_r+\zeta}^{\mathcal{I}\mathcal{W}} \rightarrow L_{\mu_r+\zeta}^{\mathcal{I}\mathcal{W}} \rightarrow \nabla_{\mu_r+\zeta}^{\mathcal{I}\mathcal{W}}$$

are also isomorphisms. In particular, the IC-sheaf $L_{\mu_r+\zeta}^{\mathcal{I}\mathcal{W}} \in \mathrm{Perv}_{\mathcal{I}\mathcal{W}}(\mathrm{Gr})$ is clean.

Notice that the map

$$f : \nu'' \mapsto \mu_r - (\nu - \nu'')$$

induces a bijection between the sets $\{\nu'', \nu'' \leq \nu\}$ and $\{\mu'', \mu'' \leq \mu_r\}$. Put $\mu'_r := f(\nu')$. The stalk of $L_{\mu_r+\zeta}^{\mathcal{I}\mathcal{W}} \simeq L_{\zeta}^{\mathcal{I}\mathcal{W}} \star \mathcal{I}\mathcal{C}^{\mu_r}$ at $[t^{\mu'_r}] \in Y_{\mu'_r}$ can be written as an extension of certain complexes $\widetilde{K}_{\mu''}$, for $\mu'' \leq \mu_r$ (only a finite number of complexes will be non-zero). Each $\widetilde{K}_{\mu''}$ is defined as the stalk at $[t^{\mu'_r}]$ of the direct image under the multiplication morphism

$$(7.1) \quad m : I^- \cdot (w_0(\zeta)) \times^{G((t))} \mathrm{Gr}^{\mu''} \rightarrow \mathrm{Gr}$$

of the restriction of $L_{\zeta}^{\mathcal{I}\mathcal{W}} \widetilde{\boxtimes} \mathcal{I}\mathcal{C}^{\mu_r}$ to $I^- \cdot (w_0(\zeta)) \times^{G((t))} \mathrm{Gr}^{\mu''}$. By [ABB⁺05, Remark 5.3.3], the preimage of $[t^{\mu'_r}]$ by the morphism (7.1) is isomorphic to $N((t)) \cdot w_0(\mu'_r - \mu'') \cap I^- \cdot (w_0(\zeta))$ when r is large enough (so that μ_r is dominant enough). Thus

$$\begin{aligned} \widetilde{K}_{\mu''} &\simeq H_c^{\bullet}(N((t)) \cdot w_0(\mu'_r - \mu'') \cap I^- \cdot (w_0(\zeta)), \mathcal{L} \otimes_{\mathbb{k}} \mathcal{I}\mathcal{C}^{\mu_r}|_{\mathrm{Gr}^{\mu''}}) \\ &\simeq H_c^{\bullet}(N((t)) \cdot w_0(\mu'_r - \mu'') \cap I^- \cdot (w_0(\zeta)), \mathcal{L}) \otimes_{\mathbb{k}} \mathcal{I}\mathcal{C}^{\mu_r}|_{\mathrm{Gr}^{\mu''}} \end{aligned}$$

where $\mathcal{I}\mathcal{C}^{\mu_r}|_{\mathrm{Gr}^{\mu''}}$ is meant to be the stalk of $\mathcal{I}\mathcal{C}^{\mu_r}$ at any point of $\mathrm{Gr}^{\mu''}$.

Since $\mu_r - f(\nu'') = \nu - \nu''$ by construction, we can take r large enough so that

$$\mathcal{I}\mathcal{C}_{\nu}^{\infty}|_{\nu'' \mathrm{Bun}_{N^-}} \simeq \mathcal{I}\mathcal{C}^{\mu_r}|_{\mathrm{Gr}^{f(\nu'')}}$$

for all $\nu'' < \nu$ such that $K_{\nu''}$ is non-zero by Corollary 5.15. Moreover $\mu'_r - f(\nu'') = \nu' - \nu''$ by definition of f . So we get an isomorphism

$$K_{\nu''} \simeq \tilde{K}_{f(\nu'')} \quad \forall \nu'' \leq \nu.$$

So the Euler characteristic of the stalk of $\mathrm{IC}_{\nu+\zeta}^{\psi}$ at any point of $\mathrm{ev}_{\nu'}^{-1}(N^- \cdot w_0)$ coincides with the Euler characteristic of the stalk of $\mathrm{L}_{\mu_r+\zeta}^{\mathcal{W}}$ at $[t^{\mu'_r}]$. But $\mu'_r < \mu_r + \zeta$, so that the latter stalk is zero by cleanness. This concludes the proof. \square

7.2. Partial integrability and consequences. We denote by

$$(7.2) \quad \mathrm{av}_{!(N^-, \psi)}, \mathrm{av}_{*(N^-, \psi)} : \mathrm{D}(\mathbb{1}_{\infty} \overline{\mathrm{Bun}}_{N^-}) \rightarrow \mathrm{D}_{(I_{\bar{u}}^-, \psi)}(\mathbb{1}_{\infty} \overline{\mathrm{Bun}}_{N^-})$$

the functors which are respectively left and right adjoint to the forgetful functor. They are defined by

$$\mathrm{av}_{!(N^-, \psi)}(\mathcal{F}) := a_!(\psi^* \mathcal{L}_{\mathrm{AS}} \boxtimes \mathcal{F})[2\dim(N^-)], \quad \mathrm{av}_{*(N^-, \psi)}(\mathcal{F}) := a_*(\psi^* \mathcal{L}_{\mathrm{AS}} \boxtimes \mathcal{F}),$$

where $a : N^- \times \mathbb{1}_{\infty} \overline{\mathrm{Bun}}_{N^-} \rightarrow \mathbb{1}_{\infty} \overline{\mathrm{Bun}}_{N^-}$ is the action morphism. Moreover we denote by

$$\mathrm{Av}_{!(I_{\bar{u}}, \psi)}, \mathrm{Av}_{*(I_{\bar{u}}, \psi)} : \mathrm{D}(\mathbb{1}_{\infty}^{\bar{u}} \overline{\mathrm{Bun}}_{N^-}) \rightarrow \mathrm{D}_{(I_{\bar{u}}^-, \psi)}(\mathbb{1}_{\infty} \overline{\mathrm{Bun}}_{N^-})$$

the functors obtained by composing (7.2) with the forgetful functor $\mathrm{D}(\mathbb{1}_{\infty}^{\bar{u}} \overline{\mathrm{Bun}}_{N^-}) \rightarrow \mathrm{D}(\mathbb{1}_{\infty} \overline{\mathrm{Bun}}_{N^-})$.

Let us denote by

$$\mathrm{For}_{I_{\bar{u}}}^I : \mathrm{D}(\mathbb{1}_{\infty}^{\bar{u}} \overline{\mathrm{Bun}}_{N^-}) \rightarrow \mathrm{D}(\mathbb{1}_{\infty} \overline{\mathrm{Bun}}_{N^-})$$

the forgetful functor. From the definitions, we have isomorphisms of functors from $\mathrm{D}(\mathbb{1}_{\infty}^{\bar{u}} \overline{\mathrm{Bun}}_{N^-})$ to $\mathrm{D}_{(I_{\bar{u}}^-, \psi)}(\mathbb{1}_{\infty} \overline{\mathrm{Bun}}_{N^-})$:

$$(7.3) \quad \mathrm{Av}_{!(I_{\bar{u}}, \psi)}[-2\dim(N^-)] \circ \mathrm{For}_{I_{\bar{u}}}^I \simeq \Delta_{\psi} \star (-), \quad \mathrm{Av}_{*(I_{\bar{u}}, \psi)} \circ \mathrm{For}_{I_{\bar{u}}}^I \simeq \nabla_{\psi} \star (-)$$

where $\Delta_{\psi} \in \mathrm{Perv}_{(N^-, \psi)}(G/B)$ (resp. ∇_{ψ}) is the $!$ -extension (resp. $*$ -extension) of the rank one Whittaker local system on the dense N^- -orbit in G/B (and the convolution product \star is defined as in Remark 5.20).

Lemma 7.3. [ABB⁺05, Lemma 4.4.12] *The canonical morphism of functors*

$$\mathrm{Av}_{!(I_{\bar{u}}, \psi)}[-\dim(N^-)] \rightarrow \mathrm{Av}_{*(I_{\bar{u}}, \psi)}[\dim(N^-)]$$

is an isomorphism. Moreover, the resulting functor

$$\mathrm{D}(\mathbb{1}_{\infty}^{\bar{u}} \overline{\mathrm{Bun}}_{N^-}) \rightarrow \mathrm{D}_{(I_{\bar{u}}^-, \psi)}(\mathbb{1}_{\infty} \overline{\mathrm{Bun}}_{N^-})$$

is t -exact.

We will denote by

$$\mathrm{Av}_{(I_{\bar{u}}, \psi)} : \mathrm{D}(\mathbb{1}_{\infty}^{\bar{u}} \overline{\mathrm{Bun}}_{N^-}) \rightarrow \mathrm{D}_{(I_{\bar{u}}^-, \psi)}(\mathbb{1}_{\infty} \overline{\mathrm{Bun}}_{N^-})$$

the obtained t -exact functor.

Proof. First, notice that the category $\mathrm{D}(\mathbb{1}_{\infty}^{\bar{u}} \overline{\mathrm{Bun}}_{N^-})$ is generated (as a triangulated category) by the essential image of the forgetful functor

$$\mathrm{For}_{I_{\bar{u}}}^I : \mathrm{D}(\mathbb{1}_{\infty}^{\bar{u}} \overline{\mathrm{Bun}}_{N^-}) \rightarrow \mathrm{D}(\mathbb{1}_{\infty} \overline{\mathrm{Bun}}_{N^-}).$$

Thus, the first point can be proved for the corresponding functors composed with $\mathrm{For}_{I_{\bar{u}}}^I$. In this case, we have the isomorphisms of functors (7.3), so that the morphism

$$\mathrm{Av}_{!(I_{\bar{u}}, \psi)}[-\dim(N^-)] \circ \mathrm{For}_{I_{\bar{u}}}^I \rightarrow \mathrm{Av}_{*(I_{\bar{u}}, \psi)}[\dim(N^-)] \circ \mathrm{For}_{I_{\bar{u}}}^I$$

is induced by the canonical morphism $\Delta_\psi \rightarrow \nabla_\psi$. But the latter morphism is an isomorphism, as both objects are easily seen to be isomorphic to the simple perverse sheaf $\psi_{G/B}$.

Finally, we deduce that the resulting functor is t -exact. Indeed, $\text{av}_{*,(N^-, \psi)}$ is right t -exact since it is defined as the $*$ -pushforward along an affine morphism [BBDG18, Théorème 4.1.1], and $\text{av}_{!,(N^-, \psi)}$ is left t -exact since it is defined as the $!$ -pushforward along an affine morphism [BBDG18, Corollaire 4.1.2]. \square

An object of $\text{Perv}_{I_u}(\mathcal{Fl}_{\frac{\infty}{2}})$ is called *partially integrable* if all of its irreducible subquotients are of the form $\text{IC}_{wt_\nu}^{\frac{\infty}{2}}$, with $\nu \in \mathbf{Y}$ and $w \neq w_0$. We let ${}^f\text{Perv}_{I_u}(\mathcal{Fl}_{\frac{\infty}{2}})$ denote the resulting Serre quotient category obtained by quotienting $\text{Perv}_{I_u}(\mathcal{Fl}_{\frac{\infty}{2}})$ by the subcategory of partially integrable objects. The next result parallels Proposition 3.26.

Proposition 7.4. *For any $wt_\nu \in W_{\text{ext}}$ we have isomorphisms*

$$\text{Av}_{I_u, \psi}(\nabla_{wt_\nu}^{\frac{\infty}{2}}) \simeq \nabla_\psi, \quad \text{Av}_{I_u, \psi}(\Delta_{wt_\nu}^{\frac{\infty}{2}}) \simeq \Delta_\psi.$$

Proof. By (7.3), we have

$$\text{Av}_{I_u, \psi}(\nabla_{wt_\nu}^{\frac{\infty}{2}}) \simeq \nabla_\psi \star \nabla_{wt_\nu}^{\frac{\infty}{2}},$$

where $\nabla_{wt_\nu}^{\frac{\infty}{2}}$ is seen as an object of $\text{D}(\infty \overline{\text{Bun}}_{N^-})$. Recall that, by definition

$$\nabla_{wt_\nu}^{\frac{\infty}{2}} = (\bar{i}_\nu)_* \left(\tilde{\nabla}_w \boxtimes \text{IC}_{\nu \overline{\text{Bun}}_{N^-}} \right).$$

So one can check that

$$\nabla_\psi \star \nabla_{wt_\nu}^{\frac{\infty}{2}} \simeq (\bar{i}_\nu)_* \left((\nabla_\psi \star \tilde{\nabla}_w) \boxtimes \text{IC}_{\nu \overline{\text{Bun}}_{N^-}} \right).$$

As in [BY13, Lemma 4.4.8], we have an isomorphism $\nabla_\psi \star \tilde{\nabla}_w \simeq \psi_{G/B}$, concluding the proof of the first isomorphism. The proof of the second one is similar. \square

The proof of the next result is done exactly as in [ABB⁺05, Proposition 4.4.13].

Proposition 7.5. (1) *The functor*

$$\text{Av}_{(I_u, \psi)} : \text{Perv}_{I_u}(\mathcal{Fl}_{\frac{\infty}{2}}) \rightarrow \text{Perv}_{(I_u^-, \psi)}(\mathcal{Fl}_{\frac{\infty}{2}})$$

factors through ${}^f\text{Perv}_{I_u}(\mathcal{Fl}_{\frac{\infty}{2}})$.

(2) *The resulting functor*

$${}^f\text{Perv}_{I_u}(\mathcal{Fl}_{\frac{\infty}{2}}) \rightarrow \text{Perv}_{(I_u^-, \psi)}(\mathcal{Fl}_{\frac{\infty}{2}})$$

is exact and faithful.

8. THE CONVOLUTION FUNCTOR

8.1. Construction of the functor. Here we follow [ABB⁺05, §6.1.1]. We recall the construction of a functor

$$\text{Conv}^{\text{Hecke}} : \text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R}) \rightarrow \text{Ind}(\text{Perv}_{I_u}(\mathcal{Fl}_{\frac{\infty}{2}})),$$

which will (conjecturally) restrict to an equivalence on the categories of finite length objects. Let $\mathcal{S} \in \text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$. For any $\lambda, \mu \in \mathbf{Y}$, consider the morphisms

$$\alpha_{\lambda, \mu} : \mathcal{S}_\lambda \star \mathcal{R}_{-\mu} \star \text{IC}_{\mu-\lambda}^{\frac{\infty}{2}} \rightarrow \mathcal{S}_{\lambda-\mu} \star \text{IC}_{\mu-\lambda}^{\frac{\infty}{2}}$$

$$\beta_{\lambda, \mu} : \mathcal{S}_\lambda \star \mathcal{R}_{-\mu} \star \text{IC}_{\mu-\lambda}^{\frac{\infty}{2}} \rightarrow \mathcal{S}_\lambda \star \text{IC}_{-\lambda}^{\frac{\infty}{2}},$$

induced respectively by the right \mathcal{R} -module structure on \mathcal{S} and the left \mathcal{R} -module structure on $\text{IC}_{\frac{\infty}{2}}$ (cf. §6.6).

We then define $\text{Conv}^{\text{Hecke}}(\mathcal{S})$ as the coequalizer of the following diagram:

$$\bigoplus_{\lambda, \mu} \mathcal{S}_\lambda \star \mathcal{R}_{-\mu} \star \mathcal{IC}_{\mu-\lambda}^{\frac{\infty}{2}} \rightrightarrows \bigoplus_{\nu} \mathcal{S}_\nu \star \mathcal{IC}_{-\nu}^{\frac{\infty}{2}},$$

where the two arrows correspond to $\alpha_{\lambda, \mu}$ and $\beta_{\lambda, \mu}$.

In other words, the object $\text{Conv}^{\text{Hecke}}(\mathcal{S})$ represents the covariant functor which, to any $\mathcal{F} \in \text{Ind}(\text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}}))$, associates the set of collections of morphisms $(\mathcal{S}_\nu \star \mathcal{IC}_{-\nu}^{\frac{\infty}{2}} \rightarrow \mathcal{F})_{\nu \in \mathbf{Y}}$ such that, for any $\lambda, \mu \in \mathbf{Y}$, the following diagram commutes:

$$(8.1) \quad \mathcal{S}_\lambda \star \mathcal{R}_{-\mu} \star \mathcal{IC}_{\mu-\lambda}^{\frac{\infty}{2}} \rightrightarrows \mathcal{F},$$

where the top (resp. bottom) arrow is the composition of $\alpha_{\lambda, \mu}$ (resp. $\beta_{\lambda, \mu}$) with the morphism $\mathcal{S}_{\lambda-\mu} \star \mathcal{IC}_{\mu-\lambda}^{\frac{\infty}{2}} \rightarrow \mathcal{F}$ (resp. $\mathcal{S}_\lambda \star \mathcal{IC}_{-\lambda}^{\frac{\infty}{2}} \rightarrow \mathcal{F}$).

8.2. First properties. Recall that $\Phi : \text{Perv}_{I_u}(\text{Gr}) \rightarrow \text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ was defined in §3.10.

Proposition 8.1. [ABB⁺05, Proposition 3.1.2] *For any $\mathcal{S} \in \text{Perv}_{I_u}(\text{Gr})$ and $\mu \in \mathbf{Y}$, we have a canonical isomorphism*

$$\text{Conv}^{\text{Hecke}}(\Phi(\mathcal{S})\langle\mu\rangle) \simeq \mathcal{S} \star \mathcal{IC}_{-\mu}^{\frac{\infty}{2}}.$$

Proof. We first construct a morphism of functors

$$(8.2) \quad \text{Hom}_{\text{Ind}(\text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}}))} \left(\text{Conv}^{\text{Hecke}}(\Phi(\mathcal{S})\langle\mu\rangle), - \right) \rightarrow \text{Hom}_{\text{Ind}(\text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}}))} \left(\mathcal{S} \star \mathcal{IC}_{-\mu}^{\frac{\infty}{2}}, - \right).$$

For any morphism $\text{Conv}^{\text{Hecke}}(\Phi(\mathcal{S})\langle\mu\rangle) \rightarrow \mathcal{F}$, we take the component

$$(\Phi(\mathcal{S})\langle\mu\rangle)_\mu \star \mathcal{IC}_{-\mu}^{\frac{\infty}{2}} \rightarrow \mathcal{F}.$$

Composing with the canonical morphism $\mathcal{S} \rightarrow \mathcal{S} \star \mathcal{R}_0 = (\Phi(\mathcal{S})\langle\mu\rangle)_\mu$, we obtain the desired map $\mathcal{S} \star \mathcal{IC}_{-\mu}^{\frac{\infty}{2}} \rightarrow \mathcal{F}$.

Next, we construct an inverse of (8.2). Consider a morphism $\varphi : \mathcal{S} \star \mathcal{IC}_{-\mu}^{\frac{\infty}{2}} \rightarrow \mathcal{F}$. For any $\lambda \in \mathbf{Y}$, we construct a morphism $(\Phi(\mathcal{S})\langle\mu\rangle)_\lambda \star \mathcal{IC}_{-\lambda}^{\frac{\infty}{2}} \rightarrow \mathcal{F}$ in the following way:

$$(\Phi(\mathcal{S})\langle\mu\rangle)_\lambda \star \mathcal{IC}_{-\lambda}^{\frac{\infty}{2}} = \mathcal{S} \star \mathcal{R}_{\lambda-\mu} \star \mathcal{IC}_{-\lambda}^{\frac{\infty}{2}} \xrightarrow{\text{id} \star H_{\lambda-\mu, -\lambda}} \mathcal{S} \star \mathcal{IC}_{-\mu}^{\frac{\infty}{2}} \xrightarrow{\varphi} \mathcal{F}.$$

The fact that this collection of morphisms makes the diagram (8.1) commute follows from the associativity property of the action of \mathcal{R} on $\mathcal{IC}^{\frac{\infty}{2}}$ (Proposition 6.6). The fact that we indeed obtain an inverse of (8.2) is readily checked. \square

Corollary 8.2. *For any $w \in W_{\text{ext}}$, we have an isomorphism*

$$\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{L}}_w) \simeq \text{IC}_w^{\frac{\infty}{2}}.$$

Proof. Let $w \in W_{\text{ext}}$, and write it as $w = yt_\lambda$ for $(y, \lambda) \in W_{\text{ext}}^{\text{res}} \times \mathbf{Y}$. By construction, $\widehat{\mathcal{L}}_w = \Phi(L_y)\langle-\lambda\rangle$, so that Proposition 8.1 yields an isomorphism

$$\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{L}}_w) \simeq L_y \star \mathcal{IC}_\lambda^{\frac{\infty}{2}}.$$

By Theorem 6.9, the right-hand side is isomorphic to $\text{IC}_{yt_\lambda}^{\frac{\infty}{2}}$. \square

Proposition 8.3. [ABB⁺05, Proposition 6.2.2] *The functor $\text{Conv}^{\text{Hecke}}$ is exact.*

Proof. By construction, the functor $\text{Conv}^{\text{Hecke}}$ is right exact. So it remains to check that it is left exact. Since $\text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ is the ind-completion of $\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$, it suffices to check left exactness for the restriction of $\text{Conv}^{\text{Hecke}}$ to $\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$. So let

$$0 \rightarrow \mathcal{S}_1 \xrightarrow{u} \mathcal{S}_2 \rightarrow \mathcal{S} \rightarrow 0$$

be an exact sequence in $\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$. We have to prove that $\text{Conv}^{\text{Hecke}}(\mathcal{S}_1) \rightarrow \text{Conv}^{\text{Hecke}}(\mathcal{S}_2)$ is injective. By an inductive argument on the length of \mathcal{S}_2 , it is sufficient to prove the result in the case where \mathcal{S} is simple. In particular (thanks to Theorem 3.15), we may and will assume that there exist $\mathcal{F} \in \text{Perv}_{I_u}(\text{Gr})$ and $\mu \in \mathbf{Y}$ such that $\mathcal{S} \simeq \Phi(\mathcal{F})\langle\mu\rangle$. Consider the morphism

$$\mathcal{F} \rightarrow \mathcal{F} \star \mathcal{R}_0 \simeq \mathcal{S}_{-\mu}.$$

Since $(\mathcal{S}_2)_{-\mu}$, $\mathcal{S}_{-\mu}$ are ind-objects of the category $\text{Perv}_{I_u}(\text{Gr})$, there exists $\mathcal{F}' \in \text{Perv}_{I_u}(\text{Gr})$ together with a morphism $\varphi : \mathcal{F}' \rightarrow (\mathcal{S}_2)_{-\mu}$ and a surjective morphism $\mathcal{F}' \rightarrow \mathcal{F}$ such that the following diagram commutes

$$\begin{array}{ccc} \mathcal{F}' & \longrightarrow & \mathcal{F} \\ \downarrow & & \downarrow \\ (\mathcal{S}_2)_{-\mu} & \longrightarrow & \mathcal{S}_{-\mu}. \end{array}$$

In particular, we obtain a morphism $a : \Phi(\mathcal{F}')\langle\mu\rangle \rightarrow \mathcal{S}_2$ by Proposition 3.14. We put $\mathcal{F}'' := \ker(\mathcal{F}' \rightarrow \mathcal{F})$. We have a commutative diagram

$$(8.3) \quad \begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ & & \Phi(\mathcal{F}'')\langle\mu\rangle & \xrightarrow{\text{id}} & \Phi(\mathcal{F}'')\langle\mu\rangle & & \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathcal{S}_1 & \longrightarrow & \mathcal{S}_1 \oplus (\Phi(\mathcal{F}')\langle\mu\rangle) & \longrightarrow & \Phi(\mathcal{F}')\langle\mu\rangle \longrightarrow 0 \\ & & \downarrow \text{id} & & \downarrow u \oplus a & & \downarrow \\ 0 & \longrightarrow & \mathcal{S}_1 & \longrightarrow & \mathcal{S}_2 & \longrightarrow & \Phi(\mathcal{F})\langle\mu\rangle \longrightarrow 0 \\ & & & & \downarrow & & \downarrow \\ & & & & 0 & & 0. \end{array}$$

To prove the desired result, we only need to prove that the maps $\text{Conv}^{\text{Hecke}}(\mathcal{S}_1) \rightarrow \text{Conv}^{\text{Hecke}}(\mathcal{S}_1 \oplus (\Phi(\mathcal{F}')\langle\mu\rangle))$ and $\text{Conv}^{\text{Hecke}}(\Phi(\mathcal{F}'')\langle\mu\rangle) \rightarrow \text{Conv}^{\text{Hecke}}(\mathcal{S}_1 \oplus (\Phi(\mathcal{F}')\langle\mu\rangle))$ are injective. For the first one this is obvious, since $\text{Conv}^{\text{Hecke}}$ transforms direct sums into direct sums. For the second one, it suffices to check that

$$\text{Conv}^{\text{Hecke}}(\Phi(\mathcal{F}'')\langle\mu\rangle) \rightarrow \text{Conv}^{\text{Hecke}}(\Phi(\mathcal{F}')\langle\mu\rangle)$$

is injective. But one deduces from Proposition 8.1 that the latter morphism coincides with

$$\mathcal{F}'' \star \mathcal{IC}_{\mu}^{\infty} \rightarrow \mathcal{F}' \star \mathcal{IC}_{\mu}^{\infty},$$

obtained by convolving the injection $\mathcal{F}'' \rightarrow \mathcal{F}'$ with $\mathcal{IC}_{\mu}^{\infty}$. This morphism is injective thanks to exactness of convolution (Theorem 6.4). \square

Notice that, thanks to Proposition 8.3 and Corollary 8.2, we immediately obtain:

Proposition 8.4. *The functor $\text{Conv}^{\text{Hecke}}$ sends finite length objects to finite length objects.*

So restriction yields a functor

$$\text{Conv}^{\text{Hecke}} : \text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R}) \rightarrow \text{Perv}_{I_u}(\mathcal{Fl}_{\frac{\infty}{2}}).$$

Notice that, by replacing $\text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ with $\text{Mod}_{\mathcal{I}\mathcal{W}}^{\mathbf{Y}}(\mathcal{R})$ in the definition of $\text{Conv}^{\text{Hecke}}$, we obtain a functor

$$\text{Conv}_{\mathcal{I}\mathcal{W}}^{\text{Hecke}} : \text{Mod}_{\mathcal{I}\mathcal{W}}^{\mathbf{Y}}(\mathcal{R}) \rightarrow \text{Ind}(\text{Perv}_{(I_u^-, \psi)}(\mathcal{Fl}_{\frac{\infty}{2}}))$$

satisfying properties analogous to the previously stated results concerning $\text{Conv}^{\text{Hecke}}$. In particular, $\text{Conv}_{\mathcal{I}\mathcal{W}}^{\text{Hecke}}$ sends simple objects to simple objects.

Proposition 8.5. *The functor $\text{Conv}^{\text{Hecke}}$ commutes with averaging functors. Namely, for any $\mathcal{S} \in \text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ we have a functorial isomorphism*

$$\text{Av}_{(I_u, \psi)}(\text{Conv}^{\text{Hecke}}(\mathcal{S})) \simeq \text{Conv}_{\mathcal{I}\mathcal{W}}^{\text{Hecke}}(\text{Av}_{(I_u, \psi)}(\mathcal{S})).$$

Proof. First, one can easily check from the constructions of the averaging functors that, for any $\lambda \in \mathbf{Y}$, we have an isomorphism functorial in $\mathcal{S} \in \text{Mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$:

$$\text{Av}_{(I_u, \psi)}(\mathcal{S} \star \mathcal{IC}_{\lambda}^{\frac{\infty}{2}}) \simeq \text{Av}_{(I_u, \psi)}(\mathcal{S}) \star \mathcal{IC}_{\lambda}^{\frac{\infty}{2}}.$$

The conclusion then follows from the fact that $\text{Av}_{(I_u, \psi)}$ is an exact functor, so it preserves coequalizers. \square

Lemma 8.6. *The functor $\text{Conv}^{\text{Hecke}}$ commutes with Verdier duality.*

Proof. Let $\mathcal{S} \in \text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$. Recall the construction of the Verdier duality on $\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ from subsection 3.11, and let $\alpha : \mathcal{S}_1 \rightarrow \mathcal{S}_2$ be a morphism between two free graded \mathcal{R} -modules such that \mathcal{S} is the cokernel of α . By Remark 3.20, $\mathbb{D}(\mathcal{S})$ coincides with the kernel of $\mathbb{D}(\alpha) : \mathbb{D}(\mathcal{S}_2) \rightarrow \mathbb{D}(\mathcal{S}_1)$. Let us spell out how $\mathbb{D}(\alpha)$ was constructed. By additivity of \mathbb{D} , we may and will assume that $\mathcal{S}_1 \simeq \Phi(\mathcal{F}_1)$ and $\mathcal{S}_2 \simeq \Phi(\mathcal{F}_2)\langle\mu\rangle$, for some $\mathcal{F}_1, \mathcal{F}_2 \in \text{Perv}_{I_u}(\text{Gr})$, and $\mu \in \mathbf{Y}$. So (using Proposition 3.14) α comes from a map

$$\tilde{\alpha} : \mathcal{F}_1 \rightarrow \mathcal{F}_2 \star \mathcal{I}_*^{-w_0(\mu)+\nu} \star \mathcal{I}_*^{-w_0(\nu)}$$

in $\text{Perv}_{I_u}(\text{Gr})$, for some large enough $\nu \in \mathbf{Y}^+ \cap (w_0(\mu) + \mathbf{Y}^+)$. Thus (up to taking a larger ν), $\mathbb{D}(\alpha)$ comes from the dual morphism

$$\beta : \mathbb{D}(\mathcal{F}_2) \rightarrow \mathbb{D}(\mathcal{F}_1) \star \mathcal{I}_*^{\mu-w_0(\nu)} \star \mathcal{I}_*^{\nu}.$$

Now, the object $\text{Conv}^{\text{Hecke}}(\mathbb{D}(\mathcal{S}))$ is the kernel of $\text{Conv}^{\text{Hecke}}(\mathbb{D}(\alpha))$ thanks to Proposition 8.3. By Proposition 8.1, the latter morphism is a morphism between the objects $\mathbb{D}(\mathcal{F}_2) \star \mathcal{IC}_{-\mu}^{\frac{\infty}{2}}$ and $\mathbb{D}(\mathcal{F}_1) \star \mathcal{IC}_0^{\frac{\infty}{2}}$. It is constructed as the composition

$$\begin{aligned} \mathbb{D}(\mathcal{F}_2) \star \mathcal{IC}_{-\mu}^{\frac{\infty}{2}} &\xrightarrow{\beta \star \text{id}} \mathbb{D}(\mathcal{F}_1) \star (\mathcal{I}_*^{\mu-w_0(\nu)} \star \mathcal{I}_*^{\nu} \star \mathcal{IC}_{-\mu}^{\frac{\infty}{2}}) \\ &\xrightarrow{\text{id} \star H'} \mathbb{D}(\mathcal{F}_1) \star \mathcal{IC}_0^{\frac{\infty}{2}}, \end{aligned}$$

where H' is the image of $H_{-\mu, 0}^{\nu}$ through the composition

$$\begin{aligned} \text{Hom}((\mathcal{I}_*^{-w_0(\mu)+\nu} \star \mathcal{I}_*^{-w_0(\nu)}) \star \mathcal{IC}_0^{\frac{\infty}{2}}, \mathcal{IC}_{-\mu}^{\frac{\infty}{2}}) &\xrightarrow[\sim]{\text{adjunction}} \text{Hom}(\mathcal{IC}_0^{\frac{\infty}{2}}, (\mathcal{I}_!^{\mu-w_0(\nu)} \star \mathcal{I}_!^{\nu}) \star \mathcal{IC}_{-\mu}^{\frac{\infty}{2}}) \\ &\xrightarrow[\sim]{\mathbb{D}} \text{Hom}((\mathcal{I}_*^{\mu-w_0(\nu)} \star \mathcal{I}_*^{\nu}) \star \mathcal{IC}_{-\mu}^{\frac{\infty}{2}}, \mathcal{IC}_0^{\frac{\infty}{2}}). \end{aligned}$$

On the other hand, $\text{Conv}^{\text{Hecke}}(\mathcal{S})$ is the cokernel of $\text{Conv}^{\text{Hecke}}(\alpha)$ thanks to Proposition 8.3. The latter morphism coincides with the composition

$$\begin{aligned} \mathcal{F}_1 \star \mathcal{IC}_0^{\frac{\infty}{2}} &\xrightarrow{\tilde{\alpha} \star \text{id}} \mathcal{F}_2 \star (\mathcal{I}_*^{-w_0(\mu)+\nu} \star \mathcal{I}_*^{-w_0(\nu)} \star \mathcal{IC}_0^{\frac{\infty}{2}}) \\ &\xrightarrow{\text{id} \star H_{-\mu,0}^\nu} \mathcal{F}_2 \star \mathcal{IC}_{-\mu}^{\frac{\infty}{2}}. \end{aligned}$$

To conclude, we just need to notice that $\text{Conv}^{\text{Hecke}}(\mathbb{D}(\alpha))$ and $\text{Conv}^{\text{Hecke}}(\alpha)$ are transformed into one another by Verdier duality. \square

Conjecture 8.1. *The functor*

$$\text{Conv}^{\text{Hecke}} : \text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R}) \rightarrow \text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}})$$

induces an equivalence of categories between $\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ and the subcategory of finite-length objects in $\text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}})$.

9. STRATEGY OF PROOF FOR CONJECTURE 8.1

We give a strategy of proof for Conjecture 8.1, which follows [ABB⁺05] and will work provided Conjecture 6.1 holds.

9.1. First steps. The strategy is to prove that $\text{Conv}^{\text{Hecke}}$ respects the highest weight structures on both sides.

Lemma 9.1. *Let $\gamma \in \mathbf{Y}^+$, $\nu \in \mathbf{Y}$ and $w, w' \in W$ such that $\ell(w w_0 w') = \ell(w) + \ell(w_0 w')$. We have canonical isomorphisms*

$$\mathcal{N}_\gamma \star^I \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_\nu) \simeq \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_{\gamma+\nu}), \quad \mathcal{N}_w \star^I \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_{w't_\nu}) \simeq \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_{w w' t_\nu}).$$

Proof. Recall that $\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_{\gamma+\nu})$ represents the covariant functor which, to any $\mathcal{F} \in \text{Ind}(\text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}}))$, associates the set of collections of morphisms

$$((\widehat{\mathcal{Z}}'_{\gamma+\nu})_\lambda \star \mathcal{IC}_{-\lambda}^{\frac{\infty}{2}} \rightarrow \mathcal{F})_{\nu \in \mathbf{Y}}$$

such that, for any $\lambda, \mu \in \mathbf{Y}$, the following diagram commutes (cf. (8.1)):

$$(\widehat{\mathcal{Z}}'_{\gamma+\nu})_\lambda \star \mathcal{R}_{-\mu} \star \mathcal{IC}_{\mu-\lambda}^{\frac{\infty}{2}} \rightrightarrows \mathcal{F}.$$

Applying Proposition 3.21 together with the first point of Lemma 3.4, we obtain that $\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_{\gamma+\nu})$ represents the covariant functor which, to any $\mathcal{F} \in \text{Ind}(\text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}}))$, associates the set of collections of morphisms

$$((\widehat{\mathcal{Z}}'_\nu)_\lambda \star \mathcal{IC}_{-\lambda}^{\frac{\infty}{2}} \rightarrow \mathcal{D}_{-\gamma} \star \mathcal{F})_{\nu \in \mathbf{Y}}$$

such that, for any $\lambda, \mu \in \mathbf{Y}$, the following diagram commutes (cf. (8.1)):

$$(\widehat{\mathcal{Z}}'_\nu)_\lambda \star \mathcal{R}_{-\mu} \star \mathcal{IC}_{\mu-\lambda}^{\frac{\infty}{2}} \rightrightarrows \mathcal{D}_{-\gamma} \star \mathcal{F}.$$

By the first point of Lemma 3.4 again, this functor is represented by $\mathcal{N}_\gamma \star^I \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_\nu)$.

The second isomorphism is proved in the same way, using that $\mathcal{N}_w \star^I \widehat{\mathcal{Z}}'_{w't_\nu} \simeq \widehat{\mathcal{Z}}'_{w w' t_\nu}$ (Proposition 3.22). \square

We have the following key result.

Proposition 9.2. *Assume that Conjecture 6.1 holds. There is an isomorphism*

$$(9.1) \quad \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_0) \simeq \nabla_0^{\frac{\infty}{2}}.$$

9.2. Proof of Proposition 9.2. Recall (cf. Proposition 3.23) that, for any $w \in W_{\text{ext}}$, the object $\widehat{\mathcal{Z}}_w$ has a simple head, isomorphic to $\widehat{\mathcal{L}}_w$. We start with a lemma (inspired by [ABB⁺05, Corollary 4.4.14]).

Lemma 9.3. *Let $\nu \in \mathbf{Y}$.*

(1) *The kernel of*

$$f : \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}_{w_0 t_\nu} \rightarrow \widehat{\mathcal{L}}_{w_0 t_\nu})$$

is partially integrable.

(2) *For any $w \in W$, $\text{IC}_{w_0 t_\nu}^{\frac{\infty}{2}}$ is the only non-partially integrable irreducible sub-quotient of $\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}_{w t_\nu})$.*

(3) *The object $\text{IC}_{w_0 t_\nu}^{\frac{\infty}{2}}$ is the head of $\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_\nu)$ and the socle of $\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}_\nu)$. In particular, $\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_\nu)$ and $\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}_\nu)$ are indecomposable.*

Proof. (1) By exactness of the functor $\text{Av}_{(I_u, \psi)}$ (second point of Proposition 7.5), it suffices to show that the image of f through $\text{Av}_{(I_u, \psi)}$ is an isomorphism. But we have

$$\text{Av}_{(I_u, \psi)}(f) \simeq \text{Conv}_{\mathcal{TW}}^{\text{Hecke}}(\text{Av}_{(I_u, \psi)}(\widehat{\mathcal{Z}}_{w_0 t_\nu} \rightarrow \widehat{\mathcal{L}}_{w_0 t_\nu}))$$

thanks to Proposition 8.5, and the morphism $\text{Av}_{(I_u, \psi)}(\widehat{\mathcal{Z}}_{w_0 t_\nu} \rightarrow \widehat{\mathcal{L}}_{w_0 t_\nu})$ is easily seen to be an isomorphism thanks to Proposition 3.25.

(2) Let $w' \in W$ be such that $w = w'w_0$. We have a canonical morphism $\mathcal{D}_0 \rightarrow \mathcal{D}_{w'}$, whose image through $\text{Av}_{(I_u, \psi)}$ is an isomorphism. Convoluting it with $\text{id} : \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}_{w_0 t_\nu}) \rightarrow \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}_{w_0 t_\nu})$ we obtain a morphism

$$\begin{aligned} \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}_{w_0 t_\nu}) \rightarrow \mathcal{D}_{w'} \star \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}_{w_0 t_\nu}) &\stackrel{\text{Lem. 9.1}}{\simeq} \text{Conv}^{\text{Hecke}}(\mathcal{D}_{w'} \star \widehat{\mathcal{Z}}_{w_0 t_\nu}) \\ &\stackrel{\text{Prop. 3.22}}{\simeq} \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}_{w t_\nu}). \end{aligned}$$

The image of the above map through $\text{Av}_{(I_u, \psi)}$ is an isomorphism. So we conclude thanks to the first point and the second point of Proposition 7.5.

(3) Since Verdier duality swaps $\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_\nu)$ and $\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}_\nu)$ (Lemma 8.6) and preserves $\text{IC}_{w_0 t_\nu}^{\frac{\infty}{2}}$, it suffices to show that $\text{IC}_{w_0 t_\nu}^{\frac{\infty}{2}}$ is the head of $\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_\nu)$. In view of the second point, it suffices to show that, if $\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_\nu)$ maps to an irreducible object \mathcal{S} , then \mathcal{S} must be non-partially integrable. Here we repeat the argument of [ABB⁺05, Proposition 2.3.2]. So assume that $\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_\nu)$ maps to a partially integrable irreducible object $\mathcal{S} \in \text{Perv}_{I_u}(\mathcal{Fl}^{\frac{\infty}{2}})$. Since \mathcal{S} is partially integrable and is an IC-sheaf, it must be equivariant with respect to a minimal parabolic subgroup $P_s \subset G$ associated with a simple reflection s . Then $\mathcal{N}_s \star \mathcal{S}$ lives in perverse cohomological degree 1. However, we have an isomorphism

$$\mathcal{N}_s \star \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_\nu) \simeq \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_{st_\nu})$$

by Proposition 3.22, so that the above complex is perverse. Therefore

$$\text{Hom}(\mathcal{N}_s \star \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_\nu), \mathcal{N}_s \star \mathcal{S}) = 0,$$

which is a contradiction since the functor $\mathcal{N}_s \star (-)$ is an equivalence (with inverse $\mathcal{D}_s \star (-)$ by the first point of Lemma 3.4). \square

We now construct a morphism

$$(9.2) \quad \varphi : \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_0) \rightarrow \nabla_0^{\frac{\infty}{2}}.$$

Let $\mu \in \mathbf{Y}$. Recall (§3.12) that by $(\widehat{\mathcal{Z}}'_0)_\mu = \varinjlim_{\lambda \in \mathbf{Y}^+ - \mu} \nabla_{\mu+\lambda} \star \mathcal{I}_*^{-w_0(\lambda)}$. For any $\lambda \in \mathbf{Y}^+ - \mu$ we define a morphism

$$\begin{aligned} \nabla_{\mu+\lambda} \star \mathcal{I}_*^{-w_0(\lambda)} \star \mathcal{IC}_{-\mu}^{\frac{\infty}{2}} &\xrightarrow[\sim]{\text{Prop. 6.5}} \bigoplus_{\nu} \nabla_{\mu+\lambda} \star \mathcal{IC}_{\nu-\mu}^{\frac{\infty}{2}} \otimes \text{Sat}(\mathcal{I}_*^{-w_0(\lambda)})(w_0(\nu)) \\ &\rightarrow \nabla_{\mu+\lambda} \star \mathcal{IC}_{-\mu-\lambda}^{\frac{\infty}{2}} \xrightarrow{(6.10)} \mathcal{N}_{\mu+\lambda} \star \nabla_{-\mu-\lambda}^{\frac{\infty}{2}} \\ &\xrightarrow[\sim]{\text{Prop. 6.7}} \nabla_0^{\frac{\infty}{2}}. \end{aligned}$$

Thanks to the commutativity of the diagrams (6.11) and (6.12), the above morphisms are compatible with the transition morphisms in the limit defining $(\widehat{\mathcal{Z}}'_0)_\mu$. We therefore obtain a collection of morphisms $((\widehat{\mathcal{Z}}'_0)_\mu \star \mathcal{IC}_{-\mu}^{\frac{\infty}{2}} \rightarrow \nabla_0^{\frac{\infty}{2}})_{\mu \in \mathbf{Y}}$. The fact that this collection of morphisms factors through $\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_0)$ follows from the construction of the structures of right \mathcal{R} -module on $\widehat{\mathcal{Z}}'_0$ (§3.12) and left \mathcal{R} -module on $\mathcal{IC}^{\frac{\infty}{2}}$ (§6.6). So we have defined a map $\varphi : \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_0) \rightarrow \nabla_0^{\frac{\infty}{2}}$.

Proposition 9.4. *The morphism $\text{Av}_{(I_u, \psi)}(\varphi)$ is an isomorphism.*

Proof. We have an isomorphism $\text{Av}_{(I_u, \psi)}(\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_0)) \simeq \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'^{\text{TW}})$ thanks to propositions 8.5 and 3.26. But $\widehat{\mathcal{Z}}'^{\text{TW}} \simeq \widehat{\mathcal{L}}^{\text{TW}}$, so that we have an isomorphism

$$\text{Av}_{(I_u, \psi)}(\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_0)) \simeq \text{IC}_0^\psi$$

by Corollary 8.2. On the other hand, we also have an isomorphism

$$\text{Av}_{(I_u, \psi)}(\nabla_0^{\frac{\infty}{2}}) \xrightarrow[\simeq]{\text{Prop. 7.4}} \nabla_0^\psi \xrightarrow[\simeq]{\text{Prop. 7.2}} \text{IC}_0^\psi.$$

Therefore it suffices to check that $\text{Av}_{(I_u, \psi)}(\varphi)$ is nonzero to conclude. For that, it suffices to check that the image of

$$\nabla_{\mu+\lambda} \star \mathcal{IC}_{-\lambda-\mu}^{\frac{\infty}{2}} \rightarrow \mathcal{N}_{\mu+\lambda} \star \nabla_{-\mu-\lambda}^{\frac{\infty}{2}} \simeq \nabla_0^{\frac{\infty}{2}}$$

through $\text{Av}_{(I_u, \psi)}$ is nonzero for every $\mu \in \mathbf{Y}$, $\lambda \in \mathbf{Y}^{++} - \mu$, which brings back to checking that

$$\text{Av}_{(I_u, \psi)}(\mathcal{N}_{\mu+\lambda} \star \mathcal{IC}_{-\lambda-\mu}^{\frac{\infty}{2}} \rightarrow \mathcal{N}_{\mu+\lambda} \star \nabla_{-\mu-\lambda}^{\frac{\infty}{2}})$$

is nonzero. But the latter is isomorphic to

$$\text{Av}_{(I_u, \psi)}(\mathcal{N}_{\mu+\lambda}) \star \mathcal{IC}_{-\lambda-\mu}^{\frac{\infty}{2}} \rightarrow \text{Av}_{(I_u, \psi)}(\mathcal{N}_{\mu+\lambda}) \star \nabla_{-\mu-\lambda}^{\frac{\infty}{2}},$$

where $\text{Av}_{(I_u, \psi)} : \text{Perv}_{I_u}(\text{Fl}) \rightarrow \text{Perv}_{\mathcal{I}\mathcal{W}}(\text{Fl})$ is the averaging functor. The above morphism is obviously nonzero by construction of the convolution product. \square

Next, we construct a morphism

$$\gamma : \nabla_0^{\frac{\infty}{2}} \rightarrow \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_0),$$

assuming that Conjecture 6.1 holds. Equivalently, we construct a morphism

$$\mathbb{D}(\gamma) : \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}_0) \rightarrow \Delta_0^{\frac{\infty}{2}}.$$

By Corollary 3.30, we need to construct (for all $\mu \in \mathbf{Y}$) the last arrow below

$$(9.3) \quad (\widehat{\mathcal{Z}}_0)_\mu \star \mathcal{IC}_{-\mu}^{\frac{\infty}{2}} \simeq ({}^{w_0}(\widehat{\mathcal{Z}}'_{w_0 t_{-2\zeta}}))_\mu \star \mathcal{IC}_{-\mu}^{\frac{\infty}{2}} \rightarrow \Delta_0^{\frac{\infty}{2}}.$$

One can easily deduce from Proposition 3.21 that we have an isomorphism

$$({}^{w_0}(\widehat{\mathcal{Z}}'_{w_0 t_{-2\zeta}}))_\mu \simeq \mathcal{N}_{w_0} \star^I ({}^{w_0}(\widehat{\mathcal{Z}}'_{t_{-2\zeta}}))_\mu.$$

Thus, by the first point of Lemma 3.4, constructing the morphism (9.3) is equivalent to constructing a morphism between $({}^{w_0}(\widehat{\mathcal{Z}}'_{t-2\zeta}))_\mu \star \mathcal{IC}_{-\mu}^{\frac{\infty}{2}}$ and $\mathcal{D}_{w_0} \star^I \Delta_0^{\frac{\infty}{2}} \simeq \Delta_{w_0}^{\frac{\infty}{2}}$. Now, recall that

$$({}^{w_0}(\widehat{\mathcal{Z}}'_{t-2\zeta}))_\mu := \varinjlim_{\lambda} \nabla_{w_0(\mu)+\lambda-2\zeta} \star \mathcal{I}_*^{-w_0(\lambda)}$$

and let us write $\delta := w_0(\mu) + \lambda - 2\zeta$ for concision. We define a morphism

$$\begin{aligned} \nabla_{\delta} \star \mathcal{I}_*^{-w_0(\lambda)} \star \mathcal{IC}_{-\mu}^{\frac{\infty}{2}} &\xrightarrow[\sim]{\text{Prop. 6.5}} \bigoplus_{\nu} \nabla_{\delta} \star \mathcal{IC}_{\nu-\mu}^{\frac{\infty}{2}} \otimes \text{Sat}(\mathcal{I}_*^{-w_0(\lambda)})(w_0(\nu)) \\ &\rightarrow \nabla_{\delta} \star \mathcal{IC}_{-w_0(\lambda)-\mu}^{\frac{\infty}{2}} \xrightarrow{\text{Conj. 6.1}} \mathcal{N}_{\delta} \star \Delta_{w_0 t - w_0(\lambda) - \mu + 2\zeta}^{\frac{\infty}{2}} \\ &\xrightarrow[\sim]{(6.13)} \Delta_{w_0}^{\frac{\infty}{2}}. \end{aligned}$$

Thanks to the commutativity of the diagrams (3.12) and (6.14), the above morphisms are compatible with the transition morphisms in the limit defining $(\widehat{\mathcal{Z}}_0)_\mu$. We therefore obtain a collection of morphisms $((\widehat{\mathcal{Z}}_0)_\mu \star \mathcal{IC}_{-\mu}^{\frac{\infty}{2}} \rightarrow \Delta_0^{\frac{\infty}{2}})_{\mu \in \mathbf{Y}}$. Once again, the fact that this collection of morphisms factors through $\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}_0)$ follows from the construction of the structures of right \mathcal{R} -module on $\widehat{\mathcal{Z}}'_0$ (§3.12) and left \mathcal{R} -module on $\mathcal{IC}^{\frac{\infty}{2}}$ (§6.6).

Proposition 9.5. *Assume that Conjecture 6.1 holds. The morphism φ is an isomorphism.*

Proof. First, notice that the same arguments as in the proof of Proposition 9.4 show that the morphism $\text{Av}_{(I_u, \psi)}(\gamma)$ is an isomorphism.

Thus, the morphism $\text{Av}_{(I_u, \psi)}(\varphi \circ \gamma) \simeq \text{Av}_{(I_u, \psi)}(\gamma) \circ \text{Av}_{(I_u, \psi)}(\varphi)$ is an isomorphism thanks to the previous claim and Proposition 9.4. In particular, the composition

$$\nabla_0^{\frac{\infty}{2}} \xrightarrow{\gamma} \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_0) \xrightarrow{\varphi} \nabla_0^{\frac{\infty}{2}}$$

is nonzero. Therefore $\varphi \circ \gamma$ is the identity up to a nonzero scalar.

On the other hand, the morphism $\text{Av}_{(I_u, \psi)}(\gamma \circ \varphi) \simeq \text{Av}_{(I_u, \psi)}(\varphi) \circ \text{Av}_{(I_u, \psi)}(\gamma)$ is a nonzero endomorphism of

$$\text{Av}_{(I_u, \psi)}(\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_0)) \simeq \text{Conv}_{\mathcal{I}\mathcal{W}}^{\text{Hecke}}(\text{Av}_{(I_u, \psi)}(\widehat{\mathcal{Z}}'_0)) \simeq \text{Conv}_{\mathcal{I}\mathcal{W}}^{\text{Hecke}}(\widehat{\mathcal{L}}_0^{\mathcal{I}\mathcal{W}}) \simeq \text{IC}_0^{\psi},$$

so there exists $\alpha \in \mathbb{k} - \{0\}$ such that $\text{Av}_{(I_u, \psi)}(\alpha \cdot \gamma \circ \varphi) = \text{id}$. Assume that $\gamma \circ \varphi$ is not invertible. Thus $\alpha \cdot \gamma \circ \varphi$ is not invertible. Since $\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_0)$ is indecomposable by the third point of Lemma 9.3, its endomorphism ring is local. Therefore $\text{id} - \alpha \cdot \gamma \circ \varphi$ must be invertible. But

$$\text{Av}_{(I_u, \psi)}(\text{id} - \alpha \cdot \gamma \circ \varphi) = \text{id} - \text{Av}_{(I_u, \psi)}(\alpha \cdot \gamma \circ \varphi) = 0,$$

which is a contradiction. So $\gamma \circ \varphi$ is invertible.

From the two previous paragraphs we deduce that φ has both a left and a right inverse, so that it is invertible. \square

9.3. Proof of Conjecture 8.1 assuming Conjecture 6.1. Until the end, we assume that Conjecture 6.1 holds. We can then finally conclude.

Corollary 9.6. *For all $w \in W_{\text{est}}$, we have isomorphisms*

$$\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_w) \simeq \nabla_w^{\frac{\infty}{2}}, \quad \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}_w) \simeq \Delta_w^{\frac{\infty}{2}}.$$

Proof. By Lemma 8.6 and the fact that Verdier duality sends standard objects to costandard objects, it is enough to construct isomorphisms

$$\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_w) \simeq \nabla_w^{\frac{\infty}{2}}$$

for all $w \in W_{\text{ext}}$. First, if $w = t_\lambda$ for some $\lambda \in \mathbf{Y}^+$, then the isomorphism is obtained from φ by convolution with \mathcal{N}_λ thanks to Lemma 9.1 and Proposition 6.7.

Next, let $w := xt_\lambda \in W_{\text{ext}}$ be an arbitrary element, and pick $\mu \in \mathbf{Y}^+$ dominant enough so that $\lambda + \mu \in \mathbf{Y}^+$. Then we have

$$\mathcal{N}_\mu \star \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_\lambda) \xrightarrow[\text{Lem. 9.1}]{\sim} \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_{\lambda+\mu}) \simeq \nabla_{\lambda+\mu}^{\frac{\infty}{2}},$$

where the last isomorphism follows from the first paragraph. Thanks to the first point of Lemma 3.4, we can apply $\mathcal{D}_{-\mu} \star ()$ to obtain an isomorphism

$$\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_\lambda) \simeq \mathcal{D}_{-\mu} \star \nabla_{\lambda+\mu}^{\frac{\infty}{2}}.$$

But thanks to Proposition 6.7 (and the first point of Lemma 3.4), we know that

$$\mathcal{D}_{-\mu} \star \nabla_{\lambda+\mu}^{\frac{\infty}{2}} \simeq \nabla_\lambda^{\frac{\infty}{2}},$$

so that we have an isomorphism $\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_\lambda) \simeq \nabla_\lambda^{\frac{\infty}{2}}$.

Finally, notice that we have isomorphisms $\mathcal{N}_{w_0} \star \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_{w_0 t_\lambda}) \simeq \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_\lambda)$ (Proposition 3.22) and $\mathcal{N}_{w_0} \star \nabla_{w_0 t_\lambda}^{\frac{\infty}{2}} \simeq \nabla_\lambda^{\frac{\infty}{2}}$ (equation (5.5)). Thanks to the first point of Lemma 3.4, we deduce that there exists an isomorphism $\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_{w_0 t_\lambda}) \simeq \nabla_{w_0 t_\lambda}^{\frac{\infty}{2}}$. Applying the functor $\mathcal{N}_{xw_0} \star ()$ to the latter isomorphism, we obtain the desired isomorphism thanks to Proposition 3.22 and equation (5.5). \square

We can conclude the proof of Theorem 8.1. Combining the results of the two previous subsections, we obtain that the functor $\text{Conv}^{\text{Hecke}}$ is exact and satisfies

$$\text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}_w) \simeq \Delta_w^{\frac{\infty}{2}}, \quad \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{Z}}'_w) \simeq \nabla_w^{\frac{\infty}{2}}, \quad \text{Conv}^{\text{Hecke}}(\widehat{\mathcal{L}}_w) \simeq \text{IC}_w^{\frac{\infty}{2}}$$

for all $w \in W_{\text{ext}}$. Since both categories $\text{mod}_{I_u}^{\mathbf{Y}}(\mathcal{R})$ and $\text{Perv}_{I_u}(\mathcal{F}l^{\frac{\infty}{2}})$ are Artinian and satisfy the Ext vanishing property between standard and costandard objects (cf. Proposition 5.17 and Proposition 3.24), we can apply [ABB⁺05, Lemma 6.2.7] to conclude that $\text{Conv}^{\text{Hecke}}$ is an equivalence of categories.

REFERENCES

- [ABB⁺05] S. Arkhipov, A. Braverman, R. Bezrukavnikov, D. Gaitsgory, and I. Mirković. Modules over the small quantum group and semi-infinite flag manifold. *Transform. Groups*, 10(3-4):279–362, 2005. doi:10.1007/s00031-005-0401-5.
- [Ach21] Pramod N. Achar. *Perverse Sheaves and Applications to Representation Theory*, volume 258. American Mathematical Soc., 2021.
- [AG03] Sergey Arkhipov and Dennis Gaitsgory. Another realization of the category of modules over the small quantum group. *Advances in Mathematics*, 173(1):114–143, 2003. URL: <https://www.sciencedirect.com/science/article/pii/S0001870802000166>, doi:10.1016/S0001-8708(02)00016-6.
- [AR21] Pramod N. Achar and Simon Riche. A geometric Steinberg formula. 2021. arXiv:2109.11980.
- [AR22] Pramod N. Achar and Simon Riche. A geometric model for blocks of Frobenius kernels. 2022. arXiv:2203.03530.
- [BBDG18] Alexander Beilinson, Joseph Bernstein, Pierre Deligne, and Ofer Gabber. *Faisceaux pervers. Astérisque*, 100, 2018.
- [BFGM02] A. Braverman, M. Finkelberg, D. Gaitsgory, and I. Mirković. Intersection cohomology of Drinfeld’s compactifications. *Selecta Math. (N.S.)*, 8(3):381–418, 2002. doi:10.1007/s00029-002-8111-5.
- [BG02] A. Braverman and D. Gaitsgory. Geometric Eisenstein series. *Invent. Math.*, 150(2):287–384, 2002. doi:10.1007/s00222-002-0237-8.
- [BGM⁺19] Roman Bezrukavnikov, Dennis Gaitsgory, Ivan Mirković, Simon Riche, and Laura Rider. An Iwahori-Whittaker model for the Satake category. *J. É. polytech. — Math.*, 6:707–735, 2019. doi:10.5802/jep.104.

- [BL94] Joseph Bernstein and Valery Lunts. *Equivariant sheaves and functors*, volume 1578 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1994. doi:10.1007/BFb0073549.
- [BL95] Arnaud Beauville and Yves Laszlo. Un lemme de descente. *C. R. Acad. Sci. Paris Sér. I Math.*, 320(3):335–340, 1995.
- [BR18] Pierre Baumann and Simon Riche. Notes on the geometric Satake equivalence. In D. Prasad V. Heiermann, editor, *Relative aspects in Representation Theory, Langlands Functoriality and Automorphic Forms*, volume 2221 of *Lecture Notes in Mathematics*, pages 1–134. Springer, 2018. URL: <https://hal.archives-ouvertes.fr/hal-01491529>, doi:10.1007/978-3-319-95231-4_1.
- [BR24] Roman Bezrukavnikov and Simon Riche. On two modular geometric realizations of an affine hecke algebra, 2024. arXiv:2402.08281.
- [BY13] Roman Bezrukavnikov and Zhiwei Yun. On Koszul duality for Kac-Moody groups. *Represent. Theory*, 17:1–98, 2013. doi:10.1090/S1088-4165-2013-00421-1.
- [Cia21] Joshua Ciappara. Hecke category actions via Smith-Treumann theory, 2021. arXiv:2103.07091.
- [Fal03] Gerd Faltings. Algebraic loop groups and moduli spaces of bundles. *J. Eur. Math. Soc. (JEMS)*, 5(1):41–68, 2003. doi:10.1007/s10097-002-0045-x.
- [FFKM99] Boris Feigin, Michael Finkelberg, Alexander Kuznetsov, and Ivan Mirkovic. Semi-infinite flags. II. Local and global intersection cohomology of quasimaps’ spaces. *Differential topology, infinite-dimensional Lie algebras, and applications: DB Fuchs’ 60th Anniversary Collection, 113-148. Amer. Math. Soc. Transl. Ser. 2*, 1999.
- [FGV01] E. Frenkel, D. Gaitsgory, and K. Vilonen. Whittaker patterns in the geometry of moduli spaces of bundles on curves. *Ann. of Math. (2)*, 153(3):699–748, 2001. doi:10.2307/2661366.
- [FM99] Michael Finkelberg and Ivan Mirković. Semi-infinite flags I, the case of a global curve \mathbb{P}^1 . *Differential topology, infinite-dimensional Lie algebras, and applications: DB Fuchs’ 60th Anniversary Collection, 81-112. Amer. Math. Soc. Transl. Ser. 2*, 1999.
- [FS24] Laurent Fargues and Peter Scholze. Geometrization of the local langlands correspondence, 2024. arXiv:2102.13459.
- [Gai18] D. Gaitsgory. The semi-infinite intersection cohomology sheaf. *Advances in Mathematics*, 327:789–868, 2018. Special volume honoring David Kazhdan. doi:10.1016/j.aim.2017.08.007.
- [Jan03] Jens Carsten Jantzen. *Representations of Algebraic Groups*. Math. Surveys Monogr. American Mathematical Society, 2003.
- [KS06] Masaki Kashiwara and Pierre Schapira. *Categories and sheaves*, volume 332 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 2006. doi:10.1007/3-540-27950-4.
- [Kuz96] Alexander Kuznetsov. The Laumon’s resolution of Drinfeld’s compactification is small. *Math. Res. Lett.*, 4(2-3):349–364, 1996.
- [Lau90] G. Laumon. Faisceaux automorphes liés aux séries d’Eisenstein. In *Automorphic forms, Shimura varieties, and L-functions, Vol. I (Ann Arbor, MI, 1988)*, volume 10 of *Perspect. Math.*, pages 227–281. Academic Press, Boston, MA, 1990.
- [LO09] Yves Laszlo and Martin Olsson. Perverse t -structure on Artin stacks. *Math. Z.*, 261(4):737–748, 2009. doi:10.1007/s00209-008-0348-z.
- [Lus] George Lusztig. Intersection cohomology methods in representation theory: plenary address, International Congress of Mathematicians, Kyoto, Japan, August, 1990. *Proceedings of the International Congress of Mathematicians (Kyoto, 1990)*, I,II.
- [Lus81] George Lusztig. Singularities, character formulas, and a q -analogue of weight multiplicities, Analysis and topology on singular spaces, II, III. *Astérisque*, 1981.
- [MV04] Ivan Mirković and Karl Vilonen. Geometric Langlands duality and representations of algebraic groups over commutative rings. *Ann. of Math.*, 166:95–143, 2004.
- [Ric16] Simon Riche. *Geometric Representation Theory in positive characteristic*. Habilitation à diriger des recherches, Université Blaise Pascal (Clermont Ferrand 2), 2016. URL: <https://tel.archives-ouvertes.fr/tel-01431526>.
- [RW21] Simon Riche and Geordie Williamson. A simple character formula. *Annales Henri Lebesgue*, 4:503–535, 2021.
- [RW22] Simon Riche and Geordie Williamson. Smith–Treumann theory and the linkage principle. *Publications mathématiques de l’IHÉS*, 136(1):225–292, 2022.
- [Sta22] The Stacks project authors. The stacks project. <https://stacks.math.columbia.edu>, 2022.
- [Zab22] Emilien Zabeth. Block decomposition via the geometric Satake equivalence, 2022. arXiv:2207.06184.

- [Zhu16] Xinwen Zhu. An introduction to affine grassmannians and the geometric satake equivalence, 2016. [arXiv:1603.05593](#).

UNIVERSITÉ CLERMONT AUVERGNE, CNRS, LMBP, F-63000 CLERMONT-FERRAND, FRANCE
Email address: emilien.zabeth@uca.fr