

A short proof of Tuza's conjecture for weak saturation in hypergraphs

Nikolai Terekhov*

April 8, 2025

Abstract

Given an r -uniform hypergraph H and a positive integer n , the weak saturation number $\text{wsat}(n, H)$ is the minimum number of edges in an r -uniform hypergraph F on n vertices such that the missing edges in F can be added, one at a time, so that each added edge creates a copy of H . Shapira and Tyomkyn (Proceedings of the American Mathematical Society, 2023) proved Tuza's conjecture on asymptotic behaviour of $\text{wsat}(n, H)$. In this paper we provide a significantly shorter proof of the conjecture.

1 Introduction

Given an r -uniform hypergraph H , we say that an r -uniform hypergraph F is weakly H -saturated if it is possible to arrange the edges $\binom{V(F)}{r} \setminus E(F)$ in an order e_1, \dots, e_k such that the addition of each e_i to the hypergraph $F \cup \{e_1, \dots, e_{i-1}\}$ creates a new copy of H . In this case, we write $F \in \text{wSAT}(n, H)$. Let $\text{wsat}(n, H)$ denote the smallest number of edges in such a hypergraph F .

The notion of weak saturation was introduced by Bollobás [1]. For the case of graphs ($r = 2$), Alon [2] described the asymptotic behaviour of $\text{wsat}(n, H)$ by proving the existence of the limit $\lim_{n \rightarrow \infty} \text{wsat}(n, H)/n$.

Tuza [3] proved the following generalisation for hypergraphs. For an r -uniform hypergraph H , define the *sparseness* $s(H)$ as the size of the smallest subset $S \subseteq V(H)$ such that there exists exactly one edge $U \in E(H)$ with $S \subseteq U$; note that $0 \leq s(H) \leq r$ for every non-empty r -uniform hypergraph H . Tuza proved that

$$\text{wsat}(n, H) = \Theta(n^{s(H)-1}), \quad (1)$$

and conjectured that there is, in fact, an exact limit, specifically:

*Department of Discrete Mathematics, Moscow Institute of Physics and Technology, Dolgoprudny, Russia; nikolayterek@gmail.com

Conjecture 1.1. *For any r -uniform hypergraph H with at least two edges, there exists a constant $C_H > 0$ such that*

$$\text{wsat}(n, H) = C_H \cdot n^{s(H)-1}(1 + o(1)).$$

Shapira and Tyomkyn [4] proved this conjecture by utilising the following result by Rödl [5].

Theorem 1.2. *For any $k \geq t \geq 0$ and $\delta > 0$, there exists $N_0(k, t, \delta) \geq k$ such that for any set X of size $|X| \geq N_0(k, t, \delta)$, there exists a family $\mathcal{F}_X \subseteq \binom{X}{k}$ of size $|\mathcal{F}_X| \leq (1 + \delta) \binom{|X|}{t} / \binom{k}{t}$ such that every $A \in \binom{X}{t}$ is contained in some $W \in \mathcal{F}_X$.*

They utilised Theorem 1.2 to combine r -uniform hypergraphs from $\text{wSAT}(m, H)$ for small m into an r -uniform hypergraph from $\text{wSAT}(n, H)$ with the desired estimate on asymptotic. To execute this proof strategy, they introduced a supplementary technical tool, the $T_{r,h,s}$ -template saturation process. In this paper we also derive Conjecture 1.1 from Theorem 1.2 but without the use of the $T_{r,h,s}$ -template saturation process, resulting in a significantly shorter proof of Conjecture 1.1.

2 Proof of Conjecture 1.1

Fix an r -uniform hypergraph H . The existence of at least two edges ensures that $s(H) \geq 1$. Let $v = |V(H)|$ and $s = s(H)$. The statement of Conjecture 1.1 is equivalent to the existence of the limit

$$\lim_{n \rightarrow +\infty} \frac{\text{wsat}(n, H)}{\binom{n-v}{s-1}}.$$

From (1) it follows that

$$\tilde{C}_H = \liminf_{n \rightarrow +\infty} \frac{\text{wsat}(n, H)}{\binom{n-v}{s-1}}$$

exists and $\tilde{C}_H > 0$.

Fix $\varepsilon > 0$. By the definition of \tilde{C}_H , there exists $m \geq v + s - 1$ such that

$$\text{wsat}(m, H) \leq (\tilde{C}_H + \varepsilon) \cdot \binom{m-v}{s-1}.$$

Let F_0 be a hypergraph from $\text{wSAT}(m, H)$ such that $|F_0| = \text{wsat}(m, H)$.

By Theorem 1.2, there exists $N_0 \geq m - v$ such that for any set X with $|X| \geq N_0$, there exists a family $\mathcal{F}_X \subseteq \binom{X}{m-v}$ such that every subset of X of size $s - 1$ is contained in at least one element of \mathcal{F}_X , where

$$|\mathcal{F}_X| \leq (1 + \varepsilon) \cdot \frac{\binom{|X|}{s-1}}{\binom{m-v}{s-1}}.$$

It also follows that any subset of X of size less than $s - 1$ lies inside some element of \mathcal{F}_X , as such a subset can always be extended to the size $s - 1$.

Fix $n \geq N = N_0 + v$. We construct an r -uniform hypergraph F on vertex set $[n]$ in the following way. Let $Z = [v]$ and $X = [n] \setminus Z$. For each element $W \in \mathcal{F}_X$, add to the hypergraph F a copy of F_0 on the vertex set $Z \cup W$. We get that:

$$|E(F)| \leq |\mathcal{F}_X| \cdot |E(F_0)| \leq (1 + \varepsilon) \cdot \frac{\binom{|X|}{s-1}}{\binom{m-v}{s-1}} \cdot (\tilde{C}_H + \varepsilon) \cdot \binom{m-v}{s-1} = (1 + \varepsilon)(\tilde{C}_H + \varepsilon) \cdot \binom{n-v}{s-1}.$$

Given that $\varepsilon > 0$ can be chosen arbitrarily small, to prove the theorem, it is sufficient to show that F is weakly H -saturated.

First, we complete all copies of F_0 in F to cliques. By the definition of \mathcal{F}_X , we obtain a hypergraph \tilde{F} such that it contains all edges $e \in \binom{[n]}{r}$ satisfying $|X \cap e| \leq s - 1$. This is because any such edge belongs to a clique on $W \cup Z$, where $W \in \mathcal{F}_X$ and $X \cap e \subseteq W$. Thus, to complete the proof of Conjecture 1.1, it remains to establish the following claim.

Claim 2.1. *Let H be an r -uniform hypergraph, and let F be an r -uniform hypergraph whose vertices can be partitioned into two sets Z and X , where $|Z| \geq |V(H)|$ and F contains all edges e such that $|X \cap e| \leq s(H) - 1$. Then F is weakly H -saturated.*

Proof. We will show that all missing edges can be saturated by induction on $|e \cap X|$. Let $j \in [s(H) - 1, r - 1]$, and assume that all edges e such that $|e \cap X| \leq j$ are added to F , constituting the hypergraph $F_j \supseteq F$. Let us fix any r -edge e such that $|e \cap X| = j + 1$ and show that its addition to F_j creates a copy of H . This will complete the proof.

By the definition of $s(H)$, there exists a set $S \subseteq V(H)$ of size $s(H)$ that is contained in exactly one edge $U \in E(H)$.

Construct an injection $f : V(H) \rightarrow V(F_j)$ such that $f(V(H)) \subseteq e \cup Z$, $f(U) = e$, and $f(S) \subseteq X$. The later is possible since $|e \cap X| \geq s(H)$.

Now, given that any edge $\tilde{U} \in E(H)$ distinct from U does not fully contain S and $X \cap f(V(H)) \subseteq e = f(U)$, it follows that $|f(\tilde{U}) \cap X| < |f(U) \cap X| = |e \cap X| = j + 1$. Therefore, the only missing edge in $f(E(H))$ is e , and adding it creates a new copy of H , as required. \square

References

- [1] B. BOLLOBÁS, Weakly k -saturated graphs, Beiträge zur Graphentheorie (Kolloquium, Manebach, 1967), Teubner, Leipzig, 1968, pp. 25–31.
- [2] N. ALON, An extremal problem for sets with applications to graph theory, *J. Combin. Theory Ser. A* **40** (1985), 82–89.
- [3] Z. TUZA, Asymptotic growth of sparse saturated structures is locally determined. *Discrete Math.* **108**:1-3 (1992) 397–402. Topological, algebraical and combinatorial structures. Frolík’s memorial volume.
- [4] A. SHAPIRA, M. TYOMKYN, Weakly saturated hypergraphs and a conjecture of Tuza, *Proc. Amer. Math. Soc.* **151** (2023), 2795–2805.
- [5] V. RÖDL, On a packing and covering problem, *Eur. J. Combin.* **6** (1985), 69–78.