

INVOLUTIONS OF CURVES IN ABELIAN SURFACES AND THEIR JACOBIANS

KATRINA HONIGS AND PIJUSH PRATIM SARMAH

ABSTRACT. We examine étale covers of genus two curves that occur in the linear system of a polarizing line bundle of type $(1, d)$ on a complex abelian surface. We give results counting fixed points of involutions on such curves as well as decomposing their Jacobians into isogenous products.

In [ES93], Ekedahl and Serre ask whether it is possible to find a curve of each genus whose Jacobian is isogenous to a product of elliptic curves. Much progress has been made on this question — see for example [PS25] for recent updates — but it remains open. Here, we consider a related question:

Question 1. Given a smooth curve C on a simple complex abelian surface A , its Jacobian J_C is isogenous to a product of abelian varieties $A \times B$. Is B isogenous to a product of elliptic curves?

It is shown in [CvdG92] that if C is a very general curve in the linear system $|L|$ of a very ample line bundle on A , then B is simple. However, certain curves inherit extra automorphisms from the ambient abelian surface, such as translation and the inverse of the group law $[-1]$, which tend to result in additional splitting of their Jacobians.

In this paper, we consider curves C in the linear system of a line bundle L that defines a primitive polarization on A that are furthermore étale covers of smooth genus 2 curves (Construction 1.1).

Applying [KR89, Theorem B] of Kani and Rosen to the automorphism groups of these curves allows us to recover known constructions where Question 1 has a positive answer. These occur when the polarization on A is of type $(1, d)$ for $d = 2$ [Bar87], $d = 3$ [BL95], $d = 4$ [BO19] and $d = 9$, which was analyzed in recently in [BS25]. We prove general results examining the limits of this methodology, which appears not to yield any further examples.

The following two theorems generalize the special cases where $d = 2, 3$ (Theorem 0.1) and $d = 9$ (Theorem 0.2). Theorem 0.2 also gives a new proof of the main result in [CLR21], which demonstrates that there exist curves whose Jacobian is isogenous to an arbitrarily high number of Jacobians of the same dimension.

Theorem 0.1 (Theorem 4.2). *Let C be as in Construction 1.1 for $X = \langle x \rangle \leq K(L)$ a cyclic group. We have the following isogeny relations:*

- (a) *When d is odd: $J_C \sim A \times J_{C/\langle -1 \rangle}^2$.*
- (b) *When d is even: $J_C \sim A \times J_{C/\langle -1 \rangle} \times J_{C/\langle -1 \circ t_x \rangle}$.*

Theorem 0.2 (Theorem 4.3). *Let p be an odd prime and $d = p^2$. Let C be as in Construction 1.1 for*

$$(\mathbb{Z}/p\mathbb{Z})^2 \cong X \leq K(L).$$

Then we have the following isogeny relations, where each C_i is a curve of genus $\frac{p-1}{2}$ that is a quotient of C by $[-1]$ and a translation:

$$J_C \sim A \times \prod_{i=1}^{p+1} J_{C_i}^2, \quad J_{C/\langle -1 \rangle} \sim \prod_{i=1}^{p+1} J_{C_i}.$$

We also give a class of examples where J_C contains an elliptic factor.

In order to obtain our results on Jacobian decompositions, we examine the numbers of fixed points of involutions $[-1] \circ t_x$ on C , which control how low the genus of a quotient curve can be.

Theorem 0.3 (Proposition 2.5). *Let d be odd and C a curve as in Construction 1.1. Any involution $[-1] \circ t_x$ on C has 6 fixed points.*

In [BO19], Borówka and Ortega completed an explicit construction of all possible smooth hyperelliptic curves in (generic) abelian surfaces by producing a curve that is a Klein cover of a genus 2 curve on a $(1, 4)$ -polarized abelian surface. Question 1 has a positive answer for this curve, and we generalize this construction by giving criteria for the existence of an involution $[-1] \circ t_x$ that has 12 fixed points.

Theorem 0.4 (Propositions 3.1, 3.2). *Let d be even and C a curve as in Construction 1.1, which is an X -covering of a smooth genus 2 curve for $X \leq K(L)$. There is an $x \in X$ so that the involution $[-1] \circ t_x$ on C has 12 fixed points if and only if the Sylow 2-subgroup of X is not cyclic.*

The next example where this result applies occurs when $d = 8$, and although the resulting Jacobian has quite a bit of splitting, our methods do not show that B is isogenous to a product of elliptic curves.

Outline. In section 1, we begin by presenting the curves C that we will study. In section 2, we relate fixed points of involutions $[-1] \circ t_x$ on C to translations of the line bundle L , and conclude by proving Theorem 0.3. In section 3, we prove Theorem 0.4. We then take the opportunity to briefly review the smooth hyperelliptic curves in $|L|$; when $d = 2, 4$, they arise from curves made with Construction 1.1. In section 4, we give our results on Jacobian splittings, and discuss the curves giving a positive answer to Question 1.

Notation. The polarization isogeny given by L is denoted by $\varphi_L : A \rightarrow A^\vee$, $a \mapsto t_a^* L \otimes L^{-1}$ and we write its kernel as:

$$K(L) := \ker \varphi_L \cong (\mathbb{Z}/d\mathbb{Z})^2.$$

We write the involution given by the inverse group law map on A as $[-1]$. We denote the set of fixed points of an involution ι on C by $\text{Fix}_C(\iota)$ and their number by $\#\text{Fix}_C(\iota)$.

Acknowledgements. The authors thank Jen Paulhus and Nils Bruin for helpful conversations. K. H. is supported by an NSERC Discovery Grant.

1. CONSTRUCTION

We now construct the curves that we will analyze in this paper. Let A be a simple abelian surface that is $(1, d)$ -polarized by a Néron–Severi class l .

Line bundles that have the same Néron–Severi class differ by translation. However, because we wish to identify the number of ramification points of specific automorphisms of curves in our exposition, we will fix a particular line bundle representative. Considering A as a complex torus V/Λ , we fix a decomposition $V_1 \oplus V_2$ for l and select the line bundle L of characteristic 0 (see [BL04, §3.1]) with respect to this decomposition.

Construction 1.1. Let X be a subgroup of $K(L)$ of order d and consider the quotient $\pi : A \rightarrow A/X$. By Lemma 1.3 below, A/X is principally polarized by the line bundle $\pi_* L =: M$. Furthermore by [BL04, Prop. 6.5.2], we may choose a decomposition of A/X for the Néron–Severi class of M that is compatible with the decomposition of A for l (replacing the decomposition of A if needed) and so that both L and M have characteristic 0 with respect to these decompositions.

The abelian surface A/X is simple since it is isogenous to A . Thus, since it is principally polarized, there is a smooth curve H of genus 2 so that $A/X \simeq J_H$ and $M \simeq \mathcal{O}(H)$.

The curve we construct is the preimage $C := \pi^{-1}(H)$ in $|L|$.

Remark 1.2. For each $x \in X$, the translation t_x is an automorphism of the curve C . Since $[-1]$ is an involution of H , it is an involution of C as well. Thus $\text{Aut}(C)$ contains a subgroup isomorphic to $\langle -1 \rangle \rtimes X$ whose elements are of the form $[-1] \circ t_x$ and t_x for $x \in X$.

Lemma 1.3. *Let X be a subgroup of $K(L)$ of order dividing d and consider the quotient isogeny $A \rightarrow A/X$. Then A/X is $(1, \frac{d}{|X|})$ -polarized by the pushforward of the polarization φ_L by π .*

Proof. Since $\ker(\pi) \subseteq \ker(\varphi_L)$ and is therefore isotropic under the pairing determined by φ_L , the pushforward polarization $\pi_* \varphi_L : A/X \rightarrow (A/X)^\vee$ fills

in the following diagram and corresponds to the line bundle π_*L [Mum70, Corollary, p. 231] (moreover, $L \simeq \pi^*L$):

$$\begin{array}{ccc}
 A & \xrightarrow{\varphi_L} & A^\vee \\
 \downarrow \pi & & \uparrow \pi^\vee \\
 A/X & \xrightarrow{\varphi_{\pi_*L}} & (A/X)^\vee
 \end{array} \tag{1.1}$$

That φ_{π_*L} is a $(1, \frac{d}{|X|})$ -polarization follows, for instance, from an analysis of the kernels in the above diagram. \square

2. SYMMETRIC LINEAR SYSTEMS AND RAMIFICATION OF $[-1]$

In this section, we analyze the number of fixed points of involutions $[-1] \circ t_x$ by relating them to properties of symmetric line bundles. We begin by reviewing properties of such line bundles.

Let A be a simple abelian surface that is $(1, d)$ -polarized by a symmetric line bundle L .

Since L is symmetric, there is an isomorphism $[-1]^*L \simeq L$ and thus $[-1]^*$ is a linear involution on $H^0(A, L)$. Its eigenvalues are 1 and -1 and we denote the associated eigenspaces by $H^0(A, L)^+$ and $H^0(A, L)^-$, respectively. The involution $[-1]$ of A restricts to an involution of each curve whose class is contained in $H^0(A, L)^\pm$.

The dimensions of $H^0(A, L)^\pm$ are affected by the parity of L and whether it has a symmetric theta structure (sts).

The Weil pairing on A gives a non-degenerate strictly alternating form on $A[2]$ as an \mathbb{F}_2 vector space. The quadratic forms associated to this pairing form a principal homogeneous space over $A[2]$ and correspond to the symmetric line bundles with Néron–Severi class $[L]$. These line bundles are called even or odd depending on the parity of the corresponding quadratic form. We examine further specifics on determining this parity in section 2.0.1.

A symmetric theta structure is an isomorphism between the extended theta and extended Heisenberg groups. Such an isomorphism exists if and only if a constraint on the dimension of $h^0(A, L)^+$ or $h^0(A, L)^-$ is satisfied (maximality, see [BL04, Exercise 6.10(10)]).

A line bundle chosen to have characteristic 0 with respect to a decomposition, as in Construction 1.1 has an sts and is even.

The following proposition summarizes the dimensions and base loci of $H^0(A, L)^\pm$. The base loci are contained in $A[2]$ and we call them $A[2]^\mp$, named for the points where the quadratic form associated to L takes the values -1 and $+1$, respectively. For any smooth curve $C \in H^0(A, L)^\pm$, $\text{Fix}_C[-1] = A[2]^\mp$.

Proposition 2.1 ([BL04, Prop. 4.7.5, Exercise 6.10(10)]). *Let A be an abelian surface that is $(1, d)$ -polarized by a symmetric line bundle L .*

If d is odd, then L has an sts and the linear eigensystems of L under the action of $[-1]^$ have the following properties:*

$$\begin{aligned} L \text{ even:} \quad & h^0(A, L)^+ = \frac{d+1}{2}, & h^0(A, L)^- = \frac{d-1}{2}, \\ & \#A[2]^- = 6, & \#A[2]^+ = 10. \\ L \text{ odd:} \quad & h^0(A, L)^+ = \frac{d-1}{2}, & h^0(A, L)^- = \frac{d+1}{2}, \\ & \#A[2]^- = 10, & \#A[2]^+ = 6. \end{aligned}$$

If d is even, we distinguish more cases, depending on whether L has an sts.

$$\begin{aligned} L \text{ sts, even:} \quad & h^0(A, L)^+ = \frac{d}{2} + 1, & h^0(A, L)^- = \frac{d}{2} - 1. \\ & \#A[2]^- = 4, & \#A[2]^+ = 12. \\ L \text{ sts, odd:} \quad & h^0(A, L)^+ = \frac{d}{2} - 1, & h^0(A, L)^- = \frac{d}{2} + 1, \\ & \#A[2]^- = 12, & \#A[2]^+ = 4. \\ L \text{ no sts:} \quad & h^0(A, L)^+ = \frac{d}{2}, & h^0(A, L)^- = \frac{d}{2}, \\ & \#A[2]^- = 8, & \#A[2]^+ = 8. \end{aligned}$$

Given a curve C as in Construction 1.1, for each $x \in X$, $[-1] \circ t_x$ is an involution of C . The following lemma shows that the fixed locus of $[-1] \circ t_x$ is in bijection with the fixed locus of $[-1]$ acting on a translated curve $C + y$, which lies in the linear system of t_{-y}^*L . Subsequent results in this section allow us to determine the properties of t_{-y}^*L so that we may apply Proposition 2.1 to find $\# \text{Fix}_C([-1] \circ t_x)$.

Lemma 2.2. *Let C be a curve contained in an abelian surface A . Suppose there are $x, a \in A$ so that $[-1] \circ t_{x+a}$ is an involution of C . For any $y \in A$ such that $2y = x$, $[-1] \circ t_a$ is an involution of the translated curve $C + y$ and there is a bijection of sets:*

$$\text{Fix}_C[-1 \circ t_{x+a}] \leftrightarrow \text{Fix}_{C+y}[-1 \circ t_a].$$

Proof. For any point $c \in C$, by our assumptions, $-c - x - a \in C$. Any point in $C + y$ may be written as $c + y$ for some $c \in C$. Applying $[-1] \circ t_a$ gives $-c - y - a$, and we note $(-c - y - a) - y = -c - a - x \in C$, and thus $-c - y - a$ is in $C + y$.

Any point $c \in C$ is fixed by $[-1] \circ t_{x+a}$ if $c = -c - x - a$ and $c + y$ is fixed by $[-1] \circ t_a$ if $c + y = -c - y - a$. Since $2y = x$, these conditions are identical. Thus the bijection between these fixed loci is given by translation by y . \square

Lemma 2.3. (a) *For any $y \in A$, t_y^*L is symmetric if and only if $2y \in K(L)$.*

- (b) If L has an sts, then for any $y \in A$, t_y^*L has an sts if and only if $t_y^*L \simeq t_z^*L$ (i.e. $\varphi_L(y) = \varphi_L(z)$) for some $z \in A[2]$.

Proof. (a) Since L is symmetric and $2y = x$, then t_y^*L is symmetric:

$$t_y^*t_y^*L \simeq t_x^*L \simeq L \Rightarrow t_y^*L \simeq t_{-y}^*L \Rightarrow [-1]^*t_y^*L \simeq [-1]^*t_{-y}^*L \simeq t_y^*[-1]^*L \simeq t_y^*L$$

The elements of the quotient group $\langle y \mid 2y \in K(L) \rangle / K(L)$ correspond to distinct line bundles t_y^*L and has order $|A[2]|$. Thus we obtain all $16 = |A[2]|$ symmetric line bundles in this way.

- (b) See the proof of [BL04, Theorem 6.9.5]. Moreover, the symmetric line bundles admitting symmetric theta structures correspond to the cosets $A[2]/(K(L) \cap A[2])$. \square

Lemma 2.4. *Let C be a curve produced using Construction 1.1 with the group $X \leq K(L)$.*

- (a) $[C] \in H^0(A, L)$ has eigenvalue 1 with respect to action of $[-1]^*$.
 (b) Let $a \in A$ such that $2a \in X$ and let M be the principal polarization of A/X produced in Construction 1.1. The eigenvalue of $[C - a] \in H^0(A, t_a^*L)$ is 1 or -1 if $t_{\pi(a)}^*M$ is even or odd, respectively.

Proof. (a) Let C be a curve produced using Construction 1.1 with the subgroup $X \leq K(L)$. Let $\pi : A \rightarrow A/X$ be the quotient isogeny. Then $M := \pi_*L$ principally polarizes A/X and let $H \subset A/X$ be the genus 2 curve so that $M \simeq \mathcal{O}(H)$.

Both L and M both have characteristic 0 with respect to compatible decompositions. In particular, M is even. Since π commutes with $[-1]$, and the following pullback maps $[H]$ to $[C]$, $[C]$ must also have an eigenvalue of 1 with respect to $[-1]^*$:

$$\pi^* : H^0(J_H, M) \rightarrow H^0(A, L) \quad (2.1)$$

- (b) Let $a \in A$ be such that $2a \in X$. Then $t_{\pi(a)}^*M$ is symmetric and since $X \leq K(L)$, by Lemma 2.3(a), t_a^*L is symmetric. The following pullback maps $[H - \pi(a)]$ to $[C - a]$ and is compatible with $[-1]^*$:

$$\pi^* : H^0(A/X, t_{\pi(a)}^*M) \rightarrow H^0(A, t_a^*L). \quad (2.2)$$

Thus, the eigenvalue of $[C - a]$ under the action of $[-1]^*$ is 1 if $t_{\pi(a)}^*M$ is even and -1 if it is odd. \square

Let C be any curve in Construction 1.1. Combining Proposition 2.1 and Lemma 2.2, we may find $\# \text{Fix}_C([-1] \circ t_x)$ by producing $y \in A$ so that $2y = x$ and determining whether t_{-y}^*L has an sts and if so, determining the parity of t_{-y}^*L and the eigenvalue of $[C - y] \in H^0(A, L)$ under the action of $[-1]^*$. We may determine whether t_{-y}^*L has an sts using the criteria in Lemma 2.3(b) and by Lemma 2.4, we may determine the eigenvalue of $[C - y]$ given the

parity of $t_{\pi(a)}^*M$. We use this line of reasoning to distinguish several cases in the following propositions.

Proposition 2.5. *Let d be odd and C be a curve as in Construction 1.1. For any $x \in X$, the action of $[-1] \circ t_x$ on C has 6 fixed points.*

Proof. Since the order of x is odd, we may choose $y \in \langle x \rangle$ of the same order as x so that $2y = x$. Then, $y \in X \leq K(L)$, $t_{-y}^*L \simeq L$ and $\pi(-y) = 0$, so $t_{\pi(y)}^*M \simeq M$. Both L and M are even, so

$$\# \text{Fix}_{C+y}[-1] = \# \text{Fix}_C[-1 \circ t_x] = 6. \quad \square$$

Proposition 2.6. *Let d be even and C be a curve as in Construction 1.1. For any $x \in X$, one of the following mutually exclusive conditions applies to $\text{Fix}([-1] \circ t_x)$.*

Let n be the order $x \in X$ and 2^m the highest power of 2 dividing d .

(a) *Suppose that for all $y \in A$ such that $2y = x$, $y \notin K(L)$. Then, $\# \text{Fix}([-1] \circ t_x) = 8$. This case applies, for instance, if $2^m | n$.*

Now, suppose there exists $y \in A$ such that $2y = x$ and $y \in K(L)$.

- (b) *If $t_{\pi(-y)}^*M$ is even, then $\# \text{Fix}([-1] \circ t_x) = 4$. This case applies, for instance, if $y \in X$ (hence $t_{\pi(y)}^*M = M$), which must occur if n is odd or if n is even and x is contained in a cyclic subgroup of X with order $2n$.*
- (c) *If $t_{\pi(-y)}^*M$ is odd, then $\# \text{Fix}([-1] \circ t_x) = 12$.*

Proof. (a) Let $y \in A$ such that $2y = x$. The condition for part (a) is equivalent to the statement that for any $z \in A[2]$, $z - y \notin K(L)$. By Lemma 2.3(b), t_{-y}^*L does not have a symmetric theta structure, hence $\# \text{Fix}([-1] \circ t_x) = 8$.

This condition applies if $2^m | n$: For any $y \in A$ such that $2y = x$, y has order $2n$ and therefore cannot be contained in $K(L)$.

(b) Since $y \in K(L)$, $t_{-y}^*L \simeq L$ has an sts and is even, and the result follows.

If n is odd, then there is some $y \in \langle x \rangle$ such that $2y = x$, so $y \in X$. If n is even but x is contained in a cyclic subgroup of X of order $2n$ then we may take y to be its generator.

(c) Under the above conditions, $t_{-y}^*L \simeq L$ has an sts and is even, so the result follows. \square

2.0.1. Even and odd. To apply Proposition 2.6, we examine how to determine the parity of $t_{\pi(-y)}^*M$ for $y \in K(L)$ such that $2y \in X$.

In Construction 1.1, we have fixed a decomposition of A/X for the Néron–Severi class of M . The induced decomposition on the 2-torsion subgroup is

a sum of maximal isotropic subgroups of $(A/X)[2]$ with respect to the Weil pairing:

$$(A/X)[2] \cong Z_1 \oplus Z_2.$$

This decomposition allows us to determine the parity of t_z^*M for any $z \in (A/X)[2]$, which is even or odd if the quadratic form $q_M(z)$ is 1 or -1 , respectively. We may choose dual symplectic bases $\{u_1, u_2\}$ and $\{v_1, v_2\}$ of Z_1 and Z_2 as \mathbb{F}_2 -vector spaces so that $\langle u_i, v_j \rangle = (-1)^{\delta_{ij}}$. We may write $z = u + v$ for $u \in Z_1, v \in Z_2$, and then $q_M(z) = \langle u, v \rangle$.

However, if $z = \pi(-y)$ for some $y \in K(L)$ (as in the cases we treat in Proposition 2.6), then it is contained in a smaller subspace of $(A/X)[2]$. The decomposition of A for l induces a decomposition of $K(L)$ as follows:

$$K(L) = K_1 \oplus K_2 = \langle k_1 \rangle \oplus \langle k_2 \rangle.$$

The image of $K(L)$ in A/X is $\pi(K(L)) = \langle \pi(k_1) \rangle \oplus \langle \pi(k_2) \rangle$, which, by the compatibility assumptions of Construction 1.1, respects the decomposition of A/X with respect to M . Since $\pi(K(L))$ is generated by two cyclic groups, its intersection with $(A/X)[2]$ (if nontrivial) is contained in two cyclic groups:

$$\pi(K(L)) \cap (A/X)[2] \subseteq \langle w_1 \rangle \oplus \langle w_2 \rangle$$

for some w_i such that $\mathbb{Z}/2\mathbb{Z} \cong \langle w_i \rangle \leq Z_i$ and $\langle w_1, w_2 \rangle = -1$. Then we have the following result:

If $\pi(-y)$ is 0, w_1 , or w_2 , then $t_{\pi(-y)}^*M$ is even.

If $\pi(-y)$ is $w_1 + w_2$, then $t_{\pi(-y)}^*M$ is odd.

3. COUNTING FIXED POINTS OF $[-1] \circ t_x$

In this section, let C be as in Construction 1.1 and d be even. We give further results on $\# \text{Fix}_C([-1] \circ t_x)$ for $x \in X$, particularly investigating the unusual case where the number of fixed points is 12, and then apply these results to review the number of smooth hyperelliptic curves in L .

3.1. When can $\# \text{Fix}_C([-1] \circ t_x) = 12$? In the first result in this section, we see a situation where such an involution cannot occur, and in our second result, we characterize such involutions. Let 2^m be the highest power of 2 dividing d . Let X_2 be the Sylow 2-subgroup of X . We refer to the projection of any $x \in X$ to X_2 as the 2-primary part of x .

Proposition 3.1. *If X_2 is cyclic, then, for any $x \in X$:*

- (a) $\# \text{Fix}_C([-1] \circ t_x) = 8$ if the 2-primary part of x generates X_2 .
- (b) $\# \text{Fix}_C([-1] \circ t_x) = 4$ if the 2-primary part of x does not generate X_2 .

Proof. The order of X_2 is 2^m , which is also the highest order of any elements in the 2-primary part of $K(L)$.

(a) If the 2-primary part of x generates X_2 , we may choose y such that $2y = x$ where $y \notin K(L)$ (we may choose the non 2-primary part of y to be in X). Thus Proposition 2.6(a) applies.

(b) Choose $y \in X$ so that $2y = x$ and apply Proposition 2.6(b). \square

Proposition 3.2. *If X_2 is not cyclic, then there is an $x \in X$ so that $\# \text{Fix}_C([-1] \circ t_x) = 12$.*

Moreover, for any $x \in X$, $\# \text{Fix}_C([-1] \circ t_x)$ is either 4 or 12.

Proof. We may write the decomposition of the 2-primary part of $K(L)$ for l as follows:

$$K(L)_2 := K(L) \cap A[2^m] \cong K_1 \oplus K_2 = \langle k_1 \rangle \oplus \langle k_2 \rangle.$$

Since we have chosen compatible decompositions in Construction 1.1, we may also write X_2 as a sum with respect to this decomposition for some $0 < m_1 \leq m_2$ and $m_1 + m_2 = m$:

$$X_2 \simeq \mathbb{Z}/2^{m_1}\mathbb{Z} \oplus \mathbb{Z}/2^{m_2}\mathbb{Z}.$$

Since $m_1, m_2 < m$, for any $x \in X$, there exists $y \in K(L)$ so that $2y = x$, and thus Proposition 2.6(a) cannot apply and $\# \text{Fix}_C([-1] \circ t_x)$ cannot be 8, leaving 4 and 12 as the possible numbers of fixed points.

Again, by examining orders of elements, there is a $y \in K(L) \setminus X$ so that $2y = x \in X$ and $\pi(y)$ is not contained in either group $\langle \pi(k_i) \rangle$. As shown in section 2.0.1, $t_{\pi(-y)}^*$ must be odd. \square

3.2. Smooth hyperelliptic curves in $(1, d)$ -polarizations. In this section, we review the number of smooth hyperelliptic curves in a fixed linear system of type $(1, d)$. These values are nonzero for $1 \leq d \leq 4$ and in these cases are 1, 6, 9 and 4, respectively (see [BOPY18, Table 1]).

These curves are symmetric and have hyperelliptic involution $[-1]$ or $[-1] \circ t_x$ for some $x \in K(L)$ or are translations of such a curve: by Lemma 2.2, for any curve $C \subset A$ with involution $[-1] \circ t_x$ and $b \in K(L)$, $\# \text{Fix}_C([-1] \circ t_x) = \# \text{Fix}_{C+b}([-1] \circ t_{x-2b})$.

Smooth curves in $|L|$ have genus $d + 1$ and thus a hyperelliptic involution on such a curve must have $2d + 4$ fixed points. For $d = 2$ or 4 , we produce these curves using Construction 1.1. In those cases, we may then produce $|K(L)/X|$ distinct translations of each curve.

3.2.1. d odd. In the principally polarized case ($d = 1$), we may read directly from Proposition 2.1 that any symmetric line bundle on a simple abelian surface has its linear system given by a single genus 2 curve with hyperelliptic involution $[-1]$, making the total number of hyperelliptic curves $|K(L)| = 1$.

When $d = 3$, a hyperelliptic involution must have 10 fixed points. In Proposition 2.1, we see that the systems $H^0(A, L)^\pm$ contain a single smooth

curve where $[-1]$ has 10 fixed points. By Proposition 2.5, this curve cannot occur as a cover of a genus 2 curve using Construction 1.1. Borówka and Sankaran [BS17] exhibit it in terms of theta functions and, show that, up to translation, it is the unique hyperelliptic curve in a general $(1, 3)$ -polarized abelian surface.

Given this curve C , for each $b \in K(L)$, the translation $C + b$ is a distinct hyperelliptic curve in $|L|$. The total number of hyperelliptic curves in $|L|$ produced in this way is:

$$|K(L)| = 9.$$

3.2.2. $(1, 2)$ -polarizations. In this case, a hyperelliptic involution must have 8 fixed points. Any curve C as in Construction 1.1 is formed using an order two cyclic group $\langle x \rangle = X \leq K(L)$. By Proposition 3.1, $\# \text{Fix}_C[-1] = 4$ and $\# \text{Fix}_C([-1] \circ t_x) = 8$, making $[-1] \circ t_x$ a hyperelliptic involution.

A detailed treatment of the $(1, 2)$ -polarized case where L has a symmetric theta structure is given by Barth in [Bar87]. Our argument here recovers his result that, for instance, a hyperelliptic involution of the curves in $|L|$ must be distinct from $[-1]$.

The total number of hyperelliptic curves in $|L|$ that we can produce from this construction, as well as translating, is thus:

$$\sum_{\mathbb{Z}/2\mathbb{Z} \simeq X \leq K(L)} |K(L)/X| = 3 \cdot 2 = 6.$$

3.2.3. $(1, 4)$ -polarizations. In this case a hyperelliptic involution must have 12 fixed points. Curves can be obtained using Construction 1.1 using a choice of order 4 subgroup $X \leq K(L) \simeq (\mathbb{Z}/4\mathbb{Z})^2$.

By Proposition 3.1, if $X \leq K(L)$ is cyclic, then $\# \text{Fix}([-1] \circ t_x)$ is 4 or 8, so Construction 1.1 will not produce a hyperelliptic involution.

Now, let $X \leq K(L)$ be the Klein group. We may write $K(L)$ as a sum with respect to the decomposition of A for $[L]$:

$$\langle k_1 \rangle \oplus \langle k_2 \rangle \cong K_1 \oplus K_2.$$

Using this notation, $X = \langle 2k_1 \rangle \oplus \langle 2k_2 \rangle$. By Proposition 3.2,

$$\begin{aligned} \# \text{Fix}_C[-1] &= \# \text{Fix}_C([-1] \circ t_{2k_1}) = \# \text{Fix}_C([-1] \circ t_{2k_2}) = 4, \\ \# \text{Fix}_C([-1] \circ t_{2k_1+2k_2}) &= 12, \end{aligned}$$

making $[-1] \circ t_{2k_1+2k_2}$ a hyperelliptic involution for C .

Since there is only one way to embed the Klein group V_4 into $K(L)$, we have found one hyperelliptic curve using Construction 1.1. The total number of hyperelliptic curves in $|L|$ that we may then produce by translating is:

$$|K(L)/V_4| = 4.$$

Remark 3.3. This construction coincides with that of Borówka and Ortega [BO19], and so we may think of Proposition 2.6 as a generalization of their results. However, in Construction 1.1 we have specified additional data by choosing decompositions.

The authors prove their curve is the unique hyperelliptic curve (up to translation) on a general $(1, 4)$ -polarized abelian surface. It is constructed as a cover of a genus 2 curve, $\tilde{C} \rightarrow H$ with respect to the Klein group $\{0, \eta_1, \eta_2, \eta_1 + \eta_2\} = G \leq J_H[2]$. The result [BO19, Theorem 4.7] states that if \tilde{C} is hyperelliptic then G is non-isotropic with respect to the Weil pairing on $J_H[2]$ and that, conversely, if G is non-isotropic and η_1 and η_2 may be written as the difference of two Weierstrass points, then \tilde{C} is hyperelliptic.

In the notation of this paper $\pi(K(L))$ plays the role of the group G generated by $\pi(k_1), \pi(k_2)$. It's precisely this non-isotropy condition that allows us to pick an element of $z \in \pi(K(L))$ where t_z^*M is odd. The Weierstrass points of $z \in H$ are those where $q_M(z) = -1$, but since we have arranged that $q_M(\pi(k_1)) = q_M(\pi(k_2)) = 1$, we can see that $\pi(k_1)$ and $\pi(k_2)$ must be differences of Weierstrass points.

4. DECOMPOSING JACOBIANS USING SUBGROUPS

By Poincaré reducibility, every abelian variety is isogenous to a product of simple abelian varieties that are unique, up to isogeny. In this section, we decompose the Jacobians of the curves C produced using Construction 1.1 into, up to isogeny, products of smaller abelian varieties.

In section 4.1, we give some general consequences of [KR89, Theorem B] for decomposing J_C . In section 4.2, we use our methods to examine decompositions of curves satisfying the conditions of Proposition 3.2 for $d = 4$ and 8. Finally in section 4.3 we give some examples of curves that cover an elliptic curve.

4.1. General results on decomposing Jacobians. Kani and Rosen proved the following theorem on decomposing Jacobians J_C by showing that partitions of the automorphism group of C induce idempotent relations in the endomorphism algebra of J_C . The automorphisms of the curves C produced using Construction 1.1 are described in Remark 1.2. The structure of this group affects the results that may be gained from applying this theorem. By the result [Sch94, Theorem 3.5.10] characterizing groups that admit partitions, some subgroups of the automorphism group of C do not admit a partition. The only subgroups of $\text{Aut}(C)$ that admit partitions are dihedral groups or certain p -groups.

Theorem 4.1 ([KR89, Theorem B]). *Let C be a (smooth, projective, geometrically connected) curve and G be a finite subgroup of $\text{Aut}(C)$ such that*

$G = H_1 \cup \cdots \cup H_t$ where the subgroups $H_i \leq G$ satisfy $H_i \cap H_j = 1$ if $i \neq j$. Then we have the isogeny relation

$$J_C^{t-1} \times J_{C/G}^g \sim J_{C/H_1}^{h_1} \times \cdots \times J_{C/H_t}^{h_t} \quad (4.1)$$

where $|G| = g$ and $|H_i| = h_i$.

We deduce two general results on the decomposition of Jacobians of curves C produced as in Construction 1.1 in the cases where $X \leq K(L)$ is cyclic or it is a product of two cyclic groups of odd prime order.

Theorem 4.2. *Let C be as in Construction 1.1 for $X = \langle x \rangle \leq K(L)$ a cyclic group (of order d). We have the following isogeny relations:*

- (a) *When d is odd: $J_C \sim A \times J_{C/\langle -1 \rangle}^2$.*
- (b) *When d is even: $J_C \sim A \times J_{C/\langle -1 \rangle} \times J_{C/\langle -1 \circ t_x \rangle}$.*

When $d = 2$ or 3 , Theorem 4.2 implies the answer to Question 1 is yes for any curve C as in Construction 1.1. When $d = 2$, as shown in section 3.2.2, $J_{C/\langle -1 \rangle}$ is elliptic and $J_{C/\langle -1 \circ t_x \rangle}$ is trivial. When $d = 3$, by Proposition 2.5, $J_{C/\langle -1 \rangle}$ is an elliptic curve (cf. [BL95, §5]).

Proof of Theorem 4.2. $\text{Aut}(C)$ contains the dihedral group $G := \langle -1, t_x \rangle$ of order $2d$, which admits the following partition:

$$G = \langle t_x \rangle \cup \langle -1 \rangle \cup \langle -1 \circ t_x \rangle \cup \cdots \cup \langle -1 \circ t_{(d-1)x} \rangle.$$

Applying Theorem 4.1 gives the following isogeny relation:

$$J_C^d \times J_{C/G}^{2d} \sim J_{C/\langle t_x \rangle}^d \times J_{C/\langle -1 \rangle}^2 \times J_{C/\langle -1 \circ t_x \rangle}^2 \times \cdots \times J_{C/\langle -1 \circ t_{(d-1)x} \rangle}^2.$$

We now analyze these quotients further. The quotient C/G is isomorphic to the quotient of the genus two curve $C/X \cong H$ of Construction 1.1 by its hyperelliptic involution, and thus $C/G \cong \mathbb{P}^1$, making $J_{C/G}$ trivial.

Furthermore, quotienting a curve by conjugate subgroups yields isomorphic quotient curves. When d is odd, the order two subgroups of G are precisely the 2-Sylow subgroups of G and hence are conjugate, proving (a).

When d is even, the order two subgroups generated by $-1, -1 \circ t_{2x}, \dots, -1 \circ t_{(d-2)x}$ are conjugate to one another, and the remaining order two subgroups are all conjugate to $\langle -1 \circ t_x \rangle$, proving result (b). \square

For $d = 9$, the following result shows how to construct a curve C with a positive answer to Question 1.

Theorem 4.3. *Let p be an odd prime and $d = p^2$. Let C be as in Construction 1.1 for:*

$$(\mathbb{Z}/p\mathbb{Z})^2 \cong X = \langle x_1 \rangle \times \langle x_2 \rangle \leq K(L).$$

Then we have the following isogeny relations, where each C_i is a curve of genus $\frac{p-1}{2}$ that is a quotient of C by $[-1]$ and a translation:

$$J_C \sim A \times \prod_{i=1}^{p+1} J_{C_i}^2, \quad J_{C/\langle -1 \rangle} \sim \prod_{i=1}^{p+1} J_{C_i}.$$

Proof. The subgroup $G := \langle t_{x_1}, t_{x_2} \rangle$ of $\text{Aut}(C)$ has the following partition:

$$\langle t_{x_1} \rangle \cup \langle t_{x_2} \rangle \cup \langle t_{x_1+x_2} \rangle \cup \langle t_{x_1+2x_2} \rangle \cup \cdots \cup \langle t_{x_1+(p-1)x_2} \rangle.$$

Then Theorem 4.1 gives the following isogeny relation:

$$J_C^p \times J_{C/G}^{p^2} \sim J_{C/\langle t_{x_1} \rangle} \times \cdots \times J_{C/\langle t_{x_1+(p-1)x_2} \rangle}. \quad (4.2)$$

Each curve on the right-hand side in the above isogeny relation is a quotient of C by a cyclic group of order p , which is in turn a cyclic cover of degree p of a genus 2 curve.

Since the quotient $C/\langle t_x \rangle$ is in the linear system of a $(1, p)$ -polarization on $A/\langle x_1 \rangle$, we may apply the argument used in Theorem 4.2(a):

$$J_{C/\langle t_{x_1} \rangle} \sim A/\langle x_1 \rangle \times J_{C_1}^2 \sim A \times J_{C_1}^2$$

where $C_1 = C/\langle t_{px_1}, -1 \rangle$ is of genus $\frac{p-1}{2}$. Performing an analogous decomposition for each cyclic quotient of C and observing that $J_{C/G} \sim A$, we simplify (4.2) to

$$J_C^p \sim A^p \times \prod_{i=1}^{p+1} J_{C_i}^{2p} \Rightarrow J_C \sim A \times \prod_{i=1}^{p+1} J_{C_i}^2. \quad (4.3)$$

Now, let $G := \langle t_{x_1}, t_{x_2}, [-1] \rangle \leq \text{Aut}(C)$, which has the following partition:

$$G = \left(\bigcup_{0 \leq m, n \leq p-1} \langle [-1] \circ t_{mx_1+nx_2} \rangle \right) \cup \langle t_{x_1}, t_{x_2} \rangle.$$

Applying Theorem 4.1 gives the isogeny relation:

$$J_C^{p^2} \times J_{C/G}^{p^2} \sim J_{C/\langle t_{x_1}, t_{x_2} \rangle}^{p^2} \times \prod_{0 \leq m, n \leq p-1} J_{C/\langle [-1] \circ t_{mx_1+nx_2} \rangle}^2. \quad (4.4)$$

Now, in the notation of Construction 1.1, $J_{C/\langle t_{x_1}, t_{x_2} \rangle} \cong J_H \sim A$, and hence $C/G \cong \mathbb{P}^1$. All order 2 subgroups of G are Sylow, hence conjugate to $\langle -1 \rangle$. Simplifying (4.4) and combining with (4.3) yields:

$$J_{C/\langle -1 \rangle} \sim \prod_{i=1}^{p+1} J_{C_i}. \quad \square$$

4.2. Curves satisfying Proposition 3.2. We begin by examining the Klein cover of [BO19] and recover the isogenous decomposition of its Jacobian using Theorem 4.1. Given that this curve has a positive answer to Question 1, one might ask if we may produce more such curves by generalizing this construction. It appears not: We examine the next case of a curve satisfying Proposition 3.2, where $d = 8$, and our methods do not yield curves where the answer to Question 1 is positive.

4.2.1. $(1, 4)$ -polarized surfaces. Let C be as in Construction 1.1 where we quotient by the Klein group $X = \langle x_1 \rangle \oplus \langle x_2 \rangle \cong (\mathbb{Z}/2\mathbb{Z})^2$. In this notation, $[-1] \circ t_{x_1+x_2}$ is the hyperelliptic involution (see section 3.2.3). We begin analogously to Theorem 4.3. The group $G := \langle t_{x_1}, t_{x_2} \rangle \leq \text{Aut}(C)$ admits a partition:

$$G = \langle t_{x_1} \rangle \cup \langle t_{x_2} \rangle \cup \langle t_{x_1+x_2} \rangle.$$

The isogeny relation given by Theorem 4.1 simplifies to:

$$J_C \times A^2 \sim J_{C/\langle t_{x_1} \rangle} \times J_{C/\langle t_{x_2} \rangle} \times J_{C/\langle t_{x_1+x_2} \rangle}. \quad (4.5)$$

There is a cover $C/\langle t_{x_1} \rangle \rightarrow H$ given by taking the quotient by t_{x_2} . Thus the automorphism group of $C/\langle t_{x_1} \rangle$ contains the order 4 dihedral group $\langle -1, t_{x_2} \rangle$. Applying Theorem 4.1, we have:

$$J_{C/\langle t_{x_1} \rangle} \sim A/\langle x_1 \rangle \times J_{C/\langle t_{x_1}, -1 \rangle} \times J_{C/\langle t_{x_1}, -1 \circ t_{x_2} \rangle}.$$

Since $\langle t_{x_1}, -1 \circ t_{x_2} \rangle$ contains the hyperelliptic involution, $J_{C/\langle t_{x_1}, -1 \circ t_{x_2} \rangle}$ is trivial, meaning that $J_{C/\langle t_{x_1}, -1 \rangle}$ is elliptic. Arguing similarly for $C/\langle t_{x_2} \rangle$ and $C/\langle t_{x_1+x_2} \rangle$, the isogeny relation (4.5) reduces to the following relation between J_C and the product of A with three elliptic curves (see [BO19, Theorem 5.5]), which are each the Jacobian of a quotient of C by a subgroup of $\text{Aut}(C)$ that does not contain its hyperelliptic involution:

$$J_C \sim A \times J_{C/\langle t_{x_1}, -1 \rangle} \times J_{C/\langle t_{x_2}, -1 \rangle} \times J_{C/\langle t_{x_1+x_2}, -1 \circ t_{x_1} \rangle}.$$

4.2.2. $(1, 8)$ -polarized surfaces. To produce a curve satisfying Proposition 3.2, we may write, without loss of generality, $X = \langle x_1 \rangle \oplus \langle x_2 \rangle \simeq \mathbb{Z}/4\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$, decomposed with respect to the pairing on A .

We may proceed similarly to the $d = 4$ case by first considering the partition of $\langle 2x_1, x_2 \rangle$. The decomposition of Jacobian J_C will contain the factor $J_{C/\langle t_{x_2} \rangle}$, where $C/\langle x_2 \rangle$ is a curve in a $(1, 4)$ -polarization whose automorphisms contain the dihedral group $\langle t_{x_1}, -1 \rangle$. The proof of Theorem 4.2 shows that we may decompose $J_{C/\langle t_{x_2} \rangle}$ into the product $A \times J_{C/\langle t_{x_2}, -1 \rangle} \times J_{C/\langle t_{x_2}, -1 \circ t_{x_1} \rangle}$ of two abelian surfaces and an elliptic curve. Our methods do not yield a further decomposition.

When $d = 8$, the only other choice for X is $\mathbb{Z}/8\mathbb{Z}$, and our methods again do not yield a decomposition beyond what shown in Theorem 4.2 and thus do not produce a curve where this a positive answer for Question 1.

4.3. Curves that cover an elliptic curve. We can use our methods to examine the following weaker condition: does the curve C of Construction 1.1 cover an elliptic curve? The following result gives some cases where this question has a positive answer.

Proposition 4.4. *When $d = 2k$ (resp. $d = 3k$) for some $k \in \mathbb{N}$ and $X \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/k\mathbb{Z}$ (resp. $\mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}/k\mathbb{Z}$), the curve C of Construction 1.1 covers an elliptic curve. Further the degree of the cover is d (resp. $\frac{2d}{3}$) and J_C has an elliptic factor in its isogeny class.*

Proof. Let $d = 2k$ and let $x_1, x_2 \in K(L)$ be elements of orders k and 2 so that $X = \langle x_1, x_2 \rangle \leq K(L)$ is a subgroup of order $2k$.

The curve $\tilde{C} := C/\langle t_{x_1} \rangle$ has genus 3 and quotienting by t_{x_2} gives a cover $\tilde{C} \rightarrow H$, and thus the automorphism group of \tilde{C} contains $\langle t_{x_2}, -1 \rangle \cong D_4$. Applying Theorem 4.1, we have:

$$J_{\tilde{C}} \sim A/\langle x_1 \rangle \times J_{\tilde{C}/\langle -1 \rangle} \times J_{\tilde{C}/\langle -1 \circ t_{x_2} \rangle}.$$

Comparing dimensions, we see that one of $\tilde{C}/\langle -1 \rangle$ or $\tilde{C}/\langle -1 \circ t_{x_2} \rangle$ must be an elliptic curve E and the covering map $C \rightarrow E$ has degree $2k$.

Since $J_C \sim J_{\tilde{C}} \times P(C/\tilde{C})$, where $P(C/\tilde{C})$ is the Prym variety, E appears as an isogenous factor in the decomposition of J_C .

For $d = 3k$, we make the same argument with x an element of order 3. Then $\tilde{C} := C/\langle t_{x_1} \rangle$ is a curve of genus 4 whose automorphism group contains $\langle -1, t_{x_2} \rangle \cong D_6$ and $J_{\tilde{C}} \sim A/\langle x_1 \rangle \times J_{\tilde{C}/\langle -1 \rangle}^2$, thus $\tilde{C}/\langle -1 \rangle =: E$ is elliptic, $C \rightarrow E$ has degree $3k$, and E is an isogenous factor of J_C . \square

REFERENCES

- [Bar87] Wolf Barth. Abelian surfaces with $(1, 2)$ -polarization. In *Algebraic geometry, Sendai, 1985*, volume 10 of *Adv. Stud. Pure Math.*, pages 41–84. North-Holland, Amsterdam, 1987.
- [BL95] Ch. Birkenhake and H. Lange. Moduli spaces of abelian surfaces with isogeny. In *Geometry and analysis (Bombay, 1992)*, pages 225–243. Tata Inst. Fund. Res., Bombay, 1995.
- [BL04] Ch. Birkenhake and H. Lange. *Complex abelian varieties*, volume 302 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 2004.
- [BO19] Paweł Borówka and Angela Ortega. Hyperelliptic curves on $(1, 4)$ -polarised abelian surfaces. *Math. Z.*, 292(1-2):193–209, 2019.
- [BOPY18] Jim Bryan, Georg Oberdieck, Rahul Pandharipande, and Qizheng Yin. Curve counting on abelian surfaces and threefolds. *Algebr. Geom.*, 5(4):398–463, 2018.
- [BS17] P. Borówka and G. K. Sankaran. Hyperelliptic genus 4 curves on abelian surfaces. *Proc. Amer. Math. Soc.*, 145(12):5023–5034, 2017.
- [BS25] Paweł Borówka and Anatoli Shatsila. Pryms of $\mathbb{Z}_3 \times \mathbb{Z}_3$ coverings of genus 2 curves, 2025.
- [CLR21] Angel Carocca, Herbert Lange, and Rubí E. Rodríguez. Decomposable Jacobians. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 22(4):1673–1690, 2021.

- [CvdG92] Ciro Ciliberto and Gerard van der Geer. On the Jacobian of a hyperplane section of a surface. In *Classification of irregular varieties (Trento, 1990)*, volume 1515 of *Lecture Notes in Math.*, pages 33–40. Springer, Berlin, 1992.
- [ES93] Torsten Ekedahl and Jean-Pierre Serre. Exemples de courbes algébriques à jacobienne complètement décomposable. *C. R. Acad. Sci. Paris Sér. I Math.*, 317(5):509–513, 1993.
- [KR89] E. Kani and M. Rosen. Idempotent relations and factors of Jacobians. *Math. Ann.*, 284(2):307–327, 1989.
- [Mum70] D. Mumford. *Abelian varieties*, volume 5 of *Tata Institute of Fundamental Research Studies in Mathematics*. Published for the Tata Institute of Fundamental Research, Bombay by Oxford University Press, London, 1970.
- [PS25] Jennifer Paulhus and Andrew V. Sutherland. Completely decomposable modular Jacobians, 2025.
- [Sch94] Roland Schmidt. *Subgroup lattices of groups*, volume 14 of *De Gruyter Expositions in Mathematics*. Walter de Gruyter & Co., Berlin, 1994.

DEPARTMENT OF MATHEMATICS, SIMON FRASER UNIVERSITY, 8888 UNIVERSITY DRIVE,
BURNABY, BC, V5A 1S6, CANADA