

UNIQUENESS OF EQUILIBRIUM MEASURE FOR A FAMILY OF PARTIALLY HYPERBOLIC HORSESHOES

KRERLEY OLIVEIRA, MARLON OLIVEIRA, AND EDUARDO SANTANA

ABSTRACT. In this paper, we show the uniqueness of equilibrium state for a family of partially hyperbolic horseshoes, introduced in [12] for some classes of continuous potentials. For the first class, the method used here is making use of the Sarig's theory for countable shifts. For this purpose, we study the dynamics of an induced map associated to the horseshoe map, we build a symbolic system with infinitely many symbols that is topologically conjugated to this induced map and we show that the induced potential is locally Hölder and recurrent. For the second class, by following ideas of [29] and [31], we prove that uniqueness for the horseshoe is equivalent to uniqueness for the restriction to a non-uniformly expanding map, which is a hyperbolic potential and then has uniqueness. Both classes include potentials with high variation, differently from previous results for potentials with low variation. We also prove uniqueness when the potential presents its supremum at a special fixed point and less than the pressure.

1. INTRODUCTION

The theory of equilibrium states on dynamical systems was firstly developed by Sinai, Ruelle and Bowen in the sixties and seventies. It was based on applications of techniques of Statistical Mechanics to smooth dynamics. Given a continuous map $f : M \rightarrow M$ on a compact metric space M and a continuous potential $\phi : M \rightarrow \mathbb{R}$, an *equilibrium state* is an invariant measure that satisfies a variational principle, that is, a measure μ such that

$$h_\mu(f) + \int \phi d\mu = \sup_{\eta \in \mathcal{M}_f(M)} \left\{ h_\eta(f) + \int \phi d\eta \right\},$$

where $\mathcal{M}_f(M)$ is the set of f -invariant probabilities on M and $h_\eta(f)$ is the so-called metric entropy of η .

In the context of uniform hyperbolicity, which includes uniformly expanding maps, equilibrium states do exist and are unique if the potential is Hölder continuous and the map is transitive. In addition, the theory for finite shifts was developed and used to achieve the results for smooth dynamics.

Beyond uniform hyperbolicity, the theory is still far from complete. It was studied by several authors, including Bruin, Keller, Demers, Li, Rivera-Letelier, Iommi and Todd [6, 15, 16, 19, 20] for interval maps; Denker and Urbanski [13] for rational maps; Leplaideur, Oliveira and Rios [18] for partially hyperbolic horseshoes; Buzzi, Sarig and Yuri [7, 37], for countable Markov shifts and for piecewise expanding maps in one and higher dimensions. For local diffeomorphisms with some kind of non-uniform expansion, there are results due to Oliveira [22]; Arbieto, Matheus and Oliveira [1]; Varandas and Viana [36]. All of whom proved the existence and uniqueness of equilibrium states for potentials with low oscillation. Also, for this type of maps, Ramos and

Viana [30] proved it for potentials so-called *hyperbolic*, which includes the previous ones. The hyperbolicity of the potential is characterized by the fact that the pressure emanates from the hyperbolic region. In all these studies the maps do not have the presence of critical sets and recently, Alves, Oliveira and Santana proved the existence of at most finitely many equilibrium states for hyperbolic potentials, possible with the presence of a critical set (see [2]). More recently, Santana completed this by showing uniqueness in [32]. Moreover, in this work it is proved that the class of hyperbolic potentials is equivalent to the class of continuous zooming potential (which satisfies a key inequality between free energies). It includes the null potential, which implies existence and uniqueness of measure of maximal entropy. A similar work is developed in [28], where they deal with expanding measures and potentials.

In this work, we deal with equilibrium states in the context of partially hyperbolic dynamics. More precisely, for a special family of partially hyperbolic horseshoes introduced by *Diaz et al* in [12]. For this family, *Leplaideur, Oliveira, Rios* proved in [18] that for any continuous potential we have existence of equilibrium states and for a residual subset of continuous potentials they obtained uniqueness. After that, *Arbieto, Prudente* proved uniqueness in [3] for a class of Hölder potentials that are constant along the center-stable direction. Later on, *Rios, Siqueira* obtained uniqueness in [31] for Hölder potentials with low variation and constant along the unstable direction. Finally, *Ramos, Siqueira* proved the uniqueness of equilibrium state for this horseshoe class and Hölder potentials with low variation without restrictions in the directions.

Our results prove uniqueness for classes of continuous potentials with high variation with some restrictions. First, we prove it for what we call *admissible potentials*, where the corresponding potentials for the subshift associated to the horseshoe are locally Hölder. We create a countable alphabet over the subshift and a full shift to use the Sarig's theory on thermodynamic formalism for countable shifts. We also prove uniqueness for what we call *projective hyperbolic potentials*, with a condition for the restriction to a non-uniformly expanding map which implies hyperbolicity. We do it by proving that the condition implies that the restriction is an expanding potential and based on a result in [2] we have equivalence with hyperbolic potentials. So, by using a result in [30] we obtain uniqueness.

To obtain all this, we prove that the uniqueness of equilibrium state for the horseshoe is equivalent to the uniqueness for the restriction. Moreover, we also give a result for uniqueness when the potential gives supremum at very important fixed point and it is less than the pressure.

2. SETUP AND MAIN RESULTS

We emphasize that the question concerning the uniqueness of equilibrium states and phase transitions for partially hyperbolic maps is wide open. In this paper, we make a contribution to this theme by proving the uniqueness of equilibrium states for a very interesting family of partially hyperbolic horseshoes, introduced by *Diaz et al* in [12] and two classes of continuous potentials. We reproduce here the ideas for the construction of this family of horseshoes. Consider the cube $[0, 1] \times [0, 1] \times [0, 1]$, the parallelepipeds:

$$R_0 := I \times I \times [0, 1/6] \quad \text{and} \quad R_1 := I \times I \times [5/6, 1]$$

and constants $0 < \lambda_0 < \frac{1}{3}$, $\beta_0 > 6$, $0 < \sigma < \frac{1}{3}$ and $3 < \beta_1 < 4$. Let f be the time-one map of the vector field $y' = (1 - y)y$, i.e., f is defined by

$$f^n(y) = \frac{1}{1 - (1 - \frac{1}{y})e^{-n}},$$

for every $n \in \mathbb{Z}$ and $y \neq 0$. Define the family of *horseshoes maps* $F = F_{\lambda_0, \beta_0, \beta_1, \sigma} : R_0 \cup R_1 \rightarrow \mathbb{R}^3$ by

$$F(x, y, z) = (\lambda_0 x, f(y), \beta_0 z),$$

whenever $(x, y, z) \in R_0$ and

$$F(x, y, z) = \left(\frac{3}{4} - \lambda_0 x, \sigma(1 - y), \beta_1 \left(z - \frac{5}{6} \right) \right),$$

for $(x, y, z) \in R_1$.

Let Λ be the maximal invariant set

$$\Lambda = \bigcap_{n \in \mathbb{Z}} F^n(R_0 \cup R_1).$$

The set Λ is a partially hyperbolic set and has an heterodimensional cycle, i.e., the points $Q = (0, 0, 0)$ and $P = (0, 1, 0)$ are saddles of F with indices 2 and 1 respectively, and such that

$$W^s(P) \cap W^u(Q) \neq \emptyset \quad \text{and} \quad W^s(Q) \cap W^u(P) \neq \emptyset.$$

The homoclinic class of P , defined by $H(P, F) = \overline{W^s(P) \cap W^u(P)}$, coincides with Λ and contains the homoclinic class of the point Q that is trivial, i.e., $\Lambda = H(P, F) \supset H(Q, F) = \{Q\}$.

Given $X = (x^s, x^c, x^u) \in \Lambda$, we consider the *central manifold*

$$W^c(X) = \{(x_s, y, x_u); y \in [0, 1]\},$$

tangent to the central subspace E^c at X and the invariant subset

$$\hat{\Lambda} = \{X \in \Lambda ; \Lambda \cap W^c(X) = \{X\}\},$$

composed by the points whose central manifold is trivial. The set $\hat{\Lambda}^c$ can be decomposed as disjoint uncountable union of these segments and in addition the sets $\hat{\Lambda}$ and $\hat{\Lambda}^c$ are dense on Λ . We summarize these results in the following theorem:

Theorem 2.1. ([12], *Theorem 1.*) *Consider $F : R \rightarrow \mathbb{R}^3$ the horseshoe map, then*

- (1) *The diffeomorphism F has a heterodimensional cycle associated with the saddles P and Q . Moreover, there exists a non-transverse intersection between $W^s(Q)$ and $W^u(P)$ whose orbit is contained in $H(P, F)$.*
- (2) *The homoclinic class of Q is trivial and contained in the non-trivial homoclinic class of P . In particular, $H(P, F)$ is not hyperbolic. For Σ_{11} the subshift of finite type where only the transition $1 \rightarrow 1$ is forbidden, there is a surjection*

$$h : \Lambda \rightarrow \Sigma_{11} \quad \text{with} \quad h \circ F = \sigma \circ h.$$

Moreover, there exist infinitely many non-trivial central segments $J = \{x\} \times [a, b] \times \{z\}$ contained in the homoclinic class $H(P, F)$ and h is constant on J .

The next lemma gives us important information about the invariant measures δ_Q and δ_P .

Lemma 2.1. *If μ is an F -invariant measure such that $\mu(R_1) = 0$, then μ is one of the Dirac measures supported in the points P and Q or a convex combination of these measures.*

Proof. Given an ergodic F -invariant measure μ such that $\mu(R_1) = 0$, we have $\text{supp} \mu \subset R_0$, and as the set $\text{supp} \mu$ is invariant, we conclude that it must be contained in $\{0\} \times [0, 1] \times \{0\}$, since any point in R_0 and outside this central curve, eventually leaves R_0 under F -iterations. Thus, using that any point of $\{0\} \times (0, 1] \times \{0\}$ is attracted by P , μ must be δ_Q or δ_P . If μ is an invariant measure such that $\mu(R_1) = 0$, by the Ergodic Decomposition Theorem we conclude that μ must to be a convex combination of these two Dirac measures. \square

The study of equilibrium states for the horseshoe map F started in [18]. In this work, the authors proved that every ergodic invariant measure is *hyperbolic* (in the sense that the Lyapunov exponents are all different from zero) and there is a gap in the set of the central Lyapunov exponents. In addition, they showed that every recurrent point different from Q and P belongs to $\hat{\Lambda}$. As a consequence of these results, they obtained the existence of equilibrium state for every continuous potentials and uniqueness for a residual set of potentials. Moreover, they presented a family of continuous potentials Hölder continuous that has a *phase transition*. More precisely, the following statement holds.

Theorem 2.2. ([18], *Theorem 2.1, 2.2, Proposition 3.3, Remark 4.*)

Let F be partially hyperbolic horseshoe map. Then

- (1) For any recurrent point X different from Q :

$$\liminf_{n \rightarrow +\infty} \frac{1}{n} \log |DF^n(X)|_{E^c} \leq 0.$$

Moreover, any ergodic invariant measure for F different from δ_Q has negative central Lyapunov exponent.

- (2) Let X be a point in Λ . If there exists some natural number k such that $h(F^n(X)) \in \underbrace{[10\dots 01]}_k$ for infinitely many $n \in \mathbb{Z}$, then

$$\Lambda \cap W^c(X) = \{X\}.$$

- (3) Any continuous potential $\phi : \Lambda \rightarrow \mathbb{R}$ admits an equilibrium state, and there exists a residual set in $C^0(\Lambda)$ such that the equilibrium state is unique.
- (4) The family of potentials $\phi_t = t \log |DF|_{E^c}$, for $t \in \mathbb{R}$, has a phase transition, i.e., there exists $t_0 > 0$ such that ϕ_{t_0} admits at least two different equilibrium states.

Remark 2.1. Arbieto and Prudente in [3] proved that the horseshoe map F has an unique maximal entropy measure μ_{max} . Since $h_{top}(F) = \log \frac{1+\sqrt{5}}{2} > 0$, using Lemma 2.1 we obtain that $\mu_{max}(R_1) > 0$. Moreover, by Theorem 2.2 item (2) the set $\hat{\Lambda}$ contains all recurrent points, so we conclude that $\mu_{max}(\hat{\Lambda}) = 1$.

In [3], it was obtained the uniqueness of equilibrium state for the horseshoe map F with respect to Hölder continuous potentials $\phi : \Lambda \rightarrow \mathbb{R}$ that do not depend on the center-stable directions, that is, such that there exists a Hölder continuous potential θ such that $\phi(X) = \theta(z)$ for every point $X = (x, y, z) \in \Lambda$. The uniqueness also was obtained in [31], considering Hölder continuous potentials ϕ on Λ with small variation and that do not depend on the unstable direction, that is, such that there exists a Hölder continuous potential θ such that $\phi(X) = \theta(x, y)$ for every point $X = (x, y, z) \in \Lambda$. The following result in [29] extends it without the restriction on the unstable direction.

Theorem 2.3. Let $F : \Lambda \rightarrow \Lambda$ be the horseshoe map and let $\phi : \Lambda \rightarrow \mathbb{R}$ be a Hölder continuous potential that satisfies the condition:

$$(1) \quad \sup \phi - \inf \phi < \frac{\log \omega}{2}, \quad (\text{small variation})$$

where $\omega = \frac{1+\sqrt{5}}{2}$. Then there exists a unique equilibrium state for the system F with respect to the potential ϕ .

In [20], for rational maps, it was considered a class of potential called hyperbolic that include potentials that satisfies (1) and also some potential with large variation ([20], Lemma A.1) . In [30] it was introduced an another class of hyperbolic potentials that include potentials that satisfy the condition (1). In this way, questions naturally arise:

Question 1. *Are there hyperbolic potentials in the senses of the works [20] and [30] with large variation and that admit uniqueness of equilibrium states for the Horseshoe F ?*

Our results provide answers to this question. That is, we obtain hyperbolic potentials in both contexts that admit a unique equilibrium state and with big variation.

In order to state our main results, we need some definitions.

In the Section 3, we construct an inducing scheme using a parametrized family of subsets $\Sigma_\alpha \subset \Sigma$ and a respective inducing time $\rho : \Sigma_\alpha \rightarrow \mathbb{N}$ such that the induced map $T = \sigma^\rho : \Sigma_\alpha \rightarrow \Sigma_\alpha$ is topologically conjugated to a full shift $\sigma : S^{\mathbb{Z}} \rightarrow S^{\mathbb{Z}}$ on an countable alphabet S , i.e., there exists a continuous homeomorphism $\Pi : S^{\mathbb{Z}} \rightarrow \Sigma_\alpha$ such that $\Pi \circ \sigma = T \circ \Pi$. Given a potential $\varphi : \Sigma \rightarrow \mathbb{R}$, we define the induced potential as $\varphi_\rho(w) = \sum_{k=0}^{\rho(w)-1} \varphi(\sigma^k(w))$ and $\Psi = \varphi_\rho \circ \Pi$.

Now, we define the class of potentials that we consider in our first main result.

Definition 2.1. *A continuous potential $\phi : \Lambda \rightarrow \mathbb{R}$ is called admissible if satisfies:*

- (C₁) *The potential $\Psi = \varphi_\rho \circ \Pi$ associated to $\varphi := \phi \circ h^{-1} : \Sigma \rightarrow \mathbb{R}$ is locally Hölder continuous.*
- (C₂) *There exists a natural $n \in \mathbb{N}$ such that*

$$P_{top}(\phi) > \sup_{X \in \Lambda} \frac{\phi_n(X)}{n},$$

$$\text{where } \phi_n(X) = \sum_{k=0}^{n-1} \phi(F^k(X)).$$

Remark 2.2. *If $\psi = \phi_n/n$ then $P_{top}(\psi) = P_{top}(\phi)$ and the equilibrium states for ψ and ϕ coincide. Thus, we can assume without loss of generality that $n = 1$. Another remark about admissible potentials is that since the condition (C₂) is open in the C^0 -topology and the topological entropy of F is positive, it is easy to see that ϕ satisfy (C₂) if $\sup |\phi|$ is small enough.*

Remark 2.3. *Let μ be an F -invariant probability measure on Λ . Denote by $\lambda_\mu^c = \int \log |DF|_{E^c} d\mu$ the central Lyapunov exponent associated to μ . Using arguments similar to [18], Lemma 4.3 and Corollary 4.4], it is possible to show that the central Lyapunov exponents of the equilibrium states associated to admissible potentials are negative.*

Now, we state our first main result:

Theorem A. *Let $F : \Lambda \rightarrow \Lambda$ be the horseshoe map. If $\phi : \Lambda \rightarrow \mathbb{R}$ is an admissible potential, then there exists a unique equilibrium state for ϕ .*

In [29], the authors proved that if ϕ is a Hölder continuous potential with small variation, i.e., $\sup \phi - \inf \phi$ is small enough, then ϕ has a unique equilibrium state. The result of Theorem A differs from [29] due to the strategy of the proof of uniqueness of equilibrium states and also due to the fact that the class of admissible potential contains potentials that do not satisfy the condition of small variation. Indeed, in Section 9.1, we construct new examples of potentials that are admissible and, consequently, have a unique equilibrium state. In fact, we show that if ϕ is a Hölder potential that is constant on the set $\{(x, y, z); z < c\}$, for some $5/6 < c < 1$ and $\sup \phi = \phi(Q)$, then there exists a non-empty open interval $I \subset \mathbb{R}$ such that if $t \in I$ then $t\phi$ is admissible. A precise description of I is given in Proposition 9.1.

Now we reproduce the definition of a map G , as introduced in [31], which is related to the projection of F^{-1} in two center-stable planes. We consider an abstract space consisting of three rectangles S_1 , S_2 and S_3 defined as follows:

$$\begin{aligned} S_1 &= [0, \lambda_0] \times [0, 1] \times \{0\}, \\ S_2 &= [3/4 - \lambda_0, 3/4] \times [0, \sigma] \times \{0\}, \\ S_3 &= [0, \lambda_0] \times [0, 1] \times \{5/6\}. \end{aligned}$$

The rectangles are inside two planes $P_0 = [0, 1] \times [0, 1] \times \{0\}$ and $P_1 = [0, 1] \times [0, 1] \times \{5/6\}$. Let $g_0 : [0, 1] \rightarrow \mathbb{R}$, $g_0(y) = f^{-1}(y)$ and let $g_1 : [0, 1] \rightarrow \mathbb{R}$, $g_1(y) = 1 - \sigma^{-1}y$. Let $\alpha = \lambda_0^{-1}$.

We define a map $G : \cup_{i=1}^3 S_i \rightarrow P_1 \cup P_2$ by its restrictions G_i to each rectangle S_i as follows:

$$\begin{aligned} G_1(x, y, z) &= (\alpha x, g_0(y), 0), \\ G_2(x, y, z) &= (\alpha(3/4 - x), g_1(y), 5/6), \\ G_3(x, y, z) &= (\alpha x, g_0(y), 0). \end{aligned}$$

The behavior of the map G is similar on S_1 and S_3 . In these rectangles, the vertical direction is expanding. In the horizontal direction, G sends the points from the right side to the left side, except for the extreme points whose the x coordinates are fixed. In the rectangle S_2 both directions are expanding.

We consider Ω the maximal invariant set of the union of the rectangles:

$$\Omega := \bigcap_{n \in \mathbb{N}} G^{-n} \left(\bigcup_{i=1}^3 S_i \right),$$

and from now on we call G the restriction $G|_{\Omega} : \Omega \rightarrow \Omega$. See details of the map G in [31] and [29], as the proof that this map is non-uniformly expanding. We also observe that G is strongly mixing.

The map G is very important because we can use results for non-uniformly expanding maps and transfer them to the horseshoe F , as in the following main result, which allows potentials with big variation. This is a novelty with respect to previous results.

Theorem B. *Let $F : \Lambda \rightarrow \Lambda$ be the horseshoe map and $G : \Omega \rightarrow \Omega$ be the projection of F^{-1} as defined above. The system (F^{-1}, ϕ) has uniqueness of equilibrium state if, and only if, the system $(G, \phi|_{\Omega})$ does so.*

We define a class of potentials that we will use to obtain uniqueness of equilibrium states in our third main result. They are related to the non-uniformly expanding map G as follows.

Definition 2.2. *A continuous potential $\phi : \Lambda \rightarrow \mathbb{R}$ is called projective hyperbolic if the following holds:*

- (D₁) *The restriction $\varphi := \phi|_{\Omega}$ is Hölder and there exists a measurable function $u : \Omega \rightarrow \mathbb{R}$ such that $\varphi \geq u - u \circ G$.*
- (D₂) *If \mathcal{E} denotes the set of G -invariant expanding measures on Ω , we have that*

$$\sup_{\nu \in \mathcal{E}^c} \left\{ \int \varphi d\nu \right\} < h(G) - \sup_{\nu \in \mathcal{E}^c} \{h_{\nu}(G)\}.$$

The class of projective hyperbolic potentials contains some potentials with big variation, as will be seen in Section 9.2.

We can now state our third main result as a class of potentials that includes potentials that do not satisfy the condition of small variation and also constant on the fibers in the z -axis:

Theorem C. *Let $F : \Lambda \rightarrow \Lambda$ be the horseshoe map. If $\phi : \Lambda \rightarrow \mathbb{R}$ is a projective hyperbolic potential, then there exists a unique equilibrium state μ_ϕ for the system G with respect to the potential $\varphi = \phi|_\Omega$. It is equivalent to existing a unique equilibrium state for the horseshoe F .*

We observe that in Section 7 we show that if ϕ is a projective hyperbolic potential, then $\varphi = \phi|_\Omega$ is a hyperbolic potential (Proposition 7.1) and since G is a non-uniformly expanding map that is also strongly mixing (with density of pre-images of the points in the expanding set), we can obtain uniqueness of equilibrium states by [30][Theorem 2].

In the following, we state our fourth main result, which gives a condition for uniqueness of potentials with supremum at the fixed point Q and it is less than the pressure.

Theorem D. *Let $F : \Lambda \rightarrow \Lambda$ be the horseshoe, η the unique measure of maximal entropy for the non-uniformly expanding map $G : \Omega \rightarrow \Omega$ and ϕ a Hölder potential such that it holds that $\sup \phi|_\Omega = \phi|_\Omega(Q) < P(\phi|_\Omega)$. Then, we obtain uniqueness of equilibrium state. Additionally, we have either every Hölder potential φ such that $\sup \varphi|_\Omega < P(\varphi|_\Omega)$ has uniqueness of equilibrium state or there exists a Hölder potential ϕ such that $\phi|_\Omega(Q) < \sup \phi|_\Omega < P(\phi|_\Omega)$ and ϕ does not have uniqueness of equilibrium states.*

The Theorems A and C of this article differs from Theorem 2.3 because of the potentials in Theorem A and C include examples with big variation and also the strategy used to obtain uniqueness can be applied for other classes of systems. The same can be said for our Theorem C in this article, by comparing with the Theorem B in [31].

The work is organized as follows: Section 3 is devoted to the construction of the symbolic structures that allow us to make use of Sarig's theory for countable shifts and show our first main theorem. In the Section 4 we prove the first main result and we describe new classes of potentials that are admissible in Section 9.1 and fits in the hypothesis of Theorem A. Section 5 is devoted to the proof of Theorem B. In Section 6 we prove an equivalence between hyperbolic and expanding potentials (see definitions there), which will be useful to show that the restrictions of projective hyperbolic potentials are hyperbolic potentials. In Section 7 we prove our second main result. Section 8 is devoted to prove our third main result. Finally, in Section 9.1 we give a family of examples of admissible potentials and Section 9.2 gives a construction of projective hyperbolic potentials.

3. CONSTRUCTING THE INDUCING SCHEME

In this section, we construct an inducing scheme associated to the horseshoe map F that admits a symbolic representation as a shift over a countable alphabet.

We define the sequences

$$d_n^+(w) = \frac{\#\{k ; w_k = 1, 0 \leq k \leq n-1\}}{n}.$$

For each $\alpha > 0$ we consider the subset

$$(2) \quad \Sigma_\alpha^+ = \{w \in [1]; \overline{\lim}_n d_n^+(w) > \alpha\}.$$

The set Σ_α^+ is composed by the sequences with upper frequency of digits 1 greater than α . We define on Σ_α^+ the α -return function $\rho : \Sigma_\alpha^+ \rightarrow \mathbb{N}$ by

$$\rho(w) = \min\{k > 1 ; w_{k-1} = 1 \text{ and } d_k^+(w) > \alpha\}.$$

Note that Σ_α^+ can be decomposed into level sets of the function ρ ,

$$\Sigma_\alpha^+ = \bigcup_i \Sigma_i,$$

where Σ_i is given by $\Sigma_i = \{w \in \Sigma_\alpha^+; \rho(w) = i\}$.

Now, we consider the set

$$\Sigma_\alpha^- = \{w \in [1]; \exists n_k \rightarrow +\infty, \sigma^{-(n_1+n_2+\dots+n_k)}(w) \in \Sigma_{n_k}, \forall k \geq 1\}.$$

We define the set $\Sigma_\alpha := \Sigma_\alpha^+ \cap \Sigma_\alpha^-$, which is invariant under σ^ρ . By using Item 2 of Theorem 2.2, we have that for any $\omega \in \Sigma_\alpha$, the set $h^{-1}(\omega)$ is a single point, i.e., $\Sigma_\alpha \subset \Sigma = h(\hat{\Lambda})$.

We define the *induced map* $T : \Sigma_\alpha \rightarrow \Sigma_\alpha$ associated to ρ as

$$T(\omega) = \sigma^{\rho(\omega)}(\omega).$$

Consider the *tower* associated to ρ , defined by

$$(3) \quad W = \bigcup_{i>1} \bigcup_{k=0}^{i-1} \sigma^k(\Sigma_i).$$

Let ν be a T -invariant measure such that $\int \rho d\nu < \infty$, then we define the σ -invariant measure, that will be called *lifted* of ν by

$$(4) \quad \mathcal{L}(\nu)(A) := \left(\int \rho d\nu \right)^{-1} \sum_{i>1} \sum_{k=0}^{i-1} \nu(\sigma^{-k}(A) \cap \Sigma_i)$$

for $A \subset \Sigma_{11}$.

The map \mathcal{L} that acts on $\mathcal{M}_T(W)$ is not necessarily surjective over $\mathcal{M}_\sigma(\Sigma_{11})$. Given a measure $\mu \in \mathcal{M}_\sigma(\Sigma_{11})$, if there exists ν in $\mathcal{M}_T(W)$ such that $\mathcal{L}(\nu) = \mu$, then we say that μ is a *liftable measure*. We denote this class of measures by

$$(5) \quad \mathcal{M}_L(\sigma, W) := \{\mu \in \mathcal{M}_\sigma; \exists \nu, \mathcal{L}(\nu) = \mu, \mu(W) = 1\}.$$

Given a measurable set $A \subset \Sigma$ and $i > 1$, we define

$$e(i, A) := \frac{\#\{0 \leq k \leq i-1; \sigma^k(\Sigma_i) \cap A \neq \emptyset\}}{i}.$$

The following lemma will be useful to ensure that a measure is liftable.

Theorem 3.1 ([26], Theorem 3.1). *An ergodic σ -invariant measure μ that satisfies $\mu(W) > 0$ is a liftable measure if there exists a number $N \geq 0$ and a subset $A \subset \Sigma$ such that*

$$\mu(A) > \sup_{i>N} e(i, A).$$

Let $\varphi : \Sigma \rightarrow \mathbb{R}$ be a potential, we define the *induced potential* associated to φ by

$$(6) \quad \varphi_\rho(w) := \sum_{k=0}^{i-1} \varphi(\sigma^k(w))$$

whenever $w \in \Sigma_i$.

The generalized Kač-Abramov formula gives an interesting relation between the entropy of the original system and the entropy of the induced system.

Proposition 3.1 (see [38], Theorem 5.1 and [25], Theorem 2.3). *If ν is a measure in $\mathcal{M}_T(W)$ such that $\int \rho d\nu < \infty$, then*

$$h_{\mathcal{L}(\nu)}(\sigma) \int \rho d\nu = h_\nu(T).$$

Let φ be a potential and φ_ρ its induced potential. If $\int \varphi_\rho d\nu < \infty$ then

$$\int \varphi d\mathcal{L}(\nu) \int \rho d\nu = \int \varphi_\rho d\nu.$$

We define the quantity

$$P_L(\varphi) := \sup_{\mu \in \mathcal{M}_L(\sigma, W)} \{h_\mu(\sigma) + \int_W \varphi d\mu\},$$

that will be called *relative pressure*. An invariant probability measure that attains the supremum is called *relative equilibrium state*.

Now, it will be established a relation between the induced system and a countable Markov shift.

Firstly, note that each Σ_k can be decomposed as a disjoint finite union of k -cylinders, that is, $\Sigma_k = \bigcup_{i=1}^{r_k} D_i^k$, where $D_i^k = [w_0 w_1 \dots w_{k-1}]$. Let $\hat{\pi}$ be the map that sends a cylinder to a word

$$\hat{\pi}(D_i^k) = w_0 w_1 \dots w_{k-1}.$$

Using the new alphabet $S = \{D_i^k; k \geq 1, 1 \leq i \leq r_k\}$ we define the *coding map* $\Pi : S^{\mathbb{Z}} \rightarrow \Sigma_\alpha$ by the following amalgamation

$$\Pi(\hat{w}) = (\dots \hat{\pi}(D_{i_{k-1}}^{k-1}) \hat{\pi}(D_{i_{k_0}}^{k_0}) \hat{\pi}(D_{i_{k_1}}^{k_1}) \dots),$$

where $\hat{w} = (D_{i_{k_n}}^{k_n})_{n \in \mathbb{Z}} \in S^{\mathbb{Z}}$.

Observe that by construction Π is well defined, and it is a conjugacy between T and σ (the countable Markov shift that acts on $S^{\mathbb{Z}}$). In fact, given a sequence $w \in \Sigma_\alpha$, by definition we have that there exists a (unique) sequence $\hat{w} = (D_{i_{k_n}}^{k_n})_{n \in \mathbb{Z}}$ in $S^{\mathbb{Z}}$, such that $\sigma^{k_n}(w) \in D_{i_{k_{n+1}}}^{k_{n+1}}$ for $n \geq 0$ and $\sigma^{-k_n}(w) \in \Sigma_{k_n}$ for $n < 0$ i.e., $\Pi(\hat{w}) = w$.

Now, we discuss the thermodynamic formalism of countable Markov shifts. The main references are [27], [33], [34] and [35].

Let σ be the full shift on $S^{\mathbb{Z}}$ and $\Psi : S^{\mathbb{Z}} \rightarrow \mathbb{R}$ a potential, the k -variation is defined by

$$\text{Var}_k(\Psi) = \sup_{[i_{-k+1} \dots i_{k-1}]} \sup_{\hat{w}, \hat{s} \in [i_{-k+1} \dots i_{k-1}]} \{|\Psi(\hat{w}) - \Psi(\hat{s})|\},$$

where $[i_{-k+1} \dots i_{k-1}]$ is the two-sided cylinder of all the sequences $w = (w_i)_{i \in \mathbb{Z}}$ with $w_{-k+1} = i_{-k+1}, \dots, w_0 = i_0, \dots, w_{k-1} = i_{k-1}$.

We say that Ψ has *strongly summable variations* if

$$\sum_{k \geq 1} k \text{Var}_k(\Psi) < \infty,$$

and Ψ is *locally Hölder continuous* if there exists $C > 0$ and $a \in (0, 1)$ such that for all $k \geq 1$

$$\text{Var}_k(\Psi) \leq C a^k.$$

If $\Psi : S^{\mathbb{Z}} \rightarrow \mathbb{R}$ is locally Hölder continuous then the *Gurevich pressure* of Ψ is defined by

$$(7) \quad P_G(\Psi, a) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \sum_{\substack{\sigma^n(\hat{w}) = \hat{w} \\ \hat{w} \in [a]}} \exp(\Psi_n(\hat{w})) \mathcal{X}_{[a]}(\hat{w}),$$

where $\mathcal{X}_{[a]}$ is the indicator function on the cylinder set $[a]$.

In [33] it was proved that the limit (7) exists and it is independent from $a \in S$. We denote it by $P_G(\Psi)$.

Let $\mathcal{M}_\sigma(S^{\mathbb{Z}})$ be the set of σ -invariant Borel probabilities on $S^{\mathbb{Z}}$ and

$$\mathcal{M}_\sigma(\Psi) := \left\{ \eta \in \mathcal{M}_\sigma(S^{\mathbb{Z}}); \int \Psi d\eta > -\infty \right\}.$$

A measure η_Ψ in $\mathcal{M}_\sigma(S^{\mathbb{Z}})$ is an equilibrium state for a potential Ψ if satisfies

$$h_{\eta_\Psi}(\sigma) + \int \Psi d\eta_\Psi = \sup_{\eta \in \mathcal{M}_\sigma(\Psi)} \left\{ h_\eta(\sigma) + \int \Psi d\eta \right\}.$$

We say that η is a Gibbs measure for Ψ if there exists a constant C such that for a cylinder $[i_0 \dots i_{n-1}]$ in $S^{\mathbb{Z}}$ and $\hat{w} \in [i_0 \dots i_{k-1}]$ we have

$$C^{-1} \leq \frac{\eta([i_0 \dots i_{n-1}])}{e^{(\Psi_n(\hat{w}) - nP_G(\Psi))}} \leq C.$$

The next result can be found in [27][Theorem 3.1].

Theorem 3.2. *Assume that $\sup_{\hat{w} \in S^{\mathbb{Z}}} \Psi(\hat{w}) < \infty$ and Ψ has strongly summable variations. Then*

(1) *the variational principle for Ψ holds*

$$P_G(\Psi) = \sup_{\eta \in \mathcal{M}_\sigma(\Psi)} \left\{ h_\eta(\sigma) + \int \Psi d\eta \right\}.$$

(2) *If $P_G(\Psi) < \infty$, then there exists a unique σ -invariant ergodic Gibbs measure η_Ψ for Ψ .*

(3) *If furthermore, $h_{\eta_\Psi}(\sigma) < \infty$, then $\eta_\Psi \in \mathcal{M}_\sigma(\Psi)$ and it is the unique equilibrium measure for Ψ .*

We say that an invariant Borel probability measure η associated to the continuous transformation T has *exponential decay of correlations* for a class \mathcal{H} of functions if there exists $0 < \theta < 1$ such that, for any $h_1, h_2 \in \mathcal{H}$,

$$\left| \int h_1(T^n(x))h_2(x) d\eta - \int h_1(x) d\eta \int h_2(x) d\eta \right| \leq K\theta^n,$$

for some $K = K(h_1, h_2) > 0$.

The measure η satisfies the *central limit theorem* (CLT) for functions in \mathcal{H} if there exist $\sigma \in \mathbb{R}$ such that $\frac{1}{\sqrt{n}} \sum_{i=0}^{n-1} (h(T^i(x)) - \int h d\eta)$ converges in law to a normal distribution $\mathcal{N}(0, \sigma)$ for any $h \in \mathcal{H}$.

Proposition 3.2 ([27] Theorem 4.7). *Assume that $P_G(\Psi)$, $\sup_{\hat{w} \in S^{\mathbb{Z}}} (\Psi(\hat{w})) < \infty$ and that Ψ is locally Hölder continuous. If $h_{\eta_\Psi}(\sigma) < \infty$, then the measure η_Ψ has exponential decay of correlations and satisfies the CLT with respect to the class of bounded locally Hölder continuous.*

4. PROOF OF THEOREM A

The proof of Theorem A is divided into two steps. The first one is to show the uniqueness of equilibrium state for the induced map and the second one is to obtain the uniqueness for the horseshoe map.

Consider the constant $\alpha \in (0, \frac{2}{3})$ and the inducing scheme (ρ, Σ_α) , obtained as in the Section 3, the next result gives us sufficient condition for a potential φ to admit a unique relative equilibrium state.

Theorem 4.1. *Let $\varphi : \Sigma \rightarrow \mathbb{R}$ be a continuous potential and φ_ρ be the induced potential satisfying the following conditions:*

- (P₁) *The potential $\Psi := \varphi_\rho \circ \Pi$ is locally Hölder continuous.*
- (P₂) *There exist a natural $n \in \mathbb{N}$ such that $P_L(\varphi) > \sup \frac{\varphi_n}{n}$.*

Then, there exists a unique relative equilibrium state ν_φ . Moreover, the measure ν_φ has exponential decay of correlations and satisfies the (CLT) with respect to class of functions whose induced functions on Σ_α are bounded Hölder continuous.

Note that the pressure $P_L(\varphi)$ is finite, so we take $\varphi_\rho = \overline{\varphi - P_L(\varphi)}$.

Claim 4.1. *If φ satisfies (P₂), then there exists a constant $\epsilon > 0$ such that*

$$(8) \quad \sum_{i>1} i \sup_{w \in \Sigma_i} e^{\varphi_\rho(w)+i\epsilon} < \infty.$$

Proof of Claim. By the property (P₂) and Remark 2.2 we get $\epsilon_0 > 0$ such that $P_L(\varphi) > \sup \varphi + \epsilon_0$ and thus for $P := P_L(\varphi)$, $w \in \Sigma_i$ and all i it occurs that

$$-\epsilon_0 > \frac{\sum_{k=0}^{i-1} \varphi(\sigma^k(w))}{i} - P.$$

This implies that $\sum_{k=0}^{i-1} \varphi(\sigma^k(w)) - iP < -i\epsilon_0$ and $e^{\varphi_\rho(w)} < e^{-i\epsilon_0}$. Since the estimate is independent of w , it is possible to obtain ϵ small enough such that

$$(9) \quad \sup\{e^{\varphi_\rho(w)+i\epsilon}; w \in \Sigma_i\} \leq e^{-i\epsilon_0}.$$

Hence, by summing the expression (9) over i we get

$$\sum_i i \sup\{e^{\varphi_\rho(w)+i\epsilon}; w \in \Sigma_i\} \leq \sum_i i e^{-i\epsilon_0} < +\infty.$$

Thus proving the Claim 4.1. □

We now consider the potential $\Psi = \varphi_\rho \circ \Pi$.

Claim 4.2. *If φ satisfies (P₁) and (P₂), then Ψ the Gurevich pressure is finite.*

Proof of Claim. By the condition (P₁) the Gurevich Pressure $P_G(\Psi)$ is well defined. Let n be positive integer and a cylinder $[D_{i_1}^{l_1} \dots D_{i_n}^{l_n}]$ in $S^{\mathbb{Z}}$, there exists a unique sequence $w \in$

$\Pi([D_{i_1}^{l_1} \dots D_{i_n}^{l_n}])$ such that $T^n(w) = w$. Thus, by fixing $a > 1$ and $1 \leq j \leq r_a$ we get

$$\begin{aligned} \left(\sum_{i>1} \sup_{w \in \Sigma_i} e^{\varphi_\rho(w)} \right)^n &= \sum_{i_1, \dots, i_n} C_{i_1, \dots, i_n} \sup_{w \in \Sigma_{i_1}} e^{\varphi_\rho(w)} \dots \sup_{w \in \Sigma_{i_n}} e^{\varphi_\rho(w)} \\ &\geq \sum_{i_2, \dots, i_n} C_{a, i_2, \dots, i_n} \sup_{w \in \Sigma_a} e^{\varphi_\rho(w)} \sup_{w \in \Sigma_{i_2}} e^{\varphi_\rho(w)} \dots \sup_{w \in \Sigma_{i_n}} e^{\varphi_\rho(w)} \\ &\geq \sum_{\substack{T^n(w)=w \\ w \in D_j^a}} e^{\sum_{k=0}^{n-1} \varphi_\rho(T^k w)}. \end{aligned}$$

Then by (8) and (12), we have

$$\begin{aligned} P_G(\Psi) &= \lim_{n \rightarrow \infty} \frac{1}{n} \log \sum_{\substack{\sigma^n(\hat{w})=\hat{w} \\ \hat{w} \in [D_j^a]}} e^{\sum_{j=0}^{n-1} \Psi(\sigma^j(\hat{w}))} \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \log \sum_{\substack{T^n(w)=w \\ w \in D_j^a}} e^{\sum_{k=0}^{n-1} \varphi_\rho(T^k(w))} \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{n} \log \left(\sum_{i>1} \sup_{w \in \Sigma_i} e^{\varphi_\rho(w)} \right)^n < +\infty. \end{aligned}$$

□

Note that by using the same arguments we prove that the Gurevich pressure of the potential Ψ_δ on $S^{\mathbb{Z}}$, given by $\Psi_\delta(\hat{w}) = \varphi_\rho(w) + \delta\rho(w)$ is finite for some $\delta > 0$ small enough. A potential Ψ that satisfies this is called *positive recurrent potential*.

Proof of Theorem 4.1. By Theorem 3.2, we have the variational principle

$$P_G(\Psi) = \sup_{\eta \in \mathcal{M}_\sigma(\Psi)} \{h_\eta(\sigma) + \int \Psi d\eta\}.$$

Moreover, there exists a unique Gibbs measure η_Ψ for Ψ , so the measure $\nu = \Pi_* \eta_\Psi$ also has this property with respect to the potential φ_ρ , that is, there exists a constants $K > 0$ such that for $n > 1$ and $\Sigma_n = \cup_{i=1}^{r_n} D_i^n$ we have

$$(10) \quad K^{-1} \leq \frac{\nu(D_i^n)}{e^{\varphi_\rho(w) - P}} \leq K,$$

for $w \in D_i^n$ and $P = P_G(\Psi)$.

Now, we define

$$c(\alpha) = \limsup \frac{1}{n} \log \left[\sum_{k \leq \alpha n} \binom{n}{k} \right].$$

By using standing theorem for $n!$, we can easily check that $\lim_{\alpha \rightarrow 0} c(\alpha) = 0$. Choose $\alpha > 0$, such that $c(\alpha) < P_L(\varphi) - \sup(\varphi)$. We observe that

$$(11) \quad \begin{aligned} \limsup \frac{1}{n} \log r_n &\leq \limsup \frac{1}{n} \log \#\{w = (w_0 w_1 \dots); d_{n-1}(w) \leq \alpha\} \\ &< c(\alpha) < P_L(\varphi) - \sup(\varphi). \end{aligned}$$

Thus, for n large enough we have that $r_n < e^{\epsilon n}$. Hence by summing (10) with respect to $n > 1$ and using (8), we have

$$\begin{aligned}
 \int_{\Sigma_\alpha} \rho d\nu &= \sum_{n>1} n\nu(\Sigma_n) \\
 &= \sum_{n>1} n \sum_{i=1}^{r_n} \nu(D_i^n) \\
 (12) \quad &\leq \frac{K}{e^P} \sum_{n>1} nr_n \sup_{w \in D_i^n} \{e^{\varphi_\rho(w) + \epsilon n}\} < +\infty.
 \end{aligned}$$

By using (12) and Abramov-Kac formulas, we have

$$\begin{aligned}
 h_{\eta_\Psi}(\sigma) &= h_\nu(T) \\
 &= \int \rho d\nu \cdot h_{\mathcal{L}(\nu)}(\sigma) < +\infty.
 \end{aligned}$$

Since $P_L(\varphi) > -\infty$ and $\int \varphi d\mathcal{L}(\nu) > -\infty$, we have

$$\begin{aligned}
 \int \Psi d\eta_\Psi &= \int \varphi_\rho d\nu \\
 &= \left(\int \rho d\nu \right) \cdot \int \varphi - P_L(\varphi) d\mathcal{L}(\nu) > -\infty.
 \end{aligned}$$

Hence $\eta_\Psi \in \mathcal{M}_\sigma(\Psi)$, and by using Theorem 3.2 once again, we obtain that the measure η_Ψ is the unique equilibrium state of Ψ . However, we can define the measure $\mathcal{L}(\nu)$ that is contained in $\mathcal{M}_L(\sigma, W)$. By using that Ψ is a positive recurrent potential and [[25], Theorem 4.4] we obtain that the measure $\mathcal{L}(\nu)$ is the unique relative equilibrium state for $P_L(\varphi)$.

In order to complete the proof of Theorem 4.1, we just need to check the properties of Proposition 3.2. We make use of the following lemma.

Lemma 4.1. *The measure ν has exponential tail, that is, there exist $C > 0$ and $0 < \vartheta < 1$ such that for all $n > 0$,*

$$\nu(\{w \in \Sigma_\alpha; \rho(w) \geq m\}) \leq C\vartheta^n.$$

Proof. Note that $\nu(\{w \in \Sigma_\alpha; \rho(w) \geq m\}) = \sum_{n \geq m}^\infty \sum_{i=1}^{r_n} \nu(D_i^n)$, then we obtain the result in a similar way to (12). \square

It follows that by Lemma 4.1 and the results [[37], Theorem 2, Theorem 3] that the measure $\mathcal{L}(\nu)$ has exponential decay of correlations and satisfies the (CLT) with respect to class of functions whose induced functions on Σ_α are bounded Hölder continuous. \square

Proof of Theorem A. By using Theorem 2.2, we have that every continuous potential ϕ has an equilibrium state. Assume that μ_1 and μ_2 are ergodic equilibrium states for F and let us prove that $\mu_1 = \mu_2$.

By hypothesis (C_2), there exists a natural n such that $P(\phi) > \sup \frac{\phi_n}{n}$, so

$$\phi(Q) \leq \sup \frac{\phi_n}{n} < P(\phi).$$

Then, the Dirac measure δ_Q , can not be an equilibrium state for ϕ . Analogously, we can obtain the same for δ_P . Therefore, by Lemma 2.1, we can consider a constant $\alpha > 0$ small enough such that $\mu_i(R_1) > \alpha$ for $i = 1, 2$.

Now, we consider the inducing scheme (ρ, Σ_α) such that $\Sigma_\alpha \subset \Sigma_{11}$ is the subset composed by the sequences with frequency of symbols 1's at least α , as defined in (2), and the associated tower W as in (3).

For each $i = 1, 2$, we denote by ν_i the push-forward of μ_i with respect to semi-conjugacy h , i.e., the measure defined on $A \subset \Sigma_{11}$ by

$$\nu_i(A) = \mu_i(h^{-1}(A)).$$

The measure ν_i is ergodic and satisfies $\nu_i([1]) = \mu_i(R_1) > \alpha > 0$, so since W is an σ -invariant set, by ergodicity of ν_i we have $\nu_i(W) = 1$. Hence, applying the Lemma 3.1 for $A = [1]$, we conclude that ν_i is a liftable measure, moreover it is a relative equilibrium state for the potential $\varphi = h_*\phi$. In the other hand, the potential ϕ satisfies the conditions (C_1) and (C_2) , so φ satisfies the properties (P_2) and (P_1) . Then, by Theorem 4.1, there exists a unique relative equilibrium state, so $\nu_1 = \nu_2$. Therefore, we conclude that $\mu_1 = \mu_2$. The uniqueness is proven. \square

5. PROOF OF THEOREM B

In this section, we prove that uniqueness of equilibrium state for the restriction of the potential to Ω and the map $G : \Omega \rightarrow \Omega$ is equivalent to uniqueness of equilibrium state for the horseshoe map $F : \Lambda \rightarrow \Lambda$. The technique is based on the strategy given in the works [31] and [29]. First, we prove that potentials ϕ with uniqueness for the restriction $\phi|_\Omega$ and such that they do not depend on the z coordinate have uniqueness. After that, we prove that potentials without this condition have cohomologous satisfying this and the uniqueness is obtained in the general case.

In order to make the paper self contained, we will prove our result following the proofs in the works [31] and [29] along the same lines. All the strategy works here because the unique property used with respect to the restriction to Ω is the uniqueness obtained from the small variation of the potential and the non-uniform expansion of the map G .

As it is said in [31], some of the following ideas are based in the constructions in [8]. We define the projection of the parallelepipeds R_0 and R_1 onto the planes P_0 and P_1 , $\pi : R_0 \cup R_1 \rightarrow P_0 \cup P_1$ by

$$\pi(x, y, z) = \begin{cases} (x, y, 0), & \text{if } (x, y, z) \in R_0, \\ (x, y, \frac{5}{6}), & \text{if } (x, y, z) \in R_1. \end{cases}$$

It is straightforward to check that π is continuous, surjective and

$$\pi \circ F^{-1} = G \circ \pi.$$

Lemma 5.1 ([31], Lemma 5.1). *For each $X \in \Omega$ we have $h(F^{-1}, \pi^{-1}(X)) = 0$.*

Since the projection π is actually a semiconjugacy between the inverse horseshoe F^{-1} and the map G , we have (see [4])

$$h_{\text{top}}(F^{-1}) \leq h_{\text{top}}(G) + \sup\{h(F^{-1}, \pi^{-1}(X)) \mid X \in \Omega\}.$$

By Lemma 5.1 we get

$$h_{\text{top}}(F^{-1}) \leq h_{\text{top}}(G).$$

On the other hand, because π is a semiconjugacy, we have the other inequality immediately (see [5]), which gives us

$$h_{\text{top}}(F^{-1}) = h_{\text{top}}(G).$$

Thus, we have proved that the topological entropies of these two maps are equal.

Proposition 5.1 ([31], Proposition 5.2). *For $\varphi = \phi|_\Omega \circ \pi$ we have $P_{\text{top}}(F^{-1}, \varphi) = P_{\text{top}}(G, \phi|_\Omega)$.*

Proof. Given μ a F^{-1} -invariant probability measure, taking $\nu = \mu \circ \pi^{-1}$ a result by Ledrappier and Walters (see [17]) guarantees that

$$h_\mu(F^{-1}) + \int \phi_{|\Omega} \circ \pi d\mu \leq h_\nu(G) + \int \phi_{|\Omega} d\nu + \int_\Omega h(F^{-1}, \pi^{-1}(X)) d\mu(X) = h_\nu(G) + \int \phi_{|\Omega} d\nu \leq P_{\text{top}}(G),$$

because by Lemma 5.1 we have $h(F^{-1}, \pi^{-1}(X)) = 0$ for each $X \in \Omega$ and by semiconjugacy ν is G -invariant.

Going back to the inequality, we have for $\varphi = \phi_{|\Omega} \circ \pi$ and any F^{-1} -invariant probability μ

$$h_\mu(F^{-1}) + \int \varphi d\mu \leq P_{\text{top}}(G) \implies P_{\text{top}}(F^{-1}, \varphi) \leq P_{\text{top}}(G, \phi_{|\Omega}).$$

In the other hand, π is a semiconjugacy, which implies (see [5])

$$P_{\text{top}}(G, \phi_{|\Omega}) \leq P_{\text{top}}(F^{-1}, \varphi) \implies P_{\text{top}}(G, \phi_{|\Omega}) = P_{\text{top}}(F^{-1}, \varphi).$$

□

Theorem 5.1. *Given a G -invariant Borel probability μ , there exists an F^{-1} -invariant Borel probability $\bar{\mu}$ such that $h_{\bar{\mu}}(F^{-1}) \geq h_\mu(G)$.*

We divide the proof of Theorem 5.2 into five lemmas.

The first lemma constructs the measure and a σ -algebra.

Lemma 5.2. *Denote by \mathcal{A}_Ω the Borel σ -algebra in Ω . Given a G -invariant probability μ , there exists a σ -algebra \mathcal{A} and a measure $\bar{\mu} : \mathcal{A} \rightarrow [0, 1]$ depending on μ .*

Proof. Let \mathcal{A}_Ω be the Borel σ -algebra in Ω and define $\mathcal{A}_0 := \pi^{-1}(\mathcal{A}_\Omega)$. Thus, \mathcal{A}_0 is a σ -algebra on the horseshoe, whose elements are given by $A = \pi^{-1}(B)$, where B is a Borelian in Ω . Note that $\mathcal{A}_0 \subset F^{-1}(\mathcal{A}_0)$. If for each $n \in \mathbb{N}$ we define $\mathcal{A}_n := F^{-n}(\mathcal{A}_0)$, we have an increasing sequence of σ -algebras

$$\mathcal{A}_0 \subset \mathcal{A}_1 \subset \mathcal{A}_2 \subset \dots \subset \mathcal{A}_n \subset \dots$$

For each n define a probability measure $\bar{\mu}_n : \mathcal{A}_n \rightarrow [0, 1]$ by

$$\bar{\mu}_n(F^{-n}(A_0)) = \mu(\pi(A_0)) \text{ for all } A_0 \in \mathcal{A}_0.$$

Note that $\bar{\mu}_n$ depends also on $\phi_{|\Omega}$. Using the semiconjugacy it is straightforward to prove that each $\bar{\mu}_n$ is invariant under F^{-1} . Finally, we consider $\mathcal{A} := \bigcup_{n=0}^{\infty} \mathcal{A}_n$, which is a σ -algebra on Λ . Define $\bar{\mu} : \mathcal{A} \rightarrow [0, 1]$ by

$$\bar{\mu}(A) = \bar{\mu}_n(A) \text{ if } A \in \mathcal{A}_0.$$

□

It is easy to see that $\bar{\mu}$ is well-defined. The second lemma gives a criterium to show that the measure is σ -additive, which is classical in measure theory.

Lemma 5.3. *Let \mathcal{A} be a σ -algebra and $\bar{\mu} : \mathcal{A} \rightarrow [0, 1]$ such that*

$$\bar{\mu}(A_1 \cup \dots \cup A_n) = \sum_{i=1}^n \bar{\mu}(A_i),$$

for every finite family $\{A_i\}_{i=1}^n$ of disjoint sets in \mathcal{A} . If there exists a family $\mathcal{C} \subset \mathcal{A}$ such that

- \mathcal{C} is compact: if $C_1 \supset C_2 \supset \dots \supset C_n \supset \dots$ is a sequence of sets in \mathcal{C} then $\bigcap_{i=1}^{\infty} C_i \neq \emptyset$.

- \mathcal{C} has the approximation property: for all $A \in \mathcal{A}$ we have

$$\bar{\mu}(A) = \sup\{\bar{\mu}(C) \mid C \subset A, C \in \mathcal{C}\};$$

then $\bar{\mu}$ is a probability measure on \mathcal{A} .

The third lemma constructs a family satisfying the criterium.

Lemma 5.4. *For the σ -algebra \mathcal{A} in Lemma 5.2 there exists a family \mathcal{C} satisfying the criterium in Lemma 5.3 which warrants that the measure $\bar{\mu}$ in Lemma 5.2 is a probability on \mathcal{A} .*

Proof. We consider K_Ω the collection of all compact sets in Ω and we define $K_0 := \pi^{-1}(K_\Omega)$, which is a subfamily of the σ -algebra \mathcal{A}_0 . We define $\mathcal{C} : \bigcup_{n=0}^{\infty} F^{-n}(K_0)$ which is contained in \mathcal{A} and it is a compact class with the approximation property.

- Let $C_1 \supset C_2 \supset \dots \supset C_n \supset \dots$ be sets of \mathcal{C} . If for every $i \in \mathbb{N}$ there exists $D_i \subset K_\Omega$ and there exists $n_i \in \mathbb{N}$ such that $C_i = F^{-n_i}(\pi^{-1}(D_i))$. Since $\pi^{-1}(D_i)$ is a closed set in a compact space Ω , $\pi^{-1}(D_i)$ is compact. By the continuity of F^{-1} , $(C_i)_{i \in \mathbb{N}}$ is a family of nested compact sets, therefore $\bigcap_{n=1}^{\infty} C_n \neq \emptyset$.
- Let $A \in \mathcal{A}$. Using the regularity of the measure μ on Ω we have $\mu(B) = \sup\{\mu(D) \mid D \subset B, D \in K_\Omega\}$ for all $D \in \mathcal{A}_\Omega$. By the definition of \mathcal{C} , for all $C \in \mathcal{C}$ there exists $D \in K_\Omega$ and $n \in \mathbb{N}$ such that $C = F^{-n}(\pi^{-1}(D))$. Thus, $\bar{\mu}(C) = \bar{\mu}(F^{-n}(\pi^{-1}(D))) = \mu(\pi(\pi^{-1}(D))) = \mu(D)$. Then

$$\sup\{\mu(D) \mid D \subset B, D \in K_\Omega\} = \sup\{\bar{\mu}(C) \mid C \subset A, C \in \mathcal{C}\}.$$

To finish it is enough to note that for all $A \in \mathcal{A}$, there exists $B \in K_\Omega$ such that $\bar{\mu}(A) = \mu(B)$.

Then, by the criterion, we proved that $\bar{\mu}$ is a probability measure on \mathcal{A} . Since every $\bar{\mu}_n$ is invariant under F^{-1} , the measure $\bar{\mu}$ is also F^{-1} -invariant. \square

The fourth lemma shows that the measure is F^{-1} -invariant on the Borel σ -algebra.

Lemma 5.5. *The extension of the σ -algebra \mathcal{A} coincides with the Borel σ -algebra \mathcal{A}_Λ on Λ .*

Proof. Since π is continuous we have that \mathcal{A} is contained in the Borel σ -algebra. It is enough to prove that for all $X \in \Lambda$ there exists a fundamental system of neighborhoods of X contained in \mathcal{A} . Let $X \in \Lambda$ and remember that $\rho < 1$ is the contraction factor in the direction of the Z -axis. For each $n \in \mathbb{N}$, by the uniform continuity of G^n there exists $\delta_n > 0$ such that $d(z, w) < \delta_n$ implies $d(G^n(z), G^n(w)) < 1/n$. We define $B_n(X) = F^{-n}(\pi^{-1}(B(\pi(F^n(X)), \delta_n)))$. Therefore, for every $n \in \mathbb{N}$ and $X \in B_n(X)$, we have $\text{diam} B_n(X) = \rho^n + 1/n$ which goes to zero when n goes to infinity. \square

The fifth lemma shows the inequality between the entropies.

Lemma 5.6. *We have $h_{\bar{\mu}}(F^{-1}) \geq h_\mu(G)$.*

Proof. Using the well-known Brin-Katok theorem, we can compute the entropy by the formula:

$$h_\mu(G) = \lim_{\epsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{n} \log \frac{1}{\mu(B_G(X, \epsilon, n))},$$

for μ -almost every $X \in \Omega$, where $B_G(X, \epsilon, n)$ denotes the dynamical ball of G at X , radius ϵ and length n . We note that, given $\epsilon > 0$, by the continuity of π (on a compact space), there exists $0 \leq \delta \leq \epsilon$ such that

$$\pi(B(Z, \delta)) \subset B(\pi(Z), \epsilon) \text{ for every } Z \in \Lambda.$$

Using this fact it is straightforward to check that

$$B_{F^{-1}}(Y, \delta, n) \subset \pi^{-1}(B_G(X, \epsilon, n)) \text{ for every } Y \in \pi^{-1}(X).$$

By the definition of the measure $\bar{\mu}$ we have

$$\bar{\mu}(B_{F^{-1}}(Y, \delta, n)) \leq \bar{\mu}(\pi^{-1}(B_G(X, \epsilon, n))) = \mu(B_G(X, \epsilon, n)).$$

If we consider $A \subset \Omega$ a full measure set with respect to μ , then $\pi^{-1}(A) \subset \Lambda$ satisfies $\bar{\mu}(\pi^{-1}(A)) = \mu(A) = 1$. Thus,

$$h_{\bar{\mu}}(F^{-1}) = \lim_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{n} \log \frac{1}{\bar{\mu}(B_{F^{-1}}(Y, \delta, n))} \geq \lim_{\epsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{n} \log \frac{1}{\mu(B_G(X, \epsilon, n))} = h_{\mu}(G).$$

□

Proposition 5.2. *If the measure μ is an equilibrium state for the system $(G, \phi|_{\Omega})$ then $\bar{\mu}$ given by Theorem 5.2 is an equilibrium state for the system (F^{-1}, φ) .*

Proof. Given an equilibrium state μ for the system $(G, \phi|_{\Omega})$, Theorem 5.2 gives an F^{-1} -invariant Borel probability $\bar{\mu}$ such that $h_{\bar{\mu}}(F^{-1}) \geq h_{\mu}(G)$. Hence,

$$P_{\text{top}}(G) = h_{\mu}(G) + \int \phi_{\Omega} d\mu \leq h_{\bar{\mu}}(F^{-1}) + \int \varphi d\bar{\mu} \leq P_{\text{top}}(F^{-1}).$$

Once $P_{\text{top}}(G) = P_{\text{top}}(F^{-1})$ and by a change of variables we get

$$\int \phi_{\Omega} d\mu = \int \phi_{\Omega} \circ \pi d\bar{\mu} = \int \varphi d\bar{\mu},$$

then we have that $\bar{\mu}$ is an equilibrium state for the system (F^{-1}, φ) . □

Proposition 5.3. *If the measure μ is an equilibrium state for the system (F^{-1}, φ) then the measure given by $\nu = \mu \circ \pi^{-1}$ is an equilibrium state for the system $(G, \phi|_{\Omega})$.*

Proof. Let μ an equilibrium state for the system (F^{-1}, φ) and define $\nu(A) = \mu(\pi^{-1}(A))$ for all Borelian A in Ω . Again by semiconjugacy, the probability ν is G -invariant. We have that

$$P_{\text{top}}(F^{-1}, \varphi) = h_{\mu}(F^{-1}) + \int \varphi d\mu \leq h_{\nu}(G) + \int \phi_{\Omega} d\nu \leq P_{\text{top}}(G),$$

By Proposition 5.1 we have $P_{\text{top}}(G, \phi|_{\Omega}) = P_{\text{top}}(F^{-1}, \varphi)$, which implies that ν is an equilibrium state for the system $(G, \phi|_{\Omega})$. □

Proposition 5.4. *The system (F^{-1}, φ) has uniqueness of equilibrium state if, and only if, the system $(G, \phi|_{\Omega})$ does so.*

Proof. Assume that the system $(G, \phi|_{\Omega})$ has uniqueness of equilibrium state μ . Proposition 5.2 guarantees that $\bar{\mu}$ is an equilibrium state for the system (F^{-1}, φ) . Let us suppose, by contradiction, that there exists another equilibrium state $\bar{\nu}$. We will show that $\nu := \bar{\nu} \circ \pi^{-1}$ is different from the equilibrium state μ .

In fact, since $\bar{\nu} \neq \bar{\mu}$, we can find $A \in \mathcal{A}$ such that $\bar{\nu}(A) \neq \bar{\mu}(A)$. The σ -algebra \mathcal{A} , which coincides with the Borelians on Λ , was obtained as $\mathcal{A} = \bigcup_{n=0}^{\infty} F^{-n}(\mathcal{A}_0)$. Thus, there exist $A_0 \in \mathcal{A}_0$ and $n_0 \in \mathbb{N}$ such that $A = F^{-n_0}(A_0)$.

Remember also that $\mathcal{A}_0 = \pi^{-1}(\mathcal{A}_{\Omega})$ and therefore there exists $B \in \mathcal{A}_{\Omega}$ such that $A_0 = \pi^{-1}(B)$. Using that $\bar{\mu}$ is invariant under F^{-1} we have

$$\nu(B) = \bar{\nu}(\pi^{-1}(B)) = \bar{\nu}(A_0) = \bar{\nu}(F^{-n_0}(A_0)) = \bar{\nu}(A) \neq$$

$$\bar{\mu}(A) = \bar{\nu}(F^{-n_0}(A_0)) = \bar{\nu}(A_0) = \bar{\mu}(\pi^{-1}(B)) = \mu(B).$$

It means that $\nu \neq \mu$ and the first part is proven.

Now, assuming that the system (F^{-1}, φ) has uniqueness, let us suppose, by contradiction, that the system $(G, \phi|_{\Omega})$ admits two distinct equilibrium states μ and ν . Proposition 5.3 guarantees that $\bar{\mu}$ and $\bar{\nu}$ are equilibrium states for the system (F^{-1}, φ) . By uniqueness, we have $\bar{\mu} = \bar{\nu}$. Given $B \in \mathcal{A}_{\Omega}$, analogously to what is given above, we can consider the following

$$\begin{aligned} \nu(B) &= \bar{\nu}(\pi^{-1}(B)) = \bar{\nu}(A_0) = \bar{\nu}(F^{-n_0}(A_0)) = \bar{\nu}(A) = \\ \bar{\mu}(A) &= \bar{\nu}(F^{-n_0}(A_0)) = \bar{\nu}(A_0) = \bar{\mu}(\pi^{-1}(B)) = \mu(B). \end{aligned}$$

Once B is arbitrary, we conclude that $\mu = \nu$. This contradiction proves the uniqueness. The Proposition is proven. \square

Finally, we prove that potentials which depends on the z -coordinate have cohomologous which does not depend on it. We follow ideas of [31].

Proposition 5.5. ([31] Proposition 3.1) *Let $\phi : R_0 \cup R_1 \rightarrow \mathbb{R}$ be a Hölder continuous potential. There exist a Hölder continuous potential $\varphi : R_0 \cup R_1 \rightarrow \mathbb{R}$ satisfying:*

- φ is cohomologous to ϕ , that is, there exists $u : R_0 \cup R_1 \rightarrow \mathbb{R}$ such that $\varphi = \phi + u - u \circ F^{-1}$.
- φ does not depend on the z -coordinate.

Proof. They take

$$u(X) = \sum_{j=0}^{\infty} (\phi \circ F^j - \phi \circ F^j \circ \pi)(X).$$

and prove that φ like above does not depend on the z -coordinate. \square

6. HYPERBOLIC POTENTIALS VERSUS EXPANDING POTENTIALS

In the works [2] and [30] the authors prove existence of equilibrium state for non-uniformly expanding maps and the so-called hyperbolic potentials. While in [30] they also prove uniqueness with a mild condition, in [2] prove an equivalence between the so-called expanding potentials and the hyperbolic ones. It will allows us to obtain uniqueness for the horseshoe in some cases where the restriction of the potential is expanding (or hyperbolic) by using Theorem B. We recall the definitions of hyperbolic and expanding potentials as it is given in [2].

6.1. Topological pressure. We recall the definition of relative pressure for non-compact sets by means of dynamical balls. Let M be a compact metric space. Consider $f : M \rightarrow M$ and $\phi : M \rightarrow \mathbb{R}$ both continuous. Given $\delta > 0$, $n \in \mathbb{N}$ and $x \in M$, we define the *dynamical ball* $B_{\delta}(x, n)$ as

$$B_{\delta}(x, n) = \{y \in M \mid d(f^i(x), f^i(y)) < \delta, \text{ for } 0 \leq i \leq n\}.$$

Consider for each $N \in \mathbb{N}$, the set

$$\mathcal{F}_N = \{B_{\delta}(x, n) \mid x \in M \text{ and } n \geq N\}.$$

Given $\Lambda \subset M$, denote by $\mathcal{F}_N(\Lambda)$ the finite or countable families of elements in \mathcal{F}_N which cover Λ . Define for $n \in \mathbb{N}$

$$S_n \phi(x) = \phi(x) + \phi(f(x)) + \cdots + \phi(f^{n-1}(x)),$$

and

$$R_{n,\delta} \phi(x) = \sup_{y \in B_{\delta}(x,n)} S_n \phi(y).$$

Given a f -invariant set $\Lambda \subset M$, not necessarily compact, define for each $\gamma \in \mathbb{R}$

$$m_f(\phi, \Lambda, \delta, \gamma, N) = \inf_{\mathcal{U} \in \mathcal{F}_N(\Lambda)} \left\{ \sum_{B_\delta(x, n) \in \mathcal{U}} e^{-\gamma n + R_{n, \delta} \phi(x)} \right\}.$$

Define also

$$m_f(\phi, \Lambda, \delta, \gamma) = \lim_{N \rightarrow +\infty} m_f(\phi, \Lambda, \delta, \gamma, N)$$

and

$$P_f(\phi, \Lambda, \delta) = \inf\{\gamma \in \mathbb{R} \mid m_f(\phi, \Lambda, \delta, \gamma) = 0\}.$$

Finally, define the *relative pressure* of ϕ on Λ as

$$P_f(\phi, \Lambda) = \lim_{\delta \rightarrow 0} P_f(\phi, \Lambda, \delta).$$

The *topological pressure* of ϕ is, by definition, $P_f(\phi) = P_f(\phi, M)$. It satisfies

$$(13) \quad P_f(\phi) = \sup\{P_f(\phi, \Lambda), P_f(\phi, \Lambda^c)\},$$

where Λ^c denotes the complement of Λ on M . We refer the reader to [24, Theorem 11.2] for the proof of (13) and for additional properties of the pressure.

6.2. Expanding measures and hyperbolic potentials. Given $0 < \sigma < 1$ and $\varepsilon > 0$, we say that $n \in \mathbb{N}$ is a *hyperbolic time* for $x \in M$ if

- there exists a neighbourhood $V_n(x)$ of x such that f^n maps $V_n(x)$ homeomorphically onto the ball $B_\varepsilon(f^n(x))$;
- $d(f^i(y), f^i(z)) \leq \sigma^{n-i} d(f^n(y), f^n(z))$, for all $y, z \in V_n(x)$ and all $0 \leq i \leq n-1$.

We say that $x \in M$ has *positive frequency* of (σ, ε) -hyperbolic times if

$$d(x) := \limsup_{n \rightarrow \infty} \frac{1}{n} \#\{0 \leq j \leq n-1 \mid j \text{ is a } \sigma\text{-hyperbolic time for } x\} > 0,$$

and define the *expanding set*

$$H = \{x \in M \mid d(x) > 0\}.$$

We say that a Borel probability measure μ (not necessarily f -invariant) on M is *expanding* if $\mu(H) = 1$, and that a continuous function $\phi : M \rightarrow \mathbb{R}$ is a *hyperbolic potential* if the topological pressure $P_f(\phi)$ is located on H , i.e.

$$P_f(\phi, H^c) < P_f(\phi).$$

Proposition 6.1. *Let ϕ be a hyperbolic potential. If μ is an ergodic probability measure such that $h_\mu(f) + \int \phi d\mu > P_f(\phi, H^c)$, then $\mu(H) = 1$.*

Proof. Since H is an invariant set and μ is an ergodic probability measure, we have $\mu(H) = 0$ or $\mu(H) = 1$. Since the potential ϕ is hyperbolic, we get

$$h_\mu(f) + \int \phi d\mu > P_f(\phi, H^c) \geq \sup_{\nu(H^c)=1} \left\{ h_\nu(f) + \int_{H^c} \phi d\nu \right\}.$$

(For the second inequality, see [24, Theorem A2.1]) So, we cannot have $\mu(H^c) = 1$ and we obtain $\mu(H) = 1$, i.e. μ is an expanding measure. \square

6.3. Expanding potentials. Let $\phi : M \rightarrow \mathbb{R}$ be a continuous potential. We say that ϕ is an *expanding potential* if the following inequality holds

$$\sup_{\mu \in \mathcal{E}^c} \left\{ h_\mu(f) + \int \phi d\mu \right\} < \sup_{\mu \in \mathcal{E}} \left\{ h_\mu(f) + \int \phi d\mu \right\},$$

where \mathcal{E} denotes the set of all expanding measures for f .

We observe that Proposition 6.1 shows that every hyperbolic potential is an expanding potential. Since for an expanding potential ϕ we can find a sequence μ_n of expanding measures such that we have $h_{\mu_n}(f) + \int \phi d\mu_n \rightarrow P_f(\phi)$, if ϕ is also Hölder, we also can find finitely many ergodic equilibrium states for ϕ which are expanding measures. In fact, all the proof is similar from the point where Proposition 6.1 is proved. We obtain the following Theorem:

Theorem 6.1. ([2][Theorem B]) *Let $f : M \rightarrow M$ be a continuous map with an expanding measure and $\phi : M \rightarrow \mathbb{R}$ a continuous potential. Then ϕ is a hyperbolic potential if, and only if, it is an expanding potential. In particular, if ϕ is hyperbolic and Hölder with finite pressure $P_f(\phi)$, there exist finitely many equilibrium states which are expanding measures.*

With Theorem 6.1 we can use the following result for hyperbolic potentials.

Theorem 6.2. ([30][Theorem 2]) *Let $f : M \rightarrow M$ be a local homeomorphism and let $\phi : M \rightarrow \mathbb{R}$ be a hyperbolic Hölder continuous potential. Then, there exist finitely many ergodic equilibrium states associated with (f, ϕ) . In addition, the equilibrium state is unique if the pre-orbit $\{f^{-n}(x)\}_{n \geq 0}$ of every point $x \in H$ is dense in M .*

We observe that in [32] the zooming measures and continuous zooming potentials in the context of exponential contractions are what we call here expanding measures and expanding potentials. Hence, our Theorem 6.1 is a corollary of [32][Theorem 8.2.1]. We state this result in the following:

Theorem 6.3. ([32][Theorem 8.2.1]) *Let $f : M \rightarrow M$ a continuous zooming map and the contraction $(\alpha_n)_n$ satisfying $\alpha_n(r) \leq ar$ for some $a \in (0, 1)$, every $n \in \mathbb{N}$ and every $r \in [0, +\infty)$ (Lipschitz contractions, for example) and $\phi : M \rightarrow \mathbb{R}$ a continuous potential. Then ϕ is a hyperbolic potential if, and only if, it is a zooming potential. In particular, if ϕ is hyperbolic and Hölder with finite pressure $P_f(\phi)$, there exist finitely many equilibrium states which are zooming measures.*

7. PROOF OF THEOREM C

In the work [31] the authors prove that the map G , the projection of F^{-1} , is a non-uniformly expanding map and satisfy our assumption of hyperbolic times in section 6. We can now prove our Theorem C.

Proposition 7.1. *If a potential $\phi : \Lambda \rightarrow \mathbb{R}$ is projective hyperbolic, then its restriction $\varphi = \phi|_\Omega$ is an expanding potential (or hyperbolic) with respect to G .*

Proof. Once every potential with low variation is hyperbolic (or expanding), we have that the null potential is hyperbolic and then there exists a measure of maximal entropy $\mu \in \mathcal{E}$, that is, $h(G) = h_\mu(G)$. Since φ is a potential such that $\varphi(x) \geq u(x) - u(G(x))$ for some measurable function $u : \Omega \rightarrow \mathbb{R}$ and every $x \in H$ we have that $\int \phi d\eta \geq 0$ for every $\eta \in \mathcal{E}$. It implies that

$$\sup_{\nu \in \mathcal{E}^c} \left\{ h_\nu(G) + \int \phi d\nu \right\} \leq \sup_{\nu \in \mathcal{E}^c} \{h_\nu(G)\} + \sup_{\nu \in \mathcal{E}^c} \left\{ \int \phi d\nu \right\} < h_\mu(G) \leq$$

$$h_\mu(G) + \int \phi d\mu \implies \sup_{\nu \in \mathcal{E}^c} \left\{ h_\nu(G) + \int \phi d\nu \right\} < \sup_{\mu \in \mathcal{E}} \left\{ h_\mu(G) + \int \phi d\mu \right\}.$$

It shows that the potential ϕ is expanding (or hyperbolic). \square

Corollary 7.1. *If $\phi : \Lambda \rightarrow \mathbb{R}$ is a projective hyperbolic potential, then there exists a unique equilibrium state for G associated to the potential $\varphi = \phi|_\Omega$.*

Proof. By using of the Proposition 7.1 the potential φ is expanding and by Theorem 6.1 it is also a hyperbolic potential. In this way, since the map G is strongly topologically mixing and non-uniformly expanding, by [[30], Theorem 2] we have that there exists a unique equilibrium state for the pair (G, φ) . \square

Hence, by Theorem B we obtain uniqueness for the horseshoe F .

8. PROOF OF THEOREM D

Lemma 8.1. *Let $F : \Lambda \rightarrow \Lambda$ be the horseshoe, η the unique measure of maximal entropy for the non-uniformly expanding map $G : \Omega \rightarrow \Omega$ and $\phi : \Lambda \rightarrow \mathbb{R}$ a Hölder potential and $X \in \Lambda$ such that $\sup \phi|_\Omega = \phi(X) < P(\phi|_\Omega) - h_\eta(G)$. Then the restriction of $\phi|_\Omega$ is an expanding potential, that is, it is a hyperbolic potential.*

Proof. The existence and uniqueness of the measure of maximal entropy η is given by Ramos-Viana [30] because Santana [32] gives that the null potential is zooming (expanding), then hyperbolic. So, denoting by \mathcal{E} the set of expanding measures, since $\sup \phi|_\Omega = \phi(X) < P(\phi|_\Omega) - h_\eta(G)$, we have

$$\begin{aligned} \sup_{\nu \in \mathcal{E}^c} \left\{ h_\nu(G) + \int (\phi|_\Omega - P(\phi|_\Omega)) d\nu \right\} &< h_\eta(G) + \sup_{\nu \in \mathcal{E}^c} \left\{ \int (\phi|_\Omega - \phi(X) - h_\eta(G)) d\nu \right\} = \\ &h_\eta(G) + \sup_{\nu \in \mathcal{E}^c} \left\{ \int (\phi|_\Omega - \phi(X)) d\nu \right\} - h_\eta(G) \leq 0 = \\ &P(\phi|_\Omega - P(\phi|_\Omega)) = \sup_{\nu \in \mathcal{E}} \left\{ h_\nu(G) + \int (\phi|_\Omega - P(\phi|_\Omega)) d\nu \right\}. \end{aligned}$$

Hence, $\phi|_\Omega - P(\phi|_\Omega)$ is expanding and so is $\phi|_\Omega$. Then, $\phi|_\Omega$ is hyperbolic. Ramos-Viana gives a unique equilibrium state because G is topologically exact and the pre-images of points in the expanding set are dense. \square

Lemma 8.2. *Let $F : \Lambda \rightarrow \Lambda$ be the horseshoe, η the unique measure of maximal entropy for the non-uniformly expanding map $G : \Omega \rightarrow \Omega$ and a Hölder potential such that it holds that $\sup \phi|_\Omega = 0 = \phi|_\Omega(Q) < P(\phi|_\Omega)$. Then, there exists a Hölder potential $\varphi : M \rightarrow \mathbb{R}$ such that $\sup \varphi|_\Omega < P(\varphi|_\Omega) - h_\eta(G)$ and $\varphi|_\Omega$ is cohomologous to $\phi|_\Omega$.*

Proof. Once Q is a fixed point, we have that

$$\sup \phi|_\Omega = 0 = \phi(Q) = h_{\delta_Q}(G) + \int \phi|_\Omega d\delta_Q \leq P(\phi|_\Omega) \implies P(\phi|_\Omega) = h_\eta(G).$$

Define for $t \in \mathbb{R}$ and $u = -t\phi$

$$\varphi(x) = (1-t)\phi(x) + t\phi(F^{-1}(x)) = \phi(x) + (-t\phi)(x) - (-t\phi(F^{-1}(x))) = \phi(x) + u(x) - u(F^{-1}(x)).$$

We claim that there exists $t \in \mathbb{R}$ such that $\sup \varphi_{|\Omega} < P(\varphi_{|\Omega}) - h_\eta(G)$. Since it holds that $\sup \phi_{|\Omega} = 0 = \phi_{|\Omega}(Q) < P(\phi_{|\Omega}) = h_\eta(G)$, we can take $0 < t < 1 \approx s$ such that it holds that $h_\eta(G) < sP(\phi_{|\Omega})$, $s < 1 + (t-1) \sup \phi_{|\Omega}/P(\phi_{|\Omega})$ and for every $x \in \Omega$

$$t\phi(F^{-1}(x)) = t\phi(G(x)) \leq t \sup \phi_{|\Omega} \leq 0 < sP(\phi_{|\Omega}) - h_\eta(G), (1-t) \sup \phi_{|\Omega} < (1-s)P(\phi_{|\Omega}).$$

So,

$$\begin{aligned} \varphi_{|\Omega}(x) &= (1-t)\phi_{|\Omega}(x) + t\phi_{|\Omega}(G(x)) < (1-s)P(\phi_{|\Omega}) + sP(\phi_{|\Omega}) - h_\eta(G) = \\ &P(\phi_{|\Omega}) - h_\eta(G) = P(\varphi_{|\Omega}) - h_\eta(G), \end{aligned}$$

because ϕ and φ are cohomologous and have same pressures. Then $\sup \varphi_{|\Omega} < P(\varphi_{|\Omega}) - h_\eta(G)$ as we wished. \square

By Lemma 1 $\varphi_{|\Omega}$ is hyperbolic or expanding and it means that it has uniqueness of equilibrium state. Once by Lemma 2 $\phi_{|\Omega}$ is cohomologous to $\varphi_{|\Omega}$, $\phi_{|\Omega}$ also has uniqueness of equilibrium state.

Remark 8.1. *We observe that if $\phi_{|\Omega}(Q) \neq 0$ we can take the potential $\varphi = \phi - \phi_{|\Omega}(Q)$ and the result follows.*

Lemma 8.3. *We have either every Hölder potential φ such that $\sup \varphi_{|\Omega} < P(\varphi_{|\Omega})$ has uniqueness of equilibrium state or there exists a Hölder potential ϕ such that $\phi_{|\Omega}(Q) < \sup \phi_{|\Omega} < P(\phi_{|\Omega})$ and ϕ does not have uniqueness of equilibrium states.*

Proof. Let us suppose that not every Hölder potential φ like that has uniqueness of equilibrium state. We can suppose that $\varphi_{|\Omega} \geq 0$. Also, $\varphi_{|\Omega}$ does not have uniqueness. By Lemma 8.2, we have either $\sup \varphi_{|\Omega} \geq P(\varphi_{|\Omega})$ or $\varphi_{|\Omega}(Q) < \sup \varphi_{|\Omega}$. Once $\sup \varphi_{|\Omega} < P(\varphi_{|\Omega})$ by hypothesis, we then have $\varphi_{|\Omega}(Q) < \sup \varphi_{|\Omega}$

Since $\sup \varphi_{|\Omega} < P(\varphi_{|\Omega})$, denote by $\text{supp} \mu$ the support of the measure μ . Consider the following union

$$X := \bigcup_{\mu \text{ invariant} \neq \delta_Q} \text{supp} \mu.$$

Take $t < 0$ such that $(1 + td(Q, X))\varphi(Q) < \sup \varphi_{|X}$ and define

$$\phi(x) := (1 + td(x, X))\varphi(x) \implies \phi_{|\Omega}(Q) = (1 + td(Q, X))\varphi_{|\Omega}(Q) < \sup \varphi_{|X} \leq \sup \varphi_{|\Omega} < P(\varphi_{|\Omega}).$$

We used that $d(x, X) = 0$ for every $x \in X$ and $\phi_{|X} = \varphi_{|X}$. Also, we have that for $\mu \neq \delta_Q$ an invariant measure

$$\int \phi_{|\Omega} d\mu = \int_{\text{supp} \mu} \phi_{|\Omega} d\mu = \int_{\text{supp} \mu} (1 + d(x, X))\varphi_{|\Omega}(x) d\mu(x) = \int \varphi_{|\Omega} d\mu \implies P(\varphi_{|\Omega}) = P(\phi_{|\Omega}),$$

because

$$\begin{aligned} h_{\delta_Q}(G) + \int \phi_{|\Omega} d\delta_Q &= \phi_{|\Omega}(Q) < P(\varphi_{|\Omega}) = \sup_{\mu} \left\{ h_{\mu}(G) + \int \varphi_{|\Omega} d\mu \right\} = \\ &\sup_{\mu} \left\{ h_{\mu}(G) + \int \phi_{|\Omega} d\mu \right\} = P(\phi_{|\Omega}). \end{aligned}$$

Hence, $\phi_{|\Omega}(Q) < P(\varphi_{|\Omega}) = P(\phi_{|\Omega})$. Also, once $\phi_{|X} \equiv \varphi_{|X}$, we have $\phi_{|\Omega}(Q) < \sup \varphi_{|X} = \sup \phi_{|X} \leq \sup \phi_{|\Omega}$. It means that μ is an equilibrium state for ϕ and since φ has at least 3 equilibrium states, ϕ does not have uniqueness of equilibrium state. Moreover, ϕ is Hölder because φ and $d(x, X)$ are Hölder. \square

It means that among the Hölder potentials φ such that $\sup \varphi|_{\Omega} < P(\Omega)$ either all of them has uniqueness or the condition $\varphi|_{\Omega}(Q) = \sup \varphi|_{\Omega}$ is needed to have uniqueness.

9. EXAMPLES

9.1. Examples of Admissible Potentials. In this section it will be given examples of admissible potentials with respect to the horseshoe map F .

Firstly, we present some preliminary results and definitions used to get the conditions (C_1) and (C_2) .

Let n be positive integer and denote the words

$$b_n = \underbrace{0 \dots 01}_{n \text{ times}}$$

Given a sequence $w \in [1]$ and a positive integer i , we denote its i -segment by

$$[w]_i := w_0 w_1 \dots w_{i-1}.$$

If $\sigma^i(w) \in [1]$, so the i -segment of w is the concatenation of words b_n , that is,

$$1b_{n_1}b_{n_2} \dots b_{n_r}.$$

We denote the cardinality of the words b_n in the i -segment of w by $a(w, i, n)$. Note that if $w \in \Sigma_{\alpha}$, then it is a concatenation of infinite words of the type b_n .

Lemma 9.1. *Let $\tau < \alpha$ and $N = \lfloor \frac{1}{\alpha - \tau} \rfloor$ be fixed constants. Then*

$$(14) \quad 1 + \sum_{k=1}^{N-1} a(w, i, k) \geq \tau(i + 1),$$

for $w \in \Sigma_i$.

Proof. By contradiction, assume that

$$(15) \quad 1 + \sum_{k=1}^{N-1} a(w, i, k) < \tau(i + 1).$$

Note that the number of digits 1's in $[w]_i$ is given by

$$1 + \sum_{k=1}^s a(w, i, k),$$

where s is the largest number n such that b_n is in $[w]_i$, and it is possible that some terms of the sum are null. Since $\rho(w) = i$ we have

$$(16) \quad 1 + \sum_{k=1}^s a(w, i, k) > \alpha(i + 1).$$

On the other hand

$$(17) \quad 1 + \sum_{k=1}^s (k + 1)a(w, i, k) = (i + 1).$$

Then

$$a(w, i, N) + a(w, i, N + 1) + \dots + a(w, i, s) > (\alpha - \tau)(i + 1),$$

we conclude that

$$\begin{aligned} \sum_{k=N}^s (k+1)a(w, i, k) &\geq (N+1) \left(\sum_{k=N}^s a(w, i, k) \right) \\ &> (N+1)(\alpha - \tau)(i+1) \\ &> (i+1), \end{aligned}$$

which contradicts the expression (17). \square

A way to understand the Lemma 9.1 is that, if i is a α -return time of a sequence w to $[1]$, then the words b_1, \dots, b_N represent at least a part τi of $[w]_i$.

Given a point $X = (x, y, z)$ in Λ , such that $h(X) = w$, we use the follow notation

$$\Phi_{[w]_n}(y) = f_{w_{n-1}} \circ \dots \circ f_{w_0}(y),$$

and also the inverse case

$$\Phi_{[w]_n}^-(y) = f_{w_{-1}} \circ \dots \circ f_{w_{-n}}(y).$$

The next result gives us non-uniform contraction

Lemma 9.2. *[[18], Lemma 3.1] Let $w \in \Sigma_{11}^+$ be sequence with infinitely many 1's. Assume $w_0 = 1$. Let n_0, n_1, \dots be the successive positions of the symbol 1 in w . Then, there exist a sequence of positive real numbers $(\delta_j)_{j \geq 0}$ and a positive real number C such that the following holds.*

- (i) C depends only on n_0 .
- (ii) Each δ_j depends only on n_i , for $i \leq j$ and belongs to the $[0, \sigma]$.
- (iii) For every $i > 0$ and for every $y \in [0, 1]$,

$$|\Phi'_{[w]_{n_i}}(y)| \leq C_{n_0} \prod_{j=1}^{i-1} \theta_j$$

$$\text{where } \theta_j = \frac{1 - \frac{\delta_j}{\sigma}}{1 - \delta_j}.$$

Remark 9.1. *The Lemma 9.2 is obtained by using the contraction provided by the block of the form $0\dots 01$. Since the factors of the product in (iii) are all strictly less than 1, if for a block b_n it appears k times in the segment $[w]_{n_i}$, we have that*

$$|\Phi'_{[w]_{n_i}}(y)| \leq C_{n_0} \theta^k,$$

where θ depends only of b_n .

Given points X and Y in Λ , we use the follow norm

$$(18) \quad \|X - Y\| = \|X - Y\|_s + \|X - Y\|_c + \|X - Y\|_u,$$

where $\|X - Y\|_*$ is the distance with respect to direction $*$ for $* \in \{s, c, u\}$.

By the hyperbolic behavior of F on the stable and unstable directions and using the distance (18), given two points $X, Y \in \Lambda$ such that $h(X), h(Y) \in [w_{-n+1} \dots w_{n-1}]$ we have

$$(19) \quad \|X - Y\|_s \leq \lambda^n \quad \text{e} \quad \|X - Y\|_u \leq \beta^{-n}.$$

Now we are able to start the result of this section.

Proposition 9.1. *Let $c_0 \in (5/6, 1)$ be a constant and $\mathcal{Q}_{c_0} = \{X = (x, y, z) \in \Lambda ; z \leq c_0\}$. If $\phi : \Lambda \rightarrow \mathbb{R}$ is a Hölder continuous potential constant on \mathcal{Q}_{c_0} with $\sup \phi = \phi(Q)$, then there exists a non-empty open interval $I \subset \mathbb{R}$ such that for every $t \in I$ the potential $\phi_t := t\phi$ is admissible.*

Proof. First, note that given $5/6 < c_0 < 1$, we take $m = m(c_0)$ by

$$m = \max\{k ; \beta_0^k \beta_1(c_0 - 5/6) > 1/6\}.$$

So, for every $X \in \Lambda \cap R_1 - \mathcal{Q}_{c_0}$ we have $h(X) \in [1\underbrace{0\dots 0}_m 1]$. Now, consider the potential $\varphi = \phi \circ h^{-1}$ and its induced φ_ρ with respect to the inducing scheme of section 3. Given sequences w and v in Σ_{11} such that $\Pi^{-1}(w)$ and $\Pi^{-1}(v)$ are in the cylinder $[D_{i_{-n+1}}^{l_{-n+1}} \dots D_{i_0}^{l_0} \dots D_{i_{n-1}}^{l_{n-1}}]$ of spaces $S^{\mathbb{Z}}$, so w and v belong to

$$D_{i_0}^{l_0} = [\overbrace{10\dots 01}^{n_1} \dots 10\dots 01],$$

where the numbers n_1, \dots, n_r are the positions of the digit 1. Let $X_0 = h^{-1}(w)$ and $Y_0 = h^{-1}(v)$ be points in Λ and n_{r_k} the k th return of points X_0 and Y_0 to the set $\Lambda \cap (R_1 - \mathcal{Q}_{c_0})$, then we have

$$(20) \quad \begin{aligned} |\varphi_\rho(w) - \varphi_\rho(v)| &\leq \sum_{i=0}^{l_0-1} |\phi(F^i(X_0)) - \phi(F^i(Y_0))| \\ &= C \sum_{k=1}^{\bar{r}} \|F^{n_{r_k}}(X_0) - F^{n_{r_k}}(Y_0)\|^\xi, \end{aligned}$$

where \bar{r} is number of return of the points X_0 and Y_0 to the set $\Lambda \cap (R_1 - \mathcal{Q}_{c_0})$. Note that the terms of the sum (20) related to the points in \mathcal{Q}_{c_0} , are equal to zero, since ϕ is constant in this set. Moreover, by (19), it is enough to analyze the central direction. By using Lemma 9.1, we have that in each segment $[w]_{s_n}$, where $s_n = \sum_{k=1}^{n-1} l_{-k}$, the words b_1, b_2, \dots, b_N appear at least with frequency τs_n . Then, by Lemma 9.2 and Remark 9.1, we conclude

$$(21) \quad |\Phi'_{[w]_{s_n}^-}(y)| \leq C_N \theta^n,$$

for $y \in [0, 1]$ and constants $C_N > 0$ and $\theta \in (0, 1)$ that depends only on the words b_1, b_2, \dots, b_N . Therefore, let X^* and Y^* be points in Λ such that $F^{s_n}(X^*) = X_0$ and $F^{s_n}(Y^*) = Y_0$, by the Lemma 9.2 and the inequality (21) we have

$$(22) \quad \begin{aligned} \|F^{n_{r_k}}(X_0) - F^{n_{r_k}}(Y_0)\|_c &= \|F^{s_n+n_{r_k}}(X^*) - F^{s_n+n_{r_k}}(Y^*)\|_c \\ &\leq |\Phi'_{[\sigma^{n_{r_k}}(w)]_{s_n+n_{r_k}}^-}| \\ &\leq C_N \theta^n \prod_{j=1}^{r_k} \theta_{n_j} \\ &\leq C_N \theta^n \gamma^k, \end{aligned}$$

where $1 \leq k \leq \bar{r}$ and the constant $\gamma \in (0, 1)$ depends only on word b_m . Hence, by using (22) in the expression (20), we have

$$|\varphi_\rho(w) - \varphi_\rho(v)| \leq \tilde{C} \Theta^n,$$

where $\Theta = \max\{\lambda, \beta, \theta\}$ and $\tilde{C} = C.C_N.\Upsilon$, for $\Upsilon = \sum_{k=0}^{\infty} \gamma^k$. Therefore, the potential ϕ satisfies the condition (C_1) .

Consider the family of the potentials $\phi_t : \Lambda \rightarrow \mathbb{R}$ given by

$$\phi_t(X) = t\phi(X),$$

where $t \in \mathbb{R}$. By using an analogous argument we can show that ϕ_t satisfies the condition (C_1) for every $t \in \mathbb{R}$. We denote the topological pressure of ϕ_t by $P(t)$. Note that the function $t \mapsto P(t)$ is convex, so it is also a continuous function on $t \in \mathbb{R}$, since $h_{top}(F) > 0$. Consider the set

$$\Delta = \{\vartheta; t \in [0, \vartheta), P(t) > t\phi(Q)\}.$$

It is a non-empty set, since $P_{top}(0) > 0$. The supremum of Δ is given by

$$t_1 = \sup_{\substack{\mu \in \mathcal{M}_F(\Lambda) \\ \mu \neq \delta_Q}} \left\{ \frac{h_\mu(F)}{\phi(Q) - \int \phi d\mu} \right\}.$$

Since the maximal entropy measure of F is not the Dirac measure δ_Q , we have

$$t_1 > \frac{h_{top}(F)}{\phi(Q) - \int \phi d\mu_{max}}.$$

By hypothesis on the potential ϕ , its variation is given by

$$Var(\phi_t) = t\phi(Q) - t \inf \phi.$$

Then, we take

$$t_0 = \sup \left\{ t; Var(\phi_t) < \frac{h_{top}(F)}{2} \right\}.$$

Since

$$t_1 > \frac{h_{top}(F)}{\phi(Q) - \inf \phi},$$

the interval $I = (t_0, t_1)$ is non-empty. Note that $Var(\phi_t) \geq \frac{h_{top}(F)}{2}$, that is, ϕ_t does not have small variation.

Moreover, by construction, for every $t \in (t_0, t_1)$ the potential ϕ_t satisfies the condition (C_2) for $n = 1$.

Therefore, for every $t \in (t_0, t_1)$ the potential ϕ_t is an admissible potential. \square

9.2. Examples of Projective Hyperbolic Potentials. As examples of projective hyperbolic potentials, we have the null one and the set of coboundaries. We will give other examples in this section. We need to find potentials ϕ such that

(D₁) The restriction $\varphi := \phi|_\Omega$ is Hölder and there exists a measurable function $u : \Omega \rightarrow \mathbb{R}$ such that $\varphi \geq u - u \circ G$.

(D₂) If \mathcal{E} denotes the set of G -invariant expanding measures on Ω , we have that

$$\sup_{\nu \in \mathcal{E}^c} \left\{ \int \varphi d\nu \right\} < h(G) - \sup_{\nu \in \mathcal{E}^c} \{h_\nu(G)\}.$$

Example 9.1. Let u be a Hölder function such that $\sup u - \inf u > h(G)$. Let v be a Hölder function such that $v > u$ and $\sup(v - u \circ G) < h(G) - \sup_{\nu \in \mathcal{E}^c} \{h_\nu(G)\}$. It implies that

$$u - u \circ G < v - u \circ G \leq \sup(v - u \circ G) < h(G) - \sup_{\nu \in \mathcal{E}^c} \{h_\nu(G)\}.$$

By taking $\phi = v - u \circ G$, we get a potential which is projective admissible.

In order to obtain projective hyperbolic potentials ϕ with big variation, that is, such that $\sup \phi - \inf \phi > h(F)$, we use the fact that $h(F) = h(G)$ (see [31]) and Example 9.1. We take a coboundary $u - u \circ F$ with big variation and, since the Hölder functions are dense in the space of continuous functions, we can take ϕ such that $\phi \geq u - u \circ G$ and they are close enough to have the integrals close enough. Since the integrals of $u - u \circ G$ are all zero, we can take ϕ such that $\sup_{\nu \in \mathcal{E}^c} \{ \int \phi d\nu \} < h(G) - \sup_{\nu \in \mathcal{E}^c} \{ h_\nu(f) \}$. This finishes the proof.

REFERENCES

1. ARBIETO, A., MATHEUS, C., OLIVEIRA, K., *Equilibrium States for Random Non-Uniformly Expanding Maps*, **Nonlinearity**, 17, (2004), 581-593.
2. ALVES, J. F., K. OLIVEIRA, K., SANTANA, E., *Equilibrium States for Hyperbolic Potentials via Inducing Schemes*, **Nonlinearity**, 37, (2024), 095030.
3. ARBIETO, A., PRUDENTE, L., *Uniqueness of equilibrium states for some partially hyperbolic horseshoes*, **Discrete and Continuous Dynamical Systems**, 32 (2012), pp. 27-40.
4. BOWEN, R., *Entropy for group endomorphisms and homogeneous spaces*, **Trans. Amer. Math. Soc.** 153 (1971), pp. 401-414.
5. BOWEN, R. *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*, **Lecture Notes in Mathematics** , vol. 470, Springer-Verlag (1975).
6. BRUIN, H., KELLER, G., *Equilibrium states for S-unimodal maps*, **Ergodic Theory and Dynamical Systems**, 18, (1998), pp. 765-789.
7. BUZZI, J., SARIG, O., *Uniqueness of Equilibrium Measures for Countable Markov Shifts and Multidimensional Piecewise Expanding Maps*, **Ergodic Theory and Dynamical Systems**, 23, (2003), 1383-1400.
8. CASTRO, A., NASCIMENTO, T., *Statistical properties of the maximal entropy measure for partially hyperbolic attractors* , **Ergodic Theory and Dynamical Systems**,(2016), pp 1-42.
9. CLIMENHAGA, V., FISHER, T., THOMPSON, J., *Unique equilibrium states for Bonatti-Viana diffeomorphisms*, **Nonlinearity** 31(6), (2018), pp 2532-2570.
10. CRISOSTOMO, J., TAHZIBI, T., *Equilibrium states for partially hyperbolic diffeomorphisms with hyperbolic linear part*, **Nonlinearity**, 32(2), (2019).
11. DÍAZ, L., FISHER, T., *Symbolic extensions for partially hyperbolic diffeomorphisms*, **Discrete and Continuous Dynamical Systems**, 29 (2011), pp. 1419-1441.
12. DÍAZ, L., HORITA, V., RIOS, I., SAMBARINO, M., *Destroying horseshoes via heterodimensional cycles: generating bifurcations inside homoclinic classes*, **Ergodic Theory and Dynamical Systems**, 29, (2009), pp. 433-474.
13. DENKER, M., URBANSKI, M., *Ergodic Theory of Equilibrium States for Rational Maps*, **Nonlinearity**, 4, (1991), 103-134.
14. GODIN, *On the central limit theorem for stationary processes*, **Akademi Nauk SSSR**, 188:4 (1969), pp. 739-741.
15. IOMMI, G., TODD, M., *Thermodynamic Formalism for Interval Maps: Inducing Schemes*, **Dynamical Systems**, 28, 3, (2013), pp. 354-380.
16. IOMMI, G., TODD, M., *Natural Equilibrium States for Multimodal Maps*, **Communications in Mathematical Physics**, 300, (2010), 65-94.
17. LEDRAPPIER, F., WALTERS, P. *A relativised variational principle for continuous transformations*, **J. Lond. Math. Soc.** (2) 16 (1977), pp. 568-579.
18. LEPLAIDEUR, R., OLIVEIRA, K., RIOS, I., *Equilibrium States for partially hyperbolic horseshoes*, **Ergodic Theory and Dynamical Systems**, 31 (2011), pp 179-195.
19. LI, H., RIVERA-LETELIER, J. *Equilibrium States of Weakly Hyperbolic One-Dimensional Maps for Hölder Potentials*, **Commun. Math. Phys.**, 31 (2011), pp 328-397.
20. LI, H., RIVERA-LETELIER, J. *Equilibrium states of interval maps for hyperbolic potentials*, **Nonlinearity**, 27 (2014).
21. OLIVEIRA, K., VIANA M., *Foundations of ergodic theory (Cambridge Studies in Advanced Mathematics, 151)* , Cambridge University Press, Cambridge, 2016.

22. OLIVEIRA, K., *Equilibrium States for Non-Uniformly Expanding Maps*, **Ergodic Theory and Dynamical Systems**, 23, 06, (2003), 1891-1905.
23. OLIVEIRA, K., VIANA M., *Thermodynamical formalism for robust classes of potentials and nonuniformly hyperbolic maps*, **Ergodic Theory and Dynamical Systems**, 28 (2008), pp 501-533.
24. PESIN, YA., *Dimension Theory in Dynamical Systems: Contemporary Views and Applications*, The University of Chicago Press, (1997).
25. PESIN, Y., SENTI, S., *Equilibrium measures for maps with inducing schemes*, **J. Mod. Dyn.**, 2(3) (2008), pp 397-430.
26. PESIN, Y., SENTI, S., ZHANG, K., *Lifting measures to inducing schemes*, **Ergodic Theory and Dynamical Systems**, 28 (2008), pp 553-574.
27. PESIN, Y., SENTI, S., ZHANG, K., *Thermodynamics of towers of hyperbolic type* **Trans. Amer. Math. Soc.** 368 (2016), pp 8519-8552.
28. PINHEIRO, V., VARANDAS, P., *Thermodynamic Formalism for Expanding Measures*, preprint, arXiv:2202.05019v2
29. RAMOS, V., SIQUEIRA, J., *On Equilibrium States for Partially Hyperbolic Horseshoes: Uniqueness and Statistical Properties*, **Bulletin of the Brazilian Mathematical Society**, New Series, (2017), pp 1-29.
30. RAMOS, V. , VIANA, M., *Equilibrium states for hyperbolic potentials*, **Nonlinearity**, 30 (2017), pp 825.
31. RIOS, I., SIQUEIRA, J., *On equilibrium state for partially hyperbolic horseshoes*, **Ergod. Theor. Dynam. Syst.** 38, (2018), pp 301-35.
32. SANTANA. E., *Equilibrium States for Open Zooming Systems*, preprint arXiv:2010.08143.
33. SARIG, O., *Thermodynamic formalism for countable Markov shifts*, **Ergodic Theory Dynam. Systems**, 19 (1999), pp 1565-1593.
34. SARIG, O., *Characterization of the existence of Gibbs measure for countable Markov shift*, **Proc of AMS**, 131:6 (2003), pp 1751-1758.
35. SARIG, O., *Thermodynamic formalism for countable Markov shifts*, **Proc. of Symposia in Pure Math**, 89 (2015), pp 81-117 1751-1758.
36. VARANDAS, P., VIANA, M. *Existence, uniqueness and stability of equilibrium states for non-uniformly expanding maps* **Annales Inst. Henri Poincaré - Analyse Non-Linéaire** 27 (2009), 555-593.
37. YOUNG, L-S., *Statistical properties of dynamical systems with some hyperbolicity*, **Ann. of Math.** (2), 147(3) (1998), pp 585-650.
38. ZWEIMÜLLER, R., *Invariant measures for general(ized) induced transformations*, **Proc. Amer.Math.Soc.** , 133 (2005), pp 2283-2295(electronic).

KRERLEY OLIVEIRA, INSTITUTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE ALAGOAS, 57072-090 MACEIÓ, BRAZIL

Email address: krerley@gmail.com

MARLON OLIVEIRA, DEPARTAMENTO DE MATEMÁTICA E INFORMÁTICA, UNIVERSIDADE ESTADUAL DO MARANHÃO, 65055-310 SÃO LUÍS, BRAZIL

Email address: marlon.cs.oliveira@gmail.com

EDUARDO SANTANA, UNIVERSIDADE FEDERAL DE ALAGOAS, 57200-000 PENEDO, BRAZIL

Email address: jemsmath@gmail.com