Model Consistency of Iterative Regularization for Low-Complexity Regularization

Jie Gao^{*} Cesare Molinari[†] Silvia Villa[†] Jingwei Liang^{*}

Abstract. Regularization is a core component of modern inverse problems as it allows to establish well-posedness to the solution of interests. Popular regularization approaches include variational regularization and iterative regularization. The former one can be tackled by solving a variational optimization problem, which is the sum of a regularization term and a data-fidelity term balanced by a proper weight, while the latter one chooses a proper stopping time to avoid overfitting to the noise. In the study of regularization, an important topic is the relation between the solution obtained by regularization and the original ground truth. When the ground truth has low-complexity structure which is encoded as the "model", a sensitivity property shows that the solution obtained from proper regularization that promotes the same structure is robust to small perturbations, this is called "model consistency". For variational regularization, the existence of model consistency is an open problem. In this paper, based on a recent development of partial smoothness which is also considered in [1], we show that if the noise level is sufficiently small and a proper stopping time is chosen, the solution by iterative regularization also achieves model consistency and more exhibit local linear convergence behavior. Numerical simulations are provided to verify our theoretical findings.

Key words. linear inverse problems, iterative regularization, dual gradient descent, partial smoothness, model consistency, local linear convergence.

AMS subject classifications. $47A52 \cdot 49J52 \cdot 60H50 \cdot 65K10 \cdot 90C31$

1 Introduction

Linear inverse problems are widely encountered in many fields through science and engineering, such as signal processing, compressed sensing, (medical) image processing, remote sensing, etc. In (linear) inverse problems, the target of interests, say w^{\dagger} , is not directly available, and one can only observe it indirectly which eventually can be modelled by a linear operator. In the finite dimensional setting, let $X \in \mathbb{R}^{n \times p}$ represent the linear mapping and $y^{\dagger} \in \mathbb{R}^{n}$ the observation without noise perturbation, then

$$y^{\dagger} = Xw^{\dagger},$$

Retrieving y^{\dagger} is called the *forward problem*, due to the imperfect of the observation procedure, the observation very often is degraded by noise which leads to the following form

$$y_{\delta} = Xw^{\dagger} + \varepsilon, \tag{1.1}$$

where ε is additive white Gaussian noise with noise level δ .

Inverse problems mean recovering or approximating w^{\dagger} from the observation data. In the *noiseless* case, the problem can be described as

find
$$w \in \mathbb{R}^p$$
 such that $Xw = y^{\dagger}$. (1.2)

^{*}School of Mathematical Sciences and Institute of Natural Sciences, Shanghai Jiao Tong University, Shanghai, China. E-mail: {zjgaojie, jingwei.liang}@sjtu.edu.cn

[†]MaLGa, DIMA, Dipartimento di Eccellenza 2023-2027, Università degli Studi di Genova, Genoa, Italy. E-mail: cecio.molinari@gmail.com; silvia.villa@unige.it

However, the above problem in general is ill-posed mostly due to the bad condition or degeneracy of the forward operator X. The problem becomes even more difficult if the observation is noisy.

1.1 Regularization

However, the ground truth often poses certain low-complexity prior property, also called the "model", which can be captured explicitly by some functions R(w) named regularization term or regularizer. Popular priors include sparsity (ℓ_1 -norm [2]), group sparsity ($\ell_{1,2}$ -norm [3]), gradient sparsity (total variation [4]) and low rank (nuclear norm [5]), we also refer to [6] for more discussion on regularization. Combine with regularization, problem (1.2) turns into

$$\min_{w \in \mathbb{P}^p} R(w) \quad \text{such that} \quad Xw = y^{\dagger}. \tag{P}$$

More generally, one may consider a data-fit function $\ell : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R} \cup \{+\infty\}$,

$$\min_{w \in \mathbb{R}^p} R(w) \quad \text{such that} \quad w \in \arg\min\left\{\ell(Xw, y^{\dagger})\right\}. \tag{P_0}$$

Problem (P_0) recovers (P) when ℓ is the indicator function of the set $\{w \in \mathbb{R}^p \mid Xw = y^{\dagger}\}$. Similar to (P_0) , in the noisy setting, one can consider the following new constraint

$$\min_{w \in \mathbb{R}^p} R(w) \quad \text{such that} \quad \ell(Xw, y_{\delta}) \le \delta, \qquad \qquad (\widehat{P}_{\delta})$$

where δ is the noise strength. Problem (\widehat{P}_{δ}) is more difficult to tackle as the constraint requires knowing/estimating δ , a simpler approach is putting the constraint into the objective with a weight $\lambda > 0$, this leads to the following variational regularization model

$$\min_{w \in \mathbb{R}^p} \left\{ p_{\lambda,\delta}(w) := R(w) + \frac{1}{\lambda} \ell(Xw, y_{\delta}) \right\}.$$
 (P_{\lambda,\delta})

The optimal choice λ_{δ} can be determined through validation criterion such as discrepancy principles [7], cross-validation [8] and SURE [9, 10]. Recently, data-driven approaches have also been designed to determine the proper regularization parameter [11]. The above formulations are referred as *variational regularization*, once the formulation is chosen, proper numerical scheme will be chosen and ran until convergence.

In parallel, another popular regularization approach is called *iterative regularization*, which is realized by properly choosing the stopping time. More precisely, for the noisy observation y_{δ} , instead of considering $(P_{\lambda,\delta})$, one can directly consider (P_0) by replacing y^{\dagger} with y_{δ} which leads to the following problem

$$\min_{w \in \mathbb{R}^p} R(w) \quad \text{such that} \quad w \in \arg\min\left\{\ell(Xw, y_{\delta})\right\}. \tag{P_{δ}}$$

Given that the constraint is nonempty, one can solve the problem with certain numerical scheme that generates a sequence $\{w_{\delta}^{(k)}\}_{k\in\mathbb{N}}$. Apparently, as the constraint contains noise, the limiting point of $\{w_{\delta}^{(k)}\}_{k\in\mathbb{N}}$ overfits to the noise, hence a proper stopping time, denoted as k_{δ} , should be chosen to avoid the overfitting. Therefore, in iterative regularization, stopping time is used to control the regularization.

In practice, the selection of the stopping time in iterative regularization is based on validation criteria similar to those used for variational regularization. However, notable differences still exist between them. On the one side, the choice of λ_{δ} in the variational regularization approach is generally computationally expensive while k_{δ} controls at the same time the accuracy of the solution and the computational cost. For large scale problems, iterative regularization performs more efficiently because it avoids solving various optimization problems. On the other hand, the conditions for obtaining convergence and convergence rate results are more complex than the ones needed in the analysis of variational regularization. Because iterative regularization problem.

1.2 Model consistency

The premise of regularization is that the ground truth w^{\dagger} has low-complexity structure, which is also called the "model" of it. Such a structure can often be captured by certain smooth manifold $\mathcal{M}_{w^{\dagger}}$, that is $w^{\dagger} \in \mathcal{M}_{w^{\dagger}}$. This includes the popular low-complexity structures, such as (group) sparsity, low-rank and analysis sparsity (e.g. piecewise constant). In the same time, the outcome of the regularization approaches above enforces the solution to have similar low-complexity structure. Denote the outcome of the variational regularization or iterative regularization as $w_{\delta}^{(\lambda_{\delta}/k_{\delta})}$. Then, a natural question to ask is whether $w_{\delta}^{(\lambda_{\delta}/k_{\delta})}$ has the same structure as w^{\dagger} , or if $w_{\delta}^{(\lambda_{\delta}/k_{\delta})} \in \mathcal{M}_{w^{\dagger}}$ also holds? This is referred as "model consistency".

Model consistency reflects the robustness of the problem to perturbations. For linear inverse problems and variational regularization, model consistency property is studied in [1, 12, 13], where it is shown that $\ell(Xw, y_{\delta}) = \frac{1}{2} ||Xw - y_{\delta}||^2$ takes the least-square form, then if the noise in y_{δ} is small enough, with properly chosen regularization parameter λ , the solution of $(P_{\lambda,\delta})$ has model consistency property. However, the model consistency of iterative regularization is not considered in the literature.

1.3 Related work

Iterative regularization The study of iterative regularization properties of gradient descent, or the Landweber iteration, dates back to the 1950s. The classical result shows that gradient descent applied to least squares and initialized at zero converges to the minimal norm solution and early stopping leads to Tikhonov regularization [14, Chapter 6]. Results in the stochastic setting is also considered [15]. For strongly convex regularization functions, iterative regularization of dual gradient descent, linearized Bregman iterations and mirror descent are studied in [16, 17, 18], with corresponding stability and convergence estimates available. For merely convex regularization, Bregman operator splitting, linearized/preconditioned ADMM, and primal-dual method and its data-driven version are studied in [19, 20, 21].

If other data fit functions, rather than the least-square, are used, the above approach neglects them. One approach is to employ a diagonal strategy [16, 22], which combines an optimization algorithm with a sequence of approximations λ_k of the original problem that changes at each iteration. For only convex regularization, a technique known as exact regularization [23, 24] exists. This involves solving

$$\min_{w \in \mathbb{R}^p} \left\{ R_{\alpha}(w) := R(w) + \frac{\alpha}{2} \|w\|^2 \right\} \text{ subject to } Xw = y^{\dagger}, \tag{1.3}$$

where there exists small enough $\alpha > 0$ such that the minimizer of the new problem coincides with the minimizer of the original problem (*P*). Therefore, one can consider strongly convex R_{α} by adding a quadratic term.

Model consistency Using the tool of partial smoothness [25, 26], it can be proven that in variational regularization the model can be identified when δ is sufficiently small and proper λ_{δ} is chosen [1, 12, 13], which implies the model consistency property. A related problem is the local linear convergence analysis, see for instance [27, 28, 29, 30, 31] and the references therein for results on first-order optimization algorithms.

1.4 Contribution of this work

In this paper, motivated by a recent development of partial smoothness [32], we study the model consistency property of iterative regularization. Under the assumption that the low-complexity regularization function is partly smooth and strongly convex, we show that if the problem (P_{δ}) solved by dual gradient descent (DGD) method as studied in [16], we are able to show the following results:

- We show that if the noise level is sufficiently small and proper early stopping time is chosen, the obtained point from iterative regularization has model consistency property; See Theorem 4.2 (i). Moreover, we also show that with a fixed sufficiently small noise level, there exists an interval of stopping time such that model consistency occurs; See Theorem 4.2 (ii). We also give a similar result for the accelerated dual gradient descent.
- We discuss that the optimization method can be viewed as a special setting of continuous dynamic system and the model consistency property remains.
- With a given small enough noise level such that model consistency occurs, we further prove local linear convergence if the underlying manifold $\mathcal{M}_{w^{\dagger}}$ is linear.

Finally, we present numerical results on ℓ_1 -norm, one dimensional total variation (1d-TV), $\ell_{1,2}$ -norm, and nuclear norm, both model consistency and local linear convergence are verified.

Paper organization The remainder of the paper is organized as follows: Section 2 provides some mathematical background, including basic concepts and assertions from convex analysis and the definition of partial smoothness. We present the details of the specific problem and the dual gradient descent (DGD) and accelerated dual gradient descent (ADGD) in Section 3. A comprehensive analysis of the main theoretical results of model consistency of iterative regularization is then performed in Section 4. Section 5 provides the results for the local linear convergence rate. Finally, Section 6 presents numerical examples that demonstrate the model consistency and local linear convergence.

2 Mathematical background

In this section, we collect some necessary background materials, which are frequently used in our subsequent theoretical analysis.

2.1 Notations and definitions

We refer to [33, 34] for more details of the contents below. Let \mathbb{R}^p be a Euclidean space with inner product $\langle \cdot, \cdot \rangle$ and norms $\|\cdot\|$. Id denotes the identity matrix. Given $w \in \mathbb{R}^p, \epsilon > 0$, let $\mathcal{B}(w, \epsilon)$ be the open ball of center w and radius ϵ .

Let S be a non-empty closed convex set of \mathbb{R}^p , denote $int(\cdot)$, $bdy(\cdot)$ the interior and boundary of a set, respectively. The relative interior of S is defined by

$$\operatorname{ri}(S) = \{ w \in S \mid \mathcal{B}(w, \epsilon) \cap \operatorname{aff} S \subseteq S \text{ for some } \epsilon > 0 \}.$$

Given a point $w \in \mathbb{R}^p$, the distance of w to S and the projection of w to S are defined by

dist
$$(w, S) = \inf_{w' \in S} ||w - w'||$$
 and $P_S(w) = \arg \min_{w' \in S} ||w - w'||$.

We denote by $\Gamma_0(\mathbb{R}^p)$ the set of proper, convex, and lower semicontinuous functions from \mathbb{R}^p to $\mathbb{R} \cup \{+\infty\}$. We say R is σ -strongly convex if $R - \frac{\sigma}{2} \|\cdot\|^2$ is convex and R is L-smooth if it is differentiable and ∇R is L-Lipschitz continuous. The subdifferential of $R \in \Gamma_0(\mathbb{R}^p)$ at point w is defined as

$$\partial R(w) = \left\{ u \in \mathbb{R}^p \mid R(w') - R(w) - \langle u, w' - w \rangle \ge 0, \forall w' \in \mathbb{R}^p \right\}.$$

Any element in $\partial R(w)$ is called a subgradient of R at w. The conjugate function R^* of $R \in \Gamma_0(\mathbb{R}^p)$ is defined by

$$R^*(u) = \sup_{w \in \mathbb{R}^p} \left\{ \langle u, w \rangle - R(w) \right\}.$$

If $R \in \Gamma_0(\mathbb{R}^p)$, then $R^* \in \Gamma_0(\mathbb{R}^p)$. Moreover, if $R \in \Gamma_0(\mathbb{R}^p)$ is σ -strongly convex, $\sigma > 0$, the conjugate function R^* is $\frac{1}{\sigma}$ -smooth. Given $\gamma > 0$, the proximity operator of R is defined by

$$\operatorname{prox}_{\gamma R}(w) = \operatorname{arg\,min}_{w' \in \mathbb{R}^p} \Big\{ R(w') + \frac{1}{2\gamma} \|w' - w\|^2 \Big\},$$

and is firmly nonexpansive.

2.2 Partial smoothness

Given $S \subseteq \mathbb{R}^p$ a closed convex set, the smallest linear subspace of \mathbb{R}^p that contains S is denoted by span(S). Let $\mathcal{M} \subset \mathbb{R}^p$ be a C^2 -manifold around $w \in \mathbb{R}^p$, denote $\mathcal{T}_{\mathcal{M}}(w)$ the tangent space of \mathcal{M} at w. For any vector $w \in \mathbb{R}^p$ with $\partial R(w) \neq \emptyset$, denote

$$T_w = \big(\operatorname{span}(\partial R(w))\big)^{\perp}.$$

The following definition of partial smoothness is adapted from [25] to the convex scenario.

Definition 2.1 (Partly smooth functions). Let $R \in \Gamma_0(\mathbb{R}^p)$ and $w^{\dagger} \in \mathbb{R}^p$ such that $\partial R(w^{\dagger}) \neq \emptyset$. *R* is said to be partly smooth at w^{\dagger} relative to a set $\mathcal{M}_{w^{\dagger}} \subset \mathbb{R}^p$ if

- (Smoothness) $\mathcal{M}_{w^{\dagger}}$ is a manifold about w^{\dagger} and $R|_{\mathcal{M}_{w^{\dagger}}}$ is a C^2 function near w^{\dagger} ;
- (Sharpness) The tangent space $\mathcal{T}_{\mathcal{M}_{w^{\dagger}}}(w^{\dagger})$ is the model tangent subspace $T_{w^{\dagger}}$;
- (Continuity) ∂R restricted to $\mathcal{M}_{w^{\dagger}}$ is continuous at w^{\dagger} .

Remark 2.2. If *R* is non-convex, proper regularity is needed [25]. When *R* is partly smooth, $R_{\alpha} = R(w) + \frac{\alpha}{2} ||w||^2, \alpha > 0$ is also partly smooth relative to the same manifold, according to the smooth perturbation rule of partial smoothness [25].

In Table 1 below, we summarize several popular examples of partly smooth functions that are widely used in inverse problems.

Table 1: Examples of partly smooth function: $\operatorname{supp}(w) := \{i : w_i \neq 0\}; D_{\text{DIF}} \text{ stands for the finite differences operator; for } \ell_{1,2}\text{-norm}, w_{\mathcal{I}} \text{ is the restriction of } w \text{ to the entries indexed in } \mathcal{I}, \cup_{i=1}^{G} \mathcal{I}_i = \{1, \cdots, p\}, \operatorname{supp}_{\mathcal{I}}(w) := \{i : w_{\mathcal{I}_i} \neq 0\}.$

Function	Expression	Partial smooth manifold
ℓ_1 -norm	$\sum_{i=1}^{p} w_i $	$\mathcal{M}_{w^{\dagger}} = T_{w^{\dagger}} = \{ w \in \mathbb{R}^p \mid \mathrm{supp}(w) \subset \mathrm{supp}(w^{\dagger}) \}$
TV semi-norm	$\ D_{\text{DIF}}w\ _1$	$\mathcal{M}_{w^{\dagger}} = T_{w^{\dagger}} = \{ w \in \mathbb{R}^p \mid \operatorname{supp}(D_{\operatorname{DIF}}w) \subset \operatorname{supp}(D_{\operatorname{DIF}}w^{\dagger}) \}$
$\ell_{1,2}$ -norm	$\sum_{i=1}^m \ w_{\mathcal{I}_i}\ $	$\mathcal{M}_{w^{\dagger}} = T_{w^{\dagger}} = \{ w \in \mathbb{R}^p \mid \operatorname{supp}_{\mathcal{I}}(w) \subset \operatorname{supp}_{\mathcal{I}}(w^{\dagger}) \}$
Nuclear norm	$\sum_{i=1}^r \sigma_i(w)$	$\mathcal{M}_{w^{\dagger}} = \{ w \in \mathbb{R}^{n \times p} \mid \operatorname{rank}(w) = \operatorname{rank}(w^{\dagger}) \}$

Remark: The first three partly smooth functions are with linear manifold, while for nuclear norm is not. Moreover, the first two are polyhedra functions, the subdifferential is locally constant around w^{\dagger} along $w^{\dagger} + T_{w^{\dagger}}$.

A crucial property of partial smoothness is the identifiability of the smooth manifold, which means if there exists a sequence $w^{(k)}$ and $z^{(k)} \in \partial R(w^{(k)})$ and $\operatorname{dist}(z^{\dagger}, \partial R(w^{(k)})) \to 0$ for $z^{\dagger} \in \operatorname{ri}(\partial R(w^{\dagger}))$, then $w^{(k)} \in \mathcal{M}_{w^{\dagger}}$ for all large k. This property is called "finite identification" [35]. Below, we provide a recent generalization of this result in [32] which does not require the convergence of the sequence.

Lemma 2.3 ([32, Proposition 4.1, Theorem 4.3]). Let function $R \in \Gamma_0(\mathbb{R}^p)$ be a partly smooth at w^{\dagger} relative to a C^2 -manifold $\mathcal{M}_{w^{\dagger}} \subset \mathbb{R}^p$. For small enough $\epsilon > 0$, define the local union as

$$\mathcal{U}_{\epsilon} = \bigcup_{w \in \mathcal{M}_{w^{\dagger}} \cap \mathcal{B}(w^{\dagger}, \epsilon)} (w + \partial R(w)).$$

Then there holds $\operatorname{span}(\mathcal{U}) = \mathbb{R}^p$, and

1. If $z^{\dagger} \in \operatorname{ri}(\partial R(w^{\dagger}))$, then $w^{\dagger} + z^{\dagger} \in \operatorname{int}(\mathcal{U}_{\epsilon})$.

2. If there is a sequence $(\{w^{(k)}, z^{(k)})\}_{k \in \mathbb{N}}$ such that $z^{(k)} \in \partial R(w^{(k)})$ and

$$\limsup_{k \to \infty} \|(w^{(k)} + z^{(k)}) - (w^{\dagger} + z^{\dagger})\| < \operatorname{dist}(w^{\dagger} + z^{\dagger}, \operatorname{bdy}(\mathcal{U}_{\epsilon})),$$

then fore all k large enough there holds $w^{(k)} \in \mathcal{M}_{w^{\dagger}}$.

3 Iterative Regularization

In this part, we briefly recall the iteration regularization of dual gradient descent method and accelerated dual gradient descent method studied in [16, 36]. For problem (P_{δ}) , suppose y_{δ} belongs to the range of X, then we can specify the constraint as $Xw = y_{\delta}$. Regarding the regularization, strongly convex functions is considered as in (1.3), leading to the following optimization problem

$$\min_{w \in \mathbb{R}^p} R_{\alpha}(w) \quad \text{subject to} \quad Xw = y_{\delta}. \tag{$\overline{P_{\delta}}$}$$

Dual problem By the Fenchel-Rockafellar duality, the dual problem of $(\overline{P_{\delta}})$ is

$$\min_{v \in \mathbb{R}^n} \Big\{ \phi(v) := R^*_{\alpha}(-X^\top v) + \langle y_{\delta}, v \rangle \Big\}, \qquad (\overline{D_{\delta}})$$

where $R^*_{\alpha}(-X^{\top}v)$ is the conjugate of R_{α} and reads

$$R_{\alpha}^{*}(\cdot) = \frac{1}{2\alpha} \|\cdot\|^{2} - \inf_{w} \Big\{ R(w) + \frac{\alpha}{2} \|w - \cdot/\alpha\|^{2} \Big\}.$$

Note that $\inf_{w} \{R(w) + \frac{\alpha}{2} ||w - \frac{\alpha}{2}||w| + \frac{\alpha}{2} ||w| + \frac{\alpha}{2} ||w|$

$$\nabla R^*_{\alpha}(-X^{\top}v) = \frac{1}{\alpha}(-X^{\top}v) - \frac{1}{\alpha}\Big(\alpha\big(\frac{1}{\alpha}(-X^{\top}v) - \operatorname{prox}_{\alpha^{-1}R}(\alpha^{-1}(-X^{\top}v))\big)\Big)$$
$$= \operatorname{prox}_{\alpha^{-1}R}\big(-\alpha^{-1}X^{\top}v\big),$$

Moreover, we have $\nabla \phi(v) = -X \operatorname{prox}_{\alpha^{-1}R} \left(-\alpha^{-1} X^{\top} v \right) + y_{\delta}$ is $\frac{\|X\|^2}{\alpha}$ -Lipschitz continuous. Hence we can apply gradient descent method to solve $(\overline{D_{\delta}})$, as considered in [16, 36].

Dual gradient descent The dual gradient descent (DGD) method for solving $(\overline{P_{\delta}})$ is described below.

Algorithm 1 Dual gradient descent (DGD)

The iterate $z_{\delta}^{(k)}$ is a subgradient of R_{α} at $w_{\delta}^{(k)}$, as from the optimality condition of proximal operator we have

$$\frac{1}{\alpha} z_{\delta}^{(k)} - w_{\delta}^{(k)} \in \frac{1}{\alpha} \partial R(w_{\delta}^{(k)}) \quad \Longleftrightarrow \quad z_{\delta}^{(k)} \in \partial R_{\alpha}(w_{\delta}^{(k)}).$$

The iteration can be written only in terms of $v_{\delta}^{(k)}$ as

$$v_{\delta}^{(k+1)} = v_{\delta}^{(k)} - \gamma \nabla \phi(v_{\delta}^{(k)}), \qquad (3.1)$$

which is a gradient descent step for the dual problem $(\overline{D_{\delta}})$. According to [34, 37], we have the following convergence result.

Lemma 3.1. For the DGD Algorithm 1, there exists an $v^{\dagger} \in \arg\min(\phi)$ such that $v_{\delta}^{(k)} \to v_{\delta}^{\dagger}$ as $k \to \infty$. Moreover, there exists a constant e > 0 such that

$$\|v_{\delta}^{(k)} - v_{\delta}^{(k-1)}\| \le \frac{e}{\sqrt{k}}.$$

In iteration regularization, we are interested in the relation between the early stopping point of $w_{\delta}^{(k)}$ and the ground truth w^{\dagger} . We impose the following assumption and notion

• The ground truth w^{\dagger} is the unique solution of $(\overline{P_{\delta}})$ for $\delta = 0$.

• The early stopping time is denoted as k_{δ} , and the corresponding point is $w_{\delta}^{(k_{\delta})}$.

From [16, 36], we have the following result.

Lemma 3.2 ([36, Theorem 4.1]). Let $\delta \in]0, 1[$, and consider the DGD Algorithm 1. Suppose there exists $\bar{v} \in \mathbb{R}^n$ such that $-X^{\top}\bar{v} \in \partial R_{\alpha}(w^{\dagger})$. Set $a = 2\|X\|^{-1}$ and $b = \|X\|\|v^{\dagger}\|\alpha^{-1}$. Then there exists $k_{\delta} \in \{\lfloor c\delta^{-1} \rfloor, \cdots, 2\lfloor c\delta^{-1} \rfloor\}$ and some c > 0 such that

$$\|w_{\delta}^{(k_{\delta})} - w^{\dagger}\| \le (a(c^{1/2} + 1) + bc^{-1/2})\delta^{1/2},$$

Denote $c_{\delta} > 0$ such that $k_{\delta} = c_{\delta} \delta^{-1}$.

The above stability result implies that the early stopping point $w_{\delta}^{(k_{\delta})}$ converges to w^{\dagger} at the rate of $\delta^{1/2}$. Moreover, we have $k_{\delta} \to \infty$ as $\delta \to 0$.

In [16, 36], they also consider an accelerate version of DGD, denoted as ADGD, which is applying the Nesterov's accelerated gradient method [38] to DGD; see the algorithm below.

Algorithm 2 Accelerated dual gradient descent (ADGD)

The above iteration is slightly different from the scheme considered in [36] in terms of the inertial parameter $\frac{k-1}{k+\theta}$, while in [36], they use $\frac{t_k-1}{t_k}$, $t_k = \frac{1+\sqrt{1+4t_{k-1}^2}}{2}$. The reason of choosing this new choice is that it not only maintains the $O(1/k^2)$ convergence rate on the dual objective, but also allows to prove convergence of the sequence. For the original choice, one can also consider the modified accelerated scheme in [39].

Lemma 3.3. For the ADGD Algorithm 2, let $\theta > 2$, then there exists an $v^{\dagger} \in \arg\min(\phi)$ such that $u_{\delta}^{(k)} \to v_{\delta}^{\dagger}$ as $k \to \infty$. Moreover, there holds

$$\sum_{k=1}^{\infty} k \| u_{\delta}^{(k)} - u_{\delta}^{(k-1)} \|^2 < +\infty.$$

The above result implies that $\|u_{\delta}^{(k)} - u_{\delta}^{(k-1)}\| = o(1/k)$. Different from DGD, ADGD is not a monotone scheme in the sense $\|u_{\delta}^{(k)} - v^{\dagger}\| \leq \|u_{\delta}^{(k-1)} - v^{\dagger}\|$ no longer holds. Also note that the result holds for any $\theta > 2$.

Lemma 3.4 ([36, Theorem 4.2]). Let $\delta \in]0,1[$. Let $\{w_{\delta}^{(k)}\}$ be the sequence generated by Algorithm 2. Suppose there exists $\bar{v} \in \mathbb{R}^n$ such that $-X^{\top}\bar{v} \in \partial R_{\alpha}(w^{\dagger})$. Set $a = 4||X||^{-1}$ and $b = 2||X|| ||v^{\dagger}||\alpha^{-1}$, where v^{\dagger} is a solution of the dual problem. Then

$$\|w_{\delta}^{(k)} - w^{\dagger}\| \le ak\delta + bk^{-1}.$$

In particular, choosing $k_{\delta} = \lceil c\delta^{-1/2} \rceil$, then for some c > 0 there holds $\|w_{\delta}^{(k_{\delta})} - w^{\dagger}\| \leq (a(c+1) + bc^{-1})\delta^{1/2}.$

4 Model Consistency of Iterative Regularization

In this section, we present our main result, the model consistency of iterative regularization. From the results above, for small enough $\delta > 0$, the early stopping point $w_{\delta}^{(k_{\delta})}$ will be sufficiently close to w^{\dagger} . If we can further control the dual variable, then Lemma 2.3 can be applied to show that $w_{\delta}^{(k_{\delta})}$ is model consistent.

In terms of the dual variable, similar to the work [1], we also rely on the following nondegenerate source condition of w^{\dagger} . As we have imposed that w^{\dagger} is the unique solution of $(\overline{P_{\delta}})$ for $\delta = 0$. For the DGD Algorithm 1, denote the iterates as $w_0^{(k)}$ and $v_0^{(k)}$. The dual solution is then denoted as v^{\dagger} .

Definition 4.1 (Nondegenerate source condition). For the ground truth w^{\dagger} , let v^{\dagger} be a dual solution such that the following relative inclusion holds

$$z^{\dagger} := -X^{\top} v^{\dagger} \in \operatorname{ri}(\partial R_{\alpha}(w^{\dagger})). \qquad (\overline{\operatorname{SC}}_{w^{\dagger}})$$

When R is partly smooth at w^{\dagger} relative to $\mathcal{M}_{w^{\dagger}}$, so is R_{α} . According to Lemma 2.3, for small enough $\epsilon > 0$, define

$$\mathcal{U}_{\epsilon} = \bigcup_{w \in \mathcal{M}_{w^{\dagger}} \cap \mathcal{B}(w^{\dagger}, \epsilon)} (w + \partial R_{\alpha}(w)).$$

There holds $w^{\dagger} + z^{\dagger} \in \operatorname{int}(\mathcal{U}_{\epsilon})$. Denote $r := \operatorname{dist}(w^{\dagger} + z^{\dagger}, \operatorname{bdy}(\mathcal{U}_{\epsilon}))$ the distance of $w^{\dagger} + z^{\dagger}$ to the boundary of \mathcal{U}_{ϵ} . Back to the iterates of DGD, if we have the distance bound

$$\|(w_{\delta}^{(k_{\delta})} + z_{\delta}^{(k_{\delta})}) - (w^{\dagger} + z^{\dagger})\| < r,$$
(4.1)

holds for all large enough k, then we can conclude that $w_{\delta}^{(k_{\delta})} \in \mathcal{M}_{w^{\dagger}}$. This is the main idea behind the model consistency of iterative regularization. For the rest of the section, we shall elaborate the details of the proof.

4.1 Model consistency of iterative regularization

As discussed above, the key of showing model consistency of iterative regularization is to bound the distance in (4.1). Directly handling the distance in (4.1) is rather difficult, mostly due to the subgradient $z_{\delta}^{(k)}$ (or the dual variable $v_{\delta}^{(k)}$). Follow the idea of [16, 36], and we need an intermediate point. For $\delta = 0$, denote the iterates of DGD Algorithm 1 as $z_0^{(k)}, w_0^{(k)}$ and $v_0^{(k)}$, then

$$\|(w_{\delta}^{(k_{\delta})} + z_{\delta}^{(k_{\delta})}) - (w^{\dagger} + z^{\dagger})\| \le \|w_{\delta}^{(k_{\delta})} - w^{\dagger}\| + \|z_{\delta}^{(k_{\delta})} - z_{0}^{(k)}\| + \|z_{0}^{(k)} - z^{\dagger}\|.$$
(4.2)

The first term on the right hand side (rhs) can be directly handled by Lemma 3.2, while the later two terms need some technicality which is left to the proof.

As long as we bound the rhs of (4.2), we obtain the model consistency of iterative regularization with dual gradient descent.

Theorem 4.2 (Model consistency of DGD). Consider the problems $(\overline{P_{\delta}})$ and $(\overline{D_{\delta}})$. Suppose the following assumptions hold

- w^{\dagger} is the unique solution of $(\overline{P_{\delta}})$ for $\delta = 0$, and $v_0^{(k)} \to v^{\dagger}$ with v^{\dagger} being a dual solution.
- The nondegeneracy source condition $(\overline{SC}_{w^{\dagger}})$ holds for w^{\dagger} and v^{\dagger} .
- R is partly smooth at w^{\dagger} relative to $\mathcal{M}_{w^{\dagger}}$.

• The strong convexity α is small enough such that

$$\alpha c_{\delta} \|X\|^{-1} < r = \operatorname{dist} \left(w^{\dagger} + z^{\dagger}, \operatorname{bdy}(\mathcal{U}_{\epsilon}) \right), \tag{4.3}$$

where c_{δ} refers to the constant in Lemma 3.2.

Let k_{δ} be the optimal early stopping time of DGD, and denote the corresponding point as $w_{\delta}^{(k_{\delta})}$, then when $\delta > 0$ is small enough, there holds

$$w_{\delta}^{(k_{\delta})} \in \mathcal{M}_{w^{\dagger}}$$

Moreover, there exists $0 < \underline{k} \le k_{\delta} \le \overline{k}$ such that $w_{\delta}^{(k)} \in \mathcal{M}_{w^{\dagger}}$ holds for all $k \in [\underline{k}, \overline{k}]$.

Proof. Our proof is motivated by the proofs in [16, 36]. From the updates of DGD, the optimality condition of

$$w_{\delta}^{(k)} = \operatorname{prox}_{\alpha^{-1}R}(-\alpha^{-1}X^{\top}v_{\delta}^{(k)}),$$

yields

$$\frac{-\alpha^{-1}X^{\top}v_{\delta}^{(k)} - w_{\delta}^{(k)}}{\alpha^{-1}} = -X^{\top}v_{\delta}^{(k)} - \alpha w_{\delta}^{(k)} \in \partial R(w_{\delta}^{(k)}).$$

which can be further written as

$$z_{\delta}^{(k)} := -X^{\top} v_{\delta}^{(k)} \in \partial R(w_{\delta}^{(k)}) + \alpha w_{\delta}^{(k)} = \partial R_{\alpha}(w_{\delta}^{(k)}).$$

Recall that $z_{\delta}^{(k)} = -X^{\top} v_{\delta}^{(k)}$ and $z_{0}^{(k)} = -X^{\top} v_{0}^{(k)}$, then from (4.2) $\|(w_{\delta}^{(k)} + z_{\delta}^{(k)}) - (w^{\dagger} + z^{\dagger})\| \le \|w_{\delta}^{(k)} - w^{\dagger}\| + \|z_{\delta}^{(k)} - z_{0}^{(k)}\| + \|z_{0}^{(k)} - z^{\dagger}\|$

$$\begin{aligned} \|w_{\delta}^{(k)} - (w^{\dagger} + z^{\dagger})\| &\leq \|w_{\delta}^{(k)} - w^{\dagger}\| + \|z_{\delta}^{(k)} - z_{0}^{(k)}\| + \|z_{0}^{(k)} - z^{\dagger}\| \\ &\leq \|w_{\delta}^{(k)} - w^{\dagger}\| + \|X\| \|v_{\delta}^{(k)} - v_{0}^{(k)}\| + \|X\| \|v_{0}^{(k)} - v^{\dagger}\|. \end{aligned}$$

$$(4.4)$$

Let $k = k_{\delta}$ the early stopping time of iterative regularization. From the previous convergence result, we have

• From Lemma 3.2, we directly have

$$\|w_{\delta}^{(k_{\delta})} - w^{\dagger}\| \le \left(a(c^{1/2} + 1) + bc^{-1/2}\right)\delta^{1/2}.$$

Denote $f = a(c^{1/2} + 1) + bc^{-1/2}$.

• For the term $\|v_{\delta}^{(k)} - v_{0}^{(k)}\|$. Since $R_{\alpha}^{*}(-X^{\top}v)$ is convex differentiable with Lipschitz continuous gradient, we have from [34] that the operator

$$\mathcal{F} = \mathrm{Id} + \gamma X \nabla R^*_\alpha(-X^\top \cdot)$$

is nonexpansive (i.e. 1-Lipschitz), and $\mathcal{F}(v_{\delta}^{(k)}) = v_{\delta}^{(k)} + \gamma X w_{\delta}^{(k)}$. As a result, we get

$$\begin{aligned} \|v_{\delta}^{(k)} - v_{0}^{(k)}\| &= \|(v_{\delta}^{(k-1)} + \gamma(Xw_{\delta}^{(k-1)} - y_{\delta})) - (v_{0}^{(k-1)} + \gamma(Xw_{0}^{(k-1)} - y^{\dagger}))\| \\ &\leq \|(v_{\delta}^{(k-1)} + \gamma Xw_{\delta}^{(k-1)}) - (v_{0}^{(k-1)} + \gamma Xw_{0}^{(k-1)})\| + \gamma \|y_{\delta} - y^{\dagger}\| \\ &\leq \|\mathcal{F}(v_{\delta}^{(k-1)}) - \mathcal{F}(v_{0}^{(k-1)})\| + \gamma \|y_{\delta} - y^{\dagger}\| \\ &\leq \|v_{\delta}^{(k-1)} - v_{0}^{(k-1)}\| + \gamma \delta. \end{aligned}$$
(4.5)

Tracing back to k = 0 yields

$$\|v_{\delta}^{(k)} - v_0^{(k)}\| \le \gamma k\delta.$$

Letting $k = k_{\delta}$ leads to

$$\|v_{\delta}^{(k_{\delta})} - v_{0}^{(k_{\delta})}\| \leq \gamma k_{\delta} \delta = \alpha \|X\|^{-2} k_{\delta} \delta = \alpha c_{\delta} \|X\|^{-2}$$

• Lastly for $||v_0^{(k)} - v^{\dagger}||$, as we impose $v_0^{(k)} \to v^{\dagger}$, then $||v_0^{(k)} - v^{\dagger}|| \to 0$. Now for $k = k_{\delta}$, since k_{δ} increases as δ decreases, we have $\xi_{\delta} = ||v_{\delta}^{(k_{\delta})} - v^{\dagger}|| \to 0$ as $\delta \to 0$.

Assemble the above cases, we arrive at

$$\begin{aligned} |(w_{\delta}^{(k_{\delta})} + z_{\delta}^{(k_{\delta})}) - (w^{\dagger} + z^{\dagger})|| &\leq ||w_{\delta}^{(k_{\delta})} - w^{\dagger}|| + ||X|| ||v_{\delta}^{(k_{\delta})} - v_{0}^{(k)}|| + ||X|| ||v_{0}^{(k)} - v^{\dagger}|| \\ &\leq f \delta^{1/2} + \alpha c_{\delta} ||X||^{-1} + ||X|| \xi_{\delta} \end{aligned}$$
(4.6)

Apparently, if

$$\alpha c_{\delta} \|X\|^{-1} < r = \operatorname{dist}(w^{\dagger} + z^{\dagger}, \operatorname{bdy}(\mathcal{U}_{\epsilon}))$$

holds, then for small enough $\delta > 0$, we have

$$\|(w_{\delta}^{(k_{\delta})} + z_{\delta}^{(k_{\delta})}) - (w^{\dagger} + z^{\dagger})\| < r$$

which in turn implies $w_{\delta}^{(k_{\delta})} \in \mathcal{M}_{w^{\dagger}}$, and we obtain model consistency.

Next we show that there exists an interval $[\underline{k}, \overline{k}]$ with $k_{\delta} \in [\underline{k}, \overline{k}]$ such that $w_{\delta}^{(k)} \in \mathcal{M}_{w^{\dagger}}$ holds for all $k \in [\underline{k}, \overline{k}]$. Consider

$$\begin{aligned} &\|(w_{\delta}^{(k)} + z_{\delta}^{(k)}) - (w^{\dagger} + z^{\dagger})\| \\ &\leq \|(w_{\delta}^{(k)} + z_{\delta}^{(k)}) - (w_{\delta}^{(k_{\delta})} + z_{\delta}^{(k_{\delta})})\| + \|(w_{\delta}^{(k_{\delta})} + z_{\delta}^{(k_{\delta})}) - (w^{\dagger} + z^{\dagger})\| \end{aligned}$$

Apparently, the second term above is already discussed, and suppose

$$\|(w_{\delta}^{(k_{\delta})} + z_{\delta}^{(k_{\delta})}) - (w^{\dagger} + z^{\dagger})\| < r$$

holds. For the first term, we have

$$\begin{split} &\|(w_{\delta}^{(k)} + z_{\delta}^{(k)}) - (w_{\delta}^{(k_{\delta})} + z_{\delta}^{(k_{\delta})})\| \\ &\leq \|w_{\delta}^{(k)} - w_{\delta}^{(k_{\delta})}\| + \|z_{\delta}^{(k)} - z_{\delta}^{(k_{\delta})}\| \\ &= \|\operatorname{prox}_{\alpha^{-1}R}(\alpha^{-1}z_{\delta}^{(k)}) - \operatorname{prox}_{\alpha^{-1}R}(\alpha^{-1}z_{\delta}^{(k_{\delta})})\| + \|z_{\delta}^{(k)} - z_{\delta}^{(k_{\delta})}\| \\ &\leq (1 + \frac{1}{\alpha})\|z_{\delta}^{(k)} - z_{\delta}^{(k_{\delta})}\| \\ &\leq (1 + \frac{1}{\alpha})\|X\|\|v_{\delta}^{(k)} - v_{\delta}^{(k_{\delta})}\|. \end{split}$$

From Lemma 3.1, when $k = k_{\delta} - 1$, we have for some constant e > 0,

$$\|v_{\delta}^{(k)} - v_{\delta}^{(k_{\delta})}\| \le \frac{e}{k^{1/2}}$$

As a result, for small enough δ such that k_{δ} is large enough and

$$\left(1 + \frac{1}{\alpha}\right) \|X\| \frac{e}{k^{1/2}} + \|(w_{\delta}^{(k_{\delta})} + z_{\delta}^{(k_{\delta})}) - (w^{\dagger} + z^{\dagger})\| < r$$

also holds, we have $w_{\delta}^{(k)} \in \mathcal{M}_{w^{\dagger}}$ for $k = k_{\delta} - 1$. Similarly we can show that $w_{\delta}^{(k)} \in \mathcal{M}_{w^{\dagger}}$ for $k = k_{\delta} + 1$. We can continue with $k = k_{\delta} \pm 2, 3, ...$ until the above inequality fails. As a result, there exists $\underline{k} \leq k_{\delta} \leq \overline{k}$ such that $w_{\delta}^{(k)} \in \mathcal{M}_{w^{\dagger}}$ holds for all $k \in [\underline{k}, \overline{k}]$. We conclude the proof.

Remark 4.3. Our result is similar to that of [1] for variational regularization in the sense that the model consistency holds for small enough noise level δ .

For the ADGD algorithm, we can also obtain its model consistency property.

Corollary 4.4 (Model consistency of ADGD). Consider the problems $(\overline{P_{\delta}})$ and $(\overline{D_{\delta}})$. Suppose the assumptions of Theorem 4.2 hold. Let k_{δ} be the optimal early stopping time of ADGD, and denote the corresponding point as $w_{\delta}^{(k_{\delta})}$, then when $\delta > 0$ is small enough and θ is large enough, there holds

$$w^{(k_{\delta})}_{\delta} \in \mathcal{M}_{w^{\dagger}}$$

Moreover, there exists $0 < \underline{k} \le k_{\delta} \le \overline{k}$ such that $w_{\delta}^{(k)} \in \mathcal{M}_{w^{\dagger}}$ holds for all $k \in [\underline{k}, \overline{k}]$.

Below we provide a brief proof mainly highlighting the difference compared to the proof of Theorem 4.2, the main difficult part in this case is that the sequence generated by ADGD is not monotonic.

Proof. From the updates of ADGD, denote $z_{\delta}^{(k)} := -X^{\top} u_{\delta}^{(k)}$, then $z_{\delta}^{(k)} = \partial R_{\alpha}(w_{\delta}^{(k)}).$

Again from (4.2) we have

$$\begin{aligned} \|(w_{\delta}^{(k)} + z_{\delta}^{(k)}) - (w^{\dagger} + z^{\dagger})\| &\leq \|w_{\delta}^{(k)} - w^{\dagger}\| + \|z_{\delta}^{(k)} - z^{\dagger}\| \\ &= \|w_{\delta}^{(k)} - w^{\dagger}\| + \|X\| \|u_{\delta}^{(k)} - v^{\dagger}\| \\ &\leq \|w_{\delta}^{(k)} - w^{\dagger}\| + \|X\| \|u_{\delta}^{(k)} - u_{0}^{(k)}\| + \|X\| \|u_{0}^{(k)} - v^{\dagger}\|. \end{aligned}$$

$$(4.7)$$

The first and third terms of the last line above can be handled as in the proof of Theorem 4.2: term $\|w_{\delta}^{(k)} - w^{\dagger}\|$ is bounded by Lemma 3.4, while $\|u_{0}^{(k)} - v^{\dagger}\|$ converges to 0 due to the convergence. Next we discuss term $\|u_{\delta}^{(k)} - u_{0}^{(k)}\|$. Follow the derivation of (4.5), we have

$$\begin{split} |u_{\delta}^{(k)} - u_{0}^{(k)}|| &= \|(v_{\delta}^{(k)} + \gamma(Xr_{\delta}^{(k)} - y_{\delta})) - (v_{0}^{(k)} + \gamma(Xr_{0}^{(k)} - y^{\dagger}))\| \\ &\leq \|\mathcal{F}(v_{\delta}^{(k)}) - \mathcal{F}(v_{0}^{(k)})\| + \gamma\|y_{\delta} - y^{\dagger}\| \\ &\leq \|v_{\delta}^{(k)} - v_{0}^{(k)}\| + \gamma\delta \\ &= \|(u_{\delta}^{(k-1)} + \frac{k-1}{k+\theta}(u_{\delta}^{(k-1)} - u_{\delta}^{(k-2)})) - (u_{0}^{(k-1)} + \frac{k-1}{k+\theta}(u_{0}^{(k-1)} - u_{0}^{(k-2)}))\| + \gamma\delta \\ &\leq \|u_{\delta}^{(k-1)} - u_{0}^{(k-1)}\| + \frac{k-1}{k+\theta}(\|u_{\delta}^{(k-1)} - u_{\delta}^{(k-2)}\| + \|u_{0}^{(k-1)} - u_{0}^{(k-2)}\|) + \gamma\delta \\ &\leq \sum_{i=1}^{k} \frac{i-1}{i+\theta}(\|u_{\delta}^{(i-1)} - u_{\delta}^{(i-2)}\| + \|u_{0}^{(i-1)} - u_{0}^{(i-2)}\|) + \gamma k\delta. \end{split}$$

Denote

$$\beta_k = \sum_{i=1}^k \frac{i-1}{i+\theta} \big(\|u_{\delta}^{(i-1)} - u_{\delta}^{(i-2)}\| + \|u_0^{(i-1)} - u_0^{(i-2)}\| \big).$$

From the convergence result Lemma 3.4, $\beta_k < +\infty$ for any $k \in \mathbb{N}$. When θ is large enough, we can control β_k to be small enough. As a consequence, we can bound $\|(w_{\delta}^{(k_{\delta})} + z_{\delta}^{(k_{\delta})}) - (w^{\dagger} + z^{\dagger})\|$ and obtain the model consistency of ADGD.

Remark 4.5. Though we can obtain model consistency for ADGD, the result somehow is counter intuitive, in the sense that

- When δ is very small, k_δ is very large. In turn, β_{k_δ} will be large.
 To bound ||(w_δ^(k_δ) + z_δ^(k_δ)) − (w[†] + z[†])||, we need θ to be even large, which makes ADGD behaves more like DGD despite it still converges at $O(1/k^2)$ rate.

The main reason, as we emphasized above, is that the sequence generated by ADGD is not monotonic, and the inertial term introduces the composition of the current point $u_{\delta}^{(k)}$ and its history $u_{\delta}^{(k-1)}$. While in practice, model consistency of ADGD are observed for small value of θ ; see Section 6 for more details.

Model consistency of continuous time dynamics 4.2

In [22], the authors considered the continuous time dynamics of DGD and its accelerated version, and demonstrate their iterative regularization property. We can also extend our model consistency result to continuous time dynamics. In this part, for simplicity we only consider the continuous time dynamics of DGD. For the DGD iteration, it can be reformulated as

$$\begin{split} w_{\delta}^{(k)} &= \nabla R_{\alpha}^*(-X^{\top}v_{\delta}^{(k)}),\\ \frac{v_{\delta}^{(k+1)} - v_{\delta}^{(k)}}{\gamma} &= Xw_{\delta}^{(k)} - y_{\delta} = -\nabla \phi(v_{\delta}^{(k)}). \end{split}$$

Suppose let γ be time step and $t_k = k\gamma$, then $v_{\delta}^{(k)}$ and $w_{\delta}^{(k)}$ are the samples of the continuous time dynamics. Let $\gamma \to 0$ we get the following continuous time dynamics of DGD

$$v_{\delta}(0) = v_0 = 0, \quad \begin{cases} w_{\delta}(t) = \nabla R^*_{\alpha}(-X^{\top}v_{\delta}(t)) \\ \dot{v}_{\delta}(t) = -\nabla \phi(v_{\delta}(t)). \end{cases}$$
(4.8)

Corollary 4.6 (Model consistency of continuous time DGD). Consider the problems $(\overline{P_{\delta}})$ and $(\overline{D_{\delta}})$. Suppose the assumptions of Theorem 4.2 hold. Let t_{δ} be the optimal early stopping time of (4.8), and denote the corresponding point as $w_{\delta}(t_{\delta})$, then when $\delta > 0$ is small enough, there holds

$$w_{\delta}(t_{\delta}) \in \mathcal{M}_{w^{\dagger}}$$

Moreover, there exists $0 < \underline{t} \le t_{\delta} \le \overline{t}$ such that $w_{\delta}(t) \in \mathcal{M}_{w^{\dagger}}$ holds for all $t \in [\underline{t}, \overline{t}]$.

In Figure 4 we provide a numerical demonstration between the comparison of DGD and continuous time DGD.

5 Local linear convergence

In Theorem 4.2 we have an interval $[\underline{k}, \overline{k}]$ where model consistency holds. Since along the manifold $\mathcal{M}_{w^{\dagger}}$ the R(x) is smooth differentiable, plus that $\mathcal{M}_{w^{\dagger}}$ has low dimension, these imply that the problem becomes better conditioned along $\mathcal{M}_{w^{\dagger}}$ and the algorithm DGD could achieve faster convergence speed, which turns out to be linear.

As local linear convergence is a byproduct of model consistency, for the sake of simplicity, we only discuss the case that $\mathcal{M}_{w^{\dagger}}$ is linear, which includes ℓ_1 -norm, TV semi-norm and $\ell_{1,2}$ -norm in Table 1. For linear $\mathcal{M}_{w^{\dagger}}$, it can be represented as

$$\mathcal{M}_{w^{\dagger}} = w^{\dagger} + T_{w^{\dagger}}$$

where $T_{w^{\dagger}}$ is the tangent space of $\mathcal{M}_{w^{\dagger}}$ at w^{\dagger} . Moreover, for any $w \in \mathcal{M}_{w^{\dagger}}$, there holds

$$w - w^{\dagger} = P_{T_{w^{\dagger}}}(w - w^{\dagger}),$$

where $P_{T_{w^{\dagger}}}$ is the projection matrix onto $T_{w^{\dagger}}$. For X, we need the following "restricted injectivity" condition.

Definition 5.1 (Restricted injectivity). Let $T_{w^{\dagger}}$ be the tangent space of $\mathcal{M}_{w^{\dagger}}$ at w^{\dagger} . The restricted injectivity condition is defined as

$$\ker(X) \cap T_{w^{\dagger}} = \{0\},\tag{INJ}_{T_{w^{\dagger}}}$$

where $\ker(X)$ is the null space of X.

Let $X_{T_{w^{\dagger}}} := X P_{T_{w^{\dagger}}}$, condition $(INJ_{T_{w^{\dagger}}})$ implies $X_{T_{w^{\dagger}}}^{\top} X_{T_{w^{\dagger}}}$ is positive definite along $T_{w^{\dagger}}$.

Remark 5.2. Condition $(INJ_{T_w^{\dagger}})$ is also considered in [28] where the authors study the local linear convergence of the proximal gradient method type methods.

Since $\mathcal{M}_{w^{\dagger}}$ is linear, we have the Riemannian gradient and Hessian of R at $w_{\delta}^{(\underline{k})}$ as

$$\nabla_{\mathcal{M}_{w^{\dagger}}} R(w_{\delta}^{(\underline{k})}) = P_{T_{w^{\dagger}}} \partial R(w_{\delta}^{(\underline{k})}) \quad \text{and} \quad \nabla^{2}_{\mathcal{M}_{w^{\dagger}}} R(w_{\delta}^{(\underline{k})}) = P_{T_{w^{\dagger}}} \nabla^{2} R(w_{\delta}^{(\underline{k})}) P_{T_{w^{\dagger}}}.$$

Along $T_{w^{\dagger}}$, that $P_{T_{w^{\dagger}}} + \frac{1}{\alpha} \nabla^2_{\mathcal{M}_{w^{\dagger}}} R(w^{(\underline{k})}_{\delta})$ is invertible, and define the following matrix

$$M_{\text{DGD}} := P_{T_{w^{\dagger}}} - \frac{1}{\|X\|^2} \left(P_{T_{w^{\dagger}}} + \nabla^2_{\mathcal{M}_{w^{\dagger}}} R(w_{\delta}^{(\underline{k})}) \right)^{-1} X_{T_{w^{\dagger}}}^{\top} X_{T_{w^{\dagger}}}.$$

Denote $\rho_{_{M_{\rm DGD}}}$ the spectral radius of $M_{\rm DGD},$ we have the following local linear convergence rate result.

Theorem 5.3 (Local linear convergence). Let R(w) be partly smooth at w^{\dagger} relative to $\mathcal{M}_{w^{\dagger}}$ such that $\mathcal{M}_{w^{\dagger}}$ is linear. Let the assumptions in Theorem 4.2 and condition (INJ_{T_{w^{\dagger}}) hold, then}

$$\rho_{M_{\text{DGD}}} < 1.$$

If $[\underline{k}, \overline{k}]$ contains more than three points, then for all $k \in]\underline{k}, \overline{k}[$ and any $\rho \in]\rho(M_{\text{DGD}}), 1[$ we have

$$||w_{\delta}^{(k+1)} - w_{\delta}^{(k)}|| \le O(\rho^{k-\underline{k}}).$$

Remark 5.4. Local linear convergence of first-order method along the identified manifold is well studied in the literature, see for instance [27, 28, 29, 31, 40] and the references therein.

Proof. Denote $W = P_{T_{w^{\dagger}}} + \frac{1}{\alpha} \nabla^2_{\mathcal{M}_{w^{\dagger}}} R(w_{\delta}^{(\underline{k})})$, then along $T_{w^{\dagger}}$ it is invertible, and all the eigenvalues of W^{-1} are real and less than or equal to 1. Moreover, as W is symmetric, then

$$W^{-1}X_{T_{w^{\dagger}}}^{\top}X_{T_{w^{\dagger}}} = W^{-1/2} \left(W^{-1/2}X_{T_{w^{\dagger}}}^{\top}X_{T_{w^{\dagger}}} W^{-1/2} \right) W^{1/2}$$

which means $W^{-1}X_{T_{w^{\dagger}}}^{\top}X_{T_{w^{\dagger}}}$ is similar to $W^{-1/2}X_{T_{w^{\dagger}}}^{\top}X_{T_{w^{\dagger}}}W^{-1/2}$. Note that $X_{T_{w^{\dagger}}}^{\top}X_{T_{w^{\dagger}}}$ is positive definite along $T_{w^{\dagger}}$ and $\|X_{T_{w^{\dagger}}}\| \leq \|X\|$. Let $\sigma_i > 0, i = 1, ..., d$ be the eigenvalues of $W^{-1/2}X_{T_{w^{\dagger}}}^{\top}X_{T_{w^{\dagger}}}W^{-1/2}$ where $d = \dim(T_{w^{\dagger}})$ is the dimension of $T_{w^{\dagger}}$. Then the eigenvalues of M_{DGD} along $T_{w^{\dagger}}$ are $1 - \frac{1}{\|X\|^2}\sigma_i, i = 1, ..., d$ which are strictly less than 1.

Denote $g_{\delta}^{(k)} := -\frac{1}{\alpha} X^{\top} v_{\delta}^{(k)}$, then from the update of $w_{\delta}^{(k)}$,

$$g_{\delta}^{(k)} - w_{\delta}^{(k)} \in \alpha^{-1} \partial R(w_{\delta}^{(k)}).$$

For $k \in]\underline{k}, \overline{k}[$, we have $w_{\delta}^{(k)}, w_{\delta}^{(k+1)} \in \mathcal{M}_{w^{\dagger}}$ and

$$\begin{aligned} P_{T_{w^{\dagger}}}(g_{\delta}^{(k)} - w_{\delta}^{(k)}) &= \alpha^{-1} P_{T_{w^{\dagger}}}\left(\partial R(w_{\delta}^{(k)})\right) \\ P_{T_{w^{\dagger}}}(g_{\delta}^{(k+1)} - w_{\delta}^{(k+1)}) &= \alpha^{-1} P_{T_{w^{\dagger}}}\left(\partial R(w_{\delta}^{(k+1)})\right). \end{aligned}$$

Take the difference of the above two equations, we get

$$\begin{split} P_{T_{w^{\dagger}}}(w_{\delta}^{(k+1)} - w_{\delta}^{(k)}) + \alpha^{-1}P_{T_{w^{\dagger}}}\left(\partial R(w_{\delta}^{(k+1)})\right) &- \alpha^{-1}P_{T_{w^{\dagger}}}\left(\partial R(w_{\delta}^{(k)})\right) \\ &= P_{T_{w^{\dagger}}}(w_{\delta}^{(k+1)} - w_{\delta}^{(k)}) \\ &+ \alpha^{-1}\left(\nabla_{\mathcal{M}_{w^{\dagger}}}^{2}R(w_{\delta}^{(k)})(w_{\delta}^{(k+1)} - w_{\delta}^{(k)}) + o(\|w_{\delta}^{(k+1)} - w_{\delta}^{(k)}\|)\right) \\ &- \alpha^{-1}\left(\nabla_{\mathcal{M}_{w^{\dagger}}}^{2}R(w_{\delta}^{(k)})(w_{\delta}^{(k)} - w_{\delta}^{(k)}) + o(\|w_{\delta}^{(k)} - w_{\delta}^{(k)}\|)\right) \\ &= \left(P_{T_{w^{\dagger}}} + \frac{1}{\alpha}\nabla_{\mathcal{M}_{w^{\dagger}}}^{2}R(w_{\delta}^{(k)})\right)(w_{\delta}^{(k+1)} - w_{\delta}^{(k)}) + o(\|w_{\delta}^{(k+1)} - w_{\delta}^{(k)}\|) + o(\|w_{\delta}^{(k)} - w_{\delta}^{(k)}\|) \\ &= P_{T_{w^{\dagger}}}(g_{\delta}^{(k+1)} - g_{\delta}^{(k)}) \\ &= -\frac{1}{\alpha}P_{T_{w^{\dagger}}}X^{\top}(v_{\delta}^{(k+1)} - v_{\delta}^{(k)}) \\ &= -\frac{\gamma}{\alpha}P_{T_{w^{\dagger}}}X^{\top}(Xw_{\delta}^{(k)} - y_{\delta}). \end{split}$$

Similarly, we have

$$W(w_{\delta}^{(k)} - w_{\delta}^{(k-1)}) + o(\|w_{\delta}^{(k)} - w_{\delta}^{(\underline{k})}\|) + o(\|w_{\delta}^{(k-1)} - w_{\delta}^{(\underline{k})}\|) = -\frac{\gamma}{\alpha} P_{T_{w^{\dagger}}} X^{\top} (Xw_{\delta}^{(k-1)} - y_{\delta}).$$

The difference of the above two equations gives

$$\begin{split} W(w_{\delta}^{(k+1)} - w_{\delta}^{(k)}) - W(w_{\delta}^{(k)} - w_{\delta}^{(k-1)}) &= -\frac{\gamma}{\alpha} P_{T_{w^{\dagger}}} X^{\top} X(w_{\delta}^{(k)} - w_{\delta}^{(k-1)}) + \eta_k \\ &= -\frac{\gamma}{\alpha} P_{T_{w^{\dagger}}} X^{\top} X P_{T_{w^{\dagger}}}(w_{\delta}^{(k)} - w_{\delta}^{(k-1)}) + \eta_k \\ &= -\frac{\gamma}{\alpha} X_{T_{w^{\dagger}}}^{\top} X_{T_{w^{\dagger}}}(w_{\delta}^{(k)} - w_{\delta}^{(k-1)}) + \eta_k, \end{split}$$

where

$$\eta_k = o(\|w_{\delta}^{(k+1)} - w_{\delta}^{(\underline{k})}\|) + o(\|w_{\delta}^{(k)} - w_{\delta}^{(\underline{k})}\|) + o(\|w_{\delta}^{(k)} - w_{\delta}^{(\underline{k})}\|) + o(\|w_{\delta}^{(k-1)} - w_{\delta}^{(\underline{k})}\|).$$

Since $\gamma = \alpha / \|X\|^2$, we have

$$w_{\delta}^{(k+1)} - w_{\delta}^{(k)} = W^{-1} \Big(W - \frac{1}{\|X\|^2} X_{T_{w^{\dagger}}}^{\top} X_{T_{w^{\dagger}}} \Big) (w_{\delta}^{(k)} - w_{\delta}^{(k-1)}) + W^{-1} \eta_k$$
$$= M_{\text{DGD}} (w_{\delta}^{(k)} - w_{\delta}^{(k-1)}) + \bar{\eta}_k,$$

with $\bar{\eta}_k = W^{-1}\eta_k$. Tracking back to k_{δ} , we have $w_{\delta}^{(k+1)} - w_{\delta}^{(k)} = M_{\text{DGD}}(w_{\delta}^{(k)} - w_{\delta}^{(k)})$

from which we get

$$\|w_{\delta}^{(k+1)} - w_{\delta}^{(k)}\| \le \|M_{\text{DGD}}^{k-\underline{k}}\| \| w_{\delta}^{(\underline{k}+1)} - w_{\delta}^{(\underline{k})}\| + \|\sum_{i=\underline{k}+1}^{k} M_{\text{DGD}}^{k-i} \bar{\eta}_{i}\|.$$

The spectral radius theorem there exists c > 0 such that

$$\|M_{\rm DGD}^{k-\underline{k}}\| \le c\rho^{k-\underline{k}},$$

for any $\rho \in]\rho(M_{\text{DGD}}), 1[$. Since $k \leq \overline{k}$ is finite, $\|\sum_{i=\underline{k}+1}^{k} M_{\text{DGD}}^{k-i} \overline{\eta}_i\|$ is also finite, then

$$\left\|\sum_{i=\underline{k}+1}^{k} M_{\mathrm{DGD}}^{k-i} \bar{\eta}_{i}\right\| = \rho^{k-\underline{k}} \times \frac{1}{\rho^{k-\underline{k}}} \left\|\sum_{i=\underline{k}+1}^{k} M_{\mathrm{DGD}}^{k-i} \bar{\eta}_{i}\right\|.$$

Let

$$d = \max\left\{c, \ \frac{1}{\rho^{\overline{k}-\underline{k}}} \|\sum_{i=\underline{k}+1}^{\overline{k}} M_{\mathrm{DGD}}^{k-i} \bar{\eta}_i\|\right\},\$$

we have

$$\|w_{\delta}^{(k+1)} - w_{\delta}^{(k)}\| \le d\rho^{k-\underline{k}}$$

which concludes the proof.

Remark 5.5. As Theorem 5.3 requires the manifold $\mathcal{M}_{w^{\dagger}}$ to be linear, it does not hold for nuclear norm whose manifold is curved. By incorporating more Riemannian manifold arguments (such as parallel transportation [28]), we can extend Theorem 5.3 to more general $\mathcal{M}_{w^{\dagger}}$ that includes nuclear norm. However, for the sake of simplicity, we decide to omit the discussion here and only provide numerical illustration in the next section.

6 Numerical Experiments

In this section we provide numerical experiments to support our theoretical findings. We first demonstrate the model consistency result, and then the local linear convergence behavior. Four low-complexity regularization functions R are considered: $\ell_1, \ell_{1,2}$ -norms, one dimensional total variation (1d-TV) and nuclear norm.

6.1 Model consistency

Problem settings Recall that in the forward model (1.1), $X \in \mathbb{R}^{n \times p}$ is the forward operator, $\varepsilon \sim \mathcal{N}(0, \delta^2)$ is the noise, For each choice of R, the corresponding problem settings are

- $\ell_1, \ell_{1,2}$ -norms $(n, p) = (100, 500), w^{\dagger} \in \mathbb{R}^p$ has 5 non-zero elements, $X \in \mathbb{R}^{n \times p}$ with element sampled from standard normal distribution normalized by \sqrt{n} . For $\ell_{1,2}$ -norm, the group size is 5, without overlapping.
- 1d-TV $(n,p) = (20,50), w^{\dagger} \in \mathbb{R}^p$ and $\|D_{\text{DIF}}w^{\dagger}\|_1 = 1$. $X \in \mathbb{R}^{n \times p}$ with element sampled from standard normal distribution normalized by \sqrt{n} .
- Nuclear norm $w^{\dagger} \in \mathbb{R}^{20 \times 20}$ is a square matrix with rank $(w^{\dagger}) = 1, X \in \mathbb{R}^{20 \times 20}$ is a 0-1 projection mask with only 50% entries equal 1.

For noise ε , the strength of the noise determined by signal-to-noise-ratio (SNR), which is defined by

$$\operatorname{SNR} = 20 \log \left(\frac{\|y\|}{\|\varepsilon\|} \right).$$

In our experiment, SNR = 40 is considered.

In the model $(\overline{P_{\delta}})$, we fix $\alpha = 0.01$. For both DGD and ADGD methods, the step-size is $\gamma = \alpha / \|X\|^2$ and the parameter θ for ADGD is set as 5. The starting point of both algorithms is set as 0, as indicated in the algorithms.

Case ℓ_1 -norm We first report the results on ℓ_1 -norm, which is shown in Figure 1: the *left* column is for DGD, while the *right* one is for ADGD. In each figure, the *blue* line stands for the normalized error $\frac{\|w_{\delta}^{(k)}-w^{\dagger}\|}{\|w^{\dagger}\|}$, and the *red* line represents the support size of $w_{\delta}^{(k)}$. We obtain the following observations:

- Support size $|\operatorname{supp}(w_{\delta}^{(k)})|$: As the initial point is 0, the support of $w_{\delta}^{(k)}$ increases from 0. For certain range of k, we have $|\operatorname{supp}(w_{\delta}^{(k)})| = 5$ which corresponds to the model or the regularization: For the error ^{||w_δ^(k)|/||w[†]||}, it can be seen that the minimal value
 Iterative regularization: For the error ^{||w_δ^(k)-w[†]||}/_{||w[†]||}, it can be seen that the minimal value
- is obtained when model consistency occurs. Note that the error is flat inside the model consistency interval, we can use model consistency as guide to find stopping time.



Figure 1: Model consistency result of ℓ_1 -norm.

For this problem, we also conduct experiment to test the smallest possible SNR value such that model consistency occurs. To this end, we let $SNR \in [10, 60]$, for each value we run DGD and record the support size of the optimal $w_{\delta}^{(k_{\delta})}$ in terms of the distance to w^{\dagger} . The value

 $|\operatorname{supp}(w_{\delta}^{(k_{\delta})})|$ over SNR value is plotted in Figure 2, from which we observe that model consistency holds for SNR greater than about 21.



Figure 2: Model consistency of DGD over different SNR values.

Other cases In Figure 3 we provide the model consistency result for $\ell_{1,2}$ -norm (1st row), 1d-TV (2nd row) and nuclear norm (3rd row). The observations are quite similar to those of ℓ_1 -norm. Moreover,

- For $\ell_{1,2}$ -norm, the non-zero groups of $w_{\delta}^{(k)}$ in the model consistency interval are also identical to those of w^{\dagger} .
- The 1d-TV is more sensitive to noise, though for the present example model consistency holds, there are also cases where it fails under same SNR value. We refer to [1] for more discussion on the structure dependency of noise.

Model consistency of the continuous dynamics We also verify the model consistency of the continuous dynamical system (4.8) using ℓ_1 -norm with SNR = 40. The 4-order Runge-Kutta method is used to solve the system (4.8). The result is provided in Figure 4, accompanied with DGD method. It can be observed that model consistency is also obtained by the continuous dynamical system.

6.2 Local linear convergence

From the above illustration, model consistency indeed occurs in an interval. In this part, we show that $\|w_{\delta}^{(k+1)} - w_{\delta}^{(k)}\|$ converges linearly.

Due to the numerical precision problem, the problem size considered in this part is smaller than the above. In particular, the following settings are considered

- ℓ_1 , $\ell_{1,2}$ -norms (n,p) = (20,100), $w^{\dagger} \in \mathbb{R}^p$ has 2 non-zero elements, $X \in \mathbb{R}^{n \times p}$ with element sampled from standard normal distribution normalized by \sqrt{n} . For $\ell_{1,2}$ -norm, the group size is 2, without overlapping.
- Nuclear norm $w^{\dagger} \in \mathbb{R}^{10 \times 10}$ is a square matrix with $\operatorname{rank}(w^{\dagger}) = 1, X \in \mathbb{R}^{10 \times 10}$ is a 0-1 projection mask with only 10% entries equal 1.

For noise ε , SNR = 40 is considered. In the model $(\overline{P_{\delta}})$, same as before we fix $\alpha = 0.01$. The step-size is $\gamma = \alpha/\|X\|^2$ and the starting point of both algorithms is set as 0, as indicated in the algorithms.

In Figure 5 we provide the observations of local linear convergence for ℓ_1 -norm and $\ell_{1,2}$ -norm, the left column shows the global behavior of $||w_{\delta}^{(k+1)} - w_{\delta}^{(k)}||$, and the right column displays the zoom-in region of model consistency interval.



Figure 3: Model consistency result of $\ell_{1,2}\text{-norm},$ 1d-TV and nuclear norm.



Figure 4: Model consistency of the dynamical system (4.8).



Figure 5: Local linear convergence of DGD in terms of $\|w_{\delta}^{(k+1)} - w_{\delta}^{(k)}\|$ for ℓ_1 -norm and $\ell_{1,2}$ -norm.

We also provide an example for the nuclear norm in Figure 6, thought our theory does not cover this case, local linear convergence is observed.



Figure 6: Numerical results for Nuclear norm

7 Conclusions

In this paper, we study the model consistency and local linear convergence of the iterative regularization methods including dual gradient descent (DGD) and its accelerated scheme. Relying on a recent development of the partial smoothness, we show that the output of iterative regularization can identify the model of the ground truth in the small noise regime. We also show that the model consistency holds for an interval of stopping time, in which the residual error of the DGD scheme converges linearly. Numerical results are provided to validate our theoretical findings.

Acknowledgements JL is supported by the National Natural Science Foundation of China (No. 12201405), the "Fundamental Research Funds for the Central Universities", the National Science Foundation of China (BC4190065) and the Shanghai Municipal Science and Technology Major Project (2021SHZDZX0102).

References

- Samuel Vaiter, Gabriel Peyré, and Jalal Fadili. Model consistency of partly smooth regularizers. *IEEE Transactions on Information Theory*, 64(3):1725–1737, 2017.
- [2] Emmanuel J Candès, Justin Romberg, and Terence Tao. Robust uncertainty principles: Exact signal reconstruction from highly incomplete frequency information. *IEEE Transactions* on information theory, 52(2):489–509, 2006.
- [3] Junzhou Huang and Tong Zhang. The benefit of group sparsity. The Annals of Statistics, 38(4):1978–2004, 2010.
- [4] Leonid I Rudin, Stanley Osher, and Emad Fatemi. Nonlinear total variation based noise removal algorithms. *Physica D: nonlinear phenomena*, 60(1-4):259–268, 1992.
- [5] Emmanuel J Candes and Benjamin Recht. Exact matrix completion via convex optimization. Communications of the ACM, 55(6):111–119, 2012.
- [6] Martin Benning and Martin Burger. Modern regularization methods for inverse problems. Acta numerica, 27:1–111, 2018.
- [7] V Morozov. On the solution of functional equations by the method of regularization. Doklady Akademii Nauk SSSR, 167(3):510, 1966.

- [8] Gene H Golub and Urs Von Matt. Generalized cross-validation for large-scale problems. Journal of Computational and Graphical Statistics, 6(1):1–34, 1997.
- [9] Charles M Stein. Estimation of the mean of a multivariate normal distribution. *The annals of Statistics*, pages 1135–1151, 1981.
- [10] Charles-Alban Deledalle, Samuel Vaiter, Jalal Fadili, and Gabriel Peyré. Stein unbiased gradient estimator of the risk (sugar) for multiple parameter selection. SIAM Journal on Imaging Sciences, 7(4):2448–2487, 2014.
- [11] Jonathan Chirinos-Rodríguez, Ernesto De Vito, Cesare Molinari, Lorenzo Rosasco, and Silvia Villa. On learning the optimal regularization parameter in inverse problems. *Inverse Problems*, 40(12):125004, 2024.
- [12] Samuel Vaiter, Mohammad Golbabaee, Jalal Fadili, and Gabriel Peyré. Model selection with low complexity priors. *Information and Inference: A Journal of the IMA*, 4(3):230– 287, 2015.
- [13] Jalal Fadili, Guillaume Garrigos, Jérôme Malick, and Gabriel Peyré. Model consistency for learning with mirror-stratifiable regularizers. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pages 1236–1244. PMLR, 2019.
- [14] Heinz Werner Engl, Martin Hanke, and Andreas Neubauer. Regularization of inverse problems, volume 375. Springer Science & Business Media, 1996.
- [15] Bangti Jin and Xiliang Lu. On the regularizing property of stochastic gradient descent. Inverse Problems, 35(1):015004, 2018.
- [16] Guillaume Garrigos, Lorenzo Rosasco, and Silvia Villa. Iterative regularization via dual diagonal descent. Journal of Mathematical Imaging and Vision, 60:189–215, 2018.
- [17] Martin Burger, Elena Resmerita, and Lin He. Error estimation for bregman iterations and inverse scale space methods in image restoration. *Computing*, 81:109–135, 2007.
- [18] Lorenzo Rosasco and Silvia Villa. Learning with incremental iterative regularization. Advances in Neural Information Processing Systems, 28, 2015.
- [19] Cesare Molinari, Mathurin Massias, Lorenzo Rosasco, and Silvia Villa. Iterative regularization for convex regularizers. In *International conference on artificial intelligence and statistics*, pages 1684–1692. PMLR, 2021.
- [20] Cesare Molinari, Mathurin Massias, Lorenzo Rosasco, and Silvia Villa. Iterative regularization for low complexity regularizers. *Numerische Mathematik*, pages 1–49, 2024.
- [21] Cristian Vega, Cesare Molinari, Lorenzo Rosasco, and Silvia Villa. Fast iterative regularization by reusing data. Journal of Inverse and Ill-posed Problems, 32(4):733–759, 2024.
- [22] Luca Calatroni, Guillaume Garrigos, Lorenzo Rosasco, and Silvia Villa. Accelerated iterative regularization via dual diagonal descent. SIAM Journal on Optimization, 31(1):754– 784, 2021.
- [23] Michael P Friedlander and Paul Tseng. Exact regularization of convex programs. SIAM Journal on Optimization, 18(4):1326–1350, 2008.
- [24] Wotao Yin. Analysis and generalizations of the linearized bregman method. SIAM Journal on Imaging Sciences, 3(4):856–877, 2010.
- [25] Adrian S Lewis. Active sets, nonsmoothness, and sensitivity. SIAM Journal on Optimization, 13(3):702–725, 2002.
- [26] Adrian S Lewis and Tonghua Tian. Identifiability, the kl property in metric spaces, and subgradient curves. Foundations of Computational Mathematics, pages 1–38, 2024.
- [27] Jingwei Liang, Jalal Fadili, and Gabriel Peyré. Local linear convergence of forward-

backward under partial smoothness. Advances in neural information processing systems, 27, 2014.

- [28] Jingwei Liang, Jalal Fadili, and Gabriel Peyré. Activity identification and local linear convergence of forward-backward-type methods. SIAM Journal on Optimization, 27(1):408– 437, 2017.
- [29] Cesare Molinari, Jingwei Liang, and Jalal Fadili. Convergence rates of forward-douglasrachford splitting method. *Journal of Optimization Theory and Applications*, 182:606–639, 2019.
- [30] Julie Nutini, Issam Laradji, and Mark Schmidt. Let's make block coordinate descent converge faster: faster greedy rules, message-passing, active-set complexity, and superlinear convergence. *Journal of Machine Learning Research*, 23(131):1–74, 2022.
- [31] Quentin Klopfenstein, Quentin Bertrand, Alexandre Gramfort, Joseph Salmon, and Samuel Vaiter. Local linear convergence of proximal coordinate descent algorithm. *Optimization Letters*, 18(1):135–154, 2024.
- [32] Ziqi Qin and Jingwei Liang. Partial smoothness, subdifferentials and set-valued operators. arXiv preprint arXiv:2501.15540, 2025.
- [33] Amir Beck. First-order methods in optimization. SIAM, 2017.
- [34] Heinz H Bauschke, Patrick L Combettes, Heinz H Bauschke, and Patrick L Combettes. Convex Analysis and Monotone Operator Theory in Hilbert Spaces. Springer, 2017.
- [35] Warren L Hare and Adrian S Lewis. Identifying active constraints via partial smoothness and prox-regularity. *Journal of Convex Analysis*, 11(2):251–266, 2004.
- [36] Silvia Villa, Simon Matet, Bang Cong Vũ, and Lorenzo Rosasco. Implicit regularization with strongly convex bias: Stability and acceleration. Analysis and Applications, 21(01):165–191, 2023.
- [37] Jingwei Liang, Jalal Fadili, and Gabriel Peyré. Convergence rates with inexact nonexpansive operators. *Mathematical Programming*, 159(1):403–434, 2016.
- [38] Y. Nesterov. A method for solving the convex programming problem with convergence rate $O(1/k^2)$. Dokl. Akad. Nauk SSSR, 269(3):543-547, 1983.
- [39] Jingwei Liang, Tao Luo, and Carola-Bibiane Schönlieb. Improving "fast iterative shrinkagethresholding algorithm": Faster, smarter, and greedier. SIAM Journal on Scientific Computing, 44(3):A1069–A1091, 2022.
- [40] Clarice Poon and Jingwei Liang. Trajectory of alternating direction method of multipliers and adaptive acceleration. Advances in Neural Information Processing Systems, 32, 2019.