SUBLINEARLY MORSENESS IN HIGHER RANK SYMMETRIC SPACES

ROU WEN

ABSTRACT. The goal of this paper is to develop a theory of "sublinearly Morse boundary" and prove a corresponding sublinearly Morse lemma in higher rank symmetric space of non-compact type. This is motivated by the work of Kapovich–Leeb–Porti and the theory of sublinearly Morse quasi-geodesics developed in the context of CAT(0) geometry.

1. INTRODUCTION

The Morse lemma, also known as the fellow traveling property of quasi-geodesics, in hyperbolic space is a fundamental tool for studying such spaces. However, it fails in higher rank symmetric spaces due to the existence of isometrically embedded copies of Euclidean spaces.

In order to deal with this, Kapovich–Leeb–Porti defined uniformly regular sequences. Let X be a higher rank symmetric space of non-compact type, d_X a Riemannian symmetric distance on X, and θ be a subset of the simple roots that correspond to a root system of the Lie algebra associated to X. Let [xy] denote the geodesic segment in X that joins x and y, [xy] is called θ -uniform regular if the Cartan projection of [xy] stays uniformly away from the Weyl chamber walls defined by elements in θ .

Definition 1.1. A discrete path $q : \mathbb{Z} \to X$ is θ -uniformly regular if there exists a constant D such that for any $m < n \in \mathbb{Z}$ with $(n - m) \ge D$, the geodesic segment [q(m)q(n)] is θ -uniformly regular. In addition, if q is a quasi-isometry, such sequences are called undistorted θ -uniformly regular.

Among many other things, the undistorted θ -uniform regular sequences provide an equivalent characterization of Anosov subgroups: a discrete subgroup $\Gamma \subset G$, is P_{θ} -Anosov if and only if an(y) orbit of every geodesic in the Cayley graph of Γ with respect to a word metric is a undistorted θ -uniformly regular sequence (Theorem 5.5; [14]).

In this paper, we consider relative P_{θ} -Anosov groups and P_{θ} -transverse groups, which generalize P_{θ} -Anosov groups. Undistorted uniformly regular sequences might seem like a good candidate to capture the hyperbolic-like behaviors in these subgroups. However, we will show that once we leave the Anosov setting, such sequences are rarely found. To make "rarely found" precise, we use Patterson–Sullivan measures on the flag manifold \mathcal{F}_{θ} associated to P_{θ} .

Theorem (Theorem 4.1). For a P_{θ} -transverse subgroup Γ satisfying certain conditions, there exists a Patterson–Sullivan measure μ associated to Γ . If Γ is not P_{θ} -Anosov, then the set of endpoints of the undistorted θ -uniform regular sequences in Γ has measure zero with respect to μ .

A quick way to see the theorem hold for relative P_{θ} -Anosov subgroups is that because a generic geodesic in X can spend arbitrarily large amount of time in horoballs centered at bounded parabolic limit points, the undistorted θ -uniform regular sequences are rare in such subgroups.

The lack of Morse directions also occurs in general CAT(0) metric spaces. In order to capture hyperbolic-like behaviors in this setting, Qing–Rafi–Tiozzo [19] defined sublinearly Morse geodesics, and constructed a new class of boundaries for CAT(0) spaces, and later for general proper metric spaces [18], called *sublinearly Morse boundary*, consisting of endpoints of the sublinearly Morse geodesics. Gekhtman–Qing–Rafi [10] proved this boundary has full measure in several sensible measures on the visual boundary (the Patterson–Sullivan measure, stationary measure of random walk etc.), when the CAT(0) space has a rank one axis (i.e. a geodesic fixed by an element in the isometry group that does not bound any flat).

Inspired by the Qing–Rafi–Tiozzo construction, we generalize the Kapovich–Leeb–Porti notion of uniformly regular sequences to what we call *sublinearly Morse sequences*.

Throughout the paper, we will work under the case that G is a connected semi-simple real Lie group of non-compact type. Let θ be a subset of the simple roots, and P_{θ} the parabolic subgroup associated with this choice of θ (the precise definition can be found in Section 2.2, Definition 2.4). Let $X \coloneqq G/K$, where K is a maximal compact subgroup of G, be the symmetric space associated to G, κ be a Cartan projection, and d_X a Riemannian symmetric distance on X.

Definition 1.2 (P_{θ} -Sublinearly Morse sequence). A sequence $\{g_n\}_n \subset G$ is called P_{θ} -sublinearly Morse if:

(1) there exists some sublinear function η such that

$$d_X(g_nK,g_{n+1}K) \le \eta(d_X(g_0K,g_nK));$$

- (2) there exists a constant $C \ge 1$ and a sublinear function $\bar{\eta} = O(\eta)$, such that the path obtained by concatenating the geodesic segments $(\bigcup_i [g_i K, g_{i+1}K])$ is a $(C, \bar{\eta})$ sublinear ray based at $g_0 K$ (see Definition 2.12);
- (3) there exist a > 0 and a sublinear function $\eta' = O(\eta)$, such that for all n and $\alpha \in \theta$,

$$\alpha(\kappa(g_n^{-1}g_{n+i})) \ge a \cdot d_X(g_nK, g_{n+i}K) - \eta'(d_X(g_0K, g_{n+i}K)).$$

We will explain in Section 2.1 in more details of the meaning of symbols appearing in this definition, but for readers familiar with the structure of the Lie algebra associated to such G, condition (3) is equivalent to requiring the Cartan projection of the sequence to stay away from certain walls of the Weyl chamber of the Cartan sub-algebra associated to elements in θ .

Moreover, in Section 8 we prove a geometric interpretation of the sublinearity of a P_{θ} sublinearly Morse sequence, of which we called the sublinearly Morse lemma in higher rank
symmetric spaces. Let V be the Weyl cone defined by g_0K and the "end point" of the
sequence (we will make this precise in Section 5 and 8).

Theorem (Corollary 8.10). If $\{g_n\} \subset G$ is P_{θ} -sublinearly Morse, then $\{g_n\}$ stays in a sublinear neighborhood of V. It is worth noting that there is a sublinearly Morse lemma [17, Lemma 3.4] proven by Pallier in hyperbolic spaces. One can see that our Theorem 8.8 is a generalization of this result because condition (3) in Definition 1.2 becomes vacuous in rank one setting, our sublinearly Morse sequences then coincide with the sublinear rays that Pallier worked with.

The definition of sublinearly Morse sequences gives rise to a well-defined boundary in the partial flag manifold \mathcal{F}_{θ} . Our main theorem is that for a Patterson–Sullivan measure defined by a P_{θ} -transverse subgroup Γ of G on the limit set $\Lambda_{\theta}(\Gamma) \subset \mathcal{F}_{\theta}$, the subset consisting P_{θ} -sublinearly Morse points in the boundary has full measure. The exact statement of the main theorem requires specifying some technical conditions. However, there is one corollary (see Theorem 7.3) for readers familiar with *relative* P_{θ} -Anosov subgroups.

Corollary (Theorem 7.3). Let (Γ, \mathcal{P}) be a relatively P_{θ} -Anosov pair in G. Let μ be a Patterson–Sullivan measure of certain dimension supported on $\Lambda_{\theta}(\Gamma) \subset \mathcal{F}_{\theta}$. Then the P_{θ} -sublinearly Morse boundary associated to Γ has full measure in the limit set with respect to μ .

The corollary shows that sublinearly Morse points are abundant in the limit set in this more general setting, in contrast to Theorem 4.1.

Acknowledgments. This project was partially supported by grants DMS-2105580 and DMS-2104381 from the National Science Foundation. I would like to thank my advisor, Andrew Zimmer, for his support. I would also like to thank Mitul Islam, Gabriel Pallier, Joan Porti, Yulan Qing, and Feng Zhu for the helpful and encouraging discussions we exchanged.

2. Settings and Preliminaries

Let X be a higher rank symmetric space of non-compact type with a Riemannian symmetric distance d_X . By a theorem of Cartan, X can be seen as the quotient of a connected semi-simple Lie group G by a maximal compact Lie subgroup K that is invariant under a Cartan involution. Note that G caan be supposed to be the identity component of Isom(X).

2.1. Lie group theory. Let \mathfrak{g} and \mathfrak{k} be the Lie algebras associated to G and K respectively. Fix the Cartan involution τ of \mathfrak{g} associated to K, it gives rise to the Cartan decomposition

$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$

where \mathfrak{k} and \mathfrak{p} are the 1 and -1 eigenspaces of τ respectively.

There is a maximal abelian subspace $\mathfrak{a} \subset \mathfrak{p}$ with respect to the Lie bracket, called a Cartan subalgebra. The dimension of \mathfrak{a} equals the rank of the corresponding symmetric space, throughout the paper we assume $rank(X) = dim(\mathfrak{a}) \geq 2$.

For a linear functional $\alpha \in \mathfrak{a}^*$, the *root space* associated to α is

$$g_{\alpha} \coloneqq \{x \in \mathfrak{g} \mid [y, x] = \alpha(y)x \text{ for all } y \in \mathfrak{a}\},\$$

and the root system associated to \mathfrak{a} to be

$$\Sigma := \{ \alpha \in \mathfrak{a}^* \mid \alpha \neq 0 \text{ and } \mathfrak{g}_\alpha \neq 0 \}.$$

Then

$$\mathfrak{g} = \mathfrak{g}_0 \oplus \left(\bigoplus_{\alpha \in \Sigma} \mathfrak{g}_\alpha \right)$$

is called the *Cartan decomposition* of \mathfrak{g} associated to \mathfrak{a} and the corresponding root system Σ . Note that by fixing a $H_0 \in \mathfrak{a} - \bigcup_{\alpha \in \Sigma} \ker(\alpha)$, one can split Σ into a positive and a negative part:

$$\Sigma^+ := \{ \alpha \in \Sigma \mid \alpha(H_0) > 0 \} \text{ and } \Sigma^- := -\Sigma^+.$$

Let $\Delta \subset \Sigma^+$ denote the set of simple roots. The kernels of $\alpha \in \Delta$ divide the Cartan subalgebra into Weyl chambers. We can also view these kernels as the orthogonal complements of certain elements in \mathfrak{a} with respect to the non-degenerate bilinear form $\langle ., . \rangle$ on \mathfrak{a} . More specifically, for each $\alpha \in \Delta$, we can find $H_{\alpha} \in \mathfrak{a}$ such that for all $X \in \mathfrak{a}$

$$\langle H_{\alpha}, X \rangle = \alpha(X).$$

Then we can define

$$H'_{\alpha} \coloneqq \frac{H_{\alpha}}{\langle H_{\alpha}, H_{\alpha} \rangle}$$

to be the coroot corresponding to α , and

$$\omega_{\alpha}(H_{\beta}') \coloneqq \begin{cases} 1 & \alpha = \beta \\ 0 & \alpha \neq \beta \end{cases}$$

the fundamental weight in \mathfrak{a} associated to α .

We can pick a model Weyl chamber \mathfrak{a}^+ in the Cartan subspace, i.e.

$$\mathfrak{a}^{+} \coloneqq \{ x \in \mathfrak{a} \mid \alpha(x) \ge 0 \text{ for all } \alpha \in \Delta \}$$
$$= \{ x \in \mathfrak{a} \mid \langle H_{\alpha}, x \rangle \ge 0 \text{ for all } \alpha \in \Delta \},$$

and define the corresponding Weyl group

$$W \coloneqq N_K(\mathfrak{a})/Z_K(\mathfrak{a}),$$

where $N_K(\mathfrak{a})$ is the stablizer of \mathfrak{a} in K, and $Z_K(\mathfrak{a})$ is the centralizer of \mathfrak{a} in K. W acts on \mathfrak{a} , and each orbit of W intersects \mathfrak{a}^+ exactly once. The Weyl group also acts on the set of Weyl chambers transitively. In W, there is an element w_0 such that $w_0(\mathfrak{a}^+) = -\mathfrak{a}^+$. Using w_0 we can define a map $\iota : \mathfrak{a} \to \mathfrak{a}$ called the *opposite involution*, with $\iota(x) = -w_0.x$.

Fact 2.1. ι induces a dual map on \mathfrak{a}^* , denoted by ι^* , and $\iota^*(\Delta) = \Delta$.

We can now fix a KAK decomposition of an element in G such that

$$g = m_g e^{\kappa(g)} l_g,$$

where $\kappa : G \to \mathfrak{a}^+$ is the *Cartan projection* that sends g to the unique element in the model chamber \mathfrak{a}^+ such that the above decomposition hold, and m, l are elements in K.

Fact 2.2. $\iota(\kappa(g)) = \kappa(g^{-1})$ for all $g \in G$.

The Cartan projection defines a vector-valued distance d_{Δ} on the symmetric space X in the following manner: for $gK, hK \in X$, let

$$d_{\Delta}(gK, hK) \coloneqq \kappa(h^{-1}g).$$

One thing to keep in mind is that this metric is not necessarily symmetric as $\kappa(h^{-1}g) = \iota(\kappa(g^{-1}h))$ which does not necessarily equal to $\kappa(g^{-1}h)$. Nevertheless, this distance satisfies triangle inequalities [13, Equation 2.7] if we fix $||\cdot||$ to be a *W*-invariant norm on the Cartan subalgebra viewed as a vector space,

$$\|d_{\Delta}(xK, yK) - d_{\Delta}(xK, y'K)\| \le \|d_{\Delta}(yK, y'K)\| = c \cdot d_X(yK, y'K)$$

and

$$\|d_{\Delta}(xK, yK) - d_{\Delta}(x'K, yK)\| \le \|d_{\Delta}(xK, x'K)\| = c \cdot d_X(xK, x'K)$$

for some $c \ge 0$.

One should note that the Cartan projection is invariant under action of K on both side, so the distance is well-defined under the choice of representatives $g, h \in G$ of elements gK, hKin the symmetric space X. Moreover, by composting d_{Δ} and $\|\cdot\|$ we get a distance that is equivalent to the Riemannian distance d_X we fixed at the beginning on X = G/K. To abuse notation, we will omit the K when writing the distance between two cosets, and assume $d_X(g,h) = \|d_{\Delta}(g,h)\|.$

Remark 2.3. In the case of $PSL(d, \mathbb{R})$, if we fix $K = PSO(d, \mathbb{R})$, then the maximal flat based at the identity matrix can be realized as the space of diagonal matrices with positive entries in $PSL(d, \mathbb{R})$. The Cartan projection in this case is composed of the singular values of the matrix, i.e. for all $g \in PSL(d, \mathbb{R})$

$$\kappa(g) = [\log \sigma_1(g), ..., \log \sigma_d(g)] \in \mathfrak{a}^+,$$

where the singular values are ordered from the largest to the smallest.

The set $\Delta = \{\alpha_k\}$ where $k \in \{1, ..., d-1\}$ consists of simple roots in the form of

$$\alpha_k(\kappa(g)) = \log \frac{\sigma_k}{\sigma_{k+1}}(g),$$

and the corresponding fundamental weights satisfy

$$\omega_k(\kappa(g)) = \log \prod_{i=1}^k \sigma_i(g),$$

and the opposite involution ι^* sends α_k to α_{d-k} .

In this paper, we will look at a subset $\theta \subset \Delta$ instead of the whole Δ . Such θ is called symmetric if $\iota^*(\theta) = \theta$. We can define \mathfrak{a}_{θ} and \mathfrak{a}_{θ}^+ correspondingly:

$$\mathfrak{a}_{\theta} \coloneqq \{ x \in \mathfrak{a} \mid \alpha(x) = 0 \text{ for all } \alpha \notin \theta \}$$
$$\mathfrak{a}_{\theta}^{+} \coloneqq \{ x \in \mathfrak{a}_{\theta} \mid \alpha_{k}(x) > 0 \text{ for all } k \in \theta \},$$

and the dual of \mathfrak{a}_{θ} , \mathfrak{a}_{θ}^* , can be seen as:

$$\mathfrak{a}_{\theta}^* = \operatorname{Span}\{\omega_{\alpha} \text{ for } \alpha \in \theta\} \subset \mathfrak{a}^*.$$

where ω_{α} are the fundamental weights associated to the coroot H'_{α} of $\alpha \in \theta$.

Moreover, there exist a unique map p_{θ} projecting **a** onto \mathbf{a}_{θ} such that

$$\omega_{\alpha}(p_{\theta}(x)) = \omega_{\alpha}(x)$$

for all $x \in \mathfrak{a}$ and all $\alpha \in \theta$. With this partial Cartan projection we have $\phi(p_{\theta}(x)) = \phi(x)$ for all $\phi \in \mathfrak{a}_{\theta}^{*}$, this will be useful when we define Patterson–Sullivan measures on the partial flag manifold in Section 2.3.

2.2. Parabolic Subgroup and Flag Manifold. In order to define the partial flag manifold associated to θ , we first need to define the corresponding *parabolic subgroup* P_{θ} .

Definition 2.4 (Parabolic Subgroup). Let

$$\mathfrak{u}_{\theta} \coloneqq \bigoplus_{\alpha \in \Sigma_{\theta}^{+}} \mathfrak{g}_{\alpha}$$

where $\Sigma_{\theta}^{+} \coloneqq \Sigma^{+} \setminus \text{Span}(\Delta \setminus \theta)$. The normalizer of \mathfrak{u}_{θ} in *G*, denoted by P_{θ} , is called the *parabolic* subgroup associated to θ .

Similarly, one can also define the parabolic subgroup *opposite* to θ , denoted by P_{θ}^{opp} , as the normalizer of

$$\mathfrak{u}_{\theta}^{-} \coloneqq \bigoplus_{\alpha \in \Sigma_{\theta}^{+}} \mathfrak{g}_{-\iota^{*}(\alpha)}$$

This gives rise to a partial flag manifold $\mathcal{F}_{\theta} \coloneqq G/P_{\theta}$, and a partial flag manifold opposite to θ , $\mathcal{F}_{\theta}^{opp} \coloneqq G/P_{\theta}^{opp}$. Note that if we pick a representative k_0 of the element with longest word length in the Weyl group W, then

$$k_0 P_{\theta} k_0^- = k_0^- P_{\theta} k_0 = P_{\iota^*(\theta)}^{opp}$$

Let us now assume that θ is symmetric from now on, we can then identify \mathcal{F}_{θ} with $\mathcal{F}_{\theta}^{opp}$ by sending mP_{θ} to $mk_0P_{\theta}^{opp}$. Using the relation above, one can verify easily that the map is well defined, i.e. it is independent of the choice of the representative m because if $g^{-1}m$ belongs to P_{θ} , then $k_0^{-1}g^{-1}mk_0 \in k_0^{-1}P_{\theta}k_0 = P_{\iota(\theta)}^{opp} = P_{\theta}^{opp}$.

Moreover, we can define a map U_{θ} that maps every $g \in G$ into a partial flag

$$U_{\theta}(g) = m_g P_{\theta},$$

where $m_g \in K$ is as in the fixed KAK decomposition of g. Note U_{θ} is uniquely defined if $\alpha(\kappa(g)) > 0$ for all $\alpha \in \theta$.

Remark 2.5. In $PSL(d, \mathbb{R})$, the parabolic subgroups correspond to block upper triangular matrices, and the partial flag manifold \mathcal{F}_{α_k} can be identified with the Grassmannians $Gr_k(\mathbb{R}^d)$ in \mathbb{R}^d . The map $U_k(g) \coloneqq U_{\alpha_k}(g) = m < e_1, ..., e_k >$ is sending group elements in G to a k dimensional subspace in \mathbb{R}^d .

There are certain type of discrete subgroups of G whose action on X would give rise to a well defined limit set, namely P_{θ} -divergent groups. Let $\Gamma \subset G$ be a discrete subgroup.

Definition 2.6 (P_{θ} -divergent). Γ is P_{θ} -divergent if

$$\min_{\alpha\in\theta}\alpha(\kappa(g_n))\to+\infty$$

holds for any sequence $\{g_n\} \subset \Gamma$ of pairwise distinct elements.

$$\Lambda_{\theta}(\Gamma) \coloneqq \left\{ \xi \in \mathcal{F}_{\theta} \mid \exists \{g_n\} \subset \Gamma \text{ and } \lim_{n} U_{\theta}(g_n) = \xi \right\}.$$

Two distinct elements gP_{θ} and hP_{θ} in the partial flag manifold \mathcal{F}_{θ} are *transverse* if $(gP_{\theta}, hk_0P_{\theta}^{opp})$ lies in the orbit of $(P_{\theta}, P_{\theta}^{opp})$ under the action of G. With this, we can define:

Definition 2.7 (P_{θ} -transverse). A discrete subgroup $\Gamma \subset G$ is P_{θ} -transverse if it is P_{θ} -divergent, and any two distinct points in the limit set $\Lambda_{\theta}(\Gamma)$ are transverse.

Remark 2.8. For $\Gamma \subset PSL(d, \mathbb{R})$, Γ is P_{α_k} -divergent (and P_{α_k} -transverse) if and only if it is $P_{\alpha_{d-k}}$ -divergent (and $P_{\alpha_{d-k}}$ -transverse respectively), so for simplicity we will assume $k \leq d-k$ and use P_k -divergent (P_k -transverse) to denote $P_{\alpha_k,\alpha_{d-k}}$ -divergent ($P_{\alpha_k,\alpha_{d-k}}$ -transverse).

2.3. **Patterson–Sullivan Measures.** Given a discrete subgroup Γ of G, $\theta \subset \Delta$, and $\phi \in \mathfrak{a}_{\theta}^*$, we can define the *critical exponent* associated to ϕ , denoted by δ_{ϕ} , as

$$\delta_{\phi} \coloneqq \inf \left\{ s \in \mathbb{R}^+ : Q_{\Gamma}^{\phi}(s) < \infty \right\},\$$

where the sum $Q^{\phi}_{\Gamma}(s) \coloneqq \sum_{\gamma \in \Gamma} \exp(-s\phi(p_{\theta} \circ \kappa(\gamma))) = \sum_{\gamma \in \Gamma} \exp(-s\phi(\kappa(\gamma)))$ is the Poincaré series associated to Γ and ϕ .

Let $B_{\theta}: G \times \mathcal{F}_{\theta} \to \mathfrak{a}$ be the *partial Iwasawa cocycle* defined in Section 2.1.5 of [7], which is a higher rank analog of the usual Busemann cocycle in hyperbolic geometry. A probability measure μ is a *Patterson–Sullivan measure* if it is supported on $\Lambda_{\theta}(\Gamma)$, and is ϕ -conformal of dimension β , i.e. $\gamma_*\mu$ and μ are absolutely continuous, and for all $F \in \Lambda_{\theta}(\Gamma)$ and $\gamma \in \Gamma$,

$$\frac{d\gamma_*\mu}{d\mu}(F) = e^{-\beta \cdot B_\theta(\gamma^{-1},F)}$$

By Proposition 3.2 in [7], for P_{θ} -divergent groups, there always exists a Patterson–Sullivan measures.

Proposition 2.9. [7, Proposition 3.2] If $\theta \subset \Delta$ is symmetric, $\Gamma \subset G$ is P_{θ} -divergent, $\phi \in \mathfrak{a}_{\theta}^*$ and $\delta_{\phi} < \infty$, then there is a ϕ -Patterson–Sullivan measure μ for Γ of dimension δ_{ϕ} .

The limit set $\Lambda_{\theta}(\Gamma)$ can be used to define several flow spaces on which a Bowen–Margulis– Sullivan measure can be constructed. Kim and Oh [15] considered the space $\tilde{\Omega}_{\Gamma} \coloneqq \Lambda_{\theta}(\Gamma)^{(2)} \times \mathfrak{a}_{\theta}$, where $\Lambda_{\theta}(\Gamma)^{(2)}$ consists of the pairs of partial flags in $\mathcal{F}_{\theta} \times \mathcal{F}_{\theta}^{opp}$ that are transverse. They showed that Γ acts properly discontinuously on $\tilde{\Omega}_{\Gamma}$. After picking a linear functional $\phi \in \mathfrak{a}_{\theta}^{*}$, $\Gamma \setminus \tilde{\Omega}_{\Gamma}$ fibers over a one dimensional flow space $\tilde{\Omega}_{\Gamma,\phi} \coloneqq \Lambda_{\theta}(\Gamma)^{(2)} \times \mathbb{R}$ by sending (ξ, η, v) to $(\xi, \eta, \phi(v))$. The action of Γ descends and is properly discontinuous on $\tilde{\Omega}_{\Gamma,\phi}$, thus we can quotient out the action of Γ and get a flow space $\Gamma \setminus (\Lambda_{\theta}(\Gamma)^{(2)} \times \mathbb{R}) \cong \Omega_{\Gamma,\phi}$ with a natural translational flow defined.

However, we will not focus on this flow space in our paper. Instead, we map the subgroup Γ into the automorphism group of a properly convex domain, and use the geodesic flow there to prove our results. In order to define the flow space and the associated Bowen–Margulia–Sullivan measure, we first need to introduce some terminologies.

2.4. **Properly Convex Domain.** Let $\mathbb{P}(\mathbb{R}^n)$ be the projectivization of $\mathbb{R}^n - \{0\}$. A open subset Ω of $\mathbb{P}(\mathbb{R}^n)$ is called a *convex domain* if for any two points $x, y \in \Omega$, there exists a projective line segment that connects x and y and is fully contained in Ω . Furthermore, we call Ω a *properly convex domain* if there exists a affine chart that fully contains Ω . With such affine chart, we can pick the unique projective line segment connecting x and y that's contained in this chart and use [x, y] to represent it.

There exists a natural distance d_H , called the Hilbert distance, on such Ω defined as follows: let \mathbb{A}^{n-1} be the affine chart that contains Ω and is equipped with the usual Euclidean norm $|\cdot|$, for any two points $x, y \in \Omega$ let a, b be the two points that lies in $\mathbb{P}(\operatorname{span}(x, y)) \cap \partial\Omega$ and are in the order of a, x, y, b, then

$$d_H(x,y) \coloneqq \frac{1}{2} \log \left(\frac{|b-x||y-a|}{|b-y||a-x|} \right).$$

Note that (Ω, d_H) is a proper, complete, geodesic metric space, but is not necessarily uniquely geodesic.

The automorphism group of Ω is

$$Aut(\Omega) \coloneqq \{g \in PGL(n, \mathbb{R}) \text{ with } g\Omega = \Omega\}.$$

Elements in $Aut(\Omega)$ act by isometries on Ω because the cross ratio between any four points is invariant under projective linear transformations.

We have the following fact that relates the Hilbert distance on Ω to the singular values of $g \in Aut(\Omega)$.

Fact 2.10 (Prop. 10.1; [9]). For any $o \in \Omega$, there exists a constant E such that for any $g \in Aut(\Omega)$,

(2.1)
$$\left| d_H(g.o,o) - \frac{1}{2} \log \frac{\sigma_1}{\sigma_n}(g) \right| \le E.$$

Let d_{Haus} denote the Hausdorff distance between subsets of Ω . For $A, B \subset \Omega$

$$d_{Haus}(A,B) \coloneqq \max\{\min_{x \in A} d_H(x,B), \min_{y \in B} (d_H(y,A))\}$$

The following fact gives us a way to bound d_{Haus} .

Fact 2.11. [11, Proposition 5.3] For p_1 , $p_2 \in \Omega$, and q_1 , $q_2 \in \overline{\Omega}$, if

- (1) $q_1, q_2 \in \Omega$, then $d_{Haus}([p_1q_1], [p_2q_2]) \le \max\{d_H(p_1, p_2), d_H(q_1, q_2)\};$
- (2) $q_1 = q_2 \in \partial \Omega$, then $d_{Haus}([p_1, q_1), [p_2, q_2)) \leq d_H(p_1, p_2)$.

These estimates will be useful in proving our main theorem.

2.5. Sublinear Rays. A sublinear function $\eta:[0,\infty) \to \mathbb{R}$ is a function such that

$$\lim_{t \to \infty} \frac{\eta(t)}{t} = 0$$

Definition 2.12 (Sublinear Ray). A path $c : [0, \infty) \to X$ is a (C, η) sublinear ray if exists a constant $C \ge 1$ and a sublinear function η such that for all $s, t \in [0, \infty)$

$$\frac{1}{C}|s-t| - \eta(\max(s,t)) \le d_X(c(s),c(t)) \le C|s-t| + \eta(\max(s,t)).$$

Remark 2.13. Without loss of generality, we can assume that η is non decreasing and concave (Section 2.2.1; [10]), that is

$$\eta(a) \leq \eta(b)$$
, for $a \leq b$

and

$$(1-t)\eta(s) + t\eta(r) \le \eta((1-t)s + tr).$$

In particular, the second equation has an useful derivation

$$\eta(as) \le a\eta(s)$$
, for $a > 1$.

3. TRANSVERSE REPRESENTATIONS

Let G be the semi-simple Lie group defined in Section 2, and Γ a discrete subgroup of G. A theorem in Canary–Zhang–Zimmer [7] states that whenever Γ is P_{θ} -transverse, we can push everything to a properly convex domain Ω . To state the theorem precisely, we need to define the following things.

Let $\Gamma_0 \subset Aut(\Omega)$ be a discrete subgroup. The *full orbital limit set*, denoted by $\Lambda_{\Omega}(\Gamma_0)$, is defined to be the accumulation points of all orbits of Γ_0 in Ω . A subgroup in $Aut(\Omega)$ is *projectively visible* if for any two points $x, y \in \Lambda_{\Omega}(\Gamma_0)$, the open projective line segment (x, y)is fully contained in Ω , and every point in $\Lambda_{\Omega}(\Gamma_0)$ is a C^1 -smooth point of $\partial\Omega$.

A representation $\rho : \Gamma_0 \to G$ is P_{θ} -transverse if it induces a continuous ρ -equivariant embedding $\xi : \Lambda_{\Omega}(\Gamma_0) \to \mathcal{F}_{\theta}$ such that $\xi(\Lambda_{\Omega}(\Gamma_0))$ is a transverse subset of \mathcal{F}_{θ} and for all sequences $\{g_n\} \subset \Gamma_0$ with $g_n(a) \to x \in \Lambda_{\Omega}(\Gamma_0)$ and $g_n^{-1}(a) \to y \in \Lambda_{\Omega}(\Gamma_0)$ for $a(ny) \ a \in \Omega$, we have $\rho(g_n)(F) \to \xi(x)$ for all $F \in \mathcal{F}_{\theta}$ transverse to $\xi(y)$.

Theorem 3.1 ([6, Theorem 4.2]; [7, Theorem 6.2]). Suppose G has trivial center, $\theta \in \Delta$ is symmetric and P_{θ} contains no simple factors of G. If $\Gamma \subset G$ is P_{θ} -transverse, then there exist $D \in \mathbb{N}$, and $\Omega \subset \mathbb{P}(\mathbb{R}^{D})$ a properly convex domain, $\Gamma_{0} \subset Aut(\Omega)$ a projectively visible subgroup, and $\rho \colon \Gamma_{0} \to G$ a faithful P_{θ} -transverse representation such that:

- (1) $\rho(\Gamma_0) = \Gamma;$
- (2) $\min_{\alpha \in \theta} \alpha(\kappa(\rho(g))) = \log \frac{\sigma_1}{\sigma_2}(g)$ for all $g \in \Gamma_0$;
- (3) the limit map $\xi : \Lambda_{\Omega}(\Gamma_0) \to \Lambda_{\theta}(\Gamma)$ induced by ρ is a homeomorphism;
- (4) Fix a base point $o \in \Omega$, then there exists $L \ge 1$, $l \ge 0$ such that for all $g \in \Gamma_0$

$$\frac{1}{L} \cdot d_X(\rho(g), e) - l \le d_H(g.o, o) \le L \cdot d_X(\rho(g), e) + l$$

Proof of Theorem 3.1. We first map G into a linear group $PSL(d, \mathbb{R})$ by the following proposition.

Proposition 3.2. [7, Proposition B.1] There exist $d \in \mathbb{N}$, a linear representation $\Phi : G \to SL(d,\mathbb{R})$, and a Φ -equivariant smooth embedding $\xi_{\Phi} : \mathcal{F}_{\theta} \to \mathcal{F}_{1,d-1}(\mathbb{R}^d)$ such that:

- (1) F_1, F_2 are transverse in \mathcal{F}_{θ} if and only if $\xi_{\Phi}(F_1)$ and $\xi_{\Phi}(F_2)$ are;
- (2) $\log \frac{\sigma_1}{\sigma_2}(\Phi(g)) = \min_{\alpha \in \theta} \alpha(\kappa(g))$ for all $g \in G$;
- (3) If $\alpha(g) > 0$ for all $\alpha \in \theta$, then

$$\xi_{\Phi}(U_{\theta}(g)) = U_{1,d-1}(\Phi(g));$$

(4) $\Gamma \subset G$ is P_{θ} -transverse if and only if $\Phi(\Gamma)$ is $P_{1,d-1}$ -transverse, and ξ_{Φ} induces a homeomorphism between $\Lambda_{\theta}(\Gamma)$ and $\Lambda_{1,d-1}(\Phi(\Gamma))$.

The assumptions that G has trivial center and P_{θ} does not contain any simple factor of G assure the injectiveness of Φ , so Φ is faithful.

By property (4) in Proposition 3.2, $\Phi(\Gamma)$ is $P_{1,d-1}$ -transverse. Following the construction of Theorem 4.2 in [6], there exist a properly convex domain $\Omega \subset \mathbb{P}(\mathbb{R}^D)$, a projectively visible subgroup $\Gamma_0 \subset Aut(\Omega) \subset PSL(D,\mathbb{R})$, and a faithful representation $\bar{\rho} : \Phi(\Gamma) \to Aut(\Omega)$ such that:

- (a) $\bar{\rho}(\Phi(\Gamma)) = \Gamma_0;$
- (b) $\alpha_1(\kappa(\bar{\rho}(\Phi(g)))) = \alpha_1(\kappa(\Phi(g)))$ for all $g \in \Gamma$;
- (c) $\bar{\rho}$ induces a limit map $\xi_{\bar{\rho}} : \mathcal{F}_{1,d-1} \to \mathcal{F}_{1,D-1}$ such that ξ is a homeomorphism between the limit sets $\Lambda_{1,d-1}(\Phi(\Gamma))$ and $\Lambda_{1,D-1}(\Gamma_0)$, $\bar{\rho}$ -equivariant, and sends transverse pairs in $\Lambda_{1,d-1}(\Phi(\Gamma))$ to transverse pairs in $\Lambda_{1,D-1}(\Gamma_0)$.

By property (b) of $\bar{\rho}$ and Remark 2.8, Γ_0 is P_1 -transverse. According to Lemma 3.4 in [6], $\pi_1(\Lambda_{1,D-1}(\Gamma_0)) = \Lambda_{\Omega}(\Gamma_0)$, where π_1 is projecting onto the first component in the flag manifold $\mathcal{F}_{1,D-1}$. Moreover, π_1 has a well defined inverse on $\Lambda_{\Omega}(\Gamma_0)$

$$\pi_1^{-1}(x)$$
 = $(x, T_\Omega x)$

where $T_{\Omega}x$ is the unique supporting hyperplane of Ω at x (the uniqueness is guaranteed because Γ_0 is projectively visible). Composing π_1 with $\xi_{\bar{\rho}}$ we get a limit map from $\Lambda_{1,d-1}(\Phi(\Gamma))$ to $\Lambda_{\Omega}(\Gamma_0)$ that is a homeomorphism, $\bar{\rho}$ -equivariant, and send transverse pairs in $\Lambda_{1,d-1}(\Phi(\Gamma))$ to distinct points in $\Lambda_{\Omega}(\Gamma_0)$.

Let $\phi = \bar{\rho} \circ \Phi|_{\Gamma}$, this gives us a faithful representation of Γ into $Aut(\Omega)$ with image $\rho(\Gamma) = \Gamma_0$. Since ϕ is bijective, we can take

$$\rho \coloneqq (\phi|_{\Gamma})^{-1}$$

and show that ρ satisfies all properties in Theorem 3.1.



Condition (1) is satisfied by the construction of ρ , and condition (2) is true by combining property (2) of Φ and property (b) of $\bar{\rho}$. The limit map induced by ρ is of the form $\xi := \xi_{\phi}^{-1} \circ \xi_{\bar{\rho}}^{-1} \circ \pi_1^{-1}$. By construction, ξ is ρ -equivariant, and sends distinct pair of points in $\Lambda_{\Omega}(\Gamma_0)$ to transverse flags in \mathcal{F}_{θ} , hence ξ is a P_{θ} -transverse representation. ξ is also a homeomorphism between $\Lambda_{\Omega}(\Gamma_0)$ and $\Lambda_{\theta}(\Gamma)$ as $\xi_{\phi}^{-1}, \xi_{\bar{\rho}}^{-1}, \pi_1^{-1}$ are, hence condition (3) is satisfied.

For any $g \in \Gamma_0$, $\log \frac{\sigma_1}{\sigma_D}(g)$ is bi-Lipschitz to $\log \frac{\sigma_1}{\sigma_d}(\bar{\rho}^{-1}(g))$ by construction in [6] and $\log \frac{\sigma_1}{\sigma_d}(\bar{\rho}^{-1}(g))$ is bi-Lipchitz to $d_X(\rho(g), e)$. The latter is because Φ is injective, we can pull back the Riemannian symmetric metric on the tangent bundle over $PSL(d, \mathbb{R})/PSO(d, \mathbb{R})$ that induces the distance $\log \frac{\sigma_1}{\sigma_d}$ to get another Riemannian metric on the tangent bundle

of X, which has to be equivalent to the one that induces d_X . Substituting $\log \frac{\sigma_1}{\sigma_D}(g)$ with $d_X(\rho(g), e)$ in Fact 2.10 shows ρ satisfies property (4), and hence proves the theorem. \Box

3.1. Bowen–Margulis–Sullivan Measure. Fix $\Gamma \subset G$ a P_{θ} -transverse subgroup. Let Ω and Γ_0 be as in Theorem 3.1. Once we equip Ω with the Hilbert distance d_H , and focus only on the *straight* geodesics, i.e. the geodesics formed by intersecting projective lines in $\mathbb{P}(\mathbb{R}^n)$ with Ω , there is a natural geodesic flow ψ^t on the unit tangent bundle

 $S\Omega := \{ \text{unit tangent vectors along straight geodesics in } \Omega \}.$

The geodesic flow ψ^t is commutative with the action of $Aut(\Omega)$, hence the flow descends to a geodesic flow $\psi^t_{\Gamma_0}$ on $\Gamma_0 \setminus S\Omega$.

Now in order to put measures on $S\Omega$ and $\Gamma_0 \setminus S\Omega$, we fix a ϕ -Patterson–Sullivan measure μ of dimension δ_{ϕ} on $\Lambda_{\theta}(\gamma)$, and pull it back to $\Lambda_{\Omega}(\Gamma_0)$ by the limit map ξ in Theorem 3.1. Using the Hopf parametrization of

$$S_{\Gamma_0}\Omega \coloneqq \{v \in S\Omega | \pi_{fp}(\psi^{\pm\infty}(v)) \in \Lambda_{\Omega}(\Gamma_0)\}$$
$$\cong (\Lambda_{\Omega}(\Gamma_0) \times \Lambda_{\Omega}(\Gamma_0) \setminus \{(x, y) | x \neq y\}) \times \mathbb{R},$$

we get a Bowen–Margulis–Sullivan measure on $S_{\Gamma_0}\Omega$

$$dm(x,y,t) = e^{-\delta_{\phi}\phi([\xi(x),\xi(y)]_{\theta})} d(\xi_*\bar{\mu})(x) \otimes d(\xi_*\mu)(y) \otimes ds(t)$$

where $[.,.]_{\theta} : \mathcal{F}_{\theta}^2 \to \mathfrak{a}_{\theta}$ satisfies certain conditions to make m a invariant measure under action of Γ_0 (the explicit construction can be found in section 6 of [7]), and $\bar{\mu}$ is a $\bar{\phi}$ Patterson– Sullivan measure of dimension $\delta_{\bar{\phi}}$ with $\bar{\phi} := \iota^*(\phi)$. Notice that $\delta_{\bar{\phi}} = \delta_{\phi}$. Moreover, m is also ψ^t invariant. Then m descends to a ψ^t invariant measure \bar{m} on $\Gamma_0 \backslash S\Omega$.

3.2. Hopf–Tsuji–Sullivan Dichotomy. One important result in Canary–Zimmer–Zhang [7] is the Hopf–Tsuji–Sullivan dichotomy of the action of Γ . In order to state the dichotomy we need to define a few terms first. Γ is ϕ -divergent if the Poincare series diverge at δ_{ϕ} , i.e. $Q_{\Gamma}^{\phi}(\delta_{\phi}) = \infty$. The P_{θ} -transverse subgroup Γ acts on $\Lambda_{\theta}(\Gamma)$ as a convergence subgroup (Proposition 3.3; [6]). With respect to this action, a point $x \in \Lambda_{\theta}(\Gamma)$ is a *conical limit point* if there exist $a \neq b \in \Lambda_{\theta}(\Gamma)$ and a sequence $\{\gamma_n\} \subset \Gamma$ such that $\gamma_n(x)$ converges to a, and $\gamma_n(y)$ converges to b for all $y \in \Lambda_{\theta}(\Gamma) \setminus \{x\}$. The *conical limit set* $\Lambda_{\theta}^{con}(\Gamma) \subset \Lambda_{\theta}(\Gamma)$ is the set of all conical limit points of the convergence group action.

Theorem 3.3. [7, Theorem. 8.1, Theorem. 10.1, Theorem 11.1] Let $\Omega \subset \mathbb{P}(\mathbb{R}^D)$, Γ_0 , ρ be as defined in Theorem 3.1. Let $\phi \in \mathfrak{a}_{\theta}^*$, and μ and $\bar{\mu}$ be some Patterson–Sullivan measures for $\Gamma = \rho(\Gamma_0)$ of dimension δ_{ϕ} associated to ϕ and $\phi \circ \iota$ respectively, let \bar{m} be the Bowen–Margulis–Sullivan measure on $\Gamma_0 \backslash S_{\Gamma_0} \Omega$ associated to μ and $\bar{\mu}$. We have the following dichotomy:

- (1) If $Q^{\phi}_{\Gamma}(\delta_{\phi}) = +\infty$, then the action of the geodesic flow ψ^t on $\Gamma_0 \setminus S_{\Gamma_0}\Omega$ is ergodic with respect to \bar{m} , and $\mu(\Lambda^{con}_{\theta}(\Gamma)) = \bar{\mu}(\Lambda^{con}_{\theta}(\Gamma)) = 1$. Moreover, μ and $\bar{\mu}$ have no atom.
- (2) If $Q^{\phi}_{\Gamma}(\delta_{\phi}) < +\infty$, then the action of the geodesic flow ψ^t on $\Gamma_0 \setminus S_{\Gamma_0}\Omega$ is non-ergodic with respect to \bar{m} , and $\mu(\Lambda^{con}_{\theta}(\Gamma)) = 0$.

One may notice that the theorem above did not require \bar{m} to be finite. In this case, ergodicity of the geodesic flow means that invariant set has zero measure or the complement of it has zero measure.

4. The Morse Boundary

Kapovich–Leeb–Porti proved a higher rank Morse lemma ([13, Theorem 5.16, Corollary 5.23]) which states that, for a discrete subgroup $\Gamma \subset G$, any geodesic ray in the Cayley graph of $\Gamma =: Cay(\Gamma)$ whose orbit in X is undistorted and θ –uniformly regular has to stay uniformly close to a Weyl cone. Such sequence *flag converges conically* to the unique element in the flag manifold \mathcal{F}_{θ} that defines the associated Weyl cone ([12, Definition 2.58]).

This allows us to define a Morse boundary associated to Γ , denoted by $\partial_{M,\theta}(\Gamma)$ as follows: a point $\zeta \in \Lambda_{\theta}(\Gamma)$ is a Morse boundary point if there exists a geodesic ray $\{g_n\}_{n\in\mathbb{N}}$ in $Cay(\Gamma)$ and a quasi-isometric embedding $q: \mathbb{N} \to X$ that sends g_n to $g_n K$ and the image forms a θ -uniform regular sequence in X and flag converges conically to ζ . We will again omit K, the cosets. Notice that the quasi isometry condition assures that there exists a constant Csuch that $d_X(g_i, g_{i+1}) \leq C$ for all $i \in \mathbb{N}$.

As discussed in the introduction, we want to show that if Γ is not P_{θ} -Anosov, this boundary has zero measure in the limit set with respect to a Patterson–Sullivan measure μ associated to Γ . We prove the contrapositive statement, which also gives an alternative characterization of Anosov subgroups.

Theorem 4.1. Let Γ be a P_{θ} -transverse subgroup, $\phi \in \mathfrak{a}_{\theta}^*$ with $\delta_{\phi} < \infty$, and μ is a Patterson– Sullivan measure of dimension δ_{ϕ} associated to Γ . If $\mu(\partial_{M,\theta}(\Gamma)) > 0$, then Γ is P_{θ} -Anosov.

Remark 4.2. It is worth noting that when reducing to the rank one case, the Morse boundary $\partial_{M,\theta}(\Gamma)$ of the subgroup Γ we define here does not coincide exactly with the intersection of the conical limit set and the usual Morse boundary of the underlying hyperbolic space X. This is mainly because we require the existence of a quasi-isometrically embedded sequence (regularity is satisfied automatically in rank one case) that converges to $\xi \in \Lambda$ for ξ to be a Morse boundary point. However, such sequences could be rare if Γ does not admit a convex cocompact action on the convex hull of the limit set.

We will use the following characterization of P_{θ} -Anosov groups to prove the theorem.

Fact 4.3. [3, Theorem 8.4] If there exist a, b > 0 such that for every $\gamma \in \Gamma$ we have

(4.1) $\alpha(\kappa(\gamma)) \ge a \cdot |\gamma| - b$

for all $\alpha \in \theta$, and $|\cdot|_S$ a word metric on $Cay(\Gamma)$ associated to a generating set S, then Γ is word hyperbolic and P_{θ} -Anosov.

Our goal is to show Equation (4.1) holds for all $\gamma \in \Gamma$. In order to do this, we need another fact to relate an arbitrary γ to a loxodromic element.

Fact 4.4. [1, Lemma 2.4] Let $H \subset Homeo(M)$ be a convergence group acting on a compact metrizable space M. Then there exist a finite set $F \subset H$ and $\epsilon > 0$ such that for any $\gamma \in H$, there exists $f \in F$ with γf loxodromic, i.e. γf has distinct fixed points in M, denoted $(\gamma f)^{\pm}$, and $d((\gamma f_{\gamma})^{+}, (\gamma f_{\gamma})^{-}) > \epsilon$.

Let Ω , Γ_0 , ρ , and ξ be as defined in Theorem 3.1. Γ_0 is a convergence group acting on the compact space $\partial\Omega$. Moreover, because of property (2) of ρ , Equation (4.1) holds for all $\gamma \in \Gamma_0$ with respect to $\alpha = \alpha_1$ if and only if it holds for all $\rho(\gamma) \in \Gamma = \rho(\Gamma_0)$ with respect to all $\alpha \in \theta$.

For arbitrary $\gamma \in \Gamma_0$, let f_{γ} be the element in F in Fact 4.4 such that γf_{γ} is loxodromic. Because F is finite, there exist a constant L independent of γ , such that

$$\max\left(\max_{f\in F} d_X(\rho(\gamma),\rho(\gamma f)), \max_{f\in F} |f|_{\rho^{-1}(S)}\right) \le L.$$

Moreover, by triangle inequalities and the geometric interpretation of d_{Δ} , γ satisfies Equation (4.1) if and only if γf_{γ} does.

We have reduced the proof of Theorem 4.1 to find lower bounds of $\alpha_1(\kappa(\gamma f_{\gamma}))$. We will do this by approximating γf_{γ} with θ -uniformly regular sequences for $\theta = \{\alpha_1, \alpha_{D-1}\}$.

Now fix a base point $o \in \Omega$. Due to Fact 4.4, there exists a constant $R < +\infty$ such that for all $\gamma \in \Gamma_0$,

$$d_H(o,((\gamma f_{\gamma})^+(\gamma f_{\gamma})^-)) \le R.$$

Let $o_{\gamma} \in ((\gamma f_{\gamma})^+ (\gamma f_{\gamma})^-)$ be the point such that $d_H(o_{\gamma}, o) = d_H(o, ((\gamma f_{\gamma})^+ (\gamma f_{\gamma})^-)), v \in S_{\Gamma_0}\Omega$ be the directional vector based at o_{γ} and point toward $(\gamma f_{\gamma})^+$.

Kapovich–Leeb showed in [12, Theorem 3.18] that when Γ is P_{θ} -transverse, a point in the limit set is conical in the convergence group sense if and only if there exists a sequence in X that flag converges to it conically. By construction, $\partial_{M,\theta}(\Gamma) \subset \Lambda_{\theta}^{con}(\Gamma)$. The assumption that $\mu(\partial_{M,\theta}(\Gamma)) > 0$ in Theorem 4.1 implies $\mu(\Lambda_{\theta}^{con}(\Gamma)) > 0$. By the dichotomy in Theorem 3.3, Γ is ϕ -divergent and the geodesic flow acts ergodically on $\Gamma_0 \setminus S_{\Gamma_0} \Omega$. The ergodicity ensures that for \overline{m} almost every $w \in \Gamma_0 \setminus S_{\Gamma_0} \Omega$, the forward orbit $\{\psi^t(v)\}_{t \in [0,\infty)}$ is dense in $\supp(\overline{m}) \subset \Gamma_0 \setminus S_{\Gamma_0} \Omega$. Because $\mu(\partial_{M,\theta}(\Gamma)) > 0$, there exist a point $g_{\infty} \in \xi^{-1}(\partial_{M,\theta}(\Gamma))$, and a θ -uniform regular sequence $\{\rho(g_n)\}_{n \in \mathbb{N}} \subset \Gamma$ such that $g_n.o \to g_{\infty}$, and the forward orbit of $w \in S_{\Gamma_0} \Omega$ based at o and defines the direction towards g_{∞} , is dense. Then we can find $t_j \in [0, \infty)$ and $a_j \in \Gamma_0$ such that $a_j \psi^{t_j}(w) \to v$ in $S_{\Gamma_0} \Omega$.



Proposition 4.5. There exists $R' < \infty$ such that the orbit $\{g_n.o\}$ stays in the R' neighborhood of $[o, g_{\infty})$.

Proof. First note that the backward limit $(g_n)^{-1} \cdot o \to g_{-\infty}$ of the sequence exists by Remark 2.8, as $\{g_n^{-1}\}$ is also θ -uniform regular. We also need the following lemma:

Lemma 4.6. [3, Lemma 2.5] There exist $l \in \mathbb{N}$ and $\delta > 0$ such that if $n < k < m \in \mathbb{N}$ and $\min\{k - n, m - k\} \ge l$, then

(4.2)
$$\leq \left(U_1(g_k^{-1}g_n), U_{D-1}(g_k^{-1}g_m) \right) \geq \delta.$$

Remark 4.7. In Bochi–Potrie–Sambarino [3], Lemma 4.6 is stated for sequences $\{A_i\}_{i\in I} \in \mathcal{D}(K, 1, \mu, c, I)$. One can show the equivalence between the two statements by setting $A_0 = g_0$, and $A_i = g_{i+1}^{-1}g_i$, and realizing the condition that $(A_i)_{i\in I} \in \mathcal{D}(K, 1, \mu, c, I)$ is equivalent to $\{g_n\}$ being $\{\alpha_1, \alpha_{D-1}\}$ -uniformly regular.

If we take $\min\{k - n, m - k\}$ to go to infinity, then

$$U_1(g_k^{-1}g_n) \to U_1(g_{-\infty}),$$

and

$$U_{D-1}(g_k^{-1}g_m) \to U_{D-1}(g_k^{-1}g_\infty).$$

Let $\zeta \in \Lambda_{\Omega}(\Gamma_0)$ be any limit point of $\{g_k^{-1}g_\infty\}$. Lemma 4.6 assures that $U_1(g_k^{-1}g_m) \in U_{D-1}(g_k^{-1}g_m)$ and $U_1(g_k^{-1}g_n)$ have an uniform angle separation ε for all n < k < m, so $\zeta \neq g_{-\infty}$.

We prove Proposition 4.5 by contradiction. Assume there exist $i \to \infty$ such that

$$d_H(g_i.o, [o, g_\infty)) = d_H(o, [g_i^{-1}.o, g_i^{-1}g_\infty)) \to \infty$$

Upto taking subsequences, we can assume $g_i^{-1}g_{\infty} \to \zeta$, then we have

$$d_H(o, g_i^{-1}.[o, g_\infty)) \rightarrow d_H(o, (g^-, \zeta)) = \infty.$$

This is a contradiction because Γ_0 is a projectively visible group, so $(g^-, \zeta) \subset \Omega$.

Proof of Theorem 4.1. By Proposition 4.5 and the uniform upper bound C on $d_H(g_{i.o}, g_{i+1.o})$, there exists n_j such that $d_H(g_{n_j.o}, \pi_{fp}(\psi^{t_j}(w))) \leq R' + C$, where π_{fp} is the footpoint projection that projects a unit tangent vector to its footpoint in Ω . Moreover, there exist $a_j \in \Gamma_0$ and $o' \in B_{R'+C+R}(o)$ such that $a_j g_{n_j.o} \to o'$ as $j \to \infty$.

Let $d \coloneqq d_H(o_\gamma, \gamma f_\gamma. o_\gamma)$. Since γf_γ preserves $(\gamma f_\gamma)^{\pm}$, $(\gamma f_\gamma^-, \gamma f_\gamma^+)$ is a axis of γf_γ , i.e. $\gamma f_{\gamma}. o_\gamma \in (\gamma f_\gamma^-, \gamma f_\gamma^+)$. Then $\psi^d a_j \psi^{t_j}(w) = a_j \psi^{t_j+d}(w) \to v'$, where v' is the unit tangent vector based at $\gamma f_\gamma. o_\gamma$ and point towards $(\gamma f_\gamma)^+$. Similarly we can find m_j such that $g_{m_j}. o$ is R' + Cclose to the footpoint of $a_j \psi^{t_j+d}(w)$, and $a_j g_{m_j}. o$ converges to $x \in B_{R'+C+R}(\gamma f_\gamma. o)$.

Using triangle inequality of the Cartan projection κ , for j large enough we have

(4.3)
$$\alpha_1(\kappa(\gamma f_{\gamma})) \ge \alpha_1\left(\kappa((a_j g_{n_j})^{-1}(a_j g_{m_j})) - 2E\right)$$
$$= \alpha_1(\kappa(g_{n_j}^{-1} g_{m_j})) - 2E$$
$$\ge a \cdot (m_j - n_j) - (b + 2E)$$

for a, b depending only on the uniform regular sequence $\{g_n\}$, and E > 0 depending only on R', R, and C. Note that $|g_{m_j}^{-1}g_{n_j}| = (m_j - n_j)$ is quasi-isometric to $d_H(a_jg_{m_j}.o, a_jg_{n_j}.o)$, and hence to $d_H(o, \gamma f_{\gamma}.o)$. Thus $(m_j - n_j)$ is bounded from above and below for all j. Moreover,

there exist C' depending only on R' + C + R such that $|a_j g_{n_j}|, |(\gamma f_{\gamma})^{-1} a_j g_{m_j}| \leq C'$ for j large enough, then the equation above turns into

(4.4)

$$\alpha_1(\kappa(\gamma f_{\gamma})) \ge a \cdot |(a_j g_{n_j})^{-1}(a_j g_{m_j})| - (b + 2E)$$

$$\ge a \cdot (|\gamma f_{\gamma}| - |a_j g_{n_j}| - |(\gamma f_{\gamma})^{-1}(a_j g_{m_j})|) - (b + 2E)$$

$$\ge a \cdot |\gamma f_{\gamma}| - b'$$

with b' = 2aC' + b + 2E. Since a, b' and the finite set F are independent of the arbitrary γ we pick, we obtain Equation (4.1) for every $\gamma \in \Gamma_0$ for a uniform a, b. By Fact 4.3, Γ_0 has to be $P_{1,D-1}$ -Anosov, and $\Gamma = \rho(\Gamma_0)$ is P_{θ} -Anosov.

5. The sublinearly Morse Boundary

In this section, let G, X, Γ be as defined before. Let $e = K \in X$ be a base point in X. We use the following theorem to define the P_{θ} -sublinearly Morse boundary associated to Γ .

Theorem 5.1. Given a P_{θ} -sublinearly Morse sequence $\{g_n\} \subset G$, $\{U_{\theta}(g_n)\}$ converges in \mathcal{F}_{θ} .

Definition 5.2 (sublinearly Morse boundary). The sublinearly Morse boundary of a discrete subgroup $\Gamma \subset G$ is:

 $\partial_{SM,\theta}(\Gamma) \coloneqq \{\eta \in \mathcal{F}_{\theta} \mid \exists \{g_n\}_n \subset \Gamma \text{ s.t. } U_{\theta}(g_n) \to \eta \text{ and } \{g_n\}_n \text{ is } P_{\theta}\text{-sublinearly Morse} \}.$

We prove Theorem 5.1 by proving the result for $\Phi(\Gamma) \subset GL(d,\mathbb{R})$, and then draw the connection between the general case and linear case by the following lemma.

Proposition 5.3. For a sequence $\{g_n\}$ in G, then for Φ defined in Proposition 3.2 the image $\{\Phi(g_n)\}$ is P_1 -sublinearly Morse if and only if $\{g_n\}$ is P_{θ} -sublinearly Morse.

Proof. We need to show that the three conditions in Definition 1.2 are satisfied for $\{g_n\}$ and $\{\Phi(g_n)\}$ simultaneously. By property (4) of Theorem 3.1, we can replace $\log \frac{\sigma_1}{\sigma_d}(\Phi(g_n)) =: d_Y(\Phi(g_n), I_d)$ with $d_X(g_n, e)$ up to some multiplicative constants. Since conditions (1) and (2) for sublinear Morseness are purely geometric in terms of the Riemannian distances, and the sublinearity is unaffected by taking scalar multiples of the argument (due to the remark in Section 2.5), they are satisfied for $\{g_n\}$ if and only if the same hold for $\{\Phi(g_n)\}$.

With property (2) in Proposition 3.2, condition (3) in Definition 1.2 turns into

$$\alpha(\kappa(g_0^{-1}g_n)) \ge \alpha_1(\kappa(\Phi(g_0^{-1}g_n))) \ge a \cdot d_Y(\Phi(g_n), \Phi(g_0)) - \eta'(d_Y(\Phi(g_n), \Phi(g_0)))$$

for all $\alpha \in \theta$. Again, by interchanging d_Y with d_X up to the multiplicative constant (i.e. modifying a), we obtain the desired inequality for $\{g_n\}$ and $\{\Phi(g_n)\}$ simultaneously, and this concludes the proof.

Now we can prove Theorem 5.1:

Proof of Theorem 5.1. $\{U_{\theta}(g_n)\}$ converges in \mathcal{F}_{θ} if and only if $\{U_{1,d-1}(\Phi(g_n))\}$ converges in $\mathcal{F}_{1,d-1}$. Thus, it suffices to show $\{U_{1,d-1}(\Phi(g_n))\}$ is Cauchy.

A matrix M has a gap of index k if $\sigma_k(M) > \sigma_{k+1}(M)$. For any M with a gap of index k, the map $U_k(M) \in Gr_k(\mathbb{R}^d)$ is well-defined. Let d_k be the natural angle metric defined on the Grassmannian $Gr_k(\mathbb{R}^d)$.

By Lemma A.4 in [3], for $A, AB \in GL(d, \mathbb{R})$ that have gap of index k, the distance $d_k(U_k(A), U_k(AB))$ is bounded above by $\frac{\sigma_1}{\sigma_d}(B^{-1})\frac{\sigma_{k+1}}{\sigma_k}(A)$. When n is large, $\Phi(g_n)$ has a gap of index 1 (and respectively d-1) by construction. Then assuming n < m and omitting the representation Φ we have

(5.1)

$$d_{1}(U_{1}(g_{n}), U_{1}(g_{m})) \leq \sum_{i=n}^{m-1} d_{1}\left(U_{1}(g_{i}), U_{1}(g_{i}(g_{i})^{-1}g_{i+1})\right)$$

$$\leq \sum_{i=n}^{m-1} \exp(d_{Y}(g_{i+1}^{-1}g_{i})) \cdot \exp(-\alpha_{1}(\kappa(g_{i})))$$

$$\leq \sum_{i=n}^{m-1} \exp(\eta(d_{Y}(g_{0}, g_{i}))) \cdot \exp(-\alpha_{1}(\kappa(g_{0}^{-1}g_{i})) + d_{Y}(g_{0}, e))$$

Then $\{\Phi(g_n)\}$ being P_1 -sublinearly Morse due to Proposition 5.3 and the assumption that $\{g_n\}$ is P_{θ} -sublinearly Morse implies

$$d_1(U_1(g_n), U_1(g_m)) \le \sum_{i=n}^{m-1} C' \cdot \exp\left(-a \cdot d_Y(g_i, g_0) + (\eta + \eta')(d_Y(g_i, g_0))\right)$$

for a, b the constants in the definition of the sublinearly Morse sequence.

In order to see that the partial sum in Equation (5.1) converges, we need to show the distance $d_Y(g_i, g_0)$ grows fast enough, more precisely, it has to grow linearly in terms of *i*.

Claim. There exists a constant d' such that $d_Y(g_i, g_0) \ge d' \cdot i$ when i is large enough.

Proof of claim. Because the concatenated path $\bigcup_i [g_i, g_{i+1}]$ parametrized by arclength is a sublinear ray by definition, we have

$$d_Y(g_i, g_0) \ge \frac{1}{C} \left(\sum_{n=0}^{i-1} d_Y(g_n, g_{n+1}) \right) - \bar{\eta} \left(\sum_{n=0}^{i-1} d_Y(g_n, g_{n+1}) \right).$$

The quantity $\sum_{n=0}^{i-1} d_Y(g_n, g_{n+1})$ is increasing as Γ is discrete: there exits a lower bound $d := \inf_{g \neq h, \in \Gamma} d_Y(g, h) > 0$. For all $\epsilon \in (0, \frac{1}{C})$, there exists *i* large enough, such that the sublinear function $\eta(t)$ is bounded above by $\varepsilon(t)$ for all $t \ge d \cdot i$, hence

$$d_Y(g_i, g_0) \ge \left(\frac{1}{C} - \varepsilon\right) \left(\sum_{n=0}^{i-1} d_Y(g_n, g_{n+1})\right)$$
$$\ge d' \cdot i,$$

with $d' = \left(\frac{1}{C} - \varepsilon\right) \cdot d$.

The claim shows that the partial sum of the exponentials in the last line of Equation (5.1) converges, and since η and η' are sublinear, they do not affect the convergence of the series as n and m tend to infinity.

Note that convergence of the quantity in Equation (5.1) suffices to ensure the convergence of $\{U_{1,d-1}(\Phi(g_n))\}$ because the metric on $\mathcal{F}_{1,d-1} \subset Gr_1(\mathbb{R}^d) \oplus Gr_{d-1}(\mathbb{R}^d)$ is composed of the individual d_i , and hence we proved the proposition.

 \diamond

6. Main Theorem

With all the background knowledge introduced, we can now state our main theorem. Let G be as in Theorem 3.1 and X the symmetric space associated to G.

Theorem 6.1 (Main Theorem). Let Γ be a non-elementary, discrete, P_{θ} -transverse, and ϕ -divergent subgroup of G for some $\phi \in \mathfrak{a}_{\theta}^*$ with $\delta_{\phi} < \infty$. Let μ and $\bar{\mu}$ be a ϕ and $\bar{\phi} := \iota^*(\phi)$ -Patterson–Sullivan measure of dimension δ_{ϕ} supported on $\Lambda_{\theta}(\Gamma) \subset \mathcal{F}_{\theta}$, and the induced Bowen–Margulis–Sullivan measure \bar{m} on $\Gamma_0 \backslash S\Omega$ is finite. Then $\mu(\partial_{SM,\theta}(\Gamma)) = 1$.

For Γ as in the Main Theorem, let us fix ρ , $\Omega \subset \mathbb{P}(\mathbb{R}^D)$, and $\Gamma_0 \subset Aut(\Omega)$ as in Theorem 3.1. Again we omit the representation ρ when the context is clear.

6.1. Equivalent Condition of sublinearly Morseness. Given the identification between $\Lambda_{\Omega}(\Gamma_0)$ and $\Lambda_{\theta}(\Gamma)$ mentioned in Section 3, we can find an equivalent condition of Definition 1.2 for P_{θ} -sublinearly Morse sequences. In order to do this, we need to define several things first. Let C_{Γ_0} denote the convex hull associated to $\Lambda_{\Omega}(\Gamma_0)$. Fix a base point $o \in C_{\Gamma_0} \subset \Omega$.

Definition 6.2 (Compact part of γ). Let $\gamma : [0, \infty) \to \Omega$ be a straight geodesic ray. Fix r > 0 and let $\Gamma_0.B_r(o)$ denote the set of metric balls with radius r around the orbit of o under the action of Γ_0 . Then we can define the *compact part of* γ to be

$$Cpct_{\gamma}(T) \coloneqq \{t \in [0,T] \mid \gamma(t) \in \Gamma_0.B_r(o)\}.$$

Theorem 6.3. If

(6.1)
$$\lim_{T \to \infty} \frac{Leb(Cpct_{\gamma}(T))}{T} = M > 0,$$

then

$$\gamma(\infty) \coloneqq \lim_{t \to \infty} \gamma(t) \in \xi^{-1}(\partial_{SM,\theta}(\Gamma)).$$

Fix γ that satisfies Equation (6.1). We prove the theorem by constructing a P_{θ} -sublinearly Morse sequence that converges to $\xi(\gamma(\infty))$ explicitly.

Definition 6.4. Define:

- (1) $A_{\gamma} \coloneqq \{g \in \Gamma_0 \mid \gamma \cap B_r(g.o) \neq \emptyset\};$
- (2) Let $A_{\gamma,C} \subseteq A_{\gamma}$ be a maximal subset such that for all $g, h \in A_{\gamma,C}, d_H(g.o, h.o) \ge C$.

Remark 6.5. For all $x \in \Omega$, let

$$\pi_{\gamma}(x) \coloneqq \{\gamma(t) \mid d_H(x,\gamma) = d_H(x,\gamma(t))\}$$

be the nearest point projection. Let $t_x := \min_{\gamma(t) \in \pi_{\gamma}(x)} t$. We can put an order on A_{γ} (and $A_{\gamma,C}$ respectively) as follow: for all $g, h \in A_{\gamma}$, if $t_g := t_{g,o} \neq t_h$, then g < h if and only if $t_g < t_h$; if $t_g = t_h = t'$, then put any order on the set $S_{t'} := \{g \in \Gamma_0 \mid t_g = t'\}$. This is turns A_{γ} and $A_{\gamma,C}$ into totally ordered set, as each $S_{t'}$ is finite because Ω is proper and Γ_0 is discrete and acts properly discontinuously on Ω . With this ordering of A_{γ} and $A_{\gamma,C}$, we can index the sets by $A_{\gamma} = \{g_n\}$ and $A_{\gamma,C} = \{g_{n_i}\}_i$ as a subsequence.

Moreover, we can also describe the maximality of $A_{\gamma,C}$ explicitly. Fix $g_{n,0} = g_0 \in A_{\gamma}$ to be the element such that $t_{g_0} = \min_{g \in A_{\gamma}} t_g$. For all $i \in \mathbb{N}$, define $g_{n_{i+1}}$ to be the smallest element in A_{γ} after g_{n_i} with $d_H(g_{n_i}.o, g_{n_{i+1}}.o) \ge C$. We will show that $\{\rho(g_{n_i})\}_i$ is P_{θ} -sublinearly Morse.

Without loss of generality, we can also assume that $t_{g_0} \leq r$, i.e. the geodesic ray starts within $B_r(g_0.o)$. This can be seen by truncating the initial segment of the ray that is not in $B_r(g_0.o)$, and denoting the new geodesic ray by $\bar{\gamma}$. By construction, $Cpct_{\gamma}(T) = Cpct_{\bar{\gamma}}(T - t_{g_0} + r)$, then we have

$$\lim_{T \to \infty} \frac{Leb(Cpct_{\gamma}(T))}{T} \le \lim_{T \to \infty} \frac{Cpct_{\bar{\gamma}}(T - t_{g_0} + r)}{T - t_{g_0} + r} = \lim_{T \to \infty} \frac{Cpct_{\bar{\gamma}}(T)}{T}$$

The above inequality shows that truncating the geodesic ray does not change the positivity of the limit in Equation (6.1).

We break the proof of Theorem 6.3 into three parts.

Proposition 6.6. There exists a sublinear function η such that the sequence $\{g_n\}$ satisfies the equation

(6.2)
$$d_X(g_i, g_{i+1}) \le \eta \left(d_X(g_i, g_0) \right).$$

Proof. Let $t_i := t_{q_i}$ as defined above. By construction we have

$$Leb(Cpct_{\gamma}(t_{i+1})) = Leb(Cpct_{\gamma}(t_i)) + Leb(\{t \in [t_i, t_{i+1}] \mid \gamma(t) \cap A_{\gamma}.B_r(o) \neq \emptyset\})$$

Notice that the second term on the right hand side is at most 2r.

Because the limit in (6.1) goes to M, which is strictly bigger than 0, then for all $\epsilon > 0$, we can find *i* big enough such that

$$Leb(Cpct_{\gamma}(t_{i+1})) \ge (m-\epsilon)t_{i+1}$$

and

$$Leb(Cpct_{\gamma}(t_i)) \leq (M + \epsilon)t_i.$$

This implies

$$(M-\epsilon)t_{i+1} \le (M+\epsilon)t_i + 2r.$$

Hence

$$M(t_{i+1} - t_i) \le \epsilon(t_{i+1} + t_i) + 2r \le 2\epsilon t_{i+1} + 2r$$

and

 $(t_{i+1} - t_i) \le \epsilon' t_{i+1} + r'$

with $\epsilon' = \frac{2\epsilon}{M}$ and $r' = \frac{2r}{M}$. Now, ϵ can be chosen arbitrarily small, then $\eta_0(t_{i+1}) = \epsilon' t_{i+1} + r' + 2r$ is a sublinear function in t_i , and we have

(6.3)
$$d_H(g_{i,o}, g_{i+1,o}) \le (t_{i+1} - t_i) + 2r \le \eta_0(t_{i+1}).$$

Using triangle inequalities, we have

$$t_{i+1} \le d_H(g_0.o, g_{i+1}.o) + 2r.$$

Along with the monotonicity and convexity of the sublinear function, Equation (6.3) turns into

$$d_H(g_{i.o}, g_{i+1.o}) \le \eta_0(d_H(g_{0.o}, g_{i+1.o}) + 2r).$$

One simplification is that any constant added to the argument or of η or the function itself would not affect its sublinearity, i.e. we can find another sublinear function that bounds it above. Denote this new function by η_0 again, and we have

(6.3*)
$$d_H(g_i.o, g_{i+1}.o) \le \eta_0(d_H(g_0.o, g_{i+1}.o))$$

Note that this is close in forms to the relations we want to have in the first condition of the P_{θ} -sublinearly Morseness, but we still need to find ways to related $d_H(g_{0.0}, g_{i+1.0})$ with $d_H(g_{0.0}, g_{i.0})$. To do this we need the following fact:

Fact 6.7. [18, Proposition 3.2] There exists $d_1, d_2 > 0$, depending on η such that for all $x, y \in \Omega$,

$$d_H(x,y) \le \eta(d_H(x,o)) \Rightarrow d_1\eta(d_H(y,o)) \le \eta(d_H(x,o)) \le d_2\eta(d_H(y,o)).$$

We can then rewrite Equation (6.3*) by letting $x = g_0^{-1}g_{i+1}$ o and $y = g_0^{-1}g_i$ o

(6.4)
$$d_H(g_{i.o}, g_{i+1.o}) \le d_2 \bar{\eta}(d_H(g_{i.o}, g_{0.o})).$$

Then, by Fact 2.10

$$\log \frac{\sigma_1}{\sigma_D} (g_{i+1}^{-1} g_i) \le 2E + 2d_2 \bar{\eta} \left(\frac{1}{2} \log \frac{\sigma_1}{\sigma_D} (g_0^{-1} g_i) + E \right)$$
$$\le 2E + 2d_2 \bar{\eta} \left(\log \frac{\sigma_1}{\sigma_D} (g_0^{-1} g_i) + 2E \right).$$

Let $\eta(\log \frac{\sigma_1}{\sigma_D}(g_0^{-1}g_i))$ be a sublinear function that bounds $2E + 2d_2\bar{\eta}(\log \frac{\sigma_1}{\sigma_D}(g_0^{-1}g_i) + 2E)$ from above. This gives us

$$\log \frac{\sigma_1}{\sigma_D} \left(g_{i+1}^{-1} g_i \right) \le \eta \left(\log \frac{\sigma_1}{\sigma_D} \left(g_0^{-1} g_i \right) \right).$$

By construction of ρ in Section 3, $\log \frac{\sigma_1}{\sigma_D}(g)$ is bi-Lipschitz with respect to $d_X(\rho(g), e)$ for all $g \in \Gamma_0$. Substituting this into the equation above proves the proposition.

Proposition 6.6 shows $\{g_n\}$ satisfies condition (1) in Definition 1.2. However, $\{g_n\}$ does not necessarily satisfy condition (2). This can be resolved by looking at the subsequence $\{g_{n_i}\}$, but we then need to check whether $\{g_{n_i}\}$ satisfies Equation (6.2).

Corollary 6.8. If Proposition 6.6 hold for $\{g_n\}$, then there exists another sublinear function $\tilde{\eta} = O(\eta)$, such that Proposition 6.6 is also true for $\{g_{n_i}\}$ with respect to $\tilde{\eta}$.

Proof. By triangle inequality

$$d_{H}(g_{n_{i}}.o, g_{n_{i+1}}.o) \leq \sum_{k=n_{i}}^{n_{i+1}-1} d_{H}(g_{k}.o, g_{k+1}.o)$$

$$\leq \sum_{k=n_{i}}^{n_{i+1}-1} \bar{\eta}(d_{H}(g_{k}.o, g_{0}.o)) \quad \text{by Proposition 6.6}$$

$$\leq \sum_{k=n_{i}}^{n_{i+1}-1} d_{2}^{k-n_{i}} \bar{\eta}(d_{H}(g_{n_{i}}.o, g_{0}.o)) \quad \text{by Fact 6.7.}$$

Let $F := \#\{g \in \Gamma_0 \mid d_H(o, g.o) \leq C\}$. F is finite as Ω is a proper metric space and Γ_0 is discrete and acts properly discontinuously on Ω . Moreover, F + 1 is a uniform upper

bounded on how many elements are omitted in $\{g_i\}$ between consecutive g_{n_i} . If not, i.e. $n_{i+1} - n_i > F + 1$ for some i, then

$$d_H(g_{n_i}.o, g_{n_{i+1}}.o) > C$$

By maximality of $\{g_{n_i}\}_i$, we also have

$$d_H(g_{n_i}.o, g_{n_{i+1}-1}.o) < C$$

This is a contradiction because $n_{i+1} - n_i - 1 > F$, then by definition of F

$$d_H(g_{n_i}.o, g_{n_{i+1}-1}.o) > C.$$

Then the above inequality turns into

$$d_H(g_{n_i}.o, g_{n_{i+1}}.o) \le \left(\sum_{k=0}^{F-1} d_2^k\right) \bar{\eta}(d_H(g_{n_i}.o, g_0.o))$$

but the coefficient in front of $\bar{\eta}(d_H(g_n.o, g_0.o))$ is now just a constant. Hence, using the same argument in the proof of Proposition 6.6 we get a sublinear function $\tilde{\eta}$ that satisfies condition (1) and thus proves the corollary.

To simplify notation, let $h_i \coloneqq g_{n_i}$ for the rest of this section.

Proposition 6.9. There exist a constant a and a sublinear function $\eta' = O(\eta)$ such that the following hold for all *i*, *j*, and $\alpha \in \theta$:

(6.5)
$$\alpha(\kappa(h_i^{-1}h_{i+j})) \ge a \cdot d_X(h_i, h_{i+j}) - \eta'(d_X(h_0, h_{i+j})).$$

Proof. Let γ_i be the straight geodesic ray that connects o and $h_i^{-1}\gamma_{\infty}$, where γ_{∞} is the end point of γ . Due to Fact 2.11, $d_{Haus}(\gamma_i, h_i^{-1}\gamma|_{[t_{n_i},\infty]}) \leq d_H(o, h_i^{-1}\gamma(t_{n_i})) < r$. Then by construction and triangle inequality, $h_i^{-1}h_{i+j}.o$ is 2r close to γ_i for all j. Let $a_{i,k} \coloneqq \pi_{\gamma_i}(h_i^{-1}h_k.o)$ be a point on γ_i that realizes the shortest distance between $h_0^{-1}h_k.o$ and γ_i . Then

$$d_{H}(h_{i}^{-1}h_{i+j}.o, [o, h_{i}^{-1}h_{i+j+1}.o]) \leq d_{H}(h_{i}^{-1}h_{i+j}.o, a_{i,i+j}) + d_{H}(a_{i,i+j}, \pi_{[o, h_{i}^{-1}h_{i+j+1}.o]}(h_{i}^{-1}h_{i+j}.o))$$

$$\leq 2r + d_{Haus}([o, a_{i,I+j+1}], [o, h_{i}^{-1}h_{i+j+1}.o]).$$

Using Fact 2.11 again, we can bound the last term in the above inequality by 2r and obtain

$$d_H(h_i^{-1}h_{i+j}.o, [o, h_i^{-1}h_{i+j+1}.o]) \le 4r.$$

Next we have a lemma from [6] to relate the singular values of elements in $Aut(\Omega)$ going towards the boundary "without backtracking":

Lemma 6.10 (Lemma 6.6, [6]). For any $b_0 \in \Omega$ and R > 0 there exists H > 0 such that: for all $g, h \in \Gamma_0$ with

$$d_H(gb_0, [b_0, hb_0]) \le R,$$

and for all $\alpha \in \theta$ we have

$$\alpha(\kappa(\rho(h))) \ge \alpha(\kappa(\rho(g))) + \alpha(\kappa(\rho(g^{-1}h))) - H.$$

Applying Lemma 6.10 to $b_0 = o$ and R = 4r and iterating this inequality gives

(6.6)
$$\alpha(\kappa(h_i^{-1}h_{i+j})) \ge \sum_{k=1}^j \alpha(\kappa(h_{i+k-1}^{-1}h_{i+k})) - j \cdot H.$$

Now pick the constant C in $A_{\gamma,C}$ large enough so that for all $g, h \in Aut(\Omega)$, if $d_H(g.o, h.o) \geq C$, then $\alpha(\kappa(g^{-1}h)) > H + 1$. This turns Equation (6.6) into

(6.7)
$$\alpha(\kappa(h_i^{-1}h_{i+j})) \ge j.$$

Now all that is left to do to prove Proposition 6.9 is to relate the index j with the distance $d_X(h_i, h_{i+j})$.

For all $M > \varepsilon > 0$ small, we can find T large enough such that

$$(M - \varepsilon) \cdot t \leq Leb(Cpct_{\gamma}(t)) \leq (M + \varepsilon) \cdot t$$

for all $t \ge T$ due to Equation (6.1). Moreover, for all *i* such that $t_{n_i} > T$, we have

$$2r \cdot (F+1) \cdot j \ge 2r \cdot (n_{i+j} - n_i) \ge Leb(Cpct_{\gamma}(t_{n_{i+j}}) - Cpct_{\gamma}(t_{n_i}))$$

$$> (M - \varepsilon) \cdot t_{n_{i+j}} - (M + \varepsilon) \cdot t_{n_i}$$

$$> M \cdot (t_{n_{i+j}} - t_{n_i}) - 2\varepsilon t_{n_{i+j}}$$

$$> M \cdot (d_H(h_i.o, h_{i+j}.o) - 2r) - 2\varepsilon \cdot (d_H(h_0.o, h_{i+j}.o) + 2r)$$

Again, because ε is arbitrarily small as $t_{n_{i+i}} \to \infty$,

$$\eta''(d_H(h_{0.0}, h_{i+j.0})) := 2\varepsilon \cdot (d_H(h_{0.0}, h_{i+j.0}) + 2r) + 2mr$$

is sublinear as r, m are fixed constants.

Using η'' we can rewrite Equation (6.7) as

$$\alpha(\kappa(h_i^{-1}h_{i+j})) \ge \frac{M}{2r(F+1)} \cdot d_H(h_i.o, h_{i+j}.o) - \frac{\eta''(d_H(h_0.o, h_{i+j}.o))}{2r(F+1)}.$$

By Equation (2.1) and again the fact that $\log \frac{\sigma_1}{\sigma_D}(g)$ is bi-Lipschitz to $d_X(\rho(g), e)$, we obtain the desired inequality

$$\alpha(\kappa(h_i^{-1}h_{i+j})) \ge a \cdot d_X(h_i, h_{i+j}) - \eta'(d_X(h_{i+j}, h_0))$$

with a depend on M, F, r, E, and the bi-Lipschitz constant, and $\eta' = O(\eta'')$.

Moreover, by construction, $\eta' = O(\eta)$ as they are derived from the same limit in Equation (6.1). This concludes the proof of Proposition 6.9.

Proof of Thm. 6.3. Proposition 6.6 and Proposition 6.9 show that $\{h_i\}$ satisfies condition (1) and (3) in Definition 1.2. All that is left to do for proving $\{h_i\}$ is P_{θ} -sublinearly Morse is to show that the concatenated path $(\bigcup_i [h_i, h_{i+1}])$ is a sublinear ray.

Let $c: [0, \infty) \to X$ be the unit speed parametrization of $(\bigcup_i h_0^{-1}[h_i, h_{i+1}])$, it follows from the triangle inequality of the symmetric space distance d_X that for all $s \leq w \in [0, \infty)$,

$$d_X(c(s), c(w)) \le d_X(c(s), h_{i_s+1}) + \sum_{n=i_s+1}^{i_w-1} d_X(h_n, h_{n+1}) + d_X(h_{i_w}, c(w)) = |w - s|$$

with $h_{i_s}, h_{i_w} \in \rho(A_{\gamma,C})$ the elements such that $c(s) \in [h_{i_s}, h_{i_s+1}]$ and $c(w) \in [h_{i_w}, h_{i_w+1}]$. Hence, to show c is a sublinear ray, we only need to find a suitable lower bound for $d_X(c(s), c(w))$. We do this by cases:

Case 1: If $c(s) = h_{i_s}, c(w) = h_{i_w}$, then

$$d_X(c(s), c(w)) \ge 2L \cdot \frac{1}{2} \log \frac{\sigma_1}{\sigma_D} (\rho^{-1}(h_{i_s}^{-1}h_{i_w})) \ge 2L \cdot (d_H(h_{i_s} \cdot o, h_{i_w} \cdot o) - E)$$

for L the bi-Lipschitz constant relating $\log \frac{\sigma_1}{\sigma_D}$ to d_X and E in Fact 2.10.

By triangle inequality

$$d_H(h_{i_s}.o, h_{i_w}.o) \ge (t_{i_w} - t_{i_s}) - 2r \ge \sum_{k=i_s}^{i_w - 1} (d_H(h_k.o, h_{k+1}.o) - 2r) - 2r$$

By construction, $d_H(h_k.o, h_{k+1}.o) \ge C$ for all k, hence

(6.8)
$$d_X(c(s), c(w)) \ge c' \sum_{k=i_s}^{i_w-1} d_H(h_k.o, h_{k+1}.o) - 2r \ge c'' \sum_{k=i_s}^{i_w-1} d_X(h_k.o, h_{k+1}.o) - E'$$

with c', c'' and E' depending only on C, E, r and L. Note that we can always enlarge the constant C that defines $A_{\gamma,C}$ to ensure that c'' is positive.

Case 2: If c(s), c(w) do not belong to the orbit $\{h_n\}$, then without loss of generality assume $i_s \leq i_w - 1$ (if they were equal then c(s), c(w) lie on the same geodesic segment and the statement is trivial), then by triangle inequality and Proposition 6.6

$$d_X(c(s), c(w)) \ge d_X(h_{i_s}, h_{i_w+1}) - \eta(d_X(h_{i_s}, h_0)) - \eta(d_X(h_{i_w}, h_0))$$

$$\ge c'' \sum_{k=i_s}^{i_w} d_X(h_k, h_{k+1}) - 2\eta(d_X(h_{i_w}, h_0))$$

$$\ge c''|w - s| - 2\eta(w).$$

This proves that the concatenated path c is a $(\frac{1}{c''}, \bar{\eta})$ sublinear ray, for $\bar{\eta} = 2\eta$ and thus finishes the proof of Theorem 6.3.

6.2. **Proof of Main Theorem.** Because we are working with a ϕ -divergent group, i.e. $Q^{\phi}_{\Gamma}(\delta_{\phi}) = +\infty$, Theorem 3.3 tells us the geodesic flow on $\Gamma_0 \backslash S_{\Gamma_0} \Omega$ is ergodic.

By Birkhoff Ergodic Theorem, let $V \subset \Gamma_0 \backslash S_{\Gamma_0} \Omega$ be any Borel measurable set, then for \overline{m} a.e. $v \in \Gamma_0 \backslash S_{\Gamma_0} \Omega$, we have

$$\lim_{T \to \infty} \frac{Leb(\{t \in [0, T] \mid [\psi^t(v)] \in V\})}{T} = \frac{\overline{m}(V)}{\overline{m}(\Gamma_0 \setminus S_{\Gamma_0} \Omega)}.$$

By assumption, $\overline{m}(\Gamma_0 \setminus S_{\Gamma_0}\Omega) < \infty$, so we can normalize \overline{m} to be a probability measure. Furthermore, if $W \subset S_{\Gamma_0}\Omega$ is Γ_0 -invariant, the limit can be reformulated as

(6.9)
$$\lim_{T \to \infty} \frac{Leb(\{t \in [0,T] \mid \psi^t(v) \in W\})}{T} = \bar{m}(\Gamma_0 \backslash W).$$

Fix W to be the collection of vectors in $S_{\Gamma_0}\Omega$ whose base points are in $\Gamma_0.B_r(o)$. Then W is Γ_0 -invariant.

Let $\pi: S\Omega \to \Omega$ be the map that sends a unit tangent vector to its base point. Geometrically, $\psi^t(v) \in W$ means $\pi(\psi^t(v)) \in \Gamma_0.\overline{B_r(o)}$. The limit then turns into

$$\lim_{T \to \infty} \frac{Leb(Cpct_{\gamma_v}(T))}{T} = \bar{m}(\Gamma_0 \backslash W)$$

where γ_v is the geodesic ray $\pi(\psi^t(v))$.

Since $\Gamma_0 \setminus W$ contains non-empty compact sets, namely $\pi^{-1}(\overline{B_r(o)})$, and \overline{m} has no atom, we can conclude that $\overline{m}(\Gamma_0 \setminus W) > 0$ if we choose r to be big enough. Then for \overline{m} a.e. $v \in \Gamma_0 \setminus S_{\Gamma_0} \Omega$, the limit above converges to a positive number. By Theorem 6.3, there exist a sequence of elements $\{h_i\}_i = A_{\gamma_v}, C$ that is P_{θ} -sublinearly Morse.

We finish the proof of our main theorem by assuming the set

 $U := \{\zeta \in \Lambda_{\theta}(\Gamma) \mid \text{there is no } P_{\theta}\text{-sublinearly Morse sequence converging to } \zeta\}$ has positive measure in $\Lambda_{\theta}(\Gamma)$. This means for all $\zeta \in U$, any geodesic ray $l \subset \Omega$ with $l(+\infty) = \xi^{-1}(\zeta)$, we have

$$\lim_{T \to \infty} \frac{Leb(Cpct_l(T))}{T} = 0$$

To abuse notation, we ignore the representation ξ^{-1} . We can then define

 $V \coloneqq \{ v \in S_{\Gamma_0} \Omega \mid \lim_{t \to \infty} \pi(\psi^t(v)) = \zeta \}.$

Let \tilde{V} be the projection of V to the quotient space $\Gamma_0 \setminus S_{\Gamma_0} \Omega$. Note that m(V) is proportional to $\bar{\mu}(\Lambda_{\Omega}(\Gamma_0)) \cdot \mu(U) \cdot s(\mathbb{R}) > 0$ by Fubini's Theorem, then $\bar{m}(\tilde{V}) > 0$. By construction, for all $v \in V$

$$\lim_{T \to \infty} \frac{Leb(Cpct_{\gamma_v}(T))}{T} = 0.$$

This is a contradiction because on the positive measured set \tilde{V} , the limit above converges to zero instead of a positive number. Hence, we conclude the proof of the genericity of P_{θ} -sublinearly Morse points in $\Lambda_{\theta}(\Gamma)$.

7. Relatively Anosov Representations

In this section, we prove a corollary of our main theorem in the relative Anosov setting. In order to make relative Anosovness precise, we need to define a few things first. Let G be the higher rank Lie group defined in Section 2.

Definition 7.1 (Geometrically Finiteness). Let M be a compact perfect metrizable space and $H \subset Homeo(M)$ be a convergence group acting on M. H is geometrically finite if for all $\eta \in M$, η is either

• conical in the convergence group sense, as defined in Section 3.2,

or

• bounded parabolic, that is $Stab_{\Gamma}(\eta)$ is a parabolic subgroup of Γ , and that $Stab_{\Gamma}(\eta)$ acts on $M \setminus \{\eta\}$ cocompactly.

Let $\Gamma \subset G$ be a finitely generated group, \mathcal{P} a collection of finitely generated subgroups of Γ . The pair (Γ, \mathcal{P}) is *relative hyperbolic* if Γ acts on a compact perfect metrizable space M as

a geometrically finite convergence subgroup and \mathcal{P} contains all possible maximal parabolic subgroups up to conjugating by elements in Γ .

Fixing a generating set S of Γ , we can construct the Groves-Manning cusp space by gluing combinatorial horoballs to $Cay(\Gamma, S)$. One can refer to Section 3.4 in Zhu–Zimmer [21] for the explicit construction. The Groves–Manning cusp space is Gromov hyperbolic, and its Gromov boundary, denoted by $\partial(\Gamma, \mathcal{P})$, serves as a compact perfect metrizable space where Γ act on as a geometrically finite convergence group.

There are various characterizations of relative Anosov subgroups presented in Section 4 of [21], we will present just one here.

Definition 7.2. A subgroup $\Gamma \subset G$ is P_{θ} -Anosov relative to \mathcal{P} if it is P_{θ} -transverse, (Γ, \mathcal{P}) is a relatively hyperbolic pair, and the induced boundary map

 $\xi': \partial(\Gamma, \mathcal{P}) \to \mathcal{F}_{\theta}$

is Γ -equivariant and a homeomorphism onto $\Lambda_{\theta}(\Gamma)$.

Fix (Γ, \mathcal{P}) to be relatively P_{θ} -Anosov. Let Ω , $\Gamma_0 \subset Aut(\Omega)$, and ρ be as in Theorem 3.1. They exist because Γ is P_{θ} -transverse by definition. Note that $\xi'(\partial(\Gamma, \mathcal{P}))$ is homeomorphic to $\Lambda_{\Omega}(\rho^{-1}(\Gamma))$. We will omit the representations in the rest of this section when the context is clear. Our main theorem can be restated as follow.

Theorem 7.3. For $\phi \in \mathfrak{a}_{\theta}^*$, and μ the unique Patterson–Sullivan measure of dimension $\delta_{\phi} < \infty$ supported on $\Lambda_{\theta}(\Gamma) \subset \mathcal{F}_{\theta}$, $\mu(\partial_{SM,\theta}(\Gamma)) = 1$.

The key differences here are that we can drop the assumptions in our main theorem where 1) Γ is ϕ -divergent; 2) the induced Bowen–Margulis–Sullivan measure \bar{m} as defined in Section 3.1 is finite. The fact that Γ is ϕ -divergent, i.e. the Poincaré series diverges at δ_{ϕ} , follows directly from Theorem 8.1 in [8], when Γ is P_{θ} -Anosov relative to \mathcal{P} . The finiteness of \bar{m} is shown by the following proposition.

Proposition 7.4 (Finiteness of Bowen–Margulis–Sullivan measure). Let (Γ, \mathcal{P}) be the pair defined as above. Then $\overline{m}(\Gamma_0 \setminus S_{\Gamma_0} \Omega)$ is finite.

Remark 7.5. In their paper [15], Kim–Oh proved a similar result (Theorem 1.1) of finiteness of the Bowen–Margulis–Sullivan measure when Γ is a relative Anosov subgroup. However, as we noted in section 2.3, their result applies to a different flow space than the ones we are focusing on in this paper, thus we cannot apply their result directly.

We prove the proposition mainly by following methods used in Blayac–Zhu [2, Theorem 8.1].

We will first show that $S_{\Gamma_0}\Omega$ can be decomposed into a compact part and a family of disjoint horoballs based at the bounded parabolic points (which, after quotienting out by action of Γ_0 , are referred as the cusps). Then one can show that \bar{m} is finite on each of the cusps. Since there are only finitely many conjugacy classes of maximal parabolic subgroups [4], the number of orbits of the bounded parabolic points is finite. Moreover, because the support of \bar{m} is compact outside of these cusp regions, and after quotienting, there are only finitely many cusp regions with finite Bowen–Margulis–Sullivan measure, we can then conclude that $\bar{m}(\Gamma_0 \backslash S\Omega)$ is finite. Before we proceed to the actual proof, there is one last technical detail. The conical limit points used in [2] are different from the conical limit points of a convergence group. In [2], a point $\eta \in \Lambda_{\Omega}(\Gamma_0)$ is *conical* if there exists a sequence $\{\gamma_n\} \subset \Gamma_0$ that converges to η and $\gamma_n.o$ is uniformly bounded away from the straight geodesic ray $[o, \eta)$.

This gap is bridged by the following lemma.

Lemma 7.6. [6, Lemma 3.6] For all $x \in \Lambda_{\Omega}(\Gamma_0)$, $x \in \xi^{-1} \circ \xi'(\partial^{con}(\Gamma, \mathcal{P}))$ if and only if there is $a(ny) \ b_0 \in \Omega$ and a sequence $\{\gamma_n\}$ in Γ_0 such that $\gamma_n \to x$ and

$$\sup_{n\geq 1} d_{\Omega}(\gamma_n.b_0, [b_0, x)) < +\infty.$$

Here $\partial^{con}(\Gamma, \mathcal{P})$ denotes the set of conical limit points of the convergence group action in $\partial(\Gamma, \mathcal{P})$. Since conicality in the convergence group sense is a purely topological condition, and ξ and ξ' are homeomorphisms, the conicality is preserved under $\xi^{-1} \circ \xi'$ (similarly, one can check that the bounded parabolic points are also preserved). The lemma above then says that the two notions of conical limit points coincide.

Other concepts involved in the proof of Proposition 7.4 are horofunctions and hororballs associated to Ω . For any $x \in \Omega$, the *horofunction* $\beta_x : \Omega \times \Omega \to \mathbb{R}$ is defined as

$$\beta_x(a,b) = d_H(a,x) - d_H(b,x).$$

Note that after fixing a basepoint $o \in \Omega$, the horofunctions provide a way to embed Ω into $\mathcal{C}(\Omega)$, the space of continuous functions defined on Ω , by sending $x \in \Omega$ to $\beta_x(\cdot, o)$. The image of Ω , denoted by $\beta(\Omega)$, is relatively compact with respect to the topology of uniform convergence, hence we can view the closure of $\beta(\Omega)$ as a compactification of Ω , denoted by $\overline{\Omega^h}$, and let $\partial_h \Omega := \overline{\Omega^h} \setminus \beta(\Omega)$ be the *horoboundary* of Ω . For any $\beta \in \partial_h \Omega$ and $x \in \Omega$, we define the *horoball* and *horosphere* centered at β and passing through x, denoted by $\mathcal{H}_{\beta}(x)$ and $\partial \mathcal{H}_{\beta}(x)$, to be

$$\mathcal{H}_{\beta}(x) \coloneqq \{ y \in \Omega \mid \beta(x, y) > 0 \},\$$

$$\partial \mathcal{H}_{\beta}(x) \coloneqq \{ y \in \Omega \mid \beta(x, y) = 0 \}.$$

Moreover, a theorem of Walsh [20, Theorem 1.3] states that for all $\beta \in \partial_h \Omega$ and $\{x_k\} \subset \Omega$ such that $\beta_{x_k} \to \beta$, there exists $\xi \in \partial \Omega$ such that $x_k \to \xi$ in $\overline{\Omega}$. Hence there is a natural surjective projection $\pi_h : \partial_h \Omega \to \partial \Omega$. For any $\xi \in \partial \Omega$, the preimage $\pi_h^{-1}(\xi)$ contains exactly one point if and only if ξ is a C^1 -smooth in $\partial \Omega$ [5, Lemma 3.2]. Note that for all $\xi \in \Lambda_\Omega(\Gamma_0)$, ξ is C^1 -smooth as Γ_0 is projectively visible, hence π_h^{-1} is well-defined on $\Lambda_\Omega(\Gamma_0)$. We will use β_{ξ} to denote the preimage of $\xi \in \Lambda_\Omega(\Gamma_0)$, and shorthand \mathcal{H}_{ξ} for $\mathcal{H}_{\beta_{\xi}}$.

With the above discussion, $\rho^{-1}(\Gamma)$ acts geometrically finitely on Ω as in Definition 1.10 in [2], then we have the following.

Lemma 7.7. [2, Lemma 8.11] If x, y are two bounded parabolic points in $\Lambda_{\Omega}(\Gamma_0)$, and \mathcal{H}' a horoball centered at y. Then there exists a horoball \mathcal{H} centered at x such that for all $\gamma \in \Gamma_0$, either $\mathcal{H}' \cap \gamma \mathcal{H} = \emptyset$ or $\gamma x = y$.

Remark 7.8. In [2], Blayac–Zhu worked with smooth domains. Although we cannot assume the same for Ω , we can still apply their results because all they used is the smoothness and strict convexity between points in the limit points, which is given in our case as Γ_0

is projectively visible. Moreover, this lemma allows the existence of a family of disjoint horoballs \mathcal{H}_x centered at bounded parabolic points, which are Γ_0 -equivariant and each are preserved by the parabolic subgroup $Stab_{\Gamma_0}(x)$. This will be the set of horoballs stated in the next lemma.

Lemma 7.9. For each bounded parabolic point $\alpha \in \Lambda_{\Omega}(\Gamma_0)$, fix an open horoball \mathcal{H}_{α} centered at α , such that $\gamma \mathcal{H}_{\alpha} = \mathcal{H}_{\gamma\alpha}$ for all $\gamma \in \Gamma_0$, and the distinct horoballs are mutually disjoint. Then the set of unit tangent vectors $v \in S_{\Gamma_0}\Omega$ whose base point projection $\pi(v)$ does not belong to $\bigcup \{\mathcal{H}_{\alpha} \mid \alpha \text{ is bounded parabolic}\}$ is compact.

Proof. The proof is mainly based on the proof of Lemma 8.12 in [2]. Define

$$D \coloneqq \{x \in \Omega \mid d_H(x, \gamma.o) \ge d_H(x, o) \text{ for all } \gamma \in \Gamma_0\},\$$

then showing the set

$$A \coloneqq D \bigcap \left(\bigcup_{\beta, \eta \in \Lambda_{\Omega}(\Gamma_0)} (\beta, \eta) \right) \setminus (\bigcup \mathcal{H}_{\alpha})$$

is compact suffices to prove the lemma, because the subset of interest can be seen as a bundle over A with compact fibers.

Assume the contrapositive that A is not compact, then by Lemma 7.6 there exist a unbounded sequence $\{x_n\} \subset A$, and, up to taking subsequence, x_n converges to some $\alpha \in \Lambda_{\Omega}(\Gamma_0)$. Since Γ_0 acts geometrically finitely on $\partial\Omega$, a is either conical or bounded parabolic.

If α is conical (in the convex domain sense), then there exists a sequence $\{\gamma_n\} \subset \Gamma_0$ such that $\sup_{n\geq 1} d_H(\gamma_n.o, [o, \alpha)) < +\infty$ and $\gamma_n.o$ converges to α . However, this would imply that

$$\infty = \lim_{n \to \infty} \beta_{\alpha}(o, \gamma_n.o) \coloneqq \lim_{n \to \infty} \lim_{k \to \infty} \beta_{x_k}(o, \gamma_n.o) = \lim_{n \to \infty} \lim_{k \to \infty} d_H(x_k, o) - d_H(x_k, \gamma_n.o) \le 0,$$

which gives us a contradiction. So α has to be bounded parabolic.

By construction of A, there exists $\{\zeta_n\}$ and $\{\eta_n\}$ in $\Lambda_{\Omega}(\Gamma_0)$ such that $x_n \in (\eta_n, \zeta_n)$ and $x_n \notin \mathcal{H}_{\alpha}$. Since \mathcal{H}_{α} is convex (because Ω is a properly convex domain), the intersection $\mathcal{H}_{\alpha} \cap (\eta_n, \zeta_n)$ is connected. Then up to exchanging η_n and ζ_n , we can assume $[x_n, \eta_n)$ is disjoint from the open horoball \mathcal{H}_{α} . After picking subsequences, we can also assume that $\zeta_n \to \zeta$ and $\eta_n \to \eta \in \Lambda_{\Omega}(\Gamma_0)$.

Since α is bounded parabolic, we can also pick a diverging sequence $\{\gamma_n\} \subset Stab_{\Gamma_0}(\alpha)$ such that $\gamma_n\eta_n$ converges to $\bar{\eta} \neq \alpha$. This is possible because $Stab_{\Gamma_0}(\alpha)$ act cocompactly on $\Lambda_{\Omega}(\Gamma_0) \setminus \{\alpha\}$ and $\eta_n \neq \alpha$ for infinitely many n since $\mathcal{H}_{\alpha} \cap [x_n, \eta_n) = \emptyset$ by construction. Note that β_{α} is preserved under the action of $Stab_{\Gamma_0}(\alpha)$, so for all $\gamma_n \in Stab_{\Gamma_0}(\alpha)$, we have

$$[\gamma_n x_n, \gamma_n \eta_n) \cap \mathcal{H}_{\alpha} = \emptyset.$$

By taking a subsequence, we can assume that $\gamma_n x_n$ converges to $x \in \overline{\Omega}$. We claim that $x \neq \alpha$. If false, then $[\gamma_n x_n, \gamma_n \eta_n)$ converges to (α, η) that's also disjoint from \mathcal{H}_{α} . This gives us a contradiction, because $(\alpha, \eta) \subset \Omega$ and thus has to intersect all the horoballs centered at α . Then

$$\infty = \lim_{n \to \infty} d_H(o, \gamma_n.o) \le \lim_{n \to \infty} \left[d_H(x_n, \gamma_n^{-1}o) + d_H(\gamma_n o, o) - d_H(x_n, o) \right]$$
$$= \lim_{n \to \infty} \left[d_H(\gamma_n x_n, o) + d_H(\gamma_n o, o) - d_H(\gamma_n x_n, \gamma_n o) \right]$$
$$= \langle x, \alpha \rangle_o < \infty$$

where \langle , \rangle_o is the Gromov product based at o, and it's finite because (x, α) intersects Ω . This gives us a contradiction and thus α can not be bounded parabolic either, so $\{x_n\} \subset A$ is bounded, hence A is compact.

Proof of Proposition 7.4. After having the decomposition of $S_{\Gamma_0}\Omega$ in Lemma 7.9, all that is left to show is that \bar{m} is finite on each cusp.

For any $\eta \in \Lambda_{\Omega}(\Gamma_0)$ bounded parabolic, take the horoball \mathcal{H}_{η} defined as above, and $P := Stab_{\Gamma_0}(\eta) \in \mathcal{P}$ the corresponding parabolic subgroup fixing η and \mathcal{H}_{η} . Let \mathcal{C}_P be a strict fundamental domain of the action of P on \mathcal{H}_{η} , that is, for any $x \in \mathcal{H}_{\eta}$, there exists a unique $p \in P$ such that $px \in \mathcal{C}_P$. Moreover, since η is bounded parabolic, we can choose a relatively compact strict fundamental domain F for the action of P on $\Lambda_{\Omega}(\Gamma_0) \setminus \{\eta\}$.

Since the ϕ -Patterson–Sullivan density μ we used to obtain the Bowen–Margulis–Sullivan measure m has no atom [6, Proposition 9.1], we can write the measure of the quotient of the horoball as

(7.1)
$$\bar{m}\left(\Gamma_0\backslash S\mathcal{H}_\eta\right) = \sum_{p,q\in P} \int_{pF\times qF} e^{-\delta_\phi\phi\left(\left[\xi(x),\xi(y)\right]_\theta\right)} d\bar{\mu}(\xi(x)) d\mu(\xi(y)) \cdot \int_{(x,y)\cap\mathcal{C}_P} ds dx$$

Because m is Γ_0 invariant, so is \overline{m} , we can then rewrite the equation above as

(7.1*)
$$\bar{m}\left(\Gamma_{0}\backslash S\mathcal{H}_{\eta}\right) = \sum_{p,q\in P} \int_{F\times p^{-1}qF} e^{-\delta_{\phi}\phi\left(\left[\xi(x),\xi(y)\right]_{\theta}\right)} d\bar{\mu}(\xi(x)) d\mu(\xi(y)) \cdot \int_{(x,y)\cap\mathcal{H}_{\eta}} ds$$
$$= \sum_{p\in P} \int_{F\times pF} e^{-\delta_{\phi}\phi\left(\left[\xi(x),\xi(y)\right]_{\theta}\right)} d\bar{\mu}(\xi(x)) d\mu(\xi(y)) \cdot \int_{(x,y)\cap\mathcal{H}_{\eta}} ds,$$

where the second equality is because C_P is a strict fundamental domain for the action of P on \mathcal{H}_{η} and the action is transitive.

Geometrically, the second term in the product measures how much time the geodesic (x, y) spends in \mathcal{H}_{η} , and we show this is bounded.

Claim. There exists an open neighborhood $U \subset \overline{\Omega}$ of η such that for all $\zeta \in F$, all $w \in U$, we have $[w, \zeta) \cap \mathcal{H}_{\eta} \neq \emptyset$

Proof of claim. Assume the claim fails, then there exists a sequence of shrinking open neighborhoods $\{U_n\}$ of η such that there exist $\{w_n \in U_n\}$ and $\{\zeta_n \in F\}$ with $[w_n, \zeta_n) \cap \mathcal{H}_{\eta} = \emptyset$. Because F is relatively compact and U_n are shrinking, $\zeta_n \to \zeta \in \overline{F}$ and $w_n \to \eta$. This would then give us a contradiction because (ζ, η) would be a geodesic line that doesn't intersect \mathcal{H}_{η} .

With this neighborhood U, we can define $R \coloneqq d_H(o, \partial \mathcal{H}_\eta \setminus U) < \infty$. Fix $x \in F$ and $y \in pF$ such that $(x, y) \cap \mathcal{H}_\eta \neq \emptyset$. Let a, b be the unique intersection points of (x, y) and $\partial \mathcal{H}_\eta$ such that the ordering of them on (x, y) is x, a, b, y. By construction, [a, x) and [b, y)are disjoint from the open horoball \mathcal{H}_η , hence a and $p^{-1}b$ are not in U. We then have $d_H(a, o) \leq R, d_H(p^{-1}.b, o) = d_H(b, p.o) \leq R$, and we can bound the second integral in (7.1*) from above

$$\int_{(x,y)\cap\mathcal{H}_{\eta}}^{\cdot} ds \leq 2R + d_H(o,po).$$

We can find an open neighborhood $V \in \Omega \cup \partial \Omega$ of \overline{F} , such that $\eta \notin V$. Since the pair (Γ, \mathcal{P}) is relative hyperbolic, the Bowditch boundary $\partial(\Gamma, \mathcal{P}) \simeq \Lambda_{\Omega}(\Gamma_0)$ is a compact perfect metrizable

space, hence $\bar{F} \times (\Lambda_{\Omega}(\Gamma_0) \setminus V)$ is also compact. Because ξ , ϕ , and the Gromov product are both continuous and defined on this space, as it doesn't contain any non transverse pair, $\phi([\xi(x),\xi(y)]_{\theta})$ is bounded on $\bar{F} \times (\Lambda_{\Omega}(\Gamma_0) \setminus V)$, i.e. there exist C_1, C_2 such that $C_1 \leq \phi([\xi(x),\xi(y)]_{\theta}) \leq C_2$ for $(x,y) \in \bar{F} \times (\Lambda_{\Omega}(\Gamma_0) \setminus V)$ and $x \neq y$, and Equation (7.1*) turns into $\bar{m}(\Gamma_0 \setminus S\mathcal{H}_n) \leq \exp(-C_1) : \mu_1(F) \sum (2B + d\mu(\rho, n, \rho)) : \mu_2(pF)$

$$\bar{m}\left(\Gamma_{0}\backslash S\mathcal{H}_{\eta}\right) \leq \exp\left(-C_{1}\right) \cdot \mu_{o}(F) \sum_{p \in P} \left(2R + d_{H}(o, p.o)\right) \cdot \mu_{o}(pF).$$

Note here that we are able to rewrite the whole sum because although $e^{-\delta_{\phi}\phi([\xi(x),\xi(y)]_k)}$ is not uniformly bounded below on $\overline{F} \times V$, we can omit all the terms where $(x,y) \in \overline{F} \times V$ by choosing the horoballs \mathcal{H}_{η} in Lemma 7.9 to be small enough at the beginning so that $(x,y) \cap \mathcal{H}_{\eta} = \emptyset$ and with the same geometric interpretation of the second integral, these terms does not contribute to the sum.

Now we just have to find ways to control $\mu_o(pF)$. Since \overline{F} and $P.o \cup \{\eta\}$ are disjoint and compact, and for all $x \in \overline{F}, y \in P.o \cup \{\eta\}$, $[x, y] \cap \Omega$, then there exist R' > 0 such that $[x, y] \cap B_{R'}(o) \neq \emptyset$. This implies that $F \subset \mathcal{O}_{R'}(p.o, o)$ for all $p \in P$, equivalently $pF \subset \mathcal{O}_{R'}(o, p.o)$ for all $p \in P$. Hence, applying a shadow lemma in this setting [6, Proposition 7.1], there exist a constant C such that

$$\mu_o\left(\xi(\mathcal{O}_{R'}(o, p.o) \cap \Lambda_{\Omega}(\Gamma))\right) \le C \cdot \exp\left(-\delta_\phi \phi(\kappa(p))\right)$$

for all $p \in P$.

Then we have

$$\bar{m}\left(\Gamma_{0}\backslash S\mathcal{H}_{\eta}\right) \leq C'\sum_{p\in P}\left(2R + d_{H}(o, p.o)\right) \cdot \exp\left(-\delta_{\phi}\phi(\kappa(p))\right)$$

with $C' = C \cdot \exp(-C_1) \cdot \mu_o(F)$.

Theorem 10.1 in [8] states that for any $\phi \in \mathfrak{a}_{\theta}^*$ and $p \in \Gamma$, $\phi(\kappa(p))$ is quasi-isometric to $d_X(p,e)$, and thus to $d_H(g.o,o)$ by property (4) of Theorem 3.1. Up to changing some constants, the equation above turns into

$$\bar{m}\left(\Gamma_{0}\backslash S\mathcal{H}_{\eta}\right) \leq C' \sum_{p \in P} \exp\left\{-\delta_{\phi} \cdot \left[\phi(\kappa(p)) - \epsilon_{0}\log(\phi(\kappa(p)))\right]\right\}$$

for some $\epsilon_0 > 0$. Since the logarithm function is sublinear, then as argued before, it does not affect the convergence of the sequence. Then the series is bounded by the Poincare series of P, $Q_P^{\phi}(\delta_{\phi} \cdot (1 - \epsilon')) = \sum_{p \in P} \exp(\delta_{\phi} \cdot (1 - \epsilon') \cdot \phi(\kappa(p)))$ for all $\epsilon' > 0$. By Theorem 7.1 and Corollary 7.2 in [8], there is an entropy drop of peripheral subgroups $P \in \mathcal{P}$ of the relative pair (Γ, \mathcal{P}) , i.e. $\delta_{\phi}(\Gamma) > \delta_{\phi}(P)$. Then for ϵ' small enough, $Q_P^{\phi}(\delta_{\phi}(\Gamma) \cdot (1 - \epsilon'))$ converges, hence $\overline{m} (\Gamma_0 \setminus S\mathcal{H}_{\eta})$ is finite. This concludes the proof of Proposition 7.4, and Theorem 7.3 follows from Theorem 6.1.

8. A Geometric Interpretation

Let X be a higher rank symmetric space defined in Section 2 and θ a symmetric subset of the set of simple roots. As mentioned in Section 4, the higher rank Morse lemma in [13] links regularity of a quasi-geodesic ray to the existence of a uniform upper bound on its distance to a certain Weyl cone. We want to generalize this result to sublinear rays with the following property. **Definition 8.1.** A path $c: I \to X$ is (q, χ) P_{θ} -sublinearly Morse if there exist a constant q > 0 and a sublinear function χ such that for all $s, t \in I$ and all $\alpha \in \theta$ we have

$$\alpha(\kappa(c(s)^{-1}c(t))) \ge q \cdot d_X(c(s), c(t)) - \chi(\max\{d_X(c(0), c(s)), d_X(c(0), c(t))\}).$$

We can show that given a P_{θ} -sublinearly Morse sequence $\{g_n\}$, the concatenated sublinear ray as defined in property (2) in Definition 1.2 is P_{θ} -sublinearly Morse.

Lemma 8.2. Let c be the $(C, \bar{\eta})$ sublinear ray associated to $\{g_n\}$, and a the constant in property (3) of Definition 1.2. There exists $\bar{\eta}' = O(\bar{\eta})$ such that c is $(a, \bar{\eta}') P_{\theta}$ -sublinearly Morse.

Proof. Let g_s , g_t be such that $c(s) \in [g_s, g_{s+1}]$ and $c(t) \in [g_t, g_{t+1}]$. By triangle inequality and property (1) of the sequence g_n

$$||d_{\Delta}(c(s), c(t)) - d_{\Delta}(g_s, g_t)|| \le 2\eta \left(\max\{d_X(g_0, g_s), d_X(g_0, g_t)\} \right).$$

This implies that there exists a constant B such that for all $\alpha \in \theta$

$$\alpha(\kappa(c(s)^{-1}c(t))) \ge \alpha(\kappa(g_s^{-1}g_t)) - B \cdot \eta(\max\{d_X(g_0, g_s), d_X(g_0, g_t)\})$$

then by property (3) of $\{g_n\}$ we have

$$\alpha(\kappa(c(s)^{-1}c(t))) \ge a \cdot d_X(g_s, g_t) - (\eta' + B \cdot \eta) \left(\max\{d_X(g_0, g_s), d_X(g_0, g_t)\} \right).$$

By triangle inequality again, and the fact that $\eta' = O(\eta)$, we can find another sublinear function $\bar{\eta}' = O(\eta)$ such that

$$\alpha(\kappa(c(s)^{-1}c(t)) \ge a \cdot d_X(c(s), c(t)) - \bar{\eta}'(\max\{d_X(g_0, g_s), d_X(g_0, g_t)\})$$

and thus finishes the proof.

8.1. Diamonds, Weyl cones, and Parallel Sets. In order to state our theorem precisely, we need to define diamonds, Weyl cones, and parallel sets in X. One can view them as the higher rank analogies of geodesic segments, rays, and paths in hyperbolic spaces.

Recall that in Section 2.2, we defined a map $U_{\theta} : G \to \mathcal{F}_{\theta}$ such that $U_{\theta}(g) = m_g P_{\theta}$. Let $U \subset G$ consist of points where U_{θ} is uniquely defined, i.e. for all $g \in U$ and $\alpha \in \theta$, $\alpha(\kappa(g)) > 0$.

Definition 8.3. A Weyl cone associated to some $\zeta \in \mathcal{F}_{\theta}$ with tip at $g \in X$, denoted by $V(g,\zeta)$, is defined as

$$V(g,\zeta) \coloneqq \{h \in X \mid g^{-1}h \in U \text{ and } U_{\theta}(g^{-1}h) = \zeta\}.$$

We can also define a parallel set in X associated to a pair of opposite flags. A pair of flags $(\zeta^+, \zeta^-) \in \mathcal{F}_{\theta} \times \mathcal{F}_{\theta}$ is *opposite* if there exists $g \in G$ such that $(\zeta^+, \zeta^-) = (gP_{\theta}, gP_{\theta}^{opp})$. For \mathfrak{a}^+_{θ} as defined in Section 2.1, let $H \in \mathfrak{a}^+_{\theta}$, then we obtain a bi-infinite geodesic $l : (-\infty, \infty) \to X$ of the form

$$l(t) = g \exp(tH).$$

We will use $l(\pm\infty)$ to denote the points that $U_{\theta}(l(t))$ converges to in \mathcal{F}_{θ} as $t \to \pm\infty$.

For a pair of opposite flags (ζ^+, ζ^-) , we can define the parallel set associated to it. Let l be the bi-infinite geodesic as above.

Definition 8.4. A parallel set associated to (ζ^+, ζ^-) , denoted by $P(\zeta^+, \zeta^-)$, is defined to be the union of all maximal flats in X that contains l.

Definition 8.5. For any $x \neq y \in X$ with $x^{-1}y \in U$, we can define the *diamond* with tips at x and y, denoted by $\diamond(x, y)$, to be the collection of points $z \in X$ such that $U_{\theta}(z^{-1}x)$ and $U_{\theta}(z^{-1}y)$ are opposite.

Remark 8.6. In Kapovich–Leeb–Porti [13], the diamonds, Weyl cones, and parallel sets are defined using the spherical building structure on $\partial_{\infty} X$, the visual boundary of X. By fixing a spherical model apartment in $\partial_{\infty} X$ and a model chamber σ_{mod} in this apartment defined with respect to the Weyl group action, they construct a type map $\theta_{type} : \partial_{\infty} X \to \sigma_{mod}$. Since the *G*-orbit of a point in $\partial_{\infty} X$ intersects the model chamber σ exactly once, the type map θ_{type} is well defined.

For any face $\tau \in \sigma_{mod}$, they defined the open stars in σ_{mod} associated to τ to be the union of all open faces in σ_{mod} whose closure contains τ_{mod} . If we pick τ_{mod} defined by the intersection of the reflection hyperplanes for all $\beta \notin \theta$, the connected components of the pre-image of the open star of τ_{mod} under θ_{type} one to one correspond to the visual boundary of pre-image of ζ under $U_{\theta}(\cdot)$ for all $\zeta \in \mathcal{F}_{\theta}$. To abuse notation, we will use ζ to denote both the element in \mathcal{F}_{θ} and the simplex in the corresponding connected component with type τ_{mod} . With this identification, one can show that our definitions of diamonds, Weyl cones, and parallel sets agree with those in [13].

Moreover, regularity of a geodesic segment [x, y] in [13] means $x^{-1}y \in U$, and uniform regularity translates to there exist c > 0 such that for all $\alpha \in \theta$

$$\frac{\alpha(\kappa(x^{-1}y))}{d_X(x,y)} > c$$

we will use c-regular to describe segments that satisfy the above inequality.

8.2. Asymptotic Cones. Another major tool used by Kapovich–Leeb–Porti in [13] is the asymptotic cone of symmetric spaces. We only introduce here the basic construction and properties of the asymptotic cone we need to prove Theorem 8.8. We refer the reader to Sections 2.7 and 3.9 in [13] for a thorough discussion on the object.

Let ω be a non-principal ultrafilter on the set of natural numbers \mathbb{N} . Let λ_n be a sequence of positive numbers whose ω -lim is 0. Let \star_n be a sequence of points in X. Then we can define *asymptotic cone* of X, denoted by X_{ω} , with respect to \star_n and λ_n to be the ultralimit of the sequence of pointed spaces (X, \star_n) equipped with a distance function defined as

$$d_n = \lambda_n d_X$$

A theorem of Kleiner-Leeb [16, Chapter 5] says that X_{ω} is a Euclidean building of the same rank and type as the original symmetric space X. Let $F = \omega - \lim_{n \to \infty} \lambda_n F_n$, where F_n are the maximal flats in X_n associated to the Cartan subalgebra. F is a maximal flat in X_{ω} based at e_{ω} , then we can define a "Cartan projection" κ_{ω} from X_{ω} to the tangent space of F based at e_{ω} where for all $x = (x_n), y = (y_n) \in X_{\omega}$,

$$\kappa_{\omega}(x^{-1}y) \coloneqq \omega \operatorname{-lim} \lambda_n \kappa(x_n^{-1}y_n)$$

With κ_{ω} we can analogously define a subset $U_{\omega} \subset X_{\omega}$ such that $x = (x_n) \in U_{\omega}$ if for all $\alpha \in \theta$

$$\alpha(\kappa_{\omega}(x)) = \omega - \lim_{n \to \infty} \alpha(\lambda_n \kappa(x_n)) > 0.$$

The following proposition shows that if we have a sequence of $(q, \chi_1) P_{\theta}$ -sublinear (k, χ_2) sublinear rays c_n such that $c_n(0) = \star_n$, then the rescaled path $\lambda_n c_n$ where

$$\lambda_n c_n(t) \coloneqq c_n(\lambda_n^{-1}t)$$

ultraconverge to a bilipschitz ray $c_{\omega} \coloneqq \omega - \lim_{n \to \infty} \lambda_n c_n$ in X_{ω} with some regularity.

Proposition 8.7. The ultralimit c_{ω} is a k-bilipschitz ray and satisfies the following: for all $0 \le s < r < \infty$, the geodesic segment $[c_{\omega}(s), c_{\omega}(r)]$ is q-regular, i.e. for all $\alpha \in \theta$

$$\alpha(\kappa_{\omega}(c_{\omega}(s)^{-1}c_{\omega}(r))) \ge q \cdot d_{\omega}(c_{\omega}(s), c_{\omega}(r)).$$

Proof. First we show c_{ω} is k-bilipschitz. Let $s_n < r_n \in [0, \infty)$ be such that

 $s = \omega - \lim \lambda_n s_n$ and $r = \omega - \lim \lambda_n r_n$,

then

$$d_{\omega}(c_{\omega}(s), c_{\omega}(r)) = \omega - \lim_{n} d_{n}(c_{n}(s_{n}), c_{n}(r_{n}))$$

and because each c_n is a (k, χ_2) -sublinear ray,

$$\lambda_n \cdot \left[\frac{1}{k}(r_n - s_n) - \chi_2(r_n)\right] \le d_n(c_n(s_n), c_n(r_n)) \le \lambda_n \cdot \left[k(r_n - s_n) + \chi_2(r_n)\right].$$

Since r > 0, $r_n \to \infty$. Then for arbitrary $\varepsilon > 0$, we can find n large such that

$$\lambda_n \cdot \left[\frac{1}{k}(r_n - s_n) - \varepsilon r_n\right] \le d_n(c_n(s_n), c_n(r_n)) \le \lambda_n \cdot \left[k(r_n - s_n) + \varepsilon r_n\right]$$

Taking the ultralimit of the inequalities above gives

$$\frac{1}{k}(r-s) - \varepsilon r \le d_{\omega}(c_{\omega}(s), c_{\omega}(r)) \le k(r-s) + \varepsilon r.$$

Since this equation hold for all ε , we get the desired bi-Lipschitz inequality.

The second assertion is equivalent to say that c_{ω} is a uniformly regular ray in X_{ω} in the sense of Kapovich–Leeb–Porti [13]. We rewrite the singular values to be

(8.1)
$$\alpha(\kappa_{\omega}(c_{\omega}(s)^{-1}c_{\omega}(r))) = \omega - \lim_{n} \alpha(\lambda_{n}\kappa(c_{n}(s_{n})^{-1}c_{n}(r_{n}))).$$

Since c_n are all (q, χ_1) P_{θ} -sublinearly Morse, we have for all $\alpha \in \theta$

$$\alpha(\kappa(c_n(s_n)^{-1}c_n(r_n))) \ge q \cdot d_X(c_n(s_n), c_n(r_n)) - \chi_1(d_X(e, c_n(r_n)))$$

Plug this into (8.1) gives

$$\alpha(\kappa_{\omega}(c_{\omega}(s)^{-1}c_{\omega}(r))) \geq \omega \lim_{n} \lambda_{n} \left[q \cdot d_{X}(c_{n}(s_{n}), c_{n}(r_{n})) - \chi_{1}(d_{X}(e, c_{n}(r_{n}))) \right]$$
$$\geq \omega \lim_{n} \lambda_{n} \left[q \cdot d_{X}(c_{n}(s_{n}), c_{n}(r_{n})) - \chi_{1}(qr_{n} + \chi_{2}(r_{n})) \right].$$

Again, since $r_n \to \infty$, we have

$$\alpha(\kappa_{\omega}(c_{\omega}(s)^{-1}c_{\omega}(r))) \ge \omega - \lim_{n} \lambda_{n}(q \cdot d_{X}(c_{n}(s_{n}), c_{n}(r_{n})) - \varepsilon r_{n})$$
$$= q \cdot d_{\omega}(c_{\omega}(s), c_{\omega}(r)) - \varepsilon r$$

for all $\varepsilon > 0$, hence the proposition follows.

8.3. sublinearly Morse Lemma. Before we can state our sublinearly Morse lemma for P_{θ} -sublinearly Morse sublinear rays in X, we need to show any P_{θ} -sublinearly Morse ray converges to a unique point in the flag manifold.

Let $c: [0, \infty) \to X$ be a P_{θ} -sublinearly Morse (k, χ_2) -sublinear ray. Adapting the proof of Theorem 5.1 by picking a sequence of points $c(t_n)$ with $t_0 = 0$ on c with

$$d_X(c(t_n), c(t_{n+1})) \le \chi_2(d_X(c(0), c(t_n)))$$
 and $|t_n - t_{n+1}| > C > 0$,

we can show that $U_{\theta}(c(0)^{-1}c(t_n))$ flag converges to a unique point $\zeta \in \mathcal{F}_{\theta}$. Similarly one can show that for all $s \in [t_n, t_{n+1}]$, $U_{\theta}(c(0)^{-1}c(s))$ is arbitrarily close to $U_{\theta}(c(0)^{-1}c(t_n))$ as n goes to infinity. Hence, $U_{\theta}(c(t)) \to \zeta$.

Theorem 8.8 (sublinearly Morse Lemma). Suppose c is a (q, χ_1) -P_{θ}-sublinearly Morse (k, χ_2) -sublinear ray, let $\zeta \in \mathcal{F}_{\theta}$ be the unique limit point of $U_{\theta}(c(t))$. Then there exists a sublinear function $\eta \coloneqq \eta(q, \chi_1, k, \chi_2)$ such that the image of c is contained in the η -sublinear neighborhood of $V \coloneqq V(c(0), \zeta)$ defined as

$$\mathcal{N}_{\eta}(V) \coloneqq \{ x \in X \mid d_X(x, V) \le \eta(d_X(c(0), x)) \}$$

We will prove the theorem by the following stronger statement.

Lemma 8.9. There exists a sublinear function $\eta' := \eta'(q, k, \chi_1, \chi_2)$ such that for all $a \in [0, \infty)$ we have for all $t \in [0, a]$

$$d_X(c(t), \diamondsuit(c(0), c(a))) \le \eta'(d_X(c(0), c(a))).$$

Proof. We prove this claim by contradiction. Assume that there exists a sequence of (q, χ_1) - P_{θ} -sublinearly Morse (k, χ_2) -sublinear rays $c_n : [0, \infty) \to X$ (note that c_n are not necessarily distinct) such that there exist $a_n \in [0, \infty)$ with

$$\max_{t \in [0,a_n]} \{ d_X(c_n(t), \diamondsuit(c_n(0), c_n(a_n))) \} = D_n$$

and $D_n \to \infty$ as $n \to \infty$, and there exist $t_n \in (0, a_n) =: I_n$ such that $c_n(t_n)$ realizes the maximum above, and

$$\liminf_{n \to \infty} \frac{D_n}{d_X(c_n(t_n), c_n(0))} = m > 0.$$

Because c_n are (k, χ_2) sublinear rays, we have

$$\liminf_{n \to \infty} \frac{D_n}{\frac{1}{k}t_n - \chi_2(t_n)} \ge \liminf_{n \to \infty} \frac{D_n}{d_X(c_n(t_n), c_n(0))} = m > 0,$$

and since $t_n \to \infty$, we can rewrite the left hand side and get

(8.2)
$$\liminf_{n \to \infty} \frac{D_n}{t_n} \ge k \cdot m > 0.$$

Without loss of generality, we can assume that for all n

$$c_n(0) = e \text{ and } \lim_{t \to \infty} U_\theta(c_n(t)) = \zeta \in \mathcal{F}_\theta.$$

This can be done because G acts transitively on X and \mathcal{F}_{θ} , and \mathcal{P}_{θ} -sublinearly Morseness and sublinear rays are preserved under this action. Let $\diamondsuit_n \coloneqq \diamondsuit(e, c_n(a_n))$, then by construction

$$c_n(I_n) \subset \mathcal{N}_{D_n}(\diamondsuit_n)$$
 but $c_n(I_n) \notin \mathcal{N}_{(1-\varepsilon)D_n}(\diamondsuit_n)$

for any $\varepsilon > 0$.

Also, for *n* large enough, $d_X(e, c_n(a_n)) \ge \frac{1}{k}a_n - \chi_2(a_n) \to \infty$. Then the geodesic segment $[e, c_n(a_n)]$ is in *U* since

(8.3)
$$\alpha(\kappa(c_n(a_n))) \ge q \cdot d_X(e, c_n(a_n)) - \chi_1(d_X(e, c_n(a_n))) \\ \ge (q - \varepsilon) \cdot d_X(e, c_n(a_n))$$

for all $\alpha \in \theta$ Note that this shows that $[e, c_n(a_n))$ are uniform regular in the sense of Kapovich–Leeb–Porti for *n* large. Then the pairs of flags

$$(U_{\theta}(c_n(a_n)), U_{\theta}(c_n(a_n)^{-1})) =: (\zeta_n^+, \zeta_n^-)$$

is well defined and opposite for n large. Each \diamond_n is embedded in the parallel set $P(\zeta_n^+, \zeta_n^-) = P_n$.

Pick $\lambda_n = D_n^{-1}$, we obtain a asymptotic cone X_{ω} . Then the paths c_n are rescaled to $D_n^{-1}c_n: D_n^{-1}I_n \to X_n$. Passing to the ultralimit we get a path $c_{\omega}: I_{\omega} \to X_{\omega}$ such that

$$c_{\omega}(s) \coloneqq (D_n^{-1}c_n(s_n))$$

for $s \coloneqq \omega - \lim_n D_n^{-1} s_n$, and $I_\omega \coloneqq [0, a_\omega] \cap \mathbb{R}$ with $a_\omega \coloneqq \omega - \lim_n D_n^{-1} a_n$. According to Equation (8.2), there exist $t_\omega < \infty$ in I_ω such that $t_\omega \coloneqq \omega - \lim_n D_n^{-1} t_n$.

The parallel sets P_n ultraconverge to a parallel set in X_{ω} , i.e.

$$P_{\omega} \coloneqq \omega - \lim_{n} D_n^{-1} P_n$$

exists [13, Lemma 3.81]. Since $\zeta_n^{\pm} = U_{\theta}(c_n(a_n)^{\pm})$ are opposite, let l_n be geodesic lines such that $l_n(\pm \infty) \in \zeta_n^{\pm}$. Note that l_n ultraconverge to a geodesic line $l_{\omega} \subset P_{\omega}$, and $l_{\omega}(\pm \infty) =: \zeta_{\omega}^{\pm}$ is a pair of opposite flags of type τ_{mod} that defines P_{ω} . Moreover,

$$\diamondsuit_{\omega} \coloneqq \omega - \lim_{n} D_n^{-1} \diamondsuit_n$$

is contained in P_{ω} as a convex closed subset. Note that \diamond_{ω} is not always a diamond in X_{ω} . By construction

(8.4)
$$c_{\omega} \in \mathcal{N}_1(\diamondsuit_{\omega}) \text{ but } c_{\omega} \notin \diamondsuit_{\omega}.$$

By Proposition 8.7, c_{ω} is a bilipschitz regular path in the definition of Kapovich–Leeb– Porti. Then Lemma 5.22 in [13] implies

 $c_{\omega} \subset \diamondsuit_{\omega}.$

We have a contradiction.

Proof of Theorem 8.8. Since $U_{\theta}(c(t))$ flag converges to ζ as $t \to \infty$, the truncated Weyl cones Hausdorff converge [13, Lemma 3.65], i.e. for all R > 0

$$d_{Haus}\left(V(e, U_{\theta}(c(t))) \cap \bar{B}_{R}(e), V(e, \zeta) \cap \bar{B}_{R}(e)\right) \to 0$$

as $t \to \infty$.

Let \bar{c} be the projection of c to the Weyl cone $V(e,\zeta)$, and for all $t \in [0,\infty)$ let c_t be the projection of c([0,t]) to $V(e,U_{\theta}(c(t)))$. Fix an arbitrary $s \in [0,\infty)$, let $R_s := \max\{d_X(c_t(s),e), d_X(\bar{c}(s),e)\}$. Then for $\varepsilon > 0$ we can find t_s large enough such that

$$d_{Haus}\left(V(e, U_{\theta}(c(t_s))) \cap B_{R_s}(e), V(e, \zeta) \cap B_{R_s}(e)\right) < \varepsilon.$$

Then

$$d_X(c(s), V(e, \zeta)) \le d_X(c(s), V(e, U_\theta(c(t_s)))) + \varepsilon.$$

Since $\diamond(e, c(t_s)) \in V(e, U_{\theta}(c(t_s)))$, Lemma 8.9 implies

$$d_X(c(s), V(e, \zeta)) \le \eta'(d_X(e, c(s))) + \varepsilon.$$

Because we chose s arbitrarily, and $\eta(\cdot) \coloneqq \eta'(\cdot) + \varepsilon$ is again a sublinear function, we then have for all $s \in [0, \infty)$

$$c(s) \in \mathcal{N}_{\eta}(V(e,\zeta))$$

and this concludes the proof of Theorem 8.8.

Corollary 8.10. Let $\{g_n\} \subset G$ be a P_{θ} -sublinearly Morse sequence, and let $\zeta = \lim_{n \to \infty} U_{\theta}(g_n) \in \mathcal{F}_{\theta}$ be its limit point. Then there exist a sublinear function τ such that for all n,

$$d_X(g_n, V(g_0, \zeta)) \le \tau(d_X(g_0, g_n)).$$

Proof. The corollary follows directly from Lemma 8.2 and Theorem 8.8.

References

- Pierre-Louis Blayac, Richard Canary, Feng Zhu, and Andrew Zimmer. Patterson-Sullivan theory for coarse cocycles, 2024.
- [2] Pierre-Louis Blayac and Feng Zhu. Ergodicity and equidistribution in Hilbert geometry. arXiv e-prints, page arXiv:2106.08079, June 2021.
- [3] Jairo Bochi, Rafael Potrie, and Andrés Sambarino. Anosov representations and dominated splittings. arXiv e-prints, page arXiv:1605.01742, May 2016.
- [4] B. H. Bowditch. Relatively hyperbolic groups. International Journal of Algebra and Computation, 22(03):1250016, 2012.
- [5] Harrison Bray. Ergodicity of Bowen-Margulis measure for the Benoist 3-manifolds. arXiv e-prints, page arXiv:1705.08519, May 2017.
- [6] Richard Canary, Tengren Zhang, and Andrew Zimmer. Entropy rigidity for cusped Hitchin representations. arXiv e-prints, page arXiv:2201.04859, January 2022.
- [7] Richard Canary, Tengren Zhang, and Andrew Zimmer. Patterson-Sullivan measures for transverse subgroups. arXiv e-prints, page arXiv:2304.11515, April 2023.
- [8] Richard Canary, Andrew Zimmer, and Tengren Zhang. Patterson-Sullivan measures for relatively Anosov groups. *arXiv e-prints*, page arXiv:2308.04023, August 2023.
- [9] Jeffrey Danciger, François Guéritaud, and Fanny Kassel. Convex cocompact actions in real projective geometry. arXiv e-prints, page arXiv:1704.08711, April 2017.
- [10] Ilya Gekhtman, Yulan Qing, and Kasra Rafi. Genericity of sublinearly Morse directions in CAT(0) spaces and the Teichmüller space. *arXiv e-prints*, page arXiv:2208.04778, August 2022.
- [11] Mitul Islam and Andrew Zimmer. A flat torus theorem for convex co-compact actions of projective linear groups. arXiv e-prints, page arXiv:1907.03277, July 2019.
- [12] Michael Kapovich and Bernhard Leeb. Discrete isometry groups of symmetric spaces. arXiv e-prints, page arXiv:1703.02160, March 2017.
- [13] Michael Kapovich, Bernhard Leeb, and Joan Porti. A Morse Lemma for quasigeodesics in symmetric spaces and euclidean buildings. *arXiv e-prints*, page arXiv:1411.4176, November 2014.
- [14] Michael Kapovich, Bernhard Leeb, and Joan Porti. Anosov subgroups: Dynamical and geometric characterizations. arXiv e-prints, page arXiv:1703.01647, March 2017.
- [15] Dongryul M. Kim and Hee Oh. Relatively Anosov groups: finiteness, measure of maximal entropy, and reparameterization. arXiv e-prints, page arXiv:2404.09745, April 2024.

- [16] Bruce Kleiner and Bernhard Leeb. Rigidity of quasi-isometries for symmetric spaces and Euclidean buildings. Publications Mathématiques de l'IHÉS, 86:115–197, 1997.
- [17] Gabriel Pallier. Large-scale sublinearly Lipschitz geometry of hyperbolic spaces. arXiv e-prints, page arXiv:1801.05163, January 2018.
- [18] Yulan Qing, Kasra Rafi, and Giulio Tiozzo. Sublinearly Morse Boundary I: CAT(0) Spaces. arXiv e-prints, page arXiv:1909.02096, September 2019.
- [19] Yulan Qing, Kasra Rafi, and Giulio Tiozzo. Sublinearly Morse Boundary II: Proper geodesic spaces. arXiv e-prints, page arXiv:2011.03481, November 2020.
- [20] Cormac Walsh. The horofunction boundary of the Hilbert geometry. arXiv Mathematics e-prints, page math/0611920, November 2006.
- [21] Feng Zhu and Andrew Zimmer. Relatively Anosov representations via flows I: theory. *arXiv e-prints*, page arXiv:2207.14737, July 2022.