Matt Visser[®]

School of Mathematics and Statistics, Victoria University of Wellington, PO Box 600, Wellington 6140, New Zealand.

E-mail: matt.visser@sms.vuw.ac.nz

Abstract:

From known effective bounds on the prime counting function of the form

$$|\pi(x) - \operatorname{Li}(x)| < a \ x \ (\ln x)^b \ \exp\left(-c \ \sqrt{\ln x}\right); \qquad (x \ge x_0);$$

it is possible to establish exponentially tight effective upper and lower bounds on the prime number theorem: For $x \ge x_*$ where $x_* \le \max\{x_0, 17\}$ we have:

$$\frac{\operatorname{Li}(x)}{1+a\;(\ln x)^{b+1}\;\exp\left(-c\;\sqrt{\ln x}\right)} < \pi(x) < \frac{\operatorname{Li}(x)}{1-a\;(\ln x)^{b+1}\;\exp\left(-c\;\sqrt{\ln x}\right)}$$

Furthermore, it is possible to establish exponentially tight effective upper and lower bounds on the location of the n^{th} prime. Specifically:

$$p_n < \text{Li}^{-1} \left(n \left[1 + a \, (\ln[n \ln n])^{b+1} \, \exp\left(-c \, \sqrt{\ln[n \ln n]} \right) \right] \right); \qquad (n \ge n_*).$$
$$p_n > \text{Li}^{-1} \left(n \left[1 - a \, (\ln[n \ln n])^{b+1} \, \exp\left(-c \, \sqrt{\ln[n \ln n]} \right) \right] \right); \qquad (n \ge n_*).$$

Here the range of validity is explicitly bounded by some n_* satisfying

$$n_* \le \max\left\{\pi(x_0), \pi(17), \pi\left((1+e^{-1})\exp\left(\left[\frac{2(b+1)}{c}\right]^2\right)\right)\right\}.$$

Many other fully explicit bounds along these lines can easily be developed.

DATE: 20 April 2025; LATEX-ed April 22, 2025

KEYWORDS: Prime counting function $\pi(x)$; n^{th} prime p_n ; effective bounds.

Contents

1	Introduction: Known bounds on $ \pi(x) - Li(x) $	1
2	Some elementary variants on these bounds	2
3	Exponentially bounding the prime number theorem	4
4	Localizing the n^{th} prime:	5
5	Special case	9
6	Further developments	10
7	Conclusions	10

1 Introduction: Known bounds on $|\pi(x) - \text{Li}(x)|$

Work over the last decade or so has developed a number of fully effective bounds on the prime counting function $\pi(x)$ of the general form [1–3]:

$$|\pi(x) - \operatorname{Li}(x)| < a \ x \ (\ln x)^b \ \exp\left(-c \ \sqrt{\ln x}\right); \qquad (x \ge x_0).$$
(1.1)

For some widely applicable effective bounds of this type see Table I below. (A straightforward elementary numerical computation is required to determine the numerical coefficients in the Trudgian [1] bounds.)

a	b	С	x_0	Source	Notes
0.4394	-3/4	0.32115	59	Trudgian [1]	Eq(7)
0.2795	-3/4	0.3936	229	Trudgian [1]	Th 2
9.2211	1/2	0.8476	2	Fiori–Kadiri–Swidinsky [3]	Eq (3) , Eq (43)
9.59	0.515	0.8274	2	Johnston–Yang [2]	Eq (1.6)

Table 1. Some widely applicable effective bounds on $\pi(x)$.

2 Some elementary variants on these bounds

There are various strategies for relaxing these bounds to make them more tractable. The most simple technique is to simply increase either of the coefficients $\{a, b\}$, or decrease the coefficient c, in the Trudgian bounds so as to increase the domain of validity (decrease x_0). (The FKS and JY bounds are already maximal in this regard.)

A more subtle technique (based on a variant of the discussion in reference [4]) is as follows: Let $\tilde{b} \leq b$ and $\tilde{c} < c$. Write

$$a x (\ln x)^{b} \exp\left(-c \sqrt{\ln x}\right)$$
$$= a x \left\{ (\ln x)^{b-\tilde{b}} \exp\left(-[c-\tilde{c}] \sqrt{\ln x}\right) \right\} (\ln x)^{\tilde{b}} \exp\left(-\tilde{c} \sqrt{\ln x}\right).$$
(2.1)

Then the quantity in braces is bounded, reaching a global maximum at

$$x_{max} = \exp\left(\frac{4[b-\tilde{b}]^2}{[c-\tilde{c}]^2}\right),\tag{2.2}$$

where it takes on the value

$$\left\{ (\ln x)^{b-\tilde{b}} \exp\left(-[c-\tilde{c}] \sqrt{\ln x}\right) \right\}_{max} = \left(\frac{4[b-\tilde{b}]^2}{[c-\tilde{c}]^2}\right)^{b-b} \exp\left(-2[b-\tilde{b}]\right) \quad (2.3)$$

Then defining

$$\tilde{a} = a \left(\frac{4[b-\tilde{b}]^2}{[c-\tilde{c}]^2}\right)^{b-\tilde{b}} \exp\left(-2[b-\tilde{b}]\right) = a \left(\frac{2[b-\tilde{b}]}{e[c-\tilde{c}]}\right)^{2(b-\tilde{b})}.$$
(2.4)

we see that

$$a x (\ln x)^b \exp\left(-c \sqrt{\ln x}\right) \le \tilde{a} x (\ln x)^{\tilde{b}} \exp\left(-\tilde{c} \sqrt{\ln x}\right); \qquad (x > 1); \qquad (2.5)$$

with equality only at $x = x_{max}$. Consequently any bound of the form

$$|\pi(x) - \operatorname{Li}(x)| < a \ x \ (\ln x)^b \ \exp\left(-c \ \sqrt{\ln x}\right); \qquad (x \ge x_0);$$
 (2.6)

implies bounds of the form

$$|\pi(x) - \operatorname{Li}(x)| < \tilde{a} \ x \ (\ln x)^{\tilde{b}} \ \exp\left(-\tilde{c} \ \sqrt{\ln x}\right); \qquad (x \ge x_0).$$

$$(2.7)$$

Here the coefficients $\{\tilde{a}, \tilde{b}, \tilde{c}\}$ are as defined above. (Both $\tilde{b} < b$ and $\tilde{c} < c$ are free to be chosen, while $\tilde{a} > a$ is to be derived.) As long as x_0 is not too large, one can use explicit computations to find a larger domain of validity (some smaller \tilde{x}_0).

Specifically there will be some $\tilde{x}_0 \leq x_0$ such that

$$|\pi(x) - \operatorname{Li}(x)| < \tilde{a} \ x \ (\ln x)^{\tilde{b}} \ \exp\left(-\tilde{c} \ \sqrt{\ln x}\right); \qquad (x \ge \tilde{x}_0; \ \tilde{x}_0 \le x_0).$$
 (2.8)

In previous work, see reference [4], I had focussed on the case $\tilde{b} = 0$ which is most appropriate to finding explicit bounds (for $\vartheta(x)$) of the de la Vallé Poussin form [5]. Herein we shall soon see that for technical reasons when considering $\pi(x)$ it is more useful to consider the case $\tilde{b} = -1$. Table II below presents a number of derived widely applicable effective bounds of this general type.

				()	1 (/	
a	b	С	x_0	ã	\tilde{b}	\tilde{c}	x_{max}	\tilde{x}_0
0.4394	-3/4	0.32115	59	0.4680	-7/8	1/4	203 931	41
0.4394	-3/4	0.32115	59	0.4795	-1	1/6	$35 \ 439$	41
0.2795	-3/4	0.3936	229	0.2804	-5/6	1/3	2 097	227
0.2795	-3/4	0.3936	229	0.3164	-1	1/4	184 165	223
9.2211	1/2	0.8476	2	9.7590	0	1/2	3930	2
9.2211	1/2	0.8476	2	11.9026	-1/2	1/5	$13\ 874$	2
9.2211	1/2	0.8476	2	29.6698	-1	1/10	9 849 130	2
9.59	0.515	0.8274	2	11.148	0	1/2	19 877	2
9.59	0.515	0.8274	2	13.659	-1/2	1/5	35 206	2
9.59	0.515	0.8274	2	34.955	-1	1/10	34 331 213	2

Table 2. Several derived widely applicable effective bounds on $\pi(x)$.

(Input parameters on the left; output parameters on the right.)

(Note the bounds are closest to each other in the vicinity of x_{max} .)

 $|\pi(x) - \operatorname{Li}(x)| < \tilde{a} \ x \ (\ln x)^{\tilde{b}} \ \exp\left(-\tilde{c} \ \sqrt{\ln x}\right).$

With Table II in hand, it is now possible to increase the parameter a to expand the range of validity. Several examples of this are given in Table III.

Table 3. More derived and very widely applicable effective bounds on $\pi(x)$.

(Here instead a is adjusted so as to maximize the region of validity, $x_0 = 2$.)

 $|\pi(x) - \operatorname{Li}(x)| < a \ x \ (\ln x)^b \ \exp\left(-c \sqrt{\ln x}\right).$

a	b	С	x_0
3/4	-3/4	0.32115	2
7/8	-7/8	0.32115	2
0.93	-1	0.32115	2
0.8935	-3/4	0.3936	2
0.94	-7/8	0.3936	2
1.05	-1	0.3936	2

We shall subsequently apply these bounds (both Tables II and III) in various ways — specifically to bounding the location of the n^{th} prime.

3 Exponentially bounding the prime number theorem

Note that for any $a > 0, b \ge -1, c > 0$, we can write:

$$a x (\ln x)^{b} \exp\left(-c \sqrt{\ln x}\right) = a \frac{x}{\ln x} (\ln x)^{b+1} \exp\left(-c \sqrt{\ln x}\right).$$
(3.1)

It is useful to define

$$f(x) = (\ln x)^{b+1} \exp\left(-c \sqrt{\ln x}\right).$$
 (3.2)

Note that $\frac{x}{\ln x} < \pi(x)$ for x > 17, so that

$$a x (\ln x)^b \exp\left(-c \sqrt{\ln x}\right) < a \pi(x) f(x); \qquad (x > 17).$$
 (3.3)

This is already somewhat interesting since it implies that the various bounds on $|\pi(x) - \text{Li}(x)|$ discussed above all lead to

$$|\pi(x) - \operatorname{Li}(x)| < a \ \pi(x) \ f(x); \qquad (x \ge \max\{x_0, 17\}). \tag{3.4}$$

When feasible, $(x_0 \text{ sufficiently small})$, direct computation might potentially increase the domain of validity

$$|\pi(x) - \operatorname{Li}(x)| < a \ \pi(x) \ f(x); \qquad (x \ge x_*; \ x_* \le \max\{x_0, 17\}). \tag{3.5}$$

This can be rearranged to yield

$$\frac{\text{Li}(x)}{1+a f(x)} < \pi(x) < \frac{\text{Li}(x)}{1-a f(x)}; \qquad (x \ge x_*).$$
(3.6)

Note that $f(x) \to 0$ with exponential rapidity in $\sqrt{\ln x}$.

To be more explicit, for $x \ge x_*$:

$$\frac{\text{Li}(x)}{1+a\ (\ln x)^{b+1}\ \exp\left(-c\ \sqrt{\ln x}\right)} < \pi(x) < \frac{\text{Li}(x)}{1-a\ (\ln x)^{b+1}\ \exp\left(-c\ \sqrt{\ln x}\right)}.$$
 (3.7)

These inequalities provide a remarkably clean and effective version of the prime number theorem.

4 Localizing the n^{th} prime:

Note that for $b \ge -1$ the function $f(x) = (\ln x)^{b+1} \exp\left(-c \sqrt{\ln x}\right)$ achieves a global maximum at $x_{peak} = \exp\left(\left[2(b+1)/c\right]^2\right) \ge 1$, above which f'(x) < 0, and below which f'(x) > 0.

Concentrating on the region above the peak we shall first re-write f(x) in terms of $p_{\pi(x)}$, yielding a bound in terms of $f(p_{\pi(x)})$. We will then convert this into an explicit function of $\pi(x)$, yielding a bound in terms of $f(\pi(x) \ln \pi(x))$. Doing so will require some careful attention to the domain of validity of the various inequalities. (One wishes the various manipulations to bring one closer to the peak without overshooting the peak.)

Observe that in all generality $p_{\pi(x)} \leq x < p_{\pi(x)+1}$, and more specifically one has $p_{\pi(x_{peak})} \leq x_{peak} < p_{\pi(x_{peak})+1}$ So if we now choose $x > p_{\pi(x_{peak})+1} > x_{peak}$ then we can safely write

$$f(x) < f(p_{\pi(x)});$$
 $(x > p_{\pi(x_{peak})+1}).$ (4.1)

In 1952 Nagura [6] (among other results) showed

$$\vartheta\left(\frac{4}{3}x\right) - \vartheta(x) > 0; \qquad (x \ge 109 = p_{29}). \tag{4.2}$$

This bound is by no means optimal, but it will be good enough for current purposes. This implies

$$p_{n+1} < \frac{4}{3} p_n; \qquad (n \ge 29).$$
 (4.3)

Explicitly checking smaller integers yields

$$p_{n+1} < \frac{4}{3} p_n; \qquad (n \ge 5 = \pi(11)).$$
 (4.4)

But then let us choose $x > \frac{4}{3} x_{peak}$ and note

$$x > \frac{4}{3}x_{peak} \ge \frac{4}{3}p_{\pi(x_{peak})} > p_{\pi(x_{peak})+1}$$
(4.5)

Then certainly

$$f(x) < f(p_{\pi(x)});$$
 $\left(x > \max\left\{11, \frac{4}{3}x_{peak}\right\}\right).$ (4.6)

Now invoke Rosser's theorem $p_n > n \ln n$, $(n \ge 1)$ [7], then

$$f(x) < f(\pi(x) \ln \pi(x));$$
 (4.7)

where in addition to the prior constraints $x > \max\{11, \frac{4}{3}x_{peak}\}$ we now need to check that $\pi(x)\ln(\pi(x)) > x_{peak}$.

To get a more computationally useful grasp on the domain of validity use the fact that $p_n < n \ln(n \ln n)$ for $n \ge 6$ [8] and note

$$p_n < n \ln(n \ln n) = n(\ln n + \ln \ln n) = n \ln n \left\{ 1 + \frac{\ln \ln n}{\ln n} \right\}; \quad (n \ge 6). \quad (4.8)$$

The quantity in braces is bounded and takes on the maximum value $1+e^{-1}$ at $x=e^{e}$. Thence

$$p_n < (1 + e^{-1}) \ n \ln n = 1.367879441 \ n \ln n; \qquad (n \ge 4).$$
 (4.9)

So we see

$$\pi(x)\ln(\pi(x)) > \frac{p_{\pi(x)}}{1+e^{-1}}; \qquad (x \ge p_4 = 7)$$
(4.10)

This implies that the condition $\pi(x) \ln(\pi(x)) > x_{peak}$ relevant to inequality (4.7) is certainly satisfied for $p_{\pi(x)} > (1 + e^{-1})x_{peak}$, which in turn is certainly satisfied for $x > (1 + e^{-1})x_{peak} > \frac{4}{3}x_{peak}$, so that we can safely write

$$f(x) < f(\pi(x) \ln \pi(x));$$
 (x > max{11, (1 + e⁻¹) x_{peak}}). (4.11)

Finally, combine this result with the conditions for the validity of the bound (3.4), to yield

$$|\pi(x) - \operatorname{Li}(x)| < a \ \pi(x) \ f(\pi(x) \ln \pi(x)); \qquad (x \ge \max\{x_0, 17, (1 + e^{-1})x_{peak}\}).$$
(4.12)

We remind the reader that for current purposes we are interested in

$$x_{peak} = \exp\left(\left[\frac{2(b+1)}{c}\right]^2\right) \ge 1.$$
(4.13)

Because we have made various approximations getting to this stage, the actual range of the validity of the final inequality (4.12) may be somewhat larger than naively advertised max{ $x_0, 17, (1 + e^{-1})x_{peak}$ }. That is, while the bounds (4.12) are certainly guaranteed to hold for values of x sufficiently far above the peak, they might still hold both at the peak and for some region below the peak.

Whenever feasible (meaning whenever x_{peak} is not too large) this should be checked by explicit computation to find a suitable x_* such that:

$$|\pi(x) - \operatorname{Li}(x)| < a \ \pi(x) \ f(\pi(x) \ln \pi(x)); \qquad (x \ge x_*; \ x_* \le \max\{x_0, 17, (1 + e^{-1})x_{peak}\})$$
(4.14)

Sometimes x_* is significantly lower than the naive bound $\max\{x_0, 17, (1+e^{-1})x_{peak}\}$, and sometimes $(1+e^{-1})x_{peak} \leq 17$ and so that the peak can be safely ignored. But sometimes we simply need to keep track of this complication. If we evaluate this final inequality (4.14) at $x = p_n$ then

$$|\mathrm{Li}(p_n) - n| < a \ n \ f(n \ln n); \qquad (n \ge n_*; \ n_* \le \max\{\pi(x_0), 7, \pi((1 + e^{-1}) \ x_{peak})\}).$$
(4.15)

Finally, as promised in the abstract,

$$\operatorname{Li}^{-1}(n[1 - a \ f(n \ln n)]) < p_n < \operatorname{Li}^{-1}(n[1 + a \ f(n \ln n)]); \qquad (n \ge n_*).$$
(4.16)

To be more explicit

$$p_n < \text{Li}^{-1} \left(n \left[1 + a \, (\ln[n \ln n])^{b+1} \, \exp\left(-c \, \sqrt{\ln[n \ln n]}\right) \right] \right); \qquad (n \ge n_*);$$
$$p_n > \text{Li}^{-1} \left(n \left[1 - a \, (\ln[n \ln n])^{b+1} \, \exp\left(-c \, \sqrt{\ln[n \ln n]}\right) \right] \right); \qquad (n \ge n_*).$$

Here the range of validity is explicitly bounded by some n_* with

$$n_* \le \max\left\{\pi(x_0), 7, \pi\left((1+e^{-1})\exp\left(\left[\frac{2(b+1)}{c}\right]^2\right)\right)\right\}.$$

These effective bounds can be viewed as fully explicit versions of the usual Cippola asymptotic expansion — see for instance the discussion in reference [9].

See Table IV below (based on Tables I, II, and III) for some explicit examples. The outputs from Table II, $\{\tilde{a}, \tilde{b}, \tilde{c}, \tilde{x}_0\}$, have been relabelled as $\{a, b, c, x_0\}$, and used as inputs for new computations of x_* and n_* for the new inequalities discussed in this section.

a	b	С	x_0	x_{peak}	x_*	n_0	n_{peak}	n_*	
0.4394	-3/4	0.32115	59	11.3	41	17	5	13	
0.4680	-7/8	1/4	41	e	37	13	1	12	
0.4795	-1	1/6	41	1	37	13	0	12	
0.2795	-3/4	0.3936	229	5.022	149	50	3	35	
0.2804	-5/6	1/3	227	e	149	49	1	35	
0.3164	-1	1/4	223	1	97	48	0	25	
9.2211	1/2	0.8476	2	275 789	2	1	24 104	1	
9.7590	0	1/2	2	e^{16}	2	1	$595 \ 341$	1	
11.9026	-1/2	1/5	2	e^{25}	*	1	$\frac{1}{25}e^{25}$	*	
29.6698	-1	1/10	2	1	2	1	0	1	
9.59	0.515	0.8274	2	$667\ 161$	2	1	$54\ 105$	1	
11.148	0	1/2	2	e^{16}	2	1	$595\ 431$	1	
13.659	-1/2	1/5	2	e^{25}	*	1	$\frac{1}{25}e^{25}$	*	
34.955	-1	1/10	2	1	2	1	0	1	

Table 4. Some widely applicable effective bounds on $|\operatorname{Li}(x) - \pi(x)|$ and p_n : $|\operatorname{Li}(x) - \pi(x)| < a \pi(x) \quad (\ln[\pi(x) \ln \pi(x)])^{b+1} \exp\left(-c \sqrt{\ln[\pi(x) \ln \pi(x)]}\right);$ $p_n \leq \operatorname{Li}^{-1}\left(n \left[1 \pm a \ (\ln[n \ln n])^{b+1} \ \exp\left(-c \sqrt{\ln[n \ln n]}\right)\right]\right).$

Explicit evaluation of x_* and n_* in the situations labelled * proved computationally infeasible due to the magnitude of x_{peak} and n_{peak} . All indications are that $x_* = 2$ and $n_* = 1$ in both cases.

5 Special case

As previously indicated, the special case b = -1 is particularly appealing.

Starting from any bound of the form

$$|\pi(x) - \operatorname{Li}(x)| < a \ x \ (\ln x)^{-1} \ \exp\left(-c \ \sqrt{\ln x}\right); \qquad (x \ge x_0);$$
 (5.1)

we have the particularly simple result that:

$$|\pi(x) - \operatorname{Li}(x)| < a \ \pi(x) \ \exp\left(-c \ \sqrt{\ln x}\right); \qquad (x \ge x_*; \ x_* \le \max\{x_0, 17\}).$$
 (5.2)

Thence in the range $x \ge x_*$ we have

$$\frac{\operatorname{Li}(x)}{1+a \, \exp\left(-c \,\sqrt{\ln x}\right)} < \pi(x) < \frac{\operatorname{Li}(x)}{1-a \, \exp\left(-c \,\sqrt{\ln x}\right)}.$$
(5.3)

Furthermore, (noting that in this case we always have $x_{peak} = 1$),

$$p_n < \operatorname{Li}^{-1}\left(n\left[1 + a\exp\left(-c\sqrt{\ln[n\ln n]}\right)\right]\right); \qquad (n \ge n_*); \tag{5.4}$$

$$p_n > \operatorname{Li}^{-1}\left(n\left[1 - a\exp\left(-c\sqrt{\ln[n\ln n]}\right)\right]\right); \quad (n \ge n_*),$$
 (5.5)

subject to the simple constraint

$$n_* \le \max\{\pi(x_0), 7\}.$$
 (5.6)

For several explicit effective examples of this specific behaviour see Table V (which is a subset of Table IV and a minor extension of parts of Table III). These bounds are not necessarily optimal but, (given the initial input information in Table I), they are relatively simple to derive.

Table 5. Widely applicable effective bounds on $|\operatorname{Li}(x) - \pi(x)|$ and p_n for b = -1: $|\operatorname{Li}(x) - \pi(x)| < a \pi(x) \exp\left(-c \sqrt{\ln[\pi(x) \ln \pi(x)]}\right);$ $p_n \leq \operatorname{Li}^{-1}\left(n \left[1 \pm a \exp\left(-c \sqrt{\ln[n \ln n]}\right)\right]\right).$

	• /			(*	L	·'/.]/	
a	b	С	x_0	x_{peak}	x_*	n_0	n_{peak}	n_*
0.4795	-1	1/6	41	1	37	13	0	12
0.3164	-1	1/4	223	1	97	48	0	25
29.6698	-1	1/10	2	1	2	1	0	1
34.955	-1	1/10	2	1	2	1	0	1
1	-1	0.32115	2	1	2	1	0	1
1.1	-1	0.3936	2	1	2	1	0	1

6 Further developments

Another useful trick is to evaluate the inequality

$$|\pi(x) - \text{Li}(x)| < a \ \pi(x) \ f(\pi(x) \ln \pi(x)); \qquad (x \ge x_*)$$
(6.1)

both at $x = p_n$ and at $x = p_{n+1}^-$, just below p_{n+1} . Then since $\pi(p_{n+1}^-) = n = \pi(p_n)$ we have both

$$|n - \operatorname{Li}(p_n)| < a \ n \ f(n \ln n); \qquad (n \ge n_*); \tag{6.2}$$

and

$$|n - \operatorname{Li}(p_{n+1})| < a \ n \ f(n \ln n); \qquad (n \ge n_*).$$
 (6.3)

But then by the triangle inequality

$$\operatorname{Li}(p_{n+1}) - \operatorname{Li}(p_n) < 2a \ n \ f(n \ln n); \qquad (n \ge n_*);$$
 (6.4)

Implying

$$p_{n+1} < \operatorname{Li}^{-1} \left(\operatorname{Li}(p_n) + 2a \ n \ f(n \ln n) \right); \qquad (n \ge n_*).$$
 (6.5)

This observation can be used to develop yet more asymptotic bounds along the lines of the Cippola expansion, as discussed in [9], and can also be related to the effective bounds developed using the first Chebyshev $\vartheta(x)$ function as described in reference [10].

7 Conclusions

With some hindsight, an ineffective version of the bounds considered herein could have been developed some 125 years ago. The current analysis is interesting in two respects: First, using developments made over the past decade, the bounds have been made explicitly and fully effective. Secondly, because the bounds are asymptotically exponential, they will eventually always overtake and outperform bounds based on the Cippola expansion. Finally, I do not claim any of these bounds are in any way optimal. As always there is a three-way trade-off between stringency of the bound, simplicity of the bound, and the region of validity. Herein I have largely focussed on simplicity of the bound and a wide (sometimes maximal) region of validity.

References

- [1] Tim Trudgian, "Updating the error term in the prime number theorem", The Ramanujan Journal **39** (2016) 225–236. doi:10.1007/S11139-014-9656-6.
 [arXiv: 1401.2689 [math.NT]]
- [2] Daniel R. Johnston, Andrew Yang,
 "Some explicit estimates for the error term in the prime number theorem", Journal of Mathematical Analysis and Applications 527 # 2 (2023), 127460. doi:10.1016/j.jmaa.2023.127460 [arXiv: 2204.01980 [math.NT]]
- [3] Andrew Fiori, Habiba Kadiri, Joshua Swindisky,
 "Sharper bounds for the error term in the Prime Number Theorem", Research in Number Theory 9 (2023) 63. doi:10.1007/s40993-023-00454-w
 [arXiv: 2206.12557 [math.NT]]
- [4] Matt Visser,
 "Effective de la Vallé Poussin style bounds on the first Chebyshev function", International Mathematical Forum 18 # 4 (2023) 133–139 doi:10.12988/imf.2023.912416 [arXiv: 2211.00840 [math.NT]]
- [5] Charles Jean de la Vallé Poussin,
 "Recherches analytiques sur la théorie des nombres premiers",
 Ann. Soc. Scient. Bruxelles, deuxiéme partie 20, (1896), pp. 183–256
- [6] Jitsuro Nagura, "On The Interval Containing At Least One Prime Number", Proc. Japan Acad. 28 # 4 (1952) 177-181. doi:10.3792/pja/1195570997
- [7] Rosser, J. B. "The *n*-th Prime is Greater than $n \log n$ ". Proceedings of the London Mathematical Society **45** (1939) 21-44, doi:10.1112/plms/s2-45.1.21
- [8] Rosser, J. B. "Explicit bounds for some functions of prime numbers". Amer. J. Math. 63 (1941), 211-232. doi:10.2307/2371291 https://www.jstor.org/stable/237129
- [9] Juan Arias de Reyna, Toulisse Jeremy,
 "The n-th prime asymptotically", J. Théor. Nombres Bordeaux 25 (2013) 521–555, doi:10.5802/jtnb.847 arXiv: 1203.5413 [math.NT]
- [10] Matt Visser,"Effective exponential bounds on the prime gaps", arXiv: 2211.06469 [math.NT]