

ON THE TURÁN NUMBER OF THE $G_{3 \times 3}$ IN LINEAR HYPERGRAPHS

JÓZSEF SOLYMOSI

ABSTRACT. We show a construction for dense 3-uniform linear hypergraphs without 3×3 grids, improving the lower bound on its Turán number.

1. INTRODUCTION

Finding the Turán number of hypergraphs is challenging. The typical question asks for the maximum number of edges that can be avoided in a given hypergraph. There are a few examples only when sharp bounds are known. Such questions are even more challenging in an important subfamily called *linear hypergraphs*. In an r -uniform linear hypergraph, every hyperedge has r vertices, and any pair of edges have at most one common vertex. An r -uniform linear hypergraph on r^2 vertices is called an r by r grid if it is isomorphic to a pattern of r horizontal and r vertical lines.

Answering a question by Füredi and Ruszinkó [2] about hypergraphs avoiding 3×3 grids (denoted by $G_{3 \times 3}$), Gishboliner and Shapira gave a construction for dense linear 3-uniform hypergraphs not containing $G_{3 \times 3}$.




FIGURE 1. A $G_{3 \times 3}$ hypergraph on nine vertices with six edges

Extremal problems about $G_{3 \times 3}$ are important due to the connection to the Brown-Erdős-Sós conjecture, coding theory and geometric application (see examples in [2] [1] and [3]).

Theorem 1.1 (Gishboliner and Shapira [1]). *For infinitely many n , there exists a linear $G_{3 \times 3}$ -free 3-uniform hypergraph with n vertices and $(\frac{1}{16} - o(1))n^2$ edges.*

Füredi and Ruszinkó conjectured that there are arbitrarily large Steiner triple systems avoiding $r \times r$ grids for any $r \geq 3$. We are unsure about this conjecture (particularly the $r = 3$ case), but at least we improve the Gishboliner-Sharipa bound.

2. THE NEW BOUND

As in many extremal constructions, we are using objects in finite geometries.

Theorem 1. *For infinitely many n , there exists a linear $G_{3 \times 3}$ -free 3-uniform hypergraph with n vertices and $(\frac{1}{12} - o(1)) n^2$ edges.*

Proof. First, we reprove the Gishboliner-Shapira bound with the constant $\frac{1}{16}$ in a different way. Let q be a prime power, let \mathbb{F}_q be the finite field of order q , and let $AG(2, q)$ denote the Desarguesian affine plane of order q .

The vertex set of the hypergraph \mathcal{H} consists of the points of two parabolas, $V_1 = \{(x, x^2) : x \in \mathbb{F}_q\}$ and $V_2 = \{(x, x^2 + 1) : x \in \mathbb{F}_q\}$. The edge set is defined by collinear triples on the lines of the secants of V_1 that have a non-empty intersection with V_2 . If the line has two intersection points in V_2 , choose only one of them to form an edge. Let's count the number of edges.

There are q points in V_1 , so the number of its secants is $\binom{q}{2}$. For two distinct points (a, a^2) and (b, b^2) the equation of the line connecting them is $y = (a + b)x - ab$. It intersects V_2 iff the discriminant $(a - b)^2 - 4$ is a quadratic residue or zero (we assumed here that q is odd, although one could perform similar calculations for even q as well). It is quadratic residue or zero for $q(q - \chi(-1))/2$ ordered (a, b) , $a \neq b$ pairs, where χ is the quadratic character of -1 in \mathbb{F}_q . The number of intersecting secants, and therefore the number of edges is

$$E(\mathcal{H}) = \frac{q(q - \chi(-1))}{4} \approx \frac{q^2}{4} = \left(\frac{|V(\mathcal{H})|}{2}\right)^2 \frac{1}{4} = \frac{1}{16} |V(\mathcal{H})|^2$$

for large q .

This 3-uniform linear hypergraph doesn't contain 3×3 grids. If there were one in \mathcal{H} , then six of its vertices were in V_1 , so by Pappus' theorem*, the three vertices in V_2 were collinear. On the other hand, the parabola, V_2 , has no collinear triples.

Most of the intersecting lines of the secants of V_1 intersect V_2 in two points, and we selected only one of them. Now we select a random subset, hoping that many lines will still intersect at a point as the number of points reduces. Let's select points of V_2 independently at random with probability p into a set S . The expected size of S is pq . The number of secants of V_1 with two intersection points in V_2 is $\frac{q(q - \chi(-1) - 4)}{4}$, since we have to discard the tangents, the cases when the discriminant is zero, i.e. when $a - b = \pm 2$. We are looking for an asymptotic bound, so we will continue counting with $\approx q^2/4$ such lines. The expected number of lines with at least one point in S is $(2p - p^2)q^2/4$. Select p to maximize

$$\frac{(2p - p^2)q^2/4}{(q + pq)^2} = \frac{2p - p^2}{4(1 + p)^2}.$$

The maximum is achieved when $p = 1/2$. Then the ratio is $1/12$ as required. As the final step, we select the edges of the random subgraph of \mathcal{H} . If the line of the secant has four points, choose one of the two from S . The secants with one point in S determine a unique edge. Standard probability estimates (like Chernoff's bounds and the union bound) show that there are sets $S \in V_2$ that provide values close to the expected values. □

*Pappus' theorem holds over finite fields (or more precisely over projective geometries).

3. ACKNOWLEDGEMENTS

The research was supported in part by an NSERC Discovery grant and by the National Research Development and Innovation Office of Hungary, NKFIH, Grant No. KKP133819 and Excellence 151341.

REFERENCES

- [1] L. Gishboliner and A. Shapira, *Constructing dense 3-grid-free linear graphs*, Proceedings of the American Mathematical Society **150** (2022), 69–74. <https://doi.org/10.1090/proc/15086>
- [2] Z. Füredi and M. Ruszinkó, *Uniform hypergraphs containing no grids*, Advances in Mathematics **240** (2013), 302–324. <https://doi.org/10.1016/j.aim.2013.01.005>
- [3] J. Solymosi, *On the Structure of Pointsets with Many Collinear Triples*, Discrete & Computational Geometry **72** (2024), 986–1009. <https://doi.org/10.1007/s00454-023-00579-w>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC, CANADA,
AND OBUDA UNIVERSITY, BUDAPEST, HUNGARY
Email address: solymosi@math.ubc.ca